# VAROPOULOS EXTENSIONS IN DOMAINS WITH AHLFORS-REGULAR BOUNDARIES AND APPLICATIONS TO BOUNDARY VALUE PROBLEMS FOR ELLIPTIC SYSTEMS WITH  $L^{\infty}$  COEFFICIENTS

MIHALIS MOURGOGLOU AND THANASIS ZACHAROPOULOS

ABSTRACT. Let  $\Omega \subset \mathbb{R}^{n+1}$ ,  $n \geq 1$ , be an open set with s-Ahlfors regular boundary  $\partial \Omega$ , for some  $s \in (0, n]$ , such that either  $s = n$  and  $\Omega$  is a corkscrew domain with the pointwise John condition, or  $s < n$  and  $\Omega = \mathbb{R}^{n+1} \setminus E$ , for some s-Ahlfors regular set  $E \subset \mathbb{R}^{n+1}$ . In this paper we provide a unifying method to construct Varopoulos type extensions of BMO and  $L^p$  boundary functions. In particular, we show that a) if  $f \in \text{BMO}(\partial \Omega)$ , there exists  $F \in C^{\infty}(\Omega)$  such that  $dist(x, \Omega^c)|\nabla F(x)|$  is uniformly bounded in  $\Omega$  and the Carleson functional of dist $(x, \Omega^c)^{s-n} |\nabla F(x)|$  as well the sharp non-tangential maximal function of F are uniformly bounded on  $\partial\Omega$  with norms controlled by the BMO-norm of f, and  $F \to f$  in a certain non-tangential sense  $\mathcal{H}^s|_{\partial\Omega}$ -almost everywhere; b) if  $\bar{f} \in L^p(\partial\Omega)$ ,  $1 < p \leq \infty$ , there exists  $\overline{F} \in C^{\infty}(\Omega)$  such that the non-tangential maximal functions of  $\bar{F}$  and dist $(\cdot, \Omega^c) |\nabla \bar{F}|$  as well as the Carleson functional of dist $(\cdot, \Omega^c)^{s-n} |\nabla \bar{F}|$  are in  $L^p(\partial\Omega)$  with norms controlled by the  $L^p$ -norm of  $\bar{f}$ , and  $\bar{F} \to \bar{f}$  in some non-tangential sense  $\mathcal{H}^{s}$  |∂Ω-almost everywhere. If, in addition, the boundary function is Lipschitz with compact support, then both  $F$  and  $F$  can be constructed so that they are also Lipschitz on  $\overline{\Omega}$  and converge to the boundary data continuously. The latter results hold without the additional assumption of the pointwise John condition. Finally, for elliptic systems of equations in divergence form with merely bounded complex-valued coefficients, we show some connections between the solvability of Poisson problems with interior data in the appropriate Carleson or tent spaces and the solvability of Dirichlet problem with  $L^p$  and BMO boundary data.

### **CONTENTS**



*Key words and phrases.* Extensions, trace, Carleson spaces, Tent spaces, Non-tangential maximal function, BMO, Campanato space, Boundary Value Problems in rough domains, Poisson Problems, divergence form elliptic PDEs, elliptic systems, complex coefficients.

M.M. was supported by IKERBASQUE and partially supported by the grant PID2020-118986GB-I00 of the Ministerio de Economía y Competitividad (Spain) and by the grant IT-1615-22 (Basque Government). T.Z. was supported by the FPI grant PRE2018-084984 of the Ministerio de Ciencia e Innovación (Spain) and partially supported by the grant PID2020-118986GB-I00 of the Ministerio de Economía y Competitividad (Spain).

#### 2 MIHALIS MOURGOGLOU AND THANASIS ZACHAROPOULOS



### 1. INTRODUCTION

<span id="page-1-0"></span>In the present manuscript we are concerned with open sets  $\Omega \subset \mathbb{R}^{n+1}$ ,  $n \geq 1$ , which satisfy one of the following assumptions:

- (a)  $\Omega$  satisfies the corkscrew condition and its boundary  $\partial\Omega$  is n-Ahlfors regular (see Definitions [2.1](#page-9-3) and [2.10\)](#page-17-1), or
- (b)  $\Omega = \mathbb{R}^{n+1} \setminus E$ , for some s-Ahlfors regular set  $E \subset \mathbb{R}^{n+1}$  with  $s < n$ .

We will call such domains  $AR(s)$  *domains* for  $s \in (0, n]$ . We also define  $\sigma_s := \mathcal{H}^s|_{\partial\Omega}$  to be the "surface" measure of  $\Omega$ , where  $\mathcal{H}^s$  is the s-dimensional Hausdorff measure.

Our first goal is to construct, in AR(s) domains, smooth extensions  $u : \Omega \to \mathbb{R}$  of boundary functions that are in BMO( $\sigma_s$ ) (resp. in  $L^p(\sigma_s)$  for  $p \in (1,\infty]$ ) so that their sharp nontangential maximal function defined in  $(2.10)$  (resp. their non-tangential maximal functions defined in  $(2.9)$ ) and the modified Carleson functionals (see  $(2.12)$ ) for the definition) of their "weighted" gradients are uniformly bounded (resp. in  $L^p(\sigma_s)$ ) with norms controlled by the BMO( $\sigma_s$ ) (resp.  $L^p(\sigma_s)$ ) norms of the boundary functions. The identification on the boundary is in the non-tangential convergence sense (up to a set of measure zero on the boundary). To do so, when  $s = n$ , we assume that  $\Omega$  satisfies the *pointwise John condition* (see Definition [2.13\)](#page-18-0), while no additional connectivity assumption is required for  $s < n$ . This is the first time that such results are proved in so general geometric setting and also when the co-dimension is larger than 1.

Our second goal is to construct such extensions of Lipschitz functions with compact support on the boundary of an AR(s) domain so that they are Lipschitz on  $\overline{\Omega}$  and in the weighted Sobolev space  $\dot{W}^{1,2}(\Omega;\omega_s)$  $\dot{W}^{1,2}(\Omega;\omega_s)$  $\dot{W}^{1,2}(\Omega;\omega_s)$ , where  $\omega_s(x) := \delta_{\Omega}(x)^{s-n}$ . In fact, the construction of those extensions is even more important due to their applications to Boundary Value Problems given in Section [8](#page-53-0) and also in [\[GalMT23,](#page-69-1) [MP24\]](#page-70-0). Finally, we also prove similar extensions of boundary functions in the Campanato space  $\Lambda_{\beta}(\partial\Omega)$  for  $\beta \in (0,1)$ .

The original result of Varopoulos [\[Var77,](#page-70-1) [Var78\]](#page-70-2) entails an extension property for BMO functions in  $\mathbb{R}^n$ . Specifically, for  $f \in BMO(\mathbb{R}^n)$ , there exists an extension F defined in the

<span id="page-1-1"></span><sup>&</sup>lt;sup>1</sup>When  $s = n$ , this is the standard homogeneous Sobolev space  $\dot{W}^{1,2}(\Omega)$ .

upper half-space  $\mathbb{R}^{n+1} := \{(x,t) \in \mathbb{R}^n \times (0,\infty)\}\)$ , such that  $F(x,t) = f$  non-tangentially, with  $F \in C^{\infty}(\mathbb{R}^{n+1})$ , satisfying a size condition given by

<span id="page-2-1"></span>(1.1) 
$$
\sup_{(x,t)\in\mathbb{R}^{n+1}_+} t |\nabla F(x,t)| < \infty,
$$

and a  $L^1$  Carleson measure condition

<span id="page-2-0"></span>(1.2) 
$$
\sup_{r>0, x\in\mathbb{R}^n} r^{-n} \iint_{\{|x-y|
$$

where the implicit constants are purely dimensional. It is crucial to note that the Carleson condition [\(1.2\)](#page-2-0) involves an  $L^1$  bound, in contrast to the more standard  $L^2$  estimate (with respect to the weighted measure  $tdydt$ :

<span id="page-2-2"></span>
$$
(1.3) \qquad \sup_{r>0, x \in \mathbb{R}^n} r^{-n} \iint_{\{|x-y| < r\} \cap \mathbb{R}^{n+1}_+} |\nabla F(y,t)|^2 \, dy \, t dt \le C \|f\|_{\text{BMO}}^2.
$$

In the presence of the size condition  $(1.1)$ , it is evident that  $(1.2)$  implies  $(1.3)$ . However, the former is strictly stronger. An example illustrating this distinction is provided by Garnett in  $[Gar81]$ . In this example,  $(1.3)$  holds, but  $(1.2)$  fails. Garnett's example involves the standard Poisson extension of a suitably constructed (bounded) function  $f$ . In this case, Fefferman and Stein [\[FS72\]](#page-69-3) established  $(1.3)$  for the Poisson extension of f, and this bound played a crucial role in his proof of the  $H^1$ -BMO duality.

Varopoulos initially sought to establish his extension result with the aim of extending Carleson's Corona Theorem [\[Car82\]](#page-69-4) to  $\mathbb{C}^n$  with  $n \geq 2$ . While this particular endeavor did not succeed, the outcome, known as the "Varopoulos extension", has proven to be of significant interest for other reasons. Notably, employing  $(1.2)$  instead of  $(1.3)$  allows for a simplification in one of the key steps of Fefferman's duality theorem proof.

Varopoulos based his extension theorem on a deep property of harmonic functions (or solutions of divergence form elliptic equations more broadly), now termed ε-*approximability*. This property asserts that for any bounded harmonic function  $u$  (initially defined in the half-space but extendable to more general settings), normalized so that  $||u||_{\infty} \leq 1$ , and for a fixed  $\varepsilon > 0$ , there exists an " $\varepsilon$ -approximator"  $\varphi_{\varepsilon} \in C^{\infty}(\mathbb{R}^{n+1}_+)$  such that

$$
||u - \varphi_{\varepsilon}||_{L^{\infty}(\mathbb{R}^{n+1}_+)} < \varepsilon
$$

and  $\|\mathscr{C}_n(\nabla\varphi_{\varepsilon})\|_{L^{\infty}(\partial\Omega)} < C_{\varepsilon}$ . In particular, he demonstrated that there exists  $\varepsilon \in (0,1)$ such that every harmonic function u, as described above, can be  $\varepsilon$ -approximated. Garnett later established that this holds for every  $0 < \varepsilon < 1$ . Thus, even though Garnett's example demonstrates that [\(1.2\)](#page-2-0) can fail for bounded harmonic functions, these functions can still be approximated closely in the  $L^{\infty}$  norm by a function  $\varphi_{\varepsilon}$  that does satisfy this property.

It is noteworthy that, for example, in Lipschitz domains, the  $\varepsilon$ -approximability of bounded solutions for divergence form elliptic equations is connected to the solvability of the Dirichlet problem with  $L^p$  data<sup>[2](#page-2-3)</sup>. Furthermore, in the case of harmonic functions in more general

<span id="page-2-3"></span><sup>&</sup>lt;sup>2</sup>The relationships outlined have been explored and utilized in the works of Dahlberg [\[Da80\]](#page-69-5) and Kenig, Koch, Pipher, and Toro [\[KKPT00\]](#page-70-3), with significant contributions from Garnett [\[Gar81\]](#page-69-2).

domains, the  $\varepsilon$ -approximation property has significant geometric implications. In particular, if  $\Omega \in \text{AR}(n)$ , it *characterizes* uniform rectifiability of the boundary<sup>[3](#page-3-0)</sup>. This equivalence is derived from the combined work of Hofmann, Martell, and Mayboroda [\[HMM16\]](#page-69-6) and Garnett, Tolsa, and the first named author [\[GarMT18\]](#page-69-7) (see also [\[AGMT23\]](#page-69-8)).

Varopoulos established his extension theorem by first iterating  $\varepsilon$ -approximability to obtain the extension property for  $f \in L^{\infty}(\mathbb{R}^n)$ . Subsequently, he used a "Corona" type decomposition of BMO( $\mathbb{R}^n$ ) functions, credited to Garnett [\[Gar81\]](#page-69-2), to establish the general case. Varopoulos's approach is both powerful and innovative; however, it has a fundamental limitation. Due to its reliance on the  $\varepsilon$ -approximability property of bounded harmonic functions, the Varopoulos method cannot treat domains whose boundaries fail to be uniformly rectifiable (see [\[HT21\]](#page-70-4) for the construction of Varopoulos extensions in such domains).

In the present manuscript, we overcome this geometric obstacle and clarify the true nature of the important extension property of Varopoulos, and the ingredients that go into its proof. We also disprove a conjecture of Hofmann and Tapiola (originally stated in the preprint version of [\[HT21\]](#page-70-4) on arXiv), which was saying that the existence of Varopoulos extensions of BMO functions in  $AR(n)$  domains implies uniform *n*-rectifiability of the boundary. Indeed, one can easily find sets which are n-Ahlfors regular and purely unrectifiable so that their complements are uniform domains (i.e., they satisfy Definitions [2.10](#page-17-1) and [2.11\)](#page-17-2) and so they satisfy the local John (and thus the pointwise John) condition (see Definitions [2.13](#page-18-0) and [2.15\)](#page-18-1). The 4-corner Cantor set of Garnett is such an example in  $\mathbb{R}^2$ .

<span id="page-3-1"></span>Our first main result is the following.

**Theorem 1.1.** *Let*  $\Omega \in AR(s)$  *for*  $s \in (0, n]$ *, which, for*  $s = n$ *, satisfies the pointwise John condition.* If  $f \in \text{BMO}(\sigma_s)$ , there exist  $u : \Omega \to \mathbb{R}$  and  $c_0 \in (0, \frac{1}{4})$  $\frac{1}{4}$  *such that, for any*  $c \leq c_0$ , it holds that

- (i)  $u \in C^{\infty}(\Omega)$ ,
- (ii) sup  $\sup_{\xi \in \partial \Omega} \mathcal{N}_{\sharp,c}(u)(\xi) + \sup_{x \in \Omega} \delta_{\Omega}(x) |\nabla u(x)| \lesssim ||f||_{\text{BMO}(\sigma_s)},$
- (iii) sup  $\sup_{\xi \in \partial \Omega} C_{s,c}(\nabla u)(\xi) \lesssim ||f||_{\text{BMO}(\sigma_s)},$
- (iv) *For*  $\sigma_s$ -almost every  $\xi \in \partial \Omega$ ,

$$
\oint_{B(x,\delta_{\Omega}(x)/2)} u(y) dy \to \begin{cases} f(\xi) & \text{non-tangentially,} \\ f(\xi) & \text{quasi-non-tangentially,} \end{cases} \text{ if } s < n
$$

*Moreover, if*  $f \in L^{\infty}(\sigma_s)$ *, we also obtain the stronger estimate* 

(v) 
$$
\sup_{x \in \Omega} |u(x)| \lesssim ||f||_{L^{\infty}(\sigma_s)}
$$
.

*The constant*  $c_0$  *only depends on dimension, the Ahlfors regularity constants, and the corkscrew condition. Moreover, in the case that*  $s = n$  *and*  $\Omega$  *satisfies the pointwise John condition but not the local John condition, we also assume that* f *is compactly supported.*

Remark 1.2. Note that the second term on the left hand side of the estimate (ii) of Theorem [1.1](#page-3-1) can also be written in terms of the non-tangential maximal function. Namely, (ii) is

<span id="page-3-0"></span> $3$ Uniform rectifiability is a quantitative, scale-invariant version of the classical notion of rectifiability whose theory has been developed extensively in the deep work of David and Semmes [\[DS91,](#page-69-9) [DS93\]](#page-69-10).

equivalent to the estimate

$$
\sup_{\xi \in \partial \Omega} \mathcal{N}_{\sharp,c}(u)(\xi) + \sup_{\xi \in \partial \Omega} \mathcal{N}(\delta_{\Omega} \nabla u)(\xi) \lesssim ||f||_{\text{BMO}(\sigma_s)}.
$$

Our proof of Theorem [1.1,](#page-3-1) under the additional assumption that the local John condition is satisfied when  $s = n$ , does not involve a decomposition of the boundary data, as in the Varopoulos argument. Instead, it relies on a direct approach, which is novel in the case of  $L^{\infty}$  and BMO functions, and hinges on three crucial components. First, we apply Theorem [5.4,](#page-36-0) demonstrating that the regularized dyadic extension of  $f \in BMO(\sigma_s)$  (see [\(3.1\)](#page-21-0) for its definition) is *uniformly* ε*-approximable* (see subsection [2.4](#page-14-0) for the definition). Second, we establish a trace theorem (see Definition [2.16](#page-18-2) and Proposition [6.4\)](#page-41-0). Finally, we employ an iteration argument inspired by Varopoulos.

It is worth noting that the local John condition, representing a scale-invariant connectivity condition between boundary points and corkscrew points at all scales and locations, is necessary only for the trace theorem and specifically in the case where  $\Omega \in AR(s)$  and  $s = n$ , while it is always satisfied in AR(s) for  $s < n$ . A significant challenge in the iteration argument is the need to introduce the Sobolev-type space  $N_{\text{sum}}(\Omega)$  (see [\(2.20\)](#page-13-0) for its definition) and demonstrate its sequential completeness. This constitutes a non-trivial task, occupying the majority of the Appendix (see Corollary [A.9\)](#page-68-0).

The aforementioned method does not seem to work when  $s = n$  and the domain satisfies a connectivity condition weaker than the local John condition called the *pointwise John condition*. So in the case that  $\Omega$  satisfies the pointwise John condition but *not* the local John condition, we resort to a proof closer in spirit to that of Varopoulos. Specifically, we decompose the boundary data f into the sum of a 'good' function  $g \in L^{\infty}$  and a 'bad' function  $b = \sum_j a_j \chi_{Q_j}$ , where  $\sup_{j \ge 1} |a_j| \lesssim ||f||_{\text{BMO}(\sigma)}$  and  $\{Q_j\}_{j \ge 1}$  is a family of dyadic cubes on the boundary satisfying a Carleson packing condition (see [\(4.1\)](#page-26-1) for the definition). The construction of the extension of  $b$  is more standard, although, in rough domains, it is technically involved and requires quite some effort to be accomplished (see [\[HT21,](#page-70-4) Proposition 1.3]). The challenging part is how to build the extension of  $q$  without relying on the  $\varepsilon$ -approximability property of the harmonic extension of g (and, thus, avoiding the restriction to domains with uniformly rectifiable boundaries). Interestingly, the (direct) proof described above works for  $L^{\infty}$  functions in domains merely satisfying the pointwise John condition, allowing us to construct a Varopoulos extension of  $q$ . The reason this occurs is that, rather than obtaining an estimate for  $\mathcal{N}_{\sharp,c}(u)$  for the approximating function of  $v_q$ , we are able to establish a stronger estimate like  $(v)$ . The extension of f is just the sum of the extensions of  $g$  and  $b$ .

We also prove a version of Theorem [1.1](#page-3-1) for boundary functions that belong to  $L^p$ . This result was previously shown in  $\mathbb{R}^{n+1}_+$  by Hytönen and Rosén [\[HR18\]](#page-70-5).

<span id="page-4-0"></span>**Theorem 1.3.** Let  $\Omega \in \text{AR}(s)$  for  $s \in (0, n]$ , which, for  $s = n$ , satisfies the pointwise John *condition.* If  $f \in L^p(\sigma_s)$  with  $p \in (1,\infty)$ , there exist  $u : \Omega \to \mathbb{R}$  and  $c_0 \in (0, \frac{1}{4})$  $\frac{1}{4}$  *such that, for any*  $c \in (0, c_0]$ *, it holds that* 

- (i)  $u \in C^{\infty}(\Omega)$ ,
- (ii)  $\|\mathcal{N}(u)\|_{L^p(\sigma_s)} + \|\mathcal{N}(\delta_{\Omega} \nabla u)\|_{L^p(\sigma_s)} \lesssim \|f\|_{L^p(\sigma_s)}$
- (iii)  $\|\mathcal{C}_{s,c}(\nabla u)\|_{L^p(\sigma_s)} \lesssim \|f\|_{L^p(\sigma_s)}$

(iv) *For*  $\sigma_s$ -almost every  $\xi \in \partial \Omega$ ,<sup>[4](#page-5-0)</sup>

$$
\oint_{B(x,\delta_\Omega(x)/2)} u(y) dy \to \begin{cases} f(\xi) & \text{non-tangentially,} \\ f(\xi) & \text{quasi-non-tangentially,} \end{cases} \quad \text{if } s < n
$$

*The constant*  $c_0$  *only depends on dimension, the Ahlfors regularity constants, and the corkscrew condition.*

In the proof of Theorem [1.3,](#page-4-0) we utilize the regularized dyadic extension of the boundary function and establish its  $\varepsilon$ -approximability in  $L^p$  for  $p \in (1,\infty)$  (see Subsection [2.4](#page-14-0) for the definition). Subsequently, we demonstrate a trace theorem (see Proposition [6.4\)](#page-41-0) and, finally, emloy an iteration argument. This approach follows the general scheme presented in [\[HR18\]](#page-70-5), where the  $\varepsilon$ -approximability in  $L^p$  for the dyadic average extension operator in  $L^p$  for  $p \in (1, \infty)^5$  $p \in (1, \infty)^5$  first appeared.

In fact, this is a key feature of our approach in both Theorems [1.1](#page-3-1) and [1.3,](#page-4-0) which is inspired by, but significantly advances Hytönen and Rosén's work (even in  $\mathbb{R}^{n+1}_+$  for the endpoint spaces). Let us highlight that we encounter significant challenges due to the geometry of our domains. For example, in [\[HR18\]](#page-70-5), it is crucially used the separation of variables  $(x, t) \in \mathbb{R}^n \times \mathbb{R}_+$  to reduce the case to estimating  $\mathscr{C}_n(\partial_t w)$ , where  $\partial_t w$  stands for the partial derivative in the transversal direction. In higher co-dimensions, even if  $\Omega = \mathbb{R}^3 \setminus \mathbb{R}$ , such a reduction does not seem to work, let alone in "rough" domains. Instead, we resort to multiscale analysis to construct the approximating functions. An important component is the proof of the packing condition of the top cubes, which was not shown in  $[HR18]$  (see Proposition [4.2\)](#page-27-0), while the trace theorem and the iteration, also require subtle arguments in our case.

Our second main goal is to establish Varopoulos extensions of Lipschitz functions with compact supports, which are Lipschitz on  $\overline{\Omega}$  and also belong to  $\dot{W}^{1,2}(\Omega;\omega_s)$ . This is crucial due to its applications in Boundary Value Problems for elliptic equations (and systems) with merely bounded coefficients (see Theorem [1.7\)](#page-7-0). In fact, the second part of the theorem has been employed without explicit verification in connection with Boundary Value Problems (see, e.g., [\[DaKe87\]](#page-69-11) and [\[MiTa01\]](#page-70-6)). To the best of our knowledge, a comprehensive proof of this theorem is not available in the literature. However, it should not be considered folklore since establishing it is far from trivial, at least in our setting, and it has neither been documented nor been known among experts.

<span id="page-5-2"></span>**Theorem 1.4.** Let  $\Omega \in AR(s)$  for  $s \in (0, n]$ . If  $f \in Lip<sub>c</sub>(\partial \Omega)$ , there exist a function  $F: \overline{\Omega} \to \mathbb{R}$  and  $c_0 \in (0, 1/2]$ *, such that for any*  $c \in (0, c_0]$ *, it holds that* 

- (i)  $F \in C^{\infty}(\Omega) \cap \text{Lip}(\overline{\Omega}) \cap \dot{W}^{1,2}(\Omega; \omega_s)$ ,
- (ii)  $\|\mathcal{N}(F)\|_{L^p(\sigma_s)} + \|\mathcal{N}(\delta_{\Omega} \nabla F)\|_{L^p(\sigma_s)} \lesssim \|f\|_{L^p(\sigma_s)}$ , for  $p \in (1, \infty]$ ,
- (iii)  $\|\mathcal{C}_{s,c}(\nabla F)\|_{L^p(\sigma_s)} \lesssim \|f\|_{L^p(\sigma_s)}$
- (iv)  $F|_{\partial\Omega} = f$  *continuously*,

<span id="page-5-0"></span> $^{4}$ For the definitions of non-tangential and quasi-non-tangential convergence, see Definition [2.16.](#page-18-2)

<span id="page-5-1"></span><sup>&</sup>lt;sup>5</sup>The concept of  $\varepsilon$ -approximability in  $L^p$  for  $p \in (1,\infty)$  was introduced by Hytönen and Rosén in [\[HR18\]](#page-70-5) who showed that the dyadic average extension operator as well as any weak solution to certain elliptic PDEs in  $\mathbb{R}^{n+1}_+$  are  $\varepsilon$ -approximable in  $L^p$  for every  $\varepsilon \in (0,1)$  and  $p \in (1,\infty)$ . The second part of that work was extended by Hofmann and Tapiola in [\[HT20\]](#page-69-12) to harmonic functions in  $\Omega = \mathbb{R}^{n+1} \setminus E$  where  $E \subset \mathbb{R}^{n+1}$  is a uniformly n-rectifiable set. The converse direction was shown by Bortz and Tapiola in [\[BT19\]](#page-69-13).

(v) 
$$
\|[\mathcal{C}_s(\delta_\Omega|\nabla F|^2)]^{1/2}\|_{L^q(\sigma_s)} \lesssim \|f\|_{L^q(\sigma_s)}, \quad \text{for } q \in [2, \infty).
$$

*Moreover, there exist a function*  $\overline{F} : \overline{\Omega} \to \mathbb{R}$  *and*  $c_0 \in (0, 1/2]$  *such that for any*  $c \in (0, c_0]$ *, it holds that*

- (i)  $\bar{F} \in C^{\infty}(\Omega) \cap \text{Lip}(\overline{\Omega}) \cap \dot{W}^{1,2}(\Omega; \omega_s)$ ,
- (ii) sup  $\sup_{\xi \in \partial \Omega} \mathcal{N}_{\sharp,c}(\overline{F})(\xi) + \sup_{x \in \Omega} \delta_{\Omega}(x) |\nabla \overline{F}(x)| \lesssim ||f||_{\text{BMO}(\sigma_s)},$
- (iii)  $\sup_{\xi \in \partial \Omega} C_{s,c}(\nabla \bar{F})(\xi) \lesssim ||f||_{\text{BMO}(\sigma_s)}$
- (iv)  $\bar{F}|_{\partial\Omega} = f$  *continuously*, (v)  $\sup [\mathcal{C}_s(\delta_\Omega|\nabla \bar{F}|^2)(\xi)]^{1/2} \lesssim ||f||_{\text{BMO}(\sigma_s)}$ . ξ∈∂Ω

*The constant*  $c_0$  *only depends on dimension, the Ahlfors regularity constants, and the corkscrew condition.*

To prove Theorem [1.4,](#page-5-2) considering  $f \in \text{Lip}_c(\partial \Omega)$ , we apply Theorems [5.4](#page-36-0) and [5.5](#page-36-1) to generate approximating functions for the regularized dyadic extension of  $f$ . We define the extension to be equal to the approximation everywhere except in a neighborhood of the boundary with a "width"  $\delta > 0$ , where it is set to equal the regularized dyadic extension. Subsequently, we choose  $\delta$  as  $||f||_{L^p(\sigma_s)}/||f||_{\dot{M}^{1,p}(\sigma_s)}$  in the case of  $p \in (1,\infty)$  (resp.  $||f||_{L^{\infty}(\sigma_s)}$ / Lip f for  $p = \infty$ ) and  $||f||_{BMO(\sigma_s)}$ / Lip f in the case of BMO( $\sigma_s$ ), achieving the desired estimates.

Notably, we don't construct an extension *a priori* and subsequently modify it to obtain the Lipschitz extension; instead, we directly modify the  $\varepsilon$ -approximator of  $v_f$ . This is why we don't need to impose any connectivity condition, as in Theorems [1.1](#page-3-1) and [1.3.](#page-4-0) Instead, the existence of the "trace" is readily ensured by the continuity of  $v_f \in \text{Lip}(\Omega)$ .

The theorem presented below is a variation of Theorem [1.4](#page-5-2) for boundary functions belonging to the Campanato space  $\Lambda_{\beta}(\partial\Omega)$  for  $\beta \in (0,1)$ , as well as the space of Hölder continuous functions  $\text{Lip}_\beta(\partial\Omega)$ . In our setting, any function in  $\Lambda_\beta(\partial\Omega)$  coincides  $\sigma_s$ -almost everywhere with a Hölder continuous function, and the two semi-norms are comparable (refer to Remark [2.2\)](#page-10-0).

<span id="page-6-0"></span>**Theorem 1.5.** Let  $\Omega \in \text{AR}(s)$  for  $s \in (0, n]$ . If  $f \in \Lambda_{\beta}(\partial \Omega)$  for  $\beta \in (0, 1)$ , there exists a *function*  $F : \overline{\Omega} \to \mathbb{R}$  *and*  $c_0 \in (0, 1/2]$ *, such that for any*  $c \in (0, c_0]$ *, it holds that* 

- (i)  $F \in C^{\infty}(\Omega)$ ,
- (ii) sup ξ∈∂Ω  $\mathcal{N}_{\sharp,c}^{(\beta)}(F)(\xi) + \sup_{x \in \Omega} \delta_{\Omega}(x)^{1-\beta} |\nabla F(x)| \lesssim ||f||_{\Lambda_{\beta}(\partial \Omega)},$
- (iii)  $\sup_{\mathcal{S}\in\mathcal{S}} C_{s,c}^{(\beta)}(\nabla F)(\xi) \lesssim ||f||_{\Lambda_{\beta}(\partial\Omega)},$ ξ∈∂Ω
- (iv) nt-lim<sub>x→ξ</sub>  $F|_{\partial\Omega}(x) = f(\xi)$  *for*  $\sigma_s$ -*a.e*  $\xi \in \partial\Omega$ *.*

*Moreover, if*  $f \in \text{Lip}_{\beta}(\partial \Omega)$ , then  $F \in \text{Lip}_{\beta}(\Omega)$  and  $F|_{\partial \Omega} = f$  continuously. The con*stant*  $c_0$  *only depends on dimension, the Ahlfors regularity constants, and the corkscrew condition.*

Remarkably, this theorem marks the first appearance of such a result, despite  $\Lambda_{\beta}(\partial\Omega)$ being a natural endpoint in the interpolation scale that contains  $L^p(\sigma_s)$  and  ${\rm BMO}(\sigma_s)$ . Its proof is a lot easier compared to those for  $L^p(\sigma_s)$  and  $\text{BMO}(\sigma_s)$  boundary functions, as

the regularized version of the dyadic extension of the boundary data already satisfies the desired properties, eliminating the need for  $\varepsilon$ -approximability.

<span id="page-7-1"></span>Remark 1.6. The proof of the existence of extensions of complex-valued boundary functions is exactly the same but for the sake of simplicity we prefer to state and prove our results for real-valued boundary functions. Moreover, if  $\vec{f} : \partial\Omega \to \mathbb{C}^m$  with  $\vec{f} = (f_1, \dots, f_m)$ , then its extension is just the vector field  $\vec{F} = (F_1, \ldots, F_m)$ , where  $F_j$  is the extension of  $f_j$ for each  $j \in \{1, 2, ..., m\}$ .

Lastly, we use Theorem [1.4](#page-5-2) to obtain connections between Poisson and Boundary Value Problems (see Definitions [2.6,](#page-16-0) [2.8,](#page-17-3) and [2.9\)](#page-17-4) for systems of elliptic equations in divergence form with merely bounded complex-valued coefficients. The estimates derived for solutions of elliptic boundary value problems can be perceived as far-reaching extensions of the aforementioned alternative approach to Fefferman's duality theorem [\[FS72\]](#page-69-3). In particular, we will prove the following.

<span id="page-7-0"></span>**Theorem 1.7.** Let  $\Omega \in AR(n)$  and L be defined in [\(2.31\)](#page-15-1). If  $L^*$  is its formal adjoint, then *the following hold:*

- *(1)* If  $(\text{PR}_{p}^{L})$  is solvable in  $\Omega$  for some  $p > 1$ , then  $(D_{p'}^{L*})$  is also solvable, where  $1/p + 1/p' = 1.$
- (2) If  $(\widetilde{\text{PR}}_q^L)$  *with*  $H = 0$  *is solvable in*  $\Omega$  *for*  $q \in [1, 2]$ *, then both*  $(\widetilde{\text{PD}}_{q'}^{L^*})$  *with*  $H = 0$  $and \ (\widetilde{\mathbf{D}}_{q'}^{L^*})$  *are solvable in*  $\Omega$ *.*
- *(3)* If  $(PD_{p}^{L})$  for  $p \in (1, \infty)$  is solvable in  $\Omega$  with  $H = 0$ , then the Dirichlet problem  $(D_p^L)$  *is also solvable in*  $\Omega$ *.*
- *(4) If*  $(\widetilde{PD}_{q}^{L})$  *is solvable in*  $\Omega$  *with*  $H = 0$  *for some*  $q \in [2, \infty]$ *, then*  $(\widetilde{D}_{q}^{L})$  *is also solvable in* Ω*.*

Recently, the first named author and Tolsa [\[MT22\]](#page-70-7) constructed an *almost harmonic extension* of functions in the Hajłasz Sobolev space  $\dot{M}^{1,p}(\sigma_n)$ , which is the correct analogue of the  $L^p$  version of Varopoulos extension for one "smoothness level" up. To be precise, it was proved in  $[MT22]$  that the Carleson functional defined in  $(2.11)$  of the distributional Laplacian of the almost harmonic extension is in  $L^p(\sigma_n)$  and in [\[MPT22\]](#page-70-8) that the non-tangential maximal function of its gradient is in  $L^p(\sigma_n)$  with norms controlled by the  $\dot{M}^{1,p}(\sigma_n)$  seminorm of the boundary function. The almost harmonic extension and its elliptic analogue (see [\[MPT22\]](#page-70-8)) were very important since they turned out to be the main ingredients for the solution of the  $L^p$ -Regularity problem in domains with interior big pieces of chord-arc domains ([\[AHMMT20,](#page-69-14) Definition 2.12, p. 892]) for the Laplace operator [\[MT22\]](#page-70-7) and for elliptic operators satisfying the Dahlberg-Kenig-Pipher (DKP) condition [\[MPT22\]](#page-70-8) respectively. This solved a 30 year-old question of Kenig.

The Poisson Dirichlet problem  $(\mathrm{PD}_p^L)$  (resp. the Poisson regularity problem  $(\mathrm{PR}_p^L)$ ) with interior data in suitable Carleson spaces for  $p > 1$  with scale-invariant estimates for the nontangential maximal function of the (resp. gradient of the) solution (see Definitions [2.8](#page-17-3) and [2.9\)](#page-17-4), was first defined in [\[MPT22\]](#page-70-8) in order to overcome the obstacle that it is not known if elliptic operators satisfying the DKP condition have bounded layer potentials. In particular, the authors show that, for such operators,  $(PD_{p'}^{L^*}) \Rightarrow (R_p^L)$  for any  $p > 1$ . To do so, they

use the almost elliptic extension. One of their results states the following equivalences for elliptic equations (not systems) with merely bounded coefficients:

$$
(\mathcal{D}_{p'}^{L^*}) \Leftrightarrow (\mathcal{P}\mathcal{D}_{p'}^{L^*}) \Leftrightarrow (\mathcal{P}\mathcal{R}_p^L) \qquad \text{if } p \in (1, \infty).
$$

The equivalence  $(PD_{p'}^{L^*}) \Leftrightarrow (PR_p^L)$  holds for systems as well, as the proof in [\[MPT22\]](#page-70-8) does not utilize tools like the maximum principle or the elliptic measure. However, the remaining results rely significantly on the connection between the weak- $A_{\infty}$  condition of the elliptic measure and the solvability of  $(D_{p'}^{L^*})$ , and thus, they only hold for real equations. Inspired by the use of the almost elliptic extension in [\[MPT22\]](#page-70-8), we employ the Varopoulos extension constructed in Theorem [1.4](#page-5-2) to extend some of these results to elliptic systems. We also obtain endpoint results, which are new even for real equations.

Moreover, we introduce the notions of solvability for the Dirichlet and Poisson problems with square function estimates and for the Poisson regularity problem  $(\widetilde{PR}_{q}^{L})$  with data in Coifman-Meyer-Stein tent spaces and establish connections between these problems. In the recent work of Gallegos, Tolsa, and the first named author [\[GalMT23\]](#page-69-1), the authors, partially inspired by the introduction of  $(\widetilde{PR}_q^L)$  in the present manuscript, show extrapolation of solvability of the regularity problem  $(R_p^L)$  (see Definition [2.7\)](#page-17-5) and the (modified) Poisson regularity problem  $(\widetilde{\text{PR}}_q^L)$  in  $\text{AR}(n)$  domains for elliptic operators with real and merely bounded coefficients. Theorem [1.7](#page-7-0) played a crucial role in the proof of the extrapolation of  $(\widetilde{\text{PR}}_q^L)$  as it was used to demonstrate that  $(\widetilde{\text{PR}}_q^L) \Rightarrow \omega_L \in \text{weak-}A_\infty$ . For a detailed history of work in this area, we refer to the introduction of [\[MPT22\]](#page-70-8).

It is worth noting that Theorem [1.4](#page-5-2) plays a crucial role in a forthcoming work by Poggi and the first named author [\[MP24\]](#page-70-0). In that work, it is established that, for real equations, not only  $(D_p^L) \Leftrightarrow (PD_p^L)$ , but the unique solution of the continuous Dirichlet problem with boundary data g is well approximated by a sequence of solutions of  $(PD<sub>p</sub><sup>L</sup>)$  with  $H = 0$  in the following sense: If  $g \in \text{Lip}_c(\partial \Omega)$  and u is the unique solution of  $(D_p^L)$ , then there exists a sequence  $\{\vec{F}_j\}_{j\geq 1} \subset \text{Lip}_c(\Omega;\mathbb{R}^{n+1})$  with the following properties: (i)  $\sup_{j\geq 1} \|C_q(|\vec{F}_j|)\|_{L^p(\sigma)} \ \lesssim \ \|g\|_{L^p(\sigma)};$  (ii) If  $w_j \ \in \ Y_0^{1,2}$  $\chi_0^{1,2}(\Omega)$  is the unique solution of  $Lw_j = -\text{div }\vec{F}_j$ , it satisfies  $w_j \to u$  in the following topologies: a) in the strong topology of  $C^{\alpha}_{loc}(\Omega)$  for some  $\alpha \in (0,1)$ ; b) in the strong topology of  $W^{1,2}_{loc}(\Omega)$ ; c) in the weak-\* topology of  $N_{r,p}(\Omega)$ , for each  $r, p \in (1,\infty)$ . Moreover, the weak-\* convergence cannot be improved to convergence in the weak-topology of  $N_{r,p}(\Omega)$  for any  $r, p \in (1,\infty)$  unless  $g=0.$ 

<span id="page-8-0"></span>1.1. Related results. While writing this paper, we were informed by Bruno Poggi and Xavier Tolsa that in collaboration with Simon Bortz and Olli Tapiola, they have indepen-dently obtained in [\[BOPT23\]](#page-69-15), as a corollary of their main result studying  $\varepsilon$ -approximability of solutions to arbitrary elliptic partial differential equations, a less general version of Theo-rem [1.1](#page-3-1) which holds for uniform domains with  $n$ -Ahlfors regular boundaries such that there is an elliptic measure which is  $A_{\infty}$  with respect to surface measure. Their assumptions hold, in particular, for the complement of the 4-corner Cantor set in  $\mathbb{R}^2$ , thus they also show that uniform rectifiability is not a necessary condition in order to construct Varopoulos-type extensions.

#### <span id="page-9-0"></span>10 MIHALIS MOURGOGLOU AND THANASIS ZACHAROPOULOS

#### 2. PRELIMINARIES AND NOTATION

We will write  $a \lesssim b$  if there is a constant  $C > 0$  so that  $a \leq Cb$  and  $a \approx b$  if  $\alpha \lesssim b$  and  $b \lesssim a$ . If we want to indicate the dependence of C on a certain quantity s, we write  $a \lesssim_s b$ . For a function space X we denote by  $X_c$  the space of all the compactly supported functions in  $X$ .

<span id="page-9-1"></span>2.1. Preliminaries. In  $\mathbb{R}^{n+1}$  and for  $s \in [0, n+1]$ , we denote by  $\mathcal{H}^s$  the s-dimensional Hausdorff measure and assume that  $\mathcal{H}^{n+1}$  is normalized so that it coincides with  $\mathcal{L}^{n+1}$ , the  $(n + 1)$ -dimensional Lebesgue measure in  $\mathbb{R}^{n+1}$ . We also denote by  $\sigma_s := \mathcal{H}^s|_{\partial\Omega}$ the "surface" measure of  $\Omega$ . When the dimension is clear from the context we drop the dependence on s and just write  $\sigma$ .

<span id="page-9-3"></span>**Definition 2.1.** If  $s \in (0, n + 1]$ , a measure  $\mu$  in  $\mathbb{R}^{n+1}$  is called *s-Ahlfors regular* if there exists some constant  $C_0 > 0$  such that

$$
C_0^{-1} r^s \le \mu(B(x, r)) \le C_0 r^s
$$

for all  $x \in \text{supp } \mu$  and  $0 < r < \text{diam}(\text{supp }\mu)$ . If  $E \subset \mathbb{R}^{n+1}$  is a closed set we say that  $E$ is *s*-*Ahlfors regular* if  $\mathcal{H}^s|_E$  is *s*-Ahlfors regular.

<span id="page-9-2"></span>2.2. Function spaces. We write  $2^* = \frac{2(n+1)}{n-1}$  $\frac{(n+1)}{n-1}$  and  $2_* = (2^*)' = \frac{2(n+1)}{n+3}$ . Recall that  $C_c^{\infty}(\Omega)$  is the space of compactly supported smooth functions in  $\Omega$ . For  $p \in [1,\infty)$  and a non-negative function  $w \in L^1_{loc}(\Omega)$  we define the homogeneous weighted Sobolev space  $\dot{W}^{1,p}(\Omega; w)$  to be the space consisted of  $L^1_{loc}(\Omega)$  functions whose weak gradients exist in  $\Omega$  and are in  $L^p(\Omega; w)$ . We also define the inhomogeneous weighted Sobolev space  $W^{1,p}(\Omega; w)$  to be the space of functions in  $L^p(\Omega; w)$  whose weak derivatives exist in  $\Omega$ and are also in  $L^p(\Omega; w)$  and  $W_0^{1,p}$  $C_0^{1,p}(\Omega; w)$  to be the completion of  $C_c^{\infty}(\Omega)$  under the norm  $||u||_{W^{1,p}(\Omega; w)} := ||u||_{L^p(\Omega; w)} + ||\nabla u||_{L^p(\Omega; w)}$ . Finally, we let  $Y_0^{1,2}$  $\chi_0^{1,2}(\Omega; w)$  be the completion of  $C_c^{\infty}(\Omega)$  under the norm  $||u||_{Y^{1,2}(\Omega;w)} := ||u||_{L^{2^*}(\Omega;w)} + ||\nabla u||_{L^2(\Omega;w)}$ .

Let  $\Sigma$  be a metric space equipped with a non-atomic doubling measure  $\sigma$  on  $\Sigma$ , which means that there is a uniform constant  $C_{\sigma} \ge 1$  such that  $\sigma(B(x, 2r)) \le C_{\sigma} \sigma(B(x, r))$ , for all  $x \in \Sigma$  and  $r > 0$ . If  $E \subset \Sigma$  is a Borel set such that  $0 < \sigma(E) < \infty$  and  $f \in L^1_{loc}(\sigma)$ , we denote the average of  $f$  over  $E$  by

(2.1) 
$$
m_{\sigma,E}f := \int_E f \, d\sigma := \frac{1}{\sigma(E)} \int_E f d\sigma.
$$

If  $\sigma$  is the Lebesgue measure then we simply write  $m_E f$ .

For  $\beta \in [0, 1)$  we define  $\Lambda_{\beta}(\partial \Omega)$  to be the *Campanato space* consisting of the functions  $f \in L^1_{loc}(\sigma)$  satisfying

$$
(2.2) \t\t ||f||_{\Lambda_{\beta}(\partial\Omega)} := \sup_{\substack{x \in \text{supp }\sigma \\ r \in (0,2\text{ diam }\partial\Omega)}} \frac{1}{r^{\beta}} \int_{B(x,r)} |f(y) - m_{\sigma,B(x,r)}f| d\sigma(y) < \infty.
$$

Note that  $Λ_0(σ) = BMO(σ)$ , the space of functions of *bounded mean* σ-*oscillation*. We also define the space of functions of *vanishing mean oscillation*<sup>[6](#page-9-4)</sup>, which we denote by

<span id="page-9-4"></span> $6$ VMO was originally introduced by Sarason in [\[Sar75\]](#page-70-9).

VMO( $\sigma$ ), to be the closure of the space of continuous functions with compact support  $C_c(\Sigma)$  in the BMO( $\sigma$ ) norm.

We say that  $\alpha$  is a 2-*atom* if there exists  $x \in \Sigma$  and  $0 < r < \text{diam}(\Sigma)$  such that

$$
\operatorname{supp} \alpha \subset B(x,r), \quad \|\alpha\|_{L^2(\sigma)} \lesssim \sigma(B(x,r))^{-1/2} \quad \text{and} \quad \int \alpha \, d\sigma = 0.
$$

We define the *atomic Hardy space*  $H^1(\sigma)$  as follows:  $f \in H^1(\sigma)$  if there exist a sequence  $\lambda_j \in \mathbb{C}$  and a sequence of 2-atoms  $\alpha_j$  such that  $f = \sum_j \lambda_j \alpha_j$  in  $L^1(\sigma)$ . We say that f has an atomic decomposition.  $H^1(\sigma)$  is a subspace of  $L^1(\sigma)$  and is a Banach space with norm

$$
||f||_{H^1(\sigma)} := \inf \left\{ \sum_j |\lambda_j| : \text{all atomic decompositions } f = \sum_j \lambda_j \alpha_j \right\}
$$

By the work of Coifmann and Weiss, [\[CW77\]](#page-69-16), we have that  $(H^1(\sigma))^* = \text{BMO}(\sigma)$  and  $(VMO(\sigma))^* = H^1(\sigma).$ 

For  $\beta \in (0,1]$  we define  $\text{Lip}_{\beta}(\Sigma)$  to be the space of measurable functions that satisfy

(2.3) 
$$
||f||_{\text{Lip}_{\beta}(\Sigma)} := \sup_{\substack{x,y \in \Sigma \\ x \neq y}} \frac{|f(x) - f(y)|}{|x - y|^{\beta}} < \infty.
$$

When  $\beta = 1$  we simply write  $\text{Lip}(\Sigma)$  since it is the space of Lipschitz functions. If  $\Sigma$ is locally compact then it holds that  $\text{Lip}_c(\Sigma)$  is dense in  $C_c(\Sigma)$  in the supremum norm. Therefore, it is easy to see that in that case

$$
\overline{\text{Lip}_c(\Sigma)}^{\text{BMO}(\sigma)} = \text{VMO}(\sigma).
$$

<span id="page-10-0"></span>Remark 2.2. By a simple inspection of the proof of [\[MS79,](#page-70-10) Theorem 4], it is easy to see that if  $\Sigma$  is a metric space equipped with a measure  $\sigma$  which is s-Ahlfors regular, then, if  $\beta \in (0,1)$ , for every  $f \in \Lambda_{\beta}(\Sigma)$  there exists  $g \in \text{Lip}_{\beta}(\Sigma)$  such that  $f(x) = g(x)$  for  $\sigma$ -a.e.  $x \in \Sigma$  and  $||f||_{\Lambda_{\beta}(\sigma)} \approx ||f||_{\mathrm{Lip}_{\beta}(\Sigma)}.$ 

Following [\[Ha96\]](#page-69-17), we will introduce the *Hajłasz's Sobolev space* on Σ. For a Borel function  $f : \Sigma \to \mathbb{R}$ , we say that a non-negative Borel function  $g : \Sigma \to \mathbb{R}$  is a *Hajłasz upper gradient of* f if

(2.4) 
$$
|f(x) - f(y)| \le |x - y| (g(x) + g(y))
$$
 for  $\sigma$ -a.e.  $x, y \in \Sigma$ .

We denote the collection of all the Hajłasz upper gradients of f by  $D(f)$ .

For  $p > 0$ , we denote by  $\dot{M}^{1,p}(\sigma)$  the space of Borel functions f which have a Hajłasz upper gradient in  $L^p(\sigma)$ , and we let  $M^{1,p}(\sigma)$  be the space of functions  $f \in L^p(\sigma)$  which have a Hajłasz upper gradient in  $L^p(\sigma)$ , i.e.,  $M^{1,p}(\sigma) = \dot{M}^{1,p}(\sigma) \cap L^p(\sigma)$ . We define the semi-norm (as it annihilates constants)

<span id="page-10-1"></span>(2.5) 
$$
||f||_{\dot{M}^{1,p}(\sigma)} = \inf_{g \in D(f)} ||g||_{L^p(\sigma)}.
$$

If  $\Sigma$  is bounded, then we define the norm

<span id="page-10-2"></span>(2.6) 
$$
||f||_{M^{1,p}(\sigma)} = (\text{diam}\,\Sigma)^{-1} ||f||_{L^p(\sigma)} + \inf_{g \in D(f)} ||g||_{L^p(\sigma)},
$$

.

while if  $\Sigma$  is unbounded, we consider the space  $M^{1,p}(\sigma) := \dot{M}^{1,p}(\sigma)/\mathbb{R}$ . Observe that, from the uniform convexity of  $L^p(\sigma)$  for  $p \in (1,\infty)$ , one easily deduces that the infimum in the definition of the norm  $\|\cdot\|_{\dot{M}^{1,p}(\Sigma)}$  and  $\|\cdot\|_{M^{1,p}(\Sigma)}$ , in [\(2.5\)](#page-10-1) and [\(2.6\)](#page-10-2) respectively, is attained and is unique. We denote by  $\nabla_{H,p} f$  the function g which attains the infimum which we will call the *least Hajłasz upper gradient* of f.

<span id="page-11-0"></span>2.3. Maximal operators and Carleson functionals. Set  $\delta_{\Omega}(\cdot) := \text{dist}(\cdot, \Omega^c)$ ,  $B^x :=$  $B(x, \delta_{\Omega}(x))$ , and  $cB^x := B(x, c \delta_{\Omega}(x))$ , for  $c \in (0, \frac{1}{2})$  $\frac{1}{2}$ . For  $f \in L^1_{loc}(\mu)$  and  $x \in \Omega$ , we denote

$$
m_{q,c}(f)(x) := \begin{cases} m_{cB^x}(|f|^q)^{1/q} & \text{if } 1 \le q < \infty, \\ \sup_{y \in cB^x} |f(y)| & \text{if } q = \infty, \end{cases}
$$

and

(2.7) 
$$
m_{\sharp,c}(f)(x) := m_{\infty,c}(f - m_{cB^x}f)(x).
$$

We define the *centered Hardy-Littlewood maximal operator* for a function  $f \in L^1_{loc}(\sigma)$ as

$$
\mathcal{M}(f)(x) := \sup_{r>0} m_{\sigma,B(x,r)}(|f|), \qquad x \in \Sigma
$$

while the *non-centered Hardy-Littlewood maximal operator* is defined to be

$$
\mathcal{M}(f)(x) := \sup_{B \ni x} m_{\sigma,B}(|f|),
$$

where the supremum is taken over all balls B containing  $x \in \Sigma$ . The *dyadic Hardy*-*Littlewood maximal operator* with respect to a dyadic lattice  $\mathcal{D}_{\sigma}$  on  $\Sigma^7$  $\Sigma^7$  will be denoted

$$
\mathcal{M}_{\mathcal{D}_{\sigma}}f(x) := \sup_{Q \in \mathcal{D}_{\sigma}, Q \ni x} m_{\sigma, Q}(|f|),
$$

and if the measure is clear from the context, we will just write  $M_{\mathcal{D}}f$ . We also set

(2.8) 
$$
Mf(Q) := \sup_{\substack{R \in \mathcal{D}_{\sigma} \\ Q \subset R}} m_{\sigma,R}(|f|)
$$

to be a *truncated* version of  $M_{\mathcal{D}}f(x)$ .

From now on, we assume that  $\Omega \in \text{AR}(s)$  in  $\mathbb{R}^{n+1}$  and  $\sigma = \mathcal{H}^s|_{\partial \Omega}$ .

For  $\alpha > 0$  and  $\xi \in \partial \Omega$  we define the *cone* with vertex  $\xi$  and aperture  $\alpha > 0$  to be the set

<span id="page-11-1"></span>
$$
\gamma_{\alpha}(\xi) := \{ x \in \Omega : |x - \xi| < (1 + \alpha) \text{dist}(x, \partial \Omega) \};
$$

The *non-tangential maximal operator* of a measurable function  $f : \Omega \to \mathbb{R}$  for a fixed aperture  $\alpha > 0$  by

(2.9) 
$$
\mathcal{N}_{\alpha}(f)(\xi) := \sup_{x \in \gamma_{\alpha}(\xi)} |f(x)|, \qquad \xi \in \partial \Omega.
$$

By a straightforward modification of the classical proof of Feffermann and Stein [\[FS72,](#page-69-3) Lemma 1], one can show the following.

<span id="page-11-2"></span><sup>&</sup>lt;sup>7</sup>For the construction of dyadic lattices in this setting, see e.g. subsection [2.7.](#page-19-0)

<span id="page-12-3"></span>**Lemma 2.3.** *For*  $\Omega \in AR(s)$  *and for*  $s \in (0, n]$ *, there holds*  $\|\mathcal{N}_{\alpha}(f)\|_{L^{p}(\sigma)} \approx_{\alpha, \beta, s}$  $\|\mathcal{N}_{\beta}(f)\|_{L^{p}(\sigma)}$  for all  $\alpha, \beta > 0$  and  $p \in (0, \infty)$ .

For a fixed aperture  $\alpha > 0$ ,  $\beta \in [0, 1)$  and a constant  $c \in (0, \frac{1}{2})$  $\frac{1}{2}$ , we also define the *sharp non-tangential maximal opeartor* applied to a measurable function  $f : \Omega \to \mathbb{R}$  by

<span id="page-12-0"></span>(2.10) 
$$
\mathcal{N}_{\sharp,\alpha,c}^{(\beta)}(f)(\xi) := \sup_{x \in \gamma_\alpha(\xi)} \delta_{\Omega}(x)^{-\beta} m_{\sharp,c}(f)(x), \quad \xi \in \partial \Omega.
$$

Setting  $\omega_s(x) := \delta_{\Omega}(x)^{s-n}$  for  $x \in \Omega$ , we define the *Carleson functional* of a function  $F \in L^1_{\rm loc}(\Omega, \omega_s(x) dx)$  by

<span id="page-12-2"></span>
$$
(2.11) \t\t\mathscr{C}_{s}^{(\beta)}(F)(\xi) := \sup_{r>0} \frac{1}{r^{s+\beta}} \int_{B(\xi,r)\cap\Omega} |F(x)| \,\omega_{s}(x) \,dx, \quad \xi \in \partial\Omega.
$$

We define the *modified Carleson functional* of a locally bounded function F by means of

<span id="page-12-1"></span>(2.12) 
$$
\mathcal{C}_{s,c}^{(\beta)}(F)(\xi) := \mathscr{C}_s^{(\beta)}(m_{\infty,c}(F))(\xi), \quad \xi \in \partial \Omega.
$$

For any  $q \in [1, \infty)$ , the *q-Carleson functional* of a function  $F \in L^q_{loc}(\Omega, dx)$  is defined to be

$$
(2.13) \qquad \mathcal{C}_{s,q,c}^{(\beta)}(F)(\xi) := \sup_{r>0} \frac{1}{r^{s+\beta}} \int_{B(\xi,r)\cap\Omega} m_{q,\sigma,c} B^x(|F|) \,\omega_s(x) \,dx, \quad \xi \in \partial\Omega.
$$

<span id="page-12-4"></span>**Lemma 2.4.** *If*  $\Omega \in \text{AR}(s)$  *for*  $s \in (0, n]$ *,*  $\beta \in [0, 1)$ *,*  $q \in [1, \infty)$ *,*  $F \in L_{\text{loc}}^q(\Omega, dx)$ *, and*  $0 < c_1 < c_2 \leq \frac{1}{2}$  $\frac{1}{2}$ , then

(2.14) 
$$
\mathcal{C}_{s,q,c_2}^{(\beta)}(F)(\xi) \lesssim_{c_1,c_2} \mathcal{C}_{s,q,c_1}^{(\beta)}(F)(\xi), \text{ for any } \xi \in \partial \Omega.
$$

*Proof.* The case  $s = n$  and  $\beta = 0$  was proved in [\[MPT22,](#page-70-8) Lemma 2.2], while the proof in the other cases follow by a routine adaptation of the same arguments.  $\Box$ 

If it is clear from the context and in view of Lemmas [2.3](#page-12-3) and [2.4,](#page-12-4) most of the times we will drop the dependence of  $\mathcal{N}_{\alpha}$ ,  $\mathcal{N}_{\sharp,\alpha,c}^{(\beta)}$ ,  $\mathcal{C}_{s,q,c}^{(\beta)}$ , and  $\mathcal{C}_{s,c}^{(\beta)}$  on  $\alpha$  and  $c$ , and write  $\mathcal{N}$ ,  $\mathcal{N}_{\sharp}^{(\beta)}$ 'ι<sup>μ</sup>',<br>♯  $\mathcal{C}_{s,q}^{(\beta)}$ , and  $\mathcal{C}_{s}^{(\beta)}$ . If  $s=n$  and/or  $\beta=0$ , we will drop the dependence on s and/or  $\beta$  as well.

For  $p \in (1,\infty)$  we introduce the Banach spaces

(2.15) 
$$
N^{p}(\Omega) := \{w : \Omega \to \mathbb{R} : w \text{ is measurable and } \mathcal{N}(w) \in L^{p}(\sigma)\},
$$

$$
(2.16) \tC_{s,\infty}^p(\Omega) := \{ w \in L_{\text{loc}}^{\infty}(\Omega) : \mathcal{C}_s(w) \in L^p(\sigma) \},
$$

equipped with the respective norms

 $\|w\|_{\mathrm{N}^p(\Omega)} := \|\mathcal{N}(w)\|_{L^p(\sigma)} \quad \text{and} \quad \|\vec{F}\|_{\mathrm{C}^p_{s,\infty}(\Omega)} := \|\mathcal{C}_s(|\vec{F}|)\|_{L^p(\sigma)}.$ 

For  $p = \infty$  we define

(2.17) 
$$
N^{\infty}(\Omega) := \{ w \in C(\Omega) : \sup_{\xi \in \partial \Omega} \mathcal{N}(w)(\xi) < \infty \}
$$

(2.18) 
$$
C_{s,\infty}^{\infty}(\Omega) := \{ w \in C(\Omega) : \sup_{\xi \in \partial \Omega} C_s(w)(\xi) < \infty \}
$$

(2.19) 
$$
N^{\infty}_{\sharp}(\Omega) := \{ w \in C(\Omega) : \sup_{\xi \in \partial \Omega} \mathcal{N}_{\sharp,c}(w)(\xi) < \infty \}
$$

and equip them with the respective norms

$$
\|w\|_{\mathrm{N}^\infty(\Omega)}:=\sup_{\xi\in\partial\Omega}\mathcal{N}(w)(\xi)\quad\text{and}\quad \|\vec{F}\|_{{\mathbf C}^\infty_{s,\infty}(\Omega)}:=\sup_{\xi\in\partial\Omega}\mathcal{C}_s(|\vec{F}|)(\xi),
$$

and the semi-norm

<span id="page-13-0"></span>
$$
||w||_{\mathrm{N}_{\sharp}^{\infty}(\Omega)}:=\sup_{\xi\in\partial\Omega} \mathcal{N}_{\sharp,c}(w)(\xi)=\sup_{x\in\Omega} m_{\sharp,c}(w)(x).
$$

We also define the space

(2.20) 
$$
N_{sum}(\Omega) := \{ u \in C^{1}(\Omega) : (u, \delta_{\Omega} \nabla u) \in N_{\sharp}^{\infty}(\Omega) \times N^{\infty}(\Omega) \}
$$

and equip it with the semi-norm

$$
||u||_{\mathrm{N}_{\mathrm{sum}(\Omega)}} := ||u||_{\mathrm{N}_{\sharp}^{\infty}(\Omega)} + ||\delta_{\Omega}\nabla u||_{\mathrm{N}^{\infty}(\Omega)}.
$$

In Corollary [A.9,](#page-68-0) we will show that  $(N_{sum}(\Omega), \| \cdot \|_{N_{sum}(\Omega)})$  is sequentially complete. We set

$$
\mathrm{C}^{1,p}_{s,\infty}(\Omega):=\{u\in C^1(\Omega):\nabla u\in \mathrm{C}^p_{s,\infty}(\Omega)\},\quad\text{for }p\in(1,\infty]
$$

and equip it with the semi-norm  $||u||_{C^{1,p}_{s,\infty}(\Omega)} := ||\nabla u||_{C^p_{s,\infty}(\Omega)}$ .

If  $G : \Omega \to \mathbb{R}$  is a measurable function in  $\Omega$ , we define the *area functional* of G, for a fixed aperture  $\alpha > 0$  as

(2.21) 
$$
\mathcal{A}^{(\alpha)}G(\xi) := \int_{\gamma_{\alpha}(\xi)} |G(x)| \delta_{\Omega}(x)^{-n} dx, \quad \xi \in \partial \Omega.
$$

The following lemma is proved in the Appendix.

**Lemma 2.5.** *Let*  $\Omega \in \text{AR}(s)$  *for*  $s \in (0, n]$ *,*  $u \in L^1_{\text{loc}}(\Omega, \omega_s)$ *,*  $p \in [1, \infty)$ *, and*  $\alpha \geq 1$ *. Then there exists*  $C \geq 1$  *such that for any*  $\xi \in \partial\Omega$  *and*  $r \in (0, 2 \text{ diam}(\partial\Omega))$ *, it holds that* 

<span id="page-13-2"></span>
$$
(2.22) \t\t\t\t\|\mathcal{A}^{(\alpha)}(u\mathbf{1}_{B(\xi,r)})\|_{L^p(\sigma,B(\xi,r))} \lesssim r^{\beta} \|\mathscr{C}_s^{(\beta)}(u\mathbf{1}_{B(\xi,Cr)})\|_{L^p(\sigma,B(\xi,Cr))}.
$$

*If*  $\beta = 0$ *, it also holds* 

$$
(2.23) \t\t\t\t\t\|\mathcal{A}^{(\alpha)}(u)\|_{L^p(\sigma)} \lesssim \|\mathscr{C}_s(u)\|_{L^p(\sigma)}.
$$

*Moreover for*  $\beta = 0$  *and*  $1 < p \le \infty$  *we have* 

(2.24) kCs(u)kLp(σ) . kA(α) (u)kLp(σ) .

We also introduce the *modified non-tangential maximal operator*  $\widetilde{N}_{\alpha,c,r}$  for a given aperture  $\alpha > 0$ , a parameter  $c \in (0, 1/2]$  and  $r \ge 1$ : for any  $u \in L_{loc}^r(\Omega)$  it is defined as

<span id="page-13-1"></span>
$$
\widetilde{\mathcal{N}}_{\alpha,c,r}u(\xi) := \sup_{x \in \gamma_{\alpha}(\xi)} \Big( \operatorname{\int}_{B(x,c\delta_{\Omega}(x))} |u(y)|^{r} dy \Big)^{1/r}, \quad \xi \in \partial \Omega.
$$

The  $L^p$ -norms of these non-tangential maximal functions with different aperture  $\alpha$  or averaging parameter  $c$  are comparable (see [\[MPT22,](#page-70-8) Lemma 2.1]), and for ease the notation we will just write  $\mathcal{N}_r = \mathcal{N}_{\alpha,c,r}$  when we do not need to specify neither  $\alpha$  nor c.

For any  $q \ge 1$  and  $p > 1$ , we define the Banach spaces

$$
C_{s,q,p}(\Omega) := \{ H \in L^q_{loc}(\Omega) : \mathcal{C}_{s,q}(H) \in L^p(\sigma) \},
$$

with norm  $\|H\|_{{\rm C}_{s,q,p}}=\|{\mathcal C}_{s,q}(H)\|_{L^p(\sigma)},$  and for  $r\in[1,\infty],$   $p>1$  let

$$
N_{r,p}(\Omega) := \{ u \in L^r_{loc}(\Omega) : \widetilde{\mathcal{N}}_r(u) \in L^p(\sigma) \},
$$

(where we identify  $\mathcal{N}_{\infty} = \mathcal{N}$ ) with norm  $||u||_{N_{r,p}(\Omega)} = ||\mathcal{N}_r(u)||_{L^p(\sigma)}$ . By (the proof of) [\[MPT22,](#page-70-8) Proposition 2.4], it holds that if either  $\Omega$  is bounded or  $\partial\Omega$  is unbounded,  $N_{q,p}(\Omega) = (C_{s,q',p'}(\Omega))^*$ . When  $s = n$  we drop the subscript s from  $C_{s,q,p}$ .

If  $\Omega \in AR(s)$ , we define the *tent spaces* 

(2.25) 
$$
T_{s,2}^{\infty}(\Omega) := \{ f \in L_{\text{loc}}^2(\Omega) : \mathscr{C}_s(f^2 \,\delta_{\Omega}^{-1}) \in L^{\infty}(\sigma) \}
$$

and

(2.26) 
$$
T_2^p(\Omega) := \{ g \in L^2_{\text{loc}}(\Omega) : \left( \mathcal{A}(g^2 \, \delta_{\Omega}^{-1}) \right)^{1/2} \in L^p(\sigma) \}, \text{ for } p \in (0, \infty),
$$

and we equip them with the respective norms

$$
||f||_{T_{s,2}^{\infty}(\Omega)} = ||\mathscr{C}_s(f^2 \,\delta_{\Omega}^{-1})^{1/2}||_{L^{\infty}(\sigma)} \quad \text{and} \quad ||g||_{T_2^p(\Omega)} = ||(\mathcal{A}(g^2 \,\delta_{\Omega}^{-1})^{1/2}||_{L^p(\sigma)}.
$$

When  $s = n$ , we drop the subscript s from  $T_{2,s}^{\infty}$  and just write  $T_2^{\infty}$ . It is not hard to see that  $L_c^2(\Omega)$  is a dense subspace of  $T_2^p$  $\mathcal{D}_2^p(\Omega)$  for any  $p \in [1,\infty)$ .

The tent spaces were first introduced and studied in [\[CMS85\]](#page-69-18) in the upper-half space  $\mathbb{R}^{n+1}_+$  and was extended to  $AR(n)$  domains in  $[MPT13]^8$  $[MPT13]^8$ . An important result in this theory is the duality between tent spaces. Namely, if  $\Omega \in AR(n)$ , the pairing

$$
\langle f, g \rangle = \int_{\Omega} f(x) g(x) \frac{dx}{\delta_{\Omega}(x)}
$$

realizes  $T_2^{\infty}(\Omega)$  as the Banach dual of  $T_2^1(\Omega)$ . Moreover, for  $p \in (1,\infty)$ , the same pairing realizes  $T_2^{p'}$  $T_2^{p'}(\Omega)$  as the Banach dual of  $T_2^p$  $2^p_2(\Omega)$ , where  $1/p + 1/p' = 1$ . In this generality, this follows from the proof of Theorem 4.2 and Remarks 4.3 and 4.4 in [\[MPT13\]](#page-70-11). By an inspection of the proofs, one can easily show that if  $\Omega \in AR(s)$  for  $s \in (0, n]$ , then the pairing

$$
\langle f, g \rangle := \int_{\Omega} f(x) g(x) \frac{dx}{\delta_{\Omega}(x)^{n+1-s}}
$$

realizes  $T_{s,2}^{\infty}(\Omega)$  as the Banach dual of  $T_2^1(\Omega)$ . Analogously, for  $p \in (1,\infty)$ , the same pairing realizes  $T_2^{p'}$  $T_2^{p'}(\Omega)$  as the Banach dual of  $T_2^p$  $2^{p}(\Omega)$ , where  $1/p + 1/p' = 1$ .

<span id="page-14-0"></span>2.4.  $L^p$  and uniform  $\varepsilon$ -approximation. If  $\varepsilon > 0$ , we say that a function w is *uniformly ε*-*approximable*, if there exist a constant  $C_\varepsilon > 0$  and a function  $\varphi = \varphi^\varepsilon \in C^\infty(\Omega)$  such that

(2.27) 
$$
\sup_{x \in \Omega} |w(x) - \varphi(x)| + \sup_{x \in \Omega} \delta_{\Omega}(x) |\nabla(w - \varphi)(x)| \lesssim \varepsilon
$$

and

(2.28) 
$$
\sup_{\xi \in \partial \Omega} C_{s,c}(\nabla \varphi)(\xi) \lesssim \varepsilon^{-2},
$$

where  $\delta_{\Omega}(\cdot) = \text{dist}(x \cdot, \Omega^c)$  and the implicit constants are independent of  $\varepsilon$ .

<span id="page-14-1"></span><sup>&</sup>lt;sup>8</sup>Note that the results are stated in chord-arc domains but an easy inspection of the proofs in [\[MPT13\]](#page-70-11) reveals that neither the Harnack chain condition nor the exterior corkscrew condition are necessary.

If  $\Omega \in AR(s)$ , then, for fixed  $p \in (1,\infty)$ , we say that a function w is  $\varepsilon$ -*approximable in*  $L^p(\sigma_s)$  if there exists a function  $\varphi = \varphi^{\varepsilon} \in C^{\infty}(\Omega)$  such that

$$
(2.29) \qquad \|\mathcal{N}(w-\varphi)\|_{L^p(\sigma_s)} + \|\mathcal{N}(\delta_{\Omega}\nabla(w-\varphi))\|_{L^p(\sigma_s)} \lesssim_p \varepsilon \|\mathcal{N}w\|_{L^p(\sigma_s)}
$$

and

$$
(2.30) \t\t\t ||C_{s,c}(\nabla \varphi)||_{L^p(\sigma_s)} \lesssim_p \varepsilon^{-2} ||\mathcal{N}w||_{L^p(\sigma_s)}.
$$

<span id="page-15-0"></span>2.5. Elliptic systems and Boundary value problems. In this section we consider domains  $\Omega \in AR(n), n \geq 1$ . Let L be an elliptic operator acting on column vector-fields  $u =$  $(u^1, \ldots, u^m)^T$ , where  $u^{\beta} : \Omega \to \mathbb{C}$  for  $\beta = 1, 2, \ldots, m$ , defined as follows:

<span id="page-15-1"></span>(2.31) 
$$
Lu(x) = -\sum_{i,j=1}^{n+1} \partial_i (A_{ij}(x)\partial_j u(x)) = -\sum_{\alpha,\beta=1}^m \sum_{i,j=1}^{n+1} \partial_i (a_{ij}^{\alpha\beta}(x)\partial_j u^{\beta}(x)),
$$

where  $\partial_i = \frac{\partial}{\partial x}$  $\frac{\partial}{\partial x_i}$ ,  $1 \le i \le n+1$  and  $A_{ij}$  are  $m \times m$  matrix-valued functions on  $\mathbb{R}^{n+1}$  with entries  $a_{ij}^{\alpha\beta}:\Omega\to\mathbb{C}, \alpha,\beta\in\{1,\ldots,m\}$  for which there exists  $\lambda\in(0,1]$  such that

(2.32) 
$$
\sum_{\alpha,\beta=1}^m \sum_{i,j=1}^{n+1} |a_{ij}^{\alpha\beta}(x)|^2 \leq \lambda^{-2}, \text{ for a.e. } x \in \Omega \text{ and}
$$

<span id="page-15-2"></span>
$$
(2.33) \qquad \Re \epsilon \sum_{\alpha,\beta=1}^m \sum_{i,j=1}^{n+1} a_{ij}^{\alpha\beta}(x) \xi_j^{\beta} \overline{\xi_i^{\alpha}} \ge \lambda \sum_{\alpha=1}^m \sum_{i=1}^{n+1} |\xi_i^{\alpha}|^2 \quad \text{for a.e. } x \in \Omega.
$$

For  $m = 1$  and  $a_{ij} : \Omega \to \mathbb{R}$ , [\(2.33\)](#page-15-2) amounts to the standard accretivity condition

(2.34) 
$$
\lambda |\xi|^2 \leq \sum_{i,j=1}^{n+1} a_{ij}(x)\xi_i \dot{\xi}_j, \quad \text{for a.e. } x \in \Omega, \ \xi \in \mathbb{R}^{n+1}.
$$

Notice that the  $\alpha$ -th component of the column vector  $Lu$  coincides with

(2.35) 
$$
(Lu)^{\alpha}(x) := -\sum_{\beta=1}^{m} \sum_{i,j=1}^{n+1} \partial_i (a_{ij}^{\alpha\beta}(x) \partial_j u^{\beta}(x))
$$

We also define its adjoint operator of  $L$  by

$$
L^*u(x) = -\sum_{\alpha,\beta=1}^m \sum_{i,j=1}^{n+1} \partial_i \big( \overline{a_{ji}^{\beta\alpha}}(x) \partial_j u^{\beta}(x) \big) \big),
$$

i.e.,  $L^* = - \operatorname{div} A^* \nabla$ , where  $A^* = (\overline{A}_{ij})^T$  or equivalently  $(a_{ij}^{\alpha \beta})^* = a_{ji}^{\beta \alpha}$ .

We assume that  $H: \Omega \to \mathbb{C}^m$  is given by  $H = (H^1, \dots, H^m)$  and  $\Xi: \Omega \to \mathbb{C}^{m(n+1)}$  is given by  $\Xi := (\vec{\Xi}^1, \dots, \vec{\Xi}^m)$ , where  $\vec{\Xi}^{\alpha} : \Omega \to \mathbb{C}^{n+1}$  and  $\vec{\Xi}^{\alpha} = (\Xi_1^{\alpha}, \dots, \Xi_{n+1}^{\alpha})$  for  $\alpha =$ 1, . . . m. We are interested in solutions of the inhomogeneous equation  $Lu = -\text{div }\mathbf{E} + H$ in  $\Omega$  in the sense

$$
Lu(x) = -\sum_{\alpha=1}^{m} \sum_{i=1}^{n+1} \partial_i \Xi_i^{\alpha}(x) + \sum_{\alpha=1}^{m} H^{\alpha}(x), \text{ for } x \in \Omega.
$$

For  $H \in L^2_{loc}(\Omega; \mathbb{C}^m)$  and  $\Xi \in L^2_{loc}(\Omega; \mathbb{C}^{m(n+1)})$ , we say that the vector field  $w \in$  $W^{1,2}_{loc}(\Omega;\mathbb{C}^m)$  solves  $Lw = H - \text{div }\mathbf{\Xi}$  in the *weak sense*, or that w is a *weak solution* to the equation  $Lw = H - \text{div }\mathbf{\Xi}$ , if for any  $\Phi \in C_c^{\infty}(\Omega; \mathbb{C}^m)$ , we have that

<span id="page-16-4"></span>
$$
(2.36) \qquad \sum_{\alpha,\beta=1}^m \sum_{i,j=1}^{n+1} \int_{\Omega} a_{ij}^{\alpha\beta}(x) \partial_j w^{\beta} \overline{\partial_i \Phi^{\alpha}} = \sum_{\alpha=1}^m \sum_{i=1}^{n+1} \int_{\Omega} \Xi_i^{\alpha} \overline{\partial_i \Phi^{\alpha}} + \sum_{\alpha=1}^m \int_{\Omega} H^{\alpha} \overline{\Phi^{\alpha}}.
$$

We say that the *variational Poisson-Dirichlet problem* for L is solvable in Ω if for every  $H \in L^{2_*}(\Omega; \mathbb{C}^m)$  and  $\Xi \in L^2(\Omega; \mathbb{C}^{m(n+1)})$  there exists  $u \in W^{1,2}_{loc}(\Omega; \mathbb{C}^m)$  such that

<span id="page-16-1"></span>(2.37) 
$$
(PD_{v}^{L}) = \begin{cases} Lu = -\operatorname{div} A \nabla u = -\operatorname{div} \Xi + H & \text{weakly in } \Omega \\ u \in Y_{0}^{1,2}(\Omega). \end{cases}
$$

By Lax-Milgram's theorem, this problem is always solvable and its solution is unique. Moreover,

<span id="page-16-2"></span>
$$
\|u\|_{Y_0^{1,2}(\Omega)}\lesssim \|H\|_{L^{2*}(\Omega;{\mathbb C}^m)}+\|\Xi\|_{L^2(\Omega;{\mathbb C}^{m(n+1)})}.
$$

We say that the *variational Dirichlet problem* for L is solvable in  $\Omega$  if, for every  $\varphi \in$  $\mathrm{Lip}(\partial\Omega;\mathbb{C}^m)$  and  $\Phi \in \dot{W}^{1,2}(\Omega;\mathbb{C}^m) \cap \mathrm{Lip}(\overline{\Omega};\mathbb{C}^m)$  satisfying  $\Phi|_{\partial\Omega} = \varphi$ , there exists  $w \in \dot{W}^{1,2}(\Omega; \mathbb{C}^m)$  such that

(2.38) 
$$
(D_v^L) = \begin{cases} Lw = -\operatorname{div} A \nabla w = 0 & \text{weakly in } \Omega \\ w - \Phi \in Y_0^{1,2}(\Omega). & \text{on } \partial \Omega. \end{cases}
$$

If u is the solution of [\(2.37\)](#page-16-1) for  $\mathbf{\Xi} = -A\nabla\Phi \in L^2(\Omega; \mathbb{C}^{m(n+1)})$  and  $H = 0$ , then, it is easy to see that  $w = u + \Phi$  is the solution of [\(2.38\)](#page-16-2).

Since the set of compactly supported Lipschitz functions on  $\partial\Omega$ , is dense in the set of compactly supported continuous functions on  $\partial\Omega$ , we can extend the definition of the Dirichlet problem to  $C_c(\partial\Omega)$ . Namely, for any  $\varphi \in C_c(\partial\Omega; \mathbb{C}^m)$  and  $\Phi \in \dot{W}^{1,2}(\Omega; \mathbb{C}^m)$  $C(\overline{\Omega}; \mathbb{C}^m)$  satisfying  $\Phi|_{\partial \Omega} = \varphi$ , there exists  $w \in \dot{W}^{1,2}(\Omega; \mathbb{C}^m)$  satisfying [\(2.38\)](#page-16-2).

We set

$$
Y^q(\sigma) = \begin{cases} L^q(\sigma) & \text{if } q \in (1, \infty) \\ \text{BMO}(\sigma) & \text{if } q = \infty, \end{cases}
$$

<span id="page-16-0"></span>**Definition 2.6.** For any  $q \in (1, \infty)$ , we say that the *Dirichlet problem* with  $L^q$  boundary data is solvable for L in  $\Omega$  (write  $(D_q^L)$  is solvable in  $\Omega$ ), if there exists  $C > 0$  so that for each  $g \in \text{Lip}_c(\partial \Omega)$ , the solution u of [\(2.38\)](#page-16-2) for L with boundary data g satisfies the estimate

(2.39) 
$$
\|\widetilde{\mathcal{N}}_{2^*}(u)\|_{L^q(\sigma)} \leq C \|g\|_{L^q(\sigma)},
$$

where  $2^* := \frac{2(n-1)}{n+1}$ . Similarly, we will say that  $(\widetilde{D}_q^L)$  is solvable for L in  $\Omega$  if there exists  $C > 0$  so that for each  $g \in \text{Lip}_{c}(\partial \Omega)$ , the solution u of [\(2.38\)](#page-16-2) for L with boundary data g satisfies the estimate

<span id="page-16-3"></span>(2.40) kδΩ∇uk<sup>T</sup> q 2 (Ω) ≤ Ckgk<sup>Y</sup> <sup>q</sup>(σ) .

<span id="page-17-5"></span>**Definition 2.7.** For any  $p \in (0, \infty)$ , we say that the (homogeneous) *Dirichlet regularity problem* (or just *regularity problem*) with boundary data in  $\dot{M}^{1,p}(\sigma)$  is solvable for L in  $\Omega$ (write  $(R_p^L)$  is solvable in  $\Omega$ ), if there exists  $C > 0$  so that for each  $f \in \text{Lip}_c(\partial \Omega)$ , the solution  $u$  of [\(2.38\)](#page-16-2) with boundary data  $f$  satisfies the estimate

(2.41) 
$$
\|\mathcal{N}_2(\nabla u)\|_{L^p(\sigma)} \leq C \|f\|_{\dot{M}^{1,p}(\sigma)}.
$$

<span id="page-17-9"></span><span id="page-17-3"></span>Following [\[MPT22\]](#page-70-8), we introduce the Poisson-regularity problem with data in  $C_{q,p}(\Omega)$ .

**Definition 2.8.** For any  $p \in (1,\infty)$ , we say that the *Poisson-Dirichlet problem*  $(PD_p^L)$ is solvable in  $\Omega$  if there exists  $C > 0$  so that for each  $H \in L_c^{\infty}(\Omega; \mathbb{C}^m)$  and  $\Xi \in$  $L_c^{\infty}(\Omega; \mathbb{C}^{m(n+1)})$ , the solution v of the problem [\(2.37\)](#page-16-1) satisfies the estimate

<span id="page-17-8"></span>
$$
(2.42) \qquad \qquad \|\widetilde{\mathcal{N}}_{2^*}(u)\|_{L^p(\sigma)} \leq C \left( \|\mathcal{C}_{2_*}(\delta_{\Omega} H)\|_{L^p(\sigma)} + \|\mathcal{C}_2(\Xi)\|_{L^p(\sigma)} \right).
$$

Similarly, for  $p \in (1, \infty)$ , we say that the *Poisson-Dirichlet problem*  $(\widetilde{PD}_{p}^{L})$  for  $H = 0$  is solvable in  $\Omega$  if there exists  $C > 0$  so that for each  $\Xi \in L_c^{\infty}(\Omega; \mathbb{C}^{m(n+1)})$ , the solution u of the problem  $(2.37)$  for  $H = 0$  satisfies the estimate

<span id="page-17-7"></span>(2.43) kδΩ∇uk<sup>T</sup> p 2 (Ω) ≤ C kC2(Ξ)kLp(∂Ω).

<span id="page-17-4"></span>**Definition 2.9.** For any  $p \in (1, \infty)$ , we say that the *Poisson-regularity problem*  $(\text{PR}_{p}^{L})$ is solvable in  $\Omega$  if there exists  $C > 0$  so that for each  $H \in L_c^{\infty}(\Omega; \mathbb{C}^m)$  and  $\Xi \in$  $L_c^{\infty}(\Omega; \mathbb{C}^{m(n+1)})$ , the solution v of the problem [\(2.37\)](#page-16-1) satisfies the estimate

$$
(2.44) \qquad \qquad \|\widetilde{\mathcal{N}}_2(\nabla v)\|_{L^p(\sigma)} \leq C \left( \|\mathcal{C}_{2*}(H)\|_{L^p(\sigma)} + \|\mathcal{C}_2(|\boldsymbol{\Xi}|/\delta_{\Omega})\|_{L^p(\sigma)} \right).
$$

Similarly, for  $p \in [1, \infty)$ , we say that the *Poisson-regularity problem*  $(\widetilde{\text{PR}}_p^L)$  for  $H = 0$  is solvable in  $\Omega$  if there exists  $C > 0$  so that for each  $\Xi \in L^2_c(\Omega; \mathbb{C}^{m(n+1)})$ , the solution v of the problem  $(2.37)$  for  $H = 0$  satisfies the estimate

<span id="page-17-6"></span>
$$
(2.45) \t\t\t\t\|\widetilde{\mathcal{N}}_2(\nabla v)\|_{L^p(\sigma)} \leq C \|\Xi\|_{T_2^p(\Omega)}.
$$

<span id="page-17-0"></span>2.6. Geometry of domains. Following Jerison and Kenig  $JK82$ , we introduce the notions of corkscrew and Harnack chain conditions.

<span id="page-17-1"></span>**Definition 2.10.** We say that an open set  $\Omega \subset \mathbb{R}^{n+1}$  satisfies the *c-corkscrew condition* for  $c \in (0, 1/2)$ , if for every ball  $B(\xi, r)$  with  $\xi \in \partial\Omega$  and  $0 < r < \text{diam}(\Omega)$ , there exists a point  $x \in \Omega \cap B(\xi, r)$  such that  $B(x, cr) \subset \Omega \cap B(\xi, r)$ .

<span id="page-17-2"></span>**Definition 2.11.** Given two points  $x, x' \in \Omega$ , and a pair of numbers  $M, N \geq 1$ , an  $(M, N)$ -*Harnack Chain connecting* x *to* x', is a chain of open balls  $B_1, \ldots, B_N \subset \Omega$ , with  $x \in$  $B_1, x' \in B_N$ ,  $B_k \cap B_{k+1} \neq \emptyset$  and  $M^{-1}$  diam $(B_k) \leq$  dist $(B_k, \partial \Omega) \leq M$  diam $(B_k)$ . We say that Ω satisfies the *Harnack Chain condition* if there is a uniform constant M such that for any two points  $x, x' \in \Omega$ , there is an  $(M, N)$ -Harnack Chain connecting them, with N depending only on M and the ratio  $|x - x'| / (\min (\delta_{\Omega}(x), \delta_{\Omega}(x')))$ .

It is not hard to see that if  $E \subset \mathbb{R}^{n+1}$  is s-Ahlfors regular for  $s \in (0, n]$ , then  $\mathbb{R}^{n+1} \setminus E$ satisfies the c-corkscrew condition for some  $c \in (0, 1/2)$  depending only on the Ahlfors regularity constants. In the case that  $s < n$ ,  $\mathbb{R}^{n+1} \setminus E$  satisfies the Harnack chain condition as well (see [\[DFM21,](#page-69-19) Lemma 2.2]).

**Definition 2.12.** A connected rectifiable curve  $\gamma : [0, \ell] \to \overline{\Omega}$  connecting  $\xi \in \partial\Omega$  and  $x \in$  $Ω$ , parametrized by the arc-length  $s ∈ [0, ℓ]$  and such that  $γ(0) = ξ$  and  $γ(ℓ) = x$ , is called  $\lambda$ *-good curve* (or  $\lambda$ *-carrot path*), for some  $\lambda \in (0, 1]$ , if  $\gamma \setminus \{\xi\} \subset \Omega$  and  $\delta_{\Omega}(\gamma(s)) > \lambda s$ , for every  $s \in (0, \ell]$ .

<span id="page-18-0"></span>**Definition 2.13.** An open set  $\Omega \subset \mathbb{R}^{n+1}$  is said to satisfy the *pointwise John condition*, if there exists a constant  $\theta \in (0,1)$  such that for  $\sigma_n$ -a.e.  $\xi \in \partial\Omega$ , there exist  $x_{\xi} \in \Omega$  and  $r_{\xi} > 0$  such that  $x_{\xi} \in B(\xi, 2r_{\xi})$  and  $\delta_{\Omega}(x_{\xi}) \geq \theta r_{\xi}$ , and also there exists a  $\theta$ -good curve  $\gamma_{\xi} \subset \Omega \cap B(\xi, 2r_{\xi})$  connecting the points  $\xi$  and  $x_{\xi}$  such that  $\ell(\gamma_{\xi}) \leq \theta^{-1}r_{\xi}$ . We will write that  $\xi \in JC(\theta)$  if the pointwise John condition holds for the point  $\xi$  with constant  $\theta \in (0,1)$ .

**Remark 2.14.** Any domain  $\Omega \in \text{AR}(n)$  with *n*-rectifable boundary satisfies the pointwise John condition.

Following [\[HMT10\]](#page-69-20) we also introduce the notion of local John domains, which are also examples of domains satisfying the pointwise John condition.

<span id="page-18-1"></span>**Definition 2.15.** An open set  $\Omega \subset \mathbb{R}^{n+1}$  is said to satisfy the *local John condition* if there is  $\theta \in (0, 1)$  such that the following holds: For all  $x \in \partial\Omega$  and  $r \in (0, 2 \text{diam}(\Omega))$  there is  $y \in B(x,r) \cap \Omega$  such that  $B(y,\theta r) \subset \Omega$  with the property that for all  $z \in B(x,r) \cap \partial \Omega$ one can find a rectifiable path  $\gamma_z: [0,1] \to \overline{\Omega}$  with length at most  $\theta^{-1}|x-y|$  such that

 $\gamma_z(0) = z$ ,  $\gamma_z(1) = y$ , dist $(\gamma_z(t), \partial \Omega) \ge \theta |\gamma_z(t) - z|$  for all  $t \in [0, 1]$ .

If  $\Omega \in AR(s)$  for  $0 \leq s \leq n$ , then it clearly satisfies the local John condition as it satisfies the corkscrew and the Harnack chain conditions. If  $s = n$ , any semi-uniform (and thus any uniform) domain has the local John condition.

<span id="page-18-2"></span>**Definition 2.16.** Let  $\Omega$  be a corkscrew domain,  $F : \Omega \to \mathbb{R}$ , and  $f : \partial \Omega \to \mathbb{R}$ . We say that F *converges non-tangentially* to f at  $\xi \in \partial\Omega$  (write  $F \to f$  n.t. at  $\xi$ ) if there exists  $\alpha > 0$  such that for every sequence  $x_k \in \gamma_\alpha(\xi)$  for which  $x_k \to \xi$  as  $k \to \infty$ , it holds that  $F(x_k) \to f(\xi)$  as  $k \to \infty$ . We will also write

$$
\operatorname*{nt-lim}_{x \to \xi} F(x) = f(\xi).
$$

We will say that  $F \to f$  *quasi-non-tangentially* at  $\xi \in \partial \Omega$  (write  $F \to f$  q.n.t. at  $\xi$ ) if there exist  $r_{\xi} > 0$ , a corkscrew point  $x_{\xi} \in \Omega \cap B(\xi, 2r_{\xi})$ , and a  $\theta$ -good curve  $\gamma_{\xi} \in B(\xi, 2r_{\xi})$ connecting  $\xi$  and  $x_{\xi}$ , such that for any  $x_k \in \gamma_{\xi}$  converging to  $\xi$  as  $k \to \infty$ , it holds that  $\lim_{k\to\infty} F(x_k) = f(\xi)$ . We will also write

$$
\operatorname*{qrt-lim}_{x \to \xi} F(x) = f(\xi).
$$

<span id="page-19-0"></span>2.7. Dyadic lattices. Given an s-Ahlfors-regular measure  $\mu$  in  $\mathbb{R}^{n+1}$ , we consider the dyadic lattice of "cubes" built by David and Semmes in [\[DS93,](#page-69-10) Chapter 3 of Part I]. The properties satisfied by  $\mathcal{D}_{\mu}$  are the following. Assume first, for simplicity, that diam(supp  $\mu$ ) =  $\infty$ ). Then for each  $j \in \mathbb{Z}$  there exists a family  $\mathcal{D}_{\mu,j}$  of Borel subsets of supp  $\mu$  (the dyadic cubes of the  $j$ -th generation) such that:

- (a) each  $\mathcal{D}_{\mu,j}$  is a partition of supp  $\mu$ , i.e. supp  $\mu = \bigcup_{Q \in \mathcal{D}_{\mu,j}} Q$  and  $Q \cap Q' = \emptyset$ whenever  $Q, Q' \in \mathcal{D}_{\mu,j}$  and  $Q \neq Q'$ ;
- (b) if  $Q \in \mathcal{D}_{\mu,j}$  and  $Q' \in \mathcal{D}_{\mu,k}$  with  $k \leq j$ , then either  $Q \subset Q'$  or  $Q \cap Q' = \emptyset$ ;
- (c) for all  $j \in \mathbb{Z}$  and  $Q \in \mathcal{D}_{\mu,j}$ , we have  $2^{-j} \lesssim \text{diam}(Q) \le 2^{-j}$  and  $\mu(Q) \approx 2^{-js}$ ;
- (d) there exists  $C > 0$  such that, for all  $j \in \mathbb{Z}$ ,  $Q \in \mathcal{D}_{\mu,j}$ , and  $0 < \tau < 1$ ,

<span id="page-19-2"></span>
$$
(2.46) \qquad \mu\big(\big\{x \in Q : \text{dist}(x, \text{supp}\,\mu \setminus Q) \leq \tau 2^{-j}\big\}\big) \\qquad \qquad + \mu\big(\big\{x \in \text{supp}\,\mu \setminus Q : \, \text{dist}(x, Q) \leq \tau 2^{-j}\big\}\big) \leq C\tau^{1/C}2^{-js}.
$$

This property is usually called the *small boundaries condition*. From [\(2.46\)](#page-19-2), it follows that there is a point  $x_Q \in Q$  (the center of Q) such that  $dist(x_Q, \text{supp }\mu \setminus \mathcal{Q})$  $Q \geq 2^{-j}$  (see [\[DS93,](#page-69-10) Lemma 3.5 of Part I]).

We set

$$
\mathcal{D}_\mu := \bigcup_{j\in \mathbb{Z}} \mathcal{D}_{\mu,j}.
$$

In case that diam(supp  $\mu$ ) <  $\infty$ , the families  $\mathcal{D}_{\mu,j}$  are only defined for  $j \geq j_0$ , with  $2^{-j_0} \approx \text{diam}(\text{supp}\,\mu)$ , and the same properties above hold for  $\mathcal{D}_{\mu} := \bigcup_{j \ge j_0} \mathcal{D}_{\mu,j}$ .

Given a cube  $Q \in \mathcal{D}_{\mu,j}$ , we say that its *side-length* is  $2^{-j}$ , and we denote it by  $\ell(Q)$ . Notice that  $\text{diam}(Q) \leq \ell(Q)$ . We also denote

(2.47) 
$$
B(Q) := B(x_Q, c_1 \ell(Q)), \qquad B_Q = B(x_Q, \ell(Q)),
$$

where  $c_1 > 0$  is some fix constant so that  $B(Q) \cap \text{supp }\mu \subset Q$ , for all  $Q \in \mathcal{D}_{\mu}$ . Clearly, we have  $Q \subset B_Q$ . For  $\lambda > 1$ , we write

$$
\lambda Q = \{ x \in \mathrm{supp}\,\mu : \, \mathrm{dist}(x, Q) \le (\lambda - 1) \,\ell(Q) \}.
$$

The side-length of a "true cube"  $P \subset \mathbb{R}^{n+1}$  is also denoted by  $\ell(P)$ . On the other hand, given a ball  $B \subset \mathbb{R}^{n+1}$ , its radius is denoted by  $r(B)$ . For  $\lambda > 0$ , the ball  $\lambda B$  is the ball concentric with B with radius  $\lambda r(B)$ .

<span id="page-19-1"></span>2.8. The Whitney decomposition. Recall that a domain is a connected open set. In the whole paper,  $\Omega$  will be an open set in  $\mathbb{R}^{n+1}$ , with  $n \geq 1$ . We will denote the *n*-Hausdorff measure on  $\partial\Omega$  by  $\sigma$ .

We consider the following Whitney decomposition of  $\Omega$  (assuming  $\Omega \neq \mathbb{R}^{n+1}$ ): we have a family  $W(\Omega)$  of dyadic cubes in  $\mathbb{R}^{n+1}$  with disjoint interiors such that

$$
\bigcup_{P \in \mathcal{W}(\Omega)} P = \Omega,
$$

and moreover there are some constants  $\Lambda > 20$  and  $D_0 \ge 1$  such the following holds for every  $P \in \mathcal{W}(\Omega)$ :

(i)  $10P \subset \Omega$ ; (ii)  $\Lambda P \cap \partial \Omega \neq \emptyset$ ; (iii) there are at most  $D_0$  cubes  $P' \in W(\Omega)$  such that  $10P \cap 10P' \neq \emptyset$ . Further, for such cubes P', we have  $\frac{1}{2}\ell(P') \leq \ell(P) \leq 2\ell(P')$ .

From the properties (i) and (ii) it is clear that  $dist(P, \partial \Omega) \approx \ell(P)$  and so there exists  $\Lambda' > 20$ such that

<span id="page-20-3"></span>(2.48) 
$$
\text{dist}(x, \partial \Omega) \le \Lambda' \ell(P), \quad \text{for every } x \in P.
$$

We assume that the Whitney cubes are small enough so that

(2.49) 
$$
\text{diam}(P) < \frac{1}{20} \text{dist}(P, \partial \Omega).
$$

The arguments to construct a Whitney decomposition satisfying the properties above are standard and can be found for example in [\[ST70\]](#page-70-13).

Suppose that  $\partial\Omega$  is s-Ahlfors-regular and consider the dyadic lattice  $\mathcal{D}_{\sigma}$  defined above. Then, for each Whitney  $P \in \mathcal{W}(\Omega)$  there is some cube  $Q \in \mathcal{D}_{\sigma}$  such that

<span id="page-20-1"></span>(2.50) ℓ(Q) = ℓ(P) and dist(P, Q) ≈ ℓ(Q),

with the implicit constant depending on the parameters of  $\mathcal{D}_{\sigma}$  and on the Whitney decomposition. For every  $P \in \mathcal{W}(\Omega)$  there is a uniformly bounded number of cubes  $Q \in \mathcal{D}_{\sigma}$ depending on *n* and the s-Ahlfors regularity of  $\partial\Omega$  that satisfy [\(2.50\)](#page-20-1). To each  $P \in \mathcal{W}(\Omega)$ we associate precisely on such cube  $Q \in \mathcal{D}_{\sigma}$  and denote it by  $Q = b(P)$ . We say that Q is the *boundary cube* of P.

Conversely, given  $Q \in \mathcal{D}_{\sigma}$ , we let

(2.51) 
$$
w(Q) = \bigcup_{P \in \mathcal{W}(\Omega): Q=b(P)} P.
$$

In the case of *n*-Ahlfors regular boundary, it is immediate to check that  $w(Q)$  is made up at most of a uniformly bounded number of cubes P, but it may happen that  $w(Q) = \emptyset$ .

In higher co-dimensions where  $s < n$  it is also true that for every boundary cube  $Q \in \partial \Omega$ , there exist a uniformly bounded number of Whitney cubes  $P \in \mathcal{W}(\Omega)$  such that  $b(P) = Q$ . For the proof of this fact one can see [\[MaPo21,](#page-70-14) Lemma 4.16, Lemma 4.18].

We also denote the "fattened" Whitney region of  $Q$  by

(2.52) 
$$
\widetilde{w}(Q) = \bigcup_{P \in \mathcal{W}(\Omega): Q = b(P)} 1.1P.
$$

<span id="page-20-2"></span>**Remark 2.17.** If  $x \in \overline{P} \in \mathcal{W}(\Omega)$ , then there exists a constant  $C_w > 1$  depending only on  $n$ , the constants of the Whitney decomposition, and the  $s$ -Alhlfors regularity, so that for every  $P' \in \mathcal{W}(\Omega)$  that has the property  $x \in 1.1P'$ , it holds that

$$
b(P') \subset B(x_{b(P)}, C_w \ell(P)) =: B_P.
$$

#### 3. REGULARIZED DYADIC EXTENSION OF FUNCTIONS ON THE BOUNDARY

<span id="page-20-0"></span>Let  $\Omega \subset \mathbb{R}^{n+1}$ ,  $n \geq 1$ , be an open set and let  $\mathcal{W}(\Omega)$  be the collection of Whitney cubes in which  $\Omega$  is decomposed as in Subsection [2.8.](#page-19-1) Let  $\{\varphi_P\}_{P \in \mathcal{W}(\Omega)}$  be a partition of unity subordinate to the open cover  $\{1.1P\}_{P \in \mathcal{W}(\Omega)}$  such that

$$
\varphi_P \in C^{\infty}(\mathbb{R}^n), \ |\nabla \varphi_P| \lesssim \frac{1}{\ell(P)}, \ \text{supp}\,\varphi \subseteq 1.1P, \text{ and } \sum_{P \in \mathcal{W}(\Omega)} \varphi_P(x) = \mathbf{1}_{\Omega}(x), x \in \Omega.
$$

For  $f \in L^1_{loc}(\sigma)$ , we define the *regularized dyadic extension of* f in  $\Omega$  by

<span id="page-21-0"></span>(3.1) 
$$
v_f(x) := \begin{cases} \sum_{P \in \mathcal{W}(\Omega)} m_{\sigma, b(P)} f \varphi_P(x) & \text{if } x \in \Omega \\ f(x) & \text{if } x \in \partial \Omega. \end{cases}
$$

where, in the case that  $\Omega$  is an unbounded domain with compact boundary, we set  $b(P)$  =  $\partial\Omega$  for every  $P \in \mathcal{W}(\Omega)$  with  $\ell(P) \geq \text{diam}(\partial\Omega)$ .

The fact that  $v_f$  is indeed an extension of f in  $\Omega$  (in the non-tangential sense) is proved in the following lemma.

<span id="page-21-2"></span>**Lemma 3.1.** Let 
$$
\Omega \in \text{AR}(s)
$$
 for  $s \in (0, n]$  and  $f \in L^1_{loc}(\sigma_s)$ . There exists  $\alpha > 0$  such that 
$$
\underset{x \to \xi}{\text{nt-lim}} v_f(x) = f(\xi), \quad \text{for } \sigma\text{-a.e. } \xi \in \partial\Omega,
$$

*for some cone*  $\gamma_{\alpha}$  *where*  $\alpha > 0$  *only depends on n, the Ahlfors regularity constant, and the constants of the corkscrew condition.*

*Proof.* By [\[EG15,](#page-69-21) Theorem 1.33], it holds that

(3.2) 
$$
\lim_{r \to 0} m_{\sigma, B(\xi, r)}(|f - f(\xi)|) = 0, \text{for } \sigma\text{-a.e. } \xi \in \partial\Omega.
$$

Fix  $\varepsilon > 0$  and  $\xi \in \partial \Omega$  be a point such that [\(3.2\)](#page-21-1) holds. Let  $0 < \varepsilon' < \varepsilon$  to be picked later. Then there exists  $\delta = \delta(\varepsilon', \xi) > 0$  such that  $m_{\sigma, B(\xi, r)}(|f - f(\xi)|) < \varepsilon'$  for every  $r < \delta$ . Let now  $0 < \delta' < \delta$  be a small constant which will be chosen momentarily. For fixed  $x \in \gamma_\alpha(\xi) \cap B(\xi, \delta')$ , we have that

<span id="page-21-1"></span>
$$
|v_f(x) - f(\xi)| \le \sum_{P \in \mathcal{W}(\Omega)} |m_{\sigma, b(P)}f - f(\xi)| \varphi_P(x).
$$

Let  $P_0 \in \mathcal{W}(\Omega)$  be a fixed cube so that  $x \in \overline{P}_0$ . Then the only cubes  $P \in \mathcal{W}(\Omega)$  that contribute to the sum are the ones that  $x \in 1.1P$  and, by Remark [2.17,](#page-20-2)  $b(P) \subset B_{P_0}$  $B(x_{P_0}, C_w \ell(P_0))$ . By the properties of Whitney cubes, there exists a constant  $C'_w > C_w$ such that  $B_P \subset B(\xi, C'_w \ell(P))$ . Therefore, choosing  $\delta' > 0$  sufficiently small so that  $B(\xi, C'_w \ell(P)) \subset B(\xi, \delta/2)$ , we get that for any such P,

$$
|m_{\sigma,b(P)}f-f(\xi)|\lesssim m_{\sigma,B_{P_0}}(|f-f(\xi)|)\lesssim \varepsilon',
$$

which, in turn, by the bounded overlaps of the Whitney cubes, infers that there exists a constant  $C > 1$  such that

$$
|v_f(x) - f(\xi)| < C \,\varepsilon',
$$

concluding the proof of the lemma once we choose  $\varepsilon = C \varepsilon'$ .

. — Первый процесс в после производите в собстании и производите в собстании и производите в собстании и произ<br>В собстании и производите в собстании и производите в собстании и производите в собстании и производите в собс

<span id="page-22-4"></span>**Lemma 3.2.** Let  $\Omega \in AR(s)$  for  $s \in (0, n]$ . Assume that  $f \in L^1_{loc}(\sigma)$  and  $v_f$  is the *extension defined in* [\(3.1\)](#page-21-0)*. For any*  $P \in \mathcal{W}(\Omega)$ *, we have that* 

<span id="page-22-0"></span>(3.3) 
$$
\sup_{x \in \overline{P}} |\nabla v_f(x)| \lesssim \ell(P)^{-1} m_{\sigma, B_P}(|f|).
$$

*If, additionally,*  $f \in \Lambda_{\beta}(\partial \Omega)$  *for*  $\beta \in [0,1)$ *, then it holds that* 

<span id="page-22-1"></span>(3.4) 
$$
\sup_{x \in \overline{P}} |\nabla v_f(x)| \lesssim \ell(P)^{\beta - 1} ||f||_{\Lambda_\beta(\partial\Omega)},
$$

*while, if*  $f \in \dot{M}^{1,p}(\sigma)$ *, we get that* 

$$
|\nabla v_f(x)| \lesssim m_{\sigma, B_P}(\nabla_{H, p} f),
$$

*where*  $\nabla_{H,p} f$  *is the least Hajłasz upper gradient of f. Moreover, for any*  $\xi \in \partial \Omega$ *,* 

$$
\mathcal{N}_{\alpha}(v_f)(\xi) \lesssim_{\alpha} \mathcal{M}f(\xi),
$$

*Proof.* Fix  $x \in \overline{P} \in \mathcal{W}(\Omega)$ . For the proof of the estimate [\(3.3\)](#page-22-0), just note that

<span id="page-22-3"></span><span id="page-22-2"></span>
$$
|\nabla v_f(x)| \lesssim \sum_{\substack{P' \in \mathcal{W}(\Omega) \\ x \in 1.1P'}} m_{\sigma, b(P')}(|f|) \,\ell(P')^{-1} \lesssim_{D_0} m_{\sigma, B_P}(|f|) \,\ell(P)^{-1},
$$

where we used that  $\ell(P) \approx \ell(P')$ . If, in addition,  $f \in \Lambda_{\beta}(\partial\Omega)$ , then, using the fact that  $\nabla \big( \sum_{P \in \mathcal{W}(\Omega)} \varphi_P(x) \big) = 0$ , we get

$$
|\nabla v_f(x)| = \Big| \sum_{P' \in \mathcal{W}(\Omega)} (m_{\sigma,b(P')}f - m_{\sigma,b(P)}f) \nabla \varphi_{P'}(x) \Big|
$$
  
\$\lesssim \ell(P)^{-1} \sum\_{P' \in \mathcal{W}(\Omega) \atop x \in 1.1P'} |m\_{\sigma,b(P')}f - m\_{\sigma,b(P)}f|\$  
\$\lesssim \ell(P)^{-1} \sum\_{P' \in \mathcal{W}(\Omega) \atop x \in 1.1P'} m\_{\sigma,b(P')}(|f - m\_{\sigma,B\_P}f|) \lesssim\_{D\_0} ||f||\_{\Lambda\_\beta(\partial\Omega)} \ell(P)^{\beta-1}\$.

This obviously implies estimate  $(3.4)$ .

Now, in order to prove [\(3.5\)](#page-22-2), let  $f \in \dot{M}^{1,p}(\sigma)$  and  $x \in \overline{P}$ . Then we have

$$
|\nabla v_f(x)| = \left| \sum_{P' \in \mathcal{W}(\Omega)} m_{\sigma,b(P')} f \nabla \varphi_{P'}(x) \right| \lesssim \sum_{\substack{P' \in \mathcal{W}(\Omega) \\ x \in 1.1P'}} \frac{1}{\ell(P')} |m_{\sigma,b(P')} f - m_{\sigma,b(P)} f|
$$
  

$$
\lesssim \sum_{\substack{P' \in \mathcal{W}(\Omega) \\ x \in 1.1P'}} \frac{1}{\ell(P')} \int_{b(P')} \int_{b(P)} |x - y| [\nabla_{H,p} f(x) + \nabla_{H,p} f(y)] d\sigma(x) d\sigma(y)
$$
  

$$
\lesssim \sum_{\substack{P' \in \mathcal{W}(\Omega) \\ x \in 1.1P'}} (m_{\sigma,b(P')} (\nabla_{H,p} f) + m_{\sigma,b(P)} (\nabla_{H,p} f)) \lesssim_{D_0} m_{\sigma,B_P} (\nabla_{H,p} f),
$$

where  $B_P$  was defined in Remark [2.17.](#page-20-2)

Finally, fox fixed  $\xi \in \partial \Omega$ , if  $x \in \gamma_{\alpha}(\xi)$ , we have

$$
|v_f(x)| \leq \sum_{\substack{P' \in \mathcal{W}(\Omega) \\ x \in 1.1P'}} m_{\sigma,b(P)}(|f|) \varphi_P(x) \lesssim m_{\sigma,B_P}(|f|) \lesssim m_{\sigma,B(\xi,C'_w\ell(P))}(|f|) \leq \mathcal{M}f(\xi),
$$

for some constant  $C'_w > C_w$  depending on  $\alpha$  and the Whitney constants. This readily proves [\(3.6\)](#page-22-3) by taking supremum over all  $x \in \gamma_\alpha(\xi)$ .

<span id="page-23-2"></span>**Lemma 3.3.** *If*  $f \in \Lambda_{\beta}(\partial \Omega)$  *for*  $\beta \in [0,1)$ *, then it holds that* 

(3.7) 
$$
\sup_{\xi \in \partial \Omega} \mathcal{N}_{\sharp}^{(\beta)}(\nu_f)(\xi) \lesssim \|f\|_{\Lambda_{\beta}(\partial \Omega)}.
$$

*Proof.* Fix  $\xi \in \partial\Omega$  and take  $x \in \gamma_\alpha(\xi)$  with  $x \in \overline{P}_0$ , for some  $P_0 \in \mathcal{W}(\Omega)$ . It is enough to bound the difference

<span id="page-23-0"></span>
$$
\sup_{y \in B(x,c\delta_{\Omega}(x))} \int_{B(x,c\delta_{\Omega}(x))} |v_{f}(y) - v_{f}(z)| dz.
$$

To this end, fix a point  $z \in B(x, c\delta_{\Omega}(x))$ , with  $z \in \overline{P}_1 \in \mathcal{W}(\Omega)$  and a point  $y \in$  $B(x, c\delta_{\Omega}(x))$  with  $y \in \overline{P}_2 \in \mathcal{W}(\Omega)$ . Since  $c > 0$  is small enough, the Whitney cubes  $P_1, P_2$  and  $P_0$  are "close-enough" to each other, in the sense that the intersection of a dilation of these cubes is non-empty. Then, by the properties of Whitney cubes, we get that  $\ell(P_1) \approx \ell(P_2) \approx \ell(P_0) \approx \delta_{\Omega}(x)$ . Thus, there exist a large enough constant  $\Lambda_0 > 1$ , such that for any  $P \in \mathcal{W}(\Omega)$  with  $1.1P \cap (P_1 \cup P_2) \neq \emptyset$ , we have that

$$
B_{b(P)} \subset B_0 := B(x_{b(P_0)}, \Lambda_0 \ell(P_0)).
$$

It holds that

$$
\sum_{\substack{P \in \mathcal{W}(\Omega) \\ P \cap B(x,c\delta_{\Omega}(x)) \neq \emptyset}} (\varphi_P(z) - \varphi_P(y)) m_{\sigma,B_0} f = 0,
$$

and so

$$
|v_f(y) - v_f(z)| \leq \sum_{\substack{P \in \mathcal{W}(\Omega) \\ P \cap B(x, c\delta_{\Omega}(x)) \neq \emptyset}} |\varphi_P(y) - \varphi_P(z)| |m_{\sigma, b(P)} f - m_{\sigma, B_0} f|
$$
  

$$
\lesssim \sum_{\substack{P \in \mathcal{W}(\Omega) \\ P \cap B(x, c\delta_{\Omega}(x)) \neq \emptyset}} ||\nabla \varphi_P||_{L^{\infty}} |y - z||m_{\sigma, b(P)} f - m_{\sigma, B_0} f|
$$
  

$$
\lesssim \sum_{\substack{P \in \mathcal{W}(\Omega) \\ P \cap B(x, c\delta_{\Omega}(x)) \neq \emptyset}} m_{\sigma, b(P)} (|f - m_{\sigma, B_0} f|) \lesssim ||f||_{\Lambda_{\beta}(\partial \Omega)} \ell(P_0)^{\beta},
$$

since there are uniformly bounded many Whitney cubes  $P \cap B(x, c \delta_{\Omega}(x)) \neq \emptyset$ . This readily implies  $(3.7)$ .

<span id="page-23-3"></span><span id="page-23-1"></span>**Lemma 3.4.** *Let*  $\Omega \in \text{AR}(s)$  *for*  $s \in (0, n]$ *. If*  $f \in \Lambda_{\beta}(\partial \Omega)$  *for*  $\beta \in (0, 1)$ *, then* (3.8) sup ξ∈∂Ω  $\mathcal{C}^{(\beta)}_s(\nabla v_f)(\xi) \lesssim \|f\|_{\Lambda_\beta(\partial\Omega)}.$ 

*Proof.* By [\(3.4\)](#page-22-1), it is easy to see that for every  $\xi \in \partial \Omega$  and  $r > 0$ , it holds that

$$
\int_{B(\xi,r)\cap\Omega} |\nabla v_f| \,\omega_s(x) \,dx \lesssim \|f\|_{\Lambda_\beta(\partial\Omega)} \sum_{\substack{P \in \mathcal{W}(\Omega) \\ P \cap B(\xi,r) \neq \emptyset}} \ell(P)^{\beta-1} \ell(P)^{s+1}
$$
\n
$$
\lesssim \|f\|_{\Lambda_\beta(\partial\Omega)} \sum_{\substack{P \in \mathcal{W}(\Omega) \\ P \cap B(\xi,r) \neq \emptyset}} \ell(b(P))^{\beta} \sigma(b(P))
$$
\n
$$
\lesssim \|f\|_{\Lambda_\beta(\partial\Omega)} \sum_{k=0}^{\infty} 2^{-\beta k} r^{\beta} \sum_{\substack{Q \in \mathcal{D}_{k,\sigma} \\ Q \subset B(\xi, M_0 r)}} \sigma(Q) \lesssim \|f\|_{\Lambda_\beta(\partial\Omega)} r^{\beta} r^s,
$$

which implies  $(3.8)$ .

If the boundary function is in  $\text{Lip}_{\beta}(\partial \Omega)$  for  $\beta \in (0,1]$ , then we can show that  $v_f$  is  $\text{Lip}_{\beta}(\Omega)$  by arguments similar to the ones in [\[MT22,](#page-70-7) Lemma 4.2].

<span id="page-24-0"></span>**Lemma 3.5.** If  $f \in \text{Lip}_{\beta}(\partial \Omega)$  for  $\beta \in (0,1]$  and  $v_f$  is the regularized dyadic extension defined in [\(3.1\)](#page-21-0), then it holds that  $v_f\in \mathrm{Lip}_\beta(\Omega)$  with  $\mathrm{Lip}_\beta(v_f)\lesssim \mathrm{Lip}_\beta(f).$ 

*Proof.* We start by proving that  $v_f \in \text{Lip}_{\beta}(\Omega)$ . Fix  $x, y \in \Omega$  and let  $P_1, P_2 \in \mathcal{W}(\Omega)$  such that  $x \in \overline{P}_1$  and  $y \in \overline{P}_2$ . We split into cases.

**Case 1:** Suppose that  $2P_1 \cap 2P_2 \neq \emptyset$ . In this case let  $P_0 \in \mathcal{W}(\Omega)$  be the smallest cube such that it holds

$$
2B_{b(P)} \subset 2B_{b(P_0)}
$$
, for every  $P \in \mathcal{W}(\Omega)$ , with  $1.1P \cap (P_1 \cup P_2) \neq \emptyset$ .

Then, by the properties of Whitney cubes, we get that  $\ell(P_1) \approx \ell(P_2) \approx \ell(P_0)$ . As

$$
v_f(x) - v_f(y) = \sum_{P \in \mathcal{W}(\Omega)} m_{\sigma, b(P)} f(\varphi_P(x) - \varphi_P(y))
$$

and

$$
\sum_{P \in \mathcal{W}(\Omega)} (\varphi_P(x) - \varphi_P(y)) m_{\sigma, b(P_0)} f = 0,
$$

we can write

$$
v_f(x) - v_f(y)
$$
  
= 
$$
\sum_{P \in \mathcal{W}(\Omega)} m_{\sigma,b(P)} f(\varphi_P(x) - \varphi_P(y)) - \sum_{P \in \mathcal{W}(\Omega)} m_{\sigma,b(P_0)} f(\varphi_P(x) - \varphi_P(y))
$$
  
= 
$$
\sum_{P \in \mathcal{W}(\Omega)} (\varphi_P(x) - \varphi_P(y)) (m_{\sigma,b(P)}) f - m_{\sigma,b(P_0)} f)
$$

and thus we get

$$
|v_f(x) - v_f(y)| \leq \sum_{P \in \mathcal{W}(\Omega)} |\varphi_P(x) - \varphi_P(y)| \left| m_{\sigma, b(P)} f - m_{\sigma, b(P_0)} f \right|.
$$

Observe now that

$$
|\varphi_P(x) - \varphi_P(y)| \le |\nabla \varphi_P||x - y| \lesssim \ell(P)^{-1}|x - y|
$$

while, for fixed  $w \in 2B_{b(P_0)}$ , we can estimate

$$
\left| m_{\sigma,b(P)} f - m_{\sigma,b(P_0)} f \right| \leq m_{\sigma,b(P)} (|f(z) - f(w)|) + m_{\sigma,b(P_0)} (|f(z) - f(w)|)
$$
  

$$
\lesssim \mathrm{Lip}_{\beta}(f) \, \ell(P_0)^{\beta}.
$$

To deal with the sum, we may assume that the cubes  $P$  appearing in the sum are such that either  $1.1P \cap P_1 \neq \emptyset$  or  $1.1P \cap P_2 \neq \emptyset$ , since otherwise the associated summand vanishes. We denote by  $I_0$  the family of such cubes. So the cubes from  $I_0$  are such that  $B_{b(P)} \subset 2B_{b(P_0)}$  and they satisfy  $\ell(P) \approx \ell(P_0)$ . Combining this observation with the last two estimates and the fact that  $|x - y| \lesssim \ell(P_0)$ , we obtain

$$
(3.9) \quad |v_f(x) - v_f(y)| \lesssim \sum_{P \in \mathcal{W}(\Omega)} \frac{1}{\ell(P)} |x - y| \text{Lip}_{\beta}(f) \, \ell(P_0)^{\beta} \lesssim_{D_0} \text{Lip}_{\beta}(f) \, |x - y|^{\beta}.
$$

**Case 2:** Suppose that  $2P_1 \cap 2P_2 = \emptyset$ . In this case we have

$$
v_f(x) - v_f(y) = \sum_{P \in \mathcal{W}(\Omega)} m_{\sigma, b(P)} f(\varphi_P(x) - \varphi_P(y))
$$
  
= 
$$
\sum_{P \in \mathcal{W}(\Omega)} \varphi_P(x) (m_{\sigma, b(P)} f - m_{\sigma, b(P_1)} f) + \sum_{P \in \mathcal{W}(\Omega)} \varphi_P(y) (m_{\sigma, b(P_2)} f - m_{\sigma, b(P)}) f
$$
  
+ 
$$
(m_{\sigma, b(P_1)} f - m_{\sigma, b(P_2)} f) =: S_1 + S_2 + S_3.
$$

If we use that  $\ell(P_i) \lesssim |x - y|$  and the fact that  $|m_{\sigma,b(P)}f - m_{\sigma,b(P_i)}f| \lesssim \mathrm{Lip}_\beta(f)\,\ell(P_i)^\beta$ , for  $i = 1, 2$ , we can show that

$$
|S_1|+|S_2|\lesssim \mathrm{Lip}_\beta(f)|x-y|^\beta.
$$

It remains to bound  $S_3$ . If  $w_1 \in b(P_1)$  and  $w_2 \in b(P_2)$ ,

$$
|S_3| \le |m_{\sigma,b(P_1)}f - f(w_1)| + |f(w_1) - f(w_2)| + |f(w_2) - m_{\sigma,b(P_2)}f|.
$$

Since  $f \in \text{Lip}_{\beta}(\partial \Omega)$ ,

$$
|f(w_1) - f(w_2)| \le \mathrm{Lip}_{\beta}(f) |w_1 - w_2|^{\beta} \le \mathrm{Lip}_{\beta}(f) (|w_1 - x|^{\beta} + |x - y|^{\beta} + |y - w_2|^{\beta})
$$
  

$$
\lesssim \mathrm{Lip}_{\beta}(f) (\ell(P_1)^{\beta} + |x - y|^{\beta} + \ell(P_2)^{\beta}) \lesssim \mathrm{Lip}_{\beta}(f) |x - y|^{\beta},
$$

while, for  $i = 1, 2$ , once again using that  $f \in \text{Lip}_{\beta}(\partial \Omega)$ , it is easy to see that

$$
\left| m_{\sigma,b(P_i)}(f - f(w_i)) \right| \leq m_{\sigma,b(P_i)}(|f - f(w_i)|) \leq \mathrm{Lip}_{\beta}(f) \,\ell(P_i)^{\beta} \lesssim \mathrm{Lip}_{\beta}(f) \,|x - y|^{\beta}.
$$

implying that  $|S_3| \lesssim \mathrm{Lip}_\beta(f)|x-y|^\beta$ . If we combine the above estimates, we get

$$
|v_f(x) - v_f(y)| \le |S_1| + |S_2| + |S_3| \lesssim \mathrm{Lip}_\beta(f) |x - y|^\beta
$$

,

in the case 2 as well and thus for all  $x, y \in \Omega$  with  $x \neq y$ . This readily implies that  $\upsilon_f \in \mathrm{Lip}_\beta(\Omega)$  with  $\mathrm{Lip}_\beta(\upsilon_f) \leq \mathrm{Lip}_\beta(f)$ .

<span id="page-25-0"></span>It remains to prove that

(3.10) 
$$
|v_f(x) - v_f(y)| \lesssim \mathrm{Lip}_{\beta}(f) |x - y|^{\beta}, \quad \text{for any } x \in \partial \Omega \text{ and } y \in \Omega.
$$

To this end, we fix such x and y and estimate

$$
|v_f(x) - v_f(y)| = |f(x) - v_f(y)| = \left| f(x) - \sum_{P \in \mathcal{W}(\Omega)} m_{\sigma, b(P)} f \varphi_P(y) \right|
$$
  

$$
\leq \sum_{P \in \mathcal{W}(\Omega)} \varphi_P(y) |f(x) - m_{\sigma, b(P)} f| \leq \sum_{P \in \mathcal{W}(\Omega) : \atop 1, 1P \ni y} \varphi_P(y) |f(x) - m_{\sigma, b(P)} f|.
$$

As  $f \in \text{Lip}_{\beta}(\partial \Omega)$ , for every  $P \in \mathcal{W}(\Omega)$  such that  $y \in 1.1P$ , we have that

$$
|f(x) - m_{\sigma,b(P)}f| \lesssim \mathrm{Lip}_{\beta}(f) \,\ell(P)^{\beta} \approx \mathrm{Lip}_{\beta}(f) \,\delta_{\Omega}(y)^{\beta} \le \mathrm{Lip}_{\beta}(f) \,|x-y|^{\beta},
$$

which implies  $(3.10)$  by the bounded overlaps of the Whitney cubes that contain y, concluding the proof of the lemma.

*proof of Theorem [1.5.](#page-6-0)* It follows by combining Lemmas [3.1,](#page-21-2) [3.2,](#page-22-4) [3.3,](#page-23-2) [3.4,](#page-23-3) and [3.5](#page-24-0) (see also Remark [2.2\)](#page-10-0).

# <span id="page-26-1"></span>4. A CORONA DECOMPOSITION FOR FUNCTIONS IN  $L^p$  or BMO

<span id="page-26-0"></span>In this section we will assume that  $\Omega \in AR(s)$  for  $s \in (0, n]$ .

We say that a family of cubes  $\mathcal{F} \subset \mathcal{D}_{\sigma}$  satisfies a *Carleson packing condition* with constant  $M > 0$ , and we write  $\mathcal{F} \in \text{Car}(M)$ , if for any  $S \in \mathcal{D}_{\sigma}$ , it holds that

(4.1) 
$$
\sum_{R \in \mathcal{F}: R \subset S} \sigma(R) \leq M \sigma(S),
$$

A family  $T \subset \mathcal{D}_{\sigma}$  is a *tree* if it verifies the following properties:

- (1)  $T$  has a maximal element (with respect to inclusion)  $Q(T)$  which contains all the other elements of  $\mathcal T$  as subsets of  $\mathbb R^{n+1}$ . The cube  $Q(\mathcal T)$  is the "root" of  $\mathcal T$  and we will call it "top" cube.
- (2) If  $Q, Q'$  belong to  $\mathcal T$  and  $Q \subset Q'$ , then any  $\mu$ -cube  $P \in \mathcal D_{\sigma}$  such that  $Q \subset P \subset Q'$ also belongs to  $\mathcal T$ .

For a tree T, if  $R = Q(\mathcal{T})$  is a top cube, we will write  $\mathcal{T} = \text{Tree}(R)$ .

**Definition 4.1.** A *corona decomposition* of  $\sigma$  is a partition of  $\mathcal{D}_{\sigma}$  into a family of "good cubes", which we denote by  $\mathcal{G}$ , and a family of "bad cubes", which we denote by  $\mathcal{B}$ , so that the following hold:

- (1)  $\mathcal{D}_{\sigma} = \mathcal{G} \cup \mathcal{B}$ ;
- (2) There is a partition of  $G$  into trees, that is,

$$
\mathcal{G} = \bigcup_{\mathcal{T} \subset \mathcal{G}} \mathcal{T};
$$

- (3) The collections of maximal cubes  $Q(\mathcal{T})$  of the trees  $\mathcal T$  satisfies [\(4.1\)](#page-26-1) for some  $M_0 > 0;$
- (4) The collection of cubes B satisfies [\(4.1\)](#page-26-1) for some  $M_1 > 0$ .

We can also define a localized Corona decomposition in a cube  $R_0 \in \mathcal{D}_{\sigma}$  if, in the definition above, we replace  $\mathcal{D}_{\sigma}$  by  $\mathcal{D}_{\sigma}(R_0)$ .

We recall the definition of the truncated (at large scales) dyadic Hardy-Littlewood maximal function

$$
Mf(Q) = \sup_{\substack{R \in \mathcal{D}_{\sigma} \\ Q \subset R}} m_{\sigma,R} |f|, \quad f \in L^{1}_{loc}(\sigma).
$$

Given any  $R \in \mathcal{D}_{\sigma}$  and for fixed  $\varepsilon > 0$ , we define the collection  $\text{Stop}(R) \subset \mathcal{D}(R)$  consisting of cubes  $S \in \mathcal{D}(R)$  which are maximal (thus disjoint) with respect to the condition

<span id="page-27-1"></span>(4.2) 
$$
|m_{\sigma,R}f - m_{\sigma,S}f| \geq \begin{cases} \varepsilon Mf(S) & ,\text{if } f \in L^1_{\text{loc}}(\sigma) \\ \varepsilon \|f\|_{\text{BMO}(\sigma)} & ,\text{if } f \in \text{BMO}(\sigma). \end{cases}
$$

We fix a cube  $R_0 \in \mathcal{D}_{\sigma}$  and we define the family of the top cubes with respect to  $R_0$  as follows: first we define the families  $\text{Top}_k(R_0)$  for  $k \geq 0$  inductively. We set

$$
\mathsf{Top}_0(R_0) = \{R_0\}.
$$

Assuming that  $\mathsf{Top}_k(R_0)$  has been defined, we set

$$
\mathsf{Top}_{k+1}(R_0) = \bigcup_{R \in \mathsf{Top}_k(R_0)} \mathsf{Stop}(R),
$$

and then we define

$$
\mathsf{Top}(R_0) = \bigcup_{k \geq 0} \mathsf{Top}_k(R_0).
$$

We also set

$$
\text{Tree}(R) := \{ Q \in \mathcal{D}_{\sigma}(R) : \nexists \ S \in \text{Stop}(R) \text{ such that } Q \subset S \}.
$$

and notice that

$$
\mathcal{D}_{\sigma}(R_0) = \bigcup_{R \in \mathsf{Top}(R_0)} \mathrm{Tree}(R),
$$

and this union is disjoint. This is a localized Corona decomposition in  $R_0$  and notice that, in this case,  $\mathcal{B} = \emptyset$ .

In the rest of this section, we will devote all our efforts to proving that  $Top(R_0)$  satisfies a Carleson packing condition.

<span id="page-27-0"></span>**Proposition 4.2.** *For any*  $R_0 \in \mathcal{D}_{\sigma}$ , the family of cubes  $\textsf{Top}(R_0) \in \text{Car}(C\varepsilon^{-2})$  for some C > 0 *depending on the Ahlfors-regularity constants.*

For the proof of proposition [4.2](#page-27-0) we consider the cases  $f \in \text{BMO}(\sigma)$  and  $f \in L^1_{\text{loc}}(\sigma)$ separately.

*Proof of Proposition* [4.2](#page-27-0) *when*  $f \in BMO(\sigma)$ . For any  $R \in Top(R_0)$  it holds

$$
|m_{\sigma,S}f - m_{\sigma,R}f| > \varepsilon ||f||_{\text{BMO}(\sigma)}.
$$

Define

$$
f_R(x) := \sum_{Q \in \text{Tree}(R)} \Delta_Q f(x) = \sum_{Q \in \text{Tree}(R)} \sum_{Q' \in \text{ch}(Q)} (m_{\sigma, Q'} f - m_{\sigma, Q} f) 1_{Q'}(x).
$$

If  $x \in P \in \textsf{Stop}(R)$ , we have that  $f_R(x) = m_{\sigma,P} f - m_{\sigma,R} f$  and so,  $|f_R(x)| > \varepsilon \|f\|_{\mathrm{BMO}(\sigma)}$ . This implies that

$$
\varepsilon^{2} \|f\|_{\text{BMO}(\sigma)}^{2} \sum_{P \in \text{Stop}(R)} \sigma(P) \leq \sum_{P \in \text{Stop}(R)} \int_{P} |f_{R}(x)|^{2} d\sigma(x)
$$

$$
= \int_{\bigcup_{P \in \text{Stop}(R)}} |f_{R}|^{2} d\sigma \leq \int |f_{R}|^{2} d\sigma.
$$

By the above estimate and the orthogonality of  $\Delta_Q f$ ,

$$
\sum_{R \in \mathsf{Top}(R_0)} \sum_{P \in \mathsf{Stop}(R)} \sigma(P) \le \frac{1}{\varepsilon^2} \frac{1}{\|f\|_{\mathrm{BMO}(\sigma)}^2} \sum_{R \in \mathsf{Top}(R_0)} \|f_R\|_{L^2(\sigma)}^2
$$
\n
$$
= \frac{1}{\varepsilon^2} \frac{1}{\|f\|_{\mathrm{BMO}(\sigma)}^2} \sum_{R \in \mathsf{Top}(R_0)} \sum_{Q \in \mathrm{Tree}(R)} \|\Delta_Q f\|_{L^2(\sigma)}^2
$$
\n
$$
\le \frac{1}{\varepsilon^2} \frac{1}{\|f\|_{\mathrm{BMO}(\sigma)}^2} \sum_{Q \in \mathcal{D}_{\sigma}(R_0)} \|\Delta_Q f\|_{L^2(\sigma)}^2
$$
\n
$$
= \frac{1}{\varepsilon^2} \frac{1}{\|f\|_{\mathrm{BMO}(\sigma)}^2} \|\mathbf{1}_{R_0}(f - m_{\sigma, R_0}f)\|_{L^2(\sigma)}^2 \lesssim \varepsilon^{-2} \sigma(R_0)
$$

which proves [\(4.1\)](#page-26-1) in the case that  $S \in Top(R_0)$ . By the same argument as in the end of the proof of Proposition [4.2](#page-27-0) when  $f \in L^1_{loc}(\sigma)$ , we obtain [\(4.1\)](#page-26-1) for any  $S \in \mathcal{D}_{\sigma}(R_0)$ .  $\Box$ 

<span id="page-28-0"></span>To prove the proposition for  $f \in L^1_{loc}$ , we first need some auxiliary lemmas.

**Lemma 4.3.** *Let*  $f \in L^1_{loc}(\sigma)$  *and*  $Q \in \mathcal{D}_{\sigma}$ *. Then it holds* 

(4.3) 
$$
\frac{\sigma(Q)}{Mf(Q)^2} \le 8 \int_Q \frac{1}{\left(\mathcal{M}_{\mathcal{D}_\sigma}f(x)\right)^2} d\sigma(x)
$$

*Proof.* This was shown in [\[HR18,](#page-70-5) Lemma 4.1] for the Lebesgue measure but the same proof works for any non-atomic Radon measure and so we skip the details.  $\Box$ 

Let  $\mathcal{F} \subset \mathcal{D}_{\sigma}$  be any collection of dyadic cubes. Given any cube  $Q \in \mathcal{D}_{\sigma}$ , define its stopping parent  $Q^*$  to be the minimal  $Q^* \in \mathcal{F}$  such that  $Q \subsetneq Q^*$ . If no such  $Q^*$  exists, we set  $Q^* := Q$ . Define the stopped square function

(4.4) 
$$
\mathcal{S}_{\mathcal{F}}f(x) := \Big(\sum_{Q \in \mathcal{F}} |m_{\sigma,Q}f - m_{\sigma,Q^*}f|^2 \mathbf{1}_Q(x)\Big)^{1/2}.
$$

<span id="page-28-1"></span>In the special case  $\mathcal{F} = Top(R_0)$ , we will simply write  $\mathcal{S}f$ .

**Lemma 4.4.** *If*  $w \in A_{\infty}(\sigma)$  *and*  $1 \leq p < \infty$ *, then* 

$$
\|\mathcal S_{\mathcal F} f\|_{L^p(\partial\Omega;w)}\lesssim \|M_{\mathcal D} f\|_{L^p(\partial\Omega;w)}
$$

*uniformly for any collection of dyadic cubes* F*.*

*Proof.* This was proved in [\[HR18,](#page-70-5) Proposition 3.2] for the Lebesgue measure but the same proof works verbatim for  $\sigma$ .

We will now proceed to the proof of Proposition [4.2](#page-27-0) for  $f \in L^1_{loc}(\sigma)$ , which is based on the one of  $[HR18, Theorem 1.2(3)].$ 

*Proof of Proposition* [4.2](#page-27-0) *when*  $f \in L^1_{loc}(\sigma)$ . We first fix  $S \in Top(R_0)$ . As for any  $R \in$  $\textsf{Top}(R_0)$  it holds that

$$
|m_{\sigma,R}f - m_{\sigma,R^*}f| > \varepsilon Mf(R)
$$

we have that

$$
\sum_{R \in \mathsf{Top}(R_0)} \sigma(R) \leq \sum_{R \in \mathsf{Top}(R_0)} \frac{|m_{\sigma,R}f - m_{\sigma,R^*}f|^2}{\varepsilon^2 Mf(R)^2} \sigma(R)
$$
\n
$$
\lesssim \sum_{R \in \mathsf{Top}(R_0)} \frac{|m_{\sigma,R}f - m_{\sigma,R^*}f|^2}{\varepsilon^2} \int_S \frac{\mathbf{1}_R(x)}{\mathcal{M}_{\mathcal{D}_{\sigma}}f(x)^2} d\sigma = \int_S \frac{\mathcal{S}f(x)^2}{\varepsilon^2} \frac{d\sigma(x)}{\mathcal{M}_{\mathcal{D}_{\sigma}}f(x)^2},
$$

where, in the second inequality, we used Lemma [4.3.](#page-28-0) We write

$$
(\mathcal{M}_{\mathcal{D}_{\sigma}}f)^{-2} = 1 \cdot ((\mathcal{M}_{\mathcal{D}_{\sigma}}f)^{2\gamma})^{1-q}
$$

for  $\gamma \in (0, 1/2)$  and  $q = 1 + \frac{1}{\gamma} > 3$ . Since  $f \in L^1_{loc}(\sigma)$  and  $2\gamma \in (0, 1)$ , using for example [\[CG85,](#page-69-22) Theorem 3.4, p.158] (whose proof works for doubling Borel measures), we get that  $(\mathcal{M}_{\mathcal{D}_{\sigma}}f)^{2\gamma} \in A_1(\sigma)$ . As  $1 \in A_1$  and  $q > 1$  it follows from [\[CG85,](#page-69-22) Theorem 2.16, p.407] (whose proof also works for doubling Borel measures) that  $1 \cdot \left( (\mathcal{M}_{\mathcal{D}} f)^{2\gamma} \right)^{1-q} \in A_q(\sigma)$ . Therefore,  $(M_{\mathcal{D}_{\sigma}}f)^{-2} \in A_q(\sigma) \subset A_{\infty}(\sigma)$ . We now apply Lemma [4.4](#page-28-1) with the collection of cubes  $\widetilde{\mathcal{F}} := \{ R \in \text{Top}(R_0) : R \subset S \}$  to the function

$$
\tilde{f}(x) := \begin{cases} f(x) - m_{\sigma,S}f & \text{, if } x \in S \\ 0 & \text{, if } x \notin S, \end{cases}
$$

for the weight  $w := (\mathcal{M}_{\mathcal{D}_{\sigma}}f)^{-2}$  and  $p = 2$ , and obtain

$$
\int |S_{\tilde{\mathcal{F}}} \tilde{f}|^2 w d\sigma \lesssim \int |\mathcal{M}_{\mathcal{D}} \tilde{f}|^2 w d\sigma = \int_S |\mathcal{M}_{\mathcal{D}} \tilde{f}|^2 w d\sigma \lesssim \int_S |\mathcal{M}_{\mathcal{D}} f|^2 w d\sigma.
$$

Thus, since  $|S_{\tilde{\mathcal{F}}} \tilde{f}(x)|^2 = |S_{\tilde{\mathcal{F}}} f(x)|^2$  for all  $x \in S$ , we infer that

$$
\int_{S} |\mathcal{S}f|^{2}w d\sigma \lesssim \int_{S} |\mathcal{S}_{\widetilde{\mathcal{F}}}f|^{2}w d\sigma + \int_{S} |\mathcal{M}_{D}f|^{2}w d\sigma \le \int |\mathcal{S}_{\widetilde{\mathcal{F}}}f|^{2}w d\sigma + \int_{S} |\mathcal{M}_{D}f|^{2}w d\sigma
$$
  

$$
\lesssim \int_{S} |\mathcal{M}_{D}f|^{2}w d\sigma = \int_{S} |\mathcal{M}_{D}f|^{2} \frac{d\sigma}{(\mathcal{M}_{D}f)^{2}} = \sigma(S),
$$

proving [\(4.1\)](#page-26-1) in the case that  $S \in Top(R_0)$ .

If  $S \in \mathcal{D}_{\sigma}(R_0) \setminus \text{Top}(R_0)$ , we can find a maximal collection  $\mathcal{F}_0$  of cubes  $\widetilde{S} \in \text{Top}(R_0)$ such that

$$
S = \bigcup_{\widetilde{S} \in \mathcal{F}_0} \widetilde{S}.
$$

Then,

$$
\sum_{\substack{R \in \mathsf{Top}(R_0) \\ R \subset S}} \sigma(R) = \sum_{\widetilde{S} \in \mathcal{F}_0} \sum_{\substack{R \in \mathsf{Top}(\widetilde{S}):\\ R \subset \widetilde{S}}} \sigma(R) \lesssim \sum_{\widetilde{S} \in \mathcal{F}_0} \sigma(\widetilde{S}) = \sigma(S)
$$

and the proof is now complete.

<span id="page-30-3"></span>**Remark 4.5.** If supp  $\sigma$  is bounded, we can pick  $R_0 = \text{supp }\sigma$ . In the case that supp  $\sigma$  is not bounded we apply a technique described in p. 38 of [\[DS91\]](#page-69-9): we consider a family of cubes  $\{R_i\}_{i\in J}\subset \mathcal{D}_{\sigma}$  which are pairwise disjoint, whose union is all of supp  $\sigma$ , and which have the property that for each k there at most C cubes from  $\mathcal{D}_{\sigma,k}$  not contained in any cube  $R_j$ . For each  $R_j$  we construct a family  $\textsf{Top}(R_j)$  analogous to  $\textsf{Top}(R_0)$ . Then we set

$$
\mathsf{Top}:=\bigcup_{j\in J}\mathsf{Top}(R_j)
$$

and

$$
\mathcal{B} := \{ S \subset \mathcal{D}_{\sigma} : \text{ there does not exist } j \in J \text{ such that } S \subset R_j \in \mathsf{Top} \}.
$$

One can easily check that the families  $Top$  and  $B$  satisfy a Carleson packing condition. See [\[DS91,](#page-69-9) p. 38] for the construction of the family  $\{R_i\}$  and additional details.

# <span id="page-30-0"></span>5.  $L^p$  and uniform  $\varepsilon$ -approximability of the regularized dyadic extension

Given  $A > 1$ , we say that two cubes  $Q_1, Q_2$  are A-close if

$$
\frac{1}{A}\operatorname{diam} Q_1 \le \operatorname{diam} Q_2 \le A \operatorname{diam} Q_1
$$

and

$$
dist(Q_1, Q_2) \le A(\text{diam } Q_1 + \text{diam } Q_2).
$$

<span id="page-30-1"></span>The following lemma was proved in [\[DS93,](#page-69-10) p. 60].

**Lemma 5.1.** *If we have a Corona decomposition such that*  $\text{Top} \in \text{Car}(M_0)$  *for some* M<sup>0</sup> > 0*, then, the collection of cubes*

 $\mathcal{A}_0 := \{Q \in \mathcal{D}_{\sigma} : Q \in \text{Tree}(R) \text{ for } R \in \text{Top} \text{ and }$  $\exists Q' \in \text{Tree}(R')$  for some  $R' \neq R \in \text{Top}$ , such that  $Q'$  A-close to  $Q$ 

*is in*  $\text{Car}(M_1)$  *for some*  $M_1 > 0$  *depending on* M, A, and the Ahlfors-regularity constants.

<span id="page-30-2"></span>**Lemma 5.2** (Lemma I.3.27, p. 59 in [\[DS93\]](#page-69-10) ). *If*  $\mathcal{F} \subset \mathcal{D}_{\sigma}$  *is in* Car( $M_1$ ) *for some*  $M_1 > 0$ , *then the family*

$$
\mathcal{F}_A := \{ Q \in \mathcal{D}_{\sigma} : Q \text{ is } A \text{-close to some } Q' \in \mathcal{F} \}
$$

*is in*  $\text{Car}(M_2)$  *for some*  $M_2 > 0$  *depending on*  $M_1$ *, A, and the Ahlfors-regularity constants.* 

Let us define the family of Whitney cubes

 $\mathcal{P}_0 := \{ P \in \mathcal{W}(\Omega) : \text{there exists } P' \in \mathcal{W}(\Omega) \text{ such that } 1.2P \cap 1.2P' \neq \emptyset \text{ and there exist } \Omega' \in \mathcal{W}(\Omega) \text{ such that } 1.2P \cap 1.2P' \neq \emptyset \text{ and there exist } \Omega' \in \mathcal{W}(\Omega) \text{ such that } 1.2P \cap 1.2P' \neq \emptyset \text{ and there exist } \Omega' \in \mathcal{W}(\Omega) \text{ such that } \Omega' \neq \emptyset \text{ and there exist } \Omega' \in \mathcal{W}(\Omega) \text{ such$  $R, R' \in \text{Top with } R \neq R'$  such that  $b(P) \in \text{Tree}(R)$  and  $b(P') \in \text{Tree}(R')$ .

Then, by the properties of Whitney cubes, for every  $P \in \mathcal{P}_0$ , the cubes  $P' \in \mathcal{W}(\Omega)$  such that  $b(P')$  is not in the same tree as  $b(P)$  have the following properties:

$$
\bullet \ \ell(b(P))/2 \leq \ell(b(P')) \leq 2\ell(b(P))
$$

• dist $(b(P), b(P')) \leq C_1 \ell(b(P)).$ 

If, for fixed  $R \in \text{Top}$ , we define

 $\partial \text{Tree}(R) := \{ Q \in \text{Tree}(R) : \text{there exists } P \in \mathcal{P}_0 \text{ such that } b(P) = Q \}$ 

then, there exists  $A > 1$  large enough depending only on  $C_1$  and n such that

$$
\bigcup_{R\in\mathsf{Top}}\partial\mathrm{Tree}(R)\subset\mathcal{A}_0,
$$

and, by Lemma [5.1,](#page-30-1)  $\bigcup_{R \in \mathsf{Top}} \partial \text{Tree}(R) \in \text{Car}(M_1)$ . If F is a family of "true" dyadic cubes in  $\mathbb{R}^{n+1}$ , we also define

$$
\mathcal{N}(\mathcal{F}) := \{ P \in \mathcal{W}(\Omega) : \text{there exists } P' \in \mathcal{F} \text{ such that } 1.2P \cap 1.2P' \neq \emptyset \}.
$$

So, for  $\mathcal{F} = \mathcal{P}_0$ , we set

$$
\partial \text{Tree}^*(R) := \{ Q \in \mathcal{D}_\sigma : \exists P \in \mathcal{N}(\mathcal{P}_0) \text{ such that } Q = b(P) \in \text{Tree}(R) \}.
$$

It is easy to see that

$$
\mathcal{J}:=\bigcup_{R\in\mathsf{Top}}\partial\mathrm{Tree}^*(R)\subset\Big(\bigcup_{R\in\mathsf{Top}}\partial\mathrm{Tree}(R)\Big)_A
$$

and, by Lemma [5.2,](#page-30-2)  $\mathcal{J} \in \text{Car}(M_2)$  for some  $M_2 > 0$ . Finally, recall that  $\mathcal{B}$  is a collection of bad cubes satisfying a Carleson packing condition [\(4.1\)](#page-26-1) and define

$$
\mathcal{B}_0 := \{ P \in \mathcal{W}(\Omega) : b(P) \in \mathcal{B} \}.
$$

We are now ready to define the *approximating function* of  $v_f$  by

<span id="page-31-0"></span>
$$
(5.1) \ u(x) = \sum_{S \in \mathcal{B}_0} m_{\sigma, b(S)} f \varphi_S(x)
$$
  
+ 
$$
\sum_{R \in \text{Top}} \Big[ \sum_{\substack{P \in \mathcal{W}(\Omega) \setminus \mathcal{P}_0 \\ b(P) \in \text{Tree}(R)}} m_{\sigma, R} f \varphi_P(x) + \sum_{\substack{P \in \mathcal{P}_0 \\ b(P) \in \text{Tree}(R)}} m_{\sigma, b(P)} f \varphi_P(x) \Big],
$$

using the Corona decomposition constructed in Section [4.](#page-26-0) Note that when  $\Omega$  is bounded, Top = Top( $\partial\Omega$ ) and  $\mathcal{B} = \emptyset$ , while if  $\partial\Omega$  is unbounded, Top and  $\mathcal{B}$  are the families con-structed in Remark [4.5.](#page-30-3) Finally, when  $\Omega$  is an unbounded domain with compact boundary  $\partial\Omega$ , we modify the definition of the approximating function as follows.

(5.2) 
$$
u(x) = \sum_{P \in \mathcal{W}(\Omega): \ell(P) \ge \text{diam}(\partial \Omega)} m_{\sigma, \partial \Omega} f \varphi_P(x) + \sum_{R \in \text{Top}} \left[ \sum_{\substack{P \in \mathcal{W}(\Omega) \setminus \mathcal{P}_0 \\ b(P) \in \text{Tree}(R)}} m_{\sigma, R} f \varphi_P(x) + \sum_{\substack{P \in \mathcal{P}_0 \\ b(P) \in \text{Tree}(R)}} m_{\sigma, b(P)} f \varphi_P(x) \right].
$$

<span id="page-32-9"></span>**Theorem 5.3.** Let  $f \in L^1_{loc}(\sigma)$  and  $\varepsilon > 0$ . There exists  $\alpha_0 \ge 1$  such that for any  $\alpha \ge \alpha_0$ *and any*  $\xi \in \partial \Omega$ *,* 

<span id="page-32-0"></span>(5.3)  
\n
$$
\mathcal{N}_{\alpha}(u - v_f)(\xi) + \mathcal{N}_{\alpha}(\delta_{\Omega} \nabla (u - v_f))(\xi) \lesssim \varepsilon \mathcal{M}f(\xi),
$$
\n(5.4)  
\n
$$
\mathcal{N}_{\alpha}(\delta_{\Omega} \nabla u)(\xi) \lesssim \mathcal{M}f(\xi) + \mathcal{M}(\mathcal{M}f)(\xi),
$$

<span id="page-32-5"></span><span id="page-32-2"></span>(5.5) 
$$
\mathcal{C}_s(\nabla u)(\xi) \lesssim_{\varepsilon} \mathcal{M}(\mathcal{M}f)(\xi) + \mathcal{M}(\mathcal{M}(\mathcal{M}f))(\xi),
$$

<span id="page-32-4"></span>(5.6) 
$$
\mathcal{C}_s(\delta_{\Omega}|\nabla u|^2)(\xi) \lesssim_{\varepsilon} \mathcal{M}((\widetilde{\mathcal{M}}f)^2)(\xi) + \mathcal{M}(\widetilde{\mathcal{M}}((\mathcal{M}f)^2))(\xi).
$$

*Here*  $c_{\varepsilon}$  *is a positive constant depending on*  $\varepsilon$  *and*  $\alpha_0$  *depend only on n and the Ahlfors regularity, the corkscrew condition, and the Whitney constants.*

Let  $f \in BMO(\sigma)$  and  $\varepsilon > 0$ . Then, for any  $x \in \Omega$ , it holds that

<span id="page-32-8"></span>(5.7) 
$$
|u(x) - v_f(x)| + \delta_{\Omega}(x)|\nabla(u - v_f)(x)| \leq \varepsilon ||f||_{\text{BMO}(\sigma)}
$$

<span id="page-32-6"></span>
$$
\delta_{\Omega}(x)|\nabla u(x)| \lesssim ||f||_{\text{BMO}(\sigma)},
$$

*and for any*  $\xi \in \partial\Omega$ *,* 

<span id="page-32-3"></span>(5.9) 
$$
\mathcal{C}_s(\nabla u)(\xi) + \left[\mathcal{C}_s(\delta_{\Omega}|\nabla u|^2)(\xi)\right]^{1/2} \lesssim_{\varepsilon} ||f||_{\text{BMO}(\sigma)},
$$

*where the implicit constants depend only on* n *and the Ahlfors regularity, the corkscrew condition, and the Whitney constants.*

<span id="page-32-7"></span>*Moreover, if*  $f \in \text{Lip}_c(\partial \Omega)$ *, then*  $u \in \text{Lip}_{loc}(\Omega)$  *and for any*  $x \in \Omega$ *,* 

(5.10) 
$$
\delta_{\Omega}(x)|\nabla u(x)| \lesssim \text{Lip}(f)\,\text{diam}(\text{supp }f).
$$

*Proof.* We will only deal with the case that both  $\Omega$  and  $\partial\Omega$  are unbounded as the other cases can be treated in a similar but easier way. We remark first that if we choose  $\alpha_0$ large enough, depending on  $n$ , the constants of the corkscrew condition and the Whitney decomposition, the cone is always non-empty and for every  $Q \in \mathcal{D}_{\sigma}$  such that  $\xi \in Q$ , there exists  $P \in \mathcal{W}(\Omega)$  such that  $b(P) = Q$  and  $P \subset \gamma_{\alpha_0}(\xi)$ .

For fixed  $\xi \in \partial\Omega$  we let  $x \in \gamma_\alpha(\xi)$  for  $a \ge \alpha_0$ . There exists  $P_0 \in \mathcal{W}(\Omega)$  such that  $x \in \bar{P}_0$  and we either have that  $P_0 \in \mathcal{B}_0$  or that there is a unique  $R_0 \in \text{Top}$  such that  $b(P_0) \in Tree(R_0)$ . If either  $P_0 \in \mathcal{P}_0$  and there does not exist any  $P \in \mathcal{N}(\mathcal{P}_0) \setminus \mathcal{P}_0$  such that  $x \in 1.1P$ , or  $P_0 \in \mathcal{B}_0$ , it is easy to see that  $u(x) - v_f(x) = 0$ , while, if  $P_0 \in \mathcal{P}_0$  and there exists some  $\widetilde{P} \in \mathcal{N}(\mathcal{P}_0) \setminus \mathcal{P}_0$  such that  $x \in 1.1\widetilde{P}$ , then

$$
u(x) - v_f(x) = \sum_{\substack{P \in \mathcal{N}(\mathcal{P}_0) \backslash \mathcal{P}_0 \\ b(P) \in \text{Tree}(R_0)}} (m_{\sigma, R_0} f - m_{\sigma, b(P)} f) \varphi_P(x).
$$

The same is true if  $P_0 \in \mathcal{N}(\mathcal{P}_0) \setminus \mathcal{P}_0$  and there is  $P \in \mathcal{P}_0$  such that  $x \in 1.1P$ . In any other case, we have that

<span id="page-32-1"></span>(5.11) 
$$
u(x) - v_f(x) = \sum_{\substack{P \in \mathcal{W}(\Omega) \\ b(P) \in \text{Tree}(R_0)}} (m_{\sigma, R_0} f - m_{\sigma, b(P)} f) \varphi_P(x).
$$

Therefore, since  $b(P) \in Tree(R_0)$ , by [\(4.2\)](#page-27-1),

$$
|u(x) - v_f(x)| \le \varepsilon \sum_{\substack{P \in \mathcal{W}(\Omega) \\ b(P) \in \text{Tree}(R_0)}} Mf(b(P)) \varphi_P(x).
$$

For any  $P \in \mathcal{W}(\Omega)$  such that  $x \in 1.1P \cap \gamma_\alpha(\xi)$ , since  $|x - \xi| \approx \delta_\Omega(x) \approx \ell(P)$ , it holds that  $P \subset B(\xi, M\ell(P))$ . The same is true for any  $S \in \mathcal{D}_{\sigma}$  such that  $b(P) \subset S$ , i.e.,  $S \subset B(\xi, M'(\ell(S)))$ , for a possibly larger constant  $M' > 0$  depending also on the Ahlfors regularity constants. Thus,

<span id="page-33-0"></span>
$$
(5.12) \qquad |u(x) - v_f(x)| \lesssim \varepsilon \sup_{S \supset b(P)} m_{\sigma, B(\xi, M'\ell(S))}(|f|) \lesssim \varepsilon \sup_{r \gtrsim \delta_{\Omega}(x)} m_{\sigma, B(\xi, r)}(|f|).
$$

which, by taking supremum over all  $x \in \gamma_\alpha(\xi)$ , implies [\(5.3\)](#page-32-0). By the same arguments and the fact the  $\nabla \varphi_P(x) \lesssim \ell(P)^{-1} \approx \delta_\Omega(x)^{-1}$ , we infer that

$$
\nabla (u - v_f)(x) \lesssim \varepsilon \mathcal{M}(f)(\xi) \delta_{\Omega}(x)^{-1},
$$

which implies

(5.13) 
$$
\mathcal{N}_{\alpha}(\delta_{\Omega}\nabla(u-v_f))(\xi) \lesssim \varepsilon \mathcal{M}(f)(\xi).
$$

In the case that  $f \in \text{BMO}(\sigma)$ , in view of [\(5.11\)](#page-32-1) and  $\nabla \varphi_P(x) \lesssim \ell(P)^{-1} \approx \delta_{\Omega}(x)^{-1}$ , we have that

(5.14) 
$$
|u(x) - v_f(x)| + \delta_{\Omega} |\nabla (u - v_f)(x)| \lesssim \varepsilon ||f||_{\text{BMO}}.
$$

We now turn our attention to the proof of [\(5.5\)](#page-32-2) and [\(5.9\)](#page-32-3). Let  $x \in \overline{P}_0 \in \mathcal{W}(\Omega)$ . Then, once again, either there exists a unique  $R_0 \in \text{Top}$  such that  $b(P_0) \in \text{Tree}(R_0)$ , or there exists  $B_0 \in \mathcal{B}_0$  such that  $b(P_0) = B_0$ . For the sake of brevity, we denote

(5.15) 
$$
B_x := c B^x = B(x, c \delta_{\Omega}(x)),
$$

for a small enough constant  $c > 0$  to be chosen. Fix  $y \in B_x$  and if  $P \in \mathcal{W}(\Omega)$  is such that  $y \in 1.1P$ , then  $x \in 1.2P$ . Indeed, by [\(2.48\)](#page-20-3), we always have that  $dist(x, 1.1P) \le |x-y| \le$  $c \,\delta_{\Omega}(x) \leq c \,\Lambda' \,\ell(P_0)$ , and if there exists  $P \in \mathcal{W}(\Omega)$  such that  $y \in 1.1P$  and  $x \notin 1.2P$  it also holds that dist $(x, 1.1P) \ge 0.1\ell(P)$ . Now, note that since  $\frac{1}{2}\ell(P_0) \le \ell(P) \le 2\ell(P_0)$ , we get that  $\frac{1}{2}\ell(P_0) \le c \Lambda' \ell(P_0)$  and if we choose  $c = \frac{1}{4\Lambda'}$  we reach a contradiction.

It is easy to see that  $\nabla u(y) = 0$  if there does not exist any cube  $P \in \mathcal{P}_0$  or  $P \in \mathcal{B}_0$  such that  $y \in 1.1P$ . Using that  $\sum \nabla \varphi_P (y) = 0$ , we have that

<span id="page-33-1"></span>(5.16) 
$$
\nabla u(y) = \left(\sum_{P \in \mathcal{B}_0} + \sum_{\substack{P \in \mathcal{N}(\mathcal{P}_0) \\ b(P) \notin \text{Tree}(R_0)}} \right) (m_{\sigma, b(P)} f - m_{\sigma, b(P_0)} f) \nabla \varphi_P(y) + \sum_{\substack{P \in \mathcal{P}_0 \\ b(P) \in \text{Tree}(R_0)}} (m_{\sigma, b(P)} f - m_{\sigma, b(P_0)} f) \nabla \varphi_P(y) + \sum_{\substack{P \in \mathcal{N}(\mathcal{P}_0) \setminus \mathcal{P}_0 \\ b(P) \in \text{Tree}(R_0)}} (m_{\sigma, R_0} f - m_{\sigma, b(P)} f) \nabla \varphi_P(y).
$$

Therefore, by Remark [2.17,](#page-20-2) arguing as in the proof of  $(5.12)$  and using  $(5.16)$ , the fact that  $\ell(P_0) \approx \ell(P)$  for any  $P \in \mathcal{W}(\Omega)$  such that  $1.1P \ni y$ , and the Carleson packing of the cubes in  $\mathcal{B}_0$ , for fixed  $\xi \in \partial \Omega$  and  $r > 0$ , we can estimate

$$
\int_{B(\xi,r)\cap\Omega} \sup_{y\in B_{x}} |\nabla u(y)| \frac{dx}{\delta_{\Omega}(x)^{n-s}}\n\lesssim \left(\sum_{P\in\mathcal{B}_{0}} + \sum_{R\in\text{Top}} \sum_{P\in\mathcal{N}(P_{0})} \right) \int_{P\cap B(\xi,r)} m_{\sigma,B(x_{b(P)},C_{w}\ell(P))} (|f|) \frac{dx}{\ell(P)^{n+1-s}}\n+ 2\varepsilon \sum_{R\in\text{Top}} \sum_{P\in\mathcal{P}_{0}} \int_{P\cap B(\xi,r)} \sup_{\rho\gtrsim\ell(P)} m_{\sigma,B(x_{b(P)},\rho)} (|f|) \frac{dx}{\ell(P)^{n+1-s}}\n+ \varepsilon \sum_{R\in\text{Top}} \sum_{P\in\mathcal{N}(P_{0})\backslash\mathcal{P}_{0}} \int_{P\cap B(\xi,r)} \sup_{\rho\gtrsim\ell(P)} m_{\sigma,B(x_{b(P)},\rho)} (|f|) \frac{dx}{\ell(P)^{n+1-s}}\n+ \varepsilon \sum_{R\in\text{Top}} \sum_{P\in\mathcal{N}(P_{0})\backslash\mathcal{P}_{0}} \sup_{P\cap B(\xi,r)\geqslant\ell(P)} \sup_{\rho\gtrsim\ell(P_{0})} m_{\sigma,B(x_{b(P)},C_{w}\ell(P))} (|f|)\n+ \sum_{P\in\mathcal{B}_{0}} + \sum_{R\in\text{Top}} \sum_{P\in\mathcal{N}(P_{0})} \sigma(b(P)) \inf_{\rho\in\mathcal{B}(x_{b(P)},M\ell(P))} \mathcal{M}f(\zeta)\n+ \sum_{R\in\text{Top}} \sum_{P\in\mathcal{N}(P_{0})} \sigma(b(P)) \inf_{\rho\in\mathcal{B}(x_{b(P)},M\ell(P))} \mathcal{M}f(\zeta)\n+ \sum_{Q\in\mathcal{B}(\mathcal{F},r)\neq\emptyset} \sigma(Q) m_{\sigma,B(x_{Q},C_{w}\ell(Q))} (|f|) + \sum_{Q\in\mathcal{J}} \sigma(Q) m_{\sigma,B(x_{Q},M\ell(Q))} (\mathcal{M}f) \frac{dx}{\rho_{C\in\mathcal{B}(\xi,C')}}\n\lesssim \int_{B(\xi,Cr)} \sup_{Q\ni z} m_{\sigma,B(x_{
$$

for  $M > 1$  possible larger than  $C_w$  and where in the antepenultimate inequality we used that if  $P \cap B(\xi, r) \neq \emptyset$ , then  $b(P) \subset B(\xi, C'r)$  for some large constant  $C' > 0$  depending on Ahlfors-regularity and the Whitney constants, while the penultimate inequality follows from Carleson's embedding theorem (see [\[To14,](#page-70-15) Theorem 5.8, p. 144]) since the families  $\mathcal{J} = \cup_{R \in \text{Top}} \partial^* \text{Tree}(R)$  and B are Carleson families. This readily concludes [\(5.5\)](#page-32-2), while the proof of  $(5.6)$  follows by similar arguments. We omit the details.

If  $f \in \text{BMO}(\sigma)$ , using [\(5.16\)](#page-33-1), for  $\xi \in \partial \Omega$  and  $r > 0$ ,

$$
\int_{B(\xi,r)\cap\Omega} \sup_{y\in B_x} |\nabla u(y)|\frac{dx}{\delta_\Omega(x)^{n-s}}\lesssim_\varepsilon \sum_{R\in\mathsf{Top}} \sum_{\substack{P\in\mathcal{N}(\mathcal{P}_0)\\ b(P')\in\mathrm{Tree}(R)}} \int_{P\cap B(\xi,r)} \|f\|_{\mathrm{BMO}(\sigma)}\, \omega_s(x)\, dx
$$

36 MIHALIS MOURGOGLOU AND THANASIS ZACHAROPOULOS

$$
+ \Big( \sum_{P \in \mathcal{B}_0} + \sum_{R \in \text{Top}} \sum_{\substack{P \in \mathcal{N}(\mathcal{P}_0) \\ b(P) \notin \text{Tree}(R)}} \Big) \int_{P \cap B(\xi,r)} \|f\|_{\text{BMO}(\sigma)} \frac{\omega_s(x)}{\ell(P)} dx
$$
  

$$
\lesssim \|f\|_{\text{BMO}(\sigma)} \sum_{\substack{P \in \mathcal{N}(\mathcal{P}_0) \cup \mathcal{B}_0 \\ P \cap B(\xi,r) \neq \emptyset}} \sigma(b(P)) \le \|f\|_{\text{BMO}(\sigma)} \sum_{\substack{Q \in \mathcal{J} \cup \mathcal{B} \\ Q \subset B(\xi, Mr)}} \sigma(Q)
$$
  

$$
\lesssim \|f\|_{\text{BMO}(\sigma)} r^s,
$$

for  $M > 1$  large enough constant depending on the Ahlfors regularity and the Whitney constants. For the last inequality we used that the families of surface cubes  $\mathcal J$  and  $\mathcal B$  satisfy Carleson packing condition from Lemma [5.2.](#page-30-2) Similarly, we can show that

$$
\int_{B(\xi,r)\cap\Omega} \sup_{y\in B_x} \delta_{\Omega}(y) |\nabla u(y)|^2 \frac{dx}{\delta_{\Omega}(x)^{n-s}} \lesssim_{\varepsilon} ||f||_{\operatorname{BMO}(\sigma)}^2 r^s
$$

The two estimates above obviously imply  $(5.9)$ .

Let  $\xi \in \partial\Omega$  and  $x \in \gamma_{\alpha}(\xi)$ . There exists  $P_0 \in \mathcal{W}(\Omega)$  such that  $x \in \overline{P}_0$ . Then using [\(5.16\)](#page-33-1) and the bounded overlaps of the Whitney cubes, it holds

<span id="page-35-0"></span>
$$
|\nabla u(x)| \lesssim \ell(P_0)^{-1} (m_{\sigma, B_{P_0}}(|f|) + \varepsilon \sup_{\rho \gtrsim \ell(P_0)} m_{\sigma, B(x_{b(P_0)}, \rho)}(|f|))
$$
  
(5.17)  

$$
\leq \delta_{\Omega}(x)^{-1} \big( m_{\sigma, B(\xi, C' \ell(P_0))}(|f|) + \inf_{\zeta \in B(\xi, C' \ell(P_0))} \mathcal{M}f(\zeta) \big)
$$
  

$$
\lesssim \delta_{\Omega}(x)^{-1} \big( \mathcal{M}f(\xi) + \mathcal{M}(\mathcal{M}f)(\xi) \big).
$$

By a similar but easier argument, we can show that

<span id="page-35-1"></span>(5.18) 
$$
|\nabla u(x)| \lesssim ||f||_{\text{BMO}(\sigma)} \delta_{\Omega}(x)^{-1}.
$$

Since  $\sup_{x \in \Omega} \delta_{\Omega}(x) |\nabla u(x)| = \sup_{\xi \in \partial \Omega} \sup_{x \in \gamma_{\alpha}(\xi)} \delta_{\Omega}(x) |\nabla u(x)|$ , it easily follows that the estimates  $(5.17)$  and  $(5.18)$  imply the estimates  $(5.4)$  and  $(5.8)$  respectively.

It remains to prove [\(5.10\)](#page-32-7) in the case that  $f \in \text{Lip}_{c}(\partial \Omega)$ . Using [\(5.16\)](#page-33-1) and the bounded overlaps of the Whitney cubes, it holds that for any  $x \in \Omega$ ,

<span id="page-35-2"></span>(5.19) 
$$
|\nabla u(x)| \lesssim \text{Lip}(f) \operatorname{diam}(\text{supp } f) \ell(P)^{-1} + \varepsilon \sum_{\substack{P \in \mathcal{N}(\mathcal{P}_0) \backslash \mathcal{P}_0 \\ b(P) \in \text{Tree}(R_0)}} Mf(b(P)) \ell(P)^{-1}.
$$

Since f has compact support, if  $\xi_0 \notin \text{supp } f$ , for every  $Q \supset b(P)$ , we have that

$$
m_{\sigma,Q}(|f|) = m_{\sigma,Q}(|f - f(\xi_0)|) \lesssim \text{diam}(\text{supp } f) \operatorname{Lip}(f).
$$

Taking supremum over all cubes  $Q \supset b(P)$  and using again the bounded overlaps of the Whitney cubes, by [\(5.19\)](#page-35-2) and the fact that  $\delta_{\Omega}(x) \approx \ell(P)$  for all  $P \in \mathcal{W}(\Omega)$  such that  $x \in 1.1P$ , we infer that

$$
|\nabla u(x)| \lesssim \text{Lip}(f) \text{ diam}(\text{supp } f) \delta_{\Omega}(x)^{-1},
$$

and this ends the proof.  $\Box$ 

As a corollary we get that if  $f \in L^p(\sigma)$ ,  $p \in (1,\infty)$ , (resp.  $f \in \text{BMO}(\sigma)$ ), then  $v_f$  is  $\varepsilon$ -approximable in  $L^p$  (uniformly  $\varepsilon$ -approximable).

<span id="page-36-0"></span>**Theorem 5.4.** *If*  $f \in \text{BMO}(\sigma)$ *, for any*  $\varepsilon > 0$ *, there exists*  $u = u_{\varepsilon} \in C^{\infty}(\Omega)$ *,*  $\alpha_1 \geq 1$ *, and a constant*  $c_{\varepsilon} > 1$ *, such that for any*  $\alpha \geq \alpha_1$ *, it holds that* 

<span id="page-36-6"></span>(5.20) 
$$
\sup_{x \in \Omega} |v_f(x) - v_f)(x)| + \sup_{x \in \Omega} \delta_{\Omega}(x) |\nabla (v_f - v_f))(x)| \lesssim \varepsilon \|f\|_{\text{BMO}(\sigma)},
$$

<span id="page-36-7"></span>(5.21) 
$$
\sup_{\xi \in \partial \Omega} C_s(\nabla u)(\xi) + \sup_{\xi \in \partial \Omega} \left[ C_s(\delta_{\Omega} |\nabla u|^2)(\xi) \right]^{1/2} \lesssim_{\varepsilon} ||f||_{\text{BMO}(\sigma)},
$$

<span id="page-36-8"></span>(5.22) 
$$
\sup_{x \in \Omega} \delta_{\Omega}(x) |\nabla u(x)| \lesssim ||f||_{\text{BMO}(\sigma)},
$$

*where the implicit constants depend on* s*,* n*, and the constants of the Ahlfors regularity, the corkscrew condition, and the Whitney decomposition.*

*Proof.* The result follows immediatelly by the estimates [\(5.7\)](#page-32-8), [\(5.8\)](#page-32-6) and [\(5.9\)](#page-32-3).  $\Box$ 

<span id="page-36-1"></span>**Theorem 5.5.** *If*  $f \in L^p(\sigma)$ *, for*  $p \in (1,\infty]$ *, then for any*  $\varepsilon > 0$ *, there exists*  $u = u_{\varepsilon} \in$  $C^{\infty}(\Omega)$ ,  $\alpha_0 \geq 1$ , and a constant  $c_{\varepsilon} > 1$ , such that for any  $\alpha \geq \alpha_0$ , it holds that

<span id="page-36-3"></span>
$$
(5.23) \quad \|\mathcal{N}_{\alpha}(u-v_f)\|_{L^p(\sigma)} + \|\mathcal{N}_{\alpha}(\delta_{\Omega}\nabla(u-v_f)\|_{L^p(\sigma)} \lesssim \varepsilon \|f\|_{L^p(\sigma)},
$$

<span id="page-36-4"></span>(5.24) 
$$
\|\mathcal{N}_{\alpha}(\delta_{\Omega}\nabla u)\|_{L^{p}(\sigma)} \lesssim \|f\|_{L^{p}(\sigma)},
$$

<span id="page-36-5"></span>(5.25) 
$$
\|\mathcal{C}_s(\nabla u)\|_{L^p(\sigma)} \lesssim \varepsilon^{-2} \|f\|_{L^p(\sigma)},
$$

<span id="page-36-2"></span>
$$
(5.26) \t\t ||\left[\mathcal{C}_s(\delta_{\Omega}|\nabla u|^2)\right]^{1/2}||_{L^q(\sigma)} \lesssim \varepsilon^{-2} ||f||_{L^q(\sigma)}, \quad q \in [2,\infty),
$$

*where the implicit constants depend on* s*,* n*,* p*,* q*, and the constants of the Ahlfors regularity, the corkscrew condition, and the Whitney decomposition.*

*Proof.* The proof is an immediate consequence of  $(5.3)$ ,  $(5.4)$ ,  $(5.5)$ ,  $(5.6)$ , Lemma [2.3,](#page-12-3) and the fact that M and M are  $L^p(\sigma) \to L^p(\sigma)$ -bounded for any  $p \in (1,\infty)$ . In order to prove  $(5.26)$  for  $q = 2$ , we should estimate it directly using  $(5.16)$  as in the proof of  $(5.5)$  along with the fact that

<span id="page-36-9"></span>
$$
(5.27) \qquad \|\mathcal{C}_s(\delta_\Omega |\nabla F|^2)^{1/2})\|_{L^2(\sigma)}^2 \approx \int_\Omega \delta_\Omega(x) \sup_{y \in B_x} |\nabla F(y)|^2 \frac{dx}{\delta_\Omega(x)^{n-s}}.
$$

Indeed,

$$
\int_{\Omega} \delta(x) \sup_{y \in B_x} |\nabla u(y)|^2 \frac{dx}{\delta_{\Omega}(x)^{n-s}} \lesssim \sum_{P \in \mathcal{W}(\Omega)} \int_{P} \ell(P) \sup_{y \in B_x} |\nabla u(y)|^2 \frac{dx}{\delta_{\Omega}(x)^{n-s}}
$$
  

$$
\lesssim \Big( \sum_{P \in \mathcal{B}_0} + \sum_{R \in \mathsf{Top}} \sum_{\substack{P \in \mathcal{N}(\mathcal{P}_0) \\ b(P) \notin \text{Tree}(R)}} \Big) \int_{P} m_{\sigma, B(x_{b(P)}, C_w \ell(P))} (|f|)^2 \frac{dx}{\ell(P)^{n+1-s}}
$$
  

$$
+ 2\varepsilon^2 \sum_{R \in \mathsf{Top}} \sum_{\substack{P \in \mathcal{P}_0 \\ b(P) \in \text{Tree}(R)}} \int_{P} \sup_{\rho \gtrsim \ell(P)} m_{\sigma, B(x_{b(P)}, \rho)} (|f|)^2 \frac{dx}{\ell(P)^{n+1-s}}
$$

$$
+ \varepsilon^2 \sum_{R \in \text{Top}} \sum_{P \in \mathcal{N}(P_0) \backslash P_0} \int_P \sup_{\rho \gtrsim \ell(P)} m_{\sigma, B(x_{b(P)}, \rho)}(|f|)^2 \frac{dx}{\ell(P)^{n+1-s}} \n\lesssim \Big( \sum_{P \in \mathcal{B}_0} + \sum_{R \in \text{Top}} \sum_{P \in \mathcal{N}(P_0)} \int \sigma(b(P)) m_{\sigma, B(x_{b(P)}, C_w\ell(P))}(|f|)^2 \n+ \sum_{R \in \text{Top}} \sum_{P \in \mathcal{N}(P_0)} \sigma(b(P)) \sup_{\rho \gtrsim \ell(P)} m_{\sigma, B(x_{b(P)}, \rho)}(|f|)^2 \n\lesssim \sum_{Q \in \mathcal{B} \cup \mathcal{J}} \sigma(Q) \sup_{\rho \gtrsim \ell(Q)} m_{\sigma, B(x_Q, \rho)}(|f|)^2 \n\lesssim \int_{\partial \Omega} \sup_{Q \ni z} \sum_{\rho \gtrsim \ell(Q)} m_{\sigma, B(x_Q, \rho)}(|f|)^2 d\sigma(z) \lesssim \int_{\partial \Omega} \widetilde{\mathcal{M}} f^2 d\sigma \lesssim \|f\|_{L^2(\sigma)}^2.
$$

**Remark 5.6.** Note that since  $L^{\infty}(\sigma) \subset BMO(\sigma)$ , the estimates [\(5.23\)](#page-36-3), [\(5.24\)](#page-36-4), and [\(5.25\)](#page-36-5) for  $p = \infty$ , follow from [\(5.20\)](#page-36-6), [\(5.21\)](#page-36-7), and [\(5.22\)](#page-36-8).

 $\Box$ 

## <span id="page-37-0"></span>6. CONSTRUCTION OF VAROPOULOS-TYPE EXTENSIONS OF  $L^p$  and BMO **FUNCTIONS**

We will first construct extensions of boundary functions that belong to BMO.

# Hypothesis  $[\tilde{T}]$

<span id="page-37-1"></span>(i) There exists a bounded linear *trace* operator

$$
\text{Tr}: \mathrm{C}^{1,\infty}_{s,\infty}(\Omega) \to \text{BMO}(\sigma)
$$

such that  $|| \text{Tr}(w) ||_{\text{BMO}(\sigma)} \lesssim || \nabla w ||_{\text{C}^{\infty}_{s,\infty}(\Omega)}$ .

<span id="page-37-2"></span>(ii) If  $v_f$  is the regularized dyadic extension of  $f \in BMO(\sigma)$ , then  $Tr(v_f)(\xi) = f(\xi)$ for  $\sigma$ -a.e.  $\xi \in \partial \Omega$ , and for any  $w \in C^{\infty}_{s,\infty}(\Omega)$ , it holds that

(6.1) 
$$
||f - \text{Tr}(w)||_{\text{BMO}(\sigma)} = || \text{Tr}(v_f - w)||_{\text{BMO}(\sigma)} \lesssim \sup_{x \in \Omega} |v_f(x) - w(x)|.
$$

<span id="page-37-3"></span>**[T](#page-37-1)heorem 6.1.** *Let*  $\Omega \in AR(s)$  *for*  $s \in (0, n]$  *satisfying the* Hypothesis  $[\tilde{T}]$ *. If*  $f \in$ BMO( $\sigma$ ), then there exist a function  $u: \Omega \to \mathbb{R}$  and a constant  $c_0 \in (0, \frac{1}{2})$  $\frac{1}{2}$  *such that for every*  $c \in (0, c_0]$ *, it holds that* 

- (i)  $u \in C^1(\Omega)$ . (ii) sup  $\sup_{\xi \in \partial \Omega} \mathcal{N}_{\sharp,c}(u)(\xi) + \sup_{x \in \Omega} \delta_{\Omega}(x) |\nabla u(x)| \lesssim ||f||_{\text{BMO}(\sigma)},$
- (iii)  $\sup_{s,c} \mathcal{C}_{s,c}(\nabla u)(\xi) \lesssim ||f||_{\text{BMO}(\sigma)}$ , and ξ∈∂Ω
- (iv)  $\text{Tr}(u)(\xi) = f(\xi)$  *for*  $\sigma$ *-a.e.*  $\xi \in \partial\Omega$ *.*

*Proof.* If  $f \in \text{BMO}(\sigma)$  and  $v_f$  its regularized dyadic extension, we apply Theorem [5.5](#page-36-1) and construct the  $\varepsilon$ -approximating function of  $v_f$ , which we denote by  $u_0$ . In light of [\(5.21\)](#page-36-7), and Hypothesis  $[\tilde{T}]$  $[\tilde{T}]$  $[\tilde{T}]$ , we have that the trace  $\text{Tr}(u_0)$  exists and it is in  $\text{BMO}(\sigma)$ . We set

$$
f_1 := f - \text{Tr}(u_0).
$$

Inductively, for every  $k \geq 1$ , we define  $u_k$  to be the  $\varepsilon$ -approximating function of  $v_{f_k}$  and set

$$
f_{k+1} := f_k - \text{Tr}(u_k),
$$

Therefore by  $(6.1)$  and  $(5.20)$ , we have that

<span id="page-38-0"></span>
$$
||f_{k+1}||_{\text{BMO}(\sigma)} \lesssim \sup_{x \in \Omega} |v_{f_k}(x) - u_k(x)| \lesssim \varepsilon ||f_k||_{\text{BMO}(\sigma)},
$$

which implies that

(6.2) 
$$
||f_{k+1}||_{\text{BMO}(\sigma)} \leq C\varepsilon ||f_k||_{\text{BMO}(\sigma)} \leq \cdots \leq (C\varepsilon)^{k+1} ||f||_{\text{BMO}(\sigma)}.
$$

Assume that  $C\varepsilon \leq 1/2$  and set  $S_k := \sum_{j=0}^k u_j$ , for any positive integer k. Using [\(5.20\)](#page-36-6) and  $(3.7)$ , and finally  $(6.2)$ , we can estimate

$$
||S_{k} - S_{m}||_{N_{sum}^{\infty}(\Omega)} = ||S_{k} - S_{m}||_{N_{\sharp}^{\infty}(\Omega)} + ||\delta_{\Omega}\nabla S_{k} - \delta_{\Omega}\nabla S_{m}||_{N^{\infty}(\Omega)}
$$
  
\n
$$
\lesssim \sum_{j=k+1}^{m} \left( \sup_{x \in \Omega} |v_{f_{j}}(x) - u_{j}(x)| + \sup_{x \in \Omega} \delta_{\Omega}(x)|\nabla(v_{f_{j}} - u_{j})(x)| \right)
$$
  
\n
$$
+ \sum_{j=k+1}^{m} \left( \sup_{\xi \in \partial\Omega} \mathcal{N}_{\sharp}(v_{f_{j}})(\xi) + \sup_{x \in \Omega} \delta_{\Omega}(x)|\nabla u_{j}(x)| \right)
$$
  
\n
$$
\lesssim \sum_{j=k+1}^{m} \left( \varepsilon ||f_{j}||_{\text{BMO}(\sigma)} + ||f_{j}||_{\text{BMO}(\sigma)} + ||f_{j}||_{\text{BMO}(\sigma)} \right) \lesssim \sum_{j=k+1}^{m} (C\varepsilon)^{j} ||f||_{\text{BMO}(\sigma)}
$$
  
\n
$$
\leq (2^{-k} - 2^{-m}) ||f||_{\text{BMO}(\sigma)}.
$$

Thus,  $S_k$  is Cauchy sequence in the  $(N_{sum}(\Omega), \| \cdot \|_{sum})$  and so, by Corollary [A.9,](#page-68-0) there exists  $u \in N_{sum}(\Omega)$  such that  $S_k \to u$  in  $N_{sum}(\Omega)$  (this implies that  $\nabla S_k \to \nabla u$  locally uniformly in  $\Omega$ ). By [\(5.21\)](#page-36-7) and [\(5.22\)](#page-36-8), for any  $m > k$ , we have that

$$
\|\nabla S_k - \nabla S_m\|_{C^{\infty}_{s,\infty}(\Omega)} \leq \sum_{j=k+1}^m \sup_{\xi \in \partial \Omega} C_{s,c}(\nabla u_j)(\xi)
$$
  

$$
\lesssim \sum_{j=k+1}^m \|f_j\|_{\text{BMO}(\sigma)} \leq (2^{-k} - 2^{-m})\|f\|_{\text{BMO}(\sigma)}.
$$

Therefore, since  $\mathrm{C}_{s,\infty}^{\infty}(\Omega)$  is a Banach space, there exists  $\vec{F}\in\mathrm{C}_{s,\infty}^{\infty}(\Omega)$  such that  $\nabla S_k\to\bar{F}$ in  $C_{s,\infty}^{\infty}(\Omega)$  and, by Lemma [A.3,](#page-60-0) for any fixed  $x \in \Omega$ ,

$$
\sup_{y \in B_x} |\nabla S_k(y) - \vec{F}(y)| \delta_{\Omega}(y) \to 0, \quad \text{as } k \to \infty,
$$

which implies that  $\vec{F} = \nabla u$  in  $\Omega$ . Therefore,

$$
\nabla S_k \to \nabla u(x) \quad \text{in } \mathrm{C}_{s,\infty}^{\infty}(\Omega) \cap \mathrm{N}_{\mathrm{sum}}^{\infty}(\Omega).
$$

Furthermore, we have that

(6.3) 
$$
\sup_{\xi \in \partial \Omega} C_s(\nabla u)(\xi) \leq \sum_{k=0}^{\infty} \sup_{\xi \in \partial \Omega} C_s(\nabla u_k)(\xi) \lesssim_{\varepsilon} \sum_{k=0}^{\infty} ||f_k||_{\text{BMO}(\sigma)} \lesssim ||f||_{\text{BMO}(\sigma)},
$$

and, similarly,

$$
||u||_{N^{\infty}_{sum}(\Omega)} \leq \sum_{k=0}^{\infty} ||u_k||_{N^{\infty}_{sum}(\Omega)} \lesssim \sum_{k=0}^{\infty} ||f_k||_{BMO(\sigma)} \lesssim ||f||_{BMO(\sigma)}.
$$

Finally, it holds that

$$
0 = \lim_{k \to \infty} ||f_{k+1}||_{\text{BMO}(\sigma)} = ||f - \sum_{j=0}^{k} \text{Tr}(u_j)||_{\text{BMO}(\sigma)}
$$

and since, by construction,  $\sum_{j=0}^{k} u_j - u \in C_{s,\infty}^{1,\infty}(\Omega)$ , using the linearity of the trace and Hypothesis  $[\tilde{T}]$  $[\tilde{T}]$  $[\tilde{T}]$ , we have that

$$
\left\| \text{Tr}(u) - \sum_{j=0}^{k} \text{Tr}(u_j) \right\|_{\text{BMO}(\sigma)} = \left\| \text{Tr}\left( u - S_k \right) \right\|_{\text{BMO}(\sigma)} \lesssim \left\| \nabla u - \nabla S_k \right\|_{\text{C}_{s,\infty}^{\infty}(\Omega)} \stackrel{k \to \infty}{\longrightarrow} 0.
$$

This gives that u is an extension of  $f \in BMO(\sigma)$  with  $Tr(u) = f$  in  $BMO(\sigma)$  and so

(6.4) 
$$
\text{Tr}(u)(\xi) = f(\xi) + c \text{ for } \sigma\text{-almost every } \xi \in \partial\Omega.
$$

If we set  $\tilde{u} = u - c$ , it is clear that  $\tilde{u}$  satisfies (i), (ii), (iii), and (iv) concluding the proof of the theorem. the theorem.

**Remark 6.2.** Note that by the proof of Theorem [6.1,](#page-37-3) it is immediate that the extension u satisfies the estimate

(6.5) 
$$
\sup_{x \in \Omega} |v_f(x) - u(x)| \lesssim \|f\|_{\text{BMO}(\sigma)}.
$$

We shall now deal with boundary data in  $L^p(\sigma)$ .

### Hypothesis [T]

<span id="page-39-0"></span>(i) For  $p \in (1, \infty]$ , there exists a bounded linear *trace* operator

<span id="page-39-2"></span>Tr: 
$$
\mathcal{N}^p(\Omega) \cap C^{1,p}_{s,\infty}(\Omega) \to L^p(\sigma)
$$

<span id="page-39-1"></span>such that  $|| \text{Tr}(w) ||_{L^p(\sigma)} \leq ||w||_{N^p(\Omega)}$ .

(ii) If  $v_f$  the regularized dyadic extension of  $f \in L^p(\sigma)$ , then  $\text{Tr}(v_f)(\xi) = f(\xi)$  for  $\sigma$ -a.e.  $\xi \in \partial \Omega$ , and for any  $w \in N^p(\Omega) \cap C^p_{s,\infty}(\Omega)$ , it holds that

(6.6) 
$$
||f - \text{Tr}(w)||_{L^p(\sigma)} = || \text{Tr}(v_f - w)||_{L^p(\sigma)} \le ||v_f - w||_{N^p(\Omega)}.
$$

<span id="page-39-3"></span>**[T](#page-39-0)heorem 6.3.** *Let*  $\Omega \in AR(s)$  *for*  $s \in (0, n]$  *satisfying the* Hypothesis [T]. *If*  $f \in L^p(\sigma)$ *with*  $p \in (1, \infty]$ *, then there exists a function*  $u : \Omega \to \mathbb{R}$  *such that* 

 $(i)$   $u \in C^1(\Omega)$ , *(ii)*  $\text{Tr}(u)(\xi) = f(\xi)$  *for*  $\sigma$ *-a.e.*  $\xi \in \partial\Omega$ *,*  $(iii)$   $\|\mathcal{N}(u)\|_{L^p(\sigma)} + \|\mathcal{N}(\delta_{\Omega}\nabla u)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)},$ 

$$
(iv) \|\mathcal{C}_s(\nabla u)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)}.
$$

*Proof.* Fix  $\varepsilon > 0$  to be chosen. By Theorem [5.5,](#page-36-1) we construct  $u_0$ , the  $\varepsilon$ -approximating function of  $v_f$ , and by Hypothesis [[T](#page-39-0)], the trace of  $u_0$  exists and  $\text{Tr}(u_0) \in L^p(\sigma)$ . We set

$$
f_1 := f - \text{Tr}(u_0) \in L^p(\sigma).
$$

We then let  $u_1$  be the  $\varepsilon$ -approximating function of  $v_{f_1}$  and set

$$
f_2 := f_1 - \text{Tr}(u_1) \in L^p(\sigma).
$$

Inductively, for every  $k \geq 1$ , we define  $u_k$  to be the  $\varepsilon$ -approximating function of  $v_{f_k}$  and set  $f_{k+1} := f_k - \text{Tr}(u_k)$ . Therefore, by [\(6.6\)](#page-39-1) and [\(5.23\)](#page-36-3), we have that

$$
||f_{k+1}||_{L^p(\sigma)} \leq ||v_{f_k} - u_k||_{\mathcal{N}^p(\Omega)} \leq C \varepsilon ||f_k||_{L^p(\sigma)},
$$

which implies that

<span id="page-40-0"></span>(6.7) 
$$
||f_{k+1}||_{L^p(\sigma)} \leq C \varepsilon ||f_k||_{L^p(\sigma)} \leq \cdots \leq (C\varepsilon)^{k+1} ||f||_{L^p(\sigma)}.
$$

Thus, if we choose  $\varepsilon$  so that  $C\varepsilon \leq \frac{1}{2}$  and set  $S_k := \sum_{j=0}^k u_j$ , then for  $k < m$ , using [\(5.23\)](#page-36-3),  $(3.6)$ , and  $(6.7)$ , it holds that

$$
||S_{k} - S_{m}||_{N^{p}(\Omega)} \leq \sum_{j=k+1}^{m} (||\mathcal{N}(u_{j} - v_{f_{j}})||_{L^{p}(\sigma)} + ||\mathcal{N}v_{f_{j}}||_{L^{p}(\sigma)})
$$
  

$$
\lesssim \sum_{j=k+1}^{m} (\varepsilon ||f_{j}||_{L^{p}(\sigma)} + ||f_{j}||_{L^{p}(\sigma)}) \leq (1 + \varepsilon) \sum_{j=k+1}^{m} (C\varepsilon)^{j} ||f||_{L^{p}(\sigma)}
$$
  

$$
\leq (2^{-k+1} - 2^{-m+1}) ||f||_{L^{p}(\sigma)}.
$$

Thus,  $S_k$  is a Cauchy sequence in  $N^p(\Omega)$ , and since  $N^p(\Omega)$  is a Banach space, there exists  $u \in N^p(\Omega)$  such that  $S_k \to u$  in  $N^p(\Omega)$ . It is easy to see that  $S_k \to u$  uniformly in  $B_x$ , for any  $x \in \Omega$ , and so we define

(6.8) 
$$
u(x) = \sum_{k=0}^{\infty} u_k(x) \text{ for all } x \in \Omega.
$$

Similarly, we can show that  $\nabla S_k = \sum_{j=0}^k \nabla u_j$  is convergent in the Banach space  $C_{s,\infty}^p(\Omega)$ (resp.  $N^p(\Omega)$ ), since, by [\(5.25\)](#page-36-5) (resp. [\(5.24\)](#page-36-4)) and [\(6.7\)](#page-40-0),

$$
\|\nabla S_k - \nabla S_m\|_{\mathcal{C}^p_{s,\infty}(\Omega)} + \|\delta_{\Omega} \nabla S_k - \delta_{\Omega} \nabla S_m\|_{\mathcal{N}^p(\Omega)}\n\n\leq \sum_{j=k+1}^m \|\mathcal{C}_s(\nabla u_j)\|_{L^p(\sigma)} + \sum_{j=k+1}^m \|\mathcal{N}(\delta_{\Omega}|\nabla u_j|)\|_{L^p(\sigma)} \leq C \varepsilon^{-2} \sum_{j=k+1}^m \|f_j\|_{L^p(\sigma)}.
$$

Thus, there exists  $\vec{F}_1 \in C_{s,\infty}^p(\Omega)$  (resp.  $\vec{F}_2$  so that  $\delta_{\Omega} \vec{F}_2 \in N^p(\Omega)$ ) such that  $\nabla S_k \to \vec{F}_1$ in  $C_{s,\infty}^p(\Omega)$  (resp.  $\delta_{\Omega} \nabla S_k \to \delta_{\Omega} \vec{F}_2$  in  $N^p(\Omega)$ ), and, by Lemma [A.3,](#page-60-0) we have that for any fixed  $x \in \Omega$ ,

$$
\sup_{y \in B_x} |\nabla S_k - F_i| \delta_{\Omega}(y) \to 0, \quad \text{for } i = 1, 2,
$$

which readily implies that  $\vec{F}_1 = \vec{F}_2 =: \vec{F}$  in  $\Omega$ . Hence,  $\nabla S_k$  converges to  $\sum_{k=0}^{\infty} \nabla u_k$ uniformly in  $B_x$  for every  $x \in \Omega$ , and by Lemma [A.2,](#page-58-0) we deduce that  $u \in C^1(\Omega)$  and

$$
\sum_{k=0}^{\infty} \nabla u_k(x) = \nabla u(x) \quad \text{for all } x \in \Omega.
$$

In fact,

$$
\|\mathcal{N}(u)\|_{L^p(\sigma)} + \|\mathcal{N}(\delta_{\Omega}\nabla u)\|_{L^p(\sigma)} + \|\mathcal{C}_s(\nabla u)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)}.
$$

To show that u is an extension of f, notice first that, in light of  $(6.7)$ ,

$$
0 = \lim_{k \to \infty} ||f_{k+1}||_{L^p(\sigma)} = \lim_{k \to \infty} ||f - \text{Tr}\left(\sum_{j=0}^k u_j\right)||_{L^p(\sigma)}.
$$

Since, by construction,  $\sum_{j=0}^{k} u_j - u \in N^p(\Omega)$ , in light of Hypothesis [[T](#page-39-0)] (i), we get that

$$
\| \operatorname{Tr} \left( \sum_{j=0}^{k} u_j \right) - \operatorname{Tr}(u) \|_{L^p(\sigma)} = \| \operatorname{Tr} \left( \sum_{j=0}^{k} u_j - u \right) \|_{L^p(\sigma)} \leq \| \sum_{j=0}^{k} u_j - u \|_{N^p(\Omega)} \stackrel{k \to \infty}{\longrightarrow} 0,
$$

 $\Box$ 

which entails that  $\text{Tr}(u)(\xi) = f(\xi)$  for  $\sigma$ -almost every  $\xi \in \partial \Omega$ .

<span id="page-41-0"></span>**Proposition 6.4.** *Let*  $\Omega \in AR(s)$  *for*  $s \in (0, n]$ *, which for*  $s = n$  *satisfies the pointwise John condition. Then, for any*  $p \in (1,\infty]$ *, there exists a bounded linear trace operator*  $\text{Tr}\,\Omega:\mathcal{N}^p(\Omega)\cap\mathrm{C}^{1,p}_{s,\infty}(\Omega)\to L^p(\sigma)$  $\text{Tr}\,\Omega:\mathcal{N}^p(\Omega)\cap\mathrm{C}^{1,p}_{s,\infty}(\Omega)\to L^p(\sigma)$  $\text{Tr}\,\Omega:\mathcal{N}^p(\Omega)\cap\mathrm{C}^{1,p}_{s,\infty}(\Omega)\to L^p(\sigma)$  satisfying the Hypothesis [T]. Moreover, if  $\Omega$  satisifies *the local John condition for*  $s = n$ , there exists a bounded linear trace operator  $Tr \Omega$ :  $C^{1,\infty}_{s,\infty}(\Omega) \to \text{BMO}(\sigma).$ 

*Proof.* For any  $x \in \Omega$  and fixed  $c \in (0, 1/2]$ , we define

(6.9) 
$$
E(x) := \int_{B(x,c\delta_{\Omega}(x))} u(z) dz.
$$

Fix  $\xi \in \partial \Omega$  such that  $\xi \in JC(\theta)$  (see Definition [2.13\)](#page-18-0). Then there exist  $r_{\xi} > 0$  and  $x_{\xi} \in B(\xi, 2r_{\xi}) \cap \Omega$  such that  $\delta_{\Omega}(x_{\xi}) \ge \theta r_{\xi}$ , and also there exists a good curve  $\gamma : [0, 1] \to \mathbb{R}$ in  $B(\xi, 2r_{\xi}) \cap \Omega$  connecting the points  $\xi$  and  $x_{\xi}$  such that  $|\dot{\gamma}(t)| = 1 \ \forall t \in [0, 1]$ . For any fixed points  $x_1, x_2 \in \gamma$ , there exist  $t_1, t_2 \in [0, 1]$  such that  $x_1 = \gamma(t_1)$  and  $x_2 = \gamma(t_2)$ . Applying a change of variables and applying the mean value theorem, we estimate

$$
|E(x_1) - E(x_2)| = \Big| \int_{B(0,1)} \left( u(x_1 + wc\delta_{\Omega}(x_1)) - u(x_2 + wc\delta_{\Omega}(x_2)) \right) dw \Big|
$$
  
\n
$$
= \Big| \int_{B(0,1)} \left( u(\gamma(t_1) + wc\delta_{\Omega}(\gamma(t_1))) - u(\gamma(t_2) + wc\delta_{\Omega}(\gamma(t_2))) \right) dw \Big|
$$
  
\n
$$
= \Big| \int_{B(0,1)} \int_{t_1}^{t_2} \nabla u(\gamma(t) + wc\delta_{\Omega}(\gamma(t))) \cdot \nabla \delta_{\Omega}(\gamma(t)) \dot{\gamma}(t) dt dw \Big|
$$
  
\n
$$
\leq \int_{t_1}^{t_2} \int_{B(0,1)} \Big| \nabla u(\gamma(t) + wc\delta_{\Omega}(\gamma(t)) \Big| dw dt
$$

where we used that  $|\dot{\gamma}(t)| = 1$  and  $|\nabla \delta_{\Omega}(\gamma(t))| \leq 1$  since the function dist( $\cdot$ ,  $\partial \Omega$ ) is 1-Lipschitz. Note that there exists  $M_j \in \mathbb{N}$  such that  $2^{-M_j} \le t_j \le 2^{-M_j+1}$ , for  $j = 1, 2$ . By the Fundamental Theorem of Calculus, since  $u \in C^1(\Omega)$  and  $\gamma(t) + w c \delta_{\Omega}(\gamma(t))$  is  $(1 + c)$ -Lipschitz in t for any  $w \in B(0, 1)$ , we have that there exists  $s_k \in [2^{-k}, 2^{1-k}]$  such that

$$
\int_{t_1}^{t_2} \int_{B(0,1)} \left| \nabla u(\gamma(t) + wc \delta_{\Omega}(\gamma(t)) \right| dw dt
$$
\n
$$
= \sum_{k=M_1}^{M_2} \int_{2^{-k}}^{2^{-k+1}} \int_{B(0,1)} \left| \nabla u(\gamma(t) + wc \delta_{\Omega}(\gamma(t)) \right| dw dt
$$
\n
$$
= \sum_{k=M_1}^{M_2} 2^{-k} \int_{B(0,1)} \left| \nabla u(\gamma(s_k) + wc \delta_{\Omega}(\gamma(s_k)) \right| dw
$$
\n
$$
= \sum_{k=M_1}^{M_2} 2^{-k} \int_{B(\gamma(s_k), c\delta_{\Omega}(\gamma(s_k)))} \left| \nabla u(y) \right| \frac{dy}{(c\delta_{\Omega}(\gamma(s_k)))^{n+1}}
$$
\n
$$
\lesssim \sum_{k=M_1}^{M_2} \int_{B(\gamma(s_k), c\delta_{\Omega}(\gamma(s_k)))} \left| \nabla u(y) \right| \frac{dy}{\delta_{\Omega}(y)^n},
$$

where is the last inequality we used that  $\delta_\Omega(\gamma(s_k)) \approx s_k \approx 2^{-k}$  and that  $\delta_\Omega(y) \lesssim \delta_\Omega(\gamma(s_k))$ for any  $y \in B(\gamma(s_k), c\delta_{\Omega}(\gamma(s_k)))$ . Therefore, there exists a cone  $\gamma_{\alpha}(\xi)$  with apperture depending on c and  $\theta$  such that  $B(\gamma(s_k), c\delta_{\Omega}(\gamma(s_k))) \subset \gamma_{\alpha}(\xi)$ , and, by the bounded overlaps of the balls  $B(\gamma(s_k), c\delta_{\Omega}(\gamma(s_k)))$ , we infer that

$$
|E(x_1) - E(x_2)| \lesssim \int_{\gamma_\alpha(\xi) \cap B(\xi, C\delta_\Omega(x_2))} |\nabla u(y)| \frac{dy}{\delta_\Omega(y)^n}.
$$

By [\(2.23\)](#page-13-1), we have that  $\mathcal{A}_s^{(\alpha)}(\nabla u) \in L^p(\sigma)$  for  $p \in (1,\infty)$  and  $\mathcal{A}_s^{(\alpha)}(\nabla u) \in L^q_{loc}(\sigma)$  for any  $q \in (1,\infty)$  when  $p = \infty$ . Thus,  $\mathcal{A}_s^{(\alpha)}(\nabla u)(\xi) < \infty$  for  $\sigma$ -almost every  $\xi \in \partial\Omega$  and using the fact that the above estimate holds for any points  $x_1, x_2 \in \gamma_{\xi}$ , we can assume that  $x_1, x_2 \in B(\xi, \varepsilon)$  for some  $\varepsilon > 0$  small compared to  $r_{\xi}$ . Therefore,

$$
|E(x_1)-E(x_2)| \lesssim \int_{\gamma_\alpha(\xi)\cap B(\xi,C\varepsilon)} |\nabla u(y)| \, \delta_\Omega(y)^{-n} \, dy \lesssim \mathcal{A}^{(\alpha)}(\nabla u)(\xi) < \infty.
$$

By the dominated convergence theorem, we get that  $|E(x_1) - E(x_2)| \to 0$  as  $\varepsilon \to 0$ , i.e.,  $E(x)$  is Cauchy on  $\gamma_{\xi}$  and thus convergent. This shows that the quasi-non-tangential limit of  $E(x)$  at  $\xi \in \partial \Omega$  exists for  $\sigma$ -a.e.  $\xi \in \partial \Omega$  and we can define the desired trace operator by

(6.10) 
$$
\operatorname{Tr}_{\Omega}(u)(\xi) := \operatorname{qnt-lim}_{x \to \xi} E(x) \quad \text{for } \sigma\text{-a.e. } \xi \in \partial\Omega.
$$

In the case  $s < n$ , we just define  $\text{Tr}_{\Omega}(u)(\xi) = nt\text{-}\lim_{x\to\xi} E(x)$  since  $\Omega$  has only one connected component and any  $\xi \in \partial \Omega$  can be connected to a corkscrew point by a good curve. It is clear that Tr :  $C_{s,\infty}^{1,p}(\Omega) \to L^p(\sigma)$  is a linear operator, while the fact that  $\text{Tr}: \mathbb{N}^p(\Omega) \cap \mathbb{C}^{1,p}_{s,\infty}(\Omega) \to L^p(\sigma)$  is bounded if  $p \in (1,\infty]$  can be proved pretty easily.

Indeed, let  $\xi \in \partial \Omega$  such that  $\xi \in JC(\theta)$ . For fixed  $\varepsilon > 0$ , there exists  $\delta > 0$  such that, if  $x \in B(\xi, \delta) \cap \gamma_{\xi}$ , it holds that

$$
|\operatorname{Tr}_{\Omega}(u)(\xi)| \leq |\operatorname{Tr}_{\Omega}(u)(\xi) - E(x)| + m_{\sigma, B(x, c\delta_{\Omega}(x))}(|u|) < \varepsilon + \mathcal{N}(u)(\xi).
$$

Letting  $\varepsilon \to 0$  we infer that  $|\text{Tr}_{\Omega}(u)(\xi)| \leq \mathcal{N}(u)(\xi)$  for  $\sigma$ -a.e.  $\xi \in \partial\Omega$ , which readily concludes Hypothesis  $[T]$  $[T]$  $[T]$  (i), while (ii) readily follows from Lemma [3.1.](#page-21-2)

Assume now that  $\Omega$  satisfies the local John condition when  $s = n$  (in the case  $s < n$ ) this is automatic). Fix  $\xi \in \partial \Omega$  and  $r > 0$ , and, by the local John condition, there exists a corkscrew point  $x_r \in B(\xi, r)$  such that any  $\zeta \in B(\xi, r) \cap \partial \Omega$  can be connected to  $x_r$  by a good curve. The existence of the trace operator follows by the same argument as above and we define it the same way. It remains to show that  $\text{Tr}: C_{s,\infty}^{1,\infty}(\Omega) \to \text{BMO}(\sigma)$  is bounded. For  $u \in C^{1,\infty}_{s,\infty}(\Omega)$ , if  $B_r := B(x_r, c\delta_{\Omega}(x_r))$  is a corkscrew ball centered at  $x_r$  with radius  $c\delta_{\Omega}(x_r) \approx r$ , by the same proof as above, we can show that

$$
|\operatorname{Tr}_{\Omega}(u)(\zeta)-\int_{B_r}u(y)\,dy|\lesssim\mathcal{A}_s(\nabla u\mathbf{1}_{B(\xi,C'r)})(\zeta),\quad\forall\zeta\in B(\xi,r)\cap\partial\Omega.
$$

Thus, taking averages over the ball  $B(\xi, r)$  and applying [\(2.22\)](#page-13-2) in  $L^1(B(\xi, C'r))$ , we conclude that

$$
\int_{B(\xi,r)} \left| \text{Tr}_{\Omega}(u) - \int_{B_r} u(y) \, dy \right| d\sigma \lesssim \int_{B(\xi,C''r)} C_s (\nabla u \mathbf{1}_{B(\xi,C''r)}) \, d\sigma \leq \|\mathcal{C}_s(\nabla u)\|_{L^{\infty}(\sigma)}.
$$

This readily implies that  $|| \text{Tr}_{\Omega}(u) ||_{\text{BMO}(\sigma)} \leq ||\mathcal{C}_s(\nabla u)||_{L^{\infty}(\sigma)}$ , which, combined with  $||f||_{\text{BMO}(\sigma)} \le 2||f||_{L^{\infty}(\sigma)}$ , [\(6.5\)](#page-39-2), and Hypothesis [[T](#page-39-0)] (ii) (we have already proved it above), proves Hypothesis  $[T]$  $[T]$  $[T]$ .

<span id="page-43-0"></span>We state [\[ST70,](#page-70-13) Theorem 2, p. 171] in the following lemma.

**Lemma 6.5.** Let  $E \subset \mathbb{R}^{n+1}$  be a closed set and  $\delta_E$  be the distance function with respect *to* E. Then there exist positive constants  $m_1$  and  $m_2$  and a function  $\beta_E$  defined in  $E^c$  such *that*

- (i)  $m_1 \delta_E(x) \leq \beta_E(x) \leq m_2 \delta_E(x)$ , for every  $x \in E^c$ , and
- (ii)  $\beta_E$  *is smooth in*  $E^c$  *and*

$$
\left|\frac{\partial^{\alpha}}{\partial x^{\alpha}}\beta_E(x)\right| \leq C_{\alpha}\beta_E(x)^{1-|\alpha|}.
$$

*In addition, the constants*  $m_1$ ,  $m_2$  *and*  $C_\alpha$  *are independent of E*.

Following [\[HT21,](#page-70-4) Section 3], we define a kernel  $\Lambda(\cdot, \cdot) : \Omega \times \Omega \to [0, \infty]$ , which will be necessary in the proof of Theorem [6.1.](#page-37-3) To this end, let  $\beta = \beta_{\Omega^c}$  be the function constructed in Lemma [6.5](#page-43-0) and let  $\zeta \ge 0$  be a smooth non-negative function supported on  $B(0, \frac{c}{4m})$  $\frac{c}{4m_2}$ ), satisfying  $\zeta \leq 1$  and  $\int \zeta = 1$ . For every  $\lambda > 0$ , we set

$$
\zeta_{\lambda}(x) := \lambda^{-(n+1)} \zeta(x/\lambda),
$$

and define the mollifier

$$
\Lambda(x,y) := \zeta_{\beta(x)}(x-y) = \frac{1}{\beta(x)^{n+1}} \zeta\left(\frac{2(x-y)}{\beta(x)}\right).
$$

Observe that, by construction, for every  $x \in \Omega$ ,

<span id="page-44-1"></span>(6.11) 
$$
\text{supp}(\Lambda(x,\cdot)) \subset \widetilde{B}_x := B(x,c\,\delta_{\Omega}(x)/8)
$$
 and  $\int_{\Omega} \Lambda(x,y) \,dy = 1.$ 

Moreover, it is easy to prove that

<span id="page-44-2"></span>(6.12) 
$$
\sup_{y \in \widetilde{B}_x} \Lambda(x, y) \lesssim \delta_{\Omega}(x)^{-n-1} \quad \text{and} \quad \sup_{y \in \widetilde{B}_x} |\nabla_x \Lambda(x, y)| \lesssim \delta_{\Omega}(x)^{-n-2}.
$$

For any  $F : \Omega \to \mathbb{R}$ , we define the *smooth modification* of F by

(6.13) 
$$
\widetilde{F}(x) := \int_{\Omega} \Lambda(x, y) F(y) dy.
$$

The next lemma was essentially proved in [\[HT21,](#page-70-4) Section 3] but we provide its proof for the reader's convenience.

<span id="page-44-4"></span>**Lemma 6.6.** Let  $\Omega \subset \mathbb{R}^{n+1}$  be an open set satisfying the corkscrew condition. If  $F \in$  $C^1(\Omega;\mathbb{R}^{n+1})$  and  $\tilde{F}$  is the smooth modification of F as defined in [\(6.13\)](#page-44-0), then we have that

(a) *For any*  $x \in \Omega$ ,

<span id="page-44-0"></span>
$$
|\widetilde{F}(x)| \lesssim \sup_{\widetilde{B}_x} |F(y)|.
$$

(b) *For any*  $x \in \Omega$ ,

$$
|\widetilde{F}(x_1)-\widetilde{F}(x_2)| \lesssim |x_1-x_2|\,\delta_{\Omega}(x)^{-1}\,m_{\sharp,c}(F)(x), \quad \text{for all } x_1, x_2 \in B_x.
$$

(c) *For any*  $x \in \Omega$ ,

$$
m_{\sharp,c}(\widetilde{F})(x) \lesssim m_{\sharp,c}(F)(x).
$$

(d) *For any*  $\xi \in \partial \Omega$ *,* 

$$
\sup_{x \in \gamma_{\alpha}(\xi)} \delta_{\Omega}(x) |\nabla \widetilde{F}(x)| \lesssim \mathscr{C}_s(\nabla \widetilde{F})(\xi).
$$

(e) *For any*  $\xi \in \partial \Omega$ ,

<span id="page-44-3"></span>
$$
\mathcal{C}_{s,c}(\nabla \widetilde{F})(\xi) \lesssim \mathscr{C}_s(\nabla F)(\xi).
$$

(f) *If* qnt-lim<sub>x→ξ</sub>  $F(x) = f(\xi)$  (resp. nt-lim<sub>x→ξ</sub>  $F(x) = f(\xi)$ ) *for*  $\sigma$ -a.e.  $\xi \in \partial\Omega$ , *then* qnt-lim<sub>x→ξ</sub>  $\widetilde{F}(x) = f(\xi)$  (resp. nt-lim<sub>x→ξ</sub>  $\widetilde{F}(x) = f(\xi)$ ) *for*  $\sigma$ -*a.e.*  $\xi \in \partial\Omega$ *.* 

*Proof.* (a) follows by definition. For (b), we let  $x_1, x_2 \in B_x$ . Then, by triangle inequality,

(6.14) 
$$
B(x_1, c\,\delta_{\Omega}(x_1)/8) \cup B(x_2, c\,\delta_{\Omega}(x_2)/8) \subset \frac{5}{4}B_x.
$$

Combining  $(6.11)$ ,  $(6.12)$ , and  $(6.14)$ , we get that

$$
\begin{aligned} |\widetilde{F}(x_1) - \widetilde{F}(x_2)| &= \left| \int (\Lambda(x_1, y) - \Lambda(x_2, y)) \left( F(y) - m_{\frac{5}{4}B_x} F \right) dy \right| \\ &\leq |x_1 - x_2| \sup_{z \in \frac{5}{4}B_x} \sup_{w \in B_x} |\nabla_w \Lambda(w, z)| \, m_{\frac{5}{4}B_x} \left( |F - m_{\frac{5}{4}B_x} F| \right) \, |\frac{5}{4}B_x| \\ &\lesssim |x_1 - x_2| \, \delta_{\Omega}(x)^{-1} \, m_{\sharp,c}(F)(x), \end{aligned}
$$

which proves (b) and thus (c). We turn our attention to (d) and fix  $\xi \in \partial\Omega$  and  $r > 0$ . For any  $z \in B(\xi, r) \cap \Omega$  and  $x \in B_z$ , using [\(6.11\)](#page-44-1), [\(6.12\)](#page-44-2), and Poincaré inequality, we can write

$$
|\nabla \widetilde{F}(x)| = \Big| \int \nabla_x \Lambda(x, y) F(y) \, dy \Big| = \Big| \int \nabla_x \Lambda(x, y) \big( F(y) - m_{\widetilde{B}_x} F \big) \, dy \Big|
$$
  

$$
\lesssim \delta_{\Omega}(x)^{-n-2} \int_{\widetilde{B}_x} |F(y) - m_{\widetilde{B}_x} F| \, dy \lesssim \int_{\widetilde{B}_x} |\nabla F| \, dy,
$$

which immediately implies (d). To prove (e), we first define

$$
A_k(\xi, r) := \{ x \in B(\xi, r) \cap \Omega : 2^{-k-1}r \le \delta_{\Omega}(x) < 2^{-k}r \}
$$
\n
$$
A_k^*(\xi, r) := \{ x \in B(\xi, r) \cap \Omega : 2^{-k-2}r \le \delta_{\Omega}(x) < 2^{-k+1}r \}
$$

and write

$$
\int_{B(\xi,r)\cap\Omega} \sup_{y\in B_x} |\nabla \widetilde{F}(y)| dy \leq \sum_{k=0}^{\infty} \int_{A_k(\xi,r)} \sup_{y\in B_x} |\nabla \widetilde{F}(y)| dy.
$$

As  $\bigcup_{y \in B_x} B_y \subset 2B_x$ , by Fubini's theorem,

$$
\int_{A_k(\xi,r)} \sup_{y \in B_x} |\nabla \widetilde{F}(y)| dy \lesssim \int_{A_k(\xi,r)} \int_{2B_x} |\nabla F(y)| dy \lesssim \int_{A_k^*(\xi,r)} |\nabla F(y)| dy,
$$

Summing over k and using that  $A_k^*(\xi, r)$  have bounded overlap, we get (e). Finally, (f) follows from [\[HT21,](#page-70-4) Lemma 3.14].

*proof of Theorem [1.3.](#page-4-0)* It is an immediate consequence of Theorem [6.3,](#page-39-3) Proposition [6.4,](#page-41-0) and Lemma [6.6.](#page-44-4)

Let us now turn our attention to the proof of Theorem [1.1.](#page-3-1) When  $s = n$  and  $\Omega$  satisfies the pointwise John condition but not the local John condition, we will need the following generalization of Garnett's Lemma, which was proved in [\[HT21,](#page-70-4) Lemma 10.1].

<span id="page-45-0"></span>**Lemma 6.7.** *Let*  $\Omega \in \text{AR}(n)$ ,  $Q_0 \in \mathcal{D}_{\sigma}$ , and let  $f \in \text{BMO}(\sigma)$  which vanishes on  $\partial \Omega \setminus Q_0$ *(if it is non-empty). Then, there exists a collection of cubes*  $\widetilde{S}(Q_0) = \{Q_i\}_i \subset \mathcal{D}(Q_0)$  *and coefficients*  $\alpha_i$  *such that* 

*(1)*  $\sup_j |\alpha_j| \lesssim ||f||_{\text{BMO}(\sigma)}$ *(2)*  $f = g + \sum_{j} \alpha_j \mathbf{1}_{Q_j}$ , where  $g \in L^{\infty}(\sigma)$  with  $||g||_{L^{\infty}(\sigma)} \lesssim ||f||_{\text{BMO}(\sigma)}$  and (3)  $\widetilde{S}(Q_0)$  *satisfies a Carleson packing condition.* 

*proof of Theorem [1.1.](#page-3-1)* Recall that if  $s < n$  then  $\Omega \in AR(s)$  is uniform and thus it satisfies the local John condition. Therefore, by Theorem [6.1,](#page-37-3) Proposition [6.4,](#page-41-0) and Lemma [6.6,](#page-44-4) we can construct the desired extension of Theorem [1.1](#page-3-1) when either  $s < n$ , or  $s = n$  and  $\Omega$ satisfies, in addition, the local John condition. We are left with the case  $s = n$  so that  $\Omega$  satisfies the pointwise John condition but not the local John condition. By Lemma [6.7,](#page-45-0) if  $f \in \text{BMO}(\sigma)$  with compact support in  $Q_0 \in \mathcal{D}_{\sigma}$ , there exists  $g \in L^{\infty}(\sigma)$  and  $b =$  $\sum_j \alpha_j \mathbf{1}_{Q_j} \in \text{BMO}(\sigma)$  such that  $f = g + b$ . We construct an extension G of g by Theorem  $1.\overline{3}$  and so it remains to prove the existence of the extension of b. By [\[HT21,](#page-70-4) Proposition 1.3], there exists  $B_0: \Omega \to \mathbb{R}$  such that  $\sup_{\xi \in \partial \Omega} \mathscr{C}_n(\nabla B_0)(\xi) + \sup_{x \in \Omega} \delta_{\Omega}(x) |\nabla B_0(x)| \lesssim$   $||f||_{\text{BMO}}$  and  $B_0 \rightarrow b$  non-tangentially for  $\sigma$ -a.e.  $\xi \in \partial \Omega$ . By Lemma [6.6,](#page-44-4) if we set  $B = B_0$  (as defined in [\(6.13\)](#page-44-0)), we get the desired extension of b. The extension of f is given by  $G + B$ . given by  $G + B$ .

### <span id="page-46-0"></span>7. VAROPOULOS-TYPE EXTENSIONS OF COMPACTLY SUPPORTED LIPSCHITZ FUNCTIONS

<span id="page-46-2"></span>We begin by constructing an extension of  $L^p$ -boundary functions in the next theorem.

**Theorem 7.1.** Let  $\Omega \in AR(s)$  for  $s \in (0, n]$ . If  $f \in Lip_c(\partial \Omega)$ , there exists a function  $F: \overline{\Omega} \to \mathbb{R}$  *such that for every*  $p \in (1,\infty]$ ,

- (i)  $F \in C^{\infty}(\Omega) \cap \text{Lip}(\overline{\Omega})$ , (ii)  $F|_{\partial\Omega} = f$  *continuously*,
- (iii)  $\|\mathcal{N}(F)\|_{L^p(\sigma)} + \|\mathcal{N}(\delta_{\Omega}\nabla F)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)},$
- (iv)  $\|\mathcal{C}_s(\nabla F)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)},$
- (v)  $\|[\mathcal{C}_s(\delta_\Omega|\nabla F|^2)]^{1/2}\|_{L^q(\sigma)} \lesssim \|f\|_{L^q(\sigma)}, \quad \text{for} \ \ q \in [2,\infty).$

*When*  $p = \infty$  *the norms on left hand-side of* (iii) *and* (iv) *are the* sup-norms instead of  $L^{\infty}$ .

*Proof.* Let  $\{\varphi_P\}_{P \in \mathcal{W}(\Omega)}$  be a partition of unity of  $\Omega$  so that each  $\varphi_P$  is supported in 1.1P and  $\|\nabla \varphi_P\|_{\infty} \lesssim 1/\ell(P)$ . For each  $\delta \in (0, \text{diam}(\Omega))$ , set

$$
\mathcal{W}_{\delta}(\Omega) = \{ P \in \mathcal{W}(\Omega) : \ell(P) \ge \delta \}
$$

and

$$
\varphi_{\delta} = \sum_{P \in \mathcal{W}_{\delta}(\Omega)} \varphi_P.
$$

From the properties of the Whitney cubes, there exists  $C > 0$  (depending on the parameters of the construction of the Whitney cubes) such that

$$
\varphi_{\delta}(x) = 0
$$
, if  $dist(x, \partial \Omega) \le \delta/C$ 

and

$$
\varphi_{\delta}(x) = 1
$$
, if  $dist(x, \partial \Omega) \ge C\delta$ .

Consequently, for a suitable constant  $C'$  depending on  $C$ , we infer that

(7.1) 
$$
\mathrm{supp}\,\nabla\varphi_{\delta}\subset\{x\in\Omega:\delta/C\le\mathrm{dist}(x,\partial\Omega)\le C\delta\}=:S_{\delta}\subset\bigcup_{P\in\mathcal{I}_{\delta}}P,
$$

where

<span id="page-46-3"></span>
$$
\mathcal{I}_\delta := \{P \in \mathcal{W}(\Omega): \frac{1}{2^{N_0+1}2^{N_1}} \le \ell(P) \le \frac{2^{N_1}}{2^{N_0}}\}
$$

with  $N_0 \in \mathbb{N}$  such that  $\frac{1}{2^{N_0+1}} \le \delta \le \frac{1}{2^{N_0}}$  and  $N_1 \in \mathbb{N}$  satisfies  $2^{N_1} \le C \le 2^{N_1+1}$ . We define

(7.2) 
$$
F(x) := v_f(x)(1 - \varphi_\delta(x)) + u(x)\varphi_\delta(x),
$$

where u is the approximation function of  $v_f$  as constructed in Theorem [5.5.](#page-36-1) It holds that

<span id="page-46-1"></span>(7.3) 
$$
\mathcal{C}_s(\nabla F) \leq \mathcal{C}_s(\nabla u) + \mathcal{C}_s(|\nabla \varphi_\delta| (u - v_f)) + \mathcal{C}_s(|\nabla v_f| (1 - \varphi_\delta))
$$

and

<span id="page-47-2"></span>
$$
(7.4) \ \mathcal{C}_s(\delta_{\Omega}|\nabla F|^2) \lesssim \mathcal{C}_s(\delta_{\Omega}|\nabla v_f|^2 (1-\varphi_\delta)^2) + \mathcal{C}_s(\delta_{\Omega}|u-v_f|^2 |\nabla \varphi|^2) + \mathcal{C}_s(\delta_{\Omega}|\nabla u|^2).
$$

For fixed  $\xi \in \partial \Omega$  and  $r > 0$ , we have

$$
\int_{B(\xi,r)\cap\Omega} |\nabla \varphi_{\delta}| \, |u - v_{f}| \, \frac{dx}{\delta_{\Omega}(x)^{n-s}} \lesssim \sum_{\substack{P \in \mathcal{I}_{\delta} \\ P \cap B(\xi,r) \neq \emptyset}} \int_{P} |u - v_{f}| \, \frac{dx}{\delta_{\Omega}(x)^{n+1-s}}
$$
\n
$$
\lesssim \sum_{\substack{P \in \mathcal{I}_{\delta} \\ P \cap B(\xi,r) \neq \emptyset}} \ell(P)^{s} \inf_{\zeta \in b(P)} \mathcal{N}_{\alpha}(u - v_{f})(\zeta) \lesssim \sum_{\substack{P \in \mathcal{I}_{\delta} \\ P \cap B(\xi,r) \neq \emptyset}} \int_{b(P)} \mathcal{N}(u - v_{f}) \, d\sigma
$$
\n
$$
\lesssim \sum_{k=-(N_{1}+1)}^{N_{1}} \sum_{\substack{\ell(P)=2^{k}/2^{N_{0}} \\ P \subset B(\xi,Mr)}} \int_{b(P)} \mathcal{N}_{\alpha}(u - v_{f}) \, d\sigma \lesssim_{C} \int_{B(\xi,Mr)} \mathcal{N}_{\alpha}(u - v_{f}) \, d\sigma,
$$

for suitably chosen constants  $\alpha > 1$  and  $M > 1$  large enough. Thus, when  $p \in (1, \infty)$ , we will get that

<span id="page-47-0"></span>(7.5) 
$$
\mathcal{C}_s(|\nabla \varphi_\delta|(u-v_f))(\xi) \lesssim \mathcal{M}(\mathcal{N}_\alpha(u-v_f))(\xi).
$$

By similar arguments we can also infer that

<span id="page-47-3"></span>(7.6) 
$$
\mathcal{C}_s(\delta_{\Omega}(u-v_f)^2|\nabla\varphi_{\delta}|^2)(\xi) \lesssim \mathcal{M}(\mathcal{N}_{\alpha}(u-v_f)^2)(\xi).
$$

When  $p = \infty$ , by [\(5.7\)](#page-32-8), we get  $\sup_{x \in \Omega} \sup_{y \in B_x} |u(y) - v_f(y)| \leq 2\varepsilon ||f||_{L^{\infty}(\sigma)}$ , which, arguing as above, implies

<span id="page-47-4"></span>
$$
(7.7) \quad \int_{B(\xi,r)\cap\Omega} |\nabla \varphi_{\delta}| \, |u - v_{f}| \frac{dx}{\delta_{\Omega}(x)^{n-s}} \lesssim \|f\|_{L^{\infty}(\sigma)} \sum_{\substack{P \in \mathcal{I}_{\delta} \\ P \cap B(\xi,r) \neq \emptyset}} \ell(P)^{s}
$$

$$
\lesssim \|f\|_{L^{\infty}(\sigma)} \sum_{\substack{k=-(N_{1}+1) \\ b(P) \subset B(\xi,Mr)}}^{\infty} \sigma(b(P)) \lesssim r^{s} \|f\|_{L^{\infty}(\sigma)}
$$

and thus

<span id="page-47-1"></span>(7.8) 
$$
\sup_{\xi \in \partial \Omega} C_s (|\nabla \varphi_\delta| (u - v_f))(\xi) \lesssim ||f||_{L^\infty(\sigma)}.
$$

For the last term on the right hand side of [\(7.3\)](#page-46-1), when  $p \in (1,\infty)$ , since  $f \in \text{Lip}_{c}(\partial \Omega)$ , we have that  $f \in \dot{M}^{1,p}(\sigma)$ . So, for fixed  $\xi \in \partial\Omega$  and  $r > 0$ , if  $\nabla_H f$  is the least upper gradient of  $f$ , in view of  $(3.5)$ , we estimate

$$
\int_{B(\xi,r)\cap\Omega}|\nabla v_f(x)||1-\varphi_\delta(x)|\frac{dx}{\delta_\Omega(x)^{n-s}}\lesssim \sum_{\substack{P\in \mathcal{W}(\Omega)\\P\cap B(\xi,r)\neq\emptyset\\ \ell(P)\lesssim \delta}}\int_P|\nabla v_f(x)|\frac{\ell(P)^s}{\ell(P)^n}\,dx
$$

$$
\lesssim \sum_{\substack{P \in \mathcal{W}(\Omega) \\ P \cap B(\xi,r) \neq \emptyset \\ \ell(P) \leq C\delta}} m_{\sigma,b(P)} (\nabla_H f) \ell(P)^{s+1} \lesssim \sum_{\substack{P \in \mathcal{W}(\Omega) \\ P \cap B(\xi,r) \neq \emptyset \\ \ell(P) \leq C\delta}} \ell(P) \sigma(b(P)) \inf_{\substack{\zeta \in b(P) \\ \zeta \in b(P)}} \mathcal{M}(\nabla_H f) (\zeta)
$$
  

$$
\leq \sum_{k \geq N_0 - N_1} 2^{-k} \sum_{\substack{Q \in \mathcal{D}_{\sigma} \\ Q \subset B(\xi, Mr)}} \int_Q \mathcal{M}(\nabla_H f) d\sigma \lesssim \delta m_{\sigma,B(\xi, Mr)} (\mathcal{M}(\nabla_H f)) r^s,
$$

which shows that

(7.9) 
$$
\mathcal{C}_s\big(|\nabla v_f|(1-\varphi_\delta)\big)(\xi)\lesssim \delta \mathcal{M}(\mathcal{M}(\nabla_H f))(\xi).
$$

In view of  $(3.3)$  and  $(3.5)$ ,

<span id="page-48-2"></span><span id="page-48-1"></span><span id="page-48-0"></span>
$$
\sup_{x \in P} |\nabla v_f(x)|^2 \lesssim \ell(P)^{-1} m_{\sigma, B_P}(|f|) m_{\sigma, B_P}(\nabla_H f).
$$

Arguing as above and using the latter estimate, we get

(7.10) 
$$
\mathcal{C}_s(\delta_{\Omega}|\nabla v_f|^2(1-\varphi_{\delta})^2)(\xi) \lesssim \delta \mathcal{M}(\mathcal{M}(\nabla_H f)\mathcal{M}f)(\xi).
$$

For  $p = \infty$ , we use Lemma [3.5](#page-24-0) to get

(7.11) 
$$
\mathcal{C}_s(|\nabla v_f|(1-\varphi_\delta))(\xi) \lesssim \mathrm{Lip}(f)\delta.
$$

Indeed, for  $\xi \in \partial \Omega$  and  $r > 0$ ,

$$
\int_{B(\xi,r)\cap\Omega} |\nabla v_f||1 - \varphi_{\delta}| \frac{dx}{\delta_{\Omega}(x)^{n-s}} \lesssim \mathrm{Lip}(f) \sum_{\substack{P \in \mathcal{W}(\Omega) \\ \ell(P) \leq C\delta}} \int_P \frac{\ell(P)^s}{\ell(P)^n} dx
$$
  

$$
\lesssim \mathrm{Lip}(f) \sum_{\substack{P \in \mathcal{W}(\Omega) \\ P \cap B(\xi,r) \neq \emptyset \\ \ell(P) \leq C\delta}} \ell(P)^{s+1} \leq \mathrm{Lip}(f) \sum_{\substack{k \geq N_0 - N_1 \\ \ell(P) \leq C\delta}} 2^{-k} \sum_{\substack{P \in \mathcal{W}(\Omega) \\ \ell(P) = 2^{-k} \\ \ell(P) = 2^{-k}}} \sigma(b(P))
$$
  

$$
\lesssim \mathrm{Lip}(f) \sum_{\substack{k \geq N_0 - N_1 \\ \ell \geq N_0 - N_1}} 2^{-k} \sum_{\substack{Q \in \mathcal{D}_{\sigma} \\ Q \subset B(\xi, Mr) \\ \ell(Q) = 2^{-k}}} \sigma(Q) \lesssim \mathrm{Lip}(f) \delta r^s,
$$

for  $M > 1$  large enough constant depending on Ahlfors regularity and Whitney constants. Combining [\(5.5\)](#page-32-2), [\(7.3\)](#page-46-1), [\(7.5\)](#page-47-0), [\(7.8\)](#page-47-1), [\(7.9\)](#page-48-0), and [\(7.11\)](#page-48-1), and choosing

$$
\delta = \begin{cases} ||f||_{L^{\infty}(\sigma)} / \operatorname{Lip}(f) & , \text{if } p = \infty \\ ||f||_{L^p(\sigma)} / ||f||_{\dot{M}^{1,p}(\sigma)} & , \text{if } p \in (1, \infty), \end{cases}
$$

it follows that

$$
\|\mathcal{C}_s(\nabla F)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)}, \quad \text{for } p \in (1,\infty].
$$

Moreover, combining [\(5.6\)](#page-32-4), [\(7.4\)](#page-47-2), [\(7.6\)](#page-47-3), [\(7.10\)](#page-48-2) and choosing  $\delta = ||f||_{L^q(\sigma)}/||f||_{\dot{M}^{1,q}(\sigma)}$  it follows

$$
\|\mathcal{C}_s(\delta_{\Omega}|\nabla F|^2)^{1/2}\|_{L^q(\sigma)} \lesssim \|f\|_{L^q(\sigma)} \quad \text{for } q \in (2,\infty).
$$

The last estimate can be proved for  $q = 2$  by direct estimates using [\(5.27\)](#page-36-9). We leave the proof as an exercise.

For the non-tangential estimate note that, since  $v_f = \varphi_\delta v_f + (1 - \varphi_\delta)v_f$ , we can write

$$
v_f - F = \varphi_\delta(v_f - u).
$$

So, for every  $\xi \in \partial \Omega$  we have

$$
\mathcal{N}(v_f - F)(\xi) = \sup_{x \in \gamma(\xi)} |v_f(x) - F(x)| \le \sup_{x \in \gamma(\xi)} |v_f(x) - u(x)| = \mathcal{N}(v_f - u)(\xi)
$$

and thus, by  $(5.23)$ , we get

$$
\|\mathcal{N}(v_f - F)\|_{L^p(\sigma)} \lesssim \varepsilon \|f\|_{L^p(\sigma)}, \quad \text{for } p \in (1, \infty).
$$

Using this and [\(3.6\)](#page-22-3), for  $p \in (1, \infty)$ , we get that

$$
\|\mathcal{N}(F)\|_{L^p(\sigma)} \lesssim \|\mathcal{N}(v_f - F)\|_{L^p(\sigma)} + \|\mathcal{N}(v_f)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)}.
$$

Moreover, combining [\(3.3\)](#page-22-0), [\(5.3\)](#page-32-0), [\(5.17\)](#page-35-0), the fact that  $|\nabla \varphi_{\delta}(x)| \lesssim \delta_{\Omega}(x)^{-1}$ , and using the  $L^p$ -boundedness of the Hardy-Littlewood maximal function, we can easily infer that

$$
\|\mathcal{N}(\delta_{\Omega}\nabla F)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)}, \quad \text{for } p \in (1, \infty).
$$

The estimates for  $p = \infty$  can be proved similarly and the routine details are omitted.

Note that the extension F is Lipschitz in  $\overline{\Omega}$ . Indeed, if  $E := \text{supp } f$ , in light of Lemmas [3.2](#page-22-4) and [3.5,](#page-24-0) and Theorem [5.3,](#page-32-9) we infer that for every  $x \in \Omega$ ,

$$
|\nabla v_f(x)(1-\varphi_{\delta}(x))| \lesssim \text{Lip}(f),
$$
  
\n
$$
|\nabla u(x)\varphi_{\delta}(x)| \lesssim \delta_{\Omega}(x)^{-1} ||f||_{L^{\infty}(\sigma)} |\varphi_{\delta}(x)| \lesssim \frac{1}{\delta} \text{Lip}(f) \text{ diam } E,
$$
  
\n
$$
|(u(x)-v_f(x))\nabla\varphi_{\delta}(x)| \lesssim \frac{1}{\delta} \inf_{\zeta \in B(\xi_x, 2\delta_{\Omega}(x)) \cap \partial\Omega} \mathcal{N}(u-v_f)(\zeta) \lesssim_{\varepsilon} \frac{1}{\delta} ||\mathcal{M}(f)||_{L^{\infty}} \lesssim \frac{||f||_{L^{\infty}(\sigma)}}{\delta} \lesssim \frac{\text{Lip}(f)}{\delta} \text{ diam } E.
$$

These estimates imply that  $\|\nabla F\|_{L^{\infty}(\Omega)} \lesssim_{\delta,\text{diam } E} \text{Lip}(f)$ . Moreover, since  $v_f \in \text{Lip}(\overline{\Omega})$ ,  $F = v_f$  around  $\partial\Omega$ , and  $v_f|_{\partial\Omega} = f$ , we deduce that  $F \in \text{Lip}(\overline{\Omega})$  with  $F|_{\partial\Omega} = f$  and  $\text{Lip}(F) \lesssim_{\delta, \text{diam } E} \text{Lip}(f)$ , concluding the proof of the theorem.  $\Box$ 

**Remark 7.2.** Note that the convergence to the boundary is inherited from the one of  $v_f$  and there is no need for an iteration argument as in the proof of Theorem [1.3.](#page-4-0)

We now turn our attention to the construction of an extension of BMO-boundary functions.

<span id="page-49-0"></span>**Theorem 7.3.** *Let*  $\Omega \in AR(s)$  *for*  $s \in (0, n]$ *. If*  $f \in Lip<sub>c</sub>(\partial \Omega)$ *, then there exists an extension*  $F: \overline{\Omega} \to \mathbb{R}$  *such that* 

- $(i)$   $F \in C^{\infty}(\Omega) \cap \text{Lip}(\overline{\Omega}),$
- *(ii)*  $F|_{\partial\Omega} = f$  *continuously,*
- *(iii)* sup  $\sup_{\xi \in \partial \Omega} \mathcal{N}_{\sharp}(F)(\xi) + \sup_{x \in \Omega} \delta_{\Omega}(x) |\nabla F(x)| \lesssim ||f||_{\text{BMO}(\sigma)},$
- $(iv) \ \sup_{s \in \mathcal{C}_{s}} \mathcal{C}_{s}(\nabla F)(\xi) \lesssim ||f||_{\text{BMO}(\sigma)},$ ξ∈∂Ω
- (*v*)  $\sup [\mathcal{C}_s(\delta_\Omega |\nabla F|^2)(\xi)]^{1/2} \lesssim ||f||_{\text{BMO}(\sigma)}.$ ξ∈∂Ω

*Proof.* Let 
$$
w
$$
 be the approximation of  $v_f$  given by Theorem 5.5 and define

<span id="page-50-4"></span>(7.12) 
$$
F(x) := v_f(x)(1 - \varphi_\delta(x)) + w(x)\varphi_\delta(x).
$$

Then, for any  $\xi \in \partial \Omega$ ,

<span id="page-50-0"></span>(7.13) 
$$
\mathcal{C}_s(\nabla F)(\xi) \leq \mathcal{C}_s(\nabla v_f(1-\varphi_\delta))(\xi) + \mathcal{C}_s(\nabla w)(\xi) + \mathcal{C}_s((w-v_f)\nabla\varphi_\delta)(\xi)
$$

and

$$
(7.14) \mathcal{C}_s(\delta_{\Omega}|\nabla F|^2) \lesssim \mathcal{C}_s(\delta_{\Omega}|\nabla v_f|^2(1-\varphi_\delta)^2) + \mathcal{C}_s(\delta_{\Omega}|w-v_f|^2|\nabla \varphi|^2) + \mathcal{C}_s(\delta_{\Omega}|\nabla w|^2).
$$

For the second term on the right hand side of  $(7.13)$  we just use  $(5.21)$ , while for the first one, by  $(7.11)$ , we have that

<span id="page-50-1"></span>
$$
(7.15) \t\t\t\t\mathcal{C}_s(\nabla v_f(1-\varphi_\delta))(\xi) \lesssim \delta \operatorname{Lip}(f).
$$

The third term can be bounded as in  $(7.7)$  and get

<span id="page-50-2"></span>(7.16) 
$$
\sup_{\xi \in \partial \Omega} C_s (\nabla \varphi_\delta(w - v_f))(\xi) \lesssim \|f\|_{\text{BMO}}.
$$

Combining [\(7.13\)](#page-50-0), [\(7.15\)](#page-50-1), and [\(7.16\)](#page-50-2), and choosing  $\delta = ||f||_{\text{BMO}} / \text{Lip}(f)$ , we obtain (iv). Moreover, (v) follows from (iv) and the second estimate of (iii).

For the sharp non-tangential maximal function estimate, note that since  $F - v_f = \varphi_\delta(w - \xi)$  $v_f$ ), using [\(3.7\)](#page-23-0) and [\(5.7\)](#page-32-8), we get that for every  $\xi \in \partial \Omega$  it holds that

$$
\mathcal{N}_{\sharp}(F)(\xi) \leq 2\mathcal{N}(F - \nu_f)(\xi) + \mathcal{N}_{\sharp}(\nu_f)(\xi) \leq 2\mathcal{N}(w - \nu_f)(\xi) + \mathcal{N}_{\sharp}(\nu_f)(\xi) \lesssim ||f||_{\text{BMO}(\sigma)}.
$$

It remains to prove that  $F \in \text{Lip}(\Omega)$ . We first show that

$$
||\nabla F||_{L^{\infty}(\sigma)} \lesssim \text{Lip}(f).
$$

To this end, for every  $x \in \Omega$ , by Lemma [3.5,](#page-24-0) we have that

<span id="page-50-3"></span>
$$
|\nabla v_f(x)(1-\varphi_\delta(x))| \lesssim \text{Lip}(f).
$$

By [\(5.7\)](#page-32-8) and the fact that  $\delta_{\Omega}(x) \approx \delta$  in the support of  $\nabla \varphi_{\delta}$ , we obtain

$$
|(w(x) - v_f(x))\nabla\varphi_\delta(x)| \lesssim_{\varepsilon} \delta_{\Omega}(x)^{-1} \|f\|_{\text{BMO}(\sigma)} \approx \delta^{-1} \|f\|_{\text{BMO}(\sigma)} = \text{Lip}(f).
$$

while, by  $(5.8)$  it holds that

$$
|\nabla w(x)\varphi_{\delta}(x)| \lesssim ||f||_{\text{BMO}(\sigma)} \delta_{\Omega}(x)^{-1} |\varphi_{\delta}(x)| \lesssim \delta^{-1} ||f||_{\text{BMO}(\sigma)} = \text{Lip}(f),
$$

which implies [\(7.17\)](#page-50-3). By construction F is continuous around the boundary and  $F|_{\partial\Omega} = f$ continuously, which implies that  $F \in \text{Lip}(\overline{\Omega}) \cap C^{\infty}(\overline{\Omega})$  with  $\text{Lip}(F) \lesssim \text{Lip}(f)$ . Moreover, combining the last two estimates above and  $(3.4)$ , we get that

$$
\sup_{x \in \Omega} \delta_{\Omega}(x) |\nabla F(x)| \lesssim ||f||_{\text{BMO}(\sigma)},
$$

which concludes the proof of the theorem.  $\Box$ 

Our last goal is to modify that the extensions constructed in Theorems [7.1](#page-46-2) and [7.3](#page-49-0) so that they are also in  $\dot{W}^{1,2}(\Omega;\omega_s)$ . This will conclude the proof of Theorem [1.4.](#page-5-2)

**Theorem 7.4.** *Let*  $\Omega \in AR(s)$  *for*  $s \in (0, n]$ *. If*  $f \in Lip<sub>c</sub>(\partial \Omega)$ *, then there exists an*  $extension\ F_0\in \dot W^{1,2}(\Omega;\omega_s)$  (resp.  $\bar F_0\in \dot W^{1,2}(\Omega;\omega_s))$  that satisfies the conclusions (i)-(iv) *of Theorem [7.1](#page-46-2) (resp. Theorem [7.3\)](#page-49-0).*

*Proof.* Let  $f \in \text{Lip}_c(\partial \Omega)$ ,  $E := \text{supp } f$ , and  $r_0 := \text{diam } E$ . Without loss of generality we may assume that  $0 \in E$  and so  $E \subset B(0, r_0)$ . Now let  $B = B(0, Mr_0)$  for some  $M > 1$ large enough depending on the Whitney constants so that for every  $P \in \mathcal{W}(\Omega)$  satisfying  $\ell(P) \leq M^{-1}r_0$  and  $1.2P \cap (2B \setminus B) \neq \emptyset$  it holds that  $b(P) \cap E = \emptyset$ . We denote the collection of all such Whitney cubes by  $\mathscr{P}_s(E)$  ("s" stands for "small"). We also denote by  $\mathscr{P}_l(E)$  the collection of  $P \in \mathcal{W}(\Omega)$  satisfying  $\ell(P) > M^{-1}r_0$  and  $1.2P \cap (2B \setminus B) \neq \emptyset$ ("l" stands for "large"). It is easy to see that

(7.18) 
$$
\sum_{\substack{Q \subset R: Q = b(P) \\ P \in \mathcal{P}_l(E)}} \sigma(Q) \lesssim r_0^s \lesssim \sigma(R).
$$

Note that if  $x \in (2B \setminus B) \cap \Omega$  and there exists  $P \in \mathscr{P}_s(E)$  such that  $x \in 1.1P$ , then the extension F of Theorem [7.1](#page-46-2) satisfies  $F(x) = 0$  (resp.  $\overline{F}$  of Theorem [7.3](#page-49-0) satisfies  $\bar{F}(x) = 0$ ). We define now the cut-off function  $\psi_{r_0} \in C_c^{\infty}(\mathbb{R}^{n+1})$  such that  $0 \le \psi_{r_0} \le 1$ ,  $\psi_{r_0} = 1$  in  $\overline{B}$ ,  $\psi_{r_0} = 0$  in  $\mathbb{R}^{n+1} \setminus 2B$ , and  $|\nabla \psi_{r_0}| \lesssim 1/r_0$ . Then we define

<span id="page-51-1"></span>
$$
F_0(x) := F(x) \psi_{r_0}(x)
$$
 and  $\bar{F}_0(x) := \bar{F}(x) \psi_{r_0}(x)$ 

It is clear that  $F_0|_{\partial\Omega} = f$  (resp.  $\bar{F}_0|_{\partial\Omega} = f$ ) and remark that

(7.19) 
$$
\begin{aligned} \sup_{\text{supp}(\bar{F}\nabla\psi_{r_0})} \Big\} &\subset T_{r_0} := \{x \in \Omega \cap (2B \setminus B) : \text{dist}(x, \partial\Omega) \ge c_0 r_0\}, \end{aligned}
$$

for some  $c_0 \in (0, 1)$  small enough depending on M and the Whitney constants. Therefore, for any  $\xi \in \partial\Omega$ , if  $B(\xi, r) \cap \text{supp}(F\nabla \psi_{r_0}) \neq \emptyset$ , then  $r \geq c_1 \max\{r_0, \text{dist}(\xi, 2B \setminus B)\}$  for some constant  $c_1 \in (0, 1)$  depending on  $c_0$ . Moreover,

<span id="page-51-0"></span>
$$
(7.20) \quad |F(x)| |\nabla \psi_{r_0}(x)| \lesssim \delta_{\Omega}(x)^{-1} |F(x)| \quad \text{and} \quad |\bar{F}(x)| |\nabla \psi_{r_0}(x)| \lesssim \delta_{\Omega}(x)^{-1} |\bar{F}(x)|.
$$

We will only prove the theorem for  $F_0$  and unbounded domains with unbounded boundary since for domains with compact boundary the arguments are similar.

We first prove that  $F_0$  satisfies the conclusions of Theorem [7.1.](#page-46-2) It is easy to see that  $||\mathcal{N}(F_0)||_{L^p(\sigma)} \lesssim ||f||_{L^p(\sigma)}$  since  $|F_0| \leq |F|$  and the same estimate holds for F. We have that  $\nabla F_0 = \nabla F \psi_{r_0} + F \nabla \psi_{r_0}$  and it is easy to see that  $\|\mathcal{N}(\delta_\Omega \nabla F_0)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)}$ by  $(7.20)$  and the estimates in (ii) and (iii) for F in Theorem [7.1.](#page-46-2) To prove the estimate  $||C_s(\nabla F_0)||_{L^p(\sigma)} \lesssim ||f||_{L^p(\sigma)}$ , it is enough to show that  $||C_s(F \nabla \psi_{r_0})||_{L^p(\sigma)} \lesssim ||f||_{L^p(\sigma)}$ . Thus, for any such  $r$ , we have that

$$
\int_{B(\xi,r)\cap\Omega} \sup_{y\in B_x} |F(y)| |\nabla \psi_{r_0}(y)| \, \omega_s(x) \, dx \lesssim r_0^{-1} \int_{B(\xi,r)\cap(2B\setminus B)\cap\Omega} \sup_{y\in B_x} |F(y)| \, \omega_s(x) \, dx.
$$

By [\(5.1\)](#page-31-0), [\(4.2\)](#page-27-1), [\(7.2\)](#page-46-3), and the choice of the constant M, for any  $x \in B(\xi, r) \cap (2B \setminus B) \cap \Omega$ and for any  $y \in B_x$ , it holds that

$$
\begin{aligned} |F(y)| \leq & \sum_{R \in \mathsf{Top}} \sum_{P \in \mathscr{P}_l(E) \backslash \mathcal{P}_0} \left| m_{\sigma,R} f - m_{\sigma,b(P)} f \right| \varphi_P(y) + \sum_{P \in \mathscr{P}_l(E)} \left| m_{\sigma,b(P)} f \right| \varphi_P(y) \\ \leq & \varepsilon \sum_{R \in \mathsf{Top}} \sum_{\substack{P \in \mathscr{P}_l(E) \backslash \mathcal{P}_0 \\ b(P) \in \mathsf{Tree}(R)}} M f(b(P)) \, \varphi_P(y) + \sum_{P \in \mathscr{P}_l(E)} \left| m_{\sigma,b(P)} f \right| \varphi_P(y), \end{aligned}
$$

which, in turn, implies that

$$
r_0^{-1} \int_{B(\xi,r) \cap (2B\backslash B) \cap \Omega} \sup_{y \in B_x} |F(y)| \, \omega_s(x) \, dx \lesssim r_0^{-1} \sum_{P \in \mathscr{P}_l(E)} \ell(P)\sigma(b(P))Mf(b(P))
$$
  

$$
\lesssim \int_{CB} \mathcal{M}(f) \, d\sigma \le \int_{B(\xi,C'r)} \mathcal{M}f \, d\sigma,
$$

for some constant  $C' > 1$  depending on C and M. This readily yields that for every  $\xi \in \partial \Omega$ ,

$$
\mathcal{C}_s(F|\nabla\psi_{r_0}|)(\xi)\lesssim \mathcal{M}(\mathcal{M}f)(\xi),
$$

and the desired estimate follows for any  $p \in (1,\infty]$ . Similarly, we can show that

$$
\mathcal{C}_s(\delta_\Omega |F|^2 |\nabla \psi_{r_0}|^2)(\xi) \lesssim \mathcal{M}((\mathcal{M}f)^2)(\xi)
$$

which implies (v) of Theorem [7.1](#page-46-2) for  $q > 2$ . The case  $q = 2$  once again can be treated separately using [\(5.27\)](#page-36-9). We omit the details.

It remains to show that  $F_0 \in \dot{W}^{1,2}(\Omega;\omega_s)$  since it is clear that  $F_0 \in C^{\infty}(\Omega) \cap \text{Lip}(\overline{\Omega})$ such that  $F_0|_{\partial\Omega} = f$ . To this end, by the definition of F and the proof of its Lipschitz property, we get that

$$
\int_{\Omega} |\nabla F_0|^2 \omega_s(x) dx \le \int_{2B \cap \Omega} |\nabla F|^2 \omega_s(x) dx
$$
  

$$
\lesssim \left( \left( 1 + \frac{r_0^2}{\delta^2} \right) \min(r_0, \delta) + \frac{r_0^3}{\delta^2} \right) r_0^s (\text{Lip } f)^2.
$$

Moreover, using [\(7.19\)](#page-51-1) and the fact that supp  $f = E$ , we can show that

$$
\int_{\Omega} |F \nabla \psi_{r_0}|^2 \omega_s(x) dx \lesssim r_0^{-2-n+s} \int_{T_{r_0}} |F|^2 dx \lesssim r_0^{s-1} \|f\|_{L^{\infty}(\partial \Omega)}^2 \le r_0^{s+1} (\text{Lip } f)^2,
$$

concluding the proof of the Theorem for  $L^p$ ,  $p \in (1,\infty]$ .

To demonstrate the theorem for  $\bar{F}_0$ , it is enough to prove the Carleson estimate as the estimates for the non-tangential maximal functions are easy to prove and we leave them as an exercise. By  $(7.19)$ ,  $(5.1)$ ,  $(4.2)$ ,  $(7.12)$ , and the choice of the constant M, for any  $x \in B(\xi, r) \cap (2B \setminus B) \cap \Omega$  and every  $y \in B_x$ , it holds that

$$
\begin{aligned} |\bar F(y)|\leq & \sum_{R\in \text{Top}}\sum_{\substack{P\in \mathscr{P}_l(E)\backslash \mathcal{P}_0\\ b(P)\in \text{Tree}(R)}} \left|m_{\sigma,R}f-m_{\sigma,b(P)}f\right|\varphi_P(y)+\sum_{P\in \mathscr{P}_l(E)} \left|m_{\sigma,b(P)}f\right|\varphi_P(y)\\ \leq & \varepsilon \|f\|_{\text{BMO}(\sigma)}+\sum_{P\in \mathscr{P}_l(E)} \left|m_{\sigma,b(P)}f\right|\varphi_P(y). \end{aligned}
$$

It is not hard to see that for every  $P \in \mathcal{P}_l(E)$ , there exists  $P^* \in \mathcal{W}(\Omega)$  such that  $\ell(P^*) \approx$  $dist(P^*, P) \approx r_0$  and  $b(P^*) \subset \partial\Omega \setminus E$ , and it holds that  $m_{\sigma, b(P^*)}f = 0$ . Thus, for any  $x \in B(\xi, r) \cap (2B \setminus B) \cap \Omega$ ,

$$
|\bar{F}(y)| \leq \varepsilon ||f||_{\text{BMO}(\sigma)} + \sum_{P \in \mathscr{P}_l(E)} |m_{\sigma,b(P)}f - m_{\sigma,b(P^*)}f| \varphi_P(x) \lesssim ||f||_{\text{BMO}(\sigma)},
$$

which, arguing as above, implies that  $\sup_{\xi \in \partial \Omega} C_s(F \nabla \psi_{r_0})(\xi) \lesssim ||f||_{\text{BMO}(\sigma)}$ . This finishes the proof of the theorem since the same proof as above shows that  $\bar{F}_0 \in \dot{W}^{1,2}(\Omega;\omega_s)$ .  $\Box$ 

## <span id="page-53-0"></span>8. APPLICATIONS TO BOUNDARY VALUE PROBLEMS FOR SYSTEMS OF ELLIPTIC **EQUATIONS**

We define the *variational co-normal derivative* of a solution  $v \in \dot{W}^{1,2}(\Omega; \mathbb{C}^m)$  of  $Lv =$  $-$  div  $\Xi$  + H in  $\Omega$ , and denote it by  $\partial_{\nu_A} v$ , to be the linear functional defined in terms of the sesquilinear form associated to  $L$  as follows:

$$
\langle \partial_{\nu_A} v, \varphi \rangle := \ell_v(\varphi) := \mathbf{B}(v, \Phi) = \sum_{\alpha, \beta=1}^m \sum_{i,j=1}^{n+1} \int_{\Omega} a_{ij}^{\alpha \beta}(x) \partial_j v^{\beta}(x) \partial_i \Phi^{\alpha} dx - \sum_{\alpha=1}^m \sum_{i=1}^{n+1} \int_{\Omega} \Xi_i^{\alpha}(x) \partial_i \Phi^{\alpha}(x) dx - \sum_{\alpha=1}^m \int_{\Omega} H^{\alpha}(x) \Phi(x)^{\alpha} dx,
$$

<span id="page-53-2"></span>where  $\varphi \in \text{Lip}_c(\partial \Omega; \mathbb{C}^m)$  and  $\Phi \in \dot{W}^{1,2}(\Omega; \mathbb{C}^m) \cap \text{Lip}(\overline{\Omega}; \mathbb{C}^m)$  such that  $\Phi|_{\partial \Omega} = \varphi$ .

## **Lemma 8.1.**  $\ell_v$  *is unambiguously defined.*

*Proof.* If  $\Phi^1, \Phi^2 \in \dot{W}^{1,2}(\Omega; \mathbb{C}^m) \cap \text{Lip}(\overline{\Omega}; \mathbb{C}^m)$  such that  $\Phi^1|_{\partial\Omega} = \Phi^2|_{\partial\Omega} = \varphi$  and  $\Phi^1 \neq \emptyset$  $\Phi^2$ , then  $\Psi := \Phi^1 - \Phi^2 \in \dot{W}^{1,2}(\Omega : \mathbb{C}^m) \cap \text{Lip}(\overline{\Omega}; \mathbb{C}^m)$  such that  $\Psi|_{\partial\Omega} = 0$ , which implies that  $\Psi \in Y_0^{1,2}$  $\mathcal{L}_0^{1,2}(\Omega)$ . Since  $Lv = -\operatorname{div} \mathbf{\Xi} + H$ , we have that  $\mathbf{B}(v, \Psi) = 0$  and thus,  $\mathbf{B}(v, \Phi^1) = \mathbf{B}(v, \Phi^2)$ . So any extension of  $\varphi$  belonging to  $\dot{W}^{1,2}(\Omega; \mathbb{C}^m) \cap \text{Lip}(\overline{\Omega}; \mathbb{C}^m)$ defines the same linear functional  $\ell_{\nu}$ .

From now on, we assume that  $\Omega \in \text{AR}(n)$ ,  $n \geq 2$ , and that either  $\Omega$  is bounded or  $\partial \Omega$ unbounded. This is because we will use the duality  $N_{q,p}(\Omega) = (C_{s,q',p'}(\Omega))^*$ .

In the sequel, for simplicity, we will prove our results specifically for real elliptic equations (i.e., when  $m = 1$ ). Nevertheless, the proofs for  $m > 1$  and complex-valued coefficients are identical (see also Remark [1.6\)](#page-7-1).

## <span id="page-53-1"></span>8.1. Some connections between Poisson Problems and Boundary Value Problems. We set

$$
X^q(\sigma) = \begin{cases} L^q(\sigma) & \text{if } q \in (1, \infty) \\ H^1(\sigma) & \text{if } q = 1, \end{cases}
$$

<span id="page-53-5"></span>where  $H^1(\sigma)$  is the atomic Hardy space with respect to  $\sigma$ .

**Proposition 8.2.** If  $(\text{PR}_p^L)$  is solvable in  $\Omega$  for some  $p > 1$ , then its solution u satisfies the *one-sided Rellich-type inequality*

<span id="page-53-3"></span>(8.1) 
$$
\|\partial_{\nu_A} u\|_{L^p(\sigma)} \lesssim \|H\|_{\mathcal{C}_{2*,p}(\Omega)} + \|\Xi/\delta_{\Omega}\|_{\mathcal{C}_{2,p}(\Omega)}.
$$

*Moreover, if*  $(\widetilde{\text{PR}}_q^L)$ ,  $q \in [1,2]$ , is solvable in  $\Omega$  with data  $H = 0$  and  $\Xi \in L_c^2(\Omega)$ , then its *solution satisfies the one-sided Rellich-type inequality*

<span id="page-53-4"></span>(8.2) 
$$
\|\partial_{\nu_A} u\|_{X^q(\sigma)} \lesssim \|\Xi\|_{T_2^p(\Omega)}, \quad q \in [1,2].
$$

*Proof.* Suppose that u is the solution of  $(\text{PR}_p^L)$ . Let  $\varphi \in \text{Lip}_c(\partial \Omega)$  and  $F \in \dot{W}^{1,2}(\Omega) \cap$ Lip( $\overline{\Omega}$ ) be the Varopoulos extension of the L<sup>p</sup>-boundary data  $\varphi$  constructed in Theorem [1.4.](#page-5-2) Then, by Lemma [8.1,](#page-53-2) we get

$$
|\ell_u(\varphi)| = |\mathbf{B}(u,F)| \le ||A||_{L^{\infty}(\Omega)} \int_{\Omega} |\nabla u||\nabla F| + \int_{\Omega} |H||F| + \int_{\Omega} |\Xi| |\nabla F|.
$$

By duality (see [\[MPT22,](#page-70-8) Proposition 2.4]),  $(2.45)$ , and the properties of the extension F, we infer that

$$
\int_{\Omega} |\nabla u| |\nabla F| \lesssim \|\widetilde{\mathcal{N}}_2(\nabla u)\|_{L^p(\sigma)} \|\mathcal{C}_2(\nabla F)\|_{L^{p'}(\sigma)} \lesssim (\|H\|_{\mathcal{C}_{2*,p}} + \|\Xi/\delta_{\Omega}\|_{\mathcal{C}_{2,p}(\Omega)}) \|\varphi\|_{L^{p'}}.
$$

By duality and using Theorem [1.4](#page-5-2) (ii) and (iii), we infer that

$$
\int_{\Omega} |H||F| + \int_{\Omega} |\Xi| |\nabla F|
$$
\n
$$
\lesssim ||H||_{\mathcal{C}_{2*,p}} ||\widetilde{\mathcal{N}}_2(F)||_{L^{p'}(\sigma)} + ||\Xi/\delta_{\Omega}||_{\mathcal{C}_{2,p}(\Omega)} ||\mathcal{N}_{2^*}(\delta_{\Omega} \nabla F)||_{L^{p'}(\sigma)}
$$
\n
$$
\lesssim (||H||_{\mathcal{C}_{2*,p}} + ||\Xi/\delta_{\Omega}||_{\mathcal{C}_{2,p}(\Omega)}) ||\varphi||_{L^{p'}(\sigma)}.
$$

Thus, by the above estimates, the density of  $\text{Lip}_c(\partial \Omega)$  in  $L^p(\sigma)$ , and duality, we infer [\(8.1\)](#page-53-3).

Let now v be the solution of  $(\widetilde{PR}_1^L)$  and let  $\varphi \in \text{Lip}_c(\partial \Omega)$ . If  $\widetilde{F}$  is the Varopoulos extension of  $\varphi$  constructed in the second part of Theorem [1.4,](#page-5-2) by Lemma [8.1,](#page-53-2) we get that

$$
|\ell_v(\varphi)| \leq ||A||_{L^{\infty}(\Omega)} \int_{\Omega} |\nabla v| |\nabla \widetilde{F}| + \int_{\Omega} |\Xi| |\nabla \widetilde{F}|.
$$

By duality,  $(2.45)$ , and Theorem [1.4](#page-5-2) (iii) (for BMO), we have that

$$
\int_{\Omega}|\nabla v||\nabla \widetilde{F}| \lesssim \|\widetilde{\mathcal{N}}_2(\nabla v)\|_{L^1(\sigma)}\sup_{\xi \in \partial \Omega}\mathcal{C}(\nabla \widetilde{F})(\xi) \lesssim \|\boldsymbol{\Xi}\|_{T_2^1(\Omega)}\|\varphi\|_{{\rm BMO}(\sigma)}.
$$

For the second term, by  $T_2^{\infty}(\Omega) = (T_2^1(\Omega))^*$  and Theorem [1.4](#page-5-2) (v) (for BMO), it holds that

$$
\int_{\Omega} |\Xi| |\nabla \widetilde{F}| \lesssim ||\Xi||_{T_2^1(\Omega)} ||\delta_{\Omega} \nabla \widetilde{F}||_{T_2^{\infty}(\Omega)} \lesssim ||\Xi||_{T_2^1(\Omega)} ||\varphi||_{\text{BMO}(\sigma)}.
$$

Thus

 $|\ell_v(\varphi)| \lesssim ||\boldsymbol{\Xi}||_{T_2^1(\Omega)} ||\varphi||_{\mathrm{BMO}(\sigma)},$ 

which, since  $\overline{\text{Lip}_c(\partial\Omega)}^{\text{VMO}(\sigma)} = \text{VMO}(\sigma) = (H^1(\sigma))^*$ , implies [\(8.2\)](#page-53-4). The proof of (8.2) for  $q \in (1, 2]$ , is similar and is omitted.

<span id="page-54-0"></span>**Theorem 8.3.** *If*  $(\text{PR}_{p}^{L})$  *with*  $\Xi = 0$  *is solvable in*  $\Omega$  *for some*  $p > 1$ *, then*  $(\text{D}_{p'}^{L^*})$  $_{p'}^{L^*}$ ) is also *solvable in*  $\Omega$ *, where*  $1/p + 1/p' = 1$ *.* 

*Proof.* Let  $f \in \text{Lip}_c(\partial \Omega)$  and let u be the solution to [\(2.38\)](#page-16-2) for  $L^*$  with data f. Due to the density of  $L_c^{\infty}(\Omega)$  in  $\mathrm{C}_{2_*,p}(\Omega)$  and duality, we have that

$$
\|\mathcal{N}_{2^*}(u)\|_{L^{p'}(\sigma)} \lesssim \sup_{\substack{H \in L^{\infty}_c(\Omega): \\ \|H\|_{\mathcal{C}_{2_*, p}(\Omega)} = 1}} \Big| \int_{\Omega} u H \Big|.
$$

Fix such an  $H \in L_c^{\infty}(\Omega)$  and let  $w \in Y_0^{1,2}$  $\chi_0^{-1,2}(\Omega)$  be the solution to  $(\text{PR}_p^L)$  with data  $\Xi = 0$ and H. Then, using the fact that  $L^*u = 0$  and [\(8.1\)](#page-53-3), we estimate

$$
\Big|\int_{\Omega} uH\Big| = \Big| - \int_{\Omega} \nabla u A \nabla w + \int_{\partial \Omega} \partial_{\nu_A} w f\Big| = \Big| \int_{\partial \Omega} \partial_{\nu_A} w f\Big| \lesssim \|C_{2*}(H)\|_{L^p(\sigma)} \|f\|_{L^{p'}(\sigma)},
$$
 which readily implies that

$$
\|\mathcal{N}_{2^*}(u)\|_{L^{p'}(\sigma)} \lesssim \|f\|_{L^{p'}(\sigma)}.
$$

<span id="page-55-0"></span>**Theorem 8.4.** *If*  $(\widetilde{\text{PR}}_q^L)$  *with*  $H = 0$  *is solvable in*  $\Omega$  *for*  $q \in [1,2]$ *, then both*  $(\widetilde{\text{PD}}_{q'}^{L^*})$  *with*  $H = 0$  and  $(\widetilde{D}_{q'}^{L^*})$  are solvable in  $\Omega$ .

*Proof.* Let  $v_1$  be the solution of [\(2.37\)](#page-16-1) with data  $\Xi \in L^2_c(\Omega; \mathbb{R}^{n+1})$  and  $H = 0$  and  $v_2$  be the solution of [\(2.38\)](#page-16-2) with data  $\varphi \in \text{Lip}_{c}(\partial \Omega)$ , and we define  $w = v_1 + v_2$ . Using the tent space duality  $(T_2^{q'} )$  $T_2^{q'}(\Omega))^* = T_2^q$  $L_2^q(\Omega)$  along with the density of  $L_c^2(\Omega)$  functions in  $T_2^q$  $\mathfrak{z}^q_2(\Omega)$ , it holds that

$$
\|\delta_{\Omega}\nabla w\|_{T_2^{q'}(\Omega)} \approx \sup_{\substack{\Psi \in L_c^2(\Omega):\\ \|\Psi\|_{T_2^q(\Omega)}=1}} \Big|\int_{\Omega} \delta_{\Omega}(x) \nabla w \, \Psi \frac{dx}{\delta_{\Omega}(x)}\Big| = \sup_{\substack{\Psi \in L_c^2(\Omega):\\ \|\Psi\|_{T_2^q(\Omega)}=1}} \Big|\int_{\Omega} \nabla w \, \Psi \Big|.
$$

Then, if  $u \in Y_0^{1,2}$  $\mathcal{O}_0^{1,2}(\Omega)$  is the solution of  $(\text{PR}^L_q)$ , with data  $\Psi \in L^2_c(\Omega)$  and  $H = 0$ , by duality and  $(8.2)$ , we have

$$
\left| \int_{\Omega} \Psi \nabla w \right| = \left| - \int_{\Omega} A \nabla u \nabla w + \int_{\partial \Omega} \partial_{\nu_A} u f \right| \le \left| \int_{\Omega} A^* \nabla w \nabla u \right| + \left| \int_{\partial \Omega} \partial_{\nu_A} u f \right|
$$
  
\n
$$
= \left| \int_{\Omega} \Xi \nabla u \right| + \left| \int_{\partial \Omega} \partial_{\nu_A} u f \right|
$$
  
\n
$$
\lesssim ||C_2(\Xi)||_{L^{q'}(\sigma)} ||\widetilde{\mathcal{N}}_2(\nabla u)||_{L^q(\sigma)} + ||\Psi||_{T_2^q(\Omega)} ||f||_{Y^{q'}(\sigma)}
$$
  
\n
$$
\lesssim \left( \|\Xi\|_{\mathcal{C}_{2,q'}(\Omega)} + ||f||_{Y^{q'}(\sigma)} \right) ||\Psi||_{T_2^q(\Omega)},
$$

which proves the desired estimates  $(2.40)$  and  $(2.43)$ .

<span id="page-55-1"></span>**Theorem 8.5.** (i) If 
$$
(PD_p^L)
$$
 is solvable in  $\Omega$  with  $H = 0$  for some  $p \in (1, \infty)$ , then  $(D_p^L)$  is also solvable in  $\Omega$ .

(ii) If  $(\widetilde{PD}_{q}^{L})$  is solvable in  $\Omega$  with  $H = 0$  for some  $q \in [2, \infty]$ , then  $(\widetilde{D}_{q}^{L})$  is also *solvable in* Ω*.*

*Proof.* Let  $f \in \text{Lip}_c(\partial \Omega)$  and let F be the Varopoulos extensions of the  $L^p$ -boundary data f given by Theorem [1.4.](#page-5-2) In the construction of the solution of  $(2.38)$  with data f, we can use F as the Lipschitz extension of f. So if u is the aforementioned solution, then  $u = w + F$ , where w is the solution of [\(2.37\)](#page-16-1) with  $\Xi = -A\nabla F \in L^2(\Omega)$  and  $H = 0$ . Then, by Theorem [1.4](#page-5-2) (ii) and  $(2.42)$ , we have that

$$
\|\widetilde{\mathcal{N}}_{2^*}(u)\|_{L^p(\sigma)} \leq \|\mathcal{N}(F)\|_{L^p(\sigma)} + \|\widetilde{\mathcal{N}}_{2^*}(w)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)} + \|C_2(\boldsymbol{\Xi})\|_{L^p(\sigma)}
$$
  

$$
\leq \|f\|_{L^p(\sigma)} + \|A\|_{L^\infty(\Omega)} \|\mathcal{C}(\nabla F)\|_{L^p(\sigma)} \lesssim \|f\|_{L^p(\sigma)},
$$

showing (i). Similarly, assuming  $(\widetilde{PD}_{q}^{L})$  is solvable in  $\Omega$  with  $H = 0$  for some  $q \in [2, \infty]$ , we obtain

$$
\|\delta_{\Omega}\nabla u\|_{T_2^q(\Omega)} \le \|\delta_{\Omega}\nabla F\|_{T_2^q(\Omega)} + \|\delta_{\Omega}\nabla w\|_{T_2^q(\Omega)} \lesssim \|f\|_{Y^q(\sigma)} + \|C_2(\boldsymbol{\Xi})\|_{L^q(\sigma)}
$$
  

$$
\le \|f\|_{Y^q(\sigma)} + \|A\|_{L^\infty(\Omega)} \|\mathcal{C}(\nabla F)\|_{L^p(\sigma)} \lesssim \|f\|_{Y^q(\sigma)},
$$

where we used Theorem [1.4](#page-5-2) (v) (when  $q = \infty$ , we use the extension  $\tilde{F}$  of Theorem [1.4\)](#page-5-2) and (2.43), which finishes the proof of the Theorem. and [\(2.43\)](#page-17-7), which finishes the proof of the Theorem.

*Proof of Theorem [1.7.](#page-7-0)* It follows by combining Theorems [8.3,](#page-54-0) [8.4,](#page-55-0) and [8.5.](#page-55-1) □

### <span id="page-56-0"></span>8.2. Conditional one-sided Rellich-type inequalities.

**Proposition 8.6.** Suppose that  $(R_p^L)$  is solvable in  $\Omega$  for some  $p \geq 1$ . If u is the solution of  $(2.38)$  *for*  $L^*$  *in*  $\Omega$  *with data*  $f \in \text{Lip}_c(\partial \Omega)$ *, it holds that* 

(8.3) 
$$
\|\partial_{\nu_{A^*}} u\|_{(\dot{M}^{1,p}(\sigma))^*} \lesssim \|f\|_{Y_{p'}(\sigma)},
$$

 $w$ here  $(\dot{M}^{1,p}(\partial\Omega))^*$  stands for the Banach space dual of  $\dot{M}^{1,p}(\partial\Omega)/\mathbb{R}$  and  $Y_{p'}(\sigma)$  is equal *to*  $L^{p'}(\sigma)$  *if*  $p > 1$  *and*  $BMO(\sigma)$  *if*  $p = 1$ *.* 

*Proof.* By definition,

<span id="page-56-1"></span>
$$
\|\partial_{\nu_{A^*}} u\|_{(\dot{M}^{1,p}(\partial\Omega))^*} = \sup_{\substack{\varphi \in \text{Lip}_c(\partial\Omega):\\ \|\varphi\|_{\dot{M}^{1,p}(\sigma)} = 1}} |\langle \partial_{\nu_{A^*}} u, \varphi \rangle|.
$$

Fix  $\varphi \in \text{Lip}_c(\partial \Omega)$  such that  $\|\varphi\|_{\dot{M}^{1,p}(\sigma)} = 1$  and let w be the solution of  $(R_p^L)$  with data  $\varphi$ . Let also  $F \in \text{Lip}(\overline{\Omega})$  be the  $L^{p'}$ -Varopoulos extension of f as constructed in Theorem [1.4.](#page-5-2) Then, by Lemma [8.1,](#page-53-2) we have that

$$
\langle \partial_{\nu_{A^*}} u, \varphi \rangle = \int_{\Omega} A^* \nabla u \nabla w = \int_{\Omega} A \nabla w \nabla (u - F) + \int_{\Omega} A \nabla w \nabla F = \int_{\Omega} A \nabla w \nabla F,
$$

since  $u - F \in Y_0^{1,2}$  $U_0^{1,2}(\Omega)$  and  $Lw = 0$ . Therefore, by duality, Theorem [1.4](#page-5-2) (ii), and [\(2.41\)](#page-17-9), we infer that

$$
|\langle \partial_{\nu_{A^*}} u, \varphi \rangle| = \Big| \int_{\Omega} A \nabla w \nabla F \Big| \leq \|A\|_{L^{\infty}(\Omega)} \|\widetilde{\mathcal{N}}_2(\nabla u)\|_{L^p(\sigma)} \|\mathcal{C}_2(\nabla F)\|_{L^{p'}} \lesssim \|\varphi\|_{\dot{M}^{1,p}(\sigma)} \|f\|_{L^{p'}(\sigma)},
$$

which shows [\(8.3\)](#page-56-1) for  $p > 1$ . The proof in the case  $p = 1$  is similar and we leave it as an exercise.  $\Box$ 

**Proposition 8.7.** Let  $q \geq 1$ . If u is a solution of [\(2.36\)](#page-16-4) for  $H \in L_c^{\infty}(\Omega)$  and  $\Xi \in$  $L_c^{\infty}(\Omega;\mathbb{R}^{n+1})$  such that  $\mathcal{N}_q(\nabla u) \in L^p(\sigma)$  for  $p > 1$ , it holds that

$$
(8.4) \t\t ||\partial_{\nu_{A^*}} u||_{L^p(\sigma)} \lesssim ||\nabla u||_{N_{q,p}(\Omega)} + ||H||_{C_{1,p}(\Omega)} + ||\Xi/\delta_{\Omega}||_{C_{1,p}(\Omega)}.
$$

*If* u is a solution of [\(2.36\)](#page-16-4) for  $H = 0$  and  $\Xi \in L^{\infty}_c(\Omega; \mathbb{R}^{n+1})$  such that  $\mathcal{N}_q(\nabla u) \in L^1(\sigma)$ ,

(8.5) 
$$
\|\partial_{\nu_{A^*}} u\|_{H^1(\sigma)} \lesssim \|\nabla u\|_{N_{q,1}(\Omega)} + \|\Xi\|_{T_2^1(\Omega)}.
$$

<span id="page-57-0"></span>*Proof.* It follows by the same arguments that prove Proposition [8.2.](#page-53-5) We skip the details.  $\square$ 

### <span id="page-57-2"></span>APPENDIX A.

**Lemma A.1.** *Let*  $\Omega \in \text{AR}(s)$  *for*  $s \in (0, n]$ *,*  $u \in L^1_{\text{loc}}(\Omega, \omega_s)$ *,*  $p \in [1, \infty)$ *, and*  $\alpha \ge 1$ *. Then there exists*  $C \geq 1$  *such that for any*  $\xi \in \partial \Omega$  *and*  $r \in (0, 2 \text{ diam}(\partial \Omega))$ *, it holds that* 

<span id="page-57-1"></span>(A.1) 
$$
\|\mathcal{A}^{(\alpha)}(u\mathbf{1}_{B(\xi,r)})\|_{L^p(\sigma|_{B(\xi,r)})} \lesssim r^{\beta} \|\mathscr{C}_s^{(\beta)}(u\mathbf{1}_{B(\xi,Cr)})\|_{L^p(\sigma|_{B(\xi,r)})}.
$$

*If*  $\beta = 0$ *, it also holds* 

(A.2) 
$$
\|\mathcal{A}^{(\alpha)}(u)\|_{L^p(\sigma)} \lesssim \|\mathscr{C}_s(u)\|_{L^p(\sigma)}.
$$

*Moreover for*  $\beta = 0$  *and*  $1 < p \le \infty$  *we have* 

$$
\|\mathscr{C}_s(u)\|_{L^p(\sigma)} \lesssim \|\mathcal{A}^{(\alpha)}(u)\|_{L^p(\sigma)}.
$$

*Proof.* We adapt the proof of [\[HR18,](#page-70-5) Proposition 2.4] and argue by duality. Indeed, if  $1/p + 1/p' = 1$ , we let  $h \in L^{p'}(\sigma)$  be a non-negative function supported in  $B(\xi, r)$  with  $\|h\|_{L^{p'}(\sigma)} = 1$  and such that  $\|\mathcal{A}(u\mathbf{1}_{B(\xi,r)})\|_{L^p(\sigma|_{B(\xi,r)})} \approx \int \mathcal{A}(u\mathbf{1}_{B(\xi,r)})h\,d\sigma$ . Thus,

$$
\|\mathcal{A}(u\mathbf{1}_{B(\xi,r)})\|_{L^{p}(\sigma|_{B(\xi,r)})} \approx \int_{\partial\Omega} \Big( \int_{\gamma_{\alpha}(\xi)\cap B(\xi,r)} |u(y)| \delta_{\Omega}(y)^{-n} dy \Big) h(\xi) d\sigma(\xi)
$$
  
\n
$$
\leq \int_{\Omega\cap B(\xi,r)} |u(y)| \delta_{\Omega}(y)^{s-n} \Big( \delta_{\Omega}(y)^{-s} \int_{B(y,\alpha\delta_{\Omega}(y))} h(\xi) d\sigma(\xi) \Big) dy
$$
  
\n
$$
= \int_{\Omega\cap B(\xi,r)} |u(y)| \delta_{\Omega}(y)^{s-n} H(y) dy
$$
  
\n
$$
= \int_{0}^{\infty} \int_{\Omega\cap B(\xi,r)\cap\{H(y)>\lambda\}} |u(y)| \delta_{\Omega}(y)^{s-n} dy d\lambda,
$$

where, we have set

$$
H(y) := \delta_{\Omega}(y)^{-s} \int_{B(y,\alpha\delta_{\Omega}(y))} h(\xi) d\sigma(\xi).
$$

For any  $y \in \Omega \cap B(\xi, r)$  we let  $\hat{y}$  to be a point in  $B(\xi, r) \cap \partial \Omega$  such that  $|y - \hat{y}| = \delta_{\Omega}(y)$ and set  $B_{\hat{y}} := B(\hat{y}, (\alpha + 1)\delta(y)) \supset B(y, \alpha \delta_{\Omega}(y))$ . Define

$$
E_{\lambda} := \{ y \in \Omega \cap B(\xi, r) : H(y) > \lambda \}
$$

and note that for any  $y \in E_\lambda$ , it holds that  $m_{\sigma,B_0}h > c\lambda$  for some  $c \in (0,1)$  depending on  $\alpha$ . If we set

$$
\widehat{E}_{\lambda} := \{ \zeta \in \partial \Omega : \zeta = \hat{y} \text{ for some } y \in E_{\lambda} \} \text{ and } \mathscr{B}_{\lambda} = \{ B_{\hat{y}} : y \in E_{\lambda} \},
$$

then, there exists large enough  $C > 1$ , such that

$$
\bigcup_{\zeta \in \widehat{E}_{\lambda}} B_{\zeta} \cap \partial \Omega \subset \{ \zeta \in \partial \Omega : \mathcal{M}h(\zeta) > c\lambda \} \cap B(\xi, Cr).
$$

So, by Vitali's covering lemma, there exists a subcollection  $\mathscr{G}_{\lambda} \subset \mathscr{B}_{\lambda}$  of pairwise disjoint balls such that

$$
\bigcup_{B'\in \mathscr B_\lambda}B'\subset \bigcup_{B\in \mathscr G_\lambda}5B.
$$

It is clear that

$$
E_\lambda\subset\bigcup_{B\in\mathscr{G}_\lambda}5B
$$

and thus,

$$
\int_{E_{\lambda}} |u(y)| \delta_{\Omega}(y)^{s-n} dy \leq \sum_{B \in \mathscr{G}_{\lambda}} \int_{5B} |u(y)| \delta_{\Omega}(y)^{s-n} dy
$$
  
\$\lesssim \sum\_{B \in \mathscr{G}\_{\lambda}} \sigma(B) r(B)^{\beta} \inf\_{\zeta \in B \cap \partial \Omega} \mathscr{C}\_{s}^{(\beta)} (u \mathbf{1}\_{B(\xi,Cr)})(\zeta)\$  
\$\lesssim r^{\beta} \int\_{B(\xi,Cr) \cap \{\mathcal{M}h > \lambda\}} \mathscr{C}\_{s}^{(\beta)} (u \mathbf{1}\_{B(\xi,Cr)})(\zeta) d\sigma(\zeta)\$.

Therefore, since  $||h||_{L^{p'}(\sigma)} = 1$ ,

$$
\|\mathcal{A}_{s,\alpha}(u\mathbf{1}_{B(\xi,r)})\|_{L^{p}(\sigma|_{B(\xi,r)})} \lesssim r^{\beta} \int_{0}^{\infty} \int_{B(\xi,Cr)\cap\{\mathcal{M}h>\lambda\}} \mathscr{C}_{s}^{(\beta)}(u\mathbf{1}_{B(\xi,Cr)})(\zeta) d\sigma(\zeta) d\lambda
$$
  

$$
\lesssim r^{\beta} \int_{B(\xi,Cr)} \mathscr{C}_{s}^{(\beta)}(u\mathbf{1}_{B(\xi,Cr)})(\zeta) \mathcal{M}h(\zeta) d\sigma(\zeta)
$$
  

$$
\leq r^{\beta} \|\mathscr{C}_{s}^{(\beta)}(u\mathbf{1}_{B(\xi,Cr)})\|_{L^{p}(\sigma|_{B(\xi,r)})},
$$

proving  $(A.1)$ . The proof of  $(A.2)$  is similar and we omit the details.

Finally, for  $1 < p \leq \infty$ , we have

$$
\mathcal{M}(\mathcal{A}^{(\alpha)}(u))(x) = \sup_{r>0} \frac{1}{\mu(B(x,r))} \int_{B(x,r) \cap \partial\Omega} \mathcal{A}^{(\alpha)}u(\xi) d\sigma(\xi)
$$
  
\n
$$
\gtrsim \sup_{r>0} \frac{1}{r^s} \int_{B(x,r) \cap \partial\Omega} \int_{\gamma_{\alpha}(\xi)} |u(y)| \delta_{\Omega}(y)^{-n} dy d\sigma(\xi)
$$
  
\n
$$
\gtrsim \sup_{r>0} \frac{1}{r^s} \int_{B(x,r) \cap \Omega} \left( \delta_{\Omega}(y)^{-s} \int_{B(y,(1+\alpha)\delta_{\Omega}(y))} d\sigma(\xi) \right) |u(y)| \delta_{\Omega}(y)^{s-n} dy
$$
  
\n
$$
\gtrsim \sup_{r>0} \frac{1}{r^s} \int_{B(x,r) \cap \Omega} |u(y)| \delta_{\Omega}(y)^{s-n} dy = \mathscr{C}_{s}(u)(x).
$$

This implies that

$$
\|\mathscr{C}_s(u)\|_{L^p(\sigma)} \lesssim \|\mathcal{A}^{(\alpha)}(u)\|_{L^p(\sigma)}
$$

and the proof is now complete.  $\Box$ 

<span id="page-58-0"></span>**Lemma A.2.** Let  $B \subset \mathbb{R}^{n+1}$  be a compact and convex set and let  $\{f_n\}_{n\geq 1}$  sequence of differentiable functions in B. Let  $x_0 \in B$  such that the sequence  $\{f_n(x_0)\}_{n=1}^{\infty}$  is convergent. If  $\nabla f_n \to \vec{F}$  uniformly on  $B$ , then, there exists a function  $f : B \to \mathbb{R}$  which is differentiable *at*  $x_0$  *such that* 

$$
f_n \to f
$$
 uniformly on B and  $\vec{F}(x_0) = \nabla f(x_0)$ .

*Moreover, if for every*  $x \in B$ ,  $\{f_n(x)\}_{n=1}^{\infty}$  *is convergent and*  $\nabla f_n$  *is continuous on* B, then ∇f *is continuous on* B *as well.*

*Proof.* <sup>[9](#page-59-0)</sup> For any  $x \in B$ , we write

$$
\frac{f(x) - f(x_0) - \vec{F}(x_0)(x - x_0)}{|x - x_0|} = \frac{f(x) - f(x_0) - (f_n(x) - f_n(x_0))}{|x - x_0|} \n+ \frac{f_n(x) - f_n(x_0) - \nabla f_n(x_0)(x - x_0)}{|x - x_0|} + \frac{(\nabla f_n(x_0) - \vec{F}(x_0))(x - x_0)}{|x - x_0|} \n=: I(x) + II(x) + III(x).
$$

In order to control  $I(x)$ , first note that, since  $\nabla f_n$  converges uniformly on B, for fixed  $\varepsilon > 0$ , there exists  $n_1 = n_1(\varepsilon, B) \in \mathbb{N}$  such that for every  $n, m > n_1$  it holds

<span id="page-59-1"></span>(A.4) 
$$
|\nabla f_n(x) - \nabla f_m(x)| < \frac{\varepsilon}{3} \min\{1, \operatorname{diam}(B)^{-1}\}, \text{ for all } x \in B.
$$

As  $f_n$  is differentiable at  $x_0$ , we have that there exists  $\delta = \delta(\varepsilon, B, x_0)$  such that

<span id="page-59-2"></span>(A.5) 
$$
\frac{|f_n(x) - f_n(x_0) - \nabla f_n(x_0) \cdot (x - x_0)|}{|x - x_0|} < \frac{\varepsilon}{3} \min\{1, \operatorname{diam}(B)^{-1}\},
$$

for all  $x \in (B(x_0, \delta) \setminus \{x_0\}) \cap B$ . By [\(A.4\)](#page-59-1) and [\(A.5\)](#page-59-2), we have that, for any  $n > n_1$ ,

<span id="page-59-3"></span>(A.6) 
$$
\frac{|f_n(x) - f_n(x_0) - f_m(x) + f_m(x_0)|}{|x - x_0|} < \varepsilon \min\{1, \operatorname{diam}(B)^{-1}\},
$$

for all  $x \in (B(x_0, \delta) \setminus \{x_0\}) \cap B$ .

For every  $x, y \in B$ , we set  $\gamma : [0, 1] \to B$  to be the line segment  $\gamma(t) = x + t(y - x)$ for  $t \in [0, 1]$ . Then  $\gamma(0) = x$ ,  $\gamma(1) = y$  and  $\dot{\gamma}(t) = y - x$ . Since B is convex, the line  $\gamma(t)$ lies entirely within B for every  $t \in [0, 1]$ . For every  $n, m > n_1$  we estimate

$$
|f_n(x) - f_n(y) - f_m(x) + f_m(y)| = \left| \int_0^1 (\nabla f_n(\gamma(t)) \dot{\gamma}(t) - \nabla f_m(\gamma(t)) \dot{\gamma}(t)) dt \right|
$$
  

$$
\leq \int_0^1 |\nabla f_n(\gamma(t)) - \nabla f_m(\gamma(t))| |\dot{\gamma}(t)| dt \leq \frac{\varepsilon}{2} \min\{1, \operatorname{diam}(B)^{-1}\} |x - y| \leq \frac{\varepsilon}{3}.
$$

Since the sequence  $\{f_n\}$  converges at  $x_0$ , there exists  $n_2 = n_2(\varepsilon, x_0) \in \mathbb{N}$  such that  $n_2 \geq$  $n_1$  and for every  $n > n_2$ ,

$$
|f_m(x) - f_n(x)| \le |f_n(x) - f_n(x_0) - f_m(x) + f_m(x_0)| + |f_n(x_0) - f_m(x_0)| < \varepsilon,
$$

for every  $x \in B$ . That is, the sequence  $\{f_n\}$  is uniformly Cauchy and so it converges to some function f uniformly in B. Therefore, letting  $m \to \infty$  in [\(A.6\)](#page-59-3) and using [\(A.5\)](#page-59-2), we get that, for every  $n > n_2$ ,

$$
|I(x)|+|II(x)|<2\varepsilon, \quad \text{for all } x\in (B(x_0,\delta)\setminus\{x_0\})\cap B.
$$

Moreover, the sequence  ${\nabla f_n}_{n>1}$  converges to  $\vec{F}$  uniformly in B and thus, there exists  $n_3 \in \mathbb{N}$  such that, for all  $n > n_3$ ,

$$
|III(x)| \leq |\nabla f_n(x_0) - \vec{F}(x_0)| < \varepsilon.
$$

<span id="page-59-0"></span><sup>&</sup>lt;sup>9</sup>This proof was given by Professor Giovanni Leoni at [math.stackexchange/gradient convergence.](https://math.stackexchange.com/questions/2255618/generalization-of-theorem-from-mathbb-r-to-mathbb-rn)

Setting  $n_0 = \max\{n_1, n_2, n_3\}$  and using the above estimates we get that, for all  $n > n_0$ , it holds

$$
\left|\frac{f(x)-f(x_0)-\vec{F}(x_0)(x-x_0)}{|x-x_0|}\right|<3\varepsilon,
$$

for all  $x \in (B(x_0, \delta) \setminus \{x_0\}) \cap B$ , which implies that f if differentiable at  $x_0$  with  $\nabla f(x_0) =$  $\vec{F}(x_0)$ . Finally, if  $f_n$  is pointwisely convergent on B and  $\nabla f_n$  is continuous on B, then  $\nabla f$ is continuous on B as the uniform limit of a sequence of continuous functions.  $\Box$ 

<span id="page-60-0"></span>**Lemma A.3.** *If*  $F \in L^1_{loc}(\Omega)$ *, then for any*  $x \in \Omega$  *it holds that* 

$$
|F(x)| \lesssim \frac{1}{\delta_{\Omega}(x)^{1+n/p}} \|\mathcal{C}_s(F)\|_{L^p(\sigma)}, \quad \text{for every } p \in (1, \infty)
$$

*and*

$$
|F(x)| \lesssim \frac{1}{\delta_{\Omega}(x)} \sup_{\xi \in \partial \Omega} C_s(F)(\xi).
$$

*Proof.* Fix  $x \in \Omega$  and note that if  $c' = \frac{c}{c+1}$ , then for any  $z \in B(x, c' \delta_{\Omega}(x))$  we have that

$$
|z - x| \le c' \operatorname{dist}(x, \partial \Omega) \le c' \operatorname{dist}(z, \partial \Omega) + c' |z - x|,
$$

which implies that  $|z - x| \leq c \delta_{\Omega}(z)$ , i.e.,  $x \in B_z$ . If  $\xi_x \in \partial \Omega$  is a point such that  $\delta_{\Omega}(x) = |x - \xi_x|$ , it is clear that  $B_x \subset B(\xi, 3\delta_{\Omega}(x)) \cap \Omega$  for every  $\xi \in B(\xi_x, \delta_{\Omega}(x))$ . So

$$
\begin{split} \|\mathcal{C}_{s}(F)\|_{L^{p}(\sigma)} &= \Big(\int_{\partial\Omega}\Big[\sup_{r>0}\frac{1}{r^{s}}\int_{B(\xi,r)\cap\Omega}\sup_{y\in B_{z}}|F(y)|\,\omega_{s}(z)\,dz\Big]^{p}\,d\sigma(\xi)\Big)^{1/p} \\ &\gtrsim \Big(\int_{B(\xi_{x},\delta_{\Omega}(x))\cap\partial\Omega}\Big[\frac{1}{\delta_{\Omega}(x)^{s}}\int_{B(\xi,3\delta_{\Omega}(x))}\sup_{y\in B_{z}}|F(y)|\,\omega_{s}(z)\,dz\Big]^{p}\,d\sigma(\xi)\Big)^{1/p} \\ &\gtrsim \Big(\int_{B(\xi_{x},\delta_{\Omega}(x))\cap\partial\Omega}\Big[\frac{1}{\delta_{\Omega}(x)^{n}}\int_{B(x,c'\delta_{\Omega}(x))}\sup_{y\in B_{z}}|F(y)|\,dz\Big]^{p}\,d\sigma(\xi)\Big)^{1/p} \\ &\gtrsim \delta_{\Omega}(x)^{1+\frac{n}{p}}|F(x)|. \end{split}
$$

Note that for  $p = \infty$ , by the same argument, we can directly infer that

$$
\sup_{\xi \in \partial \Omega} C_s(F)(\xi) \gtrsim \delta_{\Omega}(x)|F(x)|.
$$

 $\Box$ 

### Definition A.4. We define the spaces

(A.7) 
$$
N_{\sharp,\mathcal{D}}^{\infty}(\Omega) := \{ w \in C(\Omega) : \sup_{P \in \mathcal{W}(\Omega)} \sup_{x \in P} |w(x) - \int_{P} w| < \infty \}
$$

(A.8) 
$$
N_{\mathcal{D}}^{\infty}(\Omega) := \{ w \in C(\Omega) : \sup_{P \in \mathcal{W}(\Omega)} \sup_{x \in P} |w(x)| < \infty \}
$$

and equip them with

$$
||w||_{N_{\sharp,\mathcal{D}}^{\infty}(\Omega)} := \sup_{P \in \mathcal{W}(\Omega)} \sup_{x \in P} |w(x) - \int_{P} w|, \text{ and}
$$

$$
||w||_{N_{\mathcal{D}}^{\infty}(\Omega)} := \sup_{P \in \mathcal{W}(\Omega)} \sup_{x \in P} |w(x)|,
$$

respectively. Note that  $||w||_{N^{\infty}_{\sharp,D}(\Omega)}$  is a semi-norm, while  $||w||_{N^{\infty}_{D}(\Omega)}$  is a norm. We also define the space

(A.9)  $N_{\text{sum},\mathcal{D}}(\Omega) := \{ u \in C^1(\Omega) : (u, \delta_{\Omega} \nabla u) \in N_{\sharp,\mathcal{D}}^{\infty}(\Omega) \times N_{\mathcal{D}}^{\infty}(\Omega) \}$ 

and equip it with the semi-norm

$$
||u||_{N_{\text{sum},\mathcal{D}}(\Omega)} := ||u||_{N_{\sharp,\mathcal{D}}^{\infty}(\Omega)} + ||\delta_{\Omega} \nabla u||_{N_{\mathcal{D}}^{\infty}(\Omega)}.
$$

<span id="page-61-0"></span>**Lemma A.5.** If  $\Omega\subset\mathbb{R}^{n+1}$  is an open and connected set, then  $\mathrm{N}_{\sharp,\mathcal{D}}^{\infty}(\Omega)/\mathbb{R}$  and  $\mathrm{N}_{\mathrm{sum},\mathcal{D}}^{\infty}(\Omega)/\mathbb{R}$ *are normed spaces.*

*Proof.* Since it is easy to see that  $\|\cdot\|_{N_{\sharp,\mathcal{D}}^{\infty}(\Omega)}$  and  $\|\cdot\|_{N_{\text{sum},\mathcal{D}}^{\infty}}$  are semi-norms, we will only show that if  $||u||_{N^{\infty}_{\sharp,\mathcal{D}}(\Omega)} = 0$ , then there exists  $c \in \mathbb{R}$  such that  $u = c$  in  $\Omega$ . Indeed, if  $\sup_{P \in \mathcal{W}(\Omega)} ||u||_{\Lambda_P} = 0$ , then  $\max_{x \in \overline{P}} |u(x) - f_P u| = 0$ , for every  $P \in \mathcal{W}(\Omega)$ , which implies  $u = f_P u$  on  $\overline{P}$  for every  $P \in \mathcal{W}(\Omega)$ . If  $P_1, P_2 \in \mathcal{W}(\Omega)$  are such that  $\overline{P}_1 \cap \overline{P}_2 \neq \emptyset$ , we have that  $u = f_{P_1} u$  on  $\bar{P}_1$  and  $u = f_{P_2} u$  on  $\bar{P}_2$ . As there exists  $\xi \in \partial P_1 \cap \partial P_2$ and u is continuous in  $\Omega$ , it holds that  $f_{P_1} u = f_{P_2} u$  for every  $P_1, P_2 \in \mathcal{W}(\Omega)$  such that  $\bar{P}_1 \cap \bar{P}_2 \neq \emptyset$ . So, if  $\sup_{P \in \mathcal{W}(\Omega)} ||u||_{\Lambda_P} = 0$ , there exists a constant  $c \in \mathbb{R}$  so that  $u = c$  in  $\Omega$ , which implies that  $N^{\infty}_{\sharp,\mathcal{D}}(\Omega)/\mathbb{R}$  and  $N^{\infty}_{sum,\mathcal{D}}(\Omega)/\mathbb{R}$  are normed spaces.

**Remark A.6.** By Lemma [A.5,](#page-61-0) we have that if  $\Omega$  is open and connected set, and a sequence converges in  $N^{\infty}_{\sharp,\mathcal{D}}(\Omega)/\mathbb{R}$  or  $N_{sum,\mathcal{D}}(\Omega)$ , then the limit is unique modulo constants.

<span id="page-61-1"></span>**Lemma A.7.** Let  $\Omega \subset \mathbb{R}^{n+1}$  be an open set. Then both  $(N^{\infty}_{\sharp,\mathcal{D}}(\Omega), \|\cdot\|_{N^{\infty}_{\sharp,\mathcal{D}}(\Omega)})$  and  $(N_{sum,\mathcal{D}}(\Omega), \|\cdot\|_{N_{sum,\mathcal{D}}(\Omega)})$  *are sequentially complete.* 

*Proof.* Let us first assume that  $\Omega$  is connected. We define the space

$$
\Lambda_P := \{ u : \Omega \to \mathbb{R} : u \in C(\bar{P}) \text{ and } \sup_{x \in P} |u(x) - \int_P u| < \infty \},
$$

and equip it with the semi-norm  $||u||_{\Lambda_P} := \sup_{x \in P} |u(x) - f_P u|$ . In fact, by the continuity of  $u$  on  $P$ , we have that

$$
||u||_{\Lambda_P} = \max_{x \in \overline{P}} |u(x) - \int_P u|.
$$

We will first prove that the space  $\Lambda_P$  is sequentially complete with respect to the seminorm  $\|\cdot\|_{\Lambda_P}$  for any fixed  $P \in \mathcal{W}(\Omega)$ . To this end, let  $\{u_n\}_{n\in\mathbb{N}}$  be a Cauchy sequence in  $\Lambda_P$  and fix  $\varepsilon > 0$ . Then, there exists  $n_0 = n_0(\varepsilon, P) \in \mathbb{N}$  such that for every  $n, m > n_0$  it holds that  $||u_n - u_m||_{\Lambda_P} < \varepsilon$ . Consequently, for any  $y \in P$ ,

$$
\left| u_n(y) - u_m(y) - \int_P (u_n(z) - u_m(z)) dz \right| < \varepsilon,
$$

which means that the sequence  $\{u_n - f_p u_n\}_{n \in \mathbb{N}}$  is uniformly Cauchy on  $\overline{P}$  and so it converges uniformly on  $\overline{P}$  to some  $u_P$ . Thus, there exists a positive integer  $n_1 = n_1(\varepsilon, P)$ such that for any  $n > n_1$ ,

(A.10) 
$$
\max_{y \in \overline{P}} \left| u_n(y) - \int_P u_n - u_P(y) \right| < \varepsilon/2,
$$

and so, for any  $n > n_1$ ,

(A.11) 
$$
\max_{y \in \overline{P}} \left| u_n(y) - u_P(y) - \oint_P (u_n - u_P) \right|
$$
  
\n
$$
\leq \max_{y \in \overline{P}} \left| u_n(y) - \oint_P u_n - u_P(y) \right| + \left| \oint_P (u_P(z) + u_n(z) - u_n(z)) dz \right|
$$
  
\n
$$
\leq \max_{y \in \overline{P}} \left| u_n(y) - \oint_P u_n - u_P(y) \right| + \int_P \left| u_n(z) - \oint_P u_n - u_P(z) \right| dz < \varepsilon,
$$

concluding that  $\Lambda_P$  is sequentially complete. Moreover,

<span id="page-62-0"></span>
$$
\left| \int_{P} u_{P} \right| \leq \int_{P} \left| u_{P} - u_{n}(x) + \int_{P} u_{n} \right| \leq \max_{x \in \overline{P}} \left| u_{n}(x) - \int_{P} u_{n} - u_{P}(x) \right| \stackrel{n \to \infty}{\longrightarrow} 0,
$$

which implies that

$$
(A.12)\qquad \qquad \int_P u_P = 0.
$$

It is easy to see that since the half-open Whitney cubes  $P \in \mathcal{W}(\Omega)$  are disjoint, the countable direct sum  $\bigoplus_{P \in \mathcal{W}(\Omega)} \Lambda_P$  equipped with the sup norm is sequentially complete. Indeed, if  $\{u_n\}$  be a Cauchy sequence in  $\bigoplus_{P \in \mathcal{W}(\Omega)} \Lambda_P$ , then for  $\varepsilon > 0$ , there exists  $n_0 =$  $n_0(\varepsilon) \in \mathbb{N}$  such that

$$
\max_{\bar{P}} \left| u_n - \oint_P u_n - u_m + \oint_P u_m \right| < \varepsilon, \quad \text{for all } m, n > n_0 \text{ and all } P \in \mathcal{W}(\Omega).
$$

Using that  $\lim_{m\to\infty} (u_m - f_p u_m) = u_p$  uniformly on  $\bar{P}$ , we take limits as  $m \to \infty$  in the last inequality and infer that

$$
\max_{\bar{P}} \left| u_n(x) - \oint_P u_n - u_P(x) \right| \le \varepsilon, \quad \text{for all } n > n_0 \text{ and all } P \in \mathcal{W}(\Omega).
$$

Then, the function defined by  $v(x) = u_P(x)$  for every  $x \in P$  and every  $P \in \mathcal{W}(\Omega)$ , satisfies  $u_n \to v$  in  $\bigoplus_{P \in \mathcal{W}(\Omega)} \Lambda_P$ .

We shall now prove that  $\dot{N}_{\sharp,\mathcal{D}}(\Omega)$  is sequentially complete with respect to the semi-norm  $\| \cdot \|_{N^{\infty}_{\sharp,D}(\Omega)}$ . Let  $\{u_n\}$  be a Cauchy sequence in  $N_{\sharp,D}(\Omega)$ . For fixed  $\varepsilon > 0$ , there exists  $n_0 = n_0(\varepsilon) \in \mathbb{N}$  such that for every  $n, m > n_0$ , we have that  $||u_n - u_m||_{N_{\sharp D}(\Omega)} < \varepsilon$ . Note that

$$
\mathcal{N}_{\sharp,\mathcal{D}}^{\infty}(\Omega)=\bigoplus_{P\in\mathcal{W}(\Omega)}\Lambda_{P}\cap C(\Omega)
$$

and as  $\bigoplus_{P \in \mathcal{W}(\Omega)} \Lambda_P$  is sequentially complete, there exists v (as defined above) such that  $u_n \to v$  in the  $\|\cdot\|_{N_{\sharp,\mathcal{D}}^{\infty}(\Omega)}$  semi-norm. So there exists  $\{u_P\}_{P\in\mathcal{W}(\Omega)}$  satisfying [\(A.12\)](#page-62-0) such that  $u_n \to u_P$  in the  $\|\cdot\|_{\Lambda_P}$  semi-norm, uniformly in  $P \in \mathcal{W}(\Omega)$ . Note that  $u_P \in C(\overline{P})$ for each  $P \in \mathcal{W}(\Omega)$  as the uniform limit of a sequence of continuous functions.

Our main goal is to modify each  $u_p$  adding a suitable constant so that we can define a new function  $u \in C(\Omega)$  which satisfies  $u_n \to u$  in  $\|\cdot\|_{N^{\infty}_{\sharp,\mathcal{D}}(\Omega)}$ . To do so, it is important to show that if  $P_1, P_2 \in \mathcal{W}(\Omega)$  with  $\partial P_1 \cap \partial P_2 \neq \emptyset$ , then, for every  $x \in \partial P_1 \cap \partial P_2$ ,

 $(A.13)$  $(x) + C(P_1, P_2) = u_{P_2}(x).$ 

If

<span id="page-63-0"></span>
$$
C_n(P_1, P_2) := \int_{P_1} u_n - \int_{P_2} u_n,
$$

since  $\{u_n\}$  is Cauchy in  $N^{\infty}_{\sharp,\mathcal{D}}(\Omega)$ , for any  $x \in \partial P_1 \cap \partial P_2$ , it holds that  $|C_n(P_1, P_2) - C_m(P_1, P_2)|$ 

$$
\begin{aligned} \n\mathcal{L}_n(1,1,1,2) &= \mathcal{L}_m(1,1,1,2) \\ \n&\le \left| u_n(x) - u_m(x) - \int_{P_1} (u_n - u_m) \right| + \left| u_n(x) - u_m(x) - \int_{P_2} (u_n - u_m) \right| \\ \n&\le 2 \| u_n - u_m \|_{\mathcal{N}^{\infty}_{\sharp, \mathcal{D}}(\Omega)} \to 0, \quad \text{as } m, n \to \infty. \n\end{aligned}
$$

Therefore, there exists  $C(P_1, P_2) \in \mathbb{R}$  such that  $\lim_{n\to\infty} C_n(P_1, P_2) = C(P_1, P_2)$ . So, for every  $x \in \partial P_1 \cap \partial P_2$ ,

$$
u_{P_2}(x) - u_{P_1}(x) = \left(u_{P_2}(x) - u_n(x) + \int_{P_2} u_n\right) - \left(u_{P_1}(x) - u_n(x) + \int_{P_1} u_n\right) + C_n(P_1, P_2) \stackrel{n \to \infty}{\longrightarrow} C(P_1, P_2),
$$

which shows  $(A.13)$ .

If  $\partial\Omega$  is compact, we fix a (starting) cube  $P_0 \in \mathcal{W}(\Omega)$  such that  $\ell(P_0) \leq \text{diam}(\partial\Omega)$  and  $\ell(P) \leq \ell(P_0)$  for every  $P \in \mathcal{W}(\Omega)$ . If  $\partial\Omega$  is unbounded, we pick as a starting cube some  $P_0 \in \mathcal{W}(\Omega)$  such that  $\ell(P_0) = 1$ . Once we have fixed such a cube  $P_0$ , we define

$$
\mathcal{G}_1(P_0) := \{ P \in \mathcal{W}(\Omega) : \partial P \cap \partial P_0 \neq \emptyset \}.
$$

For every  $P \in \mathcal{G}_1(P_0)$ , we let  $u_P$  be the limit of  $\{u_n\}$  in the  $\Lambda_P$  semi-norm and set

(A.14) 
$$
u_P^1 := u_P + C(P, P_0).
$$

It is clear that  $u_n \to u_P^1$  in the  $\Lambda_P$  semi-norm as well, while, in view of [\(A.13\)](#page-63-0),  $u_P^1 = u_{P_0}$ on  $\partial P_0 \cap \partial P$ . Observe that, repeating the proof of [\(A.13\)](#page-63-0), we can show that for every  $\widetilde{P} \in \mathcal{G}_1(P) \cap \mathcal{G}_1(P_0) \setminus \{P_0\}$ , there exists a constant  $C(P, \widetilde{P})$  such that

(A.15) 
$$
u_P^1 + C(P, \widetilde{P}) = u_{\widetilde{P}}^1 \quad \text{on } \partial P \cap \partial \widetilde{P}.
$$

Since  $P, \widetilde{P} \in \mathcal{G}_1(P_0)$  and  $\widetilde{P} \in \mathcal{G}_1(P)$  we have

<span id="page-63-1"></span>
$$
u_{\widetilde{P}}^1(z) = u_{P_0}(z) = u_P^1(z) \quad \forall z \in \partial P_0 \cap \partial P \cap \partial \widetilde{P},
$$

which implies that  $C(P, \tilde{P}) = 0$  and so, by [\(A.15\)](#page-63-1),

(A.16) 
$$
u_P^1 = u_{\tilde{P}}^1 \qquad \text{on } \partial P \cap \partial \tilde{P}.
$$

Setting  $C_P := C(P, P_0)$ , we can write  $u_P^1 := u_P + C_P$  and define

$$
v^1 := \begin{cases} u_P^1, & P \in \mathcal{G}_1(P_0) \\ u_P, & P \in \mathcal{W}(\Omega) \setminus \mathcal{G}_1(P_0). \end{cases}
$$

Notice that

$$
p^1 \in C(\mathcal{O}_1), \quad \text{where} \quad \mathcal{O}_1 := \bigcup_{P \in \mathcal{G}_1(P_0)} \bar{P},
$$

and  $\lim_{n\to\infty} \|u_n - v^1\|_{\mathcal{N}^{\infty}_{\sharp,\mathcal{D}}(\Omega)} = 0.$ 

Moving forward, for every  $P \in \mathcal{G}_1(P_0)$ , we set

v

$$
\widetilde{\mathcal{G}}_1(P) := \mathcal{G}_1(P) \setminus \mathcal{G}_1(P_0) \text{ and } \mathcal{G}_2(P_0) := \bigcup_{P \in \mathcal{G}_1(P_0)} \widetilde{\mathcal{G}}_1(P).
$$

If  $P \in \mathcal{G}_1(P_0)$  and  $P' \in \mathcal{G}_1(P)$ , we define

$$
u_{P'}^2 := u_{P'} + C(P', P),
$$

which also satisfies  $u_n \to u_{P'}^2$  in the  $\Lambda_{P'}$  semi-norm. By the same arguments as above, we can show that

$$
u_{P'}^2 = u_P^1 \qquad \text{on } \partial P \cap \partial P'
$$

and, for every  $P'' \in \widetilde{\mathcal{G}}_1(P)$  such that  $\partial P' \cap \partial P'' \neq \emptyset$ ,

$$
u_{P'}^2 = u_{P''}^2 \qquad \text{on } \partial P' \cap \partial P''
$$

.

Moreover, if  $P \in \mathcal{G}_1(P_0)$  such that  $\partial P \cap \partial P \neq \emptyset$  and  $P' \in \mathcal{G}_1(P) \cap \mathcal{G}_1(P)$ , we define  $\widetilde{u}_{P'}^2 := u_{P'} + C(P', \widetilde{P})$ . Then, for every  $x \in \partial P' \cap \partial P \cap \partial \widetilde{P}$ , it holds that

$$
u_P^1(x) = u_{\widetilde{P}}^1(x), \quad u_{P'}^2(x) = u_P^1(x), \quad \text{and} \quad \widetilde{u}_{P'}^2(x) = u_{\widetilde{P}}^1(x),
$$

which we may combine to deduce that  $C(P', P) = C(P', \tilde{P})$  and consequently

$$
u_{P'}^2 = \widetilde{u}_{P'}^2.^{10}
$$

The same is true for every  $\tilde{P} \in \mathcal{G}_1(P_0)$  such that  $P' \in \tilde{\mathcal{G}}_1(P) \cap \tilde{\mathcal{G}}_1(\tilde{P})$  since, for every such cube, there is a chain of cubes  $P = P_1, P_2, \ldots, P_N = P$  such that  $P_k \in \mathcal{G}_1(P_0)$  and  $\partial P_k \cap \partial P_{k+1} \neq \emptyset$  for  $k \in \{1, ..., N\}$ , where N is a dimensional constant. Therefore, we can unambiguously set  $C_{P'} := C(P', P)$  for any  $P \in \mathcal{G}_1(P_0)$  such that  $P' \in \widetilde{\mathcal{G}}_1(P)$  and so  $u_{P'}^2 := u_{P'} + C_{P'}$ . If

$$
\mathcal{F}_2(P_0) := \mathcal{G}_1(P_0) \cup \mathcal{G}_2(P_0),
$$

we define

$$
v^2 := \begin{cases} u_P^1, & P \in \mathcal{G}_1(P_0) \\ u_P^2, & P \in \mathcal{G}_2(P_0) \\ u_P, & P \in \mathcal{W}(\Omega) \setminus \mathcal{F}_2(P_0) \end{cases}
$$

and notice that

$$
v^2 \in C(\mathcal{O}_2), \quad \text{where} \quad \mathcal{O}_2 := \bigcup_{P \in \mathcal{F}_2(P_0)} \bar{P},
$$

and  $\lim_{n\to\infty} \|u_n - v^2\|_{\mathcal{N}_{\sharp,\mathcal{D}}^{\infty}(\Omega)} = 0.$ 

<span id="page-64-0"></span><sup>&</sup>lt;sup>10</sup>The part of the argument dealing with two descendants and one ancestor is better suited to cases where the descendants have, at most, equal side-lengths with the ancestor. Meanwhile, the part involving two ancestors and one descendant is more relevant for cases where the descendant has a larger side-length than the ancestor. The terms "ancestor" and "descendant" are used in relation to the selection process.

We proceed by iteration and, for every  $P \in \mathcal{G}_{k-1}(P_0)$ ,  $k \geq 2$ , we set

$$
\widetilde{\mathcal{G}}_1(P) := \mathcal{G}_1(P) \setminus \mathcal{G}_{k-1}(P_0) \quad \text{and} \quad \mathcal{G}_k(P_0) := \bigcup_{P \in \mathcal{G}_{k-1}(P_0)} \widetilde{\mathcal{G}}_1(P).
$$

If  $P \in \mathcal{G}_{k-1}(P_0)$  and  $P' \in \widetilde{\mathcal{G}}_1(P)$ , we define

$$
u_{P'}^k := u_{P'} + C(P', P)
$$

which also satisfies that  $u_n \to u_{P'}^k$  in the  $\Lambda_{P'}$  semi-norm. Arguing as above, we can prove that

$$
u_{P'}^k = u_P^{k-1} \qquad \text{on } \partial P \cap \partial P'
$$

and, for every  $P'' \in \widetilde{\mathcal{G}}_1(P)$  such that  $\partial P' \cap \partial P'' \neq \emptyset$ ,

$$
u_{P'}^k = u_{P''}^k \qquad \text{on } \partial P' \cap \partial P''.
$$

Moreover, we can unambiguously set  $C_{P'} := C(P', P)$  for any  $P \in \mathcal{G}_{k-1}(P_0)$  such that  $P' \in \widetilde{\mathcal{G}}_1(P)$  and so  $u_{P'}^k := u_{P'} + C_{P'}$ . If

$$
\mathcal{F}_k(P_0) := \bigcup_{j=1}^k \mathcal{G}_j(P_0)
$$

and

$$
v^{k} := \begin{cases} u_{P}^{1}, & P \in \mathcal{G}_{1}(P_{0}) \\ u_{P}^{2}, & P \in \mathcal{G}_{2}(P_{0}) \\ \vdots & \vdots \\ u_{P}^{k}, & P \in \mathcal{G}_{k}(P_{0}) \\ u_{P}, & P \in \mathcal{W}(\Omega) \setminus \mathcal{F}_{k}(P_{0}), \end{cases}
$$

it is easy to see that

$$
v^k \in C(\mathcal{O}_k), \quad \text{where} \quad \mathcal{O}_k := \bigcup_{P \in \mathcal{F}_k(P_0)} \bar{P},
$$

and  $\lim_{n\to\infty} \|u_n - v^k\|_{\mathrm{N}^\infty_{\sharp,\mathcal{D}}(\Omega)} = 0.$ 

If we define  $u(x) := \lim_{k \to \infty} v^k(x)$  for  $x \in \Omega$ , then, by construction and the fact that  $\Omega$ is connected, it is clear that  $u \in C(\Omega)$  and  $\lim_{n \to \infty} ||u_n - u||_{N^{\infty}_{\sharp,\mathcal{D}}(\Omega)} = 0$ . Hence,  $N^{\infty}_{\sharp,\mathcal{D}}(\Omega)$ is sequentially complete.

It remains to prove that  $N_{sum,\mathcal{D}}(\Omega)$  is sequentially complete. To this end, let  $\{u_n\}$  be a Cauchy sequence in  $N_{sum,\mathcal{D}}(\Omega)$ . Then for fixed  $\varepsilon > 0$ , there exists  $n_0 = n_0(\varepsilon) \in \mathbb{N}$  such that for every  $n, m \ge n_0$  we have  $||u_n - u_m||_{N_{\text{sum},\mathcal{D}}(\Omega)} < \varepsilon$ , which implies that

<span id="page-65-0"></span>
$$
||u_n - u_m||_{N^{\infty}_{\sharp,\mathcal{D}}(\Omega)} < \varepsilon
$$
 and  $||\delta_{\Omega}(\nabla u_n - \nabla u_m)||_{N^{\infty}_{\mathcal{D}}(\Omega)} < \varepsilon$ .

Since  $N^{\infty}_{\sharp,\mathcal{D}}(\Omega)$  is sequentially complete, if  $u = \lim_{n\to\infty} u_n \in N^{\infty}_{\sharp,\mathcal{D}}(\Omega)$ , it suffices to show that

(A.17) 
$$
u \in C^{1}(\Omega) \text{ and } \lim_{n \to \infty} ||\delta_{\Omega}(\nabla u_{n} - \nabla u)||_{N_{\mathcal{D}}^{\infty}(\Omega)} = 0.
$$

Since  $\{\delta_{\Omega} \nabla u_n\}_{n=1}^{\infty} \subset C(\Omega)$  is Cauchy in  $N_{\mathcal{D}}^{\infty}(\Omega)$ , then it is uniformly Cauchy on  $\overline{P}_{\Omega}$ (uniformly in  $P \in W(\Omega)$ ) and so there exists  $\vec{w}_P^0 \in C(\overline{P})$  such that  $\delta_{\Omega} \nabla u_n \to \vec{w}_P^0$ uniformly on  $\overline{P}$  for every  $P \in \mathcal{W}(\Omega)$ . If  $P_1, P_2 \in \mathcal{W}(\Omega)$  so that  $\partial P_1 \cap \partial P_2 \neq \emptyset$ , for every  $x \in \partial P_1 \cap \partial P_2$ , it holds that

$$
\vec{w}_{P_1}^0(x) - \vec{w}_{P_2}^0(x) = (\vec{w}_{P_1}^0(x) - \delta_{\Omega}(x)\nabla u_n(x)) - (\vec{w}_{P_1}^0(x) - \delta_{\Omega}(x)\nabla u_n(x)) \to 0, \text{ as } n \to \infty,
$$

which implies that  $\vec{w}_{P_1}^0(x) = \vec{w}_{P_2}^0(x)$  for every  $x \in \partial P_1 \cap \partial P_2$ . Therefore, if we define  $\vec{w}_0(x) := \vec{w}_P^0(x)$  for every  $x \in P$  and all  $P \in \mathcal{W}(\Omega)$ , it is evident that  $\vec{w}_0 \in C(\Omega)$  and  $\lim_{n\to\infty} ||\delta_{\Omega}\nabla u_n - \vec{w}_0||_{N_{\mathcal{D}}^{\infty}(\Omega)} = 0.$  Set now  $\vec{w} := \delta_{\Omega}^{-1}$  $\overline{\Omega}^{-1} \vec{w}_0 \in C(\Omega)$  and

$$
w_P^n := u_n - \oint_P u_n \quad \text{on } \bar{P},
$$

and note that

$$
w_P^n \to u_P
$$
 uniformly on  $\overline{P}$  and  $\nabla w_P^n = \nabla u_n \to \overrightarrow{w}$  uniformly on  $\overline{P}$ .

We can now apply Lemma [A.2](#page-58-0) on each  $\bar{P}$  to the sequence  $\{w_P^n\}_{n=1}^{\infty} \subset C^1(\bar{P})$  and deduce that, for every  $P \in \mathcal{W}(\Omega)$ ,

$$
u_P
$$
 is differentiable on  $\overline{P}$  and  $\nabla u_P = \overline{w} \in C(\overline{P})$ .

In particular,  $\nabla u_P(z) = \vec{w}(z) = \nabla u_{P'}(z)$  for every  $z \in \partial P \cap \partial P'$ . If  $P \in \mathcal{W}(\Omega)$  and  $y \in \mathcal{W}(\Omega)$  $\partial P$ , for fixed  $\varepsilon > 0$ , there exist  $\delta_P = \delta_P(\varepsilon, y) > 0$  such that for every  $x \in \overline{P} \cap B(y, \delta_P)$ ,

(A.18) 
$$
\frac{|u_P(x) - u_P(y) - \vec{w} \cdot (x - y)|}{|x - y|} < \varepsilon.
$$

Let us fix a cube  $P \in \mathcal{W}(\Omega)$  and a point  $z \in \partial P$ . We define Z to be the family of Whitney cubes P' such that  $z \in \partial P'$  (the number of such cubes is at most a fixed dimensional constant) and set

<span id="page-66-0"></span>
$$
\delta = \min \Big\{ \text{dist}\Big(z,\Omega \setminus \bigcup_{P' \in \mathcal{Z}} \bar{P}'\Big), \min_{P' \in \mathcal{Z}} \delta_{P'} \Big\}.
$$

Note that for every  $x \in B(z, \delta) \setminus \partial P$ , there exists a unique  $P_x \in \mathcal{Z}$  such that  $x \in P_x^o$  and, by [\(A.18\)](#page-66-0),

$$
\frac{|u(x)-u(z)-\vec{w}\cdot(x-z)|}{|x-z|}=\frac{|u_{P_x}(x)-u_{P_x}(z)-\vec{w}\cdot(x-z)|}{|x-z|}<\varepsilon.
$$

If  $x \in B(z, \delta) \cap \partial P$ , since  $u(x) = u_P(x) + C_P$ , by [\(A.18\)](#page-66-0), it holds that

$$
\frac{|u(x)-u(z)-\vec{w}\cdot(x-z)|}{|x-z|}=\frac{|u_P(x)-u_P(z)-\vec{w}\cdot(x-z)|}{|x-z|}<\varepsilon,
$$

showing that u is differentiable at any  $z \in \bigcup_{P \in \mathcal{W}(\Omega)} \partial P$ . Therefore, since  $u_P$  is differentiable in  $P^o$  for any  $P \in \mathcal{W}(\Omega)$  and  $u = u_P + C_P$  in  $P^o$ , we deduce that u is differentiable in  $\Omega$  and  $\nabla u = \vec{w} \in C(\Omega)$ , proving [\(A.17\)](#page-65-0) and concluding the proof of the lemma when  $\Omega$ is connected.

If  $\Omega$  is not connected, it can be written as the union of at most countably many disjoint connected components, i.e.,  $\Omega = \bigcup_{i=1}^{\infty} \Omega_i$ . For every  $i \in \mathbb{N}$ , we have proved that  $N_{\sharp,\mathcal{D}}(\Omega_i)$  and  $N_{sum,\mathcal{D}}(\Omega_i)$  are sequentially complete and, as the connected components  $\Omega_i$  are mutually disjoint, we have that

$$
N_{\sharp,\mathcal{D}}(\Omega)=\bigoplus_{i=1}^{\infty}N_{\sharp,\mathcal{D}}(\Omega_i)\ \ \text{and}\ \ N_{\text{sum},\mathcal{D}}(\Omega)=\bigoplus_{i=1}^{\infty}N_{\text{sum},\mathcal{D}}(\Omega_i)\quad\text{are sequentially complete},
$$

<span id="page-67-0"></span>concluding the proof of Lemma  $A.7$ .

**Lemma A.8.** Let  $\Omega \subset \mathbb{R}^{n+1}$  be an open set. The semi-norms  $\|\cdot\|_{N_{\text{sum}}(\Omega)}$  and  $\|\cdot\|_{N_{\text{sum},\mathcal{D}}(\Omega)}$ *are equivalent and the implicit constants only depend on* n*.*

*Proof.* If  $x \in \Omega$  and  $B_x = B(x, c \delta_{\Omega}(x))$ , for  $c \in (0, 1/2]$ , then, by easy volume considerations, one can prove that there exists a uniformly bounded number of Whitney cubes  $P \in \mathcal{W}(\Omega)$  that cover the ball  $B_x$ . We denote this collection of cubes by  $\mathcal{B}_x$ . By the mean value theorem, for any  $P \in \mathcal{B}_x$ , we estimate

$$
\left| \oint_{B_x} u - \oint_P u \right| \leq \oint_{B_x} \oint_P |u(z) - u(\zeta)| \, dz \, d\zeta \lesssim \delta_\Omega(x) \max_{P \in \mathcal{B}_x} \max_{z \in \overline{P}} |\nabla u(z)|
$$
  

$$
\lesssim \max_{P \in \mathcal{B}_x} \max_{z \in \overline{P}} \delta_\Omega(z) |\nabla u(z)| \lesssim ||\delta_\Omega \nabla u||_{N_\mathcal{D}^\infty(\Omega)}.
$$

Consequently,

$$
||u||_{N_{\sharp}^{\infty}(\Omega)} \leq \sup_{x \in \Omega} \sup_{P \in \mathcal{B}_x} \sup_{y \in P} |u(y) - \int_{B_x} u|
$$
  

$$
\lesssim \sup_{x \in \Omega} \sup_{P \in \mathcal{B}_x} \sup_{y \in P} |u(y) - \int_P u| + ||\delta_{\Omega} \nabla u||_{N_{\mathcal{D}}^{\infty}(\Omega)}
$$
  

$$
\lesssim ||u||_{N_{\sharp,\mathcal{D}}^{\infty}(\Omega)} + ||\delta_{\Omega} \nabla u||_{N_{\mathcal{D}}^{\infty}(\Omega)}.
$$

Since for any  $x \in \Omega$  there exists a unique half-open cube  $P \in \mathcal{W}(\Omega)$  such that  $x \in P$ , we have that

$$
\delta_{\Omega}(x)|\nabla u(x)| \leq \sup_{z \in P} \delta_{\Omega}(z)|\nabla u(z)| \leq ||\delta_{\Omega}\nabla u||_{N_{\mathcal{D}}^{\infty}(\Omega)}
$$

and so  $\|\delta_\Omega\nabla u\|_{\mathrm{N}^\infty(\Omega)}\lesssim \|\delta_\Omega\nabla u\|_{\mathrm{N}^\infty_D(\Omega)}.$  Therefore,

(A.19) 
$$
||u||_{N_{sum}(\Omega)} = ||u||_{N_{\sharp}^{\infty}(\Omega)} + ||\delta_{\Omega} \nabla u||_{N^{\infty}(\Omega)}
$$

$$
\lesssim ||u||_{N_{\sharp, \mathcal{D}}(\Omega)} + ||\delta_{\Omega} \nabla u||_{N^{\infty}_{\mathcal{D}}(\Omega)} = ||u||_{N_{sum, \mathcal{D}}(\Omega)}.
$$

For the converse direction, let us fix  $c \in (0, 1/2]$  so that for every  $P \in \mathcal{W}(\Omega)$ , if  $x_P$  is the center of P, it holds that  $P \subset B(x_P, c \, \delta(x_P)) =: B_P$ . By the mean value theorem, for any  $P \in \mathcal{W}(\Omega)$ , we have that

$$
\left| \int_{B_P} u - \int_P u \right| \le \max_{z \in B_P} \delta_{\Omega}(z) |\nabla u(z)| \le ||\delta_{\Omega} \nabla u||_{N^{\infty}(\Omega)}
$$

and so

$$
\sup_{x \in P} |u(z) - \oint_P u| \le \sup_{x \in P} |u(z) - \oint_{B_P} u| + \|\delta_{\Omega} \nabla u\|_{N^{\infty}(\Omega)} \lesssim \|u\|_{N^{\infty}_{\sharp}(\Omega)} + \|\delta_{\Omega} \nabla u\|_{N^{\infty}(\Omega)}.
$$
\nThis implies that ||u||<sub>N</sub>  $\le$   $\le$   $\le$  ||u||<sub>N</sub>  $\le$   $\le$ 

This implies that  $||u||_{N_{\sharp,\mathcal{D}}(\Omega)} \gtrsim ||u||_{N_{\text{sum}}(\Omega)}$  and as

$$
\|\delta_{\Omega}\nabla u\|_{N_{\mathcal{D}}^{\infty}(\Omega)} \leq \sup_{P \in \mathcal{W}(\Omega)} \sup_{z \in B_P} \delta_{\Omega}(z) |\nabla u(z)| \leq \|\delta_{\Omega}\nabla u\|_{N^{\infty}(\Omega)},
$$

we infer that  $||u||_{N_{\text{sum},\mathcal{D}}(\Omega)} \lesssim ||u||_{N_{\text{sum}}(\Omega)}$  for any  $c \in (0, 1/2)$  such that  $P \subset B_P$ . It remains to prove that if  $0 < c_0 < c$ , then

<span id="page-68-1"></span>
$$
\sup_{x \in \Omega} \sup_{z \in B(x, c\delta_{\Omega}(x))} \left| u(z) - \int_{B(x, c\delta_{\Omega}(x))} \right| \lesssim \sup_{y \in \Omega} \sup_{z \in B(y, c_0\delta_{\Omega}(y))} \left| u(z) - \int_{B(y, c_0\delta_{\Omega}(y))} \right|
$$
\n(A.20)

and

<span id="page-68-2"></span>(A.21) 
$$
\sup_{y \in \Omega} \sup_{z \in B(y,c\delta_{\Omega}(y))} \delta_{\Omega}(z) |\nabla u(z)| \lesssim \sup_{y \in \Omega} \sup_{z \in B(y,c_0\delta_{\Omega}(y))} \delta_{\Omega}(z) |\nabla u(z)|,
$$

where the implicit constants are independent of  $c_0$  and  $c$ . For any  $x \in \Omega$ , there exists a uniformly bounded number of balls  ${B_j^x}_{j=1}^N$  so that  $B_j^x = B(x_j, c_0 \delta_{\Omega}(x_j))$ , where  $x_j \in$  $B(x, c\delta_{\Omega}(x))$  and  $B(x, c\delta_{\Omega}(x)) \subset \cup_{j=1}^{N} B_j^x$ . Therefore, by the mean value theorem, we have that for any  $x \in \Omega$ ,

$$
\left| \int_{B(x,c\delta_{\Omega}(x))} u - \int_{B_j} u \right| \lesssim \sup_{\substack{j \in \{1,\ldots,N\} \\ y \in \Omega}} \sup_{z \in B_{j}^{x}} \delta_{\Omega}(z) |\nabla u(z)|
$$
  
\$\leq \sup\_{y \in \Omega} \sup\_{z \in B(y,c\_0\delta\_{\Omega}(y))} \delta\_{\Omega}(z) |\nabla u(z)|\$.

Therefore,

$$
\sup_{x \in \Omega} \sup_{z \in B(x,c\delta_{\Omega}(x))} \left| u(z) - \int_{B(x,c\delta_{\Omega}(x))} u \right| \leq \sup_{x \in \Omega} \sup_{j \in \{1,\dots,N\}} \sup_{z \in B_j^x} \left| u(z) - \int_{B(x,c\delta_{\Omega}(x))} u \right|
$$
  
\n
$$
\leq \sup_{x \in \Omega} \sup_{j \in \{1,\dots,N\}} \left( \sup_{z \in B_j^x} \left| u(z) - \int_{B_j} u \right| + \left| \int_{B(x,c\delta_{\Omega}(x))} u - \int_{B_j} u \right| \right)
$$
  
\n
$$
\lesssim \sup_{y \in \Omega} \sup_{z \in B(y,c_0\delta_{\Omega}(y))} \left| u(z) - \int_{B(y,c_0\delta_{\Omega}(y))} u \right| + \sup_{y \in \Omega} \sup_{z \in B(y,c_0\delta_{\Omega}(y))} \delta_{\Omega}(z) |\nabla u(z)|,
$$

which shows  $(A.20)$ . By similar considerations, it is easy to prove  $(A.21)$ , concluding the proof of the lemma.  $\Box$ 

<span id="page-68-0"></span>**Corollary A.9.** *The space*  $(N_{sum}(\Omega), \| \cdot \|_{N_{sum}(\Omega)})$  *is sequentially complete.* 

*Proof.* The result readily follows from Lemmas [A.7](#page-61-1) and [A.8.](#page-67-0)

An immediate corollary of Lemmas [A.5](#page-61-0) and [A.7](#page-61-1) is the following.

**Corollary A.10.** *If*  $\Omega \subset \mathbb{R}^{n+1}$  *is an open set and*  $\Omega = \bigcup_{i=1}^{\infty} \Omega_i$ *, where*  $\{\Omega_i\}_{i=1}^{\infty}$  *are the connected components of*  $\Omega$ , then  $\bigoplus_{i=1}^{\infty} [N_{sum}(\Omega_i)/\mathbb{R}]$  *is a Banach space.* 

### <span id="page-69-0"></span>**REFERENCES**

<span id="page-69-22"></span><span id="page-69-21"></span><span id="page-69-20"></span><span id="page-69-19"></span><span id="page-69-18"></span><span id="page-69-17"></span><span id="page-69-16"></span><span id="page-69-15"></span><span id="page-69-14"></span><span id="page-69-13"></span><span id="page-69-12"></span><span id="page-69-11"></span><span id="page-69-10"></span><span id="page-69-9"></span><span id="page-69-8"></span><span id="page-69-7"></span><span id="page-69-6"></span><span id="page-69-5"></span><span id="page-69-4"></span><span id="page-69-3"></span><span id="page-69-2"></span><span id="page-69-1"></span>

VAROPOULOS EXTENSIONS IN DOMAINS WITH AHLFORS-REGULAR BOUNDARIES 71

<span id="page-70-14"></span><span id="page-70-12"></span><span id="page-70-11"></span><span id="page-70-10"></span><span id="page-70-6"></span><span id="page-70-5"></span><span id="page-70-4"></span><span id="page-70-3"></span><span id="page-70-0"></span>[HT21] S. Hofmann and O. Tapiola, *Uniform rectifiability implies Varopoulos extensions.* Adv. Math., 390: 107961, 2021. [HR18] T. Hytönen and A. Rosén, *Bounded variation approximation of*  $L_p$  *dyadic martingales and solutions to elliptic equations.* J. Eur. Math. Soc. (JEMS), 20(8): 1819–1850, 2018. [JK82] D. S. Jerison and C. E. Kenig, Boundary behavior of harmonic functions in nontangentially accessible domains. Adv. Math., 46(1): 80–147, 1982. [KKPT00] C. Kenig, H. Koch, J. Pipher, and T. Toro, *A new approach to absolute continuity of elliptic measure, with applications to non-symmetric equations.* Adv. Math., 153(2): 231–298, 2000. [MS79] R. A. Macias and C. Segovia, *Lipschitz functions on spaces of homogeneous type.* Adv. Math., 33(3):257–270, 1979. [MaPo21] S. Mayboroda and B. Poggi, *Carleson perturbations of elliptic operators on domains with low dimensional boundaries.* J. Funct. Anal. , 280(8): 108930, 2021. [MPT13] E. Milakis, J. Pipher, and T. Toro, *Harmonic analysis on chord arc domains.* J. Geom. Anal., 23(4): 2091–2157, 2013. [MiTa01] M. Mitrea and M. Taylor, *Potential theory on Lipschitz domains in Riemannian manifolds:*  $L^p$ Hardy, and Hölder space results. Commun. Anal. Geom., 9(2): 369-421, 2001. [MP24] M. Mourgoglou and B. Poggi, *Replacing rough external temperature data with internal heat flux sources: Approximation of the Dirichlet problem with data in* L p *by a sequence of Poisson problems.* In preparation. [MPT22] M. Mourgoglou, B. Poggi, and X. Tolsa,  $L^p$ -solvability of the Poisson-Dirichlet problem and its *applications to the regularity problem.* Preprint, July 2022, arXiv:2207.10554. [MT22] M. Mourgoglou and X. Tolsa, *The regularity problem for the Laplace equation in rough domains.* Duke Math. J., 173(9): 1731–1837, 2024. [Sar75] D. Sarason, *Functions of vanishing mean oscillation.* Trans. Amer. Math. Soc. , 207: 391–405, 1975. [ST70] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions.* Princeton University Press, 1970. [To14] X. Tolsa, *Analytic capacity, the Cauchy transform, and non-homogeneous Calderón-Zygmund theory, volume 307 of Progress in Mathematics. Birkhäuser Verlag, Basel, 2014.* [Var77] N. Th. Varopoulos, *BMO functions and the* ∂*-equation.* Pacific J. Math., 71(1): 221–273, 1977. [Var78] N. Th. Varopoulos, *A remark on functions of bounded mean oscillation and bounded harmonic functions. Addendum to: " BMO functions and the* ∂*-equation"* ( Pacific J. Math. 71 (1977), no.1, 221–273). Pacific J. Math., 74(1): 257–259, 1978.

<span id="page-70-15"></span><span id="page-70-13"></span><span id="page-70-9"></span><span id="page-70-8"></span><span id="page-70-7"></span><span id="page-70-2"></span><span id="page-70-1"></span>DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DEL PAÍS VASCO, BARRIO SARRIENA S/N 48940 LEIOA, SPAIN AND, IKERBASQUE, BASQUE FOUNDATION FOR SCIENCE, BILBAO, SPAIN. *Email address*: michail.mourgoglou@ehu.eus

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DEL PAÍS VASCO, BARRIO SARRIENA S/N 48940 LEIOA, SPAIN.

*Email address*: athanasios.zacharopoulos@ehu.eus