

VARIATIONS ON A RESULT OF GRANVILLE ABOUT POLYNOMIAL INVARIANCE

LABIB HADDAD AND CHARLES HELOU

ABSTRACT

In 2004, Andrew Granville proved the following: A polynomial in variables x, y, z , invariant under the map $\varphi : z \rightarrow -(x + y + z)$ is a polynomial in $b = z(x + y + z)$. We tell, curtly, its history and then give a variation on this theme.

HISTORY

In March 2004, the second author received an inquiry from a colleague about the factorization of some symmetric polynomials, with power monomials, in two or three variables. There was a statement about a factorization in $\mathbb{Z}[x, y]$ of the polynomial $F_p(x, y) = (x + y)^p - x^p - y^p$, for a prime number $p > 3$, namely

$$F_p(x, y) = pxy(x + y) (x^2 + xy + y^2)^e C_p(x, y),$$

where $e = 1$ or 2 according to whether $p \equiv -1$ or $1 \pmod{6}$, respectively, and $C_p(x, y) \in \mathbb{Z}[x, y]$ is a so-called Cauchy polynomial, described as a well-known elementary result. In fact, $F_p(x, y)$ is a homogeneous polynomial of degree p , which, upon division by y^p , gives a polynomial $K_p(t)$, of degree $p - 1$ in $t = \frac{x}{y}$, namely

$$\frac{F_p(x, y)}{y^p} = \left(\frac{x}{y} + 1\right)^p - \left(\frac{x}{y}\right)^p - 1 = (t + 1)^p - t^p - 1 = \sum_{k=1}^{p-1} \binom{p}{k} t^k = K_p(t).$$

Clearly, $K_p(0) = K_p(-1) = 0$ and $K_p'(0) = p$, $K_p'(-1) = -p \neq 0$, so 0 and 1 are simple roots of K_p , so that K_p is divisible by $t(t + 1)$ in $\mathbb{Z}[t]$, and also by p since the binomial coefficients $\binom{p}{k}$ are divisible by p for $1 \leq k \leq p - 1$. Moreover, letting $\zeta_3 = \exp\left(\frac{2i\pi}{3}\right)$ be a primitive cubic root of unity, we have

$$K_p(\zeta_3) = (\zeta_3 + 1)^p - \zeta_3^p - 1 = -(1 + \zeta_3^p + \zeta_3^{2p}) = 0$$

and

$$K_p'(\zeta_3) = p((\zeta_3 + 1)^{p-1} - \zeta_3^{p-1}) = p\zeta_3^{p-1}(\zeta_3^{p-1} - 1),$$

so that $K'_p(\zeta_3) = 0$ if and only if $p \equiv 1 \pmod{3}$, i.e., $p \equiv 1 \pmod{6}$. Thus ζ_3 is a root of $K_p(t)$, and it's a double root if and only if $p \equiv 1 \pmod{6}$, i.e., $K_p(t)$ is divisible by $t^2 + t + 1$ if $p \equiv -1 \pmod{6}$ and by $(t^2 + t + 1)^2$ if $p \equiv 1 \pmod{6}$, in $\mathbb{Z}[t]$. Thus

$$K_p(t) = pt(t+1)(t^2+t+1)^e C_p(t),$$

where

$$e = \begin{cases} 1, & \text{if } p \equiv -1 \pmod{6} \\ 2, & \text{if } p \equiv 1 \pmod{6}, \end{cases}$$

and $C_p(t) \in \mathbb{Z}[t]$, of degree $p - 3 - 2e$, is called a Cauchy polynomial, because it was introduced by A. L. Cauchy in 1839 and reconsidered in 1841 ([2, 3, 5]). The Cauchy polynomials were studied by D. Mirimanoff in 1903 ([4]), who conjectured their irreducibility over \mathbb{Q} , a still open conjecture. For more details, one can consult the 1989 paper of G. Terjanian ([6]). This factorization of $K_p(t)$ implies the factorization of $F_p(x, y)$ cited above. To be accurate, $C_p(t)$ should be $C_p(t, 1)$ in the factorization of $K_p(t)$, but by "abuse of notation", we identified the two.

This leads to introduce the polynomial

$$E_2 = \frac{(x+y)^p - x^p - y^p}{pxy(x+y)} = (x^2 + xy + y^2)^e C_p(x, y),$$

which lies in $Z[x, y]$, and, by analogy, in the case of 3 variables x, y, z , the polynomial

$$E_3 = \frac{(x+y+z)^p - x^p - y^p - z^p}{p(x+y)(y+z)(z+x)},$$

which lies in $Z[x, y, z]$. Indeed, letting

$$H_p(x, y, z) = (x+y+z)^p - x^p - y^p - z^p = \sum_{\substack{0 \leq j, k, l < p \\ j+k+l=p}} \frac{p!}{j!k!l!} x^j y^k z^l,$$

it is easily seen that all the coefficients of H_p are divisible by p , and that $H_p(x, -x, z) = H_p(x, y, -y) = H_p(x, y, -x) = 0$, so that H_p is divisible by $p, x+y, y+z$ and $z+x$, in $Z[x, y, z]$, which are pairwise coprime. Therefore H_p is divisible by their product $p(x+y)(y+z)(z+x)$, i.e., E_3 has its coefficients in \mathbb{Z} . Moreover, it is noted that

$$E_3 = E_2 + Q(x, y, z),$$

where $Q \in z\mathbb{Z}[x, y, z]$. Indeed,

$$\begin{aligned} E_3 - E_2 &= \frac{(x+y+z)^p - x^p - y^p - z^p}{p(x+y)(y+z)(z+x)} - \frac{(x+y)^p - x^p - y^p}{pxy(x+y)} = \\ &= \frac{xy((x+y+z)^p - x^p - y^p - z^p) - (y+z)(z+x)((x+y)^p - x^p - y^p)}{pxy(x+y)(y+z)(z+x)} = \\ &= \frac{xy \left(\left(\sum_{k=1}^{p-1} \binom{p}{k} (x+y)^k z^{p-k} + (x+y)^p - x^p - y^p \right) - (xy + xz + yz + z^2)((x+y)^p - x^p - y^p) \right)}{pxy(x+y)(y+z)(z+x)} = \\ &= \frac{xy \sum_{k=1}^{p-1} \binom{p}{k} (x+y)^k z^{p-k} - z(x+y+z)((x+y)^p - x^p - y^p)}{pxy(x+y)(y+z)(z+x)}, \end{aligned}$$

which is a polynomial in $\mathbb{Z}[x, y, z]$ all of whose terms are multiples of z .

Then, came the main question. Setting $b = z(x + y + z)$, my correspondent made the observation that

$E_3 = a_n + b(a_{n-1} + b(a_{n-2} + \dots + b(a_2 + b(a_1 + b)) \dots)) = b^n + a_1 b^{n-1} + \dots + a_{n-2} b^2 + a_{n-1} b + a_n$, where $n = \frac{p-3}{2}$ and all the $a_k \in \mathbb{Z}[x, y]$, with $a_n = E_2$. This amounts to stating that E_3 is a polynomial in b with coefficients in $\mathbb{Z}[x, y]$, and asking for a proof.

In April 2004, a second message came, extending the first one. In it, he makes the observation that $a_n + xy a_{n-1} = (x + y)^{p-3}$, and he conjectures that, for $0 \leq m \leq n - 1$, where $n = \frac{p-3}{2}$,

$$a_{n-m} + xy a_{n-m-1} = a_{n-m}^{(1)} (x + y)^{p-2m-3},$$

where $a_k^{(j)}$ is the coefficient of the j -th term in the polynomial a_k . Then it is claimed that a result of E. Catalan, cited in L. E. Dickson's *History of the Theory of Numbers, vol. II*, gives the coefficients of $C_p(x, y) = \sum_{k=0}^{p-3-2e} c_k x^{p-3-2e-k} y^k$, namely (with a correction)

$$c_k = \frac{1}{p} \left(\binom{p-1}{k} - (-1)^k \right).$$

He then claims to deduce that $a_{n-1}^{(1)} = a_n^{(2)}$ and $a_n^{(2)} = \frac{p-3}{2}$. All computations were done, using *Mathematica*.

In May 2004, a third message came saying that he sent his query to Andrew Granville, who replied that any polynomial $F(x, y, z)$ which is invariant under the map $\varphi : z \mapsto -(x + y + z)$ (as is E_3) can be written as a polynomial in $b = z(x + y + z)$, providing a simple proof by induction on the degree of F . Indeed, setting $a_0 = F(x, y, 0)$, Granville notes that a_0 and $F(x, y, z) - a_0$ are invariant under φ . But $F(x, y, z) - a_0$ is divisible by z , and, applying φ , it is also divisible by $-(x + y + z)$, and therefore it is divisible by b . Noting that b is also invariant under φ , he deduces that $G(x, y, z) = \frac{F(x, y, z) - a_0}{b}$ is invariant under ϕ , and has lower degree than F . Thus, by the induction hypothesis, G can be written as a polynomial in b , and therefore so can F .

It is this brilliantly simple argument of Granville that we attempt to generalize below.

Remarks. The polynomials denoted by E_2 and E_3 will henceforth be designated by $E_{2,p}$ and $E_{3,p}$ to emphasize their dependence on p . Note that

$$\begin{aligned} \frac{(x + y)^p - (x^p + y^p)}{x + y} &= (x + y)^{p-1} - \frac{x^p + y^p}{x + y} = \sum_{k=0}^{p-1} \binom{p-1}{k} x^k y^{p-1-k} - \sum_{k=0}^{p-1} (-1)^k x^k y^{p-1-k} \\ &= \sum_{k=0}^{p-1} \left(\binom{p-1}{k} - (-1)^k \right) x^k y^{p-1-k} = \sum_{k=1}^{p-2} \left(\binom{p-1}{k} - (-1)^k \right) x^k y^{p-1-k}, \end{aligned}$$

since $\binom{p-1}{k} - (-1)^k = 0$ for $k = 0$ and $k = p - 1$. Hence

$$\begin{aligned} E_{2,p}(x, y) &= \frac{(x + y)^p - (x^p + y^p)}{pxy(x + y)} = \frac{1}{pxy} \sum_{k=1}^{p-2} \left(\binom{p-1}{k} - (-1)^k \right) x^k y^{p-1-k} = \\ &= \sum_{k=1}^{p-2} \frac{\binom{p-1}{k} - (-1)^k}{p} x^{k-1} y^{p-k-2}, \end{aligned}$$

where the coefficients $\frac{\binom{p-1}{k} - (-1)^k}{p}$ are integers, for $1 \leq k \leq p-2$, since

$$\binom{p-1}{k} = \frac{(p-1)(p-2)\cdots(p-k)}{k!} \equiv \frac{(-1)^k k!}{k!} \equiv (-1)^k \pmod{p}.$$

On the other hand,

$$\begin{aligned} H_p(x, y, z) &= (x+y+z)^p - (x^p + y^p + z^p) = \sum_{m=0}^p \binom{p}{m} (x+y)^m z^{p-m} - (x^p + y^p + z^p) = \\ &= \sum_{m=1}^p \binom{p}{m} (x+y)^m z^{p-m} - (x^p + y^p), \end{aligned}$$

so that

$$\begin{aligned} E_{3,p}(x, y, z) &= \frac{H_p(x, y, z)}{p(x+y)(y+z)(z+x)} = \frac{1}{p(y+z)(z+x)} \left(\sum_{m=1}^p \binom{p}{m} (x+y)^{m-1} z^{p-m} - \frac{x^p + y^p}{x+y} \right) \\ &= \frac{1}{p(y+z)(z+x)} \left(\sum_{m=1}^p \binom{p}{m} (x+y)^{m-1} z^{p-m} - \sum_{k=0}^{p-1} (-1)^k x^k y^{p-1-k} \right) = \\ &= \frac{1}{p(y+z)(z+x)} \sum_{k=0}^{p-1} \left(\binom{p}{k+1} (x+y)^k z^{p-k-1} - (-1)^k x^k y^{p-k-1} \right), \end{aligned}$$

the last equality being obtained by setting $k = m-1$ in the first of the two sums.

Furthermore, multiplying and dividing by $(x+y)$, distributing in the numerator, and setting $l = k+1$, we get

$$\begin{aligned} E_{3,p}(x, y, z) &= \frac{1}{p(x+y)(y+z)(z+x)} \left(\sum_{l=1}^p \binom{p}{l} (x+y)^l z^{p-l} + \sum_{l=1}^p (-1)^l x^l y^{p-l} - \sum_{k=0}^{p-1} (-1)^k x^k y^{p-k} \right) \\ &= \frac{1}{p(x+y)(y+z)(z+x)} \left(\sum_{l=0}^p \binom{p}{l} (x+y)^l z^{p-l} - x^p - y^p - z^p \right). \end{aligned}$$

Expanding the binomial $(x+y)^l$, and replacing all binomial coefficients by their defining expressions, then reversing the order of summations, and factoring out the common factors, we get

$$E_{3,p}(x, y, z) = \frac{1}{p(x+y)(y+z)(z+x)} \left(p! z^p \sum_{k=0}^p \frac{1}{k!} \left(\frac{x}{y} \right)^k \sum_{l=k}^p \frac{1}{(p-l)!(l-k)!} \left(\frac{y}{z} \right)^l - x^p - y^p - z^p \right).$$

Hence the following two expressions for $E_{3,p}$, established above,

$$\begin{aligned} E_{3,p}(x, y, z) &= \frac{1}{p(y+z)(z+x)} \sum_{k=0}^{p-1} \left(\binom{p}{k+1} (x+y)^k z^{p-k-1} - (-1)^k x^k y^{p-k-1} \right) = \\ &= \frac{1}{p(x+y)(y+z)(z+x)} \left(p! z^p \sum_{k=0}^p \frac{1}{k!} \left(\frac{x}{y} \right)^k \sum_{l=k}^p \frac{1}{(p-l)!(l-k)!} \left(\frac{y}{z} \right)^l - x^p - y^p - z^p \right). \end{aligned}$$

Similarly, exchanging $x + y$ and $y + z$, then $x + y$ and $x + z$, gives, respectively,

$$\begin{aligned} E_{3,p}(x, y, z) &= \frac{1}{p(x+y)(x+z)} \sum_{k=0}^{p-1} \left(\binom{p}{k+1} (y+z)^k x^{p-k-1} - (-1)^k y^k z^{p-k-1} \right) = \\ &= \frac{1}{p(x+y)(y+z)(z+x)} \left(p! x^p \sum_{k=0}^p \frac{1}{k!} \left(\frac{y}{z} \right)^k \sum_{l=k}^p \frac{1}{(p-l)!(l-k)!} \left(\frac{z}{x} \right)^l - x^p - y^p - z^p \right), \end{aligned}$$

and

$$\begin{aligned} E_{3,p}(x, y, z) &= \frac{1}{p(x+y)(y+z)} \sum_{k=0}^{p-1} \left(\binom{p}{k+1} (x+z)^k y^{p-k-1} - (-1)^k x^k z^{p-k-1} \right) = \\ &= \frac{1}{p(x+y)(y+z)(z+x)} \left(p! y^p \sum_{k=0}^p \frac{1}{k!} \left(\frac{z}{x} \right)^k \sum_{l=k}^p \frac{1}{(p-l)!(l-k)!} \left(\frac{x}{y} \right)^l - x^p - y^p - z^p \right). \end{aligned}$$

Moreover, after a search for the result of E. Catalan mentioned above, we located it in an article ([1]), originally published in 1861, then taken up again, and complemented, in 1885, by Catalan, where he gives an explicit expression for the polynomials

$$E_{3,n}(x, y, z) = \frac{(x+y+z)^n - x^n - y^n - z^n}{(x+y)(y+z)(z+x)},$$

for odd integers $n > 3$. He defines the homogeneous polynomials of degree $m \geq 0$, with all coefficients equal to 1

$$H_m = H_m(x, y, z) = \sum_{j,k,l \in \mathbb{N}: j+k+l=m} x^j y^k z^l = \sum_{j=0}^m \sum_{k=0}^{m-j} x^j y^k z^{m-j-k},$$

with $H_0 = 1$, he sets $P = x + y + z$, and he establishes that, for odd $n > 3$,

$$E_{3,n}(x, y, z) = \sum_{m=0}^{n-3} H_m(x, y, z) P^{n-3-m} + 2H_{\frac{n-3}{2}}(x^2, y^2, z^2).$$

THE VARIATION

Let K be a commutative field. Take $n + 1$ variables $x = (x_1, \dots, x_n)$ and z , then consider the $K[x]$ -algebra of polynomials $A = K[x, z]$. Choose a polynomial, $p(x, z)$, in A and define the following transformation T of A into A : To each polynomial $F(x, z)$, it matches the polynomial

$$(1) \quad TF(x, z) = F(x, p(x, z)).$$

This transformation T is a $K[x]$ -homomorphism of the algebra A into itself. Let B be the set of polynomial invariant under T , i.e.,

$$(2) \quad B = \{F \in A : TF = F\}.$$

Of course, B is a $K[x]$ -subalgebra of A and it contains $K[x]$. Then set

$$(3) \quad C = B \setminus K[x].$$

How can the elements of B be characterized?

What is needed for C not to be empty?

The case $p(x, z) = z$ is trivial since $T = \text{id}_A$ and $B = A$.

We shall say that polynomial $p(x, z)$ is *à la Granville* when

$$(4) \quad p(x, z) = -z + r(x).$$

Notice that T is an involution (i.e., $T^2 = \text{id}_A$) in that case since we will have $Tp(x, z) = z$.

The case studied by Granville is that in which $p(x, z) = -(x + y + z)$.

Now, on to the “variation”!

1 Preparation. Introduce the sequence T^k of iterates of the transformation T , and the related sequence of polynomials $p_k(x, z) = T^k z$.

Of course, T^0 is id_A , and $T^1 = T$.

Similarly, $p_0(x, z) = z$ and $p_1(x, z)$ is $p(x, z)$ itself.

For short, we say that $p(x, z)$ is 1-adequate when $p(x, z) = z$.

Let $m > 1$ be an integer. Henceforth, we assume that K contains the m -th roots of unity.

We say that a polynomial $p(x, z)$ is m -adequate when it has the following form:

$$(*) \quad p(x, z) = qz + r(x),$$

where q is a primitive m -th root of unity, and $r(x)$ is in $K[x]$.

So, $p(x, z)$ is “à la Granville” if and only if it is 2-adequate.

When $p(x, z)$ is m -adequate for some $m \geq 1$, just say it is adequate.

2 Remark. When $p(x, z)$ is m -adequate, we have $p_m(x, z) = z$ and therefore $T^m = \text{id}_A$. More precisely, we have

$$\begin{aligned}
 p_0(x, z) &= z \\
 p_1(x, z) &= qz + r(x) \\
 p_2(x, z) &= q^2z + (q + 1).r(x) \\
 &\dots\dots\dots \\
 p_k(x, z) &= q^kz + (q^{k-1} + \dots + q + 1).r(x) \\
 &\dots\dots\dots \\
 p_m(x, z) &= z.
 \end{aligned}$$

When $r(x) \neq 0$, polynomials p_0, p_1, \dots, p_{m-1} are two by two mutually prime.

Let $d(P)$ be the degree in z of each polynomial $P(x, z)$.]

3 Theorem. Let $p(x, z)$ be m -adequat. Let b be the product of polynomials $\{p_k : 0 \leq k < m\}$. A polynomial $F(x, z)$ is in B if and only if it has the following form:

$$(6) \quad F(x, z) = G(b), \text{ with } G(y) \in K[x][y].$$

Proof. Observe that polynomial b is in B , so $TG(b) = G(b)$, and every polynomial which has the form (6) is then in B .

For the converse, distinguish two cases.

If $r(x) = 0$, then F is in B means $F(x, z) = F(x, qz)$; since $q^m = 1$, we conclude that F is a polynomial in z^m .

Otherwise, proceed by induction on the degree, $d(F)$, of F in z .

The case $d(F) = 0$ is trivial.

In four steps.

Lemma 1. If $F(x, z)$ is in B , each of the $p_k(x, z)$, $k \geq 0$, divides the polynomial $W(x, z) := F(x, z) - F(x, 0)$.

Indeed, $W(x, z)$ also is in B and is divisible by z . Therefore, we must have $W(x, z) = z.U(x, z)$, and for $k \geq 1$, we have

$$W(x, z) = T^k U(x, z) = p_k(x, z).T^k U(x, z).$$

If $F(x, z)$ is in B , polynomial $F(x, z) - F(x, 0)$ is divisible by b , i.e., we have $F(x, z) - F(x, 0) = b(x, z).V(x, z)$. So, $V(x, z)$ is in B , moreover we have $d(V) < d(F)$. Whence the conclusion. \square

Complement. The following three conditions are equivalent.

- (i) The set C is not empty.
- (ii) There exists an integer $m \geq 1$ such that $T^m = \text{id}_A$.
- (iii) Polynomial $p(x, z)$ is adequate.

Proof. If $p(x, z)$ is m -adequate, we already know that C is not empty, and that $T^m = \text{id}_A$. \square

Lemma 2. If $d(p) \neq 1$, then $B = K[x]$.

Indeed, $d(TF) = d(F).d(p)$. If F is in B , i.e., $TF = F$, we have $d(F) = d(F).d(p)$, and if, moreover, $d(p) \neq 1$, then $d(F) = 0$. \square

Lemma 3. If C is not empty, the polynomial $p(x, z)$ is adequate.

Proof. By the previous lemma, we know that $p(x, z)$ has the following form:

$$(7) \quad p(x, z) = q(x).z + r(x), \text{ with } q(x) \neq 0.$$

Take $F \in B$ such that $d(F) = m \geq 1$.

Ordering F in decreasing powers of z , write

$$F = u.z^m + v.z^{m-1} + \dots$$

whence

$$\begin{aligned} TF &= u.(qz + r)^m + v.(qz + r)^{m-1} + \dots \\ &= u.q^m z^m + (u.m.q^{m-1}.r + v.q^{m-1}z^{m-1} + \dots \end{aligned}$$

then, identifying the first two coefficients, obtain both equalities

$$(8) \quad q^m = 1$$

$$(9) \quad q^{m-1}.(m.r + v) = v.$$

For $q = 1$, we therefore have $r = 0$. \square

Lemma 4. If $T^m = \text{id}_A$ for some $m \geq 1$, the product b of polynomials in $\{p_k : 0 \leq k < m\}$ is invariant by T , and $d(b) \geq 1$.

This ends the proof

$\square \square$

REFERENCES

1. E. Catalan, *Sur le dernier théorème de Fermat (1861)*, Mém. Soc. Roy. Sci. Liège (2) **12** (1885), 179-187.
2. A. L. Cauchy, *Rapport sur un Mémoire de M. Lamé, relatif au dernier théorème de Fermat*, C. R. Acad. Sci. Paris **9** (1839), 359-363.
3. A. L. Cauchy, *Note sur Quelques Théorèmes d'Algèbre*, Exerc. d'Analyse et de Phys. Math., vol. 2, Bachelier, Paris, 1841, pp. 137-144.
4. D. Mirimanoff, *Sur l'équation $(x + 1)^l - x^l - 1 = 0$* , Nouv. Ann. Math. **3** (1903), 385-397.
5. P. Ribenboim, *13 Lectures on Fermat's Last Theorem*, Springer, 1979.
6. G. Terjanian, *Sur la loi de réciprocité des puissances l -èmes*, Acta Arith. **54** (1989), 87-125.

120 RUE DE CHARONNE, 75011 PARIS, FRANCE; EMAIL: LABIB.HADDAD@WANADOO.FR

PENN STATE UNIV., 25 YEARSLEY MILL RD, MEDIA, PA 19063, USA; E-MAIL: CXH22@PSU.EDU