Convergence rates for sums-of-squares hierarchies with correlative sparsity

Milan Korda

Victor Magron

Rodolfo Ríos Zertuche

March 28, 2023

Abstract

This work derives upper bounds on the convergence rate of the moment-sum-of-squares hierarchy with correlative sparsity for global minimization of polynomials on compact basic semialgebraic sets. The main conclusion is that both sparse hierarchies based on the Schmüdgen and Putinar Positivstellensätze enjoy a polynomial rate of convergence that depends on the size of the largest clique in the sparsity graph but not on the ambient dimension. Interestingly, the sparse bounds outperform the best currently available bounds for the dense hierarchy when the maximum clique size is sufficiently small compared to the ambient dimension and the performance is measured by the running time of an interior point method required to obtain a bound on the global minimum of a given accuracy.

Contents

| 1 | Introduction | | | | |
|----------|-----------------------------|---|-----------|--|--|
| | 1.1 Notations | | | | |
| | 1.2 | Results | 4 | | |
| | | 1.2.1 Sparse Schmüdgen-type representation on $[-1,1]^n$ | 4 | | |
| | | 1.2.2 Sparse Putinar-type representation on arbitrary domains | 5 | | |
| 2 | The sparse Jackson kernel | | | | |
| 3 | Sparse approximation theory | | | | |
| 4 | Pro | fs | 14 | | |
| | 4.1 | Proof of Theorem 2 | 14 | | |
| | 4.2 | Proof of Theorem 4 | 16 | | |
| | 4.3 | An asymptotic lemma | 22 | | |

1 Introduction

This work provides rates of convergence for the sums-of-squares hierarchy with correlative sparsity. For a positive $n \in \mathbb{N}$, consider the polynomial optimization problem

$$f_{\min} \coloneqq \min_{x \in S(\mathbf{g})} f(x)$$

where f is an element of the ring $\mathbb{R}[x]$ of polynomials in $x = (x_1, \ldots, x_n)$, and $S(\mathbf{g})$ is a basic compact semialgebraic set determined by a finite collection of polynomials $\mathbf{g} = \{g_1, \ldots, g_{\bar{k}}\}$ by $S(\mathbf{g}) = \{x \in \mathbb{R}^n : g_i(x) \ge 0, i = 1, \ldots, \bar{k}\}$. An approach to attack this problem, first proposed by Lasserre [8] and Parrilo [18], is as follows: Imagine we knew that $f(x) - \lambda$ could be written as

$$f(x) - \lambda = \sum_{j=0}^{\bar{k}} \sigma_j g_j(x) \quad \text{or} \quad f(x) - \lambda = \sum_{J \subseteq \{1, \dots, \bar{k}\}} \sigma_J \prod_{j \in J} g_j(x),$$

with $g_0(x) = 1$ and σ_j and σ_J being sum-of-squares (SOS) polynomials. Then the right-hand sides of each of these equations would be clearly nonnegative on $S(\mathbf{g})$, so we would know that $f_{\min} \geq \lambda$. By bounding the degree of the SOS polynomials, we obtain the following two hierarchies of lower bounds:

$$\begin{aligned} \mathrm{lb}_q(f,r) &= \max\{\lambda \in \mathbb{R} : f - \lambda = \sum_{j=0}^k \sigma_j g_j, \\ \mathrm{deg}(\sigma_j g_j) &\leq 2r, \ \sigma_j \in \Sigma[x]\}, \\ \mathrm{lb}_p(f,r) &= \max\{\lambda \in \mathbb{R} : f - \lambda = \sum_{J \subseteq \{1, \dots, \bar{k}\}} \sigma_J \prod_{j \in J} g_j, \\ \mathrm{deg}\left(\sigma_J \prod_{j \in J} g_j\right) &\leq 2r, \ \sigma_J \in \Sigma[x]\}, \end{aligned}$$

where $\Sigma[x]$ is the convex cone of all sum-of-squares polynomials. These satisfy $lb_q(f,r) \leq lb_p(f,r) \leq f_{\min}$. The lower bound $lb_q(f,r)$ is associated to a so-called quadratic module certificate, while $lb_p(f,r)$ corresponds to a preordering certificate; this terminology is justified by the definitions in Section 1.2. The well-known Putinar and Schmüdgen Positivstellensätze [19, 21], respectively, guarantee that these bounds converge to f_{\min} as $r \to +\infty$, the former with the additional assumption that the associated quadratic module be Archimedian¹. Here we will prove sparse quantitative versions of these results.

Polynomial optimization schemes have generated substantial interest due to their abundant fields of application; see for example [11, 10]. The first proof of convergence, without a convergence rate, was given by Lasserre [8] using the Archimedian positivstellensatz due to Putinar [19]. Eventually, rates of convergence were obtained; initially in [16] these were logarithmic in the degree of the polynomials involved, and later on they were improved [3, 12, 23, 1] (using ideas of [20, 2, 17]) to polynomial rates; refer to Table 1. The crux of the argument used to obtain those rates is a bound of the deformation incurred by a polynomial strictly-positive on the domain of interest, as it passes through an integral operator that closely approximates the identity and is associated to a strictly-positive polynomial kernel that is itself composed of sums of squares and similar to the Christoffel-Darboux and Jackson kernels (see Definition 9).

The techniques used to obtain these results generally involve linear operators on the space of polynomials (mostly Christoffel-Darboux kernel operators; see [23]) that are close to the identity and that, for positive polynomials, are easily (usually, by construction) proved to output polynomials that are sums of squares and/or of their products with the functions in \mathbf{g} . All of these results deal, however, with the dense case.

| domain $S(\mathbf{g})$ | error | certificate | ref. |
|------------------------|------------------|---|------|
| Archimedean | $O(1/\log(r)^c)$ | quadratic module | [16] |
| Archimedean | $O(1/r^c)$ | quadratic module | [1] |
| general | $O(1/r^c)$ | preordering | [22] |
| general | $O(1/r^c)$ | quadratic module & uniform denominators | [14] |
| S^{n-1} | $O(1/r^2)$ | quadratic module / preordering | [3] |
| $\{0,1\}^n$ | see $[24]$ | quadratic module / preordering | [24] |
| B^n | $O(1/r^2)$ | quadratic module / preordering | [23] |
| $[-1,1]^n$ | $O(1/r^2)$ | preordering | [12] |
| Δ^n | O(1/r) | preordering | [7] |
| Δ^n | $O(1/r^2)$ | preordering | [23] |

Table 1: Known results on the asymptotic error of Lasserre's hierarchies of lower bounds; based in part on [23, Table 1]. The domain $S(\mathbf{g})$ is assumed to be compact in all cases, $\epsilon \in [0, 1/2), c > 0, S^{n-1}$ is the unit sphere, B^n is the unit ball, Δ^n is the standard simplex.

In this work, we treat the case where the problem possesses the so-called correlative sparsity, where each function g_i depends only on a certain subset of variables and the function f decomposes as a sum of functions depending only on these subsets of variables. This structure can be exploited in order to define sparse lower bounds that are cheaper to compute but possibly weaker. Nevertheless, these sparse lower bounds allow one to tackle large-scale polynomial optimization problems arising from various applications including roundoff error bounds in computer arithmetic, quantum correlations and robustness certification of deep networks; see the recent survey [13]. In [9] Lasserre proved that these sparse lower bounds converge as the degree of the SOS multipliers tends to infinity provided the variable groups satisfy the so-called running intersection property (RIP). A shorter and more direct proof was

¹This means that there are R > 0 and $\sigma_j \in \Sigma[x]$ such that $R - ||x||^2 = \sum_j \sigma_j g_j(x)$.

provided in [4], and was adapted in [15] to obtain a sparse variant of Reznick's Positivstellensatz. In this work, we show polynomial rates of convergence for sparse hierarchies based on both Schmüdgen and Putinar Positivstellensätze. Importantly, we obtain rates that depend only on the size of the largest clique in the sparsity graph rather than the overall ambient dimension. This allows the perhaps surprising conclusion that, asymptotically, the sparse hierarchy is more accurate than the dense hierarchy for a given computation time of an optimization method, provided that the size of the largest clique is no more than the square root of the ambient dimension. This assumes that the running time of the optimization method is governed by the size of the largest PSD block and the number of such blocks in the semidefinite programming reformulations of the dense and sparse SOS problems which is the case for the interior point method as well as the most commonly used first-order methods.

To the best of our knowledge, these are the first quantitative results of this kind. Our proof techniques rely on an adaption of [4] and utilize heavily the recent results from [12] and [1], and can thus be seen as a generalization of these works to the sparse setting.

The results will be detailed below in Section 1.2 and further discussed in Section 1.2.1, after a brief interlude to establish some notations in Section 1.1. Some machinery will be developed in Sections 2 and 3, regarding variants of the Jackson kernel and some approximation theory, respectively, and the proofs of the main theorems are presented in Section 4.

1.1 Notations

Denote by \mathbb{R} the set of real numbers, by \mathbb{N} the set of positive integers, and by $\mathbb{N}_0 = \{0, 1, ...\}$ the set of nonnegative integers. Denote by $e_1, ..., e_n$ the vectors of the standard basis of Euclidean space \mathbb{R}^n .

For a Lipschitz continuous function $f: [-1,1]^n \to \mathbb{R}$, we set

Lip
$$f = \max\left(1, \sup_{x,y \in [-1,1]^n} \frac{|f(x) - f(y)|}{\|x - y\|}\right).$$

We take this to be at least 1 to simplify estimates below.

A multi-index $I = (i_1, \ldots, i_n) \in \mathbb{N}_0^n$ is an *n*-tuple of nonnegative integers i_k , and its weight is denoted by

$$|I| = \sum_{k=1}^{n} i_k.$$

For a multi-index $I = (i_1, \ldots, i_n) \in \mathbb{N}_0^n$ and $J \subset \{1, \ldots, n\}$, we will write $I \subseteq J$ to indicate that for all $1 \leq k \leq n$ if $i_k > 0$ then $k \in J$. Similarly, given a multi-index $I \in \mathbb{N}_0^n$ and a subset $J \subseteq \mathbb{N}_0$, we let I_J be the multi-index whose k-th entry is either i_k if $k \in J$ or 0 if $k \notin J$. For two multi-indices I and I', we will write $I \leq I'$ if the entrywise inequalities $i_k \leq i'_k$ hold for all $1 \leq k \leq n$. We will distinguish two special multi-indices:

$$\mathbf{1} = (1, 1, \dots, 1)$$
 and $\mathbf{2} = (2, 2, \dots, 2).$

We will denote $x^{I} = x_{1}^{i_{1}} x_{2}^{i_{2}} \dots x_{n}^{i_{n}}$. Also, we denote the Hamming weight of $I \in \mathbb{N}_{0}^{n}$ by

$$w(I) = \#\{k : i_k > 0, \ 1 \le k \le n\}.$$

In other words, w(I) is the number of nonzero entries in I.

We will denote the space of polynomials in n variables by $\mathbb{R}[x]$, and within this set we will distinguish the subspace $\mathbb{R}[x]_d$ of polynomials of total degree at most d. We will denote, for a polynomial $p(x) = \sum_I c_I x^I$, by $\overline{\deg p}$ the vector whose *i*-th entry is the degree of p in x_i ,

$$\overline{\deg} p = \left(\max_{c_I \neq 0} i_1, \max_{c_I \neq 0} i_2, \dots, \max_{c_I \neq 0} i_n\right).$$

Observe that deg $p \leq \left| \overline{\deg} p \right| = \sum_{k=1}^{n} \max_{c_I \neq 0} i_k$. Set also

$$\mathcal{I}_p = \{ I \in \mathbb{N}_0^n : c_I \neq 0 \}.$$

Given a subset $J \subset \{1, \ldots, n\}$, we let $\mathbb{R}[x_J]$ denote the set of polynomials in the variables $\{x_j\}_{j \in J}$. For a multi-index $\mathbf{r} = (r_1, \ldots, r_n) \in \mathbb{N}_0^n$, we let $\mathbb{R}[x]_{\mathbf{r}}$ denote the set of polynomials p such that, if $p(x) = \sum_I c_I x^I$ for some real numbers $c_I \in \mathbb{R}$, then for each $I = (i_1, \ldots, i_n)$ with $c_I \neq 0$ we also have $i_k \leq r_k$ for $1 \leq k \leq n$. Finally, we let $\mathbb{R}[x_J]_{\mathbf{r}} = \mathbb{R}[x_J] \cap \mathbb{R}[x]_{\mathbf{r}}$; in other words, $\mathbb{R}[x_J]_{\mathbf{r}}$ is the set of polynomials p with deg $p \leq \mathbf{r}$ in the variables $\{x_j : j \in J\} \subseteq \{x_1, \ldots, x_n\}$. Given a set X, we will write X^n to denote the product

$$X^n = \underbrace{X \times X \times \cdots \times X}_n.$$

We will denote by $\|\cdot\|_{\infty}$ the supremum norm on $[-1,1]^n$. The notation $\lceil s \rceil$ stands for the least integer $\geq s$.

1.2 Results

Let $\Sigma[x_J]$ denote the set of polynomials p that are sums of squares of polynomials in $\mathbb{R}[x_J]$, that is, of the form $p = p_1^2 + \cdots + p_\ell^2$ for $p_1, \ldots, p_\ell \in \mathbb{R}[x_J]$.

Let $\bar{k} \in \mathbb{N}$ and let $\mathbf{g} = \{g_1, \ldots, g_{\bar{k}}\}$ be a collection of polynomials $g_i \in \mathbb{R}[x]$ defining a set

$$S(\mathbf{g}) = \{ x \in \mathbb{R}^n : g_i(x) \ge 0, \ i = 1, \dots, \bar{k} \}.$$

For convenience, denote also $g_0 = 1$. To the collection **g**, a multi-index **r**, we associate the *(variable- and degree-wise truncated) quadratic module* associated to the collection **g** and a multi-index **r** be

$$\mathcal{Q}_{\mathbf{r},J}(\mathbf{g}) = \{\sum_{i=0}^{\bar{k}} \sigma_i g_i : \sigma_i \in \Sigma[x_J], \overline{\deg}(\sigma_i g_i) \le \mathbf{r}\}.$$

Similarly, we have a (variable- and degree-wise truncated) preordering

$$\mathcal{P}_{\mathbf{r},J}(\mathbf{g}) = \mathcal{Q}_{\mathbf{r},J}(\{g_K : K \subseteq \{1,\dots,\bar{k}\}\}) = \{\sum_{K \subseteq \{1,\dots,\bar{k}\}} \sigma_K g_K : \sigma_K \in \Sigma[x_J], \ \overline{\operatorname{deg}}(\sigma_K g_K) \le \mathbf{r}\}$$

where

$$g_K = \prod_{i \in K} g_i.$$

Definition 1. A collection $\{J_1, \ldots, J_\ell\}$ of subsets of $\{1, \ldots, n\} \supset J_j$ satisfies the running intersection property if for all $1 \le k \le \ell - 1$ we have

$$J_{k+1} \cap \bigcup_{j=1}^{k} J_j \subset J_s \text{ for some } s \leq k.$$

Denote, for $j = 2, 3, \ldots, \ell$,

$$\mathcal{J}_j = J_j \cap \bigcup_{k < j} J_k. \tag{1}$$

1.2.1 Sparse Schmüdgen-type representation on $[-1,1]^n$

Let $\overline{L} := \sum_{k=1}^{\ell} \operatorname{Lip} p_k$, $M := \max_{\substack{1 \le k \le \ell \\ 1 \le m \le n}} (\overline{\deg} \, p_k)_m$, and $\overline{J} := \max_{1 \le k \le \ell} |J_k|$.

Theorem 2. Let n > 0 and $\ell \ge 2$, and let $\mathbf{r}_1, \mathbf{r}_2, \ldots, \mathbf{r}_\ell \in \mathbb{N}^n$, $\mathbf{r}_j = (r_{j,1}, \ldots, r_{j,n})$, be nowhere-vanishing multi-indices. Let also J_1, \ldots, J_ℓ be subsets of $\{1, \ldots, n\}$ satisfying the running intersection property. Let $p = p_1 + p_2 + \cdots + p_\ell$ be a polynomial that is the sum of finitely many polynomials $p_j \in \mathbb{R}[x_{J_j}]_{\mathbf{r}_j}$. Then if $p \ge \varepsilon$ on $[-1, 1]^n$, we have

$$p \in \mathcal{P}_{\mathbf{r}_1, J_1}(\{1 - x_i^2\}_{i \in J_1}) + \dots + \mathcal{P}_{\mathbf{r}_\ell, J_\ell}(\{1 - x_i^2\}_{i \in J_\ell})$$

as long as, for all $1 \leq j \leq \ell$ and all $1 \leq i \leq n$,

$$r_j^2 \ge \frac{2^{\overline{J}+3}(\ell+2)n\pi^2 \|p\|_{\infty}}{\varepsilon} \left(\max\left(M, 4C_{\operatorname{Jac}}(\ell+2)\frac{\overline{J}\overline{L}}{\varepsilon}\right) + 2 \right)^{J+2}.$$

For small enough $0 < \varepsilon < 4C_{\text{Jac}}(\ell+2)\overline{J}\overline{L}/M$, this boils down to

$$r_j \ge \sqrt{\frac{A \|p\|_{\infty}}{\varepsilon^{\overline{J}+3}}} = O(\varepsilon^{-\frac{\overline{J}+3}{2}}),$$

with

$$A = n\pi^2 (4C_{\text{Jac}}\overline{J}\,\overline{L} + 2)^{\overline{J}+2} (2(\ell+2))^{\overline{J}+3}$$

The proof will be presented in Section 4.1.

Discussion. Solving the dense problem considered by [12] using the sum-of-squares hierarchy reduces to a semidefinite program with the largest PSD block of size $\binom{n+r}{r}$ that typical optimization methods (e.g., interior point or first order) can solve in an amount of time proportional to a power of

$$\binom{n+r}{r} \approx \binom{n+C\varepsilon^{-1/2}}{C\varepsilon^{-1/2}} =: B_{\text{dense}}(\varepsilon).$$

The bounds we find in Theorem 2 —in the case in which J_j is the largest of the sets J_1, \ldots, J_ℓ give a bound for the complexity of the leading term as (the same power of)

$$\ell\binom{|J_j| + |\mathbf{r}_j|}{|\mathbf{r}_j|} \leq \ell\binom{|J_j|(1 + C'\varepsilon^{-\frac{|J_j| + 3}{2}})}{|J_j|C'\varepsilon^{-\frac{|J_j| + 3}{2}}} =: B_{\text{sparseSchm}}(\varepsilon).$$

The reason we have $|\mathbf{r}_j| \leq |J_j| C' \varepsilon^{-|J_j|-3}$ is that $r_{j,i} \leq O(\varepsilon^{-\frac{|J_j|+3}{2}})$ and there are at most $|J_j|$ values of i with $r_{j,i} \neq 0$.

Proposition 3. If $n > |J_j|(|J_j| + 3)$ for all $j = 1, \ldots, \ell$, then we have

$$\lim_{\varepsilon \searrow 0} \frac{B_{\text{sparseSchm}}(\varepsilon)}{B_{\text{dense}}(\varepsilon)} = 0.$$

Thus, if the size of the largest clique is of the order of square root of the ambient dimension n or smaller, the sparse bound outperforms the best available dense bound if the performance is measured by the amount of time required by an optimization method to find a bound of a given accuracy ε .

Proof of Proposition 3. By Lemma 16 we have, as $\varepsilon \searrow 0$,

$$\frac{B_{\text{sparseSchm}}(\varepsilon)}{B_{\text{dense}}(\varepsilon)} = \frac{\binom{|J_j|(1+C'\varepsilon^{-\frac{|J_j|+3}{2}})}{|J_j|C'\varepsilon^{-\frac{|J_j|+3}{2}}}}{\binom{n+C\varepsilon^{-1/2}}{C\varepsilon^{-1/2}}} = O(\varepsilon^{\frac{1}{2}(n-|J_j|(|J_j|+3))}),$$

and this tends to 0 if the sparsity of the polynomial p is such that $n > |J_j|(|J_j| + 3)$.

1.2.2 Sparse Putinar-type representation on arbitrary domains

Let n > 0, k > 0, $\ell \ge 2$, $J_1, \ldots, J_\ell \subset \{1, \ldots, n\}$, $\mathbf{r}_1, \ldots, \mathbf{r}_\ell \in \mathbb{N}^n$, and $p = p_1 + \cdots + p_\ell$ with $p_j \in \mathbb{R}[x_{J_j}]_{\mathbf{r}_j}$. Assume that the sets J_1, \ldots, J_ℓ satisfy the running intersection property (Definition 1). Let $K_1, \ldots, K_\ell \subset \{1, \ldots, \bar{k}\}$ and let $\mathbf{g} = \{g_1, \ldots, g_{\bar{k}}\} \subset \mathbb{R}[x]$ be a collection of \bar{k} polynomials such that, if $i \in K_j$ for some $1 \le j \le \ell$, then $g_i \in \mathbb{R}[x_{J_j}]$. Let

$$S(\mathbf{g}) = \{ x \in \mathbb{R}^n : g_i(x) \ge 0 \text{ for all } i = 1, \dots, k \}.$$

Denote

$$\mathbf{g}_{K_i} = \{g_i : i \in K_j\}.$$

Let $c_1, \ldots, c_\ell \geq 1$ and $L_1, \ldots, L_\ell \geq 1$ be constants such that

$$dist(x, S(\mathbf{g}_{K_j}))^{\mathsf{L}_j} \le -\mathsf{c}_j \min\{\{0\} \cup \{g_i(x) : i \in K_j\}\}, \quad x \in [-1, 1]^n;$$

this is a version of the *Lojasiewicz inequality*, and its validity (with appropriate constants c_j, L_j) for semialgebraic functions is justified in [1, Thm. 2.3] and the papers cited therein.

Theorem 4. Assume that

$$||g_j||_{\infty} \le \frac{1}{2}, \quad j = 1, 2, \dots, \bar{k}.$$
 (2)

Assume that $S(\mathbf{g}) \subset [-1,1]^n$ and that there exist polynomials $s_{j,i} \in \mathbb{R}[x_{J_j}]_{\mathbf{r}_j}$, $j = 1, \ldots, \ell$, $i \in \{0\} \cup K_j$, such that the Archimedean conditions

$$1 - \sum_{i \in J_j} x_i^2 = s_{j,0}(x)^2 + \sum_{i \in K_j} s_{j,i}(x)^2 g_i(x), \quad j = 1, \dots, \ell,$$
(3)

hold; that is to say, we assume that $1 - \sum_{i \in J_j} x_i^2 \in \mathcal{Q}_{\mathbf{r}_j, J_j}(\mathbf{g}_{K_j})$.

Then there are constants $\mathbf{C}_j > 0$, depending only on \mathbf{g} , J_1, \ldots, J_ℓ , such that, if $p \ge \varepsilon > 0$ on $S(\mathbf{g})$, we have

$$p \in \mathcal{Q}_{\mathbf{r}_1+\mathbf{2},J_j}(\mathbf{g}_{K_j}) + \dots + \mathcal{Q}_{\mathbf{r}_\ell+\mathbf{2},J_\ell}(\mathbf{g}_{K_\ell})$$

as long as, for all $1 \leq j \leq \ell$ and $1 \leq k \leq n$,

$$(r_{j,k}+2)^{2} \geq \mathbf{C}_{j} \frac{4(\ell+2)\left(\sum_{i} \|p_{i}\|_{\infty}\right)^{\mathsf{L}_{j}+1} \left(\deg p_{j} \sum_{i=1}^{\ell} \operatorname{Lip} p_{i}\right)^{(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{\mathsf{SL}_{j}}{3})}}{\varepsilon^{1+\mathsf{L}_{j}+\frac{4\mathsf{L}_{j}+1}{3}(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{\mathsf{SL}_{j}}{3})}},$$
(4)

$$(r_{j,k}+2)^{2} \ge \mathbf{C}_{j} \left(\frac{\left(\sum_{i=1}^{\ell} \|p_{i}\|_{\infty} \right)^{\frac{4L_{j}+1}{3}} (\deg p_{j} \sum_{i=1}^{\ell} \operatorname{Lip} p_{i})^{\frac{8L_{j}}{3}}}{\varepsilon^{\frac{12L_{j}+1}{3}}} \right)^{2}$$
(5)

The proof of the theorem can be found in Section 4.2.

Discussion. By the same arguments we used in the discussion at the end of the previous section, if we assume $L_1 = \cdots = L_{\bar{k}} = 1$, the bounds we find in Theorem 4 give a bound for the complexity of the leading term as (a power of)

$$\ell \binom{|J_j| + |\mathbf{r}_j|}{|\mathbf{r}_j|} \approx \binom{|J_j|(1 + C''\varepsilon^{-\frac{26}{3} - \frac{5}{3}|J_j|})}{|J_j|C''\varepsilon^{-\frac{26}{3} - \frac{5}{3}|J_j|}}) =: B_{\text{sparsePut}}$$

The assumption $\mathsf{L}_1 = \cdots = \mathsf{L}_{\bar{k}} = 1$ is realized for example when the so-called *constraint qualification* condition that, at each point $x \in S(\mathbf{g})$ all the active constraints g_{i_1}, \ldots, g_{i_l} (i.e., those satisfying $g_{i_j}(x) =$ 0) have linearly independent gradients $\nabla g_{i_1}(x), \ldots, \nabla g_{i_l}(x)$), holds; this latter statement is proved in [1, Thm 2.11].

In this case, we have:

Proposition 5. If $\frac{n}{2} > |J_j|(\frac{26}{3} + \frac{5}{3}|J_j|)$ for all $j = 1, \ldots, \ell$ and if $L_1 = \cdots = L_{\bar{k}} = 1$, then we have

$$\lim_{\varepsilon \searrow 0} \frac{B_{\text{sparsePut}}}{B_{\text{dense}}} = 0.$$

Again the implication is that the sparse bound asymptotically outperforms the dense bound provided that the largest clique is sufficiently small.

Proof of Proposition 5. Lemma 16 gives

$$\frac{B_{\text{sparsePut}}}{B_{\text{dense}}} = \frac{\binom{|J_j|(1+C'\varepsilon^{-\frac{26}{3}-\frac{5}{3}|J_j|})}{|J_j|C'\varepsilon^{-\frac{26}{3}-\frac{5}{3}|J_j|}}}{\binom{n+C\varepsilon^{-1/2}}{C\varepsilon^{-1/2}}} = O(\varepsilon^{\frac{n}{2}-|J_j|(\frac{26}{3}+\frac{5}{3}|J_j|)}),$$

which tends to 0 if $\frac{n}{2} > |J_j|(\frac{26}{3} + \frac{5}{3}|J_j|).$

Organization of the paper. The proof of Theorem 2 can be seen as a variable-separated version of the proof in [12], which relies on the Jackson kernel. Therefore in Section 2 we derive the suitable ingredients for sparse Jackson kernels while carefully taking into account each variable separately.

A strategy is also required to write a positive polynomial p that is known to be a sum $p = p_1 + \cdots + p_\ell$ with $p_i \in \mathbb{R}[x_{J_i}]$ as a similar sum $p = h_1 + \cdots + h_\ell$ but now with $h_j \in \mathbb{R}[x_{J_j}]$ and $h_j \ge 0$ on $[-1, 1]^{|J_j|}$; this is done in Section 3.

Section 4 gives the proofs of Theorems 2 and 4, together with the statement and proof of Lemma 16, which was used in the proofs of Propositions 3 and 5 above.

The sparse Jackson kernel $\mathbf{2}$

The measure μ_n on the box $[-1,1]^n$ defined by

$$d\mu_n(x) \coloneqq \frac{dx_1}{\pi\sqrt{1-x_1^2}} \cdots \frac{dx_n}{\pi\sqrt{1-x_n^2}}, \quad x = (x_1, \dots, x_n) \in [-1, 1]^n,$$

is known as the *(normalized) Chebyshev measure*; it is a probability measure on $[-1, 1]^n$. It induces the inner product

$$\langle f,g \rangle_{\mu_n} \coloneqq \int_{[-1,1]^n} f(x)g(x) \, d\mu_n(x)$$

and the norm $||f||_{\mu_n} = \sqrt{\langle f, f \rangle_{\mu_n}}$. For $k = 0, 1, \ldots$, we let $T_k \in \mathbb{R}[x]$ be the univariate *Chebyshev polynomial* of degree k, defined by

$$T_k(\cos\theta) \coloneqq \cos(k\theta), \quad \theta \in \mathbb{R}$$

The Chebyshev polynomials satisfy $|T_k(x)| \leq 1$ for all $x \in [-1, 1]$, and

$$\langle T_a, T_b \rangle_{\mu_1} = \int_{-1}^1 \frac{T_a(x)T_b(x)}{\pi\sqrt{1-x^2}} dx = \begin{cases} 0, & a \neq b, \\ 1, & a = b = 0, \\ \frac{1}{2}, & a = b \neq 0. \end{cases}$$

For a multi-index $I = (i_1, \ldots, i_n)$, we let

$$T_I(x_1,\ldots,x_n) \coloneqq T_{i_1}(x_1)T_{i_2}(x_2)\cdots T_{i_n}(x_n)$$

be the multivariate Chebyshev polynomials, which then satisfy (see for example [25, §II.A.1]), for multiindices I and I',

deg
$$T_I = |I|$$
 and $\langle T_I, T_{I'} \rangle_{\mu_n} = \begin{cases} 0, & I \neq I', \\ 2^{-w(I)}, & I = I'. \end{cases}$ (6)

Thus $p \in \mathbb{R}[x]_d$ can be expanded as $p = \sum_{|I| \leq d} 2^{w(I)} \langle p, T_I \rangle_{\mu_n} T_I$. If we let, for a finite collection $\Lambda \subseteq \mathbb{R} \times \mathbb{N}_0^n$ of pairs (λ, I) of a real number λ and a multi-index I,

$$K^{\Lambda}(x,y) = \sum_{(\lambda,I)\in\Lambda} 2^{w(I)}\lambda T_I(x)T_I(y), \quad x,y\in\mathbb{R}^n,$$

then, for any $p \in \mathbb{R}[x]$, we have

$$\mathbf{K}^{\Lambda}(p)(x) \coloneqq \int_{[-1,1]^n} K^{\Lambda}(x,y) p(y) \, d\mu_n(y) = \sum_{\substack{(\lambda,I) \in \Lambda \\ |I| \le d}} 2^{w(I)} \lambda \langle p, T_I \rangle_{\mu_n} T_I(x).$$

This means that, if we set all the nonzero numbers λ equal to 1, then \mathbf{K}^{Λ} is the identity operator in the linear span of $\{T_I : \exists \lambda \neq 0 \text{ s.t. } (\lambda, I) \in \Lambda\} \subseteq \mathbb{R}[x]_d$.

We let, for $r, k \in \mathbb{N}$,

$$\lambda_k^r = \frac{1}{r+2} \left((r+2-k) \cos \frac{\pi k}{r+2} + \frac{\sin \frac{\pi k}{r+2}}{\sin \frac{\pi}{r+2}} \cos \frac{\pi}{r+2} \right), \quad 1 \le k \le r.$$

and

$$\lambda_0^r = 1$$

We set, for $\mathbf{r} = (r_1, \ldots, r_n) \in \mathbb{N}_0^n$,

$$\lambda_I^{\mathbf{r}} = \prod_{j=1}^n \lambda_{i_j}^{r_j}$$

and

$$\Lambda_{\mathbf{r}} = \{ (\lambda_I^{\mathbf{r}}, I) : I \le \mathbf{r} \}.$$

Then $K_{\mathbf{r}}^{\text{Jac}} = K^{\Lambda_{\mathbf{r}}}$ is the (**r**-adapted) Jackson kernel, and its associated linear operator $\mathbf{K}^{\Lambda_{\mathbf{r}}}$ will be denoted $\mathbf{K}_{\mathbf{r}}^{\text{Jac}}$.

Theorem 6. We have $\mathbf{K}_{\mathbf{r}}^{\text{Jac}}(\mathbb{R}[x_J]_{\mathbf{r}}) \subseteq \mathbb{R}[x_J]_{\mathbf{r}}$, and if $p(x) \ge 0$ on $[-1,1]^n$ then $\mathbf{K}_{\mathbf{r}}^{\text{Jac}}(p) \ge 0$ on $[-1,1]^n$. Also, we have:

P1. If $p \in \mathbb{R}[x_J]_{\mathbf{r}}$ satisfies $p(x) \ge 0$ for all $x \in [-1, 1]^n$,

$$\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}}(p) \in \mathcal{P}_{\mathbf{r},J}(\{1-x_i^2\}_{i \in J})$$

P2. Let $p \in \mathbb{R}[x_J]_{\mathbf{r}}$ be a polynomial that satisfies $0 \leq p(x) \leq 1$ for all $x \in [-1,1]^n$, and for all $I = (i_1, \ldots, i_n) \in \mathcal{I}_p$, assume that $\mathbf{r} = (r_1, \ldots, r_n)$ verifies

$$\frac{i_j^2}{(r_j+2)^2} \le \frac{1}{2\pi^2 n}, \quad 1 \le j \le n.$$
(7)

Assume that

$$\varepsilon \ge 2n\pi^2 \left(\prod_{1 \le k \le n} ((\overline{\deg}\,p)_k + 1) \right) \max_{I \in \mathcal{I}_p} \left[2^{w(I)/2} \max_j \frac{i_j^2}{(r_j + 2)^2} \right] > 0.$$

Then

$$\|(\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}})^{-1}(p+\varepsilon) - (p+\varepsilon)\|_{\infty} \leq \varepsilon.$$

Proof. The first statement of the theorem corresponds to Lemma 9(i) and 9(ix). Property P2 follows from Lemma 9(vii).

Let us prove property P1. Take a finite subset $\{z_i\}_i$ of $[-1,1]^n$ and a corresponding set of positive weights $\{w_i\}_i \subset \mathbb{R}$ giving a quadrature rule for integration of polynomials $q \in \mathbb{R}[x_J]_r$, so that

$$\int_{[-1,1]^n} q(x) \, d\mu_n(x) = \sum_i w_i q(z_i), \quad q \in \mathbb{R}[x_J]_{\mathbf{r}}.$$

Then we have, for p as in the statement of P1,

$$\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}}(p)(x) = \sum_{i} w_{i} p(z_{i}) K_{\mathbf{r}}^{\mathrm{Jac}}(z_{i}, x),$$

with $w_i p(z_i) \ge 0$. Since, by Lemma 9(*viii*) and Theorem 8 below, $K_{\mathbf{r}}^{\text{Jac}}(z_i, x)$ is in $\mathcal{P}_{\mathbf{r},J}(\{1 - x_i^2\}_{i \in J})$, so is $\mathbf{K}_{\mathbf{r}}^{\text{Jac}}(p)$.

Corollary 7. If $p \in \mathbb{R}[x_J]_{\mathbf{r}}$ satisfies $0 \le p(x) \le 1$ for all $x \in [-1, 1]^n$, then

$$p + \varepsilon \in \mathcal{P}_{\mathbf{r},J}(\{1 - x_i^2\}_{i \in J})$$

for all multi-indices \mathbf{r} satisfying (7) and

$$\varepsilon \ge 2n\pi^2 \left(\prod_{1 \le k \le n} ((\overline{\deg} \, p)_k + 1) \right) \max_{I \in \mathcal{I}_p} \left(2^{w(I)/2} \max_{1 \le j \le n} \frac{i_j^2}{(r_j + 2)^2} \right).$$

Here, $\mathbf{r} = (r_1, ..., r_n)$ and $I = (i_1, ..., i_n)$.

Proof. By property P2 in Theorem 6,

$$\|(\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}})^{-1}(p+\varepsilon) - (p+\varepsilon)\|_{\infty} \leq \varepsilon$$

Thus, $(\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}})^{-1}(p+\varepsilon) \geq 0$ on $[-1,1]^n$. By property P1 and Lemma 9(i),

$$p + \varepsilon = \mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}} \circ (\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}})^{-1} (p + \varepsilon) \in \mathcal{P}_{\mathbf{r},J}(\{1 - x_i^2\}_{i \in J}).$$

The rest of this section is devoted to results used in the proof of Theorem 6.

Theorem 8 ([6, Th. 10.3]). If $p \in \mathbb{R}[y]$ is a univariate polynomial of degree d nonnegative on the interval $[a,b] \subset \mathbb{R}$, then

$$\begin{cases} p = \sigma_0 + \sigma_1(b - y)(y - a), & \sigma_0 \in \Sigma_d[y], & \sigma_1 \in \Sigma_{d-2}[y] & d \text{ even,} \\ p = \sigma_0(y - a) + \sigma_1(b - y), & \sigma_0 \in \Sigma_{d-1}[y], & \sigma_1 \in \Sigma_{d-1}[y] & d \text{ odd,} \end{cases}$$

where Σ_d is the cone of sum-of-squareds of polynomials of degree at most d.

Lemma 9. Let $\mathbf{r} \in \mathbb{N}_0^n$ be a multi-index. The operator $\mathbf{K}_{\mathbf{r}}^{\text{Jac}}$ defined above has the following properties:

- *i.* $\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}}(\mathbb{R}[x_J]_{\mathbf{r}}) \subseteq \mathbb{R}[x_J]_{\mathbf{r}}.$
- ii. We have

$$\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}}(T_{I}) = \begin{cases} \lambda_{I}^{\mathbf{r}} T_{I}, & I \leq \mathbf{r}, \\ 0, & otherwise. \end{cases}$$

In particular, $\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}}(1) = 1$.

- *iii.* $\mathbf{K}_{\mathbf{r}}^{\text{Jac}}$ is invertible in $\mathbb{R}[x_J]_{\mathbf{r}}$ with $J = \{i : 1 \leq i \leq n, r_i > 0\}$.
- iv. $0 < \lambda_I^{\mathbf{r}} \leq 1$ for all $0 \leq I \leq \mathbf{r}$.
- v. For $I = (i_1, \ldots, i_n)$ and $\mathbf{r} = (r_1, \ldots, r_n)$ in \mathbb{N}_0^n ,

$$|1 - \lambda_I^{\mathbf{r}}| = 1 - \lambda_I^{\mathbf{r}} \le n\pi^2 \max_j \frac{i_j^2}{(r_j + 2)^2}.$$

vi. For $I = (i_1, \ldots, i_n)$ and $\mathbf{r} = (r_1, \ldots, r_n)$ in \mathbb{N}_0^n that verify (7), we have

$$\left|1 - \frac{1}{\lambda_I^{\mathbf{r}}}\right| \le 2n\pi^2 \max_j \frac{i_j^2}{(r_j + 2)^2}.$$

vii. Let $p \in \mathbb{R}[x]_{J,\mathbf{r}}$ with $p(x) \geq 0$ for all $x \in [-1,1]^n$ and $\|p\|_{\infty} \leq 1$. Let $\mathbf{r} = (r_1,\ldots,r_n)$ be a multiindex such that $I \leq \mathbf{r}$ for all $I \in \mathcal{I}_p$, and assume that, for all $I = (i_1, \ldots, i_n) \in \mathcal{I}_p$, condition (7) is verified. Then we have

$$\|(\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}})^{-1}(p) - p\|_{\infty} \le 2n\pi^2 \left(\prod_{1\le k\le n} ((\overline{\deg p})_k + 1)\right) \max_{\substack{I\in\mathcal{I}_p\\1\le j\le n}} \left[2^{w(I)/2} \frac{i_j^2}{(r_j+2)^2}\right].$$

viii. $K_{\mathbf{r}}^{\text{Jac}}(x,y) \geq 0$ for all $x, y \in [-1,1]^n$.

ix. If
$$p \in \mathbb{R}[x]$$
 is such that $p(x) \ge 0$ for $x \in [-1,1]^n$, then $\mathbf{K}^{\mathrm{Jac}}_{\mathbf{r}}(p)(x) \ge 0$ for all $x \in [-1,1]^n$.

Proof. Throughout, we follow [12].

Item (ii) is immediate from the definitions and (6). Item (i) follows from item (ii) and the fact that

 $\{T_I : I \leq \mathbf{r}, I \subseteq J\}$ is a basis for $\mathbb{R}[x_J]_{\mathbf{r}}$. Observe that item (*ii*) means that $\mathbf{K}_{\mathbf{r}}^{\text{Jac}}$ is diagonal in $\mathbb{R}[x_J]_{\mathbf{r}}$, so in order to prove item (*iii*) it suffices to show that $\lambda_I^{\mathbf{r}} > 0$ for all $I \leq \mathbf{r}, I \subseteq J$. This follows immediately from item (*iv*), which in turn follows from the definition of $\lambda_I^{\mathbf{r}}$ and [12, Proposition 6(ii)], which shows that $0 < \lambda_k^r \leq 1$ for all $0 \leq k \leq r$.

Similarly, by [12, Proposition 6(iii)] we have that, if $k \leq r$, then

$$|1 - \lambda_k^r| = 1 - \lambda_k^r \le \frac{\pi^2 k^2}{(r+2)^2}$$

Thus, if $\gamma_j = 1 - \lambda_{i_j}^{r_j} \leq \pi^2 i_j^2 / (r_j + 2)^2$ and $\gamma = \max_j \gamma_j$, we also have, using Bernoulli's inequality [12, Lemma 11]

$$1 - \lambda_I^{\mathbf{r}} = 1 - \prod_{j=1}^n \lambda_{i_j}^{r_j}$$
$$= 1 - \prod_{j=1}^n (1 - \gamma_j)$$
$$\leq 1 - (1 - \gamma)^n$$
$$\leq n\gamma$$
$$\leq n\pi^2 \max_j \frac{i_j^2}{(r_j + 2)^2}$$

This shows item (v). Using it, we can prove item (vi) as follows: condition (7) implies, by item (v), that $|1 - \lambda_I^{\mathbf{r}}| \leq 1/2$, and hence $|\lambda_I^{\mathbf{r}}| \geq 1/2$, so

$$\left|1 - \frac{1}{\lambda_I^{\mathbf{r}}}\right| = \frac{|1 - \lambda_I^{\mathbf{r}}|}{|\lambda_I^{\mathbf{r}}|} \le 2n\pi^2 \max_j \frac{i_j^2}{(r_j + 2)^2},$$

leveraging item (v) again.

Let us show item (vii). From items (ii) and (iii), we have

$$\begin{aligned} \|(\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}})^{-1}(p) - p\|_{\infty} &= \left\| \sum_{I} \left[\frac{1}{\lambda_{I}^{\mathbf{r}}} 2^{w(I)} \langle p, T_{I} \rangle_{\mu_{n}} T_{I} - 2^{w(I)} \langle p, T_{I} \rangle_{\mu_{n}} T_{I} \right] \right\|_{\infty} \\ &\leq \sum_{I} 2^{w(I)} |\langle p, T_{I} \rangle_{\mu_{n}} | \left| 1 - \frac{1}{\lambda_{I}^{\mathbf{r}}} \right|, \end{aligned}$$

because $|T_I(x)| \leq 1$ for all $x \in [-1,1]^n$. Plugging in the estimate from item (vi), we get

$$\begin{aligned} \|(\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}})^{-1}(p) - p\|_{\infty} &\leq \sum_{I} 2^{w(I)/2+1} n\pi^{2} \max_{j} \frac{i_{j}^{2}}{(r_{j}+2)^{2}} \\ &\leq 2n\pi^{2} \left(\prod_{1 \leq k \leq n} ((\overline{\deg} p)_{k} + 1) \right) \max_{I \in \mathcal{I}_{p}} \left[2^{w(I)/2} \max_{j} \frac{i_{j}^{2}}{(r_{j}+2)^{2}} \right] \end{aligned}$$

where we have also used

$$|\langle p, T_I \rangle_{\mu_n}| \le ||p||_{\mu_n} ||T_I||_{\mu_n} \le ||T_I||_{\mu_n} = 2^{-w(I)/2},$$

which follows from (6).

To prove item (viii), let, for fixed \mathbf{r} ,

$$\Lambda_k = \{ (\lambda_I^r, I) : I \le (0, \dots, 0, r_k, 0, \dots, 0) \} = \{ (\lambda_{i_k}^{r_k}, (0, \dots, 0, i_k, 0, \dots, 0)) : i_k \le r_k \}, \quad 1 \le k \le n,$$

and observe that

$$\mathbf{K}_{\mathbf{r}}^{\mathrm{Jac}} = \mathbf{K}^{\Lambda_{\mathbf{r}}} = \mathbf{K}_{x_{1}}^{\Lambda_{1}} \circ \mathbf{K}_{x_{2}}^{\Lambda_{2}} \circ \cdots \circ \mathbf{K}_{x_{n}}^{\Lambda_{n}}$$

$$\tag{8}$$

where $\mathbf{K}_{x_k}^{\Lambda_k}$ is the operator \mathbf{K}^{Λ_k} acting in the variable x_k , i.e.,

$$\mathbf{K}_{x_k}^{\Lambda_k}(p)(x) = \int_{-1}^1 K^{\Lambda_k}(x_k, y) \, p(x_1, \dots, x_{k-1}, y, x_{k+1}, \dots, x_n) \, d\mu_1(y).$$

Equation (8) follows from the identity

$$K^{\Lambda_{\mathbf{r}}}(x,y) = K^{\Lambda_{1}}(x_{1},y_{1})K^{\Lambda_{2}}(x_{2},y_{2})\cdots K^{\Lambda_{n}}(x_{n},y_{n})$$

= $K^{\mathrm{Jac}}_{(r_{1})}(x_{1},y_{1})K^{\mathrm{Jac}}_{(r_{2})}(x_{2},y_{2})\cdots K^{\mathrm{Jac}}_{(r_{n})}(x_{n},y_{n})$

that can be checked from the definitions. Item (viii) then follows from the well-known fact that $K_{(r)}^{\text{Jac}}(x,y) \ge 0$ for all $r \in \mathbb{N}_0$ and all $x, y \in [-1,1]$; see for example [25, §II.C.2–3]. Item (ix) follows immediately from item (viii).

Sparse approximation theory 3

For $1 \leq i \leq n$ and a function $f: [-1,1]^n \to \mathbb{R}$, let

$$\operatorname{Lip}_{i} f = \sup_{\substack{x \in [-1,1]^{n} \\ y \in [-1,1]}} \frac{|f(x) - f(x_{1}, \dots, x_{i-1}, y, x_{i+1}, \dots, x_{n})|}{|x_{i} - y|}.$$

Theorem 10. There is a constant $C_{\text{Jac}} > 0$ such that the following is true. Let $f \in C^0([-1,1]^n)$ be a Lipschitz function with variable-wise Lipschitz constants $\text{Lip}_1 f, \ldots, \text{Lip}_n f$. Then for each multi-index $\mathbf{m} = (m_1, \ldots, m_n) \in \mathbb{N}^n$ there is a polynomial $p \in \mathbb{R}[x]_{\mathbf{m}}$ such that

$$\sup_{x \in [-1,1]^n} |f(x) - p(x)| \le C_{\operatorname{Jac}} \sum_{i=1}^n \frac{\operatorname{Lip}_i f}{m_i}$$

and

$$\operatorname{Lip}_i p \le 2 \operatorname{Lip}_i f.$$

Proof. Jackson [5, p. 2–6] proved that there is a constant C > 0 such that, if $g: \mathbb{R} \to \mathbb{R}$ is Lipschitz and π -periodic, $g(0) = g(\pi)$, then

$$\left| g(\theta) - \frac{\int_{-\pi/2}^{\pi/2} g(\theta - \vartheta) \left(\frac{\sin m\vartheta}{m \sin \vartheta}\right)^4 d\vartheta}{\int_{-\pi/2}^{\pi/2} \left(\frac{\sin m\vartheta}{m \sin \vartheta}\right)^4 d\vartheta} \right| \le \frac{C \operatorname{Lip} g}{m}, \quad m \in \mathbb{N}, x \in \mathbb{R}.$$
(9)

For a multivariate Lipschitz function $g: \mathbb{R}^n \to \mathbb{R}$ and a multi-index $\mathbf{m} = (m_1, \dots, m_n) \in \mathbb{N}^n$, let

$$L_{i}(g)(\theta) = \frac{\int_{-\pi/2}^{\pi/2} \cdots \int_{-\pi/2}^{\pi/2} g(\theta_{1} - \vartheta_{1}, \dots, \theta_{i} - \vartheta_{i}, \theta_{i+1}, \dots, \theta_{n}) \prod_{j=1}^{i} \left(\frac{\sin m_{j}\vartheta_{j}}{m_{j}\sin \vartheta_{j}}\right)^{4} d\vartheta_{1} \dots d\vartheta_{i}}{\prod_{j=1}^{i} \int_{-\pi/2}^{\pi/2} \left(\frac{\sin m_{j}\vartheta_{j}}{m_{j}\sin \vartheta_{j}}\right)^{4} d\vartheta_{j}}$$

Then we have, using the triangle inequality and the single-variable inequality (9) at each step,

$$\begin{aligned} |g(\theta) - L_n(g)(\theta)| \\ &\leq |g(\theta) - L_1(g)(\theta)| + |L_1(g)(\theta) - L_2(g)(\theta)| + \dots + |L_{n-1}(g)(\theta) - L_n(g)(\theta)| \\ &\leq C\left(\frac{\operatorname{Lip}_1 g}{m_1} + \dots + \frac{\operatorname{Lip}_n g}{m_n}\right). \end{aligned}$$

The function $\prod_j (\sin m_j \theta_j / m_j \sin \theta_j)^4$ is a polynomial of degree m_j in $\cos \theta_j$ (cf. [5, p. 3]). If we replace f with its Lipschitz extension to $[-2, 2]^n$ and apply the results above to $g(\theta) = f(2\cos\theta_1, \ldots, 2\cos\theta_n)$ we get a polynomial $L_n(g)(\theta)$ in $\cos\theta_1, \ldots, \cos\theta_n$ satisfying the above inequality. Thus

$$p(x) = L_n(g)(\arccos(x_1/2), \dots, \arccos(x_n/2)), \quad x \in [-2, 2],$$

is a polynomial with $\overline{\deg} p \leq \mathbf{m}$ that satisfies (cf. [5, p. 13–14])

$$|f(x) - p(x)| \le C\left(\frac{\operatorname{Lip}_1 g}{m_1} + \dots + \frac{\operatorname{Lip}_n g}{m_n}\right) \le 2C\left(\frac{\operatorname{Lip}_1 f}{m_1} + \dots + \frac{\operatorname{Lip}_n f}{m_n}\right),$$

since $\operatorname{Lip}_i g \leq 2 \operatorname{Lip}_i f$. This proves the first statement, setting $C_{\operatorname{Jac}} = 2C$. We also have

$$\left|\frac{d}{dx}\arccos(x/2)\right| = \frac{1}{2\sqrt{1-(x/2)^2}} \le \frac{1}{\sqrt{3}} \quad \text{for} \quad x \in [-1,1]$$

and, by linearity and monotonicity of L_n ,

 $|L_n g(\theta) - L_n g(\theta_1, \dots, \theta_{i-1}, \theta_i + t, \theta_{i+1}, \dots, \theta_n)| \leq |L_n(|t| \operatorname{Lip}_i g)(\theta)| = |t| \operatorname{Lip}_i g \leq 2|t| \operatorname{Lip}_i f,$

whence

$$\operatorname{Lip}_{i} p = \operatorname{Lip}_{i} L_{n}(g)(\operatorname{arccos}(x_{1}/2), \dots, \operatorname{arccos}(x_{n}/2))$$
$$\leq \operatorname{Lip}_{i} L_{n}(g) \left| \frac{d}{dx_{i}} \operatorname{arccos} \frac{x_{i}}{2} \right| \leq \operatorname{Lip}_{i} L_{n}(g) \leq 2\operatorname{Lip}_{i} f.$$

Lemma 11 (a version of [4, Lemma 3]). Let J_1, \ldots, J_ℓ be subsets of $\{1, \ldots, n\}$ satisfying the running intersection property. Suppose $f = f_1 + \cdots + f_\ell$ with $\ell \ge 2$, $f_j \in \mathbb{R}[x_{J_j}]$. Let $\varepsilon > 0$ be such that $f \ge \varepsilon$ on $S(\mathbf{g}) \subseteq [-1, 1]^n$. Pick numbers $\epsilon, \eta > 0$ so that

$$\varepsilon = (\ell - 1)\epsilon - (\ell - 2)\eta \quad and \quad \epsilon > 2\eta$$

Set, for $2 \leq l \leq \ell$,

$$D_{l,m} = \left\lceil \frac{2 C_{\text{Jac}} \left| \mathcal{J}_l \right| \sum_{k=l}^m \text{Lip} f_k}{\epsilon - 2\eta} \right\rceil$$
(10)

with \mathcal{J}_l as in (1), and $D_{1,m} = D_{2,m}$.

Then $f = h_1 + \cdots + h_\ell$ for some $h_j \in \mathbb{R}[x_{J_j}]$ with $h_j \ge \eta$ on $S(\mathbf{g}) \subseteq [-1, 1]^n$ and

$$\overline{\deg} h_j \le \max(\overline{\deg} f_j, \overline{D}_{j,\ell}, \overline{D}_{j+1,\ell}, \dots, \overline{D}_{\ell,\ell})_{J_j}$$
(11)

where $\overline{D}_{j,m}$ is the multi-index whose k-th entry equals $D_{j,m}$ if $k \in \mathcal{J}_j = J_j \cap \bigcup_{k < j} J_i$ and 0 otherwise, and the maximum is taken entry-wise.

Additionally, if Lip f denotes the Lipschitz constant of f on $[-1,1]^n$, then

$$\operatorname{Lip} h_j \le 3 \sum_{k=j}^{\ell} \operatorname{Lip} f_k$$

Finally, we have

$$||h_j||_{\infty} \le 3 \times 2^{\ell-1} \sum_{j=1}^{\ell} ||f_j||_{\infty}.$$

Remark 12. If $S(\mathbf{g}) = [-1, 1]^n$, we also have the obvious estimate $||h_j||_{\infty} \leq ||f||_{\infty}$, that follows from $0 \leq h_j \leq f$.

Proof. In order to prove the result by induction, let us first consider the case $\ell = 2$. In this case, $\varepsilon = \epsilon$ and $\epsilon > 2\eta$. Assume that $J_1 \cap J_2 \neq \emptyset$. For a subset $J \subset \{1, \ldots, n\}$, let π_J denote the projection onto the variables with indices in J, that is, $\pi_J(x) = (x_i)_{i \in J} \in [-1, 1]^J$ for $x \in [-1, 1]^n$. Define $g: [-1, 1]^{J_1 \cap J_2} \to \mathbb{R}$ by

$$g(x) := \min_{y \in \pi_{J_1 \setminus J_2}(S(\mathbf{g})) \subseteq [-1,1]^{J_1 \setminus J_2}} f_2(x,y) - \frac{\varepsilon}{2}, \qquad x \in [-1,1]^{J_1 \cap J_2}$$

The function g is Lipschitz continuous on $[-1,1]^{J_1 \cap J_2}$. To see why, let $x, x' \in [-1,1]^{J_1 \cap J_2}$ and pick $y, y' \in \pi_{J_1 \setminus J_2}(S(\mathbf{g})) \subseteq [-1,1]^{J_1 \setminus J_2}$ minimizing $f_2(x,y)$ and $f_2(x',y')$, respectively. Then

$$|g(x) - g(x')| = |f_2(x, y) - f_2(x', y')| \le \max(|f_2(x, y) - f_2(x', y)|, |f_2(x, y') - f_2(x', y')|) \le \operatorname{Lip}(f_2)|x - x'|,$$

where $\operatorname{Lip}(f_2)$ denotes the Lipschitz constant of f_2 on $[-1,1]^n$.

The function g also satisfies

$$f_1 + g \ge \frac{\varepsilon}{2}$$
 and $f_2 - g \ge \frac{\varepsilon}{2}$

on $S(\mathbf{g})$. The second inequality follows from the definition of g, and the first one can be shown taking $(x, y, z) \in S(\mathbf{g})$ with $x \in [-1, 1]^{J_1 \cap J_2}$, $y \in [-1, 1]^{J_1 \setminus J_2}$, and $z \in [-1, 1]^{J_2 \setminus J_1}$, taking care to pick y only after x has been chosen, in such a way that the minimum is in the definition of g is realized there, that is, $g(x) = f_2(x, y) - \varepsilon/2$ holds (this is possible by compactness of $S(\mathbf{g})$ and continuity of f); then we have

$$f_1(x,z) + g(x) = f_1(x,z) + f_2(x,y) - \frac{\varepsilon}{2} = f(x,y,z) - \frac{\varepsilon}{2} \ge \frac{\varepsilon}{2}$$

For $j \in J_1 \cap J_2$, let

$$m_j = D_{2,2} = \left\lceil \frac{2C_{\text{Jac}} |J_1 \cap J_2| \operatorname{Lip}(f_2)}{\varepsilon - 2\eta} \right\rceil.$$

Set $m_j = 0$ for all other $0 \le j \le n$, and $\mathbf{m} = (m_1, \ldots, m_n) = \overline{D}_{2,2}$. Then Theorem 10 gives a polynomial p_2 such that

$$\|g - p_2\|_{\infty} = C_{\text{Jac}} \sum_{j \in J_1 \cap J_2} \frac{\operatorname{Lip}_j g}{m_j} \le C_{\text{Jac}} |J_1 \cap J_2| \operatorname{Lip}(f_2) \frac{2}{D_{2,2}} \le \frac{\varepsilon}{2} - \eta.$$

Also,

$$\overline{\deg} \, p_2 \le \mathbf{m} = \bar{D}_{2,2}.\tag{12}$$

Let

$$h_1 \coloneqq f_1 + p_2$$
 and $h_2 \coloneqq f_2 - p_2$

so that $f = h_1 + h_2$, $h_1 \ge \eta$ and $h_2 \ge \eta$ on $S(\mathbf{g})$, and $h_j \in \mathbb{R}[x_{J_j}]$. The bound (11) follows from the definition of h_j and (12). Observe also that, by the last part of Theorem 10,

$$\operatorname{Lip} p_2 \le 2\operatorname{Lip} g \le 2\operatorname{Lip}(f_2)$$

Finally, we have

$$\|p_2\|_{\infty} \le \|g\|_{\infty} + \frac{\varepsilon}{2} - \eta \le \|f_2\|_{\infty} + \varepsilon - \eta \le 2\|f_2\|_{\infty},\tag{13}$$

 \mathbf{SO}

$$||h_j||_{\infty} \le ||f_j||_{\infty} + ||p_2||_{\infty} \le ||f_j||_{\infty} + 2||f_2||_{\infty} \le 3(||f_1||_{\infty} + ||f_2||_{\infty}).$$

For the induction step, let $\ell \geq 3$ and set $\tilde{f} = f_1 + \cdots + f_{\ell-1} - (\ell-2)(\epsilon - \eta)$, so that we have $f - (\ell-2)(\epsilon - \eta) = \tilde{f} + f_\ell \geq \epsilon$ since $f \geq \epsilon = (\ell-1)\epsilon - (\ell-2)\eta$. The proof for the case $\ell = 2$ with $\epsilon = \epsilon_{\ell-1}$ gives a polynomial $p_\ell \in \mathbb{R}[x_{\mathcal{J}_\ell}]$ such that

$$f - p_{\ell} \ge \eta$$
 and $f_{\ell} + p_{\ell} \ge \eta$

on $S(\mathbf{g})$, and with $\overline{\deg} p_{\ell} = \overline{D}_{\ell,\ell}$, $\operatorname{Lip} p_{\ell} \leq 2 \operatorname{Lip} f_{\ell}$. and, analogously to (13),

$$\|p_{\ell}\|_{\infty} \le 2\|f_{\ell}\|_{\infty}.$$
 (14)

Write

$$f_1' + \dots + f_{\ell-1}' = f_1 + \dots + f_{\ell-1} - p_\ell$$

where $f'_j = f_j - p_\ell$ for the largest j with $\mathcal{J}_\ell \subset J_j$ (which must happen for some j, by the running intersection property; see Definition 1) and $f'_k = f_k$ for all other $k \neq j$. Thus $f'_j \in \mathbb{R}[x_{J_j}]$,

$$\overline{\deg} f'_j \le \max(\overline{\deg} f_j, \overline{\deg} p_\ell) = \max(\overline{\deg} f_j, \overline{D}_{\ell,\ell}), \qquad \operatorname{Lip} f'_j \le \operatorname{Lip} f_j + \operatorname{Lip} p_\ell, \tag{15}$$

The induction hypothesis applies to the polynomial

$$f'_1 + \dots + f'_{\ell-1} + (\ell-2)(\epsilon - \eta) = \tilde{f} + (\ell-2)(\epsilon - \eta) - p_\ell \ge (\ell-2)\epsilon - (\ell-3)\eta$$

This means that there are polynomials $h_1, \ldots, h_{\ell-1}$ such that

- $f'_1 + \dots + f'_{\ell-1} = f_1 + \dots + f_{\ell-1} p_\ell = h_1 + \dots + h_{\ell-1},$
- $h_j \in \mathbb{R}[x_{J_j}]$ for all $1 \le j \le \ell 1$,
- $h_j \ge \eta$ for all $1 \le j \le \ell 1$,
- We have, for all $1 \le j \le \ell 1$,

$$\overline{\deg} h_j \le \max(\overline{\deg} f'_j, \overline{D}_{j,\ell} \dots, \overline{D}_{\ell-1,\ell}) \le \max(\overline{\deg} f_j, \overline{\deg} p_\ell, \overline{D}_{j,\ell}, \dots, \overline{D}_{\ell-1,\ell}) = \max(\overline{\deg} f_j, \overline{D}_{j,\ell}, \dots, \overline{D}_{\ell,\ell}).$$

Observe that the second index in each $\bar{D}_{k,\ell}$ is ℓ because of the accumulation of Lipschitz constants resulting from the estimate (15).

• We have, for all $1 \le j \le \ell - 1$, again because of (15),

$$\operatorname{Lip} h_j \le 3\sum_{k=j}^{\ell} \operatorname{Lip} f_k$$

• We have, for all $1 \le j \le \ell - 1$, using (14),

$$\|h_j\|_{\infty} \le 3 \times 2^{\ell-2} \sum_{k=1}^{\ell-1} \|f'_k\|_{\infty} \le 3 \times 2^{\ell-2} \left(\sum_{k=1}^{\ell-1} \|f_k\|_{\infty} + \|p_\ell\|_{\infty} \right) \le 3 \times 2^{\ell-1} \sum_{k=1}^{\ell} \|f_k\|_{\infty}.$$

Let $h_{\ell} = f_{\ell} + p_{\ell}$. Then again $f_1 + \dots + f_{\ell} = h_1 + \dots + h_{\ell}, h_{\ell} \in \mathbb{R}[x_{J_{\ell}}], h_{\ell} \ge \eta$ on $S(\mathbf{g}), \overline{\deg} h_j \le \max(\overline{\deg} f_{\ell}, \overline{D}_{\ell,\ell}), \operatorname{Lip} h_{\ell} \le \operatorname{Lip} f_{\ell} + \operatorname{Lip} p_{\ell} \le 3 \operatorname{Lip} f_{\ell},$

$$\|h_{\ell}\|_{\infty} \le \|f_{\ell}\|_{\infty} + \|p_{\ell}\| \le 3\|f_{\ell}\| \le 3 \times 2^{\ell-1} \sum_{j=1}^{\ell} \|f_j\|_{\infty}$$

so the lemma is proven.

4 Proofs

4.1 Proof of Theorem 2

Theorem 2 will follow from Theorem 13, which presents a more detailed bound, together with the definitions of $\overline{L}, M, \overline{J}$.

Theorem 13. Let n > 0 and $\ell \ge 2$, and let $\mathbf{r}_1, \mathbf{r}_2, \ldots, \mathbf{r}_\ell \in \mathbb{N}^n$, $\mathbf{r}_j = (r_{j,1}, \ldots, r_{j,n})$, be nowhere-vanishing multi-indices. Let also J_1, \ldots, J_ℓ be subsets of $\{1, \ldots, n\}$ satisfying the running intersection property. Let $p = p_1 + p_2 + \cdots + p_\ell$ be a polynomial that is the sum of finitely many polynomials $p_j \in \mathbb{R}[x_{J_j}]_{\mathbf{r}_j}$. Then if $p \ge \varepsilon$ on $[-1, 1]^n$, we have

$$p \in \mathcal{P}_{\mathbf{r}_1, J_1}(\{1 - x_i^2\}_{i \in J_1}) + \dots + \mathcal{P}_{\mathbf{r}_\ell, J_\ell}(\{1 - x_i^2\}_{i \in J_\ell})$$

as long as, for all $1 \leq j \leq \ell$ and all $1 \leq i \leq n$,

$$(r_{j,k}+2)^{2} \geq \frac{2^{\frac{|J_{j}|}{2}+2}(\ell+2)\|p\|_{\infty}n\pi^{2}}{\varepsilon} \\ \cdot \prod_{1\leq m\leq n} \left(\max\left[(\overline{\deg p_{j}})_{m}, \max_{\substack{j\leq l\leq \ell\\m\in\mathcal{J}_{l}}} \frac{4C_{\mathrm{Jac}}(\ell+2)|\mathcal{J}_{l}|\sum_{t=l}^{\ell}\mathrm{Lip}\,p_{t}}{\varepsilon} \right] + 2 \right) \\ \cdot \max_{l\in J_{j}} \left[(\overline{\deg p_{j}})_{l}, \max_{\substack{j\leq q\leq \ell\\l\in\mathcal{J}_{q}}} \frac{4C_{\mathrm{Jac}}(\ell+2)|\mathcal{J}_{q}|\sum_{t=q}^{\ell}\mathrm{Lip}\,p_{t}}{\varepsilon} \right]^{2},$$
(16)

and

$$(r_{j,i}+2)^2 \ge 2\pi^2 n \max\left[\max_{1\le m\le n} (\overline{\deg}\,p_j)_m, \max_{j\le k\le \ell} \frac{4C_{\operatorname{Jac}}(\ell+2)|\mathcal{J}_k|\sum_{t=k}^{\ell}\operatorname{Lip} p_t}{\varepsilon}\right]^2,$$
(17)

Proof of Theorem 13. Let

$$\epsilon = \frac{\varepsilon + (\ell - 2)\eta}{\ell} \quad \text{and} \quad \eta = \frac{\varepsilon}{2(\ell + 2)}$$
(18)

and apply Lemma 11 (with $\mathbf{g} = 0$, so that $S(\mathbf{g}) = [-1, 1]^n$) to get polynomials h_1, \ldots, h_ℓ with $h_j \in \mathbb{R}[x_{J_j}]$, $p = h_1 + \cdots + h_\ell$ and, for all $1 \leq j \leq \ell, h_j \geq \eta$ on $[-1, 1]^n$ and

$$\overline{\deg} h_j \le \max(\overline{\deg} p_j, \overline{D}_j, \overline{D}_{j+1}, \dots, \overline{D}_\ell),$$

where $\bar{D}_l := (\delta_{j \in \mathcal{J}_l} D_l)_{j=1}^{\ell}, \, \delta_{j \in \mathcal{J}_l}$ equals 1 if $j \in \mathcal{J}_l$ and 0 otherwise, and

$$D_l := \left\lceil \frac{2C_{\text{Jac}} |\mathcal{J}_l| \sum_{t=l}^{\ell} \text{Lip} p_t}{\varepsilon_{l-1} - 2\eta} \right\rceil = \left\lceil \frac{4C_{\text{Jac}} \ell |\mathcal{J}_l| \sum_{t=l}^{\ell} \text{Lip} p_t}{\eta(\ell+2)} \right\rceil, \qquad l = 2, 3, \dots, \ell,$$

(since $\varepsilon_j - 2\eta = \eta(\ell + 2)/2\ell$) and $D_1 := D_2$. Thus $|\bar{D}_l| = |\mathcal{J}_l|D_l$ for $2 \le l \le \ell$. Apply Corollary 7 to each of the polynomials

$$H_j = \frac{h_j - \min_{[-1,1]^n} h_j}{\max_{[-1,1]^n} h_j - \min_{[-1,1]^n} h_j}$$

to see that, for

$$\varepsilon_j \ge 2n\pi^2 \left(\prod_{1 \le k \le n} \left((\overline{\deg} H_j)_k + 1 \right) \right) \max_{I \in \mathcal{I}_{H_j}} \left(2^{w(I)/2} \max_{1 \le k \le n} \frac{i_k^2}{(r_{j,k} + 2)^2} \right), \tag{19}$$

(recall that \mathcal{I}_{H_j} is the set of multiindices $I = (i_1, \ldots, i_n)$ corresponding to exponents of x_1, \ldots, x_n in the terms appearing in H_j and w(I) is the number of nonzero entries in I) we have

$$H_j + \epsilon_j \in \mathcal{P}_{\mathbf{r}_j, J_j}(\{1 - x_i^2\}_{i \in J_j});$$

$$\tag{20}$$

when applying the corollary, note that (17) implies (7) in this case because, if $I = (i_1, \ldots, i_n) \in \mathcal{I}_{H_j}$, then

$$i_{k} \leq (\overline{\deg} H_{j})_{k} \leq (\overline{\deg} h_{j})_{k} \leq \max(\overline{\deg} p_{j}, \overline{D}_{j}, \dots, \overline{D}_{\ell})$$
$$\leq \max\left[\max_{1 \leq m \leq n} (\overline{\deg} p_{j})_{m}, \max_{j \leq k \leq \ell} \frac{4C_{\operatorname{Jac}}(\ell+2)|\mathcal{J}_{k}| \sum_{t=k}^{\ell} \operatorname{Lip} p_{t}}{\varepsilon}\right],$$

by the definition of \overline{D}_l . Observe that (20) means also that

$$h_j - \min_{[-1,1]^n} h_j + \epsilon_j \left(\max_{[-1,1]^n} h_j - \min_{[-1,1]^n} h_j \right) \in \mathcal{P}_{\mathbf{r}_j,J_j}(\{1 - x_i^2\}_{i \in J_j}).$$
(21)

Note that we have $\overline{\deg} H_j = \overline{\deg} h_j$, $\mathcal{I}_{H_j} \setminus \mathcal{I}_{h_j} = \emptyset$, and $\mathcal{I}_{h_j} \setminus \mathcal{I}_{H_j} \subseteq \{(0, \ldots, 0)\}$ since the powers of all terms in h_j and in H_j are the same, with the only possible exception of the constant term, which may appear in one of these and vanish in the other. Now, going back to our choice (18) of η and using (16), we have

$$\begin{split} \eta &= \frac{\varepsilon}{2(\ell+2)} \\ &\geq \|p\|_{\infty} n\pi^2 \left(\max\left[\max_{1 \leq m \leq n} (\overline{\deg} \, p_j)_m, \max_{j \leq k \leq \ell} \frac{2C_{\text{Jac}} \, |\mathcal{J}_k| \sum_{t=k}^{\ell} \operatorname{Lip} p_t}{\eta} \right] + 2 \right)^{|J_j|+2} \\ &\quad \cdot \frac{2^{\frac{|J_j|}{2}+1}}{\min_{1 \leq k \leq n} (r_{j,k}+2)^2} \\ &\geq \|p\|_{\infty} n\pi^2 \left(\max\left[\max_{1 \leq m \leq n} (\overline{\deg} \, p_j)_m, \max_{j \leq k \leq \ell} \left[\frac{2C_{\text{Jac}} \, \ell \, |\mathcal{J}_k| \sum_{t=k}^{\ell} \operatorname{Lip} p_t}{\eta(\ell+2)} \right] \right] + 1 \right)^{|J_j|+2} \\ &\quad \cdot \frac{2^{\frac{|J_j|}{2}}+1}}{\min_{1 \leq k \leq n} (r_{j,k}+2)^2}. \end{split}$$

for all $j \in \{1, ..., \ell\}$. Notice that after separating two of the $|J_j| + 2$ terms in the product and removing the +1 factor from them, we obtain

$$\begin{split} \eta &\geq \|p\|_{\infty} n\pi^2 \left(\max\left[\max_{1 \leq m \leq n} (\overline{\deg} \, p_j)_m, \max_{j \leq k \leq \ell} \left[\frac{2C_{\operatorname{Jac}} \,\ell \,|\mathcal{J}_k| \sum_{t=k}^{\ell} \operatorname{Lip} p_t}{\eta(\ell+2)} \right] \right] + 1 \right)^{|\mathcal{J}_j|} 2^{\frac{|\mathcal{J}_j|}{2} + 1} \\ &\cdot \frac{\max\left[\max_{1 \leq m \leq n} (\overline{\deg} \, p_j)_m, \max_{j \leq k \leq \ell} \left[\frac{2C_{\operatorname{Jac}} \,\ell \,|\mathcal{J}_k| \sum_{t=k}^{\ell} \operatorname{Lip} p_t}{\eta(\ell+2)} \right] \right]^2}{\min_{1 \leq k \leq n} (r_{j,k} + 2)^2} \\ &\geq \|p\|_{\infty} n\pi^2 \left(\prod_{k \in \mathcal{J}_j} \left(\left[\max\left(\overline{\deg} \, p_j, \bar{D}_j, \dots, \bar{D}_\ell \right) \right]_k + 1 \right) \right) 2^{\frac{|\mathcal{J}_j|}{2} + 1} \\ &\cdot \frac{\max_{1 \leq k \leq n} \left[\max\left(\overline{\deg} \, p_j, \bar{D}_j, \dots, \bar{D}_\ell \right) \right]_k^2}{\min_{1 \leq k \leq n} (r_{j,k} + 2)^2}, \end{split}$$

where we have used the definition of \overline{D}_l , as well as the fact that each factor has been replaced by one that is smaller or equal, the original expression containing the maximum of them on each factor. Next,

use $\overline{\deg} H_j \leq \max(\overline{\deg} p_j, \overline{D}_j, \dots, \overline{D}_\ell)$ as well as $w(I) \leq |J_j|$ for every multi-index I in \mathcal{I}_{H_j} , which is true because $H_j \in \mathbb{R}[x_{J_j}]$, yielding

$$\begin{split} \eta &\geq \|p\|_{\infty} \left(2n\pi^{2} \left(\prod_{1 \leq k \leq n} \left(\left(\overline{\deg} \, H_{j} \right)_{k} + 1 \right) \right) \max_{I \in \mathcal{I}_{H_{j}}} \left(2^{w(I)/2} \max_{1 \leq k \leq n} \frac{i_{k}^{2}}{(r_{j,k} + 2)^{2}} \right) \right) \\ &\geq \left(\max_{[-1,1]^{n}} h_{j} - \min_{[-1,1]^{n}} h_{j} \right) \\ &\cdot \left(2n\pi^{2} \left(\prod_{1 \leq k \leq n} \left(\left(\overline{\deg} \, H_{j} \right)_{k} + 1 \right) \right) \max_{I \in \mathcal{I}_{H_{j}}} \left(2^{w(I)/2} \max_{1 \leq k \leq n} \frac{i_{k}^{2}}{(r_{j,k} + 2)^{2}} \right) \right), \end{split}$$

since we have $\max_{[-1,1]^n} h_j - \min_{[-1,1]^n} h_j \leq ||p||_{\infty}$. With this bound for η , together with the fact that $\min_{[-1,1]^n} h_j \geq \eta$, we get

$$h_{j} \geq h_{j} - \min_{[-1,1]^{n}} h_{j} + \eta$$

$$\geq h_{j} - \min_{[-1,1]^{n}} h_{j} + \left(\max_{[-1,1]^{n}} h_{j} - \min_{[-1,1]^{n}} h_{j} \right)$$

$$\cdot \left(2n\pi^{2} \left(\prod_{1 \leq k \leq n} \left((\overline{\deg} H_{j})_{k} + 1 \right) \right) \max_{I \in \mathcal{I}_{H_{j}}} \left(2^{w(I)/2} \max_{1 \leq k \leq n} \frac{i_{k}^{2}}{(r_{j,k} + 2)^{2}} \right) \right),$$

so that, by (19) and (21), $h_j \in \mathcal{P}_{\mathbf{r},J_j}(\{1-x_i^2\}_{i \in J_j})$ and hence

$$p = h_1 + \dots + h_\ell \in \mathcal{P}_{\mathbf{r}, J_1}(\{1 - x_i^2\}_{i \in J_1}) + \dots + \mathcal{P}_{\mathbf{r}, J_\ell}(\{1 - x_i^2\}_{i \in J_\ell}).$$

4.2 Proof of Theorem 4

Overview. For this proof, we will first use the sparse approximation theory developed in Section 3 to represent the sparse polynomial p as a sum of positive polynomials $h_1 + \cdots + h_\ell$, each of them depending on a clique of variables J_j . We then work with each of these polynomials h_j using the tools developed by Baldi–Mourrain [1] to write $h_j = \hat{f}_j + \hat{q}_j$, where \hat{q}_j is by construction obviously an element of the corresponding quadratic module, and \hat{f}_j is strictly positive on $[-1, 1]^n$. Thus Corollary 7 can be applied to \hat{f}_j , which shows that it belongs to the preordering, and then one argues (also following the ideas of [1]) that the preordering is contained in the quadratic module, hence giving that \hat{f}_j is contained in the latter as well. In sum, this shows that h_j is in the quadratic module, which is what want. Most of the heavy lifting goes to estimating the minimum of \hat{f}_j to justify the application of Corollary 7.

Proof of Theorem 4. For each $j = 1, \ldots, \ell$, pick $\mathbf{C}_j > 0$ such that the following two bounds are satisfied:

$$\begin{aligned} \mathbf{C}_{j} &\geq 2\pi^{2} |J_{j}|^{1+\frac{16\mathsf{L}_{j}}{3}} C_{d}^{2} C_{\mathrm{Jac}}^{\frac{16\mathsf{L}_{j}}{3}} 2^{1+2(4+3\frac{8}{3})\mathsf{L}_{j}} 3^{\frac{(16+8\ell)\mathsf{L}_{j}+2}{3}} \bar{k}^{-\frac{2}{3}} c_{j}^{\frac{8}{3}} (\max_{i\in K_{j}} \deg g_{i})^{2} (2(\ell+2))^{8\mathsf{L}_{j}}, \end{aligned} \tag{22} \\ \mathbf{C}_{j} &\geq C_{f} (C_{\mathrm{Jac}} C_{m})^{(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} |J_{j}| \pi^{2} 2^{4\mathsf{L}_{j}+\frac{|J_{j}|}{2}+1+(1+\frac{4\mathsf{L}_{j}+1}{3})(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} \\ &\times 3^{\ell(\mathsf{L}_{j}+1)+(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} (\ell+2)^{1+\mathsf{L}_{j}+\frac{4\mathsf{L}_{j}+1}{3}(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} \bar{k} \\ &\times \mathsf{c}_{j}^{1+\frac{3}{4}(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} \left(\sum_{i=j}^{\ell} |\mathcal{J}_{i}|^{2}(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3}) \right) \\ &\cdot (\max_{k\in K_{j}} \deg g_{k}+1)^{(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})}. \end{aligned} \tag{23}$$

Note that these only depend on \mathbf{g} and J_1, \ldots, J_ℓ .

Apply Lemma 11 to f = p, $f_i = p_i$, $\epsilon = 3\varepsilon/2\ell$, $\eta = \varepsilon/2(\ell+2)$ to get polynomials h_1, \ldots, h_ℓ such that

$$p = h_1 + \dots + h_\ell, \qquad h_i \in \mathbb{R}[x_{J_i}], \qquad h_i(x) \ge \eta = \frac{\varepsilon}{2(\ell+2)} \text{ for } x \in S(\mathbf{g}),$$
 (24)

and

$$\overline{\deg} h_i \le \max(\overline{\deg} p_i, \overline{D}_{i,\ell}, \dots, \overline{D}_{\ell,\ell})_{J_i}.$$
(25)

In the dense setting, Baldi–Mourrain [1] construct a family of single-variable polynomials

$(\mathsf{h}_{t,m})_{(t,m)\in\mathbb{N}\times\mathbb{N}}$

providing useful approximation properties that we have adapted to the (separated-variables) sparse setting and collected in Lemma 14. To state this, we set, for all $j = 1, \ldots, \ell$ and for $(t_j, m_j) \in \mathbb{N} \times \mathbb{N}$ as well as for $s_j > 0$,

$$q_{j,t_j,m_j}(x) := \sum_{i \in K_i} \mathsf{h}_{t_i,m_i} \left(g_i(x) \right)^2 g_i(x), \tag{26}$$

$$f_{j,s_j,t_j,m_j}(x) := h_j(x) - s_j \, q_{j,t_j,m_j}(x).$$
(27)

Let us give an idea of what these functions do. The single-variable polynomial \mathbf{h}_{t_j,m_j} is of degree m_j and roughly speaking approximates the function that equals 1 on $(-\infty, 0)$ and $1/t_j$ elsewhere. Thus q_{j,t_j,m_j} almost vanishes (for large t_j) on $S(\mathbf{g}_{K_j})$, and outside of this domain it is roughly a sum of multiples of the negative parts of \mathbf{g}_{K_j} 's entries. The definition of f_{j,s_j,t_j,m_j} is engineered to obtain a polynomial that is almost equal to h_j in $S(\mathbf{g}_{K_j})$ yet remains positive throughout $[-1, 1]^n$. Instead of going into the details of the construction, we record the properties we need in the following lemma.

Lemma 14 (a version of [1, Props. 2.13, 3.1, and 3.2, Lem. 3.5]). Assume (2) and the Archimedean conditions (3) are satisfied. Then for each $j = 1, ..., \ell$ there are values s_j, t_j, m_j of the parameters involved in Definition (26) and Definition (27), such that the following holds with the shorthands

$$\hat{f}_j = f_{j,s_j,t_j,m_j}$$
 and $\hat{q}_j = s_j q_{j,t_j,m_j}$: (28)

i. [1, Prop. 3.1] gives

$$\hat{f}_j(x) \ge \frac{1}{2} \min_{y \in S(\mathbf{g}_{K_j})} h_j(y) \ge \frac{\eta}{2} = \frac{\varepsilon}{4(\ell+2)} \quad \text{for all} \quad x \in [1,1]^n.$$

ii. We have $\hat{q}_j \in \mathcal{Q}_{\mathbf{r},J_j}(\mathbf{g})$ for all multi-indices $\mathbf{r} = (r_1, \ldots, r_n)$ with

$$r_i \ge (2m_j + 1) \max_{k \in K_j} (\overline{\deg} g_k)_i$$

iii. [1, eq. (20)] gives the existence of a constant $C_m > 0$ such that

$$m_j \le C_m \mathsf{c}_j^{\frac{4}{3}} \bar{k}^{\frac{1}{3}} 2^{4\mathsf{L}_j} (\deg h_j)^{\frac{\mathtt{sL}_j}{3}} \left(\frac{\min_{x \in S(\mathbf{g})} h_j(x)}{\|h_j\|_{\infty}}\right)^{-\frac{\mathtt{sL}_j + 1}{3}}$$

iv. [1, eq. (16)] gives the existence of a constant $C_f > 0$ such that

$$\|\hat{f}_j\|_{\infty} \le C_f \|h_j\|_{\infty} 2^{3L_j} \bar{k} \mathsf{c}_j (\deg h_j)^{2L_j} \left(\frac{\min_{x \in S(\mathbf{g})} h_j(x)}{\|h_j\|_{\infty}}\right)^{-L_j}.$$

v. [1, eq. (17)] gives the existence of a constant $C_d > 0$ such that

$$\deg \hat{f}_j \le C_d 2^{4\mathsf{L}_j} \bar{k}^{\frac{1}{3}} \mathsf{c}_j^{\frac{4}{3}} (\max_{i \in K_j} \deg g_i) (\deg h_j)^{\frac{\mathsf{sL}_j}{3}} \left(\frac{\min_{x \in S(\mathbf{g})} h_j(x)}{\|h_j\|_{\infty}} \right)^{-\frac{\mathsf{sL}_j+1}{3}}$$

Item (ii) follows ² from deg $h_{t_j,m_j} = m_j$ and the definition of q_{j,t_j,m_j} . The proofs of the other items can be found in the indicated sources.

Take $s_j, t_j, m_j, \hat{f}_j, \hat{q}_j$ for $j = 1, ..., \ell$ satisfying the properties (i)-(v) collected in Lemma 14. Continuing with the proof of Theorem 4, denote

$$F_j \coloneqq \frac{\hat{f}_j - \min_{[-1,1]^n} \hat{f}_j}{\max_{[-1,1]^n} \hat{f}_j - \min_{[-1,1]^n} \hat{f}_j}$$

²This calculation is slightly different to the one in [1, Lem. 3.5] because the definition of q_{j,t_j,m_j} (or in their notations, f - p) differs from the one given there in that the functions h_j are squared here, an idea we take from the exposition of the results of [1] in the dissertation of L. Baldi and that is advantageous because then $q_{j,t_j,m_j} \in \mathcal{Q}_{\mathbf{r},J_j}(\mathbf{g})$ automatically. This requires taking m_j twice as large, and we absorb this difference into the constant C_m .

Since $\hat{f}_j \ge \varepsilon/4(\ell+2)$ on $[-1,1]^n$, we may apply Corollary 7 with $p = F_j$ to get that

$$F_j + \epsilon_j \in \mathcal{P}_{\mathbf{r}_j, J_j}(\{1 - x_i^2\}_{i \in J_j})$$

$$\tag{29}$$

as long as

$$\epsilon_j \ge 2|J_j|\pi^2 \left(\prod_{i \in J_j} ((\overline{\deg} F_j)_i + 1) \right) \max_{I = (i_1, \dots, i_n) \in \mathcal{I}_{F_j}} \left(2^{w(I)/2} \max_{1 \le k \le n} \frac{i_k^2}{(r_{j,k} + 2)^2} \right)$$
(30)

and (7) are verified. In this context, the condition (7) required in Corollary 7 is equivalent to the theorem's assumption (5); let us show how this works: First, using $i_k \leq (\overline{\deg} F_j)_k$, $\overline{\deg} F_j \leq \overline{\deg} \hat{f}_j \leq (\deg \hat{f}_j)\mathbf{1}$, Lemma 14(v), we get

$$2\pi^{2}|J_{j}|i_{k}^{2} \leq 2\pi^{2}|J_{j}| \left(\overline{\deg} F_{j}\right)_{k}^{2} \leq 2\pi^{2}|J_{j}| (\deg \hat{f}_{j})^{2}$$

$$\leq 2\pi^{2}|J_{j}| \left(C_{d}2^{4\mathsf{L}_{j}}\bar{k}^{\frac{1}{3}}\mathsf{c}_{j}^{\frac{4}{3}}(\max_{i\in K_{j}}\deg g_{i})(\deg h_{j})^{\frac{\mathsf{sL}_{j}}{3}} \left(\frac{\min_{x\in S(\mathbf{g})}h_{j}(x)}{\|h_{j}\|_{\infty}}\right)^{-\frac{4\mathsf{L}_{j}+1}{3}}\right)^{2}.$$

Now use equation (25) to get that this is

$$\leq 2\pi^{2} |J_{j}| \\ \cdot \left(C_{d} 2^{4L_{j}} \bar{k}^{\frac{1}{3}} \mathsf{c}_{j}^{\frac{4}{3}} (\max_{i \in K_{j}} \deg g_{i}) \max \left(\overline{\deg} p_{j}, \overline{D}_{j,\ell}, \dots, \overline{D}_{\ell,\ell} \right)^{\frac{8L_{j}}{3}} \left(\frac{\min_{x \in S(\mathbf{g})} h_{j}(x)}{\|h_{j}\|_{\infty}} \right)^{-\frac{4L_{j}+1}{3}} \right)^{2} \\ \leq 2\pi^{2} |J_{j}| \left(C_{d} 2^{4L_{j}} \bar{k}^{\frac{1}{3}} \mathsf{c}_{j}^{\frac{4}{3}} (\max_{i \in K_{j}} \deg g_{i}) \max \left(\overline{\deg} p_{j}, \overline{D}_{j,\ell}, \dots, \overline{D}_{\ell,\ell} \right)^{\frac{8L_{j}}{3}} \left(\frac{3^{\ell} \sum_{i} \|p_{i}\|_{\infty}}{\varepsilon/2(\ell+2)} \right)^{\frac{4L_{j}+1}{3}} \right)^{2},$$

where we have also used the fact that

$$\|h_j\|_{\infty} \le 3 \times 2^{\ell-1} \sum_{i=1}^{\ell} \|p_i\|_{\infty} \le 3^{\ell} \sum_{i=1}^{\ell} \|p_i\|_{\infty}$$
(31)

and the last estimate from (24). Next, use (10), $|\mathcal{J}_j| \leq |J_j|$, $\varepsilon_i - 2\eta_i = \frac{\ell+6}{2\ell(\ell+2)}\varepsilon$ to get

$$2\pi^{2}|J_{j}|i_{k}^{2} \leq 2\pi|J_{j}| \left(C_{d}2^{4\mathsf{L}_{j}}\bar{k}^{-\frac{1}{3}}\mathsf{c}_{j}^{\frac{4}{3}}(\max_{i\in K_{j}}g_{i})(\deg p_{j})^{\frac{\mathsf{sL}_{j}}{3}} \left(\frac{2C_{\mathsf{Jac}}|J_{j}|(3\sum_{i=1}^{\ell}\operatorname{Lip}p_{i})}{\frac{\ell+6}{2\ell(\ell+2)}\varepsilon}\right)^{\frac{\mathsf{sL}_{j}}{3}} \cdot \left(\frac{3^{\ell}\sum_{i}\|p_{i}\|_{\infty}}{\varepsilon/2(\ell+2)}\right)^{\frac{\mathsf{sL}_{j+1}}{3}} \right)^{2} \leq C_{j} \left(\left(\sum_{i=1}^{\ell}\|p_{i}\|_{\infty}\right)^{\frac{\mathsf{sL}_{j+1}}{3}} (\deg p_{j})^{\frac{\mathsf{sL}_{j}}{3}} \frac{\left(\sum_{i=1}^{\ell}\operatorname{Lip}p_{i}\right)^{\frac{\mathsf{sL}_{j}}{3}}}{\varepsilon^{\frac{(\mathsf{s}+4)\mathsf{L}_{j+1}}{3}}} \right)^{2} \leq (r_{j,k}+2)^{2},$$

where we have additionally used equation (22) and our assumption (5); this is precisely (7).

We would next like to show that

$$\hat{f}_j \in \mathcal{P}_{\mathbf{r}_j, J_j}(\{1 - x_i^2\}_{i \in J_j}).$$
 (32)

Let us first explain why this will be enough to prove the theorem. Once we have (32), by Lemma 15 \hat{f}_j is also contained in $\mathcal{Q}_{\mathbf{r}_j+\mathbf{2},J_j}(1-\sum_{i\in J_j}x_i^2)$, and it is our assumption (3) that $\mathcal{Q}_{\mathbf{r}_j+\mathbf{2},J_j}(1-\sum_{i\in J_j}x_i^2) \subseteq \mathcal{Q}_{\mathbf{r}_j+\mathbf{2},J_j}(\mathbf{g}_{K_j})$. In other words, we have

$$\hat{f}_j \in \mathcal{Q}_{\mathbf{r}_j+\mathbf{2},J_j}(\mathbf{g}_{K_j}).$$

By Lemma 14(ii), \hat{q}_j also belongs to $\mathcal{Q}_{\mathbf{r}_j+\mathbf{2},J_j}(\mathbf{g}_{K_j})$, so we can conclude that

$$h_j \in \mathcal{Q}_{\mathbf{r}_j+\mathbf{2},J_j}(\mathbf{g}_{K_j}),$$

which is equivalent to the conclusion of the theorem.

Thus we need to prove (32). Let us explain why, in order to obtain this conclusion we just need to show that

$$\frac{\eta}{2} = \frac{\varepsilon}{4(\ell+2)} \ge \left(\max_{[-1,1]^n} \hat{f}_j - \min_{[-1,1]^n} \hat{f}_j\right) 2|J_j| \pi^2 \left(\prod_{i \in J_j} \left((\overline{\deg} F_j)_i + 1\right)\right) \\ \times \max_{I=(i_1,\dots,i_n) \in \mathcal{I}_{F_j}} \left(2^{w(I)/2} \max_{1 \le k \le n} \frac{i_k^2}{(r_{j,k}+2)^2}\right).$$
(33)

Observe that (29) is equivalent to

$$\hat{f}_j - \min_{[-1,1]^n} \hat{f}_j + \epsilon_j (\max_{[-1,1]^n} \hat{f}_j - \min_{[-1,1]^n} \hat{f}_j) \in \mathcal{P}_{\mathbf{r}_j,J_j}(\{1 - x_i^2\}_{i \in J_j})$$
(34)

If (33) were true, we would then have

$$\begin{split} \hat{f}_{j} &\geq \hat{f}_{j} - \min_{[-1,1]^{n}} \hat{f}_{j} + \frac{\varepsilon}{4(\ell+2)} \\ &\geq \hat{f}_{j} - \min_{[-1,1]^{n}} \hat{f}_{j} + (\max_{[-1,1]^{n}} \hat{f}_{j} - \min_{[-1,1]^{n}} \hat{f}_{j}) \ 2|J_{j}|\pi^{2} \left(\prod_{i \in J_{j}} ((\overline{\deg} F_{j})_{i} + 1) \right) \\ &\times \max_{I = (i_{1}, \dots, i_{n}) \in \mathcal{I}_{F_{j}}} \left(2^{w(I)/2} \max_{1 \leq k \leq n} \frac{i_{k}^{2}}{(r_{j,k} + 2)^{2}} \right). \end{split}$$

So in view of (30) and (34), we would indeed have $\hat{f}_j \in \mathcal{P}_{\mathbf{r}_j, J_j}(\{1 - x_i^2\}_{i \in J_j})$, which is (32). Let us now collect some preliminary estimates that will help us to prove (33). For $I \in \mathcal{I}_{F_j}$ we have $w(I) \leq |J_j|$ so we estimate

$$2^{w(I)/2} \le 2^{|J_j|/2}.$$
(35)

We also estimate

$$\frac{i_k^2}{(r_{j,k}+2)^2} \le \frac{\left(\overline{\deg}\,\hat{f}_j\right)_k^2}{\min_{1\le l\le n}(r_{j,l}+2)^2}.$$
(36)

Now we will estimate $\max_{[-1,1]^n} \hat{f}_j - \min_{[-1,1]^n} \hat{f}_j$ from above. Using Lemma 14(i) and (iv), we get

$$\max_{[-1,1]^n} \hat{f}_j - \min_{[-1,1]^n} \hat{f}_j \le C_f \|h_j\|_{\infty} 2^{3\mathsf{L}_j} \bar{k} \mathsf{c}_j (\deg h_j)^{2\mathsf{L}_j} \left(\frac{\min_{x \in S(\mathbf{g})} h_j(x)}{\|h_j\|_{\infty}}\right)^{-\mathsf{L}_j} - \frac{\varepsilon}{4(\ell+2)}$$

Use (24), (25) and (31) to see that this is

$$\max_{[-1,1]^{n}} \hat{f}_{j} - \min_{[-1,1]^{n}} \hat{f}_{j} \\
\leq C_{f} \left(3^{\ell} \sum_{i} \|p_{i}\|_{\infty} \right) 2^{3L_{j}} \bar{k} c_{j} \max(\deg p_{j}, |\bar{D}_{j,\ell}|, \dots, |\bar{D}_{\ell,\ell}|)^{2L_{j}} \\
\cdot \left(\frac{\varepsilon}{2(\ell+2)3^{\ell} \sum_{i} \|p_{i}\|_{\infty}} \right)^{-L_{j}} \\
= C_{f} \left(3^{\ell} \sum_{i} \|p_{i}\|_{\infty} \right) 2^{3L_{j}} \bar{k} c_{j} \max(\deg p_{j}, |\mathcal{J}_{j}|D_{j,\ell}, \dots, |\mathcal{J}_{\ell}|D_{\ell,\ell})^{2L_{j}} \\
\cdot \left(\frac{\varepsilon}{2(\ell+2)3^{\ell} \sum_{i} \|p_{i}\|_{\infty}} \right)^{-L_{j}}$$
(37)

For the last line, we have used the definition of $\overline{D}_{l,m}$ as in Lemma 11.

Additionally, we obtain the following estimate

$$\overline{\deg}\,\hat{f}_{j} \leq \max\left(\overline{\deg}\,h_{j},\overline{\deg}\,\hat{q}_{j}\right) \\
\leq \max\left(\overline{\deg}\,p_{j},\bar{D}_{j,\ell},\ldots,\bar{D}_{\ell,\ell},(2m_{j}+1)\max_{k\in K_{j}}\overline{\deg}\,g_{k}\right) \\
\leq \max\left(\overline{\deg}\,p_{j},\bar{D}_{j,\ell},\ldots,\bar{D}_{\ell,\ell},\left(2\left(C_{m}\mathsf{c}_{j}^{\frac{3}{4}}\bar{k}^{-\frac{1}{3}}2^{4\mathsf{L}_{j}}(\deg h_{j})^{\frac{\mathsf{sL}_{j}}{3}}\left(\frac{\min_{x\in S(\mathbf{g})}h_{j}(x)}{\|h_{j}\|_{\infty}}\right)^{-\frac{4\mathsf{L}_{j}+1}{3}}\right) + 1)\max_{k\in K_{j}}\overline{\deg}\,g_{k}\right) \\
\leq \max\left(\overline{\deg}\,p_{j},\bar{D}_{j,\ell},\ldots,\bar{D}_{\ell,\ell},\left(C_{m}2^{4\mathsf{L}_{j}+1}\mathsf{c}^{\frac{3}{4}}\bar{k}^{-\frac{1}{3}}(\deg h_{j})^{\frac{\mathsf{sL}_{j}}{3}}\left(\frac{3^{\ell}\sum_{i}\|p_{i}\|_{\infty}}{\varepsilon/2(\ell+2)}\right)^{\frac{4\mathsf{L}_{j}+1}{3}} + 1\right)\max_{k\in K_{j}}\overline{\deg}\,g_{k}\right). \tag{38}$$

The first inequality comes from (27), the second one from (25) and Lemma 14(ii), the third one from Lemma 14(iii), and the last one from (24), (31), and (25). Compare with Lemma 14(v).

With those estimates under our belt, we now turn to showing that (33) is true. Using (37), (35), (36), as well as $\overline{\deg} F_j \leq \overline{\deg} \hat{f}_j$, we can start to estimate the right-hand side of (33) by

$$\begin{split} (\max_{[-1,1]^n} \hat{f}_j - \min_{[-1,1]^n} \hat{f}_j) \ 2|J_j| \pi^2 \left(\prod_{i \in J_j} ((\overline{\deg} F_j)_i + 1) \right) \\ & \times \max_{I = (i_1, \dots, i_n) \in \mathcal{I}_{F_j}} \left(2^{w(I)/2} \max_{1 \le k \le n} \frac{i_k^2}{(r_{j,k} + 2)^2} \right) \\ & \leq C_f \left(3^\ell \sum_i \|p_i\|_{\infty} \right) 2^{3\mathsf{L}_j} \bar{k} \mathsf{c}_j \max(\deg p_j, |\mathcal{J}_j| D_{j,\ell}, \dots, |\mathcal{J}_\ell| D_{\ell,\ell})^{2\mathsf{L}_j} \left(\frac{\varepsilon}{2(\ell+2)3^\ell \sum_i \|p_i\|_{\infty}} \right)^{-\mathsf{L}_j} \\ & \times 2|J_j| \pi^2 \left(\max_{1 \le i \le n} (\overline{\deg} \hat{f}_j)_i + 1 \right)^{|J_j|} \frac{2^{|J_j|/2} \max_{1 \le k \le n} (\overline{\deg} \hat{f}_j)_i^2}{\min_{1 \le k \le n} (r_{j,k} + 2)^2}. \end{split}$$

Next we denote

$$m_{j,\ell} = \max(\deg p_j, |\mathcal{J}_j| D_{j,\ell}, \dots, |\mathcal{J}_\ell| D_{\ell,\ell})$$

and we reorganize and consolidate the terms and then we use (38) to see that this is

$$\begin{split} &\leq \varepsilon^{-\mathsf{L}_{j}} C_{f} |J_{j}| \pi^{2} \left(3^{\ell} \sum_{i} ||p_{i}||_{\infty} \right)^{\mathsf{L}_{j}+1} 2^{4\mathsf{L}_{j} + \frac{|J_{j}|}{2} + 1} (\ell + 2)^{\mathsf{L}_{j}} \bar{k} \mathsf{c}_{j} m_{j,\ell}^{2\mathsf{L}_{j}} \\ &\qquad \times \left(\max(\overline{\deg} f_{j})_{i} + 1 \right)^{|J_{j}|+2} \frac{1}{\min_{1 \leq k \leq n} (r_{j,k} + 2)^{2}} \\ &\leq \varepsilon^{-\mathsf{L}_{j}} C_{f} |J_{j}| \pi^{2} \left(3^{\ell} \sum_{i} ||p_{i}||_{\infty} \right)^{\mathsf{L}_{j}+1} 2^{4\mathsf{L}_{j} + \frac{|J_{j}|}{2} + 1} (\ell + 2)^{\mathsf{L}_{j}} \bar{k} \mathsf{c}_{j} m_{j,\ell}^{2\mathsf{L}_{j}} \\ &\qquad \times \left(\max\left(\overline{\deg} p_{j}, \bar{D}_{j,\ell}, \dots, \bar{D}_{\ell,\ell}, \right. \right. \\ \left(C_{m} 2^{4\mathsf{L}_{j}+1} \mathsf{c}^{\frac{3}{4}} \bar{k}^{-\frac{1}{3}} (\deg h_{j})^{\frac{\mathsf{s}\mathsf{L}_{j}}{3}} \left(\frac{3^{\ell} \sum_{i} ||p_{i}||_{\infty}}{\varepsilon/2(\ell + 2)} \right)^{\frac{4\mathsf{L}_{j}+1}{3}} + 1 \right) \\ &\qquad \times \max_{k \in K_{j}} \overline{\deg} g_{k} \right) + 1 \Big)^{|J_{j}|+2} \frac{1}{\min_{1 \leq k \leq n} (r_{j,k} + 2)^{2}} \\ &\leq \varepsilon^{-\mathsf{L}_{j}} C_{f} |J_{j}| \pi^{2} \left(3^{\ell} \sum_{i} ||p_{i}||_{\infty} \right)^{\frac{\mathsf{L}_{j}+1}{2}} 2^{4\mathsf{L}_{j} + \frac{|J_{j}|}{2} + 1} (\ell + 2)^{\mathsf{L}_{j}} \bar{k} \mathsf{c}_{j} m_{j,\ell}^{2\mathsf{L}_{j}} \\ &\qquad \times \left(\max\left(\overline{\deg} p_{j}, \bar{D}_{j,\ell}, \dots, \bar{D}_{\ell,\ell}, \right)^{\mathsf{L}_{j}+1} 2^{4\mathsf{L}_{j} + \frac{|J_{j}|}{2} + 1} (\ell + 2)^{\mathsf{L}_{j}} \bar{k} \mathsf{c}_{j} m_{j,\ell}^{2\mathsf{L}_{j}} \\ &\qquad \times \left(\max\left(\overline{\deg} p_{j}, \bar{D}_{j,\ell}, \dots, \bar{D}_{\ell,\ell}, \right)^{\mathsf{L}_{j}+1} 2^{4\mathsf{L}_{j} + \frac{|J_{j}|}{2} + 1} (\ell + 2)^{\mathsf{L}_{j}} \bar{k} \mathsf{c}_{j} m_{j,\ell}^{2\mathsf{L}_{j}} \\ &\qquad \times \left(\max\left(\overline{\deg} p_{j}, \bar{D}_{j,\ell}, \dots, \bar{D}_{\ell,\ell}, \right)^{\mathsf{L}_{j}+1} 2^{4\mathsf{L}_{j} + \frac{|J_{j}|}{2} + 1} (\ell + 2)^{\mathsf{L}_{j}} \bar{k} \mathsf{c}_{j} m_{j,\ell}^{2\mathsf{L}_{j}} \right)^{\mathsf{L}_{j}+1} \right) \\ &\qquad \times \max_{k \in K_{j}} \overline{\deg} g_{k} \right) + 1 \right)^{|J_{j}|+2} \frac{1}{\min_{1 \leq k \leq n} (r_{j,k} + 2)^{2}}. \end{split}$$

Now use (10) as well as $\varepsilon_i - 2\eta_i = \frac{\ell+6}{2\ell(\ell+2)}\varepsilon$ to see that the above is bounded by

$$\begin{split} \varepsilon^{-\mathsf{L}_{j}} C_{f} |J_{j}| \pi^{2} \left(3^{\ell} \sum_{i} ||p_{i}||_{\infty} \right)^{\mathsf{L}_{j}+1} 2^{4\mathsf{L}_{j} + \frac{|J_{j}|}{2} + 1} (\ell + 2)^{\mathsf{L}_{j}} \bar{k} \mathsf{c}_{j} \\ & \times \left(\max \left[\deg p_{j}, |\mathcal{J}_{j}|^{2} \frac{2\ell(\ell + 2)}{\ell + 6} \frac{2C_{\mathrm{Jac}} \sum_{k=j}^{\ell} \operatorname{Lip} \hat{f}_{k}}{\varepsilon} + 1, \ldots, \right. \\ & \left| \mathcal{J}_{\ell} \right|^{2} \frac{2\ell(\ell + 2)}{\ell + 6} \frac{2C_{\mathrm{Jac}} \sum_{i} ||p_{i}||_{\infty}}{\varepsilon} \right)^{\frac{4\mathsf{L}_{j}+1}{3}} + 1, \\ & \left(C_{m} 2^{4\mathsf{L}_{j}+1} \mathsf{c}^{\frac{3}{4}} \bar{k}^{-\frac{1}{3}} \left(\frac{3^{\ell} \sum_{i} ||p_{i}||_{\infty}}{\varepsilon/2(\ell + 2)} \right)^{\frac{4\mathsf{L}_{j}+1}{3}} + 1 \right) \max_{k \in K_{j}} \overline{\deg} g_{k} \right] \\ & + 1 \Big)^{(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} \frac{1}{\min_{1 \le k \le n}(r_{j,k} + 2)^{2}} \\ \leq \varepsilon^{-\mathsf{L}_{j}-\frac{4\mathsf{L}_{j}+1}{3}} (2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} C_{f}(C_{\mathrm{Jac}}C_{m})^{(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} |J_{j}| \pi^{2} \left(3^{\ell} \sum_{i} ||p_{i}||_{\infty} \right)^{\mathsf{L}_{j}+1} \\ & \times 2^{4\mathsf{L}_{j}+\frac{|J_{j}|}{2}} + 1 + (1+\frac{4\mathsf{L}_{j}+1}{3})(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} (\ell + 2)^{1+\mathsf{L}_{j}+\frac{4\mathsf{L}_{j}+1}{3}} (2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3}) \bar{k}} \\ & \times \mathsf{C}_{j}^{1+\frac{3}{4}} (2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} \left(\sum_{i=j}^{\ell} |\mathcal{J}_{i}|^{2(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} \right) \\ & \times (3\deg p_{j} \sum_{i=1}^{\ell} \operatorname{Lip} p_{i})^{(2\mathsf{L}_{j}+|J_{j}|+2)(1+\frac{8\mathsf{L}_{j}}{3})} \frac{1}{\min_{1 \le k \le n}(r_{j,k} + 2)^{2}}. \end{split}$$

Finally use (23) and then (4) to get that the above is less than

$$\mathbf{C}_{j} \frac{\varepsilon^{-\mathsf{L}_{j} - \frac{4\mathsf{L}_{j} + 1}{3}(2\mathsf{L}_{j} + |J_{j}| + 2)(1 + \frac{8\mathsf{L}_{j}}{3})}{\min_{1 \le k \le n} (r_{j,k} + 2)^{2}} \leq \frac{\varepsilon}{4(\ell + 2)}.$$

This shows that (33) holds, and hence also (32), which proves the theorem.

Lemma 15 ([1, Lemma 3.8]). Let $J \subset \{1, \ldots, n\}$, and let $\mathbf{r} = (r_1, \ldots, r_n)$ be a multi-index such that $r_i > 0$ only if $i \in J$. The quadratic module $\mathcal{Q}_{\mathbf{r+2},J}(1 - \sum_{i \in J} x_i^2)$ contains the preordering $\mathcal{P}_{\mathbf{r},J}(\{1 - x_i^2\}_{i \in J})$,

$$\mathcal{P}_{\mathbf{r},J}(\{1-x_i^2\}_{i\in J})\subseteq \mathcal{Q}_{\mathbf{r}+2,J}(1-\sum_{i\in J}x_i^2).$$

Proof. This follows from

$$1 \pm x_i = \frac{1}{2}(1 - x_i^2 + (1 \pm x_i)^2) = \frac{1}{2}((1 - \|x\|^2) + \sum_{\substack{j \in J \\ j \neq i}} x_j^2 + (1 \pm x_i)^2)$$

and

$$1 - x_i^2 = (1 - x_i)(1 + x_i)$$

= $\frac{1}{4}(1 - ||x||_J^2) \left(2\sum_{j \neq i} x_j^2 + (1 - x_i)^2 + (1 + x_i)^2\right)$
+ $\frac{1}{4}(1 - ||x||_J^2)^2 + \left(\sum_{j \neq i} x_j^2\right)^2 + \sum_{j \neq i} x_j^2((1 - x_i)^2 + (1 + x_i)^2).$

The increase of **2** in **r** stems from the fact that $deg(1 - x_i^2) = 2$ while the degree of the right-hand side above is 4.

4.3 An asymptotic lemma

Lemma 16. For a, b, c, d, p, q > 0, with $cq - ap \neq 0$,

$$\limsup_{\varepsilon \searrow 0} \left| \frac{\log \left[\binom{a+b\varepsilon^{-p}}{b\varepsilon^{-p}} \middle/ \binom{c+d\varepsilon^{-q}}{d\varepsilon^{-q}} \right]}{(cq-ap)\log \varepsilon} \right| < +\infty.$$

Proof. Using the Stirling approximation, which states that there exist $C' \ge 0$ such that

$$|\log(n!) - n\log n + n| \le C'\log n$$

for all large n > 0, we get that,

$$\begin{split} \limsup_{\varepsilon \searrow 0} \left| \frac{\log \left[\binom{a+b\varepsilon^{-p}}{b\varepsilon^{-p}} \right] / \binom{c+d\varepsilon^{-q}}{d\varepsilon^{-q}} \right]}{(cq-ap)\log\varepsilon} \right| \\ &= \limsup_{\varepsilon \searrow 0} \left| \frac{\log \left[\frac{(a+b\varepsilon^{-p})!}{a!(b\varepsilon^{-p})!} / \frac{(c+d\varepsilon^{-q})!}{c!(d\varepsilon^{-q})!} \right]}{(cq-ap)\log\varepsilon} \right| \\ &= \limsup_{\varepsilon \searrow 0} \left| \frac{\log(a+b\varepsilon^{-p})! - \log a! - \log(b\varepsilon^{-p})! - \log(c+d\varepsilon^{-q})! + \log c! + \log(d\varepsilon^{-q})!}{(cq-ap)\log\varepsilon} \right| \\ &= \limsup_{\varepsilon \searrow 0} \left| \frac{\log(a+b\varepsilon^{-p})! - \log(b\varepsilon^{-p})! - \log(c+d\varepsilon^{-q})! + \log(d\varepsilon^{-q})!}{(cq-ap)\log\varepsilon} \right| \\ &= \limsup_{\varepsilon \searrow 0} \left| \frac{\log(a+b\varepsilon^{-p})! - \log(b\varepsilon^{-p})! - \log(c+d\varepsilon^{-q})! + \log(d\varepsilon^{-q})!}{(cq-ap)\log\varepsilon} \right| \end{split}$$
(39)

$$+ (c + d\varepsilon^{-q})\log(c + d\varepsilon^{-q}) + d\varepsilon^{-q}\log(d\varepsilon^{-q})$$

$$\tag{40}$$

$$-\log a! + \log c! - (a + b\varepsilon^{-p}) + (b\varepsilon^{-p}) + (c + d\varepsilon^{-q}) - (d\varepsilon^{-q}))$$

$$\tag{41}$$

$$+ O\left(\left|\log(a+b\varepsilon^{-p})\right| + \left|\log(b\varepsilon^{-p})\right| + \left|\log(c+d\varepsilon^{-q})\right| + \left|\log(d\varepsilon^{-q})\right|\right)\right|$$
(42)

Notice that the terms in (41) are asymptotically much smaller than the denominator, which tends to $+\infty$. The last line (42) is, as $\varepsilon \searrow 0$, of the order of

$$\frac{O(|\log(a+b\varepsilon^{-p})|+|\log(b\varepsilon^{-p})|+|\log(c+d\varepsilon^{-q})|+|\log(d\varepsilon^{-q})|)}{|cq-ap||\log\varepsilon|}$$
$$=\frac{O(2|\log b|+2p|\log\varepsilon|+2|\log d|+2q|\log\varepsilon|)}{|cq-ap||\log\varepsilon|}$$
$$=\frac{(p+q)O(|\log\varepsilon|)}{|cq-ap||\log\varepsilon|}=O\left(\frac{p+q}{|cq-ap|}\right)=O(1).$$

Let us now show that the remaining two lines (39)-(40) tend to 1 in the limit. Now,

$$\lim_{\varepsilon \searrow 0} b\varepsilon^{-p} \left(\log(a + b\varepsilon^{-p}) - \log(b\varepsilon^{-p}) \right) = a \quad \text{and} \quad \lim_{\varepsilon \searrow 0} d\varepsilon^{-p} \left(\log(c + d\varepsilon^{-p}) - \log(d\varepsilon^{-p}) \right) = c,$$

so we get

$$\lim_{\varepsilon \searrow 0} \frac{1}{(cq-ap)\log\varepsilon} \Big((a+b\varepsilon^{-p})\log(a+b\varepsilon^{-p}) - a\log a - b\varepsilon^{-p}\log(b\varepsilon^{-p}) \\ - (c+d\varepsilon^{-q})\log(c+d\varepsilon^{-q}) + c\log c + d\varepsilon^{-q}\log(d\varepsilon^{-q}) \Big) \\ = \lim_{\varepsilon \searrow 0} \frac{-a\log a + c\log c + a - c + a\log(a+b\varepsilon^{-p}) - c\log(c+d\varepsilon^{-q})}{(cq-ap)\log\varepsilon} \\ = \lim_{\varepsilon \searrow 0} \frac{a\log(a+b\varepsilon^{-p}) - c\log(c+d\varepsilon^{-q})}{(cq-ap)\log\varepsilon}.$$

In this quotient, both the numerator and the denominator tend to $\pm \infty$, so we can apply a version of the l'Hôpital rule, which states that, if the limit of the quotient of their derivatives exists, then the original limit above equals that limit. Taking the limit of the quotient of the derivatives gives

$$\lim_{\varepsilon \searrow 0} \frac{\frac{cdq\varepsilon^{-1-q}}{c+d\varepsilon^{-q}} - \frac{abp\varepsilon^{-1-p}}{a+b\varepsilon^{-p}}}{(cq-ap)/\varepsilon}$$

$$= \lim_{\varepsilon \searrow 0} \frac{cdq\varepsilon^{-q}(a+b\varepsilon^{-p}) - abp\varepsilon^{-p}(c+d\varepsilon^{-q})}{(cq-ap)(a+b\varepsilon^{-p})(c+d\varepsilon^{-q})}$$

$$= \lim_{\varepsilon \searrow 0} \frac{cdq(a\varepsilon^{p}+b) - abp(c\varepsilon^{q}+d)}{(cq-ap)(a\varepsilon^{q}+b)(c\varepsilon^{p}+d)}$$

$$= \lim_{\varepsilon \searrow 0} \frac{cdqb - abpd}{(cq-ap)bd} = 1.$$

References

- [1] Lorenzo Baldi and Bernard Mourrain. On the effective Putinar's Positivstellensatz and moment approximation. *Mathematical Programming*, pages 1–33, 2022.
- [2] Andrew C Doherty and Stephanie Wehner. Convergence of SDP hierarchies for polynomial optimization on the hypersphere. arXiv preprint arXiv:1210.5048, 2012.
- [3] Kun Fang and Hamza Fawzi. The sum-of-squares hierarchy on the sphere and applications in quantum information theory. *Mathematical Programming*, 190(1):331–360, 2021.
- [4] David Grimm, Tim Netzer, and Markus Schweighofer. A note on the representation of positive polynomials with structured sparsity. Archiv der Mathematik, 89(5):399–403, 2007.
- [5] Dunham Jackson. The theory of approximation, volume 11 of Colloquium Publications. American Mathematical Society, 1930.
- [6] Samuel Karlin and Lloyd S Shapley. Geometry of moment spaces, volume 12. American Mathematical Society, 1953.
- [7] Felix Kirschner and Etienne De Klerk. Convergence rates of RLT and Lasserre-type hierarchies for the generalized moment problem over the simplex and the sphere. *Optimization Letters*, pages 1–18, 2022.
- [8] Jean B Lasserre. Global optimization with polynomials and the problem of moments. SIAM Journal on optimization, 11(3):796–817, 2001.
- [9] Jean B Lasserre. Convergent SDP-relaxations in polynomial optimization with sparsity. SIAM Journal on Optimization, 17(3):822–843, 2006.
- [10] Jean Bernard Lasserre. Moments, positive polynomials and their applications, volume 1. World Scientific, 2009.
- [11] Monique Laurent. Sums of squares, moment matrices and optimization over polynomials. In Emerging applications of algebraic geometry, pages 157–270. Springer, 2009.
- [12] Monique Laurent and Lucas Slot. An effective version of Schmüdgen's Positivstellensatz for the hypercube. Optimization Letters, 2023.
- [13] V. Magron and J. Wang. Sparse polynomial optimization: theory and practice. Series on Optimization and Its Applications, World Scientific Press, 2023. Accepted for publication.
- [14] Ngoc Hoang Anh Mai and Victor Magron. On the complexity of putinar-vasilescu's positivstellensatz. Journal of Complexity, page 101663, 2022.
- [15] Ngoc Hoang Anh Mai, Victor Magron, and Jean Lasserre. A sparse version of Reznick's Positivstellensatz. Mathematics of Operations Research, 2022.
- [16] Jiawang Nie and Markus Schweighofer. On the complexity of Putinar's Positivstellensatz. Journal of Complexity, 23(1):135–150, 2007.
- [17] P.A. Parrilo. Approximation quality of SOS relaxations. Talk at ICCOPT 2013, 2013.
- [18] Pablo A Parrilo. Semidefinite programming relaxations for semialgebraic problems. Mathematical programming, 96(2):293–320, 2003.
- [19] Mihai Putinar. Positive polynomials on compact semi-algebraic sets. Indiana University Mathematics Journal, 42(3):969–984, 1993.
- [20] Bruce Reznick. Uniform denominators in Hilbert's seventeenth problem. Mathematische Zeitschrift, 220(1):75–97, 1995.
- [21] Konrad Schmüdgen. The K-moment problem for compact semi-algebraic sets. Mathematische Annalen, 289(1):203–206, 1991.

- [22] Markus Schweighofer. On the complexity of Schmüdgen's Positivstellensatz. *Journal of Complexity*, 20(4):529–543, 2004.
- [23] Lucas Slot. Sum-of-squares hierarchies for polynomial optimization and the Christoffel–Darboux kernel. SIAM Journal on Optimization, 32(4):2612–2635, 2022.
- [24] Lucas Slot and Monique Laurent. Sum-of-squares hierarchies for binary polynomial optimization. Mathematical Programming, pages 1–40, 2022.
- [25] Alexander Weiße, Gerhard Wellein, Andreas Alvermann, and Holger Fehske. The kernel polynomial method. *Reviews of modern physics*, 78(1):275, 2006.