Twin Brownian particle method for the study of Oberbeck-Boussinesq fluid flows

Jiawei Li^{*}, Zhongmin Qian[†] and Mingyu Xu[‡]

Abstract

We establish stochastic functional integral representations for solutions of Oberbeck-Boussinesq equations in the form of McKean-Vlasov-type mean field equations, which can be used to design numerical schemes for calculating solutions and for implementing Monte-Carlo simulations of Oberbeck-Boussinesq flows. Our approach is based on the duality of conditional laws for a class of diffusion processes associated with solenoidal vector fields, which allows us to obtain a novel integral representation theorem for solutions of some linear parabolic equations in terms of the Green function and the pinned measure of the associated diffusion. We demonstrate via numerical experiments the efficiency of the numerical schemes, which are capable of revealing numerically the details of Oberbeck-Boussinesq flows within their thin boundary layer, including Bénard's convection feature.

Key words: Boussinesq approximation, conditional laws, diffusion processes, incompressible fluid flow, Monte-Carlo simulation, random vortex method.

MSC classifications: 76M35, 76M23, 60H30, 65C05, 68Q10

[∗]School of Mathematics, University of Edinburgh, James Clerk Maxwell Building, Peter Guthrie Tait Rd, Edinburgh, United Kingdom, EH9 3FD. Email: jiawei.li@ed.ac.uk

[†]Mathematical Institute, University of Oxford, Oxford, United Kingdom, OX2 6GG, and Oxford Suzhou Centre for Advanced Research, Suzhou, China. Email: qianz@maths.ox.ac.uk

[‡]Department of Mathematics, Fudan University, Shanghai, China. Email: xumy@fudan.edu.cn

Contents

1 Introduction

By Oberbeck-Boussinesq flows, we mean fluid flows governed by approximation equations of motion for heat-conducting fluid flows (see e.g. Landau-Lifshitz $[44,$ Chapters II and V $]$), for details, the reader may refer to Chandrasekhar [\[12\]](#page-41-0) and Drazin-Reid [\[21,](#page-41-1) Chapter 2]. These approximation equations in the form of partial differential equations were proposed independently by Oberbeck [\[54\]](#page-43-1) and Boussinesq [\[8\]](#page-41-2), cf. also Rayleigh [\[62\]](#page-44-0), in which the approximation equations were derived under the assumption that the fluid density ρ is almost constant.

The primary goal of the paper is to develop Monte-Carlo-type numerical methods for the study of Oberbeck-Boussinesq flows based on exact stochastic formulations of the Oberbeck-Boussinesq flows to be established in the paper. This will be achieved by establishing the functional integral representations for solutions of the Oberbeck-Boussinesq equations. These stochastic integral representations, as well as the approach presented in this work, appear to have independent interests on their own. Indeed we hope these ideas will be useful in the study, theoretically and numerically, of other non-linear systems of partial differential equations.

The Oberbeck-Boussinesq equations are composed of the Navier-Stokes equations for the velocity $u(x, t)$ coupled with a transport equation for the temperature $\theta(x, t)$:

$$
\frac{\partial}{\partial t}u + (u \cdot \nabla)u = \nu \Delta u - \nabla P + f(\theta),\tag{1.1}
$$

$$
\nabla \cdot u = 0,\tag{1.2}
$$

and

$$
\frac{\partial}{\partial t}\theta + (u \cdot \nabla)\theta = \kappa \Delta \theta, \tag{1.3}
$$

where $u(x, t)$ is subject to the no-slip condition, i.e. $u(x, t)$ vanishes along a solid boundary, and $\theta(x, t)$ the temperature generally possesses a non-trivial value since the heat is supplied to the fluid system from the solid boundary. $\nu > 0$ is the kinematic viscosity, and $\kappa > 0$ is the thermal diffusivity. Hence the Oberbeck-Boussinesq equations are more sophisticated than the Navier-Stokes equations alone. Nevertheless, they still serve as approximation equations for the much more complicated motion equations governing viscous fluid flows with thermal conduction when the fluid density remains relatively constant during the heating process. The Oberbeck-Boussinesq model provides a good explanation for the regular cellular pattern of the fluid motion - when the fluid at the bottom receives heat and expands with increasing temperature to the level at which the buoyancy dominates over the viscosity effect, a phenomenon called the Bénard convection occurs, reported first by Bénard [\[7\]](#page-41-3).

The Oberbeck-Boussinesq model has been investigated as one of the very successful examples in the theory of hydrodynamic stability, see Chandrasekhar [\[12\]](#page-41-0), Joseph [\[36,](#page-42-0) [37,](#page-42-1) [38\]](#page-42-2) and Drazin-Reid [\[21\]](#page-41-1). In recent years, with the development of computational power, various numerical methods have been employed in the study of various fluid flows, including the questions of hydrodynamic stability and transition to turbulence, see [\[20\]](#page-41-4) for example.

In the study of fluid dynamics over a century, statistical and probabilistic ideas have penetrated gradually into the research area of fluid mechanics, though at the beginning, only very primitive concepts in statistics were borrowed to the study of isotropic and homogeneous turbulent flows, as seen in the seminal work by Taylor [\[67\]](#page-44-1). In fact, in the statistical theory of turbulence put forward by Taylor [\[67\]](#page-44-1), Von Kármán [\[68\]](#page-44-2), and etc., only the idea of averaging was adopted to describe the mean motions of turbulent flows. Later in K41 theory, Kolmogorov [\[41,](#page-43-2) [42\]](#page-43-3) applied more sophisticated concept of conditional laws for random fields and introduced the concept of locally isotropic turbulent flows. Moreover, the ideas of random walks and diffusions in fluid flows emerged as means to describe turbulent flows. Taylor [\[66\]](#page-44-3) formally introduced Brownian fluid particles into the study of fluid dynamics and made the study of diffusions in turbulence a useful tool in the description of various aspects of turbulent flows, see [\[58\]](#page-43-4), [\[52\]](#page-43-5) and

[\[26\]](#page-42-3) for a very detailed review. These probabilistic studies of fluid dynamics, i.e. statistical fluid mechanics, occurred before the major development in probability theory, such as the creation of stochastic calculus by Doob, Itô, etc. Since then, stochastic calculus has been gradually applied to the study of fluid mechanics too. For example, in LeJan and Sznitman [\[45\]](#page-43-6), the energy dissipation cascades in turbulence were interpreted in terms of random walks, while in LeJan and Raimond [\[46,](#page-43-7) [47\]](#page-43-8), Brownian particles were studied.

Particularly in the study of incompressible fluid flows, vorticity has been singled out as a crucial fluid dynamical variable, in addition to the flow velocity. It seems that Helmholtz [\[33\]](#page-42-4) was the first person who emphasised the significance of vortex motions in the study of fluid mechanics. Since then, motions of vortices in fluid flows have been studied throughout the history of fluid dynamics. The random vortex method, originated by Chorin [\[13\]](#page-41-5), is a probabilistic method for incompressible flows developed based on the following simple but fundamental observation. The motion of vortices in turbulent flows in nature exhibits (approximately) statistical independence, leading to a potentially easier description of the fluid flow via its vortex motion. The vorticity equations governing the evolution of the vorticity, which appears as a parabolic transport equation, demonstrate that the vortices are transported along the fluid flow. The random vortex method, though limited to two-dimensional fluid flows, based on the exact fluid dynamic equations, was first discovered by Goodman [\[31\]](#page-42-5) (see also Long [\[49\]](#page-43-9)). The relatively novel applications of vortex dynamics in numerical schemes for solving fluid dynamic equations have been rapidly established as an important branch of fluid mechanics, see e.g. [\[5,](#page-41-6) [6\]](#page-41-7), [\[18\]](#page-41-8), [\[52\]](#page-43-5) and etc. for excellent reviews on the vortex method.

According to Feynman [\[27\]](#page-42-6) and Kac [\[39\]](#page-42-7), it is possible to express solutions of certain linear parabolic and elliptic equations in terms of path integrals. In fact, the idea of Feynman-Kac has been generalised to a class of semi-linear parabolic equations by Pardoux and Peng in [\[56\]](#page-43-10), where a nonlinear version of Feynman-Kac formula was obtained via backward stochastic differential equations. In recent years, the random vortex methods have been greatly enhanced under the name of the stochastic Lagrangian (vorticity) approach and great progress has been made in Holm [\[34\]](#page-42-8), Busnello [\[9\]](#page-41-9), Busnello, Flandoli and Romito [\[10\]](#page-41-10), Constantin [\[15,](#page-41-11) [16\]](#page-41-12). In particular Constantin and Iyer [\[17\]](#page-41-13) obtained a stochastic integral representation for solutions of the Navier-Stokes equations, cf. Zhang [\[75\]](#page-44-4) as well in which stochastic integral representation has been established for solutions to backward stochastic Navier-Stokes equations. Their representations were established by using the version of Itô's formula for stochastic flows established in [\[45\]](#page-43-6). These stochastic integral representations are applicable to incompressible fluid flows on the whole space or flows with periodic boundary conditions. Later on in Constantin and Iyer [\[17\]](#page-41-13) the stochastic Lagrangian approach has been generalised to incompressible fluid flows constrained in a domain with boundary, and further extended by Iyer [\[35\]](#page-42-9) to inviscid fluid flows (with a stochastic perturbation). These stochastic integral representations not only utilise Taylor's diffusion driven by the fluid flow velocity but also its backward flows.

The stochastic Lagrangian approach, more precisely, various stochastic formulations and stochastic integral representations (though implicit) for solutions of incompressible fluid flows are very useful in the study of fluid dynamics, in particular in gaining information through numerical simulations. In a series of remarkable papers by Drivas and Eyink [\[22,](#page-42-10) [23\]](#page-42-11), Eyink, Gupta and Zaki [\[24,](#page-42-12) [25\]](#page-42-13), the stochastic Lagrangian approach has been applied successfully to the study of isotropic turbulent flows, verifying the small-scale theory, such as those proposed in $[41, 42]$ $[41, 42]$ $[41, 42]$, $[55]$, $[19]$ and $[43]$, also Wung-Tseng $[73]$, Yu et al. $[74]$ and the literature therein.

There are other interesting applications of stochastic calculus in the mathematical study of the Navier-Stokes equations, and in the descriptions of fluid dynamics including turbulent flows, let us mention only a few of them: $[11]$, $[30]$, $[32]$, $[57]$, $[64]$, $[69]$, $[70, 71]$ $[70, 71]$ $[70, 71]$, and certainly there are more interesting works the present authors must apologize for their ignorance not due to less significance of the work we are not aware of.

In this paper, based on the approach developed in recent works [\[59\]](#page-43-14) and [\[60\]](#page-44-11), we aim to deal

with highly complicated yet important fluid flows with heat transfer by bringing in several new ideas. From the perspective of solving the Navier-Stokes equations numerically for fluid flows with heat conduction, we devise the following approach, inspired mainly by Taylor [\[66\]](#page-44-3), the fundamental ideas in Goodman [\[31\]](#page-42-5) and Long [\[49\]](#page-43-9). For simplicity, let us illustrate our method for two-dimensional fluid flows. To determine the velocity $u(x, t)$ of the flow, it is equivalent to describe the "fictitious" Brownian motion particles with velocity $u(x, t)$, departed from all possible site ξ . These Brownian fluid particles are denoted by $X^{\xi} = (X_t^{\xi})$ ζ_t^{ζ} _t \geq ₀, which are diffusion processes defined as the weak solutions of Itô's stochastic differential equations

$$
dX_t^{\xi} = u(X_t^{\xi}, t)dt + \sqrt{2\nu}dB_t, \quad X_0^{\xi} = \xi,
$$

(called Taylor's diffusion (cf. Taylor [\[66\]](#page-44-3))). By borrowing the idea from the mean field theory, the key step in our approach is to turn the preceding stochastic differential equation into a McKean-Vlasov type stochastic differential equations (cf. [\[53\]](#page-43-15)) by using the law of the Brownian particles X^{ξ} and the governing fluid dynamical equations [\(1.1,](#page-2-1) [1.2](#page-2-2) and [1.3\)](#page-2-3). In random vortex methods, this is achieved by using the vorticity transport equation for $\omega(x, t)$. Indeed, by taking curl operation on both sides of the Navier-Stokes equation [\(1.1\)](#page-2-1), $\omega(x, t)$ evolves according to a linear (considering the velocity and the temperature θ as known fluid dynamical variables) parabolic equation

$$
\nu \Delta \omega - (u \cdot \nabla)\omega - \frac{\partial}{\partial t}\omega + g = 0.
$$

where for simplicity we use q to denote the curl of f. While handling the boundary condition imposed on the vorticity ω for wall-bounded flows poses a technical issue (for example see [\[3\]](#page-41-16) and [\[14\]](#page-41-17)), we shall for now focus on unbounded flows on the plane for the sake of clarity. Nonetheless, we should emphasize the importance of the interaction term q in the vorticity equation - which always occurs for wall-bounded flows, and the underlying reasons will be explained in the subsequent sections. Let $p(s, x, t, y)$ denote the transition probability density function of the diffusion process X^{ξ} . Since $\nabla \cdot u = 0$, $p(s, x, t, y)$ (for $s < t$ and $x, y \in \mathbb{R}^2$) is also the Green function of the forward parabolic equation

$$
\left(\nu\Delta - u \cdot \nabla - \frac{\partial}{\partial t}\right)v = 0.
$$
\n(1.4)

Therefore, according to the vorticity transport equation, the following representation (see [\[29\]](#page-42-16)) holds:

$$
\omega(x,t) = \int \omega(y,0)p(0,y,t,x)dy + \int_0^t \int g(y,s)p(s,y,t,x)dy.
$$
\n(1.5)

According to the Biot-Savart law,

$$
u(x,t) = \int K(y,x) \wedge \omega(y,t) dy
$$

=
$$
\int \mathbb{E} \left[K(X_t^z, x) \wedge \omega(z,0) \right] dz + \int_0^t \iint K(y,x) \wedge g(\xi,s) p(s,\xi,t,y) dy d\xi ds \qquad (1.6)
$$

where $K(y, x) = (2\pi)^{-1}(y - x)/|y - x|^2$ is the Biot-Savart singular integral kernel. The second integral on the right-hand side of [\(1.6\)](#page-4-0) may be written in terms of an expectation

$$
\int_0^t \int \mathbb{E}\left[K(X_t^{\xi,s}, x) \wedge g(\xi, s)\right] d\xi ds,
$$

where $X^{\xi,s}$ is the Taylor's diffusion starting from all possible site ξ and at instance s (for all $s \geq 0$: √

$$
dX_t^{\xi,s} = u(X_t^{\xi,s},t)dt + \sqrt{2\nu}dB_t, \quad X_\tau^{\xi,s} = \xi, \text{ for } \tau \le s. \tag{1.7}
$$

Therefore

$$
u(x,t) = \int \mathbb{E}\left[K(X_t^z, x) \wedge \omega(z,0)\right] \mathrm{d}z + \int_0^t \int \mathbb{E}\left[K(X_t^{\xi,s}, x) \wedge g(\xi,s)\right] \mathrm{d}\xi \mathrm{d}s,\tag{1.8}
$$

which allows us to reformulate the stochastic differential equation (1.7) into a McKean-Vlasovtype mean field equation. More precisely,

$$
dX_t^{\xi,s} = \int \mathbb{E}\left[K(X_t^{z,s},x) \wedge \omega(z,0)\right]dz + \int_0^t \int \mathbb{E}\left[K(X_t^{\xi,s},x) \wedge g(\xi,s)\right]\Big|_{x=X_t^{\xi,s}}d\xi ds
$$

+ $\sqrt{2\nu}dB_t, \quad X_\tau^{\xi,s} = \xi \quad \text{for all } \tau \le s.$ (1.9)

For the case where the external force f can be computed separately or it is known, then the McKean-Vlasov type mean field equation [\(1.9\)](#page-5-0), in combining the strong law of large number, can be used to design Monte-Carlo type numerical schemes for solving the velocity of [\(1.9\)](#page-5-0) accordingly. The velocity $u(x,t)$ may be determined by X^{ξ} , so the previous equation, which is the kind of ordinary stochastic differential equations involving the law of the solution X^{ξ} , has an advantage for numerically calculating the velocity $u(x, t)$, and hence provides a method for numerically solving the Navier-Stokes equations. There is extensive literature on numerical solutions of both ordinary stochastic differential equations and McKean-Vlasov type mean field equations, see [\[40\]](#page-43-16) for example. Analogous stochastic integral representations may be established for a 3D flow in a domain with or without boundary in terms of the Taylor's diffusion [\(1.7\)](#page-4-1) alone, as seen in [\[60\]](#page-44-11) and [\[59\]](#page-43-14). In these works, new stochastic integral representation theorems were established by using the duality of conditional distributions of a class of diffusion processes and a forward-type Feynman-Kac formula for solutions of parabolic equations.

There is however a serious disadvantage, and indeed, it is an obstacle to implementing the numerical schemes for computing numerically the solutions of the nonlinear mean field equation [\(1.9\)](#page-5-0) which is established based on the classical representation [\(1.5\)](#page-4-2). In fact, numerical methods for solving the mean field equation [\(1.9\)](#page-5-0) require numerically simulating Brownian fluid particles $X^{\xi,s}$ starting not only from any site ξ in the region of fluid, but also for every instance $s \geq 0$. This becomes unavoidable (at least under the current technology) when the interaction force f is not trivial – unfortunately, it is the case for wall-bounded flows and also for fluid flows with internal interaction force applying to the underlying fluid. The requirement for simulating diffusion paths starting at every instance substantially increases the computational cost for computing the solutions of the mean field equation [\(1.9\)](#page-5-0). In this paper, to overcome this obstacle for implementing the random vortex approach to the Oberbeck-Boussinesq flows where a conducting force is an essential feature, we utilise the divergence-free condition that $\nabla \cdot u = 0$ and the duality of conditional laws established in the previous work [\[60\]](#page-44-11), and derive a new integral representation theorem for solutions of the linear parabolic equation [\(1.4\)](#page-4-3).

To the best knowledge of the present authors, vortex methods have been studied for the Navier-Stokes equations without the consideration of energy transfer through heat or other fields. Indeed, substantial novel ideas need to be introduced in order to extend the random vortex method to important fluid flows appearing in applications, such as fluid flows with thermal conduction described by [\(1.1,](#page-2-1) [1.2,](#page-2-2) [1.3\)](#page-2-3).

In this paper, our goal is to develop new numerical schemes by establishing stochastic functional integral representation theorems for solutions to Oberbeck-Boussinesq fluid flows with thermal conduction in unbounded and wall-bounded domains.

Of course we must point out that the Computational Fluid Dynamics (CFD) is a huge subject, and there is a large volume of literature, see for example [\[28,](#page-42-17) [72\]](#page-44-12) for a small sample. CFD by default covers all aspects of computational techniques for calculating numerically solutions of all interesting fluid flows. In the past, however, CFD mainly concerns with the finite difference and finite element methods applying to fluid flows in science and engineering. Numerical simulations for turbulent flows have become popular too, due to the increasing computational capability over recent years. Various simulation tools have been developed in recent years, such as Direct Numerical Simulations (DNS), Large Eddy Simulations (LES), Probability Density Function (PDF) method and etc. The reader may refer to [\[28,](#page-42-17) [48,](#page-43-17) [58,](#page-43-4) [63\]](#page-44-13) for an overview of these aspects.

There is also good literature about the numerical solutions of Oberbeck-Boussinseq flows, and in general about Monte-Carlo simulations for fluid flows, see for example [\[1,](#page-40-1) [2\]](#page-41-18).

The paper is organised as follows: in Section 2, some preliminary results on Taylor's diffusions associated with solenoidal vector fields are recalled, and a new functional integral representation formula is derived for the solution of the parabolic equation associated with Taylor's diffusion. In section 3, we review the Biot-Savart law and introduce a variation of the classical Biot-Savart law to link the flow velocity field to its vorticity field for the latter sections. To handle the coupled equations governing the fluid flow with thermal conduction, we introduce the twin Brownian particles in Section 4, which serve as Taylor's diffusions in the representation formula discovered in section 2. We establish the results for the unbounded case first. In Section 5, by applying the representation formula with the twin Brownian particles, the probabilistic representations for the fluid dynamical variables in Oberbeck-Boussinesq flows in \mathbb{R}^2 and \mathbb{R}^3 are established, and this allows us to formulate a closed random vortex dynamical system for Oberbeck-Boussinesq flows on unbounded domains. In section 6, we handle the flows in wall-bounded domains. With the twin Brownian particles in bounded domains, we first derive the representation results for Oberbeck-Boussinesq flows in half-spaces using dynamical variables mollified in a thin layer adjacent to the boundary. Additionally, the random vortex dynamics are characterised using these representations for wall-bounded flows in dimensions two and three. Next, we send the thickness of the layer to zero and find limiting representations of these variables. Finally, in section 7, we provide numerical schemes based on the representation formulae in Sections 5 and 6, for unbounded and wall-bounded Oberbeck-Boussinesq flows, along with some numerical experiment results with different Prandtl numbers.

Notations and Conventions: Unless otherwise specified, Einstein's summation convention over repeated indices is assumed throughout the paper. For two-dimensional vectors $a = (a_1, a_2)$ and $b = (b_1, b_2), a \wedge b = a_1b_2 - a_2b_1$. For a real number c, $a \wedge c$ is identified with the vector $(a_2c, -a_1c)$.

2 Divergence-free vector fields

In this section, we recall several results on divergence-free vector fields on \mathbb{R}^d , where the dimension $d \geq 2$, although we are only interested in the case where $d = 2$ or $d = 3$.

Suppose that $b(x, t)$ is a time-dependent, bounded and Borel measurable vector field on \mathbb{R}^d , such that $\nabla \cdot b = 0$ on \mathbb{R}^d in the sense of distribution for every t. Let $\lambda > 0$ be a constant. Then

$$
L_{\lambda,b} = \lambda \Delta + b \cdot \nabla
$$

is a second-order elliptic operator on \mathbb{R}^d . Furthermore, since $\nabla \cdot b = 0$, the adjoint operator $L^*_{\lambda,b}$ of $L_{\lambda,b}$ is $L_{\lambda,-b}$.

For every $\tau \geq 0$ and $\xi \in \mathbb{R}^d$, there is a unique weak solution, denoted by $X^{\xi,\tau} = (X^{\xi,\tau}_t)$ $(t^{5,7})_{t\geq0},$ of the stochastic differential equation (SDE):

$$
dX_t = b(X_t, t)dt + \sqrt{2\lambda}dB_t, \quad X_\tau = \xi, \quad \text{ for } t \ge \tau,
$$

where B is a d-dimensional Brownian motion on some probability space. If $\tau = 0$, then $X^{\xi,0}$ will be denoted by X^{ξ} for simplicity. $X^{\xi,\tau}_t$ may be also denoted by $X(\xi,\tau;t)$ (for $t \geq 0$). The distribution of $X^{\xi,\tau}$ is denoted by $\mathbb{P}^{\xi,\tau}$ and by \mathbb{P}^{ξ} if $\tau=0$, which are probability measures on the path space $C([0,\infty);\mathbb{R}^d)$. $X^{\xi,\tau}$ (also $\mathbb{P}^{\xi,\tau}$) is called the diffusion with infinitesimal generator $L_{\lambda,b}$, or called the $L_{\lambda,b}$ -diffusion.

It is known that the law of $X_t^{\xi,\tau}$ $t_t^{\xi,\tau}$ for every $t > \tau \geq 0$ has a positive and continuous probability density function with respect to the Lebesgue measure, denoted by $p_{\lambda b}(\tau, \xi, t, x)$. The function $p_{\lambda,b}(\tau,\xi,t,x)$ for $t > \tau \geq 0$ and $\xi, x \in \mathbb{R}^d$ is the transition probability density function of the $L_{\lambda,b}$ -diffusion. Since $b(x,t)$ is a Borel measurable, bounded vector field, $p_{\lambda,b}(\tau,\xi,t,x)$ is jointly Hölder continuous in $t > \tau$ and $\xi, x \in \mathbb{R}^d$. Moreover, if $b(x, t)$ is smooth, so is $p_{\lambda, b}$.

The conditional law $\mathbb{P}^{\xi,\tau}[\cdot | X_T^{\xi,\tau} = \eta]$ (also called the pinned measure) is denoted by $\mathbb{P}^{\xi,\tau \to \eta,T}$, and by $\mathbb{P}^{\xi \to \eta}$ if $\tau = 0$ and $T > 0$ is given. For the construction of the conditional laws, cf. [\[60\]](#page-44-11) and [\[59\]](#page-43-14).

It is known that $p_{\lambda,b}(\tau,\xi,t,x)$ is the Green function of the backward parabolic operator $L_{\lambda,b} + \frac{\partial}{\partial t}$ (cf. [\[65\]](#page-44-14)), so that $p_{\lambda,b}(\tau,\xi,t,x)$ coincides with the Green function of the forward parabolic operator $L^*_{\lambda,b} - \frac{\partial}{\partial t}$ (cf. [\[29,](#page-42-16) Chapter 1, Theorem 15]). Since $b(x,t)$ is divergence-free on \mathbb{R}^d in the distribution sense, $L_{\lambda,b}^* = L_{\lambda,-b}$, therefore $p_{\lambda,b}(\tau,\xi,t,x)$ is the Green function of the forward parabolic operator $L_{\lambda,-b} - \frac{\partial}{\partial t}$ on \mathbb{R}^d .

Suppose that $D \subset \mathbb{R}^d$ is a domain with a Lipschitz continuous boundary ∂D . Let $p_{\lambda,b}^D(\tau,\xi,t,x)$ (for $t > \tau \geq 0$, $\xi, x \in D$) be the transition density function of the diffusion X killed on leaving the domain D, that is

$$
\mathbb{E}\left[\varphi(X_t^{\xi,\tau})1_{\{t<\zeta(X^{\xi,\tau})\}}\right] = \int_D p_{\lambda,b}^D(\tau,\xi,t,x)\varphi(x)dx
$$

for any bounded and Borel measurable function φ , where $\zeta(\psi) = \inf\{t \geq 0 : \psi(t) \notin D\}$. Since $\nabla \cdot b = 0$ in the distribution sense in \mathbb{R}^d , $p_{\lambda,b}^D(\tau,\xi,t,x)$ is the Green function to the Dirichlet problem of the forward parabolic equation operator

$$
\left(L_{\lambda,-b} - \frac{\partial}{\partial t}\right)w(x,t) = 0 \quad \text{in } D \times (0,\infty)
$$
\n(2.1)

subject to the Dirichlet boundary condition that

$$
w(x,t) = 0 \quad \text{for } x \in \partial D. \tag{2.2}
$$

Next, we establish the main technical tool for the present work.

Let $\Psi(x,t) = (\Psi^1(x,t), \cdots, \Psi^n(x,t))$ be a smooth solution of the parabolic equations

$$
\left(\lambda \Delta - b \cdot \nabla - \frac{\partial}{\partial t}\right) \Psi^i + q_l^i \Psi^l + F^i = 0 \quad \text{in } D \times (0, \infty), \tag{2.3}
$$

$$
\Psi(x,t) = 0 \quad \text{for } x \in \partial D,\tag{2.4}
$$

for $i = 1, \dots, n$, where $q(x, t) = (q_i^i(x, t))_{1 \leq i,l \leq n}$ is a bounded, Borel measurable $n \times n$ matrixvalued function on D. $q_l^i(x,t) = 0$ for $x \notin D$, or otherwise we replace $q(x,t)$ by $1_D(x)q(x,t)$ instead. $F(x,t) = (F^i(x,t))_{1 \leq i \leq n}$ is a family of functions which are $C^{2,1}(D \times [0,\infty))$.

Let $\Omega = C([0,\infty);\mathbb{R}^d)$, and let τ_T denote the time reversal at T on Ω . That is, $\psi \circ \tau_T(s) =$ $\psi(T-s)$ for $s \in [0, T]$. Also, as defined before, $\zeta(\psi) = \inf \{t : \psi(t) \notin D\}$, the first exit time from the region D, and $\gamma_T(\psi) = \sup \{t \in (0,T) : \psi(t) \notin D\}$, the last exit time before time T from D.

Theorem 2.1. Let $T > 0$ and $\eta \in \mathbb{R}^d$. Let $t \mapsto Q^i_j(\eta, T; t)$ (for $t \in [0, T]$) be the solutions to the ordinary differential equations

$$
\frac{d}{dt}Q_j^i(\eta, T; t) = -Q_k^i(\eta, T; t)q_j^k(X_t^{\eta}, t), \quad Q_j^i(\eta, T; T) = \delta_{ij}
$$
\n(2.5)

for $i, j = 1, \cdots, n$. Then

$$
\Psi^{i}(\xi,T) = \int_{D} \mathbb{E} \left[1_{\{T < \zeta(X^{\eta})\}} Q_{j}^{i}(\eta,T;0) \Psi^{j}(\eta,0) \Big| X_{T}^{\eta} = \xi \right] p_{\lambda,b}(0,\eta,T,\xi) d\eta
$$
\n
$$
+ \int_{0}^{T} \int_{D} \mathbb{E} \left[1_{\{t > \gamma_{T}(X^{\eta})\}} Q_{j}^{i}(\eta,T;t) F^{j}(X_{t}^{\eta},t) \Big| X_{T}^{\eta} = \xi \right] p_{\lambda,b}(0,\eta,T,\xi) d\eta dt \tag{2.6}
$$
\nFigure 6.5. D, and T > 0, i = 1, ...

for every $\xi \in D$ and $T > 0$, $i = 1, \dots, n$.

Proof. The subscript λ will be omitted in the proof. For each $T > 0$, let \tilde{X}^{ξ} be the solution to the stochastic differential equation

$$
d\tilde{X}_t^{\xi} = -b(\tilde{X}_t^{\xi}, T - t)dt + \sqrt{2\lambda}dB_t, \quad \tilde{X}_0^{\xi} = \xi
$$

for every $\xi \in \mathbb{R}^d$ and B is a standard Brownian motion in \mathbb{R}^d on some probability space, where $b(x, t) = 0$ for $t < 0$. Define

$$
d\tilde{Q}_j^i(t) = \tilde{Q}_k^i(t)q_j^k(\tilde{X}_t^{\xi}, T - t)dt, \quad \tilde{Q}_j^i(0) = \delta_{ij}
$$

(note that we assume that $q_j^i(x,t) = 0$ for $x \notin D$), where $i, j = 1, \dots, n$. Let

$$
Y_t = \Psi(\tilde{X}_{t \wedge T_{\xi}}^{\xi}, T - t) = 1_{\{t < T_{\xi}\}} \Psi(\tilde{X}_t^{\xi}, T - t),
$$

where $T_{\xi} = \inf \{ t : \tilde{X}_t^{\xi} \notin D \}.$ Here the second equality is ensured as Ψ vanishes along the boundary ∂D . Let $M^i = \tilde{Q}_j^i Y^j$. Then by Itô's formula and Eq. [\(2.3,](#page-7-0) [2.4\)](#page-7-1) we obtain that

$$
M_t^i = Y_0^i + \sqrt{2\lambda} \int_0^t \mathbf{1}_{\{s < T_\xi\}} \tilde{Q}_j^i(s) \nabla \Psi^j(\tilde{X}_s^\xi, T - s) \cdot \mathrm{d}B_s
$$

$$
- \int_0^t \mathbf{1}_{\{s < T_\xi\}} \tilde{Q}_j^i(s) F^j(\tilde{X}_s^\xi, T - s) \mathrm{d}S.
$$

Taking expectation on both sides and using the fact that Ψ vanishes along the boundary, we obtain that

$$
\Psi^{i}(\xi,T) = \mathbb{E}\left[\tilde{Q}_{j}^{i}(T)\Psi^{j}(\tilde{X}_{T}^{\xi},0)1_{\{T\n(2.7)
$$

Note that T_{ξ} is a stopping time with respect to the filtration generated by \tilde{X} , so that $\tilde{Q}^i_j(t)1_{\{t < T_{\xi}\}}$ is therefore measurable with respect to \tilde{X} running up to time t, which allows us to take conditional expectation on giving $\tilde{X}_T = \eta$, to obtain that

$$
J_2^i = \int_0^T \int_{\mathbb{R}^d} \mathbb{E} \left[\tilde{Q}_j^i(t) 1_{\{t < T_\xi\}} F^j(\tilde{X}_t^\xi, T - t) \middle| \tilde{X}_T^\xi = \eta \right] \mathbb{P} \left[\tilde{X}_T^\xi \in \mathrm{d}\eta \right] \mathrm{d}t
$$

=
$$
\int_0^T \int_D \mathbb{E} \left[\tilde{Q}_j^i(t) 1_{\{t < T_\xi\}} F^j(\tilde{X}_t^\xi, T - t) \middle| \tilde{X}_T^\xi = \eta \right] p_{-b_T}(0, \xi, T, \eta) \mathrm{d}\eta \mathrm{d}t
$$

where $p_{-b_T}(\tau, \xi, t, \eta)$ is the transition probability density function of the diffusion \tilde{X}^{ξ} , and the second equality follows from the fact that the conditional expectation is zero if $\eta \notin D$, so we may restrict the integral for $\eta \in D$ only. Similarly

$$
J_1^i = \int_D \mathbb{E}\left[\tilde{Q}_j^i(T)1_{\{T < T_{\xi}\}}\middle|\tilde{X}_T^{\xi} = \eta\right]\Psi^j(\eta,0)p_{-b_T}(0,\xi,T,\eta)d\eta.
$$

Since $\nabla \cdot b = 0$, so that we can replace $p_{-b_T}(0,\xi,T,\eta)$ by $p_b(0,\eta,T,\xi)$. Hence

$$
J_1^i = \int_D \mathbb{E}\left[\tilde{Q}_j^i(T)1_{\{T < T_\xi\}} \middle| \tilde{X}_T^\xi = \eta\right] \Psi^j(\eta, 0) p_b(0, \eta, T, \xi) d\eta \tag{2.8}
$$

and

$$
J_2^i = \int_0^T \int_D \mathbb{E}\left[\tilde{Q}_j^i(t)1_{\{t < T_\xi\}} F^j(\tilde{X}_t^\xi, T - t)\Big|\tilde{X}_T^\xi = \eta\right] p_b(0, \eta, T, \xi) d\eta dt \tag{2.9}
$$

for $i = 1, \dots, n$. Next we utilise the approach in [\[60\]](#page-44-11) and rewrite the conditional expectations in terms of the diffusion process with infinitesimal generator $\lambda \Delta + b \cdot \nabla$. To this end, we first introduce a few notations. Let $\tilde{\mathbb{P}}^{\xi}$ denote the law of \tilde{X}^{ξ} and $\tilde{\mathbb{P}}^{\xi\to\eta}$ denote the conditional law of \tilde{X}^{ξ} given the terminal value that $\tilde{X}^{\xi}_{T} = \eta$. Let $\tilde{Q}(\psi; t)$ denote the solution to the linear system of ordinary differential equations

$$
\frac{\mathrm{d}}{\mathrm{d}t}\tilde{Q}_j^i(\psi;t) = \tilde{Q}_k^i(\psi;t)q_j^k(\psi(t),T-t), \quad \tilde{Q}_j^i(\psi;0) = \delta_{ij}
$$

for every $\psi \in C([0,T]; \mathbb{R}^d)$, where $i, j = 1, \dots, n$. The representations $(2.8, 2.9)$ $(2.8, 2.9)$ $(2.8, 2.9)$ can be rewritten as:

$$
J_1^i = \int_D \tilde{\mathbb{P}}^{\xi \to \eta} \left[\tilde{Q}_j^i(\psi;T) 1_{\{T < \zeta(\psi)\}} \right] \Psi^j(\eta,0) p_b(0,\eta,T,\xi) d\eta
$$

and

$$
J_2^i = \int_0^T \int_D \tilde{\mathbb{P}}^{\xi \to \eta} \left[\tilde{Q}_j^i(\psi; t) F^j(\psi(t), T - t) \mathbb{1}_{\{t < \zeta(\psi)\}} \right] p_b(0, \eta, T, \xi) d\eta dt.
$$

Since $\nabla \cdot b = 0$, according to the duality for the conditional laws (cf. [\[60,](#page-44-11) Theorem 3.1]), the conditional law $\tilde{\mathbb{P}}^{\xi \to \eta}$ coincides with the conditional law $\mathbb{P}^{\eta \to \xi} \circ \tau_T$, where τ_T denotes the time reversal at T. Therefore we may rewrite

$$
J_1^i = \int_D \mathbb{P}^{\eta \to \xi} \left[1_{\{T < \zeta(\psi \circ \tau_T)\}} \tilde{Q}_j^i(\psi \circ \tau_T; T) \right] \Psi^j(\eta, 0) p_b(0, \eta, T, \xi) d\eta
$$

and

$$
J_2^i = \int_0^T \int_D \mathbb{P}^{\eta \to \xi} \left[1_{\{t < \zeta(\psi \circ \tau_T)\}} \tilde{Q}_j^i(\psi \circ \tau_T; t) F^j(\psi(T - t), T - t) \right] p_b(0, \eta, T, \xi) d\eta dt.
$$

It remains to identify the flow $\tilde{Q}_j^i(\psi \circ \tau_T;t)$ for $t \in [0,T]$. Consider $Q_j^i(\psi, T; s) = \tilde{Q}_j^i(\psi \circ \tau_T; T - s)$ for $0 < s \leq T$. Then one can easily verify that $Q_j^i(\psi, T; s)$ is exactly the unique solution of Eq. [\(2.5\)](#page-7-2). Therefore

$$
J_1^i = \int_D \mathbb{P}^{\eta \to \xi} \left[1_{\{T < \zeta(\psi \circ \tau_T)\}} Q_j^i(\psi, T; 0) \right] \Psi^j(\eta, 0) p_b(0, \eta, T, \xi) d\eta
$$

and

$$
J_2^i = \int_0^T \int_D \mathbb{P}^{\eta \to \xi} \left[1_{\{t < \zeta(\psi \circ \tau_T)\}} Q_j^i(\psi, T; T - t) F^j(\psi(T - t), T - t) \right] p_b(0, \eta, T, \xi) d\eta dt
$$

=
$$
\int_0^T \int_D \mathbb{P}^{\eta \to \xi} \left[1_{\{T - t < \zeta(\psi \circ \tau_T)\}} Q_j^i(\psi, T; t) F^j(\psi(t), t) \right] p_b(0, \eta, T, \xi) d\eta dt.
$$

Hence, by using equalities in (2.7) , we obtain

$$
\Psi^{i}(\xi,T) = \int_{D} \mathbb{P}^{\eta \to \xi} \left[1_{\{T < \zeta(\psi \circ \tau_T)\}} Q^i_j(\psi,T;0) \right] p_b(0,\eta,T,\xi) \Psi^j(\eta,0) d\eta \n+ \int_0^T \int_D \mathbb{P}^{\eta \to \xi} \left[1_{\{T-t < \zeta(\psi \circ \tau_T)\}} Q^i_j(\psi,T;t) F^j(\psi(t),t) \right] p_b(0,\eta,T,\xi) d\eta dt.
$$

Now we make the following observation. For $\xi, \eta \in D$, then for a path $\psi \in C([0, T]; \mathbb{R}^d)$ with $\psi(0) = \eta$ and $\psi(T) = \xi$, then $T < \zeta(\psi \circ \tau)$ is equivalent to say $\psi(T - t) \in D$ for all $t \in (0,T)$, and therefore the condition that $T < \zeta(\psi \circ \tau)$ almost surely under the conditional law $\mathbb{P}^{\eta \to \xi}$ is equivalent to that $T < \zeta(\psi)$ almost surely w.r.t. $\mathbb{P}^{\eta \to \xi}$. Therefore the indicator function $1_{\{T < \zeta(\psi \circ \tau_T)\}}$ can be replaced by $1_{\{T < \zeta(\psi)\}}$. The treatment for the random function $1_{\{T-t<\zeta(\psi\circ\tau_T)\}}$ (for $t\in(0,T)$) in the second term is more subtle. Under the conditional law $\mathbb{P}^{\eta \to \xi}$ for $\xi, \eta \in D$, we may replace

$$
\zeta(\psi \circ \tau_T) = \inf \{ s > 0 : \psi(T - s) \in \partial D \} \wedge T
$$

= $\inf \{ T - s > 0 : \psi(s) \in \partial D \} \wedge T$
= $T - \sup \{ s \in (0, T) : \psi(s) \in \partial D \}$

with the convention that $\sup \emptyset = 0$. Thus $T - t < \zeta(\psi \circ \tau)$ is equivalent to

$$
t > \sup\left\{s \in (0, T) : \psi(s) \in \partial D\right\}.
$$

Therefore it is useful to introduce the notation that

$$
\gamma_T(\psi) = \sup \{ s \in (0, T) : \psi(s) \in \partial D \}
$$

for every path ψ and $T > 0$. Then $1_{\{T-t < \zeta(\psi \circ \tau_T)\}}$ can be replaced by $1_{\{t > \gamma_T(\psi)\}}$. Hence (2.6) follows immediately. \Box

Remark 2.2. In handling the term J_2^i $(i = 1, \dots, n)$, one may use the conditional law given $\tilde{X}_t = \eta$ in place of $\tilde{X}_T = \eta$, so that

$$
J_2^i = \int_0^T \int_{\mathbb{R}^d} \mathbb{E} \left[\tilde{Q}_j^i(t) 1_{\{t < T_\xi\}} F^j(\tilde{X}_t^\xi, T - t) \Big| \tilde{X}_t^\xi = \eta \right] \mathbb{P} \left[\tilde{X}_t^\xi \in \mathrm{d}\eta \right] \mathrm{d}t
$$

\n
$$
= \int_0^T \int_D \mathbb{E} \left[\tilde{Q}_j^i(t) 1_{\{t < T_\xi\}} \Big| \tilde{X}_t^\xi = \eta \right] F^j(\eta, T - t) p_{-b_T}(0, \xi, t, \eta) \mathrm{d}\eta \mathrm{d}t
$$

\n
$$
= \int_0^T \int_D \mathbb{E} \left[\tilde{Q}_j^i(t) 1_{\{t < T_\xi\}} \Big| \tilde{X}_t^\xi = \eta \right] F^j(\eta, T - t) p_b(T - t, \eta, T, \xi) \mathrm{d}\eta \mathrm{d}t.
$$

This leads to the similar formula in the case where $D = \mathbb{R}^d$ and $q_j^i \equiv 0$. Indeed, for this case $T_{\xi} = \infty$ and $\tilde{Q}^i_j = \delta_{ij}$ so that

$$
\mathbb{E}\left[\tilde{Q}_j^i(t)1_{\{t < T_{\xi}\}}\middle|\tilde{X}_t^{\xi} = \eta\right] = \delta_{ij}
$$

and therefore

$$
J_2^i = \int_0^T \int_{\mathbb{R}^d} F^i(\eta, T - t) p_b(T - t, \eta, T, \xi) d\eta dt
$$

=
$$
\int_0^T \int_{\mathbb{R}^d} F^i(\eta, t) p_b(t, \eta, T, \xi) d\eta dt.
$$

Remark 2.3. The duality of conditional laws among certain diffusion processes used in the proof of the preceding theorem is the main tool developed in [\[60\]](#page-44-11) in order to formulate a random vortex method for three dimensional incompressible fluid flows by using only the forward Taylor diffusion of the flow velocity. The conditional law techniques for diffusions have their origin from the study of symmetric diffusion processes and Dirichlet forms. The notion of duality of diffusion distributions was certainly developed from the well-known concept of self-adjoint operators, and for diffusion semigroups, the path-space version of the self-adjointness was first discovered in a seminal work by Lyons and Zheng [\[50\]](#page-43-18). In fact, Lyons-Zheng [\[51\]](#page-43-19) has utilised the conditional laws of symmetric diffusions to the study of heat kernels for a class of non-symmetric diffusion processes. The reader may find a detailed account in [\[59\]](#page-43-14) about conditional law duality and its applications to forward type Feynman-Kac formulas.

While in the presence of a non-trivial gauge function $q(x, t)$, or the presence of non-empty boundary, it seems that the formulation is more useful by conditioning on \tilde{X}_T . Due to these are important cases, we may formulate the following theorem, which is indeed an extension of the theorem proved in [\[60\]](#page-44-11).

Theorem 2.4. Under the same assumptions and notations as in Theorem [2.1,](#page-7-4) suppose that $\Psi = (\Psi^i)_{1 \leq i \leq n}$ is a solution to Eq. [\(2.3\)](#page-7-0) on \mathbb{R}^d , then

$$
\Psi^{i}(\xi,T) = \int_{\mathbb{R}^{d}} \mathbb{E}\left[Q_{j}^{i}(\eta,T;0)\Psi^{j}(\eta,0)\Big|X_{T}^{\eta}=\xi\right]p_{b}(0,\eta,T,\xi)d\eta
$$

$$
+\int_{0}^{T}\int_{\mathbb{R}^{d}} \mathbb{E}\left[Q_{j}^{i}(\eta,T;t)F^{j}(X_{t}^{\eta},t)\Big|X_{T}^{\eta}=\xi\right]p_{b}(0,\eta,T,\xi)d\eta dt
$$

for every $\xi \in \mathbb{R}^d$, $T > 0$ and $i = 1, \dots, n$.

3 The Biot-Savart laws

In this section we formulate the Biot-Savart laws we need in this work for velocity, vorticity and temperature and its gradient.

Recall that the Green function in \mathbb{R}^d is given by the following formula

$$
\Gamma_d(y, x) = \begin{cases} -\frac{1}{(d-2)s_{d-1}} \frac{1}{|y-x|^{d-2}}, & \text{if } d > 2, \\ \frac{1}{2\pi} \ln|y-x|, & \text{if } d = 2, \end{cases}
$$

where s_{d-1} is the surface area of a unit sphere in \mathbb{R}^d , so $s_2 = 4\pi$. We are only interested in the cases where $d = 2$ and $d = 3$. The Biot-Savart singular integral kernels $K_d(y, x) = \nabla_y \Gamma_d(y, x)$, so that

$$
K_2(y, x) = \frac{1}{2\pi} \frac{y - x}{|y - x|^2}
$$
 for $y \neq x$

and

$$
K_3(y,x) = \frac{1}{4\pi} \frac{y-x}{|y-x|^3}
$$
 for $y \neq x$.

Recall that under our convention for two-dimensional vectors, $\omega = \nabla \wedge u$ is identified with the scalar function $\frac{\partial}{\partial x_1}u^2 - \frac{\partial}{\partial x}$ $\frac{\partial}{\partial x_2}u^1$, and for $a=(a_1,a_2)$, $a \wedge \omega$ is identified with $(a_2\omega, -a_1\omega)$.

Lemma 3.1 (The Biot-Savart law). Let $d = 2$ or $d = 3$.

(1) If u is a vector field on \mathbb{R}^d such that $\nabla \cdot u = 0$ and $\omega = \nabla \wedge u$, then

$$
u(x) = \int_{\mathbb{R}^d} K_d(y, x) \wedge \omega(y) dy, \quad \forall x \in \mathbb{R}^d.
$$
 (3.1)

(2) If θ is a function on \mathbb{R}^d and $\Theta = \nabla \cdot \theta$, then

$$
\theta(x) = -\int_{\mathbb{R}^d} K_d(y, x) \cdot \Theta(y) dy, \quad \forall x \in \mathbb{R}^d.
$$
 (3.2)

These formulae follow immediately from the Green function and integration by parts.

The Green function for the upper half space $\mathbb{R}^d_+ = \{x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d : x_d > 0\}$ is given by $G_d(y, x) = \Gamma_d(y, x) - \Gamma_d(y, \overline{x})$

$$
G_d(y,x)=\Gamma_d(y,x)-\Gamma_d(y,\overline{x})
$$

for every $x, y \in \mathbb{R}^d_+$, and $\overline{x} = (x_1, \dots, x_{d-1}, -x_d)$ is the reflection of x about the hyperplane ${x_d = 0}$. Then the Green formula for \mathbb{R}^d_+ implies that

$$
\varphi(x) = \int_{\mathbb{R}_+^d} G_d(y, x) \Delta \varphi(y) dy - \int_{\{y_d = 0\}} \varphi(y) \frac{\partial}{\partial y_d} G_d(y, x) dy_1 \cdots dy_{d-1}
$$
(3.3)

for $x \in \mathbb{R}_+^d$, where φ is C^2 , continuous up to the boundary where $x_d = 0$, and vanishes at the infinity.

Similarly, we define $A_d(y, x) = \nabla_y G_d(y, x)$, which may be called the Biot-Savart singular kernel on \mathbb{R}^d_+ . Then

$$
A_2(y,x) = \frac{1}{2\pi} \left(\frac{y-x}{|y-x|^2} - \frac{y-\overline{x}}{|y-\overline{x}|^2} \right) \quad \text{for } y \neq x \text{ or } \overline{x}
$$
 (3.4)

and

$$
A_3(y,x) = \frac{1}{4\pi} \left(\frac{y-x}{|y-x|^3} - \frac{y-\overline{x}}{|y-\overline{x}|^3} \right) \quad \text{for } y \neq x \text{ or } \overline{x}.
$$

The Biot-Savart laws we need in this paper follow from the Green formula [\(3.3\)](#page-12-0), and are stated as two lemmas below.

Lemma 3.2. Let $d = 2$ or $d = 3$. If u is a vector field on \mathbb{R}^d_+ (continuous up to the boundary, decays to zero at infinity) such that $\nabla \cdot u = 0$ in \mathbb{R}^d_+ and $u(x) = 0$ when $x_d = 0$. Let $\omega = \nabla \wedge u$. Then

$$
u(x) = \int_{\mathbb{R}_+^d} A_d(y, x) \wedge \omega(y) dy, \quad \text{for } x \in \mathbb{R}_+^d.
$$
 (3.5)

Proof. Since $\nabla \cdot u = 0$ and $\nabla \wedge u = \omega$,

$$
\Delta u = -\nabla \wedge \nabla \wedge u = -\nabla \wedge \omega.
$$

Now as $u(x) = 0$ when $x_d = 0$, by Green's formula we obtain that

$$
u(x) = \int_{\mathbb{R}_+^d} G_d(y, x) \Delta u(y) dy = - \int_{\mathbb{R}_+^d} G_d(y, x) \nabla \wedge \omega(y) dy
$$

and the claim follows immediately after applying integration by parts.

Next, we formulate a similar law of Biot-Savart's for the temperature.

Lemma 3.3. Let θ be a scalar function on \mathbb{R}^d_+ and $\Theta = \nabla \theta$ be its gradient with its components $\Theta_i = \frac{\partial}{\partial x}$ $\frac{\partial}{\partial x_i}\theta$ for $i=1,\cdots,d.$ Suppose

$$
\theta(x) = \theta_0(x_1, \dots, x_{d-1}), \quad \forall x = (x_1, \dots, x_{d-1}, 0)
$$

is the trace of θ along the boundary $\{x_d = 0\}.$ (1) If $d = 3$, then

$$
\theta(x) = -\int_{\mathbb{R}^3_+} \Lambda_3(y, x) \cdot \Theta(y) dy + \frac{x_3}{2\pi} \int_{\mathbb{R}^2} \frac{\theta_0(y_1, y_2)}{|y_1 - x_2|^2 + |y_2 - x_2|^2 + |x_3|^2} dy_1 dy_2
$$

for $x = (x_1, x_2, x_3)$ with $x_3 > 0$.

(2) If
$$
d = 2
$$
, then

$$
\theta(x) = -\int_{\mathbb{R}_+^2} A_2(y, x) \cdot \Theta(y) dy + \frac{x_2}{\pi} \int_{-\infty}^{\infty} \frac{\theta_0(y_1)}{|y_1 - x_1|^2 + x_2^2} dy_1
$$

for $x = (x_1, x_2)$ with $x_2 > 0$.

Proof. By definition, $\Delta \theta = \nabla \cdot \Theta$, so according to the Green formula,

$$
\theta(x) = -\int_{\mathbb{R}_+^d} \nabla_y G_d(y,x) \cdot \Theta(y) dy - \int_{\{y_d=0\}} \theta_0(y) \frac{\partial}{\partial y_d} G_d(y,x) dy_1 \cdots dy_{d-1}.
$$

The claims follow immediately.

 \Box

 \Box

4 Twin Brownian particles

In the remainder of the paper, $u(x,t)$ is a time-dependent vector field on \mathbb{R}^d (where $d=2$ or $d = 3$. $u(x, t)$ may be the velocity of an incompressible fluid flow in \mathbb{R}^d or the appropriate (divergence-free) extension of the velocity of an incompressible fluid flow in $D \subset \mathbb{R}^d$. Let $\nu > 0$ be the kinematic viscosity constant, and $\kappa > 0$ the thermal diffusivity constant. We then introduce two families of random particles X and Y , defined by the following stochastic differential equations: √

$$
dX_t^{\xi} = u(X_t^{\xi}, t)dt + \sqrt{2\nu}dB_t^{\nu}, \quad X_0^{\xi} = \xi
$$
\n(4.1)

and

$$
dY_t^{\xi} = u(Y_t^{\xi}, t)dt + \sqrt{2\kappa}dB_t^{\kappa}, \quad Y_0^{\xi} = \xi
$$
\n(4.2)

for every $\xi \in \mathbb{R}^d$. Here B^{ν} and B^{κ} are two independent standard d-dimensional Brownian motions on some probability space.

The transition probability density functions for X (resp. for Y) are denoted by $p_{\nu}(s, x, t, y)$ and $p_{\kappa}(s, x, t, y)$ respectively.

Let $A_j^i = \frac{\partial}{\partial x_i^i}$ $\frac{\partial}{\partial x_j}u^i$ be the entries of Jacobian matrix of $u(x,t)$. Given a domain $D \subset \mathbb{R}^d$, we introduce two gauge functionals $Q = (Q_j^i(\eta, t; s))$ and $R = (R_j^i(\eta, t; s))$ (for $0 \le s \le t$ and $\eta \in \mathbb{R}^d$, where $i, j = 1, \dots, d$, defined by the following ordinary differential equations:

$$
\frac{\mathrm{d}}{\mathrm{d}s}Q_j^i(\eta, t; s) = -Q_k^i(\eta, t; s)1_D(X_s^{\eta})A_j^k(X_s^{\eta}, s), \quad Q_j^i(\eta, t; t) = \delta_{ij}
$$
\n(4.3)

and

$$
\frac{d}{ds}R_i^j(\eta, t; s) = R_i^l(\eta, t; s)1_D(Y_s^{\eta})A_l^j(Y_s^{\eta}, s), \quad R_i^j(\eta, t; t) = \delta_{ij}
$$
\n(4.4)

respectively.

The diffusions $X^{\xi,\tau}$ and $Y^{\eta,\tau}$ (where $\xi, \eta \in \mathbb{R}^d$ and $\tau \geq 0$) are defined as the (weak) solutions of the stochastic differential equations:

$$
dX_t^{\xi,\tau} = u(X_t^{\xi,\tau},t)dt + \sqrt{2\nu}dB_t^{\nu}, \quad X_s^{\xi,\tau} = \xi \quad \text{for } s \le \tau
$$
\n(4.5)

and

$$
dY_t^{\eta,\tau} = u(Y_t^{\eta,\tau}, t)dt + \sqrt{2\kappa}dB_t^{\kappa}, \quad Y_s^{\eta,\tau} = \eta \quad \text{for } s \le \tau
$$
\n(4.6)

for $t \geq 0$.

5 Unbounded fluid flows

In this section, we consider an ideal fluid flow in \mathbb{R}^d , $d=2$ or $d=3$, with an external source that supplies heat to the fluid.

5.1 The Oberbeck-Boussinesq equations in \mathbb{R}^d

In the Oberbeck-Boussinesq model, the fluid flow has no space constraint, so such a model serves as a model of homogeneous turbulent flows with heat transfer. Suppose the temperature gradient is small so that the density of the fluid is nearly constant. Therefore the model is described by the Boussinesq equations on the whole space \mathbb{R}^d . Suppose the heat source is represented by temperature $\theta(x)$, so that the flow is described by [\(1.1,](#page-2-1) [1.2,](#page-2-2) [1.3\)](#page-2-3) in \mathbb{R}^d . The Navier-Stokes equations may be written as

$$
\left(\nu\Delta - u \cdot \nabla - \frac{\partial}{\partial t}\right)u - \nabla P + f(\theta) = 0 \quad \text{in } \mathbb{R}^d
$$
\n(5.1)

where the dimension $d = 2$ or 3, and the interaction force $f = (f_1, \dots, f_d)$. u is divergence-free, i.e. $\nabla \cdot u = 0$, and the temperature transport equation may be written as

$$
\left(\kappa \Delta - u \cdot \nabla - \frac{\partial}{\partial t}\right) \theta = 0 \quad \text{in } \mathbb{R}^d. \tag{5.2}
$$

Let $\omega = \nabla \wedge u$ and $\Theta = \nabla \theta$. If $d = 3$, the vorticity transport equation for ω is the following PDE

$$
\left(\nu\Delta - u\cdot\nabla - \frac{\partial}{\partial t}\right)\omega + A\omega + F = 0 \quad \text{in } \mathbb{R}^3,
$$

where $(A\omega)^i = A^i_l \omega^l$ and

$$
F(x,t) = \Theta(x,t) \wedge f'(\theta(x,t)),
$$
 with $f' = (f'_1, f'_2, f'_3).$

For example, we may take $f_i(\theta) = (\theta - \theta_0)\delta_{i3}, i = 1, 2, 3$, then

$$
F^{i} = \delta_{i1}\Theta_{2} - \delta_{i2}\Theta_{1} - \delta_{i1}\frac{\partial\theta_{0}}{\partial x_{2}} + \delta_{i2}\frac{\partial\theta_{0}}{\partial x_{1}}.
$$

For two-dimensional flows,

$$
\left(\nu\Delta - u \cdot \nabla - \frac{\partial}{\partial t}\right)\omega + F = 0 \quad \text{in } \mathbb{R}^2
$$

where $F = \Theta_2 f'_1(\theta) - \Theta_1 f'_2(\theta)$.

The temperature gradient Θ satisfies the following transport equations

$$
\left(\kappa \Delta - u \cdot \nabla - \frac{\partial}{\partial t}\right) \Theta - A\Theta = 0 \quad \text{in } \mathbb{R}^d \tag{5.3}
$$

where $(A\Theta)_j = A_j^l \Theta_l$.

5.2 Representations of the vorticity and the temperature gradient

In this part, we shall work out the functional integral representations for the vorticity and the temperature gradient. We simplify our notation and omit u in the notation of transition probability densities, using $p_{\nu}(s, \eta, t, \xi)$ and $p_{\kappa}(s, \eta, t, \xi)$ instead of $p_{\nu, u}(s, \eta, t, \xi)$ and $p_{\kappa, u}(s, \eta, t, \xi)$.

Lemma 5.1. The temperature gradient $\Theta = \nabla \theta$ has the following integral representation:

$$
\Theta(\xi, t) = \int_{\mathbb{R}^d} \mathbb{E} \left[R(\eta, t; 0) \Theta(\eta, 0) \middle| Y_t^{\eta} = \xi \right] p_{\kappa}(0, \eta, t, \xi) d\eta
$$

for $\xi \in \mathbb{R}^d$ and $t > 0$, where R are defined by (4.4) with $D = \mathbb{R}^d$ $(d = 2 \text{ or } d = 3)$. Here we have used the convention that $(R\Theta)_j = R_j^l \Theta_l$.

Proof. Apply Theorem [2.4](#page-11-1) to the parabolic equations [\(5.3\)](#page-14-1).

Lemma 5.2. (1) Suppose $d = 3$. The vorticity $\omega = \nabla \wedge u$ possesses the following integral representation:

$$
\omega(\xi, t) = \int_{\mathbb{R}^3} \mathbb{E} \left[Q(\eta, t; 0) \omega(\eta, 0) \middle| X_t^{\eta} = \xi \right] p_{\nu}(0, \eta, t, \xi) d\eta
$$

$$
+ \int_0^t \int_{\mathbb{R}^3} \mathbb{E} \left[Q(\eta, t; s) F(X_s^{\eta}, s) \middle| X_t^{\eta} = \xi \right] p_{\nu}(0, \eta, t, \xi) d\eta ds \tag{5.4}
$$

 \Box

for $\xi \in \mathbb{R}^3$ and $t > 0$, where Q is defined by (4.3) with $D = \mathbb{R}^3$. Here $(Q\omega)^i = Q^i_l \omega^l$ and $(QF)^i = Q_l^i F^l$.

(2) Suppose $d=2$, so that $\omega = \frac{\partial}{\partial x}$ $\frac{\partial}{\partial x_1}u^2 - \frac{\partial}{\partial x}$ $\frac{\partial}{\partial x_2}u^1$, and $F = \Theta_1 f'_2(\theta) - \Theta_2 f'_1(\theta)$. ω has the following representation:

$$
\omega(\xi, t) = \int_{\mathbb{R}^2} p_\nu(0, \eta, t, \xi) \omega(\eta, 0) d\eta
$$

+
$$
\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[F(X_s^{\eta}, s) \right] X_t^{\eta} = \xi \left] p_\nu(0, \eta, t, \xi) d\eta ds \tag{5.5}
$$

for $\xi \in \mathbb{R}^2$ and $t > 0$.

The proof also follows from Theorem [2.4](#page-11-1) immediately.

5.3 Random vortex dynamics

Using the integral representations derived above, we are able to identify the random vortex dynamical systems associated with the heat-conducting incompressible fluid flows in \mathbb{R}^2 and \mathbb{R}^3 .

5.3.1 Random vortex dynamics in two-dimensional case

Now we are in a position to formulate the vortex dynamics for the model. Let us start with the two-dimensional model. As we have pointed out that for two-dimensional flows the vorticity ω is a scalar function, and $A\omega$ vanishes, so that one does not need the gauge functional Q.

Let $u(x, t)$ and $\theta(x, t)$ be the regular solutions to the model [\(5.1,](#page-13-5) [5.2\)](#page-14-2) when $d = 2$.

Theorem 5.3. The following representation holds:

$$
u(x,t) = \int_{\mathbb{R}^2} \mathbb{E}\left[K_2(X_t^{\eta}, x) \wedge \omega(\eta, 0)\right] d\eta
$$

$$
+ \int_0^t \int_{\mathbb{R}^2} \mathbb{E}\left[K_2(X_t^{\eta}, x) \wedge F(X_s^{\eta}, s)\right] d\eta ds
$$

for $x \in \mathbb{R}^2$ and $t > 0$.

Proof. By using the Biot-Savart law (see Lemma [3.1,](#page-11-2) Eq. [\(5.5\)](#page-15-2)), we obtain that

$$
u(x,t) = \int_{\mathbb{R}^2} K_2(\xi, x) \wedge \omega(\xi, t) d\xi
$$

=
$$
\int_{\mathbb{R}^2} K_2(\xi, x) \wedge \left(\int_{\mathbb{R}^2} p_{\nu}(0, \eta, t, \xi) \omega(\eta, 0) d\eta \right) d\xi
$$

+
$$
\int_{\mathbb{R}^2} K_2(\xi, x) \wedge \left(\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[F(X_s^{\eta}, s) \right] X_t^{\eta} = \xi \right] p_{\nu}(0, \eta, t, \xi) d\eta ds \right) d\xi
$$

and the conclusion then follows from an application of Fubini's theorem.

Similarly, we have the following theorem.

Theorem 5.4. The following representation for the temperature $\theta(x, t)$ holds:

$$
\theta(x,t) = -\int_{\mathbb{R}^2} \mathbb{E}\left[K_2(Y_t^{\eta},x) \cdot R(\eta,t;0)\Theta(\eta,0)\right] d\eta
$$

for $x \in \mathbb{R}^2$ and $t > 0$, where $R = (R_i^j)$ \mathcal{L}_i^j is defined by Eq. [\(4.4\)](#page-13-3), and $R\Theta = (R_i^l \Theta_l)$. Proof. This follows from [\(3.1\)](#page-11-2) and Lemma [5.1](#page-14-3) immediately.

 \Box

 \Box

Theorem [5.3,](#page-15-3) Theorem [5.4,](#page-15-4) the X particle equation (4.1) , the Y particle equation (4.2) , the gauge equation [\(4.4\)](#page-13-3) (with $D = \mathbb{R}^2$), together with the equations

$$
A_j^i = \frac{\partial}{\partial x_j} u^i, \quad \Theta_i = \frac{\partial}{\partial x_i} \theta \tag{5.6}
$$

and

$$
F(x,t) = \Theta_2(x,t)f'_1(\theta(x,t)) - \Theta_1(x,t)f'_2(\theta(x,t))
$$

give rise to a closed random vortex system for the two-dimensional model.

5.3.2 Random vortex dynamics in three-dimensional case

In this part, we work out the random vortex system for three-dimensional flows defined by the equations of motion $(5.1, 5.2)$ $(5.1, 5.2)$ $(5.1, 5.2)$ where $d = 3$, and the details of computations will be omitted.

Let $u(x, t)$ and $\theta(x, t)$ be the regular solutions to [\(5.1,](#page-13-5) [5.2\)](#page-14-2) with $d = 3$.

Theorem 5.5. The following representation formula for the velocity $u(x, t)$ holds:

$$
u(x,t) = \int_{\mathbb{R}^3} \mathbb{E}\left[K_3(X_t^{\eta}, x) \wedge Q(\eta, t; 0)\omega(\eta, 0)\right] d\eta
$$

$$
+ \int_{\mathbb{R}^3} \int_0^t \mathbb{E}\left[K_3(X_t^{\eta}, x) \wedge Q(\eta, t; s) F(X_s^{\eta}, s)\right] d\eta ds
$$

for $x \in \mathbb{R}^3$ and $t > 0$, where $Q\omega = (Q^i_l\omega^l)$ and $QF = (Q^i_lF^l)$, $Q = (Q^i_j)$ is defined by Eq. [\(4.3\)](#page-13-4) with $D = \mathbb{R}^3$.

This formula follows from Eq. [\(3.2\)](#page-11-3) and Eq. [\(5.4\)](#page-14-4).

Theorem 5.6. The temperature $\theta(x,t)$ possesses the following representation:

$$
\theta(x,t) = -\int_{\mathbb{R}^3} \mathbb{E}\left[K_3(Y_t^{\eta},x) \cdot R(\eta,t;0)\Theta(\eta,0)\right] d\eta
$$

for every $x \in \mathbb{R}^3$ and $t > 0$, where $R = (R_i^j)$ $i_j^{(j)}$ is defined by [\(4.4\)](#page-13-3) with $D = \mathbb{R}^3$, and $R\Theta = (R_i^l \Theta_l)$.

Theorem [5.5,](#page-16-2) Theorem [5.6,](#page-16-3) the particle defining equations [\(4.1,](#page-13-6) [4.2\)](#page-13-7), the gauge functional defining equations [\(4.3,](#page-13-4) [4.4\)](#page-13-3), together with the relations

$$
A_j^i = \frac{\partial}{\partial x_j} u^i, \quad \Theta_i = \frac{\partial}{\partial x_i} \theta, \quad F = \Theta \wedge f'(\theta)
$$

give rise to the random vortex dynamic system for three-dimensional flows.

6 Wall-bounded flows

In this section, we study the random vortex dynamics associated with the Oberbeck-Boussinesq flows along a flat plate, which is very important examples of wall-bounded fluid flows, in particular, wall-bounded turbulent flows with thermal conduction from the solid wall.

6.1 The Oberbeck-Boussinesq equations in \mathbb{R}^d_+

In the Oberbeck-Boussinesq model, it is assumed that the fluid density ρ diverges little from the fluid density ρ_0 at the fluid temperature θ_0 :

$$
\rho = \rho_0 \left(1 - \alpha \left(\theta - \theta_0 \right) \right),
$$

where θ is the fluid temperature and α is a very small constant. Therefore the density ρ is considered a constant. The Oberbeck-Boussinesq model is composed of the following partial differential equations on \mathbb{R}^d_+ (where the dimension $d=3$ or $d=2$). The Navier-Stokes equations with interaction from the heat supply on the boundary

$$
\left(\nu\Delta - u\cdot\nabla - \frac{\partial}{\partial t}\right)u - \nabla\left(\frac{P}{\rho_0} + gx_d\right) + f(\theta - \theta_0) = 0 \quad \text{in } \mathbb{R}^d_+,
$$

where the interaction force $f(\theta - \theta_0)$, with its components

$$
f_i(\theta - \theta_0) = \alpha g(\theta - \theta_0)\delta_{id}, \quad i = 1, \cdots, d.
$$

Here q is the gravity constant, and P denotes the pressure. The continuity equation

$$
\nabla \cdot u = 0 \quad \text{ in } \mathbb{R}^d_+,
$$

and the equation of energy in terms of the temperature θ :

$$
\left(\kappa \Delta - u \cdot \nabla - \frac{\partial}{\partial t}\right) \theta = 0 \quad \text{in } \mathbb{R}^d_+,
$$
\n(6.1)

where $\kappa = k/(\rho_0 c_P)$ (perfect gas) or $k/(\rho_0 c)$ (for liquid) is a positive constant.

The velocity $u(x, t)$ satisfies the no-slip condition, i.e. $u(x, t) = 0$ when $x_d = 0$, so that $u(x,t)$ is extended via reflection about $\{x_d = 0\}$ to a time-dependent vector field on \mathbb{R}^d :

$$
u^{i}(x,t) = u^{i}(\bar{x},t)
$$
 for $i = 1, \dots, d-1$, $u^{d}(x,t) = -u^{d}(\bar{x},t)$,

so that $u(\cdot, t)$ is divergence-free on the whole space \mathbb{R}^d in the distribution sense.

Let $\omega = \nabla \wedge u$ and $\Theta = \nabla \theta$. We assume that they are continuous up to the boundary ${x_d = 0}.$

The vorticity transport equation for three-dimensional flows is given as the following:

$$
\left(\nu\Delta - u\cdot\nabla - \frac{\partial}{\partial t}\right)\omega + A\omega + F = 0 \quad \text{in } \mathbb{R}^3_+,
$$

where $A = (A_l^i)$ with $A_l^i = \frac{\partial}{\partial x_l}$ $\frac{\partial}{\partial x_l} u^i$ for $i, l = 1, 2, 3, (\mathcal{A} \omega)^i = A^i_l \omega^l$ and

$$
F = (\Theta - \nabla \theta_0) \wedge f'(\theta - \theta_0).
$$

For two-dimensional flows, the non-linear vorticity stretching term $A\omega$ vanishes identically. Therefore

$$
\left(\nu\Delta - u\cdot\nabla - \frac{\partial}{\partial t}\right)\omega + F = 0 \quad \text{in } \mathbb{R}^2_+,
$$

where

$$
F = (\Theta_1 - \frac{\partial}{\partial x_1} \theta_0) f_2' (\theta - \theta_0) - (\Theta_2 - \frac{\partial}{\partial x_2} \theta_0) f_1' (\theta - \theta_0).
$$

Similarly, by differentiating the heat equation [\(6.1\)](#page-17-1) one obtains the evolution equations for the temperature gradient

$$
\left(\kappa \Delta - u \cdot \nabla - \frac{\partial}{\partial t}\right) \Theta - A\Theta = 0 \quad \text{in } \mathbb{R}^d_+.
$$

Here we have used the convention that $(A\Theta)_j = A_j^l \Theta_l$.

An essential difference exists between this bounded case and the case discussed in Section 4, where there is no physical boundary for the fluid flow. For wall-bounded flows, it is impossible to determine the boundary value of the vorticity ω and the temperature gradient Θ , although the boundary vorticity can be identified as the normal stress at the boundary, cf. Anderson [?]. Nevertheless, the boundary the temperature gradient can not be specified either, while by the definition of the Oberbeck-Boussinesq motion, the boundary temperature gradient is considered small, so it can be treated as zero in numerical experiments.

6.2 Diffusions in \mathbb{R}^d_+

Let $D = \mathbb{R}^d_+$. The reflection $\mathscr R$ about the hyperplane $\{x_d = 0\}$ on \mathbb{R}^d is the linear map which sends $x = (x_1, \dots, x_d)$ to $\Re x = \overline{x} = (x_1, \dots, x_{d-1}, -x_d)$ for every $x \in \mathbb{R}^d$.

The velocity $u(x, t)$ is extended to a time-dependent and divergence-free vector field on \mathbb{R}^d such that $\mathscr{R}(b(x,t)) = b(\mathscr{R}x, t)$ for every $x \in \mathbb{R}^d$ and $t \geq 0$. That is

$$
u^{i}(\overline{x}, t) = u^{i}(x, t)
$$
 for $i = 1, ..., d - 1$ and $u^{d}(\overline{x}, t) = -u^{d}(x, t)$)

for all $x \in \mathbb{R}^d$. Then $\overline{X^{\xi,\tau}}$ and $X^{\overline{\xi},\tau}$ have the same distribution, so that $p_{\lambda,u}(\tau,\overline{\xi},t,\overline{x}) =$ $p_{\lambda,u}(\tau,\xi,t,x)$, and

$$
p_{\lambda,u}^D(\tau,\xi,t,x) = p_{\lambda,u}(\tau,\xi,t,x) - p_{\lambda,u}(\tau,\overline{\xi},t,x)
$$
\n(6.2)

for any $\xi, x \in \mathbb{R}^d_+$. The last equality may be verified by checking that the right-hand side is indeed the Green function of the backward parabolic operator $\lambda\Delta + b\cdot\nabla + \frac{\partial}{\partial t}$. In particular

$$
\mathbb{E}\left[\varphi(X_t^{\xi,\tau})1_{\{t<\zeta(X^{\xi,\tau})\}}\right] = \mathbb{E}\left[1_D(X_t^{\xi,\tau})\varphi(X_t^{\xi,\tau})\right] - \mathbb{E}\left[1_D(X_t^{\overline{\xi},\tau})\varphi(X_t^{\overline{\xi},\tau})\right] \tag{6.3}
$$

for any bounded and Borel measurable function φ . This approach has been put forward in [\[61\]](#page-44-15) for simulating the solutions of the Navier-Stokes equations within the boundary layer.

6.3 Representations of the vorticity and the temperature gradient

We observe that the vorticity ω , the temperature θ and the temperature gradient Θ possess non-homogeneous Dirichlet boundary conditions, i.e.

$$
\omega(x,t) = \sigma(x_1, \cdots, x_{d-1}, t) \quad \text{on } \{x_d = 0\},
$$

$$
\Theta(x,t) = \gamma(x_1, \cdots, x_{d-1}, t) \quad \text{on } \{x_d = 0\}
$$

and

$$
\theta(x,t) = \theta_0(x_1, \dots, x_{d-1})
$$
 on $\{x_d = 0\}$

for all $t \geq 0$, where θ_0 is considered the external heat supply on the boundary. so we use the cutting-off technique to reduce their boundary conditions to homogeneous ones.

Let $\phi : [0, \infty) \to [0, 1]$ be a smooth cut-off function such that $\phi(r) = 1$ for $r \in [0, 1/3)$ and $\phi(r) = 0$ for $r \geq 2/3$. Let

$$
\omega^{\varepsilon}(x,t) = \omega(x,t) - \sigma(x_1,\cdots,x_{d-1},t)\phi\left(\frac{x_d}{\varepsilon}\right)
$$

and

$$
\theta^{\varepsilon}(x,t) = \theta(x,t) - \theta_0(x_1,\cdots,x_{d-1})\phi\left(\frac{x_d}{\varepsilon}\right)
$$

for every $\varepsilon > 0$. It is clear from the definition that

$$
\lim_{\varepsilon \to \infty} \omega^{\varepsilon}(x,t) = \omega(x,t) - 1_{\{x_d \ge 0\}} \sigma(x_1, \cdots, x_{d-1}, t),
$$

$$
\lim_{\varepsilon \to \infty} \theta^{\varepsilon}(x, t) = \theta(x, t) - 1_{\{x_d \ge 0\}} \theta_0(x_1, \dots, x_{d-1}),
$$

$$
\lim_{\varepsilon \to 0+} \omega^{\varepsilon}(x, t) = \omega(x, t) - 1_{\{x_d = 0\}} \sigma(x_1, \dots, x_{d-1}, t)
$$

and

$$
\lim_{\varepsilon \to 0+} \theta^{\varepsilon}(x,t) = \theta(x,t) - 1_{\{x_d=0\}} \theta_0(x_1,\cdots,x_{d-1}).
$$

Then ω^{ε} and θ^{ε} satisfy the homogeneous Dirichlet boundary conditions $\omega^{\varepsilon}(x,t) = 0$ and $\theta^{\varepsilon}(x,t) = 0$ on $\{x_d = 0\}$. ω^{ε} and θ^{ε} , for every $\varepsilon > 0$, evolve according to the following parabolic equations:

$$
\left(\nu\Delta - u \cdot \nabla - \frac{\partial}{\partial t}\right)\omega^{\varepsilon} + A\omega^{\varepsilon} + F + \chi_{\varepsilon} = 0 \quad \text{in } \mathbb{R}^{d}_{+},\tag{6.4}
$$

(here if the dimension $d = 2$, then the non-linear stretching term $A\omega^{\varepsilon}$ may be dropped),

$$
\left(\kappa \Delta - u \cdot \nabla - \frac{\partial}{\partial t}\right) \Theta^{\varepsilon} - A \Theta^{\varepsilon} + \alpha_{\varepsilon} = 0 \quad \text{in } \mathbb{R}^d_+ \tag{6.5}
$$

and

$$
\left(\kappa\Delta - u \cdot \nabla - \frac{\partial}{\partial t}\right)\theta^{\varepsilon} + \beta_{\varepsilon} = 0 \quad \text{in } \mathbb{R}^d_+,
$$
\n(6.6)

where the correction terms are given by the following formulae:

$$
\chi_{\varepsilon}(x,t) = \phi\left(\frac{x_d}{\varepsilon}\right) \left(\nu\Delta - \sum_{j=1}^{d-1} u^j(x,t) \frac{\partial}{\partial x_j} - \frac{\partial}{\partial t} + A\right) \sigma(x_1, \dots, x_{d-1}, t) \n- \frac{1}{\varepsilon} \phi'\left(\frac{x_d}{\varepsilon}\right) \sigma(x_1, \dots, x_{d-1}, t) u^d(x,t) + \nu \frac{1}{\varepsilon^2} \phi''\left(\frac{x_d}{\varepsilon}\right) \sigma(x_1, \dots, x_{d-1}, t), \quad (6.7)
$$

(if the dimension $d = 2$, then the term involving A may be dropped),

$$
\alpha_{\varepsilon}(x,t) = \phi\left(\frac{x_d}{\varepsilon}\right) \left(\kappa \Delta - \sum_{j=1}^{d-1} u^j(x,t) \frac{\partial}{\partial x_j} - \frac{\partial}{\partial t} - A\right) \gamma(x_1, \dots, x_{d-1}, t) \n- \frac{1}{\varepsilon} \phi'\left(\frac{x_d}{\varepsilon}\right) \gamma(x_1, \dots, x_{d-1}, t) u^d(x,t) + \kappa \frac{1}{\varepsilon^2} \phi''\left(\frac{x_d}{\varepsilon}\right) \gamma(x_1, \dots, x_{d-1}, t),
$$

and

$$
\beta_{\varepsilon}(x,t) = \phi\left(\frac{x_d}{\varepsilon}\right) \left(\kappa \Delta - \sum_{j=1}^{d-1} u^j \frac{\partial}{\partial x_j}\right) \theta_0(x_1,\dots,x_{d-1})
$$

$$
-\frac{1}{\varepsilon} \phi'\left(\frac{x_d}{\varepsilon}\right) \theta_0(x_1,\dots,x_{d-1}) u^d(x,t) + \kappa \frac{1}{\varepsilon^2} \phi''\left(\frac{x_d}{\varepsilon}\right) \theta_0(x_1,\dots,x_{d-1}).
$$

Here $x = (x_1, \dots, x_d)$ and the Laplacian Δ on σ and θ_0 is the (boundary) $(d-1)$ -dimensional Laplacian $\sum_{j=1}^{d-1} \frac{\partial^2}{\partial x^2}$ $\frac{\partial^2}{\partial x_j^2}$.

It is clear from the definition that the following limits

$$
\lim_{\varepsilon \to \infty} \chi_{\varepsilon}(x, t) = 1_{\{x_d \ge 0\}} \left(\nu \Delta - \sum_{j=1}^{d-1} u^j(x, t) \frac{\partial}{\partial x_j} - \frac{\partial}{\partial t} + A \right) \sigma(x_1, \dots, x_{d-1}, t),
$$

$$
\lim_{\varepsilon \to \infty} \alpha_{\varepsilon}(x, t) = 1_{\{x_d \ge 0\}} \left(\kappa \Delta - \sum_{j=1}^{d-1} u^j(x, t) \frac{\partial}{\partial x_j} - \frac{\partial}{\partial t} - A \right) \gamma(x_1, \dots, x_{d-1}, t),
$$

and

$$
\lim_{\varepsilon \to \infty} \beta_{\varepsilon}(x, t) = 1_{\{x_d \ge 0\}} \left(\kappa \Delta - \sum_{j=1}^{d-1} u^j(x, t) \frac{\partial}{\partial x_j} \right) \theta_0(x_1, \dots, x_{d-1})
$$

exist, where $x = (x_1, \dots, x_d)$. Unfortunately, these limits involve the velocity of the outer layer flow, and the limits as $\varepsilon \to 0+$ do not exist in the ordinary sense, so we need to evaluate them under integration. However, since $u(x, t)$ satisfies the no-slip condition, so that

$$
\lim_{\varepsilon \to 0+} \phi\left(\frac{x_d}{\varepsilon}\right) \left(\nu \Delta - \sum_{j=1}^{d-1} u^j \frac{\partial}{\partial x_j} - \frac{\partial}{\partial t} + A\right) \sigma = 1_{\{x_d = 0\}} \left(\nu \Delta - \frac{\partial}{\partial t} + A\right) \sigma, \tag{6.8}
$$

$$
\lim_{\varepsilon \to 0+} \phi\left(\frac{x_d}{\varepsilon}\right) \left(\kappa \Delta - \sum_{j=1}^{d-1} u^j \frac{\partial}{\partial x_j} - \frac{\partial}{\partial t} - A\right) \gamma = 1_{\{x_d = 0\}} \left(\kappa \Delta - \frac{\partial}{\partial t} - A\right) \gamma,
$$

and

$$
\lim_{\varepsilon \to 0+} \phi\left(\frac{x_2}{\varepsilon}\right) \left(\kappa \Delta - \sum_{j=1}^{d-1} u^j \frac{\partial}{\partial x_j}\right) \theta_0 = \kappa \mathbb{1}_{\{x_d = 0\}} \Delta \theta_0,
$$

which vanish almost surely in $x = (x_1, \dots, x_d)$ with respect to the Lebesgue measure on \mathbb{R}^d .

By using the forward Feynman-Kac formula, we may work out the functional integral representations for θ^{ε} , Θ^{ε} and ω^{ε} respectively. Again, as in previous sections, we drop u in the notation of the transition probability density functions.

Proposition 6.1. For every $\varepsilon > 0$, the temperature θ has the following representation

$$
\theta(\xi, t) = \phi\left(\frac{\xi_d}{\varepsilon}\right) \theta_0(\xi_1, \dots, \xi_{d-1}) + \int_{\mathbb{R}_+^d} \mathbb{E}\left[1_{\{t < \zeta(Y^{\eta})\}} \middle| Y_t^{\eta} = \xi\right] p_{\kappa}(0, \eta, t, \xi) \theta^{\varepsilon}(\eta, 0) d\eta
$$

$$
+ \int_0^t \int_{\mathbb{R}_+^d} \mathbb{E}\left[1_{\{s > \gamma_t(Y^{\eta})\}} \beta_{\varepsilon}(Y_s^{\eta}, s) \middle| Y_t^{\eta} = \xi\right] p_{\kappa}(0, \eta, t, \xi) d\eta dt
$$

 \Box

for every $\xi \in \mathbb{R}^d_+$ and $t > 0$, where $d = 2$ or $d = 3$.

Proof. It follows from the PDE [\(6.6\)](#page-19-0) and Theorem [2.1](#page-7-4) immediately.

There is a similar representation for the temperature gradient.

Proposition 6.2. For every $\varepsilon > 0$, the following representation holds:

$$
\Theta(\xi, t) = \phi \left(\frac{\xi_d}{\varepsilon}\right) \Theta(\xi_1, \dots, \xi_{d-1}, 0, t)
$$

+
$$
\int_{\mathbb{R}_+^d} \mathbb{E} \left[1_{\{t < \zeta(Y^{\eta})\}} R(\eta, t; 0) \Theta^\varepsilon(\eta, 0) \Big| Y_t^{\eta} = \xi\right] p_\kappa(0, \eta, t, \xi) d\eta
$$

+
$$
\int_0^t \int_{\mathbb{R}_+^d} \mathbb{E} \left[1_{\{s > \gamma_t(Y^{\eta})\}} R(\eta, t; s) \alpha_\varepsilon(Y_s^{\eta}, s) \Big| Y_t^{\eta} = \xi\right] p_\kappa(0, \eta, t, \xi) d\eta ds
$$

for all $\xi \in \mathbb{R}^d_+$ and $t > 0$, where $d = 2$ and $d = 3$, and $(R\Theta^{\varepsilon})_j = R^l_j \Theta^{\varepsilon}_l$.

The representation for the temperature follows immediately from [\(6.5\)](#page-19-1) and Theorem [2.1.](#page-7-4)

Finally, the functional integral representation for the vorticity in dimension two takes a simpler form, so we state it separately.

Proposition 6.3. Suppose $d = 2$. Then the vorticity ω possesses the following integral representation:

$$
\omega(\xi, t) = \phi\left(\frac{\xi_2}{\varepsilon}\right) \sigma(\xi_1, t) + \int_{\mathbb{R}_+^2} \mathbb{E}\left[1_{\{t < \zeta(X^{\eta})\}} \Big| X_t^{\eta} = \xi\right] p_{\nu}(0, \eta, t, \xi) \omega^{\varepsilon}(\eta, 0) d\eta
$$

$$
+ \int_0^t \int_{\mathbb{R}_+^2} \mathbb{E}\left[1_{\{s > \gamma_t(X^{\eta})\}} F(X_s^{\eta}, s) \Big| X_t^{\eta} = \xi\right] p_{\nu}(0, \eta, t, \xi) d\eta ds
$$

$$
+ \int_0^t \int_{\mathbb{R}_+^2} \mathbb{E}\left[1_{\{s > \gamma_t(X^{\eta})\}} \chi_{\varepsilon}(X_s^{\eta}, s) \Big| X_t^{\eta} = \xi\right] p_{\nu}(0, \eta, t, \xi) d\eta ds
$$

for every $\xi = (\xi_1, \xi_2) \in \mathbb{R}^2_+$ and $t > 0$.

In particular, for two-dimensional flow, we do not need to introduce the gauge functional Q. However, for three-dimensional flows, we need both gauge functionals Q and R , defined by (4.3) and [\(4.4\)](#page-13-3) respectively with $D = \mathbb{R}^3_+$.

Proposition 6.4. Suppose $d = 3$. Then the vorticity ω has the functional integral representation given by

$$
\omega(\xi, t) = \phi\left(\frac{\xi_3}{\varepsilon}\right) \sigma(\xi_1, \xi_2, t) + \int_{\mathbb{R}_+^3} \mathbb{E}\left[\left[1_{\{t < \zeta(X^{\eta})\}} Q(\eta, t; 0)\omega^{\varepsilon}(\eta, 0)\right| X_t^{\eta} = \xi\right] p_{\nu}(0, \eta, t, \xi) d\eta
$$

$$
+ \int_0^t \int_{\mathbb{R}_+^3} \mathbb{E}\left[1_{\{s > \gamma_t(X^{\eta})\}} Q(\eta, t; s) F(X_s^{\eta}, s)\right| X_t^{\eta} = \xi\right] p_{\nu}(0, \eta, t, \xi) d\eta ds
$$

$$
+ \int_0^t \int_{\mathbb{R}_+^3} \mathbb{E}\left[1_{\{s > \gamma_t(X^{\eta})\}} Q(\eta, t; s) \chi_{\varepsilon}(X_s^{\eta}, s)\right| X_t^{\eta} = \xi\right] p_{\nu}(0, \eta, t, \xi) d\eta ds
$$

for every $\xi = (\xi_1, \xi_2, \xi_3) \in \mathbb{R}^3_+$ and $t > 0$, where $(Q\omega^{\varepsilon})^i = Q^i_j(\omega^{\varepsilon})^j$ and $(QF)^i = Q^i_j F^j$.

These functional integral representations are consequences of the vorticity transport equation [\(6.4\)](#page-19-2) and Theorem [2.1](#page-7-4) immediately.

6.4 Representations of the velocity and the temperature

To avoid repetition, unlike in the unbounded domain, we state the representations of velocity and temperature for $d = 2$ and $d = 3$ together and establish random vortex systems in the next section separately.

Combining with the Biot-Savart laws, we may establish various integral functional representations for the velocity and the temperature.

Firstly, using Proposition [6.2](#page-20-0) and the Biot-Savart law (cf. Lemma [3.3\)](#page-12-1), we may deduce the following.

Theorem 6.5. For the temperature θ , the following functional integral representation holds: (1) When $d = 2$,

$$
\theta(x,t) = \frac{x_2}{\pi} \int_{-\infty}^{\infty} \frac{\theta_0(\xi_1)}{|\xi_1 - x_1|^2 + x_2^2} d\xi_1 \n- \int_{0}^{\varepsilon} \left[\phi\left(\frac{\xi_2}{\varepsilon}\right) \int_{\mathbb{R}} A_2(\xi, x) \cdot \Theta((\xi_1, 0), t) d\xi_1 \right] d\xi_2 \n- \int_{\mathbb{R}_+^2} \mathbb{E} \left[1_{\{t < \zeta(Y^{\eta})\}} 1_{\mathbb{R}_+^2} (Y_t^{\eta}) A_2(Y_t^{\eta}, x) \cdot R(\eta, t; 0) \Theta^{\varepsilon}(\eta, 0) \right] d\eta \n- \int_{0}^t \int_{\mathbb{R}_+^2} \mathbb{E} \left[1_{\{s > \gamma_t(Y^{\eta})\}} 1_{\mathbb{R}_+^2} (Y_t^{\eta}) A_2(Y_t^{\eta}, x) \cdot R(\eta, t; s) \alpha_{\varepsilon}(Y_s^{\eta}, s) \right] d\eta ds \qquad (6.9)
$$

for $x \in \mathbb{R}_+^2$ and $t > 0$, where $(R\Theta^{\varepsilon})_j = R_j^l \Theta_l^{\varepsilon}$.

(2) When $d = 3$,

$$
\theta(x,t) = \frac{x_3}{2\pi} \int_{\mathbb{R}^2} \frac{\theta_0(\xi_1, \xi_2)}{|\xi_1 - x_2|^2 + |\xi_2 - x_2|^2 + |x_3|^2} d\xi_1 d\xi_2 \n- \int_0^{\varepsilon} \left[\phi \left(\frac{\xi_3}{\varepsilon} \right) \int_{\mathbb{R}^2} A_3(\xi, x) \cdot \Theta(\xi_1, \xi_2, 0, t) d\xi_1 d\xi_2 \right] d\xi_3 \n- \int_{\mathbb{R}^3_+} \mathbb{E} \left[1_{\{t < \zeta(Y^{\eta})\}} 1_{\mathbb{R}^3_+} (Y^{\eta}_t) A_3(Y^{\eta}_t, x) \cdot R(\eta, t; 0) \Theta^{\varepsilon}(\eta, 0) \right] d\eta \n- \int_0^t \int_{\mathbb{R}^3_+} \mathbb{E} \left[1_{\{s > \gamma_t(Y^{\eta})\}} 1_{\mathbb{R}^3_+} (Y^{\eta}_t) A_3(Y^{\eta}_t, x) \cdot R(\eta, t; s) \alpha_{\varepsilon} (Y^{\eta}_s, s) \right] d\eta ds \qquad (6.10)
$$

for $x \in \mathbb{R}^3_+$ and $t > 0$, where $(R\Theta^{\varepsilon})_j = R^l_j \Theta^{\varepsilon}_l$.

Similarly, by using the Biot-Savart law (cf. Lemma [3.2\)](#page-12-2), Proposition [6.3](#page-21-1) and Proposition [6.4,](#page-21-2) we may establish the following functional integral representations.

Theorem 6.6. Suppose $d = 2$. Then for every $\varepsilon > 0$

$$
u(x,t) = \int_0^{\varepsilon} \left[\phi\left(\frac{\xi_2}{\varepsilon}\right) \int_{-\infty}^{\infty} \Lambda_2(\xi, x) \wedge \sigma(\xi_1, t) d\xi_1 \right] d\xi_2
$$

+
$$
\int_{\mathbb{R}_+^2} \mathbb{E} \left[1_{\{t < \zeta(X^{\eta})\}} 1_{\mathbb{R}_+^2} (X_t^{\eta}) \Lambda_2(X_t^{\eta}, x) \wedge \omega^{\varepsilon}(\eta, 0) \right] d\eta
$$

+
$$
\int_0^t \int_{\mathbb{R}_+^2} \mathbb{E} \left[1_{\{s > \gamma_t(X^{\eta})\}} 1_{\mathbb{R}_+^2} (X_t^{\eta}) \Lambda_2(X_t^{\eta}, x) \wedge F(X_s^{\eta}, s) \right] d\eta ds
$$

+
$$
\int_0^t \int_{\mathbb{R}_+^2} \mathbb{E} \left[1_{\{s > \gamma_t(X^{\eta})\}} 1_{\mathbb{R}_+^2} (X_t^{\eta}) \Lambda_2(X_t^{\eta}, x) \wedge \chi_{\varepsilon}(X_s^{\eta}, s) \right] d\eta ds \qquad (6.11)
$$

for $x \in \mathbb{R}_+^2$ and $t > 0$, and $u(x, t) = \mathscr{R}(u(\overline{x}, t))$ if $x_2 < 0$.

In dimension 3, the representation of the velocity is much more complicated.

Theorem 6.7. Suppose $d = 3$. For every $\varepsilon > 0$, the velocity $u(x, t)$ has the following functional integral representation:

$$
u(x,t) = \int_0^{\varepsilon} \left(\phi\left(\frac{\xi_3}{\varepsilon}\right) \int_{\mathbb{R}^2} \Lambda_3(\xi, x) \wedge \sigma(\xi_1, \xi_2, t) d\xi_1 d\xi_2 \right) d\xi_3
$$

+
$$
\int_{\mathbb{R}^3_+} \mathbb{E}\left[1_{\{t < \zeta(X^{\eta})\}} 1_{\mathbb{R}^3_+} (X_t^{\eta}) \Lambda_3(X_t^{\eta}, x) \wedge Q(\eta, t; 0) \omega^{\varepsilon}(\eta, 0)\right] d\eta
$$

+
$$
\int_0^t \int_{\mathbb{R}^3_+} \mathbb{E}\left[1_{\{s > \gamma_t(X^{\eta})\}} 1_{\mathbb{R}^3_+} (X_t^{\eta}) \Lambda_3(X_t^{\eta}, x) \wedge Q(\eta, t; s) F(X_s^{\eta}, s)\right] d\eta ds
$$

+
$$
\int_0^t \int_{\mathbb{R}^3_+} \mathbb{E}\left[1_{\{s > \gamma_t(X^{\eta})\}} 1_{\mathbb{R}^3_+} (X_t^{\eta}) \Lambda_3(X_t^{\eta}, x) \wedge Q(\eta, t; s) \chi_{\varepsilon}(X_s^{\eta}, s)\right] d\eta ds.
$$
 (6.12)

for $x \in \mathbb{R}^3_+$ and $t > 0$, and $u(x,t) = \mathscr{R}(u(\overline{x}, t))$ if $x_3 < 0$, where the gauge functional Q is defined by [\(4.3\)](#page-13-4) with $D = \mathbb{R}^3_+$, $(Q\omega^{\varepsilon})^i = Q^i_j(\omega^{\varepsilon})^j$ and $(QF)^i = Q^i_jF^j$.

6.5 Random vortex dynamics

With the stochastic representation formulae derived in the previous subsections, we are ready to establish a system of random vortex dynamics for wall-bounded flows.

6.5.1 2D random vortex for wall-bounded flows

In this part, let us work out the simpler scenario, the two-dimensional case. When $d = 2$, the interaction force due to the heat conduction is given by

$$
f_1(x,t) = 0
$$
, and $f_2(x,t) = \theta(x,t) - \theta_0(x_1)$

(the constant factor αq is absorbed into the temperature θ for simplicity), so that

$$
F(x,t) = \Theta_1(x,t) - \frac{\partial}{\partial x_1} \theta_0(x_1).
$$
\n(6.13)

The random vortex dynamics may be defined as the following:

- Eq. (4.1) for X particles and Eq. (4.2) for Y particles, where in both equations, the velocity is extended by reflection so that $u(x,t) = 0$ when $x_2 = 0$, and $u^1(x,t) = u^1(\overline{x},t)$, $u^2(x,t) = -u^2(\overline{x},t).$
- Eq. [\(4.4\)](#page-13-3) with $D = \mathbb{R}^2_+$ that defines the gauge functional R.
- The representation (6.11) for the velocity and the representation (6.9) for the temperature $\theta(x,t)$.
- Other dynamical variables given by

$$
A_j^l(x,t) = \frac{\partial}{\partial x_j} u^l(x,t), \quad F(x,t) = \frac{\partial}{\partial x_1} \theta(x,t) - \frac{\partial}{\partial x_1} \theta_0(x_1), \tag{6.14}
$$

which lead to the stochastic representations for A_j^l and F, obtained by differentiating the functional integral representations for u^l and θ respectively.

During numerical experiments, we may choose $\varepsilon > 0$ sufficiently small, so that the contribution from

$$
- \int_0^{\varepsilon} \left[\phi\left(\frac{\xi_2}{\varepsilon}\right) \int_{-\infty}^{\infty} \Lambda_2(\xi, x) \cdot \Theta((\xi_1, 0), t) d\xi_1 \right] d\xi_2
$$

is negligible. Also, by the definition of the Oberbeck-Boussinesq model, the temperature gradient at the boundary is small, so it is reasonable to neglect the contribution from

$$
- \int_0^t \int_{\mathbb{R}_+^2} \mathbb{E} \left[1_{\{s > \gamma_t(Y^{\eta})\}} 1_{\mathbb{R}_+^2} (Y_t^{\eta}) A_2(Y_t^{\eta}, x) \cdot R(\eta, t; s) \alpha_{\varepsilon} (Y_s^{\eta}, s) \right] d\eta ds.
$$

Therefore, by sending $\varepsilon \downarrow 0$, we may calculate the temperature via the formula:

$$
\theta(x,t) \sim \frac{x_2}{\pi} \int_{-\infty}^{\infty} \frac{\theta_0(\xi_1)}{|\xi_1 - x_1|^2 + x_2^2} d\xi_1 \n- \int_{\mathbb{R}_+^2} \mathbb{E} \left[1_{\{t < \zeta(Y^{\eta})\}} 1_{\mathbb{R}_+^2} (Y_t^{\eta}) A_2(Y_t^{\eta}, x) \cdot R(\eta, t; 0) \Theta(\eta, 0) \right] d\eta.
$$
\n(6.15)

Similarly, for the velocity $u(x, t)$, the contribution, when $\varepsilon > 0$ is small enough, we may ignore the minor contribution from

$$
\int_0^{\varepsilon} \left[\phi\left(\frac{\xi_2}{\varepsilon}\right) \int_{-\infty}^{\infty} \Lambda_2(\xi, x) \wedge \sigma(\xi_1, t) \mathrm{d}\xi \right] \mathrm{d}\xi_2.
$$

Furthermore, by sending $\varepsilon \downarrow 0$, we obtained the limiting representation for $u(x,t)$:

$$
u(x,t) = \int_{\mathbb{R}^2_+} \mathbb{E} \left[1_{\{t < \zeta(X^{\eta})\}} 1_{\mathbb{R}^2_+} (X_t^{\eta}) A_2(X_t^{\eta}, x) \wedge \omega(\eta, 0) \right] d\eta
$$

$$
+\int_0^t \int_{\mathbb{R}_+^2} \mathbb{E}\left[\mathbf{1}_{\{s>\gamma_t(X^{\eta})\}} \mathbf{1}_{\mathbb{R}_+^2}(X_t^{\eta}) A_2(X_t^{\eta}, x) \wedge F(X_s^{\eta}, s)\right] d\eta ds
$$

+
$$
\lim_{\varepsilon \downarrow 0} \int_0^t \int_{\mathbb{R}_+^2} \mathbb{E}\left[\mathbf{1}_{\{s>\gamma_t(X^{\eta})\}} \mathbf{1}_{\mathbb{R}_+^2}(X_t^{\eta}) A_2(X_t^{\eta}, x) \wedge \chi_{\varepsilon}(X_s^{\eta}, s)\right] d\eta ds,
$$
(6.16)

where χ_{ε} can be replaced by

$$
\tilde{\chi}_{\varepsilon}(x,t) = \nu \frac{1}{\varepsilon^2} \phi''\left(\frac{x_2}{\varepsilon}\right) \sigma(x_1,t),\tag{6.17}
$$

and σ is the trace of ω on the boundary $\{x_2 = 0\}.$

We will justify this procedure in Subsection [6.6](#page-24-1) below by showing that the limit of each term on the right-hand side of [\(6.16\)](#page-24-2) exists.

6.5.2 3D random vortex for wall-bounded flows

The computation is analogous to the previous part, except we need to include the gauge functional Q now.

Hence the random vortex dynamics in dimension three can be characterised by the following system of equations:

- The equation for X particle by (4.1) , the equation for Y particle by (4.2) , where in both equations, the velocity is extended by reflection so that $u(x,t) = 0$ when $x_3 = 0$, and $u^{1}(x,t) = u^{1}(\overline{x},t), u^{1}(x,t) = u^{1}(\overline{x},t), u^{3}(x,t) = -u^{3}(\overline{x},t).$
- The equation that defines the gauge functional Q and R, [\(4.3\)](#page-13-4) and [\(4.4\)](#page-13-3) with $D = \mathbb{R}^3_+$.
- The representation (6.12) for the velocity and the representation (6.10) for the temperature $\theta(x,t)$.
- Other dynamical variables given by

$$
A_j^l(x,t) = \frac{\partial}{\partial x_j} u^l(x,t),
$$

\n
$$
F^1(x,t) = \frac{\partial}{\partial x_2} \theta(x,t) - \frac{\partial}{\partial x_2} \theta_0(x_1,x_2),
$$

\n
$$
F^2(x,t) = -\frac{\partial}{\partial x_1} \theta(x,t) + \frac{\partial}{\partial x_1} \theta_0(x_1,x_2),
$$

\n
$$
F^3(x,t) = 0,
$$

which give rise to the representations for A_j^l and F by differentiating the functional integral representations for u^l and θ .

6.6 Limiting representations

In the previous parts, we have used the families of perturbations of the vorticity, the temperature and the temperature gradient to establish the random vortex dynamical systems. For the purpose of numerical simulations, we need to choose the parameter $\varepsilon > 0$ small enough so that the outer layer velocity of the fluid flow can be avoided. In this part, we demonstrate that the limiting representations as $\varepsilon \downarrow 0$ exist, which will justify the approximations [\(6.15\)](#page-23-1) and [\(6.16\)](#page-24-2).

Let us consider the two-dimensional case. Recall that the velocity $u(x, t)$ of the fluid flow is extended for all $x \in \mathbb{R}^2$, so that $\mathscr{R}(u(x,t)) = u(\overline{x}, t)$. Hence

$$
p_{\lambda,u}^D(s,\xi,t,x) = p_{\lambda,u}(\tau,\xi,t,x) - p_{\lambda,u}(\tau,\overline{\xi},t,x)
$$

is the Green function to the Dirichlet problem of the forward parabolic operator $L_{\lambda,-u} - \frac{\partial}{\partial t}$, where $D = \mathbb{R}^2_+$. In the computation below, we drop u in the notation of transition probabilities for simplicity. By using the vorticity transport equation [\(6.4\)](#page-19-2) for $\omega^{\epsilon}(x,t)$ and the temperature transport equation [\(6.6\)](#page-19-0) for $\theta^{\varepsilon}(x,t)$, according to [\[29,](#page-42-16) Theorem 12 on page 25], we obtain that

$$
\omega^{\varepsilon}(x,t) = \int_{D} p_{\nu}^{D}(0,\eta,t,x)\omega^{\varepsilon}(\eta,0) d\eta + \int_{0}^{t} \int_{D} p_{\nu}^{D}(s,\eta,t,x)F(\eta,s) d\eta ds + \int_{0}^{t} \int_{D} p_{\nu}^{D}(s,\eta,t,x)\chi_{\varepsilon}(\eta,s) d\eta ds
$$
\n(6.18)

and

$$
\theta^{\varepsilon}(x,t) = \int_{D} p_{\kappa}^{D}(0,\eta,t,x) \theta^{\varepsilon}(\eta,0) d\eta + \int_{0}^{t} \int_{D} p_{\kappa}^{D}(s,\eta,t,x) \beta_{\varepsilon}(\eta,s) d\eta ds \tag{6.19}
$$

for every $x \in D$ and $t > 0$. Before we present the proof, let us first introduce a new notation and a helpful result that apply to all dimensions.

Definition 6.8. If φ is a Borel measurable function on \mathbb{R}^d_+ , then define

$$
\hat{\varphi}(x) = 1_{\{x_d > 0\}} \varphi(x) - 1_{\{x_d < 0\}} \varphi(\overline{x})
$$
\n(6.20)

 \Box

for every $x \in \mathbb{R}^d$.

Lemma 6.9. Let $D = \mathbb{R}^d_+$ and $\lambda > 0$. Then it holds that

$$
\int_D p_\lambda^D(s,\eta,t,x)\varphi(\eta) d\eta = \int_{\mathbb{R}^d} p_{\lambda,u}(s,\eta,t,x)\hat{\varphi}(\eta) d\eta
$$

for every $x \in D$ and $t > s \geq 0$.

Proof. Since $\nabla \cdot u = 0$ on \mathbb{R}^d , by Eq. [\(6.2\)](#page-18-2),

$$
\int_{D} p_{\lambda}^{D}(s,\eta,t,x)\varphi(\eta) d\eta = \int_{\mathbb{R}^{d}} p_{\lambda}(s,\eta,t,x) 1_{\{\eta_{d}>0\}}\varphi(\eta) d\eta - \int_{\mathbb{R}^{d}} p_{\lambda}(s,\overline{\eta},t,x) 1_{\{\eta_{d}>0\}}\varphi(\eta) d\eta
$$

$$
= \int_{\mathbb{R}^{d}} p_{\lambda}(s,\eta,t,x) 1_{\{\eta_{d}>0\}}\varphi(\eta) d\eta - \int_{\mathbb{R}^{d}} p_{\lambda}(s,\eta,t,x) 1_{\{\eta_{d}<0\}}\varphi(\overline{\eta}) d\eta,
$$

which completes the proof.

Therefore we may rewrite the representations [\(6.18\)](#page-25-0) and [\(6.19\)](#page-25-1) and obtain the following lemma.

Lemma 6.10. For every $\varepsilon > 0$ we have

$$
\omega^{\varepsilon}(x,t) = \int_{\mathbb{R}^2} p_{\nu}(0,\eta,t,x)\hat{\omega^{\varepsilon}}(\eta,0) d\eta + \int_0^t \int_{\mathbb{R}^2} p_{\nu}(s,\eta,t,x)\hat{F}(\eta,s) d\eta ds + \int_0^t \int_{\mathbb{R}^2} p_{\nu}(s,\eta,t,x)\hat{\chi_{\varepsilon}}(\eta,s) d\xi ds
$$
\n(6.21)

and

$$
\theta^{\varepsilon}(x,t) = \int_{\mathbb{R}^2} p_{\kappa}(0,\eta,t,x) \hat{\theta^{\varepsilon}}(\eta,0) d\eta + \int_0^t \int_{\mathbb{R}^2} p_{\kappa}(s,\eta,t,x) \hat{\beta}_{\varepsilon}(\eta,s) d\eta ds
$$

for every $x \in \mathbb{R}^2_+$ and $t > 0$.

In order to avoid calculating the outer layer velocity, which appears in the error terms $\hat{\chi}_{\varepsilon}(\xi, s)$ and $\hat{\beta}_{\varepsilon}(\xi, s)$, we consider their limiting representations as $\varepsilon \downarrow 0$. We may apply the same technique used in [\[61\]](#page-44-15).

Proposition 6.11. The following integral representations hold:

$$
\omega(x,t) = 1_{\{x_2=0\}} \sigma(x_1,t) + \int_{\mathbb{R}^2} p_{\nu}(0,\eta,t,x) \hat{\omega}(\eta,0) d\eta
$$

+
$$
\int_0^t \int_{\mathbb{R}^2} p_{\nu}(s,\eta,t,x) \hat{F}(\eta,s) d\eta ds
$$

+
$$
2\nu \int_0^t \int_{-\infty}^{\infty} \sigma(\eta_1,s) \frac{\partial}{\partial \eta_2} \Big|_{\eta_2=0} p_{\nu}(s,(\eta_1,\eta_2),t,x) d\eta_1 ds
$$
(6.22)

and

$$
\theta(x,t) = 1_{\{x_2=0\}} \theta_0(x_1) + \int_{\mathbb{R}^2} p_{\kappa}(0,\eta,t,x) \hat{\theta}(\eta,0) d\eta
$$

+
$$
2\kappa \int_{-\infty}^{\infty} \theta_0(\eta_1) \frac{\partial}{\partial \eta_2} \Big|_{\eta_2=0} \left(\int_0^t p_{\kappa}(s,\eta,t,x) ds \right) d\eta_1
$$
(6.23)

for $x \in \mathbb{R}^2_+$.

Proof. Let us prove [\(6.22\)](#page-26-0) in detail, and the argument for θ^{ε} is similar. Since $\lim_{\varepsilon \to 0+} \omega^{\varepsilon}(x, t) =$ $\omega(x, t)$ for $x_2 > 0$, it follows that

$$
\lim_{\varepsilon \to 0+} \int_{\mathbb{R}^2} p_{\nu}(0, \eta, t, x) \widehat{\omega}^{\varepsilon}(\eta, 0) d\eta = \int_{\mathbb{R}^2} p_{\nu}(0, \eta, t, x) \widehat{\omega}(\eta, 0) d\eta.
$$

Hence we only need to deal with the limit of the error term

$$
J(x,\varepsilon) = \int_0^t \int_{\mathbb{R}^2} p_{\nu}(s,\xi,t,x) \hat{\chi}_{\varepsilon}(\xi,s) \mathrm{d}\xi \mathrm{d}s
$$

appearing on the right-hand side of [\(6.21\)](#page-25-2), where χ_{ε} is given in [\(6.7\)](#page-19-3). According to [\(6.8\)](#page-20-1), there is no contribution as $\varepsilon \downarrow 0$ towards the limit of $J(x,\varepsilon)$ from the first term in χ_{ε} . Therefore we only need to calculate the contributions from the error terms

$$
E_1^{\varepsilon}(x,t) = -\frac{1}{\varepsilon} \phi' \left(\frac{x_2}{\varepsilon}\right) \sigma(x_1,t) u^2(x,t)
$$

and

$$
E_2^{\varepsilon}(x,t) = \nu \frac{1}{\varepsilon^2} \phi''\left(\frac{x_2}{\varepsilon}\right) \sigma(x_1,t).
$$

To this end, we may choose a concrete cut-off function ϕ . Let us set

$$
\phi(r) = \begin{cases} 1 & \text{for } r \in [0, \frac{1}{3}), \\ \frac{1}{2} + 54 \left(r - \frac{1}{2} \right)^3 - \frac{9}{2} \left(r - \frac{1}{2} \right) & \text{for } r \in [\frac{1}{3}, \frac{2}{3}], \\ 0 & \text{for } r \ge \frac{2}{3} \end{cases}
$$
(6.24)

Then $-54 \le \phi'' \le 54$, $-\frac{9}{2} \le \phi' \le 0$ on $\left[\frac{1}{3}\right]$ $\frac{1}{3}, \frac{2}{3}$ $\frac{2}{3}$ and $\phi' = 0$ for $r \leq \frac{1}{3}$ $\frac{1}{3}$ or $r \geq \frac{2}{3}$ $\frac{2}{3}$. In fact,

$$
\phi'(r) = \begin{cases} 162\left(r - \frac{1}{2}\right)^2 - \frac{9}{2}, & \text{for } r \in [\frac{1}{3}, \frac{2}{3}], \\ 0, & \text{otherwise.} \end{cases}
$$

and

$$
\phi''(r) = \begin{cases} 324\left(r - \frac{1}{2}\right), & \text{for } r \in [\frac{1}{3}, \frac{2}{3}], \\ 0, & \text{otherwise.} \end{cases}
$$

Let

$$
J_i(x,\varepsilon) = \int_0^t \int_{\mathbb{R}^2} p_{\nu}(s,\xi,t,x) \hat{E}_i^{\varepsilon}(\xi,s) \mathrm{d}\xi \mathrm{d}s
$$

where $i = 1, 2$. Then

$$
J_i(x,\varepsilon) = \int_0^t \int_{-\infty}^\infty \left[\int_0^\infty p_\nu(s,(\xi_1,\xi_2),t,x) E_i^\varepsilon(\xi,s) d\xi_2 \right] d\xi_1 ds
$$

-
$$
\int_0^t \int_{-\infty}^\infty \left[\int_0^\infty p_\nu(s,(\xi_1,-\xi_2),t,x) E_i^\varepsilon(\xi,s) d\xi_2 \right] d\xi_1 ds.
$$

Let us consider the integral

$$
I_1(\varepsilon) := \int_0^\infty p_\nu(s, (\xi_1, \xi_2), t, x) E_1^{\varepsilon}(\xi, s) d\xi_2
$$

= $-\sigma(\xi_1, s) \int_0^{\varepsilon} u^2(\xi, s) p_\nu(s, (\xi_1, \xi_2), t, x) \frac{1}{\varepsilon} \phi' \left(\frac{\xi_2}{\varepsilon}\right) d\xi_2$
= $\sigma(\xi_1, s) \int_0^{\varepsilon} \phi \left(\frac{\xi_2}{\varepsilon}\right) \frac{\partial}{\partial \xi_2} \left(u^2(\xi, t) p_\nu(s, (\xi_1, \xi_2), t, x)\right) d\xi_2$
 $\to 0 \text{ as } \varepsilon \downarrow 0,$

where for the third equality, we have used the fact that $u(x, t)$ vanishes when $x_2 = 0$. Hence

$$
\lim_{\varepsilon \to 0+} \int_0^t \int_{\mathbb{R}^2} p_{\nu}(s,\xi,t,x) \hat{E}_1^{\varepsilon}(\xi,s) \mathrm{d}\xi \mathrm{d}s = 0.
$$

Now we consider $J_2(x, \varepsilon)$. To this end, we observe that

$$
J_2(x,\varepsilon) = \int_0^t \int_{-\infty}^{\infty} \left[\int_0^{\infty} (p_\nu(s,(\xi_1,\xi_2),t,x) - p_\nu(s,(\xi_1,-\xi_2),t,x)) E_2^{\varepsilon}(\xi,s) d\xi_2 \right] d\xi_1 ds
$$

and integrate by parts twice to deduce that

$$
I_{2}(\varepsilon) := \int_{0}^{\infty} (p_{\nu}(s,(\xi_{1},\xi_{2}),t,x) - p_{\nu}(s,(\xi_{1},-\xi_{2}),t,x)) E_{2}^{\varepsilon}(\xi,s) d\xi_{2}
$$

\n
$$
= \nu \sigma(\xi_{1},t) \int_{0}^{\varepsilon} (p_{\nu}(s,(\xi_{1},\xi_{2}),t,x) - p_{\nu}(s,(\xi_{1},-\xi_{2}),t,x)) \frac{1}{\varepsilon^{2}} \phi''\left(\frac{\xi_{2}}{\varepsilon}\right) d\xi_{2}
$$

\n
$$
= -\nu \sigma(\xi_{1},t) \int_{0}^{\varepsilon} \frac{\partial}{\partial \xi_{2}} (p_{\nu}(s,(\xi_{1},\xi_{2}),t,x) - p_{\nu}(s,(\xi_{1},-\xi_{2}),t,x)) \frac{1}{\varepsilon} \phi'\left(\frac{\xi_{2}}{\varepsilon}\right) d\xi_{2}
$$

\n
$$
= 2\nu \sigma(\xi_{1},t) \frac{\partial}{\partial \xi_{2}} \Big|_{\xi_{2}=0} p_{\nu}(s,(\xi_{1},\xi_{2}),t,x)
$$

\n
$$
+ \nu \sigma(\xi_{1},t) \int_{0}^{\varepsilon} \phi\left(\frac{\xi_{2}}{\varepsilon}\right) \frac{\partial^{2}}{\partial \xi_{2}^{2}} (p_{\nu}(s,(\xi_{1},\xi_{2}),t,x) - p_{\nu}(s,(\xi_{1},-\xi_{2}),t,x)) d\xi_{2}.
$$

As a consequence, we have

$$
I_2(\varepsilon) \to 2\nu\sigma(\xi_1, t) \frac{\partial}{\partial \xi_2}\Big|_{\xi_2=0} p_\nu(s, (\xi_1, \xi_2), t, x) \quad \text{as } \varepsilon \downarrow 0
$$

and therefore

$$
\lim_{\varepsilon \to 0+} J_2(x, \varepsilon) = 2\nu \int_0^t \int_{-\infty}^\infty \sigma(\xi_1, t) \left. \frac{\partial}{\partial \xi_2} \right|_{\xi_2 = 0} p_\nu(s, (\xi_1, \xi_2), t, x) \, d\xi_1 ds
$$

which completes the proof.

By using the Biot-Savart law, we may then deduce the following functional integral representation theorem.

 \Box

Theorem 6.12. Let X and Y be defined as in (4.5) and (4.6) respectively. The following functional integral representations hold:

$$
u(x,t) = \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta}) A_2(X_t^{\eta}, x) \wedge \hat{\omega}(\eta, 0) \right] d\eta
$$

+
$$
\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta,s}) A_2(X_t^{\eta,s}, x) \wedge \hat{F}(\eta, s) \right] d\eta ds
$$

+
$$
2\nu \int_0^t \int_{-\infty}^{\infty} \frac{\partial}{\partial \eta_2} \left| \sum_{\eta_2=0} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta,s}) A_2(X_t^{\eta,s}, x) \wedge \sigma(\eta, s) \right] d\eta_1 ds
$$

and

$$
u(x,t) = I_{2,1}(x,t) + \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta}) A_2(X_t^{\eta}, x) \wedge \hat{\omega}(\eta, 0) \right] d\eta
$$

-
$$
\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta,s}) A_2(X_t^{\eta,s}, x) \wedge \frac{\partial \theta_0}{\partial \eta_1}(\eta_1, s) \right] d\eta ds
$$

+
$$
2\nu \int_0^t \int_{-\infty}^{\infty} \frac{\partial}{\partial \eta_2} \left| \sum_{\eta_2=0} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta,s}) A_2(X_t^{\eta,s}, x) \wedge \sigma(\eta, s) \right] d\eta_1 ds
$$

for every $x \in \mathbb{R}^2_+$ and $t > 0$, where

$$
I_{2,1}(x,t) = -\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} \left(Y_s^{\xi} \right) H \left(Y_s^{\xi}, s; t, x \right) \wedge \hat{\theta}(\xi, 0) \right] d\xi ds
$$

+
$$
\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} \left(\overline{Y_s^{\xi}} \right) H \left(Y_s^{\overline{\xi}}, s; t, x \right) \wedge \hat{\theta}(\xi, 0) \right] d\xi ds
$$

-
$$
2\kappa \int_0^t \int_{-\infty}^{\infty} \frac{\partial}{\partial \xi_2} \left|_{\xi_2=0} \left(\int_0^s \mathbb{E} \left[1_{\mathbb{R}_+^2} \left(Y_s^{\xi, \tau} \right) H \left(Y_s^{\xi, \tau}, s; t, x \right) \wedge \theta_0(\xi_1) \right] d\tau \right) d\xi_1 ds
$$

+
$$
2\kappa \int_0^t \int_{-\infty}^{\infty} \frac{\partial}{\partial \xi_2} \left|_{\xi_2=0} \left(\int_0^s \mathbb{E} \left[1_{\mathbb{R}_+^2} \left(\overline{Y_s^{\xi, \tau}} \right) H \left(Y_s^{\overline{\xi}, \tau}, s; t, x \right) \wedge \theta_0(\xi_1) \right] d\tau \right) d\xi_1 ds
$$

and

$$
H(\eta, s; t, x) = \frac{\partial}{\partial \eta_1} \mathbb{E} \left[1_{\mathbb{R}^2_+} (X_t^{\eta, s}) A_2(X_t^{\eta, s}, x) \right]
$$

for $x \in \mathbb{R}_+^2$ and $t > s \geq 0$.

Proof. By [\(3.5\)](#page-12-3) and [\(6.22\)](#page-26-0), we obtain that

$$
u(x,t) = \int_{\mathbb{R}_+^2} A_2(y,x) \wedge \omega(y,t) dy
$$

\n
$$
= \int_{\mathbb{R}_+^2} A_2(y,x) \wedge 1_{\{y_2=0\}} \sigma(y_1,t) dy
$$

\n
$$
+ \int_{\mathbb{R}^2} \int_{\mathbb{R}_+^2} A_2(y,x) \wedge \hat{\omega}(\eta,0) p_{\nu}(0,\eta,t,y) dy d\eta
$$

\n
$$
+ \int_0^t \int_{\mathbb{R}^2} \int_{\mathbb{R}_+^2} A_2(y,x) \wedge \hat{F}(\eta,s) p_{\nu}(s,\eta,t,y) dy d\eta ds
$$

\n
$$
+ 2\nu \int_0^t \int_{-\infty}^\infty \frac{\partial}{\partial \eta_2} \Big|_{\eta_2=0} \int_{\mathbb{R}_+^2} A_2(y,x) \wedge \sigma(\eta_1,s) p_{\nu}(s,(\eta_1,\eta_2),t,y) dy d\eta_1 ds.
$$

Hence

$$
u(x,t) = \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta}) A_2(X_t^{\eta}, x) \wedge \hat{\omega}(\eta, 0) \right] d\eta
$$

+
$$
\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta,s}) A_2(X_t^{\eta,s}, x) \wedge \hat{F}(\eta, s) \right] d\eta ds
$$

+ $2\nu \int_0^t \int_{-\infty}^\infty \frac{\partial}{\partial \eta_2} \Big|_{\eta_2=0} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta,s}) A_2(X_t^{\eta,s}, x) \wedge \sigma(\eta, s) \right] d\eta_1 ds$,

where

$$
F(x,t) = \frac{\partial}{\partial x_1} \theta(x,t) - \frac{\partial}{\partial x_1} \theta_0(x_1),
$$

which yields in particular the first representation for $u(x, t)$. To prove the second representation, we need to handle the second term

$$
I_2 = \int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta,s}) A_2(X_t^{\eta,s}, x) \wedge \hat{F}(\eta, s) \right] d\eta ds
$$

=
$$
\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta,s}) A_2(X_t^{\eta,s}, x) \wedge \frac{\partial \theta}{\partial \eta_1}(\eta, s) \right] d\eta ds
$$

-
$$
\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta,s}) A_2(X_t^{\eta,s}, x) \wedge \frac{\partial \theta_0}{\partial \eta_1}(\eta_1, s) \right] d\eta ds,
$$

where the first term on the right-hand side is denoted by

$$
I_{2,1} = \int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta,s}) A_2(X_t^{\eta,s}, x) \right] \wedge \overline{\frac{\partial \theta}{\partial \eta_1}}(\eta, s) d\eta ds.
$$

Next we notice that by definition

$$
\widehat{\frac{\partial \theta}{\partial \eta_1}}(\eta, s) = \frac{\partial}{\partial \eta_1} \widehat{\theta}(\eta, s)
$$

so that we may rewrite

$$
I_{2,1} = -\int_0^t \int_{\mathbb{R}^2} H(\eta, s; t, x) \wedge \hat{\theta}(\eta, s) \mathrm{d}\eta \mathrm{d}s,\tag{6.25}
$$

where for simplicity, we have introduced the following kernel

$$
H(\eta, s; t, x) = \frac{\partial}{\partial \eta_1} \mathbb{E}\left[1_{\mathbb{R}_+^2}(X_t^{\eta, s}) \Lambda_2(X_t^{\eta, s}, x)\right].
$$

Using the representation [\(6.23\)](#page-26-1), we have

$$
\theta(\eta, s) = 1_{\{\eta_2 = 0\}} \theta_0(\eta_1) + \int_{\mathbb{R}^2} p_{\kappa}(0, \xi, s, \eta) \hat{\theta}(\xi, 0) d\xi + 2\kappa \int_{-\infty}^{\infty} \theta_0(\xi_1) \left. \frac{\partial}{\partial \xi_2} \right|_{\xi_2 = 0} \left(\int_0^s p_{\kappa}(\tau, \xi, s, \eta) d\tau \right) d\xi_1,
$$

so that

$$
\hat{\theta}(\eta,s) = \int_{\mathbb{R}^2} 1_{\mathbb{R}_+^2}(\eta) p_{\kappa}(0,\xi,s,\eta) \hat{\theta}(\xi,0) d\xi - \int_{\mathbb{R}^2} 1_{\mathbb{R}_+^2}(\overline{\eta}) p_{\kappa}(0,\overline{\xi},s,\eta) \hat{\theta}(\xi,0) d\xi \n+ 2\kappa \int_{-\infty}^{\infty} \theta_0(\xi_1) \frac{\partial}{\partial \xi_2} \bigg|_{\xi_2=0} \left(\int_0^s 1_{\mathbb{R}_+^2}(\eta) p_{\kappa}(\tau,\xi,s,\eta) d\tau \right) d\xi_1 \n- 2\kappa \int_{-\infty}^{\infty} \theta_0(\xi_1) \frac{\partial}{\partial \xi_2} \bigg|_{\xi_2=0} \left(\int_0^s 1_{\mathbb{R}_+^2}(\overline{\eta}) p_{\kappa}(\tau,\overline{\xi},s,\eta) d\tau \right) d\xi_1.
$$

Substituting this expression into [\(6.25\)](#page-29-0) we then deduce that

$$
I_{2,1} = -\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} \left(Y_s^{\xi} \right) H \left(Y_s^{\xi}, s; t, x \right) \wedge \hat{\theta}(\xi, 0) \right] d\xi ds
$$

+
$$
\int_0^t \int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} \left(\overline{Y_s^{\xi}} \right) H \left(Y_s^{\overline{\xi}}, s; t, x \right) \wedge \hat{\theta}(\xi, 0) \right] d\xi ds
$$

-
$$
- 2\kappa \int_0^t \int_{-\infty}^{\infty} \frac{\partial}{\partial \xi_2} \Big|_{\xi_2=0} \left(\int_0^s \mathbb{E} \left[1_{\mathbb{R}_+^2} \left(Y_s^{\xi, \tau} \right) H \left(Y_s^{\xi, \tau}, s; t, x \right) \wedge \theta_0(\xi_1) \right] d\tau \right) d\xi_1 ds
$$

+
$$
2\kappa \int_0^t \int_{-\infty}^{\infty} \frac{\partial}{\partial \xi_2} \Big|_{\xi_2=0} \left(\int_0^s \mathbb{E} \left[1_{\mathbb{R}_+^2} \left(\overline{Y_s^{\xi, \tau}} \right) H \left(Y_s^{\overline{\xi}, \tau}, s; t, x \right) \wedge \theta_0(\xi_1) \right] d\tau \right) d\xi_1 ds.
$$

Putting together, we obtain the second representation.

$$
\qquad \qquad \Box
$$

7 Numerical schemes and experiment results

In this section, we formulate several numerical schemes based on the representations for flows in two dimensional space established in the previous sections, and demonstrate the numerical results.

Let us review our notations for the sake of comprehensibility. There are four fluid dynamical variables we are going to calculate by means of numerical simulations: the velocity $u(x, t)$, the vorticity $\omega(x, t)$, the temperature $\theta(x, t)$ and the temperature gradient $\Theta(x, t)$. We are given the initial velocity $u(x, 0)$, and hence the initial vorticity $\omega(x, 0)$ as well as the initial temperature $\theta(x, 0)$, which has a small gradient, i.e., the magnitude of the gradient $\Theta(x, 0) = \nabla \theta(x, 0)$ is small so can be ignored in numerical schemes. The kinematic viscosity $\nu > 0$ and the heat diffusivity constant $\kappa > 0$ depend on the nature of the fluid. The fluid density ρ is almost a constant, so we may choose it to be the unit.

7.1 Oberbeck-Boussinesq flows in \mathbb{R}^2

The numerical experiments in this part are carried out for two-dimensional fluid flows on the whole space, and therefore, the Biot-Savart singular kernel is a vector kernel given by $K_2(y, x) =$ $(2\pi)^{-1}(y-x)/|y-x|^2$, and the interaction force $f(\theta)(x,t) = (0, \alpha(\theta(x,t)-\theta(x,0)))$ where $\alpha > 0$ is a constant. Thus

$$
F(x,t) = \alpha \frac{\partial}{\partial x_1} \theta(x,t) - \alpha \frac{\partial}{\partial x_1} \theta(x,0),
$$

where $\theta(x, 0)$ is the given initial temperature distribution whose gradient is small.

Choose a lattice mesh $h > 0$. For $i_1, i_2 \in \mathbb{Z}$, denote $x^{i_1, i_2} = (i_1, i_2)h$, the lattice points, $\omega_{i_1,i_2} = \omega(x^{i_1,i_2},0)$, and $\Theta_{i_1,i_2} = \Theta(x^{i_1,i_2},0)$. Let $\delta > 0$ be the step length of the time variable and $t_i = i\delta$, $i = 0, 1, 2, \cdots$.

Note that in the numerical schemes based on the functional integral representations, the interaction term F involves the temperature gradient, therefore, we have to calculate the derivatives of $\theta(x, t)$. Since the derivative of the Biot-Savart kernel is no longer locally integrable, we need to replace K_2 with its regularisation measured via a positive parameter $\epsilon > 0$, and introduce

$$
K_{2,\epsilon}(y,x) = \left(1 - e^{-\frac{|y-x|^2}{\epsilon}}\right) K_2(y,x).
$$

Let us describe the numerical scheme by adopting the functional integral representations in Subsection [5.3.1.](#page-15-1)

7.1.1 One copy scheme

In this numerical scheme, we drop the expectation using independent copies of Brownian motion. We discretise the stochastic differential equations using the Euler scheme: for $i_1, i_2 \in \mathbb{Z}$, $k =$ $1, 2, \cdots,$

$$
X_0^{i_1, i_2} = x^{i_1, i_2}, \quad X_{t_k}^{i_1, i_2} = X_{t_{k-1}}^{i_1, i_2} + \delta u \left(X_{t_{k-1}}^{i_1, i_2}, t_{k-1} \right) + \sqrt{2\nu} (B_{t_k}^1 - B_{t_{k-1}}^1), \tag{7.1}
$$

$$
Y_0^{i_1,i_2} = x^{i_1,i_2}, \quad Y_{t_k}^{i_1,i_2} = Y_{t_{k-1}}^{i_1,i_2} + \delta u \left(Y_{t_{k-1}}^{i_1,i_2}, t_{k-1} \right) + \sqrt{2\kappa} (B_{t_k}^2 - B_{t_{k-1}}^2), \tag{7.2}
$$

where we use $X_t^{i_1,i_2}$ to denote $X_t^{x^{i_1,i_2}}$ to simplify our notation, and B^1 , B^2 are two independent two-dimensional Brownian motions.

The integral representations in Theorems [5.3](#page-15-3) and [5.4](#page-15-4) are approximated by the following discretisation:

$$
u(x, t_k) = \sum_{i_1, i_2} h^2 K_{2, \epsilon} \left(X_{t_k}^{i_1, i_2}, x \right) \wedge \omega_{i_1, i_2}
$$

+
$$
\sum_{i_1, i_2} \sum_{j=1}^k \delta h^2 K_{2, \epsilon} \left(X_{t_k}^{i_1, i_2}, x \right) \wedge F \left(X_{t_{j-1}}^{i_1, i_2}, t_{j-1} \right), \tag{7.3}
$$

and

$$
\theta(x, t_k) = -\sum_{i_1, i_2} h^2 K_{2,\epsilon} \left(Y_{t_k}^{i_1, i_2}, x \right) \cdot R(x^{i_1, i_2}, t_k; 0) \Theta_{i_1, i_2},\tag{7.4}
$$

where $R(x^{i_1,i_2}, t_k; t_k) = I$, and

$$
R(x^{i_1,i_2}, t_k; t_i) = I - \sum_{l=i+1}^k \delta A\left(Y^{i_1,i_2}_{t_l}, t_l\right) R(x^{i_1,i_2}, t_k; t_l)
$$

for $i = 0, \dots, k-1$. As for $F(x, t)$ and $A(x, t)$, we update them in each iteration by formally differentiating the equations [\(7.3\)](#page-31-1) and [\(7.4\)](#page-31-2) respectively, so that

$$
A(x, t_k) = \sum_{i_1, i_2} h^2 \nabla_x K_{2, \epsilon} \left(X_{t_k}^{i_1, i_2}, x \right) \wedge \omega_{i_1, i_2}
$$

+
$$
\sum_{i_1, i_2} \sum_{j=1}^k \delta h^2 \nabla_x K_{2, \epsilon} \left(X_{t_k}^{i_1, i_2}, x \right) \wedge F \left(X_{t_{j-1}}^{i_1, i_2}, t_{j-1} \right),
$$

$$
\Theta(x, t_k) = - \sum_{i_1, i_2} h^2 \nabla_x K_{2, \epsilon} \left(Y_{t_k}^{i_1, i_2}, x \right) \cdot R(x^{i_1, i_2}, t_k; 0) \Theta_{i_1, i_2},
$$
 (7.5)

and

$$
F(X_{t_k}^{i_1,i_2},t_k) = \alpha \Theta_1(X_{t_k}^{i_1,i_2},t_k) - \alpha \frac{\partial \theta}{\partial x_1}(X_{t_k}^{i_1,i_2},0),
$$

where the gradient of the Biot-Savart kernel is replaced by

$$
\nabla_x K_{2,\epsilon}(y,x) = \nabla_x \left[\left(1 - e^{-\frac{|y-x|^2}{\epsilon}} \right) K_2(y,x) \right]. \tag{7.6}
$$

Remark 7.1. This scheme is still quite computationally expensive as it requires storing the values of A at all times. Since in any simulation for non-linear dynamics, the time duration cannot be long, and one can use their approximations of the time integral. That is, equations (7.3) and [\(7.5\)](#page-31-3) can be substituted with the following iterations:

$$
u(x, t_k) = \sum_{i_1, i_2} h^2 K_{2, \epsilon} \left(X_{t_k}^{i_1, i_2}, x \right) \wedge \omega_{i_1, i_2}
$$

+
$$
\sum_{i_1,i_2} k \delta h^2 K_{2,\epsilon} \left(X_{t_k}^{i_1,i_2}, x \right) \wedge F \left(X_{t_{k-1}}^{i_1,i_2}, t_{k-1} \right)
$$

and

$$
A(x, t_k) = \sum_{i_1, i_2} h^2 \nabla_x K_{2,\epsilon} \left(X_{t_k}^{i_1, i_2}, x \right) \wedge \omega_{i_1, i_2}
$$

+
$$
\sum_{i_1, i_2} k \delta h^2 \nabla_x K_{2,\epsilon} \left(X_{t_k}^{i_1, i_2}, x \right) \wedge F \left(X_{t_{k-1}}^{i_1, i_2}, t_{k-1} \right)
$$

respectively.

7.1.2 Multi-copy scheme

We introduce the second numerical scheme, where the expectation is substituted with the empirical mean based on the strong law of large numbers. Take 2N independent copies of Brownian motions $(B_t^{1,m})$ $(t_t^{1,m})$ and $(B_t^{2,m})$ $t^{2,m}$, $m = 1, 2, \cdots, N$.

We repeat the diffusion processes of the twin particle N times by running $2N$ independent copies of Brownian motion and replacing the expectations with their averages. That is, for $1 \leq m \leq N$, define

$$
X_0^{m,i_1,i_2} = x^{i_1,i_2}, \quad X_{t_k}^{m,i_1,i_2} = X_{t_{k-1}}^{m,i_1,i_2} + \delta u \left(X_{t_{k-1}}^{m,i_1,i_2}, t_{k-1} \right) + \sqrt{2\nu} (B_{t_k}^{1,m} - B_{t_{k-1}}^{1,m})
$$

and

$$
Y_0^{m,i_1,i_2} = x^{i_1,i_2}, \quad Y_{t_k}^{m,i_1,i_2} = Y_{t_{k-1}}^{m,i_1,i_2} + \delta u \left(Y_{t_{k-1}}^{m,i_1,i_2}, t_{k-1} \right) + \sqrt{2\kappa} (B_{t_k}^{2,m} - B_{t_{k-1}}^{2,m}).
$$

Then the velocity and temperature are approximated by

$$
u(x, t_k) = \frac{1}{N} \sum_{m=1}^{N} \sum_{i_1, i_2} h^2 K_{2,\epsilon} \left(X_{t_k}^{m, i_1, i_2}, x \right) \wedge \omega_{i_1, i_2}
$$

+
$$
\frac{1}{N} \sum_{m=1}^{N} \sum_{i_1, i_2} \sum_{j=1}^{k} \delta h^2 K_{2,\epsilon} \left(X_{t_k}^{m, i_1, i_2}, x \right) \wedge F \left(X_{t_{j-1}}^{m, i_1, i_2}, t_{j-1} \right),
$$

$$
\theta(x, t_k) = -\frac{1}{N} \sum_{m=1}^{N} \sum_{i_1, i_2} h^2 K_{2,\epsilon} \left(Y_{t_k}^{m, i_1, i_2}, x \right) \cdot R(x^{i_1, i_2}, t_k; 0) \Theta_{i_1, i_2},
$$

where similar to the first scheme,

$$
A(x, t_k) = \frac{1}{N} \sum_{m=1}^{N} \sum_{i_1, i_2} h^2 \nabla_x K_{2,\epsilon} \left(X_{t_k}^{m, i_1, i_2}, x \right) \wedge \omega_{i_1, i_2}
$$

+
$$
\frac{1}{N} \sum_{m=1}^{N} \sum_{i_1, i_2} \sum_{j=1}^{k} \delta h^2 \nabla_x K_{2,\epsilon} \left(X_{t_k}^{m, i_1, i_2}, x \right) \wedge F \left(X_{t_{j-1}}^{m, i_1, i_2}, t_{j-1} \right),
$$

$$
\Theta(x, t_k) = -\frac{1}{N} \sum_{m=1}^{N} \sum_{i_1, i_2} h^2 \nabla_x K_{2,\epsilon} \left(Y_{t_k}^{m, i_1, i_2}, x \right) \cdot R(x^{i_1, i_2}, t_k; 0) \Theta_{i_1, i_2},
$$

where R depends on m such that for each m, $R(x^{i_1,i_2}, t_k; t_k) = I$, and

$$
R(x^{i_1,i_2}, t_k; t_i) = I - \sum_{l=i+1}^k \delta A\left(Y_{t_l}^{m,i_1,i_2}, t_l\right) R(x^{i_1,i_2}, t_k; t_l).
$$

The rest of the scheme remains the same as in the one-copy scheme.

7.1.3 Numerical experiments

Following the one-copy numerical scheme described above, we carried out several simple numerical experiments. Here, we present the results of experiments with different Prandtl numbers. In the first experiment, we set $\nu = 1$ and $\kappa = 0.15$, so that the Prandtl number Pr = 0.15. In the second experiment, we swap these two values and consider the case when $Pr = 6.67$.

We choose the typical length scale $L = 2\pi$, and assume that $\alpha = 0.0005$. The parameter we use to smooth out the Biot-Savart kernel is chosen to be $\epsilon = 0.1$.

In the experiment presented, we set the initial velocity to be of the form

$$
u(x,0) = (-10\sin(x_2),0),
$$

and the initial temperature is given by

$$
\theta_0(x) = \theta(x,0) = 0.01(8\pi^2 - x_1^2 - x_2^2).
$$

Thus, $\omega(x, 0) = 10 \cos(x_2)$, and

$$
\omega_{i_1,i_2} = 10 \cos(i_2 h).
$$

The time step is $\delta = 0.01$ with mesh size $h = 2\pi/40$. The numerical experiment results at times $t = 0.6, t = 1.2, t = 1.8$ with Pr = 0.15 and Pr = 6.67 are shown in the Figure [7.1](#page-33-1) and Figure [7.2,](#page-34-1) respectively.

Figure 7.1: Temperature and velocity fields of Oberbeck-Boussinesq flows on \mathbb{R}^2 with Prandtl number $Pr = 0.15$.

The figures show how the buoyancy from small temperature variations accelerates the flow. Besides, the growth of temperature is different from the result of a linear parabolic equation the nonlinearity revealed in the temperature evolution reflects the velocity in the drift term is, in turn, driven by the thermal convection.

Figure 7.2: Temperature and velocity fields of Oberbeck-Boussinesq flows on \mathbb{R}^2 with Prandtl number $Pr = 6.67$.

7.2 Oberbeck-Boussinesq flows in wall-bounded domains

This series of numerical experiments are based on the wall-bounded fluid flows in Subsection [6.5.1.](#page-23-0) Therefore the fluid is heated from the bottom with an external source of heat at temperature $\theta_0(x) = \theta_0(x_1)$, which depends only on the first coordinate.

The Biot-Savart kernel for this case is given by

$$
\Lambda_2(y,x) = \frac{1}{2\pi} \left(\frac{y-x}{|y-x|^2} - \frac{y-\overline{x}}{|y-\overline{x}|^2} \right) \quad \text{for } y \neq x \text{ or } \overline{x}
$$

and for the same reason as in the previous case, we take its regularisation via a real parameter $\epsilon > 0$, and replace the singular integral kernel by

$$
\Lambda_{2,\epsilon}(y,x) = \left(1 - e^{-\frac{|y-x|^2}{\epsilon}}\right) \Lambda_2(y,x).
$$

The numerical scheme is based on the functional integral representations [\(6.16,](#page-24-2) [6.15\)](#page-23-1). Choose $\varepsilon > 0$ small. The approximated representations we will use here are then given by

$$
u(x,t) \sim \int_{\mathbb{R}^2_+} \mathbb{E} \left[1_{\{t < \zeta(X^{\eta})\}} 1_{\mathbb{R}^2_+} (X_t^{\eta}) \Lambda_2(X_t^{\eta}, x) \wedge \omega(\eta, 0) \right] d\eta
$$

+
$$
\int_0^t \int_{\mathbb{R}^2_+} \mathbb{E} \left[1_{\{s > \gamma_t(X^{\eta})\}} 1_{\mathbb{R}^2_+} (X_t^{\eta}) \Lambda_2(X_t^{\eta}, x) \wedge F(X_s^{\eta}, s) \right] d\eta ds
$$

+
$$
\int_0^t \int_{\mathbb{R}^2_+} \mathbb{E} \left[1_{\{s > \gamma_t(X^{\eta})\}} 1_{\mathbb{R}^2_+} (X_t^{\eta}) \Lambda_2(X_t^{\eta}, x) \wedge \tilde{\chi}_{\varepsilon}(X_s^{\eta}, s) \right] d\eta ds \qquad (7.7)
$$

and

$$
\theta(x,t) \sim \frac{x_2}{\pi} \int_{-\infty}^{\infty} \frac{\theta_0(\xi_1)}{|\xi_1 - x_1|^2 + x_2^2} d\xi_1
$$

$$
- \int_{\mathbb{R}^2_+} \mathbb{E} \left[1_{\{t < \zeta(Y^{\eta})\}} 1_{\mathbb{R}^2_+} (Y^{\eta}_t) A_2(Y^{\eta}_t, x) \cdot R(\eta, t; 0) \Theta(\eta, 0) \right] d\eta. \tag{7.8}
$$

for $x \in \mathbb{R}^2_+$ and $t > 0$. According to [\(6.3\)](#page-18-3), the first term on the right-hand side of [\(7.7\)](#page-34-2) can be written as

$$
\int_{\mathbb{R}^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta}) A_2(X_t^{\eta}, x) \wedge \hat{\omega}(\eta, 0) \right] d\eta,
$$

where $\hat{\omega}$ can be computed using [\(6.20\)](#page-25-3) so that

$$
u(x,t) \sim \int_{\mathbb{R}_+^2} \mathbb{E} \left[1_{\mathbb{R}_+^2} (X_t^{\eta}) A_2(X_t^{\eta}, x) \wedge \hat{\omega}(\eta, 0) \right] d\eta
$$

+
$$
\int_0^t \int_{\mathbb{R}_+^2} \mathbb{E} \left[1_{\{s > \gamma_t(X^{\eta})\}} 1_{\mathbb{R}_+^2} (X_t^{\eta}) A_2(X_t^{\eta}, x) \wedge F(X_s^{\eta}, s) \right] d\eta ds
$$

+
$$
\int_0^t \int_{\mathbb{R}_+^2} \mathbb{E} \left[1_{\{s > \gamma_t(X^{\eta})\}} 1_{\mathbb{R}_+^2} (X_t^{\eta}) A_2(X_t^{\eta}, x) \wedge \tilde{\chi}_{\varepsilon}(X_s^{\eta}, s) \right] d\eta ds.
$$
 (7.9)

Now we are in a position to discretise [\(7.9\)](#page-35-1) and [\(7.8\)](#page-35-2) and obtain the numerical schemes. Namely, set $\hat{\omega}_{i_1,i_2} = \hat{\omega}(x^{i_1,i_2},0)$ for $i_2 \geq 0$. Let $\epsilon > 0$ be another small constant to take care of the differentiation of the singular kernel. Choose $\varepsilon \ll h$, and constants h, ε and ϵ are fixed through numerical experiments, but they can be adjusted.

We assume that the bounded boundary layer is $B = \{x \in \mathbb{R}^2 : 0 < x_2 < \gamma\}$, and use a different mesh size h_w for the boundary layer. Let h_{i_1,i_2} be the vertical mesh size such that

$$
h_{i_1,i_2} = \begin{cases} h_w, & \text{in } B, \\ h, & \text{otherwise.} \end{cases}
$$

7.2.1 One copy scheme

In this scheme, the expectations in the integral representations are treated by using the onecopy scheme, i.e., simply drop the expectation sign in the random vortex system. Due to the no-slip condition, the velocity and thus the heat conduction decreases rapidly within a thin layer adjoining the wall, which is commonly known as the boundary layer (see e.g. Chapter 4, [\[44\]](#page-43-0)).

Similar to the whole plane case, we discretise the SDE system using the Euler scheme:

$$
\begin{aligned}\nX_0^{i_1,i_2} &= x^{i_1,i_2}, & X_{t_k}^{i_1,i_2} &= X_{t_{k-1}}^{i_1,i_2} + \delta u(X_{t_{k-1}}^{i_1,i_2},t_{k-1}) + \sqrt{2\nu}(B_{t_k}^1 - B_{t_{k-1}}^1), \\
Y_0^{i_1,i_2} &= x^{i_1,i_2}, & Y_{t_k}^{i_1,i_2} &= Y_{t_{k-1}}^{i_1,i_2} + \delta u(Y_{t_{k-1}}^{i_1,i_2},t_{k-1}) + \sqrt{2\kappa}(B_{t_k}^2 - B_{t_{k-1}}^2),\n\end{aligned}
$$

Furthermore, we set the force F to be

$$
F(x,t) = \alpha \left(\frac{\partial \theta}{\partial x_1}(x,t) - \frac{\partial \theta_0}{\partial x_1}(x_1) \right),\,
$$

where $\theta_0(x_1)$ represents the external source of heat imposed on the boundary, and it is assumed to be almost constant $T_w > 0$ on $[-L, L]$, vanish everywhere else, and has a small derivative.

The equation [\(7.9\)](#page-35-1) is approximated by

$$
u(x, t_k) = \sum_{i_1} \sum_{i_2 \ge 0} h h_{i_1, i_2} \prod_{j=1}^k 1_{\mathbb{R}_+^2} (X_{t_j}^{i_1, i_2}) \Lambda_{2, \epsilon}(X_{t_k}^{i_1, i_2}, x) \wedge \hat{\omega}_{i_1, i_2}
$$

+
$$
\sum_{i_1} \sum_{i_2 \ge 0} \delta h h_{i_1, i_2} \sum_{j=1}^k \prod_{l=j}^k 1_{\mathbb{R}_+^2} (X_{t_l}^{i_1, i_2}) \Lambda_{2, \epsilon}(X_{t_k}^{i_1, i_2}, x) \wedge F(X_{t_{j-1}}^{i_1, i_2}, t_{j-1})
$$

$$
+\sum_{i_1}\delta hh_w \sum_{j=1}^k \prod_{l=j}^k 1_{\mathbb{R}_+^2} (X_{t_l}^{i_1,0}) \Lambda_{2,\epsilon}(X_{t_k}^{i_1,0}, x) \wedge \tilde{\chi}_{\varepsilon}(X_{t_{j-1}}^{i_1,0}, t_{j-1})
$$
\n(7.10)

for $x = (x_1, x_2)$ with $x_2 > 0$, and

$$
u(x,t) = (u^1((x_1, -x_2), t), -u^2((x_1, -x_2), t))
$$

for $x = (x_1, x_2)$ with $x_2 < 0$, where $\tilde{\chi}_{\varepsilon}$ is given by [\(6.17\)](#page-24-3),

$$
\tilde{\chi}_{\varepsilon}(x,t) = -\nu \frac{1}{\varepsilon^2} \phi''\left(\frac{x_2}{\varepsilon}\right) \left. \frac{\partial}{\partial x_2} \right|_{x_2=0+} u^1(x,t)
$$
\n
$$
= -\nu \frac{1}{\varepsilon^2} \phi''\left(\frac{x_2}{\varepsilon}\right) A_2^1((x_1,0),t)
$$

with ϕ given by [\(6.24\)](#page-26-2), whose second derivative is

$$
\phi''(r) = \begin{cases} 324(r - \frac{1}{2}), & \frac{1}{3} \le r \le \frac{2}{3}, \\ 0, & \text{otherwise.} \end{cases}
$$

The boundary integral is replaced with the third term on the right-hand side of [\(7.10\)](#page-36-0) since we assume $\varepsilon < h_w \ll h$ and $\tilde{\chi}_{\varepsilon}(x, t)$ only supports on $\frac{\varepsilon}{3} \leq x_2 \leq \frac{2\varepsilon}{3}$ $\frac{2\varepsilon}{3}$, and thus it is sufficient to consider the diffusion starting within the boundary layer.

To discretise [\(7.8\)](#page-35-2), we use the approximation

$$
\frac{x_2}{\pi} \int_{-\infty}^{\infty} \frac{\theta_0(\xi_1)}{|\xi_1 - x_1|^2 + x_2^2} d\xi_1 \approx T_w \frac{1}{\pi} \left(\tan^{-1} \left(\frac{L - x_1}{x_2} \right) + \tan^{-1} \left(\frac{L + x_1}{x_2} \right) \right)
$$

and thus

$$
\theta(x,t_k) = \frac{T_w}{\pi} \left(\tan^{-1} \left(\frac{L - x_1}{x_2} \right) + \tan^{-1} \left(\frac{L + x_1}{x_2} \right) \right)
$$

$$
- \sum_{i_1} \sum_{i_2 \ge 0} h h_{i_1, i_2} \prod_{j=1}^k 1_{\mathbb{R}_+^2} (Y_{t_j}^{i_1, i_2}) A_{2,\epsilon}(Y_{t_k}^{i_1, i_2}, x) \cdot R(x^{i_1, i_2}, t_k; 0) \Theta_{i_1, i_2}
$$
(7.11)

and the gauge functional equation [\(4.4\)](#page-13-3) by $R(x^{i_1,i_2}, t_k; t_k) = I$, and

$$
R(x^{i_1,i_2}, t_k; t_i) = I - \sum_{l=i+1}^k \delta A\left(Y^{i_1,i_2}_{t_l}, t_l\right) R(x^{i_1,i_2}, t_k; t_l) 1_{\mathbb{R}^2_+}(Y^{i_1,i_2}_{t_l})
$$

for $i = 0, \dots, k-1$. Again, $\nabla \theta(x, t)$ and $\nabla u(x, t)$ (which then update the values of $A(x, t)$ and $F(x, t)$ may be calculated by differentiating the iterations [\(7.10,](#page-36-0) [7.11\)](#page-36-1), which are defined by

$$
A(x,t_k) = \sum_{i_1} \sum_{i_2 \ge 0} h h_{i_1,i_2} \prod_{j=1}^k 1_{\mathbb{R}_+^2} (X_{t_j}^{i_1,i_2}) \nabla_x \Lambda_{2,\epsilon} (X_{t_k}^{i_1,i_2}, x) \wedge \hat{\omega}_{i_1,i_2}
$$

+
$$
\sum_{i_1} \sum_{i_2 \ge 0} \delta h h_{i_1,i_2} \sum_{j=1}^k \prod_{l=j}^k 1_{\mathbb{R}_+^2} (X_{t_l}^{i_1,i_2}) \nabla_x \Lambda_{2,\epsilon} (X_{t_k}^{i_1,i_2}, x) \wedge F(X_{t_{j-1}}^{i_1,i_2}, t_{j-1})
$$

+
$$
\sum_{i_1} \delta h h_w \sum_{j=1}^k \prod_{l=j}^k 1_{\mathbb{R}_+^2} (X_{t_l}^{i_1,0}) \nabla_x \Lambda_{2,\epsilon} (X_{t_k}^{i_1,0}, x) \wedge \tilde{\chi}_{\epsilon} (X_{t_{j-1}}^{i_1,0}, t_{j-1})
$$

and

$$
\frac{\partial}{\partial x_1} \theta(x, t_k) = \frac{T_w}{\pi} \left(\frac{x_2}{x_2^2 + (L + x_1)^2} - \frac{x_2}{x_2^2 + (L - x_1)^2} \right)
$$

$$
-\sum_{i_1}\sum_{i_2\geq 0} hh_{i_1,i_2}\prod_{j=1}^k 1_{\mathbb{R}_+^2}(Y_{t_j}^{i_1,i_2})\frac{\partial}{\partial x_1} A_{2,\epsilon}(Y_{t_k}^{i_1,i_2},x)\cdot R(x^{i_1,i_2},t_k;0)\Theta_{i_1,i_2}
$$

for $x = (x_1, x_2)$ with $x_2 > 0$, where the singular integral kernel is given by

$$
\nabla_x \Lambda_{2,\epsilon}(y,x) = \nabla_x \left(\left(1 - e^{-\frac{|y-x|^2}{\epsilon}} \right) \Lambda_{2,\epsilon}(y,x) \right).
$$

Of course, we need [\(7.1\)](#page-31-4) and [\(7.2\)](#page-31-5), which take precisely the same forms, except for $u(x,t)$ $\mathscr{R}(u(\overline{x}, t))$ if $x_2 < 0$. This completes the scheme. Finally, we would like to note that this scheme is designed for laminar flows.

Remark 7.2. It should be highlighted that the cost of computing the wall-bounded case is only marginally higher than the whole plane case. Though it seems that we need to store both diffusion paths X and Y , as well as the path of A , indeed, we only need to keep track of the latter, similar to the whole plane case. In each iteration, only the spot values of X and Y are needed.

7.2.2 Multi-copy scheme

Again, we may introduce another numerical scheme via the strong law of large numbers. By taking 2N independent Brownian particles that are modelled by the following discretised stochastic differential equations using the Euler scheme: for $1 \leq m \leq N$,

$$
X_0^{m,i_1,i_2} = x^{i_1,i_2}, \quad X_{t_k}^{m,i_1,i_2} = X_{t_{k-1}}^{m,i_1,i_2} + \delta u \left(X_{t_{k-1}}^{m,i_1,i_2}, t_{k-1} \right) + \sqrt{2\nu} (B_{t_k}^{1,m} - B_{t_{k-1}}^{1,m}),
$$

\n
$$
Y_0^{m,i_1,i_2} = x^{i_1,i_2}, \quad Y_{t_k}^{m,i_1,i_2} = Y_{t_{k-1}}^{m,i_1,i_2} + \delta u \left(Y_{t_{k-1}}^{m,i_1,i_2}, t_{k-1} \right) + \sqrt{2\kappa} (B_{t_k}^{2,m} - B_{t_{k-1}}^{2,m}),
$$

where $u(x,t) = \mathcal{R}(u(\overline{x}, t))$ if $x_2 < 0$, and the velocity u and temperature θ are approximated by

$$
u(x,t_k) = \frac{1}{N} \sum_{m=1}^{N} \sum_{i_1} \sum_{i_2 \ge 0} h h_{i_1,i_2} \prod_{j=1}^k 1_{\mathbb{R}_+^2} (X_{t_j}^{m,i_1,i_2}) \Lambda_{2,\epsilon}(X_{t_k}^{m,i_1,i_2}, x) \wedge \hat{\omega}_{i_1,i_2}
$$

+
$$
\frac{1}{N} \sum_{m=1}^{N} \sum_{i_1} \sum_{i_2 \ge 0} \delta h h_{i_1,i_2} \sum_{j=1}^k \prod_{l=j}^k 1_{\mathbb{R}_+^2} (X_{t_l}^{m,i_1,i_2}) \Lambda_{2,\epsilon}(X_{t_k}^{m,i_1,i_2}, x) \wedge F(X_{t_{j-1}}^{m,i_1,i_2}, t_{j-1})
$$

+
$$
\frac{1}{N} \sum_{m=1}^{N} \sum_{i_1} \delta h h_w \sum_{j=1}^k \prod_{l=j}^k 1_{\mathbb{R}_+^2} (X_{t_l}^{m,i_1,0}) \Lambda_{2,\epsilon}(X_{t_k}^{m,i_1,0}, x) \wedge \tilde{\chi}_{\epsilon}(X_{t_{j-1}}^{m,i_1,0}, t_{j-1})
$$

and

$$
\theta(x,t_k) = \frac{T_w}{\pi} \left(\tan^{-1} \left(\frac{L - x_1}{x_2} \right) + \tan^{-1} \left(\frac{L + x_1}{x_2} \right) \right) \n- \frac{1}{N} \sum_{m=1}^N \sum_{i_1} \sum_{i_2 \ge 0} h h_{i_1, i_2} \prod_{j=1}^k 1_{\mathbb{R}^2_+} (Y_{t_j}^{m, i_1, i_2}) A_2(Y_{t_k}^{m, i_1, i_2}, x) \cdot R(x^{i_1, i_2}, t_k; 0) \Theta_{i_1, i_2}.
$$

Similarly, for each m, the gauge functional equation is given by $R(x^{i_1,i_2}, t_k; t_k) = I$, and

$$
R(x^{i_1,i_2}, t_k; t_i) = I - \sum_{l=i+1}^k \delta A\left(Y_{t_l}^{m,i_1,i_2}, t_l\right) R(x^{i_1,i_2}, t_k; t_l) 1_{\mathbb{R}^2_+} (Y_{t_l}^{m,i_1,i_2})
$$

for $i = 0, \dots, k-1$. As for the derivatives, similar to the whole plane case, we have

$$
A(x,t_k) = \frac{1}{N} \sum_{m=1}^{N} \sum_{i_1} \sum_{i_2 \ge 0} h h_{i_1,i_2} \prod_{j=1}^{k} 1_{\mathbb{R}^2_+} (X^{m,i_1,i_2}_{t_j}) A_{2,\epsilon}(X^{m,i_1,i_2}_{t_k}, x) \wedge \hat{\omega}_{i_1,i_2}
$$

$$
+\frac{1}{N}\sum_{m=1}^{N}\sum_{i_1}\sum_{i_2\geq 0}\delta h h_{i_1,i_2}\sum_{j=1}^k\prod_{l=j}^k 1_{\mathbb{R}_+^2}(X_{t_l}^{m,i_1,i_2})\nabla_x A_{2,\epsilon}(X_{t_k}^{m,i_1,i_2},x)\wedge F(X_{t_{j-1}}^{m,i_1,i_2},t_{j-1})
$$

+
$$
\frac{1}{N}\sum_{m=1}^{N}\sum_{i_1}\delta h h_w\sum_{j=1}^k\prod_{l=j}^k 1_{\mathbb{R}_+^2}(X_{t_l}^{m,i_1,0})\nabla_x A_{2,\epsilon}(X_{t_k}^{m,i_1,0},x)\wedge \tilde{\chi}_{\varepsilon}(X_{t_{j-1}}^{m,i_1,0},t_{j-1})
$$

and

$$
\frac{\partial}{\partial x_1} \theta(x, t_k) = \frac{T_w}{\pi} \left(\frac{x_2}{x_2^2 + (L + x_1)^2} - \frac{x_2}{x_2^2 + (L - x_1)^2} \right) \n- \frac{1}{N} \sum_{m=1}^N \sum_{i_1} \sum_{i_2 \ge 0} h h_{i_1, i_2} \prod_{j=1}^k 1_{\mathbb{R}_+^2} (Y_{t_j}^{m, i_1, i_2}) \frac{\partial}{\partial x_1} A_{2,\epsilon}(Y_{t_k}^{m, i_1, i_2}, x) \cdot R(x^{i_1, i_2}, t_k; 0) \Theta_{i_1, i_2}.
$$

The rest of the scheme remains the same as in the one-copy case. This completes the SLN scheme for the wall-bounded Oberbeck-Boussinesq flows.

7.2.3 Numerical experiments

Here we carried out two sets of numerical experiments based on the one-copy scheme for the half-plane case. We consider two different Prandtl numbers: when the kinematic viscosity $\nu = 1$, and thermal diffusivity $\kappa = 0.15$, and when $\nu = 0.15$, $\kappa = 1$.

We choose the typical length scale $L = 2\pi$, consider the flow in the rectangle region $[-L, L] \times$ $[0, L]$ and assume that $\alpha = 0.0005$. The parameter we use to smooth out the Biot-Savart kernel is chosen to be $\epsilon = 0.1$.

In the experiment presented, we set the initial velocity to be of the form

$$
u(x,0) = (-10\sin(2x_2),0),
$$

and the initial temperature is given by

$$
\theta(x,0) = 0.01(8\pi^2 - x_1^2 - x_2^2).
$$

Thus, $\omega(x, 0) = 20 \cos(2x_2)$, and

$$
\hat{\omega}_{i_1, i_2} = \begin{cases} 20 \cos(2i_2 h), & i_2 > 0, \\ 0, & i_2 = 0. \end{cases}
$$

The time step is $\delta = 0.01$, and the boundary layer thickness is taken to be $\gamma = \pi/40$. The mesh size is set to be $h = 2\pi/40 \approx 0.157$ and $h_w = \gamma/20 \approx 0.00393$, and $\varepsilon = h_w/2$. We assume that the external heat source is at temperature $T_w = 0.1$, and

$$
\theta_0'(x_1) = -0.2x_1.
$$

The numerical experiment results at times $t = 0.2$, $t = 1.0$, $t = 1.8$ with Prandtl number $Pr = 0.15$ are shown in Figure [7.3,](#page-39-0) and the results at the given times with $Pr = 6.67$ are shown in Figure [7.4.](#page-40-2)

Figure 7.3: Temperature and velocity fields of Oberbeck-Boussinesq flows on \mathbb{R}^2_+ with Prandtl number $Pr = 0.15$.

Figure 7.4: Temperature and velocity fields of Oberbeck-Boussinesq flows on \mathbb{R}^2_+ with Prandtl number $Pr = 6.67$.

The simulations demonstrate well the regular pattern known as the Bénard convection, and also reveal the detailed hairy type of flows within the thin boundary layer, confirming the theoretical results and observations.

Data Availability Statement

No data are used in this article to support the findings of this study.

Acknowledgement

ZQ is supported partially by the EPSRC Centre for Doctoral Training in Mathematics of Random Systems: Analysis, Modelling and Simulation (EP/S023925/1).

Bibliography

[1] Alanko, S. 2016 Stability of Regression-Based Monte Carlo Methods for Solving Nonlinear PDEs. Communications on Pure and Applied Mathematics 69 no.5, 958-980.

- [2] Alda, W., Dzwinel, W., Witowski, J., Mościński, J., Pogoda, M. and Yuen, D. 1996 Rayleigh-Taylor instabilities simulated for a large system using molecular dynamics. Tech. Rep. UMSI 96/104, Supercomputer Institute, University of Minnesota, Minneapolis.
- [3] Anderson, C. 1986 Vorticity boundary conditions and boundary vorticity generation for two-dimensional viscous incompressible, J. Comput. Phys. 80, 72-97.
- [4] Anderson, C. and Greengard, C. 1985 On vortex methods. SIAM J. Numer. Anal. 22 (3), 413-440.
- [5] Anderson, C. and Greengard, C. eds. 1988 Vortex methods, Lecture Notes in Math. Vol. 1360, Springer-Verlag, Berlin.
- [6] Anderson, C. and Greengard, C., eds. 1991 Vortex dynamics and vortex methods, Lectures in Appl. Math. Vol. 28, AMS.
- [7] Bénard, H. 1900 Les tourbillons cellulaires dans une nappe liquide. Revue Gén. Sci. Pur. Appl. 11, 1261-1271 and 1309-1328.
- [8] Boussinesq, J. 1903 Theéorie analytique de la chaleur, vol. 2, Paris: Gauthier-Villars.
- [9] Busnello, B. 1999 A probabilistic approach to the two-dimensional Navier-Stokes equations. Ann. Probab. 27, no.4, 1750-1780.
- [10] Busnello, B., Flandoli, F. and Romito, M. 2005 A probabilistic representation for the vorticity of a three dimensional viscous fluid and for general systems of parabolic equations. Proc. Edinb. Math. Soc. 48, no 2, 295-336.
- [11] Celani, A., Cencini, M., Mazzino, A. and Vergassola, M. 2004 Active and passive fields face to face. New J. Phys, 6 72 (2004).
- [12] Chandrasekhar, S. 1961 Hydrodynamic and hydromagnetic stability. Oxford: Clarendon Press.
- [13] Chorin, A. J. 1973 Numerical study of slightly viscous flow. *J. Fluid Mech.* 57, 785-796.
- [14] Chorin, A. J. 1980 Vortex models and boundary layer instability. SIAM J. Sci. Statist. Comput. 1, no. 1, 1-21.
- [15] Constantin, P. 2001 An Eulerian-Lagrangian approach for incompressible fluids: local theory. J. Amer. Math. Soc. 14 no. 2, 263-278 (electronic).
- [16] Constantin, P. 2001 An Eulerian-Lagrangian approach to the Navier-Stokes equations. Comm. Math. Phys. 216, no. 3, 663-686.
- [17] Constantin, P. and Iyer, G. 2011 A stochastic-Lagrangian approach to the Navier-Stokes equations in domains with boundary, Ann. Appl. Probab. 21, 1466-1492 (2011).
- [18] Cottet, G. -H., and Koumoutsakos, P. D. 2000 Vortex Methods: Theory and Practice. Cambridge University Press.
- [19] Corrsin, S. 1951 On the spectrum of isotropic temperature fluctuations in isotropic turbulence, J. Appl. Phys. 22 469.
- [20] Criminale, W. O., Jackson, T. L. and Joslin, R. D. 2003 Theory and computation in hydrodynamic stability. Cambridge University Press.
- [21] Drazin, P. G. and Reid, W. H. 2004. Hydrodynamic stability. Second Edition. First Edition in 1981. Cambridge University Press.
- [22] Drivas, T.D. and Eyink, G.L. 2017 A Lagrangian fluctuation-dissipation relation for scalar turbulence. Part I. Flows with no boundary walls. Journal of Fluid Mechanics, Volume 829, 25 October 2017 , pp. 153 - 189 DOI: https://doi.org/10.1017/jfm.2017.567
- [23] Drivas, T. D. and Eyink, G.L. 2017 A Lagrangian fluctuation-dissipation relation for scalar turbulence. Part II. Wall-bounded flows. Journal of Fluid Mechanics. 829, 236-279 (2017).
- [24] Eyink, G., Gupta, A., and Zaki, T. 2020 Stochastic Lagrangian dynamics of vorticity. Part 1. General theory for viscous, incompressible fluids. Journal of Fluid Mechanics, 901, A2. doi:10.1017/jfm.2020.491
- [25] Eyink, G., Gupta, A., and Zaki, T. 2020 Stochastic Lagrangian dynamics of vorticity. Part 2. Application to near-wall channel-flow turbulence. Journal of Fluid Mechanics, 901, A3. doi:10.1017/jfm.2020.492
- [26] Falkovich, G., Gawędzki, K. and Vergassola, M. 2001 Particles and fields in fluid turbulence, Rev. Mod. Phys. 73 913-975.
- [27] Feynman, R. P. 1948 Space-time approach to non-relativistic quantum mechanics. Rev. Mod. Phys. Vol. 20, No. 2, 367-387.
- [28] Fletcher, C. A. J. 1991 Computational techniques for fluid dynamics, Vol. I and II, second edition. Springer-Verlag.
- [29] Friedman, A. 1964 Partial differential equations of parabolic type. Prentice-Hall, Inc.
- [30] Freidlin, M. 1985 Functional integration and partial differential equations. Princeton University Press.
- [31] Goodman, J. 1987 Convergence of the random vortex method. Comm. Pure Appl. Math. 40(2), 189-220.
- [32] Griebel, M., Knapek, S. and Zumbusch, G. 2007 Numerical simulation in molecular dynamics. Springer.
- [33] Helmholtz, H. 1858 Über die integrale der hydrodynamischen Gleichungen, welche den Wirbelbewegungen entsprechen, Jour. für die reine unde angewandte Math. also in Hermann von Helmholtz, Wissenschaftliche Abhandlungen, Vol. 1, pp. 101-104.
- [34] Holm, D., Marsden, J. and Ratiu, T. 1998 The Euler-Poincaré equations and semidirect products with applications to continuum theories. Adv. Math. 137 no.1, 1-81.
- [35] Iyer, G. 2006 A stochastic perturbation of inviscid flows. Comm. Math. Phys. 266 no. 3, 631-645.
- [36] Joseph, D. D. 1965 On the stability of the Boussibesq equations. Arch. Rat. Mech. Anal. 20, 59-71.
- [37] Joseph, D. D. 1966 Non-linear stability of the Boussibesq equations by the method of energy. Arch. Rat. Mech. Anal. 22, 163-84.
- [38] Joseph, D. D. 1976 Stability of fluid motions. I and II. Springer Tracts in Natural Philosophy, vol. 28, Springer-Verlag, New York.
- [39] Kac, M. 1949 On Distributions of Certain Wiener Functionals. Transactions of the American Mathematical Society , Jan., 1949, Vol. 65, No. 1 (Jan., 1949), pp. 1-13
- [40] Kloeden, P. E. and Platen, E. 1992 Numerical Solutions to Stochastic Differential Equations. Springer-Verlag Berlin Heidelberg.
- [41] Kolmogorov, A. N. 1941 The local structure of turbulence in incompressible viscous fluid for very large reynolds numbers. Comptes Rendus de l'Académie des Sciences de l'URSS, 30:301–305, 1941 (reprinted in Proc. R. Soc. Lond. A 434, 9-13, 1991).
- [42] Kolmogorov A. N. 1941 Dissipation of energy in the locally isotropic turbulence. Comptes Rendus de l'Académie des Sciences de l'URSS, 32:16–18, 1941 (reprinted in Proc. R. Soc. Lond. A 434, 15-17, 1991).
- [43] Kraichnan, R. H. 1968 Small-scale structure of a scalar field convected by turbulence. Phys. Fluids 11 945-953.
- [44] Landau, L. D. and Lifshitz, E. M. 1987 Fluid mechanics. Second edition. Pergamon Press.
- [45] LeJan, Y. and Sznitman, A. S. 1997 Stochastic cascades and 3-dimensional Navier-Stokes equations. Probab. Theory Related Fields 109 no. 3, 343-366.
- [46] LeJan, Y. and Raimond, O. 2002 Integration of Brownian vector fields, Ann. Probab. 30, 826-873.
- [47] LeJan, Y. and Raimond, O. 2004 Flows, coalescence and noise, Ann. Probab. 32, 1247-1315.
- [48] Lesieur, M., Métais, O. and Comte, P. 2005 Large-Eddy simulations of turbulence. Cambridge University Press.
- [49] Long, D. G. 1988 Convergence of the random vortex method in two dimensions. J. of Amer. Math. Soc. **1**(4), 779-804.
- [50] Lyons, T. J. and Zheng, W. 1988 A crossing estimate for the canonical process on a Dirichlet space and a tightness result. Astérisque, tome 157-158, p. 249-271.
- [51] Lyons, T. J. and Zheng, W. 1990 On conditional diffusion processes. Proceedings of the Royal Society of Edinburgh Section A: Mathematics Vol. 115, Issue 3-4, 243-255.
- [52] Majda, A. J. and Bertozzi A. L. 2002 Vorticity and incompressible flow. Cambridge University Press.
- [53] McKean Jr, H. P. 1966 A class of Markov processes associated with nonlinear parabolic equations. Proceedings of the National Academy of Sciences of the United States of America, 56(6), 1907. [MR0221595](https://mathscinet.ams.org/mathscinet/search/publdoc.html?arg3=&co4=AND&co5=AND&co6=AND&co7=AND&dr=all&pg4=AUCN&pg5=TI&pg6=PC&pg7=ALLF&pg8=ET&r=1&review_format=html&s4=&s5=A%5C%20Class%5C%20of%5C%20Markov%5C%20Processes%5C%20Associated%5C%20with%5C%20Nonlinear%5C%20Parabolic%5C%20Equations&s6=&s7=&s8=All&sort=Newest&vfpref=html&yearRangeFirst=&yearRangeSecond=&yrop=eq)<https://doi.org/10.1073/pnas.56.6.1907>
- [54] Oberbeck, A. 1879 Ueber die Wärmleitung der Flüssigkeiten bei Berücksichtigung der Strömungen infolge von Temperaturdifferenzen. Ann. Phys. Chem. 7, 271-92.
- [55] Oboukhov, A. M. 1949 Structure of the temperature field in turbulent flows, Izv. Akad. Nauk. SSSR, Geogr. and Geophyys 13 58.
- [56] Pardoux, E. and Peng, S. G. 1990 Adapted solution of a backward stochastic differential equation. Systems $\mathcal C$ Control Letters Vol. 14 no.1, 55-61.
- [57] Peskin, C. 1985 A random-walk interpretation of the incompressble Navier-Stokes equations. Comm. Pure Appl. Math. 38 no. 6, 845-852.
- [58] Pope, S. B. 2000 Turbulent Flows. Cambridge University Press.
- [59] Qian, Z. 2022 Stochastic formulation of incompressible fluid flows in wall-bounded regions. arXiv:2206.05198
- [60] Qian Z. Süli E. and Zhang Y. 2022 Random vortex dynamics via functional stochastic differential equations. Proc. R. Soc. A 478: 20220030. https://doi.org/10.1098/rspa.2022.0030
- [61] Qian, Z., Qiu, Y., Zhao, L. and Wu, J. 2022 Monte-Carlo simulations for wall-bounded fluid flows via random vortex method. arXiv:2208.13233
- [62] Rayleigh, L. 1916 On convection currents in a horizontal layer of fluid, when the higher temperature is on the under side. Phil. Mag. (6) 32, 529-46.
- [63] Sengupta, T. K. and Bhaumik, S. 2019 DNS of wall-bounded turbulent flows – A first principle approach. Springer Nature Singapore Pte Ltd.
- [64] Shlesinger, M. F., West, B. J. and Klafter, J. 1987 Lévy dynamics of enhanced diffusion: application to turbulence. Phys. Rev. Lett. 58 (11), 1100.
- [65] Stroock, D. and Varadhan, S. R. S. 1979 Multidimensional diffusion processes. Springer-Verlag Berlin Heidelberg.
- [66] Taylor, G. I. 1921 Diffusion by continuous movements. Proc. Lond. Math. Soc. 20, 196.
- [67] Taylor, G. I. 1935 Statistical theory of turbulence. Parts 1-4. Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences, 151(873):421–478, 1935. https://doi.org/10.1098/rspa.1935.0159.
- [68] Von Kármán, T. 1931 Mechanical Similitude and Turbulence. National Advisory Committee for Aeronautics, 1931.
- [69] Thalabard, S., Krstu;ovic, G. and Bec, J. 2014 Turbulent pair dispersion as a continuoustime random walk. J. Fluid Mech. 755, R4.
- [70] Vanden-Eijnden, E. W. and Vanden-Eijnden, E. 2000 Generalized flows, intrinsic stochasticity and turbulent transport. Proc. Natl Acad. Sci. USA 97, 8200-8205.
- [71] Vanden-Eijnden, E. W. and Vanden-Eijnden, E. 2001 Turbulent Prandtl number effect on passive scalar advection. Physic D, $152 - 153$, 636-645.
- [72] Wesseling, P. 2001 Principles of computational fluid dynamics. Springer-Verlag Berlin Heidelberg.
- [73] Wung, T. and Tseng, F. 1992 A color-coded particle tracking velocimeter with application to natural convection, Experimental Fluids, 13, 217-223.
- [74] Yu, H., Kanov, K., Perlman, E., Graham, J., Frederix, E., Burns, R., Szalay, A., Eyink, G. L. and Meneveau, C. 2012 Studying Lagrangian dynamics of turbulence using on-demand fluid particle tracking in a public turbulence database. J. Turbul. **12**, N13.
- [75] Zhang, X. 2010 A stochastic representation for backward incompressible Navier-Stokes equations. Probab. Theory Related Fields 148, 305-332.