Learning from Similar Linear Representations: Adaptivity, Minimaxity, and Robustness

γ e Tian γ e.t@columbia.edu

Department of Statistics Columbia University New York, NY 10027, USA

Yuqi Gu yuqi.gu@columbia.edu

Department of Statistics Columbia University New York, NY 10027, USA

Department of Biostatistics, School of Global Public Health New York University New York, NY 10003, USA

Yang Feng yang. Tenggal yang terletak di terletak yang yang yang kerama yang dikenali berada yang dikelui dala

Abstract

Representation multi-task learning (MTL) has achieved tremendous success in practice. However, the theoretical understanding of these methods is still lacking. Most existing theoretical works focus on cases where all tasks share the same representation, and claim that MTL almost always improves performance. Nevertheless, as the number of tasks grows, assuming all tasks share the same representation is unrealistic. Furthermore, empirical findings often indicate that a shared representation does not necessarily improve single-task learning performance. In this paper, we aim to understand how to learn from tasks with similar but not exactly the same linear representations, while dealing with outlier tasks. Assuming a known intrinsic dimension, we proposed a penalized empirical risk minimization method and a spectral method that are *adaptive* to the similarity structure and *robust* to outlier tasks. Both algorithms outperform single-task learning when representations across tasks are sufficiently similar and the proportion of outlier tasks is small. Moreover, they always perform at least as well as single-task learning, even when the representations are dissimilar. We provided information-theoretic lower bounds to demonstrate that both methods are nearly *minimax* optimal in a large regime, with the spectral method being optimal in the absence of outlier tasks. Additionally, we introduce a thresholding algorithm to adapt to an unknown intrinsic dimension. We conducted extensive numerical experiments to validate our theoretical findings.

1. Introduction

1.1 Representation Multi-task Learning

With the increase in computational power, machine learning systems can now process datasets of a large scale. However, for each machine learning task, we may not have access to a large amount of data due to data privacy restrictions and high cost of data acquisition. This motivated the idea of multi-task learning (MTL), where we learn many similar but not exactly the same tasks jointly to enhance model performance [\(Zhang and Yang,](#page-40-0) [2018,](#page-40-0) [2021\)](#page-40-1).

Related concepts include transfer learning (TL), learning-to-learn, and meta-learning, where model structures learned from multiple tasks can be transferred to new incoming tasks to improve their performances [\(Weiss et al.,](#page-39-0) [2016;](#page-39-0) [Hospedales et al.,](#page-37-0) [2021\)](#page-37-0). Among numerous multi-task and transfer learning approaches, representation learning has been one of the most popular and successful methods over the past few years, where a data representation is jointly learned from multiple similar data sets and can be shared across them [\(Rostami](#page-38-0) [et al.,](#page-38-0) [2022\)](#page-38-0). A successful example of multi-task and transfer representation learning is learning the weights of a few initial layers of neural networks from ImageNet pre-training, then retraining final layers on new image classification tasks [\(Donahue et al.,](#page-35-0) [2014;](#page-35-0) [Goyal](#page-36-0) [et al.,](#page-36-0) [2019\)](#page-36-0). Other applications include multilingual knowledge graph completion [\(Chen](#page-35-1) [et al.,](#page-35-1) [2020\)](#page-35-1) and reinforcement learning [\(Gupta et al.,](#page-36-1) [2017\)](#page-36-1).

While representation learning has been successful in practice, its theoretical understanding in the context of multi-task and transfer learning remains limited. Most existing theoretical works assume that the same representation is shared across all tasks, which is not always realistic in scenarios with a large number of tasks [\(Rostami et al.,](#page-38-0) [2022\)](#page-38-0). Furthermore, empirical studies have shown that freezing a representation across tasks from different contexts may not improve model performance and can even be harmful. For example, [Raghu et al.](#page-38-1) [\(2019\)](#page-38-1) found that pre-training on ImageNet offered little help to target medical tasks, and [Wang et al.](#page-39-1) [\(2019\)](#page-39-1) found that different target tasks might benefit from different pre-training in natural language understanding. Both suggested that a frozen representation may not always work well. Additionally, there may be outlier tasks that are dissimilar to other tasks [\(Zhang and Yang,](#page-40-1) [2021\)](#page-40-1) or may be contaminated with adversarial attacks on the data [\(Qiao,](#page-38-2) [2018;](#page-38-2) [Qiao and Valiant,](#page-38-3) [2018;](#page-38-3) [Konstantinov et al.,](#page-37-1) [2020\)](#page-37-1). If left unaddressed, such issues could severely impact the machine learning system's overall performance.

This paper investigates the effective learning of tasks with *similar representations* in the presence of potential outlier tasks or adversarial attacks. Specifically, we consider the following linear model with linear representations. Suppose there are T tasks in total, and we have collected sample $\{\boldsymbol{x}_i^{(t)}\}$ $\binom{t}{i}, y_{i}^{(t)}$ $\{a_i^{(t)}\}_{i=1}^n$ from the t-th task, where $\mathbf{x}_i^{(t)} \in \mathbb{R}^p$, $y_i^{(t)} \in \mathbb{R}$, and $t \in [T] = \{1, 2, \ldots, T\}$. There exists an unknown subset $S \subseteq [T]$, such that for all $t \in S$,

$$
y_i^{(t)} = (\boldsymbol{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*} + \epsilon_i^{(t)}, \quad i = 1:n,
$$
\n(1)

where the regression coefficient $\boldsymbol{\beta}^{(t)*} = \boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}$, the representation $\boldsymbol{A}^{(t)*} \in \mathcal{O}^{p \times r} = \{ \boldsymbol{A} \in \mathcal{A}^{(t)} \}$ $\mathbb{R}^{p \times r}$: $\mathbf{A}^{\top} \mathbf{A} = \mathbf{I}_r$, low-dimensional parameter $\boldsymbol{\theta}^{(t)*} \in \mathbb{R}^r$, $r \leq p$, and $\{\epsilon_i^{(t)}\}$ $\binom{t}{i}$ _{i=1} are random noises. Here r represents the intrinsic dimension of the problem, which is usually much smaller than p. The data $\{\boldsymbol{x}_i^{(t)}\}$ $\binom{t}{i}, y_{i}^{(t)}$ $\{e^{(t)}\}_{i=1}^n$ for $t \notin S$ can be arbitrarily distributed in the worst case, and the corresponding tasks in $S^c = T|\S$ are outlier or contaminated tasks. We call $\epsilon := |S^c|/T$ the *contamination proportion* or the *proportion of outlier tasks*. To ensure effective learning from similar representations, we assume that ${A^{(t)*}}_{t \in S}$ are similar to each other, in the sense that $\min_{\overline{\mathbf{A}} \in \mathcal{O}^{p \times r}} \max_{t \in S} ||\mathbf{A}^{(t)*}(\mathbf{A}^{(t)*})^{\top} - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}||_2 \leq h$, where $\overline{A} \in \mathcal{O}^{p \times r}$ achieving the minimum can be understood as a "central representation" and h is the similarity measure. Our goal is to explore the upper and lower error bounds in estimating $\{\beta^{(t)*}\}_{t\in S}$ for all possible cases of S under certain conditions. Furthermore, when the tasks in S^c also satisfy the linear model (1) , we aim to ensure the effective estimation of $\{\boldsymbol{\beta}^{(t)*}\}_{t\in S^c}$ as well.

It is worth pointing out that we allow the scales of $\{\theta^{(t)*}\}_{t\in S}$, i.e. $\{\|\theta^{(t)*}\|_2\}_{t\in S}$, to differ across tasks in S, and $\|\theta^{(t)*}\|_2$ can also diverge as $n \to \infty$. Here, $\|\theta^{(t)*}\|_2 = \|\beta^{(t)*}\|_2$ can be viewed as the signal strength of the t-th task. It turns out that the performance of representation MTL on each task is highly related to $\|\boldsymbol{\theta}^{(t)*}\|_2$. In contrast, existing literature generally assumes $\|\theta^{(t)*}\|_2 \approx 1$ for most of $t \in [T]$ and ignores the impact of $\|\theta^{(t)*}\|_2$ on the model performance.

1.2 Related Works

1.2.1 Multi-task and Transfer Learning

Representation MTL and TL: [Baxter](#page-34-0) [\(2000\)](#page-34-0) is among the earliest works to study the theory of representation MTL under general function classes, where all tasks are generated from the same distribution. [Maurer et al.](#page-38-4) [\(2016\)](#page-38-4) improved their results by using the analysis based on Rademacher complexity. [Ando et al.](#page-34-1) [\(2005\)](#page-34-1) explored the case of semi-supervised learning. More recently, [Du et al.](#page-36-2) [\(2020\)](#page-36-2) and [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2) studied linear model [\(1\)](#page-1-0) with $S = [T]$ and $h = 0$, i.e., under the assumption that there are no outlier tasks and all tasks share the same representation. They proposed the so-called task diversity condition, under which the learning rate can be significantly improved by a non-convex emprical risk minimization (ERM) algorithm. [Tripuraneni et al.](#page-39-3) [\(2020\)](#page-39-3) extended the analysis to general non-linear models and provided general results. [Thekumparampil et al.](#page-39-4) [\(2021\)](#page-39-4) proposed a polynomial-time alternating gradient descent algorithm that achieves similar performance as ERM but avoids solving the non-concex optimization directly. Other related works include federated representation learning [\(Collins et al.,](#page-35-2) [2021;](#page-35-2) [Duchi et al.,](#page-36-3) [2022\)](#page-36-3), tensor representation meta-learning [\(Deng et al.,](#page-35-3) [2022\)](#page-35-3), conditional meta-learning [\(Denevi et al.,](#page-35-4) [2020\)](#page-35-4), and matrix completion via representation MTL [\(Zhou et al.,](#page-40-2) [2021\)](#page-40-2). Note that MTL under the assumption that $\beta^{(t)*}$'s in [\(1\)](#page-1-0) share the same or similar support sets [\(Lounici](#page-37-2) [et al.,](#page-37-2) [2009,](#page-37-2) [2011;](#page-38-5) [Jalali et al.,](#page-37-3) [2010;](#page-37-3) [Li et al.,](#page-37-4) [2021;](#page-37-4) [Xu and Bastani,](#page-40-3) [2021\)](#page-40-3) can also be viewed as a special case of the general representation MTL.

Distance-based MTL and TL: There has been much literature in the statistics community studying model [\(1\)](#page-1-0) under the assumption that Euclidean distance or ℓ_1 -distance between $\beta^{(t)*}$'s are small [\(Bastani,](#page-34-2) [2021;](#page-34-2) [Li et al.,](#page-37-5) [2022b;](#page-37-5) [Duan and Wang,](#page-36-4) [2022;](#page-36-4) [Gu et al.,](#page-36-5) [2023\)](#page-36-5), which is called "distance-based" MTL and TL in [Gu et al.](#page-36-6) [\(2022\)](#page-36-6). Some extensions include high-dimensional GLMs [\(Tian and Feng,](#page-39-5) [2022\)](#page-39-5), graphical models [\(Li et al.,](#page-37-6) [2022a\)](#page-37-6), functional regression [\(Lin and Reimherr,](#page-37-7) [2022\)](#page-37-7), semi-supervised classification [\(Zhou et al.,](#page-40-4) [2022\)](#page-40-4), and unsupervised Gaussian mixture models [\(Tian et al.,](#page-39-6) [2022\)](#page-39-6). Recently, [Gu et al.](#page-36-6) [\(2022\)](#page-36-6) proposed the "angle-based" TL where they assume the angle between every pair of $\beta^{(t)*}$'s is small. As we will discuss in the next section, their setting is a special case of [\(1\)](#page-1-0) when $r = 1$.

Other related literature: Other relative literature include the non-parametric TL [\(Hanneke and Kpotufe,](#page-36-7) [2019;](#page-36-7) [Cai and Wei,](#page-35-5) [2021\)](#page-35-5), the hardness of MTL [\(Hanneke and Kpotufe,](#page-36-8) [2022\)](#page-36-8), adversarial robustness of MTL [\(Qiao,](#page-38-2) [2018;](#page-38-2) [Qiao and Valiant,](#page-38-3) [2018;](#page-38-3) [Konstantinov](#page-37-1) [et al.,](#page-37-1) [2020\)](#page-37-1), gradient-based meta-learning [\(Finn et al.,](#page-36-9) [2017;](#page-36-9) [Nichol et al.,](#page-38-6) [2018;](#page-38-6) [Finn et al.,](#page-36-10) [2019\)](#page-36-10), and theory of MTL based on distributional measure [\(Ben-David and Borbely,](#page-34-3) [2008;](#page-34-3) [Ben-David et al.,](#page-34-4) [2010\)](#page-34-4).

To help readers better understand the difference between some settings in literature with our setting under linear model [\(1\)](#page-1-0), we drew Figure [1](#page-3-0) as a simple visualization corresponding to the case where $p = 3$ and $r = 2$.

Figure 1: A simple visualization of four different settings under linear model [\(1\)](#page-1-0).

1.2.2 Beyond the Assumption of the Same Representation

Several works have studied similar problems to the current work but in different formulations. [Chua et al.](#page-35-6) [\(2021\)](#page-35-6) explored linear model [\(1\)](#page-1-0) but with the assumption that $\mathbf{A}^{(t)*} = \overline{\mathbf{A}} + \mathbf{\Delta}^{(t)*}$ with some $\overline{A} \in \mathbb{R}^{p \times r}$ and $\|\Delta^{(t)*}\|_F \leq h$ and $S = [T]$. [Duan and Wang](#page-36-4) [\(2022\)](#page-36-4) considered the same model with $\boldsymbol{\beta}^{(t)*} = \overline{\boldsymbol{A}} \boldsymbol{\theta}^{(t)*} + \boldsymbol{\delta}^{(t)*}$ with $\|\boldsymbol{\delta}^{(t)*}\|_2 \leq h$ and $S = [T]$.

To gain a deep understanding about the relationship between the problems studied in the literature and our paper, we first examine the following "most general scenario":

$$
\boldsymbol{\beta}^{(t)*} = \overline{\boldsymbol{A}} \boldsymbol{\theta}^{(t)*} + \boldsymbol{\delta}^{(t)*}, \quad \forall t \in S \subseteq [T], \tag{2}
$$

where $\overline{A} \in \mathcal{O}^{p \times r}$, $\theta^{(t)*} \in \mathbb{R}^r$, and $\delta^{(t)*} \in \mathbb{R}^p$. Note that there is no restriction on $\|\theta^{(t)*}\|_2$ and $\|\boldsymbol{\delta}^{(t)*}\|_2$, although we expect MTL to be beneficial only when $\|\boldsymbol{\delta}^{(t)*}\|_2$ is small. We call it the most general scenario because it includes most of the regimes studied in the literature, including our setting. To see this, consider the following cases.

- (i) When $S = [T]$ and $\delta^{(t)*} = 0$ for all $t \in S$, [\(2\)](#page-3-1) reduces to the case of shared representation discussed in [Du et al.](#page-36-2) [\(2020\)](#page-36-2); [Thekumparampil et al.](#page-39-4) [\(2021\)](#page-39-4); [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2). [a](#page-4-0)
- (ii) When $S = [T]$ and $||\boldsymbol{\theta}^{(t)*}||_2 \ge 1$ for all $t \in S$, [\(2\)](#page-3-1) reduces to the regime studied in [Chua](#page-35-6) [et al.](#page-35-6) [\(2021\)](#page-35-6) and [Duan and Wang](#page-36-4) [\(2022\)](#page-36-4), because we can write $\boldsymbol{\delta}^{(t)*} = \boldsymbol{\Delta}^{(t)*} \boldsymbol{e}_1$ with $\mathbf{\Delta}^{(t)*} = \big(\boldsymbol{\delta}^{(t)*} \quad \mathbf{0}_{p \times (r-1)}\big).$
- (iii) [\(2\)](#page-3-1) also includes our setting defined in Section [1.1,](#page-0-0) because by Lemma 2.6 in [Chen](#page-35-7) [et al.](#page-35-7) [\(2021b\)](#page-35-7), for any $t \in S$, there exists $\mathbf{R}^{(t)} \in \mathcal{O}^{r \times r}$ such that

$$
\|\mathbf{A}^{(t)*}-\overline{\mathbf{A}}\mathbf{R}^{(t)}\|_2 \leq \sqrt{2}\|\mathbf{A}^{(t)*}(\mathbf{A}^{(t)*})^{\top}-\overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}\|_2 \leq \sqrt{2}h.
$$

Thus, we can write $\boldsymbol{\beta}^{(t)*} = \boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}$ in our setting as $\boldsymbol{\beta}^{(t)*} = \overline{\boldsymbol{A}} \boldsymbol{R}^{(t)} \boldsymbol{\theta}^{(t)*} + \boldsymbol{\delta}^{(t)*}$, with $\boldsymbol{\delta}^{(t)*} = (\boldsymbol{A}^{(t)*} - \overline{\boldsymbol{A}} \boldsymbol{R}^{(t)}) \boldsymbol{\theta}^{(t)*}.$

We focus on the current setting defined in Section [1.1](#page-0-0) because it is simpler to analyze and leads to cleaner results. It is worth mentioning that the second algorithm we proposed – the spectral method – also works in the general scenario [\(2\)](#page-3-1). We focus on the setting in Section [1.1](#page-0-0) in the main text, and present the extension to the general regime in Section [A](#page-42-0) of the appendix.

Compared to the settings in [Chua et al.](#page-35-6) [\(2021\)](#page-35-6) and [Duan and Wang](#page-36-4) [\(2022\)](#page-36-4), our formulation leads to simpler learning algorithms, which can adapt to the unknown similarity level h (unlike [Chua et al.,](#page-35-6) [2021,](#page-35-6) where the algorithm requires tuning parameters depending on the similarity level h), with stronger theoretical results. Our formulation also inspires robust algorithms whose performance is not greatly affected by a small fraction of outlier tasks, a merit not shared by the methods following the formulations in [Chua et al.](#page-35-6) [\(2021\)](#page-35-6) and [Duan and Wang](#page-36-4) [\(2022\)](#page-36-4). Moreover, our framework is easier to generalize to an unsupervised learning setting (e.g., multiple linear/nonlinear latent factor models with similar factor loading matrices or similar factor score matrices). Additionally, a representation-based reweighting strategy was proposed in [Chen et al.](#page-35-8) [\(2021a\)](#page-35-8), which is motivated by the concern of assuming the same representation. Their similarity metric between tasks depends on the weight assigned to the objective function of each task, while our similarity metric depends on the difference between representations explicitly, which is more intuitive. Moreover, their approach can suffer from a negative transfer in the worst case, while our approaches do not. Furthermore, none of [Chen et al.](#page-35-8) [\(2021a\)](#page-35-8); [Chua et al.](#page-35-6) [\(2021\)](#page-35-6); [Duan and Wang](#page-36-4) [\(2022\)](#page-36-4) considered the presence of outlier tasks.

Finally, we present a diagram in Figure [2](#page-5-0) to summarize the relationship between the different regimes studied in various papers mentioned earlier. To compare different algorithms proposed under different settings, we can focus on the overlapping area surrounded by a dash line in Figure [2](#page-5-0) and compare the estimation errors of $\beta^{(t)*}$'s for different approaches therein. In this overlapping regime, we have our setting defined in Section [1.1](#page-0-0) with $\min_{\overline{\mathbf{A}} \in \mathcal{O}^{p \times r}} \max_{t \in S} \| \mathbf{A}^{(t)*} (\mathbf{A}^{(t)*})^{\top} - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top} \|_2 \leq h$ and $\|\boldsymbol{\theta}^{(t)*}\|_2 \approx 1$ for all $t \in S$. We summarize the estimation error of different approaches in Table [1,](#page-6-0) highlighting our two

a. [Du et al.](#page-36-2) [\(2020\)](#page-36-2) also studied the non-linear representations; here we are referring to their setting of linear representations.

proposed algorithms: penalized ERM ("pERM") and the spectral method ("Spectral"). We show that the spectral method is minimax optimal with computational efficiency when there is no contamination ($\epsilon = 0$), and pERM can handle the contaminated case more effectively.

Figure 2: The diagram illustrating the relationship between different regimes studied in different papers.^{[b](#page-5-1)}

1.3 Our Contributions

Our contributions can be summarized below.

- (i) Compared to most literature on representation MTL, we considered a more general framework. Here, the linear representations can vary across tasks, signal strengths may differ between tasks, and there can be a small fraction of unknown outlier tasks.
- (ii) We proposed two algorithms, the penalized ERM and the spectral method, to learn the regression coefficients and the representations from multiple tasks. Our algorithms were shown to have the following properties:
	- They outperform the single-task when the representations of different tasks are sufficiently similar and the proportion of outlier tasks is low.
	- They guarantee no worse performance than single-task learning (safe-net guarantee) even when task representations are dissimilar.

b. More rigorously speaking, [Du et al.](#page-36-2) [\(2020\)](#page-36-2); [Chua et al.](#page-35-6) [\(2021\)](#page-35-6); [Thekumparampil et al.](#page-39-4) [\(2021\)](#page-39-4); [Tripuraneni](#page-39-2) [et al.](#page-39-2) [\(2021\)](#page-39-2); [Duan and Wang](#page-36-4) [\(2022\)](#page-36-4) only require $\max_{t \in [T]} ||\theta^{(t)*}||_2 \lesssim 1$ and a constant proportion of $t \in [T]$ satisfies $\|\boldsymbol{\theta}^{(t)*}\|_2 \gtrsim 1$. For simplicity, we write their requirement as $\|\boldsymbol{\theta}^{(t)*}\|_2 \asymp 1$.

Regime	Algorithm	$\max_{t \in S} \ \widehat{\beta}^{(t)} - \beta^{(t)*}\ _2$	Optimal $(\epsilon = 0)$? Poly-time?	
$h=\epsilon=0$	ERM	$r\sqrt{\frac{p}{nT}}+r\sqrt{\frac{1}{n}}$	No	No
	MoM	$\frac{p}{nT} + \sqrt{\frac{r}{n}}$	No	Yes
	$Although$ $AltMinGD$	$r\sqrt{\frac{p}{nT}}+\sqrt{\frac{r}{n}}$	No	Yes
$h\neq 0$	AdaptRep	$\left r\sqrt{\frac{p}{nT}}+r\sqrt{\frac{1}{n}}+\sqrt{rh}(\frac{p}{n})^{1/4}\right \wedge\sqrt{\frac{p}{n}}$	No	No
$\epsilon = 0$	ARMUL	$\left(r\sqrt{\frac{p}{nT}}+r\sqrt{\frac{1}{n}}+rh\right)\wedge\left(r\sqrt{\frac{p}{n}}\right)$	N _o	No
$h, \epsilon \neq 0$	pERM	$\left(r\sqrt{\frac{p}{nT}}+r\sqrt{\frac{1}{n}}+\sqrt{r}h+\epsilon\frac{r^{3/2}\sqrt{p}}{\sqrt{n}}\right)\wedge\sqrt{\frac{p}{n}}$	N _o	$\rm No$
	Spectral	$\left(\sqrt{\frac{pr}{nT}}+\sqrt{\frac{r}{n}}+h+\sqrt{\epsilon r}\right)\wedge\sqrt{\frac{p}{n}}$	Yes	Yes
	Single-task		No	Yes
Lower-bound		$\sqrt{\frac{pr}{nT}} + \sqrt{\frac{r}{n}} + h + \epsilon \frac{r}{\sqrt{n}} \big) \wedge \sqrt{\frac{p}{n}}$		

Table 1: A summary of the high-probability estimation error of $\beta^{(t)*}$'s for different methods in different regimes, where the two algorithms we proposed, penalized ERM ("pERM", Algorithm [1\)](#page-11-0) and the spectral method ("Spectral", Algorithm [2\)](#page-17-0), are highlighted. Denote the contamination proportion $\epsilon = |S^c|/T$. Except for the lower bound, all the other rates are upper bounds. All the rates are up to logarithmic factors. "ERM": [Du et al.](#page-36-2) [\(2020\)](#page-36-2); [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2). "MoM": method-of-moments, [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2). "AltMinGD": alternating minimization gradient descent, [Thekumparampil et al.](#page-39-4) [\(2021\)](#page-39-4). "AdaptRep": adaptive representation learning, [Chua et al.](#page-35-6) [\(2021\)](#page-35-6). "ARMUL": adaptive and robust multi-task learning, [Duan and Wang](#page-36-4) [\(2022\)](#page-36-4). "Single-task": single-task regression by only using the local data from each task.

- (iii) We conducted a thorough analysis of the relationship between different regimes of representation MTL studied in the literature. Our derived upper bounds improve over existing rates, particularly in scenarios without task contamination. Furthermore, we are the first to examine the scenario where a few tasks are contaminated, in the context of representation MTL.
- (iv) We derived the lower bounds for model [\(1\)](#page-1-0). To the best of our knowledge, these are the first lower bound results for regression coefficient estimation under the representation MTL. Prior works such as [Duchi et al.](#page-36-3) [\(2022\)](#page-36-3) and [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2) provided analogous lower bounds for the subspace recovery problem assuming identical representations without outlier tasks. In a TL setup, [Chua et al.](#page-35-6) [\(2021\)](#page-35-6) showed that assuming the same representation can lead to worse performance than target-only learning when source representations differ from each other, but they did not provide a full lower bound which relates to the representation difference. Comparing the upper and lower bounds, we demonstrated that both algorithms we proposed are nearly minimax optimal, with the spectral method being optimal in uncontaminated settings.
- (v) We extended our analysis from linear model [\(1\)](#page-1-0) to generalized linear models (GLMs) and non-linear regression models, and obtained similar theoretical guarantees in these settings.
- (vi) We proposed a thresholding algorithm based on singular value decomposition (SVD) to estimate the unknown intrinsic dimension r . This adaptation enables our penalized ERM and spectral methods to handle cases where r is unknown, addressing a common challenge where r is not a priori known in most prior works.

1.4 Notations and Organization

Throughout the paper, we use bold capitalized and lower-case letters to denote matrices and vectors, respectively. For a real number a, |a| stands for its absolute value. For a vector u , $||u||_2$ stands for its Euclidean norm. For a matrix \mathbf{A} , $||\mathbf{A}||_2$ and $||\mathbf{A}||_F$ represent its spectral and Frobenius norm, respectively. A^{\top} denotes its transpose. $\sigma_j(A)$, $\sigma_{\max}(A)$, $\sigma_{\min}(A)$ are its j-th largest singular value, maximum singular value, and minimum (non-zero) singular value, respectively. When \boldsymbol{A} is a square matrix, we denote its maximum and minimum eigenvalues as $\lambda_{\max}(A)$ and $\lambda_{\min}(A)$, respectively. For a function $\psi: \mathcal{X} \to \mathbb{R}$, $\|\psi\|_{\infty}$ is defined to be $\max_{x \in \mathcal{X}} |\psi(x)|$. For two real numbers a and b, we denote their minimum by $\min\{a, b\}$ or $a \wedge b$ and their maximum by $\max\{a, b\}$ or $a \vee b$, respectively. For two positive real sequences ${a_n}_{n=1}^{\infty}$ and ${b_n}_{n=1}^{\infty}$, $a_n \lesssim b_n$ or $b_n \gtrsim a_n$ means there exists a universal constant $C > 0$ such that $a_n \leq C b_n$ for all n, and $a_n \ll b_n$ or $b_n \gg a_n$ means that $a_n/b_n \to 0$ as $n \to \infty$. $a_n \leq b_n$ means $a_n \leq b_n$ and $a_n \geq b_n$ hold simultaneously. Sometimes we abbreviate "with probability" as "w.p." and "with respect to" as "w.r.t.". For any $N \in \mathbb{N}_+$, [N] and 1 : N are defined to be $\{1, \ldots, N\}$. $\mathbb P$ and $\mathbb E$ are the probability measure and expectation taken over all randomness. We use $C, C', C'', C_1, C_2, C_3, c, c'$ to represent universal constants that could change from place to place.

The rest of this paper is organized as follows. In Section [2,](#page-8-0) we first proposed a penalized ERM algorithm for linear model [\(1\)](#page-1-0), established the upper bound of estimation error for global minimizers of the ERM, studied properties of the local minimizers, and discussed implementation details. Subsequently, we introduced a novel spectral method that is more computationally efficient and achieves sharper estimation error upper bounds than the penalized ERM when there is no contamination. We concluded Section [2](#page-8-0) by a lower bound analysis for the representation MTL problem. In Section [3,](#page-21-0) we extended the scope of our analysis from the linear model [\(1\)](#page-1-0) to generalized linear models (GLMs) and non-linear regression models. In Section [4,](#page-23-0) we proposed a thresholding algorithm to estimate the intrinsic dimension r and adapt our penalized ERM algorithm and the spectral method to the case where r is unknown. We conducted extensive simulation studies and analyzed a realworld dataset to demonstrate our theoretical findings in Section [5.](#page-25-0) Finally, we summarized our contributions and outlined a few potential avenues for future research in Section [6.](#page-33-0)

Due to space constraints, certain results are deferred to the appendix. In Section [A](#page-42-0) of the appendix, we extended the analysis of the spectral method to the general regime described in Section [1.2.2.](#page-3-2) In Section [B](#page-43-0) of the appendix, we studied how to transfer the knowledge to an unknown task, i.e., under the setting of transfer learning (TL) or learning-to-learn. We proposed an algorithm that leverages outputs from MTL algorithms to adapt to a new target task, presenting corresponding upper and lower bounds. All the proofs are also provided in the appendix.

2. Multi-task Learning with Similar Representations

2.1 Problem Set-up

Let us describe the problem setting introduced in Section [1.1](#page-0-0) in more detail. Suppose there are T tasks, and we have collected sample $\{\boldsymbol{x}_i^{(t)}\}$ $\binom{t}{i}, y_{i}^{(t)}$ $\{e^{(t)}\}_{i=1}^n$ from the t-th task, where $\boldsymbol{x}^{(t)}_i \in \mathbb{R}^p$, $y_i^{(t)} \in \mathbb{R}$, and $t \in [T]$. There exists an unknown subset $S \subseteq [T]$, such that for all $t \in S$,

$$
y_i^{(t)} = (\boldsymbol{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*} + \epsilon_i^{(t)}, \quad i = 1:n,
$$
\n(3)

where $\boldsymbol{\beta}^{(t)*} = \boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}, \ \boldsymbol{A}^{(t)*} \in \mathcal{O}^{p \times r} = \{ \boldsymbol{A} \in \mathbb{R}^{p \times r} : \boldsymbol{A}^\top \boldsymbol{A} = \boldsymbol{I}_r \}, \ \text{low-dimensional}$ parameter $\boldsymbol{\theta}^{(t)*} \in \mathbb{R}^r$, $r \leq p$, and $\{\epsilon_i^{(t)}\}$ $\binom{t}{i}_{i=1}^n$ are i.i.d. zero-mean sub-Gaussian variables independent of $\{\boldsymbol{x}_i^{(t)}\}$ $\binom{t}{i}_{i=1}^n$ ^{[c](#page-8-1)}. Throughout this and the next sections, we assume the intrinsic dimension r is known. The case that r is unknown will be addressed in Section [4.](#page-23-0)

Here the T tasks are divided into two groups, S and S^c . The tasks in S have "similar" representations (similarity to be defined in the following), while the tasks in S^c can be understood as *outlier* tasks or *contaminated* tasks with an arbitrary distribution. Our goal is two folds:

- 1. Improve the learning performance simultaneously on the tasks in S , when they share "similar" representations and the proportion of outlier tasks in S^c among all T tasks is small;
- 2. Maintain the single-task learning performance when the "similarity" between tasks in S is low.

It should be emphasized that if we allow the outlier tasks in S^c to be arbitrarily distributed, no guarantee can be obtained for these tasks in the worst case. However, as we will discuss later, if these tasks still follow linear models [\(3\)](#page-8-2) (without a low-dimensional representation), then the single-task linear regression estimation rate can be achieved for $\{\beta^{(t)*}\}_{t\in S^c}$.

We also want to point out that the set S is unknown. We will show that our penalized ERM algorithm and spectral method can perform well across all potential sets $S \subseteq [T]$ under certain conditions. This flexibility is crucial from both perspectives of outlier tasks and adversarial attacks. From the perspective of outlier tasks, we expect an algorithm to succeed for all possible outlier task index sets S^c as long as $|S^c|/T$ is small. In other words, the algorithm should not only work for a specific S^c and it should not rely on the indices of tasks (Otherwise, we can always drop the data from tasks in S^c to avoid the impact of outliers). From the perspective of adversarial attacks, the attacker can choose to corrupt the data from any task, which usually happens after the release of the machine learning system. Therefore, a robust learner should achieve ideal performance for all possible sets S. See Figure [3](#page-9-0) for an illustration of these two points of view.

c. This is assumed for simplicity and can be relaxed. In fact, it suffices to require $\{\epsilon_i^{(t)} | \mathbf{x}_i^{(t)} = \mathbf{x}_i\}_{i=1}^n$ to be independent zero-mean sub-Gaussian variables for almost surely $\{x_i\}_{i=1}^n$ w.r.t. the product probability measure induced by the distribution of $x_i^{(t)}$'s.

Figure 3: An illustration for two perspectives of viewing model [\(3\)](#page-8-2). In both cases, $S^c = \{4\}$.

The same setting when $\mathbf{A}^{(t)*}$'s are the same and $S = [T]$ (i.e. no outlier tasks) has been studied in [Du et al.](#page-36-2) [\(2020\)](#page-36-2), where they argued that when r is much smaller than p , a better estimation error rate of $\beta^{(t)*}$ can be achieved compared to the single-task learning. Our framework is more general and closer to the practical situation because it is difficult for all tasks to be embedded in precisely the same subspace as the number of tasks T grows [\(Rostami et al.,](#page-38-0) [2022\)](#page-38-0), and the prevalence of outlier tasks is common [\(Zhang and Yang,](#page-40-0) [2018\)](#page-40-0).

To mathematically quantify the similarity between representations ${A^{(t)*}}_{t \in S}$, we consider the maximum principal angle between subspaces spanned by the columns of these representation matrices. More specifically, we assume that there exists $h \in [0,1]$ such that

$$
\min_{\overline{\mathbf{A}} \in \mathcal{O}^{p \times r}} \max_{t \in S} \|\mathbf{A}^{(t)*}(\mathbf{A}^{(t)*})^{\top} - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}\|_2 \leq h^{d}
$$

A small h means the representations are more similar. The case when $h = 0$ reduces to the setting of the same representations in literature [\(Du et al.,](#page-36-2) [2020;](#page-36-2) [Tripuraneni et al.,](#page-39-2) [2021\)](#page-39-2). In the literature, $\|\boldsymbol{A}^{(t)*}(\boldsymbol{A}^{(t)*})^\top - \overline{\boldsymbol{A}}(\overline{\boldsymbol{A}})^\top \|_2$ is often referred to as the *maximum principal* angle between column spaces of $A^{(t)*}$ and \overline{A} . This concept and its variations have been widely used to measure the difference between subspaces in perturbation theory (e.g., [Wedin,](#page-39-7) [1972;](#page-39-7) [Cai et al.,](#page-35-9) [2013;](#page-37-8) [Kato,](#page-37-8) 2013; [Yu et al.,](#page-40-5) [2015;](#page-40-5) [Chen et al.,](#page-35-7) [2021b\)](#page-35-7). The case when $r = 1$ (where all representations are $p \times 1$ vectors and $\theta^{(t)*}$'s are scalars) reduces to the setting of [Gu et al.](#page-36-6) [\(2022\)](#page-36-6). In this case, the principle angle between subspaces becomes the angle between regression cofficient vectors.

We now make some assumptions. Without loss of generality, suppose $x^{(t)}$ is meanzero. Denote the covariance matrix $\Sigma^{(t)} = \mathbb{E}[\mathbf{x}^{(t)}(\mathbf{x}^{(t)})^{\top}]$ and the joint distribution of

d. Note that the LHS of the following inequality is always less than or equal to 1, becasue $A^{(t)*}$ and \overline{A} are orthonormal matrices. Here the minimum is achievable because of Weierstrass Theorem. Note that the objective function is continuous (over \overline{A}) under $\|\cdot\|_2$ norm and the space $(\mathcal{O}^{p\times r}, \|\cdot\|_2)$ is compact.

 $\{\{\boldsymbol{x}^{(t)}_i$ $\displaystyle{ _{i}^{(t)},y_{i}^{(t)}}$ $\{e^{(t)}\}_{t=1}^n\}_{t\in S^c}$ as \mathbb{Q}_{S^c} . For the convenience of description, define a coefficient matrix $\mathbf{B}_{S}^{*} \in \mathbb{R}^{p \times |\tilde{S}|}$, each column of which is a coefficient vector in $\{\beta^{(t)*}\}_{t \in S}$. Denote the ℓ_2 norm $\zeta^{(t)} = \|\boldsymbol{\theta}^{(t)*}\|_2 = \|\boldsymbol{\beta}^{(t)*}\|_2$, the average ℓ_2 -norm $\bar{\zeta} = \sqrt{|S|^{-1} \sum_{t \in S} (\zeta^{(t)})^2}$, and assume $\min_{t \in S} \zeta^{(t)} \gtrsim \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$. ^{[e](#page-10-0)} $\zeta^{(t)}$ can be viewed as the signal strength of the *t*-th task. In all the existing literature, it is assumed that $\zeta^{(t)} \approx 1$ for most of $t \in S = [T]$ and the impact of $\zeta^{(t)}$ on the model performance is ignored. But we allow $\zeta^{(t)}$'s to vary across tasks and we will see later how the performance of representation MTL on each task depends on $\zeta^{(t)}$.

Assumption 1 For any $t \in S$, $x^{(t)}$ is sub-Gaussian in the sense that for any $u \in \mathbb{R}^p$ and $\lambda \in \mathbb{R}, \ \mathbb{E}[e^{\lambda u^\top x^{(t)}}] \leq e^{C\lambda^2 \|u\|_2^2}$ with some constant $C > 0$. And there exist constants c, C such that $0 < c \leq \lambda_{\min}(\mathbf{\Sigma}^{(t)}) \leq \lambda_{\max}(\mathbf{\Sigma}^{(t)}) \leq C < \infty$, for all $t \in S$.

Assumption 2 There exists a constant $c > 0$ such that $\sigma_r(\mathbf{B}_S^*/\sqrt{|S|}) \geq \frac{c}{\sqrt{r}}\bar{\zeta}$, where each column of \mathbf{B}_{S}^{*} is a coefficient vector in $\{\boldsymbol{\beta}^{(t)*}\}_{t\in S}$.

Assumption 3 $n \geq C(p + \log T)$ with a sufficiently large constant $C > 0$.

Assumptions [1](#page-10-1) and [3](#page-10-2) are standard conditions in literature [\(Du et al.,](#page-36-2) [2020;](#page-36-2) [Duan and](#page-36-4) [Wang,](#page-36-4) [2022\)](#page-36-4). Assumption [2](#page-10-3) is often called the task diversity condition. When $A^{(t)*}$'s are the same, $\sigma_r(\mathbf{B}_{S}^*/\sqrt{|S|}) = \sigma_r(\mathbf{\Theta}_{S}^*/\sqrt{|S|})$, where each column of $\mathbf{\Theta}_{S}^*$ is a coefficient vector in $\{\theta^{(t)*}\}_{t\in S}$, which means that the low-dimensional task-specific parameters $\theta^{(t)*}$'s are diverse. Such a task diversity condition has been adopted in other related literature [\(Du et al.,](#page-36-2) [2020;](#page-36-2) [Chua et al.,](#page-35-6) [2021;](#page-39-2) [Tripuraneni et al.,](#page-39-2) 2021; [Duchi et al.,](#page-36-3) [2022\)](#page-36-3)^{[f](#page-10-4)}, to obtain a parametric rate which is faster than the rate without this condition [\(Maurer et al.,](#page-38-4) [2016\)](#page-38-4). The benefit of this condition is intuitive to understand, because a full exploration of all directions in the subspace is necessary to learn representations well, which is the key in representation MTL.

It is important to note that the presented assumptions are imposed on both the set of non-outlier tasks S and the model parameters. These assumptions are made from the perspective of outlier tasks. However, when considering adversarial attacks, we can replace S with the set of all tasks $[T]$ and assume that each task follows the linear model (3) . An attacker can then adversarially select a subset $S^c \subseteq [T]$ and distort the data distribution for these tasks. For simplicity, we do not distinguish between these two perspectives in the following parts of this paper.

In the following subsections, we will present two algorithms, study their properties, and derive a lower bound of the estimation error.

e. In general, we have the same results hold by defining $\zeta^{(t)} = \|\boldsymbol{\theta}^{(t)*}\|_2 \vee \sqrt{\frac{p + \log T}{n}}$. Here we define $\zeta^{(t)} = \|\boldsymbol{\theta}^{(t)*}\|_2$ and assume $\min_{t \in S} \zeta^{(t)} \gtrsim \sqrt{\frac{p + \log T}{n}}$ for presentation simplicity.

f. In [Duchi et al.](#page-36-3) [\(2022\)](#page-36-3) and [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2), the lower bound of $\sigma_r(\mathbf{B}_S^*/\sqrt{|S|})$ is defined as a parameter and appears in the estimation error. Here we follow [Du et al.](#page-36-2) [\(2020\)](#page-36-2) and impose an explicit bound on it to obtain a cleaner result, but our analysis can carry over to the analysis where the lower bound of $\sigma_r(\mathbf{B}_S^*/\sqrt{|S|})$ is denoted as a parameter.

2.2 The First Algorithm: Penalized ERM

2.2.1 Algorithm and Upper Bounds

In a special case of our setting, when $\mathbf{A}^{(t)*}$'s are the same, $S = [T]$, and $\zeta^{(t)} \approx 1$ for most of $t \in S$, [Du et al.](#page-36-2) [\(2020\)](#page-36-2) proposed an algorithm by combining the objective functions of all tasks and solving the optimization problem. A more general version accommodating various loss functions has been explored in [Tripuraneni et al.](#page-39-3) [\(2020\)](#page-39-3). Under our setting, where the representations are similar but not exactly the same, and with the inclusion of potential outlier tasks, the objective function needs to be properly adjusted. When h is large, learning $\{\beta^{(t)*}\}_{t\in S}$ by presuming similar representations may lead to a negative transfer effect. Considering these differences, we proposed a two-step learning approach in Algorithm [1](#page-11-0) that addresses these issues. Note that we will apply the same algorithm to some extended models in Section [3,](#page-21-0) so for description convenience, we introduce the algorithm with generic loss functions $f^{(t)}$ for the t-th task. For the linear model [\(3\)](#page-8-2), specifically, we set $f^{(t)}(\bm{\beta})=\frac{1}{2n}\|\bm{Y}^{(t)}-\bm{X}^{(t)}\bm{\beta}\|_2=\frac{1}{2n}$ $\frac{1}{2n}\sum_{i=1}^n[y_i^{(t)}-(\pmb{x}_i^{(t)}$ (i) ^T β ² for $\beta \in \mathbb{R}^p$, where $\boldsymbol{X}^{(t)} \in \mathbb{R}^{n \times p}$ and $Y^{(t)} \in \mathbb{R}^n$ are corresponding matrix representations of the data from the t-th task.

In Algorithm [1,](#page-11-0) Step 1 aims to learn all tasks by aggregating the data, where the penalty $||\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top} - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}||_2$ is added to force the subspaces represented by $\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^T$ to be similar. This penalty form is motivated by [Duan and Wang](#page-36-4) [\(2022\)](#page-36-4), where a similar penalty is applied on regression coefficient vectors. Note that when $||A^{(t)}(A^{(t)})^\top - \overline{A}(\overline{A})^\top||_2 = 0$, it can be shown that $A^{(t)}$ and \overline{A} are identical up to a rotation. And this distance is equivalent to other subspace distances like sin-Θ distance or the distance between two matrices up to a rotation [\(Chen et al.,](#page-35-7) [2021b\)](#page-35-7). Step 2 uses data from each task to make potential corrections to prevent negative transfer, which is often referred to as biased regularization in the literature (Schölkopf et al., [2001;](#page-39-8) [Kuzborskij and Orabona,](#page-37-9) [2013,](#page-37-9) [2017\)](#page-37-10). Such two-step methods are widely used in the distance-based MTL and TL literature to alleviate the adverse effect of negative transfer [\(Bastani,](#page-34-2) [2021;](#page-34-2) [Li et al.,](#page-37-5) [2022b;](#page-37-5) [Lin and Reimherr,](#page-37-7) [2022;](#page-37-7) [Tian and Feng,](#page-39-5) [2022\)](#page-39-5). It is important to point out that $A^{(t)*}$'s and $\theta^{(t)*}$'s are not uniquely identifiable in model [\(3\)](#page-8-2), which does not pose an issue because our focus is on estimating $\boldsymbol{\beta}^{(t)\ast}$'s.

Next, we proceed to present the upper bound on estimation errors of $\{\beta^{(t)*}\}_{t\in S}$ incurred by Algorithm [1.](#page-11-0)

Theorem [1](#page-10-1) (Upper bound for MTL) Suppose Assumptions 1[-3](#page-10-2) hold with a subset $S \subseteq$ $[T]$ satisfying $\epsilon = \frac{S^c}{T} \leq c r^{-3/2} \frac{\bar{\zeta}^2}{\max_{t \in S} (\zeta^{(t)})^2}$, where $c > 0$ is a small constant. By setting $\lambda =$ $C \frac{\max_{t \in S} (\zeta^{(t)})^2}{\min_{x \in \zeta(t)}}$ $\min_{t \in S} \zeta$ $\frac{\max_{t \in S(S)}(t)}{(t)} \sqrt{r(p + \log T)}$ and $\gamma = C' \sqrt{p + \log T}$ with sufficiently large positive constants C and C', for an arbitrary distribution \mathbb{Q}_{S^c} of $\{\{\boldsymbol{x}_i^{(t)}\}$ $\binom{t}{i},y_i^{(t)}$ $\{a_i^{(t)}\}_{i=1}^n\}_{t\in S^c}$, w.p. at least 1 – $e^{-C''(r+\log T)}$, we have

$$
\begin{aligned} \|\widehat{\beta}^{(t)}-\beta^{(t)*}\|_2\lesssim & \left\{\frac{\zeta^{(t)}}{\bar{\zeta}}r\sqrt{\frac{p}{nT}}+\left(\frac{\zeta^{(t)}\sqrt{r}}{\bar{\zeta}}\vee 1\right)\sqrt{\frac{r+\log T}{n}}+\zeta^{(t)}\sqrt{r}h\right.\\ & \left.+\frac{\zeta^{(t)}\max_{t\in S}(\zeta^{(t)})^2}{\bar{\zeta}^2\min_{t\in S}\zeta^{(t)}}r^{3/2}\sqrt{\frac{p+\log T}{n}}\epsilon\right\}\wedge\sqrt{\frac{p+\log T}{n}},\quad\forall t\in S. \end{aligned}
$$

Furthermore, if the data from tasks in S^c satisfies the linear model [\(3\)](#page-8-2) (without any latent structure assumption) and Assumption [1,](#page-10-1) then w.p. at least $1 - e^{-C'(p + \log T)}$, we also have

$$
\max_{t \in S^c} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \lesssim \sqrt{\frac{p + \log T}{n}}.
$$

In Theorem [1,](#page-12-0) the upper bound of $\|\hat{\beta}^{(t)} - \beta^{(t)*}\|_2$ for $t \in S$ is the minimum of two terms, where the first term corresponds to the rate obtained through data aggregation and the second term corresponds to the single task rate. This result shows that our algorithm is automatically adaptive to the optimal situation, whether or not aggregating data across tasks is beneficial. Furthermore, it demonstrates that Algorithm [1](#page-11-0) is robust to a small fraction of outlier tasks, in the sense that representation MTL remains beneficial when the outlier proportion ϵ is small.

Analyzing the components of the upper bound of $\|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2$ for $t \in S$ further elucidates their interpretability. The term $\frac{\zeta^{(t)}}{\overline{\zeta}}$ $\frac{d(t)}{\zeta}r\sqrt{\frac{p}{nT}}+\zeta^{(t)}\sqrt{r}h$ arises from learning similar representations, $\left(\frac{\zeta^{(t)}\sqrt{r}}{\zeta} \vee 1\right)\sqrt{\frac{r+\log T}{n}}$ $\frac{\log T}{n}$ is due to learning the representations and task-specific parameters, $\frac{\zeta^{(t)} \max_{t \in S} (\zeta^{(t)})^2}{\bar{z}^2 \min_{t \in \zeta(t)}}$ $\frac{f(t)}{\bar{\zeta}^2\min_{t\in S}(\zeta^{(t)})^2}r^{3/2}\sqrt{\frac{p+\log T}{n}}$ $\frac{\log T}{n}$ *e* accounts for outlier tasks, and $\sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$ is the error rate of single-task learning. Note that the term $\zeta^{(t)}\sqrt{r}h$ does not explicitly depend on n (although $\zeta^{(t)}$, r, h might depend on n). At the first glance, this might look stringent, but the lower bound in Section 2.4 contains a similar term that does not involve n explicitly. This phenomenon was also observed in many distance-based multi-task and transfer learning studies, such as [Li et al.](#page-37-5) [\(2022b](#page-37-5)[,a\)](#page-37-6); [Duan and Wang](#page-36-4) [\(2022\)](#page-36-4); [Lin and Reimherr](#page-37-7) [\(2022\)](#page-37-7); [Tian](#page-39-5) [and Feng](#page-39-5) [\(2022\)](#page-39-5).

To understand when Algorithm [1](#page-11-0) improves upon single-task learning, consider the scenario $\zeta^{(t)} \approx 1$ for all $t \in S$, where the upper bound simplifies to $\left(r\sqrt{\frac{p}{nT}} + \right)$ $\sqrt{r} \sqrt{\frac{r + \log T}{n}} +$ $\sqrt{r}h + r^{3/2}\sqrt{\frac{p+\log T}{n}}$ $\frac{\log T}{n} \epsilon \big) \wedge \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$. When $T \gg r^2$ (many tasks), $h \ll \sqrt{\frac{p + \log T}{nr}}$ $rac{\log T}{nr}$ (similar) representations), $p \gg r(r \vee \log T)$ (low intrinsic dimension), and $\epsilon \ll r^{-3/2}$ (a small fraction of outlier tasks), the rate is faster than the single-task error rate $\sqrt{\frac{p+\log T}{n}}$ $\frac{\log T}{n}$.

Furthermore, contrary to the results in existing literature where $\zeta^{(t)} \approx 1$ for most of $t \in S$ [\(Du et al.,](#page-36-2) [2020;](#page-36-2) [Chua et al.,](#page-35-6) [2021;](#page-39-2) [Tripuraneni et al.,](#page-39-2) 2021; [Thekumparampil et al.,](#page-39-4) [2021;](#page-39-4) [Duan and Wang,](#page-36-4) [2022\)](#page-36-4), our results indiciate that the performance of representation MTL on each task depends critically on the norm $\zeta^{(t)} = ||\boldsymbol{\theta}^{(t)*}||_2$. This insight suggests that tasks characterized by smaller coefficients may benefit more from representation MTL. We will verify this phenomenon through numerical experiments in Section [5.1.4.](#page-30-0)

As pointed out in Section [1.2,](#page-2-0) our setting reduces to the setting in [Gu et al.](#page-36-6) [\(2022\)](#page-36-6) when $r = 1$, where they considered a ridge regression by penalizing the angle between different regression coefficients. The main idea of our Algorithm [1](#page-11-0) resembles their approach, but they only considered the case when the angles between different regression coefficients are in $[0, \pi/2]$. Our result shows that representation MTL helps as long as the subspaces spanned by each coefficient (i.e., the straight line) are similar, where the angles can be either close to 0 or π . Moreover, we discovered that when $r = 1$, the estimators obtained from Step 1 of Algorithm [1](#page-11-0) already achieve a desired rate (despite of an inflation term in the single-task rate $\sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$), which means we may consider dropping Step 2. The detailed results are as follows.

Theorem 2 (No need for Step 2 when $r = 1$) Consider the case that $r = 1$. Under the same assumptions imposed in Theorem [1,](#page-12-0) for any $S \subseteq [T]$ satisfying $\epsilon = \frac{|S^c|}{T} \leq c$ with a small constant $c > 0$, and an arbitrary distribution \mathbb{Q}_{S^c} of $\{\{\boldsymbol{x}_i^{(t)}\}$ $\binom{t}{i}, y_{i}^{(t)}$ $\{e^{(t)}\}_{i=1}^{n}\}_{t\in S^{c}}, w.p.$ at least $1-e^{-C'(1+\log T)}$, we have

$$
\begin{split} \|\widehat{\beta}^{(t)}-\beta^{(t)*}\|_2 \lesssim & \Bigg\{\frac{\zeta^{(t)}}{\bar{\zeta}}\sqrt{\frac{p}{nT}} + \bigg(\frac{\zeta^{(t)}}{\bar{\zeta}} \vee 1\bigg)\sqrt{\frac{1+\log T}{n}} + \zeta^{(t)}\sqrt{r}h + \frac{\zeta^{(t)}\max_{t\in S}(\zeta^{(t)})^2}{\bar{\zeta}^2\min_{t\in S}\zeta^{(t)}}\sqrt{\frac{p+\log T}{n}}\epsilon \Bigg\} \\ & \qquad \qquad \wedge \Bigg\{\sqrt{\frac{p+\log T}{n}\frac{\max_{t\in S}(\zeta^{(t)})^2}{\min_{t\in S}(\zeta^{(t)})^2}} \Bigg\}, \quad \forall t\in S. \end{split}
$$

Furthermore, if we assume the data from tasks in S^c satisfies the linear model [\(3\)](#page-8-2) (without any latent structure assumption) and Assumption [1,](#page-10-1) then w.p. at least $1 - e^{-C'(p + \log T)}$, we also have

$$
\max_{t \in [T]} \|\widehat{\mathbf{A}}^{(t)}\widehat{\theta}^{(t)} - \beta^{(t)*}\|_2 \lesssim \sqrt{\frac{p + \log T}{n}}.
$$

Remark 3 Note that when $r = 1$, the task diversity condition in Assumption [2,](#page-10-3) i.e. $\sigma_1(\mathcal{B}_S^*/\sqrt{|S|}) \gtrsim \bar{\zeta}$, holds automatically because $\sigma_1(\mathcal{B}_S^*/\sqrt{|S|}) = \sqrt{|S|^{-1} \sum_{t \in S} (\zeta^{(t)})^2} = \bar{\zeta}$ by definition.

Before closing this section, it is worth noting that Assumption [3](#page-10-2) can be relaxed to $n \geq r + \log T$ for penalized ERM, provided that we are willing to forego the safe-net guarantee $\sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$. However, omitting the minimum with $\sqrt{\frac{p+\log T}{n}}$ $\frac{\log T}{n}$ in the estimation error can expose penalized ERM to negative transfer effects when either h or ϵ is large.

2.2.2 Discussions on Local Minimizers

It can be observed that the optimization problem in Step 1 of Algorithm [1](#page-11-0) is non-convex. Therefore, in addition to studying the global minimizers as previously examined, it is also

crucial to investigate the properties of local minimizers. Specifically, if only a local minimizer is found in Step 1, does Algorithm [1](#page-11-0) still achieve the same upper bound in Theorem [1?](#page-12-0)

Our first result below focuses on a special case where $S = [T]$ and $\lambda = +\infty$. In this scenario, Step 1 of Algorithm [1](#page-11-0) reduces to the ERM algorithm introduced in [Du et al.](#page-36-2) [\(2020\)](#page-36-2). The following theorem shows that even if we replace the global minimizer with a local minimizer in Step 1, Algorithm [1](#page-11-0) still achieves the same upper bound as stated in Theorem [1.](#page-12-0) In other words, there is no "bad" local minimizer in this case.

Theorem 4 (No bad local minimizer when $S = [T]$ and $\lambda = +\infty$) When $S = [T]$ and $\lambda = +\infty$, replacing the global minimizer in Step [1](#page-11-0) of Al[g](#page-14-0)orithm 1 with any local minimizer ^g delivers the same upper bound as in Theorem [1,](#page-12-0) i.e. w.p. at least $1 - e^{-C'(r + \log T)}$,

$$
\|\widehat{\beta}^{(t)}-\beta^{(t)*}\|_2 \lesssim \Bigg\{\frac{\zeta^{(t)}}{\bar{\zeta}} r\sqrt{\frac{p}{nT}}+\bigg(\frac{\zeta^{(t)}\sqrt{r}}{\bar{\zeta}}\vee 1\bigg)\sqrt{\frac{r+\log T}{n}}+\zeta^{(t)}\sqrt{r}h\Bigg\}\wedge\sqrt{\frac{p+\log T}{n}}, \ \forall t\in[T].
$$

[Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2) proves a similar result for the ERM when $S = [T]$, $h = 0$, and $\lambda = +\infty$. However, their result only holds for ERM with an additional regularization term, and the optimization must be conducted within a constrained set. In contrast, Theorem [4](#page-14-1) shows that there is no bad local minimizer across the *entire optimization landscape*. This result significantly supplements the existing literature in the case $S = [T]$.

Next, we present a more general result for the generic case $S \neq [T]$ with the same λ value as in Theorem [1.](#page-12-0) Unfortunately, similar to the result in [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2), in this general case, we can only establish that there is no bad local minimizer within a large regime instead of the entire landscape. This implies that we can replace the global minimizer in Step 1 by a local minimizer under some constraints to obtain the same upper bound.

Theorem 5 (No bad local minimizer in a large regime, for general $S \neq [T]$) Consider the same λ value taken in Theorem [1.](#page-12-0) Replacing the global minimizer in Step 1 of Algorithm [1](#page-11-0) with any local minimizer satisfying

$$
4(1+\sqrt{2})\max_{t\in S}\|\widehat{\boldsymbol{A}}^{(t)}(\widehat{\boldsymbol{A}}^{(t)})^\top - \overline{\boldsymbol{A}}(\overline{\boldsymbol{A}})^\top\|_2 + 4\frac{\epsilon}{1-\epsilon} < 1-\gamma;
$$

$$
\sqrt{2}\max_{t\in S}\left\{\frac{\sigma_{\max}(\boldsymbol{\Sigma}^{(t)})}{\sigma_{\min}(\boldsymbol{\Sigma}^{(t)})}\|\widehat{\boldsymbol{A}}^{(t)}(\widehat{\boldsymbol{A}}^{(t)})^\top - \boldsymbol{A}^{(t)*}(\boldsymbol{A}^{(t)*})^\top\|_2\right\} < 1-\gamma,
$$

with a universal constant $\gamma > 0$ delivers the same upper bound as in Theorem [1,](#page-12-0) i.e. w.p. at $least 1-e^{-C'(r+\log T)}$, for all $t \in S$,

$$
\|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \lesssim \left\{ \frac{\zeta^{(t)}}{\bar{\zeta}} r \sqrt{\frac{p}{nT}} + \left(\frac{\zeta^{(t)}\sqrt{r}}{\bar{\zeta}} \vee 1\right) \sqrt{\frac{r + \log T}{n}} + \zeta^{(t)}\sqrt{r}h + \frac{\zeta^{(t)}\max_{t \in S} (\zeta^{(t)})^2}{\bar{\zeta}^2 \min_{t \in S} \zeta^{(t)}} r^{3/2} \sqrt{\frac{p + \log T}{n}} \cdot \epsilon \right\} \wedge \sqrt{\frac{p + \log T}{n}}
$$

.

g. Here we say $\{\{\widehat{A}^{(t)}\}_{t=1}^T, \overline{A}, \{\widehat{\theta}^{(t)}\}_{t=1}^T\}$ is a local minimizer of function $G(\{A^{(t)}\}_{t=1}^T, \overline{A}, \{\theta^{(t)}\}_{t=1}^T) = \sum_{t=1}^T f^{(t)}(A^{(t)}\theta^{(t)}) + \frac{\lambda}{\sqrt{n}}\|A^{(t)}(A^{(t)})^T - \overline{A}(\overline{A})^T\|_2$, if there $\text{for any } \{\{\boldsymbol{A}_{\text{(t)}}^{(t)}\}_{t=1}^{T}, \boldsymbol{A}, \{\boldsymbol{\theta}^{(t)}\}_{t=1}^{T}\} \text{ with } \max_{t \in [T]} \min_{\boldsymbol{R} \in \mathcal{O}^{p \times r}} \left\{\|\widehat{\boldsymbol{A}}^{(t)} - \boldsymbol{A}^{(t)}\boldsymbol{R}\|_{2} + \|\widehat{\boldsymbol{\theta}}^{(t)} - \boldsymbol{R}^{\top}\boldsymbol{\theta}^{(t)}\|_{2}\right\} + \sum_{t=1}^{T} \left\{\|\widehat{\boldsymbol{A}}^{(t)} - \boldsymbol{R}^{\top}\boldsymbol{R$ $\min_{\mathbf{R}\in\mathcal{O}P^{\times r}}\|\overline{\mathbf{A}}-\mathbf{A}\mathbf{R}\|_2\leq \delta$, we must have $G(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^T,\overline{\mathbf{A}},\{\widehat{\theta}^{(t)}\}_{t=1}^T) < G(\{\mathbf{A}^{(t)}\}_{t=1}^T,\mathbf{A},\{\theta^{(t)}\}_{t=1}^T).$ The same definition is used in Theorem [5.](#page-14-2)

Theorem [5](#page-14-2) is similar to the results from other non-convex problems such as the EM algorithm for the mixture models [\(Balakrishnan et al.,](#page-34-5) [2017;](#page-34-5) [Cai et al.,](#page-35-10) [2019\)](#page-35-10), where a contraction basin exists within which the only local minimizer is the global one. This suggests that a good initialization can be used in the optimization problem of Step 1 to avoid bad local minima. We conjecture a similar result in Theorem [4](#page-14-1) holds in the general case as well, i.e. there is no bad local minimizer in the entire optimization lanscape. This conjecture remains a direction for future research.

The diagrams in Figure [4](#page-15-0) below summarize the results discussed in this subsection.

Figure 4: The optimization landscape of Algorithm [1](#page-11-0) proved in Theorems [4](#page-14-1) and [5.](#page-14-2) The big rectangle is the full optimization landscape.

2.2.3 Implementation

While we have provided desired upper bounds of estimation error with both local and global minimizers in Step 1 of Algorithm [1,](#page-11-0) solving the optimization problem there is challenging due to the constraint that representation matrices $\{\widehat{A}^{(t)}\}_{t=1}^T$ and \overline{A} must belong to the orthonormal space $\mathcal{O}^{p\times r} = \{ \boldsymbol{A} \in \mathbb{R}^{p\times r} : \boldsymbol{A}^\top \boldsymbol{A} = \boldsymbol{I}_r \}.$ In this subsection, we want to point out that this restriction is primarily for clarity in presentation. In practice, we can formulate the optimization problem in $\mathbb{R}^{p \times r}$ with a modified penalty term, achieving the same theoretical guarantees as with $\mathcal{O}^{p \times r}$.

Before discussing the practical formulation of the optimization problem in Step 1, we first introduce several new notations. For any non-zero matrix $\mathbf{A} \in \mathbb{R}^{p \times r}$ with $p \geq r$, we define the projection matrix onto its column space of A as $P_A = U U^{\top}$, where $U \in \mathcal{O}^{p \times r'}$ is the left singular matrix in the SVD of $\mathbf{A} = \mathbf{U}_{p \times r'} \mathbf{\Lambda}_{r' \times r'} \mathbf{V}_{r' \times r'}^{\top}$, $1 \leq r' \leq r$, and $\mathbf{\Lambda}_{r' \times r'}$ is diagonal with positive entries. When $\text{rank}(A) = r$, we have $P_A = A(A^{\top}A)^{-1}A^{\top}$. When A is zero, we define $P_A = 0_{p \times p}$. Note that while U in the SVD of A may not be unique, P_A is unique by the Hilbert projection theorem.

In practice, Step 1 of Algorithm [1,](#page-11-0) i.e.

$$
\{\widehat{\boldsymbol{A}}^{(t)}\}_{t=1}^T, \{\widehat{\boldsymbol{\theta}}^{(t)}\}_{t=1}^T, \widehat{\bar{\boldsymbol{A}}} \in \underset{\{\boldsymbol{A}^{(t)}\}_{t=1}^T, \overline{\boldsymbol{A}} \subseteq \mathcal{O}^{p \times r}, \{\boldsymbol{\theta}^{(t)}\}_{t=1}^T \subseteq \mathbb{R}^r} {\left\{\frac{1}{T} \sum_{t=1}^T \left[f^{(t)}(\boldsymbol{A}^{(t)} \boldsymbol{\theta}^{(t)}) + \frac{\lambda}{\sqrt{n}} \|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top - \overline{\boldsymbol{A}}(\overline{\boldsymbol{A}})^\top\|_2 \right] \right\}},
$$

can be replaced by

$$
\{\widehat{A}^{(t)}\}_{t=1}^{T},\{\widehat{\theta}^{(t)}\}_{t=1}^{T},\widehat{\overline{A}} \in \underset{\{A^{(t)}\}_{t=1}^{T},\overline{A} \subseteq \mathbb{R}^{p \times r},\{\theta^{(t)}\}_{t=1}^{T} \subseteq \mathbb{R}^{r}}{\arg\min} \left\{\frac{1}{T} \sum_{t=1}^{T} \left[f^{(t)}(A^{(t)}\theta^{(t)}) + \frac{\lambda}{\sqrt{n}} ||P_{A^{(t)}} - P_{\overline{A}}||_2 \right] \right\}.
$$
\n(4)

This replacement relaxes the constraint that $\{\widehat{A}^{(t)}\}_{t=1}^T$ and \overline{A} must belong to the orthonormal space $\mathcal{O}^{p\times r} = \{ \boldsymbol{A} \in \mathbb{R}^{p\times r} : \boldsymbol{A}^\top \boldsymbol{A} = \boldsymbol{I}_r \}$, allowing them to reside in $\mathbb{R}^{p\times r}$ with a modified penalty term. Intuitively, we can see why this relaxation preserves our previous theoretical guarantees. For any $A^{(t)} \in \mathbb{R}^{p \times r}$ and $\theta^{(t)} \in \mathbb{R}^r$, it can be shown that there exist $\widetilde{A}^{(t)} \in \mathcal{O}^{p \times r}$ and $\widetilde{\theta}^{(t)} \in \mathbb{R}^r$ such that $A^{(t)}\theta^{(t)} = \widetilde{A}^{(t)}\widetilde{\theta}^{(t)}$ and $P_{A^{(t)}} = P_{\widetilde{A}^{(t)}}$. Since we are interested in the product instead of $A^{(t)} \in \mathbb{R}^{p \times r}$ and $\theta^{(t)} \in \mathbb{R}^r$ individually, the replacement does not affect the estimation error of $\beta^{(t)*}$. The following theorem formally presents this result.

Theorem 6 We replace the Step [1](#page-11-0) of Algorithm 1 with the real-matrix optimization problem [\(4\)](#page-16-0), and replace $\|\widehat{A}^{(t)}(\widehat{A}^{(t)}) - \overline{A}(\overline{A})^{\top}\|_2$, $\|\widehat{A}^{(t)}(\widehat{A}^{(t)}) - A^{(t)*}(A^{(t)*})^{\top}\|_2$ in Theorem [5](#page-14-2) with $||P_{\hat{A}^{(t)}} - P_{\overline{A}}||_2$ and $||P_{\hat{A}^{(t)}} - P_{A^{(t)*}}||_2$. All theoretical results in Section [2.2.1](#page-11-1) and [2.2.2](#page-13-0) remain valid.

In all the numerical experiments, we used the automatic differentiation implemented in PyTorch [\(Paszke et al.,](#page-38-7) [2019\)](#page-38-7) along with the Adam optimizer [\(Kingma and Ba,](#page-37-11) [2015\)](#page-37-11) to solve the optimization problem [\(4\)](#page-16-0). Further implementation details will be discussed in Section [5.](#page-25-0)

2.3 The Second Algorithm: Spectral Method

In this subsection, we propose another algorithm for representation MTL, which is based on singular value decomposition (SVD), and we refer it as the spectral method.

The motivation of the spectral method arises from the special case $h = 0$ and $S = [T]$, implying $\mathbf{A}^{(t)*} \equiv \overline{\mathbf{A}}$ for all $t \in [T]$. In this scenario, $\mathbf{B}_{S}^{*} = (\beta^{(1)*}, \dots, \beta^{(T)*}) = \overline{\mathbf{A}}_{p \times r} \mathbf{\Theta}_{r \times T}^{*}$ with $\mathbf{\Theta}^* = (\theta^{(1)*}, \dots, \theta^{(T)*})$ is a rank-r matrix, and the column spaces of B_S^* and \overline{A} are the same. Thus, the left singular matrix in the SVD of \mathbf{B}_{S}^{*} can exactly recover $\overline{\mathbf{A}}$. In practice, we can use the single-task estimator to estimate each column of B_S^* and apply SVD to estimate \overline{A} . With the estimated \overline{A} , we can then perform a single-task regression to estimate $\boldsymbol{\theta}^{(t)*}$. When h is not necessarily zero and $S \neq [T]$, the same idea can still be applied. However, to avoid negative transfer when h is large, we can follow the SVD with a biased regularization step simiar to Algorithm [1.](#page-11-0) Additionally, we need to conduct SVD on a robust estimator of B_S^* to accomodate outlier tasks.

We use \prod_R to denote the projection operator to an ℓ_2 -ball centered at zero of radius R in \mathbb{R}^p , and use quantile($\{a_t\}_{t=1}^T$, $1-\bar{\epsilon}$) to denote the upper $(1-\bar{\epsilon})$ -quantile of a sequence ${a_t}_{t=1}^T \subseteq \mathbb{R}$, where $\bar{\epsilon} \in [0,1]$. We formalize the intuition into the spectral method in Algorithm [2.](#page-17-0)

Steps 1 and [2](#page-17-0) of Algorithm 2 construct a robust estimator of B^*_S by concatenating projected single-task estimators. The projection limits the impact of outlier tasks by setting the projection radius to a quantile of the ℓ_2 -norms of all single-task estimators. This projection technique, sometimes referred to as truncation or winsorization, is widely used in robust statistics (e.g., [Lugosi and Mendelson,](#page-38-8) [2021\)](#page-38-8) and differential privacy (e.g., [Dwork,](#page-36-11) [2006;](#page-36-11) [Dwork et al.,](#page-36-12) [2014\)](#page-36-12). To determine the quantile percentage, we need an upper bound $\bar{\epsilon}$ of ϵ . If we have prior knowledge of ϵ (but not S and S^c), we can set $\bar{\epsilon} = \epsilon$. Without this information, $\bar{\epsilon}$ can be chosen as a small constant, such as 0.05. In the simulations, we set $\bar{\epsilon}$ as the true ϵ value, and in the real-data study, we set $\bar{\epsilon} = 0.05$. Step 3 performs the SVD to estimate the central representation \boldsymbol{A} . Step 4 applies biased regularization as in Algorithm [1](#page-11-0) to prevent negative transfer. The spectral method avoids the complicated non-convex optimization problems present in the penalized ERM algorithm, making it easy to implement in practice. In fact, the spectral method can be solved in polynomial time with a small γ such that the objective function in Step 4 can be rewritten as a piecewise strongly convex function, consisting of a smooth part and an ℓ_2 -Lipschitz part. This is because both SVD and the minimization of such a piecewise strongly convex function can be solved in polynomial time. Besides its simplicity and efficiency, we can show that it achieves a better upper bound of estimation error for $\beta^{(t)*}$'s when $S = [T]$.

Theorem 7 (Upper bound for spectral method) Suppose Assumptions [1-](#page-10-1)[3](#page-10-2) hold with a subset $S \subseteq [T]$ satisfying $\epsilon = \frac{|S^c|}{T} \leq c r^{-1} \cdot \left(\frac{\bar{\zeta}}{\max_{t \in \bar{\zeta}} S}\right)$ $\left(\frac{\bar{\zeta}}{\max_{t \in S} \zeta^{(t)}}\right)^2$, where $c > 0$ is a small constant. By setting $\gamma = C'\sqrt{p + \log T}$ with a sufficiently large positive constant C', w.p. at least $1-e^{-C''(r+\log T)}$ and $\bar{\epsilon}$ satisfying $\epsilon \leq \bar{\epsilon} \leq \frac{c''}{r}$ $\frac{\bar{\zeta}''}{r} \cdot \left(\frac{\bar{\zeta}}{\max_{t \in \bar{\zeta}}}\right)$ $\int \frac{\bar{\zeta}}{\max_{t \in S} \zeta^{(t)}}$ with a sufficiently small positive

constant c'' , we [h](#page-18-0)ave $^{\rm h}$

$$
\begin{split} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \lesssim & \left\{ \frac{\zeta^{(t)}}{\bar{\zeta}} \sqrt{\frac{pr}{nT}} + \left(\frac{\zeta^{(t)}}{\bar{\zeta}} \vee 1 \right) \sqrt{\frac{r}{n}} + \sqrt{\frac{\log T}{n}} + \zeta^{(t)} h \cdot \left[\frac{\sigma_{\max}((\overline{A}^{\perp})^{\top} \mathbf{B}_{S}^{*})}{\sigma_{\min}((\overline{A}^{\perp})^{\top} \mathbf{B}_{S}^{*})} \wedge \sqrt{r} \right] \right. \\ &\left. + \frac{\zeta^{(t)}}{\bar{\zeta}} \max_{t \in S} \zeta^{(t)} \cdot \sqrt{r \bar{\epsilon}} \right\} \wedge \sqrt{\frac{p + \log T}{n}}, \qquad \qquad \forall t \in S, \end{split}
$$

where $B_S^* \in \mathbb{R}^{p\times |S|}$ is the coefficient matrix whose columns are $\{\boldsymbol{\beta}^{(t)*}\}_{t\in S}$, $\overline{A} \in \argmin_{\boldsymbol{A}\in \mathcal{O}^{p\times r}}$ $\max_{t\in[T]}\|A^{(t)*}(A^{(t)*})^\top - AA^\top \|_2$ is the central representation, and \overline{A}^\perp is orthogonal to \overline{A} in the sense that $\overline{A}^{\perp} \in \mathcal{O}^{p \times (p-r)}$ and $(\overline{A}^{\perp})^{\top} \overline{A} = \mathbf{0}_{(p-r) \times r}$.

Furthermore, if we assume the data from tasks in S^c satisfies the linear model [\(3\)](#page-8-2) (without any latent structure assumption) and Assumption [1,](#page-10-1) then w.p. at least $1 - e^{-C'(p + \log T)}$, we also have

$$
\max_{t\in[T]}\|\widehat{\boldsymbol{\beta}}^{(t)}-\boldsymbol{\beta}^{(t)*}\|_2\lesssim \sqrt{\frac{p+\log T}{n}}.
$$

We can similarly discuss when Algorithm [2](#page-17-0) improves single-task learning as in Theorem [1,](#page-12-0) which we do not repeat here. However, it is worth emphasizing that, like Algorithm [1,](#page-11-0) Algorithm [2](#page-17-0) is also adaptive to the unknown similarity structure and robust to a small fraction of outlier tasks. Comparing the rates in Theorems [1](#page-12-0) and [7,](#page-17-1) we can see that the estimation error rate of spectral method has a better dependence on r when $S = [T]$, although the last term related to outlier tasks is worse. In the next subsection, we will see that the spectral method is minimax optimal when $S = [T]$ and the condition number $\sigma_{\max}((\overline{\bm{A}}^{\perp})^{\top}\bm{B}_{S}^{*})/\sigma_{\min}((\overline{\bm{A}}^{\perp})^{\top}\bm{B}_{S}^{*})$ is bounded by a constant.

As mentioned, Algorithm [2](#page-17-0) requires an upper bound $\bar{\epsilon}$ of ϵ for determining the quantile percentage, and $\bar{\epsilon}$ also appears in the upper bound of estimation error. If $\bar{\epsilon} \approx \epsilon$, then we can replace $\bar{\epsilon}$ in the upper bound with the true ϵ .

Besides the intuition provided at the beginning of this subsection, Algorithm [2](#page-17-0) also has connections to average derivative estimation (ADE) and expected gradient outer product (EGOP) methods used to estimate the index space in single-index models and multi-index models (e.g., Härdle and Stoker, [1989;](#page-36-13) [Samarov,](#page-38-9) [1993;](#page-38-9) [Hristache et al.,](#page-37-12) [2001;](#page-37-12) [Yang et al.,](#page-40-6) [2017;](#page-40-6) [Yuan et al.,](#page-40-7) [2023\)](#page-40-7). For instance, in the multi-index model $\mathbb{E}[Y|X=x] := g(x) =$ $f((\boldsymbol{B}^*)^\top \boldsymbol{x}) \in \mathbb{R}$ with $\boldsymbol{B}^* \in \mathbb{R}^{p \times T}$, $\boldsymbol{x} \in \mathbb{R}^p$, g continuously differentiable and unknown, and $\mathbb{E}[\nabla g(X)(\nabla g(X))^{\top}]$ existing, we have:

$$
\mathbb{E}[\nabla g(X)(\nabla g(X))^{\top}] = (\boldsymbol{B}^*)^{\top} \mathbb{E}\{\nabla f((\boldsymbol{B}^*)^{\top} X)[\nabla f((\boldsymbol{B}^*)^{\top} X)]^{\top}\} \boldsymbol{B}^*
$$

.

Therefore, it is possible to use SVD or PCA on some estimator of $\mathbb{E}[\nabla g(X)(\nabla g(X))^{\top}]$ to recover the column space of B_S^* , and many estimators of the expected gradient outer product have been proposed in literature. In the context of representation MTL, when $S = [T]$, we can view g as a multi-variate function $x \mapsto ((\beta^{(1)*})^{\top} x, \ldots, (\beta^{(T)*})^{\top} x)^{\top}$ with f the identity function from \mathbb{R}^T to \mathbb{R}^T , implying that $\mathbb{E}[\nabla g(X)(\nabla g(X))^{\top}] = (\mathbf{B}_{S}^*)^{\top} \mathbf{B}_{S}^*$. Thus, our spectral

h. For convention, we define $0/0 = 0$, which is mainly for the case $(\overline{\mathbf{A}}^{\perp})^{\top} \mathbf{B}^* = \mathbf{0}_{(p-r)\times r}$.

method, which performs SVD on the estimated B_S^* , can be viewed as a multi-task variant of the EGOP framework.

When there are outlier tasks or contaminations, i.e., $S \neq [T]$, besides the simple projection technique, we can also borrow ideas from robust principal component analysis (PCA) literature $(e.g., Wright et al., 2009; Candès et al., 2011; Vidal et al., 2016)$ $(e.g., Wright et al., 2009; Candès et al., 2011; Vidal et al., 2016)$ $(e.g., Wright et al., 2009; Candès et al., 2011; Vidal et al., 2016)$ $(e.g., Wright et al., 2009; Candès et al., 2011; Vidal et al., 2016)$ $(e.g., Wright et al., 2009; Candès et al., 2011; Vidal et al., 2016)$ $(e.g., Wright et al., 2009; Candès et al., 2011; Vidal et al., 2016)$ $(e.g., Wright et al., 2009; Candès et al., 2011; Vidal et al., 2016)$ $(e.g., Wright et al., 2009; Candès et al., 2011; Vidal et al., 2016)$ $(e.g., Wright et al., 2009; Candès et al., 2011; Vidal et al., 2016)$ $(e.g., Wright et al., 2009; Candès et al., 2011; Vidal et al., 2016)$ to robustify the SVD procedure in Algorithm [2.](#page-17-0) For example, instead of conducting SVD on \vec{B} in Step 3 of Algorithm [2,](#page-17-0) we may conduct SVD on another $p \times T$ matrix \vec{L} which is the solution of the convex optimization [\(Liu et al.,](#page-37-13) [2012;](#page-37-13) [Xu et al.,](#page-39-11) [2012\)](#page-39-11)

$$
\min_{\mathbf{L},\mathbf{H}} \|\mathbf{L}\|_{*} + \lambda \|\mathbf{Z}\|_{2,1} \quad \text{s.t.} \quad \widehat{\mathbf{B}} = \mathbf{L} + \mathbf{Z},
$$

where $||L||_* = \sum_{j=1}^{p \wedge T} \sigma_j(L)$ is the nuclear norm, and $||Z||_{2,1} = \sum_{t=1}^{T} ||z_t||_2$ with z_t the t-th column of Z is the $L_{2,1}$ -norm. This approach might lead to a better performance when $S \neq [T]$. Given the extensive scope of the current paper, we leave the study of this approach for future research.

2.4 Lower Bound

In this subsection, we construct the lower bounds to explore the information-theoretic hardness of the representation MTL problem. Consider a collection of all subsets $S \subseteq [T]$ as

$$
\mathcal{S} = \{ S \subseteq [T] : |S^c|/T \le \epsilon \}.
$$

Given the subset S, define a coefficient matrix $B_S \in \mathbb{R}^{p \times |S|}$, each column of which corresponds to a coefficient vector in $\{\beta^{(t)}\}_{t\in S}$. Consider the parameter space for the coefficient vectors $\{\boldsymbol{\beta}^{(t)}\}_{t\in S}$ as

$$
\mathscr{B}(S,h) = \left\{ \{ \boldsymbol{\beta}^{(t)} \}_{t \in S} : \boldsymbol{\beta}^{(t)} = \boldsymbol{A}^{(t)} \boldsymbol{\theta}^{(t)} \text{ for all } t \in S, \{ \boldsymbol{A}^{(t)} \}_{t \in S} \subseteq \mathcal{O}^{p \times r}, \|\boldsymbol{\theta}^{(t)}\|_2 \le \zeta^{(t)},
$$

$$
\min_{\boldsymbol{A} \in \mathcal{O}^{p \times r}} \max_{t \in S} \|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top - \boldsymbol{\overline{A}} \boldsymbol{\overline{A}}^\top\|_2 \le h, \sigma_r \left(|S|^{-1/2} \boldsymbol{B}_S \right) \ge \frac{c}{\sqrt{r}} \sqrt{\frac{1}{|S|} \sum_{t \in S} \|\boldsymbol{\theta}^{(t)}\|_2^2} \right\}
$$

where c can be any fixed positive constant such that $\mathscr{B}(S, h) \neq \emptyset$. Given a set $S \subseteq [T]$, denote $\bar{\zeta} := \bar{\zeta}(S) = \sqrt{\frac{1}{|S|} \sum_{t \in S} (\zeta^{(t)})^2}$. We have the following lower bound.

Theorem 8 (Lower bound for MTL) Suppose $p \ge 2r$, $T \ge r^{1.01}$, $\min_{t \in [T]} \zeta^{(t)} \ge C$, and $\epsilon \leq c/r$ where C and c are some positive constants. We have the following lower bound hold:

$$
\inf_{\{\widehat{\beta}^{(t)}\}_{t=1}^T} \sup_{S \subseteq \mathcal{S}} \sup_{\{\beta^{(t)}\}_{t \in S} \in \mathcal{B}(S,h)} \mathbb{P}\Bigg(\bigcup_{t \in S} \Bigg\{\|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \gtrsim \left[\frac{\zeta^{(t)}}{\bar{\zeta}} \sqrt{\frac{pr}{nT}} + \left(\frac{\zeta^{(t)}}{\bar{\zeta}} \vee 1\right) \sqrt{\frac{r}{n}} + \frac{\log T}{n}\Bigg\}
$$

$$
+\zeta^{(t)}h + \frac{\zeta^{(t)}}{\overline{\zeta}}\frac{\epsilon r}{\sqrt{n}}\bigg] \wedge \sqrt{\frac{p + \log T}{n}}\bigg\}\bigg) \ge \frac{1}{10},
$$

where for any given $S \subseteq S$, $\mathbb{P} = \mathbb{P}_S \otimes \mathbb{Q}_{S^c}$, and \mathbb{P}_S , \mathbb{Q}_{S^c} are the probability measures on sample space of tasks in S, S^c , respectively. Furthermore, if tasks in S^c also follow the linear model (3) , then we have the following lower bound, where $\mathbb P$ is the probability measure on sample space of all tasks:

$$
\inf_{\{\widehat{\beta}^{(t)}\}_{t=1}^T}\sup_{S\subseteq\mathcal{S}}\sup_{\{\beta^{(t)}\}_{t\in S}\in\mathscr{B}(S,h)}\mathbb{P}\bigg(\max_{t\in[T]}\|\widehat{\beta}^{(t)}-\beta^{(t)*}\|_2\gtrsim \sqrt{\frac{p+\log T}{n}}\bigg)\geq \frac{1}{10}.
$$

Similar to the upper bound in Theorem [1,](#page-12-0) the lower bound of $\max_{t \in S} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2$ contains several terms reflecting the difficulty of learning different components. For example, $\zeta^{(t)}$ ζ¯ $\sqrt{\frac{pr}{nT}} + \zeta^{(t)}h$ arises from learning the similar representations in $S; (\frac{\zeta^{(t)}}{\zeta} \vee 1) \sqrt{\frac{r}{n}} + \sqrt{\frac{\log T}{n}}$ $\frac{g}{n}$ is due to learning the task-specific parameters; $\frac{\zeta^{(t)}}{\overline{\zeta}}$ $\frac{f(t)}{\zeta} \frac{\epsilon r}{\sqrt{n}}$ is caused by outlier tasks; and $\sqrt{\frac{p + \log T}{n}}$ \overline{n} is the single-task rate.

To our knowledge, this is the first lower bound for learning regression parameters in the context of representation MTL. [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2) and [Duchi et al.](#page-36-3) [\(2022\)](#page-36-3) derived lower bounds for the subspace recovery when $\zeta^{(t)} \approx 1$ for most of $t \in S = [T]$, with no outlier tasks ($\epsilon = 0$) and all tasks sharing the same representation ($h = 0$).

Comparing the upper bound for penalized ERM in Theorem [1](#page-12-0) with the lower bound, we see that the dependence on r in the upper bound is not optimal. This phenomenon has been noted in [Du et al.](#page-36-2) [\(2020\)](#page-36-2) and [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2) for both the ERM estimator and a method-of-moments estimator. The upper bounds of estimation errors for both estimators have sub-optimal dependence on r. They related this to a similar phenomenon observed in other works on linear regression models [\(Raskutti et al.,](#page-38-10) [2011\)](#page-38-10), where the upper bounds of estimation errors have sub-optimal dependence on eigenvalues of design matrices. In addition, the last term in our lower bound, $\frac{\zeta^{(t)}}{\bar{z}}$ $\frac{f(t)}{\bar{\zeta}} \frac{\epsilon r}{\sqrt{n}}$, does not depend on the full dimension p, whereas the counterpart in the upper bound does. A similar phenomenon has been noted in several papers (e.g. [Duan and Wang,](#page-36-4) [2022;](#page-36-4) [Tian et al.,](#page-39-6) [2022\)](#page-39-6). One might wonder whether it is possible to remove the dependence on p . Unfortunately, our ongoing work shows that this is impossible for the penalized ERM method like Algorithm [1](#page-11-0) with a large family of regularizers. As pointed out by [Tian et al.](#page-39-6) [\(2022\)](#page-39-6), estimators based on techniques in robust statistics like Tukey's depth function have been shown to achieve minimax rate under Huber's contamination model for location and covariance estimation [\(Chen et al.,](#page-35-12) 2018 , which might help improve the upper bound in our setting. In summary, when r is bounded, $\epsilon = 0$, and $\zeta^{(t)} \lesssim \bar{\zeta}$ for all $t \in S = [T]$, the penalized ERM is optimal. An adaptation of techniques in robust statistics to improve our estimation algorithm will be an interesting future research direction.

On the other hand, the upper bound for the spectral method in Theorem [7](#page-17-1) matches the lower bound when $\epsilon = 0$ and the condition number $\sigma_{\max}((\overline{\mathbf{A}}^{\perp})^{\top} \mathbf{B}_{S}^{*})/\sigma_{\min}((\overline{\mathbf{A}}^{\perp})^{\top} \mathbf{B}_{S}^{*})$ in Theorem [7](#page-17-1) is bounded by a constant. To our knowledge, when there is no contamination $(\epsilon = 0)$, no algorithm in the literature could achieve the optimal estimation error rate of $\beta^{(t)*}$'s before our work, even in the special case $h = 0$. This demonstrates the power of the spectral method in representation MTL. However, the term related to the outlier tasks

does not involve n and is not optimal. It remains unknown which algorithm can achieve the optimal estimation error rate when $\epsilon \neq 0$.

In Section [A](#page-42-0) of the appendix, we extend the theory of the spectral method to the most general regime described in Section [1.2.2.](#page-3-2)

3. Extensions to More General Models

In this section, we consider two extensions to the linear model [\(3\)](#page-8-2).

3.1 Generalized Linear Models

A generalization to generalized linear models (GLMs) from the linear model [\(3\)](#page-8-2) is as follows. Suppose that the conditional distribution of Y given $X = x$ for task t has density

$$
p(y_i^{(t)} = y | \boldsymbol{x}_i^{(t)} = \boldsymbol{x}) = \rho(y) \exp\left\{y \cdot \boldsymbol{x}^\top \boldsymbol{\beta}^{(t)*} - \psi(\boldsymbol{x}^\top \boldsymbol{\beta}^{(t)*})\right\}, \quad i = 1 : n,
$$
 (5)

for $t \in S$, w.r.t. some measure μ on a subset of R, where ψ is second-order continuously differentiable on \mathbb{R} , and ψ' is often called the inverse link function. More discussions on GLMs can be found in [McCullagh and Nelder](#page-38-11) [\(1989\)](#page-38-11).

Example 1 Some canonical examples of GLMs include:

- (i) Linear models: $\psi(u) = \frac{1}{2}u^2$, $\psi'(u) = u$, and μ is Lebesgue measure;
- (ii) Logistic regression models: $\psi(u) = u + \log(1 + e^{-u}), \ \psi'(u) = \frac{1}{1 + e^{-u}}, \text{ and } \mu \text{ is the }$ counting measure on $\{0,1\}$;
- (iii) Possion regression models: $\psi(u) = e^u$, $\psi'(u) = e^u$, and μ is the counting measure on $\mathbb{N} = \{0, 1, 2, \ldots\};$

We replace the linear model [\(3\)](#page-8-2) with the GLM [\(5\)](#page-21-1) and keep all the other settings the same as in Section [2.](#page-8-0) Moreover, we impose the following extra conditions for GLMs.

Assumption 4 ψ satisfies the following three conditions:

- (i) ψ is strictly convex;
- (ii) $\max_{t \in S} \max_{\|\mathbf{\Delta}\|_2 \leq 1} \psi''((\boldsymbol{\beta}^{(t)*})^\top \boldsymbol{x}^{(t)} + \boldsymbol{\Delta}^\top \boldsymbol{x}^{(t)}) \leq C$ a.s. with a constant $C > 0$;
- (iii) One of the following conditions hold:
	- (a) There exists a large constant $C' > 0$ such that $\min_{t \in S} \min_{|u| \le C' \zeta^{(t)}} \psi''(u) \ge C'$, where $C'' > 0$ is another universal constant.
	- (b) $\min_{t \in S} \min_{\|\Delta\|_2 \leq 1} \psi''((\beta^{(t)*})^\top x^{(t)} + \Delta^\top x^{(t)}) \geq C''''$ a.s., where $C''' > 0$ is a constant.

Assumption 5 $n \geq Cr(p + \log T)$ with a sufficiently large constant $C > 0$.

Assumption [4](#page-21-2) is commonly used in the non-asymptotic analysis of GLMs (e.g., see [Ne](#page-38-12)[gahban et al.,](#page-38-12) [2012;](#page-38-12) [Loh and Wainwright,](#page-37-14) [2015\)](#page-37-14). The sample size requirement in Assumption [5](#page-21-3) is more stringent than that for linear regression, serving as a technical condition needed for our proof. Specifically, the Hessian matrices of GLMs may not exhibit good spectral controls when evaluated far from the true parameter value. To ensure that our estimators avoid these problematic regions, we require the penalty term in Step 1 of the penalized ERM avoid these problematic regions, we require the penalty term in step [1](#page-11-0) of the penaltzed ERM method in Algorithm 1 (which is upper bounded by λ/\sqrt{n}) to remain reasonably small. For more details, see the proof of Lemma [52](#page-102-0) in Appendix. Note that for the spectral method in Algorithm [2,](#page-17-0) $n \geq p + \log T$ in Assumption [3](#page-10-2) is sufficient. For simplicity, we use the stronger condition in Assumption [4](#page-21-2) for both methods.

It is noteworthy that the function ψ in [\(5\)](#page-21-1) is allowed to be different across tasks. Here we assume ψ is the same for different tasks for simplicity. Additionally, we assume There we assume ψ is the same for underent tasks for simplicity. Additionally, we assume $\sqrt{r} \ge \zeta^{(t)} \ge 1$ $\sqrt{r} \ge \zeta^{(t)} \ge 1$ for all $t \in S$. For the GLM [\(5\)](#page-21-1), we apply Algorithms 1 and [2](#page-17-0) with $f^{(t)}(\boldsymbol{\beta}) = \frac{1}{n} \sum_{i=1}^{n} [-y_i^{(t)}]$ $\boldsymbol{x}_i^{(t)}\cdot(\boldsymbol{x}_i^{(t)}$ $\bm{e}_i^{(t)})^\top \bm{\beta} + \psi((\bm{x}_i^{(t)})$ $\binom{(t)}{i}^{\top}\beta$] for $\beta \in \mathbb{R}^p$. With these GLM assumptions in place, Algorithms [1](#page-11-0) and [2](#page-17-0) achieve the same upper bounds of estimation error as in linear regression models.

Theorem 9 (Upper bound for MTL under GLMs) Suppose Assumptions $4, 5$ $4, 5$ $4, 5$ and other conditions imposed in Theorems [1](#page-12-0) and γ hold, the same high-probability upper bounds in Theorems [1](#page-12-0) and [7](#page-17-1) hold for GLMs.

3.2 Non-linear Regression

In addition to GLMs, we can extend the linear model [\(3\)](#page-8-2) to a non-linear regression model as follows. Suppose

$$
y_i^{(t)} = g((\mathbf{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*}) + \epsilon_i^{(t)}, \quad i = 1 : n,
$$
\n(6)

for $t \in S$, where g is a monotone function with a continuous second-order derivative on \mathbb{R} and $\{\epsilon_i^{(t)}\}$ $\{t_i^{(t)}\}_{i=1}^n$ are i.i.d. zero-mean sub-Gaussian variables independent of $\{\boldsymbol{x}_i^{(t)}\}$ $\{e^{(t)}\}_{i=1}^{n}$. In literature, q is often referred to as the link function. More discussions for this model under a single-task learning setting can be found in [Yang et al.](#page-40-8) [\(2015\)](#page-40-8). Note that [Yang et al.](#page-40-8) [\(2015\)](#page-40-8) considered the case of a fixed design while we considered the random design case, which is more challenging. Hence, stronger conditions are necessary to guarantee the desired rate.

We replace the linear model (3) with the non-linear regression model (6) , and keep all the other settings the same as in Section [2.](#page-8-0) Furthermore, we impose the following assumptions for non-linear regression models. Recall the notation $\Sigma^{(t)} = \mathbb{E}[\boldsymbol{x}^{(t)}(\boldsymbol{x}^{(t)})^T]$ and denote $\mathbf{\Sigma}_{\epsilon}^{(t)} = \mathbb{E}[\boldsymbol{\epsilon}^{(t)}(\boldsymbol{\epsilon}^{(t)})^T].$

Assumption 6 $0 < C_1 \leq [g'(u)]^2 \leq C_2 < \infty$, for all $u \in \mathbb{R}$, and $||g''||_{\infty} \leq C_3 < \infty$, where $C_3 \leq cC_1^2 \cdot \min_{t \in S} \{\lambda_{\min}(\mathbf{\Sigma}^{(t)})/\lambda_{\max}(\mathbf{\Sigma}^{(t)})\} \min_{t \in S} \{\lambda_{\max}^{-1/2}(\mathbf{\Sigma}_{\epsilon}^{(t)})\}$ with a small constant c.

Assumption 7 $n \geq C[(p + \log T)^{3/2}] \vee [r(p + \log T)]$ with a sufficiently large constant C.

The sample size requirement is stronger than that for linear regression. The first term $(p + \log T)^{3/2}$ arises due to the heavy-tailed distributions appearing in the analysis. A similar requirement for the high-dimensional sparse non-linear regression can be found in [Yang et al.](#page-40-8) [\(2015\)](#page-40-8). The reason for the term $r(p + \log T)$ is the same as in Section [3.1](#page-21-4) for GLMs. More details can be found in the proof of Lemma [60](#page-112-0) in Appendix.

Similar to the case of GLMs, the link function q in (6) is also allowed to vary across tasks. Here we assume q is the same for different tasks for simplicity. We further assume $\sqrt{r} \gtrsim \zeta^{(t)} \gtrsim 1$ $\sqrt{r} \gtrsim \zeta^{(t)} \gtrsim 1$ for all $t \in S$. We apply Algorithms 1 and [2](#page-17-0) for the non-linear regression model [\(6\)](#page-22-0), by setting $f^{(t)}(\boldsymbol{\beta}) = \frac{1}{2n} \sum_{i=1}^{n} [y_i^{(t)} - g((\boldsymbol{x}_i^{(t)})$ $\mathbf{E}_{i}^{(t)}$ ^{$\lceil \mathbf{B}_{i} \rceil$ ² for $\mathbf{\beta} \in \mathbb{R}^{p}$. We have the same upper} bounds of estimation error for Algorithms [1](#page-11-0) and [2](#page-17-0) as in linear regression models.

Theorem 10 (Upper bound for MTL under non-linear regression models) Suppose Assumptions [6,](#page-22-1) [7](#page-22-2) and other conditions imposed in Theorems [1](#page-12-0) and [7](#page-17-1) hold, then the same high-probability upper bounds in Theorems [1](#page-12-0) and γ hold for non-linear regression models.

4. Adaptation to Unknown Intrinsic Dimension r

In Sections [2](#page-8-0) and [3,](#page-21-0) the intrinsic dimension r is assumed to be known as a priori. To our knowledge, this assumption is standard in almost all related theoretical literature on representation multi-task and transfer learning (e.g., [Ando et al.,](#page-34-1) [2005;](#page-34-1) [Maurer et al.,](#page-38-4) [2016;](#page-38-4) [Du et al.,](#page-36-2) [2020;](#page-36-2) [Thekumparampil et al.,](#page-39-4) [2021;](#page-39-4) [Tripuraneni et al.,](#page-39-2) [2021;](#page-39-2) [Chua et al.,](#page-35-6) [2021;](#page-35-6) [Collins et al.,](#page-35-2) [2021;](#page-35-2) [Deng et al.,](#page-35-3) [2022;](#page-35-3) [Duan and Wang,](#page-36-4) [2022;](#page-36-4) [Duchi et al.,](#page-36-3) [2022\)](#page-36-3), despite its potential unrealism in practice. In this section, we propose a simple yet effective algorithm to adapt the previous MTL algorithms to the case of an unknown r . Recall the notations $\zeta^{(t)} = \|\theta^{(t)*}\|_2$ and $\bar{\zeta} = \sqrt{|S|^{-1} \sum_{t \in S} (\zeta^{(t)})^2}$. Similar to the notations used in Algorithm [2,](#page-17-0) we use \prod_R to denote the projection operator to an ℓ_2 -ball centered at zero of radius R in \mathbb{R}^p , and use quantile $({a_t}_{t=1}^T, 1-\bar{\epsilon})$ to denote the upper $(1-\bar{\epsilon})$ -quantile of ${a_t}_{t=1}^T \subseteq \mathbb{R}$, where $\bar{\epsilon} \in [0, 1]$.

The algorithm is based on SVD, and the details are summarized in Algorithm [3.](#page-23-1)

column is $\prod_R(\widetilde{\boldsymbol{\beta}}^{(t)}),$ where $R = \texttt{quantile}(\{\|\widetilde{\boldsymbol{\beta}}^{(t)}\|_2\}_{t=1}^T, 1-\bar{\epsilon})$ 3 Step 3: (Thresholding) Set $\hat{r} = \max \left\{ r' \in [T] : \sigma_{r'}(\hat{B}) \right\}$ √ \overline{T}) $\geq T_1 \sqrt{\frac{p + \log T}{n}} + T_2 R$ √ $\overline{\overline{\epsilon}}$)

Algorithm [3](#page-23-1) leverages Assumption [2](#page-10-3) to determine an appropriate value for r. The underlying rationale is that when h is small, there tends to be a significant gap between the r-th largest singular value and the $(r + 1)$ -th largest singular value of $\sigma_r(B_S^*/\sqrt{T})$. For example, when $h = 0$, \mathbf{B}_{S}^{*} is a rank-r matrix, implying that $\sigma_{r'}(\mathbf{B}_{S}^{*}/\sqrt{T}) = 0$ for $r' \geq r + 1$. Therefore, thresholding on singular values of empirical version of B_S^* can be an effective strategy to estimate r. In Section [2.3,](#page-16-1) a similar approach based on SVD of the estimated B^* was used to develop the spectral method. In fact, Steps 1-[2](#page-17-0) of Algorithms 2 and [3](#page-23-1) are

the same, which construct a robust estimate of B_S^* through projected single-task estimators. Algorithm [3](#page-23-1) also shares a conceptual affinity with the thresholding method often used to determine the intrinsic dimension in principal component analysis [\(Onatski,](#page-38-13) [2010;](#page-38-13) [Fan et al.,](#page-36-14) [2021\)](#page-36-14).

Under almost identical assumptions imposed in previous sections, with proper choices of tuning parameters, Algorithm [3](#page-23-1) is shown to be consistent in estimating the true intrinsic dimension r, when the difference between representation matrices is small (i.e., h is small). As we will elaborate, this suffices to ensure the same upper bounds of estimation error for Algorithms [1](#page-11-0) and [2](#page-17-0) when r is unknown.

Theorem 11 (Consistency of the intrinsic dimension estimation) Suppose we know an $\bar{\epsilon}$ such that $\epsilon \leq \bar{\epsilon} \leq \frac{c}{r}$ $\frac{c}{r} \Big(\frac{\bar{\zeta}}{\max_{t \in \mathbb{R}}}$ $\int_{\max_{t \in S} \zeta^{(t)}}^{\overline{\zeta}} \int_{0}^{2} \text{ with a small constant } c > 0.$ Assume $\min_{t \in S} \zeta^{(t)}$. $h\left[\frac{\sigma_{\max}((\overline{\bm{A}}^{\perp})^{\top}\bm{B}^{*}_{S})}{(\sqrt{-1},\top,\mathbf{D}_{S})}\right]$ $\left[\frac{\sigma_{\max}((\overline{\bm{A}}^\perp)^\top \bm{B}^*_S)}{\sigma_{\min}((\overline{\bm{A}}^\perp)^\top \bm{B}^*_S)} \wedge \sqrt{r} \right] \leq c' \sqrt{\frac{p + \log T}{n}}$ $\mathcal{L}^{\text{domin}}(A \mid B_S)$ \rightarrow $\mathcal{L}^{\text{domin}}(A \mid B_S)$ $\frac{\log T}{n}$ with a small constant $c' > 0$, where $\mathbf{B}_{S}^{*} \in \mathbb{R}^{p \times |S|}$ is the $\bm{A}\bm{A}^\top\|_2$ is the central representation, and $\overline{\bm{A}}^\perp$ is orthogonal to $\overline{\bm{A}}$ in the sense that $\overline{\bm{A}}^\perp \in$ $\mathcal{O}^{p\times (p-r)}$ and $(\overline{\mathbf{A}}^{\perp})^{\top} \overline{\mathbf{A}} = \mathbf{0}_{(p-r)\times r}$. Further assume that:

- (i) For the linear model (3) , Assumptions [1,](#page-10-1) [2,](#page-10-3) and [5](#page-21-3) hold;
- (ii) For the GLM (5) , Assumptions [1,](#page-10-1) [2,](#page-10-3) [4,](#page-21-2) and [5](#page-21-3) hold;
- (iii) For the non-linear regression model (6) , Assumptions [1,](#page-10-1) [2,](#page-10-3) [6,](#page-22-1) [7](#page-22-2) hold;

Then there exist constants $T_1, T_2 > 0$ such that the output of Algorithm [3](#page-23-1) satisfies $\hat{r} = r \ w.p.$ at least $1 - e^{-C'(p + \log T)}$ with some constant $C' > 0$.

Remark 12 When $\min_{t \in S} \zeta^{(t)} \cdot h \left[\frac{\sigma_{\max}((\overline{A}^{\perp})^{\top} B_{\mathcal{S}}^{*})}{\sqrt{A}^{\perp} \setminus \mathbb{T}_{\max}} \right]$ $\frac{\sigma_{\max}((\overline{\bm{A}}^\perp)^\top \bm{B}^*_S)}{\sigma_{\min}((\overline{\bm{A}}^\perp)^\top \bm{B}^*_S)} \wedge \sqrt{r} \bigg] \gtrsim \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$ in the MTL problem, estimating r becomes unnecessary as the upper bounds of MTL estimation errors are dominated by the single-task rate $\sqrt{\frac{p+\log T}{n}}$ $\frac{\log T}{n}$ for both Algorithms [1](#page-11-0) and [2.](#page-17-0) In such cases, single-task learning is sufficient to achieve the minimax rate. Specifically, by the proofs of these upper bounds, the single-task rate is always guaranteed by biased regularization in Step 2 of Algorithm [1](#page-11-0) and Step 4 of Algorithm [2,](#page-17-0) regardless of the performance achieved in other steps.

According to Theorem [11](#page-24-0) and Remark [12,](#page-24-1) when r is unknown, we can first run Algorithm [3](#page-23-1) to obtain an estimate \hat{r} , then run Algorithms [1](#page-11-0) and [2](#page-17-0) with \hat{r} . All the previous results remain valid. Therefore, the entire learning procedure is **adaptive** to an unknown r .

Similar to Algorithm [2,](#page-17-0) Algorithm [3](#page-23-1) also requires $\bar{\epsilon}$ as an upper bound of the proportion of outlier tasks ϵ . If we have prior knowledge of ϵ (but not S and S^c), we can set $\bar{\epsilon} = \epsilon$. Without such information, a small constant such as 0.05 can be chosen for $\bar{\epsilon}$. Similar to Algorithm [2,](#page-17-0) we set $\bar{\epsilon}$ to the true ϵ value in simulations, while setting $\bar{\epsilon} = 0.05$ in the real-data study. We set tuning parameters $T_1 = 0.5$ and $T_2 = 0.25$ in the numerical experiments. Generally, T_1, T_2 can be chosen through cross-validation.

5. Numerical Experiments

To validate the theoretical insights discussed in previous sections, we conducted extensive simulations and one real-data study, and the results are presented in this section.

All the experiments were implemented in Python. For penalized ERM ("pERM", Algorithm [1\)](#page-11-0), we used the automatic differentiation implemented in PyTorch [\(Paszke et al.,](#page-38-7) [2019\)](#page-38-7) along with the Adam optimizer [\(Kingma and Ba,](#page-37-11) [2015\)](#page-37-11) to solve the optimization problem (4) in Step 1. We set the learning rate equal to 0.01 in torch.optim.Adam function and kept all the other parameter choices as in default. The Step 2 of pERM and Step 4 of the spectral method ("Spectral", Algorithm [2\)](#page-17-0) was also solved by the Adam optimizer with learning rate 0.01. Consistent with our theory, we set penalty parameters $\lambda = \sqrt{r(p + \log T)}$ and $\gamma = \sqrt{p + \log T}$ in pERM, and $\gamma = 0.5\sqrt{p + \log T}$ in the spectral method. As mentioned in Section [2.3,](#page-16-1) Spectral requires an upper bound $\bar{\epsilon}$ of the contamination proportion ϵ . We set $\bar{\epsilon} = \epsilon$ in simulations and $\bar{\epsilon} = 0.05$ in the real-data study. Besides pERM and Spectral, we included the following approaches as benchmarks.

- Empirical risk minimization ("ERM") in [Du et al.](#page-36-2) [\(2020\)](#page-36-2); [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2): The optimization was also solved by the Adam solver in PyTorch with learning rate 0.01;
- Method-of-moments ("MoM") in [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2);
- Adaptive representation learning ("AdaptRep") in [Chua et al.](#page-35-6) [\(2021\)](#page-35-6): We used the code included in the original paper and kept all the parameter settings as default.
- Adaptive and robust multi-task learning ("ARMUL") in [Duan and Wang](#page-36-4) [\(2022\)](#page-36-4): We used the code included in their paper (<https://github.com/kw2934/ARMUL>) and kept all the parameter settings as default. The tuning parameters were chosen using 5-fold cross-validation as default.
- Group Lasso ("GLasso") in [Yuan and Lin](#page-40-9) [\(2006\)](#page-40-9); [Lounici et al.](#page-37-2) [\(2009,](#page-37-2) [2011\)](#page-38-5): We grouped same coordinate of coefficients from different tasks and applied an $L_{2,1}$ -matrix penalty. The method was implemented in an R package RMTL [\(Cao et al.,](#page-35-13) [2019\)](#page-35-13), and we used the Python package rpy2 to call functions cv.MTL and MTL in the R package RMTL. The penalty parameter was chosen by a 5-fold cross-validation as default.
- Data pooling or pooled regression ("Pooled") in [Crammer et al.](#page-35-14) [\(2008\)](#page-35-14); [Ben-David](#page-34-4) [et al.](#page-34-4) [\(2010\)](#page-34-4): We fitted the linear regression and logistic regression models on the pooled data from all tasks. Both models were implemented in the Python module sklearn.linear models.
- Single-task regression (Single-task): In simulations and the real-data study, we run single-task linear regression and logistic regression on the data of each task, respectively.

In Section [5.1,](#page-26-0) we present the performance of different approaches under different simulation settings, such as different heterogeneity parameter h (Section [5.1.1\)](#page-26-1), different contamination proportion ϵ (Section [5.1.2\)](#page-27-0), and different number of tasks T (Section [5.1.3\)](#page-29-0). We also change the full dimension p , the intrinsic dimension r , and the per-task sample

size n from setting to setting. To verify the intuition we obtained in Section [2,](#page-8-0) where we mentioned that the performance of each task in ERM, pERM, and Spectral depends on the signal strength $\zeta^{(t)} = \|\theta^{(t)*}\|_2$, we conducted a simulation with different $\|\theta^{(t)*}\|_2$ values across tasks, and the results are presented in Section [5.1.4.](#page-30-0) Finally, in Section [5.1.5,](#page-31-0) we evaluate Algorithm [3](#page-23-1) for estimating r, compare the performance of pERM and Spectral with the estimated r and the true r, and demonstrate the effectiveness of Algorithm [3.](#page-23-1)

In Section [5.2,](#page-32-0) we compare the performance of different approaches on a real dataset. The code to reproduce the results are available at <https://github.com/ytstat/RL-MTL-TL>.

5.1 Simulations

5.1.1 SIMULATION WITH DIFFERENT HETEROGENEITY PARAMETER h

In this subsection, we investigate the linear model (1) with various values of h. We explore four distinct settings with different combinations of (n, p, r, T) :

(i)
$$
n = 100, p = 30, r = 5, T = 50;
$$

(ii) $n = 100, p = 50, r = 5, T = 50;$

(iii)
$$
n = 100, p = 80, r = 5, T = 50;
$$

(iv)
$$
n = 150, p = 80, r = 10, T = 50.
$$

In all settings, no outlier tasks are included, implying $S = [T]$ and $\epsilon = |S^c|/T = 0$. Given each (n, p, r, T) combination, we generated $\boldsymbol{x}_i^{(t)}$ $i^{(t)}_i$ i.i.d. from $N(\mathbf{0}_p, \mathbf{I}_p)$, $\epsilon_i^{(t)}$ $i^{(l)}$ i.i.d. from $N(0, 1)$, and a random $p \times r$ matrix C with i.i.d. standard normal entries. We defined \overline{A} as the first r columns of the left singular matrix of C, $\widetilde{A}^{(t)} = \overline{A} + a^{(t)}(I_{r \times r}, 0_{r \times (p-r)})^{\top}$, and $\mathbf{A}^{(t)*} = \widetilde{\mathbf{A}}^{(t)}[(\widetilde{\mathbf{A}}^{(t)})^{\top} \widetilde{\mathbf{A}}^{(t)}]^{-1}(\widetilde{\mathbf{A}}^{(t)})^{\top}$ for $t \in [T]$, where $a^{(t)}$'s are i.i.d. sampled from Unif([-h, h]). We generated each coordinate of $\theta^{(t)*} \in \mathbb{R}^r$ from Unif([-2, 2]) independently. We considered h from 0 to 0.8 in increments of 0.1 and replicated each setting 100 times.

Figure [5](#page-27-1) presents the simulation results. Across all settings, pooled regression and MoM perform worse than or comparably to single-task regression, with their performance deteriorating as h increases. AdaptRep exhibits slightly better performance than single-task regression initially but declines as h increases. GLasso performs similarly to single-task regression in settings (i), (ii), and (iv), while notably improving upon it in setting (iii). ERM significantly enhances single-task regression performance when h is small, but performs worse than single-task regression for large h . In contrast, ARMUL, pERM, and Spectral can improve the performance of single-task regression when h is small, and their performance will be comparable to single-task regression for large h . This demonstrates the adaptivity of ARMUL, pERM, and Spectral to the unknown similarity level h . An interesting phenomenon arises in setting (iii), where n is close to p . Here, ARMUL and Spectral perform worse than ERM and pERM for small h. This aligns with our discussion at the end of Section [2.2.1,](#page-11-1) emphasizing the adequacy of $n \geq r + \log T$ for pERM when h is small, while the spectral method and ARMUL require $n \gtrsim p + \log T$ and $n \gtrsim r^2(p + \log T)$, respectively.

Figure 5: Simulation with different heterogeneity parameters h: estimation error $\max_{t\in[T]} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2$ of different algorithms with different (n, p, r, T) settings. "max" error in S" in the y-axis stands for $\max_{t \in S} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2$ with $S = [T]$. Each point represents the average over 100 replications.

5.1.2 SIMULATION WITH DIFFERENT CONTAMINATION PROPORTION ϵ

In this subsection, we considered the linear model [\(1\)](#page-1-0) with varying contamination proportions $\epsilon = |S^c|/T$. We explored two settings with different values of (n, p, r, T) :

- (i) $n = 100, p = 50, r = 5, T = 100;$
- (ii) $n = 150, p = 80, r = 10, T = 100.$

For each replication in each setting, we randomly selected a subset of size $T(1 - \epsilon)$ from [T] without replacement to form S. We set $h = 0$ and generate tasks in S using the same mechanism as in Section $5.1.1$. The outlier tasks in S^c were generated by the linear model with $\boldsymbol{x}_i^{(t)}$ ^(t) i.i.d. from $N(\mathbf{0}_p, 2\mathbf{I}_p)$ and each coordinate of the coefficient $\boldsymbol{\beta}^{(t)*}$ was generated i.i.d. from Unif([-3,3]). We varied ϵ from 0 to 10% in increments of 2% and replicated each setting 100 times.

Figure 6: Simulation with different contamination proportions ϵ : within-S estimation error $\max_{t \in S} ||\widehat{\beta}^{(t)} - \beta^{(t)*}||_2$ and outlier estimation error $\max_{t \in S^c} ||\widehat{\beta}^{(t)} - \beta^{(t)*}||_2$ of different algorithms with different (n, p, r, T) settings. "max error in S" and "max error in S"" in the y-axis stand for $\max_{t \in S} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2$ and $\max_{t \in S^c} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2$, respectively. Each point represents the average over 100 replications.

The results are summarized in Figure [6,](#page-28-0) where we evaluated each method by both $\max_{t \in S} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2$ and $\max_{t \in S^c} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2$. As the outlier proportion ϵ increases, the performance of most algorithms deteriorates rapidly. Even with just 2% outlier tasks, all methods perform similarly to or worse than single-task regression on tasks in S , except for pERM and Spectral. This demonstrates the robustness of pERM and Spectral against outlier tasks. On the other hand, ARMUL does perform worse than single-task regression, even when ϵ is large. Since the outlier tasks were also generated from the linear model, our theory guarantees that pERM and Spectral can achieve the single-task performance on outlier tasks in S^c , which is indeed observed. In constrast, ARMUL performs much worse on outlier tasks in S^c compared to single-task regression.

5.1.3 Simulation with Different Number of Tasks T

In this subsection, we explore the impact of the number of tasks T on the performance of different methods in this subsection. We considered the linear model [\(1\)](#page-1-0) in two settings, with different values of (n, p, r, ϵ) :

- (i) $n = 100, p = 50, r = 5, \epsilon = 0$;
- (ii) $n = 100, p = 50, r = 5, \epsilon = 4\%$.

For each (n, p, r, ϵ) setting and T value, we set $h = 0$ and generated data in the same way as in Section [5.1.2.](#page-27-0) We increased T from 10 to 190 with in increments of 15. The performance of different methods on tasks in S is summarized in Figure [7.](#page-29-1)

Figure 7: Simulation with different number of tasks T: estimation error $\max_{t \in S} \|\hat{\beta}^{(t)} - \beta^{(t)*}\|_2$ of different algorithms with different (n, p, r, T) settings. "max error in S" in the y-axis stands for $\max_{t \in S} ||\widehat{\beta}^{(t)} - \beta^{(t)*}||_2$. Each point represents the average over 100 replications.

When there is no outlier task, ERM, pERM, ARMUL, and Spectral perform very similarly. As T increases, their estimation errors first decrease gradually then stabilize, aligning with the theoretical result. For example, when $\zeta^{(t)} = ||\boldsymbol{\theta}^{(t)*}||_2 \lesssim 1$ for all $t \in S = [T], p \gtrsim r^2$, and $h = \epsilon = 0$, Theorem [1](#page-12-0) implies that $\max_{t \in [T]} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \lesssim r \sqrt{\frac{p}{nT}} + r \sqrt{\frac{1}{n}}$ $\frac{1}{n}$, up to logarithmic factors with high probability. When T becomes large, the error is dominated by the second term $r\sqrt{\frac{1}{n}}$ $\frac{1}{n}$, which is independent of T. Besides ERM, pERM, ARMUL, and Spectral, the performance of MoM also improves as T increases. Similar to our finding in Sections [5.1.1](#page-26-1) and [5.1.2,](#page-27-0) MoM does not perform well until T becomes very large. This aligns with the empirical observations in [Tripuraneni et al.](#page-39-2) [\(2021\)](#page-39-2), where they found that ERM always outperforms MoM until T is very large, and the underlying reason is unclear.

When $\epsilon = 4\%$ tasks are contaminated, only the performance of pERM and Spectral improves as T increases, and they outperform all other methods, demonstrating their robustness against outlier tasks. ARMUL and GLasso perform slightly better than singletask regression, while the other benchmark methods suffer from severe negative transfer and are significantly impacted by the outliers.

Figure 8: Computational time of different methods with various numbers of tasks T. The y-axis is in the log-scale, and the unit is seconds. Each point represents the average over 100 replications.

Finally, we also recorded the computational time for different methods in setting (i). The experiments were run on the Terremoto HPC Cluster of Columbia University with a CPU Intel Xeon Gold 6126 2.6 GHz. We used a single core with 3 GB of memory when running each method. The computational time for different methods with different T values is plotted on a logarithmic scale in Figure [8.](#page-30-1) We can see that ARMUL and pERM are the most time-consuming methods, with pERM slightly faster than ARMUL, taking around $e^{6.25} \approx 500$ seconds for $T = 190$. In contrast, all the other methods can be run within $e^{3.5} \approx 30$ seconds for all values of T. This demonstrates the computational efficiency of the spectral method.

5.1.4 RELATIONSHIP BETWEEN TASK PERFORMANCE AND SIGNAL STRENGTH $\|\theta^{(t)*}\|_2$

In this subsection, we aim to verify our theoretical findings that the performance of ERM, pERM, and Spectral on each task can depend on the signal strength in terms of $\|\boldsymbol{\theta}^{(t)}*\|_2$, a relationship not previously discussed in the literature.

We generated data by mechanism similar to that in Section [5.1.1,](#page-26-1) with $n = 100$, $p = 50$. $r = 5, T = 10, h = \epsilon = 0$. The only difference in this section's data generation mechanism is that each $\theta^{(t)*}$ is uniformly generated from 0.5t S^{r-1} , i.e., the sphere centered at 0 in \mathbb{R}^r with a radius of 0.5t. We replicated this setting 100 times and summarized the average estimation error of the single-task regression, ERM, pERM, and Spectral on each task with different $\|\boldsymbol{\theta}^{(t)*}\|_2$ values in Figure [9.](#page-31-1)

method \triangle ERM \leftrightarrow pERM $\overline{\text{B}}$ Single−task B Spectral

Figure 9: Simulation verifying the relationship between the estimation error $\|\hat{\beta}^{(t)} - \beta^{(t)*}\|_2$ and signal strength $\|\theta^{(t)*}\|_2$ on each task for different methods. "error of each task" in the y-axis stands for $\|\hat{\beta}^{(t)} - \beta^{(t)*}\|_2$. Each point represents the average over 100 replications.

We can observe that the estimation error of ERM, pERM, and Spectral is approximately proportional to $\|\theta^{(t)*}\|_2$, which matches our theoretical findings. Specifically, the Pearson correlation coefficients between $\|\boldsymbol{\theta}^{(t)}\|_2$ and the average $\|\widehat{\boldsymbol{\beta}}^{(t)} - \boldsymbol{\beta}^{(t)*}\|_2$ for ERM, pERM, and Spectral are 0.977, 0.975, and 0.976, respectively. In contrast, single-task regression has comparable performance across different tasks, regardless of $\|\theta^{(t)*}\|_2$. This indicates that the benefit each individual task derives from representation MTL is highly dependent on the signal strength in terms of $\|\theta^{(t)*}\|_2$, unlike in single-task linear regression where the ℓ_2 -estimation error does not depend on the scale of the coefficient $\|\beta^{(t)*}\|_2$.

5.1.5 ADAPTIVITY TO THE INTRINSIC DIMENSION r

In the previous simulations, we used the true intrinsic dimension r in different representation MTL methods. In this subsection, we want to test the performance of our Algorithm [3](#page-23-1) for estimating r and its impact on helping pERM and Spectral adapt to unknown r in practice.

We consider the linear model [\(1\)](#page-1-0) under four settings, with different (n, p, r, T, ϵ) values:

- (i) $n = 100, p = 50, r = 5, T = 50, \epsilon = 0;$
- (ii) $n = 100, p = 50, r = 5, T = 50, \epsilon = 4\%$;
- (iii) $n = 150, p = 80, r = 10, T = 50, \epsilon = 0;$
- (iv) $n = 150, p = 80, r = 10, T = 50, \epsilon = 4\%.$

Given each (n, p, r, T, ϵ) , we increased h from 0 to 0.8 in increments of 0.1, generated data for tasks in S following the same machenism used in Section [5.1.1,](#page-26-1) and generated data for tasks

in S^c following the same machenism used in Section [5.1.2.](#page-27-0) We first ran Algorithm [3](#page-23-1) to obtain an estimate \hat{r} of r, then ran pERM and Spectral with \hat{r} . We refer to these two methods as pERM-adaptive and Spectral-adaptive, respectively. We also ran pERM and Spectral with the true r value as benchmarks, and we call them $pERM$ -oracle and Spectral-oracle, respectively. Each setting was replicated 100 times, and the average estimation error on tasks in S of different approaches, as well as the average \hat{r} , was plotted in Figure [10.](#page-32-1)

Figure 10: Simulation verifying the performance of Algorithm 3 for estimating r and its impact on helping pERM and Spectral adapt to unknown r . "max error in S " in the left y-axis stands for $\max_{t \in S} \|\hat{\beta}^{(t)} - \beta^{(t)*}\|_2$. The red \hat{r} in the right y-axis stands for the estimated r value. Each point represents the average over 100 replications.

We can see that the estimate \hat{r} is very close to the true r in all four settings when h is small, ensuring that pERM and Spectral using \hat{r} perform similarly to pERM and the spectral method with the true r. When h becomes large, the estimate \hat{r} may deviate from the true r. However, as we mentioned in Remark [12](#page-24-1) of Section [4,](#page-23-0) there is no need to estimate r precisely when h is large, and biased regularization in Step 2 of $pERM$ and Step 4 of Spectral can always guarantee the single-task performance.

5.2 A Real-data Study

In this subsection, we applied different approaches to a real data set, the Human Activity Recognition (HAR) Using Smartphones Data Set. This data set includes data collected from 30 volunteers performing six activities (walking, walking upstairs, walking downstairs, sitting, standing, and laying) with a smartphone [\(Anguita et al.,](#page-34-6) [2013\)](#page-34-6). Each observation has $p = 561$ time and frequency domain variables. We treated each volunteer as a task, with the sample size per task ranging from 281 to 409. The original data set is available at UCI Machine Learning Repository: [https://archive.ics.uci.edu/ml/datasets/human+](https://archive.ics.uci.edu/ml/datasets/human+activity+recognition+using+smartphones) [activity+recognition+using+smartphones](https://archive.ics.uci.edu/ml/datasets/human+activity+recognition+using+smartphones).

We focused on the binary classification problem of discriminating between walking and standing postures (i.e., walking, walking upstairs, walking downstairs, and standing) and the others (i.e., sitting and laying). We standarized the data of each task before training different algorithms. For each task, in each replication, we used 50% of the samples as training data and held 50% of the sample as test data.

We ran single-task logistic regression, pooled logistic regression, ERM, ARMUL, pERM, the spectral method, and GLasso on this problem, replicated it 100 times, and summarized the average mis-classification test error rates of different methods over $T = 30$ tasks with different r values in Table [2.](#page-33-1) We can see that $pERM$ consistently performs the best (tied with ERM when $r = 15$) among all the methods for different r values.

	r/Method Single-task Pooled	$\rm ERM$	ARMUL	pERM	Spectral	GLasso
$r=5$	$1.66(0.20)$ 1.79 (0.21) 1.62 (1.42) 2.12 (0.27) 1.33 (0.23) 1.85 (0.27) 1.44 (0.25)					
$r=10$			$1.66(0.20)$ $1.79(0.21)$ $1.42(0.23)$ $1.77(0.23)$ $1.25(0.20)$ $1.47(0.18)$ $1.44(0.25)$			
$r = 15$	$\{1.66(0.20)$ 1.79 (0.21) 1.36 (0.23) 1.68 (0.21) 1.36 (1.07) 1.50 (0.19) 1.44 (0.25)					

Table 2: The average mis-classification test error rates (standard deviations) of different methods over $T = 30$ tasks with different r values. All values are in percentages.

6. Discussions

In this work, we investigated the representation multi-task learning (MTL) problem, where most tasks share *similar* linear representations, and a small fraction of tasks can be *arbitrarily* contaminated. To address this problem, we proposed a penalized empirical risk minimization (ERM) method and a spectral method, and derived upper bounds for the estimation error. Our theory demonstrated that both algorithms are adaptive to the unknown similarity level between tasks and robust to a small fraction of outlier tasks. Additionally, the spectral method achieves a sharper estimation error bound than the penalized ERM when there is no contamination. Our theory also reveals the relationship between the performance of representation MTL methods on each task and the signal strength, which is usually ignored in the literature. We also presented the first lower bound results for estimating regression coefficients in the context of representation MTL. Our new spectral method is minimax optimal when there is no outlier task, and our penalized ERM is nearly optimal in a large regime with little impact by outlier tasks. We extended the algorithms and theory to generalized linear models and non-linear regression models. We also proposed a simple thresholding algorithm to adapt our MTL algorithms to the case of an unknown intrinsic dimension r. Finally, we conducted extensive numerical experiments to verify our theoretical findings in practice.

In Section [1.2.2,](#page-3-2) we extensively discussed the relationship between our work and the prior work beyond assuming different tasks sharing the same representation. We pointed out the most general scenario of the representation MTL problem, and this paper makes a significant step towards that direction. However, although we extended the theory of our spectral method to that general regime in the appendix, the optimal estimation error is still unknown, as well as which algorithm in that regime is optimal. We leave these interesting questions to future research.

Acknowledgments

Ye Tian is grateful to Gan Yuan (Columbia) and Yasaman Mahdaviyeh (Columbia) for a lot of valuable discussions which greatly improved the quality of this paper. Ye Tian also wants to thank Prof. Linjun Zhang (Rutgers) whose insightful discussions initially inspired this work. All numerical experiments were conducted on Ginsburg HPC Cluster and Terremoto HPC Cluster of Columbia University. Yuqi Gu acknowledges the support of the NSF Grant DMS-2210796. Yang Feng's research is partially supported by by NIH grant 1R21AG074205-01, NYU University Research Challenge Fund, and a grant from NYU School of Global Public Health. The authors are grateful to the Action Editor and four reviewers for their helpful and constructive comments.

References

- R. K. Ando, T. Zhang, and P. Bartlett. A framework for learning predictive structures from multiple tasks and unlabeled data. Journal of Machine Learning Research, 6(11), 2005.
- D. Anguita, A. Ghio, L. Oneto, X. Parra, J. Reyes-Ortiz, et al. A public domain dataset for human activity recognition using smartphones. In 21th European Symposium on Artificial Neural Networks, Computational Intelligence and Machine Learning (ESANN), pages 437–442. CIACO, 2013.
- M. Bakhshizadeh, A. Maleki, and V. H. de la Pena. Sharp concentration results for heavytailed distributions. arXiv preprint arXiv:2003.13819, 2020.
- S. Balakrishnan, M. J. Wainwright, and B. Yu. Statistical guarantees for the em algorithm: From population to sample-based analysis. The Annals of Statistics, 45(1):77–120, 2017.
- H. Bastani. Predicting with proxies: Transfer learning in high dimension. Management Science, 67(5):2964–2984, 2021.
- J. Baxter. A model of inductive bias learning. Journal of artificial intelligence research, 12: 149–198, 2000.
- S. Ben-David and R. S. Borbely. A notion of task relatedness yielding provable multiple-task learning guarantees. Machine learning, 73:273–287, 2008.
- S. Ben-David, J. Blitzer, K. Crammer, A. Kulesza, F. Pereira, and J. W. Vaughan. A theory of learning from different domains. Machine learning, 79:151–175, 2010.
- T. T. Cai and H. Wei. Transfer learning for nonparametric classification: Minimax rate and adaptive classifier. The Annals of Statistics, 49(1), 2021.
- T. T. Cai, Z. Ma, and Y. Wu. Sparse pca: Optimal rates and adaptive estimation. Annals of Statistics, 41(6):3074–3110, 2013.
- T. T. Cai, J. Ma, and L. Zhang. Chime: Clustering of high-dimensional gaussian mixtures with em algorithm and its optimality 1. Annals of Statistics, $47(3):1234-1267$, 2019.
- E. J. Candès, X. Li, Y. Ma, and J. Wright. Robust principal component analysis? Journal of the ACM (JACM), 58(3):1–37, 2011.
- H. Cao, J. Zhou, and E. Schwarz. Rmtl: an r library for multi-task learning. Bioinformatics, 35(10):1797–1798, 2019.
- M. Chen, C. Gao, and Z. Ren. Robust covariance and scatter matrix estimation under huber's contamination model. The Annals of Statistics, 46(5):1932–1960, 2018.
- S. Chen, K. Crammer, H. He, D. Roth, and W. J. Su. Weighted training for cross-task learning. $arXiv$ preprint $arXiv:2105.14095$, 2021a.
- X. Chen, M. Chen, C. Fan, A. Uppunda, Y. Sun, and C. Zaniolo. Multilingual knowledge graph completion via ensemble knowledge transfer. In Findings of the Association for Computational Linguistics: EMNLP 2020, pages 3227–3238, 2020.
- Y. Chen, Y. Chi, J. Fan, C. Ma, et al. Spectral methods for data science: A statistical perspective. Foundations and Trends \widehat{R} in Machine Learning, 14(5):566–806, 2021b.
- K. Chua, Q. Lei, and J. D. Lee. How fine-tuning allows for effective meta-learning. Advances in Neural Information Processing Systems, 34:8871–8884, 2021.
- L. Collins, H. Hassani, A. Mokhtari, and S. Shakkottai. Exploiting shared representations for personalized federated learning. In International Conference on Machine Learning, pages 2089–2099. PMLR, 2021.
- K. Crammer, M. Kearns, and J. Wortman. Learning from multiple sources. Journal of Machine Learning Research, 9(8), 2008.
- G. Denevi, M. Pontil, and C. Ciliberto. The advantage of conditional meta-learning for biased regularization and fine tuning. Advances in Neural Information Processing Systems, 33:964–974, 2020.
- S. Deng, Y. Guo, D. Hsu, and D. Mandal. Learning tensor representations for meta-learning. In International Conference on Artificial Intelligence and Statistics, pages 11550–11580. PMLR, 2022.
- J. Donahue, Y. Jia, O. Vinyals, J. Hoffman, N. Zhang, E. Tzeng, and T. Darrell. Decaf: A deep convolutional activation feature for generic visual recognition. In International conference on machine learning, pages 647–655. PMLR, 2014.
- S. S. Du, W. Hu, S. M. Kakade, J. D. Lee, and Q. Lei. Few-shot learning via learning the representation, provably. arXiv preprint arXiv:2002.09434, 2020.
- Y. Duan and K. Wang. Adaptive and robust multi-task learning. *arXiv preprint* arXiv:2202.05250, 2022.
- J. Duchi, V. Feldman, L. Hu, and K. Talwar. Subspace recovery from heterogeneous data with non-isotropic noise. arXiv preprint arXiv:2210.13497, 2022.
- C. Dwork. Differential privacy. In International colloquium on automata, languages, and programming, pages 1–12. Springer, 2006.
- C. Dwork, A. Roth, et al. The algorithmic foundations of differential privacy. Foundations and Trends \widehat{R} in Theoretical Computer Science, 9(3-4):211-407, 2014.
- J. Fan, K. Wang, Y. Zhong, and Z. Zhu. Robust high dimensional factor models with applications to statistical machine learning. Statistical science: a review journal of the Institute of Mathematical Statistics, 36(2):303, 2021.
- C. Finn, P. Abbeel, and S. Levine. Model-agnostic meta-learning for fast adaptation of deep networks. In International conference on machine learning, pages 1126–1135. PMLR, 2017.
- C. Finn, A. Rajeswaran, S. Kakade, and S. Levine. Online meta-learning. In International Conference on Machine Learning, pages 1920–1930. PMLR, 2019.
- P. Goyal, D. Mahajan, A. Gupta, and I. Misra. Scaling and benchmarking self-supervised visual representation learning. In Proceedings of the ieee/cvf International Conference on computer vision, pages 6391–6400, 2019.
- T. Gu, Y. Han, and R. Duan. Robust angle-based transfer learning in high dimensions. arXiv preprint arXiv:2210.12759, 2022.
- T. Gu, P. H. Lee, and R. Duan. Commute: Communication-efficient transfer learning for multi-site risk prediction. Journal of Biomedical Informatics, 137:104243, 2023.
- A. Gupta, C. Devin, Y. Liu, P. Abbeel, and S. Levine. Learning invariant feature spaces to transfer skills with reinforcement learning. arXiv preprint arXiv:1703.02949, 2017.
- S. Hanneke and S. Kpotufe. On the value of target data in transfer learning. Advances in Neural Information Processing Systems, 32, 2019.
- S. Hanneke and S. Kpotufe. A no-free-lunch theorem for multitask learning. The Annals of Statistics, 50(6):3119–3143, 2022.
- W. Härdle and T. M. Stoker. Investigating smooth multiple regression by the method of average derivatives. Journal of the American statistical Association, 84(408):986–995, 1989.
- T. Hospedales, A. Antoniou, P. Micaelli, and A. Storkey. Meta-learning in neural networks: A survey. IEEE transactions on pattern analysis and machine intelligence, 44(9):5149–5169, 2021.
- M. Hristache, A. Juditsky, and V. Spokoiny. Direct estimation of the index coefficient in a single-index model. Annals of Statistics, pages 595–623, 2001.
- A. Jalali, S. Sanghavi, C. Ruan, and P. Ravikumar. A dirty model for multi-task learning. Advances in neural information processing systems, 23, 2010.
- T. Kato. Perturbation theory for linear operators, volume 132. Springer Science & Business Media, 2013.
- D. P. Kingma and J. L. Ba. Adam: A method for stochastic gradient descent. In *ICLR*: international conference on learning representations, pages 1–15. ICLR US., 2015.
- N. Konstantinov, E. Frantar, D. Alistarh, and C. Lampert. On the sample complexity of adversarial multi-source pac learning. In International Conference on Machine Learning, pages 5416–5425. PMLR, 2020.
- I. Kuzborskij and F. Orabona. Stability and hypothesis transfer learning. In International Conference on Machine Learning, pages 942–950. PMLR, 2013.
- I. Kuzborskij and F. Orabona. Fast rates by transferring from auxiliary hypotheses. Machine Learning, 106:171–195, 2017.
- S. Li, T. Cai, and R. Duan. Targeting underrepresented populations in precision medicine: A federated transfer learning approach. arXiv preprint arXiv:2108.12112, 2021.
- S. Li, T. T. Cai, and H. Li. Transfer learning in large-scale gaussian graphical models with false discovery rate control. Journal of the American Statistical Association, pages 1–13, 2022a.
- S. Li, T. T. Cai, H. Li, et al. Transfer learning for high-dimensional linear regression: Prediction, estimation and minimax optimality. Journal of the Royal Statistical Society Series B, 84(1):149–173, 2022b.
- H. Lin and M. Reimherr. On transfer learning in functional linear regression. $arXiv$ preprint arXiv:2206.04277, 2022.
- G. Liu, Z. Lin, S. Yan, J. Sun, Y. Yu, and Y. Ma. Robust recovery of subspace structures by low-rank representation. IEEE transactions on pattern analysis and machine intelligence, 35(1):171–184, 2012.
- P.-L. Loh and M. J. Wainwright. Regularized m-estimators with nonconvexity: Statistical and algorithmic theory for local optima. Journal of Machine Learning Research, 16: 559–616, 2015.
- K. Lounici, M. Pontil, A. Tsybakov, and S. Van De Geer. Taking advantage of sparsity in multi-task learning. In COLT 2009-The 22nd Conference on Learning Theory, 2009.
- K. Lounici, M. Pontil, S. van de Geer, and A. B. Tsybakov. Oracle inequalities and optimal inference under group sparsity. The Annals of Statistics, 39(4):2164–2204, 2011.
- G. Lugosi and S. Mendelson. Robust multivariate mean estimation: the optimality of trimmed mean. The Annals of Statistics, 49(1):393–410, 2021.
- A. Maurer, M. Pontil, and B. Romera-Paredes. The benefit of multitask representation learning. Journal of Machine Learning Research, 17(81):1–32, 2016.
- P. McCullagh and J. A. Nelder. Generalized Linear Models, volume 37. CRC Press, 1989.
- S. N. Negahban, P. Ravikumar, M. J. Wainwright, and B. Yu. A unified framework for high-dimensional analysis of m -estimators with decomposable regularizers. Statistical Science, 27(4):538, 2012.
- A. Nichol, J. Achiam, and J. Schulman. On first-order meta-learning algorithms. arXiv preprint arXiv:1803.02999, 2018.
- A. Onatski. Determining the number of factors from empirical distribution of eigenvalues. The Review of Economics and Statistics, 92(4):1004–1016, 2010.
- A. Pajor. Metric entropy of the grassmann manifold. Convex Geometric Analysis, 34: 181–188, 1998.
- A. Paszke, S. Gross, F. Massa, A. Lerer, J. Bradbury, G. Chanan, T. Killeen, Z. Lin, N. Gimelshein, L. Antiga, et al. Pytorch: An imperative style, high-performance deep learning library. Advances in neural information processing systems, 32, 2019.
- M. Qiao. Do outliers ruin collaboration? In International Conference on Machine Learning, pages 4180–4187. PMLR, 2018.
- M. Qiao and G. Valiant. Learning discrete distributions from untrusted batches. In 9th Innovations in Theoretical Computer Science Conference (ITCS 2018). Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2018.
- M. Raghu, C. Zhang, J. Kleinberg, and S. Bengio. Transfusion: Understanding transfer learning for medical imaging. Advances in neural information processing systems, 32, 2019.
- G. Raskutti, M. J. Wainwright, and B. Yu. Minimax rates of estimation for high-dimensional linear regression over ℓ_q -balls. IEEE transactions on information theory, 57(10):6976–6994, 2011.
- M. Rostami, H. He, M. Chen, and D. Roth. Transfer learning via representation learning. In Federated and Transfer Learning, pages 233–257. Springer International Publishing Cham, 2022.
- A. M. Samarov. Exploring regression structure using nonparametric functional estimation. Journal of the American Statistical Association, 88(423):836–847, 1993.
- B. Schölkopf, R. Herbrich, and A. J. Smola. A generalized representer theorem. In International conference on computational learning theory, pages 416–426. Springer, 2001.
- K. K. Thekumparampil, P. Jain, P. Netrapalli, and S. Oh. Statistically and computationally efficient linear meta-representation learning. Advances in Neural Information Processing Systems, 34:18487–18500, 2021.
- Y. Tian and Y. Feng. Transfer learning under high-dimensional generalized linear models. Journal of the American Statistical Association, pages 1–14, 2022.
- Y. Tian, H. Weng, and Y. Feng. Unsupervised multi-task and transfer learning on gaussian mixture models. arXiv preprint arXiv:2209.15224, 2022.
- N. Tripuraneni, M. Jordan, and C. Jin. On the theory of transfer learning: The importance of task diversity. Advances in neural information processing systems, 33:7852–7862, 2020.
- N. Tripuraneni, C. Jin, and M. Jordan. Provable meta-learning of linear representations. In International Conference on Machine Learning, pages 10434–10443. PMLR, 2021.
- A. B. Tsybakov. Introduction to nonparametric estimation. Springer, New York, 2009.
- R. Vershynin. Introduction to the non-asymptotic analysis of random matrices. $arXiv$ preprint arXiv:1011.3027, 2010.
- R. Vershynin. High-dimensional probability: An introduction with applications in data science, volume 47. Cambridge university press, 2018.
- R. Vidal, Y. Ma, S. S. Sastry, R. Vidal, Y. Ma, and S. S. Sastry. Principal component analysis. Generalized principal component analysis, pages 25–62, 2016.
- M. J. Wainwright. High-dimensional statistics: A non-asymptotic viewpoint, volume 48. Cambridge university press, 2019.
- A. Wang, J. Hula, P. Xia, R. Pappagari, R. T. McCoy, R. Patel, N. Kim, I. Tenney, Y. Huang, K. Yu, et al. Can you tell me how to get past sesame street? sentence-level pretraining beyond language modeling. In Proceedings of the 57th Annual Meeting of the Association for Computational Linguistics, pages 4465–4476, 2019.
- P.-Å. Wedin. Perturbation bounds in connection with singular value decomposition. BIT Numerical Mathematics, 12:99–111, 1972.
- K. Weiss, T. M. Khoshgoftaar, and D. Wang. A survey of transfer learning. *Journal of Big* data, $3(1):1-40$, 2016 .
- J. Wright, A. Ganesh, S. Rao, Y. Peng, and Y. Ma. Robust principal component analysis: Exact recovery of corrupted low-rank matrices via convex optimization. Advances in neural information processing systems, 22, 2009.
- H. Xu, C. Caramanis, and S. Sanghavi. Robust pca via outlier pursuit. IEEE Transactions on Information Theory, 58(5):3047–3064, 2012.
- K. Xu and H. Bastani. Learning across bandits in high dimension via robust statistics. $arXiv$ preprint arXiv:2112.14233, 2021.
- Z. Yang, Z. Wang, H. Liu, Y. C. Eldar, and T. Zhang. Sparse nonlinear regression: Parameter estimation and asymptotic inference. arXiv preprint arXiv:1511.04514, 2015.
- Z. Yang, K. Balasubramanian, Z. Wang, and H. Liu. Learning non-gaussian multi-index model via second-order stein's method. Advances in Neural Information Processing Systems, 30:6097–6106, 2017.
- Y. Yu, T. Wang, and R. J. Samworth. A useful variant of the davis–kahan theorem for statisticians. Biometrika, 102(2):315–323, 2015.
- G. Yuan, M. Xu, S. Kpotufe, and D. Hsu. Efficient estimation of the central mean subspace via smoothed gradient outer products. arXiv preprint arXiv:2312.15469, 2023.
- M. Yuan and Y. Lin. Model selection and estimation in regression with grouped variables. Journal of the Royal Statistical Society Series B: Statistical Methodology, 68(1):49–67, 2006.
- Y. Zhang and Q. Yang. An overview of multi-task learning. National Science Review, 5(1): 30–43, 2018.
- Y. Zhang and Q. Yang. A survey on multi-task learning. IEEE Transactions on Knowledge and Data Engineering, 34(12):5586–5609, 2021.
- D. Zhou, T. Cai, and J. Lu. Multi-source learning via completion of block-wise overlapping noisy matrices. arXiv preprint arXiv:2105.10360, 2021.
- D. Zhou, M. Liu, M. Li, and T. Cai. Doubly robust augmented model accuracy transfer inference with high dimensional features. arXiv preprint arXiv:2208.05134, 2022.

Content of Appendices

Appendix A. Spectral Method in the General Regime

In Section [1.2.2,](#page-3-0) we introduced a general regime, which includes the settings of many papers in the literature (e.g., [Du et al.,](#page-36-0) [2020;](#page-36-0) [Tripuraneni et al.,](#page-39-0) [2021;](#page-39-0) [Thekumparampil et al.,](#page-39-1) [2021;](#page-39-1) [Chua et al.,](#page-35-0) [2021;](#page-35-0) [Duan and Wang,](#page-36-1) [2022\)](#page-36-1) and our setting described in Section [2](#page-8-0) as special cases. We mentioned in Section [2.3](#page-16-1) that the theory of the spectral method (Algorithm [2\)](#page-17-1) can be generalized to this general regime. We aim to introduce the generalization in this section.

Before we provide the upper bound of the estimation erorr for the spectral method in the general regime, let us review the setup of the general regime. Consider the linear regression model

$$
y_i^{(t)} = (\boldsymbol{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*} + \epsilon_i^{(t)}, \quad i = 1 : n, \quad t = 1 : T,
$$

where

$$
\boldsymbol{\beta}^{(t)*} = \overline{\boldsymbol{A}} \boldsymbol{\theta}^{(t)*} + \boldsymbol{\delta}^{(t)*},
$$

with $\overline{A} \in \mathcal{O}^{p \times r}$, low-dimensional parameter $\theta^{(t)*} \in \mathbb{R}^r$, $r \leq p$, and $\{\epsilon_i^{(t)}\}$ $\binom{t}{i}_{i=1}^n$ are i.i.d. zero-mean sub-Gaussian variables independent of $\{\boldsymbol{x}_i^{(t)}\}$ $\{\xi^{(t)}\}_{i=1}^n$. Denote $\zeta^{(t)} = \|\theta^{(t)*}\|_2$, $\bar{\zeta} =$ $\sqrt{1}$ $\frac{1}{|S|}\sum_{t\in S}(\zeta^{(t)})^2$, and $\mathbf{D}_{S}^* = {\delta^{(t)*}}_{t\in S} \in \mathbb{R}^{p\times |S|}$ the $p \times |S|$ matrix with t-th column $\delta^{(t)*}$. For simplicity, similar to Section [2,](#page-8-0) we assume $\min_{t \in [T]} \zeta^{(t)} \gtrsim \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$.

We replace Assumption [2](#page-10-0) with the following one.

Assumption 8 There exists a constant $c > 0$ such that $\sigma_r(\mathbf{\Theta}^*/\sqrt{T}) \geq \frac{c}{\sqrt{r}}\overline{\zeta}$, where each column of $\mathbf{\Theta}^* \in \mathbb{R}^{p \times T}$ is a coefficient vector in $\{\mathbf{\theta}^{(t)*}\}_{t=1}^T$.

Now, we are ready to introduce the upper bound of estimation error for the spectral method in the general regime.

Theorem 13 (Upper bound for spectral method in the general regime) Suppose As-sumptions [1,](#page-10-1) [3,](#page-10-2) [8](#page-43-2) hold with a subset $S \subseteq [T]$ satisfying $\epsilon = \frac{|S^c|}{T} \leq c r^{-1} \cdot \left(\frac{\bar{\zeta}}{\max_{t \in \bar{\zeta}}} \right)$ $\frac{\bar{\zeta}}{\max_{t \in S} \zeta^{(t)}}\right)^2$, where $c > 0$ is a small constant. By setting $\gamma = C' \sqrt{p + \log T}$ with a sufficiently large positive constant C' and $\bar{\epsilon}$ satisfying $\epsilon \leq \bar{\epsilon} \leq \frac{c''}{r}$ $\frac{\bar{\zeta}''}{r} \cdot \left(\frac{\bar{\zeta}}{\max_{t \in \zeta}} \right)$ $\left(\frac{\bar{\zeta}}{\max_{t \in S} \zeta^{(t)}}\right)^2$ with a sufficiently small positive constant c'', w.p. at least $1 - e^{-C''(r + \log T)}$, we have

$$
\|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \lesssim \left\{ \frac{\zeta^{(t)}}{\bar{\zeta}} \sqrt{\frac{pr}{nT}} + \left(\frac{\zeta^{(t)}}{\bar{\zeta}} \vee 1\right) \sqrt{\frac{r}{n}} + \frac{\zeta^{(t)}}{\bar{\zeta}} \sqrt{\frac{r}{T}} \|\mathbf{D}_S^*\|_2 + \|\boldsymbol{\delta}^{(t)*}\|_2 + \sqrt{\frac{\log T}{n}}
$$

$$
+ \frac{\zeta^{(t)}}{\bar{\zeta}} \max_{t \in S} \|\boldsymbol{\beta}^{(t)*}\|_2 \sqrt{r\bar{\epsilon}} \right\} \wedge \sqrt{\frac{p + \log T}{n}}, \quad \forall t \in S.
$$

Furthermore, if we assume the data from tasks in S^c satisfies the linear model [\(3\)](#page-8-1) (without any latent structure assumption) and Assumption [1,](#page-10-1) then w.p. at least $1 - e^{-C'(p + \log T)}$, we also have

$$
\max_{t \in [T]} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \lesssim \sqrt{\frac{p + \log T}{n}}.
$$

To the best of our knowledge, this is the first algorithm which is both adaptive to the unknown similarity structure and robust to a small fraction of outlier tasks in the general regime. Comparing the bound in Theorem [13](#page-43-1) with the bounds in Section [2,](#page-8-0) we can see that the impact of task heterogeneity has a more complicated form in the general regime, i.e., $\zeta^{(t)}$ $\frac{d^{(t)}}{\zeta} \sqrt{\frac{r}{T}} \|\mathcal{D}_{S}^{*}\|_{2} + \|\boldsymbol{\delta}^{(t)*}\|_{2}$. The other terms are very similar or the same as in the regime we focused on in the main text.

Unlike the spectral method, our penalized ERM algorithm does not work in the general regime, because the penalty we added in Algorithm [1](#page-11-0) only involves the maximum principal angle between representations, instead of the offset terms $\delta^{(t)*}$'s in the general regime.

Appendix B. Transferring to New Tasks (Learning-to-learn)

In this section, we extend the MTL framework discussed in the main text to a transfer learning (TL) setting.

B.1 Problem Set-up

In this section, in addition to data from the T tasks, suppose we also observe data $\{(\boldsymbol{x}^{(0)}_i$ $\binom{0}{i}, y_i^{(0)}$ $\binom{0}{i}$ $\}_{i=1}^{n_0}$ from a new task

$$
y_i^{(0)} = (\boldsymbol{x}_i^{(0)})^\top \boldsymbol{\beta}^{(0)*} + \epsilon_i^{(0)}, \quad i = 1 : n_0,
$$
\n⁽⁷⁾

where $\boldsymbol{\beta}^{(0)*} = \boldsymbol{A}^{(0)*}\boldsymbol{\theta}^{(0)*}, \ \boldsymbol{A}^{(0)*} \in \mathcal{O}^{p \times r} = \{ \boldsymbol{A} \in \mathbb{R}^{p \times r} : \ \boldsymbol{A}^\top \boldsymbol{A} = \boldsymbol{I}_r \}, \ \boldsymbol{\theta}^{(0)*} \in \mathbb{R}^r, \ \mathrm{and}$ $\{\epsilon_i^{(0)}$ ${}_{i}^{(0)}\}^{n_0}_{i=1}$ are i.i.d. zero-mean sub-Gaussian variables independent of $\{\boldsymbol{x}_{i}^{(0)}\}$ $\binom{0}{i}$ _{$i=1$}. Similar to Section [2,](#page-8-0) the intrinsic dimension r is assumed to be known. Under such a transfer learning (T_L) or learning-to-learn setting, the new task is often called the target task, and the T tasks are called source tasks. Our goal is two-fold:

- 1. Transfer knowledge from source tasks to improve the learning performance on the target task, when the source and target share "similar" representations and the number of outlier source tasks is small;
- 2. Ensure the learning performance is no worse than the target-only learning performance to avoid the negative transfer.

To describe the similarity between target and source representations, we assume

$$
\max_{t \in S} \|A^{(t)*}(\boldsymbol{A}^{(t)*})^{\top} - \boldsymbol{A}^{(0)*}(\boldsymbol{A}^{(0)*})^{\top} \|_2 \leq h,
$$

where S is a subset of $[T]$. Similar to the setting in the last section, the joint distribution \mathbb{Q}_{S^c} of data from source tasks in S^c , i.e., $\{\{\boldsymbol{x}_i^{(t)}\}$ $\binom{t}{i},y_i^{(t)}$ $\{e^{(t)}\}_{i=1}^n\}_{t\in S^c}$, is allowed to be arbitrary. For simplicity, we focus on the regime that $||\boldsymbol{\theta}^{(t)*}||_2 \leq C < \infty$ for all $t \in S$. Denote $\Sigma^{(0)} = \mathbb{E}[\boldsymbol{x}^{(0)}(\boldsymbol{x}^{(0)})^{\top}]$. We impose the following assumptions on the target task.

Assumption 9 For $u \in \mathbb{R}^p$, $u^{\top}x^{(0)}$ is sub-Gaussian in the sense that $\mathbb{E}[e^{\lambda u^{\top}x^{(0)}}] \leq$ $e^{C\lambda^2\|\mathbf{u}\|_2^2}$ for any $\lambda \in \mathbb{R}$ with some constant $C > 0$. And there exist constants c, C such that $0 < c \leq \lambda_{\min}(\mathbf{\Sigma}^{(0)}) \leq \lambda_{\max}(\mathbf{\Sigma}^{(0)}) \leq C < \infty.$

Assumption 10 $n_0 \geq Cp$ with a sufficiently large constant $C > 0$.

Remark 14 Similar to our discussion in Section [2.2.1](#page-11-1) for the penalized ERM in MTL, Assumption [10](#page-44-2) is imposed to guarantee the target-only rate $\sqrt{p/n_0}$. If we do not care about this safe-net guarantee, then it suffices to require $n_0 \geq Cr$, the same as the condition imposed in literature [\(Tripuraneni et al.,](#page-39-0) [2021;](#page-39-0) [Thekumparampil et al.,](#page-39-1) [2021\)](#page-39-1). The RHS of the following Theorem [15](#page-45-0) shall be replaced by $r\sqrt{\frac{p}{nT}} +$ $\sqrt{r}h + \sqrt{r}\sqrt{\frac{r+\log T}{n}} + \sqrt{\frac{p}{n}} \cdot \frac{|S^c|}{T}$ $\frac{S^c|}{T} r^{3/2} + \sqrt{\frac{r}{n_0}},$ which allows us for a few-shot learning when both h and $|S^c|/T$ are sufficiently small. See more details in the proof of Theorem [15](#page-45-0) in Appendix.

B.2 Upper Bounds

Similar to the Algorithm [1,](#page-11-0) a two-step transfer learning method is proposed in Algorithm [4.](#page-45-1) We introduce the algorithm with a general loss function $f^{(0)}$ for the target since the same

algorithm with different losses will be extended to other models later. For linear model [\(7\)](#page-44-3), define $f^{(0)}(\boldsymbol{\beta}) = \frac{1}{2n_0} ||\boldsymbol{Y}^{(0)} - \boldsymbol{X}^{(0)}\boldsymbol{\beta}||_2 = \frac{1}{2n_0}$ $\frac{1}{2n_0}\sum_{i=1}^{n_0}[y_i^{(0)}-(\pmb{x}_i^{(0)}$ $i^{(0)}$)[⊤] β ² for $\beta \in \mathbb{R}^p$, where $X^{(0)} \in \mathbb{R}^{n_0 \times p}$ and $Y^{(0)} \in \mathbb{R}^{n_0}$ are corresponding matrix/vector representations of target data.

Algorithm 4: Transferring to new tasks

 $\textbf{Input:} \text{ Data from a new task } (\boldsymbol{X}^{(0)}, \boldsymbol{Y}^{(0)}) = \{ \boldsymbol{x}_i^{(0)} \}$ $\binom{(0)}{i}, y_i^{(0)}$ $i^{(0)}\}_{i=1}^{n_0}$, estimator \overline{A} from Algorithm [1](#page-11-0) or [2,](#page-17-1) penalty parameter γ **Output:** Estimator $\hat{\boldsymbol{\beta}}^{(0)}$ 1 Step 1: $\widehat{\boldsymbol{\theta}}^{(0)} = \arg \min_{\boldsymbol{\theta} \in \mathbb{R}^r} \left\{ f^{(0)}(\overline{\boldsymbol{A}}\boldsymbol{\theta}) \right\}$ 2 $\text{Step 2: } \widehat{\boldsymbol{\beta}}^{(0)} = \arg \min_{\boldsymbol{\beta} \in \mathbb{R}^p} \left\{ f^{(0)}(\boldsymbol{\beta}) + \frac{\gamma}{\sqrt{n_0}} || \boldsymbol{\beta} - \overline{\boldsymbol{A}} \widehat{\boldsymbol{\theta}}^{(0)} ||_2 \right\}$

In Algorithm [4,](#page-45-1) the "central representation" $\hat{\overline{A}}$ learned by Algorithm [1](#page-11-0) is passed to Step 1 to obtain the estimator $\hat{\theta}^{(0)}$ of the target-specific low-dimensional parameter. The same step has appeared in literature when there are no outliers, and the target and source share the same representations [\(Du et al.,](#page-36-0) [2020;](#page-36-0) [Tripuraneni et al.,](#page-39-0) [2021\)](#page-39-0). Step 2 is similar to Step 2 of Algorithm [1,](#page-11-0) which guarantees the target-only rate even when the representations of the target and sources are dissimilar.

We have the following upper bounds of target estimation error for Algorithm [4.](#page-45-1)

Theorem 15 (Upper bound for TL) Suppose Assumptions [1,](#page-10-1) [2,](#page-10-0) [3,](#page-10-2) [9,](#page-44-4) and [10](#page-44-2) hold.

(i) (TL with penalized ERM) By setting $\lambda = Cr^{3/4}\sqrt{p + \log T}$ and $\gamma = C'\sqrt{p + \log T}$ with sufficiently large constants $C, C' > 0$ in Algorithm [4](#page-45-1) along with Algorithm [1](#page-11-0) to learn $\widehat{\overline{A}}$, for all $S \subseteq [T]$ satisfying $\epsilon = \frac{|S^c|}{|T|} \leq c r^{-3/2}$ with a small constant $c > 0$ and an arbitrary distribution \mathbb{Q}_{S^c} of $\{\{\boldsymbol{x}_i^{(t)}\}$ $\binom{t}{i}, y_{i}^{(t)}$ $\{e^{(t)}\}_{i=1}^n\}_{t\in S^c}$, w.p. at least 1 − $e^{-C''(r+\log T)}$, we have

$$
\|\widehat{\beta}^{(0)} - \beta^{(0)*}\|_2 \lesssim \left(r\sqrt{\frac{p}{nT}} + \sqrt{r}h + \sqrt{r}\sqrt{\frac{r+\log T}{n}} + \sqrt{\frac{p}{n}} \cdot \epsilon r^{3/2}\right) \wedge \sqrt{\frac{p}{n_0}} + \sqrt{\frac{r}{n_0}}.
$$

(ii) (TL with the spectral method) By setting $\gamma = C' \sqrt{p + \log T}$ with a sufficiently large positive constant C', for any subset $S \subseteq [T]$ satisfying $\epsilon = \frac{|S^c|}{T} \leq c r^{-1} \cdot \left(\frac{\bar{\zeta}^c}{\max_{t \in \bar{\zeta}} |S|}\right)$ $\frac{\bar{\zeta}}{\max_{t \in S} \zeta^{(t)}}\Big)^2$ with $c > 0$ a small constant, w.p. at least $1 - e^{-C''(r + \log T)}$ and $\bar{\epsilon}$ satisfying $\epsilon \leq \bar{\epsilon} \leq$ c'' $\frac{\bar{\zeta}''}{r} \cdot \left(\frac{\bar{\zeta}}{\max_{t \in \zeta}} \right)$ $\int_{\max_{t \in S} \zeta^{(t)}}^{\overline{\zeta}} \int_0^{\infty}$ with a sufficiently small positive constant c'' , we have

$$
\|\widehat{\beta}^{(0)}-\beta^{(0)*}\|_2 \lesssim \left\{\sqrt{\frac{pr}{nT}}+\sqrt{\frac{r}{n}}+h\cdot \left[\frac{\sigma_{\max}((\overline{\mathbf{A}}^{\perp})^{\top}\mathbf{B}^*_{S})}{\sigma_{\min}((\overline{\mathbf{A}}^{\perp})^{\top}\mathbf{B}^*_{S})}\wedge \sqrt{r}\right]+\sqrt{r\overline{\epsilon}}\right\}\wedge \sqrt{\frac{p}{n_0}}+\sqrt{\frac{r}{n_0}},
$$

where $B_S^* \in \mathbb{R}^{p \times |S|}$ is the coefficient matrix whose columns are $\{\beta^{(t)*}\}_{t \in S}$, $\overline{A} \in$ $\arg\min_{\bm{A}\in\mathcal{O}^{p\times r}}\max_{t\in[T]}\|\bm{A}^{(t)*}(\bm{A}^{(t)*})^\top - \bm{A}\bm{A}^\top \|_2$ is the central representation, and $\overline{\bm{A}}^\perp$ is orthogonal to \overline{A} in the sense that $\overline{A}^{\perp} \in \mathcal{O}^{p \times (p-r)}$ and $(\overline{A}^{\perp})^{\top} \overline{A} = \mathbf{0}_{(p-r) \times r}$.

Both upper bounds can be seen as the minimum of two terms which represent the rate of learning target model via data aggregation and the target-only rate $\sqrt{\frac{p}{n_0}}$, respectively. This rate entails that our algorithm is **adaptive** to the optimal situation regardless of whether transferring from source to target is beneficial. Moreover, it is robust to a small fraction of outlier source tasks, in the sense that TL is still helpful when the outlier proportion ϵ is sufficiently small.

B.3 Lower Bounds

In this subsection, we explore the lower bound of the TL problem. Consider the space for all subsets $S \subseteq [T]$ as

$$
\mathcal{S} = \{ S \subseteq [T] : |S^c|/T \le \epsilon \}.
$$

Given the subset S, consider the parameter spaces for the coefficient vectors $\{\beta^{(t)}\}_{t\in\{0\}\cup S}$ as

$$
\mathscr{B}_{0}(S,h) = \left\{ \{ \boldsymbol{\beta}^{(t)} \}_{t \in \{0\} \cup S} : \boldsymbol{\beta}^{(t)} = \boldsymbol{A}^{(t)} \boldsymbol{\theta}^{(t)} \text{ for all } t \in \{0\} \cup S, \{ \boldsymbol{A}^{(t)} \}_{t \in \{0\} \cup S} \subseteq \mathcal{O}^{p \times r}, \right\}
$$
\n
$$
\max_{t \in \{0\} \cup S} \| \boldsymbol{\theta}^{(t)} \|_{2} \le C, \max_{t \in S} \| \boldsymbol{A}^{(t)} (\boldsymbol{A}^{(t)})^{\top} - \boldsymbol{A}^{(0)} (\boldsymbol{A}^{(0)})^{\top} \|_{2} \le h, \sigma_{r} \left(|S|^{-1/2} \boldsymbol{B}_{S}^{*} \right) \ge \frac{c}{\sqrt{r}} \right\}
$$

where C and c can be any fixed positive constants such that $\mathscr{B}_0(S, h) \neq \emptyset$.

Theorem 16 (Lower bound for TL) Suppose $p \geq 2r$ and $\epsilon \leq c/r$ where c is a small constant. We have the following lower bound:

$$
\inf_{\widehat{\beta}^{(0)}} \sup_{S \subseteq \mathcal{S}} \sup_{\{\beta^{(t)}\}_{t \in \{0\}\cup S} \in \mathscr{B}_0(S,h)} \mathbb{P}\bigg(\|\widehat{\beta}^{(0)} - \beta^{(0)*}\|_2 \gtrsim \left(\sqrt{\frac{pr}{nT}} + h\right) \wedge \sqrt{\frac{p}{n_0}} + \sqrt{\frac{r}{n_0}} + \frac{\epsilon r}{\sqrt{n}} \wedge \sqrt{\frac{1}{n_0}} \bigg) \ge \frac{1}{10}.
$$

To our knowledge, this is the first lower bound for learning regression parameters under representation transfer learning. Comparing the upper and lower bounds of the representation TL problem, we can see that the upper bound of penalized ERM has suboptimal dependence on r and log T. The spectral method has a sharper upper bound when $S = [T]$ (i.e. $\epsilon = 0$) with $\bar{\epsilon} = 0$, and is minimax optimal when the condition number $\frac{\sigma_{\max}((\bar{A}^{\perp})^{\top}B_{\delta}^{*})}{\sqrt{1+\bar{\epsilon}}}\$ $\frac{\sigma_{\max}((\mathbf{A}^{\top})^{\top} \mathbf{B}_{S}^{s})}{\sigma_{\min}((\mathbf{A}^{\top})^{\top} \mathbf{B}_{S}^{s})}$ is bounded and $n \geq n_0$.

Similar to the MTL case, we can extend the upper bound from the linear model to the GLMs and non-linear regression models.

Theorem 17 (Upper bound for TL under GLMs) Suppose Assumptions $\frac{1}{4}$, [5](#page-21-1) and other conditions imposed in Theorem [15](#page-45-0) hold, then the same high-probability upper bounds in Theorem [15](#page-45-0) hold for GLMs.

Theorem 18 (Upper bound for TL under non-linear regression models) Suppose Assumptions [6,](#page-22-1) [7,](#page-22-2) and other conditions imposed in Theorem [15](#page-45-0) hold. Let $n_0 \geq Cp^{3/2}$ with $C > 0$ a sufficiently large constant. The same high-probability upper bounds in Theorem [15](#page-45-0) hold for non-linear regression models.

Appendix C. General Lemmas

C.1 Lemmas

Lemma 19 (Theorem 6.5 in [Wainwright](#page-39-2) [\(2019\)](#page-39-2)) Suppose that Assumptions [1](#page-10-1) and [3](#page-10-2) hold. Then for any $\delta > 0$ and any $t \in [T]$, w.p. at least $1 - C_1 e^{-nC_2(\delta \wedge \delta^2)}$,

$$
\|\widehat{\mathbf{\Sigma}}^{(t)} - \mathbf{\Sigma}^{(t)}\|_2 \leq C_3 \sqrt{\frac{p}{n}} + \delta,
$$

with some constants $C_1, C_2, C_3 > 0$. Note that C_1, C_2, C_3 are universal in the sense that they do not depend on t or δ . As a consequence, we have

$$
\max_{t \in [T]} \|\widehat{\mathbf{\Sigma}}^{(t)} - \mathbf{\Sigma}^{(t)}\|_2 \lesssim \sqrt{\frac{p + \log T}{n}},
$$

w.p. at least $1 - e^{-C(p + \log T)}$.

Lemma 20 (A variant of Theorem 6.5 in [Wainwright](#page-39-2) [\(2019\)](#page-39-2)) Suppose that Assump-tions [9](#page-44-4) and [10](#page-44-2) hold. Then for any $\delta > 0$, for any fixed $A, B \in \mathcal{O}^{p \times r}$, w.p. at least $1 - C_1 e^{-n_0 C_2(\delta \wedge \delta)},$

$$
\|\mathbf{A}^\top \widehat{\mathbf{\Sigma}}^{(t)} \mathbf{B} - \mathbf{A}^\top \mathbf{\Sigma}^{(t)} \mathbf{B}\|_2 \leq C_3 \sqrt{\frac{r}{n_0}} + \delta,
$$

with some constants $C_1, C_2, C_3 > 0$. Note that C_1, C_2, C_3 are universal in the sense that they do not depend on δ or t. As a consequence, for any fixed $A, B \in \mathcal{O}^{p \times r}$, we have

$$
\|\mathbf{A}^\top \widehat{\boldsymbol{\Sigma}}^{(0)} \mathbf{B} - \mathbf{A}^\top \boldsymbol{\Sigma}^{(0)} \mathbf{B}\|_2 \lesssim \sqrt{\frac{r}{n_0}},
$$

w.p. at least $1 - e^{-C(r + \log T)}$.

Lemma 21 (Lemmas 2.5 and 2.6 in [Chen et al.](#page-35-1) [\(2021b\)](#page-35-1)) Suppose $p \geq r$. Consider two matrics $\mathbf{A}, \widetilde{\mathbf{A}} \in \mathcal{O}^{p \times r}$. Suppose that $(\mathbf{A}, \mathbf{A}^{\perp}), (\widetilde{\mathbf{A}}, \widetilde{\mathbf{A}}^{\perp}) \in \mathbb{R}^{p \times p}$ are both orthonormal matrices, which means that \bm{A}^\perp and $\widetilde{\bm{A}}^\perp$ are orthonormal complements of \bm{A} and $\widetilde{\bm{A}}$, respectively. Then

$$
\|AA^{\top} - \widetilde{A}(\widetilde{A})^{\top}\|_2 = \|A^{\top}(\widetilde{A}^{\perp})\|_2 = \|(\widetilde{A})^{\top}A^{\perp}\|_2,
$$

$$
\frac{1}{\sqrt{2}}\|AA^{\top} - \widetilde{A}(\widetilde{A})^{\top}\|_F = \|A^{\top}(\widetilde{A}^{\perp})\|_F = \|(\widetilde{A})^{\top}A^{\perp}\|_F,
$$

$$
\|AA^{\top} - \widetilde{A}(\widetilde{A})^{\top}\|_2 \le \min_{\mathbf{R} \in \mathcal{O}^{r \times r}} \|A - \widetilde{A}\mathbf{R}\|_2 \le \sqrt{2}\|AA^{\top} - \widetilde{A}(\widetilde{A})^{\top}\|_2,
$$

$$
\frac{1}{\sqrt{2}}\|AA^{\top} - \widetilde{A}(\widetilde{A})^{\top}\|_F \le \min_{\mathbf{R} \in \mathcal{O}^{r \times r}} \|A - \widetilde{A}\mathbf{R}\|_F \le \|AA^{\top} - \widetilde{A}(\widetilde{A})^{\top}\|_F.
$$

Lemma 22 (Proposition 8 in [Pajor](#page-38-0) [\(1998\)](#page-38-0)) When $p \geq 2r$, for any $\delta > 0$, we have

 (i) $(C1)$ √ r $\frac{\sqrt{r}}{\delta}$)^{r(p-r)} $\leq N(\mathcal{O}^{p\times r}, \text{dist}_{\text{F}}, \delta) \leq C_2(\delta)$ √ r $\frac{\sqrt{r}}{8}$ $)^{r(p-r)}$; (*ii*) $(C_1\frac{1}{\delta})$ $\frac{1}{\delta})^{r(p-r)} \le N(\mathcal{O}^{p \times r}, \text{dist}_2, \delta) \le (C_2 \frac{1}{\delta})$ $\frac{1}{\delta})^{r(p-r)}$;

(iii) (Consequence of (i)) $M(\mathcal{O}^{p\times r}, \text{dist}_{F}, \delta) \geq C(\delta)$ √ r $\frac{\pi}{\delta}$)^{r(p-r)}.

Lemma 23 When $p \ge 2r$, for any $\delta > 0$ and $\alpha \in (0, 1)$, $\exists \overline{A} \in \mathcal{O}^{p \times r}$, such that

$$
M(\mathcal{B}_{\delta}(\overline{\mathbf{A}}, \mathcal{O}^{p\times r}, \text{dist}_2), \text{dist}_{F}, \alpha\delta) \ge \left(\frac{C\sqrt{r}}{\alpha}\right)^{r(p-r)}
$$

.

Lemma 24 (Fano's lemma, see [Tsybakov](#page-39-3) [\(2009\)](#page-39-3) and [Wainwright](#page-39-2) [\(2019\)](#page-39-2)) Suppose (Θ, d) is a metric space and each θ in this space is associated with a probability measure \mathbb{P}_{θ} . If $\{\theta_j\}_{j=1}^N$ is an s-separated set (i.e. $d(\theta_j, \theta_k) \geq s$ for any $j \neq k$), and $KL(\mathbb{P}_{\theta_j}, \mathbb{P}_{\theta_k}) \leq \alpha \log N$, then

$$
\inf_{\widehat{\theta}} \sup_{\theta \in \Theta} \mathbb{P}_{\theta}(d(\widehat{\theta}, \theta) \ge s/2) \ge 1 - \alpha - \frac{\log 2}{\log N}.
$$

Lemma 25 Consider the following generative model:

$$
y|\boldsymbol{x} \sim \mathbb{P}_{y|\boldsymbol{x},\boldsymbol{\beta}} = N(\boldsymbol{x}^\top \boldsymbol{\beta}, 1), \quad \boldsymbol{x} \sim \mathbb{P}_{\boldsymbol{x}},
$$

where \mathbb{P}_x is sub-Gaussian with $\Sigma = \mathbb{E}(xx^\top)$. Suppose there exist constants c and C such that $0 < c \leq \lambda_{\min}(\Sigma) \leq \lambda_{\max}(\Sigma) \leq C < \infty$. Define two distributions $\mathbb P$ and $\widetilde{\mathbb P}$ of (x, y) as $\mathbb{P}_{x,y} = \mathbb{P}_{y|x;\beta} \cdot \mathbb{P}_x$ and $\widetilde{\mathbb{P}}_{x,y} = \mathbb{P}_{y|x;\widetilde{\beta}} \cdot \mathbb{P}_x$. Then, the KL divergence between $\mathbb{P}_{x,y}$ and $\widetilde{\mathbb{P}}_{x,y}$ can be bounded as

$$
\mathrm{KL}(\mathbb{P}_{\boldsymbol{x},y} \| \widetilde{\mathbb{P}}_{\boldsymbol{x},y}) \leq C \|\boldsymbol{\beta} - \widetilde{\boldsymbol{\beta}}\|_2^2.
$$

Lemma 26 (Theorem 5.1 in [Chen et al.](#page-35-2) [\(2018\)](#page-35-2)) Given a family of distributions $\{\mathbb{P}_{\theta}$: $\theta \in \Theta$, which is indexed by a parameter $\theta \in \Theta$. Consider $\mathbf{x}^{(t)} \sim (1-\epsilon')\mathbb{P}_{\theta}+\epsilon'\mathbb{Q}$ independently for $t \in [T]$, and $\widetilde{\mathbf{x}} \sim \mathbb{P}_{\widetilde{\mathbf{x}}}$. Denote the joint distribution of $\{\mathbf{x}^{(t)}\}_{t=1}^T$ and \widetilde{x} as $\mathbb{P}_{(\epsilon', \theta, \mathbb{Q})} \cdot \mathbb{P}_{\widetilde{\mathbf{x}}}$.
Then Then

$$
\inf_{\widehat{\theta}} \sup_{\theta \in \Theta} (\mathbb{P}_{(\epsilon', \theta, \mathbb{Q})} \cdot \mathbb{P}_{\widetilde{\mathbf{x}}}) \left(\|\widehat{\theta} - \theta\| \geq C \varpi(\epsilon', \Theta) \right) \geq \frac{1}{2},
$$

where $\varpi(\epsilon', \Theta) \coloneqq \sup \{ \|\theta_1 - \theta_2\| : TV(\mathbb{P}_{\theta_1}, \mathbb{P}_{\theta_2}) \leq \epsilon'/(1-\epsilon'), \theta_1, \theta_2 \in \Theta \}.$

Lemma 27 (Lemma 22 in [Tian et al.](#page-39-4) [\(2022\)](#page-39-4)) Consider two data generating mechanisms:

- (i) $\mathbf{x}^{(t)} \sim (1-\epsilon') \mathbb{P}_{\theta} + \epsilon' \mathbb{Q}$ independently for $t \in [T]$, where $\epsilon' = \frac{|S^c|}{T}$ $\frac{S^c}{T}$, and $\widetilde{x} \sim \mathbb{P}_{\widetilde{x}}$;
- (ii) With a preserved set $S \subseteq [T]$, generate $\{\boldsymbol{x}^{(t)}\}_{t \in S^c} \sim \mathbb{Q}_{S^c}$ and $\boldsymbol{x}^{(t)} \sim \mathbb{P}_{\theta}$ independently for $t \in S$, and $\widetilde{x} \sim \mathbb{P}_{\widetilde{x}}$.

Denote the joint distributions of $\{\boldsymbol{x}^{(t)}\}_{t=1}^T$ and \tilde{x} in (i) and (ii) as $\mathbb{P}_{(\epsilon,\theta,\mathbb{Q})} \cdot \mathbb{P}_{\tilde{x}}$ and $\mathbb{P}_{(S,\theta,\mathbb{Q}_{S^c})} \cdot \mathbb{P}_{\tilde{x}}$, respectively. We claim that if

$$
\inf_{\widehat{\theta}} \sup_{\substack{\theta \in \Theta \\ \mathbb{Q}}} (\mathbb{P}_{(\epsilon'/50,\theta,\mathbb{Q})} \cdot \mathbb{P}_{\widetilde{\boldsymbol{x}}}) \left(\|\widehat{\theta} - \theta\| \ge C \varpi(\epsilon'/50,\Theta) \right) \ge \frac{1}{2}
$$

then

$$
\inf_{\widehat{\theta}} \sup_{S:|S|\geq T(1-\epsilon')} \sup_{\substack{\theta\in\Theta\\ \mathbb{Q}_{S^c}}} (\mathbb{P}_{(S,\theta,\mathbb{Q}_{S^c})}\cdot \mathbb{P}_{\widetilde{\boldsymbol{x}}})\left(\|\widehat{\theta}-\theta\| \geq C\varpi\left(\epsilon'/50,\Theta\right)\right) \geq \frac{1}{10},
$$

where $\varpi(\epsilon', \Theta) \coloneqq \sup \{ \|\theta_1 - \theta_2\| : TV(\mathbb{P}_{\theta_1}, \mathbb{P}_{\theta_2}) \leq \epsilon'/(1-\epsilon'), \theta_1, \theta_2 \in \Theta \}.$

Lemma 28 (Lemma 33 in [Tian et al.](#page-39-4) [\(2022\)](#page-39-4)) Given a family of distributions $\{\mathbb{P}_{\theta}$: $\theta \in \Theta$, which is indexed by a parameter $\theta \in \Theta$. Consider $\mathbf{x}^{(t)} \sim (1-\epsilon')\mathbb{P}_{\theta}+\epsilon'\mathbb{Q}$ independently for $t \in [T]$, and $\widetilde{x} \sim \mathbb{P}_{\widetilde{x}}$. Consider another family of distributions $\{\mathbb{P}_{\theta}^{(0)}\}$ $\theta_{\theta}^{(0)}$: $\theta \in \Theta$ } indexed by the same parameter set, and $\mathbf{x}^{(0)} \sim \mathbb{P}_{\theta}^{(0)}$ $\mathcal{L}_{\theta}^{(0)}$. Denote the joint distribution of $\{\boldsymbol{x}^{(t)}\}_{t=0}^T$ and $\widetilde{\boldsymbol{x}}$ as $\mathbb{P}_{(\epsilon',\theta,\mathbb{Q})}\cdot\mathbb{P}_{\widetilde{\boldsymbol{x}}}\$. Then

$$
\inf_{\widehat{\theta}}\sup_{\theta\in\Theta}(\mathbb{P}_{(\epsilon',\theta,\mathbb{Q})}\cdot\mathbb{P}_{\widetilde{\boldsymbol{x}}})\left(\|\widehat{\theta}-\theta\|\geq C\varpi(\epsilon',\Theta)\right)\geq\frac{9}{20},
$$

where $\varpi(\epsilon', \Theta) \coloneqq \sup \left\{ \|\theta_1 - \theta_2\| : TV\left(\mathbb{P}_{\theta_1}, \mathbb{P}_{\theta_2}\right) \leq \epsilon'/(1-\epsilon'), TV\left(\mathbb{P}_{\theta_1}^{(0)}\right) \right\}$ $\overset{(0)}{\theta_1}, \overset{\mathbb{P}(0)}{\theta_2}$ $\binom{0}{\theta_2} \leq 1/20, \theta_1, \theta_2 \in$ Θ .

Lemma 29 (Lemma 34 in [Tian et al.](#page-39-4) (2022)) Consider two data generating mechanisms:

- (i) $\mathbf{x}^{(t)} \sim (1-\epsilon') \mathbb{P}_{\theta} + \epsilon' \mathbb{Q}$ independently for $t \in [T]$, $\mathbf{x}^{(0)} \sim \mathbb{P}_{\theta}^{(0)}$ $_{\theta}^{(0)}$, and $\widetilde{x} \sim \mathbb{P}_{\widetilde{\bm{x}}}$, where $\epsilon' = \frac{|S^c|}{T}$ $\frac{S^{\circ}|}{T};$
- (ii) With a preserved set $S \subseteq [T]$, generate $\{\boldsymbol{x}^{(t)}\}_{t \in S^c} \sim \mathbb{Q}_{S^c}$ and $\boldsymbol{x}^{(t)} \sim \mathbb{P}_{\theta}$ independently for $t \in S$, $\boldsymbol{x}^{(0)} \sim \mathbb{P}_{\theta}^{(0)}$ $_{\theta}^{(0)}$, and $\widetilde{x} \sim \mathbb{P}_{\widetilde{\boldsymbol{x}}}.$

Denote the joint distributions of $\{\boldsymbol{x}^{(t)}\}_{t=0}^T$ and \tilde{x} in (i) and (ii) as $\mathbb{P}_{(\epsilon,\theta,\mathbb{Q})} \cdot \mathbb{P}_{\tilde{x}}$ and $\mathbb{P}_{(S,\theta,\mathbb{Q}_{S^c})} \cdot \mathbb{P}_{\tilde{x}}$, respectively. We claim that if

$$
\inf_{\widehat{\theta}}\sup_{\theta\in\Theta}(\mathbb{P}_{(\epsilon'/50,\theta,\mathbb{Q})}\cdot\mathbb{P}_{\widetilde{\bm{x}}})\left(\|\widehat{\theta}-\theta\|\geq C\varpi'\left(\epsilon'/50,\Theta\right)\right)\geq\frac{9}{20},
$$

then

$$
\inf_{\widehat{\theta}} \sup_{S:|S|\geq T(1-\epsilon')}\sup_{\substack{\theta\in\Theta\\ \mathbb{Q}_S}} (\mathbb{P}_{(S,\theta,\mathbb{Q}_S)}\cdot\mathbb{P}_{\widetilde{\boldsymbol{x}}})\left(\|\widehat{\theta}-\theta\|\geq C\varpi'\left(\epsilon'/50,\Theta\right)\right)\geq \frac{1}{10},
$$

where $\varpi(\epsilon', \Theta) \coloneqq \sup \left\{ \|\theta_1 - \theta_2\| : TV\left(\mathbb{P}_{\theta_1} \| \mathbb{P}_{\theta_2}\right) \leq \epsilon'/(1-\epsilon'), TV\left(\mathbb{P}_{\theta_1}^{(0)}\right) \right\}$ $\begin{array}{c} \left(0\right)\parallel \mathbb{P}_{\theta_2}^{\left(0\right)} \end{array}$ $\binom{00}{\theta_2} \leq 1/20, \theta_1, \theta_2 \in$ Θ .

C.2 Proofs of Lemmas

Denote $\widehat{\mathbf{\Sigma}}^{(t)} = \sum_{i=1}^{n} \frac{1}{n}$ $\frac{1}{n}\boldsymbol{x}_{i}^{\left(t\right)}$ $\binom{t}{i}(\bm{x}^{(t)}_i$ $\binom{(t)}{i}$ ^T.

C.2.1 Proof of Lemma [20](#page-47-2)

Note that for any fixed $\mathbf{A}, \mathbf{B} \in \mathcal{O}^{p \times r}$,

$$
\|\boldsymbol A^\top(\widehat{\boldsymbol \Sigma}^{(0)}-\boldsymbol \Sigma^{(0)})\boldsymbol B\|_2=\sup_{\|\boldsymbol u\|_2,\|\boldsymbol v\|_2\leq 1}(\boldsymbol{A}\boldsymbol u)^\top(\widehat{\boldsymbol \Sigma}^{(0)}-\boldsymbol \Sigma^{(0)})(\boldsymbol{B}\boldsymbol v),
$$

where $Au, Bv \in \mathbb{R}^p$ and $||Au||_2 = ||Bv||_2 = 1$. Note that both Au and Bv live in r-dimensional space (isomorphic to the unit ball in \mathbb{R}^r). Therefore, by Example 5.8 in

[Wainwright](#page-39-2) (2019) , there exist two 1/8-covers (whose components are inside the set to be covered) of $\{\beta \in \mathbb{R} : \beta = Au, ||u||_2 \le 1\}$ and $\{\beta \in \mathbb{R} : \beta = Bu, ||u||_2 \le 1\}$ under Euclidean norm, denoted as $\{\beta_j^u\}_{j=1}^{N_1}$ and $\{\beta_j^u\}_{j=1}^{N_2}$, respectively, such that $|N_1| = |N_2| \le 17^r$. Then we can proceed in the same steps as in the proof of Theorem 6.5 in [Wainwright](#page-39-2) [\(2019\)](#page-39-2) to finish the proof.

C.2.2 Proof of Lemma [23](#page-48-0)

We apply the same trick as in the proof of Proposition 3 in [Cai et al.](#page-35-3) [\(2013\)](#page-35-3). Suppose the minimum cover of $\mathcal{O}^{p\times r}$ under $dist_2(\cdot)$ corresponding to $N(\mathcal{O}^{p\times r}, dist_2, \delta)$ is N. By pigeonhole theorem, for any $\alpha \in (0,1)$,

$$
N(\mathcal{O}^{p\times r},\text{dist}_2,\alpha\delta) \leq N(\cup_{\mathbf{A}\in\mathcal{N}}\mathcal{B}_{\delta}(\mathbf{A},\mathcal{O}^{p\times r},\text{dist}_2),\text{dist}_{F},\alpha\delta)
$$

\$\leq |\mathcal{N}| \max_{\mathbf{A}\in\mathcal{N}} N(\mathcal{B}_{\delta}(\mathbf{A},\mathcal{O}^{p\times r},\text{dist}_2),\text{dist}_{F},\alpha\delta).

Then applying Lemma [22](#page-47-3) leads to

$$
\max_{\mathbf{A}\in\mathcal{N}} N(\mathcal{B}_{\delta}(\mathbf{A}, \mathcal{O}^{p\times r}, \text{dist}_2), \text{dist}_{\text{F}}, \alpha\delta) \ge \frac{N(\mathcal{O}^{p\times r}, \text{dist}_2, \alpha\delta)}{N(\mathcal{O}^{p\times r}, \text{dist}_2, \delta)} \ge \left(\frac{C\sqrt{r}}{\alpha}\right)^{r(p-r)}
$$

.

Since N has only finite elements, there must exist one $A \in \mathcal{N}$ achieving the maximum of LHS. Because the packing number is always larger than or equal to the covering number, the proof is done.

C.2.3 Proof of Lemma [25](#page-48-1)

By the form of Gaussian density function, it is straightfoward to see that

$$
\begin{split} \text{KL}(\mathbb{P}_{\bm{x},y} \| \widetilde{\mathbb{P}}_{\bm{x},y}) &= \int \log \left(\frac{\mathbb{P}_{y | \bm{x}, \bm{\beta}}}{\mathbb{P}_{y | \bm{x}, \widetilde{\bm{\beta}}}} \right) d\mathbb{P}_{y | \bm{x}, \bm{\beta}} d\mathbb{P}_{\bm{x}} \\ & \leq \underbrace{\int (y - \bm{x}^{\top} \bm{\beta}) \cdot \bm{x}^{\top} (\bm{\beta} - \widetilde{\bm{\beta}}) d\mathbb{P}_{y | \bm{x}, \bm{\beta}} d\mathbb{P}_{\bm{x}}}_{=0} + \int [\bm{x}^{\top} (\bm{\beta} - \widetilde{\bm{\beta}})]^{2} d\mathbb{P}_{\bm{x}} \\ & \lesssim \|\bm{\beta} - \widetilde{\bm{\beta}}\|_{2}^{2} . \end{split}
$$

Appendix D. Proofs for Linear Regression Models

Denote $\theta_A^{(t)} \in \arg \min_{\theta \in \mathbb{R}^r} f^{(t)}(A\theta)$ and $f^{(t)}(\beta) = \frac{1}{n} ||Y^{(t)} - X^{(t)}\beta||_2^2$. For $A, \widetilde{A} \in \mathcal{O}^{p \times r}$, define metrics $dist_2(A, \widetilde{A}) = ||AA^\top - \widetilde{A}(\widetilde{A})^\top||_2$, $dist_F(A, \widetilde{A}) = ||AA^\top - \widetilde{A}(\widetilde{A})^\top||_F$. In any metric space (\mathcal{X}, ρ) , denote the ball of radius δ with center x under metric ρ as $\mathcal{B}_{\delta}(x, \mathcal{X}, \rho)$. Denote $\zeta^{(t)} = \|\boldsymbol{\theta}^{(t)*}\|_2 \vee \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$ and $\bar{\zeta} = \sqrt{|S|^{-1} \sum_{t \in S} (\zeta^{(t)})^2}$.

D.1 Lemmas

Lemma [3](#page-10-2)0 Suppose Assumptions [1-](#page-10-1)3 hold. Then w.p. at least $1 - e^{-C(r + \log T)}$, for all $t \in [T]$ and $\mathbf{A} \in \mathcal{O}^{p \times r}$,

$$
\|\boldsymbol A\boldsymbol \theta^{(t)}_{\boldsymbol A} - \boldsymbol A^{(t)*}\boldsymbol \theta^{(t)*}\|_2 \lesssim \|\boldsymbol A\boldsymbol A^\top - \boldsymbol A^{(t)*}(\boldsymbol A^{(t)*})^\top\|_2\cdot \zeta^{(t)} + \sqrt{\frac{r+\log T}{n}}.
$$

Lemma [3](#page-10-2)1 Suppose Assumptions [1-](#page-10-1)3 hold. Then w.p. at least $1 - e^{-C(r + \log T)}$, for all $t \in S$ and $A \in \mathcal{O}^{p \times r}$, there exists a universal constant $C' > 0$ (which does not depend on t or **A**), such that $||A\theta_A^{(t)}||_2 = ||\theta_A^{(t)}||_2 \leq C'\zeta^{(t)}$.

Lemma [3](#page-10-2)2 Suppose Assumptions [1-](#page-10-1)3 hold. Then w.p. at least $1 - e^{-C(r + \log T)}$, for all $t \in [T]$ and $\mathbf{A}, \widetilde{\mathbf{A}} \in \mathcal{O}^{p \times r}$,

$$
\|\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}-\widetilde{\boldsymbol{A}}\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)}\|_{2}\lesssim\|\boldsymbol{A}\boldsymbol{A}^{\top}-\widetilde{\boldsymbol{A}}\widetilde{\boldsymbol{A}}^{\top}\|_{2}(\|\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)}\|_{2}+\|\nabla f^{(t)}(\widetilde{\boldsymbol{A}}\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)})\|_{2}).
$$

Lemma [3](#page-10-2)3 Suppose Assumptions [1-](#page-10-1)3 hold. For any $\overline{A} \in \arg \min_{\overline{A} \in \mathcal{O}^{p \times r}} \max_{t \in S} {\{\Vert A^{(t)}(A^{(t)})^\top \overline{\mathbf{A}}\overline{\mathbf{A}}^{\top} \|_{2}$, denote $\overline{\mathbf{G}} = \left\{ \nabla f^{(t)}(\overline{\mathbf{A}}\boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)}) \right\}_{t \in S}$, then w.p. at least $1 - e^{-C(p + \log T)}$,

$$
\|\overline{G}\|_2 \lesssim \sqrt{\frac{p}{n}} + \sqrt{T}\left(h\overline{\zeta} + \sqrt{\frac{r + \log T}{n}}\right).
$$

Lemma 34 Suppose $\bm{G}=\{\bm{g}_j\}_{j=1}^d\in\mathbb{R}^{p\times d}$ and $\bm{\overline{G}}=\{\bm{\overline{g}}_j\}_{j=1}^d\in\mathbb{R}^{p\times d}$ with each $\bm{g}_j,\bm{\overline{g}}_j\in\mathbb{R}^p$ and $\max_{j \in [d]} ||g_j - \overline{g}_j||_2 \leq \Delta$. Then

$$
\|\overline{G}-G\|_2\leq \sqrt{d}\Delta.
$$

Lemma [3](#page-10-2)5 Suppose Assumptions [1-](#page-10-1)3 hold. Consider any $\overline{A} \in \mathcal{O}^{p \times r}$.

- (i) When $\frac{\lambda}{\sqrt{2}}$ $\frac{\Delta_n}{\overline{n}} \geq C \zeta^{(t)} \Big[\| \widehat{\overline{A}} \boldsymbol{\theta}_{\widehat{\overline{A}}}^{(t)} \|$ $-\bm{A}^{(t)*}\bm{\theta}^{(t)*}\|_2 + \|\nabla f^{(t)}(\bm{A}^{(t)*}\bm{\theta}^{(t)*})\|_2 \Big]$ with a sufficiently large C and the minimizer $(\mathbf{A}^{(t)}, \boldsymbol{\theta}^{(t)}) = \arg \min_{\mathbf{A} \in \mathcal{O}^{p \times r}, \boldsymbol{\theta} \in \mathbb{R}^r} \{\frac{1}{T}\}$ $\frac{1}{T}f^{(t)}(\bm{A}\bm{\theta})+$ $\frac{\sqrt{n}}{nT} \lambda \| \bm{A} \bm{A}^\top \overline{A}(\overline{A})^{\top} \|_{2}$, we must have $A^{(t)}(A^{(t)})^{\top} = \overline{A}(\overline{A})^{\top}$.
- (ii) As a consequence, by Lemma [30,](#page-50-4) when $\frac{\lambda}{\sqrt{n}}$ $\frac{\lambda}{\overline{n}} \geq C \zeta^{(t)} \Big[\zeta^{(t)} \| \widehat{\overline{\bm{A}}} (\widehat{\overline{\bm{A}}})^\top - \bm{A}^{(t)*} (\bm{A}^{(t)*})^\top \|_2 + \cdots$ $\sqrt{p + \log T}$ $\left[\frac{\log T}{n}\right]$, w.p. at least $1 - e^{-C(r + \log T)}$, we have $\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top} = \widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^{\top}$.

Lemma 36 $\,Suppose\,\,\{\boldsymbol{\theta}^{(t)}\}_{t\in S}\subseteq\mathbb{R}^r\,\, satisfying \,\, |S|^{-1}\sum_{t\in S}\boldsymbol{\theta}^{(t)}(\boldsymbol{\theta}^{(t)})^\top\succeq\frac{c}{r}$ $\frac{c}{r}\bar{\zeta}I_r$, where c is a positive constant. For any subset $S' \subseteq S$ with $|S'| \geq (1 - \alpha)|S|$ and

$$
\alpha \leq \frac{c}{r} \cdot \frac{\bar{\zeta}^2}{\max_{t \in S}(\zeta^{(t)})^2},
$$

where c is the same positive constant above and $\zeta^{(t)} = ||\boldsymbol{\theta}^{(t)}||_2$, we have

$$
\frac{1}{|S'|}\sum_{t\in S'}\boldsymbol{\theta}^{(t)}(\boldsymbol{\theta}^{(t)})^{\top} \succeq \frac{c}{2r}\bar{\zeta}^2\boldsymbol{I}_r.
$$

Lemma 37 Under Assumptions [1-](#page-10-1)[3,](#page-10-2) w.p. at least $1 - e^{-C(p + \log T)}$, for all $t \in S$,

$$
\|\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \lesssim \left(\frac{\lambda}{\sqrt{n}}\right)^{1/2} + \sqrt{\frac{p + \log T}{n}}
$$

.

Lemma 38 Consider the case that $r = 1$. Suppose that $\min_{t \in S} |\theta^{(t)*}| \ge c > 0$ with some constant $c > 0$. Under Assumptions [1-](#page-10-1)[3,](#page-10-2) w.p. at least $1 - e^{-\tilde{C}'(1 + \log T)}$, for all $A \in \mathcal{O}^{p \times r}$ and $t \in S$,

$$
\|\boldsymbol A\boldsymbol \theta^{(t)}_{\boldsymbol A} - \boldsymbol A^{(t)*}\boldsymbol \theta^{(t)*}\|_2 \geq C'\zeta^{(t)}\|\boldsymbol A\boldsymbol A^\top - \boldsymbol A^{(t)*}(\boldsymbol A^{(t)*})^\top\|_2 - C\sqrt{\frac{p+\log T}{n}}.
$$

Lemma 39 (For MTL) Under Assumptions [1](#page-10-1)[-3,](#page-10-2) we have:

- (i) For all $t \in S$, when $\frac{\gamma}{\sqrt{n}} \geq \|\nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_2 + C\|\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2$, it holds that $\widehat{\boldsymbol{\beta}}^{(t)} = \widehat{\boldsymbol{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}$ w.p. at least $1 - e^{-C(p + \log T)}$;
- (ii) $\max_{t \in S} \|\widehat{\beta}^{(t)} \beta^{(t)*}\|_2 \leq C \frac{\gamma}{\sqrt{n}} + \max_{t \in S} \|\widetilde{\beta}^{(t)} \beta^{(t)*}\|_2 \ w.p.$ at least $1 e^{-C(p + \log T)}$, where $\widetilde{\beta}^{(t)} \in \arg \min_{\beta \in \mathbb{R}^p} f^{(t)}(\beta);$
- (iii) If the data from tasks in S^c satisfies the linear model [\(3\)](#page-8-1) (without any latent structure assumption) and Assumption [1,](#page-10-1) then w.p. at least $1 - e^{-C(p + \log T)}$, $\max_{e \in S^c} ||\widehat{\boldsymbol{\beta}}^{(t)} \boldsymbol{\beta}^{(t)*} \|_2 \leq C \frac{\gamma}{\sqrt{n}} + \max_{t \in S^c} \| \widetilde{\boldsymbol{\beta}}^{(t)} - \boldsymbol{\beta}^{(t)*} \|_2, \ where \ \widetilde{\boldsymbol{\beta}}^{(t)} \in \arg \min_{\boldsymbol{\beta} \in \mathbb{R}^p} f^{(t)}(\boldsymbol{\beta}).$

Lemma 40 Suppose Assumption [9](#page-44-4) hold and $n \ge Cr$. Then for **any fixed** $A \in \mathcal{O}^{p \times r}$, w.p. at least $1 - e^{-C(r + \log T)}$,

$$
\|\bm{A}\bm{\theta}^{(0)}_{\bm{A}}-\bm{A}^{(0)*}\bm{\theta}^{(0)*}\|_2 \lesssim \|\bm{A}\bm{A}^\top - \bm{A}^{(0)*}(\bm{A}^{(0)*})^\top\|_2 + \sqrt{\frac{r}{n}}.
$$

Lemma 41 Suppose Assumptions [9-](#page-44-4)[10](#page-44-2) hold. Then w.p. at least $1 - e^{-C(r + \log T)}$, for all $A \in \mathcal{O}^{p \times r}$,

$$
\|\boldsymbol{A}\boldsymbol{\theta}^{(0)}_{\boldsymbol{A}}-\boldsymbol{A}^{(0)*}\boldsymbol{\theta}^{(0)*}\|_2 \lesssim \|\boldsymbol{A}\boldsymbol{A}^{\top}-\boldsymbol{A}^{(0)*}(\boldsymbol{A}^{(0)*})^{\top}\|_2 + \sqrt{\frac{r}{n}}.
$$

Lemma 42 (For TL) Under Assumptions [9-](#page-44-4)[10,](#page-44-2) we have:

- (i) When $\frac{\gamma}{\sqrt{n_0}} \geq \|\nabla f^{(t)}(A^{(0)*}\theta^{(0)*})\|_2 + C \|\widehat{\Sigma}^{(0)}\|_2 \cdot \|\widehat{\overline{A}}\theta_{\widehat{\overline{A}}}^{(0)}\|_2$ $- A^{(0)*}\theta^{(0)*}\|_2$, it holds that $\widehat{\boldsymbol{\beta}}^{(0)}=\widehat{\bar{\boldsymbol{A}}}\boldsymbol{\theta}^{(0)}_{\widehat{\bar{\boldsymbol{A}}}}$ w.p. at least $1 - e^{-C(p + \log T)}$; [i](#page-52-7)
- (iii) $\|\widehat{\beta}^{(0)} \beta^{(0)*}\|_2 \leq C \frac{\gamma}{\sqrt{n_0}} + \|\widetilde{\beta}^{(0)} \beta^{(0)*}\|_2$ w.p. at least $1 e^{-C(p + \log T)}$, where $\widetilde{\beta}^{(0)} \in$ $\arg\min_{\boldsymbol{\beta}\in\mathbb{R}^p} f^{(0)}(\boldsymbol{\beta}).$

D.2 Proof of Theorem [1](#page-12-0)

First, we have the following proposition holds.

Proposition 43 Suppose Assumptions [1](#page-10-1)[-3](#page-10-2) hold. Further assume

(i)
$$
\min_{t \in S} \left\{ \frac{\zeta^{(t)}}{\zeta} r \sqrt{\frac{p + \log T}{nT}} + \frac{\zeta^{(t)}}{\zeta} \sqrt{r} \sqrt{\frac{r + \log T}{n}} + \zeta^{(t)} \sqrt{r} h \right\} \lesssim \sqrt{\frac{p + \log T}{n}};
$$

i. For this result to hold, it suffices to require $n_0 \geq Cr$, which is important for Remark [14](#page-44-5) to be true.

$$
(ii) |Sc|/T \le cr^{-3/2} \frac{\bar{\zeta}}{\max_{t \in S} \zeta^{(t)}} with a small constant $c > 0$.
$$

Let $\lambda \geq C \frac{\max_{t \in S} (\zeta^{(t)})^2}{\min_{x \in \zeta(t)}}$ $\frac{\max_{t \in S} (\zeta^{(t)})^2}{\min_{t \in S} \zeta^{(t)}} r^{3/4} \sqrt{(p + \log T)}$ with a sufficiently large constant C. Then w.p. at $least\ 1-e^{-C'(r+\log T)},$

$$
\|\widehat{\overline{\boldsymbol{A}}}(\widehat{\overline{\boldsymbol{A}}})^\top - \overline{\boldsymbol{A}}\overline{\boldsymbol{A}}^\top \|_2 \lesssim r \bar{\zeta}^{-1} \sqrt{\frac{p}{nT}} + \sqrt{r} h + \bar{\zeta}^{-1} \sqrt{r} \sqrt{\frac{r + \log T}{n}} + \frac{\lambda r}{\sqrt{n} \bar{\zeta}^2} \epsilon.
$$

Proposition [43](#page-52-6) provides the bound when the first term is faster than the second term in the upper bound of Theorem [1.](#page-12-0) In the other case, a direct application of Lemma [39](#page-52-2) leads to the second term in the upper bound. Combining two situations gives us the disired result for tasks in S.

Denote $\eta^{(t)} = \frac{\zeta^{(t)}}{\bar{z}}$ $\frac{f^{(t)}}{\bar{\zeta}}r\sqrt{\frac{p+\log T}{nT}}+\frac{\zeta^{(t)}}{\bar{\zeta}}$ ζ¯ $\sqrt{r} \sqrt{\frac{r + \log T}{n}} + \zeta^{(t)} \sqrt{r} h + \frac{\lambda r \zeta^{(t)}}{\sqrt{n} \overline{\zeta}^2} \epsilon.$

(i) For any t satisfying $\eta^{(t)} \leq C \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$: By Proposition [43,](#page-52-6) w.p. at least $1 - e^{-C'(r + \log T)}$,

$$
\|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - A^{(t)*}(\mathbf{A}^{(t)*})^{\top}\|_2 \lesssim r\bar{\zeta}^{-1}\sqrt{\frac{p}{nT}} + \sqrt{r}h + \bar{\zeta}^{-1}\sqrt{r}\sqrt{\frac{r+\log T}{n}} + \frac{\lambda r}{\sqrt{n}\bar{\zeta}^2}\epsilon.
$$

By Lemma [35,](#page-51-4) since $\frac{\lambda}{\sqrt{2}}$ $\frac{\lambda}{\overline{n}} \geq C \frac{\max_{t \in S} (\zeta^{(t)})^2}{\min_{t \in S} \zeta^{(t)}}$ $\frac{\max_{t\in S}(\zeta^{(t)})^2}{\min_{t\in S}\zeta^{(t)}}r^{3/4}\sqrt{\frac{(p+\log T)}{n}}\geq C\max_{t\in S}\zeta^{(t)}\eta^{(t)}\geq C\zeta^{(t)}\Big[\zeta^{(t)}\|\widehat{\overline{\pmb{A}}}(\widehat{\overline{\pmb{A}}})^\top-\zeta^{(t)}\|\widehat{\overline{\pmb{A}}}\|\widehat{\overline{\pmb{A}}}\|\widehat{\overline{\pmb{A}}}\|\widehat{\overline{\pmb{A}}}\|\widehat{\overline{\pmb{A}}}\|\widehat{\overline{\pmb{A}}}\|\widehat{\overline{\pmb{$ $\boldsymbol{A}^{(t)*} (\boldsymbol{A}^{(t)*})^\top \|_2 + \sqrt{\frac{p+\log T}{n}}$ $\left[\frac{\log T}{n}\right]$, we have $\widehat{A}^{(t)}(\widehat{A}^{(t)})^{\top} = \widehat{\overline{A}}(\widehat{\overline{A}})^{\top}$ w.p. at least $1 - e^{-C'(r + \log T)}$, which combining with Lemma [30](#page-50-4) implies that

$$
\begin{aligned}\n\|\widehat{A}^{(t)}\widehat{\theta}^{(t)} - A^{(t)*}\theta^{(t)*}\|_{2} &= \|\widehat{\overline{A}}\theta_{\widehat{\overline{A}}}^{(t)} - A^{(t)*}\theta^{(t)*}\|_{2} \\
&\lesssim \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - A^{(t)*}(A^{(t)*})^{\top}\|_{2}\zeta^{(t)} + \sqrt{\frac{r + \log T}{n}} \\
&\lesssim \eta^{(t)} + \sqrt{\frac{r + \log T}{n}},\n\end{aligned}
$$

w.p. at least $1 - e^{-C'(r + \log T)}$. Therefore by our choice $\gamma = C' \sqrt{p + \log T}$ with a large constant $C' > 0$:

$$
\|\nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_2 + C\|\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \lesssim \frac{\gamma}{\sqrt{n}}
$$

.

Then by Lemma $\frac{39}{\sqrt{1-\lambda}}$ and Proposition [43,](#page-52-6) w.p. at least $1-e^{-C'(r+\log T)}$, $\max_{t\in S}\|\hat{\beta}^{(t)}-t\|_2^2$ $\boldsymbol{\beta}^{(t)*} \|_2 \lesssim \eta^{(t)} + \sqrt{\frac{r + \log T}{n}}$ $\frac{\log T}{n}$. (ii) For any t satisfying $\eta^{(t)} > C \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$: by Lemma [39.](#page-52-2)(ii), w.p. at least $1 - e^{-C'(p + \log T)}$, $\max_{t\in S} \|\widehat{\boldsymbol{\beta}}^{(t)}-\boldsymbol{\beta}^{(t)\ast}\|_2 \lesssim \frac{\gamma}{\sqrt{n}} + \max_{t\in S} \|\widetilde{\boldsymbol{\beta}}^{(t)}-\boldsymbol{\beta}^{(t)\ast}\|_2 \lesssim \sqrt{\frac{p+\log T}{n}}$ $\frac{\log T}{n}$, where $\widetilde{\beta}^{(t)} \in \arg \min_{\beta \in \mathbb{R}^p} f^{(t)}(\beta)$. The fact that $\max_{t \in S} \|\widetilde{\beta}^{(t)} - \beta^{(t)*}\|_2 \lesssim \sqrt{\frac{p + \log T}{n}}$ w.p. at least $1 - e^{-C'(p + \log T)}$ is the standard linear regression result.

(iii) When data from tasks in S^c also satisfies the linear model [\(3\)](#page-8-1), the result comes from the same argument as in (ii).

D.3 Proof of Theorem [2](#page-13-0)

The proof essentially depends on Lemmas [37](#page-51-6) and [38.](#page-52-1) The conclusion of Lemma [37](#page-51-6) is pretty weak, which only provides a slow rate of $\|\widehat{A}^{(t)}\widehat{\theta}^{(t)} - A^{(t)*}\theta^{(t)}\|_2$. However, the proof of Lemma [37](#page-51-6) contains one important oracle inequality between $||A^{(t)*}(A^{(t)*})^{\top} - \hat{A}^{(t)}(\hat{A}^{(t)})^{\top}||_2$ and $\|\widehat{A}^{(t)}\widehat{\theta}^{(t)} - A^{(t)*}\theta^{(t)*}\|_2$. When $r = 1$, Lemma [38](#page-52-1) provides another direction of that inequality. Combining two inequalities implies a desired upper bound. We present the details in the following.

By the proof of Lemma [37,](#page-51-6) w.p. at least $1 - e^{-C'(p + \log T)}$, for all $t \in S$,

$$
\frac{\lambda}{\sqrt{n}}\|\mathbf{A}^{(t)*}(\mathbf{A}^{(t)*})^\top - \widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^\top\|_2 \geq c\|\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2^2 - C\sqrt{\frac{p + \log T}{n}}\|\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2.
$$

Then by Lemma [38,](#page-52-1) w.p. at least $1 - e^{-C'(1 + \log T)}$, for all $t \in S$,

$$
\|\widehat{A}^{(t)} \theta_{\widehat{A}^{(t)}}^{(t)} - A^{(t)*} \theta^{(t)*}\|_2 \geq C' \zeta^{(t)} \|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - A^{(t)*} (A^{(t)*})^\top\|_2 - C \sqrt{\frac{p + \log T}{n}}.
$$

Combining these two inequalities, we obtain

$$
\|\widehat{A}^{(t)} \theta_{\widehat{A}^{(t)}}^{(t)} - A^{(t)*} \theta^{(t)*} \|_2 \lesssim \sqrt{\frac{p + \log T}{n}} + \frac{\lambda^{1/2}}{n^{1/4}} \cdot \left(\frac{p + \log T}{n}\right)^{1/4} (\zeta^{(t)})^{-1/2} + \frac{\lambda}{\sqrt{n}} (\zeta^{(t)})^{-1}
$$

$$
\lesssim \sqrt{\frac{p + \log T}{n}} \cdot \frac{\max_{t \in S} (\zeta^{(t)})^2}{\min_{t \in S} (\zeta^{(t)})^2},
$$

for all $t \in S$, w.p. at least $1 - e^{-C(p + \log T)}$. The last inequality holds because $\lambda \geq$ $\max_{t \in S} (\zeta^{(t)})^2$ $\min_{t \in S} \zeta^{(t)}$ √ $p + \log T$. Combining this result with Proposition [43,](#page-52-6) we get the desired bound for $\max_{t \in S} \|\widehat{A}^{(t)} \theta_{\widehat{A}^{(t)}}^{(t)} - A^{(t)*} \theta^{(t)*} \|_2$. When the tasks in S^c also follow the linear model [\(3\)](#page-8-1), exactly the same arguments can be made for all $t \in S^c$ which completes our proof exactly the same arguments can be made for all $t \in S^c$, which completes our proof.

D.4 Proof of Theorem [4](#page-14-0)

First, note that when $\lambda = +\infty$, we have $\widehat{A}^{(t)} \equiv \overline{A}$ for all $t \in S = [T]$. We will prove that any local minimizer $(\overline{A}, {\{\hat{\theta}^{(t)}\}_{t=1}^{T}})$ which minimizes

$$
f(\boldsymbol{A},\boldsymbol{\Theta})\coloneqq \frac{1}{T}\sum_{t=1}^T f^{(t)}(\boldsymbol{A}\boldsymbol{\theta}^{(t)}).
$$

over $\mathbf{A} \in \mathcal{O}^{p \times r}$ and $\mathbf{\Theta} = (\mathbf{\theta}^{(1)}, \dots, \mathbf{\theta}^{(T)}) \in \mathbb{R}^{r \times T}$ must satisfy

$$
\|\widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^{\top} - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}\|_{\mathrm{F}} \lesssim (\bar{\zeta}^{-1}) r \sqrt{\frac{p}{nT}} + (\bar{\zeta}^{-1}) \sqrt{r} \sqrt{\frac{r + \log T}{n}} + \sqrt{r} h,
$$

simulteneously w.p. at least $1 - e^{-C'(r + \log T)}$. The following the argument in the proof of Theorem [1,](#page-12-0) we will obtain the desired result.

We prove the claim above by construction. Consider a local minimizer $(\overline{A}, \{\widehat{\theta}^{(t)}\}_{t=1}^T)$ of f and \overline{A} 's neightbor $\widetilde{A} = a_1\overline{A} + a_2\overline{A}D$, where $a_1 + a_2 = 1$, $a_1, a_2 \ge 0$, and $D \in \mathbb{R}^{r \times r}$ is invertible. Note that

$$
\|((\overline{A})^{\perp})^{\top}\widetilde{A}\|_{2} = a_{1}\|((\overline{A})^{\perp})^{\top}\overline{A}\|_{2} \le a_{1}.
$$

Define $\theta_{A}^{(t)} = \arg\min_{\theta \in \mathbb{R}^{r}} f^{(t)}(A\theta), \widehat{\Theta} = {\theta_{\widehat{A}}^{(t)}}_{I=1}^{T}, \widetilde{R} \in \mathcal{O}^{r \times r}$ by

$$
\widetilde{A}^{\top}\widetilde{A} = a_{1}^{2}I_{p} + a_{2}^{2}D^{\top}D + a_{1}a_{2}((\overline{A})^{\top}\widehat{A}D + D^{\top}(\widehat{A})^{\top}\overline{A}) = (\widetilde{R}^{-1})^{\top}\widetilde{R}^{-1}.
$$

Then define $\mathbf{A} = \widetilde{\mathbf{A}} \widetilde{\mathbf{R}}$, hence $\mathbf{A}^\top \mathbf{A} = \mathbf{I}_r$. Let \mathbf{D} satisfy $\mathbf{D} \widetilde{\mathbf{R}} \overline{\mathbf{\Theta}} = \widehat{\mathbf{\Theta}}$, where $\overline{\mathbf{\Theta}} = {\{\theta_{\overline{\mathbf{A}}}^{(t)}\}}$ $\{\frac{t}{A}\}_{t=1}^T \in$ $\mathbb{R}^{r \times T}$. Such D must exist. For example, we can take $D = (\widetilde{R} \widehat{\Theta} \widehat{\Theta}^{\top})^{-1} \widehat{\Theta} \overline{\Theta}^{\top}$. Consider Θ s.t. $\widetilde{R}\Theta = \overline{\Theta}$, then

$$
A\Theta = a_1 \overline{A} \widetilde{R}\Theta + a_2 \widehat{A} D \widetilde{R}\Theta = a_1 \overline{A} \overline{\Theta} + a_2 \widehat{A} \widehat{\Theta}.
$$
 (8)

Since $f^{(t)}$ is defined through square loss, we have

$$
f(\mathbf{A}, \Theta) - f(\widehat{\overline{\mathbf{A}}}, \widehat{\Theta}) = \frac{1}{2T} \sum_{t=1}^{T} (\mathbf{A}\theta^{(t)} - \widehat{\overline{\mathbf{A}}}\widehat{\theta}^{(t)})^{\top} \widehat{\Sigma}^{(t)} (\mathbf{A}\theta^{(t)} - \widehat{\overline{\mathbf{A}}}\widehat{\theta}^{(t)}) + \frac{1}{T} \sum_{t=1}^{T} \left[\nabla f^{(t)} (\widehat{\overline{\mathbf{A}}}\widehat{\theta}^{(t)}) \right]^{\top} (\mathbf{A}\theta^{(t)} - \widehat{\overline{\mathbf{A}}}\widehat{\theta}^{(t)}) .
$$

Therefore, [\(8\)](#page-55-0) implies that

$$
(*) \leq \frac{1}{T} \sum_{t=1}^T \left[\nabla f^{(t)}(\overline{A}\theta_{\overline{A}}^{(t)}) \right]^\top (A\theta^{(t)} - \widehat{\overline{A}}\widehat{\theta}^{(t)}) - \frac{1}{T} a_1 \sum_{t=1}^T (\widehat{\overline{A}}\widehat{\theta}^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)})^\top \widehat{\Sigma}^{(t)} (\widehat{\overline{A}}\widehat{\theta}^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)})
$$

$$
\leq \frac{1}{T} \|\{\nabla f^{(t)}(\mathbf{A}^{(t)*}\theta^{(t)*})\}_{t=1}^T\|_2 \cdot \sqrt{2r} \|\mathbf{A}\Theta - \widehat{\mathbf{A}}\widehat{\Theta}\|_{\mathrm{F}} \n+ \frac{1}{T} \|\{\nabla f^{(t)}(\mathbf{A}^{(t)*}\theta^{(t)*}) - \nabla f^{(t)}(\overline{\mathbf{A}}\theta_{\overline{\mathbf{A}}}^{(t)})\}_{t=1}^T\|_{\mathrm{F}} \|\mathbf{A}\Theta - \widehat{\mathbf{A}}\widehat{\Theta}\|_{\mathrm{F}} - \frac{C}{T} a_1 \|\widehat{\mathbf{A}}\widehat{\Theta} - \overline{\mathbf{A}}\overline{\Theta}\|_{\mathrm{F}}^2 \n\leq \frac{1}{T} \sqrt{2r} \sqrt{\frac{p+T}{n}} \|\mathbf{A}\Theta - \widehat{\mathbf{A}}\widehat{\Theta}\|_{\mathrm{F}} + \frac{C}{T} \left(\sqrt{T} \sqrt{\frac{r + \log T}{n}} + h \sqrt{\sum_{t=1}^T (\zeta^{(t)})^2} \right) \|\mathbf{A}\Theta - \widehat{\mathbf{A}}\widehat{\Theta}\|_{\mathrm{F}} \right)
$$

$$
- \frac{C}{T} a_1 \|\widehat{\overline{A}}\widehat{\Theta} - \overline{A}\overline{\Theta}\|_{\mathrm{F}}^2
$$

$$
\leq \frac{C}{T} a_1 \left(\sqrt{\frac{pr}{n}} + \sqrt{\frac{r + \log T}{n}} \sqrt{T} + h\sqrt{T}\overline{\zeta} \right) \|\widehat{\overline{A}}\widehat{\Theta} - \overline{A}\overline{\Theta}\|_{\mathrm{F}} - \frac{C}{T} a_1 \|\widehat{\overline{A}}\widehat{\Theta} - \overline{A}\overline{\Theta}\|_{\mathrm{F}}^2,
$$

w.p. at least $1 - e^{-C'(r + \log T)}$. Hence w.p. at least $1 - e^{-C'(r + \log T)}$,

$$
f(\mathbf{A}, \boldsymbol{\Theta}) - f(\widehat{\overline{\mathbf{A}}}, \widehat{\boldsymbol{\Theta}}) \le \frac{C''}{T} a_1^2 + \frac{C}{T} a_1 \left(\sqrt{\frac{pr}{n}} + \sqrt{\frac{r + \log T}{n}} \sqrt{T} + h \sqrt{T} \bar{\zeta} \right) ||\widehat{\overline{\mathbf{A}}} \widehat{\boldsymbol{\Theta}} - \overline{\mathbf{A}} \overline{\boldsymbol{\Theta}}||_{\mathrm{F}}
$$

$$
-\frac{C}{T}a_1\|\widehat{A}\widehat{\Theta}-\overline{A}\overline{\Theta}\|_{\mathrm{F}}^2.
$$
\n(9)

By [\(8\)](#page-55-0), as $a_1 \to 0^+,$

$$
\mathbf{\Theta} = a_1 \mathbf{A}^\top \overline{\mathbf{A}} \overline{\mathbf{\Theta}} + a_2 \mathbf{A}^\top \widehat{\overline{\mathbf{A}}} \widehat{\mathbf{\Theta}} \rightarrow \widetilde{\mathbf{R}}^\top \mathbf{D}^\top (\widehat{\overline{\mathbf{A}}})^\top \widehat{\overline{\mathbf{A}}} \widehat{\mathbf{\Theta}} = \widetilde{\mathbf{R}}^\top \mathbf{D}^\top \widehat{\mathbf{\Theta}},
$$

where " \rightarrow " is in the sense of $\|\cdot\|_{\max}$ (similar below). On the other hand, as $a_1 \rightarrow 0^+,$

$$
\boldsymbol{A}=a_1\overline{\boldsymbol{A}}\widetilde{\boldsymbol{R}}+a_2\overline{\widetilde{\boldsymbol{A}}}\boldsymbol{D}\widetilde{\boldsymbol{R}}\rightarrow\overline{\widetilde{\boldsymbol{A}}}\boldsymbol{D}\widetilde{\boldsymbol{R}},
$$

and $\widetilde{R}^\top D^\top D \widetilde{R} \to I_r$. Therefore, by continuity, as $a_1 \to 0^+,$

$$
\max_{t\in[T]}\min_{\boldsymbol{R}\in\mathcal{O}^{r\times r}}\left\{\|\boldsymbol{A}-\widehat{\boldsymbol{A}}\boldsymbol{R}\|_2+\|\boldsymbol{\theta}^{(t)}-\widehat{\boldsymbol{\theta}}^{(t)}\boldsymbol{R}\|_2\right\}\to 0.
$$

Due to the local optimality of $(\overline{\mathbf{A}}, \overline{\Theta})$, when a_1 is very close to 0, we must have $f(\mathbf{A}, \Theta)$ – $f(\hat{\overline{A}},\hat{\Theta}) \leq 0$, hence the RHS of [\(9\)](#page-56-1) is non-positive. Let $a_1 \lesssim \sqrt{\frac{pr}{n}} + \sqrt{\frac{r+\log T}{n}}$ \overline{n} √ $T + h$ $-\frac{1}{\sqrt{T}\bar{\zeta}},$ we have w.p. at least $1 - e^{-C'(r + \log T)}$,

$$
\|\widehat{\overline{A}}\widehat{\Theta}-\overline{A}\overline{\Theta}\|_{\rm F}\lesssim a_1+\sqrt{\frac{pr}{n}}+\sqrt{\frac{r+\log T}{n}}\sqrt{T}+h\sqrt{T}\bar{\zeta}\lesssim \sqrt{\frac{pr}{n}}+\sqrt{\frac{r+\log T}{n}}\sqrt{T}+h\sqrt{T}\bar{\zeta}.
$$

By Wedin's Theorem, w.p. at least $1 - e^{-C'(r + \log T)}$,

$$
\|\widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^\top - \overline{\mathbf{A}}(\overline{\mathbf{A}})^\top\|_{\rm F} \lesssim \sqrt{\frac{T}{r}}\bar{\zeta}\|\widehat{\overline{\mathbf{A}}}\widehat{\boldsymbol{\Theta}} - \overline{\mathbf{A}}\overline{\boldsymbol{\Theta}}\|_{\rm F} \lesssim \bar{\zeta}^{-1}r\sqrt{\frac{p}{nT}} + \bar{\zeta}^{-1}\sqrt{r}\sqrt{\frac{r+\log T}{n}} + \sqrt{r}h.
$$

This proves our previous claim. The remaining argument is the same as in the proof of Theorem [1.](#page-12-0)

D.5 Proof of Theorem [5](#page-14-1)

We prove a slightly stronger version of Theorem [5](#page-14-1) by replacing the constraint set

$$
4(1+\sqrt{2})\max_{t\in S} \|\widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)}) - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}\|_{2} + 4\frac{\epsilon}{1-\epsilon} < 1-\gamma,
$$

$$
\sqrt{2}\max_{t\in S} \left\{ \frac{\sigma_{\max}(\mathbf{\Sigma}^{(t)})}{\sigma_{\min}(\mathbf{\Sigma}^{(t)})} \|\widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^{\top} - \mathbf{A}^{(t)*}(\mathbf{A}^{(t)*})^{\top}\|_{2} \right\} < 1-\gamma.
$$
 (10)

with

$$
2(2+\sqrt{2})\max_{t\in S}\min_{\mathbf{R}\in\mathcal{O}^{r\times r}}\|\widehat{\mathbf{A}}^{(t)} - \overline{\mathbf{A}}\mathbf{R}\|_{2} + 4\frac{\epsilon}{1-\epsilon} < 1-\gamma,
$$

$$
\max_{t\in S}\left\{\frac{\sigma_{\max}(\mathbf{\Sigma}^{(t)})}{\sigma_{\min}(\mathbf{\Sigma}^{(t)})}\min_{\mathbf{R}\in\mathcal{O}^{r\times r}}\|\widehat{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*}\mathbf{R}\|_{2}\right\} < 1-\gamma.
$$
 (11)

Note that for any matrices $A, A' \in \mathcal{O}^{p \times r}$, Lemma 2.6 in [Chen et al.](#page-35-1) [\(2021b\)](#page-35-1) tells us that

$$
\|\boldsymbol{A}\boldsymbol{A}^{\top}-\boldsymbol{A}'(\boldsymbol{A}')^{\top}\|_2 \leq \min_{\boldsymbol{R}\in\mathcal{O}^{r\times r}}\|\boldsymbol{A}-\boldsymbol{A}'\boldsymbol{R}\|_2 \leq \sqrt{2}\|\boldsymbol{A}\boldsymbol{A}^{\top}-\boldsymbol{A}'(\boldsymbol{A}')^{\top}\|_2.
$$

Therefore the current constraint set (10) is indeed weaker than the constraint set (11) in the main text.

The proof follows the construction idea in the proof of Theorem [4](#page-14-0) and the argument in the proof of Proposition [43.](#page-52-6) The difference is that here we will first consider a relaxed optimization problem where we require $\widehat{A}^{(t)}$ and \overline{A} to satisfy $\|\widehat{A}^{(t)}(\widehat{A}^{(t)})^{\top} - I_r\|_2$, $\|\overline{A}(\overline{A})^{\top} - I_r\|_2 \le \delta$ with a small constant $\delta > 0$ instead of $\hat{A}^{(t)}, \overline{A} \in \mathcal{O}^{p \times r}$. Finally, we will let $\delta \to 0+$ to obtain the desired original result.

Denote

$$
G(\{\boldsymbol{A}^{(t)}\}_{t=1}^T,\overline{\boldsymbol{A}},\boldsymbol{\Theta})=\frac{1}{T}\sum_{t=1}^T f^{(t)}(\boldsymbol{A}^{(t)}\boldsymbol{\theta}^{(t)})+\sum_{t=1}^T \frac{\lambda}{\sqrt{n}}\|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top-\overline{\boldsymbol{A}}(\overline{\boldsymbol{A}})^\top\|_2,
$$

with $\mathbf{\Theta} = (\boldsymbol{\theta}^{(1)}, \dots, \boldsymbol{\theta}^{(T)}) \in \mathbb{R}^{r \times T}$. First, consider any local minimizer $\{\widehat{A}^{(t)}\}_{t=1}^T$ and \overline{A} of

$$
\min_{\{\boldsymbol{\theta}^{(t)}\}_{t=1}^T \subseteq \mathbb{R}^r} G(\{\boldsymbol{A}^{(t)}\}_{t=1}^T, \overline{\boldsymbol{A}}, \boldsymbol{\Theta}),
$$

where $\|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - I_r\|_2, \ \|\overline{A}(\overline{A})^\top - I_r\|_2 \leq \delta$ with a small constant $\delta > 0$. Define $\mathbf{A}^{(t)} = a_1 \overline{\mathbf{A}} \mathbf{D}^{(t)} + a_2 \widehat{\mathbf{A}}^{(t)}$ with $a_1 + a_2 = 1$ and $a_1, a_2 \geq 0$, $\mathbf{D}^{(t)} \in \mathcal{O}^{p \times r}$. Define a "normalization" operator $\mathcal{N}(\cdot)$ such that $\mathcal{N}(\mathbf{A}) \coloneqq \mathbf{A}(\mathbf{A}^{\top}\mathbf{A})^{-1/2} \in \mathcal{O}^{p \times r}$ for any $\mathbf{A} \in \mathbb{R}^{p \times r}$ with full column rank. Define two distances between subspaces spanned by columns of \boldsymbol{A} and $\mathbf{B} \in \mathbb{R}^{p \times r}$ as

$$
d_1(\boldsymbol{A},\boldsymbol{B}) = \|\mathcal{N}(\boldsymbol{A})(\mathcal{N}(\boldsymbol{A}))^\top - \mathcal{N}(\boldsymbol{B})(\mathcal{N}(\boldsymbol{B}))^\top\|_2, \\ d_2(\boldsymbol{A},\boldsymbol{B}) = \min_{\boldsymbol{R}\in\mathcal{O}^{r\times r}} \|\mathcal{N}(\boldsymbol{A}) - \mathcal{N}(\boldsymbol{B})\boldsymbol{R}\|_2.
$$

By Lemma 2.6 of [Chen et al.](#page-35-1) [\(2021b\)](#page-35-1), $d_1(A, B) \leq d_2(A, B) \leq$ √ $2d_1(\boldsymbol{A},\boldsymbol{B}).$ Note that

$$
\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^{\top} = a_1^2 \boldsymbol{I}_r + a_2^2 \boldsymbol{I}_r + a_1 a_2 [(\boldsymbol{D}^{(t)})^{\top} (\overline{\boldsymbol{A}})^{\top} \widehat{\boldsymbol{A}}^{(t)} + (\widehat{\boldsymbol{A}}^{(t)})^{\top} \overline{\boldsymbol{A}} \boldsymbol{D}^{(t)}].
$$

In addition, there exists a rotation matrix $\mathbf{R}^{(t)} \in \mathcal{O}^{r \times r}$ s.t.

$$
\widehat{\boldsymbol{\theta}}^{(t)} = \boldsymbol{R}^{(t)} \frac{\|\widehat{\boldsymbol{\theta}}^{(t)}\|_2}{\|\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}\|_2} \boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}.
$$

Consider $\boldsymbol{\theta}^{(t)} \in \mathbb{R}^r$ s.t.

$$
\boldsymbol{\theta}^{(t)} = \alpha \cdot \frac{\widehat{\boldsymbol{\theta}}^{(t)}}{\|\widehat{\boldsymbol{\theta}}^{(t)}\|_2},
$$

then by taking $\boldsymbol{D}^{(t)} = (\boldsymbol{R}^{(t)})^{-1}$, we have

$$
\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} = a_1\overline{\mathbf{A}}(\mathbf{R}^{(t)})^{-1}\boldsymbol{\theta}^{(t)} + a_2\widehat{\mathbf{A}}^{(t)}\boldsymbol{\theta}^{(t)} = a_1\frac{\alpha}{\|\boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)}\|_2}\overline{\mathbf{A}}\boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)} + a_2\frac{\alpha}{\|\widehat{\boldsymbol{\theta}}^{(t)}\|_2}\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}.
$$

Let $a_1 \frac{\alpha}{\ln a^{(t)}}$ $\|\theta^{(t)}_{\pi}\|$ $\frac{\alpha}{\|\vec{a}\|_2} + a_2 \frac{\alpha}{\|\hat{\theta}^{(t)}\|_2}$ $\frac{\alpha}{\|\widehat{\theta}^{(t)}\|_2} = 1$, we obtain that $\alpha = \left(\frac{a_1}{\|\theta_A^{(t)}\|_2}\right)$ $\|\theta^{(t)}_{\tau}\|$ $\frac{\frac{a_1}{(t)}\|_2}{\overline{A}} + \frac{a_2}{\|\widehat{\theta}^{(t)}\|}$ $\|\hat{\boldsymbol{\theta}}^{(t)}\|_2$ $^{-1}$, which implies that

$$
\boldsymbol{A}^{(t)}\boldsymbol{\theta}^{(t)}-\widehat{\boldsymbol{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}=\underbrace{\frac{a_1/\|\boldsymbol{\theta}_{\widehat{\boldsymbol{A}}}^{(t)}\|_2}{\frac{a_1/\|\boldsymbol{\theta}_{\widehat{\boldsymbol{A}}}^{(t)}\|_2+a_2/\|\widehat{\boldsymbol{\theta}}^{(t)}\|_2}{\cdots}_{i=a_1^{(t)}}}(\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\widehat{\boldsymbol{A}}}^{(t)}-\widehat{\boldsymbol{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}).
$$

Also define $a_2^{(t)} = 1 - a_1^{(t)}$ $1^{(l)}$. We claim that

$$
C \ge \frac{a_1^{(t)}}{a_1} = \frac{1}{a_1 + a_2 \cdot \frac{\|\theta_A^{(t)}\|_2}{\|\widehat{\theta}^{(t)}\|_2}} \ge \frac{\|\widehat{\theta}^{(t)}\|_2}{\|\theta_A^{(t)}\|_2} \ge C' > 0, \quad \forall t \in S,
$$
\n(12)

with some constants $C, C' > 0$. To see this, note that

$$
\widehat{\boldsymbol{\theta}}^{(t)} = ((\widehat{\boldsymbol{A}}^{(t)})^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \widehat{\boldsymbol{A}}^{(t)})^{-1} (\widehat{\boldsymbol{A}}^{(t)})^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*} - ((\widehat{\boldsymbol{A}}^{(t)})^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \widehat{\boldsymbol{A}}^{(t)})^{-1} (\widehat{\boldsymbol{A}}^{(t)})^{\top} \nabla f^{(t)} (\boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}).
$$

Also note that $((\widehat{\mathbf{A}}^{(t)})^{\top} \widehat{\mathbf{\Sigma}}^{(t)} \widehat{\mathbf{A}}^{(t)})^{-1}$ and $(\widehat{\mathbf{\Sigma}}^{(t)})^{-1}$ share the same eigenvalues. Since $\sigma_{\max}((\widehat{\mathbf{\Sigma}}^{(t)})^{-1}) \asymp$ $\sigma_{\min}((\hat{\Sigma}^{(t)})^{-1}) \ge 1$ w.p. at least $1 - e^{-C'(p + \log T)}$ by Lemma [19,](#page-47-4) we have

$$
\sigma_{\max}\big(((\widehat{\mathbf{A}}^{(t)})^{\top}\widehat{\mathbf{\Sigma}}^{(t)}\widehat{\mathbf{A}}^{(t)})^{-1}\big)\asymp \sigma_{\min}\big(((\widehat{\mathbf{A}}^{(t)})^{\top}\widehat{\mathbf{\Sigma}}^{(t)}\widehat{\mathbf{A}}^{(t)})^{-1}\big)\asymp 1,
$$

w.p. at least $1 - e^{-C'(p + \log T)}$. Furthermore, by Weyl's inequality and Lemma [19,](#page-47-4) with $\boldsymbol{R} \in \arg \min_{\boldsymbol{R} \in \mathcal{O}^{r \times r}} \| \mathcal{N}(\widehat{\boldsymbol{A}}^{(t)}) - \boldsymbol{A}^{(t) *} \boldsymbol{R} \|_2,$

$$
\sigma_r((\widehat{A}^{(t)})^\top \widehat{\Sigma}^{(t)} \mathbf{A}^{(t)*})
$$
\n
$$
\geq \sigma_r((\widehat{A}^{(t)})^\top \Sigma^{(t)} \mathbf{A}^{(t)*}) - \|\widehat{\Sigma}^{(t)} - \Sigma^{(t)}\|_2
$$
\n
$$
\geq \sigma_r(\mathbf{R}^\top (\mathbf{A}^{(t)*})^\top \Sigma^{(t)} \mathbf{A}^{(t)*}) - \sigma_{\max}((\mathbf{R}^\top (\mathbf{A}^{(t)*})^\top - \mathcal{N}(\widehat{\mathbf{A}}^{(t)})^\top) \Sigma^{(t)} \mathbf{A}^{(t)*}) - C\delta - C\sqrt{\frac{p + \log T}{n}}
$$

$$
\geq \sigma_r((\mathbf{A}^{(t)*})^{\top} \mathbf{\Sigma}^{(t)} \mathbf{A}^{(t)*}) - ||\mathcal{N}(\widehat{\mathbf{A}}^{(t)}) - \mathbf{A}^{(t)*} \mathbf{R}||_2 ||\mathbf{\Sigma}^{(t)}||_2 - C\delta - C\sqrt{\frac{p + \log T}{n}}
$$

\n
$$
\geq \sigma_{\min}(\mathbf{\Sigma}^{(t)}) - ||\mathcal{N}(\widehat{\mathbf{A}}^{(t)}) - \mathbf{A}^{(t)*} \mathbf{R}||_2 \cdot \sigma_{\max}(\mathbf{\Sigma}^{(t)}) - C\delta - C\sqrt{\frac{p + \log T}{n}}
$$

\n
$$
> 0,
$$

w.p. at least $1 - e^{-C'(p + \log T)}$, if $d_2(\widehat{A}^{(t)}, A^{(t)*}) \leq \frac{\sigma_{\min}(\Sigma^{(t)})}{\sigma_{\max}(\Sigma^{(t)})}$ $\frac{\sigma_{\min}(\Sigma^{(v)})}{\sigma_{\max}(\Sigma^{(t)})}(1-\gamma)$ with some $\gamma > 0$. On the other hand, by Lemma [19,](#page-47-4) w.p. at least $1 - e^{-C'(p + \log T)}$, for all $t \in S$,

$$
\sigma_{\max}((\widehat{\bm{A}}^{(t)})^{\top}\widehat{\bm{\Sigma}}^{(t)}{\bm{A}}^{(t)*})\lesssim 1.
$$

Hence w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
\|\widehat{\theta}^{(t)}\|_2 \geq C\sigma_r((\widehat{A}^{(t)})^\top \widehat{\Sigma}^{(t)} A^{(t)*})\|\theta^{(t)*}\|_2 - C'\|\nabla f^{(t)}(A^{(t)*}\theta^{(t)*})\|_2
$$

$$
\geq C'' \|\theta^{(t)*}\|_2 - C''' \sqrt{\frac{p + \log T}{n}}
$$

$$
\geq \frac{C''}{2} \|\theta^{(t)*}\|_2.
$$

In addition, note that w.p. at least $1 - e^{-C'(r + \log T)}$,

$$
|\|\boldsymbol{\theta}_{\overline{A}}^{(t)}\|_2 - \|\boldsymbol{\theta}^{(t)*}\|_2| \leq \|\boldsymbol{\theta}_{\overline{A}}^{(t)} - \boldsymbol{\theta}^{(t)*}\|_2 \lesssim h\zeta^{(t)} + \sqrt{\frac{r + \log T}{n}},
$$

implying that w.p. at least $1 - e^{-C'(r + \log T)}$,

$$
\begin{aligned} \|\pmb{\theta}_{\overline{\mathbf{A}}}^{(t)}\|_2 &\leq \|\pmb{\theta}^{(t)*}\|_2 + C\bigg(h\zeta^{(t)} + \sqrt{\frac{r+\log T}{n}}\bigg) \leq \|\pmb{\theta}^{(t)*}\|_2 + C'\bigg(r^{-1/2}\zeta^{(t)} + \sqrt{\frac{r+\log T}{n}}\bigg) \lesssim \|\pmb{\theta}^{(t)*}\|_2, \\ \|\pmb{\theta}^{(t)*}\|_2 &\lesssim \|\pmb{\theta}^{(t)*}\|_2 - C\bigg(h\zeta^{(t)} + \sqrt{\frac{r+\log T}{n}}\bigg) \leq \|\pmb{\theta}_{\overline{\mathbf{A}}}^{(t)}\|_2. \end{aligned}
$$

Hence w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
\|\widehat{\boldsymbol{\theta}}^{(t)}\|_2 \geq \frac{C''}{2}\|\boldsymbol{\theta}^{(t)*}\|_2 \gtrsim \|\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}\|_2.
$$

This proves the second half of [\(12\)](#page-58-0). On the other hand, when $a_1 \leq 1/2$, we must have

$$
\frac{a_1^{(t)}}{a_1} \le 2 \frac{\|\widehat{\boldsymbol{\theta}}^{(t)}\|_2}{\|\boldsymbol{\theta}_{\overline{A}}^{(t)}\|_2},
$$

and it is easy to see that w.p. at least $1 - e^{-C'(r + \log T)}$,

$$
\|\widehat{\theta}^{(t)}\|_2 \leq C \|\theta^{(t)*}\|_2 + C\sqrt{\frac{p+\log T}{n}} \lesssim \|\theta^{(t)*}\|_2 \lesssim \|\theta_{\overline{A}}^{(t)}\|_2,
$$

which proves the first half of (12) . Therefore, our claim (12) holds, and we will use it later in our proof.

Now similar to the proof of Proposition 43 , we divide S into the following two index sets

$$
\mathcal{A}_1 = \left\{ t \in S : \| \mathbf{A}^{(t)} (\mathbf{A}^{(t)})^\top - \widehat{\overline{\mathbf{A}}} (\widehat{\overline{\mathbf{A}}})^\top \|_2 \ge c \| \widehat{\overline{\mathbf{A}}} (\widehat{\overline{\mathbf{A}}})^\top - \overline{\mathbf{A}} \overline{\mathbf{A}}^\top \|_2 \cdot \frac{1}{\sqrt{r}} \right\},
$$

$$
\mathcal{A}_2 = \left\{ t \in S : \| \mathbf{A}^{(t)} (\mathbf{A}^{(t)})^\top - \widehat{\overline{\mathbf{A}}} (\widehat{\overline{\mathbf{A}}})^\top \|_2 < c \| \widehat{\overline{\mathbf{A}}} (\widehat{\overline{\mathbf{A}}})^\top - \overline{\mathbf{A}} \overline{\mathbf{A}}^\top \|_2 \cdot \frac{1}{\sqrt{r}} \right\},
$$

where $c > 0$ is a small constant. For any index set $\mathcal{A} \subseteq [T]$, define

$$
G_{\mathcal{A}}(\{\mathbf{A}^{(t)}\}_{t=1}^T,\overline{\mathbf{A}},\boldsymbol{\Theta})=\frac{1}{T}\sum_{t\in\mathcal{A}}f^{(t)}(\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)})+\sum_{t\in\mathcal{A}}\frac{\lambda}{\sqrt{n}T}\|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top-\overline{\mathbf{A}}(\overline{\mathbf{A}})^\top\|_2
$$

$$
= \frac{1}{2T}\sum_{t \in \mathcal{A}} \|\boldsymbol{Y}^{(t)} - \boldsymbol{X}^{(t)}\boldsymbol{A}^{(t)}\boldsymbol{\theta}^{(t)}\|_2^2 + \frac{\lambda}{T\sqrt{n}}\sum_{t \in \mathcal{A}} \|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top - \boldsymbol{A}\boldsymbol{A}^\top\|_2.
$$

Note that by basic algebra,

$$
\frac{1}{2T} \sum_{t \in \mathcal{A}} \|\mathbf{Y}^{(t)} - \mathbf{X}^{(t)} \mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)}\|_2^2 - \frac{1}{2T} \sum_{t \in \mathcal{A}} \|\mathbf{Y}^{(t)} - \mathbf{X}^{(t)} \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)}\|_2^2 \n= \frac{1}{2T} (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)})^\top \widehat{\boldsymbol{\Sigma}}^{(t)} (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)}) + \frac{1}{T} \sum_{t \in \mathcal{A}} [\nabla f^{(t)} (\widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)})]^\top (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)}) \n= \frac{1}{2T} (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)})^\top \widehat{\boldsymbol{\Sigma}}^{(t)} (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)}) + \frac{1}{T} \sum_{t \in \mathcal{A}} [\nabla f^{(t)} (\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)})]^\top (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)}) \n+ \frac{1}{T} \sum_{t \in \mathcal{A}} (\widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)} - \overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)})^\top \widehat{\boldsymbol{\Sigma}}^{(t)} (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)}).
$$

Hence

$$
G_{\mathcal{A}_{1}}(\{\mathbf{A}^{(t)}\}_{t=1}^{T}, \mathbf{A}, \Theta) - G_{\mathcal{A}_{1}}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^{T}, \widehat{\mathbf{A}}, \widehat{\Theta})
$$

\n
$$
\leq \frac{1}{2T}(\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)})^{\top} \widehat{\mathbf{\Sigma}}^{(t)}(\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)})
$$

\n
$$
+ \frac{1}{T} \sum_{t \in \mathcal{A}_{1}} \|\nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_{2} a_{1}^{(t)}(\|\overline{\mathbf{A}}\boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)} - \widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)}\|_{2} + \|\widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}\|_{2})
$$

\n
$$
+ \frac{1}{T} \left\| \left\{ \sqrt{a_{1}^{(t)}} [\nabla f^{(t)}(\overline{\mathbf{A}}\boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)}) - \nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})] \right\}_{t \in \mathcal{A}_{1}} \right\|_{\mathrm{F}} \cdot \sqrt{\sum_{t \in \mathcal{A}_{1}} a_{1}^{(t)} \|\overline{\mathbf{A}}\boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}\|_{2}^{2}}
$$

$$
-\frac{C}{T} \sum_{t \in \mathcal{A}_1} a_1^{(t)} \|\overline{A}\boldsymbol{\theta}_{\overline{A}}^{(t)} - \widehat{A}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}\|_2^2 + \frac{\lambda}{T\sqrt{n}} \sum_{t \in \mathcal{A}_1} \|A^{(t)}(A^{(t)})^\top - A A^\top\|_2
$$

$$
-\frac{\lambda}{T\sqrt{n}} \sum_{t \in \mathcal{A}_1} \|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - \widehat{A}(\widehat{A})^\top\|_2,
$$
 (13)

with

$$
[1] = \frac{1}{2T} (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)})^\top \widehat{\boldsymbol{\Sigma}}^{(t)} (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)}) + \frac{1}{T} \sum_{t \in \mathcal{A}_1} \|\nabla f^{(t)} (\mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*})\|_2 a_1^{(t)} (\|\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)} - \widehat{\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)}\|_2 + \|\widehat{\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)}\|_2) + \frac{1}{T} \left\| \left\{ \sqrt{a_1^{(t)}} [\nabla f^{(t)} (\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)}) - \nabla f^{(t)} (\mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*})] \right\}_{t \in \mathcal{A}_1} \right\|_{\mathrm{F}} \cdot \sqrt{\sum_{t \in \mathcal{A}_1} a_1^{(t)} \|\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)} - \widehat{\mathbf{A}}^{(t)} \widehat{\boldsymbol{\theta}}^{(t)}\|_2^2}
$$

$$
-\frac{C}{T}\sum_{t\in\mathcal{A}_1}a_1^{(t)}\|\overline{A}\boldsymbol{\theta}_{\overline{A}}^{(t)}-\widehat{A}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}\|_2^2,
$$

and

$$
[2] = \frac{\lambda}{T\sqrt{n}} \sum_{t \in \mathcal{A}_1} ||\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top - \boldsymbol{A}\boldsymbol{A}^\top||_2 - \frac{\lambda}{T\sqrt{n}} \sum_{t \in \mathcal{A}_1} ||\widehat{\boldsymbol{A}}^{(t)}(\widehat{\boldsymbol{A}}^{(t)})^\top - \widehat{\overline{\boldsymbol{A}}}(\widehat{\overline{\boldsymbol{A}}})^\top ||_2. \tag{14}
$$

Recall that $\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} = a_1^{(t)}$ $\frac{1}{(t)}(\overline{A}\theta_{\overline{A}}^{(t)} - \widehat{A}^{(t)}\widehat{\theta}^{(t)})$ and our previous conclusion that w.p. at least $1 - e^{-C'(r + \log T)}$, $a_1^{(t)}$ $\binom{t}{1}$ / $a_1 \simeq 1$ for all $t \in S$, hence w.p. at least $1 - e^{-C'(r + \log T)}$,

$$
[1] \leq \frac{C}{T} \sum_{t \in A_1} a_1^2 ||\overline{A} \theta_A^{(t)} - \widehat{A}^{(t)} \widehat{\theta}^{(t)}||_2^2
$$

+ $\frac{1}{T} \sum_{t \in A_1} ||\nabla f^{(t)} (A^{(t)*} \theta^{(t)*})||_2 a_1 \zeta^{(t)} \Big[d_1(\widehat{A}, \overline{A}) + d_1(\widehat{A}, \widehat{A}^{(t)}) \Big]$
+ $\frac{1}{T} \Big(\sqrt{T} \sqrt{a_1} \sqrt{\frac{r + \log T}{n}} + h \sqrt{\sum_{t \in A_1} (\zeta^{(t)})^2 a_1} \Big) \sqrt{\sum_{t \in A_1} a_1 ||\overline{A} \theta_A^{(t)} - \widehat{A}^{(t)} \widehat{\theta}^{(t)}||_2^2}$
- $\frac{C}{T} \sum_{t \in A_1} a_1 ||\overline{A} \theta_A^{(t)} - \widehat{A}^{(t)} \widehat{\theta}^{(t)}||_2^2$
 $\leq \frac{C}{T} \sum_{t \in A_1} a_1^2 ||\overline{A} \theta_A^{(t)} - \widehat{A}^{(t)} \widehat{\theta}^{(t)}||_2^2$
+ $\frac{1}{T} \sum_{t \in A_1} ||\nabla f^{(t)} (A^{(t)*} \theta^{(t)*})||_2 a_1 \zeta^{(t)} (||\widehat{A}(\widehat{A})^T - \overline{A}(\overline{A})^T ||_2 + ||\widehat{A}(\widehat{A})^T - \widehat{A}^{(t)} (\widehat{A}^{(t)})^T ||_2 + \delta)$
+ $\frac{1}{T} \Big(\sqrt{T} \sqrt{a_1} \sqrt{\frac{r + \log T}{n}} + h \sqrt{\sum_{t \in A_1} (\zeta^{(t)})^2 a_1} \Big) \sqrt{\sum_{t \in A_1} a_1 ||\overline{A} \theta_A^{(t)} - \widehat{A}^{(t)} \widehat{\theta}^{(t)}||_2^2}$
- $\frac{C}{T} \sum_{t \in A_1} a_1 ||\overline{A} \theta_A^{(t)} - \widehat{A}^{(t)} \widehat{\theta}^{($

To bound [2], we need to do some preparations. Note that

$$
\begin{split} \mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top} &= (a_1 \overline{\mathbf{A}} \mathbf{D}^{(t)} + a_2 \widehat{\mathbf{A}}^{(t)}) (a_1 \overline{\mathbf{A}} \mathbf{D}^{(t)} + a_2 \widehat{\mathbf{A}}^{(t)})^{\top} \\ &= a_2^2 \widehat{\mathbf{A}}^{(t)} (\widehat{\mathbf{A}}^{(t)})^{\top} + a_1 a_2 [\overline{\mathbf{A}} \mathbf{D}^{(t)} (\widehat{\mathbf{A}}^{(t)})^{\top} + \widehat{\mathbf{A}}^{(t)} (\mathbf{D}^{(t)})^{\top} (\overline{\mathbf{A}})^{\top}] + \mathcal{O}(a_1^2) \\ &= (1 - 2a_1) \widehat{\mathbf{A}}^{(t)} (\widehat{\mathbf{A}}^{(t)})^{\top} + a_1 [\overline{\mathbf{A}} \mathbf{D}^{(t)} (\widehat{\mathbf{A}}^{(t)})^{\top} + \widehat{\mathbf{A}}^{(t)} (\mathbf{D}^{(t)})^{\top} (\overline{\mathbf{A}})^{\top}] + \mathcal{O}(a_1^2), \end{split}
$$

where $\mathcal{O}(a_1^2)$ here refers to a $p \times p$ matrix with each entry of order $\mathcal{O}(a_1^2)$. Let $\mathbf{A} = a_1 \overline{\mathbf{A}} + a_2 \overline{\mathbf{A}}$, then similarly we have

$$
\boldsymbol{A}\boldsymbol{A}^{\top} = (1 - 2a_1)\widehat{\boldsymbol{A}}(\widehat{\boldsymbol{A}})^{\top} + a_1[\overline{\boldsymbol{A}}(\widehat{\boldsymbol{A}})^{\top} + \widehat{\boldsymbol{A}}(\overline{\boldsymbol{A}})^{\top}] + \mathcal{O}(a_1^2). \tag{15}
$$

This implies that

$$
A^{(t)}(A^{(t)})^{\top} - AA^{\top}
$$

\n
$$
= (1 - a_1)[\hat{A}^{(t)}(\hat{A}^{(t)})^{\top} - \hat{A}(\hat{A})^{\top}] - \frac{a_1}{2}(\hat{A}^{(t)}(D^{(t)})^{\top} + \hat{A})(\hat{A}^{(t)}(D^{(t)})^{\top} - \hat{A})^{\top}
$$

\n
$$
- \frac{a_1}{2}(\hat{A}^{(t)}(D^{(t)})^{\top} - \hat{A})(\hat{A}^{(t)}(D^{(t)})^{\top} + \hat{A})^{\top} + a_1\overline{A}[D^{(t)}(\hat{A}^{(t)})^{\top} - (\hat{A})^{\top}]
$$

\n
$$
+ a_1[\hat{A}^{(t)}(D^{(t)})^{\top} - \hat{A}](\overline{A})^{\top} + \mathcal{O}(a_1^2)
$$

\n
$$
= (1 - a_1)[\hat{A}^{(t)}(\hat{A}^{(t)})^{\top} - \hat{A}(\hat{A})^{\top}] + a_1[\overline{A} - \frac{1}{2}(\hat{A}^{(t)}(D^{(t)})^{\top} + \hat{A})](\hat{A}^{(t)}(D^{(t)})^{\top} - \hat{A})^{\top}
$$

\n
$$
+ a_1[\hat{A}^{(t)}(D^{(t)})^{\top} - \hat{A}]\left[\overline{A} - \frac{1}{2}(\hat{A}^{(t)}(D^{(t)})^{\top} + \hat{A})\right]^{\top} + \mathcal{O}(a_1^2)
$$

\n
$$
= (1 - a_1)[\hat{A}^{(t)}(\hat{A}^{(t)})^{\top} - \hat{A}(\hat{A})^{\top}] + a_1(\overline{A} - \hat{A})[\hat{A}^{(t)}(D^{(t)})^{\top} - \hat{A}]^{\top}
$$

\n
$$
- a_1[\hat{A}^{(t)}(D^{(t)})^{\top} - \hat{A}][\hat{A}^{(t)}(D^{(t)})^{\top} - \hat{A}]^{\top} + a_1[\hat{A}^{(t)}(D^{(t)})^{\top} - \hat{A}](\overline{A} - \hat{
$$

Note that we can multiply \overline{A} , $\widehat{A}^{(t)}$, and \overline{A} by a roration matrix from the right without changing their definitions

$$
\overline{A} \in \underset{\mathbf{A} \in \mathcal{O}^{p \times r}}{\arg \min} \|A^{(t)*} (A^{(t)*})^\top - AA^\top \|_2,
$$
\n
$$
\{\widehat{A}^{(t)}\}_{t=1}^T, \widehat{A} \in \underset{\begin{array}{c}\{\mathbf{A}^{(t)}\}_{t=1}^T, \overline{A} \in \mathbb{R}^{p \times r}\\ \|\mathbf{A}^{(t)}\}_{t=1}^T, \overline{A} \in \mathbb{R}^{p \times r}\\ \|\mathbf{A}^{(t)}\}_{t=1}^T, \overline{A} \in \mathbb{R}^{p \times r}\\ \|\mathbf{A}^{(t)}\}_{t=1}^T, \|\overline{A}\|_2^T, \|\overline{A}\|
$$

Therefore, WLOG, we assume

$$
\|\overline{A} - \mathcal{N}(\widehat{\overline{A}})\|_2 = \min_{\mathbf{R} \in \mathcal{O}^{r \times r}} \|\overline{A} - \mathcal{N}(\widehat{\overline{A}})\mathbf{R}\|_2 = d_2(\overline{A}, \widehat{\overline{A}}),
$$
(16)

$$
\|\mathcal{N}(\widehat{A}^{(t)}(\mathbf{D}^{(t)})^\top) - \mathcal{N}(\widehat{\overline{A}})\|_2 = \min_{\mathbf{R} \in \mathcal{O}^{r \times r}} \|\mathcal{N}(\widehat{A}^{(t)}) - \mathcal{N}(\widehat{\overline{A}})\mathbf{R}\|_2 = d_2(\widehat{A}^{(t)}, \widehat{\overline{A}}).
$$

Therefore, by triangle inequality,

$$
\|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top - \boldsymbol{A}\boldsymbol{A}^\top\|_2 \leq (1-a_1)\|\widehat{\boldsymbol{A}}^{(t)}(\widehat{\boldsymbol{A}}^{(t)})^\top - \widehat{\overline{\boldsymbol{A}}}(\widehat{\overline{\boldsymbol{A}}})^\top\|_2 + \max\left\{2a_1\|\overline{\boldsymbol{A}} - \widehat{\overline{\boldsymbol{A}}}\|_2\|\widehat{\boldsymbol{A}}^{(t)}(\boldsymbol{D}^{(t)})^\top - \widehat{\overline{\boldsymbol{A}}}\|_2,
$$

$$
a_1 \|\widehat{\mathbf{A}}^{(t)}(\mathbf{D}^{(t)})^\top - \widehat{\mathbf{A}}\|_2^2 \Big\} + C\sqrt{p}a_1^2
$$

$$
\leq (1 - a_1) \|\widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^\top - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^\top\|_2 + \max \Big\{2\sqrt{2}a_1 \cdot d_2(\overline{\mathbf{A}}, \widehat{\mathbf{A}})d_1(\widehat{\mathbf{A}}^{(t)}, \widehat{\mathbf{A}}),
$$

$$
\sqrt{2}a_1 \cdot d_2(\widehat{A}^{(t)}, \widehat{\overline{A}})d_1(\widehat{A}^{(t)}, \widehat{\overline{A}})\Big\} + Ca_1\delta + C\sqrt{p}a_1^2
$$

$$
\leq \|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_2 + \Big[-1 + (2\sqrt{2}d_2(\overline{A}, \widehat{\overline{A}})) \vee (\sqrt{2}d_2(\widehat{A}^{(t)}, \widehat{\overline{A}}))\Big]
$$

$$
a_1d_1(\widehat{A}^{(t)}, \widehat{\overline{A}}) + Ca_1\delta + C\sqrt{p}a_1^2
$$

$$
\leq ||\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top ||_2 + \left[-2^{-\frac{1}{2}} + (2d_2(\overline{A}, \widehat{\overline{A}})) \vee (d_2(\overline{A}, \widehat{\overline{A}}) + d_2(\widehat{A}^{(t)}, \overline{A})) \right]
$$

$$
\begin{aligned}\n&\cdot \sqrt{2}a_1d_1(\widehat{\mathbf{A}}^{(t)},\widehat{\overline{\mathbf{A}}}) + Ca_1\delta + C\sqrt{p}a_1^2 \\
&\leq \|\widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^\top - \widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^\top\|_2 + \left[-2^{-\frac{1}{2}} + \left(2d_2(\overline{\mathbf{A}},\widehat{\overline{\mathbf{A}}})\right) \vee \left(d_2(\overline{\mathbf{A}},\widehat{\overline{\mathbf{A}}}) + d_2(\widehat{\mathbf{A}}^{(t)},\overline{\mathbf{A}})\right) \right]\n\end{aligned}
$$

$$
\cdot \sqrt{2}a_1 \|\widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^\top - \widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^\top\|_2 + Ca_1\delta + C\sqrt{p}a_1^2.
$$

To further bound the RHS, we claim that $d_2(\overline{\mathbf{A}}, \hat{\overline{\mathbf{A}}}) \leq (1 + \sqrt{2}) \max_{t \in S} d_2(\widehat{\mathbf{A}}^{(t)}, \overline{\mathbf{A}}) + \sqrt{2} \frac{\epsilon}{1 - \epsilon} +$ $C' \delta$ with some constant $C' > 0$. To see this, denote $q = \max_{t \in S} d_2(\widehat{A}^{(t)}, \overline{A})$, and suppose $d_2(\overline{\mathbf{A}}, \overline{\mathbf{A}}) \ge q + b$ with some $b \ge 0$, then $\min_{t \in S} d_2(\overline{\mathbf{A}}, \widehat{\mathbf{A}}^{(t)}) \ge b$. This leads to

$$
\sum_{t=1}^T \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \widehat{A}^{(t)}(\widehat{A}^{(t)})^{\top}\|_2 \ge \frac{1}{\sqrt{2}} \sum_{t \in S} d_2(\widehat{\overline{A}}, \widehat{A}^{(t)}) - C\delta \ge \frac{|S|}{\sqrt{2}}b - CT\delta.
$$

On the other hand, we have

$$
\sum_{t \in S} d_2(\widehat{A}^{(t)}, \overline{A}) \le |S|q, \quad \sum_{t \in S^c} d_2(\widehat{A}^{(t)}, \overline{A}) \le |S^c|,
$$

which implies that

$$
\sum_{t=1}^T \|\overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top} - \widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^{\top}\|_2 \leq |S|q + |S^c| + CT\delta.
$$

By the optimization procedure, we know that $\overline{A} \in \operatorname*{arg\,min}_{A \in \mathbb{R}^{p \times r}: ||A^\top A - I_r||_2 \le \delta}$ $\sum_{t=1}^T \| \boldsymbol{A} \boldsymbol{A}^\top \widehat{A}^{(t)}(\widehat{A}^{(t)})^{\top}$ ||2. Therefore we must have $|S|q + |S^c| + CT\delta \ge \frac{|S|}{\sqrt{2}}b - CT\delta$, i.e.

$$
b \le \sqrt{2}q + \sqrt{2}\frac{|S^c|}{|S|} + C'\delta \le \sqrt{2}q + \sqrt{2}\frac{\epsilon}{1-\epsilon} + C'\delta.
$$

In other words,

$$
d_2(\overline{\mathbf{A}},\widehat{\overline{\mathbf{A}}}) \leq q + b \leq (1+\sqrt{2}) \max_{t \in S} d_2(\widehat{\mathbf{A}}^{(t)},\overline{\mathbf{A}}) + \sqrt{2} \frac{\epsilon}{1-\epsilon} + C' \delta,
$$

which proves the claim. Therefore, using this claim, we have

$$
\|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top - \boldsymbol{A}\boldsymbol{A}^\top\|_2
$$

$$
\leq \|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_2 + \left[-2^{-\frac{1}{2}} + 2(1+\sqrt{2}) \max_{t \in S} d_2(\widehat{A}^{(t)}, \overline{A}) + 2\sqrt{2} \frac{\epsilon}{1-\epsilon} \right]
$$

$$
\sqrt{2}a_1 \|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_2 + C a_1 \delta + C \sqrt{p} a_1^2
$$

$$
\leq \|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_2 - \gamma a_1 \|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_2 + C a_1 \delta + C \sqrt{p} a_1^2,
$$

when $2(2+\sqrt{2}) \max_{t \in S} d_2(\widehat{A}^{(t)}, \overline{A}) + 4 \frac{\epsilon}{1-\epsilon} < 1 - \gamma$.

Therefore, we can upper bound the term in (14) as

$$
[2] \leq -\frac{\lambda}{T\sqrt{n}} \cdot \gamma \cdot a_1 \cdot \sum_{t \in \mathcal{A}_1} \|\widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^\top - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^\top\|_2 + C\frac{\lambda}{\sqrt{n}}\gamma \cdot a_1 \delta + C\frac{\lambda}{\sqrt{n}} \cdot \gamma \cdot \sqrt{p}a_1^2.
$$

Therefore, using the bounds we have proved for [1] and [2], plugging them back in [\(13\)](#page-60-0), we have Hence

$$
G_{\mathcal{A}_1}(\{\mathbf{A}^{(t)}\}_{t=1}^T,\mathbf{A},\mathbf{\Theta}) - G_{\mathcal{A}_1}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^T,\widehat{\mathbf{A}},\widehat{\mathbf{\Theta}})
$$

\$\leq \frac{C}{T}\sum_{t\in\mathcal{A}_1}\sqrt{\frac{p+\log T}{n}}a_1\zeta^{(t)}\sqrt{r}\cdot\|\widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^\top - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^\top\|_2 + Ca_1\left(\frac{r+\log T}{n} + h^2\cdot\frac{1}{T}\sum_{t\in\mathcal{A}_1}(\zeta^{(t)})^2\right)\$

$$
+ C_{\overline{T}}^{\underline{a_1}} \delta \sqrt{\frac{p + \log T}{n}} \sum_{t \in A_1} \zeta^{(t)} - \frac{\lambda}{T\sqrt{n}} \cdot \gamma \cdot a_1 \cdot \sum_{t \in A_1} \|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_2
$$

+ $C_{\overline{\sqrt{n}}}^{\underline{\lambda}} \gamma \cdot a_1 \delta + C_{\overline{\sqrt{n}}}^{\underline{\lambda}} \cdot \gamma \cdot \sqrt{p} a_1^2$

$$
\leq C a_1 \left(\frac{r + \log T}{n} + h^2 \cdot \frac{1}{T} \sum_{t \in A_1} (\zeta^{(t)})^2\right) + C_{\overline{T}}^{\underline{a_1}} \delta \sqrt{\frac{p + \log T}{n}} \sum_{t \in A_1} \zeta^{(t)}
$$

$$
- \frac{\lambda}{T\sqrt{n}} \cdot \gamma \cdot a_1 \cdot \sum_{t \in A_1} \|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_2 + C_{\overline{\sqrt{n}}}^{\underline{\lambda}} \gamma \cdot a_1 \delta + C_{\overline{\sqrt{n}}}^{\underline{\lambda}} \cdot \gamma \cdot \sqrt{p} a_1^2.
$$

Considering
$$
G_{\mathcal{A}_2}(\{\mathbf{A}^{(t)}\}_{t=1}^T, \overline{\mathbf{A}}, \Theta) - G_{\mathcal{A}_2}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^T, \overline{\mathbf{A}}, \widehat{\Theta}),
$$
 first we have
\n
$$
\frac{1}{T} \sum_{t \in \mathcal{A}_2} [\nabla f^{(t)}(\mathbf{A}^{(t)*}\theta^{(t)*})]^\top a_1^{(t)} \cdot (\overline{\mathbf{A}}\theta_{\overline{\mathbf{A}}}^{(t)} - \widehat{\mathbf{A}}\theta_{\widehat{\mathbf{A}}}^{(t)} + \widehat{\mathbf{A}}\theta_{\widehat{\mathbf{A}}}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\theta}^{(t)})
$$
\n
$$
\leq \frac{1}{T} \left\| \left\{ \sqrt{a_1^{(t)}} \nabla f^{(t)}(\mathbf{A}^{(t)*}\theta^{(t)*}) \right\}_{t \in \mathcal{A}_2} \right\|_2 \sqrt{2r} \cdot \left\| \left\{ \sqrt{a_1^{(t)}} (\overline{\mathbf{A}}\theta_{\overline{\mathbf{A}}}^{(t)} - \widehat{\mathbf{A}}\theta_{\widehat{\mathbf{A}}}^{(t)}) \right\}_{t \in \mathcal{A}_2} \right\|_F
$$
\n
$$
+ \frac{1}{T} \sum_{t \in \mathcal{A}_2} \|\nabla f^{(t)}(\mathbf{A}^{(t)*}\theta^{(t)*})\|_2 \cdot a_1^{(t)} \cdot \|\widehat{\mathbf{A}}\theta_{\widehat{\mathbf{A}}}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\theta}^{(t)}\|_2
$$
\n
$$
\leq \frac{1}{T} \sqrt{2r} C \cdot \sqrt{\frac{p+T}{n}} a_1 \left\| \left\{ (\overline{\mathbf{A}}\theta_{\overline{\mathbf{A}}}^{(t)} - \widehat{\mathbf{A}}\theta_{\widehat{\mathbf{A}}}^{(t)}) \right\}_{t \in \mathcal{A}_2} \right\|_F + \frac{1}{T} \sum_{t \in \mathcal{A}_2} \sqrt{\frac{p + \log T}{n}} a_1 \cdot \|\widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(
$$

$$
\leq C \frac{r(p+T)}{nT}a_1 + \frac{C}{T}a_1 \left\| \left\{ (\overline{A}\boldsymbol{\theta}_{\overline{A}}^{(t)} - \widehat{\overline{A}}\boldsymbol{\theta}_{\widehat{\overline{A}}}^{(t)}) \right\}_{t \in \mathcal{A}_2} \right\|_{\mathrm{F}}^2 + \frac{1}{T} \sum_{t \in \mathcal{A}_2} \sqrt{\frac{p + \log T}{n}} a_1 \cdot \| \widehat{A}^{(t)} (\widehat{A}^{(t)})^\top - \widehat{\overline{A}} (\widehat{\overline{A}})^\top \|_2,
$$

where the second inequality holds because

$$
\left\| \left\{ \sqrt{a_1^{(t)}} \nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}) \right\}_{t \in \mathcal{A}_2} \right\|_2 \lesssim \max_{t \in \mathcal{A}_2} \sqrt{a_1^{(t)}} \cdot \sqrt{\frac{p+T}{n}},
$$

w.p. at least $1 - e^{-C(p + \log T)}$, which is due to Lemma 5.39 in [Vershynin](#page-39-5) [\(2010\)](#page-39-5). Therefore, by using the same arguments to bound $G_{A_1}(\{A^{(t)}\}_{t=1}^T, \overline{A}, \Theta) - G_{A_1}(\{\widehat{A}^{(t)}\}_{t=1}^T, \overline{A}, \widehat{\Theta})$, we can prove that

$$
G_{\mathcal{A}_2}(\{\mathbf{A}^{(t)}\}_{t=1}^T, \mathbf{A}, \Theta) - G_{\mathcal{A}_2}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^T, \widehat{\overline{\mathbf{A}}}, \widehat{\Theta})
$$

\$\leq \frac{1}{2T}(\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)})^\top \widehat{\Sigma}^{(t)}(\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)})\$

$$
+ C \frac{r(p+T)}{nT} a_1^2 + \frac{C}{T} \left\| \left\{ (\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)} - \widehat{\overline{\mathbf{A}}}\boldsymbol{\theta}_{\widehat{\overline{\mathbf{A}}}^{(t)}) \right\}_{t \in \mathcal{A}_2} \right\|_{\mathrm{F}}^2 + \frac{1}{T} \sum_{t \in \mathcal{A}_2} \sqrt{\frac{p + \log T}{n}} a_1 \cdot \| \widehat{\mathbf{A}}^{(t)} (\widehat{\mathbf{A}}^{(t)})^\top - \widehat{\overline{\mathbf{A}}} (\widehat{\overline{\mathbf{A}}})^\top \|_2
$$

$$
+\frac{C}{T}\left(\sqrt{T}\sqrt{a_1}\sqrt{\frac{r+\log T}{n}}+h\sqrt{\sum_{t\in\mathcal{A}_2}(\zeta^{(t)})^2a_1}\right)\sqrt{\sum_{t\in\mathcal{A}_2}a_1\|\bar{A}\theta_{\bar{A}}^{(t)}-\hat{A}^{(t)}\hat{\theta}^{(t)}\|_2^2}
$$

$$
-\frac{C}{T}\sum_{t\in\mathcal{A}_2}a_1\|\hat{A}^{(t)}\hat{\theta}^{(t)}-\bar{A}\theta_{\bar{A}}^{(t)}\|_2^2-\frac{\lambda}{T\sqrt{n}}\cdot\gamma a_1\cdot\sum_{t\in\mathcal{A}_2}\|\hat{A}^{(t)}(\hat{A}^{(t)})^\top-\hat{A}(\hat{A})^\top\|_2
$$

$$
\leq-\frac{C}{T}\sum_{t\in\mathcal{A}_2}a_1\|\hat{A}^{(t)}\hat{\theta}^{(t)}-\bar{A}\theta_{\bar{A}}^{(t)}\|_2^2-\frac{\lambda}{T\sqrt{n}}\cdot\gamma a_1\cdot\sum_{t\in\mathcal{A}_2}\|\hat{A}^{(t)}(\hat{A}^{(t)})^\top-\hat{A}(\hat{A})^\top\|_2
$$

$$
+Ca_1\left(\frac{pr}{nT}+\frac{r+\log T}{n}+h^2\cdot\frac{1}{T}\sum_{t\in\mathcal{A}_2}(\zeta^{(t)})^2\right).
$$
(17)

 $\text{Since } \|\widehat{A}^{(t)}\widehat{\theta}^{(t)} - \widehat{\overline{A}}\theta_{\widehat{\overline{A}}}^{(t)}\|_2 \leq \|\widehat{A}^{(t)}(\widehat{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_2 (\|\theta_{\widehat{\overline{A}}}^{(t)}\|_2)$ $\begin{aligned} &\frac{(t)}{\widehat{\bm{A}}}\|\mathbf{2}+\|\nabla f^{(t)}(\widehat{\overline{\bm{A}}}\bm{\theta}^{(t)}_{\widehat{\bm{A}}})\|_2)\leq C\zeta^{(t)}\|\widehat{\bm{A}}^{(t)}(\widehat{\bm{A}}^{(t)})^\top - \end{aligned}$ $\overline{A}(\overline{A})^{\top} \|_2 \leq C \zeta^{(t)} \|\overline{A}(\overline{A})^{\top} - \overline{A}(\overline{A})^{\top} \|_2 \frac{c}{\sqrt{r}}$, we have

$$
\begin{aligned}\n\|\widehat{\boldsymbol{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}-\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}\|_2^2 &\geq \frac{1}{2}\|\widehat{\overline{\boldsymbol{A}}}\boldsymbol{\theta}_{\widehat{\overline{\boldsymbol{A}}}}^{(t)}-\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}\|_2^2 - 2\|\widehat{\boldsymbol{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}-\widehat{\overline{\boldsymbol{A}}}\boldsymbol{\theta}_{\widehat{\overline{\boldsymbol{A}}}^{(t)}}\|_2^2 \\
&\geq \frac{1}{2}\|\widehat{\overline{\boldsymbol{A}}}\boldsymbol{\theta}_{\widehat{\overline{\boldsymbol{A}}}}^{(t)}-\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}\|_2^2 - \frac{2C^2c}{r}(\boldsymbol{\zeta}^{(t)})^2\|\overline{\boldsymbol{A}}(\overline{\boldsymbol{A}})^\top-\widehat{\overline{\boldsymbol{A}}}(\widehat{\overline{\boldsymbol{A}}})^\top\|_2^2.\n\end{aligned}
$$

This implies that

$$
G_{\mathcal{A}_2}(\{\mathbf{A}^{(t)}\}_{t=1}^T,\mathbf{A},\mathbf{\Theta})-G_{\mathcal{A}_2}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^T,\widehat{\mathbf{A}},\widehat{\mathbf{\Theta}})
$$

\n
$$
\leq -\frac{C'}{T}a_1\|\widehat{\mathbf{A}}\widehat{\mathbf{\Theta}}-\overline{\mathbf{A}}\overline{\mathbf{\Theta}}\|_{\mathrm{F}}^2+\frac{2C^2c}{r}a_1(\zeta^{(t)})^2\|\overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}-\widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^{\top}\|_2^2
$$

\n
$$
+Ca_1\left(\frac{pr}{nT}+\frac{r+\log T}{n}+h^2\cdot\frac{1}{T}\sum_{t\in\mathcal{A}_2}(\zeta^{(t)})^2\right).
$$

In addition, by triangle inequality,

$$
G_{S^c}(\{\boldsymbol{A}^{(t)}\}_{t=1}^T,\boldsymbol{A},\boldsymbol{\Theta})-G_{S^c}(\{\widehat{\boldsymbol{A}}^{(t)}\}_{t=1}^T,\widehat{\overline{\boldsymbol{A}}},\widehat{\boldsymbol{\Theta}})\leq \frac{\lambda}{\sqrt{n}T}|S^c|\|\widehat{\overline{\boldsymbol{A}}}(\widehat{\overline{\boldsymbol{A}}})^\top-\boldsymbol{A}\boldsymbol{A}^\top\|_2.
$$

Recall (15) :

$$
\boldsymbol{A}\boldsymbol{A}^{\top} = (1 - 2a_1)\widehat{\boldsymbol{A}}(\widehat{\boldsymbol{A}})^{\top} + a_1[\overline{\boldsymbol{A}}(\widehat{\boldsymbol{A}})^{\top} + \widehat{\boldsymbol{A}}(\overline{\boldsymbol{A}})^{\top}] + \mathcal{O}(a_1^2).
$$

Hecne we have

$$
\|\mathbf{A}\mathbf{A}^{\top} - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}\|_{2} \le a_{1} \|\widehat{\mathbf{A}}(\overline{\mathbf{A}} - \widehat{\mathbf{A}})^{\top}\|_{2} + a_{1} \|\overline{\mathbf{A}}(\widehat{\mathbf{A}} - \overline{\mathbf{A}})^{\top}\|_{2} + C a_{1}^{2}
$$

\n
$$
\le 2a_{1}d_{2}(\overline{\mathbf{A}}, \widehat{\mathbf{A}}) + C a_{1} \delta + C a_{1}^{2}
$$

\n
$$
\le 2\sqrt{2}a_{1} \|\overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top} - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}\|_{2} + C a_{1} \delta + C a_{1}^{2},
$$

where the second inequality is due to [\(16\)](#page-62-1). Therefore,

$$
G_{S^c}(\{\mathbf{A}^{(t)}\}_{t=1}^T,\mathbf{A},\mathbf{\Theta})-G_{S^c}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^T,\widehat{\overline{\mathbf{A}}},\widehat{\mathbf{\Theta}}) \leq 2\sqrt{2}a_1 \frac{\lambda}{\sqrt{n}} |S^c| \|\overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top} - \widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^{\top} \|_2 + Ca_1 \delta + Ca_1^2.
$$

Case 1: If $|\mathcal{A}_2| \geq \left(1 - cr^{-1} \frac{\overline{\zeta}^2}{\max_{t \in S}(\zeta^{(t)})^2}\right)|S|$:

By Wedin's sin Θ-theorem, we have By Assumption [2,](#page-10-0) Wedin's sin Θ-Theorem, and applying Lemma [36](#page-51-5) on $\{\boldsymbol{\theta}^{(t)}_{\overline{\boldsymbol{\mathcal{A}}}}\}$ $\{\boldsymbol{\bar{A}}^{(t)}\}_{t\in\mathcal{A}_2},$

$$
\|\mathcal{N}(\widehat{\overline{A}})(\mathcal{N}(\widehat{\overline{A}}))^{\top} - \overline{A}(\overline{A})^{\top}\|_{2} \leq \|\mathcal{N}(\widehat{\overline{A}})(\mathcal{N}(\widehat{\overline{A}})^{\top} - \overline{A}(\overline{A})^{\top}\|_{\mathrm{F}} \lesssim \sqrt{\frac{r}{|S|}}\bar{\zeta}^{-1} \cdot \|\{\widehat{\overline{A}}\theta_{\widehat{\overline{A}}}^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\}_{t \in \mathcal{A}_{2}}\|_{\mathrm{F}}.
$$

Here we used the fact that $\frac{1}{|S|} \sum_{t \in S} \theta_{\overline{A}}^{(t)}$ $\frac{(t)}{\overline{A}}(\bm{\theta}_{\overline{A}}^{(t)}$ $\frac{1}{\mathbf{A}}^{(t)}$ ^T $\succeq \frac{c}{r}$ $\frac{c}{r}\bar{\zeta}^2\mathbf{I}_p$. To see this, notice that

$$
\sigma_r \left(\frac{1}{\sqrt{|S|}} \{ \overline{A} \boldsymbol{\theta}_{\overline{A}}^{(t)} \}_{t \in S} \right) = \sigma_r \left(\frac{1}{\sqrt{|S|}} \{ \boldsymbol{\beta}^{(t)*} \}_{t \in S} \right) - \frac{1}{\sqrt{|S|}} || \{ \overline{A} \boldsymbol{\theta}_{\overline{A}}^{(t)} \}_{t \in S} - \{ \boldsymbol{\beta}^{(t)*} \}_{t \in S} ||_2
$$

$$
\geq \frac{c}{\sqrt{r}} \bar{\zeta} - \left(\bar{\zeta} h + \sqrt{\frac{r + \log T}{n}} \right)
$$

$$
\geq \frac{c'}{\sqrt{r}} \bar{\zeta},
$$

where by Lemmas [30](#page-50-4) and [34,](#page-51-3)

$$
\frac{1}{\sqrt{|S|}}\|\{\overline{\bm{A}}\bm{\theta}^{(t)}_{\overline{\bm{A}}}\}_{t\in S}-\{\bm{\beta}^{(t)*}\}_{t\in S}\|_2 \leq \sqrt{\sum_{t\in S}\|\overline{\bm{A}}\bm{\theta}^{(t)}_{\overline{\bm{A}}}-\bm{A}^{(t)*}\bm{\theta}^{(t)*}\|_2^2} \lesssim h\bar{\zeta}+\sqrt{\frac{r+\log T}{n}},
$$

and the last step comes from conditions (i) and (ii) by noticing that

$$
\sqrt{\frac{r + \log T}{n}} \lesssim \frac{\bar{\zeta}}{\sqrt{r}} \cdot \sqrt{\frac{p + \log T}{n}} \cdot \frac{1}{\zeta^{(t)}} \lesssim \frac{\bar{\zeta}}{\sqrt{r}}.
$$

This leads to

$$
\sigma_r\left(\frac{1}{|S|}\sum_{t\in S}\theta_{\overline{A}}^{(t)}(\theta_{\overline{A}}^{(t)})^\top\right) = \sigma_r\left(\frac{1}{|S|}\sum_{t\in S}\theta_{\overline{A}}^{(t)}(\overline{A})^\top \overline{A}(\theta_{\overline{A}}^{(t)})^\top\right) \ge \frac{c'}{r}\overline{\zeta}^2.
$$

Hence

$$
G_{[T]}(\{\mathbf{A}^{(t)}\}_{t=1}^{T}, \mathbf{A}, \Theta) - G_{[T]}(\{\hat{\mathbf{A}}^{(t)}\}_{t=1}^{T}, \hat{\mathbf{A}}, \hat{\Theta})
$$
\n
$$
\leq [G_{\mathcal{A}_{1}}(\{\mathbf{A}^{(t)}\}_{t=1}^{T}, \mathbf{A}, \Theta) - G_{\mathcal{A}_{1}}(\{\hat{\mathbf{A}}^{(t)}\}_{t=1}^{T}, \hat{\mathbf{A}}, \hat{\Theta})] + [G_{\mathcal{A}_{2}}(\{\mathbf{A}^{(t)}\}_{t=1}^{T}, \mathbf{A}, \Theta) - G_{\mathcal{A}_{2}}(\{\hat{\mathbf{A}}^{(t)}\}_{t=1}^{T}, \hat{\mathbf{A}}, \hat{\Theta})] + [G_{\mathcal{S}^{c}}(\{\hat{\mathbf{A}}^{(t)}\}_{t=1}^{T}, \hat{\mathbf{A}}, \hat{\Theta})] + [G_{\mathcal{S}^{c}}(\{\hat{\mathbf{A}}^{(t)}\}_{t=1}^{T}, \hat{\mathbf{A}}, \hat{\Theta})] \leq Ca_{1}\left(\frac{r + \log T}{n} + h^{2} \cdot \frac{1}{T} \sum_{t \in \mathcal{A}_{1}} (\zeta^{(t)})^{2}\right) + C_{T}^{\frac{a_{1}}{T}} \delta \sqrt{\frac{p + \log T}{n}} \sum_{t \in \mathcal{A}_{1}} \zeta^{(t)} + C_{\frac{\lambda}{\sqrt{n}} \gamma \cdot a_{1} \delta + C_{\frac{\lambda}{\sqrt{n}} \cdot \gamma \cdot \sqrt{p} a_{1}^{2}} \cdot \frac{1}{T} \sum_{t \in \mathcal{A}_{2}} (\zeta^{(t)})^{2}\right) + 2\sqrt{2}a_{1} \frac{\lambda}{\sqrt{n}T} |S^{c}||\mathbf{A}(\mathbf{A})^{T} - \mathbf{A}(\hat{\mathbf{A}})^{T}||_{2}
$$
\n
$$
\leq Ca_{1}\left(\frac{pr}{nT} + \frac{r + \log T}{n} + h^{2}\zeta^{2}\right) - Ca_{1}\frac{\zeta^{2}}{r} ||\mathbf{A}(\hat{\mathbf{A}})^{T} - \mathbf{A}(\mathbf{A})^{T} ||_{2}^{2} + 2\sqrt{2}a_{1} \frac{\lambda}{\sqrt{n}T} |S^{c}||
$$

w.p. at least $1 - e^{-C'(r + \log T)}$, if $\|\widehat{A}(\widehat{A})^{\top} - \overline{A}(\overline{A})^{\top}\|_2 > C\overline{\zeta}^{-1} \left(\sqrt{\frac{pr}{nT}} + \sqrt{\frac{r + \log T}{n}}\right)$ $\frac{\log T}{n}$ + Ch + $C\frac{r\lambda}{\bar{\zeta}^2\sqrt{n}}$ $|S^c|$ $\frac{S^c}{T}$ and $Ca_1\delta^2 + C\frac{a_1}{T}\delta\sqrt{\frac{p+\log T}{n}}$ $\frac{\log T}{n} \sum_{t \in S} \zeta^{(t)} + C \frac{\lambda}{\sqrt{n}}$ $\frac{\Delta}{n}\gamma\cdot a_1\delta+C\frac{\lambda}{\sqrt{n}}$ $\frac{\Delta}{n} \cdot \gamma \cdot \sqrt{p} a_1^2 < C a_1 \frac{p r}{n T} +$ $Ca_1\frac{r+\log T}{n}$ $\frac{\log T}{n}$. Note that the latter can easily hold because we are free to choose arbitrary small a_1 and δ . And as $a_1 \to 0^+$, we have $\mathbf{A}^{(t)} \to \widehat{\mathbf{A}}^{(t)}$, $\mathbf{A} \to \overline{\mathbf{A}}$, $\boldsymbol{\theta}^{(t)} \to \widehat{\boldsymbol{\theta}}^{(t)}$, in the sense of $\|\cdot\|_{\max}$. Therefore [\(18\)](#page-67-0) contradicts with the local optimality of $(\{\widehat{A}^{(t)}\}_{t=1}^T, \overline{A}, \{\widehat{\theta}^{(t)}\}_{t=1}^T)$ in terms of $G_{[T]}(\{\hat{A}^{(t)}\}_{t=1}^T, \overline{A}, \widehat{\Theta})$. Therefore, w.p. at least $1 - e^{-C'(r + \log T)}$, we must have

$$
\|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A}(\overline{A})^{\top}\|_2 \leq C\overline{\zeta}^{-1} \left(r\sqrt{\frac{p}{nT}} + \sqrt{r}\sqrt{\frac{r + \log T}{n}} \right) + C\sqrt{r}h + C\frac{r\lambda}{\overline{\zeta}^2\sqrt{n}} \frac{|S^c|}{T},
$$

when $Ca_1\delta^2 + C \frac{a_1}{T} \delta \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n} \sum_{t \in S} \zeta^{(t)} + C \frac{\lambda}{\sqrt{n}}$ $\frac{1}{n}\gamma \cdot a_1 \delta + C \frac{\lambda}{\sqrt{n}}$ $\frac{1}{n} \cdot \gamma \cdot \sqrt{p} a_1^2 < C a_1 \frac{pr}{nT} + C a_1 \frac{r + \log T}{n}$ $\frac{\log T}{n}$. **Case 2:** If $|\mathcal{A}_1| \geq cr^{-1} \frac{\bar{\zeta}^2}{\max_{t \in S} (\zeta^{(t)})^2} |S|$: Note that by [\(17\)](#page-65-0), we have

$$
G_{\mathcal{A}_2}(\{\mathbf{A}^{(t)}\}_{t=1}^T,\mathbf{A},\boldsymbol{\Theta}) - G_{\mathcal{A}_2}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^T,\widehat{\overline{\mathbf{A}}},\widehat{\boldsymbol{\Theta}})
$$

\n
$$
\leq \frac{1}{2T}(\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)})^\top \widehat{\boldsymbol{\Sigma}}^{(t)}(\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)})
$$

\n
$$
+ \frac{C}{T}a_1\sqrt{\frac{p+T}{n}}\sqrt{r}\left\|\{\overline{\mathbf{A}}\boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)} - \widehat{\overline{\mathbf{A}}}\boldsymbol{\theta}_{\widehat{\overline{\mathbf{A}}}^{(t)}\}_{t\in\mathcal{A}_2}\right\|_{\mathrm{F}} + \frac{1}{T}\sum_{t\in\mathcal{A}_2}\sqrt{\frac{p+\log T}{n}}a_1 \cdot \|\widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^\top - \widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^\top\|_2
$$

$$
+ \frac{C}{T} \left(\sqrt{T} \sqrt{a_1} \sqrt{\frac{r + \log T}{n}} + h \sqrt{\sum_{t \in A_2} (\zeta^{(t)})^2 a_1} \right) \sqrt{\sum_{t \in A_2} a_1 \|\vec{A} \theta_{\vec{A}}^{(t)} - \hat{A}^{(t)} \hat{\theta}^{(t)}\|_2^2}
$$

\n
$$
- \frac{C}{T} \sum_{t \in A_2} a_1 \|\hat{A}^{(t)} \hat{\theta}^{(t)} - \overline{A} \theta_{\vec{A}}^{(t)}\|_2^2 - \frac{\lambda}{T \sqrt{n}} \cdot \gamma a_1 \cdot \sum_{t \in A_2} \|\hat{A}^{(t)} (\hat{A}^{(t)})^\top - \hat{A} (\hat{A})^\top \|_2
$$

\n
$$
\leq - \frac{C}{T} \sum_{t \in A_2} a_1 \|\hat{A}^{(t)} \hat{\theta}^{(t)} - \overline{A} \theta_{\vec{A}}^{(t)}\|_2^2 - \frac{\lambda}{T \sqrt{n}} \cdot \gamma a_1 \cdot \sum_{t \in A_2} \|\hat{A}^{(t)} (\hat{A}^{(t)})^\top - \hat{A} (\hat{A})^\top \|_2
$$

\n
$$
+ \frac{C}{T} \left(\sqrt{T} \sqrt{a_1} \sqrt{\frac{r + \log T}{n}} + h \sqrt{\sum_{t \in A_2} (\zeta^{(t)})^2 a_1} \right) \sqrt{\sum_{t \in A_2} a_1 \|\vec{A} \theta_{\vec{A}}^{(t)} - \hat{A}^{(t)} \hat{\theta}^{(t)}\|_2^2}
$$

\n
$$
+ \frac{C}{T} a_1 \sqrt{\frac{p + T}{n}} \sqrt{r} \left\| \{\overline{A} \theta_{\vec{A}}^{(t)} - \hat{A}^{(t)} \hat{\theta}^{(t)}\}_{t \in A_2} \right\|_F + \frac{C}{T} a_1 \sqrt{\frac{p + T}{n}} \sqrt{r} \left\| \{\hat{A}^{(t)} \hat{\theta}^{(t)} - \hat{\overline{A}} \theta_{\hat{\overline{A}}}^{(t)}\}_{t \in A_2} \right\|_F
$$

\n
$$
\leq C a_1 \left(\frac{pr
$$

Therefore,

$$
G_{[T]}(\{\mathbf{A}^{(t)}\}_{t=1}^{T}, \mathbf{A}, \Theta) - G_{[T]}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^{T}, \widehat{\mathbf{A}}, \widehat{\Theta})
$$
\n
$$
\leq [G_{\mathcal{A}_{1}}(\{\mathbf{A}^{(t)}\}_{t=1}^{T}, \mathbf{A}, \Theta) - G_{\mathcal{A}_{1}}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^{T}, \widehat{\mathbf{A}}, \widehat{\Theta})] + [G_{\mathcal{A}_{2}}(\{\mathbf{A}^{(t)}\}_{t=1}^{T}, \mathbf{A}, \Theta) - G_{\mathcal{A}_{2}}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^{T}, \widehat{\mathbf{A}}, \widehat{\Theta})] + [G_{S^{c}}(\{\mathbf{A}^{(t)}\}_{t=1}^{T}, \mathbf{A}, \Theta) - G_{S^{c}}(\{\widehat{\mathbf{A}}^{(t)}\}_{t=1}^{T}, \widehat{\mathbf{A}}, \widehat{\Theta})]
$$
\n
$$
\leq C a_{1} \left(\frac{r + \log T}{n} + h^{2} \cdot \frac{1}{T} \sum_{t \in \mathcal{A}_{1}} (\zeta^{(t)})^{2}\right) + C \frac{a_{1}}{T} \delta \sqrt{\frac{p + \log T}{n}} \sum_{t \in \mathcal{A}_{1}} \zeta^{(t)} + C \frac{\lambda}{\sqrt{n}} \gamma \cdot a_{1} \delta + C \frac{\lambda}{\sqrt{n}} \cdot \gamma \cdot \sqrt{p} a_{1}^{2}
$$
\n
$$
- \frac{\lambda}{T \sqrt{n}} \cdot \gamma \cdot a_{1} \cdot \sum_{t \in \mathcal{A}_{1}} \| \widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^{T} - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{T} \|_{2} + C a_{1} \left(\frac{pr}{nT} + \frac{r + \log T}{n} + h^{2} \zeta^{2}\right)
$$
\n
$$
+ 2\sqrt{2} a_{1} \frac{\lambda}{\sqrt{nT}} |S^{c}||\overline{\mathbf{A}}(\widehat{\mathbf{A}})^{T} - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{T} \|_{2} + C a_{1} \sqrt
$$

$$
+ C \frac{\lambda}{\sqrt{n}} \gamma \cdot a_1 \delta + C \frac{\lambda}{\sqrt{n}} \cdot \gamma \cdot \sqrt{p} a_1^2 + C \frac{a_1}{T} \delta \sqrt{\frac{p + \log T}{n}} \sum_{t \in S} \zeta^{(t)} + C a_1 \sqrt{\frac{p + T}{nT}} \bar{\zeta} \cdot \|\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top} - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}\|_2
$$

\n
$$
\leq C a_1 \frac{p + \log T}{n} - C' \frac{\lambda}{\sqrt{n} T r} \cdot \gamma \frac{\bar{\zeta}^2}{\max_{t \in S} (\zeta^{(t)})^2} \cdot \|\overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top} - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}\|_2 + C \frac{\lambda}{\sqrt{n}} \gamma \cdot a_1 \delta + C \frac{\lambda}{\sqrt{n}} \cdot \gamma \cdot \sqrt{p} a_1^2
$$

\n
$$
+ C \frac{a_1}{T} \delta \sqrt{\frac{p + \log T}{n}} \sum_{t \in S} \zeta^{(t)} + C a_1 \sqrt{\frac{p + T}{n T}} \bar{\zeta} \cdot \|\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top} - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}\|_2
$$

\n
$$
\leq C a_1 \frac{p + \log T}{n} - \frac{C'}{2} \frac{\lambda}{\sqrt{n} T r} \cdot \gamma \frac{\bar{\zeta}^2}{\max_{t \in S} (\zeta^{(t)})^2} \cdot \|\overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top} - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}\|_2 + C \frac{\lambda}{\sqrt{n}} \gamma \cdot a_1 \delta + C \frac{\lambda}{\sqrt{n}} \cdot \gamma \cdot \sqrt{p} a_1^2
$$

\n
$$
+ C \frac{a_1}{T} \delta \sqrt{\frac{p + \log T}{n}} \sum_{t \in S} \zeta^{(t)}
$$

\n
$$
< 0,
$$

w.p. at least $1-e^{-C'(r+\log T)}$, when $\|\overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}-\overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}\|_2 \geq C\eta$, $C\frac{\lambda}{\sqrt{2\pi}}$ $\frac{\Delta}{n} \gamma \cdot a_1 \delta + C \frac{\lambda}{\sqrt{n}}$ $\frac{1}{\overline{n}} \cdot \gamma \cdot \sqrt{p} a_1^2 <$ $Ca_1\frac{p+\log T}{n}$ $\frac{\log T}{n}$, with a large $C'' > 0$, and $\min_{t \in S} \zeta^{(t)} \gtrsim \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$. Note that

$$
\frac{\min_{t \in S} \zeta^{(t)}}{\bar{\zeta}} \cdot r \lesssim \sqrt{p + \log T}, \quad \frac{\min_{t \in S} \zeta^{(t)}}{\bar{\zeta}} \cdot \frac{r}{\sqrt{T}} \lesssim 1. \tag{19}
$$

We used the fact that

$$
\lambda \approx \sqrt{r(p + \log T)} \cdot \frac{\max_{t \in S} (\zeta^{(t)})^2}{\min_{t \in S} \zeta^{(t)}} \times \left[\sqrt{pr} \frac{\bar{\zeta}}{\min_{t \in S} \zeta^{(t)}} + \sqrt{r(p + \log T)} \cdot \frac{\bar{\zeta}}{\min_{t \in S} \zeta^{(t)}} \right] \cdot \frac{\max_{t \in S} (\zeta^{(t)})^2}{\bar{\zeta}},
$$

where we used (19) in the second inequality. This contradicts with the **local** optimality of $({\{\widehat{A}^{(t)}\}_{t=1}^T}, \overline{A}, {\{\widehat{\theta}^{(t)}\}_{t=1}^T})$ in terms of $G_{[T]}({\{\widehat{A}^{(t)}\}_{t=1}^T}, \overline{A}, \widehat{\Theta})$. Therefore, w.p. at least $1 - e^{-C'(r + \log T)}$, we must have

$$
\|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A}(\overline{A})^{\top}\|_2 \leq C\overline{\zeta}^{-1} \left(r\sqrt{\frac{p}{nT}} + \sqrt{r}\sqrt{\frac{r + \log T}{n}} \right) + C\sqrt{r}h + C\frac{r\lambda}{\overline{\zeta}^2\sqrt{n}} \frac{|S^c|}{T},
$$

when $C \frac{a_1}{T} \delta \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n} \sum_{t \in S} \zeta^{(t)} + C \frac{\lambda}{\sqrt{n}}$ $\frac{\Delta}{n} \gamma \cdot a_1 \delta + C \frac{\lambda}{\sqrt{n}}$ $\frac{1}{n} \cdot \gamma \cdot \sqrt{p} a_1^2 < C a_1 \frac{pr}{nT} + C a_1 \frac{r + \log T}{n}$ $\frac{\log T}{n}$. Until now, we have proved that

$$
\|\widehat{\overline{\boldsymbol{A}}}(\widehat{\overline{\boldsymbol{A}}})^\top - \overline{\boldsymbol{A}}(\overline{\boldsymbol{A}})^\top\|_2 \leq C\bar{\zeta}^{-1}\bigg(r\sqrt{\frac{p}{nT}} + \sqrt{r}\sqrt{\frac{r+\log T}{n}}\bigg) + C\sqrt{r}h + C\frac{r\lambda}{\bar{\zeta}^2\sqrt{n}}\frac{|S^c|}{T},
$$

for any local minimizers $(\{\widehat{A}^{(t)}\}_{t=1}^T, \overline{A}, \{\widehat{\theta}^{(t)}\}_{t=1}^T)$ of $G_{[T]}(\{\widehat{A}^{(t)}\}_{t=1}^T, \overline{A}, \widehat{\Theta})$ satisfying $2(2+\sqrt{2})\max_{t\in S}d_2(\widehat{\boldsymbol{A}}^{(t)},\overline{\boldsymbol{A}})+4\frac{\epsilon}{1-\epsilon}<1-\gamma,$

$$
\max_{t \in S} \left\{ \frac{\sigma_{\max}(\boldsymbol{\Sigma}^{(t)})}{\sigma_{\min}(\boldsymbol{\Sigma}^{(t)})} d_2(\widehat{\boldsymbol{A}}^{(t)}, \boldsymbol{A}^{(t)*}) \right\} < 1 - \gamma.
$$

The remaining proof follows the arguments in the proof of Theorem [1](#page-12-0) and pushing $\delta \to 0$.

D.6 Proof of Theorem [6](#page-16-0)

We briefly point out how the arguments in the proofs of theorems in Section [2.2.1](#page-11-1) can be modified to prove the same results for the revised Algorithm [1](#page-11-0) which solves Step 1 in the entire $\mathbb{R}^{p \times r}$ space with the penalty on projection matrices.

We first discuss the results of global minimizers (Theorems [1](#page-12-0) and [2\)](#page-13-0).

- (i) If $\text{rank}(\widehat{A}^{(t)}) = \text{rank}(\widehat{A}) = r$ for all $t \in S$, then we can replace $\widehat{A}^{(t)}\theta_{\widehat{A}}^{(t)}$ $\hat{\vec{A}}^{(t)}$ and $\hat{\vec{A}} \theta_{\hat{\vec{A}}}^{(t)}$ with \boldsymbol{A} $\boldsymbol{\theta}^{(t)}_{\boldsymbol{A}}$ $A^{(t)}_A \coloneqq \arg\min_{\theta \in \mathbb{R}^r} f^{(t)}(A\theta)$ used in the proofs by $\mathcal{N}(\widehat{A}^{(t)})\theta_{\mathcal{N}(t)}^{(t)}$ $\mathcal{N}(\widehat{\bm{A}}^{(t)})$ and $\mathcal{N}(\widehat{\overline{\bm{A}}})\boldsymbol{\theta}_{\mathcal{N}(t)}^{(t)}$ $\overset{(\iota)}{\mathcal{N}(\widehat{\bar{\bm{A}}})},$ $\boldsymbol{A})$ where $\mathcal{N}(\widehat{A}^{(t)}) \coloneqq \widehat{A}^{(t)}[(\widehat{A}^{(t)})^{\top}\widehat{A}^{(t)}]^{-1}(\widehat{A}^{(t)})^{\top}$, and $\mathcal{N}(\overline{A}) = \overline{A}[(\overline{A})^{\top}\overline{A}]^{-1}(\overline{A})^{\top}$. It is straightforward to see that $\mathcal{N}(\widehat{\mathbf{A}}^{(t)})\boldsymbol{\theta}_{\mathcal{N}(t)}^{(t)}$ $\hat{A}^{(t)}_{\mathcal{N}(\widehat{\bm{A}}^{(t)})} = \widehat{\bm{A}}^{(t)}\bm{\theta}^{(t)}_{\widehat{\bm{A}}^{(t)}}$ $\overset{(t)}{\widehat{A}^{(t)}}$ and $\mathcal{N}(\widehat{\overline{A}})\theta^{(t)}_{\mathcal{N}(t)}$ $\mathcal{N}(\bm{A})$ $=\widehat{\overline{A}}\theta_{\widehat{\overline{A}}}^{(t)}$. We can also see that $P_{\hat{A}^{(t)}} = \mathcal{N}(\hat{A}^{(t)})(\mathcal{N}(\hat{A}^{(t)}))^{\top}$ and $P_{\hat{\overline{A}}} = \mathcal{N}(\overline{A})(\mathcal{N}(\overline{A}))^{\top}$. We can proceed the same proofs of Theorems [1](#page-12-0) and [2,](#page-13-0) with $\mathcal{N}(\widehat{A}^{(t)}), \mathcal{N}(\widehat{A}), \theta_{\widehat{A}}^{(t)}$ $\overset{(t)}{\widehat{A}^{(t)}},$ and $\overset{\boldsymbol{\theta}^{(t)}}{\widehat{A}}$ \boldsymbol{A} .
- (ii) If $\text{rank}(\hat{\mathbf{A}}^{(t)}) < r$ or $\text{rank}(\overline{\mathbf{A}}) < r$ for some $t \in S$, then for those t, we can define $\mathcal{N}(\hat{\mathbf{A}}^{(t)})$ and $\mathcal{N}(\overline{A})$ in a different way. Consider a matrix $A \in \mathbb{R}^{p \times r}$ with rank $(A) = r' < r$ and its SVD $\boldsymbol{A}_{p\times r} = \boldsymbol{U}_{p\times r'} \boldsymbol{\Lambda}_{r'\times r'} \boldsymbol{V}_{r'\times r'}^\top$, where $\boldsymbol{\Lambda}_{r'\times r'}$ is diagonal with positive entries and $\mathbf{U} \in \mathcal{O}^{p \times r'}$. Then we can define $\mathcal{N}(\mathbf{A}) = (\mathbf{U}_{p \times r'} \quad \mathbf{0}_{p \times (r-r')}) \in \mathbb{R}^{p \times r}$. Note that $\mathcal{N}(A)(\mathcal{N}(A))^{\top} = \boldsymbol{U}\boldsymbol{U}^{\top} = \boldsymbol{P}_A = \boldsymbol{P}_{\mathcal{N}(A)}$, and this definition of $\mathcal{N}(A)$ can be seen as an extension of the full-rank case. Then we can proceed the same proofs of Theorems [1](#page-12-0) and [2,](#page-13-0) with $\mathcal{N}(\widehat{\boldsymbol{A}}^{(t)}), \mathcal{N}(\widehat{\boldsymbol{A}}), \boldsymbol{\theta}_{\widehat{\boldsymbol{A}}}^{(t)}$ $\overset{(t)}{\widehat{A}^{(t)}},$ and $\overset{\boldsymbol{\theta}^{(t)}}{\widehat{A}}$ \boldsymbol{A} .

Then let us discuss the results of local minimizers (Theorems [4](#page-14-0) and [5\)](#page-14-1). Recall that we proved Theorems [4](#page-14-0) and [5](#page-14-1) by construction based on the local minimizers \overline{A} and $\widehat{A}^{(t)}$. And it suffices to replace \overline{A} and $\widehat{A}^{(t)}$ in the construction with $\mathcal{N}(\overline{A})$ and $\mathcal{N}\widehat{A}^{(t)}$ as defined above. In addition, if we want to show the stronger version of Theorem [5](#page-14-1) we proved in the proof, we need to rewrite the constraint set by replacing

$$
\label{eq:2} \begin{split} &2(2+\sqrt{2})\max_{t\in S}\min_{\mathbf{R}\in\mathcal{O}^{r\times r}}\|\widehat{\pmb{A}}^{(t)}-\overline{\pmb{A}}\pmb{R}\|_2+4\frac{\epsilon}{1-\epsilon}<1-\gamma,\\ &\max_{t\in S}\left\{\frac{\sigma_{\max}(\pmb{\Sigma}^{(t)})}{\sigma_{\min}(\pmb{\Sigma}^{(t)})}\min_{\pmb{R}\in\mathcal{O}^{r\times r}}\|\widehat{\pmb{A}}^{(t)}-\pmb{A}^{(t)*}\pmb{R}\|_2\right\}<1-\gamma. \end{split}
$$

with

$$
\begin{aligned} &2(2+\sqrt{2})\max_{t\in S}\min_{\mathbf{R}\in\mathcal{O}^{r\times r}}\|\mathcal{N}(\widehat{\boldsymbol{A}}^{(t)})-\overline{\boldsymbol{A}}\boldsymbol{R}\|_2+4\frac{\epsilon}{1-\epsilon}<1-\gamma,\\ &\max_{t\in S}\left\{\frac{\sigma_{\max}(\boldsymbol{\Sigma}^{(t)})}{\sigma_{\min}(\boldsymbol{\Sigma}^{(t)})}\min_{\mathbf{R}\in\mathcal{O}^{r\times r}}\|\mathcal{N}(\widehat{\boldsymbol{A}}^{(t)})-\boldsymbol{A}^{(t)*}\boldsymbol{R}\|_2\right\}<1-\gamma.\end{aligned}
$$

All the remaining arguments still hold.

D.7 Proof of Theorem [8](#page-19-0)

First, when tasks in S^c come from linear models, the term $\sqrt{p/n}$ in the lower bound of $\max_{t\in[T]} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2$ is the standard result in linear regression. The other part $\sqrt{\log T/n}$ can be shown using the same arguments as in [Duan and Wang](#page-36-1) [\(2022\)](#page-36-1) and [Tian et al.](#page-39-4) [\(2022\)](#page-39-4). See the proof of Theorem 4.3 in [Duan and Wang](#page-36-1) [\(2022\)](#page-36-1) and the proof of Theorem 2 in [Tian](#page-39-4) [et al.](#page-39-4) [\(2022\)](#page-39-4). We omit the proof of lower bound for $\max_{t\in[T]} \|\hat{\beta}^{(t)} - \beta^{(t)*}\|_2$ when tasks in S^c come from linear models, and only show the lower bound of $\max_{t \in S} \|\hat{\beta}^{(t)} - \beta^{(t)*}\|_2$ when tasks in S^c can be arbitrarily distributed.

Throughout this subsection, we assume the following generative model for tasks in S:

$$
y|\boldsymbol{x} \sim \mathbb{P}_{y|\boldsymbol{x}, \boldsymbol{\beta}} = N(\boldsymbol{x}^\top\boldsymbol{\beta}, 1), \quad \boldsymbol{x} \sim \mathbb{P}_{\boldsymbol{x}},
$$

where \mathbb{P}_x is sub-Gaussian with $\Sigma = \mathbb{E}(xx^\top)$. Suppose there exist constants c, C such that $0 < c \leq \lambda_{\min}(\Sigma) \leq \lambda_{\max}(\Sigma) \leq C < \infty$. Then any joint distribution $\mathbb P$ of (x, y) can be written as $\mathbb{P}_{x,y} = \mathbb{P}_{y|x;\beta} \cdot \mathbb{P}_x$.

Recall the parameter space for the coefficient vectors $\{\boldsymbol{\beta}^{(t)}\}_{t\in S}$ as

$$
\mathscr{B}(S,h) = \left\{ \{ \boldsymbol{\beta}^{(t)} \}_{t \in S} : \boldsymbol{\beta}^{(t)} = \boldsymbol{A}^{(t)} \boldsymbol{\theta}^{(t)} \text{ for all } t \in S, \{ \boldsymbol{A}^{(t)} \}_{t \in S} \subseteq \mathcal{O}^{p \times r}, \|\boldsymbol{\theta}^{(t)}\|_2 \le \zeta^{(t)},
$$

$$
\min_{\overline{\boldsymbol{A}} \in \mathcal{O}^{p \times r}} \max_{t \in S} \|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top - \overline{\boldsymbol{A}} \overline{\boldsymbol{A}}^\top \|_2 \le h, \sigma_r \left(|S|^{-1/2} \boldsymbol{B}_S \right) \ge \frac{c}{\sqrt{r}} \sqrt{\frac{1}{|S|} \sum_{t \in S} \|\boldsymbol{\theta}^{(t)}\|_2^2} \right\}
$$

where c can be any fixed positive constants such that $\mathscr{B}(S, h) \neq \emptyset$.

Given $S \subseteq [T]$, define

$$
\mathcal{A}(S,h) = \left\{ \{\mathbf{A}^{(t)}\}_{t \in S} \subseteq \mathcal{O}^{p \times r} : \min_{\overline{\mathbf{A}} \in \mathcal{O}^{p \times r}} \max_{t \in S} \|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \overline{\mathbf{A}} \overline{\mathbf{A}}^\top \|_2 \le h \right\},\
$$

$$
\Theta(S) = \left\{ \{\boldsymbol{\theta}^{(t)}\}_{t \in S} \subseteq \mathbb{R}^r : \|\boldsymbol{\theta}^{(t)}\|_2 \le \zeta^{(t)}, \frac{1}{|S|} \sum_{t \in S} \boldsymbol{\theta}^{(t)}(\boldsymbol{\theta}^{(t)})^\top \succeq \frac{c}{r} \frac{1}{|S|} \sum_{t \in S} \|\boldsymbol{\theta}^{(t)}\|_2^2 \cdot \mathbf{I}_r \right\},\
$$

$$
\bar{\zeta}_S = \sqrt{\frac{1}{|S|} \sum_{t \in S} (\zeta^{(t)})^2}.
$$

The proof proceeds as follows. We will first show

$$
\inf_{\{\widehat{\beta}^{(t)}\}_{t=1}^T} \sup_{S \subseteq \mathcal{S}} \sup_{\{\beta^{(t)}\}_{t \in S} \in \mathcal{B}(S,h)} \mathbb{P} \Bigg(\bigcup_{t \in S} \Bigg\{ \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \gtrsim \left[\frac{\zeta^{(t)}}{\bar{\zeta}_S} \sqrt{\frac{pr}{n}} + \left(\frac{\zeta^{(t)}}{\bar{\zeta}_S} \vee 1\right) \sqrt{\frac{r}{n}} + \zeta^{(t)}h \Bigg\}
$$

$$
+\frac{\zeta^{(t)}}{\bar{\zeta}_S} \frac{\epsilon r}{\sqrt{n}} \bigg] \wedge \sqrt{\frac{p}{n}} \bigg\} \bigg) \ge \frac{1}{10},\tag{20}
$$

$$
\inf_{\{\widehat{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T} \sup_{S \subseteq \mathcal{S}} \sup_{\{\boldsymbol{\beta}^{(t)}\}_{t \in S} \in \mathcal{B}([T],0)} \mathbb{P}\left(\max_{t \in S} \|\widehat{\boldsymbol{\beta}}^{(t)} - \boldsymbol{\beta}^{(t)*}\|_2 \gtrsim \sqrt{\frac{\log T}{n}}\right) \ge \frac{1}{10}.\tag{21}
$$
Then we will discuss how to get the final lower bound by the same arguments. **Part 1:** First, let us prove (20) first.

• If $h \geq \frac{1}{\bar{\zeta}_{|T|}}$ $\sqrt{\frac{pr}{nT}} + \frac{1}{\zeta_{|T|}} \sqrt{\frac{r}{n}} + \frac{1}{\zeta_{|T|}} \frac{\epsilon r}{\sqrt{n}}$: then $\zeta^{(t)} h \geq \frac{\zeta^{(t)}}{\zeta_{|T|}}$ $\bar{\zeta}_{[T]}$ $\sqrt{\frac{pr}{nT}} + \frac{\zeta^{(t)}}{\zeta_{[T]}}$ $\frac{\zeta^{(t)}}{\bar{\zeta}_{|T|}} \sqrt{\frac{r}{n}} + \frac{\zeta^{(t)}}{\bar{\zeta}_{|T|}}$ $\frac{\zeta^{(t)}}{\bar{\zeta}_{|T|}} \frac{\epsilon r}{\sqrt{n}}$ for all $t \in [T]$.

o If $\sqrt{\frac{r}{n}}$ ≤ ($\zeta^{(t_0)}$ h) ∧ $\sqrt{\frac{p}{n}}$ for some $t_0 \in [T]$: then we take $S = [T]$ and prove the lower bound

$$
\inf_{\{\widehat{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T} \sup_{\{\boldsymbol{\beta}^{(t)}\}_{t\in[T]} \in \mathscr{B}([T],h)} \mathbb{P}\bigg(\|\widehat{\boldsymbol{\beta}}^{(t_0)} - \boldsymbol{\beta}^{(t_0)}\|_2 \gtrsim (\zeta^{(t_0)}h) \wedge \sqrt{\frac{p}{n}} \bigg) \ge \frac{1}{10}.
$$

WLOG, assume $t_0 = 1$. For all $t \in [T]$, fix $\mathbf{A}^{(t)} =$ an arbitrary $\overline{\mathbf{A}} \in \mathcal{O}^{p \times r}$, and fix $\{\theta^{(t)}\}_{t=2}^T$ s.t. $\|\theta^{(t)}\|_2 \leq \zeta^{(t)}$. Let $\theta^{(1)} = \frac{1}{\sqrt{2\pi}}$ $\frac{1}{T}\zeta^{(1)}\mathbf{1}_r$. Set $\delta = \frac{1}{12} \big[\big(\frac{1}{\zeta^{(1)}} \sqrt{\frac{p}{n}} \big) \wedge h \big].$ Consider a $\delta/2$ -packing of the ball $\mathcal{B}_{\delta}(0,\mathbb{R}^p, \|\cdot\|_2)$ (denoted as M). By Example 5.8 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0), $|\mathcal{M}| \geq 5^p$. Consider

$$
\bm{A} = \overline{\bm{A}} + \frac{c'}{\sqrt{r}} \cdot \bm{u} \bm{1}_r^\top, \quad \widetilde{\bm{A}} = \overline{\bm{A}} + \frac{c'}{\sqrt{r}} \cdot \widetilde{\bm{u}} \bm{1}_r^\top,
$$

where $u \neq \tilde{u} \in \mathcal{M}$ and c' is a small constant, then

$$
\| (A - \widetilde{A}) \theta^{(1)} \|_2 = \frac{1}{r} \zeta^{(t)} \| (u - \widetilde{u}) \mathbf{1}_r^\top \mathbf{1}_r \|_2 = \| u - \widetilde{u} \|_2 \in \left[\frac{c'}{2} \zeta^{(1)} \delta, 2c' \zeta^{(1)} \delta \right] (22)
$$

$$
\| (A - \widetilde{A}) \theta^{(t)} \|_2 = \| (A - \widetilde{A}) e_{j_k} \|_2 = \frac{1}{\sqrt{r}} \zeta^{(1)} \| u - \widetilde{u} \|_2 \le \frac{\delta}{\sqrt{r}} \zeta^{(1)},
$$

where $\boldsymbol{\theta}^{(t)} = \boldsymbol{e}_{j_k}$. Hence $\mathscr{B} = \{ \boldsymbol{\beta} \in \mathbb{R}^p : \boldsymbol{\beta} = \boldsymbol{A} \boldsymbol{\theta}^{(1)}, \boldsymbol{A} = \overline{\boldsymbol{A}} + \frac{c'}{\sqrt{r}} \cdot \boldsymbol{u} \boldsymbol{1}_r^\top, \boldsymbol{u} \in \mathcal{M} \}$ is a $c'\delta/2$ -packing in \mathbb{R}^p with $|\mathscr{B}| = |\mathcal{M}| \geq 5^p$. On the other hand, for any $A \in \mathbb{R}^{p \times r}$ with $1 + 2c'\delta \geq \sigma_1(A) \geq \sigma_2(A) \geq \cdots \geq \sigma_r(A) \geq 1 - 2c'\delta > 0$ and $||A-\overline{A}||_2 \leq \delta$, consider its SVD where $A = Q\Lambda V$ where $Q \in \mathcal{O}^{p \times r}$ and $V \in \mathcal{O}^{r \times r}$. Denote $\mathbf{R} = \Lambda \mathbf{V}$. Then $0 \leq 1 - 2c' \delta \leq \sigma_{\min}(\mathbf{R}) \leq \sigma_{\max}(\mathbf{R}) \leq 1 + 2c' \delta < \infty$, and $\mathbf{A}\boldsymbol{\theta}^{(1)} = \mathbf{Q}\mathbf{R}\boldsymbol{\theta}^{(1)} = \mathbf{Q}\widetilde{\boldsymbol{\theta}}^{(1)}$. Note that $\mathbf{Q} \in \mathcal{O}^{p \times r}$ and $\widetilde{\boldsymbol{\theta}}^{(1)} = \mathbf{R}\boldsymbol{\theta}^{(1)}$ with $\|\widetilde{\theta}^{(1)}\|_2 \leq \|R\|_2 \|\theta^{(1)}\|_2 \leq C[1+2c']$. And $\|QQ^\top - \overline{A}(\overline{A})^\top\|_2 \leq \| (QR)(QR)^\top \overline{\bm{A}}(\overline{\bm{A}})^\top \|_2 \!+\! \| \bm{Q} (\bm{I}_r\!-\!\bm{R}\bm{R}^\top)\bm{Q}^\top \|_2 \leq 4 \| \bm{A} \!-\! \overline{\bm{A}}\|_2 \!+\! 4 \| \bm{I}_r \!-\! \bm{R}\|_2 \leq 4 \delta \!+\! 8 c' \delta \leq 12 \delta \leq h.$ Therefore,

$$
\inf_{\hat{\beta}^{(1)} \{\beta^{(t)}\}_{t \in S} \in \mathcal{B}([T],h)} \mathbb{P}\left(\|\hat{\beta}^{(1)} - \beta^{(1)}\|_{2} \gtrsim (\zeta^{(1)}h) \wedge \sqrt{\frac{p}{n}}\right)
$$
\n
$$
\geq \inf_{\hat{\beta}^{(1)} \text{ } 0 < c \leq \sigma_{\max}(A) \leq \sigma_{\max}(A) \leq C} \mathbb{P}\left(\|\hat{\beta}^{(1)} - A\theta^{(1)}\|_{2} \gtrsim (\zeta^{(1)}h) \wedge \sqrt{\frac{p}{n}}\right)
$$
\n
$$
:= (*)
$$

where all $\mathbf{A}^{(t)} =$ an arbitrary $\overline{\mathbf{A}} \in \mathcal{O}^{p \times r}$, and $\{\theta^{(t)}\}_{t=1}^T \subseteq \{e_j\}_{j=1}^r$ are fixed.

For any $\beta = A\theta^{(1)}$ and $\tilde{\beta} = \tilde{A}\theta^{(1)} \in \mathscr{B}$ with $A \neq \tilde{A}$, by Lemma [25](#page-48-0) and equation $(22),$ $(22),$

$$
\begin{split}\n\text{KL}\left(\prod_{t=2}^{T} \mathbb{P}_{y|\mathbf{x};\overline{A}\boldsymbol{\theta}^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(t)\otimes n} \cdot \mathbb{P}_{y|\mathbf{x};A\boldsymbol{\theta}^{(1)}}^{(1)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(1)\otimes n} \right) & \prod_{t=2}^{T} \mathbb{P}_{y|\mathbf{x};\overline{A}\boldsymbol{\theta}^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(t)\otimes n} \cdot \mathbb{P}_{y|\mathbf{x};\widetilde{A}\boldsymbol{\theta}^{(1)}}^{(1)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(1)\otimes n} \\
&= \text{KL}\left(\mathbb{P}_{y|\mathbf{x};A\boldsymbol{\theta}^{(1)}}^{(1)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(1)\otimes n} \right) & \prod_{y|\mathbf{x};\widetilde{A}\boldsymbol{\theta}^{(1)}}^{(1)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(1)\otimes n}\n\end{split}
$$
\n
$$
\leq n \|(A - \widetilde{A})\boldsymbol{\theta}^{(1)}\|_{2}^{2} \leq n(\zeta^{(1)})^{2}\delta^{2}
$$
\n
$$
\leq c \log |\mathscr{B}|.
$$

Then by Fano's lemma (Lemma [24\)](#page-48-1), $(*) \geq 1 - \frac{\log 2}{\log |\mathcal{B}|} - c \geq 1/10$.

o If $\sqrt{\frac{r}{n}}$ > ($\zeta^{(t)}h$) ∧ $\sqrt{\frac{p}{n}}$ for all $t \in [T]$: then we take $S = [T]$ and prove the lower bound

$$
\inf_{\{\widehat{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T} \sup_{\{\boldsymbol{\beta}^{(t)}\}_{t\in[T]} \in \mathcal{B}([T],0)} \mathbb{P}\bigg(\|\widehat{\boldsymbol{\beta}}^{(t_0)} - \boldsymbol{\beta}^{(t_0)}\|_2 \gtrsim \sqrt{\frac{r}{n}}\bigg) \ge \frac{1}{10},\tag{23}
$$

where $t_0 \in \arg \min_{t \in [T]} \zeta^{(t)}$. WLOG, assume $t_0 = 1$. Fix all $A^{(t)} =$ some $\mathbf{A} \in \mathcal{O}^{p \times r}$. Fixing $\mathbf{\Theta} = \{\mathbf{\theta}^{(t)}\}_{t=2}^T$ such that $\frac{1}{T} \sum_{t=2}^T \mathbf{\theta}^{(t)}(\mathbf{\theta}^{(t)})^\top \succeq \frac{c}{r}$ $\frac{c}{r}I_r$ (Hence for any $\theta^{(1)}$ with $\|\theta^{(1)}\|_2 \leq C \leq \zeta^{(1)}$, we must have $\frac{1}{T} \sum_{t=1}^T \theta^{(t)} (\theta^{(t)})^\top \succeq \frac{c}{r}$ $\frac{c}{r}I_r$). We want to show

$$
\inf_{\{\widehat{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T \|\boldsymbol{\theta}^{(1)}\|_2 \leq C} \mathbb{P}\left(\|\widehat{\boldsymbol{\beta}}^{(1)} - \boldsymbol{A}\boldsymbol{\theta}^{(1)}\|_2 \geq c\sqrt{\frac{r}{n}}\right) \geq \frac{1}{10}.
$$

Denote $\delta = \sqrt{\frac{r}{n}}$. Consider a cô-packing of $\mathcal{B}_{\delta}(\mathbf{0}, \mathbb{R}^r, \|\cdot\|_2)$ (denoted as \mathscr{T}). By Example 5.8 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0), we know that $\log |\mathcal{T}| \gtrsim r$. Note that this also defines a cô-packing of the space of $\boldsymbol{\beta}^{(1)}$ as $\mathscr{M} = \{ \boldsymbol{\beta}^{(1)} \in \mathbb{R}^p : \boldsymbol{\beta}^{(1)} = A \boldsymbol{\theta}, \boldsymbol{\theta} \in \mathscr{T} \},$ because for any $\theta^{(1)} \neq \widetilde{\theta}^{(1)} \in \mathcal{T}$ we have $||A(\theta^{(1)} - \widetilde{\theta}^{(1)})||_2 = ||\theta^{(1)} - \widetilde{\theta}^{(1)}||_2 \geq c\delta$. Denote the distribution of $\{y_i^{(t)}\}$ $\{x_i^{(t)}\}_{i=1}^n$ given $\{\boldsymbol{x}_i^{(t)}\}$ $\{(\theta_i^{(t)})\}_{i=1}^n$ and $\{\boldsymbol{\beta}^{(t)}\}_{t=1}^T$ as $\mathbb{P}_{y|\boldsymbol{x};\boldsymbol{\beta}^{(t)}}^{(t)\otimes n}$ and the distribution of $\{\boldsymbol{x}_i^{(t)}\}$ $\{\phi_i^{(t)}\}_{i=1}^n$ as $\mathbb{P}_{\bm{x}}^{(t)\otimes n}$. For any $\boldsymbol{\beta}^{(1)} \neq \widetilde{\boldsymbol{\beta}}^{(1)} \in \mathscr{M}$:

$$
\begin{aligned} \mathrm{KL}\left(\mathbb{P}_{y|\boldsymbol{x};\boldsymbol{\beta}^{(1)}}^{(1)\otimes n}\cdot \mathbb{P}_{\boldsymbol{x}}^{(1)\otimes n}\middle\Vert \mathbb{P}_{y|\boldsymbol{x};\boldsymbol{A}\widetilde{\boldsymbol{\beta}}^{(1)}}^{(1)\otimes n}\cdot \mathbb{P}_{\boldsymbol{x}}^{(1)\otimes n}\right) &\lesssim n \|\boldsymbol{\beta}^{(1)}-\widetilde{\boldsymbol{\beta}}^{(1)}\|_2^2\\ &\lesssim n\delta^2\\ &\lesssim r\\ &\leq c\log|\mathscr{M}|,\end{aligned}
$$

where c is a small constant. Finally, applying Fano's Lemma (Lemma 24), we have

$$
\inf_{\{\widehat{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T \; \|\boldsymbol{\theta}^{(1)}\|_2 \leq \delta} \mathbb{P}\left(\|\widehat{\boldsymbol{\beta}}^{(1)} - \boldsymbol{A}\boldsymbol{\theta}^{(1)}\|_2 \geq c\delta\right) \geq 1 - \frac{\log 2}{\log |\mathscr{M}|} - c \geq \frac{1}{10}.
$$

• If $h < \frac{1}{\bar{\zeta}_{[T]}}$ $\sqrt{\frac{pr}{nT}} + \frac{1}{\zeta_{|T|}} \sqrt{\frac{r}{n}} + \frac{1}{\zeta_{|T|}} \frac{\epsilon r}{\sqrt{n}}$: then $\zeta^{(t)} h < \frac{\zeta^{(t)}}{\zeta_{|T|}}$ $\bar{\zeta}_{[T]}$ $\sqrt{\frac{pr}{nT}} + \frac{\zeta^{(t)}}{\zeta_{[T]}}$ $\frac{\zeta^{(t)}}{\bar{\zeta}_{|T|}} \sqrt{\frac{r}{n}} + \frac{\zeta^{(t)}}{\bar{\zeta}_{|T|}}$ $\frac{\zeta^{(t)}}{\zeta_{|T|}} \frac{\epsilon r}{\sqrt{n}}$ for all $t \in [T]$.

 \circ If $\sqrt{\frac{r}{n}} \geq \frac{\zeta^{(t_0)}}{\overline{\zeta}_{[T]}}$ $\overline{\zeta_{|T|}}$ $\sqrt{\frac{pr}{nT}} + \frac{\zeta^{(t_0)}}{\overline{\zeta}_{[T]}}$ $\frac{\zeta^{(t_0)}}{\bar{\zeta}_{|T|}}\sqrt{\frac{r}{n}}+\frac{\zeta^{(t_0)}}{\bar{\zeta}_{|T|}}$ $\frac{\zeta(t_0)}{\zeta_{|T|}}\frac{\epsilon r}{\sqrt{n}}$ for $t_0 \in \argmin_{t \in [T]} \zeta^{(t)}$: then we take $S = |T|$ and it suffiecs to prove the lower bound

$$
\inf_{\{\widehat{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T} \sup_{\{\boldsymbol{\beta}^{(t)}\}_{t\in[T]} \in \mathcal{B}([T],0)} \mathbb{P}\bigg(\|\widehat{\boldsymbol{\beta}}^{(t_0)} - \boldsymbol{\beta}^{(t_0)}\|_2 \gtrsim \sqrt{\frac{r}{n}} \bigg) \ge \frac{1}{10}
$$

.

This has already been proved in the previous analysis.

 \circ If $\sqrt{\frac{r}{n}} < \frac{\zeta^{(t)}}{\overline{\zeta}_{[T]}}$ $\overline{\bar{\zeta}}_{[T]}$ $\sqrt{\frac{pr}{nT}} + \frac{\zeta^{(t)}}{\zeta_{[T]}}$ $\frac{\zeta^{(t)}}{\bar{\zeta}_{|T|}} \sqrt{\frac{r}{n}} + \frac{\zeta^{(t)}}{\bar{\zeta}_{|T|}}$ $\frac{\zeta^{(t)}}{\zeta_{[T]}} \frac{\epsilon r}{\sqrt{n}}$ for all $t \in [T]$: \star If $\sqrt{\frac{r}{n}} \geq \sqrt{\frac{pr}{nT}} + \epsilon \frac{r}{\sqrt{n}}$: then we take $S = [T]$ and it suffices to prove

$$
\inf_{\{\widehat{\beta}^{(t)}\}_{t=1}^T} \sup_{\{\beta^{(t)}\}_{t\in[T]} \in \mathscr{B}([T],0)} \mathbb{P}\bigg(\|\widehat{\beta}^{(t_0)} - \beta^{(t_0)}\|_2 \gtrsim \frac{\zeta^{(t_0)}}{\bar{\zeta}} \sqrt{\frac{r}{n}} \bigg) \ge \frac{1}{10},
$$

with $t_0 \in \arg \min_{t \in [T]} \zeta^{(t)}$, which is automatically true because $\zeta^{(t_0)}/\bar{\zeta} \leq 1$ and [\(23\)](#page-73-0).

★ If
$$
\sqrt{\frac{r}{n}} < \sqrt{\frac{pr}{nT}} + \epsilon \frac{r}{\sqrt{n}}
$$
:
\n>▶ If $\sqrt{\frac{pr}{nT}} \ge \epsilon \frac{r}{\sqrt{n}}$: then we take $S = [T]$ and prove

$$
\inf_{\{\widehat{\beta}^{(t)}\}_{t=1}^T} \sup_{\{\beta^{(t)}\}_{t\in[T]}\in\mathscr{B}([T],0)} \mathbb{P}\Bigg(\bigcup_{t\in[T]} \Bigg\{\|\widehat{\beta}^{(t)}-\beta^{(t)*}\|_2 \gtrsim \frac{\zeta^{(t)}}{\bar{\zeta}_{[T]}} \sqrt{\frac{pr}{nT}}\Bigg) \geq \frac{1}{10}.
$$

Note that it suffices to prove

$$
\inf_{\{\widehat{\beta}^{(t)}\}_{t=1}^T} \sup_{\{\beta^{(t)}\}_{t\in[T]}\in\mathscr{B}([T],0)} \mathbb{P}\Bigg(\frac{1}{T} \sum_{t\in[T]} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2^2 \gtrsim \frac{pr}{nT}\Bigg) \ge \frac{1}{10},
$$

because if $\frac{1}{T} \sum_{t \in [T]} \| \widehat{\beta}^{(t)} - \beta^{(t)*} \|_2^2 \gtrsim \frac{pr}{nT}$, then there must exist $t \in [T]$ such that $\|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2^2 \gtrsim (\frac{\zeta^{(t)}}{\bar{\zeta}_{[T]}})$ $\frac{\zeta^{(t)}}{\bar{\zeta}_{|T|}}$)² $\frac{pr}{nT}$. Now let us prove it. Consider an $\overline{A} \in \mathcal{O}^{p \times r}$ such that the packing number $M(\mathcal{B}_{\delta}(\overline{\mathbf{A}}, \mathcal{O}^{p\times r}, \text{dist}_2), \text{dist}_{\text{F}}, \alpha\delta) \geq (\frac{C\sqrt{r}}{\alpha})$ $\frac{\sqrt{r}}{\alpha}$)^{r(p-r)} in Lemma [23](#page-48-2) with $\delta = \frac{1}{\bar{\zeta}_{[T]}}$ $\sqrt{\frac{pr}{nT}}$. Let $\mathcal{M} = {\mathbf{A}_j}_{j=1}^{|\mathcal{M}|}$ be the maximum packing of $\mathcal{B}_\delta(\overline{\bm{A}},\mathcal{O}^{p\times r},\text{dist}_2)$ corresponding to the packing number $M(\mathcal{B}_\delta(\overline{\bm{A}},\mathcal{O}^{p\times r},\text{dist}_2),$ $\mathcal{O}_{\delta}(\mathbf{A}, \mathbf{C}^{\prime})$, disty corresponding to the packing number $M(\mathcal{O}_{\delta}(\mathbf{A}, \mathbf{C}^{\prime})$, distribution of $\epsilon > 0$. This leads to a $c' \bar{\zeta}_{[T]} \delta$ - $\text{packing } \mathscr{M} = \{ \{\bm{\beta}^{(t)}\}_{t=1}^T: \bm{\beta}^{(t)} \coloneqq \bm{A}_j \bm{R}_j \bm{\theta}^{(t)}, \bm{R}_j = \argmin_{\bm{R} \in \mathcal{O}^{r \times r}} \| \overline{\bm{A}} - \bm{\beta} \|$ $A_jR\|_2$, for all $t \in [T]$ and the same $j \in [M]$ of the space $\mathscr{B}([T],0)$ w.r.t. distance $\sqrt{\frac{1}{7}}$ $\frac{1}{T} \sum_{t \in [T]} ||\boldsymbol{\beta}^{(t)} - \tilde{\boldsymbol{\beta}}^{(t)}||_2^2$. To verify this, notice that

for any $j_1 \neq j_2 \in [\vert \mathcal{M} \vert], \ \beta^{(t)} = A_{j_1}R_{j_1}\theta^{(t)}, \ \widetilde{\beta}^{(t)} = A_{j_2}R_{j_2}\theta^{(t)}, \ \text{with}$ $\bm{R}_{j1} = \arg \min_{\bm{R} \in \mathcal{O}^{r \times r}} \| \overline{\bm{A}} - \bm{A}_{j1} \bm{R} \|_2, \, \bm{R}_{j2} = \arg \min_{\bm{R} \in \mathcal{O}^{r \times r}} \| \overline{\bm{A}} - \bm{A}_{j2} \bm{R} \|_2,$ by Wedin's sin Θ-Theorem and Assumption [2,](#page-10-0)

$$
\frac{1}{T}\sum_{t=1}^T \|\boldsymbol{\beta}^{(t)} - \widetilde{\boldsymbol{\beta}}^{(t)}\|_2^2 \geq \frac{1}{T} \|\boldsymbol{A}_{j_1}\boldsymbol{\Theta}_{j_1} - \boldsymbol{A}_{j_2}\boldsymbol{\Theta}_{j_2}\|_F^2 \geq \frac{c}{r} \|\boldsymbol{A}_{j_1}\boldsymbol{A}_{j_1}^\top - \boldsymbol{A}_{j_2}\boldsymbol{A}_{j_2}^\top\|_F^2 \geq (c')^2 \bar{\zeta}^2 \delta^2,
$$

where c' is a small constant, $\mathbf{\Theta}_{j_1} = \{\boldsymbol{R}_{j_1} \boldsymbol{\theta}^{(t)}\}_{t \in [T]},$ and $\mathbf{\Theta}_{j_2} = \{\boldsymbol{R}_{j_2} \boldsymbol{\theta}^{(t)}\}_{t \in [T]}.$ On the other hand, notice that for any $j \in [\mathcal{M}]]$, $\boldsymbol{\beta}^{(t)} = A_j R_j \boldsymbol{\theta}^{(t)}$ with $\mathbf{R}_i = \arg \min_{\mathbf{R} \in \mathcal{O}^{r \times r}} \|\overline{\mathbf{A}} - \mathbf{A}_i \mathbf{R}\|_2$, by Lemma [21,](#page-47-0) for all $t \in [T]$, we have

$$
\|\boldsymbol{\beta}^{(t)} - \overline{\boldsymbol{A}}\boldsymbol{\theta}^{(t)}\|_2 \leq \|\boldsymbol{A}_j \boldsymbol{R}_j - \overline{\boldsymbol{A}}\|_2 \|\boldsymbol{\theta}^{(t)}\|_2
$$

\n
$$
= \zeta^{(t)} \min_{\boldsymbol{R} \in \mathcal{O}^{r \times r}} \|\boldsymbol{A}_j \boldsymbol{R} - \overline{\boldsymbol{A}}\|_2
$$

\n
$$
\leq \sqrt{2} \zeta^{(t)} \|\boldsymbol{A}_j \boldsymbol{A}_j^{\top} - \overline{\boldsymbol{A}} (\overline{\boldsymbol{A}})^{\top}\|_2
$$

\n
$$
\leq \sqrt{2} \zeta^{(t)} \delta,
$$

which by triangle inequality leads to

$$
\frac{1}{T} \sum_{t \in [T]} \|\beta^{(t)} - \tilde{\beta}^{(t)}\|_2^2 \le 4C^2 \cdot \frac{1}{T} \sum_{t \in [T]} (\zeta^{(t)})^2 \cdot \delta^2 \le 4C^2 \bar{\zeta}_{[T]}^2 \delta^2, \qquad (24)
$$

for any $\{\boldsymbol{\beta}^{(t)}\}_{t=1}^T$ and $\{\widetilde{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T \in \mathcal{M}$.

By Lemma [22,](#page-47-1) $|\mathcal{M}| = |\mathcal{M}| \gtrsim C^{r(p-r)} \geq C^{rp/2}$. Denote the distribution of $\{y_i^{(t)}\}$ $\{x_i^{(t)}\}_{i=1}^n$ given $\{\boldsymbol{x}_i^{(t)}\}$ $\{e^{(t)}\}_{i=1}^n$ as $\mathbb{P}^{(t)\otimes n}_{y|\boldsymbol{x};\boldsymbol{\beta}}$ $y|x;\beta}^{(t) \otimes n}$ and the distribution of $\{x_i^{(t)}\}$ $i^{(t)}\}_{i=1}^{n}$ as $\mathbb{P}^{(t)\otimes n}_{\bm{x}}$. Then for any $\{\boldsymbol{\beta}^{(t)}\}_{t=1}^T$, $\{\widetilde{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T \in \mathcal{M}$, by Lemma [25](#page-48-0) and equation [\(24\)](#page-75-0),

$$
\text{KL}\left(\prod_{t=1}^T \mathbb{P}_{y|\boldsymbol{x};\boldsymbol{\beta}^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\boldsymbol{x}}^{(t)\otimes n} \middle\| \prod_{t=1}^T \mathbb{P}_{y|\boldsymbol{x};\widetilde{\boldsymbol{\beta}}^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\boldsymbol{x}}^{(t)\otimes n} \right) \lesssim n \sum_{t=1}^T \|\boldsymbol{\beta}^{(t)} - \widetilde{\boldsymbol{\beta}}^{(t)}\|_2^2
$$

\$\lesssim nT\zeta_{[T]}^2 \delta^2\$
\$\leq c \log |\mathcal{M}|\$,

where c is a small constant. Finally, applying Fano's Lemma (Lemma) [24\)](#page-48-1), we have

$$
\inf_{\{\hat{\beta}^{(t)}\}_{t=1}^T} \sup_{\{\beta^{(t)}\}_{t=1}^T \in \mathcal{B}([T],0)} \mathbb{P}\left(\frac{1}{T} \sum_{t \in [T]} \|\hat{\beta}^{(t)} - \beta^{(t)}\|_2^2 \ge c^2 \delta^2\right) \ge 1 - \frac{\log 2}{\log |\mathcal{B}|} - c
$$

$$
\ge \frac{1}{10}.
$$

 \triangleright If $\sqrt{\frac{pr}{nT}} < \epsilon \frac{r}{\sqrt{n}}$: then we consider $t_0 \in \arg \min_{t \in [T]} \zeta^{(t)}, S \ni t_0$, and prove

$$
\inf_{\{\widehat{\beta}^{(t)}\}_{t=1}^T} \sup_{S \subseteq \mathcal{S}} \sup_{\{\beta^{(t)}\}_{t \in [T]} \in \mathcal{B}(S,0)} \mathbb{P}\left(\|\widehat{\beta}^{(t_0)} - \beta^{(t_0)}\|_2 \gtrsim \frac{\zeta^{(t_0)}}{\bar{\zeta}_S} \epsilon \frac{r}{\sqrt{n}}\right) \ge \frac{1}{10}.
$$

Note that $\frac{\zeta^{(t_0)}}{\bar{\zeta}_s}$ $\frac{\overline{\zeta_{\mathcal{S}}}}{\overline{\zeta_{\mathcal{S}}}} \leq 1$ for any $S \subseteq [T]$. Therefore it suffices to prove

$$
\inf_{\{\widehat{\beta}^{(t)}\}_{t=1}^T} \sup_{S \subseteq \mathcal{S}} \sup_{\{\beta^{(t)}\}_{t \in [T]} \in \mathcal{B}(S,0)} \mathbb{P}\left(\|\widehat{\beta}^{(t_0)} - \beta^{(t_0)}\|_2 \gtrsim \epsilon \frac{r}{\sqrt{n}}\right) \ge \frac{1}{10}.
$$

WLOG, assume $t_0 = 1$. Fix all $\mathbf{A}^{(t)} = \text{some } \mathbf{A} \in \mathcal{O}^{p \times r}$ and $\{\mathbf{\theta}^{(t)}\}_{t=1}^T \subseteq$ $\{\boldsymbol{e}_j\}_{j=1}^r \subseteq \mathbb{R}^r \text{ s.t. } \sigma_r(T^{-1/2}\{\theta^{(t)}\}_{t=1}^T) \geq \frac{c}{\sqrt{r}} \text{ with } \#\{t : \theta^{(t)} = e_1\} =$ $[T/r]$. Without loss of generality, suppose $\boldsymbol{\theta}^{(t)} = \boldsymbol{e}_1$ when $t = [[T/r]]$. Denote $\tilde{\boldsymbol{\beta}} = \boldsymbol{A}\boldsymbol{e}_1$, and $\boldsymbol{\beta}^{(t)} = \boldsymbol{A}\boldsymbol{\theta}^{(t)}$ for $t \geq \lfloor T/r \rfloor + 1$. Consider two data generating mechanisms in Lemma [27:](#page-48-3)

- $(I) \{(x_i^{(t)}\)$ $\binom{t}{i}, y_{i}^{(t)}$ $\binom{t}{i}\}_{i=1}^n \sim (1-\epsilon')(\mathbb{P}_{\text{min}}^{\otimes n})$ $y|\bm{x};\bm{\beta}$ $\cdot \mathbb{P}_{x}^{\otimes n}$ + $\epsilon' \mathbb{Q}$ independently for $t \in$ $[|T/r|]$, where $\epsilon' = \frac{T\epsilon}{50|T|}$ $\frac{T\epsilon}{50|T/r|}, \text{and } \{\{\boldsymbol{x}_i^{(t)}% ,...,\boldsymbol{x}_i^{(t)}\}_{i=1}^{T}\}$ $\{ \widehat{u}^{(t)}_i \}_{i=1}^n \}_{t=|T/r|+1}^T \sim \mathbb{D} = \prod_{t= \lfloor T/r \rfloor +1}^T (\mathbb{P}_{\widehat{u}|\bm{x}}^{\otimes n})$ ${}^{\otimes n}_{y|\bm{x};\bm{\beta}^{(t)}}\cdot$ $\mathbb{P}_x^{\otimes n}$);
- (II) With a preserved set $S^c \subseteq [[T/r]],$ generate $\{\boldsymbol{x}^{(t)}\}_{t \in S^c} \sim \mathbb{Q}_{S^c}$ and $\{(\boldsymbol{x}_{i}^{(t)}% ,\boldsymbol{x}_{j}^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}% ^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}% ^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}% ^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}^{(t)}\boldsymbol{\beta}_{i}^{(t)}\bold$ $\binom{t}{i}, y_{i}^{(t)}$ $\binom{t}{i}$ } $\}_{i=1}^n \sim \mathbb{P}_{\mathbf{u}|\mathbf{x}}^{\otimes n}$ $y|\bm{x};\bm{\beta}$ $\cdot \mathbb{P}_{\mathbf{x}}^{\otimes n}$ independently for $t \in S \cap [[T/r]],$ and $\{\{\boldsymbol{x}^{(t)}_i$ $\{ \hat{u}^{(t)}_{i} \}_{i=1}^{n} \}_{t=|T/r|+1}^{T} \sim \mathbb{D} = \prod_{t=|T/r|+1}^{T} (\mathbb{P}_{v|x}^{\otimes n})$ $\frac{\otimes n}{y|\boldsymbol{x};\boldsymbol{\beta}^{(t)}}\cdot \mathbb{P}_{\boldsymbol{x}}^{\otimes n}$).

Denote the joint distributions of $\{\boldsymbol{x}_i^{(t)}\}$ $\binom{t}{i}_{t\in[[T/r]]}$ in (I) and (II) as $\mathbb{P}_{(\epsilon,\theta,\mathbb{Q})}$ and $\mathbb{P}_{(S,\theta,\mathbb{Q}_{S^c})}$, respectively. Note that by Lemma [25,](#page-48-0)

$$
\varpi(\epsilon', \Theta) \coloneqq \sup \{ \|\beta_1 - \beta_2\|_2 : \mathrm{TV}(\mathbb{P}_{y | \mathbf{x}; \beta_1}^{\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{\otimes n}, \mathbb{P}_{y | \mathbf{x}; \beta_2}^{\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{\otimes n}) \le \epsilon' / (1 - \epsilon') \}
$$

$$
\geq \sup\{\|\beta_1 - \beta_2\|_2 : \mathrm{KL}\big(\mathbb{P}_{y|x;\beta_1}^{\otimes n} \cdot \mathbb{P}_{x}^{\otimes n}, \mathbb{P}_{y|x;\beta_2}^{\otimes n} \cdot \mathbb{P}_{x}^{\otimes n}\big) \leq 2[\epsilon'/(1-\epsilon')]^2\}
$$

$$
\geq \sup\{\|\beta_1 - \beta_2\|_2 : n\|\beta_1 - \beta_2\|_2^2 \leq c[\epsilon'/(1-\epsilon')]^2\}
$$

$$
= c'\frac{\epsilon'}{\sqrt{n}}
$$

$$
\asymp \frac{r\epsilon}{\sqrt{n}}.
$$

Then by Lemma [26,](#page-48-4)

$$
\inf_{\widehat{\beta}^{(1)}} \sup_{\widetilde{\beta}, \mathbb{Q}} (\mathbb{P}_{(\epsilon,\theta,\mathbb{Q})} \cdot \mathbb{D}) \bigg(\| \widehat{\beta}^{(1)} - \beta^{(1)*} \|_2 \geq \varpi(\epsilon',\Theta) \bigg) \geq \frac{1}{2}.
$$

Therefore, by Lemma [27,](#page-48-3) it follows that

$$
\inf_{\{\widehat{\beta}^{(t)}\}_{t=1}^T} \sup_{S \subseteq S} \sup_{\{\beta^{(t)}\}_{t \in S} \in \mathcal{B}(S,0)} \mathbb{P}\left(\|\widehat{\beta}^{(1)} - \beta^{(1)*}\|_2 \ge c \frac{\epsilon r}{\sqrt{n}}\right)
$$
\n
$$
\ge \inf_{\widehat{\beta}^{(1)}} \sup_{S : |S| \ge T(1-\epsilon)} \sup_{\{\beta^{(t)}\}_{t \in S} \in \mathcal{B}(S,0)} (\mathbb{P}_{(S,\theta,\mathbb{Q}_{S^c})} \cdot \mathbb{D}) \left(\|\widehat{\beta}^{(1)} - \beta^{(1)*}\|_2 \ge \varpi(\epsilon/50,\Theta)\right)
$$
\n
$$
\ge \frac{1}{10}.
$$

Part 2: Next, we want to show [\(21\)](#page-71-1). Consider the case that $S = [T]$ and $h = 0$. Fix all **A**(t) = some $A \in \mathcal{O}^{p \times r}$. Fix $\{\theta^{(t)}\}_{t=1}^T \subseteq \{\sqrt{c}e_j\}_{j=1}^r$ satisfying $\frac{1}{T} \sum_{t=1}^T \theta^{(t)}(\theta^{(t)})^\top \succeq \frac{c}{r}$ $\frac{c}{r}I_r$. We want to show

$$
\inf_{\{\widehat{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T} \sup_{\{\boldsymbol{\theta}^{(t)}\}_{t=1}^T \in \Theta([T])} \mathbb{P}\left(\max_{t \in [T]} \|\widehat{\boldsymbol{\beta}}^{(t)} - \boldsymbol{A}\boldsymbol{\theta}^{(t)}\|_2 \geq c\sqrt{\frac{\log T}{n}}\right) \geq \frac{1}{10},
$$

where

$$
\Theta([T]) = \left\{ \{\boldsymbol{\theta}^{(t)}\}_{t\in[T]} \subseteq \mathbb{R}^r : \max_{t\in[T]} \|\boldsymbol{\theta}^{(t)}\|_2 \leq C, \frac{1}{T} \sum_{t=1}^T \boldsymbol{\theta}^{(t)}(\boldsymbol{\theta}^{(t)})^\top \succeq \frac{c}{r} \boldsymbol{I}_r \right\}.
$$

Denote $\delta = \sqrt{\frac{\log T}{n}}$ $\overline{\mathbf{g}T}$. Then $\mathscr{T} = {\{\{\widetilde{\boldsymbol{\theta}}^{(t)}\}_{t=1}^{T}} : \exists t_0 \text{ s.t. } \widetilde{\boldsymbol{\theta}}^{(t_0)} = {\boldsymbol{\theta}}^{(t_0)}(1 + \delta/\sqrt{c}), \widetilde{\boldsymbol{\theta}}^{(t)} =$ $θ^(t)$ for $t \neq t_0$ } is a √2δ-packing of Θ(S) w.r.t. distance max $_{t \in [T]} || \bar{θ}^{(t)} - \tilde{θ}^{(t)} ||_2$ with $\{\bar{θ}^{(t)}\}_{t=1}^T$ and $\{\widetilde{\theta}^{(t)}\}_{t=1}^T \in \mathcal{T}$. Hence $\mathscr{B} = \{\{A\widetilde{\theta}^{(t)}\}_{t=1}^T : \exists t_0 \text{ s.t. } \widetilde{\theta}^{(t_0)} = \theta^{(t_0)}(1 + \delta/\sqrt{c}), \widetilde{\theta}^{(t)} = \theta^{(t_0)}(1 + \delta/\sqrt{c})\}$ $\theta^{(t)}$ for $t \neq t_0$ is a $\sqrt{2\delta}$ -packing in \mathbb{R}^p w.r.t. distance $\max_{t \in [T]} ||\beta^{(t)} - \tilde{\beta}^{(t)}||_2$ with $\{\beta^{(t)}\}_{t=1}^T$ and $\{\widetilde{\beta}^{(t)}\}_{t=1}^T \in \mathcal{T}$. Apparently $|\mathcal{T}| = |\mathcal{B}| = T$. And for any $\{\beta^{(t)}\}_{t=1}^T$ and $\{\widetilde{\beta}^{(t)}\}_{t=1}^T \in \mathcal{T}$, they only differ by two components. WLOG, suppose the indices of different components are t_1 and t_2 . Then we have

$$
\begin{split}\n&\text{KL}\left(\prod_{t=1}^{T} \mathbb{P}_{y|\boldsymbol{x};\boldsymbol{\beta}^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\boldsymbol{x}}^{(t)\otimes n} \middle\| \prod_{t=1}^{T} \mathbb{P}_{y|\boldsymbol{x};\widetilde{\boldsymbol{\beta}}^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\boldsymbol{x}}^{(t)\otimes n}\right) \\
&= \text{KL}\left(\mathbb{P}_{y|\boldsymbol{x};\boldsymbol{A}\boldsymbol{\theta}^{(t_1)}(1+\delta/\sqrt{c})}^{(t_1)\otimes n} \cdot \mathbb{P}_{\boldsymbol{x}}^{(t_2)\otimes n} \cdot \mathbb{P}_{y|\boldsymbol{x};\boldsymbol{A}\boldsymbol{\theta}^{(t_2)}}^{(t_2)\otimes n} \cdot \mathbb{P}_{\boldsymbol{x}}^{(t_2)\otimes n} \middle\| \mathbb{P}_{y|\boldsymbol{x};\boldsymbol{A}\widetilde{\boldsymbol{\theta}}^{(t_1)}}^{(t_1)\otimes n} \cdot \mathbb{P}_{y|\boldsymbol{x};\boldsymbol{A}\widetilde{\boldsymbol{\theta}}^{(t_2)}}^{(t_2)\otimes n} \right) \\
&\lesssim n\|A\boldsymbol{\theta}^{(t_1)}(1+\delta/\sqrt{c}) - A\boldsymbol{\theta}^{(t_1)}\|_2^2 + n\|A\boldsymbol{\theta}^{(t_2)} - A\boldsymbol{\theta}^{(t_2)}(1+\delta/\sqrt{c})\|_2^2 \\
&\lesssim n\delta^2 \\
&\leq c'\log|\mathscr{T}|,\n\end{split}
$$

where c is a small constant. By Fano's Lemma (Lemma 24), we have

$$
\inf_{\{\widehat{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T} \sup_{\{\boldsymbol{\theta}^{(t)}\}_{t=1}^T \in \Theta([T])} \mathbb{P}\left(\max_{t \in [T]} \|\widehat{\boldsymbol{\beta}}^{(t)} - \boldsymbol{A}\boldsymbol{\theta}^{(t)}\|_2 \geq c\sqrt{\frac{\log T}{n}}\right) \geq 1 - \frac{\log 2}{\log |\mathcal{T}|} - c' \geq \frac{1}{10}.
$$

Part 3: Finally, let us discuss how to obtain the final desired lower bound by similar arguments to prove [\(20\)](#page-71-0) and [\(21\)](#page-71-1). Denote S' as the index set of $t \in [T]$ where $\zeta^{(t)}$ is among the largest $T(1-\frac{c'}{r})$ $\frac{c'}{r}$) ones of $\{\zeta^{(t)}\}_{t=1}^T$.

• If $\sqrt{\frac{\log T}{n}} \le \min_{t \in S'} \left\{ \left[\frac{\zeta^{(t)}}{\zeta_{S'}} \right] \right\}$ $\overline{\overline{\zeta}_{S'}}$ $\sqrt{\frac{pr}{nT}} + (\frac{\zeta^{(t)}}{\zeta_{S'}}$ $\frac{\zeta^{(t)}}{\zeta_{S'}} \vee 1$) $\sqrt{\frac{r}{n}} + \zeta^{(t)}h + \frac{\zeta^{(t)}}{\zeta_{S'}}$ $\frac{\zeta^{(t)}}{\zeta_{S'}}\epsilon \frac{r}{\sqrt{n}}$ $\left\{\wedge\sqrt{\frac{p}{n}}\right\}$: then by the same arguments we used to prove (20) , we can get a lower bound similar to (20) by replacing [T] and $\bar{\zeta}_{[T]} = \sqrt{\frac{1}{T} \sum_{t \in [T]} (\zeta^{(t)})^2}$ with S' and $\bar{\zeta}_{S'} = \sqrt{\frac{1}{|S'|} \sum_{t \in S'} (\zeta^{(t)})^2}$. Then notice that by definition of S' , $\bar{\zeta}_{S'} \gtrsim \bar{\zeta}_{[T]}$, hence we can get the bound

$$
\inf_{\{\widehat{\beta}^{(t)}\}_{t=1}^T} \sup_{S \subseteq \mathcal{S}} \sup_{\{\beta^{(t)}\}_{t \in S} \in \mathcal{B}(S,h)} \mathbb{P}\Bigg(\bigcup_{t \in S'} \Bigg\{\|\widehat{\beta}^{(t)} - \beta^{(t)}\|_2 \gtrsim \left[\frac{\zeta^{(t)}}{\bar{\zeta}_{S'}}\sqrt{\frac{pr}{nT}} + \left(\frac{\zeta^{(t)}}{\bar{\zeta}_{S'}} \vee 1\right)\sqrt{\frac{r}{n}} + \zeta^{(t)}h\Bigg\}
$$

$$
+\frac{\zeta^{(t)}}{\overline{\zeta}_{S'}}\frac{\epsilon r}{\sqrt{n}}\bigg]\wedge\sqrt{\frac{p}{n}}\Bigg\}\Bigg)\geq\frac{1}{10},
$$

which implies the desired lower bound because $\sqrt{\frac{\log T}{n}} \leq \min_{t \in S'} \left\{ \left[\frac{\zeta^{(t)}}{\zeta} \right] \right\}$ ζ¯ $\sqrt{\frac{pr}{nT}} + (\frac{\zeta^{(t)}}{\bar{\zeta}} \vee$ $1\big)\sqrt{\frac{r}{n}}+\zeta^{(t)}h+\frac{\zeta^{(t)}}{\zeta}$ $rac{r}{\zeta}$ $\epsilon \frac{r}{\sqrt{n}}$ $\bigg\} \wedge \sqrt{\frac{p}{n}}\bigg\}.$

• If $\sqrt{\frac{\log T}{n}} > \min_{t \in S'} \left\{ \left[\frac{\zeta^{(t)}}{\zeta_{S'}} \right] \right\}$ $\overline{\bar\zeta}_{S'}$ $\sqrt{\frac{pr}{nT}} + (\frac{\zeta^{(t)}}{\bar{\zeta}_{S'}}$ $\frac{\zeta^{(t)}}{\zeta_{S'}} \vee 1 \bigvee \frac{\overline{r}}{n} + \zeta^{(t)}h + \frac{\zeta^{(t)}}{\zeta_{S'}}$ $\frac{\zeta^{(t)}}{\bar{\zeta}_{S'}}\epsilon \frac{r}{\sqrt{n}}$ $\left\{\wedge \sqrt{\frac{p}{n}}\right\}$: by definition of S', automatically we have $\sqrt{\frac{\log T}{n}} > \min_{t \in [T] \setminus S'} \left\{ \left[\frac{\zeta^{(t)}}{\zeta_{s'}} \right] \right\}$ $\bar{\zeta}_{S'}$ $\sqrt{\frac{pr}{nT}} + (\frac{\zeta^{(t)}}{\bar{\zeta}_{S'}}$ $\frac{\zeta^{(t)}}{\zeta_{S'}} \vee 1 \bigvee \frac{r}{n} + \zeta^{(t)}h +$ $\zeta^{(t)}$ $\frac{\zeta^{(t)}}{\zeta_{S'}}\epsilon \frac{r}{\sqrt{n}}$ $\left\{\lambda \sqrt{\frac{p}{n}}\right\}$. Then by the same argument we used to prove [\(21\)](#page-71-1), we can get the following bound by replacing [T] in [\(21\)](#page-71-1) with $|T|\setminus S'$:

$$
\inf_{\{\widehat{\boldsymbol{\beta}}^{(t)}\}_{t=1}^T} \sup_{\{\boldsymbol{\beta}^{(t)}\}_{t\in[T]} \in \mathscr{B}([T],0)} \mathbb{P}\Bigg(\max_{t\in[T]\backslash S'} \|\widehat{\boldsymbol{\beta}}^{(t)} - \boldsymbol{\beta}^{(t)}\|_2 \gtrsim \sqrt{\frac{\log(T-|S'|)}{n}}\Bigg) \ge \frac{1}{10}.
$$

This implies the desired lower bound, because when $T \geq r^{1.01}$ we have $\log(T - |S'|) \gtrsim$ $\log(T/r) \gtrsim \log T^{\frac{0.01}{1.01}} \gtrsim \log T.$

D.8 Proof of Theorem [13](#page-43-0)

First, let us prove a useful result: When $nT \ge C''pr$, $n \ge C''r$, and $\frac{\min_{t \in S} \zeta^{(t)}}{\zeta}$ $\frac{\epsilon_S \, \zeta^{(t)}}{\bar{\zeta}} \sqrt{\frac{r}{T}} \|\boldsymbol{D}^*_S\|_2 \leq$ $\sqrt{p + \log T}$ $\frac{\log T}{n}$, we have $\|\widehat{\overline{\bm{A}}}(\widehat{\overline{\bm{A}}})^\top - \overline{\bm{A}}(\overline{\bm{A}})^\top \|_2 \lesssim \bar{\zeta}^{-1} \sqrt{\frac{pr}{nT}}$ $\frac{\overline{pr}}{nT}+\bar{\zeta}^{-1}\sqrt{\frac{r}{n}}$ $\frac{\overline{r}}{n} + \bar{\zeta}^{-1} \sqrt{\frac{\overline{r}}{T}}$ $\frac{\overline{r}}{T} \|\boldsymbol{D}^*_S\|_2 + \frac{\max_{t\in S} \|\boldsymbol{\beta}^{(t)*}\|_2}{\bar{\zeta}}$ $\overline{\bar{\zeta}}$ √ $r\bar{\epsilon},~(25)$

w.p. at least $1 - e^{-C(p + \log T)}$, and

$$
\|\widehat{\overline{A}} \boldsymbol{\theta}_{\widehat{\overline{A}}}^{(t)} - \boldsymbol{\beta}^{(t)*}\|_2 \lesssim \frac{\zeta^{(t)}}{\bar{\zeta}} \sqrt{\frac{pr}{nT}} + \left(\frac{\zeta^{(t)}}{\bar{\zeta}} \vee 1\right) \sqrt{\frac{r}{n}} + \frac{\zeta^{(t)}}{\bar{\zeta}} \sqrt{\frac{r}{T}} \|\mathbf{D}^*\|_2 + \frac{\zeta^{(t)}}{\bar{\zeta}} \max_{t \in S} \|\boldsymbol{\beta}^{(t)*}\|_2 \sqrt{r\bar{\epsilon}}
$$

$$
+ \|\boldsymbol{\delta}^{(t)*}\|_2 + \sqrt{\frac{\log T}{n}}, \quad \forall t \in S,
$$
\n
$$
(26)
$$

w.p. at least $1 - e^{-C(r + \log T)}$.

Denote $\overline{B}_S = {\overline{A}} \theta^{(t)*} t \in S \in \mathbb{R}^{p \times |S|}$ and $\widetilde{B}_S = {\{\widetilde{\beta}^{(t)}\}} t \in S$. Note that $\widetilde{\beta}^{(t)} - \beta^{(t)*} =$ $(\hat{\Sigma}^{(t)})^{-1} \frac{1}{n} (\mathbf{X}^{(t)})^{\top} \boldsymbol{\epsilon}^{(t)}$ is a sub-Gaussian vector with variance proxy $n^{-1} \sigma_{\max}((\hat{\Sigma}^{(t)})^{-1})$ given $\mathbf{X}^{(t)}$, and $\{\widetilde{\beta}^{(t)} - \beta^{(t)*}\}_{t \in [T]}$ are independent of each other. By Lemma 5.39 in [Vershynin](#page-39-1) [\(2010\)](#page-39-1), conditioned on $\{X^{(t)}\}_{t\in S}$, we have

$$
\|\widetilde{\boldsymbol{B}}_S - \boldsymbol{B}_S^*\|_2 \lesssim \sqrt{\frac{p+T}{n}} \max_{t \in S} \sigma_{\max}((\widehat{\boldsymbol{\Sigma}}^{(t)})^{-1}),
$$

w.p. at least $1 - e^{-C(p + \log T)}$. Then combining

$$
\|(\widehat{\mathbf{\Sigma}}^{(t)})^{-1}\|_2 \le \|(\mathbf{\Sigma}^{(t)})^{-1}\|_2 + \|(\widehat{\mathbf{\Sigma}}^{(t)})^{-1}\|_2 \|\widehat{\mathbf{\Sigma}}^{(t)} - \mathbf{\Sigma}^{(t)}\|_2 \|(\mathbf{\Sigma}^{(t)})^{-1}\|_2
$$

with Lemma [19,](#page-47-2) we have $\sigma_{\max}((\hat{\Sigma}^{(t)})^{-1}) = ||(\hat{\Sigma}^{(t)})^{-1}||_2 \lesssim 1$. Therefore, w.p. at least $1 - e^{-C(p + \log T)},$

$$
\|\widetilde{\boldsymbol{B}}_S-\boldsymbol{B}_S^*\|_2\lesssim \sqrt{\frac{p+T}{n}}.
$$

Note that $\hat{\bm{B}}_S - \overline{\bm{B}}_S = (\hat{\bm{B}}_S - \widetilde{\bm{B}}_S) + (\widetilde{\bm{B}}_S - \bm{B}_S^*) + (\bm{B}_S^* - \overline{\bm{B}}_S)$. By triangle inequality,

$$
\|\widehat{\boldsymbol{B}} - (\overline{\boldsymbol{B}}S \ \boldsymbol{0})_{p \times T}\|_2 \le \|\widehat{\boldsymbol{B}}_{S^c}\|_2 + \|\widehat{\boldsymbol{B}}_S - \widetilde{\boldsymbol{B}}_S\|_2 + \|\widetilde{\boldsymbol{B}}_S - \boldsymbol{B}_S^*\|_2 + \|\boldsymbol{B}_S^* - \overline{\boldsymbol{B}}_S\|_2
$$

\n
$$
\le \|\widehat{\boldsymbol{B}}_{S^c}\|_F + \|\widehat{\boldsymbol{B}}_S - \widetilde{\boldsymbol{B}}_S\|_F + \|\widetilde{\boldsymbol{B}}_S - \boldsymbol{B}_S^*\|_2 + \|\boldsymbol{B}_S^* - \overline{\boldsymbol{B}}_S\|_2
$$

\n
$$
\le R\sqrt{\epsilon}\bar{T} + \sqrt{\sum_{t \in S} \|\Pi_R(\widetilde{\boldsymbol{\beta}}^{(t)}) - \widetilde{\boldsymbol{\beta}}^{(t)}\|_2^2} + C\sqrt{\frac{p+T}{n}} + \|\boldsymbol{D}_S^*\|_2,
$$

w.p. at least $1 - e^{-C'(p + \log T)}$. Since $R = \text{quantile}(\{\|\widetilde{\beta}^{(t)}\|_2\}_{t=1}^T, 1 - \bar{\epsilon}), \bar{\epsilon} \geq \epsilon$, and

 $\max_{t\in S} \|\widetilde{\boldsymbol{\beta}}^{(t)}\|_2 \leq \max_{t\in S} \|\widetilde{\boldsymbol{\beta}}^{(t)}{-}\boldsymbol{\beta}^{(t)\ast}\|_2 + \max_{t\in S} \|\boldsymbol{\beta}^{(t)\ast}\|_2 \leq C$ $p + \log T$ $\frac{\log I}{n} + \max_{t \in S} \|\boldsymbol{\beta}^{(t)*}\|_2 \leq C \max_{t \in S} \|\boldsymbol{\beta}^{(t)*}\|_2,$ w.p. at least $1 - e^{-C'(p + \log T)}$, we have

$$
R \leq C \max_{t \in S} \|\boldsymbol{\beta}^{(t)*}\|_2,
$$

w.p. at least $1 - e^{-C'(p + \log T)}$. And there are at most $\lceil \bar{\epsilon}T \rceil$ number of $t \in S$ satisfying $\prod_{R}(\widetilde{\beta}^{(t)}) \neq \widetilde{\beta}^{(t)}$. Therefore, w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
\|\widehat{\mathbf{B}} - (\overline{\mathbf{B}}S \ \mathbf{0})_{p \times T}\|_2 \le R\sqrt{\epsilon T} + C \max_{t \in S} \|\beta^{(t)*}\|_2 \sqrt{\epsilon T} + C\sqrt{\frac{p+T}{n}} + \|\mathbf{D}_S^*\|_2
$$

$$
\le 2C \max_{t \in S} \|\beta^{(t)*}\|_2 \sqrt{\epsilon T} + C\sqrt{\frac{p+T}{n}} + \|\mathbf{D}_S^*\|_2.
$$

Similarly, by Weyl's inequality, w.p. at least $1 - e^{-C'(p + \log T)}$, $\sigma_r(\widehat{B}) - \sigma_{r+1}((\overline{B}_S \quad 0)) = \sigma_r(\widehat{B})$

$$
\geq \sigma_r(\overline{\mathbf{B}}_S) - \|\widehat{\mathbf{B}}_{S^c}\|_2 - \|\widehat{\mathbf{B}}_S - \widetilde{\mathbf{B}}_S\|_2 - \|\widetilde{\mathbf{B}}_S - \mathbf{B}_S^*\|_2 - \|\mathbf{B}_S^* - \overline{\mathbf{B}}_S\|_2
$$

\n
$$
\geq c\overline{\zeta}\sqrt{\frac{T}{r}} - 2C \max_{t \in S} \|\beta^{(t)*}\|_2 \sqrt{\overline{\epsilon}T} - C\sqrt{\frac{p+T}{n}} - \|\mathbf{D}_S^*\|_2
$$

\n
$$
\geq \frac{c}{2}\overline{\zeta}\sqrt{\frac{T}{r}},
$$

since $\max_{t \in S} ||\boldsymbol{\beta}^{(t)*}||_2 \sqrt{\min_{t \in S} ||\boldsymbol{\beta}^{(t)*}||_2^2}$ $\frac{\overline{\overline{\epsilon}}}{\overline{\epsilon}} \leq \frac{c}{60}$ $6C$ $\frac{\bar{\zeta}}{\sqrt{r}},\ \sqrt{\frac{p+T}{n}} \leq \frac{c}{60}$ $6C$ $\sqrt{p + \log T}$ $\overline{\frac{\log T}{n}} \cdot \bar{\zeta} \sqrt{\frac{T}{r}} \leq \bar{\zeta} \frac{c}{60}$ $6C$ $\sqrt{\mathcal{I}}$ $\frac{T}{r}$, and $\|\mathbf{D}_{S}^{*}\|_{2} \leq$ c $\frac{c}{6}\bar{\zeta}\sqrt{\frac{T}{r}}$ $\frac{T}{r}$, because $\bar{\epsilon} \leq \frac{c''}{r}$ $\frac{\bar{\zeta}''}{r} \cdot \left(\frac{\bar{\zeta}}{\max_{t \in S} \|} \right)$ $\max_{t\in S}\|\bm{\beta}^{(t)*}\|_2$ \int^2 , $nT \ge C''pr$, $p \ge r$, $n \ge C''r$, and $\frac{1}{\zeta}\sqrt{\frac{r}{T}} \|\mathbf{D}_S^*\|_2 \le$ $\frac{1}{\zeta}\sqrt{\frac{r}{T}}\|\mathbf{D}_{S}^{*}\|_{2}\cdot\frac{\min_{t\in S}\zeta^{(t)}}{\sqrt{(p+\log T)/n}}\leq c'',$ where $c''>0$ is a small constant and $C''>0$ is a large constant.

Then by the SVD and Wedin's sin Θ-Theorem:

$$
\begin{split} \|\widehat{\vec{A}}(\widehat{\vec{A}})^{\top} - \overline{A}(\overline{A})^{\top}\|_{2} &\lesssim \frac{\|\widehat{\bm{B}} - (\overline{\bm{B}}_{S} \quad \bm{0})_{p \times T} \|_{2}}{\sigma_{r}(\widehat{\bm{B}}) - \sigma_{r+1}((\overline{\bm{B}}_{S} \quad \bm{0}))} \\ &\lesssim \frac{\max_{t \in S} \|\beta^{(t)*}\|_{2} \sqrt{\bar{\epsilon}T} + \sqrt{\frac{p+T}{n}} + \|\bm{D}_{S}^{*}\|_{2}}{\bar{\zeta}\sqrt{\frac{T}{r}}} \\ &\lesssim \bar{\zeta}^{-1} \sqrt{\frac{pr}{nT}} + \bar{\zeta}^{-1} \sqrt{\frac{r}{n}} + \bar{\zeta}^{-1} \sqrt{\frac{r}{T}} \|\bm{D}_{S}^{*}\|_{2} + \frac{\max_{t \in S} \|\beta^{(t)*}\|_{2}}{\bar{\zeta}} \sqrt{r\bar{\epsilon}}, \end{split}
$$

w.p. at least $1 - e^{-C(p + \log T)}$. Hence (25) holds.

Next, let us prove [\(26\)](#page-79-0). By noticing that

$$
\boldsymbol{\theta}_{\widehat{\widehat{\mathbf{A}}}}^{(t)} = ((\widehat{\widehat{\mathbf{A}}})^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \widehat{\widehat{\mathbf{A}}})^{-1} (\widehat{\widehat{\mathbf{A}}})^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{\beta}^{(t)*} - ((\widehat{\widehat{\mathbf{A}}})^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \widehat{\widehat{\mathbf{A}}})^{-1} (\widehat{\widehat{\mathbf{A}}})^{\top} \nabla f^{(t)} (\boldsymbol{\beta}^{(t)*}),
$$

we have

$$
\hat{\vec{A}}\theta_{\hat{\vec{A}}}^{(t)} - \beta^{(t)*} = \left[\hat{\vec{A}}((\hat{\vec{A}})^{\top}\hat{\Sigma}^{(t)}\hat{\vec{A}})^{-1}(\hat{\vec{A}})^{\top}\hat{\Sigma}^{(t)} - I\right]\beta^{(t)*} - ((\hat{\vec{A}})^{\top}\hat{\Sigma}^{(t)}\hat{\vec{A}})^{-1}(\hat{\vec{A}})^{\top}\nabla f^{(t)}(\beta^{(t)*})
$$
\n
$$
= (\hat{\Sigma}^{(t)})^{-1/2} \Big[(\hat{\Sigma}^{(t)})^{1/2}\hat{\vec{A}}((\hat{\vec{A}})^{\top}\hat{\Sigma}^{(t)}\hat{\vec{A}})^{-1}(\hat{\vec{A}})^{\top}(\hat{\Sigma}^{(t)})^{-1/2} - I \Big] (\hat{\Sigma}^{(t)})^{1/2}\beta^{(t)*}
$$
\n
$$
- ((\hat{\vec{A}})^{\top}\hat{\Sigma}^{(t)}\hat{\vec{A}})^{-1}(\hat{\vec{A}})^{\top}\nabla f^{(t)}(\beta^{(t)*}).
$$

Denote the projection matrix of a matrix $\mathbf{A} \in \mathbb{R}^{p_1 \times p_2}$ (projection onto the column space of A) with rank $(A) = p_2 \leq p_1$ as

$$
\boldsymbol{P_A} = \boldsymbol{A} (\boldsymbol{A}^\top \boldsymbol{A})^{-1} \boldsymbol{A}^\top.
$$

Denote $\overline{B} = (\widehat{\Sigma}^{(t)})^{1/2} \overline{A} \in \mathbb{R}^{p \times r}$ and $(\overline{B})^{\perp} = (\widehat{\Sigma}^{(t)})^{-1/2} (\overline{A})^{\perp} \in \mathbb{R}^{p \times (p-r)}$. Notice that $(\overline{B})^{\top}(\overline{B})^{\perp} = 0_{r \times (p-r)}.$ By definition $P_{\widehat{\overline{B}}} + P_{(\widehat{\overline{B}})^{\perp}} = I_p$. Hence

$$
\widehat{\vec{A}}\theta_{\widehat{\vec{A}}}^{(t)} - \beta^{(t)*} = (\widehat{\Sigma}^{(t)})^{-1/2} (P_{\widehat{\vec{B}}} - I) (\widehat{\Sigma}^{(t)})^{1/2} (\overline{A}\theta^{(t)*} + \delta^{(t)*}) - ((\widehat{\vec{A}})^{\top} \widehat{\Sigma}^{(t)} \widehat{\vec{A}})^{-1} (\widehat{\vec{A}})^{\top} \nabla f^{(t)} (\beta^{(t)*})
$$

$$
= -(\widehat{\Sigma}^{(t)})^{-1/2} P_{(\widehat{\overline{B}})^{\perp}}(\widehat{\Sigma}^{(t)})^{1/2} \overline{A} \theta^{(t)*} - (\widehat{\Sigma}^{(t)})^{-1/2} P_{(\widehat{\overline{B}})^{\perp}}(\widehat{\Sigma}^{(t)})^{1/2} \delta^{(t)*} - ((\widehat{\overline{A}})^{\top} \widehat{\Sigma}^{(t)} \widehat{\overline{A}})^{-1} (\widehat{\overline{A}})^{\top} \nabla f^{(t)}(\beta^{(t)*}).
$$

Note that since $\|\widehat{\mathbf{\Sigma}}^{(t)}\|_2$ and $\|(\widehat{\mathbf{\Sigma}}^{(t)})^{-1}\|_2$ are bounded from above by constants w.p. at least $1 - e^{-C'(p + \log T)}$, we have

$$
\begin{aligned} &\|(\widehat{\boldsymbol{\Sigma}}^{(t)})^{-1/2}\boldsymbol{P}_{(\widehat{\boldsymbol{B}})^{\perp}}(\widehat{\boldsymbol{\Sigma}}^{(t)})^{1/2}\overline{\boldsymbol{A}}\boldsymbol{\theta}^{(t)*}\|_{2}\lesssim \|((\widehat{\boldsymbol{A}})^{\perp})^{\top}\overline{\boldsymbol{A}}\boldsymbol{\theta}^{(t)*}\|_{2}+\|\boldsymbol{\delta}^{(t)*}\|_{2}\lesssim \|\widehat{\boldsymbol{A}}(\widehat{\boldsymbol{A}})^{\top}-\overline{\boldsymbol{A}}(\overline{\boldsymbol{A}})^{\top}\|_{2}\zeta^{(t)}+\|\boldsymbol{\delta}^{(t)*}\|_{2},\\ &\|(\widehat{\boldsymbol{\Sigma}}^{(t)})^{-1/2}\boldsymbol{P}_{(\widehat{\boldsymbol{B}})^{\perp}}(\widehat{\boldsymbol{\Sigma}}^{(t)})^{1/2}\boldsymbol{\delta}^{(t)*}\|_{2}\lesssim \|\boldsymbol{\delta}^{(t)*}\|_{2}, \end{aligned}
$$

w.p. at least $1 - e^{-C'(p + \log T)}$. Therefore w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
\|\widehat{\overline{A}}\boldsymbol{\theta}_{\widehat{\overline{A}}}^{(t)} - \boldsymbol{\beta}^{(t)*}\|_2 \lesssim \|\widehat{\overline{A}}(\widehat{\overline{A}})^\top - \overline{A}(\overline{A})^\top\|_2 \zeta^{(t)} + \|\boldsymbol{\delta}^{(t)*}\|_2 + \|((\widehat{\overline{A}})^\top \widehat{\boldsymbol{\Sigma}}^{(t)} \widehat{\overline{A}})^{-1} (\widehat{\overline{A}})^\top \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2.
$$

Let us handle the last term. Denote $\mathbf{R} = \arg \min_{\mathbf{R} \in \mathcal{O}^{r \times r}} \|\overline{\mathbf{A}} - \overline{\mathbf{A}} \mathbf{R}\|_2$. Therefore

$$
\begin{split}\n&\|((\widehat{\overline{A}})^{\top}\widehat{\Sigma}^{(t)}\widehat{\overline{A}})^{-1}(\widehat{\overline{A}})^{\top}\nabla f^{(t)}(\beta^{(t)*})\|_{2} \\
&\leq \|((\widehat{\overline{A}})^{\top}\widehat{\Sigma}^{(t)}\widehat{\overline{A}})^{-1}(\widehat{\overline{A}} - \overline{A}\widetilde{R})^{\top}\nabla f^{(t)}(\beta^{(t)*})\|_{2} + \|((\widehat{\overline{A}})^{\top}\widehat{\Sigma}^{(t)}\widehat{\overline{A}})^{-1}(\overline{A}\widetilde{R})^{\top}\nabla f^{(t)}(\beta^{(t)*})\|_{2} \\
&\leq \|\widehat{\overline{A}} - \overline{A}\widetilde{R}\|_{2}\|\nabla f^{(t)}(\beta^{(t)*})\|_{2} + \|(\overline{A}\widetilde{R})^{\top}\nabla f^{(t)}(\beta^{(t)*})\|_{2} \\
&\lesssim \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A}(\overline{A})^{\top}\|_{2}\sqrt{\frac{p + \log T}{n}} + \sqrt{\frac{r + \log T}{n}},\n\end{split}
$$

w.p. at least $1 - e^{-C'(r + \log T)}$. Hence, w.p. at least $1 - e^{-C'(r + \log T)}$,

$$
\|\widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)} - \boldsymbol{\beta}^{(t)*}\|_{2} \lesssim \|\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top} - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}\|_{2}\zeta^{(t)} + \|\boldsymbol{\delta}^{(t)*}\|_{2} + \sqrt{\frac{r + \log T}{n}} \\
\lesssim \left(\bar{\zeta}^{-1}\sqrt{\frac{pr}{nT}} + \bar{\zeta}^{-1}\sqrt{\frac{r}{n}} + \bar{\zeta}^{-1}\sqrt{\frac{r}{T}}\|\boldsymbol{D}_{S}^{*}\|_{2} + \frac{\max_{t \in S} \|\boldsymbol{\beta}^{(t)*}\|_{2}}{\bar{\zeta}}\sqrt{r\bar{\epsilon}}\right)\zeta^{(t)} \\
+ \|\boldsymbol{\delta}^{(t)*}\|_{2} + \sqrt{\frac{r + \log T}{n}} \\
\lesssim \frac{\zeta^{(t)}}{\bar{\zeta}}\sqrt{\frac{pr}{nT}} + \left(\frac{\zeta^{(t)}}{\bar{\zeta}} \vee 1\right)\sqrt{\frac{r}{n}} + \frac{\zeta^{(t)}}{\bar{\zeta}}\sqrt{\frac{r}{T}}\|\boldsymbol{D}^{*}\|_{2} + \frac{\zeta^{(t)}}{\bar{\zeta}}\max_{t \in S} \|\boldsymbol{\beta}^{(t)*}\|_{2}\sqrt{r\bar{\epsilon}} \\
+ \|\boldsymbol{\delta}^{(t)*}\|_{2} + \sqrt{\frac{\log T}{n}} \\
:= \eta^{(t)}.
$$

Therefore, [\(26\)](#page-79-0) holds. Now let us use [\(25\)](#page-78-0) and [\(26\)](#page-79-0) to complete the proof of the theorem. (i) For $t \in S$ with $\eta^{(t)} \leq c' \sqrt{\frac{p + \log T}{n}}$ with a small $c' > 0$: we must have

$$
\|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2 + \|\widehat{\bar{\boldsymbol{A}}}\boldsymbol{\theta}_{\widehat{\bar{\boldsymbol{A}}}}^{(t)} - \boldsymbol{\beta}^{(t)*}\|_2 \lesssim \sqrt{\frac{p + \log T}{n}} + \eta^{(t)} \lesssim \frac{\gamma}{\sqrt{n}},
$$

hence by Lemma [39.](#page-52-0)(i), $\widehat{\beta}^{(t)} = \widehat{\overline{A}} \theta_{\widehat{\overline{A}}}^{(t)}$ and $\|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \lesssim \eta^{(t)}$ w.p. at least $1 - e^{-C'(r + \log T)}$. \boldsymbol{A} Note that we used [\(26\)](#page-79-0) to obtain the bound $\|\hat{\overline{A}} \theta_{\hat{\overline{A}}}^{(t)}\|$ $-\beta^{(t)*}$ |2 $\leq \eta^{(t)}$, w.p. at least 1 – $e^{-C'(r+\log T)}$. We can use it because $\eta^{(t)} \leq c' \sqrt{\frac{p+\log T}{n}}$ $\frac{\log T}{n}$ implies that $\frac{\min_{t \in S} \zeta^{(t)}}{\bar{\zeta}}$ $\frac{\epsilon_S \, \zeta^{(t)}}{\bar{\zeta}} \sqrt{\frac{r}{T}} \|\boldsymbol{D}^*\|_2 \leq$ $c' \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$, hence the condition of [\(26\)](#page-79-0) applies and (26) holds. (ii) For any $t \in S$, by Lemma [39,](#page-52-0) $\max_{t \in S} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \leq C \frac{\gamma}{\sqrt{n}} + \max_{t \in S} \|\widetilde{\beta}^{(t)} - \beta^{(t)*}\|_2$ w.p. at least $1 - e^{-C(p + \log T)}$, where $\widetilde{\beta}^{(t)} \in \arg \min_{\beta \in \mathbb{R}^p} f^{(t)}(\beta)$. Since $\gamma \asymp$ √ $\overline{p + \log T}$, this

$$
\max_{t \in S} \|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \lesssim \sqrt{\frac{p + \log T}{n}},
$$

w.p. at least $1 - e^{-C'(p + \log T)}$.

implies that

Combining (i) and (ii), we get the desired bound:

$$
\|\widehat{\beta}^{(t)} - \beta^{(t)*}\|_2 \lesssim \left\{ \frac{\zeta^{(t)}}{\bar{\zeta}} \sqrt{\frac{pr}{nT}} + \left(\frac{\zeta^{(t)}}{\bar{\zeta}} \vee 1\right) \sqrt{\frac{r}{n}} + \frac{\zeta^{(t)}}{\bar{\zeta}} \sqrt{\frac{r}{T}} \|\mathbf{D}_S^*\|_2 + \|\boldsymbol{\delta}^{(t)*}\|_2 + \sqrt{\frac{\log T}{n}} + \frac{\zeta^{(t)}}{\bar{\zeta}} \max_{t \in S} \|\boldsymbol{\beta}^{(t)*}\|_2 \sqrt{r\bar{\epsilon}} \right\} \wedge \sqrt{\frac{p + \log T}{n}},
$$

for all $t \in S$, w.p. at least $1 - e^{-C'(r + \log T)}$.

When the tasks in S^c are generated by the linear model, we can get

$$
\max_{t\in S^c}\|\widehat{\beta}^{(t)}-\beta^{(t)*}\|_2\lesssim \sqrt{\frac{p+\log T}{n}},
$$

w.p. at least $1 - e^{-C'(p + \log T)}$, similar to the argument in the proof of Theorem [1.](#page-12-0)

D.9 Proof of Theorem [7](#page-17-0)

We will transform the setting in Section [2.1](#page-8-0) into a form of the general setting in Section [A,](#page-42-0) then apply Theorem [13](#page-43-0) to obtain the result.

In Section [2.1,](#page-8-0) we have $\boldsymbol{\beta}^{(t)*} = \boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}$ for $t \in S$. Denote the projection matrix of a matrix $\mathbf{A} \in \mathbb{R}^{p_1 \times p_2}$ (projection onto the column space of \mathbf{A}) with $\text{rank}(\mathbf{A}) = p_2 < p_1$ as

$$
\boldsymbol{P_A} = \boldsymbol{A} (\boldsymbol{A}^\top \boldsymbol{A})^{-1} \boldsymbol{A}^\top.
$$

Note that $P_A + P_{A^{\perp}} = I_{p_1}$, where the orthogonal $A^{\perp} \in \mathcal{O}^{p_1 \times (p_1 - p_2)}$ satisfies $(A^{\perp})^{\top} A =$ $\mathbf{0}_{(p_1-p_2)\times p_2}$. Decompose $\boldsymbol{\beta}^{(t)*}$ as

$$
\pmb{\beta}^{(t)*}=\pmb{P}_{\overline{\pmb{A}}} \pmb{A}^{(t)*} \pmb{\theta}^{(t)*}+\pmb{P}_{(\overline{\pmb{A}})^\perp} \pmb{A}^{(t)*} \pmb{\theta}^{(t)*}=\overline{\pmb{A}}[(\overline{\pmb{A}})^\perp \pmb{A}^{(t)*} \pmb{\theta}^{(t)*}]+\pmb{P}_{(\overline{\pmb{A}})^\perp} \pmb{A}^{(t)*} \pmb{\theta}^{(t)*}.
$$

Therefore, this setting can be seen as a special case of the general setting in Section [A,](#page-42-0) by setting $\delta^{(t)*} = P_{(\overline{A})^\perp} A^{(t)*} \theta^{(t)*}$. Notice that $\sigma_{\min}((\overline{A})^\perp B_S^*) =$

Furthermore, note that

$$
\|(\overline{\bm{A}})^{\perp} \bm{A}^{(t)*} \bm{\theta}^{(t)*}\|_2 \leq \|\bm{\theta}^{(t)*}\|_2 = \zeta^{(t)},
$$

$$
\|\boldsymbol{\delta}^{(t)*}\|_{2} \leq \|\boldsymbol{P}_{(\overline{A})^{\perp}}\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2} \lesssim \|\boldsymbol{A}^{(t)*}(\boldsymbol{A}^{(t)*})^{\top} - \overline{\boldsymbol{A}}(\overline{A})^{\top}\|_{2}\zeta^{(t)} \lesssim h\zeta^{(t)},
$$

$$
\frac{1}{\sqrt{T}}\|\boldsymbol{D}_{S}^{*}\|_{2} \leq \frac{1}{\sqrt{T}}\|\boldsymbol{D}_{S}^{*}\|_{\mathrm{F}} \cdot \frac{\sigma_{\max}(\boldsymbol{D}_{S}^{*})}{\sqrt{r}\sigma_{\min}(\boldsymbol{D}_{S}^{*})} \lesssim \overline{\zeta} \cdot \frac{\sigma_{\max}(\boldsymbol{D}_{S}^{*})}{\sqrt{r}\sigma_{\min}(\boldsymbol{D}_{S}^{*})} \cdot h,
$$
(27)

where $\mathcal{D}_{S}^{*} = \mathcal{P}_{(\overline{A})^{\perp}} \mathcal{B}_{S}^{*}$ and $\|\mathcal{D}_{S}^{*}\|_{\mathrm{F}} \leq h \sqrt{\sum_{t \in S} ||\boldsymbol{\delta}^{(t)*}||_{2}^{2}} \leq h \sqrt{\sum_{t \in S} (\zeta^{(t)})^{2}}$. Regarding the last inequality (27) , there is an alternative way to bound the LHS:

$$
\frac{1}{\sqrt{T}}\|\mathbf{D}_{S}^{*}\|_{2} \leq \frac{1}{\sqrt{T}}\|\mathbf{D}_{S}^{*}\|_{\mathrm{F}} \leq \bar{\zeta}h.
$$

Therefore,

$$
\frac{1}{\sqrt{T}}\|\boldsymbol{D}_S^*\|_2 \lesssim \bar{\zeta}h\cdot \left[\frac{\sigma_{\max}(\boldsymbol{D}_S^*)}{\sqrt{r}\sigma_{\min}(\boldsymbol{D}_S^*)}\wedge 1\right].
$$

Finally, it is straightforward to see that $D_S^* = P_{(\overline{A})^\perp} B_S^*$ and $(\overline{A}^\perp)^\top B_S^*$ share the same singular values. Also note that $\|\beta^{(t)}\|_2 = \|\theta^{(t)}\|_2 = \zeta^{(t)}$ for all $t \in S$. A direct application of Theorem [13](#page-43-0) leads to the desired bound, which completes the proof.

D.10 Proof of Theorem [15](#page-45-0)

If Algorithm [4](#page-45-1) is coupled with Algotithm [1,](#page-11-0) the proof idea is very similar to the proof of Theorem [1.](#page-12-0) When the representation learning helps, an argument based on Lemma [40,](#page-52-1) Proposition [43,](#page-52-2) and Lemma [42.](#page-52-3)(i) lead to the corresponding term. In the other case, Lemma [42.](#page-52-3)(ii) guarantees that the single-task rate holds all the time. A combination of these two situations entails the final TL upper bound. The details are as follows.

Denote $\eta = r \sqrt{\frac{p}{nT}} + \sqrt{r}h + \sqrt{r} \sqrt{\frac{r + \log T}{n}} + \frac{|S^c|}{T}$ $\frac{S^c|}{T} \cdot r \cdot \frac{\lambda}{\sqrt{n}}$ $\frac{1}{n}$. (i) When $\eta \leq C \sqrt{\frac{p}{n_0}}$: note that $\hat{\overline{A}}$ is **independent** of $\{x_i^{(0)}\}$ $\binom{0}{i}, y_i^{(0)}$ ${}_{i}^{(0)}\rbrace_{i=1}^{n_0}$, which is the key for part (i) to be correct by only requiring $n \ge Cr$ (see Remark [14\)](#page-44-0). Hence by Lemma [40](#page-52-1) and an argument by first conditioning on \overline{A} then taking the expectation,

$$
\begin{aligned}\n\|\widehat{\overline{A}}\boldsymbol{\theta}_{\widehat{\overline{A}}}^{(0)} - \boldsymbol{A}^{(0)*}\boldsymbol{\theta}^{(0)*}\|_{2} &\lesssim \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \boldsymbol{A}^{(0)*}(\boldsymbol{A}^{(0)*})^{\top}\|_{2} + \sqrt{\frac{r}{n_{0}}} \\
&\leq \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{\boldsymbol{A}}(\overline{A})^{\top}\|_{2} + \|\overline{\boldsymbol{A}}(\overline{A})^{\top} - \boldsymbol{A}^{(0)*}(\boldsymbol{A}^{(0)*})^{\top}\|_{2} + \sqrt{\frac{r}{n_{0}}} \\
&\lesssim \eta + \sqrt{\frac{r}{n_{0}}} \n\end{aligned}
$$

w.p. at least $1-e^{-C(r+\log T)}$, where we used Proposision [43](#page-52-2) in the last step. Then by Lemma $42.(\text{i}), \frac{\gamma}{\sqrt{n_0}} \gtrsim \sqrt{\frac{p}{n_0}} \geq \|\nabla f^{(t)}(\bm{A}^{(0)*}\bm{\theta}^{(0)*})\|_2 + C\|\widehat{\bar{\bm{A}}}\bm{\theta}^{(0)}_{\widehat{\bar{\bm{A}}}}$ $42.(\text{i}), \frac{\gamma}{\sqrt{n_0}} \gtrsim \sqrt{\frac{p}{n_0}} \geq \|\nabla f^{(t)}(\bm{A}^{(0)*}\bm{\theta}^{(0)*})\|_2 + C\|\widehat{\bar{\bm{A}}}\bm{\theta}^{(0)}_{\widehat{\bar{\bm{A}}}}$ $- A^{(0)*} \theta^{(0)*} \|_2$, which implies that $\Vert \widehat{\boldsymbol{\beta}}^{(0)} - \boldsymbol{\beta}^{(0)\ast} \Vert_2 = \Vert \widehat{\bar{\boldsymbol{A}}}\boldsymbol{\theta}^{(0)}_{\widehat{\bar{\boldsymbol{A}}}}$ $-\boldsymbol{A}^{(0)*}\boldsymbol{\theta}^{(0)*}\|_2 \leq \eta,$

w.p. at least $1 - e^{-C(r + \log T)}$.

(ii) By Lemma [42.](#page-52-3)(ii), we always have $\|\widehat{\beta}^{(0)} - \beta^{(0)*}\|_2 \lesssim \sqrt{p/n_0}$ w.p. at least $1 - e^{-C(p + \log T)}$.

If Algorithm [4](#page-45-1) is coupled with Algotithm [2,](#page-17-1) we can follow the discussions above and the proof of Theorem [7](#page-17-0) to get the desired result.

D.11 Proof of Theorem [16](#page-46-0)

Similar to the proof of Theorem [8,](#page-19-0) we prove the following parts one by one. Combining them together entials the lower bound. Throughout this subsection, we assume the following generative model for tasks in $\{0\} \cup S$:

$$
y|\boldsymbol{x} \sim \mathbb{P}_{y|\boldsymbol{x}, \boldsymbol{\beta}} = N(\boldsymbol{x}^\top\boldsymbol{\beta}, 1), \quad \boldsymbol{x} \sim \mathbb{P}_{\boldsymbol{x}},
$$

where \mathbb{P}_x is sub-Gaussian with $\Sigma = \mathbb{E}(xx^\top)$. Suppose there exist constants c, C such that $0 < c \leq \lambda_{\min}(\Sigma) \leq \lambda_{\max}(\Sigma) \leq C < \infty$. Then any joint distribution $\mathbb P$ of (x, y) can be written as $\mathbb{P}_{x,y} = \mathbb{P}_{y|x;\beta} \cdot \mathbb{P}_x$.

Denote

$$
A_0(S, h) = \left\{ \{A^{(t)}\}_{t \in \{0\} \cup S} \subseteq \mathcal{O}^{p \times r} : \max_{t \in S} \|A^{(t)}(A^{(t)})^\top - A^{(0)}(A^{(0)})^\top \|_2 \le h \right\},
$$

$$
\Theta_0(S) = \left\{ \{ \theta^{(t)}\}_{t \in \{0\} \cup S} \subseteq \mathbb{R}^r : \max_{t \in \{0\} \cup S} \| \theta^{(t)} \|_2 \le C, \frac{1}{|S|} \sum_{t \in S} \theta^{(t)}(\theta^{(t)})^\top \ge \frac{c}{r} I_r \right\}.
$$

(i) Consider the case $S = [T]$ and $h = 0$. For any $\mathbf{A} \in \mathbb{R}^{p \times r}$ with $C \geq \sigma_1(\mathbf{A}) \geq \sigma_2(\mathbf{A}) \geq$ $\cdots \geq \sigma_r(A) \geq c > 0$, consider its SVD where $A = Q\Lambda V$ where $Q \in \mathcal{O}^{p \times r}$ and $V \in \mathcal{O}^{p \times r}$ $\mathcal{O}^{r \times r}$. Denote $\mathbf{R} = \Lambda \mathbf{V}$. Then $0 < c \leq \sigma_{\min}(\mathbf{R}) \leq \sigma_{\max}(\mathbf{R}) \leq C < \infty$, and $\mathbf{A}\boldsymbol{\theta}^{(t)} =$ $\boldsymbol{Q}\boldsymbol{R}\boldsymbol{\theta}^{(t)} = \boldsymbol{Q}\widetilde{\boldsymbol{\theta}}^{(t)}$. Note that $\boldsymbol{Q} \in \mathcal{O}^{p \times r}$ and $\widetilde{\boldsymbol{\theta}}^{(t)} = \boldsymbol{R}\boldsymbol{\theta}^{(t)}$ with $\|\widetilde{\boldsymbol{\theta}}^{(t)}\|_2 \leq \|\boldsymbol{R}\|_2\|\boldsymbol{\theta}^{(t)}\|_2 \leq C'$ and $\frac{1}{T} \sum_{t=1}^T \widetilde{\theta}^{(t)} (\widetilde{\theta}^{(t)})^{\top} = \mathbf{R} \Big[\frac{1}{T} \Big]$ $\frac{1}{T}\sum_{t=1}^T \boldsymbol{\theta}^{(t)}(\boldsymbol{\theta}^{(t)})^\top \big] \boldsymbol{R}^\top \succeq \frac{cc'}{r}$ $\frac{c'}{r}\bm{I}_r \ \ \text{if} \ \ \frac{1}{T}\sum_{t=1}^T \bm{\theta}^{(t)}(\bm{\theta}^{(t)})^\top \ \succeq \ \frac{c'}{r}$ $\frac{c'}{r} \boldsymbol{I}_r.$ Therefore, fixing $\boldsymbol{\theta}^{(t)} \in \{e_j\}_{j=1}^r$ and $\boldsymbol{\theta}^{(0)} = \frac{1}{\sqrt{n}}$ $\frac{1}{T} \mathbf{1}_r$ s.t. $\{\boldsymbol{\theta}^{(t)}\}_{t=0}^T \in \Theta_0([T]),$ we know that

$$
\inf_{\widehat{\boldsymbol{\beta}}^{(0)}} \sup_{\{\boldsymbol{\beta}^{(t)}\}_{t=1}^T \in \mathscr{B}_0([T],0)} \mathbb{P}\bigg(\|\widehat{\boldsymbol{\beta}}^{(0)} - \boldsymbol{A}^{(0)}\boldsymbol{\theta}^{(0)}\|_2 \gtrsim \sqrt{\frac{pr}{nT}} \wedge \sqrt{\frac{p}{n_0}} \bigg)
$$
\n
$$
\geq \inf_{\widehat{\boldsymbol{\beta}}^{(0)}} \sup_{0 < c \leq \sigma_{\text{max}}(\boldsymbol{A}) \leq \sigma_{\text{max}}(\boldsymbol{A}) \leq C} \mathbb{P}\bigg(\|\widehat{\boldsymbol{\beta}}^{(0)} - \boldsymbol{A}\boldsymbol{\theta}^{(0)}\|_2 \gtrsim \sqrt{\frac{pr}{nT}} \wedge \sqrt{\frac{p}{n_0}} \bigg)
$$
\n
$$
\coloneq (\ast),
$$

where $\mathbf{A}^{(t)} = \mathbf{A}$ for all $t \in \{0\} \cup [T]$. Let $\delta = \sqrt{\frac{pr}{nT}} \wedge \sqrt{\frac{p}{n_0}}$. Consider a $\delta/2$ -packing of the ball $\mathcal{B}_{\delta}(\mathbf{0}, \mathbb{R}^p, \|\cdot\|_2)$ (denoted as M). By Example 5.8 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0), $|\mathcal{M}| \geq 5^p$. Consider

$$
\mathbf{A} = \overline{\mathbf{A}} + \frac{c'}{\sqrt{r}} \cdot \mathbf{u} \mathbf{1}_r^\top, \quad \widetilde{\mathbf{A}} = \overline{\mathbf{A}} + \frac{c'}{\sqrt{r}} \cdot \widetilde{\mathbf{u}} \mathbf{1}_r^\top,
$$

where $u \neq \tilde{u} \in \mathcal{M}, c' > 0$ is a small constant such that $\sigma_{\min}(A), \sigma_{\min}(\tilde{A}) \geq 1 - c' \geq c$ and $\sigma_{\max}(\boldsymbol{A}), \sigma_{\max}(\widetilde{\boldsymbol{A}}) \leq 1 + c' \leq C.$ Then

$$
\|(\boldsymbol{A} - \widetilde{\boldsymbol{A}})\boldsymbol{\theta}^{(0)}\|_2 = \frac{1}{r} \|(\boldsymbol{u} - \widetilde{\boldsymbol{u}})\boldsymbol{1}_r^\top \boldsymbol{1}_r\|_2 = \|\boldsymbol{u} - \widetilde{\boldsymbol{u}}\|_2 \in \left[\frac{1}{2}\delta, 2\delta\right],\tag{28}
$$

$$
\|(\boldsymbol{A} - \widetilde{\boldsymbol{A}})\boldsymbol{\theta}^{(t)}\|_2 = \|(\boldsymbol{A} - \widetilde{\boldsymbol{A}})\boldsymbol{e}_{j_k}\|_2 = \frac{1}{\sqrt{r}}\|\boldsymbol{u} - \widetilde{\boldsymbol{u}}\|_2 \le \frac{\delta}{\sqrt{r}}.
$$
\n(29)

where $\boldsymbol{\theta}^{(t)} = \boldsymbol{e}_{j_k}$ with $j_t \in [r]$. Therefore $\mathscr{B} = \{ \boldsymbol{\beta} \in \mathbb{R}^p : \boldsymbol{\beta} = \boldsymbol{A} \boldsymbol{\theta}^{(0)}, \boldsymbol{A} = \overline{\boldsymbol{A}} + \frac{1}{\sqrt{n}}\}$ $\frac{1}{r}\cdot u \mathbf{1}_{r}^{\top}, u\in$ M} becomes a $\delta/2$ -packing in \mathbb{R}^p with $|\mathscr{B}| = |\mathcal{M}| \geq 5^p$. Furthermore, for any $\boldsymbol{\beta} = \boldsymbol{A}\boldsymbol{\theta}^{(0)}$ and $\widetilde{\beta} = \widetilde{A}\theta^{(0)} \in \mathscr{B}$ with $A \neq \widetilde{A}$, by Lemma [25](#page-48-0) and equations [\(28\)](#page-84-0) and [\(29\)](#page-84-1),

$$
\begin{split}\n\text{KL}\left(\prod_{t=1}^{T} \mathbb{P}_{y|\mathbf{x};\mathbf{A}\theta^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(t)\otimes n} \cdot \mathbb{P}_{y|\mathbf{x};\mathbf{A}\theta^{(0)}}^{(0)\otimes n_0} \middle\|\prod_{t=1}^{T} \mathbb{P}_{y|\mathbf{x};\widetilde{\mathbf{A}}\theta^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(t)\otimes n} \cdot \mathbb{P}_{y|\mathbf{x};\widetilde{\mathbf{A}}\theta^{(0)}}^{(t)\otimes n_0} \cdot \mathbb{P}_{\mathbf{x}}^{(0)\otimes n_0} \right) \\
&\lesssim n \sum_{t=1}^{T} \|(A-\widetilde{A})\theta^{(t)}\|_{2}^{2} + n_{0}\|(A-\widetilde{A})\theta^{(0)}\|_{2}^{2} \\
&\lesssim nT \cdot \frac{\delta^{2}}{r} + n_{0}\delta^{2} \\
&\leq c \log |\mathscr{B}|.\n\end{split}
$$

Then by Fano's lemma (Lemma [24\)](#page-48-1), $(*) \geq 1 - \frac{\log 2}{\log |\mathcal{B}|} - c \geq 1/10$. (ii) Consider the case $S = [T]$. For all $t \in [T]$, fix $\mathbf{A}^{(t)} =$ an arbitrary $\overline{\mathbf{A}} \in \mathcal{O}^{p \times r}$. Fix $\{\boldsymbol{\theta}^{(t)}\}_{t=1}^T \subseteq \{e_j\}_{j=1}^r \text{ and } \boldsymbol{\theta}^{(0)} = \frac{1}{\sqrt{n}}$ $\frac{1}{T}I_r$ s.t. $\{\boldsymbol{\theta}^{(t)}\}_{t=0}^T \in \Theta_0([T])$. Set $\delta = \frac{h}{12} \wedge \sqrt{\frac{p}{n_0}}$. Consider a $\delta/2$ -packing of the ball $\mathcal{B}_{\delta}(\mathbf{0}, \mathbb{R}^p, \|\cdot\|_2)$ (denoted as M). By Example 5.8 in [Wainwright](#page-39-0) $(2019), |\mathcal{M}| \geq 5^p$ $(2019), |\mathcal{M}| \geq 5^p$. Consider

$$
\bm{A} = \overline{\bm{A}} + \frac{c'}{\sqrt{r}} \cdot \bm{u} \bm{1}_r^\top, \quad \widetilde{\bm{A}} = \overline{\bm{A}} + \frac{c'}{\sqrt{r}} \cdot \widetilde{\bm{u}} \bm{1}_r^\top,
$$

where $u \neq \widetilde{u} \in \mathcal{M}$ and c' is a small constant, then

$$
\|(\mathbf{A} - \widetilde{\mathbf{A}})\boldsymbol{\theta}^{(0)}\|_2 = \frac{1}{r} \|(\mathbf{u} - \widetilde{\mathbf{u}})\mathbf{1}_r^\top \mathbf{1}_r\|_2 = \|\mathbf{u} - \widetilde{\mathbf{u}}\|_2 \in \left[\frac{c'}{2}\delta, 2c'\delta\right],\tag{30}
$$

hence $\mathscr{B} = \{ \boldsymbol{\beta} \in \mathbb{R}^p : \boldsymbol{\beta} = \boldsymbol{A} \boldsymbol{\theta}^{(0)}, \boldsymbol{A} = \overline{\boldsymbol{A}} + \frac{c'}{\sqrt{r}} \cdot \boldsymbol{u} \boldsymbol{1}_r^\top, \boldsymbol{u} \in \mathcal{M} \}$ is a $c' \delta/2$ -packing in \mathbb{R}^p with $|\mathscr{B}| = |\mathcal{M}| \geq 5^p$. On the other hand, for any $\mathbf{A} \in \mathbb{R}^{p \times r}$ with $1 + 2c' \delta \geq \sigma_1(\mathbf{A}) \geq \sigma_2(\mathbf{A}) \geq$ $\cdots \geq \sigma_r(A) \geq 1 - 2c'\delta > 0$ and $||A - \overline{A}||_2 \leq \delta$, consider its SVD where $A = Q\Lambda V$ where $\mathbf{Q} \in \mathcal{O}^{p \times r}$ and $\mathbf{V} \in \mathcal{O}^{r \times r}$. Denote $\mathbf{R} = \Lambda \mathbf{V}$. Then $0 < 1 - 2c' \delta \le \sigma_{\min}(\mathbf{R}) \le \sigma_{\max}(\mathbf{R}) \le$ $1+2c'\delta < \infty$, and $\mathbf{A}\boldsymbol{\theta}^{(0)} = \mathbf{Q}\mathbf{R}\boldsymbol{\theta}^{(0)} = \mathbf{Q}\widetilde{\boldsymbol{\theta}}^{(0)}$. Note that $\mathbf{Q} \in \mathcal{O}^{p \times r}$ and $\widetilde{\boldsymbol{\theta}}^{(0)} = \mathbf{R}\boldsymbol{\theta}^{(0)}$ with $\|\widetilde{\theta}^{(0)}\|_2 \leq \|R\|_2 \|\theta^{(0)}\|_2 \leq C[1+2c']$. And $\|QQ^\top - \overline{A}(\overline{A})^\top\|_2 \leq \| (QR)(QR)^\top - \overline{A}(\overline{A})^\top\|_2 +$ $\|\mathbf{Q}(\mathbf{I}_r - \mathbf{R}\mathbf{R}^\top)\mathbf{Q}^\top\|_2 \le 4\|\mathbf{A} - \overline{\mathbf{A}}\|_2 + 4\|\mathbf{I}_r - \mathbf{R}\|_2 \le 4\delta + 8c'\delta \le 12\delta \le h$. Therefore,

$$
\inf_{\hat{\beta}^{(0)}} \sup_{\{A^{(t)}\}_{t \in \{0\}\cup S} \in \mathcal{A}_0([T],h)} \mathbb{P}\left(\|\hat{\beta}^{(0)} - A^{(0)}\theta^{(0)}\|_2 \gtrsim h \wedge \sqrt{\frac{p}{n_0}}\right) \n\{\theta^{(t)}\}_{t \in \{0\}\cup S} \in \Theta_0([T]) \n\ge \inf_{\hat{\beta}^{(0)} \; 0 < c \le \sigma_{\max}(A) \le \sigma_{\max}(A) \le C} \mathbb{P}\left(\|\hat{\beta}^{(0)} - A\theta^{(0)}\|_2 \gtrsim h \wedge \sqrt{\frac{p}{n_0}}\right) \n:= (*),
$$

where all $\mathbf{A}^{(t)} =$ an arbitrary $\overline{\mathbf{A}} \in \mathcal{O}^{p \times r}$, and $\{\theta^{(t)}\}_{t=1}^T \subseteq \{e_j\}_{j=1}^r$ are fixed.

For any $\beta = A\theta^{(0)}$ and $\tilde{\beta} = \tilde{A}\theta^{(0)} \in \mathscr{B}$ with $A \neq \tilde{A}$, by Lemma [25](#page-48-0) and equation [\(30\)](#page-85-0),

$$
\begin{split}\n&\text{KL}\left(\prod_{t=1}^{T} \mathbb{P}_{y|\mathbf{x};\overline{A}\theta^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(t)\otimes n} \cdot \mathbb{P}_{y|\mathbf{x};A\theta^{(0)}}^{(0)\otimes n_0} \cdot \mathbb{P}_{\mathbf{x}}^{(0)\otimes n_0} \middle\| \prod_{t=1}^{T} \mathbb{P}_{y|\mathbf{x};\overline{A}\theta^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{y|\mathbf{x};\widetilde{A}\theta^{(0)}}^{(0)\otimes n_0} \cdot \mathbb{P}_{\mathbf{x}}^{(0)\otimes n_0}\right) \\
&= \text{KL}\left(\mathbb{P}_{y|\mathbf{x};A\theta^{(0)}}^{(0)\otimes n_0} \cdot \mathbb{P}_{\mathbf{x}}^{(0)\otimes n_0} \middle\| \mathbb{P}_{y|\mathbf{x};\widetilde{A}\theta^{(0)}}^{(0)\otimes n_0} \cdot \mathbb{P}_{\mathbf{x}}^{(0)\otimes n_0}\right) \\
&\lesssim n_0 \|(A - \widetilde{A})\theta^{(0)}\|_2^2 \\
&\lesssim n_0 \delta^2 \\
&\leq c \log |\mathcal{B}|.\n\end{split}
$$

Then by Fano's lemma (Lemma [24\)](#page-48-1), $(*) \geq 1 - \frac{\log 2}{\log |\mathcal{B}|} - c \geq 1/10$. (iii) Consider the case that $S = [T]$ and $h = 0$. Fix all $A^{(t)} =$ some $A \in \mathcal{O}^{p \times r}$. Fixing $\{\boldsymbol{\theta}^{(t)}\}_{t=1}^T$ such that $\frac{1}{T} \sum_{t=1}^T \boldsymbol{\theta}^{(t)}(\boldsymbol{\theta}^{(t)})^\top \succeq \frac{c}{r}$ $\frac{c}{r}I_r$. We want to show

$$
\inf_{\widehat{\boldsymbol{\beta}}^{(0)}} \sup_{\|\boldsymbol{\theta}^{(0)}\|_2 \leq C} \mathbb{P}\left(\|\widehat{\boldsymbol{\beta}}^{(0)} - \boldsymbol{A}\boldsymbol{\theta}^{(0)}\|_2 \geq c\sqrt{\frac{r}{n_0}}\right) \geq \frac{1}{10}.
$$

Denote $\delta = \sqrt{\frac{r}{n_0}}$. Consider a c_δ-packing of $\mathcal{B}_{\delta}(\mathbf{0}, \mathbb{R}^r, \|\cdot\|_2)$ (denoted as \mathscr{T}). By Example 5.8 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0), we know that $\log |\mathcal{F}| \gtrsim r$. Denote the distribution of $\{y_i^{(0)}\}$ $_{i}^{(0)}\}_{i=1}^{n_{0}}$ given $\{\boldsymbol{x}_i^{(0)}$ $\{e^{(0)}\}_{i=1}^{n_0}$ as $\mathbb{P}_{y|x;\boldsymbol{\beta}}^{(0)\otimes n_0}$ $y|x;\beta$ and the distribution of $\{x_i^{(0)}\}$ $\{\widehat{\theta}^{(0)}\}_{i=1}^{n_0}$ as $\mathbb{P}_{\bm{x}}^{(0)\otimes n_0}$. For any $\bm{\theta}^{(0)} \neq \widetilde{\bm{\theta}}^{(0)} \in \mathcal{F}$:

$$
\begin{split}\n\text{KL}\left(\prod_{t=1}^{T} \mathbb{P}_{y|\mathbf{x};\mathbf{A}\theta^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(t)\otimes n} \cdot \mathbb{P}_{y|\mathbf{x};\mathbf{A}\theta^{(0)}}^{(0)\otimes n_0} \cdot \mathbb{P}_{\mathbf{x}}^{(0)\otimes n_0} \middle\| \prod_{t=1}^{T} \mathbb{P}_{y|\mathbf{x};\mathbf{A}\theta^{(t)}}^{(t)\otimes n} \cdot \mathbb{P}_{\mathbf{x}}^{(t)\otimes n} \cdot \mathbb{P}_{y|\mathbf{x};\mathbf{A}\theta^{(0)}}^{(0)\otimes n_0} \cdot \mathbb{P}_{\mathbf{x}}^{(0)\otimes n_0}\right) \\
\text{KL}\left(\mathbb{P}_{y|\mathbf{x};\mathbf{A}\theta^{(0)}}^{(0)\otimes n_0} \cdot \mathbb{P}_{\mathbf{x}}^{(0)\otimes n_0} \middle\| \mathbb{P}_{y|\mathbf{x};\mathbf{A}\tilde{\theta}^{(0)}}^{(0)\otimes n_0} \cdot \mathbb{P}_{\mathbf{x}}^{(0)\otimes n_0}\right) \\
&\lesssim n_0 \| \mathbf{A}\theta^{(0)} - \mathbf{A}\tilde{\theta}^{(0)} \|_2^2 \\
&\lesssim r \\
&\leq c \log |\mathcal{T}|,\n\end{split}
$$

where c is a small constant. Finally, applying Fano's Lemma (Lemma [24\)](#page-48-1), we have

$$
\inf_{\widehat{\boldsymbol{\beta}}^{(0)} \|\boldsymbol{\theta}^{(0)}\|_2 \leq \delta} \mathbb{P}\left(\|\widehat{\boldsymbol{\beta}}^{(0)} - \boldsymbol{A}\boldsymbol{\theta}^{(0)}\|_2 \geq c\delta\right) \geq 1 - \frac{\log 2}{\log |\mathcal{T}|} - c \geq \frac{1}{10}.
$$

(iv) We follow a similar analysis in part (v) of the proof of Theorem [8.](#page-19-0) Consider the case $h = 0$. $\text{Fix all } \boldsymbol{A}^{(t)} = \text{some } \boldsymbol{A} \in \mathcal{O}^{p \times r} \text{ and } \{ \boldsymbol{\theta}^{(t)} \}_{t=1}^T \subseteq \{ \boldsymbol{e}_j \}_{j=1}^r \subseteq \mathbb{R}^r \text{ s.t. } \sigma_r(T^{-1/2} \{ \boldsymbol{\theta}^{(t)} \}_{t=1}^T) \geq \frac{c}{\sqrt{r}}$ with $\#\{t : \theta^{(t)} = e_1\} = \lfloor T/r \rfloor$. Without loss of generality, suppose $\theta^{(t)} = e_1$ when $t \in \{0\} \cup [\lfloor T/r \rfloor]$. Denote $\widetilde{\boldsymbol{\beta}} = Ae_1$, and $\boldsymbol{\beta}^{(t)} = A\boldsymbol{\theta}^{(t)}$ for $t \geq \lfloor T/r \rfloor + 1$. Consider two data generating mechanisms in Lemma [27:](#page-48-3)

(I) $\{(\bm{x}_i^{(t)}\)}$ $\binom{t}{i}, y_i^{(t)}$ $\{(\epsilon^t)\}_{i=1}^n \sim (1-\epsilon') (\mathbb{P}_{\epsilon^0 \mid \mathbf{x}^n}^{\otimes n})$ $y|\bm{x};\beta$ $\cdot \mathbb{P}_{\mathbf{x}}^{\otimes n}$ + $\epsilon' \mathbb{Q}$ independently for $t \in [[T/r]],$ where $\epsilon' =$ $T\epsilon$ $\frac{T\epsilon}{50|T/r|},\, \{(\boldsymbol{x}_i^{(0)}% ,\boldsymbol{x}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\boldsymbol{\hat{x}}_i^{(0)}\}$ $\binom{0}{i}, y_i^{(0)}$ $\{a_i^{(0)})\}_{i=1}^{n_0}$, and $\{\{\boldsymbol{x}_i^{(t)}\}$ $\{ \widehat{u}^{(t)}_i \}_{i=1}^n \}_{t=|T/r|+1}^T \sim \mathbb{D} = \prod_{t= \lfloor T/r \rfloor +1}^T (\mathbb{P}_{y \mid \bm{x}}^{\otimes n})$ $\frac{\otimes n}{y|\bm{x};\bm{\beta}^{(t)}}\cdot \mathbb{P}_{\bm{x}}^{\otimes n});$ (II) With a preserved set $S^c \subseteq [\lfloor T/r \rfloor]$, generate $\{x^{(t)}\}_{t \in S^c} \sim \mathbb{Q}_{S^c}$ and $\{(x_i^{(t)}\}_{t \in S^c})$ $\displaystyle{ _{i}^{(t)},y_{i}^{(t)}}$ $\{i^{(t)}\}\}_{i=1}^n \sim$ $\mathbb{P}^{\otimes n}$ $y|\bm{x};\!\bm{\beta}$ $\cdot \mathbb{P}_{\bm{x}}^{\otimes n}$ independently for $t \in \{0\} \cup (S \cap [[T/r]])$, and $\{\{\bm{x}_i^{(t)}\}$ $\{e^{(t)}\}_{i=1}^{n} \}_{t=|T/r|+1}^{T} \sim$ $\mathbb{D}\{\{\boldsymbol{x}_i^{(t)}% =\boldsymbol{x}_i^{(t)}\}}$ $\{ \widehat{u}^{(t)}_i \}_{i=1}^n \}_{t=|T/r|+1}^T \sim \mathbb{D} = \prod_{t= \lfloor T/r \rfloor +1}^T (\mathbb{P}_{y \mid \bm{x}}^{\otimes n})$ $\frac{\otimes n}{y|\boldsymbol{x};\boldsymbol{\beta}^{(t)}}\cdot \mathbb{P}_{\boldsymbol{x}}^{\otimes n}$).

Denote the joint distributions of $\{\boldsymbol{x}_i^{(t)}\}$ (^{t)}_i</sub> $\{e\}\$ _{t∈{0}}∪[[*T*/r]] in (I) and (II) as $\mathbb{P}_{(\epsilon,\theta,\mathbb{Q})}$ and $\mathbb{P}_{(S,\theta,\mathbb{Q}_{S^c})}$, respectively.

Note that by Lemma [25,](#page-48-0)

$$
\varpi(\epsilon', \Theta)
$$
\n
$$
\begin{split}\n&= \sup \{ \|\beta_1 - \beta_2\|_2 : TV\left(\mathbb{P}_{y|x;\beta_1}^{\otimes n} \cdot \mathbb{P}_{x}^{\otimes n}, \mathbb{P}_{y|x;\beta_2}^{\otimes n} \cdot \mathbb{P}_{x}^{\otimes n}\right) \leq \epsilon'/(1-\epsilon'), \\
&TV\left(\mathbb{P}_{y|x;\beta_1}^{\otimes n_0} \cdot \mathbb{P}_{x}^{\otimes n}, \mathbb{P}_{y|x;\beta_2}^{\otimes n_0} \cdot \mathbb{P}_{x}^{\otimes n}\right) \leq 1/20\} \\
&\geq \sup \{ \|\beta_1 - \beta_2\|_2 : KL\left(\mathbb{P}_{y|x;\beta_1}^{\otimes n} \cdot \mathbb{P}_{x}^{\otimes n}, \mathbb{P}_{y|x;\beta_2}^{\otimes n} \cdot \mathbb{P}_{x}^{\otimes n}\right) \leq 2[\epsilon'/(1-\epsilon')]^2, \\
& KL\left(\mathbb{P}_{y|x;\beta_1}^{\otimes n} \cdot \mathbb{P}_{x}^{\otimes n}, \mathbb{P}_{y|x;\beta_2}^{\otimes n}\cdot \mathbb{P}_{x}^{\otimes n}\right) \leq 1/200\} \\
&\geq \sup \{ \|\beta_1 - \beta_2\|_2 : n\|\beta_1 - \beta_2\|_2^2 \leq c[\epsilon'/(1-\epsilon')]^2, n_0\|\beta_1 - \beta_2\|_2^2 \leq c\} \\
&= c' \frac{\epsilon'}{\sqrt{n}} \wedge \frac{1}{\sqrt{n_0}} \\
&\geq \frac{r\epsilon}{\sqrt{n}} \wedge \frac{1}{\sqrt{n_0}}.\n\end{split}
$$

Then by Lemma [28,](#page-49-0)

$$
\inf_{\widehat{\boldsymbol{\beta}}^{(0)}} \sup_{\widetilde{\boldsymbol{\beta}},\mathbb{Q}} (\mathbb{P}_{(\epsilon,\theta,\mathbb{Q})} \cdot \mathbb{D}) \bigg(\| \widehat{\boldsymbol{\beta}}^{(0)} - \boldsymbol{\beta}^{(0)*} \|_2 \geq \varpi(\epsilon',\Theta) \bigg) \geq \frac{1}{2}
$$

.

Therefore, by Lemma [29,](#page-49-1) it follows that

$$
\inf_{\hat{\beta}^{(0)}} \sup_{S \subseteq S} \sup_{\{\beta^{(t)}\}_{t \in \{0\} \cup S} \in \mathcal{B}(S,0)} \mathbb{P}\left(\|\hat{\beta}^{(0)} - \beta^{(0)*}\|_2 \geq c \frac{\epsilon r}{\sqrt{n}} \wedge \frac{1}{\sqrt{n_0}}\right)
$$
\n
$$
\geq \inf_{\hat{\beta}^{(0)}} \sup_{S : |S| \geq T(1-\epsilon)} \sup_{\{\beta^{(t)}\}_{t \in \{0\} \cup S} \in \mathcal{B}(S,0)} (\mathbb{P}_{(S,\theta,\mathbb{Q}_{S^c})} \cdot \mathbb{D})\left(\|\hat{\beta}^{(0)} - \beta^{(0)*}\|_2 \geq \varpi(\epsilon/50,\Theta)\right)
$$
\n
$$
\geq \frac{1}{10}.
$$

D.12 Proofs of Lemmas and Propositions

Denote $\widehat{\mathbf{\Sigma}}^{(t)} = \sum_{i=1}^{n} \frac{1}{n}$ $\frac{1}{n}\boldsymbol{x}_{i}^{\left(t\right)}$ $\binom{t}{i}(\bm{x}^{(t)}_i$ $\binom{(t)}{i}$ ^T.

D.12.1 PROOF OF LEMMA [30](#page-50-0)

By optimality of $\boldsymbol{\theta}_A^{(t)}$ and $\boldsymbol{\theta}_A^{(t)}$ $\overset{(t)}{A^{(t)*}},$ $\boldsymbol{A}^\top \nabla f^{(t)}(\boldsymbol{A}\boldsymbol{\theta}^{(t)}_{\boldsymbol{A}}) = (\boldsymbol{A}^{(t)*})^\top \nabla f^{(t)}(\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)}_{\boldsymbol{A}^{(t)}}$ $A^{(t)}_{(t)}(31) = 0_r,$ (31) which entails

$$
\begin{aligned}\n\boldsymbol{A}^{\top} \left[\nabla f^{(t)} (\boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}) + \widehat{\boldsymbol{\Sigma}}^{(t)} (\boldsymbol{A} \boldsymbol{\theta}_{\boldsymbol{A}}^{(t)} - \boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}) \right] &= \mathbf{0}, \\
\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)} = (\boldsymbol{A}^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{A})^{-1} \boldsymbol{A}^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*} - (\boldsymbol{A}^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{A})^{-1} \boldsymbol{A}^{\top} \nabla f^{(t)} (\boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}).\n\end{aligned}
$$

Therefore,

$$
\begin{aligned}\n\|A\boldsymbol{\theta}_A^{(t)} - A^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 &\leq \underbrace{\|[I-A(A^\top\widehat{\boldsymbol{\Sigma}}^{(t)}A)^{-1}A^\top\widehat{\boldsymbol{\Sigma}}^{(t)}]A^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2}_{[1]} \\
&\quad + \underbrace{\|A(A^\top\widehat{\boldsymbol{\Sigma}}^{(t)}A)^{-1}A^\top\nabla f^{(t)}(A^{(t)*}\boldsymbol{\theta}^{(t)*})\|_2}_{[2]}\n\end{aligned}.
$$

Note that

$$
[1] \lesssim ||[I - A(A^{\top} \hat{\Sigma}^{(t)} A)^{-1} A^{\top} \hat{\Sigma}^{(t)}] A^{(t)*} ||_2 ||\theta^{(t)*}||_2
$$

\n
$$
\lesssim ||(\hat{\Sigma}^{(t)})^{-1/2}||_2 \cdot ||(\hat{\Sigma}^{(t)})^{1/2} A^{(t)*} - (\hat{\Sigma}^{(t)})^{1/2} A (A^{\top} \hat{\Sigma}^{(t)} A)^{-1} A^{\top} (\hat{\Sigma}^{(t)})^{1/2} (\hat{\Sigma}^{(t)})^{1/2} A^{(t)*} ||_2 ||\theta^{(t)*}||_2
$$

\n
$$
\lesssim ||[I - B(B^{\top} B)^{-1} B^{\top}] B^{(t)*} ||_2 ||\theta^{(t)*}||_2
$$

\n
$$
\lesssim ||P_{\mathbf{B}^{\perp}} B^{(t)*} ||_2 ||\theta^{(t)*} ||_2,
$$
\n(32)

w.p. at least $1 - e^{-Cp}$, where $B = (\hat{\Sigma}^{(t)})^{1/2} A$, $B^{(t)*} = (\hat{\Sigma}^{(t)})^{1/2} A^{(t)*}$, $A^{\perp} \in {\{\tilde{A} \in {\mathbb{R}} \}^{n \times n}}$ $(\mathcal{O}^{p\times (p-r)}) : \tilde{A}\tilde{A}^\top + AA^\top = I_p\}, \, B^\perp = (\widehat{\Sigma}^{(t)})^{-1/2}A^\perp, \, P_{\boldsymbol{B}^\perp} = B^\perp((B^\perp)^\top B^\perp)^{-1}(B^\perp)^\top = \ \mathcal{O}^{(1)}$ $I - P_B$. Further, notice that

$$
||P_{\mathbf{B}^{\perp}} \mathbf{B}^{(t)*}||_2 = ||(\widehat{\mathbf{\Sigma}}^{(t)})^{-1/2} \mathbf{A}^{\perp} [(\mathbf{A}^{\perp})^{\top} (\widehat{\mathbf{\Sigma}}^{(t)})^{-1} \mathbf{A}^{\perp}]^{-1} (\mathbf{A}^{\perp})^{\top} \mathbf{A}^{(t)*}||_2
$$

\n
$$
\leq ||(\widehat{\mathbf{\Sigma}}^{(t)})^{-1/2}||_2 \cdot \lambda_{\min}^{-1} ((\mathbf{A}^{\perp})^{\top} (\widehat{\mathbf{\Sigma}}^{(t)})^{-1} \mathbf{A}^{\perp}) \cdot ||(\mathbf{A}^{\perp})^{\top} \mathbf{A}^{(t)*}||_2
$$

\n
$$
\lesssim ||(\mathbf{A}^{\perp})^{\top} \mathbf{A}^{(t)*}||_2
$$

\n
$$
\lesssim ||\mathbf{A} \mathbf{A}^{\top} - \mathbf{A}^{(t)*} (\mathbf{A}^{(t)*})^{\top} ||_2,
$$
\n(33)

w.p. at least $1 - e^{-C(p + \log T)}$. And

$$
[2] \leq ||A(A^{\top} \hat{\Sigma}^{(t)} A)^{-1} A^{\top} \nabla f^{(t)} (A^{(t)*} \theta_{A^{(t)*}}^{(t)})||_2 + ||A(A^{\top} \hat{\Sigma}^{(t)} A)^{-1} A^{\top} [\nabla f^{(t)} (A^{(t)*} \theta_{A^{(t)*}}^{(t)}) - \nabla f^{(t)} (A^{(t)*} \theta_{A^{(t)*}}^{(t)})]||_2 \n\lesssim ||A^{\top} (A^{(t)*})^{\perp}||_2 \cdot ||\nabla f^{(t)} (A^{(t)*} \theta_{A^{(t)*}}^{(t)})||_2 + ||\theta^{(t)*} - \theta_{A^{(t)*}}^{(t)}||_2 \n\lesssim ||A A^{\top} - A^{(t)*} (A^{(t)*})^{\top} ||_2 (||\nabla f^{(t)} (A^{(t)*} \theta_{A^{(t)*}}^{(t)})||_2 + C ||\theta^{(t)*} - \theta_{A^{(t)*}}^{(t)} ||_2) + ||\theta^{(t)*} - \theta_{A^{(t)*}}^{(t)} ||_2
$$
\n(34)

$$
\lesssim \|\boldsymbol{A}\boldsymbol{A}^{\top} - \boldsymbol{A}^{(t)*}(\boldsymbol{A}^{(t)*})^{\top}\|_2 \sqrt{\frac{p + \log T}{n}} + \sqrt{\frac{r + \log T}{n}},
$$
\n(35)

j. This step holds because of Lemma [19,](#page-47-2) which shows that $\lambda_{\min}(\hat{\Sigma}^{(t)}) > 0$ (so that $\hat{\Sigma}^{(t)}$ is invertible) w.p. at least $1 - e^{-C(p + \log T)}$.

w.p. at least $1 - e^{-C(r + \log T)}$, where we used the fact that $\|\boldsymbol{\theta}^{(t)*} - \boldsymbol{\theta}_{\mathbf{\mathbf{\mathbf{\mathbf{A}}}^{(t)*}}}^{(t)} \|_2 \lesssim \sqrt{\frac{r + \log T}{n}}$ $\frac{\log T}{n}$ for all $t \in S$ w.p. at least $1 - e^{-C(r + \log T)}$ (which is the standard rate of r-dimensional linear regression). Inequality [\(34\)](#page-88-1) holds because there exists $a \in \mathbb{R}^r$ such that $\nabla f^{(t)}(A^{(t)*}\theta^{(t)*}) =$ $(A^{(t)*})^{\perp}a$ (due to [\(31\)](#page-87-0)) and $||a||_2 = ||\nabla f^{(t)}(A^{(t)*}\theta^{(t)*})||_2 \leq \sqrt{\frac{p + \log T}{n}}$ w.p. at least 1 – $e^{C(p+\log T)}$. Putting [\(32\)](#page-88-2), [\(33\)](#page-88-3), and [\(35\)](#page-88-4) together, we complete the proof.

D.12.2 PROOF OF LEMMA [31](#page-51-0)

By Lemma [30,](#page-50-0) w.p. at least $1 - e^{-C(r + \log T)}$,

$$
\|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)}\|_{2} - \|\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2} \leq \|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2} \cdot \zeta^{(t)}
$$

$$
\lesssim \|\mathbf{A}\mathbf{A}^{\top} - \mathbf{A}^{(t)*}(\mathbf{A}^{(t)*})^{\top}\|_{2} \cdot \zeta^{(t)} + \sqrt{\frac{r + \log T}{n}}
$$

$$
\leq C'\zeta^{(t)}.
$$

Note that $\|\mathbf{A}\bm{\theta}^{(t)}_{\mathbf{A}}\|_2 = \|\bm{\theta}^{(t)}_{\mathbf{A}}\|_2$ and $\|\mathbf{A}^{(t)*}\bm{\theta}^{(t)*}\|_2 = \|\bm{\theta}^{(t)*}\|_2 \leq \zeta^{(t)}.$

D.12.3 PROOF OF LEMMA [32](#page-51-1)

The proof is almost the same as the proof of Lemma [30,](#page-50-0) thus omitted.

D.12.4 PROOF OF LEMMA [33](#page-51-2)

Denote $\mathbf{G} = \left\{ \nabla f^{(t)}(\mathbf{A}^{(t)*}\theta^{(t)*}) \right\}_{t \in S}$ and $\Delta^{(t)} = \overline{\mathbf{A}}\theta_{\overline{\mathbf{A}}}^{(t)} - \mathbf{A}^{(t)*}\theta^{(t)*}$. By Lemma [30,](#page-50-0) w.p. at least $1 - e^{-C(r + \log T)}$, for all $t \in S$,

$$
\|\Delta^{(t)}\|_2 \lesssim h\zeta^{(t)} + \sqrt{\frac{r + \log T}{n}},
$$

leading to

$$
\|\{\boldsymbol{\Delta}^{(t)}\}_{t\in S}\|_{2} = \sup_{\|\boldsymbol{u}\|_{2}=\|\boldsymbol{v}\|_{2}=1} \left| \sum_{t\in S} u_{t}(\boldsymbol{\Delta}^{(t)})^{\top} \boldsymbol{v} \right|
$$

$$
\leq \sup_{\|\boldsymbol{u}\|_{2}=\|\boldsymbol{v}\|_{2}=1} \|\boldsymbol{u}\|_{2} \cdot \sqrt{\sum_{t\in S} ((\boldsymbol{\Delta}^{(t)})^{\top} \boldsymbol{v})^{2}}
$$

$$
\lesssim \sqrt{T} \left(h \bar{\zeta} + \sqrt{\frac{r + \log T}{n}} \right).
$$
 (36)

By Lemma D.3 in [Duan and Wang](#page-36-0) [\(2022\)](#page-36-0) or Theorem 5.39 in [Vershynin](#page-39-1) [\(2010\)](#page-39-1), w.p. at least $1 - e^{-C(p + \log T)}$,

$$
\|\boldsymbol{G}\|_2 \lesssim \sqrt{\frac{p+T}{n}}.\tag{37}
$$

Combining [\(36\)](#page-89-0), [\(37\)](#page-89-1), and the fact that $\|\overline{G}\|_2 \leq \|G\|_2 + \|\{\Delta^{(t)}\}_{t\in S}\|_2$ implies the desired result.

D.12.5 Proof of Lemma [34](#page-51-3)

Note that

$$
\begin{aligned}\n\|\overline{G} - G\|_2 &= \sup_{\|\boldsymbol{u}\|_2 = \|\boldsymbol{v}\|_2 = 1} \left| \sum_{j=1}^d u_j (\overline{g}_j - g_j)^\top \boldsymbol{v} \right| \\
&\leq \sup_{\|\boldsymbol{u}\|_2 = \|\boldsymbol{v}\|_2 = 1} \|\boldsymbol{u}\|_2 \cdot \sqrt{\sum_{j=1}^d [(\overline{g}_j - g_j)^\top \boldsymbol{v}]^2} \\
&\leq \sqrt{d}\Delta,\n\end{aligned}
$$

which completes the proof.

D.12.6 PROOF OF LEMMA [35](#page-51-4)

By convexity of $f^{(t)}$,

$$
f^{(t)}(\mathbf{A}^{(t)}\boldsymbol{\theta}_{\mathbf{A}^{(t)}}^{(t)}) - f^{(t)}(\widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)}) + \frac{\lambda}{\sqrt{n}} \|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^\top\|_2
$$

\n
$$
\geq \frac{\lambda}{\sqrt{n}} \|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^\top\|_2 + \nabla f^{(t)}(\widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)})^\top (\mathbf{A}^{(t)}\boldsymbol{\theta}_{\mathbf{A}^{(t)}}^{(t)} - \widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)})
$$

\n
$$
\geq \frac{\lambda}{\sqrt{n}} \|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^\top\|_2 - \|\nabla f^{(t)}(\widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)})^\top \mathbf{A}^{(t)}\boldsymbol{\theta}_{\mathbf{A}^{(t)}}^{(t)}\|_2
$$
\n(38)

$$
\geq \frac{\lambda}{\sqrt{n}} \|A^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top \|_2 - \|\nabla f^{(t)}(\widehat{\overline{A}}\boldsymbol{\theta}_{\widehat{\overline{A}}}^{(t)})\|_2 \cdot \|((\widehat{\overline{A}})^\perp)^\top \mathbf{A}^{(t)}\|_2 \cdot \|\boldsymbol{\theta}_{\mathbf{A}^{(t)}}^{(t)}\|_2 \tag{39}
$$

$$
\geq \left(\frac{\lambda}{\sqrt{n}} - C\|\nabla f^{(t)}(\widehat{\overline{A}}\theta_{\widehat{\overline{A}}}^{(t)})\|_2 \cdot \zeta^{(t)}\right) \cdot \|A^{(t)}(A^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_2
$$
\n
$$
> 0,
$$
\n(40)

when $\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top} \neq \hat{\vec{A}}(\hat{\vec{A}})^{\top}$, where [\(38\)](#page-90-0) and [\(39\)](#page-90-1) are due to the definition of $\theta_{\hat{\vec{A}}}^{(t)}$ (which $\mathcal{A}_{\mathcal{A}}$ leads to $\nabla f^{(t)}(\widehat{\overline{A}} \theta_{\widehat{\overline{A}}}^{(t)})$ $(\widehat{\overline{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)})=(\widehat{\overline{A}})^{\perp}\boldsymbol{u}$ with $\|\boldsymbol{u}\|_2=\|f^{(t)}(\widehat{\overline{A}}\boldsymbol{\theta}_{\widehat{\overline{A}}}^{(t)})\|_2$, and [\(40\)](#page-90-2) is due to Lemma [21.](#page-47-0) Note that the above strict inequality leads to a contradiction with the fact that $(\mathbf{A}^{(t)}, \mathbf{\theta}^{(t)})$ is a minimizer, therefore we must have $\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top} \neq \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}$ for all $t \in S$.

D.12.7 PROOF OF LEMMA [36](#page-51-5)

It is straightforward to see that

$$
\frac{1}{|S'|} \sum_{t \in S'} \theta^{(t)*} (\theta^{(t)*})^{\top} = \frac{|S|}{|S'|} \left(\frac{1}{|S|} \sum_{t \in S} \theta^{(t)*} (\theta^{(t)*})^{\top} - \frac{1}{|S|} \sum_{t \in S \setminus S'} \theta^{(t)*} (\theta^{(t)*})^{\top} \right)
$$
\n
$$
\succeq \frac{|S|}{|S'|} \left(\frac{c}{r} \overline{\zeta}^2 - \alpha \max_{t \in S} (\zeta^{(t)})^2 \right) \boldsymbol{I}_r
$$
\n
$$
\succeq \frac{c}{2r} \overline{\zeta}^2,
$$

because $|S|/|S'| \ge 1$ and $\alpha \le c/r \cdot \bar{\zeta}^2/[2 \max_{t \in S} (\zeta^{(t)})^2]$ with a very small $c'' > 0$.

D.12.8 PROOF OF LEMMA [37](#page-51-6)

By definition, we have

$$
0 \leq f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}) - f^{(t)}(\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}) + \frac{\lambda}{\sqrt{n}}\|\mathbf{A}^{(t)*}(\mathbf{A}^{(t)*})^\top - \widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^\top\|_2 - \frac{\lambda}{\sqrt{n}}\|\widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^\top - \widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^\top\|_2,
$$

which implies

$$
\frac{\lambda}{\sqrt{n}}\|\mathbf{A}^{(t)*}(\mathbf{A}^{(t)*})^{\top} - \widehat{\mathbf{A}}^{(t)}(\widehat{\mathbf{A}}^{(t)})^{\top}\|_{2} \geq f^{(t)}(\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)}) - f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})
$$
\n
$$
\geq \frac{1}{2}(\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})^{\top}\widehat{\boldsymbol{\Sigma}}^{(t)}(\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})
$$
\n
$$
-\|\nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_{2} \cdot \|\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2}
$$
\n
$$
\geq c\|\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2}^{2} - C\sqrt{\frac{p + \log T}{n}}\|\widehat{\mathbf{A}}^{(t)}\widehat{\boldsymbol{\theta}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2},
$$

w.p. at least $1 - e^{-C'(p + \log T)}$. Note that $||A^{(t)*}(A^{(t)*})^{\top} - \hat{A}^{(t)}(\hat{A}^{(t)})^{\top}||_2 \leq 1$. Then a simplification of above inequality entails the desired result.

D.12.9 PROOF OF LEMMA [38](#page-52-4)

The proof idea is very similar to the proof of Lemma [30.](#page-50-0) By the optimality condition, $\boldsymbol{A}^{\top} \nabla f^{(t)}(\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}) = 0$, implying that

$$
\begin{aligned}\n\boldsymbol{A}^{\top} \left[\nabla f^{(t)} (\boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}) + \widehat{\boldsymbol{\Sigma}}^{(t)} (\boldsymbol{A} \boldsymbol{\theta}_{\boldsymbol{A}}^{(t)} - \boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}) \right] &= \mathbf{0}, \\
\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)} = (\boldsymbol{A}^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{A})^{-1} \boldsymbol{A}^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*} - (\boldsymbol{A}^{\top} \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{A})^{-1} \boldsymbol{A}^{\top} \nabla f^{(t)} (\boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}).\n\end{aligned}
$$

Note that $\boldsymbol{\theta}^{(t)*} \in \mathbb{R}$ and $\boldsymbol{A}^{(t)*} \in \mathbb{R}^p$ when $r = 1$. Hence, w.p. at least $1 - e^{-C'(p + \log T)}$, we have

$$
\|A\theta_A^{(A)} - A^{(t)}\theta^{(t)*}\|_2
$$

\n
$$
\geq \| [I - A(A^{\top}\hat{\Sigma}^{(t)}A)^{-1}A^{\top}\hat{\Sigma}^{(t)}]A^{(t)*}\theta^{(t)*} \|_2 - \|A(A^{\top}\hat{\Sigma}^{(t)}A)^{-1}A^{\top}\nabla f^{(t)}(A^{(t)*}\theta^{(t)*}) \|_2
$$

\n
$$
\geq \| [I - A(A^{\top}\hat{\Sigma}^{(t)}A)^{-1}A^{\top}\hat{\Sigma}^{(t)}]A^{(t)*} \|_2 \zeta^{(t)} - \|A\|_2 \cdot \|A(A^{\top}\hat{\Sigma}^{(t)}A)^{-1} \|_2 \cdot \|A\|_2 \cdot \|\nabla f^{(t)}(A^{(t)*}\theta^{(t)*}) \|_2
$$

\n
$$
\geq \zeta^{(t)}\lambda_{\min}((\hat{\Sigma}^{(t)})^{-1/2}) \cdot \|(\hat{\Sigma}^{(t)})^{1/2}A^{(t)*} - (\hat{\Sigma}^{(t)})^{1/2}A(A^{\top}\hat{\Sigma}^{(t)}A)^{-1}A^{\top}\hat{\Sigma}^{(t)}A^{(t)*} \|_2 - C\sqrt{\frac{p + \log T}{n}}
$$

$$
\geq C' \zeta^{(t)} \| [\boldsymbol{I} - \boldsymbol{B} (\boldsymbol{B}^\top \boldsymbol{B})^{-1}) \boldsymbol{B}^\top] \boldsymbol{B}^{(t)*} \|_2 - C \sqrt{\frac{p + \log T}{n}}, \tag{41}
$$

where $\mathbf{B} = (\widehat{\boldsymbol{\Sigma}}^{(t)})^{1/2} \mathbf{A}, \, \mathbf{B}^{(t)*} = (\widehat{\boldsymbol{\Sigma}}^{(t)})^{1/2} \mathbf{A}^{(t)*}, \, \mathbf{A}^{\perp} \in \{\tilde{\mathbf{A}} \in \mathcal{O}^{p \times (p-1)}\}: \tilde{\mathbf{A}} \tilde{\mathbf{A}}^{\top} + \mathbf{A} \mathbf{A}^{\top} = \mathbf{I}_p\},$ $B^{\perp} = (\hat{\Sigma}^{(t)})^{-1/2} A^{\perp}, P_{B^{\perp}} = B^{\perp}((B^{\perp})^{\top} B^{\perp})^{-1}(B^{\perp})^{\top} = I - P_{B}.$ Further, notice that $||P_{\boldsymbol{B}^{\perp}} \boldsymbol{B}^{(t)*}||_2 = ||(\widehat{\boldsymbol{\Sigma}}^{(t)})^{-1/2} \boldsymbol{A}^{\perp} [(\boldsymbol{A}^{\perp})^{\top} (\widehat{\boldsymbol{\Sigma}}^{(t)})^{-1} \boldsymbol{A}^{\perp}]^{-1} (\boldsymbol{A}^{\perp})^{\top} \boldsymbol{A}^{(t)*}||_2$

$$
\geq \lambda_{\min}^{1/2}([(A^{\perp})^{\top}(\widehat{\Sigma}^{(t)})^{-1}A^{\perp}]^{-1}) \cdot ||(A^{\perp})^{\top}A^{(t)*}||_2
$$

=
$$
||(A^{\perp})^{\top}(\widehat{\Sigma}^{(t)})^{-1}A^{\perp}||_2^{-1/2} \cdot ||(A^{\perp})^{\top}A^{(t)*}||_2
$$

$$
\gtrsim ||AA^{\top} - A^{(t)*}(A^{(t)*})^{\top}||_2,
$$
 (42)

w.p. at least $1 - e^{-C(p + \log T)}$. Combining [\(41\)](#page-91-0) and [\(42\)](#page-92-0) completes the proof.

D.12.10 PROOF OF LEMMA [39](#page-52-0)

First, by Lemma [19,](#page-47-2) $0 < c \leq \min_{t \in S} \lambda_{\min}(\hat{\Sigma}^{(t)}) \leq \max_{t \in S} \lambda_{\max}(\hat{\Sigma}^{(t)}) \leq C < \infty$ w.p. at least $1 - e^{-C'(p + \log T)}$. Then (i) is the direct consequence of Lemmas E.3 in [Duan and Wang](#page-36-0) [\(2022\)](#page-36-0). (ii) and (iii) come from Lemma E.2 in [Duan and Wang](#page-36-0) [\(2022\)](#page-36-0).

D.12.11 PROOF OF LEMMA [40](#page-52-1)

We modify a few steps in the proof of Lemma [30.](#page-50-0) The key point here is that $A \in \mathcal{O}^{p \times r}$ is fixed.

By optimality of $\theta^{(t)}_A$ and $\theta^{(t)}_{A^{(t)}}$ $\overset{(t)}{A^{(t)*}},$

$$
\mathbf{A}^{\top} \nabla f^{(t)}(\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)}) = (\mathbf{A}^{(t)*})^{\top} \nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}_{\mathbf{A}^{(t)*}}^{(t)}) = \mathbf{0}_r,
$$

which entails

$$
\mathbf{A}^{\top} \left[\nabla f^{(t)} (\mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*}) + \hat{\mathbf{\Sigma}}^{(t)} (\mathbf{A} \boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*}) \right] = \mathbf{0},
$$

$$
\boldsymbol{\theta}_{\mathbf{A}}^{(t)} = (\mathbf{A}^{\top} \hat{\mathbf{\Sigma}}^{(t)} \mathbf{A})^{-1} \mathbf{A}^{\top} \hat{\mathbf{\Sigma}}^{(t)} \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} - (\mathbf{A}^{\top} \hat{\mathbf{\Sigma}}^{(t)} \mathbf{A})^{-1} \mathbf{A}^{\top} \nabla f^{(t)} (\mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*}).
$$

Therefore,

$$
\|\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}-\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2}\leq\underbrace{\|[\boldsymbol{I}-\boldsymbol{A}(\boldsymbol{A}^{\top}\widehat{\boldsymbol{\Sigma}}^{(t)}\boldsymbol{A})^{-1}\boldsymbol{A}^{\top}\widehat{\boldsymbol{\Sigma}}^{(t)}]\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2}}_{[1]}+\underbrace{\|\boldsymbol{A}(\boldsymbol{A}^{\top}\widehat{\boldsymbol{\Sigma}}^{(t)}\boldsymbol{A})^{-1}\boldsymbol{A}^{\top}\nabla\boldsymbol{f}^{(t)}(\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_{2}}_{[2]}.
$$

Note that

$$
[1] \lesssim \|A^{(t)*} - A(A^{\top} \hat{\Sigma}^{(t)} A)^{-1} A^{\top} \hat{\Sigma}^{(t)} A^{(t)*} \|_{2}
$$

\n= \|A^{(t)*} - A(A^{\top} \hat{\Sigma}^{(t)} A)^{-1} A^{\top} \hat{\Sigma}^{(t)} (AA^{\top} + A^{\perp} (A^{\perp})^{\top}) A^{(t)*} \|_{2}
\n\leq \| (I - AA^{\top}) A^{(t)*} \|_{2} + \|A(A^{\top} \hat{\Sigma}^{(t)} A)^{-1} A^{\top} \hat{\Sigma}^{(t)} A^{\perp} (A^{\perp})^{\top} A^{(t)*} \|_{2}
\n\lesssim \| (A^{\perp})^{\top} A^{(t)*} \|_{2} + \| (A^{\top} \hat{\Sigma}^{(t)} A)^{-1} \|_{2} \cdot \| A^{\top} \hat{\Sigma}^{(t)} A^{\perp} \|_{2} \cdot \| (A^{\perp})^{\top} A^{(t)*} \|_{2}

By Lemma [20](#page-47-3) and the condition $n_0 \geq Cr$,

$$
\begin{aligned} \|(A^\top \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{A})^{-1}\|_2 &= \lambda_{\min}^{-1} (A^\top \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{A}) \\ &\leq [\lambda_{\min} (A^\top \boldsymbol{\Sigma}^{(t)} \boldsymbol{A}) - \|A^\top \widehat{\boldsymbol{\Sigma}}^{(t)} \boldsymbol{A} - A^\top \boldsymbol{\Sigma}^{(t)} \boldsymbol{A}\|_2]^{-1} \\ &\leq [\lambda_{\min} (A^\top \boldsymbol{\Sigma}^{(t)} \boldsymbol{A}) - C \sqrt{r/n_0}]^{-1} \end{aligned}
$$

 $\lesssim C'',$

w.p. at least $1 - e^{-C'(r + \log T)}$. And similarly

$$
\|\mathbf{A}^\top \widehat{\boldsymbol{\Sigma}}^{(t)} \mathbf{A}^\perp\|_2 \lesssim \|\mathbf{A}^\top \boldsymbol{\Sigma}^{(t)} \mathbf{A}^\perp\|_2 + \|\mathbf{A}^\top (\widehat{\boldsymbol{\Sigma}}^{(t)} - \boldsymbol{\Sigma}^{(t)}) \mathbf{A}^\perp\|_2 \lesssim C,
$$

w.p. at least $1 - e^{-C'(r + \log T)}$. Therefore by Lemma [21,](#page-47-0) w.p. at least $1 - e^{-C(r + \log T)}$,

$$
[1] \lesssim \|\boldsymbol{A}\boldsymbol{A}^{\top} - \boldsymbol{A}^{(t)*}(\boldsymbol{A}^{(t)*})^{\top}\|_2.
$$
 (43)

And w.p. at least $1 - e^{-C(r + \log T)}$,

$$
[2] \lesssim ||(\mathbf{A}^\top \widehat{\mathbf{\Sigma}}^{(t)} \mathbf{A})^{-1}||_2 \cdot ||\mathbf{A}^\top \nabla f^{(t)}(\mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*})||_2 \lesssim \sqrt{\frac{r}{n_0}},
$$
\n(44)

Denote $\{\beta_j\}_{j=1}^N$ as a 1/2-cover (whose components are inside the set to be covered) of $\mathscr{B} = \{ \boldsymbol{\beta} \in \mathbb{R}^p : \boldsymbol{\beta} = A \boldsymbol{u}, \|\boldsymbol{u}\|_2 \leq 1 \}$ (which is isomorphic to the unit ball in \mathbb{R}^r), with $N \leq 5^r$ (by Example 5.8 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0)). The second inequality in [\(44\)](#page-93-0) holds because

$$
\begin{split} \|\boldsymbol{A}^{\top} \nabla f^{(t)} (\boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*})\|_{2} &= \sup_{\|\boldsymbol{u}\|_{2} \leq 1} (\boldsymbol{A} \boldsymbol{u})^{\top} \left[\frac{2}{n_{0}} (\boldsymbol{X}^{(0)})^{\top} \boldsymbol{\epsilon}^{(0)} \right] \\ &= \sup_{\boldsymbol{\beta} \in \mathcal{B}} \boldsymbol{\beta}^{\top} \left[\frac{2}{n_{0}} (\boldsymbol{X}^{(0)})^{\top} \boldsymbol{\epsilon}^{(0)} \right] \\ &\leq \sup_{\boldsymbol{\beta} \in \mathcal{B}} (\boldsymbol{\beta} - \boldsymbol{\beta}_{j_{0}}^{\top})^{\top} \left[\frac{2}{n_{0}} (\boldsymbol{X}^{(0)})^{\top} \boldsymbol{\epsilon}^{(0)} \right] + \max_{j \in [N]} \boldsymbol{\beta}_{j}^{\top} \left[\frac{2}{n_{0}} (\boldsymbol{X}^{(0)})^{\top} \boldsymbol{\epsilon}^{(0)} \right] \\ &\leq \sup_{\boldsymbol{\beta} = \boldsymbol{A} \boldsymbol{u} \in \mathcal{B}} (\boldsymbol{u} - \boldsymbol{u}_{j_{0}})^{\top} \boldsymbol{A} \nabla f^{(t)} (\boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}) + \max_{j \in [N]} \boldsymbol{\beta}_{j}^{\top} \left[\frac{2}{n_{0}} (\boldsymbol{X}^{(0)})^{\top} \boldsymbol{\epsilon}^{(0)} \right] \\ &\leq \frac{1}{2} \|\boldsymbol{A} \nabla f^{(t)} (\boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*})\|_{2} + \max_{j \in [N]} \boldsymbol{\beta}_{j}^{\top} \left[\frac{2}{n_{0}} (\boldsymbol{X}^{(0)})^{\top} \boldsymbol{\epsilon}^{(0)} \right], \end{split}
$$

leading to

$$
\|\mathbf{A}^\top \nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_2 \leq 2 \max_{j \in [N]} \beta_j^\top \left[\frac{2}{n_0}(\mathbf{X}^{(0)})^\top \boldsymbol{\epsilon}^{(0)}\right],
$$

where $\beta_{j_0} = A u_{j_0}$ with some $\| u_{j_0}\|_2 \leq 1$ is the one which is closest to $\boldsymbol \beta = A \boldsymbol u$ in the cover (depending on $\boldsymbol{\beta}$). Hence, $\|\boldsymbol{u}_{j_0}-\boldsymbol{u}\|_2 = \|\boldsymbol{\beta}_{j_0}-\boldsymbol{\beta}\|_2 \leq 1/2$. Showing $\max_{j\in[N]}\boldsymbol{\beta}_j^\top\left[\frac{2}{n_0}\right]$ $\frac{2}{n_0}(\boldsymbol{X}^{(0)})^\top \boldsymbol{\epsilon}^{(0)} \big] \lesssim$ $\sqrt{r/n_0}$ w.p. at least $1 - e^{-C(r + \log T)}$ is standard.

Finally, combining [\(43\)](#page-93-1) and [\(44\)](#page-93-0), we complete the proof.

D.12.12 PROOF OF LEMMA [41](#page-52-5)

The proof is exactly the same as the proof of Lemma 30 , so omitted.

D.12.13 PROOF OF LEMMA 42

The proof is the same as the proof of Lemma [39,](#page-52-0) so we omit it here.

D.12.14 PROOF OF PROPOSITION [43](#page-52-2)

Define the random event \mathcal{E}_1 as the event that Lemmas [30,](#page-50-0) [31,](#page-51-0) [32,](#page-51-1) [33,](#page-51-2) and [35](#page-51-4) hold, hence $\mathbb{P}(\mathcal{E}_1) \geq 1 - Ce^{-(r+\log T)}$ with some constant $C > 0$. Define another event

$$
\mathcal{E}_2 = \bigg\{\max_{t \in [T]} \|\widehat{\mathbf{\Sigma}}^{(t)} - \mathbf{\Sigma}^{(t)}\|_2 \lesssim \sqrt{\frac{p + \log T}{n}}\bigg\}.
$$

By Lemma [19,](#page-47-2) we have $\mathbb{P}(\mathcal{E}_2) \geq 1 - Ce^{-(p+\log T)}$ with some constant $C > 0$. It suffices to prove the upper bounds in Proposition [43](#page-52-2) conditioned on $\mathcal{E}_1 \cap \mathcal{E}_2$, therefore we condition on $\mathcal{E}_1 \cap \mathcal{E}_2$ in the remaining proof without stating it explicitly and all the arguments are deterministic.

Define

$$
G(\boldsymbol{A}) = \sum_{t=1}^{T} \min_{\boldsymbol{A}^{(t)} \in \mathcal{O}^{p \times r}, \boldsymbol{\theta}^{(t)} \in \mathbb{R}^{r}} \left\{ \frac{1}{T} f^{(t)}(\boldsymbol{A}^{(t)} \boldsymbol{\theta}^{(t)}) + \frac{\sqrt{n}}{nT} \lambda \|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^{\top} - \boldsymbol{A} \boldsymbol{A}^{\top}\|_{2} \right\},
$$

\n
$$
G_{S}(\boldsymbol{A}) = \sum_{t \in S} \min_{\boldsymbol{A}^{(t)} \in \mathcal{O}^{p \times r}, \boldsymbol{\theta}^{(t)} \in \mathbb{R}^{r}} \left\{ \frac{1}{T} f^{(t)}(\boldsymbol{A}^{(t)} \boldsymbol{\theta}^{(t)}) + \frac{\sqrt{n}}{nT} \lambda \|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^{\top} - \boldsymbol{A} \boldsymbol{A}^{\top}\|_{2} \right\},
$$

\n
$$
G_{S}(\boldsymbol{A}) = \sum_{t \in S} \min_{\boldsymbol{A}^{(t)} \in \mathcal{O}^{p \times r}, \boldsymbol{\theta}^{(t)} \in \mathbb{R}^{r}} \left\{ \frac{1}{T} f^{(t)}(\boldsymbol{A}^{(t)} \boldsymbol{\theta}^{(t)}) + \frac{\sqrt{n}}{nT} \lambda \|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^{\top} - \boldsymbol{A} \boldsymbol{A}^{\top}\|_{2} \right\},
$$

\n
$$
\widehat{\overline{\boldsymbol{A}}} \in \argmin_{\boldsymbol{A} \in \mathcal{O}^{p \times r}} G(\boldsymbol{A}).
$$

Our first step is to prove the same bound for $\|\widehat{\vec{A}}\vec{\vec{A}}^\top - \vec{A}\vec{A}^\top \|_2$. Let's prove it by contradiction. Denote the desired rate as

$$
\eta = \bar{\zeta}^{-1} r \sqrt{\frac{p}{nT}} + \bar{\zeta}^{-1} \sqrt{r} \sqrt{\frac{r + \log T}{n}} + \sqrt{r}h + \frac{\lambda r}{\sqrt{n} \bar{\zeta}^2} \epsilon.
$$

Suppose $\|\hat{\overline{A}}\hat{\overline{A}}^{\top} - \overline{A}\overline{A}^{\top}\|_2 > C\eta$ with some constant $C > 0$.

Consider any minimizer $(A^{(t)}, \theta^{(t)}) = \arg \min_{\mathbf{A} \in \mathcal{O}^{p \times r}, \theta \in \mathbb{R}^r} \{\frac{1}{T}\}$ $\frac{1}{T}f^{(t)}(\bm{A\theta})+$ $\frac{\sqrt{n}}{nT} \lambda \| \bm{A} \bm{A}^\top \overline{A}(\overline{A})^{\top}$ ||2}. Define two index sets

$$
\mathcal{A}_1 = \left\{ t \in S : \| \mathbf{A}^{(t)} (\mathbf{A}^{(t)})^\top - \hat{\overline{\mathbf{A}}} (\hat{\overline{\mathbf{A}}})^\top \|_2 \ge c \| \hat{\overline{\mathbf{A}}} (\hat{\overline{\mathbf{A}}})^\top - \overline{\mathbf{A}} \overline{\mathbf{A}}^\top \|_2 \cdot \frac{1}{\sqrt{r}} \right\},
$$

$$
\mathcal{A}_2 = \left\{ t \in S : \| \mathbf{A}^{(t)} (\mathbf{A}^{(t)})^\top - \hat{\overline{\mathbf{A}}} (\hat{\overline{\mathbf{A}}})^\top \|_2 < c \| \hat{\overline{\mathbf{A}}} (\hat{\overline{\mathbf{A}}})^\top - \overline{\mathbf{A}} \overline{\mathbf{A}}^\top \|_2 \cdot \frac{1}{\sqrt{r}} \right\},
$$

where $c > 0$ is a small constant. Then we have

$$
\sum_{t \in \mathcal{A}_1} \frac{1}{T} [f^{(t)}(\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)}) - f^{(t)}(\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)})] + \sum_{t \in \mathcal{A}_1} \frac{\sqrt{n}}{nT} ||\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^\top ||_2
$$

$$
\geq -\frac{1}{T} \left| \left\langle \{\nabla f^{(t)}(\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)})\}_{t \in \mathcal{A}_1}, \{\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{\mathbf{A}}}^{(t)}\}_{t \in \mathcal{A}_1} \right\rangle \right|
$$

$$
+ \frac{1}{2}\pi \|\{A^{(t)}\theta^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\}_{t \in A_1}\|_{\mathrm{F}}^2 + \sum_{t \in A_1} \frac{1}{\sqrt{n}T} \lambda \|A^{(t)}(A^{(t)})^\top - \hat{A}(\widehat{A})^\top \|_2
$$
\n
$$
\geq -\frac{1}{T} \left| \left\langle \{\nabla f^{(t)}(A^{(t)*}\theta^{(t)*})\}_{t \in A_1}, \{A^{(t)}\theta^{(t)} - \hat{A}\theta_{\widehat{A}}^{(t)} + \hat{A}\theta_{\widehat{A}}^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\}_{t \in A_1} \right\rangle \right|
$$
\n
$$
- \frac{1}{T} \|\{\nabla f^{(t)}(\overline{A}\theta_{\overline{A}}^{(t)}) - \nabla f^{(t)}(A^{(t)*}\theta^{(t)*})\}\|_{\mathrm{F}} \|\{A^{(t)}\theta^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\}_{t \in A_1} \|_{\mathrm{F}} + \frac{1}{2}\frac{1}{T} \|\{A^{(t)}\theta^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\}_{t \in A_1} \|\hat{\mathrm{F}} + \sum_{t \in A_1} \frac{1}{\sqrt{n}T} \lambda \|A^{(t)}(A^{(t)})^\top - \hat{A}(\widehat{A})^\top \|_2
$$
\n
$$
\geq -\frac{1}{T} \sum_{t \in A_1} \|\nabla f^{(t)}(A^{(t)*}\theta^{(t)*})\|_2 \cdot (\|A^{(t)}\theta^{(t)} - \hat{A}\theta_{\widehat{A}}^{(t)}\|_2 + \|\hat{A}\theta_{\widehat{A}}^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\|_2)
$$
\n
$$
- \frac{1}{T}C \left[\sqrt{T} \sqrt{\frac{r + \log T}{n}} + \hbar \sqrt{\sum_{t \in A_1} (\zeta^{(t)})^2} \right] \cdot \|\{A^{(t)}\theta^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\}_{t \in A_1} \|\mathbf{F
$$

$$
-\overline{T} \bigcup \left\{ \mathbf{V}^T \mathbf{V} - n \right\} + \mathbf{N} \bigvee_{t \in A_1} (\mathbf{S}^{\circ}) \bigcup \left\{ \mathbf{N} \mathbf{A}^{\circ} \mathbf{V} - \mathbf{A} \mathbf{U}_{\overline{A}} \mathbf{f} t \in A_1 \mathbb{I} \right\}
$$

+
$$
\frac{1}{2} \frac{1}{T} \|\{\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{A}}^{(t)}\} t \in A_1 \mathbb{I} \|^2_{\mathbf{F}} + \sum_{t \in A_1} \frac{1}{\sqrt{n} T} \lambda \|\mathbf{A}^{(t)} (\mathbf{A}^{(t)})^{\top} - \widehat{\overline{\mathbf{A}}} (\widehat{\overline{\mathbf{A}}})^{\top} \mathbb{I}_2 \tag{46}
$$

$$
\geq -C\left(h^2 \cdot \frac{1}{T} \sum_{t \in \mathcal{A}_1} (\zeta^{(t)})^2 + \frac{r + \log T}{n}\right) + \frac{|\mathcal{A}_1|}{2\sqrt{n}T} \lambda \cdot \frac{1}{\sqrt{r}} \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A} \overline{A}^{\top}\|_2, \qquad (47)
$$

where [\(45\)](#page-95-0) holds because $\|\{\nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\}_{t\in\mathcal{A}_1}\|_2 \leq \sqrt{\frac{p}{n}} + h\sqrt{\sum_{t\in\mathcal{A}_1}(\zeta^{(t)})^2} +$ √ $\overline{T}\sqrt{\frac{r+\log T}{n}}$ n (Lemma [33\)](#page-51-2) and $\|\{\nabla f^{(t)}(\overline{A}\theta_{\overline{A}}^{(t)}) - \nabla f^{(t)}(A^{(t)*}\theta^{(t)*})\}\|_{\text{F}} =$ $\sqrt{\sum_{t\in\mathcal{A}_1} \|\widehat{\mathbf{\Sigma}}^{(t)}(\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)} - \boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_2^2} \lesssim \sqrt{\sum_{t\in\mathcal{A}_1} \|\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)} - \boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2^2} \lesssim \sqrt{h^2\sum_{t\in\mathcal{A}_1}(\zeta^{(t)})^2 + T\cdot\frac{r + \log T}{n}}$ \overline{n} (Lemma [30\)](#page-50-0). Inequality [\(46\)](#page-95-1) holds because

$$
\begin{aligned}\n\|A^{(t)}\boldsymbol{\theta}^{(t)}\|_{2} &\lesssim \|A^{(t)}(A^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_{2}(\|\boldsymbol{\theta}_{\widehat{\overline{A}}}^{(t)}\|_{2} + \|\nabla f^{(t)}(\widehat{\overline{A}}\boldsymbol{\theta}_{\widehat{\overline{A}}}^{(t)})\|_{2}) \\
&\lesssim \|A^{(t)}(A^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_{2}\Big(\zeta^{(t)} + \|\widehat{\overline{A}}\boldsymbol{\theta}_{\widehat{\overline{A}}}^{(t)} - A^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2} + \|\nabla f^{(t)}(A^{(t)*}\boldsymbol{\theta}^{(t)*})\|_{2}\Big) \\
&\lesssim \|A^{(t)}(A^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top\|_{2}\zeta^{(t)},\n\end{aligned}
$$

where we used Lemmas $30, 31,$ $30, 31,$ $30, 31,$ and 32 . Inequality (47) holds because

$$
\frac{1}{\sqrt{n}T}\lambda\sum_{t\in\mathcal{A}_1}\|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top-\widehat{\overline{\boldsymbol{A}}}(\widehat{\overline{\boldsymbol{A}}})^\top\|_2
$$

$$
-\frac{1}{T}\sum_{t\in\mathcal{A}_{1}}\sqrt{\frac{p+\log T}{n}}\cdot\zeta^{(t)}(\|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top}-\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}\|_{2}+\|\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}-\mathbf{A}\mathbf{A}^{\top}\|_{2})
$$

\n
$$
\geq \frac{1}{\sqrt{n}T}\lambda\sum_{t\in\mathcal{A}_{1}}\|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top}-\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}\|_{2}-\frac{1}{T}\sum_{t\in\mathcal{A}_{1}}C\sqrt{\frac{r(p+\log T)}{n}}\cdot\zeta^{(t)}\|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top}-\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}\|_{2}
$$

\n
$$
\geq \frac{1}{2\sqrt{n}T}\lambda\sum_{t\in\mathcal{A}_{1}}\|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top}-\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}\|_{2}
$$

\n
$$
\geq \frac{|\mathcal{A}_{1}|}{2\sqrt{n}T}\lambda\cdot\frac{1}{\sqrt{r}}\|\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}-\mathbf{A}\mathbf{A}^{\top}\|_{2},
$$

when $\frac{\lambda}{4}$ $\frac{n}{\sqrt{n}} \geq 2C \max_{t \in S} \zeta^{(t)} \cdot \sqrt{\frac{r(p + \log T)}{n}}$ $rac{\log I}{n}$. $\text{For any } t \in \mathcal{A}_2 \text{, since } \|\boldsymbol{A}^{(t)}\boldsymbol{\theta}^{(t)}_{\boldsymbol{A}^{(t)}} - \widehat{\overline{\boldsymbol{A}}}\boldsymbol{\theta}^{(t)}_{\widehat{\overline{\boldsymbol{A}}}}\|_2 \leq \|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top - \widehat{\overline{\boldsymbol{A}}}(\widehat{\overline{\boldsymbol{A}}})^\top\|_2 (\|\boldsymbol{\theta}^{(t)}_{\widehat{\overline{\boldsymbol{A}}}}\|_2^2)$ \boldsymbol{A} $\frac{\hat{a}^{(t)}}{\hat{A}}\|_2 + \|\nabla f^{(t)}(\widehat{\overline{A}} \theta_{\widehat{\overline{A}}}^{(t)})\|_2) \leq$ $c \zeta^{(t)} \| \widehat{\bar{\boldsymbol{A}}} (\widehat{\bar{\boldsymbol{A}}})^\top - \overline{\boldsymbol{A}} \overline{\boldsymbol{A}}^\top \|_2 \frac{1}{\sqrt{2}}$ $\frac{1}{r}$, by triangle inequality, we have

$$
\|\boldsymbol{A}^{(t)}\boldsymbol{\theta}_{\boldsymbol{A}^{(t)}}^{(t)} - \overline{\boldsymbol{A}}\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}\|_2 \geq \|\widehat{\overline{\boldsymbol{A}}}\boldsymbol{\theta}_{\widehat{\overline{\boldsymbol{A}}}}^{(t)} - \overline{\boldsymbol{A}}\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}\|_2 - \frac{c}{\sqrt{r}}\zeta^{(t)}\|\widehat{\overline{\boldsymbol{A}}}(\widehat{\overline{\boldsymbol{A}}})^\top - \overline{\boldsymbol{A}}\overline{\boldsymbol{A}}^\top\|_2.
$$

With similar arguments to obtain (47) , here we have

$$
\sum_{t \in A_2} \frac{1}{T} [f^{(t)}(\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)}) - f^{(t)}(\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{A}}^{(t)})] + \sum_{t \in A_2} \frac{\sqrt{n}}{nT} ||\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\mathbf{A}}(\widehat{\overline{\mathbf{A}}})^\top ||_2
$$
\n
$$
\geq \frac{1}{2T} ||\{\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{A}}^{(t)}\}_{t \in A_2}||_F^2 - \frac{1}{T} \sqrt{\frac{p+T}{n}} \cdot \sqrt{2r} ||\{\widehat{\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{A}}^{(t)} - \overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{A}}^{(t)}\}_{t \in A_2}||_F
$$
\n
$$
- \frac{C'}{T} \left(\sqrt{\frac{pr}{n}} + \sqrt{T} \sqrt{\frac{r + \log T}{n}} + h \sqrt{\sum_{t \in A_2} (\zeta^{(t)})^2} \right) \cdot ||\{\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{A}}^{(t)}\}_{t \in A_2}||_F
$$
\n
$$
+ \frac{1}{T} \sum_{t \in A_2} \frac{\lambda}{\sqrt{n}} ||\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\mathbf{A}}(\widehat{\overline{\mathbf{A}}})^\top ||_2 - \frac{C}{T} \sum_{t \in A_2} \sqrt{\frac{p + \log T}{n}} \cdot \zeta^{(t)} ||\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\mathbf{A}}(\widehat{\overline{\mathbf{A}}})^\top ||_2
$$
\n(48)

$$
\geq -C\left(\frac{pr}{nT} + h^2 \cdot \frac{1}{T} \sum_{t \in A_2} (\zeta^{(t)})^2 + \frac{r + \log T}{n} \right) + \frac{C}{T} \|\{A^{(t)}\theta^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\}_{t \in A_2} \|_F^2
$$

$$
- \frac{1}{T} \sqrt{\frac{p+T}{n}} \cdot \sqrt{2r} \left\|\{\widehat{A}\theta_{\widehat{A}}^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\}_{t \in A_2} \right\|_{\mathrm{F}}
$$

$$
= -C\left(\frac{pr}{nT} + h^2 \cdot \frac{1}{T} \sum_{t \in A_2} (\zeta^{(t)})^2 + \frac{r + \log T}{n} \right) + \frac{C}{T} \sum_{t \in A_2} \|\mathbf{A}^{(t)}\theta^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\|_2^2
$$

$$
- \frac{1}{T} \sqrt{\frac{p+T}{n}} \cdot \sqrt{2r} \left\|\{\widehat{A}\theta_{\widehat{A}}^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\}_{t \in A_2} \right\|_{\mathrm{F}} \tag{49}
$$

$$
\geq -C\left(\frac{pr}{nT} + h^2 \cdot \frac{1}{T} \sum_{t \in A_2} (\zeta^{(t)})^2 + \frac{r + \log T}{n}\right) + \frac{C'}{T} \sum_{t \in A_2} ||\hat{\overline{A}} \theta_{\hat{\overline{A}}}^{(t)} - \overline{A} \theta_{\overline{A}}^{(t)}||_2^2 \n- \frac{1}{T} \sqrt{\frac{p+T}{n}} \cdot \sqrt{2r} ||\{\hat{\overline{A}} \theta_{\hat{\overline{A}}}^{(t)} - \overline{A} \theta_{\overline{A}}^{(t)}\}_{t \in A_2}||_F - \frac{c}{Tr} \sum_{t \in A_2} (\zeta^{(t)})^2 \cdot ||\hat{\overline{A}}(\hat{\overline{A}})^T - \overline{A} \overline{A}^\top ||_2^2 \n\geq -C\left(\frac{pr}{nT} + h^2 \cdot \frac{1}{T} \sum_{t \in A_2} (\zeta^{(t)})^2 + \frac{r + \log T}{n}\right) + \frac{C'}{2T} \sum_{t \in A_2} ||\hat{\overline{A}} \theta_{\hat{\overline{A}}}^{(t)} - \overline{A} \theta_{\overline{A}}^{(t)}||_2^2 \n+ \frac{C'}{2T} ||\{\hat{\overline{A}} \theta_{\hat{\overline{A}}}^{(t)} - \overline{A} \theta_{\overline{A}}^{(t)}\}_{t \in A_2}||_F^2 - \frac{1}{T} \sqrt{\frac{p+T}{n}} \cdot \sqrt{2r} ||\{\hat{\overline{A}} \theta_{\hat{\overline{A}}}^{(t)} - \overline{A} \theta_{\overline{A}}^{(t)}\}_{t \in A_2}||_F^2 \n- \frac{c}{Tr} \sum_{t \in A_2} (\zeta^{(t)})^2 \cdot ||\hat{\overline{A}}(\hat{\overline{A}})^T - \overline{A} \overline{A}^\top ||_2^2.
$$
\n
$$
\geq -C\left(\frac{pr}{nT} + h^2 \cdot \frac{1}{T} \sum_{t \in A_2} (\zeta^{(t)})^2 + \frac{r + \log T}{n}\right) + \frac{C'}{2T} \sum_{t \in A_2} ||\hat{\overline{A}} \theta_{\overline{A}}^{(
$$

Compared to the derivation of (47) , here we used a different way to bound the intermediate $\begin{split} \text{term} \enspace & \frac{1}{T} | \langle \{\nabla f^{(t)}(\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\}_{t\in\mathcal{A}_2}, \{\boldsymbol{A}^{(t)}\boldsymbol{\theta}^{(t)}-\widehat{\widehat{\boldsymbol{A}}}\boldsymbol{\theta}_{\widehat{\widehat{\boldsymbol{A}}}}^{(t)}\} \rangle, \end{split}$ $+\stackrel{\frown}{A}\theta_{\widehat{\overline{A}}}^{(t)}$ $-\overline{A}\theta_{\overline{A}}^{(t)}\}_{t\in\mathcal{A}_2}\rangle|\,$ to obtain [\(48\)](#page-96-0). More specifically,

$$
\begin{split} &\frac{1}{T}\Big|\Big\langle\{\nabla f^{(t)}(\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\}_{t\in\mathcal{A}_2}, \{\boldsymbol{A}^{(t)}\boldsymbol{\theta}^{(t)}-\widehat{\boldsymbol{A}}\boldsymbol{\theta}_{\widehat{\boldsymbol{A}}}^{(t)}+\widehat{\boldsymbol{A}}\boldsymbol{\theta}_{\widehat{\boldsymbol{A}}}^{(t)}-\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\widehat{\boldsymbol{A}}}^{(t)}\}_{t\in\mathcal{A}_2}\Big\rangle\Big|\\ &\leq \frac{1}{T}\sum_{t\in\mathcal{A}_2}\|\nabla f^{(t)}(\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_2\|\boldsymbol{A}^{(t)}\boldsymbol{\theta}^{(t)}-\widehat{\boldsymbol{A}}\boldsymbol{\theta}_{\widehat{\boldsymbol{A}}}^{(t)}\|_2+\frac{1}{T}\|\{\nabla f^{(t)}(\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\}_{t\in\mathcal{A}_2}\|_2\|\{\widehat{\boldsymbol{A}}\boldsymbol{\theta}_{\widehat{\boldsymbol{A}}}^{(t)}-\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}\}_{t\in\mathcal{A}_2}\|_*\\ &\leq \frac{1}{T}\sum_{t\in\mathcal{A}_2}\sqrt{\frac{p+\log T}{n}}\zeta^{(t)}\|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top-\widehat{\boldsymbol{A}}(\widehat{\boldsymbol{A}})^\top\|_2+\frac{1}{T}\sqrt{\frac{p+T}{n}}\cdot\sqrt{2r}\left\|\{\widehat{\boldsymbol{A}}\boldsymbol{\theta}_{\widehat{\boldsymbol{A}}}^{(t)}-\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}\}_{t\in\mathcal{A}_2}\right\|_{\rm F},\end{split}
$$

where we used the fact that $|\langle A, B \rangle| \le ||A||_2 ||B||_* \le ||A||_2 ||B||_{\mathrm{F}} \cdot \text{rank}(B)$ for any matrix **A** and **B** of the same dimension and rank $\left(\{\hat{A}\theta_{\hat{A}}^{(t)}\}\)$ $-\overline{A}\theta_{\overline{A}}^{(t)}\}_{t\in\mathcal{A}_2}$) $\leq 2r$.

Finally, notice that by triangle inequality,

$$
G_{S^c}(\widehat{\overline{A}}) - G_{S^c}(\overline{A}) \ge \sum_{t \in S^c} \min_{\mathbf{A}^{(t)} \in \mathcal{O}^{p \times r}, \mathbf{\theta}^{(t)} \in \mathbb{R}^r} \left\{ \frac{\sqrt{n}}{nT} \lambda (\|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\overline{A}}(\widehat{\overline{A}})^\top \|_2 - \|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \overline{A}(\overline{A})^\top \|_2) \right\}
$$

$$
\ge -\frac{\sqrt{n}}{nT} \lambda |S^c| \cdot \|\widehat{\overline{A}}(\widehat{\overline{A}})^\top - \overline{A}(\overline{A})^\top \|_2.
$$
 (51)

Case 1: $|\mathcal{A}_2| \geq (1 - c/r \cdot \bar{\zeta}^2 / \max_{t \in S} (\zeta^{(t)})^2) |S|$ with some small constant $c > 0$:

Then by (47) , (50) , and (51) ,

$$
G(\widehat{\overline{A}}) - G(\overline{A}) = G_S(\widehat{\overline{A}}) - G_S(\overline{A}) + G_{S^c}(\widehat{\overline{A}}) - G_{S^c}(\overline{A})
$$

\n
$$
\geq -C\left(\frac{pr}{nT} + h^2\overline{\zeta}^2 + \frac{r + \log T}{n}\right) + \frac{C'}{T} \sum_{t \in A_2} \|\widehat{\overline{A}} \theta_{\widehat{\overline{A}}}^{(t)} - \overline{A} \theta_{\overline{A}}^{(t)}\|_2^2
$$

\n
$$
- \frac{c}{Tr} \sum_{t \in A_2} (\zeta^{(t)})^2 \cdot \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A} \overline{A}^{\top}\|_2^2
$$

\n
$$
- \frac{\sqrt{n}}{nT} \lambda |S^c| \cdot \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A} \overline{A}^{\top}\|_2.
$$
 (52)

By Assumption [2,](#page-10-0) Wedin's sin Θ-Theorem, and applying Lemma [36](#page-51-5) on $\{\theta_{\overline{A}}^{(t)}\}$ $\{\frac{\lambda^{(t)}}{\boldsymbol{A}}\}_{t\in\mathcal{A}_2},$

$$
\|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A}(\overline{A})^{\top}\|_2 \le \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A}(\overline{A})^{\top}\|_{\mathrm{F}} \lesssim \sqrt{\frac{r}{|S|}}\bar{\zeta}^{-1} \cdot \|\{\widehat{\overline{A}}\theta_{\widehat{\overline{A}}}^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)}\}_{t \in A_2} \|_{\mathrm{F}}.
$$
 (53)

Here we used the fact that $\frac{1}{|S|} \sum_{t \in S} \theta_{\overline{A}}^{(t)}$ $\frac{(t)}{\overline{A}}(\bm{\theta}_{\overline{A}}^{(t)}$ $\frac{f(t)}{A}$ ^T $\geq \frac{c}{r}$ $\frac{c}{r}\bar{\zeta}^2\mathbf{I}_p$. To see this, notice that

$$
\sigma_r \left(\frac{1}{\sqrt{|S|}} \{ \overline{A} \boldsymbol{\theta}_{\overline{A}}^{(t)} \}_{t \in S} \right) = \sigma_r \left(\frac{1}{\sqrt{|S|}} \{ \boldsymbol{\beta}^{(t)*} \}_{t \in S} \right) - \frac{1}{\sqrt{|S|}} || \{ \overline{A} \boldsymbol{\theta}_{\overline{A}}^{(t)} \}_{t \in S} - \{ \boldsymbol{\beta}^{(t)*} \}_{t \in S} ||_2
$$

$$
\geq \frac{c}{\sqrt{r}} \overline{\zeta} - \left(\overline{\zeta} h + \sqrt{\frac{r + \log T}{n}} \right)
$$

$$
\geq \frac{c'}{\sqrt{r}} \overline{\zeta},
$$

where by Lemmas [30](#page-50-0) and [34,](#page-51-3)

$$
\frac{1}{\sqrt{|S|}}\|\{\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}\}_{t\in S}-\{\boldsymbol{\beta}^{(t)*}\}_{t\in S}\|_2 \leq \sqrt{\sum_{t\in S}\|\overline{\boldsymbol{A}}\boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}-\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2^2} \lesssim h\bar{\zeta}+\sqrt{\frac{r+\log T}{n}},
$$

and the last step comes from conditions (i) and (ii) by noticing that

$$
\sqrt{\frac{r + \log T}{n}} \lesssim \frac{\bar{\zeta}}{\sqrt{r}} \cdot \sqrt{\frac{p + \log T}{n}} \cdot \frac{1}{\zeta^{(t)}} \lesssim \frac{\bar{\zeta}}{\sqrt{r}}.
$$

This leads to

$$
\lambda_r\left(\frac{1}{|S|}\sum_{t\in S}\theta_{\overline{A}}^{(t)}(\theta_{\overline{A}}^{(t)})^\top\right)=\lambda_r\left(\frac{1}{|S|}\sum_{t\in S}\theta_{\overline{A}}^{(t)}(\overline{A})^\top\overline{A}(\theta_{\overline{A}}^{(t)})^\top\right)\geq \frac{c'}{r}\overline{\zeta}^2.
$$

Plugging (53) into (52) , we have

$$
G(\widehat{\overline{A}}) - G(\overline{A}) \geq -C\left(h^2\overline{\zeta}^2 + \frac{r + \log T}{n}\right) + \frac{C'}{T} \cdot \frac{|S|}{r}\overline{\zeta}^2 \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A}\overline{A}^{\top}\|_2^2
$$

$$
-\frac{c|S|}{Tr}\bar{\zeta}^{2}\|\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top} - \overline{\mathbf{A}}\overline{\mathbf{A}}^{\top}\|_{2}^{2} - C\sqrt{\frac{r(p+T)}{nT}} \cdot \|\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top} - \overline{\mathbf{A}}\overline{\mathbf{A}}^{\top}\|_{2}\sqrt{\frac{1}{T} \sum_{t \in \mathcal{A}_{2}} (\zeta^{(t)})^{2}}
$$

$$
-\frac{\sqrt{n}}{nT}\lambda|S^{c}| \cdot \|\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top} - \overline{\mathbf{A}}\overline{\mathbf{A}}^{\top}\|_{2}
$$

$$
\geq \frac{C}{T} \cdot \frac{|S|}{r}\bar{\zeta}^{2}\|\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top} - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}\|_{2}^{2} - \frac{\sqrt{n}}{nT}\lambda|S^{c}| \cdot \|\widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top} - \overline{\mathbf{A}}(\overline{\mathbf{A}})^{\top}\|_{2}
$$

$$
-C\left(\frac{pr}{nT} + h^{2}\bar{\zeta}^{2} + \frac{r + \log T}{n}\right) - C\sqrt{\frac{r(p+T)}{nT}\bar{\zeta}}.
$$

$$
> 0,
$$

because $\|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A}\overline{A}^{\top}\|_2 \ge C\eta$ with some constant $C > 0$. **Case 2:** $|\mathcal{A}_1| \ge c/r \cdot \bar{\zeta}^2 / \max_{t \in S} (\zeta^{(t)})^2 \cdot |S|$ with some small constant $c > 0$: By inequality [\(49\)](#page-96-1) and triangle inequality, we have

$$
\sum_{t \in A_2} \frac{1}{T} [f^{(t)}(\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)}) - f^{(t)}(\overline{\mathbf{A}} \boldsymbol{\theta}_{\overline{A}}^{(t)})] + \sum_{t \in A_2} \frac{\sqrt{n}}{nT} ||\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^\top ||_2
$$
\n
$$
= -C \left(\frac{pr}{nT} + h^2 \cdot \frac{1}{T} \sum_{t \in A_2} (\zeta^{(t)})^2 + \frac{r + \log T}{n} \right) + \frac{C}{T} \sum_{t \in A_2} ||\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \overline{\mathbf{A}}\boldsymbol{\theta}_{\overline{A}}^{(t)}||_2^2
$$
\n
$$
- \frac{1}{T} \sqrt{\frac{p+T}{n}} \cdot \sqrt{2r} ||\{\widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)} - \overline{\mathbf{A}}\boldsymbol{\theta}_{\overline{A}}^{(t)}\}_{t \in A_2}||_F
$$
\n
$$
\geq -C \left(\frac{pr}{nT} + h^2 \cdot \frac{1}{T} \sum_{t \in A_2} (\zeta^{(t)})^2 + \frac{r + \log T}{n} \right) + \frac{C}{T} \sum_{t \in A_2} ||\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \overline{\mathbf{A}}\boldsymbol{\theta}_{\overline{A}}^{(t)}\|_2^2
$$
\n
$$
- \frac{1}{T} \sqrt{\frac{p+T}{n}} \cdot \sqrt{2r} ||\{\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \overline{\mathbf{A}}\boldsymbol{\theta}_{\overline{A}}^{(t)}\}_{t \in A_2}||_F - \frac{1}{T} \sqrt{\frac{p+T}{n}} \cdot \sqrt{2r} ||\{\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \widehat{\mathbf{A}}\boldsymbol{\theta}_{\overline{A}}^{(t)}\}_{t \in A_2}||_F
$$

$$
\geq -C\left(\frac{pr}{nT} + h^2 \cdot \frac{1}{T} \sum_{t \in A_2} (\zeta^{(t)})^2 + \frac{r + \log T}{n} \right) + \frac{C}{T} \left| \left\{ A^{(t)} \theta^{(t)} - \overline{A} \theta_{\overline{A}}^{(t)} \right\}_{t \in A_2} \right| \Big|_{\mathrm{F}}^2
$$

$$
- \frac{1}{T} \sqrt{\frac{p+T}{n}} \cdot \sqrt{2r} \left| \left\{ A^{(t)} \theta^{(t)} - \overline{A} \theta_{\overline{A}}^{(t)} \right\}_{t \in A_2} \right| \Big|_{\mathrm{F}}^2
$$

$$
- \sqrt{\frac{p+T}{nT}} \cdot \sqrt{2r} \cdot \sqrt{\frac{1}{T} \sum_{t \in A_2} \| A^{(t)} (A^{(t)})^\top - \widehat{A} (\widehat{A})^\top \|^2_2 (\zeta^{(t)})^2}
$$

$$
\geq -C\left(\frac{pr}{nT} + h^2 \cdot \frac{1}{T} \sum_{t \in A_2} (\zeta^{(t)})^2 + \frac{r + \log T}{n} \right) + \frac{C}{T} \left| \left\{ A^{(t)} \theta^{(t)} - \overline{A} \theta_{\overline{A}}^{(t)} \right\}_{t \in A_2} \right| \Big|_{\mathrm{F}}^2
$$

$$
- \frac{1}{T} \sqrt{\frac{p+T}{n}} \cdot \sqrt{2r} \left| \left\{ A^{(t)} \theta^{(t)} - \overline{A} \theta_{\overline{A}}^{(t)} \right\}_{t \in A_2} \right| \Big|_{\mathrm{F}}^2 - \sqrt{\frac{p+T}{nT}} \cdot \overline{\zeta} \|\widehat{A}(\widehat{A})^\top - \overline{A} \overline{A}^\top \|^2_2
$$

$$
\geq -C\left(\frac{pr}{nT} + h^2\bar{\zeta}^2 + \frac{r + \log T}{n}\right) - \sqrt{\frac{p+T}{nT}} \cdot \bar{\zeta} \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A}\overline{A}^{\top}\|_2
$$
\n(54)

By the condition of the proposition, we must have

$$
\frac{\min_{t \in S} \zeta^{(t)}}{\bar{\zeta}} \cdot r \lesssim \sqrt{p + \log T}, \quad \frac{\min_{t \in S} \zeta^{(t)}}{\bar{\zeta}} \cdot \frac{r}{\sqrt{T}} \lesssim 1. \tag{55}
$$

By (47) , (51) , and (54) , we have

$$
G(\widehat{A}) - G(\overline{A})
$$

\n
$$
\geq \frac{C\lambda}{\sqrt{n}T} \left(\frac{|A_1|}{\sqrt{r}} - |S^c| \right) \cdot ||\widehat{A}(\widehat{A})^{\top} - \overline{A}\overline{A}^{\top}||_2 - \sqrt{\frac{p+T}{nT}} \cdot \overline{\zeta} ||\widehat{A}(\widehat{A})^{\top} - \overline{A}\overline{A}^{\top}||_2
$$

\n
$$
- C \left(\frac{pr}{nT} + h^2 \cdot \overline{\zeta}^2 + \frac{r + \log T}{n} \right)
$$

\n
$$
\geq \frac{C\lambda}{\sqrt{n}T} \left(\frac{|S|}{r^{3/2}} \cdot \frac{\overline{\zeta}^2}{\max_{t \in S} (\zeta^{(t)})^2} - |S^c| \right) \cdot ||\widehat{A}(\widehat{A})^{\top} - \overline{A}\overline{A}^{\top}||_2 - \sqrt{\frac{p+T}{nT}} \cdot \overline{\zeta} ||\widehat{A}(\widehat{A})^{\top} - \overline{A}\overline{A}^{\top}||_2
$$

$$
-C\left(\frac{pr}{nT} + h^2 \cdot \bar{\zeta}^2 + \frac{r + \log T}{n}\right)
$$

\n
$$
\geq \frac{C\lambda}{\sqrt{n}} r^{-3/2} \cdot \frac{\bar{\zeta}^2}{\max_{t \in S}(\zeta^{(t)})^2} \cdot \|\widehat{A}(\widehat{A})^\top - \overline{A}\overline{A}^\top\|_2 - \sqrt{\frac{p+T}{nT}} \cdot \bar{\zeta}\|\widehat{A}(\widehat{A})^\top - \overline{A}\overline{A}^\top\|_2
$$

\n
$$
-C\left(\frac{pr}{nT} + h^2 \cdot \bar{\zeta}^2 + \frac{r + \log T}{n}\right)
$$

\n
$$
\geq \frac{C\lambda}{2\sqrt{n}} r^{-3/2} \cdot \frac{\bar{\zeta}^2}{\max_{t \in S}(\zeta^{(t)})^2} \cdot \|\widehat{A}(\widehat{A})^\top - \overline{A}\overline{A}^\top\|_2 - C\left(\frac{pr}{nT} + h^2 \cdot \bar{\zeta}^2 + \frac{r + \log T}{n}\right)
$$

due to the assumption that $|S^c|/T \leq c r^{-3/2} \bar{\zeta}/\max_{t \in S} \zeta^{(t)}$ with a small constant $c > 0$ and

$$
\lambda \approx \sqrt{r(p + \log T)} \cdot \frac{\max_{t \in S} (\zeta^{(t)})^2}{\min_{t \in S} \zeta^{(t)}} \times \left[\sqrt{pr} \frac{\bar{\zeta}}{\min_{t \in S} \zeta^{(t)}} + \sqrt{r(p + \log T)} \cdot \frac{\bar{\zeta}}{\min_{t \in S} \zeta^{(t)}} \right] \cdot \frac{\max_{t \in S} (\zeta^{(t)})^2}{\bar{\zeta}},
$$

where we used [\(55\)](#page-100-1) in the second inequality. Furthermore, by $\min_{t \in S} \zeta^{(t)} \gtrsim \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$, we have

$$
G(\widehat{\overline{A}}) - G(\overline{A})
$$

\n
$$
\geq C'' \sqrt{\frac{p + \log T}{n}} r^{-1} \cdot \frac{\overline{\zeta}^2}{\min_{t \in S} \zeta(t)} \cdot \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A} \overline{A}^{\top}\|_2 - C \left(\frac{pr}{nT} + h^2 \cdot \overline{\zeta}^2 + \frac{r + \log T}{n}\right)
$$

\n
$$
\geq C'' r^{-1} \cdot \overline{\zeta}^2 \cdot \|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A} \overline{A}^{\top}\|_2 - C \left(\frac{pr}{nT} + h^2 \cdot \overline{\zeta}^2 + \frac{r + \log T}{n}\right).
$$

Therefore, if $\|\widehat{\overline{A}}(\widehat{\overline{A}})^{\top} - \overline{A} \overline{A}^{\top} \|_2 \ge C\eta$, we must have

$$
G(\widehat{\overline{\mathbf{A}}}) - G(\overline{\mathbf{A}}) > 0,
$$

which implies that $\widehat{\overline{A}}$ is not a minimizer of G and contradicts with the optimality of $\widehat{\overline{A}}$. Hence we must have $\|\widehat{\overline{\mathbf{A}}}(\widehat{\overline{\mathbf{A}}})^{\top} - \overline{\mathbf{A}}\overline{\mathbf{A}}^{\top}\|_2 \le C\eta \asymp \overline{\zeta}^{-1}r\sqrt{\frac{p}{nT}} + \overline{\zeta}^{-1}\sqrt{r}\sqrt{\frac{r+\log T}{n}} +$ $\sqrt{r}h + \frac{\lambda r}{\sqrt{n}\bar{\zeta}^2}\epsilon$ with some constant $C > 0$. This completes our proof.

Appendix E. Proofs for GLMs

E.1 Lemmas

Lemma [4](#page-21-0)4 Suppose Assumptions [1](#page-10-1) and 4 hold, and $n \geq C(p + \log T)$ with a sufficiently large C. Then when $\delta \leq C$ with some constant $C > 0$,

$$
\|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2 \lesssim \delta,
$$

w.p. at least $1 - \exp\{-Cn\delta^2 + C'p\}$. By taking $\delta \propto \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$, this implies that:

- (i) $\max_{t \in S} \|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2 \lesssim \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$, w.p. at least $1 - e^{-C(p + \log T)}$;
- $(ii) \|\nabla f^{(0)}(\boldsymbol{\beta}^{(0)*})\|_2 \lesssim \sqrt{\frac{p}{n_0}}, \ w.p. \ \text{at least} \ 1 e^{-C(p + \log T)}.$

Lemma [4](#page-21-0)5 Suppose Assumptions [1](#page-10-1) and 4 hold, and $n \geq C(p + \log T)$ with a sufficiently large C. Then when $t \leq C'$ with some constant $C' > 0$, for any $S \subseteq [T]$,

$$
\|\{\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\}_{t\in S}\|_2 \lesssim \sqrt{\frac{p+|S|+\delta}{n}},
$$

w.p. at least $1 - e^{-C'\delta}$.

Lemma 46 (A revised version of Proposition 1 in [Loh and Wainwright](#page-37-0) [\(2015\)](#page-37-0)) Suppose Assumptions [1](#page-10-1) and [4](#page-21-0) hold, and $n \geq C(p + \log T)$ with a sufficiently large C. Then there exist constants C_1, C_2 , such that w.p. at least $1 - e^{-C'(p + \log T)}$, for all $\beta \in \mathbb{R}^p$, $t \in [T]$, we have

$$
f^{(t)}(\boldsymbol{\beta}) - f^{(t)}(\boldsymbol{\beta}^{(t)*}) - \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})^{\top}(\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}) \ge \begin{cases} C_1 \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}\|_2^2, & when \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}\|_2 \le 1; \\ C_2 \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}\|_2, & when \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}\|_2 > 1. \end{cases}
$$
(56)

Lemma [4](#page-21-0)7 Suppose Assumptions [1](#page-10-1) and 4 hold, and $n \ge C(p + \log T)$ with a sufficiently large C. Then there exist constants C_1, C_2, C_3, C_4 , such that for all $\beta, \widetilde{\beta} \in \mathbb{R}^p$, $t \in [T]$, w.p. at least $1 - e^{-C(p + \log T)}$,

$$
f^{(t)}(\boldsymbol{\beta}) - f^{(t)}(\widetilde{\boldsymbol{\beta}}) - \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})^{\top}(\boldsymbol{\beta} - \widetilde{\boldsymbol{\beta}}) \geq \begin{cases} C_1 \|\boldsymbol{\beta} - \widetilde{\boldsymbol{\beta}}\|_2^2 - C_2 \|\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}^{(t)*}\|_2^2, & \text{if } \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}\|_2 \leq 1; \\ C_3 \|\boldsymbol{\beta} - \widetilde{\boldsymbol{\beta}}\|_2 - C_4 \|\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}^{(t)*}\|_2^2, & \text{if } \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}\|_2 > 1. \end{cases}
$$

Lemma [4](#page-21-0)8 Suppose Assumptions [1](#page-10-1) and 4 hold, and $n \geq C(p + \log T)$ with a sufficiently large C. Then w.p. at least $1 - e^{-C'(p + \log T)}$, for all $t \in S$, for all $A \in \mathcal{O}^{p \times r}$ with $\|\boldsymbol{A}\boldsymbol{A}^{\top}-\boldsymbol{A}^{(t)*}(\boldsymbol{A}^{(t)*})^{\top}\|_2 \leq c(\zeta^{(t)})^{-1}$, where $c > 0$ is a small constant, we have

- (i) $\|\theta_A^{(t)}\|_2 \le C'' \zeta^{(t)};$
- (ii) $||A\theta_A^{(t)} A^{(t)*}\theta^{(t)*}||_2 \leq c'$ with a small constant $c' > 0$.

Lemma [4](#page-21-0)9 Suppose Assumptions [1](#page-10-1) and 4 hold, and $n \geq C(p + \log T)$ with a sufficiently large C. Assume there exists $\widetilde{A} \in \mathcal{O}^{p \times r}$ such that $\|\widetilde{A}\theta_{\widetilde{A}}^{(t)} - A^{(t)*}\theta^{(t)*}\|_2 \leq c$ with a small constant c. Then w.p. at least $1 - e^{-C'(p + \log T)}$, for all $t \in S$, for all $A \in \mathcal{O}^{p \times r}$ with $||AA^{\top} - \widetilde{A}(\widetilde{A})^{\top}||_2 \leq c'(\zeta^{(t)})^{-1}$, where $c' > 0$ is a small constant, we have

(i) $\|\theta_A^{(t)}\|_2 \leq C'' \zeta^{(t)}$; (ii) $||A\theta^{(t)}_A - A^{(t)*}\theta^{(t)*}||_2 \leq c''$ with a small constant $c'' > 0$.

Lemma 50 Suppose Assumptions [1](#page-10-1) and [4](#page-21-0) hold, and $n \ge C(p + \log T)$ with a sufficiently large C. Then w.p. at least $1 - e^{-C'(p + \log T)}$, for all $t \in S$, for all $A, \widetilde{A} \in \mathcal{O}^{p \times r}$ with $\|\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)} - \boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2, \|\boldsymbol{\widetilde{A}}\boldsymbol{\theta}_{\boldsymbol{\widetilde{A}}}^{(t)} - \boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \leq c',$ where $c' > 0$ is a small constant, we have

$$
\|\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}-\widetilde{\boldsymbol{A}}\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)}\|_{2}\lesssim\|\boldsymbol{A}\boldsymbol{A}^{\top}-\widetilde{\boldsymbol{A}}(\widetilde{\boldsymbol{A}})^{\top}\|_{2}(\|\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)}\|_{2}+\|\nabla f^{(t)}(\widetilde{\boldsymbol{A}}\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)})\|_{2}).
$$

Lemma 5[1](#page-10-1) Suppose Assumptions 1 and [4](#page-21-0) hold, and $n \geq C(p + \log T)$ with a sufficiently large C. Then w.p. at least $1 - e^{-C'(p + \log T)}$, for all $t \in S$, for all $A, \widetilde{A} \in \mathcal{O}^{p \times r}$ with $\|\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}-\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2\leq c', \textit{ where } c'>0 \textit{ is a small constant, we have }$

$$
\|\boldsymbol A\boldsymbol \theta^{(t)}_{\boldsymbol A} - \boldsymbol A^{(t)*}\boldsymbol \theta^{(t)*}\|_2 \lesssim \|\boldsymbol A\boldsymbol A^\top - \boldsymbol A^{(t)*}(\boldsymbol A^{(t)*})^\top\|_2 \zeta^{(t)} + \sqrt{\frac{r+\log T}{n}}.
$$

Lemma 52 Suppose Assumptions [1](#page-10-1) and [4](#page-21-0) hold, and $n \ge C[(p + \log T) \vee \lambda^2]$ with a sufficiently large C. Given $\widetilde{A} \in \mathcal{O}^{p \times r}$, denote $(A^{(t)}, \theta^{(t)}) = \arg \min_{A \in \mathcal{O}^{p \times r}, \theta \in \mathbb{R}^r} \{f^{(t)}(A\theta) + \frac{\lambda}{\sqrt{p}}\}$ $\frac{\Delta}{n}\|\bm{A}\bm{A}^\top \widetilde{A}(\widetilde{A})^{\top} \|_{2}$. Then w.p. at least $1 - e^{-C(p + \log T)}$, for any $\widetilde{A} \in \mathcal{O}^{p \times r}$, we have $||A^{(t)}\theta^{(t)} \boldsymbol{\beta}^{(t)*} \|_2 \leq c$, where $c > 0$ is a small constant.

E.2 Proof of Theorem [9](#page-22-0)

For the penalized ERM, the proof logic is almost the same as the logic underlying the proofs of Proposition [43](#page-52-2) and Theorem [1.](#page-12-0) We only point out the difference and skip the details here. We need to bound $\|\overline{A}(\overline{A})^{\top} - \overline{A}(\overline{A})^{\top}\|_2$ first. Then we know that $\|\overline{A}(\overline{A})^{\top} - \overline{A}(\overline{A})^{\top}\|_2$ and $\|\hat{\vec{A}}\theta_{\hat{\vec{A}}}^{(t)} - A^{(t)*}\theta^{(t)*}\|_2$ are small w.h.p. Then a direct application of Lemma [49](#page-102-0) implies $\mathcal{A}_{\mathcal{L}_{\mathcal{L}}}$ that $\|\widehat{\vec{A}}\theta_{\widehat{\vec{A}}}^{(t)*} - A^{(t)*}\theta^{(t)*}\|_2$ is small w.h.p. Finally, applying Lemma [50,](#page-102-1) we have $\|\widehat{\vec{A}}\theta_{\widehat{\vec{A}}}^{(t)*} A \sim \sim$ $\begin{split} \overline{\bm{A}}\bm{\theta}^{(t)}_{\overline{\bm{A}}} \|_{2} \lesssim \| \widehat{\bar{\bm{A}}}(\widehat{\bar{\bm{A}}})^\top - \overline{\bm{A}}(\overline{\bm{A}})^\top \|_{2} (\zeta^{(t)}+\|\nabla f^{(t)}(\overline{\bm{A}}\bm{\theta}^{(t)}_{\overline{\bm{A}}})\|_{2}) \lesssim \| \widehat{\bar{\bm{A}}}(\widehat{\bar{\bm{A}}})^\top - \overline{\bm{A}}(\overline{\bm{A}})^\top \|_{2} \zeta^{(t)} \,\, \mathrm{w.h.p.} \end{split}$ which gives us the ideal bound for $\|\widehat{\overline{A}} \theta_{\widehat{\overline{A}}}^{(t)*}$ $-\overline{A}\theta_{\overline{A}}^{(t)}\|_2$ for all $t \in S$.

(i) In the proof of Proposition [43,](#page-52-2) for $t \in \mathcal{A}_1$, we know $||\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}||_2 \leq \text{a small}$ constant c w.h.p. by Lemma [52.](#page-102-2) Then by Lemmas $45, 47,$ $45, 47,$ $45, 47,$ and 51 , w.p. at least 1 $e^{-C'(r+\log T)},$

$$
\frac{1}{T} \sum_{t \in A_1} \left[f^{(t)}(A^{(t)}\theta^{(t)}) - f^{(t)}(\overline{A}\theta_{\overline{A}}^{(t)}) \right] \n\geq -\frac{1}{T} \left| \left\langle \{ \nabla f^{(t)}(A^{(t)*}\theta^{(t)*}) \}_{t \in A_1}, \{ A^{(t)}\theta^{(t)} - \hat{A}\theta_{\widehat{A}}^{(t)} + \hat{A}\theta_{\widehat{A}}^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)} \}_{t \in A_1} \right\rangle \right| \n+ \frac{C_1}{T} \|\{ A^{(t)}\theta^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)} \}_{t \in A_1} \|^2_{\mathrm{F}} - \frac{C_2}{T} \|\{ A^{(t)*}\theta^{(t)*} - \overline{A}\theta_{\overline{A}}^{(t)} \}_{t \in A_1} \|^2_{\mathrm{F}} \n\geq -\frac{C}{T} \sum_{t \in A_1} \sqrt{\frac{p + \log T}{n}} \cdot \zeta^{(t)} (\| A^{(t)}(A^{(t)})^\top - \hat{A}(\widehat{A})^\top \|_2 + \|\widehat{A}(\widehat{A})^\top - \overline{A}\overline{A}^\top \|_2) \n+ \frac{C_1}{T} \|\{ A^{(t)}\theta^{(t)} - \overline{A}\theta_{\overline{A}}^{(t)} \}_{t \in A_1} \|^2_{\mathrm{F}} - C \left(h^2 \cdot \frac{1}{T} \sum_{t \in A_1} (\zeta^{(t)})^2 + \frac{r + \log T}{n} \right) \n\geq -C \left(h^2 \cdot \frac{1}{T} \sum_{t \in A_1} (\zeta^{(t)})^2 + \frac{r + \log T}{n} \right) - \frac{C'}{T} \sum_{t \in A_1} \sqrt{\frac{p + \log T}{n}} \cdot \zeta^{(t)} \sqrt{r} \| A^{(t)}(A^{(t)})^\top - \widehat{A}(\widehat{A})^\top \|_2.
$$

Therefore the calculations in [\(47\)](#page-95-2) are still correct under GLMs. (ii) In the proof of Proposition [43,](#page-52-2) for $t \in \mathcal{A}_2$, we used the fact that w.p. at least 1 − $e^{-C(r+\log T)},$

$$
\|\mathbf{A}^{(t)}\boldsymbol{\theta}_{\mathbf{A}^{(t)}}^{(t)} - \widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)}\|_{2} \lesssim \|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top} - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}\|_{2}(\|\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)}\|_{2} + \|\nabla f^{(t)}(\widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)})\|_{2}).
$$
 (57)

Here this still holds because

- $||A^{(t)}\theta_{A^{(t)}}^{(t)} A^{(t)*}\theta^{(t)*}||_2 \leq \text{a small constant } c \text{ by Lemma 52};$
- For $t \in \mathcal{A}_2$, we know that $||\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top \overline{\mathbf{A}}(\overline{\mathbf{A}})^\top ||_2 \leq \frac{c}{\sqrt{r}} \leq c'(\zeta^{(t)})^{-1}$ with a small constant $c' > 0$. Then by Lemma [49,](#page-102-0) $\|\boldsymbol{\theta}_{\simeq}^{(t)}\|$ $\frac{d}{d}\left\Vert 2\right\Vert \leq C\zeta^{\left(t\right)}$ and $\Vert \widehat{\overline{A}}\theta_{\widehat{\overline{A}}}^{\left(t\right)}$ $-\bm{A}^{(t)*} \bm{\theta}^{(t)*} \|_2 \leq Cc'$ with a small constant $c' > 0$.

Then [\(57\)](#page-103-0) holds by Lemma [50.](#page-102-1)

Finally, Lemma [39](#page-52-0) still holds for the case of GLMs. Note that in Lemma [39,](#page-52-0) (i) only requires the maximum eigenvalue of Hessian to be upper bounded (see Lemma E.3 in [Duan](#page-36-0) [and Wang](#page-36-0) [\(2022\)](#page-36-0)), which is true by Assumptions [1](#page-10-1) and [4.](#page-21-0) By Lemma E.2 in [Duan and](#page-36-0) [Wang](#page-36-0) [\(2022\)](#page-36-0), (ii) and (iii) require the minimum eigenvalue of Hessian to be lower bounded when $||\boldsymbol{\beta}^{(t)} - \boldsymbol{\beta}^{(t)*}||_2 \le C$ with some constant $C > 0$ w.h.p. and $\gamma / \sqrt{n_0} \le C$, both of which are true.

For the spectral method, the proof is almost the same as the proof of Theorem [7,](#page-17-0) hence we do not repeat it here.

E.3 Proof of Theorem [17](#page-46-1)

We only outline the proof when Algorithm [4](#page-45-1) is coupled with Algorithm [1.](#page-11-0) If Algorithm 4 is coupled with Algorithm [2,](#page-17-1) we can follow the similar argument with the proof of Theorem [7.](#page-17-0)

Denote $\eta = r \sqrt{\frac{p}{nT}} + \sqrt{r}h + \sqrt{r} \sqrt{\frac{r + \log T}{n}} + \frac{|S^c|}{T}$ $\frac{S^c|}{T} \cdot r \cdot \frac{\lambda}{\sqrt{n}}$ $\frac{1}{n}$. (i) When $\eta \leq C \sqrt{\frac{p}{n_0}}$: since $\|\widehat{A}(\widehat{A})^{\top} - A^{(0)*}(A^{(0)*})^{\top}\|_2 \leq a$ small constant c, by Lemma [52,](#page-102-2) w.p. at least $1 - e^{-C(p + \log T)}$, $\|\widehat{\overline{A}} \theta_{\widehat{\overline{A}}}^{(t)}\|$ $- A^{(0)*}\theta^{(0)*} \|_2 \leq$ a small constant c' . Then by Lemma [51](#page-102-3) and Theorem [9,](#page-22-0) w.p. at least $1 - e^{-C(r + \log T)}$, $\|\hat{\vec{A}}\theta_{\hat{\vec{A}}}^{(t)}\|$ $- \, A^{(0) *} \theta^{(0) *} \|_2 \, \lesssim$ $\|\overline{A}(\overline{A})^{\top} - A^{(0)*}(A^{(0)*})^{\top}\|_2 \lesssim \eta$. As we commented in the proof of Theorem [9,](#page-22-0) Lemma [42.](#page-52-3)(i) still holds for GLMs because the maximum eigenvalue of Hessian is upper bounded w.h.p. (ii) When $\eta > C \sqrt{\frac{p}{n_0}}$: the result comes from Lemma [42.](#page-52-3)(ii), which still holds when $\gamma / \sqrt{n_0} \lesssim$ some constant C because of Lemma E.2 in [Duan and Wang](#page-36-0) (2022) .

E.4 Proofs of Lemmas

E.4.1 PROOF OF LEMMA 44

First, note that $\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*}) = \frac{2}{n} \sum_{i=1}^{n} \boldsymbol{x}_i^{(t)}$ $\psi_i^{(t)}[y_i^{(t)}] - \psi^\prime((\boldsymbol{x}_i^{(t)})$ $\left(\begin{matrix} (t) \\ i \end{matrix} \right)^{\top} \boldsymbol{\beta}^{(t)*}$]. Consider a 1/2-cover of the unit ball in \mathbb{R}^p w.r.t Euclidean norm $\{u_j\}_{j=1}^N$ with $N \leq 5^p$ (Example 5.8 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0)). By a standard argument (see the proof of Theorem 6.5 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0)), we have

$$
\|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2 \leq 2 \max_{j \in [N]} \left\{ \frac{2}{n} \sum_{i=1}^n \boldsymbol{u}_j^{\top} \boldsymbol{x}_i^{(t)} [y_i^{(t)} - \psi'((\boldsymbol{x}_i^{(t)})^{\top} \boldsymbol{\beta}^{(t)*})] \right\}.
$$

By union bound,

$$
\mathbb{E}\exp\{\lambda\|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2\} \leq N \max_{j\in[N]}\mathbb{E}\exp\left\{\frac{4\lambda}{n}\sum_{i=1}^n \boldsymbol{u}_j^\top \boldsymbol{x}_i^{(t)}[y_i^{(t)} - \psi'((\boldsymbol{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*})]\right\} = (*).
$$

Note that by the GLM density function, for any $a \in \mathbb{R}$,

$$
\mathbb{E} \exp\left\{a[y_i^{(t)} - \psi'((\mathbf{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*})]\right\} \leq \int \rho(y) \exp\left\{[a + (\mathbf{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*}]y_i^{(t)} - \psi(a + (\mathbf{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*})\right\} d\mu(y)
$$

$$
\cdot \exp\left\{\psi(a + (\mathbf{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*}) - \psi((\mathbf{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*}) - a\psi'((\mathbf{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*})\right\}
$$

$$
\leq \exp\left\{\frac{1}{2}a^2\psi''(\delta[a + (\mathbf{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*}] + (1 - \delta)(\mathbf{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*})\right\}
$$

$$
\leq \exp\left\{Ca^2\right\},
$$
(58)

where $\delta \in [0,1]$. Therefore, for any positive $\lambda \leq Cn$ with some $C > 0$,

$$
(*) \leq N \max_{j \in [N]} \mathbb{E} \exp \left\{ C' \frac{\lambda^2}{n^2} \sum_{i=1}^n (\boldsymbol{u}_j^\top \boldsymbol{x}_i^{(t)})^2 \right\} \leq \exp \left\{ \frac{C'' \lambda^2}{n} + C''' p \right\},\
$$

where the second inequality comes from the property of sub-exponential variables (see property 3 in Proposition 2.7.1 of [Vershynin](#page-39-2) [\(2018\)](#page-39-2)). Finally, for any $\delta > 0$, by Chernoff's bound,

$$
\mathbb{P}(\|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2 > \delta) \leq e^{-\lambda t} \mathbb{E} \exp\{\lambda \|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2\} \leq \exp\left\{\frac{C''\lambda^2}{n} + C'''p - \lambda \delta\right\}.
$$

Let $\lambda = C''n\delta/2$ (δ < some constant C), leading to

$$
\mathbb{P}(\|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2 > \delta) \le \exp\left\{-Cn\delta^2 + C'p\right\},\,
$$

for all $t \in S$.

E.4.2 Proof of Lemma [45](#page-101-0)

Note that $\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*}) = \frac{2}{n} \sum_{i=1}^{n} \boldsymbol{x}_i^{(t)}$ $\hat{y}_i^{(t)}[y_i^{(t)} - \psi^\prime((\boldsymbol{x}_i^{(t)})$ $[\bm{s}^{(t)}_i)^\top \bm{\beta}^{(t)*}] \coloneqq \bm{g}^{(t)} \in \mathbb{R}^p$. Denote $\{\bm{g}^{(t)}\}_{t \in S}$ as $G \in \mathbb{R}^{p \times |S|}$. Consider two 1/4-covers of the unit ball in \mathbb{R}^p and $\mathbb{R}^{|S|}$ w.r.t Euclidean norm, as $\{u^{(j_1)}\}_{j_1=1}^{N_1}$ and $\{v^{(j_2)}\}_{j_2=1}^{N_2}$ respectively, with $N_1 \leq 9^p$ and $N_2 \leq 9^{|S|}$ (Example 5.8 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0)). By a standard argument (see the proof of Theorem 6.5 in [Wainwright](#page-39-0) (2019) , we have

$$
\|\bm{G}\|_2 \lesssim \max_{\substack{j_1 \in [N_1] \\ j_2 \in [N_2]}} |(\bm{u}^{(j_1)})^\top \bm{G} \bm{v}^{(j_2)}| = \max_{\substack{j_1 \in [N_1] \\ j_2 \in [N_2]}} \left| \sum_{t \in S} (\bm{u}^{(j_1)})^\top \bm{g}^{(t)} v_k^{j_2} \right|.
$$

It follows that for any $\lambda \in \mathbb{R}$,

$$
\begin{aligned} \mathbb{E} e^{\lambda\|{\bm G}\|_2} &\lesssim \sum_{j_1=1}^{N_1}\sum_{j_2=1}^{N_2}\prod_{t=1}^{|S|} \mathbb{E} \exp\{\lambda |({\bm u}^{(j_1)})^\top {\bm g}^{(t)} v_k^{j_2}| \} \\ &\leq \sum_{j_1=1}^{N_1}\sum_{j_2=1}^{N_2}\prod_{t=1}^{|S|} \big[\mathbb{E} \exp\{\lambda ({\bm u}^{(j_1)})^\top {\bm g}^{(t)} v_k^{j_2}\} + \mathbb{E} \exp\{-\lambda ({\bm u}^{(j_1)})^\top {\bm g}^{(t)} v_k^{j_2}\} \big]. \end{aligned}
$$

Similar to the calculations in [\(58\)](#page-104-0), we can obtain that

$$
\mathbb{E} \exp\{\lambda(\boldsymbol{u}^{(j_1)})^\top \boldsymbol{g}^{(t)} v_k^{(j_2)}\}, \mathbb{E} \exp\{-\lambda(\boldsymbol{u}^{(j_1)})^\top \boldsymbol{g}^{(t)} v_k^{(j_2)}\} \lesssim \exp\left\{\frac{C}{n}\lambda^2 \cdot |v_k^{(j_2)}|^2\right\},\
$$

hence

$$
\mathbb{E}e^{\lambda||G||_2} \lesssim \sum_{j_1=1}^{N_1} \sum_{j_2=1}^{N_2} 2^{|S|} \cdot \exp\left\{\frac{C}{n}\lambda^2\right\} \lesssim \exp\left\{\frac{C}{n}\lambda^2 + C'p + C'|S|\right\}.
$$

Then similar to the proof of Lemma [44,](#page-101-2) the proof can be finished by Chernoff's bound.

E.4.3 Proof of Lemma [46](#page-101-3)

The proof of part (i) follows the idea in the proof of Theorem 9.36 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0), and the proof of part (ii) is the same as the proof of equation (39b) in Proposition 1 of [Loh](#page-37-0) [and Wainwright](#page-37-0) [\(2015\)](#page-37-0).

(i) If $\|\boldsymbol{\beta}-\boldsymbol{\beta}^{(t)*}\|_2 \leq 1$: Denote $\boldsymbol{\Delta} = \boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}, \varphi_{\tau}(u) = u^2 \mathbb{1}(|u| \leq \tau)$ with any $\tau > 0$. By Taylor expansion, with some $\delta \in [0,1]$ and any constant $T > 0$,

LHS of (56) =
$$
\frac{1}{n} \sum_{i=1}^{n} \psi''((\boldsymbol{\beta}^{(t)*})^{\top} \boldsymbol{x}_i^{(t)} + \delta \boldsymbol{\Delta}^{\top} \boldsymbol{x}_i^{(t)}) \cdot (\boldsymbol{\Delta}^{\top} \boldsymbol{x}_i^{(t)})^2
$$

\n
$$
\geq \frac{1}{n} \sum_{i=1}^{n} \psi''((\boldsymbol{\beta}^{(t)*})^{\top} \boldsymbol{x}_i^{(t)} + \delta \boldsymbol{\Delta}^{\top} \boldsymbol{x}_i^{(t)}) \varphi_{\tau}(\boldsymbol{\Delta}^{\top} \boldsymbol{x}_i^{(t)}) \mathbb{1}(|(\boldsymbol{\beta}^{(t)*})^{\top} \boldsymbol{x}_i^{(t)}| \leq T)
$$

\n
$$
\geq \min_{|u| \leq \tau + T} \psi''(u) \cdot \underbrace{\frac{1}{n} \sum_{i=1}^{n} \varphi_{\tau}(\boldsymbol{\Delta}^{\top} \boldsymbol{x}_i^{(t)}) \mathbb{1}(|(\boldsymbol{\beta}^{(t)*})^{\top} \boldsymbol{x}_i^{(t)}| \leq T)}_{(*)}
$$

It suffices to show that by fixing some $\tau, T > 0$, $(*) \ge C ||\Delta||_2^2$ w.p. at least $1 - e^{-C'n}$. In fact, we only need to consider the case $||\Delta||_2 = 1$. If the results hold when $||\Delta||_2 = 1$, then when $\|\mathbf{\Delta}\|_2 \in (0,1)$, since φ_{τ} is non-decreasing on τ , plugging in $\mathbf{\Delta}/\|\mathbf{\Delta}\|_2$ implies $(*) \ge C\|\mathbf{\Delta}\|_2^2$. So in the following analysis, we consider $||\mathbf{\Delta}||_2 = 1$.

Define

$$
Z_n = \sup_{\|\boldsymbol{\Delta}\|_2=1} \left\{ \frac{1}{n} \sum_{i=1}^n \varphi_\tau(\boldsymbol{\Delta}^\top \boldsymbol{x}_i^{(t)}) 1(|(\boldsymbol{\beta}^{(t)*})^\top \boldsymbol{x}_i^{(t)}| \leq T) - \mathbb{E}[\varphi_\tau(\boldsymbol{\Delta}^\top \boldsymbol{x}_i^{(t)}) 1(|(\boldsymbol{\beta}^{(t)*})^\top \boldsymbol{x}_i^{(t)}| \leq T)] \right\}.
$$

By bounded difference inequality (Corollary 2.21 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0)) or functional Hoeffding inequality (Theorem 3.26 in [Wainwright](#page-39-0) [\(2019\)](#page-39-0)),

$$
\mathbb{P}(Z_n \ge \mathbb{E}Z_n + \delta) \le \exp\{-Cn\delta^2/\tau^4\}.
$$
\n(59)

Denote $\{\epsilon_i\}_{i=1}^n$ as independent Rademacher variables. By symmetrization,

$$
\mathbb{E}Z_n \lesssim \mathbb{E}_{\mathbf{x}^{(t)},\epsilon} \left[\sup_{\|\mathbf{\Delta}\|_2=1} \left| \frac{1}{n} \sum_{i=1}^n \varphi_\tau (\mathbf{\Delta}^\top \mathbf{x}_i^{(t)}) \mathbb{1}(|(\boldsymbol{\beta}^{(t)*})^\top \mathbf{x}_i^{(t)}| \leq T) \cdot \epsilon_i \right| \right] \n\lesssim \tau \mathbb{E}_{\mathbf{x}^{(t)},\epsilon} \left[\sup_{\|\mathbf{\Delta}\|_2=1} \left| \frac{1}{n} \sum_{i=1}^n (\mathbf{\Delta}^\top \mathbf{x}_i^{(t)}) \epsilon_i \right| \right] \n\lesssim \tau \cdot \frac{1}{n} \mathbb{E}_{\mathbf{x}^{(t)},\epsilon} \sqrt{\sup_{\|\mathbf{\Delta}\|_2=1} \sum_{i=1}^n (\mathbf{\Delta}^\top \mathbf{x}_i^{(t)})^2 \epsilon_i^2} \n\lesssim \tau \cdot \frac{1}{\sqrt{n} \mathbb{E} \|\mathbf{x}_i^{(t)}\|_2^2}
$$
\n(61)

$$
\leq \tau \cdot \frac{n}{n} \sqrt{n} \mathbb{E} \| \mathbf{x}_i \|_2
$$
\n
$$
\lesssim \tau \sqrt{\frac{p}{n}},\tag{62}
$$

where (60) and (61) are due to Rademacher contraction inequality (equation (5.61) in [Wainwright](#page-39-0) [\(2019\)](#page-39-0), because the function $\varphi_{\tau}(\cdot) \mathbb{1}(|(\boldsymbol{\beta}^{(t)*})^{\top} \boldsymbol{x}_{i}^{(t)})$ $\left|\sum_{i=1}^{(k)}\right| \leq T$ is 2 τ -Lipschitz) and Jensen's inequality. On the other hand,

$$
\sup_{\|\mathbf{\Delta}\|_2=1} \mathbb{E}[\varphi_\tau(\mathbf{\Delta}^\top \boldsymbol{x}_i^{(t)}) \mathbb{1}(|(\boldsymbol{\beta}^{(t)*})^\top \boldsymbol{x}_i^{(t)}|] \leq T)]
$$

$$
\geq \sup_{\|\mathbf{\Delta}\|_{2}=1} \left\{ \mathbb{E}[(\mathbf{\Delta}^{\top}\mathbf{x}_{i}^{(t)})^{2} \cdot \mathbb{1}(|(\boldsymbol{\beta}^{(t)*})^{\top}\mathbf{x}_{i}^{(t)}| \leq T)] - \mathbb{E}[(\mathbf{\Delta}^{\top}\mathbf{x}_{i}^{(t)})^{2} \cdot \mathbb{1}(|\mathbf{\Delta}^{\top}\mathbf{x}_{i}^{(t)}| > \tau) \cdot \mathbb{1}(|(\boldsymbol{\beta}^{(t)*})^{\top}\mathbf{x}_{i}^{(t)}| \leq T)] \right\}
$$
\n
$$
\geq \sup_{\|\mathbf{\Delta}\|_{2}=1} \mathbb{E}(\mathbf{\Delta}^{\top}\mathbf{x}_{i}^{(t)})^{2} - \mathbb{P}(|(\boldsymbol{\beta}^{(t)*})^{\top}\mathbf{x}_{i}^{(t)}| > T) - \sup_{\|\mathbf{\Delta}\|_{2}=1} \sqrt{\mathbb{E}(\mathbf{\Delta}^{\top}\mathbf{x}_{i}^{(t)})^{4}} \cdot \sqrt{\mathbb{P}(|\mathbf{\Delta}^{\top}\mathbf{x}_{i}^{(t)}| > \tau)}
$$
\n
$$
\geq C - \exp\{-C'T^{2}/(\zeta^{(t)})^{2}\} - e^{-C''\tau^{2}}
$$
\n
$$
\geq C/2, \tag{63}
$$

when T and τ are large.

Putting [\(59\)](#page-106-2), [\(62\)](#page-106-3), and [\(63\)](#page-107-0) together, and setting $\delta =$ a small constant c, w.p. at least $1 - e^{-C''n}$, we have

$$
(*) \geq \sup_{\|\mathbf{\Delta}\|_2=1} \mathbb{E}[\varphi_\tau(\mathbf{\Delta}^\top \mathbf{x}_i^{(t)}) \mathbb{1}(|(\boldsymbol{\beta}^{(t)*})^\top \mathbf{x}_i^{(t)}|] \leq T)] - Z_n
$$

$$
\geq C/2 - C' \sqrt{\frac{p}{n}} - \delta
$$

$$
\geq C/4,
$$

when $n \geq p + \log T$.

(ii) If $\|\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}\|_2 > 1$: Denote $\boldsymbol{\Delta} = \boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}, \delta = \frac{1}{\|\boldsymbol{\Delta}\|_2^2}$ $\frac{1}{\|\mathbf{\Delta}\|_2}$. By convexity of $f^{(t)}$,

$$
f^{(t)}(\delta \beta + (1 - \delta) \beta^{(t)*}) \leq \delta f^{(t)}(\beta) + (1 - \delta) f^{(t)}(\beta^{(t)*}),
$$

which combining with part (i) implies that w.p. at least $1 - e^{-C''n}$,

$$
f^{(t)}(\boldsymbol{\beta}) - f^{(t)}(\boldsymbol{\beta}^{(t)*}) - \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})^{\top}(\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}) \ge \frac{f^{(t)}(\boldsymbol{\beta}^{(t)*} + \delta \boldsymbol{\Delta}) - f^{(t)}(\boldsymbol{\beta}^{(t)*}) - \delta \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})^{\top} \boldsymbol{\Delta}}{\delta}
$$

$$
\ge \delta C \|\boldsymbol{\Delta}\|_2^2
$$

$$
= C \|\boldsymbol{\Delta}\|_2,
$$

because $\|\delta\mathbf{\Delta}\|_2=1.$

E.4.4 Proof of Lemma [47](#page-101-1)

It is easy to see that

$$
f^{(t)}(\boldsymbol{\beta}) - f^{(t)}(\widetilde{\boldsymbol{\beta}}) - \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})^{\top}(\boldsymbol{\beta} - \widetilde{\boldsymbol{\beta}}) = \underbrace{f^{(t)}(\boldsymbol{\beta}) - f^{(t)}(\boldsymbol{\beta}^{(t)*}) - \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})^{\top}(\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*})}_{[1]} \\
 + \underbrace{f^{(t)}(\boldsymbol{\beta}^{(t)*}) - f^{(t)}(\widetilde{\boldsymbol{\beta}}) + \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})^{\top}(\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}^{(t)*})}_{[2]}.
$$
By Lemma [46,](#page-101-0) w.p. at least $1 - e^{-C''n}$,

$$
[1] \ge C \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}\|_2^2.
$$

And with some $\delta, \delta' \in [0, 1]$, by Assumption [4,](#page-21-0) w.p. at least $1 - e^{-C(p + \log T)}$,

$$
[2] \leq \nabla f^{(t)} (\delta \beta^{(t)*} + (1 - \delta) \widetilde{\beta})^{\top} (\beta^{(t)*} - \widetilde{\beta}) + \nabla f^{(t)} (\beta^{(t)*})^{\top} (\widetilde{\beta} - \beta^{(t)*})
$$

\n
$$
= (\widetilde{\beta} - \beta^{(t)*})^{\top} \nabla^2 f^{(t)} (\delta' [\delta \beta^{(t)*} + (1 - \delta) \widetilde{\beta}] + (1 - \delta') \beta^{(t)*})^{\top} (\widetilde{\beta} - \beta^{(t)*})
$$

\n
$$
\lesssim \frac{1}{n} \sum_{i=1}^n [(\widetilde{\beta} - \beta^{(t)*})^{\top} x_i^{(t)}]^2
$$

\n
$$
\lesssim ||\widetilde{\beta} - \beta^{(t)*}||_2^2,
$$

where the last inequality comes from the fact that $\|\hat{\Sigma}^{(t)}\|_2 \leq C$ w.p. at least 1 − $e^{-C'(p+\log T)}$ (by Lemma [19\)](#page-47-0). Putting all pieces together leads to the desired conclusion.

E.4.5 Proof of Lemma [48](#page-102-0)

Define $\widetilde{\theta}_{\boldsymbol{A}}^{(t)} = \boldsymbol{A}^\top \boldsymbol{A}^{(t)*} \boldsymbol{\theta}^{(t)*}$. It is easy to see that

$$
\|\mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \leq \|(\mathbf{A}\mathbf{A}^\top - \mathbf{I})\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \lesssim \|(\mathbf{A}^\perp)^\top \mathbf{A}^{(t)*}\|_2 \zeta^{(t)} \leq c.
$$

If $\|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 > 1$, then by Lemma [47,](#page-101-1) w.p. at least $1 - e^{-C(p + \log T)}$,

$$
f^{(t)}(\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)}) - f^{(t)}(\mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}) \ge -\|\nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_2 \cdot \|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}\|_2 + C\|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2
$$

$$
- C' ||A\widetilde{\theta}_{A}^{(t)} - A^{(t)*}\theta^{(t)*}||_{2}^{2}
$$

\n
$$
\geq (C - ||\nabla f^{(t)}(A^{(t)*}\theta^{(t)*})||_{2}) \cdot ||A\theta_{A}^{(t)} - A\widetilde{\theta}_{A}^{(t)}||_{2}
$$

\n
$$
- C ||A\widetilde{\theta}_{A}^{(t)} - A^{(t)*}\theta^{(t)*}||_{2} - C' ||A\widetilde{\theta}_{A}^{(t)} - A^{(t)*}\theta^{(t)*}||_{2}^{2}
$$

\n
$$
\geq \frac{1}{2}C ||A\theta_{A}^{(t)} - A\widetilde{\theta}_{A}^{(t)}||_{2} - Cc - C'c^{2}
$$

\n
$$
\geq \frac{1}{2}C ||A\theta_{A}^{(t)} - A^{(t)*}\theta^{(t)*}||_{2} - \frac{1}{2}Cc - Cc - C'c^{2}
$$

\n
$$
\geq 0,
$$

which is contradicted with the definition of $\theta^{(t)}_A$. Therefore, we must have $||A\theta^{(t)}_A A^{(t)*}\theta^{(t)*} \|_2 \leq 1$. By Lemma [47,](#page-101-1) w.p. at least $1 - e^{-C(p + \log T)}$,

$$
f^{(t)}(\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)}) - f^{(t)}(\mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}) \ge -\|\nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_2 \cdot \|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}\|_2 + C\|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}\|_2^2
$$

$$
- C' \|\widetilde{\mathbf{A}} \widetilde{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*} \theta^{(t)*} \|_{2}^{2}
$$

\n
$$
\geq -C'' \sqrt{\frac{p + \log T}{n}} + C \|\mathbf{A} \theta_{\mathbf{A}}^{(t)} - \mathbf{A} \widetilde{\theta}_{\mathbf{A}}^{(t)} \|_{2}^{2} - C' c^{2}.
$$

We know that $f^{(t)}(A\theta_A^{(t)}) - f^{(t)}(A\widetilde{\theta}_A^{(t)}) \ge 0$ by optimality, so we must have

$$
-C''\sqrt{\frac{p+\log T}{n}}+C\|\boldsymbol{A}\boldsymbol{\theta}^{(t)}_{\boldsymbol{A}}-\boldsymbol{A}\widetilde{\boldsymbol{\theta}}^{(t)}_{\boldsymbol{A}}\|_2^2-C'c^2\leq 0,
$$

i.e., $\|\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}-\boldsymbol{A}\widetilde{\boldsymbol{\theta}}_{\boldsymbol{A}}^{(t)}\|_2 \leq \sqrt{\frac{C'}{C}}$ $\frac{C'}{C}c + (C'')^{1/2}(\frac{p + \log T}{n})$ $\frac{\log T}{n}$)^{1/4} \leq a small constant c', w.p. at least $1 - e^{-C(p + \log T)}$. This implies that $\|\boldsymbol{\theta}_A^{(t)}\|_2 = \|A\boldsymbol{\theta}_A^{(t)}\|_2 \le \|A\boldsymbol{\theta}_A^{(t)} - A\widetilde{\boldsymbol{\theta}}_A^{(t)}\|_2 + \|A\widetilde{\boldsymbol{\theta}}_A^{(t)} - A\widetilde{\boldsymbol{\theta}}_A^{(t)}\|_2$ $A^{(t)*}\theta^{(t)*} \|_2 + \|A^{(t)*}\theta^{(t)*} \|_2 \leq C\zeta^{(t)},$ w.p. at least $1 - e^{-C(p + \log T)}$, which completes the proof.

E.4.6 Proof of Lemma [49](#page-102-1)

Define $\widetilde{\theta}_{\mathbf{A}}^{(t)} = \mathbf{A}^{\top} \widetilde{\mathbf{A}} \theta_{\widetilde{\mathbf{A}}}^{(t)}$. It is easy to see that

$$
\|\boldsymbol A\widetilde{\boldsymbol \theta}_{\boldsymbol A}^{(t)} - \widetilde{\boldsymbol A} \boldsymbol \theta_{\widetilde{\boldsymbol A}}^{(t)}\|_2 \leq \|(\boldsymbol A\boldsymbol A^\top - \boldsymbol I)\widetilde{\boldsymbol A} \boldsymbol \theta_{\widetilde{\boldsymbol A}}^{(t)}\|_2 \lesssim \|(\boldsymbol A^\perp)^\top \widetilde{\boldsymbol A}\|_2 \zeta^{(t)} \leq c.
$$

If $\|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 > 1$, then by Lemma [47,](#page-101-1) w.p. at least $1 - e^{-C(p + \log T)}$,

$$
f^{(t)}(\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)}) - f^{(t)}(\mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}) \ge -\|\nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_2 \cdot \|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}\|_2 + C\|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2
$$

$$
- C' || A \widetilde{\theta}_{A}^{(t)} - A^{(t)*} \theta^{(t)*} ||_2^2
$$

\n
$$
\geq (C - || \nabla f^{(t)} (A^{(t)*} \theta^{(t)*} ||_2) \cdot || A \theta_{A}^{(t)} - A \widetilde{\theta}_{A}^{(t)} ||_2
$$

\n
$$
- C || A \widetilde{\theta}_{A}^{(t)} - A^{(t)*} \theta^{(t)*} ||_2 - C' || A \widetilde{\theta}_{A}^{(t)} - A^{(t)*} \theta^{(t)*} ||_2^2
$$

\n
$$
\geq \frac{1}{2} C || A \theta_{A}^{(t)} - A \widetilde{\theta}_{A}^{(t)} ||_2 - C || A \widetilde{\theta}_{A}^{(t)} - A \theta_{\widetilde{A}}^{(t)} ||_2 - C || \widetilde{A} \theta_{\widetilde{A}}^{(t)} - A^{(t)*} \theta^{(t)*} ||_2
$$

\n
$$
- C' || A \widetilde{\theta}_{A}^{(t)} - A \widetilde{\theta}_{\widetilde{A}}^{(t)} ||_2^2 - C' || \widetilde{A} \theta_{\widetilde{A}}^{(t)} - A^{(t)*} \theta^{(t)*} ||_2^2
$$

\n
$$
\geq \frac{1}{2} C || A \theta_{A}^{(t)} - A \widetilde{\theta}_{A}^{(t)} ||_2 - Cc - C' c^2
$$

\n
$$
\geq \frac{1}{2} C || A \theta_{A}^{(t)} - A^{(t)*} \theta^{(t)*} ||_2 - \frac{1}{2} Cc - Cc - C' c^2
$$

\n
$$
> 0,
$$

which is contradicted with the definition of $\theta^{(t)}_A$. Therefore, we must have $||A\theta^{(t)}_A A^{(t)*}\theta^{(t)*} \|_2 \leq 1$. By Lemma [47,](#page-101-1) w.p. at least $1 - e^{-C(p + \log T)}$,

$$
f^{(t)}(\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)}) - f^{(t)}(\mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}) \ge -\|\nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_{2} \cdot \|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}\|_{2} + C\|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}\|_{2}^{2}
$$

$$
- C'\|\mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2}^{2}
$$

$$
\ge -\|\nabla f^{(t)}(\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*})\|_{2} \cdot \|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}\|_{2} + C\|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)}\|_{2}^{2}
$$

$$
-\frac{1}{2}C\|\boldsymbol{A}\widetilde{\boldsymbol{\theta}}_{\boldsymbol{A}}^{(t)}-\widetilde{\boldsymbol{A}}\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)}\|_{2}^{2}-C''\|\widetilde{\boldsymbol{A}}\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)}-\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_{2}^{2} \geq -C''\sqrt{\frac{p+\log T}{n}}+\frac{1}{2}C\|\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}-\boldsymbol{A}\widetilde{\boldsymbol{\theta}}_{\boldsymbol{A}}^{(t)}\|_{2}^{2}-4Cc^{2}.
$$

We know that $f^{(t)}(A\theta_A^{(t)}) - f^{(t)}(A\widetilde{\theta}_A^{(t)}) \ge 0$ by optimality, so we must have

$$
-C''\sqrt{\frac{p+\log T}{n}}+C\|\boldsymbol{A}\boldsymbol{\theta}^{(t)}_{\boldsymbol{A}}-\boldsymbol{A}\widetilde{\boldsymbol{\theta}}^{(t)}_{\boldsymbol{A}}\|_2^2-C'c^2\leq 0,
$$

i.e., $\|\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}-\boldsymbol{A}\widetilde{\boldsymbol{\theta}}_{\boldsymbol{A}}^{(t)}\|_2 \leq \sqrt{\frac{C'}{C}}$ $\frac{C'}{C}c + (C'')^{1/2}(\frac{p + \log T}{n})$ $\frac{\log T}{n}$)^{1/4} \leq a small constant c', w.p. at least $1 - e^{-C(p + \log T)}$. This implies that $\|\boldsymbol{\theta}_A^{(t)}\|_2 = \|A\boldsymbol{\theta}_A^{(t)}\|_2 \le \|A\boldsymbol{\theta}_A^{(t)} - A\widetilde{\boldsymbol{\theta}}_A^{(t)}\|_2 + \|A\widetilde{\boldsymbol{\theta}}_A^{(t)} - A\widetilde{\boldsymbol{\theta}}_A^{(t)}\|_2$ $A^{(t)*}\theta^{(t)*} \|_2 + \|A^{(t)*}\theta^{(t)*} \|_2 \leq C\zeta^{(t)}, \text{ w.p. at least } 1 - e^{-C(p + \log T)}, \text{ which completes the }$ proof.

E.4.7 Proof of Lemma [50](#page-102-2)

Similar to the proof of Lemma [32,](#page-51-0) by the optimality of $\theta_A^{(t)}$,

$$
\boldsymbol{A}^\top(\nabla f^{(t)}(\widetilde{\boldsymbol{A}}\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)})+\widetilde{\boldsymbol{\Sigma}}^{(t)}(\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}-\widetilde{\boldsymbol{A}}\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)}))=\boldsymbol{0},
$$

where $\widetilde{\Sigma}^{(t)} = \frac{1}{n}$ $\frac{1}{n}\sum_{i=1}^n\boldsymbol{x}_i^{(t)}$ $\bm{x}_i^{(t)}(\bm{x}_i^{(t)})$ $\psi^{(t)}_i)^\top \psi''\big((\boldsymbol{x}_i^{(t)}$ $(\begin{matrix} a(t) \end{matrix})^{\top} \widetilde{\bm{A}} \bm{\theta}^{(t)}_{\widetilde{\bm{A}}} + \delta(\bm{x}^{(t)}_i)$ $(\hat{i}^{(t)})^{\top} (\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \widetilde{\mathbf{A}}\boldsymbol{\theta}_{\widetilde{\mathbf{A}}}^{(t)})), \delta \in [0,1].$ With similar arguments used in the proof of Lemma [46,](#page-101-0) it can be shown that $\lambda_{\min}(\tilde{\Sigma}^{(t)}) > 0$ w.p. at least $1 - e^{-C(p + \log T)}$. The remaining steps are the same as the steps in the proof of Lemma [32.](#page-51-0)

E.4.8 Proof of Lemma [51](#page-102-3)

The proof is almost the same as the proof of Lemma [50,](#page-102-2) so we omit it here.

E.4.9 Proof of Lemma [52](#page-102-4)

If $||\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}||_2 > 1$, by Lemmas [44](#page-101-2) and [46,](#page-101-0) w.p. at least $1 - e^{-C'(p + \log T)}$, for any $\widetilde{\boldsymbol{A}} \in \mathcal{O}^{p \times r},$

$$
f^{(t)}(\boldsymbol{A}^{(t)}\boldsymbol{\theta}^{(t)})+\frac{\lambda}{\sqrt{n}}\|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top-\widetilde{\boldsymbol{A}}(\widetilde{\boldsymbol{A}})^\top\|_2-f^{(t)}(\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*})-\frac{\lambda}{\sqrt{n}}\|\boldsymbol{A}^{(t)*}(\boldsymbol{A}^{(t)*})^\top-\widetilde{\boldsymbol{A}}(\widetilde{\boldsymbol{A}})^\top\|_2
$$

$$
\geq \nabla f^{(t)} (\mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*})^{\top} (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*}) - 2 \frac{\lambda}{\sqrt{n}} + C \| \mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} \|_2
$$
\n
$$
\geq \left(C - C' \sqrt{\frac{p + \log T}{n}} \right) \| \mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} \|_2 - 2 \frac{\lambda}{\sqrt{n}}
$$
\n
$$
\geq \frac{1}{2} C - \frac{1}{4} C
$$
\n
$$
> 0,
$$

which contradicts to the definitions of $\mathbf{A}^{(t)}$ and $\boldsymbol{\theta}^{(t)}$. Hence $\|\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \leq 1$, w.p. at least $1 - e^{-C'(p + \log T)}$. Then by Lemmas [44](#page-101-2) and [46](#page-101-0) again, w.p. at least $1 - e^{-C'(p + \log T)}$, for any $\widetilde{\mathbf{A}} \in \mathcal{O}^{p \times r}$,

$$
f^{(t)}(\boldsymbol{A}^{(t)}\boldsymbol{\theta}^{(t)})+\frac{\lambda}{\sqrt{n}}\|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top-\widetilde{\boldsymbol{A}}(\widetilde{\boldsymbol{A}})^\top\|_2-f^{(t)}(\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*})-\frac{\lambda}{\sqrt{n}}\|\boldsymbol{A}^{(t)*}(\boldsymbol{A}^{(t)*})^\top-\widetilde{\boldsymbol{A}}(\widetilde{\boldsymbol{A}})^\top\|_2
$$

$$
\geq \nabla f^{(t)} (\mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*})^{\top} (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*}) - 2 \frac{\lambda}{\sqrt{n}} + C \| \mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} \|_2^2
$$

\n
$$
\geq C \| \mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} \|_2^2 - C' \sqrt{\frac{p + \log T}{n}} \| \mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} \|_2 - 2 \frac{\lambda}{\sqrt{n}}.
$$

By the optimality of $\mathbf{A}^{(t)}\theta^{(t)}$, we must have the RHS of above ≤ 0 w.p. at least 1 – $e^{-C'(p+\log T)},$ which entails that $\|\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \lesssim \sqrt{\frac{p+\log T}{n}} + \lambda^{1/2}n^{-1/4} \leq \text{a small}$ constant c, which completes the proof.

Appendix F. Proofs for Non-linear Regression Models

F.1 Lemmas

Lemma 53 Suppose Assumptions [1,](#page-10-0) [6,](#page-22-0) and [7](#page-22-1) hold. Then w.p. at least $1 - e^{-C(p + \log T)}$, for any β , $\widetilde{\beta} \in \mathbb{R}^p$, for any $t \in S$,

$$
f^{(t)}(\boldsymbol{\beta}) - f^{(t)}(\widetilde{\boldsymbol{\beta}}) - \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})^{\top}(\boldsymbol{\beta} - \widetilde{\boldsymbol{\beta}}) \ge C' ||\boldsymbol{\beta} - \widetilde{\boldsymbol{\beta}}||_2^2 - C'' ||\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}^{(t)*}||_2^2.
$$

Lemma 54 Suppose Assumptions [1,](#page-10-0) [6,](#page-22-0) and γ hold. Then when $\delta \leq C$ with some constant $C>0,$

$$
\|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2 \lesssim \delta,
$$

w.p. at least $1 - \exp\{-Cn\delta^2 + C'p\}$. By taking $t \approx \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$, this implies that:

- (i) $\max_{t \in S} \|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2 \lesssim \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$, w.p. at least $1 - e^{-C(p + \log T)}$;
- $(ii) \|\nabla f^{(0)}(\boldsymbol{\beta}^{(0)*})\|_2 \lesssim \sqrt{\frac{p}{n_0}}, \ w.p. \ \text{at least} \ 1 e^{-C(p + \log T)}.$

Lemma 55 Suppose Assumptions [1,](#page-10-0) [6,](#page-22-0) and [7](#page-22-1) hold. Then when $t \leq C$ with some constant $C > 0$, for any $S \subseteq [T]$,

$$
\|\{\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\}_{t\in S}\|_2 \lesssim \sqrt{\frac{p+|S|+\delta}{n}},
$$

w.p. at least $1 - e^{-C'\delta}$.

Lemma 56 Suppose Assumptions [1,](#page-10-0) [6,](#page-22-0) and [7](#page-22-1) hold. Then w.p. at least $1 - e^{-C'(p + \log T)}$, for all $t \in S$, for all $\mathbf{A}, \widetilde{\mathbf{A}} \in \mathcal{O}^{p \times r}$ with $\|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} \|_2$, $\|\widetilde{\mathbf{A}} \boldsymbol{\theta}_{\widetilde{\mathbf{A}}}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} \|_2 \leq c'$, where $c' > 0$ is a small constant, we have

$$
\|\boldsymbol{A}\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}-\widetilde{\boldsymbol{A}}\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)}\|_{2}\lesssim\|\boldsymbol{A}\boldsymbol{A}^{\top}-\widetilde{\boldsymbol{A}}(\widetilde{\boldsymbol{A}})^{\top}\|_{2}(\|\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)}\|_{2}+\|\nabla f^{(t)}(\widetilde{\boldsymbol{A}}\boldsymbol{\theta}_{\widetilde{\boldsymbol{A}}}^{(t)})\|_{2}).
$$

Lemma 5[7](#page-22-1) Suppose Assumptions [1,](#page-10-0) [6,](#page-22-0) and γ hold. Then w.p. at least $1 - e^{-C'(p + \log T)}$, for all $t \in S$, for all $\mathbf{A}, \widetilde{\mathbf{A}} \in \mathcal{O}^{p \times r}$ with $\|\mathbf{A}\boldsymbol{\theta}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \leq c'$, where $c' > 0$ is a small constant, we have

$$
\|\boldsymbol A\boldsymbol \theta^{(t)}_{\boldsymbol A} - \boldsymbol A^{(t)*}\boldsymbol \theta^{(t)*}\|_2 \lesssim \|\boldsymbol A\boldsymbol A^\top - \boldsymbol A^{(t)*}(\boldsymbol A^{(t)*})^\top\|_2 \zeta^{(t)} + \sqrt{\frac{r+\log T}{n}}.
$$

Lemma 58 Suppose Assumptions [1,](#page-10-0) [6,](#page-22-0) and [7](#page-22-1) hold. Then w.p. at least $1 - e^{-C'(p + \log T)}$, for all $t \in S$, for all $\boldsymbol{A} \in \mathcal{O}^{p \times r}$ with $\|\boldsymbol{A}\boldsymbol{A}^{\top} - \boldsymbol{A}^{(t)*}(\boldsymbol{A}^{(t)*})^{\top}\|_2 \leq c(\zeta^{(t)})^{-1}$, where $c > 0$ is a small constant, we have

- (i) $\|\theta_A^{(t)}\|_2 \le C'' \zeta^{(t)};$
- (ii) $||A\theta_A^{(t)} A^{(t)*}\theta^{(t)*}||_2 \leq c'$ with a small constant $c' > 0$.

Lemma 59 Suppose Assumptions [1,](#page-10-0) [6,](#page-22-0) and [7](#page-22-1) hold. Assume there exists $\tilde{A} \in \mathcal{O}^{p \times r}$ such that $\|\widetilde{A}\boldsymbol{\theta}_{\widetilde{A}}^{(t)} - A^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \leq c$, where $c > 0$ is a small constant. Then w.p. at least $1 - e^{-C'(p + \log T)}$, for all $t \in S$, for all $\boldsymbol{A} \in \mathcal{O}^{p \times r}$ with $\|\boldsymbol{A}\boldsymbol{A}^{\top} - \widetilde{\boldsymbol{A}}(\widetilde{\boldsymbol{A}})^{\top}\|_2 \leq c'(\zeta^{(t)})^{-1}$, where $c' > 0$ is a small constant, we have

- (i) $\|\theta_A^{(t)}\|_2 \leq C'' \zeta^{(t)};$
- (ii) $||A\theta^{(t)}_A A^{(t)*}\theta^{(t)*}||_2 \leq c''$ with a small constant $c'' > 0$.

Lemma 60 Suppose Assumptions [1,](#page-10-0) [6,](#page-22-0) and [7](#page-22-1) hold. Let $n \geq C\lambda^2$ with a sufficiently large $C. \quad Given \ \widetilde{\bm{A}} \ \in \ \mathcal{O}^{p \times r}, \ \ denote \ \left(\bm{A}^{(t)}, \bm{\theta}^{(t)}\right) \ = \ \argmin_{\bm{A} \in \mathcal{O}^{p \times r}, \bm{\theta} \in \mathbb{R}^r} \{f^{(t)}(\bm{A\theta}) + \frac{\lambda}{\sqrt{N}}\}$ $\frac{N}{n}\|A A^\top \widetilde{\mathbf{A}}(\widetilde{\mathbf{A}})^{\top} \|_{2}$. Then w.p. at least $1 - e^{-C(p + \log T)}$, for any $\widetilde{\mathbf{A}} \in \mathcal{O}^{p \times r}$, we have $\|\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)}\|_{2}$. $\boldsymbol{\beta}^{(t)*} \|_2 \leq c \text{ with a small constant } c > 0.$

F.2 Proof of Theorem [10](#page-23-0)

For the penalized ERM, the proof logic is almost the same as the logic underlying the proofs of Proposition [43](#page-52-0) and Theorem [1.](#page-12-0) Similar to the proof of Theorem [9,](#page-22-2) we only point out the differences and skip the details. We need to bound $\|\overline{A}(\overline{A})^{\top} - \overline{A}(\overline{A})^{\top}\|_2$ first. Then we know that $\|\widehat{\vec{A}}(\widehat{\vec{A}})^{\top} - \overline{A}(\overline{A})^{\top}\|_2$ and $\|\widehat{\vec{A}}\theta_{\widehat{\vec{A}}}\^{(t)}\|_2$ $-\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2$ are small w.h.p. Then a direct application of Lemma [59](#page-112-0) implies that $\|\widehat{A}\boldsymbol{\theta}_{\widehat{\lambda}}^{(t)*} - A^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2$ is small w.h.p. Finally, applying \overline{A} Lemma [56,](#page-111-0) we have $\|\widehat{\overline{A}} \theta_{\widehat{\overline{A}}}^{(t)*}$ $-\overline{A}\boldsymbol{\theta}_{\overline{A}}^{(t)}\|_2 \lesssim \|\widehat{\overline{A}}(\widehat{\overline{A}})^\top - \overline{A}(\overline{A})^\top \|_2 (\|\boldsymbol{\theta}_{\overline{A}}^{(t)}\|_2)$ $\frac{d^{(t)}}{A} \|_2 + \| \nabla f^{(t)}(\overline{A} \boldsymbol{\theta}_{\overline{A}}^{(t)}) \|_2) \lesssim$ $\|\widehat{\vec{A}}(\widehat{\vec{A}})^{\top} - \overline{A}(\overline{A})^{\top}\|_2 \zeta^{(t)}$ w.h.p. which gives us the ideal bound for $\|\widehat{\vec{A}} \theta_{\widehat{\vec{A}}}^{(t)*}$ $-\,\overline{A}\theta^{(t)}_{\overline{A}}\|_2$ for all $t \in S$. (i) In the proof of Proposition [43,](#page-52-0) for $t \in \mathcal{A}_1$, we know $||\mathbf{A}^{(t)}\boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}||_2 \leq c$ with a

small constant $c > 0$ w.h.p. by Lemma [60.](#page-112-1) Then by Lemmas [53,](#page-111-1) [55,](#page-111-2) and [57,](#page-112-2) w.p. at least $1 - e^{-C'(r + \log T)},$

$$
\frac{1}{T} \sum_{t \in \mathcal{A}_1} \left[f^{(t)}(\boldsymbol{A}^{(t)} \boldsymbol{\theta}^{(t)}) - f^{(t)}(\overline{\boldsymbol{A}} \boldsymbol{\theta}_{\overline{\boldsymbol{A}}}^{(t)}) \right]
$$

$$
\geq -\frac{1}{T}\left|\left\langle \{\nabla f^{(t)}(\mathbf{A}^{(t)*}\theta^{(t)*})\}_{t\in A_1}, \{\mathbf{A}^{(t)}\theta^{(t)} - \hat{\mathbf{A}}\theta_{\hat{\mathbf{A}}}^{(t)} + \hat{\mathbf{A}}\theta_{\hat{\mathbf{A}}}^{(t)} - \overline{\mathbf{A}}\theta_{\mathbf{A}}^{(t)}\}_{t\in A_1}\right\rangle\right| \n+ \frac{C_1}{T}\|\{\mathbf{A}^{(t)}\theta^{(t)} - \overline{\mathbf{A}}\theta_{\mathbf{A}}^{(t)}\}_{t\in A_1}\|_{\mathrm{F}}^2 - \frac{C_2}{T}\|\{\mathbf{A}^{(t)*}\theta^{(t)*} - \overline{\mathbf{A}}\theta_{\mathbf{A}}^{(t)}\}_{t\in A_1}\|_{\mathrm{F}}^2 \n\geq -\frac{C}{T}\sum_{t\in A_1}\sqrt{\frac{p + \log T}{n}} \cdot \zeta^{(t)}(\|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \hat{\mathbf{A}}(\hat{\mathbf{A}})^\top\|_2 + \|\hat{\mathbf{A}}(\hat{\mathbf{A}})^\top - \overline{\mathbf{A}}\mathbf{A}^\top\|_2) \n+ \frac{C_1}{T}\|\{\mathbf{A}^{(t)}\theta^{(t)} - \overline{\mathbf{A}}\theta_{\mathbf{A}}^{(t)}\}_{t\in A_1}\|_{\mathrm{F}}^2 - C\left(h^2 \cdot \frac{1}{T}\sum_{t\in A_1}(\zeta^{(t)})^2 + \frac{r + \log T}{n}\right) \n\geq -C\left(h^2 \cdot \frac{1}{T}\sum_{t\in A_1}(\zeta^{(t)})^2 + \frac{r + \log T}{n}\right) - \frac{C'}{T}\sum_{t\in A_1}\sqrt{\frac{p + \log T}{n}} \cdot \zeta^{(t)}\sqrt{r}\|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top - \widehat{\mathbf{A}}(\hat{\mathbf{A}})^\top\|_2.
$$

Therefore the calculations in [\(47\)](#page-95-0) are still correct under non-linear regression model [\(6\)](#page-22-3). (ii) In the proof of Proposition [43,](#page-52-0) for $t \in \mathcal{A}_2$, we used the fact that w.p. at least 1 − $e^{-C(r+\log T)},$

$$
\|\mathbf{A}^{(t)}\boldsymbol{\theta}_{\mathbf{A}^{(t)}}^{(t)} - \widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)}\|_{2} \lesssim \|\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^{\top} - \widehat{\mathbf{A}}(\widehat{\mathbf{A}})^{\top}\|_{2}(\|\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)}\|_{2} + \|\nabla f^{(t)}(\widehat{\mathbf{A}}\boldsymbol{\theta}_{\widehat{\mathbf{A}}}^{(t)})\|_{2}).\tag{64}
$$

Here this still holds because

- $||A^{(t)}\theta_{A^{(t)}}^{(t)} A^{(t)*}\theta^{(t)*}||_2 \leq c$ with a small constant $c > 0$, by Lemma [60;](#page-112-1)
- For $t \in \mathcal{A}_2$, we know that $||\mathbf{A}^{(t)}(\mathbf{A}^{(t)})^\top \overline{\mathbf{A}}(\overline{\mathbf{A}})^\top ||_2 \leq \frac{c}{\sqrt{r}} \leq c'(\zeta^{(t)})^{-1}$ with a small constant $c' > 0$. Then by Lemma [59,](#page-112-0) $\|\boldsymbol{\theta}_{\simeq}^{(t)}\|$ $\frac{d}{d\hat{A}}\|_2 \leq C\zeta^{(t)}$ and $\|\widehat{\overline{A}}\theta_{\widehat{\overline{A}}}^{(t)}\|_2$ $-\bm{A}^{(t)*} \bm{\theta}^{(t)*} \|_2 \leq Cc'$ with a small constant $c' > 0$.

Then (64) holds by Lemma [56.](#page-111-0)

Finally, Lemma [39](#page-52-1) still holds for non-linear regression model [\(6\)](#page-22-3). Note that in Lemma [39,](#page-52-1) (i) only requires the maximum eigenvalue of Hessian to be upper bounded when the evaluation point is close to the true $\beta^{(t)*}$ (see Lemma E.3 in [Duan and Wang](#page-36-0) [\(2022\)](#page-36-0)), which is true by Assumptions [1](#page-10-0) and [6](#page-22-0) (see our analysis in the proof of Lemma [56\)](#page-111-0). By Lemma E.2 in [Duan and Wang](#page-36-0) [\(2022\)](#page-36-0), (ii) and (iii) require the minimum eigenvalue of Hessian to be lower bounded when $||\boldsymbol{\beta}^{(t)} - \boldsymbol{\beta}^{(t)*}||_2 \le$ some constant C w.h.p. and $\gamma/\sqrt{n_0} \le C$, both of which are true.

For the spectral method, the proof is almost the same as the proof of Theorem [7,](#page-17-0) hence we do not repeat it here..

F.3 Proof of Theorem [18](#page-46-0)

We only outline the proof when Algorithm [4](#page-45-0) is coupled with Algorithm [1.](#page-11-0) If Algorithm 4 is coupled with Algorithm [2,](#page-17-1) we can follow the similar argument with the proof of Theorem [7.](#page-17-0)

.

Denote
$$
\eta = r \sqrt{\frac{p}{nT}} + \sqrt{r}h + \sqrt{r} \sqrt{\frac{r + \log T}{n}} + \frac{|S^c|}{T} \cdot r \cdot \frac{\lambda}{\sqrt{n}}
$$

(i) When $\eta \leq C \sqrt{\frac{p}{n_0}}$: since $\|\widehat{A}(\widehat{A})^{\top} - A^{(0)*}(A^{(0)*})^{\top}\|_2 \leq a$ small constant c, by Lemma [60,](#page-112-1) w.p. at least $1 - e^{-C(p + \log T)}$, $\|\widehat{A}\boldsymbol{\theta}_{\widehat{A}}^{(t)} - A^{(0)*}\boldsymbol{\theta}^{(0)*}\|_2 \leq a$ small constant c'. Then \boldsymbol{A} by Lemma [57](#page-112-2) and Theorem [10,](#page-23-0) w.p. at least $1 - e^{-C(r + \log T)}$, $\|\hat{\vec{A}}\theta_{\hat{\vec{A}}}^{(t)}\|$ $- \, A^{(0) *} \theta^{(0) *} \|_2 \, \lesssim$ $\|\overline{A}(\overline{A})^{\top} - A^{(0)*}(A^{(0)*})^{\top}\|_2 \lesssim \eta$. As we commented in the proof of Theorem [10,](#page-23-0) Lemma $42.(\text{i})$ $42.(\text{i})$ still holds for non-linear regression model (6) because the maximum eigenvalue of Hessian is upper bounded w.h.p. when the evaluation point is close to the true $\beta^{(t)*}$ (ii) When $\eta > C\sqrt{\frac{p}{n_0}}$: the result comes from Lemma [42.](#page-52-2)(ii), which still holds when $\gamma/\sqrt{n_0} \lesssim$ some constant C because of Lemma E.2 in [Duan and Wang](#page-36-0) [\(2022\)](#page-36-0).

F.4 Proofs of Lemmas

F.4.1 PROOF OF LEMMA [53](#page-111-1)

First,

$$
f^{(t)}(\boldsymbol{\beta}) - f^{(t)}(\widetilde{\boldsymbol{\beta}}) - \nabla f^{(t)}(\widetilde{\boldsymbol{\beta}})^\top (\boldsymbol{\beta} - \widetilde{\boldsymbol{\beta}}) \geq \underbrace{f^{(t)}(\boldsymbol{\beta}) - f^{(t)}(\boldsymbol{\beta}^{(t)*}) - \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})^\top (\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*})}_{[1]} + \underbrace{f^{(t)}(\boldsymbol{\beta}^{(t)*}) - f^{(t)}(\widetilde{\boldsymbol{\beta}}) - \nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})^\top (\boldsymbol{\beta}^{(t)*} - \widetilde{\boldsymbol{\beta}})}_{[2]}.
$$

By Taylor expansion,

$$
[1] = \left\langle \int_0^1 \nabla f^{(t)} (\delta \beta + (1 - \delta) \beta^{(t)*}) dt - \nabla f^{(t)} (\beta^{(t)*}), \beta - \beta^{(t)*} \right\rangle
$$

=
$$
\frac{2}{n} \sum_{i=1}^n \int_0^1 [g(\delta(\mathbf{x}_i^{(t)})^\top \beta + (1 - \delta)(\mathbf{x}_i^{(t)})^\top \beta^{(t)*}) - g((\mathbf{x}_i^{(t)})^\top \beta^{(t)*})] g'((\mathbf{x}_i^{(t)})^\top \delta \beta + (1 - \delta) \beta^{(t)*}) d\delta
$$

$$
\cdot \, [(\boldsymbol{x}_i^{(t)})^\top (\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*})] - \frac{2}{n} \sum_{i=1}^n \epsilon_i^{(t)} \cdot \int_0^1 [g' (\delta (\boldsymbol{x}_i^{(t)})^\top \boldsymbol{\beta} + (1-\delta) (\boldsymbol{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*}) - g' ((\boldsymbol{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*})] \mathrm{d}\delta
$$

$$
\begin{split}\n& \frac{\left\{\left[\left(\boldsymbol{x}_{i}^{(t)}\right)^{\top}\left(\boldsymbol{\beta}-\boldsymbol{\beta}^{(t)*}\right)\right]\right\}}{\left\|\boldsymbol{x}_{i}\right\|^{2}} \\
&= \frac{2}{n} \sum_{i=1}^{n} \int_{0}^{1} \underbrace{g'((\boldsymbol{x}_{i}^{(t)})^{\top}\boldsymbol{\beta}_{i})} \cdot \underbrace{g'(\delta(\boldsymbol{x}_{i}^{(t)})^{\top}\boldsymbol{\beta}+(1-\delta)(\boldsymbol{x}_{i}^{(t)})^{\top}\boldsymbol{\beta}^{(t)*})}_{\geq C} \delta d\delta \cdot \left[(\boldsymbol{x}_{i}^{(t)})^{\top}(\boldsymbol{\beta}-\boldsymbol{\beta}^{(t)*})\right]^{2} \\
&- \frac{2}{n} \sum_{i=1}^{n} \epsilon_{i}^{(t)} \cdot \int_{0}^{1} \int_{0}^{1} g''(w(\delta(\boldsymbol{x}_{i}^{(t)})^{\top}\boldsymbol{\beta}+(1-\delta)(\boldsymbol{x}_{i}^{(t)})^{\top}\boldsymbol{\beta}^{(t)*}) + (1-w)(\boldsymbol{x}_{i}^{(t)})^{\top}\boldsymbol{\beta}^{(t)*}) \delta d\delta dw \\
& \cdot \left[(\boldsymbol{x}_{i}^{(t)})^{\top}(\boldsymbol{\beta}-\boldsymbol{\beta}^{(t)*})\right]^{2}\n\end{split}
$$

$$
\geq C(\boldsymbol{\beta}-\boldsymbol{\beta}^{(t)\ast})^{\top}\widehat{\mathbf{\Sigma}}^{(t)}(\boldsymbol{\beta}-\boldsymbol{\beta}^{(t)\ast})-\frac{2}{n}\sum_{i=1}^n\epsilon_i^{(t)}b_i^{(t)}[(\boldsymbol{x}^{(t)}_i)^{\top}(\boldsymbol{\beta}-\boldsymbol{\beta}^{(t)\ast})]^2,
$$

where $b_i^{(t)} = \int_0^1 \int_0^1 g''(w(\delta(\bm{x}_i^{(t)}))$ $\bm{f}_i^{(t)})^\top \bm{\beta} + (1-\delta) (\bm{x}_i^{(t)})$ $\mathbf{f}_{i}^{(t)})^{\top} \boldsymbol{\beta}^{(t)\ast}) + (1-w)(\boldsymbol{x}_{i}^{(t)})$ $\delta_t^{(t)}$)[⊤]β^{(t)*})δdδdw and $|b_i^{(t)}\>$ $|i^{(t)}_{i}| \leq C'$. Therefore, by Hölder's inequality,

$$
\begin{aligned} \left| \frac{2}{n} \sum_{i=1}^n \epsilon_i^{(t)} b_i^{(t)} [(\bm{x}_i^{(t)})^\top (\bm{\beta} - \bm{\beta}^{(t)*})]^2 \right| &\leq \frac{2C'}{n} \sum_{i=1}^n |\epsilon_i^{(t)}| [(\bm{x}_i^{(t)})^\top (\bm{\beta} - \bm{\beta}^{(t)*})]^2 \\ &\leq \frac{2C'}{n} \sup_{\|\bm{u}\|_2 \leq 1} \left\| \sum_{i=1}^n |\epsilon_i^{(t)}| \cdot [(\bm{x}_i^{(t)})^\top \bm{u}]^2 \right\|_2 \cdot \|\bm{\beta} - \bm{\beta}^{(t)*}\|_2^2. \end{aligned}
$$

Suppose $\{u_j\}_{j=1}^N$ is a 1/64-packing of the unit ball in \mathbb{R}^p , with $N \lesssim 129^p$. By standard arguments,

$$
\frac{1}{n}\sup_{\|\mathbf{u}\|_2\leq 1}\left\|\sum_{i=1}^n|\epsilon_i^{(t)}|\cdot [(\mathbf{x}_i^{(t)})^\top\mathbf{u}]^2\right\|_2\lesssim \frac{1}{n}\max_{j\in[N]}\left\{\sum_{i=1}^n|\epsilon_i^{(t)}|\cdot [(\mathbf{x}_i^{(t)})^\top\mathbf{u}_j]^2\right\}.
$$

It is easy to see that for any $\delta > 0$,

$$
\mathbb{P}\left(|\epsilon_i^{(t)}|\cdot [(\boldsymbol{x}_i^{(t)})^\top \boldsymbol{u}_j]^2 - \mathbb{E}\{|\epsilon_i^{(t)}|\cdot [(\boldsymbol{x}_i^{(t)})^\top \boldsymbol{u}_j]^2\} > \delta\right) \leq \mathbb{P}\left(|\epsilon_i^{(t)}|\cdot [(\boldsymbol{x}_i^{(t)})^\top \boldsymbol{u}_j]^2 > \delta\right) \\
\leq \mathbb{P}(|\epsilon_i^{(t)}| > \delta^{1/3}) + \mathbb{P}(|(\boldsymbol{x}_i^{(t)})^\top \boldsymbol{u}_j| > \delta^{1/3})
$$

 $\lesssim \exp\{-C\delta^{2/3}\},$ and $\mathbb{E}\left\{\left|\epsilon_i^{(t)}\right.\right\}$ $_{i}^{(t)}|[(\boldsymbol{x}_{i}^{(t)}% ,\boldsymbol{x}_{i}^{(t)}% ,\boldsymbol{x}%)]$ $\sum\limits_{i}^{(t)})^{\top}\bm{u}_j]^2-\mathbb{E}[|\epsilon_i^{(t)}|$ $_{i}^{\left(t\right) }||(\boldsymbol{x}_{i}^{\left(t\right) }% ,\boldsymbol{x}_{i}^{\left(t\right) })||(\boldsymbol{x}_{i}^{\left(t\right) })||(\bold$ $\{(\epsilon_i^{(t)})^\top \bm u_j]^2]\}^2 \cdot \mathbb{1}\big(|\epsilon_i^{(t)}\big|$ $\binom{t}{i} |[({\boldsymbol{x}}_i^{(t)}% ,\boldsymbol{x}^{(t)}_{i}]]| \leq 1$ $\sum\limits_{i}^{(t)})^{\top}\bm{u}_j]^2-\mathbb{E}[|\epsilon_i^{(t)}|$ $_{i}^{\left(t\right) }||(\boldsymbol{x}_{i}^{\left(t\right) }% ,\boldsymbol{x}_{i}^{\left(t\right) })||(\boldsymbol{x}_{i}^{\left(t\right) })||(\bold$ $\left\{ \begin{matrix} (t) \\ i \end{matrix} \right\}^{\top}$ **u**_j]²] < 0) $\left\} \leq$ $\sqrt{\mathbb{E}\left\{\left|\epsilon_i^{(t)}\right.\right\}}$ $_i^{(t)}|^2[(\boldsymbol{x}_{i}^{(t)}% ,\boldsymbol{x}_{i}^{(t)}+\varepsilon]\boldsymbol{x}_{i}^{(t)}]^T$ $\{a_i^{(t)}\}^{\top} u_j]^4$ } $\leq C < \infty$. By Corollary 2 in [Bakhshizadeh et al.](#page-34-0) [\(2020\)](#page-34-0), $\mathbb{P}\left(|\epsilon_i^{(t)}\right)$ $\binom{t}{i} \cdot [(\boldsymbol{x}_i^{(t)}$ $\{(\begin{matrix} t\ i\end{matrix})^\top \bm u_j]^2-\mathbb{E}\{|\epsilon_i^{(t)}\}|$ $\binom{t}{i} \cdot [(\boldsymbol{x}_i^{(t)}$ $\left\{ \begin{array}{l} (t) \text{ } \top \textbf{u}_j]^2 \} > \delta \end{array} \right\} \leq \exp \{-Cn\delta\} + \exp \{-C(n\delta)^{2/3}\},$

which entails that w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
\frac{1}{n}\max_{j\in[N]}\left\{\sum_{i=1}^n|\epsilon_i^{(t)}|\cdot[(\boldsymbol{x}_i^{(t)})^\top\boldsymbol{u}_j]^2\right\}\lesssim\max_{j\in[N]}\mathbb{E}\{|\epsilon_i^{(t)}|\cdot[(\boldsymbol{x}_i^{(t)})^\top\boldsymbol{u}_j]^2\}+\sqrt{\frac{p+\log T}{n}}\vee\frac{(p+\log T)^{3/2}}{n}
$$

 $\leq C''$.

Hence by Lemma [19](#page-47-0) and the assumption on g'' , w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
[1] \ge C \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}\|_2^2 \ge \frac{C}{2} \|\boldsymbol{\beta} - \widetilde{\boldsymbol{\beta}}\|_2^2 - C \|\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}^{(t)*}\|_2^2. \tag{65}
$$

On the other hand, similarly, we can show that w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
[2] \lesssim \|\tilde{\boldsymbol{\beta}} - \boldsymbol{\beta}^{(t)*}\|_2^2. \tag{66}
$$

Combining [\(65\)](#page-115-0) and [\(66\)](#page-115-1) completes the proof.

F.4.2 Proof of Lemma [54](#page-111-3)

Note that $\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*}) = -\frac{2}{n}$ $\frac{2}{n}\sum_{i=1}^n \epsilon_i^{(t)}$ $\int\limits_i^{(t)}\!g'((\pmb{x}^{(t)}_i$ $\bm{g}^{(t)})^\top \bm{\beta}^{(t)*})\bm{x}^{(t)}_i$ $\binom{t}{i}$, where $\epsilon_i^{(t)}$ $\int\limits_i^{(t)} g'((\boldsymbol{x}^{(t)}_i$ $\bm{g}^{(t)})^\top\bm{\beta}^{(t)\ast})\bm{x}^{(t)}_i$ $i^{(t)}$ is $\text{sub-exponential in the sense that } \mathbb{E}\exp\{\lambda \epsilon_i^{(t)}g'((\boldsymbol{x}_i^{(t)})$ $\bm{g}^{(t)})^\top\bm{\beta}^{(t)\ast})(\bm{x}_i^{(t)}$ $\{(\boldsymbol{u}^{(t)})^{\top}\boldsymbol{u}\} \leq \exp\{C^2\lambda^2 \|\boldsymbol{u}\|_2^2\}$ for any $u \in \mathbb{R}^p$ and $\lambda \in \mathbb{R}$ with $|\lambda| ||u||_2 \leq C'$. Then the proof is very similar to the proof of Lemma [44,](#page-101-2) so we omit the details.

F.4.3 Proof of Lemma [55](#page-111-2)

The proof is very similar to the proof of Lemma [45,](#page-101-3) so omitted.

F.4.4 Proof of Lemma [56](#page-111-0)

Denote $\beta_{\delta} = \delta \beta + (1 - \delta) \tilde{\beta}$ with some $t \in [0, 1]$. It suffices to show that w.h.p, for all $t \in [0,1]$ and all $t \in S$, the minimum and maximum eigenvalues of $\nabla^2 f^{(t)}(\beta_\delta)$ are bounded away from zero and infinity.

Note that

$$
\nabla^{2} f^{(t)}(\beta_{\delta}) = \frac{1}{n} \sum_{i=1}^{n} [g'((\mathbf{x}_{i}^{(t)})^{\top} \beta_{\delta})]^{2} \cdot \mathbf{x}_{i}^{(t)}(\mathbf{x}_{i}^{(t)})^{\top} \n+ \frac{1}{n} \sum_{i=1}^{n} [g((\mathbf{x}_{i}^{(t)})^{\top} \beta_{\delta}) - y_{i}^{(t)}] \cdot g''((\mathbf{x}_{i}^{(t)})^{\top} \beta_{\delta}) \cdot \mathbf{x}_{i}^{(t)}(\mathbf{x}_{i}^{(t)})^{\top} \n= \underbrace{\frac{1}{n} \sum_{i=1}^{n} [g'((\mathbf{x}_{i}^{(t)})^{\top} \beta_{\delta})]^{2} \cdot \mathbf{x}_{i}^{(t)}(\mathbf{x}_{i}^{(t)})^{\top}}_{[1]} \n+ \underbrace{\frac{1}{n} \sum_{i=1}^{n} [g((\mathbf{x}_{i}^{(t)})^{\top} \beta_{\delta}) - g((\mathbf{x}_{i}^{(t)})^{\top} \beta_{i}^{(t)*})] \cdot g''((\mathbf{x}_{i}^{(t)})^{\top} \beta_{\delta}) \cdot \mathbf{x}_{i}^{(t)}(\mathbf{x}_{i}^{(t)})^{\top}}_{[2]} \n- \underbrace{\frac{1}{n} \sum_{i=1}^{n} \epsilon_{i}^{(t)} \cdot g''((\mathbf{x}_{i}^{(t)})^{\top} \beta_{\delta}) \cdot \mathbf{x}_{i}^{(t)}(\mathbf{x}_{i}^{(t)})^{\top}}_{[3]}.
$$

It is straightforward to see that w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
0 < C \leq \min_{t \in S} \lambda_{\min}([1]) \leq \max_{t \in S} \lambda_{\max}([1]) \leq C' < \infty,
$$

by the condition on g' and Lemma [19.](#page-47-0) And similar to the analysis in the proof of Lemma [53,](#page-111-1) w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
\lambda_{\max}([3]) \leq c,
$$

where c is small. Similarly, it can be shown that

$$
[2] = \frac{1}{n}\sum_{i=1}^n \int_0^1 g'(w(\boldsymbol{x}_i^{(t)})^\top \boldsymbol{\beta}_{\delta} + (1-w)(\boldsymbol{x}_i^{(t)})^\top \boldsymbol{\beta}^{(t)*}) \mathrm{d}w \cdot g''((\boldsymbol{x}_i^{(t)})^\top \boldsymbol{\beta}_{\delta}) \cdot \boldsymbol{x}_i^{(t)}(\boldsymbol{x}_i^{(t)})^\top \cdot (\boldsymbol{x}_i^{(t)})^\top (\boldsymbol{\beta}_{\delta} - \boldsymbol{\beta}^{(t)*}),
$$

$$
\lambda_{\max}([2]) \le C \|\boldsymbol{\beta}_{\delta} - \boldsymbol{\beta}^{(t)*}\|_2 \lesssim \|\boldsymbol{\beta} - \boldsymbol{\beta}^{(t)*}\|_2 + \|\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}^{(t)*}\|_2 \le c,
$$

w.p. at least $1 - e^{-C'(p + \log T)}$, where c is small. Putting all the pieces together, we have

$$
0 < C \leq \min_{t \in S} \lambda_{\min}(\nabla^2 f^{(t)}(\beta_\delta)) \leq \max_{t \in S} \lambda_{\max}(\nabla^2 f^{(t)}(\beta_\delta)) \leq C' < \infty,
$$

w.p. at least $1 - e^{-C(p + \log T)}$. The remaining steps are the same as in the proof of Lemma [32.](#page-51-0)

F.4.5 Proof of Lemma [57](#page-112-2)

The proof is similar to the proofs of Lemmas [56](#page-111-0) and [30,](#page-50-0) so omitted.

F.4.6 Proof of Lemma [58](#page-112-3)

Define $\widetilde{\theta}_{A}^{(t)} = A^{\top} A^{(t)*} \theta^{(t)*}$. It is easy to see that

$$
\|\mathbf{A}\widetilde{\boldsymbol{\theta}}_{\mathbf{A}}^{(t)} - \mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \leq \|(\mathbf{A}\mathbf{A}^\top - \mathbf{I})\mathbf{A}^{(t)*}\boldsymbol{\theta}^{(t)*}\|_2 \lesssim \|(\mathbf{A}^\perp)^\top \mathbf{A}^{(t)*}\|_2 \zeta^{(t)} \leq c.
$$

Then by Lemma [53,](#page-111-1) w.p. at least $1 - e^{-C(p + \log T)}$,

$$
f^{(t)}(\mathbf{A}\boldsymbol{\theta}_A^{(t)}) - f^{(t)}(\mathbf{A}\widetilde{\boldsymbol{\theta}}_A^{(t)}) \ge C\|\mathbf{A}\boldsymbol{\theta}_A^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_A^{(t)}\|_2^2 - C'\|\mathbf{A}\widetilde{\boldsymbol{\theta}}_A^{(t)} - \boldsymbol{\beta}^{(t)*}\|_2^2 - C'\|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2 \cdot \|\mathbf{A}\boldsymbol{\theta}_A^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_A^{(t)}\|_2,
$$

where the RHS must be non-positive, otherwise it is contradicted with the definition of $\theta_A^{(t)}$. Therefore, $||\mathbf{A}\boldsymbol{\theta}^{(t)}_{\mathbf{A}} - \mathbf{A}\widetilde{\boldsymbol{\theta}}^{(t)}_{\mathbf{A}}||_2 \lesssim ||\mathbf{A}\widetilde{\boldsymbol{\theta}}^{(t)}_{\mathbf{A}} - \boldsymbol{\beta}^{(t)*}||_2 + ||\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})||_2^{1/2} \leq c$, where $c > 0$ is a small constant, w.p. at least $1 - e^{-C(p + \log T)}$. This also implies that $||\boldsymbol{\theta}_{\boldsymbol{A}}^{(t)}||_2 \leq C\zeta^{(t)}$ w.p. at least $1 - e^{-C'(p + \log T)}$.

F.4.7 Proof of Lemma [59](#page-112-0)

Define $\widetilde{\theta}_{\mathbf{A}}^{(t)} = \mathbf{A}^{\top} \widetilde{\mathbf{A}} \theta_{\widetilde{\mathbf{A}}}^{(t)}$. It is easy to see that

$$
\|\boldsymbol A\widetilde {\boldsymbol \theta}^{(t)}_{\boldsymbol A} - \widetilde {\boldsymbol A} \boldsymbol \theta^{(t)}_{\widetilde {\boldsymbol A}}\|_2 \leq \|(\boldsymbol A \boldsymbol A^\top - \boldsymbol I)\widetilde {\boldsymbol A} \boldsymbol \theta^{(t)}_{\widetilde {\boldsymbol A}}\|_2 \lesssim \|(\boldsymbol A^\perp)^\top \widetilde {\boldsymbol A}\|_2\zeta^{(t)} \leq c.
$$

Then by Lemma [53,](#page-111-1) w.p. at least $1 - e^{-C(p + \log T)}$,

$$
f^{(t)}(\mathbf{A}\boldsymbol{\theta}_A^{(t)}) - f^{(t)}(\mathbf{A}\widetilde{\boldsymbol{\theta}}_A^{(t)}) \ge C\|\mathbf{A}\boldsymbol{\theta}_A^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_A^{(t)}\|_2^2 - C'\|\mathbf{A}\widetilde{\boldsymbol{\theta}}_A^{(t)} - \boldsymbol{\beta}^{(t)*}\|_2^2 - C'\|\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})\|_2 \cdot \|\mathbf{A}\boldsymbol{\theta}_A^{(t)} - \mathbf{A}\widetilde{\boldsymbol{\theta}}_A^{(t)}\|_2,
$$

where the RHS must be non-positive, otherwise it is contradicted with the denition of $\theta_A^{(t)}$. Therefore, $||\mathbf{A}\boldsymbol{\theta}^{(t)}_{\mathbf{A}} - \mathbf{A}\widetilde{\boldsymbol{\theta}}^{(t)}_{\mathbf{A}}||_2 \lesssim ||\mathbf{A}\widetilde{\boldsymbol{\theta}}^{(t)}_{\mathbf{A}} - \boldsymbol{\beta}^{(t)*}||_2 + ||\nabla f^{(t)}(\boldsymbol{\beta}^{(t)*})||_2^{1/2} \leq \text{a small constant } c$ w.p. at least $1 - e^{-C(p + \log T)}$. This also implies that $||A\theta_A^{(t)} - \beta^{(t)}||_2 \leq ||A\theta_A^{(t)} - A\widetilde{\theta}_A^{(t)}||_2 +$ $\|\widetilde{A}\widetilde{\theta}^{(t)}_{\mathbf{A}} - \widetilde{A}\theta^{(t)}_{\widetilde{\mathbf{A}}}\|_2 + \|\widetilde{A}\theta^{(t)}_{\widetilde{\mathbf{A}}} - \beta^{(t)*}\|_2 \leq \text{a small constant } c \text{ and } \|\theta^{(t)}_{\mathbf{A}}\|_2 \leq C \text{ w.p. at least}$ $\frac{A}{r}$ $1 - e^{-C'(p + \log T)}$, which completes the proof.

F.4.8 Proof of Lemma [60](#page-112-1)

By Lemmas [53](#page-111-1) and [55,](#page-111-2) w.p. at least $1 - e^{-C(p + \log T)}$, for any $\widetilde{A} \in \mathcal{O}^{p \times r}$,

$$
f^{(t)}(\boldsymbol{A}^{(t)}\boldsymbol{\theta}^{(t)})+\frac{\lambda}{\sqrt{n}}\|\boldsymbol{A}^{(t)}(\boldsymbol{A}^{(t)})^\top-\widetilde{\boldsymbol{A}}(\widetilde{\boldsymbol{A}})^\top\|_2-f^{(t)}(\boldsymbol{A}^{(t)*}\boldsymbol{\theta}^{(t)*})-\frac{\lambda}{\sqrt{n}}\|\boldsymbol{A}^{(t)*}(\boldsymbol{A}^{(t)*})^\top-\widetilde{\boldsymbol{A}}(\widetilde{\boldsymbol{A}})^\top\|_2
$$

$$
\geq \nabla f^{(t)} (\mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*})^{\top} (\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*}) - 2 \frac{\lambda}{\sqrt{n}} + C \| \mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} \|_2^2
$$

\n
$$
\geq C \| \mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} \|_2^2 - C' \sqrt{\frac{p + \log T}{n}} \| \mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*} \|_2 - 2 \frac{\lambda}{\sqrt{n}}.
$$

By definition of $A^{(t)}\theta^{(t)}$, the RHS must be non-positive, which gives us

$$
\|\mathbf{A}^{(t)} \boldsymbol{\theta}^{(t)} - \mathbf{A}^{(t)*} \boldsymbol{\theta}^{(t)*}\|_2 \lesssim \sqrt{\frac{p + \log T}{n}} + \frac{\lambda^{1/2}}{n^{1/4}} \leq c,
$$

where $c > 0$ is a small constant, w.p. at least $1 - e^{-C'(p + \log T)}$.

Appendix G. Proofs for Estimation of Intrinsic Dimension r

G.1 Proof of Theorem [11](#page-24-0)

We already mentioned the motivation of the thresholding strategy briefly in Section [4.](#page-23-1) Intuitively, there is a gap between $\sigma_r(\mathbf{B}_S^*)$ and $\sigma_{r+1}(\mathbf{B}_S^*)$ when h and $|S^c|$ are small. But in practice, we do not have access to the singular values of B_S^* . We have to estimate B_S^* first then use the singular value of this estimate. As long as the singular values of the estimate is not far from the singular values of B_S^* , thresholding can work well. We give a rigorous proof in the following. Most parts of the proof follow the arguments in the proof of Theorem [7](#page-17-0) and Theorem [13.](#page-43-0)

Consider the decomposition

$$
\boldsymbol{\beta}^{(t)*} = P_{\overline{A}}\boldsymbol{\beta}^{(t)*} + P_{\overline{A}^\perp}\boldsymbol{\beta}^{(t)*} = \overline{A}(\overline{A})^\top\boldsymbol{\beta}^{(t)*} + \overline{A}^\perp(\overline{A}^\perp)^\top\boldsymbol{\beta}^{(t)*} := \overline{A}\boldsymbol{\theta}_{\overline{A}}^{(t)*} + \boldsymbol{\delta}^{(t)*}, \quad \forall t \in S.
$$

Denote $\overline{\Theta}_S = \{ \theta_{\overline{A}}^{(t)*} \}$ $\{\overline{A}_{\overline{A}}^{(t)*}\}_{t\in S},\ \overline{B}_{S}=\{\overline{A}\theta_{\overline{A}}^{(t)*}\}_{t\in S},\ \overline{B}=\overline{A}\overline{\Theta}=(\overline{B}_{S}\hspace{0.3cm}0_{p\times |S^{c}|}),\ B_{S}^{*}=\{\beta^{(t)*}\}_{t\in S},$ $\widetilde{\mathbf{B}}_S = \{ \widetilde{\beta}^{(t)} \}_{t \in S}, \ \widehat{\mathbf{B}}_S = \{ \prod_R (\widetilde{\beta}^{(t)}) \}_{t \in S}, \ \widehat{\mathbf{B}} = \{ \prod_R (\widetilde{\beta}^{(t)}) \}_{t=1}^T, \text{ and } \mathbf{D}_S^* = \{ \delta^{(t)*} \}_{t \in S}.$ By a similar argument as in the proof of Theorem [7,](#page-17-0)

$$
\|\overline{\mathbf{B}}_{S}/\sqrt{T} - \mathbf{B}_{S}^{*}/\sqrt{T}\|_{2} = \frac{1}{\sqrt{T}} \|\mathbf{D}_{S}^{*}\|_{2}
$$

\$\leq \bar{\zeta}h \cdot \left[\frac{\sigma_{\max}(\mathbf{D}_{S}^{*})}{\sqrt{r}\sigma_{\min}(\mathbf{D}_{S}^{*})} \wedge 1\right]\$
\$\leq \frac{\bar{\zeta}}{\sqrt{r}} \left[\frac{\sigma_{\max}(\mathbf{D}_{S}^{*})}{\sigma_{\min}(\mathbf{D}_{S}^{*})} \wedge \sqrt{r}\right] \cdot h \cdot \frac{\min_{t \in S} \zeta^{(t)}}{\sqrt{(p + \log T)}/n}\$
\$\leq \frac{\bar{\zeta}}{\sqrt{r}}\$

$$
\leq \widetilde{C} \sqrt{\frac{p + \log T}{n}},
$$

when $\min_{t \in S} \zeta^{(t)} \cdot h \left[\frac{\sigma_{\max}(\mathbf{D}_{S}^{*})}{\sigma_{\min}(\mathbf{D}_{S}^{*})} \right]$ $\left[\frac{\sigma_{\max}(\boldsymbol D_S^*)}{\sigma_{\min}(\boldsymbol D_S^*)} \wedge \sqrt{r} \right] \lesssim \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$. Because $n \gtrsim p + \log T$, we have

$$
\|\overline{\mathbf{B}}_{S}/\sqrt{T} - \mathbf{B}_{S}^{*}/\sqrt{T}\|_{2} \le \widetilde{C}\sqrt{\frac{p+\log T}{n}},\tag{67}
$$

w.p. at least $1-e^{-C'(p+\log T)}$. Since $R =$ quantile $(\{\widetilde{\beta}^{(t)}\}_{t=1}^T, 1-\overline{\epsilon})$, we have $\prod_{n\geq 0}(\widetilde{\beta}^{(t)})=\widetilde{\beta}^{(t)}$ for $\overline{\epsilon}$ -proportion of S among all $t \in S$. In addition, by the fact that $\max_{t \in S} ||\widetilde{\beta}^{(t)} - \beta^{(t)*}||_2 \lesssim$ $\sqrt{p + \log T}$ $\frac{\log T}{n}$, w.p. at least $1 - e^{-C'(p + \log T)}$, the assumption $\min_{t \in S} \zeta^{(t)} \gtrsim \sqrt{\frac{p + \log T}{n}}$ $\frac{\log T}{n}$, and $\bar{\epsilon} \geq \epsilon$, we have

$$
R \le \max_{t \in S} \|\widetilde{\beta}^{(t)}\|_2 \lesssim \max_{t \in S} \zeta^{(t)},\tag{68}
$$

w.p. at least $1 - e^{-C'(p + \log T)}$. Therefore,

$$
\|\widehat{\mathbf{B}}_S/\sqrt{T}-\widetilde{\mathbf{B}}_S/\sqrt{T}\|_2 \leq \|\widehat{\mathbf{B}}_S/\sqrt{T}-\widetilde{\mathbf{B}}_S/\sqrt{T}\|_{\mathrm{F}} = \sqrt{\frac{1}{T}\sum_{t\in S} \|\Pi_R(\widetilde{\boldsymbol{\beta}}^{(t)})-\widetilde{\boldsymbol{\beta}}^{(t)}\|_2^2} \leq C\sqrt{\epsilon}R.
$$

By Lemma 5.39 in [Vershynin](#page-39-0) [\(2010\)](#page-39-0), we have

$$
\|\widetilde{\mathbf{B}}_S/\sqrt{T}-\mathbf{B}_S^*/\sqrt{T}\|_2 \leq C''\sqrt{\frac{p+T}{nT}} \leq C\sqrt{\frac{p+\log T}{n}},
$$

w.p. at least $1 - e^{-C'(p + \log T)}$. By the equation [\(67\)](#page-119-0), and the single-task rate, w.p. at least $1 - e^{-C'(p + \log T)},$

$$
\|\widehat{\mathbf{B}}_S/\sqrt{T} - \mathbf{B}_S^*/\sqrt{T}\|_2 \le \|\widehat{\mathbf{B}}_S/\sqrt{T} - \widetilde{\mathbf{B}}_S/\sqrt{T}\|_2 + \|\widetilde{\mathbf{B}}_S/\sqrt{T} - \mathbf{B}_S^*/\sqrt{T}\|_2
$$

$$
\le C\sqrt{\epsilon}R + C\sqrt{\frac{p + \log T}{n}}.
$$

Moreover, by the projection, $||\dot{\mathbf{B}}_{S^c}||$ √ $T\|_2 \leq \|\boldsymbol{B}_{S^c}/\|$ √ $\overline{T}\Vert_\text{F}\leq \sqrt{\frac{|S^c|}{T}}$ $\frac{S^c}{T} \cdot R \leq$ √ $\bar{\epsilon}R$, w.p. at least $1 - e^{-C'(p + \log T)}$. Hence w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
\|\widehat{\mathbf{B}}/\sqrt{T} - \overline{\mathbf{B}}/\sqrt{T}\|_2 \le \|\widehat{\mathbf{B}}_S/\sqrt{T} - \overline{\mathbf{B}}_S/\sqrt{T}\|_2 + \|\widehat{\mathbf{B}}_{S^c}/\sqrt{T}\|_2
$$

\n
$$
\le \|\widehat{\mathbf{B}}_S/\sqrt{T} - \mathbf{B}_S^*/\sqrt{T}\|_2 + \|\mathbf{B}_S^*/\sqrt{T} - \overline{\mathbf{B}}_S/\sqrt{T}\|_2 + \|\widehat{\mathbf{B}}_{S^c}/\sqrt{T}\|_2
$$

\n
$$
\le (\widetilde{C} + C)\sqrt{\frac{p + \log T}{n}} + (C + 1)\sqrt{\overline{\epsilon}}R.
$$

Then by Weyl's inequality, for all $r' \leq r$,

$$
\sigma_{r'}(\widehat{\mathbf{B}}/\sqrt{T}) \ge \sigma_{r'}(\overline{\mathbf{B}}/\sqrt{T}) - (\widetilde{C} + C)\sqrt{\frac{p + \log T}{n}} - (C+1)\sqrt{\epsilon}R
$$

$$
\ge \sigma_{r'}(\mathbf{B}_{S}^{*}/\sqrt{T}) - (2\widetilde{C} + C)\sqrt{\frac{p + \log T}{n}} - (C+1)\sqrt{\epsilon}R
$$

$$
\geq \frac{c}{\sqrt{r}}\bar{\zeta} - (2\tilde{C} + C)\sqrt{\frac{p + \log T}{n}} - (C+1)\sqrt{\bar{\epsilon}}R
$$

$$
\geq 2(2\tilde{C} + C)\sqrt{\frac{p + \log T}{n}} + 2(C+1)\sqrt{\bar{\epsilon}}R,
$$

w.p. at least $1 - e^{-C'(p + \log T)}$, when $n \geq \frac{36(2\tilde{C} + C)^2}{a^2\tilde{C}^2}$ $\frac{2\tilde{C}+C)^2}{c^2\tilde{\zeta}^2}r(p+\log T)$ and $\frac{c}{2\sqrt{r}}\bar{\zeta}\geq 3(C+1)\sqrt{\bar{\epsilon}}R$. The second condition holds because of $\bar{\epsilon} \leq \frac{c'}{r}$ $\frac{c'}{r}\bigg(\frac{\bar{\zeta}}{\max_{t\in\mathbb{R}}}$ $\left(\frac{\bar{\zeta}}{\max_{t \in S} \zeta^{(t)}}\right)^2$ and [\(68\)](#page-119-1).

On the other hand, by Weyl's inequality, for all $r' > r$, w.p. at least $1 - e^{-C'(p + \log T)}$,

$$
\sigma_{r'}(\widehat{\boldsymbol{B}}/\sqrt{T}) \leq \sigma_{r'}(\overline{\boldsymbol{B}}/\sqrt{T}) + (\widetilde{C}+C)\sqrt{\frac{p+\log T}{n}} + (C+1)\sqrt{\overline{\epsilon}}R = (2\widetilde{C}+C)\sqrt{\frac{p+\log T}{n}} + (C+1)\sqrt{\overline{\epsilon}}R,
$$

because $\sigma_{r'}(\overline{B}/$ \overline{T} = 0 when $r' > r$. Finally, by setting $T_1 \in (2\tilde{C} + C, 4\tilde{C} + 2C]$ and $T_2 \in (C+1, 2C+2]$, we complete the proof.