

The neutrino-antineutrino annihilation to electron-positron pair around a Schwarzschild black hole in asymptotic safety

Yuxuan Shi, Hongbo Cheng*

Department of Physics,
East China University of Science and Technology,
Shanghai 200237, China
The Shanghai Key Laboratory of Astrophysics,
Shanghai 200234, China

Abstract

We research on the neutrino pair annihilation $\nu + \bar{\nu} \rightarrow e^- + e^+$ around a massive source in asymptotic safety. The ratio $\frac{\dot{Q}}{Q_{Newt}}$ corresponding to the energy deposition per unit time over that in the Newtonian case is derived and calculated. We find that the quantum corrections to the black hole spacetime affect the emitted energy rate ratio for the annihilation. It is interesting that the more considerable quantum effect reduces the ratio value slightly. Although the energy conversion is damped because of the quantum correction, the energy deposition rate is enough during the neutrino-antineutrino annihilation. The corrected annihilation process can become a source of gamma ray burst. We also investigate the derivative $\frac{d\dot{Q}}{dr}$ relating to the star's radius r to show that the quantum effect for the black hole will drop the ratio. The more manifest quantum gravity influence leads the weaker neutrino pair annihilation.

PACS number(s):

Keywords: quantum gravity; black hole; Gamma-ray burst;

*E-mail address: hbcheng@ecust.edu.cn

I. Introduction

A lot of efforts from the astrophysical community have been contributed to explain the gamma-ray burst phenomenon with enough energy source. There are neutrinos and antineutrinos emitted by the hot accretion disk [1-14]. The process of the neutrino-antineutrino annihilation into electrons and positrons can become an energy source of gamma-ray bursts, so the gamma-ray bursts may be thought as systems powered by newborn, stellar-mass black holes accreting matter at hyper-critical rates [1-14]. The so-called gamma-ray bursts must be based on more powerful energy sources, and we should understand the origins. The annihilation $\nu + \bar{\nu} \rightarrow e^- + e^+$ augment the neutrino heating of the envelope generating a supernova explosion, further the pair e^-e^+ near the surface of collapsing neutron star sets free the gamma ray that may provide a possible explanation of the observed bursts [15]. In addition to the gravitational source in various stages respectively, the structures of backgrounds and the generalization of general relativity can put manifest influence on the energy deposition [15]. It was found that the efficiency of neutrino-antineutrino annihilation into electron-positron pair is enhanced over the Newtonian values up to a factor of more than 4 in the case of Type II supernovae, or up to a factor of 30 near the surface of collapsing neutron stars [15]. The relativistic effects on the energy deposition rate according to the neutrino pair annihilation near the rotation axis of a Kerr black hole with a thin accretion disk was studied [16, 17]. The stronger off-axis contributions to the energy-momentum deposition rate from the $\nu\bar{\nu}$ pair collisions above a Kerr black hole within thin accretion disk were also discussed [18, 19]. It was shown that the generalizations of general relativity such as charged Galileon, Einstein dilation Gauss-Bonnet, Brans-Dicke, Eddington-inspired Born-Infeld, Born-Infeld generalization of Reissner-Nordstrom solution and higher derivative gravity raise the energy deposition significantly in the neutron star and supernova envelope [20]. The authors of Ref. [21] consider the neutrino pair annihilation efficiency around a gravitational object belonging to the black hole in the presence of quintessence field to show that the quintessence powers the emitted energy rate ratio, so the enhancement could be a source for the gamma-ray burst. We also discuss the neutrino pair annihilation around a massive source with an $f(R)$ global monopole to find that this kind of stellar bodies can be well qualified as a source of gamma-ray bursts [22]. Within the frames of generalized gravities such as Hartle-Thorne spacetime, Born-Infeld Reissner-Nordstrom metric and quintessence background, it was found that the neutrino-antineutrino pair annihilation process is mediated by the Z' gauge boson from the $U(1)_{B-L}$ extension of the standard model and the maximum gamma ray bursts energy can be reached [23].

The black holes attract a lot of attentions from physical community for several decades. As solutions to Einstein's field equation in the general relativity, the black holes are also the end-results of some stars during the process of gravitational collapse [23-25]. In the case of black holes, the classical description of spacetime can not be used to define the surrounding of singularities inside the event horizon [23-25]. In order to overcome the plague from the singular background, it is necessary to generalize the general relativity up to the high energy scale [26-35]. The generalization can lead

the quantum gravity which could resolve the unphysical black hole singularities [26-35]. Several proposals such as loop quantum gravity [26-30], Horava-Lifshitz gravity [31-35], etc. have been put forward. Here we are going to focus on the powerful issue named as asymptotic safety scenario (AS) which is utilized to get rid of the singularities of the black holes [35]. The techniques of the functional renormalization group (FRG) was hired to limit the results of the theory at trans-Planckian energies, making the physical quantities to avoid the divergences at all scales [35]. The quantum effects are always shown as corrections to the classical metrics [36-44]. A series of corrected metrics include Schwarzschild [36], Kerr [37], Schwarzschild-AdS [38], Kerr-AdS [39] and Reissner-Nordstrom-AdS [40-44] without the classical singularity. More attentions have been contributed to the RGI-Schwarzschild black holes. The evaporation of the RGI-Schwarzschild black hole was investigated [45, 46], so was the mini-black hole production in colliders [47]. The radial accretion of matter onto a RGI-Schwarzschild black hole was also explored [48-50]. The shadows of black holes have been studied in the AS scenario for quantum gravity [51-54]. Recently, the quantum gravity effects on the radiation properties of a thin accretion disk surrounding a RGI-Schwarzschild black hole within the frame of IR-limit of the AS theory were discussed [55].

Here we will explore the energy deposition rate by the neutrino pair annihilation process around a Schwarzschild black hole in asymptotic safety. The metric of Schwarzschild black hole modified by the quantum effects needs to be studied in different directions. The motion of test particle around RGI-Schwarzschild black hole was described [56]. The electromagnetic ray reflection spectroscopy for the corrected black hole was also considered [25, 57-59]. The quantum gravity effects on radiation characteristics of thin accretion disks around a renormalization group improved Schwarzschild black hole were shown [55]. Besides the black hole features mentioned above, it is important to probe the annihilation energy deposition rate in the background of this kind of black holes to show the implications of asymptotic safety introducing the quantum corrections to the spacetime structures of the gravitational sources. However, to the best of our knowledge, little contributions have been paid for this topic. In this paper, we derive the integral form of the neutrino pair annihilation efficiency based on the Schwarzschild black hole with running Newton coupling $G(r)$ corresponds to the quantum effects. Secondly, we calculate the ratio of total energy deposition to total Newtonian energy deposition for parameter ξ encoding the quantum effects on the spacetime geometry. With the help of our numerical estimation, we will discuss the influence from a RGI-Schwarzschild black hole in the IR-limit of the asymptotic safety scenario on the possibility that the astrophysical bodies attract the annihilation process generating the gamma-ray burst. The results and conclusions are listed in the end.

II. The energy deposition rate by the neutrino annihilation process in the renormalization group improved Schwarzschild spacetime

We adopt the Schwarzschild metric modified by the quantum gravity in the infrared limit as follow [60-64],

$$ds^2 = f(r)dt^2 - \frac{dr^2}{f(r)} - r^2(d\theta^2 + \sin^2\theta d\varphi^2) \quad (1)$$

where

$$f(r) = 1 - \frac{2MG(r)}{r} \quad (2)$$

with mass M . For the quantum-gravity-corrected Schwarzschild metric in the infrared limit, the running coupling $G(r)$ takes the form [61],

$$G(r) = G_0(1 - \frac{\xi}{r^2}) \quad (3)$$

where ξ is a parameter with dimensions of length squared associated to the scale identification between the momentum scale and the radial coordinate. The general function of radial coordinate $G(r)$ as the generalization of gravitational constant, comes as a result of quantum effects. It should be pointed out that the running coupling $G(r)$ contains the parameter ξ standing for the renormalization group improvement and the parameter will change the spacetime structure [36, 54]. There are some upper bounds on ξ derived from a series of gravitational and nongravitational measurements [54]. We can choose $c = G_0 = 1$ in natural units. The radius of the event horizon is a root of $f(r) = 0$ [55]. According to the calculation and discussion, the dimensionless parameter $\bar{\xi} = \frac{\xi}{M^2}$ and the mass M should be limited as [55],

$$\begin{cases} 0 \leq \bar{\xi} \leq \bar{\xi}_c \\ M > M_c \end{cases} \quad (4)$$

where $\bar{\xi}_c = \frac{16}{27}$ and $M_c = \sqrt{\frac{27}{16}\xi}$, to keep the metric with two horizons [55]. It is significant to investigate the quantum gravity effects on the annihilation process within the region (4).

We start to consider the energy deposition in the spacetime governed by Eq.(2) and (3). The energy deposition per unit time and per unit volume for the neutrino pair annihilation process is given by [15],

$$\frac{dE(\mathbf{r})}{dt dV} = 2KG_F^2 F(r) \int \int n(\varepsilon_\nu) n(\varepsilon_{\bar{\nu}}) (\varepsilon_\nu + \varepsilon_{\bar{\nu}}) \varepsilon_\nu^3 \varepsilon_{\bar{\nu}}^3 d\varepsilon_\nu d\varepsilon_{\bar{\nu}} \quad (5)$$

where

$$K = \frac{1}{6\pi} (1 \pm 4\sin^2\theta_W + 8\sin^4\theta_W) \quad (6)$$

with the Weinberg angle $\sin^2\theta_W = 0.23$. For various neutrino pairs, the forms of Eq.(6) can be chosen as [15],

$$K(\nu_\mu, \bar{\nu}_\mu) = K(\nu_\tau, \bar{\nu}_\tau) = \frac{1}{6\pi} (1 - 4\sin^2\theta_W + 8\sin^4\theta_W) \quad (7)$$

and

$$K(\nu_e, \bar{\nu}_e) = \frac{1}{6\pi}(1 + 4 \sin^2 \theta_W + 8 \sin^4 \theta_W) \quad (8)$$

respectively [15]. Here the Fermi constant $G_F = 5.29 \times 10^{-44} \text{cm}^2 \text{MeV}^{-2}$. The angular integration factor is represented by [15],

$$\begin{aligned} F(r) &= \int \int (1 - \boldsymbol{\Omega}_\nu \cdot \boldsymbol{\Omega}_{\bar{\nu}})^2 d\Omega_\nu d\Omega_{\bar{\nu}} \\ &= \frac{2\pi^2}{3}(1-x)^4(x^2 + 4x + 5) \end{aligned} \quad (9)$$

where

$$x = \sin \theta_r \quad (10)$$

The angle θ_r is between the particle trajectory and the tangent vector to a circle orbit at radius r . For one kind of neutrino and antineutrino, $\Omega_\nu(\Omega_{\bar{\nu}})$ is the unit direction vector and $d\Omega_\nu(d\Omega_{\bar{\nu}})$ is a solid angle. At temperature T , $n(\varepsilon_\nu)$ and $n(\varepsilon_{\bar{\nu}})$ are number densities for neutrino and antineutrino respectively in the phase space and satisfy the Fermi-Dirac distribution [15],

$$n(\varepsilon_\nu) = \frac{2}{h^3} \frac{1}{e^{\frac{\varepsilon_\nu}{kT}} + 1} \quad (11)$$

where h is Planck constant and k is Boltzmann constant. With integrating the Eq.(5), the expression of deposition energy per unit time and unit volume is given by [15],

$$\frac{dE}{dt dV} = \frac{21\zeta(5)\pi^4}{h^6} K G_F^2 F(r) (kT)^9 \quad (12)$$

It is significant to derive the expression of $\frac{dE}{dt dV}$ which is used to further the research on the converted energy rate on the different compact bodies [15]. The expression has something to do with the position, so has the temperature $T = T(r)$ called local temperature [15].

The local temperature $T(\mathbf{r})$ measured by a local observer is defined as $T(\mathbf{r})\sqrt{g_{00}(\mathbf{r})} = \text{constant}$ with g_{00} , a component of spacetime metric [15]. The neutrino temperature at the neutrinosphere can be explained as [15],

$$T(r)\sqrt{g_{00}(r)} = T(R)\sqrt{g_{00}(R)} \quad (13)$$

where R is the radius of a gravitational source. It is elegant to replace the local temperature $T(r)$ in the future calculation according to the identity (13). The luminosity relating to the redshift can be selected as [15],

$$L_\infty = g_{00}(R_0)L(R_0) \quad (14)$$

where the luminosity for a single neutrino species at the neutrinosphere is [15],

$$L(R) = 4\pi R_0^2 \frac{7}{4} \frac{ac}{4} T^4(R) \quad (15)$$

where a is the radiation constant and c is the speed of light in vacuum. In order to replace the temperature associated with the observer's position, we combine Eq.(13), Eq.(14) and Eq.(15) and substitute them into Eq.(12) to obtain [15],

$$\frac{dE(\mathbf{r})}{dt dV} = \frac{21\zeta(5)\pi^4}{h^6} K G_F^2 k^9 \left(\frac{7}{4}\pi ac\right)^{-\frac{9}{4}} L_\infty^{\frac{9}{4}} F(r) \frac{(g_{00}(R))^{\frac{9}{4}}}{(g_{00}(r))^{\frac{9}{4}}} R_0^{-\frac{9}{2}} \quad (16)$$

In addition to the radial coordinate, the metric components for the massive source surface also appear in the expression of deposition energy per unit time and unit volume like Eq. (16). We can calculate the radiation energy power in the background of the gravitational source by means of the deposition energy density over time as Eq. (16). In order to compute the angular integration $F(r)$ from Eq.(9), we should further the discussion on the variable x in Eq.(10). We follow the procedure of Ref. [15] and solve the null geodesic in the spacetime of a spherically symmetric gravitational object [15] to show [20],

$$\begin{aligned} x^2 &= \sin^2 \theta_r |_{\theta_R=0} \\ &= 1 - \frac{R^2}{r^2} \frac{f(r)}{f(R)} \end{aligned} \quad (17)$$

Here $f(r) = g_{00}(r)$, a component of metric (2). It is useful to relate the variable $x = \sin \theta |_{\theta_R} = 0$ to the environment structure around the gravitational object [20]. According to Eq. (9), the angular integration factor become the function of the metric. We can proceed the integration of rate per unit time and unit volume from Eq.(16) on the spherically symmetric volume around the gravitational source [21],

$$\begin{aligned} \dot{Q} &= \frac{dE}{\sqrt{g_{00}} dt} \\ &= \frac{84\zeta(5)\pi^5}{h^6} K G_F^2 k^9 \left(\frac{7}{4}\pi ac\right)^{-\frac{9}{4}} L_\infty^{\frac{9}{4}} (g_{00}(R))^{\frac{9}{4}} R^{-\frac{9}{2}} \int_{R_0}^{\infty} \frac{r^2 \sqrt{-g_{11}(r)} F(r)}{g_{00}(r)} dr \end{aligned} \quad (18)$$

where $g_{11}(r) = -\frac{1}{f(r)}$, also a component of the renormalization group improved Schwarzschild metric (2). The metrics of the curved spacetime have a considerable influence on the deposition energy rate. Here \dot{Q} can reflect the total amount of energy converted from neutrinos to electron-positron pairs per unit time at any radius [15]. The conversion may become an explosion with the extremely large values of \dot{Q} . It is significant to compare the energy deposition rate (18) with the Newtonian quantities as [15, 20, 21],

$$\frac{\dot{Q}}{\dot{Q}_{Newt}} = 3(g_{00}(R))^{\frac{9}{4}} \int_1^{\infty} (x-1)^4 (x^2 + 4x + 5) \frac{y^2 \sqrt{-g_{11}(Ry)}}{g_{00}(Ry)^{\frac{9}{2}}} dy \quad (19)$$

with dimensionless variable $y = \frac{r}{R}$ and $g_{00}(r)$, $g_{11}(r)$ components of metric (1).

According to the Ref.[15], we can present $\frac{d\dot{Q}}{dr}$ as the function of radial coordinate r to exhibit the enhancement,

$$\begin{aligned} \frac{d\dot{Q}}{dr} &= 4\pi \left(\frac{dE}{dt dV} \right) \sqrt{-g_{11}(r)} r^2 \\ &= \frac{168\zeta(5)\pi^7}{3h^6} K G_F^2 k^9 \left(\frac{7}{4} \pi a c \right)^{-\frac{9}{4}} L_\infty^{\frac{9}{4}} (x-1)^4 (x^2 + 4x + 5) \left(\frac{g_{00}(R)}{g_{00}(r)} \right)^{\frac{9}{4}} R^{-\frac{5}{2}} \sqrt{-g_{11}(r)} \left(\frac{r}{R} \right)^2 \end{aligned} \quad (20)$$

Here the derivative is a function of radial coordinate with the origin at the centre of the gravitational source while involving the metric functions. It is necessary to wonder how the structure of the compact body in asymptotic safety affect the neutrino annihilation around one. The derivative function could tell us which kind of stars attracting the annihilation could become source of gamma-ray burst.

It is important to discuss the ratio (19) in the quantum-gravity-corrected Schwarzschild space-time in the infrared limit. We can relate that $g_{00}(r) = f(r)$ and $g_{11}(r) = -\frac{1}{f(r)}$ for metric (1). The ratio (19) can be reformed as [15, 60-64],

$$\frac{\dot{Q}}{\dot{Q}_{Newt}} = 3(f(R))^{\frac{9}{4}} \int_1^\infty (x-1)^4 (x^2 + 4x + 5) \frac{y^2}{(f(Ry))^5} dy \quad (21)$$

where

$$f(R) = 1 - \frac{2M}{R} + \frac{2M\xi}{R^3} \quad (22)$$

$$f(Ry) = 1 - \frac{2M}{R} \frac{1}{y} + \frac{2M\xi}{R^3} \frac{1}{y^3} \quad (23)$$

According to the metric function (2) and the running coupling (3), we rewrite the variable (17) as,

$$x^2 = 1 - \frac{1}{y^2} \frac{1 - \frac{2M}{R} \frac{1}{y} + \frac{2M\xi}{R^3} \frac{1}{y^3}}{1 - \frac{2M}{R} + \frac{2M\xi}{R^3}} \quad (24)$$

In order to show the influence from quantum gravity, we should quantify the integral expression of ratio $\frac{\dot{Q}}{\dot{Q}_{Newt}}$ in EQ.(21) and depict the dependence on $\frac{R_0}{M}$ with the effect parameter ξ in the Figures. It is necessary to gather data for the constraints over the values of ξ [54]. During our plotting, we have to magnify its values more than huge times in order to reveal the differences among the curves of the ratio $\frac{\dot{Q}}{\dot{Q}_{Newt}}$ for different magnitudes of factor ξ from quantum-gravity effects on the neutrino pair annihilation around the black hole in asymptotic safety gravity because of extremely tiny values of ξ from Ref. [54]. In Figure 1, the shape of ratio curves due to different values of parameter ξ are similar. The influence from this kind of sources consist of two parts, mass M and gravitational coupling $G(r)$ respectively. As a gravitational source, the values of ratio $\frac{\dot{Q}}{\dot{Q}_{Newt}}$ are much more than one, so this kind of stellar objects attracting the neutrino pair annihilation process may provide with a source of gamma-ray burst. For the sake of showing the asymptotic

behaviour of ratio $\frac{\dot{Q}}{Q_{Newt}}$ more clearly, we plot some curves to indicate that the ratio is decreasing function of ξ from quantum corrections in Figure 2. In contrast to the case of accretion disk around a stellar object in asymptotic safety that the quantum gravity effect can enhance the disk's thermal properties [55], the more considerable influence will reduce the ratio $\frac{\dot{Q}}{Q_{Newt}}$.

We also elaborate the derivative $\frac{d\dot{Q}}{dr}$ as a function of radius for several stellar masses denoted as $\frac{r}{R}$ with $R = 3M$ to exhibit the promotion of e^-e^+ pair energy from the neutrino annihilation in Figure 3. It is similar that a slight larger variable ξ from quantum gravity decreases the derivative $\frac{d\dot{Q}}{dr}$. The Figure 3 also indicate that the structure with smaller $\frac{r}{R}$ lead to the greater $\frac{d\dot{Q}}{dr}$. It is interesting that the increase is much larger near the surface of the neutron star similar to the case of Ref.[15] although the compact source surrounded by the annihilation includes the quantum gravity influence.

III. Conclusion

The neutrino pair annihilation $\nu\bar{\nu} \rightarrow e^-e^+$ around a Schwarzschild black hole in asymptotic safety is discussed. The quantum gravity influence on the black holes can be explored in different directions. We derivative and calculate the emitted energy rate ratio as $\frac{\dot{Q}}{Q_{Newt}}$ and the emitted energy rate \dot{Q} analyzed in the background of quantum-gravity corrected massive source. It is found that the quantum-gravity effect does not advance the ratio $\frac{\dot{Q}}{Q_{Newt}}$, but damps \dot{Q} , the energy released per unit time, in contrast to the case of accretion disk under the same conditions. The larger ξ , the parameter encoding the quantum gravity effects on the spacetime structure, will reduce the energy deposition rate. The ratio $\frac{\dot{Q}}{Q_{Newt}}$ shows numerically that the annihilation $\nu + \bar{\nu} \rightarrow e^- + e^+$ around the quantum-gravity corrected black hole can proceed as a source for the gamma-ray burst. This kind of annihilation can occur around the Type II supernova or collapsing neutron stars. Within the region of ξ , this kind of annihilation process could be proposed as a well-qualified candidate of gamma-ray bursts.

Acknowledge

This work is partly supported by the Shanghai Key Laboratory of Astrophysics 18DZ2271600.

References

- [1] D. Eichler, M. Livio, T. Piran, D. N. Schramm, *Nature* 340(1989)126
- [2] B. Paczynski, *Astrophys. J.* 363(1990)218
- [3] P. Meszaros, M. J. Rees, *MNRAS* 257(1992)29
- [4] S. E. Woosley, *Astrophys. J.* 405(1993)273
- [5] R. Mochkovitch, M. Hernanz, J. Isern, X. Martin, *Nature* 361(1993)236
- [6] M. B. Davies, W. Benz, T. Piran, F. K. Thieleman, *Astrophys. J.* 431(1994)742
- [7] M. Jaroszynski, *A A*, 305(1996)839
- [8] H. Janka, M. Ruffert, *A A*, 307(1996)L33
- [9] M. Ruffert, H. Janka, K. Takahashi, G. Schafer, *A A*, 319(1997)122
- [10] M. Ruffert, H. Janka, *A A*, 338(1998)535
- [11] R. Popham, S. E. Woosley, C. Fryer, *Astrophys. J.* 518(1999)356
- [12] A. MacFadyen, S. E. Woosley, *Astrophys. J.* 524(1999)262
- [13] M. Ruffert, H. Janka, *A A*, 344(1999)573
- [14] M. A. Aloy, E. Muller, J. M. Ibanez, J. M. Martin, A. MacFadyen, *Astrophys. J.* 531(2000)L119
- [15] J. D. Salmonson, J. R. Wilson, *Astrophys. J.* 517(1999)859
- [16] K. Asano, T. Fukuyama, *Astrophys. J.* 531(2000)949
- [17] K. Asano, T. Fukuyama, *Astrophys. J.* 546(2001)1019
- [18] W. A. Miller, N. D. George, A. Kheyfets, J. M. McGhee, *Astrophys. J.* 583(2003)833
- [19] R. Birkel, M. A. Aloy, H. Janka, E. Muller, *A A* 463(2007)51
- [20] G. Lambiase, L. Mastrototaro, *Astrophys. J.* 904(2020)1
A. Prasanna, S. Goswami, *Phys. Lett. B* 526(2002)27
- [21] G. Lambiase, L. Mastrototaro, *Eur. Phys. J.* C81(2021)932
- [22] Y. Shi, H. Cheng, *EPL* 140(2022)49001
- [23] T. K. Poddar, S. Goswami, A. K. Mishra, *Eur. Phys. J.* C83(2023)223

- [24] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*, John Wiley Sons Inc., 1972
M. Carmeli, *Classical Fields: General Relativity and Gauge Theory*, John Wiley Sons, Inc., 1982
- [25] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, Oxford University Press, Inc., 1992
- [26] A. Ashtekar, *Phys. Rev. Lett.* 57(1986)2244
- [27] A. Ashtekar, *Phys. Rev. D* 36(1987)1587
- [28] C. Rovelli, *Living Rev. Relativ.* 1(1998)1
- [29] A. Ashtekar, J. Baez, A. Corichi, K. Krasnov, *Phys. Rev. Lett.* 80(1998)904
- [30] A. Perez, *Class. Quantum Grav.* 20(2003)R43
- [31] P. Horava, *Phys. Rev. D* 79(2009)084008
- [32] P. Horava, *Phys. Rev. Lett.* 102(2009)161301
- [33] E. Kiritsis, G. Kofinas, *Nucl. Phys. B* 821(2009)467
- [34] G. Cognola, R. Myrzakulov, L. Sebastiani, S. Vagnozzi, S. Zerbini, *Class. Quantum Grav.* 33(2016)225014
- [35] M. Reuter, *Phys. Rev. D* 57(1998)971
- [36] A. Bonanno, M. Reuter, *Phys. Rev. D* 62(2000)043008
- [37] M. Reuter, E. Tuiran, *Phys. Rev. D* 83(2011)044041
- [38] B. Koch, F. Saueressig, *Class. Quantum Grav.* 31(2014)015006
- [39] J. M. Pawłowski, D. Stock, *Phys. Rev. D* 98(2018)106008
- [40] C. Gonzalez, B. Koch, *Int. J. Mod. Phys. A* 31(2016)1650141
- [41] R. Casadio, S. D. H. Hsu, B. Mirza, *Phys. Lett. B* 695(2011)317
- [42] G. Kofinas, V. Zarikas, *JCAP* 1510(2015)069
- [43] A. Bonano, B. Koch, A. Platania, *Found. Phys.* 48(2018)1393
- [44] L. Bosma, B. Knorr, F. Saueressig, *Phys. Rev. Lett.* 123(2019)101301
- [45] K. Falls, D. F. Litim, *Phys. Rev. D* 89(2014)084002
- [46] K. Falls, *Int. J. Mod. Phys. A* 27(2012)1250019

- [47] B. F. L. Ward, *Acta Phys. Polon.* B37(2006)1967
- [48] R. Yang, *Phys. Rev.* D92(2015)084011
- [49] M. U. Farooq, A. K. Ahmed, R. Yang, M. Jamil, *Chin. Phys.* C44(2020)065102
- [50] F. H. Zuluaga, L. A. Sanchez, *Chin. Phys.* C45(2021)075102
- [51] A. Held, R. Gold, A. Eichhorn, *JCAP* 1906(2019)029
- [52] X. Lu, Y. Xie, *Eur. Phys. J.* C79(2019)1016
- [53] R. Kumar, B. P. Singh, S. G. Ghosh, *Ann. Phys.* 420(2020)168252
A. Eichhorn, A. Held, *JCAP* 2105(2021)073
- [54] G. Lambiase, F. Scardigli, *Phys. Rev.* D105(2022)124054
- [55] F. H. Zuluaga, L. A. Sanchez, *Eur. Phys. J.* C81(2021)840
- [56] S. Haroon, M. P. Pavlovic, M. Sossich, A. Wang, *Eur. Phys. J.* C78(2018)519
- [57] D. N. Page, K. S. Thorne, *Astrophys. J.* 191(1974)499
- [58] B. Zhou, A. Abdikamalov, D. Ayzenberg, C. Bambi, S. Nampalliwar, A. Tripathi, *JCAP* 2101(2021)047
S. Vagnozzi, R. Roy, Y. Tsai, L. Visinelli, M. Afrin, A. Allahyari, P. Bambhaniya, D. Dey, S. G. Ghosh, P. S. Joshi, K. Jusufi, M. Khodadi, R. K. Walia, A. Ovgun, C. Bambi, "Horizon-scale tests of gravity theories and fundamental physics from the Event Horizon Telescope image of Sagittarius A*", arXiv: 2205.07787
- [59] Y. Zhang, M. Zhou, C. Bambi, *Eur. Phys. J.* C78(2018)376
- [60] D. Benedetti, P. F. Machado, F. Saueressig, *Mod. Phys. Lett.* A24(2009)2233
- [61] Y. Cai, D. A. Easson, *JCAP* 1009(2010)002
- [62] K. Falls, C. R. King, D. F. Litim, K. Nikolakopoulos, C. Rahmede, *Phys. Rev.* D97(2018)086006
- [63] K. Falls, D. F. Litim, J. Schroder, *Phys. Rev.* D99(2019)126015
- [64] Y. Kluth, D. F. Litim, arXiv: 2008.09181




Figure 1: The solid, dotted and dashed curves of the ratio $\frac{\dot{Q}}{Q_{Newt}}$ as functions of the ratio $\frac{R}{M}$ for quantum-gravity factors $\xi = 0, 0.26, 0.56$ respectively.




Figure 2: The solid, dashed and dot-dashed curves of the ratio $\frac{Q}{Q_{Newt}}$ as functions of the quantum-gravity factors ξ with $R = 4M, 4.5M, 5M$ respectively.




Figure 3: The solid, dashed, dot-dashed and dotted curves of the derivative $\frac{d\dot{Q}}{dr}$ as functions of the ratio $\frac{r}{R}$ with $R = 3M$ for quantum-gravity factors $\xi = 0, 0.4, 0.8, 1.3$ respectively.