# <span id="page-0-0"></span>Emergence of Gravitational Potential and Time Dilation from Non-interacting Systems Coupled to a Global Quantum Clock

Ashmeet  $\text{Singh}^1$  and Oliver Friedrich

Department of Physics Whitman College 345 Boyer Ave Walla Walla, WA 99362, USA

Faculty for Physics; Observatory Ludwig-Maximilians Universität München Geschwister-Scholl-Platz 1 80539 M¨unchen, Germany

#### Abstract

We study gravitational back-reaction within relational time formulations of quantum mechanics by considering two versions of time: a time coordinate, modelled as a global quantum degree of freedom, and the proper time of a given physical system, modelled via an internal degree of freedom serving as a local quantum "clock". We show that interactions between coordinate time and mass-energy in a global Wheeler-DeWitt-like constraint lead to gravitational time dilation. In the presence of a massive object this agrees with time dilation in a Schwarzchild metric at leading order in G. Furthermore, if two particles couple independently to the time coordinate we show that Newtonian gravitational interaction between those particles emerges in the low energy limit. We also observe features of renormalization of high energy divergences.

Essay written for the Gravity Research Foundation's 2023 Awards for Essays on Gravitation. Submitted March 31, 2023.

<sup>1</sup>Corresponding Author: A.S.

e-mail: [ashmeet@whitman.edu]( ashmeet@whitman.edu) , <oliver.friedrich@physik.uni-muenchen.de>

A defining feature of a complete quantum theory of gravity will be to treat space and time as quantum degrees of freedom. This will include, among other things, allowing for superpositions of classical spacetime geometries and a model for the ultraviolet degrees of freedom which make up spacetime. Treating both time and space on a quantum footing is challenging, but it allows for the possibility of quantum mechanical interactions between matter and spacetime degrees of freedom, from which gravity may emerge. With this motivation, we explore the possibility of time as a quantum degree of freedom, and show that its coupling with mass-energy leads to gravitational time dilation and an emergent Newtonian gravitational interaction potential.

Standard quantum mechanics handles time and space very differently. Time enters the Schrödinger equation as an essentially classical parameter  $t$ , while space (e.g. understood as the location of a particle) is represented through the eigenvalues and eigenspaces of a Hermitian operator. Attempts to also associate time with measurements of a time-operator have been criticised by Dirac based on the Stone-von Neumann theorem [\[1\]](#page-10-0) because the variable that is naturally conjugate to time is energy. And if the Hamiltonian operator  $\hat{H}$  and a time operator  $\hat{T}$  were to satisfy the canonical commutation relations  $[\hat{T}, \hat{H}] = i$ , then the eigenvalue spectrum of the Hamiltonian (and thus the energy) would need to be unbounded (from above and below).

Why would a time operator be desirable? In general relativity, as well as in more general theories with reparametrization invariance  $[2-4]$  $[2-4]$ , there is only a soft distinction between time and space. Any foliation of spacetime into non-intersecting space-like hypersurfaces defines a "time-coordinate". And the separation of two spacetime events can be space-like even if they are separated by a non-zero amount of coordinate time. This (literal) relativity in the definitions of time and space leads to formulations of quantum gravity, such as the Wheeler-DeWitt equation [\[5\]](#page-11-1), which are based on constraints of the form  $\hat{J}|\Psi\rangle = 0$ , where  $\hat{J}$  is a constraint operator [\[5](#page-11-1)[–7\]](#page-11-2) (see also [\[8](#page-11-3)[–10\]](#page-11-4) for a discussion of the closely-related issue of the "problem of time" in quantum gravity). Physical states  $|\Psi\rangle$  are required to satisfy the constraint and are hence in the 0-eigenspace of the constraint operator. They represent configurations of quantum gravity degrees of freedom that describe (superpositions of) full 3+1 dimensional spacetimes. In such a setting there is no notion of a Schrödinger-like time parameter, and time should instead be emergent, such as being a measurement conditional on a specified event.

#### Coordinate Time as Relational Quantum Observable

A simplified, albeit representative version of the above ideas is found in the Page-Wootters formalism  $[11-13]$  $[11-13]$  (see also related ideas by Dirac  $[6, 7]$  $[6, 7]$  $[6, 7]$ ) of relational time. Here, the global Hilbert space  $\mathcal H$  is factorized into a temporal degree of freedom  $\mathcal H_t$ , often called the "clock," and the system  $\mathcal{H}_{S}$  (what we typically describe in conventional quantum mechanics),

$$
\mathcal{H} \simeq \mathcal{H}_t \otimes \mathcal{H}_S \,.
$$

The temporal Hilbert space  $\mathcal{H}_t$  is taken isomorphic to  $\mathbb{L}_2(\mathbb{R})$  and we consider a pair of Hermitian operators  $\hat{t}$  and  $\hat{p}_t$  on that space which satisfy Heisenberg canonical commutation relation (CCR),  $[\hat{t}, \hat{p}_t] = i$  (in units with  $\hbar = 1$ ). These operators thus constitute a standard pair of conjugate operators on  $\mathcal{H}_t$ . In particular,  $\hat{t}$  has a continuous spectrum of eigenvalues t with corresponding orthonormal eigenstates  $|t\rangle$  with  $\langle t'|t\rangle = \delta(t-t')$ , such that the conjugate operator  $\hat{p}_t$  generates translations in t as  $e^{-i\hat{p}_t t'} |t\rangle = |t + t'\rangle$ . Within the Page-Wootters formalism the states  $|t\rangle$  are endowed with a temporal meaning by introducing the constraint operator  $\hat{J}$ .

<span id="page-2-1"></span>
$$
\hat{\mathbb{J}} = \hat{p}_t \otimes \hat{\mathbb{I}}_S + \hat{\mathbb{I}}_t \otimes \hat{H}_S , \qquad (2)
$$

where  $\hat{\mathbb{I}}_t$  and  $\hat{\mathbb{I}}_S$  are identity operators on  $\mathcal{H}_t$  and  $\mathcal{H}_S$ , respectively, and  $\hat{H}_S$  is the conventional Hamiltonian for the system. Physical states  $|\Psi\rangle$  in the global Hilbert space H are identified to be the ones annihilated by the constraint operator  $\mathbb{J}$ ,

$$
\hat{\mathbb{J}} \approx 0 \implies \hat{\mathbb{J}} | \Psi \rangle = 0. \tag{3}
$$

We use the double-ket notation  $|\Psi\rangle$  to stress that the state is defined on the global Hilbert space  $\mathcal{H}_t \otimes \mathcal{H}_S$ . Conventional time-dependent states of the system are obtained via a relational approach by conditioning the global, physical state  $|\Psi\rangle$  with the eigenvector  $|t\rangle$  of the time operator  $\tilde{t}$ ,

$$
|\psi(t)\rangle = \langle t|\Psi\rangle\rangle \in \mathcal{H}_S. \tag{4}
$$

The constraint equation conditioned on  $|t\rangle$  becomes

$$
\langle t|\hat{\mathbb{J}}|\Psi\rangle\rangle = \langle t|\hat{p}_t\otimes\hat{\mathbb{I}}_S|\Psi\rangle\rangle + \hat{H}_S|\psi(t)\rangle = 0, \qquad (5)
$$

which upon inserting a complete set of states on  $\mathcal{H}_t$  given by  $\int dt |t\rangle \langle t| = \hat{\mathbb{I}}_t$ , and remembering that the matrix elements of the conjugate momenta are  $\langle t|\hat{p}_t|t'\rangle = -i\frac{\partial}{\partial t}\delta(t-t')$ , gets us the time evolution equation for states  $|\psi(t)\rangle$  of the system,

$$
\hat{H}_S \left| \psi(t) \right\rangle = i \frac{\partial}{\partial t} \left| \psi(t) \right\rangle . \tag{6}
$$

This is indeed the Schrödinger equation for  $|\psi(t)\rangle \in \mathcal{H}_S$ . Thus we see that effective time evolution for states in the subfactor  $\mathcal{H}_S$  of the global Hilbert space, governed by a Hamiltonian  $\hat{H}_S$ , can be recovered from a constraint operator. One can furthermore show that any physical state can be expressed as

<span id="page-2-0"></span>
$$
|\Psi\rangle\rangle = \int dt \, |t\rangle \otimes |\psi(t)\rangle = \int dt \, |t\rangle \otimes \left(e^{-i\hat{H}_S t} \, |\psi(0)\rangle\right) \,,\tag{7}
$$

where  $\hat{U}(t) = e^{-i\hat{H}_S t}$  is the time evolution operator acting on an initial state  $|\psi(0)\rangle$  of the system. Time evolution is hence implemented in the Page-Wootters construction via entanglement between the time Hilbert space and the system Hilbert space.

#### Proper Time as an Internal Degree of Freedom

In the following we will think of the degree of freedom in  $\mathcal{H}_t$  as representing *coordinate time*. In other words, we take it to represent a very simplified version of the spacetime degrees of freedom that appear e.g. in the Wheeler-DeWitt equation. But the proper time experienced by the system  $\mathcal{H}_S$  is, in general, different than the coordinate time, and will be governed by the behavior of clocks carried by that system. To this end, we associate the system to have internal structure which serves as a "clock" to measure its proper time (imagine the system carrying a clock with it) [\[14\]](#page-11-8). Since we are focused on temporal effects in this essay, we will only consider such internal proper clocks, and we neglect spatial motion of their center of mass. On the system Hilbert space  $\mathcal{H}_S$ , we can associate a proper time observable  $\hat{\tau}_S$  and its corresponding conjugate  $\hat{\pi}_S$  which satisfies Heisenberg canonical commutation relation,  $[\hat{\tau}_S, \hat{\pi}_S] = i$ . The Hamiltonian of the system then consists of its rest mass  $(m)$  energy, and a term  $\hat{H}_S^{\text{clock}}$  which drives evolution of the internal proper time observable,

$$
\hat{H}_S = (mc^2)\hat{\mathbb{I}}_S + \hat{H}_S^{\text{clock}}\,,\tag{8}
$$

For simplicity, we take the internal proper time degree of freedom to be an ideal clock where different proper times are completely distinguishable, which leads the Hamiltonian to be the conjugate operator  $\hat{\pi}_S$  that drives translation of the proper time observable,

$$
\hat{H}_S^{\text{clock}} = \hat{\pi}_S \,. \tag{9}
$$

Such an ideal clock would in fact be subject to Dirac's criticism mentioned earlier, i.e. it would lead to an unbounded Hamiltonian. But we note that realistic clocks will always have a finite resolution and period, for which we can use Generalized Pauli Operators [\[15\]](#page-11-9) to construct a finite-dimensional version of such canonical conjugate variables. Realistic clock Hamiltonians will thus still be bounded. But we will set aside this technicality in the following and continue to assume ideal clocks. With the above system in mind, the Page-Wootters constraint operator takes the following form,

$$
\hat{\mathbb{J}}_0 = \hat{p}_t \otimes \hat{\mathbb{I}}_S + \hat{\mathbb{I}}_t \otimes ((mc^2) \hat{\mathbb{I}}_S + \hat{\pi}_S) \approx 0, \qquad (10)
$$

for which the physical states take the following form of Eq [\(7\)](#page-2-0) with the system Hamiltonian being  $\hat{H}_S = (mc^2) \hat{\mathbb{I}}_S + \hat{\pi}_S$ .

To probe the relationship between coordinate time t and the proper time  $\tau$  of the system, we consider the probability that the system's proper time  $\tau<sub>S</sub>$  reads some value  $\tau$  conditioned on the coordinate time reading  $t_c = t$ . We can compute this conditional probability from the global physical state of Eq. [\(7\)](#page-2-0) by applying the projective measurements  $\hat{E}_t(t) = |t\rangle \langle t|$  and

 $\hat{E}_S(\tau) = |\tau\rangle_S \langle \tau|$  (operators are paired with the relevant identity operators to be able to act on the full Hilbert space  $\mathcal{H}$ ) along with the Born rule [\[14\]](#page-11-8),

$$
\text{Prob}\left[\tau_S = \tau | t_c = t\right] = \frac{\text{Prob}\left[\tau_S = \tau \text{ and } t_c = t\right]}{\text{Prob}\left[t_c = t\right]},
$$
\n
$$
= \frac{\langle\langle\Psi|\hat{E}_t(t)\hat{E}_S(\tau)|\Psi\rangle\rangle}{\langle\langle\Psi|\hat{E}_t(t)|\Psi\rangle\rangle}.
$$
\n(11)

Since the quantum state of the system  $|\psi(t)\rangle$  is normalized on each time slice, which is obtained via a projective measurement with  $\hat{E}_t(t)$ , we can see that  $\langle {\langle \Psi | \hat{E}_t(t) | \Psi \rangle \rangle} = {\langle \psi(t) | \psi(t) \rangle} = 1, \forall t$ . Further, using orthonormality of the coordinate time states  $\langle t'|t\rangle = \delta(t-t')$ , and the projection  $\hat{E}_S(\tau)$ , the conditional probability can be written as matrix elements of the time evolution operator acting on the initial system state,

$$
\begin{aligned} \text{Prob}\left[\tau_S = \tau | t_c = t\right] &= \langle \psi(0) | \hat{U}^\dagger(t) \hat{E}_S(\tau) \hat{U}(t) | \psi(0) \rangle \;, \\ &= \langle \psi(t) | \hat{E}_S(\tau) | \psi(t) \rangle \\ &= \left| \langle \tau_S = \tau | e^{-i\hat{H}_S t} | \psi(0) \rangle \right|^2. \end{aligned} \tag{12}
$$

<span id="page-4-0"></span>We take the initial  $(t = 0)$  state  $|\psi(0)\rangle$  of the internal proper clock, to be a fiducial Gaussian wave packet centered around  $\tau = 0$  with a width of  $\sigma$  in the eigenbasis of the proper time observable  $\hat{\tau}_S$ ,

$$
|\psi(0)\rangle = \frac{1}{(2\pi\sigma^2)^{1/4}} \int d\tilde{\tau} \, e^{-\frac{\tilde{\tau}^2}{4\sigma^2}} \, |\tilde{\tau}\rangle \; . \tag{13}
$$

The action of  $e^{-i\hat{H}_S t}$  on  $|\psi(0)\rangle$  can be computed by using the fact that  $\hat{\pi}_S$  generates translations in eigenstates of  $\hat{\tau}_S$ , as  $e^{-i\hat{\pi}_S t} |\tilde{\tau}\rangle = |\tilde{\tau} + t\rangle$ ,

$$
e^{-i\hat{H}_S t} |\psi(0)\rangle = \frac{e^{-imc^2 t}}{(2\pi\sigma^2)^{1/4}} \int d\tilde{\tau} e^{-\frac{\tilde{\tau}^2}{4\sigma^2}} |\tilde{\tau} + t\rangle , \qquad (14)
$$

which upon taking an inner product with  $|\tau_S = \tau\rangle$  and remembering  $\langle \tau | \tilde{\tau} + t \rangle = \delta(\tilde{\tau} - (\tau - t)),$ our desired conditional probability simplifies to,

<span id="page-4-1"></span>
$$
\text{Prob}\left[\tau_S = \tau | t_c = t\right] = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(\tau - t)^2}{2\sigma^2}}.
$$
\n(15)

We can now use this conditional probability distribution to compute the average proper time as read by the system  $S$  conditioned on the coordinate time reading  $t$ ,

$$
\langle \tau_S \rangle = t \,. \tag{16}
$$

No surprises here. On average, the proper time of the system, quantum mechanical or not, is the same as the coordinate time, as one would expect from a non-gravitational, non-relativistic setup.

#### Mass-Energy Coupling with Coordinate Time

Let us now introduce an *interaction term* between the coordinate clock and the system (see also [\[16\]](#page-11-10) for a similar idea) at the level of the Wheeler-DeWitt constraint,

<span id="page-5-0"></span>
$$
\hat{\mathbb{J}} = \hat{p}_t \otimes \hat{\mathbb{I}}_S + \hat{\mathbb{I}}_t \otimes \hat{H}_S + \frac{1}{\Lambda} \left( \hat{p}_t \otimes \hat{H}_S \right) \approx 0, \qquad (17)
$$

where we have added a term  $(\hat{p}_t \otimes \hat{H}_S)$  /A to the standard Page-Wootters constraint of Eq. [\(2\)](#page-2-1) to model gravitational back reaction of the system on the coordinate degree of freedom  $\mathcal{H}_t$ . We will motivate later that the interaction strength  $\Lambda^{-1}$  of that term should be  $\mathcal{O}(G/R)$ , with G being Newton's constant and R being a typical spatial scale of the system. Physical states are now those which are annihilated by the above constraint. The rationale for our form of the interaction term comes from linearized quantum gravity which couples first order energy terms to linear metric perturbations.

The effect of this *interaction term* is in fact equivalent to modifying the generator of translations of coordinate time (the "momentum"  $\hat{p}_t$  of the time degree of freedom). To see this, we rewrite the modified constraint as,

<span id="page-5-1"></span>
$$
\hat{\mathbb{J}} = \hat{p}_t \left( \hat{\mathbb{I}}_t + \frac{\hat{p}_t}{\Lambda} \right)^{-1} \otimes \hat{\mathbb{I}}_S + \hat{\mathbb{I}}_t \otimes \hat{H}_S \approx 0, \qquad (18)
$$

which now again has the form of a non-interacting constraint. So our coupling of energy with the background metric seems to simply alter the rate of flow of coordinate time. Since the interaction term commutes with the standard Page-Wootters constraint of Eq. [\(2\)](#page-2-1),

$$
\left[ \left( \hat{p}_t \otimes \hat{\mathbb{I}}_S + \hat{\mathbb{I}}_t \otimes \hat{H}_S \right) , \frac{1}{\Lambda} \left( \hat{p}_t \otimes \hat{H}_S \right) \right] = 0 , \qquad (19)
$$

the eigenstates of this interacting constraint are of a separable, tensor product form,

$$
|\Psi\rangle\rangle_n = |p_t = \alpha\rangle |E_n\rangle , \qquad (20)
$$

where  $|p_t = \alpha\rangle$  is an eigenstate of the unmodified conjugate operator  $\hat{p}_t$  of coordinate time with eigenvalue  $\alpha \in \mathbb{R}$ , and  $|E_n\rangle$  is an eigenstate of the system Hamiltonian  $\hat{H}_S$  with eigenvalue  $E_n$  for some index  $n^1$  $n^1$ . For these eigenstates to be physical, they must be annihilated by the constraint of Eq. [\(17\)](#page-5-0), which then constrains the allowed values of coordinate clock momentum in terms of energy of the system,

<sup>&</sup>lt;sup>1</sup>We are labeling the energy eigenvalues of  $\hat{H}_S$  by a discrete index n for convenience, but we can work with a continuous spectrum just as well.

<span id="page-6-1"></span>
$$
\alpha \left( 1 + \frac{\alpha}{\Lambda} \right)^{-1} + E_n = 0 \qquad \Longrightarrow \qquad \alpha_n = -\left( \frac{E_n}{1 + E_n/\Lambda} \right) \,. \tag{21}
$$

Generic physical states (annihilated by the modified constraint of Eq. [\(18\)](#page-5-1), or equivalently by Eq. [\(17\)](#page-5-0) then have the form,

$$
|\Psi\rangle\rangle = \int dt \, |t\rangle \otimes \left(\sum_{n} c_n e^{-i\left(\frac{E_n}{1+E_n/\Lambda}\right)t}\right) |E_n\rangle \;, \tag{22}
$$

which can be equivalently expressed as (coordinate) time evolution under a modified Hamiltonian,

<span id="page-6-0"></span>
$$
|\Psi\rangle\rangle = \int dt \, |t\rangle \otimes \left(e^{-i\hat{H}_{\text{eff}}t} |\psi(0)\rangle\right) \,, \tag{23}
$$

where  $\hat{H}_{\text{eff}}$  is the effective Hamiltonian of the system's evolution induced by interaction of the system with the coordinate clock,

$$
\hat{H}_{\text{eff}} = \hat{H}_S \left( \hat{\mathbb{I}}_S + \frac{\hat{H}_S}{\Lambda} \right)^{-1} . \tag{24}
$$

Recall that the energy dependent phase, "exp  $(-iEt)$ ," in quantum mechanics is responsible for time evolution and is a measure of coherence between states. The coupling between system energy and background metric's coordinate time led to a modified generator of (coordinate) time translations, which means time runs at different rates for states with different energy content, as is a key feature of gravitation.

#### Emergent Newtonian Gravitational Potential

Let us now investigate the consequences of this modified constraint in the context of two objects, one serving as a massive body, and the other as a test particle reacting to the metric as it responds to the energy of the massive body. To do this, we assume our system to consist of two non-interacting particles A and B with rest masses  $m_A$  and  $m_B$  respectively held at a constant distance from each other. To take into account their relative motion, we would need to have a proper quantum treatment for space as well, but we are focusing on time in this essay. The system Hilbert space  $\mathcal H$  is now decomposed as,

$$
\mathcal{H}_S \simeq \mathcal{H}_A \otimes \mathcal{H}_B \,,\tag{25}
$$

where  $\mathcal{H}_A$  and  $\mathcal{H}_B$  represent the Hilbert spaces corresponding to the internal degrees of freedom of the two particles, respectively. In particular, we associate each particle to have internal structure which serves as a "clock" it carries with it to measure its proper time. On each of these Hilbert spaces, we can associate a proper time observable and its corresponding conjugate

momentum satisfying Heisenberg canonical commutation, for  $A$ ,  $[\hat{\tau}_A, \hat{\pi}_A] = i$ , and for B, we have  $[\hat{\tau}_B, \hat{\pi}_B] = i$ . The Hamiltonian for this non-interacting system of two particles is then simply,

$$
\hat{H}_{AB} = \hat{h}_A \otimes \hat{\mathbb{I}}_B + \hat{\mathbb{I}}_A \otimes \hat{h}_B \equiv \hat{H}_A + \hat{H}_B, \qquad (26)
$$

where the self-Hamiltonians of the two particles consist of their rest mass energy and a term  $\hat{H}^{\text{clock}}$  which drives evolution of their internal proper time observable,

$$
\hat{h}_A = (m_A c^2) \hat{\mathbb{I}}_A + \hat{H}_A^{\text{clock}} \,, \quad \text{and} \quad \hat{h}_B = (m_B c^2) \hat{\mathbb{I}}_B + \hat{H}_B^{\text{clock}} \,. \tag{27}
$$

For ideal clocks, as discussed before, we take the Hamiltonian of the proper time evolution of each particle to be given by the corresponding conjugate  $\hat{\pi}$ 

$$
\hat{H}_j^{\text{clock}} = \hat{\pi}_j \,, \qquad j \in \{A, B\} \,. \tag{28}
$$

The interacting Page-Wootters constraint of Eq. [\(17\)](#page-5-0) with this two-body system now takes the form,

$$
\hat{\mathbb{J}} = \hat{p}_t \otimes \hat{\mathbb{I}}_{AB} + \hat{\mathbb{I}}_t \otimes (\hat{H}_A + \hat{H}_B) + \frac{1}{\Lambda} (\hat{p}_t \otimes (\hat{H}_A + \hat{H}_B)), \qquad (29)
$$

which as we discussed in Eq. [\(23\)](#page-6-0) leads to physical states governed by a modified system Hamiltonian, in this case,

$$
\hat{H}_{\text{eff}} = \hat{H}_{AB} \left( \hat{\mathbb{I}}_{AB} + \frac{\hat{H}_{AB}}{\Lambda} \right)^{-1} . \tag{30}
$$

If the interaction strength  $\Lambda^{-1}$  is weak, as is relevant for the interesting regime of weak-field gravity, the modification to the Hamiltonian can be retained at leading order as a good approximation,

$$
\hat{H}_{\text{eff}} \approx \hat{H}_{AB} \left( \hat{\mathbb{I}}_{AB} - \frac{\hat{H}_{AB}}{\Lambda} \right) . \tag{31}
$$

When expressed in terms of the individual particle Hamiltonians, this takes the following interesting form:

<span id="page-7-0"></span>
$$
\hat{H}_{\text{eff}} = \left(\hat{H}_A - \frac{\hat{H}_A^2}{\Lambda}\right) + \left(\hat{H}_B - \frac{\hat{H}_B^2}{\Lambda}\right) - \frac{2}{\Lambda}\left(\hat{H}_A \otimes \hat{H}_B\right) \,. \tag{32}
$$

The modified Hamiltonian not only alters the self-Hamiltonians of the two particles, but more importantly, introduces a purely quantum-mechanical interaction term between the two particles which matches Newton's gravitational potential energy under the identification,

$$
\frac{1}{\Lambda} = \frac{G}{2r},\tag{33}
$$

 $r$  being the constant (classical) separation between the two particles. It is worth emphasizing that this Newtonian gravitational interaction has emerged as a consequence of interaction added at the level of the Wheeler de-Witt constraint between the coordinate clock and the system of particles. No direct, a-priori interaction was introduced between the particles, but it has emerged as a consequence of matter interacting with the coordinate clock. Note in particular that the charges of the emergent force are the self-Hamiltonians of the particles, as is appropriate for gravity.

### Gravitational Time Dilation

With all the ingredients in place, we can now ask our central question again: how do proper times correlate with coordinate time when mass-energy back reacts on the metric (as modeled by the interacting constraint)? To do so, we calculate the conditional probability of the test particle reading a certain proper time  $\tau$  given the coordinate time reads t. Our physical states now take the following form,

$$
|\Psi\rangle\rangle = \int dt \, |t\rangle \otimes \left(e^{-i\hat{H}_{\text{eff}}t} \, |\psi_{AB}(0)\rangle\right) \,, \tag{34}
$$

where  $\hat{H}_{\text{eff}}$  is the leading order modified Hamiltonian for the two-particle system given in Eq. [\(32\)](#page-7-0) as induced by the interacting constraint. To make contact with the standard Schwarzchild metric in general relativity as our working example, consider particle  $A$  to be the highly massive mass with mass  $m_A = M$ , but with no internal structure, *i.e.*  $\hat{H}_A^{\text{clock}} = 0$ . One can neglect the internal clock structure of A on the grounds that its rest mass energy dominates over any other energy scales involved in its spectrum in accordance with its role of the stationary, central mass. Particle B serves as a test mass which probes proper time dilation effects due to gravity. It therefore has a negligibly small rest mass  $m_B = m \ll M$  so that is does not significantly back-react on the coordinate clock, but only responds to the back-reaction of particle A. With this setup, the bare self-Hamiltonians of the two particles become,

$$
\hat{h}_A = Mc^2, \quad \text{and} \quad \hat{h}_B = \left( mc^2 \right) \hat{\mathbb{I}}_B + \hat{\pi}_B , \tag{35}
$$

and the interaction-induced modified Hamiltonian has the following relevant terms which act on particle B,

<span id="page-8-0"></span>
$$
\hat{H}_{\text{eff}}^{(B)} = \left(1 - \frac{2(M+m)c^2}{\Lambda}\right)\hat{\pi}_B - \frac{\hat{\pi}_B^2}{\Lambda} + \text{ mass dependent constant},\qquad(36)
$$

where the mass-dependent constant will only add an overall, irrelevant phase to the quantum state. To probe the relationship between proper time of the test mass B and the background (metric) coordinate time, we consider the probability that the proper time of particle  $B$  reads some value  $\tau$  conditioned on the coordinate time reading  $t_c = t$ . Similar to the calculation done in Eqs. [\(12\)](#page-4-0) - [\(15\)](#page-4-1), the conditional probability takes the form,

$$
\text{Prob}\left[\tau_B = \tau | t_c = t\right] = \langle \psi_B(t) | \hat{E}_B(\tau) | \psi_B(t) \rangle
$$
  
= 
$$
\left| \langle \tau_B = \tau | e^{-i\hat{H}_{\text{eff}}^{(B)}t} | \psi_B(0) \rangle \right|^2,
$$
 (37)

where  $\hat{H}_{\text{eff}}^{(B)}$  is the modified Hamiltonian on the test particle given in Eq. [\(36\)](#page-8-0). We take the initial quantum state (i.e. at  $t = 0$  coordinate time) of the test particle B (or rather: of its internal proper clock) to be a fiducial Gaussian wave packet centered around  $\tau_B = 0$  with a width of  $\sigma$  in the eigenbasis of the proper time observable,  $\hat{\tau}_B$ . Therefore, the initial quantum state of the system is simply (up to an overall phase),

$$
|\psi(0)\rangle_{AB} = |\psi(0)\rangle_B = \frac{1}{(2\pi\sigma^2)^{1/4}} \int d\tilde{\tau} e^{-\frac{\tilde{\tau}^2}{4\sigma^2}} |\tau_B = \tilde{\tau}\rangle
$$
 (38)

The conditional probability can be obtained by noting that the  $\hat{\pi}_B$  term in the effective Hamiltonian shifts the  $\hat{\tau}_B$  eigenstates, and the  $\hat{\pi}_B^2$  term makes the wave packet spread out in time,

$$
\text{Prob}\left[\tau_B = \tau | t_c = t\right] = \frac{1}{\sqrt{2\pi\sigma(t)^2}} \exp\left(-\frac{\left[\tau - t\left(1 - \frac{2Mc^2}{\Lambda}\right)\right]^2}{2\sigma(t)^2}\right). \tag{39}
$$

where  $\sigma(t) = \sigma \sqrt{1 + \left(\frac{t}{\sigma^2}\right)^2}$  $\frac{t}{\sigma^2\Lambda}$ <sup>2</sup> is now a time-dependent variance of the Gaussian probability distribution. Here, we have explicitly taken the limit  $M \gg m$  to account for the fact that particle  $B$  is a test mass with a negligibly small mass compared to the central mass  $M$ . We now use this conditional probability distribution to compute the average proper time as read by the test mass conditioned on the coordinate time reading t,

$$
\langle \tau_B \rangle = t \left( 1 - \frac{2Mc^2}{\Lambda} \right) \,. \tag{40}
$$

This, to leading order, matches the gravitational time dilation as recorded by a stationary test particle in a Schwarzchild metric with a central mass M, under our identification  $\Lambda^{-1} = G/2r$ ,

$$
d\tau = dt \sqrt{1 - \frac{2GMc^2}{r}} \approx dt \left(1 - \frac{GMc^2}{r}\right). \tag{41}
$$

With a simple model of the background coordinate time as a quantum degree of freedom coupling with mass-energy, we showed that gravitational time dilation is recovered as a consequence of a modification to the generator of coordinate time translations, as seen in Eq. [\(18\)](#page-5-1),

$$
\hat{p}_{t,\text{eff}} = \hat{p}_t \left( \hat{\mathbb{I}}_t + \frac{\hat{p}_t}{\Lambda} \right)^{-1} = \hat{p}_t \left( \hat{\mathbb{I}}_t - \frac{\hat{p}_t}{\Lambda} + \mathcal{O}\left(\frac{\hat{p}_t^2}{\Lambda^2}\right) \right) . \tag{42}
$$

This not only modifies the rate of flow of coordinate time, but also induces a deformed canonical commutation relation which is a general expectation from the theory of quantum gravity, with implications such as minimal resolution, etc. [\[17\]](#page-11-11),

$$
\left[\hat{t}, \hat{p}_{t, \text{eff}}\right] = i\left(\hat{\mathbb{I}}_t - \frac{2}{\Lambda}\hat{p}_t + \mathcal{O}\left(\frac{\hat{p}_t^2}{\Lambda^2}\right)\right). \tag{43}
$$

In addition to the Newtonian gravitational potential naturally emerging in Eq. [\(32\)](#page-7-0) as a feature of this construction, we note that the modification to the Hamiltonian, including the self-term, is reminiscent of renormalization. Following Eq. [\(21\)](#page-6-1), we see that even highly energetic states with  $E_n \to \infty$  induce time evolution at a rate governed not by their bare energy, but are rather controlled by the interaction scale as  $\alpha_n \to \Lambda$ , potentially softening out divergences at high energies.

In summary, we have shown that coupling otherwise non-interacting quantum systems to a global time degree of freedom leads to a number of features expected in (quantum) gravity: gravitational time dilation, agreeing with the relation between proper and coordinate time in the Schwarzschild metric at leading order in G; emergence of a Newtonian potential, sourced in particular by mass-energy as is appropriate for gravity; renormalization of divergent energies. This also opens the possibility of particles in superposition back-reacting on the background coordinates leading to an additional quantum correction to gravitational time dilation [\[14\]](#page-11-8), which could serve as an experimental probe (such as using atom interferometers [\[18\]](#page-11-12)) of quantum gravity in the low energy regime. What is missing from our considerations is a fully quantum treatment of time and space, as opposed to just time. This will be a target of future work and we conjecture that such a combined analysis of space and time will solidify our assumption that the coupling strength  $\Lambda^{-1}$  should scale as G divided by a characteristic spatial scale of the system.

#### Acknowledgments

AS acknowledges funding support by the MJ Murdock Trust, and by Whitman College. OF was supported by the Excellence Cluster ORIGINS which is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC-2094-390783311, as well as by a Fraunhofer-Schwarzschild Fellowship of the LMU Observatory.

## References

- <span id="page-10-0"></span>[1] F. M. Kronz and T. A. Lupher, "Unitarily inequivalent representations in algebraic quantum theory," International Journal of Theoretical Physics 44 [no. 8, \(Aug, 2005\)](http://dx.doi.org/10.1007/s10773-005-4683-0) [1239–1258.](http://dx.doi.org/10.1007/s10773-005-4683-0) <https://doi.org/10.1007/s10773-005-4683-0>.
- <span id="page-10-1"></span>[2] A. A. Deriglazov and K. E. Evdokimov, "Local symmetries and the noether identities in the hamiltonian framework," International Journal of Modern Physics A 15 no. 25, (2000) 4045–4067.
- [3] A. A. Deriglazov, "Improved extended hamiltonian and search for local symmetries," Journal of mathematical physics 50 no. 1, (2009) 012907.
- <span id="page-11-0"></span>[4] A. Deriglazov, "Notes on lagrangean and hamiltonian symmetries," arXiv preprint hep-th/9412244 (1995).
- <span id="page-11-1"></span>[5] B. S. DeWitt, "Quantum theory of gravity. ii. the manifestly covariant theory," Physical Review 162 no. 5, (1967) 1195.
- <span id="page-11-7"></span>[6] P. A. M. Dirac, "Generalized hamiltonian dynamics," Canadian journal of mathematics 2 (1950) 129–148.
- <span id="page-11-2"></span>[7] P. Dirac, "Lectures on quantum mechanics (yeshiva univ., new york, 1964); ld faddeev and r. jackiw," Phys. Rev. Lett 60 (1988) 1692.
- <span id="page-11-3"></span>[8] C. J. Isham, "Canonical quantum gravity and the problem of time," [arXiv:gr-qc/9210011 \[gr-qc\]](http://arxiv.org/abs/gr-qc/9210011). [NATO Sci. Ser. C409,157(1993)].
- [9] E. Anderson, The Problem of Time. Fundamental Theories of Physics, Vol 190. Springer International Publishing, 2017. <https://www.springer.com/us/book/9783319588469>.
- <span id="page-11-4"></span>[10] K. Kuchaˇr, "Proceedings of the 4th canadian conference on general relativity and relativistic astrophysics," 1992.
- <span id="page-11-5"></span>[11] D. N. Page and W. K. Wootters, "Evolution without evolution: Dynamics described by stationary observables," Physical Review D 27 no. 12, (1983) 2885.
- [12] W. K. Wootters, ""time" replaced by quantum correlations," International journal of theoretical physics 23 no. 8, (1984) 701–711.
- <span id="page-11-6"></span>[13] D. Page, "Itp preprint nsf-itp-89-18," Time as an inaccessible observable (1989) .
- <span id="page-11-8"></span>[14] A. R. H. Smith and M. Ahmadi, "Quantum clocks observe classical and quantum time dilation," Nature Commun. 11 [no. 1, \(2020\) 5360,](http://dx.doi.org/10.1038/s41467-020-18264-4) [arXiv:1904.12390 \[quant-ph\]](http://arxiv.org/abs/1904.12390).
- <span id="page-11-9"></span>[15] A. Singh and S. M. Carroll, "Modeling Position and Momentum in Finite-Dimensional Hilbert Spaces via Generalized Clifford Algebra," [arXiv:1806.10134 \[quant-ph\]](http://arxiv.org/abs/1806.10134).
- <span id="page-11-10"></span>[16] A. R. Smith and M. Ahmadi, "Quantizing time: Interacting clocks and systems," Quantum 3 (2019) 160.
- <span id="page-11-11"></span>[17] M. Maggiore, "A Generalized uncertainty principle in quantum gravity," [Phys.](http://dx.doi.org/10.1016/0370-2693(93)91401-8) Lett. B 304 [\(1993\) 65–69,](http://dx.doi.org/10.1016/0370-2693(93)91401-8) [arXiv:hep-th/9301067](http://arxiv.org/abs/hep-th/9301067).
- <span id="page-11-12"></span>[18] G. M. Tino, "Testing gravity with cold atom interferometry: results and prospects," Quantum Science and Technology 6 [no. 2, \(Mar, 2021\) 024014.](http://dx.doi.org/10.1088/2058-9565/abd83e) <https://dx.doi.org/10.1088/2058-9565/abd83e>.