

No long hair behaviors of ultra-compact objects

Guohua Liu^{1*} and Yan Peng^{2†}

¹ *School of Physics and Physical Engineering,*

Qufu Normal University, Qufu, Shandong, 273165, China and

² *School of Mathematical Sciences, Qufu Normal University, Qufu, Shandong 273165, China*

Abstract

We investigate distributions of matter fields outside spherically symmetric ultra-compact objects in the asymptotically flat background. Based on the dominant energy condition and the non-negative trace condition, we analytically find a no long hair behavior, which states that the effective radius of matter field hairs cannot extend beyond the outermost null circular orbit.

PACS numbers: 11.25.Tq, 04.70.Bw, 74.20.-z

arXiv:2304.04198v1 [gr-qc] 9 Apr 2023

* liuguohua1234@163.com

† yanpengphy@163.com

I. INTRODUCTION

According to the classical general relativity, a characteristic property of highly curved spacetimes is the existence of null circular orbits outside ultra-compact objects, such as black holes and horizonless ultra-compact stars [1, 2]. The null circular orbits provide valuable information on the properties of the spacetime, such as the gravitational lensing, the black hole shadow and the gravitational waves. Hence, the null circular orbits have attracted a lot of attentions from physicists and mathematicians [3]-[22].

The uniqueness theorem states that black holes are completely characterized by only three parameters: mass, electric charge and angular momentum, which are all conserved quantities defined at the infinity associated with matter fields subject to a Gauss law [23–25]. In contrast, other exterior matter fields not related to a Gauss law are usually called black hole hairs. The null circular orbit divides matter field hairs into two parts: hairs below the orbit and hairs above the orbit. It is supposed that fields below the orbit tending to fall into the horizon and fields above the orbit tending to radiate away to infinity. The authors in [26] have proposed a physical picture that self-interaction between these two parts bind together fields leading to the existence of black hole matter field hairs. So the null circular orbits may be useful in describing the distributions of matter field hairs outside black holes. In accordance with this picture, the authors in [26, 27] proposed a no short hair theorem that the effective radius of matter field hairs must extend beyond the innermost null circular orbit of spherically symmetric hairy black holes. In the horizonless ultra-compact star background, such lower bound on the effective radius of hairs also exists on the non-negative trace condition [28]. Interestingly, it was also found that over half of the matter field hair gravitational mass is contained above the null circular orbit of spherically symmetric hairy black holes [27, 29–31]. As a further step of lower bound analysis in [26–28], it is meaningful to examine whether there is any upper bound on the effective radius of matter field hairs.

This paper is to study exterior matter field hairs outside ultra-compact objects. The null circular orbits are useful in describing the distributions of hairs. We analytically find a no long hair behavior that the effective radius of matter field hairs must not extend beyond the outermost null circular orbit.

II. INVESTIGATIONS ON EFFECTIVE RADIUS OF MATTER FIELD HAIRS

We consider the gravity system with spherically symmetric ultra-compact objects surrounded by exterior matter field hairs. The curved spacetime is characterized by the line element [26, 27]

$$ds^2 = -g(r)e^{-2\chi(r)}dt^2 + \frac{dr^2}{g(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (1)$$

where $\chi(r)$ and $g(r)$ are metric functions depending on the radial coordinate r . In the asymptotically flat background, the metric functions satisfy $\chi(r \rightarrow \infty) = 0$ and $g(r \rightarrow \infty) = 1$ at the infinity.

The Einstein equations $G_{\nu}^{\mu} = 8\pi T_{\nu}^{\mu}$ yield the differential equations

$$g' = -8\pi r\rho + \frac{1-g}{r}, \quad (2)$$

$$\chi' = \frac{-4\pi r(\rho + p)}{g}, \quad (3)$$

where ρ , p and p_{τ} are interpreted as the energy density $\rho = -T_t^t$, the radial pressure $p = T_r^r$ and the tangential pressure $p_{\tau} = T_{\theta}^{\theta} = T_{\phi}^{\phi}$ respectively [27]. In this work, we take the dominant energy condition that the pressures are bounded by the non-negative energy density as [26, 27]

$$\rho > |p|, \quad |p_{\tau}| \geq 0. \quad (4)$$

In this paper, we only focus on the region above the outermost null circular orbit and it was proved that exterior null circular orbit outside black hole horizons always exists [32–36]. So we can take the condition (4) without contradiction with the fact that there is $\rho = -p$ on black hole horizons [26, 27].

We also take the non-negative trace condition [37–40]

$$T = -\rho + p + 2p_{\tau} \geq 0. \quad (5)$$

An example satisfying the relation (5) is the gravitating Einstein-Yang-Mills solitons with $T = 0$ [41]. We point out that the non-positive trace condition $T \leq 0$ is usually imposed in the curved spacetimes [26, 27].

The metric function $g(r)$ can be putted in the form [38]

$$g = 1 - \frac{2m(r)}{r}, \quad (6)$$

where $m(r)$ is the gravitational mass contained within a sphere of radial radius r .

With the equation (2), one can further express the mass term by the integral relation

$$m(r) = \int_0^r 4\pi r'^2 \rho(r') dr'. \quad (7)$$

In the following, we deduce the null circular orbit equations [27]. The energy and angular momentum are conserved on the geodesic trajectories since the metric is independent of t and ϕ . And the null circular orbit equations can be obtained from the effective potential

$$V_r = (1 - e^{2\chi})E^2 + g\frac{L^2}{r^2} \quad (8)$$

along with the characteristic relations

$$V_r = E^2 \quad \text{and} \quad V'_r = 0, \quad (9)$$

where E and L are conserved energy and conserved angular momentum respectively.

Substituting equations (2) and (3) into (8) and (9), we get the null circular orbit equation

$$N(r) = 3g(r) - 1 - 8\pi(r)^2 p(r) = 0, \quad (10)$$

where the discrete roots r_γ satisfying $N(r_\gamma) = 0$ are the radii of the null circular orbits.

From the expression (7), the finiteness of the gravitational mass implies

$$r^2 \rho(r) \rightarrow 0 \quad \text{as} \quad r \rightarrow \infty. \quad (11)$$

Since p is bounded by ρ , we obtain the asymptotical infinity behavior

$$r^2 p(r) \rightarrow 0 \quad \text{as} \quad r \rightarrow \infty. \quad (12)$$

According to (12), the null circular orbit characteristic equation asymptotically behaves

$$N(r) \rightarrow 2 \quad \text{as} \quad r \rightarrow \infty. \quad (13)$$

We define r_γ^{out} as the outermost null circular orbit radius, which corresponds to the largest positive root of $N(r) = 0$. Above the outermost null circular orbit, $N(r)$ is non-negative as

$$N(r) > 0 \quad \text{for} \quad r \in (r_\gamma^{out}, \infty). \quad (14)$$

The conservation equation $T_{\nu;\mu}^\mu = 0$ has only one nontrivial component

$$T_{r;\mu}^\mu = 0. \quad (15)$$

Substituting equations (2) and (3) into (15), we obtain the pressure equation

$$p'(r) = \frac{1}{2rg} [(3g - 1 - 8\pi r^2 p)(\rho + p) + 2gT - 8gp], \quad (16)$$

where $T = -\rho + p + 2p_\tau$ is the trace of the energy momentum tensor.

We introduce a new pressure function $P(r) = r^4 p$, which is proved to be useful in describing the distributions of matter field hairs. Then the equation (16) can be expressed as

$$P'(r) = \frac{r}{2g} [N(\rho + p) + 2gT] \quad (17)$$

with $N = 3g - 1 - 8\pi r^2 p$.

From (4), (5), (14) and (17), $P(r)$ is an increasing function above outermost null circular orbits satisfying

$$P'(r) > 0 \quad \text{for} \quad r \in (r_\gamma^{out}, \infty). \quad (18)$$

We take the condition that the energy density ρ approaches zero faster than r^{-4} , which is imposed to make sure that there are no extra conserved charges besides the ADM mass (the energy density of Einstein-Maxwell fields associated with electric charges decreases as the speed of r^{-4} at the infinity) defined at the infinity associated with matter fields in the spherically symmetric spacetimes [26, 27]. Also considering the density dominant condition (4), the effective pressure satisfies the asymptotical infinity behavior

$$P(r \rightarrow \infty) = 0. \quad (19)$$

It was found that the effective pressure $|P(r)|$ must have a local maximum value at some extremum point r_0 [26, 27]. We can define $r_m = r_0$ as the effective radii of matter fields. According to (18) and (19), $P(r)$ is an increasing function of r above the outermost null circular orbit and approaches zero at the infinity. So the effective radii $r_m = r_0$ must have an upper bound

$$r_m \leq r_\gamma^{out}. \quad (20)$$

III. CONCLUSIONS

We investigated distributions of matter fields outside spherically symmetric ultra-compact objects. We assumed the dominant energy condition and the non-negative trace condition. We defined an effective matter field radius at an extremum point, where the pressure function $|P(r)|$ possesses a local maximum value. Using analytical methods, we obtained an upper bound on the effective matter field radius expressed as $r_m \leq r_\gamma^{out}$ with r_m as the effective matter field radius and r_γ^{out} corresponding to the outermost null circular orbit radius. So we found a no long hair behavior that the effective matter field radius must not extend beyond the outermost null circular orbit.

Acknowledgments

This work was supported by the Shandong Provincial Natural Science Foundation of China under Grant No. ZR2022MA074. This work was also supported by a grant from Qufu Normal University of China under Grant No. xkjjc201906, the Youth Innovations and Talents Project of Shandong Provincial Colleges and Universities (Grant no. 201909118), Taishan Scholar Project of Shandong Province (Grant No.tsqn202103062) and the Higher Educational Youth Innovation Science and Technology Program Shandong Province (Grant No. 2020KJJ004).

-
- [1] J. M. Bardeen, W. H. Press and S. A. Teukolsky, Rotating black holes: Locally nonrotating frames, energy extraction, and scalar synchrotron radiation, *Astrophys. J.* 178,347(1972).
 - [2] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, (Oxford University Press, New York, 1983).
 - [3] Goebel, C. J., Comments on the “vibrations” of a Black Hole, *Astrophysical Journal*, vol. 172, p.L 95.
 - [4] Teo, E., Spherical Photon Orbits Around a Kerr Black Hole, *General Relativity and Gravitation* (2003) 35: 1909.
 - [5] Pedro V.P. Cunha, Carlos A.R. Herdeiro, Eugen Radu, Fundamental photon orbits: black hole shadows and spacetime instabilities, *Phys. Rev. D* 96(2017)no.2,024039.
 - [6] Jai Grover, Alexander Wittig, Black Hole Shadows and Invariant Phase Space Structures, *Phys. Rev. D* 96(2017)no.2,024045.
 - [7] Pedro V.P. Cunha, Carlos A.R. Herdeiro, Maria J. Rodriguez, Does the black hole shadow probe the event horizon geometry?, *Phys. Rev. D* 97(2018)no.8,084020.
 - [8] S. L. Shapiro and S. A. Teukolsky, *Black holes, white dwarfs, and neutron stars: The physics of compact objects*, New York, USA: Wiley(1983)645p.
 - [9] V. Cardoso, A. S. Miranda, E. Berti, H. Witek and V. T. Zanchin, Geodesic stability, Lyapunov exponents and quasinormal modes, *Phys. Rev. D* 79, 064016(2009).
 - [10] S. Hod, Spherical null geodesics of rotating Kerr black holes, *Phys. Lett. B* 718,1552(2013).
 - [11] Emanuel Gallo, J. R. Villanueva, Photon spheres in Einstein and Einstein-Gauss-Bonnet theories and circular null geodesics in axially-symmetric spacetimes, *Phys. Rev. D* 92(2015) no.6,064048.
 - [12] Zdenek Stuchlik, Jan Schee, Bobir Toshmatov, Jan Hladik, Jan Novotny, Gravitational instability of polytropic spheres containing region of trapped null geodesics: a possible explanation of central supermassive black holes in galactic halos, *JCAP* 1706(2017)no.06, 056.
 - [13] Zdenek Stuchlik, Stanislav Hledik, Jan Novotny, General relativistic polytropes with a repulsive cosmological constant, *Phys. Rev. D* 94(2016)no.10,103513.
 - [14] Ivan Zh. Stefanov, Stoytcho S. Yazadjiev, Galin G. Gyulchev, Connection between Black-Hole Quasinormal Modes and Lensing in the Strong Deflection Limit, *Phys. Rev. Lett.* 104(2010)251103.
 - [15] S. Hod, The fastest way to circle a black hole, *Physical Review D* 84, 104024 (2011).
 - [16] Yan Peng, The extreme orbital period in scalar hairy kerr black holes, *Physics Letters B* 792(2019)1-3.
 - [17] V. Cardoso, A.S. Miranda, E. Berti, H. Witek, and V.T. Zanchin, Geodesic stability, Lyapunov exponents and quasinormal modes, *Phys. Rev. D* 79, 064016 (2009).
 - [18] S. Hod, Upper bound on the radii of black-hole photonspheres, *Phys. Lett. B* 727, 345 (2013).
 - [19] P. V. P. Cunha, E. Berti, and C. A. R. Herdeiro, Light-Ring Stability for Ultracompact Objects, *Phys. Rev. Lett.* 119 (2017)251102.
 - [20] Yan Peng, On instabilities of scalar hairy regular compact reflecting stars, *JHEP* 1810(2018)185.
 - [21] Bahram Mashhoon, Stability of charged rotating black holes in the eikonal approximation, *Phys. Rev. D* 31(1985)no.2,290-293.
 - [22] S. Hod, Universal Bound on Dynamical Relaxation Times and Black-Hole Quasinormal Ringing, *Phys. Rev. D* 75(2007)064013.
 - [23] R. Ruffini and J. A. Wheeler, *Physics Today* 24, 30 (1971).
 - [24] W. Israel, *Phys. Rev.* 164, 1776 (1967) and *Commun. Math. Phys.* 8, 245 (1968).
 - [25] B. Carter, *Phys. Rev. Letters* 26, 331 (1971); R. M. Wald, *Phys. Rev. Letters* 26, 1653 (1971).
 - [26] D. Núñez, H. Quevedo, and D. Sudarsky, Black Holes Have No Short Hair, *Phys. Rev. Lett.* 76, 571(1996).
 - [27] S. Hod, Hairy Black Holes and Null Circular Geodesics, *Phys. Rev. D* 84, 124030 (2011).
 - [28] Yan Peng, No short hair behaviors of ultra-compact stars, *Eur.Phys.J.C* 81(2021)3,245.
 - [29] Yun Soo Myung, Taeyoon Moon, Hairy mass bound in the Einstein-Born-Infeld black hole, *Phys. Rev. D* 86.084047.
 - [30] Yan Peng, Hair mass bound in the black hole with nonzero cosmological constants, *Phys. Rev. D* 98(2018)104041.

- [31] Yan Peng, Hair distributions in noncommutative Einstein-Born-Infeld black holes, Nucl.Phys.B 941(2019)1-10.
- [32] S. Hod, Phys. Lett. B 727,345(2013). arXiv:1701.06587.
- [33] P.V.P.Cunha, C.A.R.Herdeiro, Phys. Rev. Lett.124,181101(2020).
- [34] Shahar Hod, Eur. Phys. J. C 82(2022)663.
- [35] S. Hod, Extremal black holes have external light rings, Phys. Rev. D 107,024028(2023), arXiv:2211.15983.
- [36] Yan Peng, The existence of null circular geodesics outside extremal spherically symmetric asymptotically flat hairy black holes, arXiv:2211.14463 [gr-qc].
- [37] S. Hod, Lower bound on the compactness of isotropic ultra-compact objects, Physical Review D 97, 084018(2018).
- [38] S. Hod, Self-gravitating field configurations: The role of the energy-momentum trace, Physics Letters B 739(2014)383.
- [39] Jan Novotný, Jan Hladík, Zdeněk Stuchlík, Polytropic spheres containing regions of trapped null geodesics, Physical Review D 95(2017)043009.
- [40] S. Hod, Analytic study of self-gravitating polytropic spheres with light rings, The European Physical Journal C 78(2018)417.
- [41] R. Bartnik and J. McKinnon, Particlelike Solutions of the Einstein-Yang-Mills Equations, Phys. Rev. Lett. 61(1998)141.