

Fundamental Sensitivity Limits for non-Hermitian Quantum Sensors

Wenkui Ding,^{1,2} Xiaoguang Wang,^{2,*} and Shu Chen^{1,3,†}

¹*Beijing National Laboratory for Condensed Matter Physics,*

Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

²*Department of Physics, Zhejiang Sci-Tech University, 310018 Zhejiang, China*

³*School of Physical Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China*

(Dated: October 30, 2023)

Considering non-Hermitian systems implemented by utilizing enlarged quantum systems, we determine the fundamental limits for the sensitivity of non-Hermitian sensors from the perspective of quantum information. We prove that non-Hermitian sensors do not outperform their Hermitian counterparts (directly couple to the parameter) in the performance of sensitivity, due to the invariance of the quantum information about the parameter. By scrutinizing two concrete non-Hermitian sensing proposals, which are implemented using full quantum systems, we demonstrate that the sensitivity of these sensors is in agreement with our predictions. Our theory offers a comprehensive and model-independent framework for understanding the fundamental limits of non-Hermitian quantum sensors and builds the bridge over the gap between non-Hermitian physics and quantum metrology.

Introduction.— Parallel with the rapid development in quantum technology, quantum metrology [1–4] and quantum sensing [5, 6] are becoming one of the focuses in quantum science. Quantum sensors exploit quantum coherence or quantum correlations to detect weak or nanoscale signals and exhibit great advantages in accuracy, repeatability and precision. Recently, a number of sensing proposals utilizing novel properties of non-Hermitian physics [7–9] have been proposed and experimentally demonstrated. For example, non-Hermitian lattice systems with skin effect [10, 11] or non-reciprocity [12] have been suggested to realize enhanced sensing. Specifically, the divergence of the susceptibility near the exceptional point (EP) is exploited to realize enhanced sensing with arbitrary precision [13–16] and it has been demonstrated using various classical (quasi-classical) physical systems [17–21] or quantum systems [22, 23]. While these early experiments claimed enhancements compared to conventional Hermitian sensors, subsequent theoretical work has cast doubt on these results [24–28], suggesting that the reported enhancements may not have fully taken into account the effects of noise. After taking into account the noise, some theoretical works show the enhancement in sensitivity provided by non-Hermitian sensors may disappear [24, 27]. However, other theoretical works have claimed that the enhancement can persist even in the presence of noise [25, 26]. While some recent experiments have demonstrated enhanced sensitivity despite the presence of noise [21, 22], others have shown no such enhancement [23]. Currently, the fundamental limitations imposed by noise on non-Hermitian sensors are still a topic of debate [29], and a definitive conclusion on whether the non-Hermitian physics is superior for sensing is still elusive.

In sensing schemes that rely on quantum systems, quantum noise always arises during the projective measurement of the parameter-dependent quantum state [30]. This noise originates from quantum mechan-

ics and cannot be eliminated, leading to the fundamental sensitivity limit. Quantum metrology focuses on how to beat the standard quantum limit by employing quantum correlations, like entanglement or squeezing [2]. While non-Hermitian systems can serve as an effective description of open system dynamics in certain situations [8, 31], the decoherence and dissipation in open systems are detrimental to the useful quantum features required for metrology [32–36]. Therefore, the sensitivity enhancement from non-Hermitian sensors, which can be embedded in open systems, is quite counter-intuitive. Various theoretical works have been devoted to analyze the effect from the noise [24–28], however, these investigations usually require modeling the effect of noise and calculating the dynamics using tools such as the quantum Langevin equation, for specific sensing schemes and probe states. Here, we provide a general conclusion on the fundamental sensitivity limit from the perspective of quantum information [37], without the requirement to solve intricate non-unitary quantum dynamics and independent of specific noise forms, probe states, and measurement regimes. We unambiguously prove that the non-Hermitian quantum sensors do not surpass the ultimate sensitivity of their Hermitian counterparts and cannot achieve arbitrary precision in realistic experimental settings with finite quantum resources.

Sensitivity bound for unitary parameter encoding.— Quantum metrology or quantum parameter estimation is to estimate the parameter λ from the parameter-dependent quantum state ρ_λ . One crucial step is to make measurements on the quantum state. The measurement can be described by a Hermitian operator Π , and the probability of obtaining the measurement outcome ξ , conditioned on the parameter λ , is $P(\xi|\lambda) = \text{Tr}(\Pi\rho_\lambda)$. We can evaluate the classical Fisher information \mathcal{I}_λ corresponding to this specific measurement as $\mathcal{I}_\lambda = \sum_\xi P(\xi|\lambda) \left(\frac{\partial \ln P(\xi|\lambda)}{\partial \lambda} \right)^2$, which reflects the amount of in-

formation about the parameter contained in the distribution of measurement outcomes. Meanwhile, the estimation uncertainty is given by $\delta^2\lambda = \left\langle \left(\frac{\lambda_{\text{est}}}{d\langle\lambda_{\text{est}}\rangle/d\lambda} - \lambda \right)^2 \right\rangle$, where λ_{est} is the estimated value when the number of probes (N) and the number of trials (ν) are finite, while λ is the true value of the parameter. For the unbiased estimator, we have $d\langle\lambda_{\text{est}}\rangle/d\lambda = 1$. In fact, the classical Fisher information bounds the estimation uncertainty achievable in this specific measurement, which fulfills the so-called Cramér-Rao bound: $\delta\lambda \geq 1/\sqrt{\nu\mathcal{I}_\lambda}$, where ν is the number of repetitions or trials. This bound can be attained asymptotically as $\nu \rightarrow \infty$. When it is optimized over all possible measurements, we can find the maximal value of the classical Fisher information, known as the quantum Fisher information (QFI) [38], $\mathcal{I}_\lambda \leq F_\lambda$. Accordingly, the ultimate precision of parameter estimation for a specific parameter-dependent quantum state can be determined using the quantum Cramér-Rao bound [39], $\delta\lambda \geq 1/\sqrt{\nu F_\lambda}$. The QFI [40] can be determined as $F_\lambda = \text{Tr}[\rho_\lambda \mathcal{L}^2]$, where \mathcal{L} is the symmetric logarithmic derivative defined by $\partial\rho_\lambda/\partial\lambda = (\mathcal{L}\rho_\lambda + \rho_\lambda\mathcal{L})/2$.

Usually, the parameter-dependent quantum state ρ_λ is obtained through time evolution governed by the parameter-dependent Hamiltonian $\hat{H}_\lambda(t)$. To be more specific, with the parameter-independent initial state (probe state) ρ_0 , the parameter encoding process can be described as $\rho_\lambda(t) = U_\lambda(0 \rightarrow t)\rho_0 U_\lambda^\dagger(0 \rightarrow t)$, where the unitary time evolution operator $U_\lambda(0 \rightarrow t) = \mathcal{T}e^{-i\int_0^t \hat{H}_\lambda(s)ds}$, with \mathcal{T} being the time-ordering operator. In the case where the initial state is a pure state, $\rho(0) = |\Psi_0\rangle\langle\Psi_0|$, the QFI can be calculated as $F_\lambda(t) = 4(\langle\Psi_0|h_\lambda^2(t)|\Psi_0\rangle - \langle\Psi_0|h_\lambda(t)|\Psi_0\rangle^2) \equiv 4\text{Var}[h_\lambda(t)|_{|\Psi_0\rangle}]$, where the Hermitian operator $h_\lambda(t) \equiv iU_\lambda^\dagger(0 \rightarrow t)\frac{\partial}{\partial\lambda}U_\lambda(0 \rightarrow t)$ is called the transformed local generator [41, 42]. We have defined $\text{Var}[\hat{A}]|_{|\Psi\rangle}$ as the variance of the Hermitian operator \hat{A} with respect to $|\Psi\rangle$. It satisfies $\text{Var}[\hat{A}]|_{|\Psi\rangle} \leq \|\hat{A}\|^2/4$ for arbitrary $|\Psi\rangle$ [43], where the seminorm is defined as $\|\hat{A}\| \equiv M_A - m_A$, with M_A (m_A) being the maximum (minimum) eigenvalue of \hat{A} . Then it follows $F_\lambda(t) \leq \|h_\lambda(t)\|^2 \equiv F_\lambda^{(c)}(t)$, where $F_\lambda^{(c)}(t)$ is defined as the channel QFI, corresponding to the maximum QFI achievable by optimizing over all possible probe states.

The triangle inequality for the seminorm of Hermitian operators [43] states that $\|\hat{A} + \hat{B}\| \leq \|\hat{A}\| + \|\hat{B}\|$. Using the definition of h_λ and the Schrödinger equation $i\partial U_\lambda/\partial t = H_\lambda U_\lambda$, we can obtain $\frac{\partial h_\lambda}{\partial t} = U_\lambda^\dagger(0 \rightarrow t)\frac{\partial H_\lambda(t)}{\partial\lambda}U_\lambda(0 \rightarrow t)$. Thus, the transformed local generator can be explicitly represented as $h_\lambda(t) = \int_0^t U_\lambda^\dagger(0 \rightarrow s)\frac{\partial H_\lambda(s)}{\partial\lambda}U_\lambda(0 \rightarrow s)ds$. By applying the triangle inequality, we obtain $\|h_\lambda(t)\| \leq \int_0^t \left\| U_\lambda^\dagger(0 \rightarrow s)\frac{\partial H_\lambda(s)}{\partial\lambda}U_\lambda(0 \rightarrow s) \right\| ds = \int_0^t \left\| \frac{\partial H_\lambda(s)}{\partial\lambda} \right\| ds$, where we have used the fact that unitary transformations

do not change the spectrum of an operator. Therefore, the upper bound of the channel QFI can be obtained as follows [44]:

$$F_\lambda^{(c)}(t) \leq \left[\int_0^t \left\| \frac{\partial H_\lambda(s)}{\partial\lambda} \right\| ds \right]^2. \quad (1)$$

Due to the convexity of QFI, the optimal probe state is always a pure state [45]. Therefore, this bound is naturally applicable for mixed probe states. Similar relations [45, 46] have been obtained using different methods and have been employed to discuss unitary parameter encoding processes governed by Hermitian time-dependent Hamiltonians. Furthermore, by utilizing the quantum Cramér-Rao bound, we obtain the lower bound for the estimation uncertainty as follows [47]:

$$\delta\lambda \geq \frac{1}{\sqrt{\nu} \int_0^t \left\| \frac{\partial H_\lambda(s)}{\partial\lambda} \right\| ds}. \quad (2)$$

Here, we realize that this relation is actually not limited to unitary parameter encoding processes. Instead, this bound can be applied to investigate non-unitary parameter encoding processes, particularly in the context of open quantum systems or dynamics governed by non-Hermitian Hamiltonians.

In this Letter, we proceed further to investigate the bound on the change rate of the QFI. By the definition of QFI, we obtain that $\frac{\partial F_\lambda}{\partial t} = 8 \text{Cov}[\frac{\partial h_\lambda}{\partial t}, h_\lambda]|_{|\Psi_0\rangle}$, where the covariance is defined as $\text{Cov}[\hat{A}, \hat{B}]|_{|\Psi\rangle} \equiv \frac{1}{2}\langle\Psi|\hat{A}\hat{B} + \hat{B}\hat{A}|\Psi\rangle - \langle\Psi|\hat{A}|\Psi\rangle\langle\Psi|\hat{B}|\Psi\rangle$. The covariance inequality deduced from the Cauchy-Schwarz inequality states that $|\text{Cov}[\hat{A}, \hat{B}]| \leq \sqrt{\text{Var}(\hat{A})\text{Var}(\hat{B})}$. Applying this inequality, we find:

$$\begin{aligned} \left| \text{Cov}\left[\frac{\partial h_\lambda}{\partial t}, h_\lambda(t)\right]|_{|\Psi_0\rangle} \right| &\leq \sqrt{\text{Var}\left[U_\lambda^\dagger\frac{\partial H_\lambda}{\partial\lambda}U_\lambda\right]|_{|\Psi_0\rangle}} \frac{F_\lambda^{1/2}(t)}{2} \\ &\leq \frac{\|\frac{\partial H_\lambda}{\partial\lambda}\| F_\lambda^{1/2}(t)}{2}. \end{aligned} \quad (3)$$

After some algebra [48], we prove the following inequality:

$$\left| \frac{\partial F_\lambda^{1/2}(t)}{\partial t} \right| \leq \left\| \frac{\partial H_\lambda(t)}{\partial\lambda} \right\|. \quad (4)$$

Namely, the change rate of the square root of QFI is only bounded by the spectral width of the derivative of the Hamiltonian with respect to the parameter. $|\partial F_\lambda^{1/2}(t)/\partial t|$ measures how fast the quantum information about the parameter flows into or out of the quantum state. It indicates that the quantum parameter encoding process cannot be accelerated by adding auxiliary parameter-independent Hamiltonian extensions.

Open system and non-Hermitian quantum sensing.—In many situations, such as dynamics in open quantum systems or systems governed by non-Hermitian Hamiltonians, the dynamical process used to encode the parameter may be non-unitary. However, it is often possible to map these non-unitary processes to equivalent unitary dynamics in an enlarged Hilbert space, by introducing extra degrees of freedom that correspond to the environment [32]. We now make this statement more rigorous for non-unitary sensing schemes. Prior to applying the perturbation that incorporates the parameter to be estimated, the dynamical process in the open quantum system or non-Hermitian system, $R_S : \rho_S(0) \rightarrow \rho_S(t)$, can be mapped from a unitary evolution in an enlarged system, $\mathcal{M}(U_{S,E}) \rightarrow R_S$. This unitary time evolution operator for the combined system corresponds to a Hermitian Hamiltonian, $U_{S,E} \rightarrow \hat{H}_{\text{tot}}$. This Hamiltonian, $\hat{H}_{\text{tot}} = H_S(t) + H_E(t) + H_{SE}(t)$, generally contains terms that describe the system $H_S(t)$, the environment $H_E(t)$ and the system-environment interaction $H_{SE}(t)$. Subsequently, we introduce the perturbation that incorporates the parameter dependence. In most scenarios, including the examples discussed in this work and various non-Hermitian sensing protocols, the parameter of interest directly couples to the degrees of freedom of the system and the perturbation can be represented by a Hermitian Hamiltonian $H_1(\lambda, t)$. As a result, the overall parameter encoding process, corresponding to the dynamical evolution in the open system or non-Hermitian system, can be mapped to a unitary dynamics governed by a Hermitian Hamiltonian $H_\lambda(t) = \hat{H}_{\text{tot}} + H_1(\lambda, t)$. By mapping the dynamics to an enlarged system, we circumvent the analysis of intricate non-unitary parameter encoding processes. By resorting to the corresponding unitary evolution in the enlarged system, we can straightforwardly apply the ultimate sensitivity bound in Eq. (2) and the QFI rate bound in Eq. (4).

Since the estimation parameter only associates with the degree of freedom of the system, we have $\partial H_\lambda / \partial \lambda = \partial H_1 / \partial \lambda$. Thus, the bounds in Eq. (2) and (4) reveal an intriguing insight: the ultimate sensitivity cannot be improved by coupling the system to the environment or by introducing auxiliary Hamiltonians. This is because these additional factors do not increase the amount of information about the parameter or the rate of information encoding. Correspondingly, the non-Hermitian sensor will not outperform its Hermitian counterpart in terms of ultimate sensitivity. We now substantiate this conclusion by analyzing some concrete examples.

Example I: single-qubit pseudo-Hermitian sensor.—A single-qubit pseudo-Hermitian [49] Hamiltonian, described by

$$\hat{H}_s = \mathcal{E}_\lambda \begin{pmatrix} 0 & \delta_\lambda^{-1} \\ \delta_\lambda & 0 \end{pmatrix}, \quad (5)$$

is employed to realize enhanced quantum sensing in

Ref. [50], where \mathcal{E}_λ and δ_λ depend on the parameter λ that is being estimated. According to the Naimark dilation theory [51, 52], a dilated two-qubit system with a properly prepared initial state can be used to simulate the dynamics of this pseudo-Hermitian Hamiltonian, conditioned on the post-selection measurement of the ancilla qubit [53]. The Hermitian Hamiltonian of this dilated two-qubit system is

$$\hat{H}_{\text{tot}} = b\hat{I}^{(a)} \otimes \hat{\sigma}_x^{(s)} - c\hat{\sigma}_y^{(a)} \otimes \hat{\sigma}_y^{(s)} + \lambda\hat{I}^{(a)} \otimes \hat{\sigma}_x^{(s)}, \quad (6)$$

where $\hat{\sigma}_{\alpha=x,y,z}^{(s)}$ ($\hat{\sigma}_{\alpha=x,y,z}^{(a)}$) represents the Pauli operators of the system qubit (ancilla qubit). The coefficients $b = 4\omega\varepsilon(1+\varepsilon)/(1+2\varepsilon)$ and $c = 2\omega\sqrt{\varepsilon(1+\varepsilon)}/(1+2\varepsilon)$, where ε and ω describe the qubit. This specific dilated Hamiltonian can be mapped to \hat{H}_s , with $\mathcal{E}_\lambda = \sqrt{(b+\lambda)^2 + c^2}$ and $\delta_\lambda = (\lambda + 2\varepsilon\omega)/\mathcal{E}_\lambda$. The time evolution of the quantum state governed by \hat{H}_s is $|\psi\rangle_s = e^{-i\hat{H}_s t}|0\rangle_s = \cos(\mathcal{E}_\lambda t)|0\rangle_s - i\delta_\lambda \sin(\mathcal{E}_\lambda t)|1\rangle_s$. Thus the normalized population in $|0\rangle_s$ is $S(\lambda, t) = 1/[1 + \delta_\lambda^2 \tan^2(\mathcal{E}_\lambda t)]$. In Fig. 1(a), we plot the susceptibility $\chi_s(\lambda) \equiv \partial S / \partial \lambda$ as a function of λ for a fixed evolution time $t = \tau \equiv \pi/[4\omega\sqrt{\varepsilon(1+\varepsilon)}]$. The result indicates that the maximal value of the susceptibility diverges as $\varepsilon \rightarrow 0$, which corresponds to the eigenstate coalescence. Based on this feature, the authors in Ref. [50] proposed the pseudo-Hermitian enhanced quantum sensing scheme.

On the other hand, for the dilated two-qubit system, the probe state should be prepared as $|\Psi_0\rangle = \left(\sqrt{\frac{1+\varepsilon}{1+2\varepsilon}}|0\rangle_a + \sqrt{\frac{\varepsilon}{1+2\varepsilon}}|1\rangle_a\right) \otimes |0\rangle_s$ in order to correctly simulate the non-Hermitian dynamics. The normalized population $S(\lambda, t)$ actually corresponds to the probability that the system qubit is in state $|0\rangle_s$, conditioned on the ancilla qubit being in state $|0\rangle_a$. Equivalently, by calculating the dynamics of the total system $|\Psi(\tau)\rangle = e^{-i\hat{H}_{\text{tot}}\tau}|\Psi_0\rangle$, we can directly evaluate the probability in state $|0\rangle_a \otimes |0\rangle_s$ as

$$P_1 = \frac{1+\varepsilon}{1+2\varepsilon} \cos^2 \left[t \sqrt{\lambda^2 + \frac{8\varepsilon(1+\varepsilon)\lambda\omega}{1+2\varepsilon} + 4\varepsilon(1+\varepsilon)\omega^2} \right]. \quad (7)$$

Due to the quantum projection noise, there is uncertainty in the determination of P_1 . This uncertainty originates from the quantum projective measurement and follows a binomial distribution. The variance of the estimated probability is $\text{Var}[\hat{P}_1] = P_1(1-P_1)/\nu$, where ν is the number of trials (repetitions) [48]. Using the error propagation formula, we can evaluate the estimation uncertainty for this specific sensing scheme as $\delta\lambda = \sqrt{\text{Var}[\hat{P}_1] / \left| \frac{\partial P_1}{\partial \lambda} \right|}$. We plot the sensitivity in Fig. 1(b), which shows no divergence at the corresponding divergent positions of $\chi_s(\lambda)$ in Fig. 1(a). This absence of divergence in the sensitivity is attributed to the fact that the divergence in $\chi_s(\lambda)$ when $\varepsilon \rightarrow 0$ is accompanied by a vanishing success probability in the




FIG. 1. (a) Susceptibility of the normalized population with respect to λ for different values of ε . It indicates that the maximal susceptibility diverges as ε approaches zero. (b) Sensitivity corresponding to the measurement of the population in the state $|0\rangle_a \otimes |0\rangle_s$. It indicates that the sensitivity at the optimal measurement point (corresponding to the maximal susceptibility) does not diverge when ε approaches zero. The blue lines represent the sensitivity bound of the Hermitian counterpart.

post-selection measurement. Namely, most experimental trails fail to provide useful information on the parameter. As a comparison, the counterpart Hermitian sensor simply employs $\hat{V} = \lambda \hat{I}^{(a)} \otimes \hat{\sigma}_x^{(s)}$ as the parameter encoding generator. The sensitivity bound in Eq. (2) indicates $\delta\lambda \geq \frac{1}{\sqrt{\nu\tau} \|\hat{\sigma}_x\|} = \frac{1}{2\sqrt{\nu\tau}}$. We plot this ultimate sensitivity bound in Fig. 1(b) as the blue lines, indicating that the non-Hermitian sensor does not outperform its Hermitian counterpart. Furthermore, the rate of dynamic QFI can be calculated exactly [48] as follows:

$$\frac{\partial F_\lambda^{1/2}(t)}{\partial t} = 2 \frac{\cos^2 \theta + \sin^2 \theta \frac{\sin(2\Omega t)}{2\Omega t}}{\sqrt{\cos^2 \theta + \sin^2 \theta \frac{\sin^2(\Omega t)}{(\Omega t)^2}}}, \quad (8)$$

where we define $(b + \lambda)/\Omega = \cos \theta$ and $c/\Omega = \sin \theta$. It follows that $-2 \leq \frac{\partial F_\lambda^{1/2}(t)}{\partial t} \leq 2$, which verifies our theory in Eq. (4).

Example II: EP based sensor using a single trapped ion.—We now consider the sensor based on exceptional point realized in a dissipative single-qubit open system in Ref. [54]. The sensing mechanism relies on an effective periodically driven [55] \mathcal{PT} -symmetric non-Hermitian Hamiltonian given by

$$\hat{H}_{\mathcal{PT}} = J[1 + \cos(\omega t)]\hat{\sigma}_x + i\Gamma\hat{\sigma}_z, \quad (9)$$

where $\hat{\sigma}_{x,z}$ are the Pauli operators, J is the coupling strength, ω is the modulation frequency of the coupling strength, and Γ is the dissipation rate. Actually, the practically implemented Hamiltonian in the experiment is $\hat{H}'_{\mathcal{PT}} = \hat{H}_{\mathcal{PT}} - i\Gamma\hat{I}$, which is a passive \mathcal{PT} -symmetric system with \hat{I} being the identity operator. The perturbation applied to the system is $\hat{H}_\delta = \frac{\delta}{2} \cos(\omega_\delta t)(\hat{I} - \hat{\sigma}_z)$, where δ and ω_δ are the amplitude and frequency of the perturbation field, respectively, while ω_δ is the parameter to be estimated. After the system evolves from specific initial states for

a duration of $T = 2\pi/\omega$, we can determine the response energy \mathcal{E}_{res} via $P_J(T) - P_\Gamma(T) = \sin^2(\mathcal{E}_{\text{res}}T)$. Here, the measurable quantities are defined as $P_J(T) = |\langle \uparrow | U(T) | \downarrow \rangle|^2$ and $P_\Gamma(T) = \left| \frac{\langle \uparrow | - \langle \downarrow |}{\sqrt{2}} U(T) \frac{|\uparrow\rangle + |\downarrow\rangle}{\sqrt{2}} \right|^2$, with $U(T) = \mathcal{T} e^{-i \int_0^T [\hat{H}_{\mathcal{PT}}(t) + \hat{H}_\delta(t)] dt}$. The absolute value of the response energy \mathcal{E}_{res} as a function of ω_δ is plotted in Fig. 2(a) [56]. As it is shown, the response energy exhibits sharp dips near the EP [57]. This characteristic feature has motivated the authors in Ref. [54] to suggest the sensing application, since a minor change in ω_δ will result in a significant variation in the response energy. Indeed, in Fig. 2(c), we present the susceptibility $|\partial \mathcal{E}_{\text{res}} / \partial \omega_\delta|$ as a function of ω_δ and it exhibits a divergence near the EP.

However, the study in Ref. [54] has neglected effects from the quantum noise. Here, since P_J and P_Γ actually correspond to projective measurements on the spin state, the quantum projection noise will result in uncertainties in their determination. The variance of the estimated \hat{P}_J and \hat{P}_Γ can be expressed as $\text{Var}[\hat{P}_i] = P_i(C_0 - P_i)/\nu$, with $i = J, \Gamma$, where ν is the number of trials and $C_0 \equiv e^{2\Gamma T}$ [58]. To avoid the complication of dealing with complex response energies, we focus on the region near the EP where $P_J - P_\Gamma > 0$. Applying the theory of uncertainty propagation, we obtain the uncertainty in the estimation of the response energy as $\text{Var}[\hat{\mathcal{E}}_{\text{res}}] = \frac{1}{4\nu T^2} \frac{C_0(P_J + P_\Gamma) - (P_J^2 + P_\Gamma^2)}{(P_J - P_\Gamma)(1 - P_J + P_\Gamma)}$, where we have used the fact that measurements on P_J and P_Γ are independent. We plot the variance of the measured response energy in Fig. 2(b) as a function of ω_δ , and it shows that the uncertainty in the determination of \mathcal{E}_{res} also diverges when ω_δ approaches the EP. The overall sensitivity can be evaluated as $\delta\omega_\delta = \sqrt{\text{Var}[\hat{\mathcal{E}}_{\text{res}}] / |\frac{\partial \mathcal{E}_{\text{res}}}{\partial \omega_\delta}|}$, and we plot it in Fig. 2(d). It shows that the divergence of the susceptibility is completely compensated by the divergence of the uncertainty, resulting in an overall sensitivity without divergence when approaching the EP. On the other hand, the Hermitian counterpart simply uses \hat{H}_δ as the parameter encoding generator. According to Eq. (2), the ultimate sensitivity bound is given by $\delta\omega_\delta \geq \frac{\omega_\delta^{\frac{1}{2}}}{\sqrt{\nu\delta^2[\sin(\omega_\delta T) - \omega_\delta T \cos(\omega_\delta T)]^2}}$. The dashed line in Fig. 2(d) corresponds to this ultimate sensitivity bound. It also demonstrates that the ultimate precision of the Hermitian sensor always exceed the corresponding non-Hermitian sensor.

Summary and discussion.—In summary, we have unveiled the fundamental sensitivity limit for non-Hermitian sensors in the context of open quantum systems. Our results indicate clearly that non-Hermitian sensors do not outperform their Hermitian counterparts. In fact, when comparing the performance of quantum sensors, it is essential to fix the quantum resources consumed by these sensors. Actually, when resources are unlimited, even ideal Hermitian sensors can theoretically achieve arbitrary precision. However, in practical sens-




FIG. 2. (a) Response energy shows sharp dips near the EP for the periodically driven non-Hermitian system. The dashed line indicates the position of the first EP. In (b-d), the range of ω corresponds to the zoom-ins of the left side of the first EP. (b) Variance of the response energy near the EP diverges. (c) Susceptibility of the response energy exhibit divergence near the EP. (d) The sensitivity, which is inversely proportional to the signal-to-noise ratio, shows no divergence. The dashed line represents the theoretical sensitivity bound of the Hermitian counterpart.

ing scenarios, resources are always limited. The number of probes, sensing time, and the number of trials are examples of limited resources. As a result, achieving arbitrary precision is not possible in practical sensing scenarios. The aforementioned instances are characterized by a single probe. Notably, although these cases exhibit divergence in certain measurable quantities, it does not imply that the sensitivity diverges, leading to ‘arbitrary precision’, since the sensitivity of their Hermitian counterparts does not diverge (even without Heisenberg scaling for only $N = 1$).

In Ref. [22], a sensing scheme utilizing an experimentally realized \mathcal{PT} -symmetric system was reported to enhance the sensitivity by a factor of 8.856 over a conventional Hermitian sensor. However, this enhancement is probably attributed to the choice of non-optimal initial probe state used for the Hermitian sensor, and similarly these seeming sensitivity enhancements in Refs. [25, 26] may not exist if making comparison over optimal probe states. Furthermore, non-Hermitian lattice systems utilizing the skin effect[10, 11] or the non-reciprocity [12], have claimed exponential scaling of sensitivity with the lattice size. However, our theory shows that the ultimate sensitivity should not depend on the lattice size, as it is solely determined by the subsystem dimension that directly couples to the parameter. Nevertheless, for non-optimal probe states or measurements, the sensitiv-

ity may still depend on the lattice size.

Although our work demonstrates that coupling to the environment cannot improve the ultimate sensitivity, when the probe state or the measurement protocol is restricted, adding appropriate auxiliary Hamiltonian may be helpful for approaching the ultimate sensitivity bound [59–61]. In fact, when the parameter couples to the environment, the bounds presented in Eq. (2) and (4) remain applicable, albeit $\partial H_\lambda/\partial\lambda$ now depends on the environment’s degrees of freedom. In addition, while our study focuses on non-Hermitian sensors implemented by full quantum systems [62–64], scrutinizing non-Hermitian sensors based on classical or quasiclassical systems [29] through the perspective of conservation of information is a compelling avenue for future research.

The work is supported by National Key Research and Development Program of China (Grant No. 2021YFA1402104), the NSFC under Grants No.12174436, No.11935012 and No.T2121001 and the Strategic Priority Research Program of Chinese Academy of Sciences under Grant No. XDB33000000.

* xgwang@zstu.edu.cn

† schen@iphy.ac.cn

- [1] V. Giovannetti, S. Lloyd, and L. Maccone, Quantum metrology, *Phys. Rev. Lett.* **96**, 010401 (2006).
- [2] L. Pezzè, A. Smerzi, M. K. Oberthaler, R. Schmied, and P. Treutlein, Quantum metrology with nonclassical states of atomic ensembles, *Rev. Mod. Phys.* **90**, 035005 (2018).
- [3] D. Braun, G. Adesso, F. Benatti, R. Floreanini, U. Marzolino, M. W. Mitchell, and S. Pirandola, Quantum-enhanced measurements without entanglement, *Rev. Mod. Phys.* **90**, 035006 (2018).
- [4] M. M. Rams, P. Sierant, O. Dutta, P. Horodecki, and J. Zakrzewski, At the limits of criticality-based quantum metrology: Apparent super-heisenberg scaling revisited, *Phys. Rev. X* **8**, 021022 (2018).
- [5] C. L. Degen, F. Reinhard, and P. Cappellaro, Quantum sensing, *Rev. Mod. Phys.* **89**, 035002 (2017).
- [6] J. F. Barry, J. M. Schloss, E. Bauch, M. J. Turner, C. A. Hart, L. M. Pham, and R. L. Walsworth, Sensitivity optimization for nv-diamond magnetometry, *Rev. Mod. Phys.* **92**, 015004 (2020).
- [7] E. J. Bergholtz, J. C. Budich, and F. K. Kunst, Exceptional topology of non-hermitian systems, *Rev. Mod. Phys.* **93**, 015005 (2021).
- [8] R. El-Ganainy, K. G. Makris, M. Khajavikhan, Z. H. Musslimani, S. Rotter, and D. N. Christodoulides, Non-hermitian physics and pt symmetry, *Nat. Phys.* **14**, 11 (2018).
- [9] X.-W. Luo, C. Zhang, and S. Du, Quantum squeezing and sensing with pseudo-anti-parity-time symmetry, *Phys. Rev. Lett.* **128**, 173602 (2022).
- [10] J. C. Budich and E. J. Bergholtz, Non-hermitian topological sensors, *Phys. Rev. Lett.* **125**, 180403 (2020).
- [11] F. Koch and J. C. Budich, Quantum non-hermitian topo-

- logical sensors, *Phys. Rev. Research* **4**, 013113 (2022).
- [12] A. McDonald and A. A. Clerk, Exponentially-enhanced quantum sensing with non-hermitian lattice dynamics, *Nat. Commun.* **11**, 1 (2020).
- [13] J. Wiersig, Enhancing the sensitivity of frequency and energy splitting detection by using exceptional points: Application to microcavity sensors for single-particle detection, *Phys. Rev. Lett.* **112**, 203901 (2014).
- [14] J. Wiersig, Sensors operating at exceptional points: General theory, *Phys. Rev. A* **93**, 033809 (2016).
- [15] J. Ren, H. Hodaiei, G. Harari, A. U. Hassan, W. Chow, M. Soltani, D. Christodoulides, and M. Khajavikhan, Ultrasensitive micro-scale parity-time-symmetric ring laser gyroscope, *Opt. Lett.* **42**, 1556 (2017).
- [16] S. Sunada, Large sagnac frequency splitting in a ring resonator operating at an exceptional point, *Phys. Rev. A* **96**, 033842 (2017).
- [17] Z.-P. Liu, J. Zhang, i. m. c. K. Özdemir, B. Peng, H. Jing, X.-Y. Lü, C.-W. Li, L. Yang, F. Nori, and Y.-x. Liu, Metrology with \mathcal{PT} -symmetric cavities: Enhanced sensitivity near the \mathcal{PT} -phase transition, *Phys. Rev. Lett.* **117**, 110802 (2016).
- [18] W. Chen, Ş. Kaya Özdemir, G. Zhao, J. Wiersig, and L. Yang, Exceptional points enhance sensing in an optical microcavity, *Nature (London)* **548**, 192 (2017).
- [19] H. Hodaiei, A. U. Hassan, S. Wittek, H. Garcia-Gracia, R. El-Ganainy, D. N. Christodoulides, and M. Khajavikhan, Enhanced sensitivity at higher-order exceptional points, *Nature (London)* **548**, 187 (2017).
- [20] Y.-H. Lai, Y.-K. Lu, M.-G. Suh, Z. Yuan, and K. Vahala, Observation of the exceptional-point-enhanced sagnac effect, *Nature (London)* **576**, 65 (2019).
- [21] R. Kononchuk, J. Cai, F. Ellis, R. Thevamaran, and T. Kottos, Exceptional-point-based accelerometers with enhanced signal-to-noise ratio, *Nature (London)* **607**, 697 (2022).
- [22] S. Yu, Y. Meng, J.-S. Tang, X.-Y. Xu, Y.-T. Wang, P. Yin, Z.-J. Ke, W. Liu, Z.-P. Li, Y.-Z. Yang, G. Chen, Y.-J. Han, C.-F. Li, and G.-C. Guo, Experimental investigation of quantum \mathcal{PT} -enhanced sensor, *Phys. Rev. Lett.* **125**, 240506 (2020).
- [23] H. Wang, Y.-H. Lai, Z. Yuan, M.-G. Suh, and K. Vahala, Petermann-factor sensitivity limit near an exceptional point in a brillouin ring laser gyroscope, *Nat. Commun.* **11**, 1 (2020).
- [24] W. Langbein, No exceptional precision of exceptional-point sensors, *Phys. Rev. A* **98**, 023805 (2018).
- [25] H.-K. Lau and A. A. Clerk, Fundamental limits and non-reciprocal approaches in non-hermitian quantum sensing, *Nat. Commun.* **9**, 1 (2018).
- [26] M. Zhang, W. Sweeney, C. W. Hsu, L. Yang, A. D. Stone, and L. Jiang, Quantum noise theory of exceptional point amplifying sensors, *Phys. Rev. Lett.* **123**, 180501 (2019).
- [27] C. Chen, L. Jin, and R.-B. Liu, Sensitivity of parameter estimation near the exceptional point of a non-hermitian system, *New J. Phys.* **21**, 083002 (2019).
- [28] R. Duggan, S. A. Mann, and A. Alu, Limitations of sensing at an exceptional point, *ACS Photonics* **9**, 1554 (2022).
- [29] J. Wiersig, Review of exceptional point-based sensors, *Photon. Res.* **8**, 1457 (2020).
- [30] W. M. Itano, J. C. Bergquist, J. J. Bollinger, J. M. Gilligan, D. J. Heinzen, F. L. Moore, M. G. Raizen, and D. J. Wineland, Quantum projection noise: Population fluctuations in two-level systems, *Phys. Rev. A* **47**, 3554 (1993).
- [31] H.-P. Breuer, F. Petruccione, *et al.*, *The theory of open quantum systems* (Oxford University Press on Demand, 2002).
- [32] B. Escher, R. de Matos Filho, and L. Davidovich, General framework for estimating the ultimate precision limit in noisy quantum-enhanced metrology, *Nat. Phys.* **7**, 406 (2011).
- [33] R. Demkowicz-Dobrzański, J. Kołodyński, and M. Guţă, The elusive heisenberg limit in quantum-enhanced metrology, *Nat. Commun.* **3**, 1 (2012).
- [34] A. W. Chin, S. F. Huelga, and M. B. Plenio, Quantum metrology in non-markovian environments, *Phys. Rev. Lett.* **109**, 233601 (2012).
- [35] S. Alipour, M. Mehboudi, and A. T. Rezakhani, Quantum metrology in open systems: Dissipative cramer-rao bound, *Phys. Rev. Lett.* **112**, 120405 (2014).
- [36] M. Beau and A. del Campo, Nonlinear quantum metrology of many-body open systems, *Phys. Rev. Lett.* **119**, 010403 (2017).
- [37] X.-M. Lu, X. Wang, and C. P. Sun, Quantum fisher information flow and non-markovian processes of open systems, *Phys. Rev. A* **82**, 042103 (2010).
- [38] S. L. Braunstein and C. M. Caves, Statistical distance and the geometry of quantum states, *Phys. Rev. Lett.* **72**, 3439 (1994).
- [39] C. W. Helstrom, Minimum mean-squared error of estimates in quantum statistics, *Phys. Lett. A* **25**, 101 (1967).
- [40] Equivalently, the quantum Fisher information can be calculated as $F_\lambda = -4 \frac{\partial^2 \mathcal{F}_Q(\rho_\lambda, \rho_{\lambda+\delta_\lambda})}{\partial \delta_\lambda^2} \Big|_{\delta_\lambda=0}$, where the quantum fidelity between quantum states is defined as $\mathcal{F}_Q(\rho_\lambda, \rho_{\lambda+\delta_\lambda}) \equiv \text{Tr} \left[\sqrt{\sqrt{\rho_{\lambda+\delta_\lambda}} \rho_\lambda \sqrt{\rho_{\lambda+\delta_\lambda}}} \right]$.
- [41] S. Pang and T. A. Brun, Quantum metrology for a general hamiltonian parameter, *Phys. Rev. A* **90**, 022117 (2014).
- [42] J. Liu, X.-X. Jing, and X. Wang, Quantum metrology with unitary parametrization processes, *Sci. Rep.* **5**, 1 (2015).
- [43] S. Boixo, S. T. Flammia, C. M. Caves, and J. Geremia, Generalized limits for single-parameter quantum estimation, *Phys. Rev. Lett.* **98**, 090401 (2007).
- [44] In particular, when H_1 is time-independent and the parameter is a multiplicative factor, i.e., $H_\lambda(t) = \lambda H_1 + H_0(t)$, we reproduce the result in Ref. [43] that, $F_\lambda \leq \|H_1\|^2 t^2$. Specifically, for the well-researched case where $H_1 = \sum_{j=1}^N h_j$, the operator seminorm $\|H_1\| = N(\varepsilon_M - \varepsilon_m)$, where ε_M (ε_m) represents the maximum (minimum) eigenvalue of the one-body Hamiltonian h_j . Consequently, the ultimate quantum Fisher information $\propto N^2$, corresponding to the well-known Heisenberg scaling. Additionally, if H_1 contains k -body interaction terms, the seminorm may scale as $\|H_1\| \propto N^k$, leading to super-Heisenberg scaling [4, 43].
- [45] L. J. Fiderer, J. M. E. Fraïsse, and D. Braun, Maximal quantum fisher information for mixed states, *Phys. Rev. Lett.* **123**, 250502 (2019).
- [46] S. Pang and A. N. Jordan, Optimal adaptive control for quantum metrology with time-dependent hamiltonians,

- Nat. Commun. **8**, 1 (2017).
- [47] In quantum sensing, the sensitivity is defined as the minimal detectable signal λ_{\min} that yields a unit signal-to-noise ratio for a unit integration time (sensing time) [5]. Therefore, the sensitivity λ_{\min} is bounded by same inequality in Eq. (2), with the substitution $t = 1$ s.
- [48] See Supplemental Material at [url] for the derivation of the bound of the change rate of the quantum Fisher information, the rate of dynamic quantum Fisher information for the pseudo-Hermitian quantum sensor and the derivation of the population fluctuation for the two-level system, which includes Ref. [30] and Ref. [43].
- [49] A non-Hermitian Hamiltonian H is said to be pseudo-Hermitian if $\eta H = H^\dagger \eta$, where η is a Hermitian invertible linear operator.
- [50] Y. Chu, Y. Liu, H. Liu, and J. Cai, Quantum sensing with a single-qubit pseudo-hermitian system, Phys. Rev. Lett. **124**, 020501 (2020).
- [51] U. Günther and B. F. Samsonov, Naimark-dilated \mathcal{PT} -symmetric brachistochrone, Phys. Rev. Lett. **101**, 230404 (2008).
- [52] K. Kawabata, Y. Ashida, and M. Ueda, Information retrieval and criticality in parity-time-symmetric systems, Phys. Rev. Lett. **119**, 190401 (2017).
- [53] Y. Wu, W. Liu, J. Geng, X. Song, X. Ye, C.-K. Duan, X. Rong, and J. Du, Observation of parity-time symmetry breaking in a single-spin system, Science **364**, 878 (2019).
- [54] L. Ding, K. Shi, Q. Zhang, D. Shen, X. Zhang, and W. Zhang, Experimental determination of \mathcal{PT} -symmetric exceptional points in a single trapped ion, Phys. Rev. Lett. **126**, 083604 (2021).
- [55] J. Li, A. K. Harter, J. Liu, L. de Melo, Y. N. Joglekar, and L. Luo, Observation of parity-time symmetry breaking transitions in a dissipative floquet system of ultracold atoms, Nat. Commun. **10**, 855 (2019).
- [56] In fact, in order to realize the EP enhanced sensing, we should first determine the location of EP. To be more specific, initially, only $\hat{H}_{\mathcal{PT}}$ is applied (by setting $\hat{H}_\delta(t) = 0$ temporarily) to determine the values of ω_{EP} and Γ_{EP} , which satisfy $P_J(T) - P_\Gamma(T) = 0$, since the sign of $P_J - P_\Gamma$ serves as an indicator of the phase boundary [54].
- [57] Due to the same $\sqrt{\nu}$ scaling in the sensitivity of the non-Hermitian sensor and its Hermitian counterpart, we plot these figures for $\nu = 1$.
- [58] The constant C_0 is given by $C_0 \equiv e^{2\Gamma T}$ because the evolution is governed by $H'_{\mathcal{PT}}$ in the experiment.
- [59] A. De Pasquale, D. Rossini, P. Facchi, and V. Giovannetti, Quantum parameter estimation affected by unitary disturbance, Phys. Rev. A **88**, 052117 (2013).
- [60] U. Mishra and A. Bayat, Driving enhanced quantum sensing in partially accessible many-body systems, Phys. Rev. Lett. **127**, 080504 (2021).
- [61] W. Ding, Y. Liu, Z. Zheng, and S. Chen, Dynamic quantum-enhanced sensing without entanglement in central spin systems, Phys. Rev. A **106**, 012604 (2022).
- [62] L. Xiao, K. Wang, X. Zhan, Z. Bian, K. Kawabata, M. Ueda, W. Yi, and P. Xue, Observation of critical phenomena in parity-time-symmetric quantum dynamics, Phys. Rev. Lett. **123**, 230401 (2019).
- [63] M. Naghiloo, M. Abbasi, Y. N. Joglekar, and K. Murch, Quantum state tomography across the exceptional point in a single dissipative qubit, Nature Physics **15**, 1232 (2019).
- [64] M. Huang, R.-K. Lee, L. Zhang, S.-M. Fei, and J. Wu, Simulating broken \mathcal{PT} -symmetric hamiltonian systems by weak measurement, Phys. Rev. Lett. **123**, 080404 (2019).

Supplementary Material: Fundamental Sensitivity Limits for non-Hermitian Quantum Sensors

Wenkui Ding,^{1,2} Xiaoguang Wang,^{2,*} and Shu Chen^{1,3,†}

¹Beijing National Laboratory for Condensed Matter Physics,

Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

²Department of Physics, Zhejiang Sci-Tech University, 310018 Zhejiang, China

³School of Physical Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China

(Dated: October 30, 2023)

THE BOUND FOR THE CHANGE RATE OF THE QUANTUM FISHER INFORMATION

Firstly, we consider the situation that the parameter to be estimated is a multiplicative factor in the Hamiltonian. The probe state is $|\Psi_0\rangle$ and the Hamiltonian to encode the parameter is $H_\lambda = \lambda H_1 + H_0$. After the time evolution, $|\Psi_f\rangle = e^{-iH_\lambda t}|\Psi_0\rangle$, the final state is dependent on the parameter λ . The quantum Fisher information for a pure quantum state is,

$$F_\lambda(t) = 4(\langle\Psi_f|\frac{\overleftarrow{\partial}}{\partial\lambda}\frac{\overrightarrow{\partial}}{\partial\lambda}|\Psi_f\rangle - |\langle\Psi_f|\frac{\overrightarrow{\partial}}{\partial\lambda}|\Psi_f\rangle|^2). \quad (1)$$

We can also calculate the quantum Fisher information in terms of the initial state as,

$$F_\lambda(t) = 4(\langle\Psi_0|h_\lambda^2|\Psi_0\rangle - |\langle\Psi_0|h_\lambda|\Psi_0\rangle|^2) \equiv 4 \text{Var}[h_\lambda]_{|\Psi_0\rangle}, \quad (2)$$

where we have defined the variance of operators and the transformed local generator is defined as,

$$h_\lambda \equiv ie^{iH_\lambda t}\frac{\partial}{\partial\lambda}e^{-iH_\lambda t} = iU_\lambda^\dagger\frac{\partial}{\partial\lambda}U_\lambda. \quad (3)$$

According to Ref. [1], we can prove that,

$$F_\lambda^{1/2} \leq t\|H_1\|, \quad (4)$$

where the seminorm $\|H_1\| \equiv u_{\max} - u_{\min}$, with u_{\max} and u_{\min} the largest and smallest eigenvalue of H_1 , respectively. Now, we want to prove that,

$$\left|\frac{\partial F_\lambda^{1/2}}{\partial t}\right| \leq \|H_1\|, \quad (5)$$

namely the change rate of the quantum Fisher information (or the magnitude of the quantum Fisher information flow) is bounded by the spectral width of the Hamiltonian that directly couples to the parameter. Calculating the derivative $\left|\frac{\partial F_\lambda^{1/2}}{\partial t}\right| = \frac{1}{2}F_\lambda^{-1/2}\left|\frac{\partial F_\lambda}{\partial t}\right|$, and using the fact that F_λ is always a positive quantity, equivalently, we are about to prove that,

$$\left|\frac{\partial F_\lambda}{\partial t}\right| \leq 2F_\lambda^{1/2}\|H_1\|. \quad (6)$$

Proof. we have the Schrödinger equation as,

$$i\frac{\partial U_\lambda}{\partial t} = H_\lambda U_\lambda, \quad (7)$$

namely, we have the relation that $\frac{\partial U_\lambda}{\partial t} = -iH_\lambda U_\lambda$ and $\frac{\partial U_\lambda^\dagger}{\partial t} = iU_\lambda^\dagger H_\lambda$. Then, according to the definition of the transformed local generator, we have,

$$\begin{aligned} \frac{\partial h_\lambda}{\partial t} &= i\frac{\partial U_\lambda^\dagger}{\partial t}\frac{\partial}{\partial\lambda}U_\lambda + iU_\lambda^\dagger\frac{\partial}{\partial\lambda}\frac{\partial U_\lambda}{\partial t} \\ &= i(iU_\lambda^\dagger H_\lambda)\frac{\partial}{\partial\lambda}U_\lambda + iU_\lambda^\dagger\frac{\partial}{\partial\lambda}(-iH_\lambda U_\lambda) \\ &= -U_\lambda^\dagger H_\lambda\frac{\partial U_\lambda}{\partial\lambda} + U_\lambda^\dagger\frac{\partial H_\lambda}{\partial\lambda}U_\lambda + U_\lambda^\dagger H_\lambda\frac{\partial U_\lambda}{\partial\lambda} \\ &= U_\lambda^\dagger\frac{\partial H_\lambda}{\partial\lambda}U_\lambda. \end{aligned} \quad (8)$$

Here, for $H_\lambda = \lambda H_1 + H_0$, we have $\frac{\partial h_\lambda}{\partial t} = U_\lambda^\dagger H_1 U_\lambda$.

According to the definition of the quantum Fisher information in the form of the variance of the transformed local generator, we obtain,

$$\begin{aligned} \frac{\partial F_\lambda}{\partial t} &= 4(\langle\Psi_0|\frac{\partial h_\lambda}{\partial t}h_\lambda + h_\lambda\frac{\partial h_\lambda}{\partial t}|\Psi_0\rangle - 2\langle\Psi_0|\frac{\partial h_\lambda}{\partial t}|\Psi_0\rangle\langle\Psi_0|h_\lambda|\Psi_0\rangle) \\ &\equiv 8 \text{Cov}\left[\frac{\partial h_\lambda}{\partial t}, h_\lambda\right]_{|\Psi_0\rangle}, \end{aligned} \quad (9)$$

where we have defined the covariance of operators as follows,

$$\text{Cov}[\hat{A}, \hat{B}]_{|\Psi\rangle} = \frac{1}{2}(\langle\Psi|\hat{A}\hat{B} + \hat{B}\hat{A}|\Psi\rangle - \langle\Psi|\hat{A}|\Psi\rangle\langle\Psi|\hat{B}|\Psi\rangle). \quad (10)$$

Using Cauchy-Schwarz inequality, we can prove the following relation (for compactness we neglect the subscript $|\Psi\rangle$ in the following discussions),

$$\left|\text{Cov}[\hat{A}, \hat{B}]\right| \leq \sqrt{\text{Var}(\hat{A})\text{Var}(\hat{B})}. \quad (11)$$

To prove this inequality, we define,

$$\hat{C} = \hat{A} - \frac{\text{Cov}[\hat{A}, \hat{B}]}{\text{Var}[\hat{B}]}\hat{B}. \quad (12)$$

Then, we have,

$$\begin{aligned}
\text{Var}[\hat{C}] &= \langle \Psi | \hat{A}^2 | \Psi \rangle + \langle \Psi | \hat{B}^2 | \Psi \rangle \left(\frac{\text{Cov}[\hat{A}, \hat{B}]}{\text{Var}[\hat{B}]} \right)^2 \\
&\quad - \langle \Psi | \hat{A}\hat{B} + \hat{B}\hat{A} | \Psi \rangle \frac{\text{Cov}[\hat{A}, \hat{B}]}{\text{Var}[\hat{B}]} - \langle \Psi | \hat{B} | \Psi \rangle^2 \left(\frac{\text{Cov}[\hat{A}, \hat{B}]}{\text{Var}[\hat{B}]} \right)^2 \\
&\quad - \langle \Psi | \hat{A} | \Psi \rangle^2 + 2\langle \Psi | \hat{A} | \Psi \rangle \langle \Psi | \hat{B} | \Psi \rangle \frac{\text{Cov}[\hat{A}, \hat{B}]}{\text{Var}[\hat{B}]} \\
&= \text{Var}[\hat{A}] + \text{Var}[\hat{B}] \left(\frac{\text{Cov}[\hat{A}, \hat{B}]}{\text{Var}[\hat{B}]} \right)^2 - 2\text{Cov}[\hat{A}, \hat{B}] \frac{\text{Cov}[\hat{A}, \hat{B}]}{\text{Var}[\hat{B}]} \\
&= \text{Var}[\hat{A}] - \frac{(\text{Cov}[\hat{A}, \hat{B}])^2}{\text{Var}[\hat{B}]} .
\end{aligned} \tag{13}$$

Since $\text{Var}[\hat{C}] \geq 0$, the inequality in Eq. 11 is proved. As a result, we have that,

$$\begin{aligned}
\left| \text{Cov} \left[\frac{\partial h_\lambda}{\partial t}, h_\lambda(t) \right] \Big|_{|\Psi_0\rangle} \right| &\leq \sqrt{\text{Var} \left[\frac{\partial h_\lambda}{\partial t} \Big|_{|\Psi_0\rangle} \text{Var} [h_\lambda(t)] \Big|_{|\Psi_0\rangle}} \\
&= \sqrt{\text{Var} [U_\lambda^\dagger H_1 U_\lambda] \Big|_{|\Psi_0\rangle} \frac{F_\lambda^{1/2}(t)}{2}} \\
&\leq \frac{\|H_1\|}{2} \frac{F_\lambda^{1/2}(t)}{2},
\end{aligned} \tag{14}$$

where we have used the relation that $F_\lambda(t) = 4 \text{Var} [h_\lambda(t)] \Big|_{|\Psi_0\rangle}$ and $\text{Var} [U_\lambda^\dagger H_1 U_\lambda] \leq \frac{\|U_\lambda^\dagger H_1 U_\lambda\|^2}{4} = \frac{\|H_1\|^2}{4}$, namely the variance of the Hermitian operator is bounded by its spectral width and the unitary evolution will not change the spectrum. Finally, we obtain,

$$\left| \frac{\partial F_\lambda}{\partial t} \right| \leq 2F_\lambda^{1/2} \|H_1\|. \tag{15}$$

The proof of the general case that the parameter is not a multiplicative factor as shown in the main text is very similar and we now briefly discuss the proof method. The Hamiltonian to encode the parameter now becomes $H_\lambda(t) = H_1(\lambda, t) + H_0(t)$ and the corresponding evolution operator becomes $U_\lambda = \mathcal{T} e^{-i \int_0^t H_\lambda(s) ds}$. The transformed local generator is defined as $h_\lambda = iU_\lambda^\dagger \frac{\partial}{\partial \lambda} U_\lambda$. We already have the bound of the quantum Fisher information as,

$$F_\lambda^{1/2}(t) \leq \int_0^t \left\| \frac{\partial H_\lambda(s)}{\partial \lambda} \right\| ds, \tag{16}$$

and we want to prove that,

$$\left| \frac{\partial F_\lambda^{1/2}(t)}{\partial t} \right| \leq \left\| \frac{\partial H_\lambda(t)}{\partial \lambda} \right\|. \tag{17}$$

To this end, we just need to follow the same procedure as above, except replacing H_1 by $\frac{\partial H_\lambda}{\partial \lambda}$.

RATE OF DYNAMIC QUANTUM FISHER INFORMATION FOR THE PSEUDO-HERMITIAN QUANTUM SENSOR

For the pseudo-Hermitian quantum sensor, the Hamiltonian to encode the parameter in the enlarged two-qubit system is as follows,

$$\hat{H}_{\text{tot}} = b\hat{I}^{(a)} \otimes \hat{\sigma}_x^{(s)} - c\hat{\sigma}_y^{(a)} \otimes \hat{\sigma}_y^{(s)} + \lambda\hat{I}^{(a)} \otimes \hat{\sigma}_x^{(s)}. \tag{18}$$

The probe state or the initial state is

$$|\Psi_0\rangle = \left(\sqrt{\frac{1+\varepsilon}{1+2\varepsilon}} |0\rangle_a + \sqrt{\frac{\varepsilon}{1+2\varepsilon}} |1\rangle_a \right) \otimes |0\rangle_s. \tag{19}$$

The dynamic quantum Fisher information can be calculated using

$$F_\lambda(t) = 4(\langle \Psi_0 | h_\lambda^2(t) | \Psi_0 \rangle - |\langle \Psi_0 | h_\lambda(t) | \Psi_0 \rangle|^2), \tag{20}$$

where the Hermitian operator

$$h_\lambda(t) \equiv iU_\lambda^\dagger(0 \rightarrow t) \frac{\partial}{\partial \lambda} U_\lambda(0 \rightarrow t) \tag{21}$$

is the transformed local generator. Since $[\hat{H}_{\text{tot}}, \hat{\sigma}_y^{(a)}] = 0$, we can represent the Hamiltonian in the two decoupled subspace as follows,

$$\hat{H}_{\text{tot}} = |\uparrow_y\rangle \langle \uparrow_y| \otimes \hat{H}_1 + |\downarrow_y\rangle \langle \downarrow_y| \otimes \hat{H}_2, \tag{22}$$

with,

$$\begin{aligned}
\hat{H}_1 &= (b + \lambda)\hat{\sigma}_x^{(s)} - c\hat{\sigma}_y^{(s)}, \\
\hat{H}_2 &= (b + \lambda)\hat{\sigma}_x^{(s)} + c\hat{\sigma}_y^{(s)}.
\end{aligned} \tag{23}$$

As a result, we can prove that,

$$\begin{aligned}
h_\lambda(t) &= i e^{i\hat{H}_{\text{tot}}t} \frac{\partial}{\partial \lambda} e^{-i\hat{H}_{\text{tot}}t} \\
&= |\uparrow_y\rangle \langle \uparrow_y| \otimes \hat{h}_1 + |\downarrow_y\rangle \langle \downarrow_y| \otimes \hat{h}_2,
\end{aligned} \tag{24}$$

where the transformed local generator in the respective subspace is as follows,

$$\begin{aligned}
\hat{h}_1 &\equiv i e^{i\hat{H}_1 t} \frac{\partial}{\partial \lambda} e^{-i\hat{H}_1 t} = h_x \hat{\sigma}_x^{(s)} + h_y \hat{\sigma}_y^{(s)} + h_z \hat{\sigma}_z^{(s)}, \\
\hat{h}_2 &\equiv i e^{i\hat{H}_2 t} \frac{\partial}{\partial \lambda} e^{-i\hat{H}_2 t} = h_x \hat{\sigma}_x^{(s)} - h_y \hat{\sigma}_y^{(s)} - h_z \hat{\sigma}_z^{(s)},
\end{aligned} \tag{25}$$

with the coefficients,

$$\begin{aligned}
h_x &= \frac{t(\lambda + b)^2}{\Omega^2} + \frac{c^2 \sin(2\Omega t)}{2\Omega^3}, \\
h_y &= \frac{c(\lambda + b)}{\Omega^2} \left[\frac{\sin(2\Omega t)}{2\Omega} - t \right], \\
h_z &= -\frac{c \sin^2(\Omega t)}{\Omega^2},
\end{aligned} \tag{26}$$

with $\Omega = \sqrt{(\lambda + b)^2 + c^2}$. Meanwhile, we can represent the initial state in the eigenbasis of $\hat{\sigma}_y^{(a)}$ as follows,

$$|\Psi_0\rangle = (\alpha |\uparrow_y\rangle + \beta |\downarrow_y\rangle) \otimes |0\rangle_s. \quad (27)$$

Then we have,

$$\begin{aligned} F_\lambda(t) &= 4 \left(\langle \Psi_0 | \hat{h}_\lambda^2(t) | \Psi_0 \rangle - |\langle \Psi_0 | h_\lambda(t) | \Psi_0 \rangle|^2 \right) \\ &= 4 \left(\alpha^2 {}_s\langle 0 | \hat{h}_1^2 | 0 \rangle_s + \beta^2 {}_s\langle 0 | \hat{h}_2^2 | 0 \rangle_s \right) \\ &\quad - 4 \left(\alpha^2 {}_s\langle 0 | \hat{h}_1 | 0 \rangle_s + \beta^2 {}_s\langle 0 | \hat{h}_2 | 0 \rangle_s \right)^2. \end{aligned} \quad (28)$$

Using the above equations, we have

$$\begin{aligned} {}_s\langle 0 | \hat{h}_1 | 0 \rangle_s &= h_z, \\ {}_s\langle 0 | \hat{h}_2 | 0 \rangle_s &= -h_z, \\ {}_s\langle 0 | \hat{h}_1^2 | 0 \rangle_s &= h_x^2 + h_y^2 + h_z^2, \\ {}_s\langle 0 | \hat{h}_2^2 | 0 \rangle_s &= h_x^2 + h_y^2 + h_z^2. \end{aligned} \quad (29)$$

Therefore, the dynamic quantum Fisher information is,

$$F_\lambda(t) = 4 [h_x^2 + h_y^2 + h_z^2 - (\alpha^2 - \beta^2)h_z^2]. \quad (30)$$

For the probe state $|\Psi_0\rangle$ in Eq. (19), we can check $\alpha^2 - \beta^2 = 0$, thus the dynamic quantum Fisher information becomes

$$F_\lambda(t) = \frac{4(b + \lambda)^2}{\Omega^2} t^2 + \frac{4c^2 \sin^2(\Omega t)}{\Omega^4}. \quad (31)$$

Finally, the change rate of the dynamic quantum Fisher information can be exactly calculated as follows,

$$\frac{\partial F_\lambda^{1/2}(t)}{\partial t} = 2 \frac{\cos^2 \theta + \sin^2 \theta \frac{\sin(2\Omega t)}{2\Omega t}}{\sqrt{\cos^2 \theta + \sin^2 \theta \frac{\sin^2(\Omega t)}{(\Omega t)^2}}}, \quad (32)$$

where we have defined $(b + \lambda)/\Omega = \cos \theta$ and $c/\Omega = \sin \theta$.

From Eq. (32), it follows that $-2 \leq \frac{\partial F_\lambda^{1/2}(t)}{\partial t} \leq 2$, which verifies our theory in the main text.

THE POPULATION FLUCTUATION FOR THE TWO-LEVEL SYSTEM

We now briefly discuss the quantum projection noise in the population measurement of the two-level system [2].

The quantum state that needs to be measured is,

$$|\psi\rangle = \alpha |\uparrow\rangle + \beta |\downarrow\rangle. \quad (33)$$

For this state, we have the population (probability) in the $|\uparrow\rangle$ state after the projection measurement as, $p_\uparrow = |\alpha|^2$. After a repetition of N times, the number of measurement outcomes to be the $|\uparrow\rangle$ state is N_\uparrow , then we can estimate the probability $p_\uparrow \approx p'_\uparrow = N_\uparrow/N$. Since N is finite, there are uncertainties in the value of N_\uparrow and equivalently there are uncertainties in the determination of p'_\uparrow . Since the repeated projection process follows the binomial distribution, the variance of the times that projected to the up state is $\text{Var}(N_\uparrow) = N p_\uparrow (1 - p_\uparrow)$. Correspondingly, the variance of the estimated population is,

$$\text{Var}(p'_\uparrow) = \text{Var}\left(\frac{N_\uparrow}{N}\right) = \frac{p_\uparrow(1 - p_\uparrow)}{N}. \quad (34)$$

The discussion above can be extended to the projective measurement of multi-level quantum systems. As an example, for a general two-qubit quantum state,

$$|\psi\rangle = \alpha_1 |00\rangle + \alpha_2 |01\rangle + \alpha_3 |10\rangle + \alpha_4 |11\rangle, \quad (35)$$

the distribution of the times N_1 , corresponding to the projected outcome in $|00\rangle$, for N repeated measurements follows the multinomial distribution. Thus, the variance of N_1 is,

$$\text{Var}(N_1) = N p_1 (1 - p_1), \quad (36)$$

where $p_1 = |\alpha_1|^2$. Accordingly, the uncertainty of the estimated population is,

$$\text{Var}(p'_1) = \text{Var}\left(\frac{N_1}{N}\right) = \frac{p_1(1 - p_1)}{N}. \quad (37)$$

* xgwang@zstu.edu.cn

† schen@iphy.ac.cn

- [1] S. Boixo, S. T. Flammia, C. M. Caves, and J. Geremia, Generalized limits for single-parameter quantum estimation, *Phys. Rev. Lett.* **98**, 090401 (2007).
- [2] W. M. Itano, J. C. Bergquist, J. J. Bollinger, J. M. Gilligan, D. J. Heinzen, F. L. Moore, M. G. Raizen, and D. J. Wineland, Quantum projection noise: Population fluctuations in two-level systems, *Phys. Rev. A* **47**, 3554 (1993).