arXiv:2304.10667v2 [cond-mat.quant-gas] 15 Jan 2024

Observation of self-oscillating supersonic flow across an acoustic horizon in two dimensions

Hikaru Tamura,¹ Sergei Khlebnikov,^{1,2} Cheng-An Chen,^{1,*} and Chen-Lung Hung^{1,2,†}

¹Department of Physics and Astronomy, Purdue University, West Lafayette, IN 47907, USA

²Purdue Quantum Science and Engineering Institute, Purdue University, West Lafayette, IN 47907, USA

(Dated: January 17, 2024)

Understanding the dynamics and stability of transonic flows in quantum fluids, especially for those beyond one spatial dimension, is an outstanding challenge, with applications ranging from nonlinear optics and condensed matter to analogue gravity. One intriguing possibility is that a system with a spatially bounded supersonic flow may evolve into a self-oscillating state that periodically emits solitons, in a process originating from the well-known Landau instability. Here, we report observation of self-oscillating supersonic flows in a two-dimensional atomic superfluid. By imposing a local particle sink with strong loss, we induce a convergent radial flow forming an acoustic analogue of a black-hole horizon and an inner horizon around the sink. The observed superflow appears to be modulated by quasi-periodic bursts of superluminal signals. We measure their frequencies and find agreement with numerical simulations of soliton oscillation frequencies within the black-hole horizon. The presented experiment demonstrates a new method for creating supersonic flows in atomic superfluids, which may find applications in quantum simulations of curved spacetime, supersonic turbulence, and self-oscillating dynamics in dissipative many-body systems.

According to Landau's criterion of superfluidity [1], a superfluid flowing past an obstacle becomes unstable with respect to production of excitations when the velocity exceeds a certain limit. For phonon excitations in a weakly interacting Bose-Einstein condensate (BEC), the critical velocity coincides with the speed of sound. In experiments, much lower critical velocities are often observed, which have been attributed to production of low energy vortex excitations [2–5]. For a one-dimensional (1D) superflow, on the other hand, the critical velocity [6] has been found to depend on the obstacle height [7, 8], and it has been suggested that the Landau criterion is violated when the local flow velocity exceeds the local sound speed [8, 9].

More generally, even without an obstacle, one expects that Landau instability plays a role as long as the translational symmetry is broken. An intriguing, yet unexplored example is a convergent two-dimensional (2D) radial flow, where an increasing flow rate at a small enough radius r grows larger than the local sound speed, and the flow could become unstable. In many related settings in 1D, Landau instability manifests itself through self-periodic emission of solitons [6–8, 10–12]. Here, we explore for the first time the stability of a 2D radial flow and report observation of self-periodic oscillations.

Intriguingly, it is precisely the transonic flows in a quantum fluid that has been theorized [13, 14] and broadly pursued (see [15–20] for examples) as a simulator of an elusive phenomenon—Hawking radiation from a black hole [21], which results from quantum fluctuations near the event horizon. An acoustic black-hole (white-hole) horizon marks the transition of a subsonic flow to (from) a supersonic region that low-frequency sound waves cannot escape (re-enter). A bounded supersonic flow, like those in a penetrable barrier or in a convergent 2D flow, is enclosed by a pair of black-hole and white-hole (inner) horizons, reminiscent of two horizons of a charged black hole in Einstein gravity. In the presence of superluminal (faster than sound) short-wave excitations, a pair of acoustic horizons can act like mirrors that form a laser cavity, further amplifying the out-going Hawking radiation via stimulated emission [22]. More generally, Hawking radiation can also be stimulated from an initial seed produced by other mechanisms [19]. Recent discussions of this effect in 1D include [9, 23–25].

In contrast to Hawking radiation, soliton and wave emissions due to Landau instability do not require initial seeds. Testing instability of supersonic flow within two horizons [9, 11, 12, 26] has so far remained an open experimental question. For instance, a recent experiment by the Technion group [15] has generated acoustic horizons by sweeping a potential step along an elongated condensate. This so-called waterfall method has led to a successful observation of Hawking radiation of phonons [15, 27] across a horizon that co-moves with the step potential. Phonons emitted following formation of an inner horizon, however, have been attributed not to the spontaneous Hawking process but to an amplification of the Cherenkov radiation emitted by a moving obstacle [28–30].

Here, we address the role of Landau instability in a 2D radial flow free from a moving obstacle. We create a particle sink at the center of an otherwise homogeneous atomic superfluid trapped in an optical box. The sink induces fast atom number loss and results in a large inward radial flow, forming an acoustic black-hole horizon and an inner horizon around the sink.

We control particle loss rate in the sink through three-body recombination [31], a dissipative process during which three atoms collide to form one bound molecule and one energetic atom, both with kinetic energy large enough to escape a shallow optical trap. Three-body recombination loss scales cubically with atomic density as $\dot{n} = -L_3 n^3$, where $L_3 \approx$ $4.3 \times 10^{-2} \ \mu m^4/s$ is the loss coefficient in our 2D geometry [32] and *n* the 2D density. In our ultracold cesium atomic samples, two-body loss is fully suppressed. We use all conservative potentials in contrast to a related proposal [33] that utilizes localized one-body loss to generate supersonic flows.

^{*} Current address: Atom Computing, Boulder, CO 80301, USA

[†] clhung@purdue.edu



FIG. 1. Supersonic flow induced by a particle sink. a, Schematics of a Gaussian attractive potential (depth V_0) addressing a 2D atomic superfluid trapped in a circular box. The high density region with a large three-body recombination loss rate serves as a particle sink, inducing strong radial flow (velocity v < 0). Dotted circles illustrate an acoustic black-hole horizon and an inner horizon. **b**, Velocity v(black) and local sound speed $c = \hbar \sqrt{ng}/m$ (red) versus radial position r for the stationary solution of the GPE in an effectively infinite system at a critical depth $V_0 = V_{\rm cr}$. Shaded region marks the supersonic flow |v(r)| > c(r). For a finite system, time-dependent simulations suggest that ramping on a sink potential beyond the critical depth triggers soliton emission. c, Density profiles at 0, 4, and 8 ms (dark to light gray curves) right after supersonic flow forms, obtained from a numerical solution of the time-dependent 2D GPE using a slow ramp of V_0 ($\Delta t = 60$ ms as illustrated in e inset). d, Sound (red) and flow velocity (gray) profiles corresponding to those in c. e-f. Full numerical time evolution of n(r) and $\Delta(r) = c(r) + v(r)$. showing initial soliton emission near the critical depth (marked by dashed lines), radial oscillations of solitons, and multiplication of soliton number following each oscillation cycle. This dynamics results in a self-oscillating supersonic flow.

As illustrated in Figs. 1a and 2, a 2D superfluid is initially trapped inside a circular box of potential height $\approx k_{\rm B} \times 60$ nK, with a uniform density $n_0 \approx 14 \,\mu \text{m}^{-2}$ and a chemical potential $\mu_0 = \hbar^2 n_0 g/m \approx k_{\rm B} \times 21$ nK [32]. Here, $g \approx 0.42$ is the interaction parameter, \hbar the reduced Planck constant, *m* the atomic mass, and k_B the Boltzmann constant. We introduce the sink by ramping on a Gaussian addressing potential of $1/e^2$ radius $r_{\rm s} \approx 6.5 \ \mu {\rm m}$ and depth $V_0 \approx k_{\rm B} \times 200 \ {\rm nK}$ at the box center. The attractive potential gives rise to a much higher peak density > $90\mu m^{-2}$ in the sink, leading to more than 250-fold increase in the local three-body loss rate and an estimated total loss rate of $\Gamma = \int_{s} |\dot{n}| d^2 r \gtrsim 6.5 \times 10^5 \text{s}^{-1}$ in the sink region. Assuming fluid continuity at $r > r_s$, one can estimate the radial velocity as $v(r) \approx -\Gamma/[2\pi rn(r)] \lesssim -1$ mm/s, indicating that v can become supersonic outside the sink.

We first perform theoretical analyses on the stability of this dissipation-induced flow. We model the process through a classical 2D Gross-Pitaevskii equation (GPE) with an additional term accounting for the three-body loss [32]. Assuming rotational symmetry, we have found stationary solutions by allowing inflow of atoms at the boundary. We expect such solutions to be close to quasi-stationary states in a large sample without an inflow. Specifically, for V_0 below a critical value $V_{\rm cr} (\approx k_{\rm B} \times 88 \text{ nK} \text{ for the chosen parameter values})$, we find a ground state solution and a transition ('droplet') state that, similarly to the saddle-point solution [34, 35] of the Ginzburg-Landau theory, can be interpreted as the fluctuation mediating a phase slip. At $V_0 = V_{cr}$, the solutions merge and disappear through a saddle-node bifurcation, in parallel to results obtained for conservative flows over obstacles in 1D [7]. A critical solution for experimentally relevant parameter values is shown in Fig. 1b. Notice that the ground state develops a small supersonic region when approaching the critical point. We have observed such a correlation also for other parameter values. We therefore interpret disappearance of the static solutions at the critical point as a Landau-type instability.

Analogies between our results and those of Refs. [7, 34, 35] suggest that ramping the potential past the critical value will induce a self-oscillation process, analogous to the soliton train in a conservative 1D flow [7] or a phase-slip center [36] in a superconducting wire. This is supported by full numerical integration of a time-dependent 2D GPE [32]. An example is shown in radial plots in Figs. 1c-f. After V_0 passes through a critical point, supersonic flow forms; see (d, f) for regions with $\Delta = c + v < 0$, where v (c) is the local flow velocity (sound speed). Coincidentally, a train of ring-shaped dark solitons [37] (3 clearly visible in this example) are emitted toward the sink center, in time separation ≤ 3 ms. They appear as left-moving dark dips in the radial plots (c, e), forcing oscillations in the supersonic flow. We find that this process is insensitive to the potential ramp speed and that soliton emission always accompanies formation of supersonic flow near the critical point.

Once initiated, a ring soliton's radial motion becomes part of a multiplication process, a remarkable effect absent in 1D black-hole lasers. We point out that a shrinking ring dark soliton cannot stop at the 'singularity' at r = 0 [37, 38]. A soliton first transmits through the inner horizon, reaches an inner turning point at $r \ge 0$, and then expands radially back towards the supersonic region [right-moving dips in the density in Fig. 1e]. Upon reaching the inner horizon, the soliton is reflected towards r = 0, leaving behind a wake of outgoing radiation. During this process, new solitons are continually emitted near the inner horizon and also become trapped at smaller radii. The system thus behaves like an amplifying, self-periodic 'soliton laser' mediating oscillating supersonic



FIG. 2. Realization of radial supersonic flow in a 2D superfluid. a, Single-shot in-situ density images measured before and at the indicated time t after the sink is fully ramped on at t = 5 ms. Dotted (dashed) circles mark $r = 26 \ \mu m$ (5 μm) radius. b, Radial density profiles n(r) measured at t = 7-63 ms (dark to light gray circles) with a time interval of 8 ms. Initial density profile (blue circles) is plotted for comparison. c, Local sound speed c(r) evaluated using profiles in b. d, Evolution of integrated atom number N(t) with (filled circles) and without (open circles) the sink, agreeing with a model assuming three-body recombination loss (red curves). Blue dashed curve is a simple exponential fit, giving the total atom decay rate γ . e, Radial flow velocity evaluated using Eq. (1).

flows while emitting sound waves out of the black-hole horizon.

In the actual experiment, we have adopted a faster ramp speed ($\Delta t = 5$ ms) to be able to observe the instability before losing many atoms. As shown in the in-situ images in Fig. 2 **a** and averaged radial density plots in **b**, shortly after the addressing potential is ramped on, atomic density slightly depletes at $r \leq 15 \,\mu$ m outside the sink, showing a strong tendency for the superfluid to flow inwards. As time increases, the density continues to decrease, indicating a continuous flow into the sink region even after the peak density has saturated. Using the radial density profiles, we evaluate the local sound speed c(r), which is nearly uniform and gradually decreases with time to < 1 mm/s as shown in **c** except within the sink where the density is high.

In Fig. 2d, we plot the total atom number N(t), excluding the central region $r \le r_s$. Decay of N(t) is consistent with atoms flowing into the sink to compensate for the loss of atoms due to three-body recombination, as described by a simple theory curve (red dashed line) in Fig. 2c [32]. The overall decay rate $\gamma \approx 27 \text{ s}^{-1}$ is determined by a fit.

Due to finite resolution of our imaging system (~ 1 μ m), we cannot clearly identify ring dark solitons in situ, as the characteristic width of their density dip is $\xi \approx 1/\sqrt{ng} = 0.2 - 0.4 \,\mu$ m. We also note that, in a superfluid with preexisting density noise [38] or imperfect rotational symmetry [39], a ring soliton suffers strong snaking instability [40] and can quickly decay into vortices [41] (also ~ ξ wide) that are challenging to measure in situ. This decay has been observed in GPE simulations as well. Generation and decay of ring dark solitons have been reported in our system but with a different experimental setting [38].

We can nevertheless identify key signatures of Landau instability and self-oscillation in the measured flow. To extract this information, we compute the local radial flow velocity using the rate of change of total atom number in an annular region bounded by (r, r_{∞}) via the expression

$$v_{\exp}(r,t) = \frac{1}{2\pi r n(r,t)} \frac{dN(r,t)}{dt},$$
(1)

where $N(r,t) = \int_{r}^{r_{\infty}} n(r',t) d^{2}r'$ and $r_{\infty} \approx 40 \ \mu \text{m}$ extends well beyond the edge of the box trap. Figure 2e plots the radial flow velocity evaluated at various times, showing mostly inward flow $v_{\exp}(r) < 0$ everywhere for $r \leq 26 \ \mu \text{m}$. The amplitude $|v_{\exp}(r)|$ increases with decreasing radial position, reaching maximum at around $r \approx 8 \ \mu \text{m}$. It then greatly decreases when approaching the sink region where density becomes high, in qualitative agreement with the flow analyses. Time-dependence of flow, on the other hand, shows an intriguing oscillatory behavior that we now discuss.

To clearly see the evolution of the superflow, we plot the full spatial-temporal dependence of $\Delta_{\exp}(r, t) = c(r, t) + v_{\exp}(r, t)$ as shown in Fig. 3a. Supersonic flow initially appears within a radial interval $(r_{in}, r_{out}) \approx (5, 10) \mu m$, enlarging to $\approx (5, 15) \mu m$ at later times. This can be viewed as a supersonic flow cavity bounded by a black-hole horizon at $r = r_{out}$ and an inner white-hole horizon at $r = r_{in}$. For comparison, we also evaluate the flow velocity v_{GPE} obtained in a GPE calculation with the same ramp speed ($\Delta t = 5 \text{ ms}$) as in the experiment and using the same Eq. (1). The result is shown in Fig. 3b-c. The local flow velocity and self-oscillation signatures are nearly the same as those found in v(r, t), the radial velocity computed directly from the GPE wavefunction.

There is however a striking difference between the experiment and the simulation results. In the experiment, at around $t \approx 10$ ms, shortly after the inward flow becomes supersonic, a sudden change to an apparent outward flow is observed $(v_{exp} > 0)$; see also Fig. 3c. At larger times, Δ_{exp} appears to display quasi-periodic short pulses with a primary time period of $t_p \approx 4$ ms. This pulsation behavior is from time-dependence of the flow velocity as the sound speed is monotonically decreasing in time. The pulse period is also longer than that of possible collective modes in the sink region, if any is excited. Most surprisingly, these pulses appear to propagate over the entire sample within a small time ≤ 2 ms, which is much shorter than the time period $\gtrsim 20$ ms required for sound waves to traverse the sample.



FIG. 3. **Observation of self-oscillating supersonic flow. a**, Measured time evolution (in 1 ms steps) of $\Delta_{exp}(r)$, showing supersonic flow ($\Delta_{exp} < 0$) at $r > r_{in} \approx 5 \ \mu m$ and $r < r_{out}$, where $r_{out} \gtrsim 10 \ \mu m$ grows slowly with time. Quasi-periodic pulses of $\Delta_{exp} > 0$ are clearly visible. **b**, Calculated $\Delta_{GPE}(r, t)$ based on flow velocity v_{GPE} evaluated using Eq. (1). **c**, Measured (top panel) and calculated (gray curve, bottom panel) flow velocities at positions as shown in the dashed lines in **a** and **b**, respectively. A 1 ms running-average (black curve) is plotted for comparing with experiment. Error bars represent the standard error of the mean. **d**, Normalized Fourier spectra of v_{exp} (black circles), v (dashed curve), and v_{GPE} (dotted curve), respectively. Gray, orange, and blue shaded bands indicate the corresponding statistical uncertainties.

The apparent short pulses could be due to outbursts of atoms from the sink region, traveling at superluminal speeds greater than 1 cm/s (kinetic energy > $k_{\rm B} \times 1 \ \mu {\rm K}$). These energetic atoms may come from three-body recombination [31, 42], with each atom carrying away 2/3 of the binding energy of the bound molecular state. The closest to the continuum, 6s state of Cs₂ [43], has a binding energy $E_{\rm b} \approx k_{\rm B} \times 20 \,\mu {\rm K}$, thus giving an estimated out-going atom velocity of ≈ 4 cm/s presumably along random directions in 3D. Some of these atoms will be imaged in our apparatus. As the recombination loss occurs primarily in the sink region, we expect it to be modulated by self-oscillations. These effects are not captured in our classical GPE calculations. Another source of energetic atoms, although likely much less prominent, may be the dynamical Casimir effect [44-47], wherein the motion of solitonic defects results in rapid density perturbations in the sink region, possibly capable of exciting short-scale fluctuations ($\leq \xi$) with superluminal speeds comparable to $2\pi\hbar/m\xi \lesssim 2$ cm/s.

To see if these fast pulses are synchronized to possible soliton motion within the horizon and are not from other systematic effects, we calculate the Fourier spectrum of $v_{exp}(t)$ and compare it with the Fourier spectra of the GPE results aver-



FIG. 4. Self-oscillation frequencies. **a**, Main peak frequencies f_p in Fourier spectra of v_{exp} from samples with two different depths $V_0 \approx k_B \times 200$ nK (filled symbols) and 125 nK (open symbols), shown in **b** and Fig. 3**d**, respectively, versus measured atom number decay rate γ . Main peak frequencies from the spectra of GPE results for v (black crosses) and v_{GPE} (gray crosses) are plotted for comparison. Error bars and shaded band reflect the uncertainty in peak detection for experiments and GPE calculations, respectively. **b**, Comparison between the experiment (symbols) and GPE spectra of v (dashed curves) and v_{GPE} (dotted curves). Gray, orange, and blue shaded bands indicate the statistical uncertainties. For **a-b**, relevant experimental parameters are $(g, n_0, r_s) \approx (0.42, 14, 6.5)$ (circles), (0.48, 13, 6.5) (squares), (0.42, 25, 5) (triangles), and (0.45, 20, 5) (diamonds); the units of length are in μ m.

aged over the sink region [32]. As shown in Fig. 3d, the most prominent frequency peak observed in the experiment is at $f_p \approx 225$ Hz ~ t_p^{-1} . This is in good agreement with the main frequency peak in the GPE result. We have verified that f_p indeed corresponds to the radial oscillation frequency of tightly trapped solitons, which are reflected upon re-entering the supersonic region.

Self-oscillations in this 2D geometry appear to be robust. Experimentally, we have observed the peak frequency to shift slightly to $f_{\rm p} \approx 250$ Hz upon a moderate increase of the atomic interaction (top panel of Fig. 4b). This is consistent with the slight variation of soliton oscillation frequency seen in the GPE simulations (Fig. SM1 and SM2). We have also adopted a shallower depth $V_0 \approx k_{\rm B} \times 125$ nK and a slightly narrower sink (bottom panel). The peak frequency is reduced to $f_{\rm p} \approx 150$ Hz, which is consistent with simulation results (Fig. SM1 and SM2) and is attributed to lower density and sound speed within the sink. In general, the oscillation frequencies should depend on the radius of the horizon, the supersonic flow speed, and the local sound speed. We summarize these results using a single parameter, the total atom number decay rate γ , which characterizes the overall dissipation rate in the sink. Figure 4 plots the measurement results, where we find the spectra to show reasonable agreement with soliton oscillation frequencies in GPE calculations. For frequency dependence on various parameters in GPE calculations, see Supplemental Figures.

In summary, by introducing a stationary, conservative local addressing potential, we observe self-induced supersonic flow initiated solely by fast local three-body recombination. Through observing periodic emission of superluminal signals, which we attribute to strong nonlinear effects in the sink, we discover that the system enters a self-oscillating state. By measuring the frequency spectrum of the oscillations, we obtain evidence that they are synchronized with the periodic soliton motion observed in our GPE simulations. Our setup may be considered as a classical analogue of a black-hole laser, with the supersonic region and the 'singularity' at the sink center acting as highly reflective cavity mirrors, a pattern distinct from the more familiar 1D black-hole lasers. More generally, our work shows that projecting arbitrary sink potentials to initiate localized dissipation processes may be a valuable way to generate complex flow patterns in atomic superfluids. Our experiments and simulations both show sustained periodic modulation signal over time, implying that energy released by three-body recombination continues to be converted to excitations and suggesting that additional instabilities may develop, such as those leading to a turbulent cascade [48-52]. We ex-

- L. Landau, Theory of the superfluidity of helium ii, Physical Review 60, 356 (1941).
- [2] R. P. Feynman, Chapter ii application of quantum mechanics to liquid helium, in *Progress in low temperature physics*, Vol. 1 (Elsevier, 1955) pp. 17–53.
- [3] T. W. Neely, E. C. Samson, A. S. Bradley, M. J. Davis, and B. P. Anderson, Observation of vortex dipoles in an oblate Bose-Einstein condensate, Physical Review Letters **104**, 160401 (2010).
- [4] A. Ramanathan, K. Wright, S. R. Muniz, M. Zelan, W. Hill III, C. Lobb, K. Helmerson, W. Phillips, and G. Campbell, Superflow in a toroidal Bose-Einstein condensate: an atom circuit with a tunable weak link, Physical Review Letters **106**, 130401 (2011).
- [5] R. Desbuquois, L. Chomaz, T. Yefsah, J. Léonard, J. Beugnon, C. Weitenberg, and J. Dalibard, Superfluid behaviour of a twodimensional Bose gas, Nature Physics 8, 645 (2012).
- [6] P. Engels and C. Atherton, Stationary and nonstationary fluid flow of a Bose-Einstein condensate through a penetrable barrier, Physical Review Letters 99, 160405 (2007).
- [7] V. Hakim, Nonlinear Schrödinger flow past an obstacle in one dimension, Physical Review E 55, 2835 (1997).
- [8] N. Pavloff, Breakdown of superfluidity of an atom laser past an obstacle, Physical Review A 66, 013610 (2002).
- [9] S. Finazzi, F. Piazza, M. Abad, A. Smerzi, and A. Recati, Instability of the superfluid flow as black-hole lasing effect, Physical Review Letters 114, 245301 (2015).
- [10] P. Leboeuf and N. Pavloff, Bose-Einstein beams: Coherent propagation through a guide, Physical Review A 64, 033602 (2001).
- [11] F. Michel and R. Parentani, Nonlinear effects in time-dependent transonic flows: An analysis of analog black hole stability, Physical Review A 91, 053603 (2015).
- [12] J. De Nova, S. Finazzi, and I. Carusotto, Time-dependent study of a black-hole laser in a flowing atomic condensate, Physical Review A 94, 043616 (2016).
- [13] W. G. Unruh, Experimental black-hole evaporation?, Physical Review Letters 46, 1351 (1981).

pect that the momentum distribution and the kinetic energy spectrum of such a cascade [48, 49] can be measured using time-of-flight [52] and in situ density noise measurements following creation of the supersonic flow. In addition, unambiguous imaging and counting of solitons and their decay into vortices can be realized in a 2D time-of-flight with increased healing length [38]. Our work calls for future studies on self-oscillations in dissipative quantum fluids and can potentially find new applications in quantum simulations of curved space-time [14, 53].

ACKNOWLEDGMENTS

We thank Chris Greene, Martin Kruczenski, and Qi Zhou for discussions. This work is supported by the W. M. Keck Foundation, the NSF (Grant # PHY-1848316), and the DOE QuantISED program through the Fermilab Quantum Consortium.

- [14] C. Barceló, S. Liberati, and M. Visser, Analogue gravity, Living reviews in relativity 14, 1 (2011).
- [15] J. R. Muñoz de Nova, K. Golubkov, V. I. Kolobov, and J. Steinhauer, Observation of thermal Hawking radiation and its temperature in an analogue black hole, Nature 569, 688 (2019).
- [16] J. Drori, Y. Rosenberg, D. Bermudez, Y. Silberberg, and U. Leonhardt, Observation of stimulated Hawking radiation in an optical analogue, Physical Review Letters **122**, 010404 (2019).
- [17] L.-P. Euvé, F. Michel, R. Parentani, T. G. Philbin, and G. Rousseaux, Observation of noise correlated by the Hawking effect in a water tank, Physical Review Letters 117, 121301 (2016).
- [18] H. S. Nguyen, D. Gerace, I. Carusotto, D. Sanvitto, E. Galopin, A. Lemaître, I. Sagnes, J. Bloch, and A. Amo, Acoustic black hole in a stationary hydrodynamic flow of microcavity polaritons, Physical Review Letters **114**, 036402 (2015).
- [19] S. Weinfurtner, E. W. Tedford, M. C. Penrice, W. G. Unruh, and G. A. Lawrence, Measurement of stimulated Hawking emission in an analogue system, Physical Review Letters **106**, 021302 (2011).
- [20] T. G. Philbin, C. Kuklewicz, S. Robertson, S. Hill, F. Konig, and U. Leonhardt, Fiber-optical analog of the event horizon, Science 319, 1367 (2008).
- [21] S. W. Hawking, Particle creation by black holes, Communications in mathematical physics 43, 199 (1975).
- [22] S. Corley and T. Jacobson, Black hole lasers, Physical Review D 59, 124011 (1999).
- [23] U. Leonhardt and T. G. Philbin, Black hole lasers revisited, Quantum analogues: from phase transitions to black holes and cosmology, 229 (2007).
- [24] S. Finazzi and R. Parentani, Black hole lasers in Bose–Einstein condensates, New Journal of Physics 12, 095015 (2010).
- [25] A. Coutant and R. Parentani, Black hole lasers, a mode analysis, Physical Review D 81, 084042 (2010).
- [26] J. De Nova, P. Palacios, I. Carusotto, and F. Sols, Long time universality of black-hole lasers, New Journal of Physics 23, 023040 (2021).

- [27] J. Steinhauer, Observation of quantum Hawking radiation and its entanglement in an analogue black hole, Nature Physics 12, 959 (2016).
- [28] M. Tettamanti, S. Cacciatori, A. Parola, and I. Carusotto, Numerical study of a recent black-hole lasing experiment, Europhysics Letters 114, 60011 (2016).
- [29] Y.-H. Wang, T. Jacobson, M. Edwards, and C. W. Clark, Mechanism of stimulated Hawking radiation in a laboratory Bose-Einstein condensate, Physical Review A 96, 023616 (2017).
- [30] V. I. Kolobov, K. Golubkov, J. R. Muñoz de Nova, and J. Steinhauer, Observation of stationary spontaneous Hawking radiation and the time evolution of an analogue black hole, Nature Physics 17, 362 (2021).
- [31] T. Weber, J. Herbig, M. Mark, H.-C. Nägerl, and R. Grimm, Three-body recombination at large scattering lengths in an ultracold atomic gas, Physical Review Letters 91, 123201 (2003).
- [32] See Supplemental Materials.
- [33] D. Sels and E. Demler, Thermal radiation and dissipative phase transition in a BEC with local loss, Annals of Physics 412, 168021 (2020).
- [34] W. A. Little, Decay of persistent currents in small superconductors, Physical Review 156, 396 (1967).
- [35] J. S. Langer and V. Ambegaokar, Intrinsic resistive transition in narrow superconducting channels, Physical Review 164, 498 (1967).
- [36] W. Skocpol, M. Beasley, and M. Tinkham, Phase-slip centers and nonequilibrium processes in superconducting tin microbridges, Journal of Low Temperature Physics 16, 145 (1974).
- [37] Y. S. Kivshar and X. Yang, Ring dark solitons, Physical Review E **50**, R40 (1994).
- [38] H. Tamura, C.-A. Chen, and C.-L. Hung, Observation of selfpatterned defect formation in atomic superfluids–from ring dark solitons to vortex dipole necklaces, Physical Review X 13, 031029 (2023).
- [39] L. Toikka and K.-A. Suominen, Snake instability of ring dark solitons in toroidally trapped Bose-Einstein condensates, Physical Review A 87, 043601 (2013).
- [40] Y. S. Kivshar and D. E. Pelinovsky, Self-focusing and transverse instabilities of solitary waves, Physics Reports 331, 117 (2000).
- [41] G. Theocharis, D. Frantzeskakis, P. Kevrekidis, B. A. Malomed, and Y. S. Kivshar, Ring dark solitons and vortex necklaces in Bose-Einstein condensates, Physical Review Letters 90, 120403 (2003).
- [42] J. Wolf, M. Deiß, A. Krükow, E. Tiemann, B. P. Ruzic, Y. Wang, J. P. D'Incao, P. S. Julienne, and J. H. Denschlag, State-to-state chemistry for three-body recombination in an ultracold rubidium gas, Science 358, 921 (2017).
- [43] M. Mark, F. Ferlaino, S. Knoop, J. G. Danzl, T. Kraemer, C. Chin, H.-C. Nägerl, and R. Grimm, Spectroscopy of ultracold trapped cesium Feshbach molecules, Physical Review A 76, 042514 (2007).
- [44] J.-C. Jaskula, G. B. Partridge, M. Bonneau, R. Lopes, J. Ruaudel, D. Boiron, and C. I. Westbrook, Acoustic analog to the dynamical Casimir effect in a Bose-Einstein condensate, Physical Review Letters 109, 220401 (2012).
- [45] C.-L. Hung, V. Gurarie, and C. Chin, From cosmology to cold atoms: observation of Sakharov oscillations in a quenched atomic superfluid, Science 341, 1213 (2013).
- [46] S. Eckel, A. Kumar, T. Jacobson, I. B. Spielman, and G. K. Campbell, A rapidly expanding Bose-Einstein condensate: an expanding universe in the lab, Physical Review X 8, 021021 (2018).
- [47] P. O. Fedichev and U. R. Fischer, "cosmological" quasiparticle production in harmonically trapped superfluid gases, Physical

Review A 69, 033602 (2004).

- [48] S. Khlebnikov, Strong acoustic turbulence and the speed of bose-einstein condensation, Physical Review A 66, 063606 (2002).
- [49] M. C. Tsatsos, P. E. Tavares, A. Cidrim, A. R. Fritsch, M. A. Caracanhas, F. E. A. dos Santos, C. F. Barenghi, and V. S. Bagnato, Quantum turbulence in trapped atomic Bose-Einstein condensates, Physics Reports 622, 1 (2016).
- [50] E. A. L. Henn, J. A. Seman, G. Roati, K. M. F. Magalhaes, and V. S. Bagnato, Emergence of turbulence in an oscillating Bose-Einstein condensate, Physical Review Letters 103, 045301 (2009).
- [51] W. J. Kwon, G. Moon, J.-y. Choi, S. W. Seo, and Y.-i. Shin, Relaxation of superfluid turbulence in highly oblate Bose-Einstein condensates, Physical Review A 90, 063627 (2014).
- [52] N. Navon, A. L. Gaunt, R. P. Smith, and Z. Hadzibabic, Emergence of a turbulent cascade in a quantum gas, Nature 539, 72 (2016).
- [53] C. Viermann, M. Sparn, N. Liebster, M. Hans, E. Kath, A. Parra-López, M. Tolosa-Simeón, N. Sánchez-Kuntz, T. Haas, H. Strobel, *et al.*, Quantum field simulator for dynamics in curved spacetime, Nature **611**, 260 (2022).
- [54] C.-A. Chen, S. Khlebnikov, and C.-L. Hung, Observation of quasiparticle pair production and quantum entanglement in atomic quantum gases quenched to an attractive interaction, Physical Review Letters **127**, 060404 (2021).
- [55] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, Feshbach resonances in ultracold gases, Reviews of Modern Physics 82, 1225 (2010).
- [56] C.-L. Hung, X. Zhang, L.-C. Ha, S.-K. Tung, N. Gemelke, and C. Chin, Extracting density–density correlations from in situ images of atomic quantum gases, New Journal of Physics 13, 075019 (2011).
- [57] X. Antoine and R. Duboscq, Gpelab, a matlab toolbox to solve gross–pitaevskii equations i: Computation of stationary solutions, Computer Physics Communications 185, 2969 (2014).
- [58] X. Antoine and R. Duboscq, Gpelab, a matlab toolbox to solve gross–pitaevskii equations ii: Dynamics and stochastic simulations, Computer Physics Communications 193, 95 (2015).

SUPPLEMENTAL MATERIALS

S1. Preparation and detection of a two-dimensional (2D) superfluid

Our 2D superfluid samples are formed by $N \approx 3 \times 10^4$ Bose-condensed cesium atoms loaded into an all repulsive optical box potential with an initial temperature T < 10 nK. Detailed descriptions on the condensate formation and trap loading procedures can be found in Refs. [38, 54]. The vertical confinement of the optical box offers a trap frequency of $\omega_z \approx 2\pi \times 1.8$ kHz. The atoms occupy the ground state with an harmonic oscillator length of $l_z = \sqrt{\hbar/m\omega_z} \approx 207$ nm, where \hbar is the reduced Planck constant and *m* is the cesium atomic mass. The horizontal box confinement is provided by a ring-shaped repulsive wall potential with an adjustable radius patterned by a digital mirror device and projected through a high numerical aperture (NA ≈ 0.6) objective lens. Crosssection of the wall has an approximate Gaussian form and the width is 5 μ m. The potential height is $\approx k_{\rm B} \times 60 n$ K, where $k_{\rm B}$ is the Boltzmann constant. In our quasi-2D geometry, the interaction parameter is $g = \sqrt{8\pi a_s}/l_z$, where a_s is the threedimensional (3D) s-wave scattering length, tunable via a Feshbach resonance [55].

After box loading, we slowly ramp the scattering length to $a_s \approx 328 \sim 368 a_0$, where a_0 is the Bohr radius, to prepare for the experiments shown in Figs. 2-4 at $g \approx 0.42 \sim 0.48$. We measure the in-situ density profiles by saturated absorption imaging through the high-NA objective. Image resolution ($\leq 1 \mu$ m) is calibrated using in-situ atomic density noise [56].

To evaluate averaged density and flow velocity profiles, we repeat each experiment condition and perform measurements ~ 50 times to record in situ density images. To minimize error due to shot-to-shot atom number fluctuations, at each hold time we select images with total atom number typically within $\pm 10\%$ deviation from a mean number for analyses. Care has been taken to evaluate the mean without being biased by large fluctuations. We estimate the error of the mean is $\lesssim 1.2\%$. Typically, around ~ 30 images are selected to evaluate the average density and flow velocity profiles.

S2. Calibration of the addressing potential

The addressing (sink) potential is formed using a far-off resonant optical beam at 876 nm wavelength, with an adjustable beam size and projected through the high-NA objective lens. We calibrate the potential depth V_0 by addressing a 2D gas at a much lower optical power to avoid fast three-body recombination loss. The depth is equal to the local chemical potential difference $\Delta \mu = \hbar^2 g \Delta n/m$ between the addressed region and the surrounding superfluid region, where Δn is the density difference. We calibrate the width r_s of the addressing beam by measuring the size of the weakly-addressed density peak. For strong addressing, we compare time-dependent GPE calculations with flow measurement results, through adjusting the width and depth of the addressing potential by 10% ~ 20% around the calibrated value, and find good agreement. In the main text, we report the adjusted values of the sink potential.

S3. Three-body recombination loss rate

We describe three-body recombination loss by the rate equation $\dot{n} = -L_3 n^3$, with a cubic dependence on the 2D density *n*. In our quasi-2D geometry, the vertical oscillator length $l_z \gg a_s$ is much bigger than the 3D scattering length and the three-body recombination process remains 3D in nature. We relate the rate coefficient L_3 in quasi-2D with the 3D loss coefficient l_3 by integrating out the ground state density distribution along the z-axis. We have $L_3 = l_3/(\pi l_z^2 \sqrt{3})$, where $l_3 = C \frac{\hbar}{m} a_s^4 \approx 0.356 \times \frac{\hbar}{m} (l_z g)^4$. $C \approx 225$ has been measured in Ref. [31] using cesium thermal gas. In the experiment, we have tuned to a larger scattering length to increase the threebody loss coefficient to $L_3 \approx 0.042 \sim 0.071 \ \mu m^4/s$. We note that our measured particle loss rate from the sink is consistent with a loss coefficient of thermal gas without the $\frac{1}{3!}$ suppression factor for identical bosons.

In Fig. 2d, we model the time dependence of the total particle number N outside the sink region $(N = \int_{r>r_0} nd^2r)$ by

$$\frac{dN}{dt} = -L_3 \int n^3 d^2 r \,, \tag{S1}$$

where $n = n_{\rm s} + n_{\rm bg}$, $n_{\rm s}(r) = n_{\rm p}e^{-2r^2/\sigma^2}$ is a saturated Gaussian density distribution at the sink, σ the radius, $n_{\rm p}$ the peak density, $n_{\rm bg} = N/(\pi R^2)$ the background uniform density, and R the sample radius. We determine $(n_p, \sigma) \approx (80 \ \mu m^{-2}, 4.8 \ \mu m)$ by fitting the sink density distribution. Time dependence of N(t) can be completely determined by Eq. (S1). In Fig. 2d, we plot the dashed curve from $t_i = 14$ ms with $n_{\rm bg}(t_i) = 12.5 \ \mu m^{-2}$ and $R = 27 \ \mu m$, and plot the dotted curve without the sink by setting $n_{\rm p} = 0$ and $t_i = 0$.

S4. Stationary solution for the supersonic flow induced by a particle sink

We look for stationary solutions to the classical 2D Gross-Pitaevskii equation (GPE) with an additional term accounting for three-body recombination:

$$i\hbar\partial_t\psi = \frac{\hbar^2}{2m}(-\nabla^2 + 2gn - i\gamma_3 n^2)\psi + [V(r) - \mu]\psi. \quad (S2)$$

Here ψ is the complex amplitude, $n = \psi^{\dagger} \psi$ is the atom density, $\gamma_3 = mL_3/\hbar$, and V(r) is the addressing potential, which we take in the Gaussian form

$$V(r) = -V_0 \exp(-2r^2/r_s^2)$$
.

At the classical level, it is consistent to assume perfect rotational symmetry, so $\psi \equiv \psi(r, t)$, and ∇^2 is the radial part of the 2D Laplacian. The radial coordinate *r* takes values in the range $0 \le r \le R$. At r = 0, we impose the standard regularity condition $\partial_r \psi = 0$. Because the system is dissipative (with a 3-body loss coefficient γ_3), a stationary solution can only exist if we allow inflow of atoms at the boundary. To achieve that, we impose different boundary conditions for the real and imaginary parts of ψ at r = R—Neumann for the real part and Dirichlet for the imaginary: $\partial_r \operatorname{Re}[\psi] = 0$, $\operatorname{Im}[\psi] = 0$. These permit a nonzero inflow velocity at r = R, which however is not specified a priori but will be found as a part of the solution. The idea is that, if *R* is sufficiently large, such a stationary solution (if any is found) will be close enough to a *quasi*-stationary state that would be obtained in a large sample without any inflow at the boundary.

To look for stationary solutions, we set the left-hand side of Eq. (S2) to zero, discretize the right-hand side on a uniform grid of N points (typically N = 500) and solve the resulting system of equations numerically by the multidimensional Newton-Raphson (NR) method. We define a real $2N \times 2N$ Jacobian matrix with elements $J_{ik} = \partial F_i / \partial u_k$, where $\{F_i\}$ is the set of the real and imaginary parts of the right-hand side of Eq. (S2) at the grid points, and $\{u_k\}$ is the set of those of ψ . It is convenient to vary the potential depth V_0 while keeping all the other parameters fixed. For V_0 below a certain critical value $V_{\rm cr}$, we find two solutions. For either, eigenvalues of the Jacobian are all real and, for one of the solutions, are in fact all positive. The other solution has one negative mode. We refer to the first solution as the ground state, and to the second as the droplet, as discussed in the main text. At $V_0 = V_{cr}$ the solutions merge and disappear through a saddle-node bifurcation. A critical solution is shown in Fig. 1b.

S5. Time-dependent GPE simulation

We perform time-dependent simulations of Eq. (S2) using code described in [57, 58] for matching experimental conditions. The initial ground state wave function is obtained by first setting $\gamma_3 = 0$ and $V_0 = 0$, while adding a confining potential of the form

$$U(r) = \frac{U_0}{2} \left[1 + \operatorname{erf}\left(\frac{r-R}{w}\right) \right] \,. \tag{S3}$$

Here $U_0 = k_B \times 60$ nK is the trap strength, $w = 2.5 \ \mu m$, and erf(r) is the error function that sets $U(r \leq R) = 0$ and $U(r \geq R) = U_0$ for us to obtain a localized initial stationary solution. Physical simulation box size is $70 \times 70 \ \mu m^2$ with typically 257×257 or 513×513 grid points.

To simulate the dynamics, we set γ_3 in Eq. (S2) to the experimental value and evolve the time-dependent GPE with a linear ramp of addressing potential to a final depth of $V_0 = k_{\rm B} \times 200$ nK (or $V_0 = k_{\rm B} \times 125$ nK). The simulation time step is 1 μ s and the time duration of ramp Δt is indicated in the main text.

We observe trains of ring dark solitons (RDSs) emitted in cases of fast ($\Delta t = 5$ ms) and slower ramps ($\Delta t > 10$ ms). Persistent self-oscillations of solitons are observed once the addressing potential depth passes through a critical value V_{cr}

close to the prediction by the stationary state solutions. More simulation results can be found in Figs. SM1 and SM2.

We note that RDSs suffer from snaking instability even in GPE calculations in the absence of preexisting perturbations. The instability can be triggered by numerical variations (deterministic error) across the simulation grids that break rotational symmetry [39, 41]. This is particularly severe when an RDS shrinks to a small radius comparable to the grid size. Eventually, an RDS will decay into a ring of vortex dipoles. While this is observed in our 2D GPE simulations, the impact of this instability on soliton self-oscillating dynamics requires further investigations.

S6. Evaluating the Fourier spectra

To test if the pulses in measured supersonic flows are synchronized with soliton oscillations, we compare the Fourier spectra of $v_{exp}(t)$ obtained in experiment and $v_{GPE}(t)$, evaluated using Eq. (1), as well as v(t), obtained using the probability current density, in GPE simulations. Due to finite experiment time and resolution (total time $t_{\text{max}} = 55 \sim 65 \text{ ms}$ with 1 ms time step), our measured Fourier spectrum has a limited frequency resolution ~ $t_{\rm max}^{-1} \approx 20$ Hz and sensitivity up to f = 500 Hz. We apply a boxcar time-window between $t = 15 \sim 65$ ms with a ± 5 ms variable length to evaluate an averaged Fourier spectrum as shown in Fig. 3(d) and Fig. 4(b). To evaluate the Fourier spectra based on GPE simulations, we evaluate a 1 ms running average of $v_{GPE}(t)$ and v(t), respectively, as shown in Fig. 3(c), and then calculate the Fourier spectra in the same window-averaging procedure as described above. Since we are testing possible synchronization with soliton motion, we average the GPE spectra over a range of radial positions within the sink. In Fig. 3(d) and the top panel of Fig. 4(b) $(V_0/k_B = 200 \text{ nK} \text{ and } r_s = 6.5 \ \mu\text{m})$, we show the spectra averaged over $r \leq 5 \ \mu m$ as these spectra clearly exhibit prominent frequency peaks and agree well with our experiment result. For the bottom panel of Fig. 4(b) $(V_0/k_{\rm B} =$ 125 nK and $r_s = 5 \ \mu m$), we average over $r \le 3.5 \ \mu m$.





Fig. SM1. Dependence of the Fourier spectra of GPE simulations on various parameters. **a-b**, Fourier spectra of calculated flow v versus addressing depth V_0 for fixed interaction parameter g = 0.42 and addressing beam radius $r_s = 7\mu$ m (**a**), and 5μ m (**b**), respectively. **c**, Fourier spectra of calculated flow v versus interaction parameter g for fixed $V_0/k_B = 200$ nK and $r_s = 6.5 \mu$ m. **d**, Fourier spectra of calculated flow v versus interaction parameter of calculated flow v versus r_s for fixed $V_0/k_B = 200$ nK and g = 0.42. In all the numerical simulations, the addressing potential is ramped on in 5 ms, as in the experiment, and the initial total atom number is set to $N = 3 \times 10^4$.

Fig. SM2. Dependence of the Fourier spectra of the flow velocity v_{GPE} , evaluated using Eq. (1) in the main text, on the parameters for the same GPE simulations as in Fig. SM1. Compared to the spectra of v in the Fig. SM1, the main oscillation peaks of v_{GPE} appear weaker due to contamination by a low frequency background signal likely caused by slow dynamics in the density profile. In Figs. SM1 and SM2, weak multi-tones other than the main peak in the Fourier spectra are attributed to anharmonicity of the oscillation signal.