Gravitational-Wave Phasing of Quasi-Circular Compact Binary Systems to the Fourth-and-a-Half post-Newtonian Order

Luc BLANCHET,^{1,*} Guillaume FAYE,^{1,2,†} Quentin HENRY,^{3,‡} François LARROUTUROU,^{4,§} and David TRESTINI^{5,1,¶}

 ${}^{1}\mathcal{GR}\varepsilon\mathbb{CO}$, Institut d'Astrophysique de Paris,

UMR 7095, CNRS, Sorbonne Université, 98^{bis} boulevard Arago, 75014 Paris, France

²Centre for Strings, Gravitation and Cosmology, Department of Physics,

Indian Institute of Technology Madras, Chennai 600036, India

³Max Planck Institute for Gravitational Physics

(Albert Einstein Institute), D-14476 Potsdam, Germany

⁴Deutsches Elektronen-Synchrotron DESY,

Notkestr. 85, 22607 Hamburg, Germany

⁵Laboratoire Univers et Théories, Observatoire de Paris,

Université PSL, Université Paris Cité, CNRS, F-92190 Meudon, France

(Dated: October 3, 2023)

The inspiral phase of gravitational waves emitted by spinless compact binary systems is derived through the fourth-and-a-half post-Newtonian (4.5PN) order beyond quadrupole radiation, and the leading amplitude mode $(\ell, m) = (2, 2)$ is obtained at 4PN order. We also provide the radiated flux, as well as the phase in the stationary phase approximation. Rough numerical estimates for the contribution of each PN order are provided for typical systems observed by current and future gravitational wave detectors.

At the time when the LIGO and Virgo gravitational-wave detectors were approved, there was no theoretical prediction available for gravitational waves (GWs) generated by compact binary systems, apart from that of the famous Einstein quadrupole formula [1–3]. However, it was soon realized that, given the frequency band and the expected sensitivity of these ground-based detectors, the waveform modeling was to be drastically improved in order to extract all the potential information from the signal, at least in the case of the inspiral of two neutron stars [4, 5]. The breakthrough came with the merging of the post-Newtonian (PN) and the multipolar post-Minkowskian (MPM) expansions into a single formalism [6–10], that was applied with success to derive step by step the waveform of compact binary systems up to 3.5PN order [11–20] (the results are also known at 4PN order for the spin-orbit coupling and the spin-spin coupling, see *e.g.* [21, 22]). Since then, many works (outlined below) have aimed at extending the precision of this result to the next level, namely 4PN or even 4.5PN order beyond the Einstein quadrupole formula.

This Letter provides the final results of these efforts, *i.e.*, the GW phasing of non-spinning compact binary systems on quasi-circular orbits up to 4.5PN order, as well as the dominant GW mode, given by $(\ell, m) = (2, 2)$, at 4PN order. Ready to be used for building accurate PN template banks for the detection and analysis of the inspiral phase of compact binaries, they should be important for third generation ground-based detectors (Einstein Telescope and Cosmic Explorer), future space-borne detectors (LISA and TianQin) and of course the current second-generation detectors (LIGO, Virgo and KAGRA). All results presented in this Letter are to be found in the ancillary file [23] associated with the companion paper [24].

Besides improving the detectors' data analysis, the motivation for computing high PN orders is also to perform high-accuracy tests of general relativity (GR), since the PN coefficients directly probe the non-linear structure of the theory. By confronting results from the PN expansion against data, one can put constraints on potential deviations from GR [25, 26]. This has already allowed for the confirmation of the signature of GW tails [27, 28], and is promising for tests with future multi-band detections between LISA and ground-based detectors [29].

We denote by f(t) the frequency of the dominant (2, 2) mode of the GW as measured by an observer in the asymptotically flat region far from the source (recall that this is twice the orbital frequency), and by $\psi(t) = \pi \int dt f(t)$ the corresponding half-phase. As usual, we define the directly-measurable PN parameter $x = \mathcal{O}(c^{-2})$ by

$$x \equiv \left(\frac{\pi Gmf}{c^3}\right)^{2/3},\tag{1}$$

where $m = m_1 + m_2$ is the binary's total mass, m_1 and m_2 being the constant masses of the progenitors. For circular orbits, x may be defined invariantly from the Killing vector of the helical symmetry in the asymptotically flat spacetime. Since compact binaries tend to have circularized by the time they enter the detector's frequency band [30], we only consider the case of quasi-circular orbits, for which the time-evolution of the frequency and phase (or "chirp") is entirely driven by the energy flux-balance equation,

$$\frac{\mathrm{d}E}{\mathrm{d}t} = -\mathcal{F}\,,\tag{2}$$

where E denotes the invariant energy of the compact binary and \mathcal{F} the total energy flux (or GW luminosity). Both Eand \mathcal{F} in the balance equation are unique functions of the PN parameter x and the two masses. They have to be evaluated with the same relative PN precision, in the present case 4.5PN ($\sim x^{9/2}$). From Eq. (2), we derive a simple ordinary differential equation for the frequency as a function of time, and, once it is solved, a further integration yields the phase as a function of frequency.

The invariant energy E follows from the conservative dynamics of the compact binary at 4PN order, which have been obtained by various groups using different methods: (i) the Arnowitt-Deser-Misner (ADM) Hamiltonian formalism [31– 34] yielded the first derivation of the 4PN energy, although with an ambiguity parameter obtained by matching the near-zone computation to results imported from gravitational self-force (GSF) [35]; (ii) the Fokker Lagrangian formalism in harmonic coordinates [36–39] derived the complete result, without ambiguity and without resorting to GSF, by using a specific regularization procedure which was proven to be equivalent to dimensional regularization; (iii) the effective field theory (EFT) approach [40–47] rederived the 4PN energy by using dimensional regularization. From this series of works, the binary's invariant energy was obtained as the Noetherian quantity associated with temporal translation, and reads at 4PN order (see *e.g.* [48, 49] for partial results up to 6PN order):

$$E = -\frac{m\nu c^2 x}{2} \left\{ 1 + \left(-\frac{3}{4} - \frac{\nu}{12} \right) x + \left(-\frac{27}{8} + \frac{19}{8}\nu - \frac{\nu^2}{24} \right) x^2 + \left[-\frac{675}{64} + \left(\frac{34445}{576} - \frac{205}{96}\pi^2 \right) \nu - \frac{155}{96}\nu^2 - \frac{35}{5184}\nu^3 \right] x^3 + \left[-\frac{3969}{128} + \left(-\frac{123671}{5760} + \frac{9037}{1536}\pi^2 + \frac{896}{15}\gamma_{\rm E} + \frac{448}{15}\ln(16x) \right) \nu + \left(-\frac{498449}{3456} + \frac{3157}{576}\pi^2 \right) \nu^2 + \frac{301}{1728}\nu^3 + \frac{77}{31104}\nu^4 \right] x^4 + \mathcal{O}(x^5) \right\}.$$
(3)

We denote by $\nu \equiv m_1 m_2/m^2$ the symmetric mass ratio (γ_E is the Euler constant). Since there are no terms of half-integer PN order for circular orbits, this expression is actually valid up to 4.5PN order (as indicated by the final error term).

The second input is the energy flux, which we have computed using the PN-MPM formalism applied to compact binaries at 4.5PN beyond the leading quadrupole formula. Crucial to this computation was the recently-completed source mass quadrupole moment at 4PN order [50–53], the source current quadrupole moment at 3PN order [54] and the non-linear tail-of-memory effect [55, 56]. We provide the technical details of the derivation in the companion paper [24], and report here only the final result:

$$\begin{split} \mathcal{F} &= \frac{32c^5}{5G}\nu^2 x^5 \Biggl\{ 1 + \left(-\frac{1247}{336} - \frac{35}{12}\nu \right) x + 4\pi x^{3/2} + \left(-\frac{44711}{9072} + \frac{9271}{504}\nu + \frac{65}{18}\nu^2 \right) x^2 + \left(-\frac{8191}{672} - \frac{583}{24}\nu \right) \pi x^{5/2} \\ &+ \left[\frac{6643739519}{69854400} + \frac{16}{3}\pi^2 - \frac{1712}{105}\gamma_{\rm E} - \frac{856}{105}\ln(16\,x) + \left(-\frac{134543}{7776} + \frac{41}{48}\pi^2 \right)\nu - \frac{94403}{3024}\nu^2 - \frac{775}{324}\nu^3 \right] x^3 \\ &+ \left(-\frac{16285}{504} + \frac{214745}{1728}\nu + \frac{193385}{3024}\nu^2 \right) \pi x^{7/2} \\ &+ \left[-\frac{323105549467}{1378375200} + \frac{232597}{4410}\gamma_{\rm E} - \frac{1369}{126}\pi^2 + \frac{39931}{294}\ln 2 - \frac{47385}{1568}\ln 3 + \frac{232597}{820}\ln x \right] \\ &+ \left(-\frac{1452202403629}{1466942400} + \frac{41478}{245}\gamma_{\rm E} - \frac{267127}{4608}\pi^2 + \frac{479062}{2205}\ln 2 + \frac{47385}{392}\ln 3 + \frac{20739}{245}\ln x \right) \nu \\ &+ \left(\frac{1607125}{6804} - \frac{3157}{384}\pi^2 \right) \nu^2 + \frac{6875}{504}\nu^3 + \frac{5}{6}\nu^4 \right] x^4 \\ &+ \left[\frac{265978667519}{745113600} - \frac{6848}{105}\gamma_{\rm E} - \frac{3424}{105}\ln(16\,x) + \left(\frac{2062241}{22176} + \frac{41}{12}\pi^2 \right) \nu \right] \end{split}$$

$$\frac{133112905}{290304}\nu^2 - \frac{3719141}{38016}\nu^3 \left[\pi x^{9/2} + \mathcal{O}(x^5)\right].$$
(4)

In the test-mass limit $\nu \to 0$, we exactly retrieve the result of linear black-hole perturbation theory [57–61]. Since BH perturbations have recently been extended numerically to second order in the mass ratio ν [62–64], it would be interesting to verify the consistency of this numerical result with the PN prediction (4). Note also that in the case of black holes, the contributions due to the absorption by the BH horizons are not included in the PN calculation, and should be added separately. The BH absorption is a 4PN effect for Schwarzschild black holes [65], and a 2.5PN effect for spinning ones [66–70].

With both (3) and (4) in hand, we apply the flux-balance equation (2) and readily obtain the time evolution of the GW frequency. Sophisticated techniques exist to increase the precision of the PN results and the overlap with numerical relativity [71–74], but we do not discuss them here and simply present the results in the form of a fully expanded Taylor PN series. We employ the time variable

$$\tau \equiv \frac{\nu c^3}{5Gm} (t_0 - t) , \qquad (5)$$

where t is the coordinate time in the asymptotic radiative coordinate system, and t_0 an integration constant. We have the freedom to redefine it as $t_0 \rightarrow t_0 + \alpha \frac{Gm}{c^3}$ where α is any constant, which amounts to the replacement $\tau \rightarrow \tau [1 + \alpha \nu / (5\tau)]$. Although t_0 is not uniquely defined, it might be formally interpreted as the instant of coalescence, when $x \rightarrow +\infty$, and then it satisfies $t_0 - t = \mathcal{O}(c^5)$ in the PN regime. At the 4PN order, using the fact that $\tau^{-1} = \mathcal{O}(c^{-8})$ is a small 4PN quantity, we conveniently adjust α so as to simplify as much as possible the result:

$$\begin{aligned} x &= \frac{\tau^{-1/4}}{4} \left\{ 1 + \left(\frac{743}{4032} + \frac{11}{48}\nu\right)\tau^{-1/4} - \frac{1}{5}\pi\,\tau^{-3/8} \\ &+ \left(\frac{19583}{254016} + \frac{24401}{193536}\nu + \frac{31}{288}\nu^2\right)\tau^{-1/2} + \left(-\frac{11891}{53760} + \frac{109}{1920}\nu\right)\pi\,\tau^{-5/8} \\ &+ \left[-\frac{10052469856691}{6008596070400} + \frac{1}{6}\pi^2 + \frac{107}{420}\gamma_{\rm E} - \frac{107}{360}\ln\left(\frac{\tau}{256}\right) \\ &+ \left(\frac{3147553127}{780337152} - \frac{451}{3072}\pi^2\right)\nu - \frac{15211}{442368}\nu^2 + \frac{25565}{331776}\nu^3\right]\tau^{-3/4} \\ &+ \left(-\frac{113868647}{433520640} - \frac{31821}{143360}\nu + \frac{294941}{3870720}\nu^2\right)\pi\tau^{-7/8} \\ &+ \left[-\frac{2518977598355703073}{3779358859513036800} + \frac{9203}{215040}\gamma_{\rm E} + \frac{9049}{258048}\pi^2 + \frac{14873}{1128960}\ln 2 + \frac{47385}{1605632}\ln 3 - \frac{9203}{3440640}\ln\tau \\ &+ \left(\frac{718143266031997}{576825222758400} + \frac{244493}{1128960}\gamma_{\rm E} - \frac{65577}{1835008}\pi^2 + \frac{15761}{47040}\ln 2 - \frac{47385}{401408}\ln 3 - \frac{244493}{18063360}\ln\tau\right)\nu \\ &+ \left(-\frac{1502014727}{5323596288} + \frac{2255}{393216}\pi^2\right)\nu^2 - \frac{258479}{33030144}\nu^3 + \frac{1195}{262144}\nu^4\right]\tau^{-1}\ln\tau \\ &+ \left[-\frac{9965202491753717}{5768252227584000} + \frac{107}{600}\gamma_{\rm E} + \frac{23}{600}\pi^2 - \frac{107}{4800}\ln\left(\frac{\tau}{256}\right)\right] \\ &+ \left(\frac{8248609881163}{2746786775040} - \frac{3157}{30720}\pi^2\right)\nu - \frac{3590973803}{20808990720}\nu^2 - \frac{520159}{1634992128}\nu^3\right]\pi\,\tau^{-9/8} + \mathcal{O}(\tau^{-5/4})\right\}. \tag{6}$$

Then, the GW half-phase ψ of the dominant harmonics is related to the binary's orbital phase ϕ by

$$\psi = \phi - \frac{2\pi G M f}{c^3} \ln\left(\frac{f}{f_0}\right),\tag{7}$$

where M is the ADM mass of the binary, and f_0 is an arbitrary unphysical scale, reflecting the different origins of time between the local coordinates covering the source and the radiative coordinates. The logarithmic phase modulation was determined in [75, 76] and is physically due to the scattering of GWs on the Schwarzschild background associated with M (*i.e.* GW tails). While the GW half-phase ψ and the corresponding GW frequency $f = \dot{\psi}/\pi$ are directly measurable, the orbital phase ϕ can only be inferred via the theoretical prediction (7). When expressing the results in terms of the GW observables ψ and f, the arbitrary scale f_0 is canceled out (see Sec. VI B in the detailed paper [24]). The explicit expression of the time-domain GW half-phase $\psi(t)$ in terms of x(t) [given by (6)] reads

$$\begin{split} \psi &= \psi_0 - \frac{x^{-5/2}}{32\nu} \Biggl\{ 1 + \left(\frac{3715}{1008} + \frac{55}{12} \nu \right) x - 10\pi x^{3/2} \\ &+ \left(\frac{15293365}{1016064} + \frac{27145}{1008} \nu + \frac{3085}{144} \nu^2 \right) x^2 + \left(\frac{38645}{1344} - \frac{65}{16} \nu \right) \pi x^{5/2} \ln x \\ &+ \left[\frac{12348611926451}{18776862720} - \frac{160}{3} \pi^2 - \frac{1712}{21} \gamma_{\rm E} - \frac{856}{21} \ln(16 x) \right. \\ &+ \left(-\frac{15737765635}{12192768} + \frac{2255}{48} \pi^2 \right) \nu + \frac{76055}{6912} \nu^2 - \frac{127825}{5184} \nu^3 \Biggr] x^3 \\ &+ \left(\frac{77096675}{2032128} + \frac{378515}{12090} \nu - \frac{74045}{6048} \nu^2 \right) \pi x^{7/2} \\ &+ \left[\frac{2550713843998885153}{2214468081745920} - \frac{9203}{126} \gamma_{\rm E} - \frac{45245}{756} \pi^2 - \frac{252755}{2646} \ln 2 - \frac{78975}{1568} \ln 3 - \frac{9203}{252} \ln x \right. \\ &+ \left(- \frac{680712846248317}{337983528960} - \frac{488986}{1323} \gamma_{\rm E} + \frac{109295}{1792} \pi^2 - \frac{1245514}{1323} \ln 2 + \frac{78975}{392} \ln 3 - \frac{244493}{1323} \ln x \right) \nu \\ &+ \left(\frac{7510073635}{24385536} - \frac{11275}{1152} \pi^2 \right) \nu^2 + \frac{1292395}{96768} \nu^3 - \frac{5975}{768} \nu^4 \Biggr] x^4 \\ &+ \left[- \frac{93098188434443}{150214901760} + \frac{1712}{21} \gamma_{\rm E} + \frac{80}{3} \pi^2 + \frac{856}{21} \ln(16x) \right] \\ &+ \left(\frac{1492917260735}{1072963584} - \frac{2255}{48} \pi^2 \right) \nu - \frac{45293335}{1016064} \nu^2 - \frac{10323755}{1596672} \nu^3 \Biggr] \pi x^{9/2} + \mathcal{O}(x^5) \Biggr\}, \tag{8}$$

where the integration constant ψ_0 is determined by initial conditions, *e.g.*, when the wave frequency enters the detector's band. The results hereabove (6)–(8) give the prediction of Einstein's general relativity for the GW frequency and phase chirp of non-spinning compact binaries up to 4.5PN precision.

Up to now we dealt with the time-domain GW half-phase $\psi(t)$. It is useful (especially for data-analysis purposes) to also control the frequency-domain GW half-phase, which we denote $\Psi(F)$. Its PN expansion is obtained by using the stationary phase approximation (SPA) [77] and reads:

$$\begin{split} \Psi_{\text{SPA}} &= 2\pi F \, T_0 + \Psi_0 \\ &+ \frac{3 \, v^{-5}}{128 \nu} \Biggl\{ 1 + \left(\frac{3715}{756} + \frac{55}{9} \nu \right) v^2 - 16\pi v^3 \\ &+ \left(\frac{15293365}{508032} + \frac{27145}{504} \nu + \frac{3085}{72} \nu^2 \right) v^4 + \left(\frac{38645}{252} - \frac{65}{3} \nu \right) \pi v^5 \ln v \\ &+ \left[\frac{11583231236531}{4694215680} - \frac{640}{3} \pi^2 - \frac{6848}{21} \gamma_{\text{E}} - \frac{6848}{21} \ln(4v) + \left(-\frac{15737765635}{3048192} + \frac{2255}{12} \pi^2 \right) \nu \right. \\ &+ \left. \frac{76055}{1728} \nu^2 - \frac{127825}{1296} \nu^3 \right] v^6 \\ &+ \left[\frac{77096675}{254016} + \frac{378515}{1512} \nu - \frac{74045}{756} \nu^2 \right] \pi v^7 \end{split}$$

$$+ \left[-\frac{2550713843998885153}{276808510218240} + \frac{90490}{189}\pi^{2} + \frac{36812}{63}\gamma_{\rm E} + \frac{1011020}{1323}\ln 2 + \frac{78975}{196}\ln 3 + \frac{18406}{63}\ln v \right. \\ \left. + \left(\frac{680712846248317}{42247941120} - \frac{109295}{224}\pi^{2} + \frac{3911888}{1323}\gamma_{\rm E} + \frac{9964112}{1323}\ln 2 - \frac{78975}{49}\ln 3 + \frac{1955944}{1323}\ln v \right) \nu \right. \\ \left. + \left(-\frac{7510073635}{3048192} + \frac{11275}{144}\pi^{2} \right) \nu^{2} - \frac{1292395}{12096}\nu^{3} + \frac{5975}{96}\nu^{4} \right] v^{8}\ln v \right. \\ \left. + \left[\frac{105344279473163}{18776862720} - \frac{640}{3}\pi^{2} - \frac{13696}{21}\gamma_{\rm E} - \frac{13696}{21}\ln(4v) \right. \\ \left. + \left(-\frac{1492917260735}{134120448} + \frac{2255}{6}\pi^{2} \right) \nu + \frac{45293335}{127008}\nu^{2} + \frac{10323755}{199584}\nu^{3} \right] \pi v^{9} + \mathcal{O}(v^{10}) \right\},$$

where $v \equiv \left(\frac{\pi GmF}{c^3}\right)^{1/3}$ with F being the Fourier frequency, and where T_0 and Ψ_0 are two integration constants. Again we have adjusted T_0 in order to simplify the result (and we have absorbed the usual $-\frac{\pi}{4}$ into Ψ_0). The coefficients up to 3.5PN, as well as the 4.5PN piece, are already in use; see *e.g.* App. A of [78].

In order to get intuition on the relative contribution of each PN order to the signal, we provide in Table I rough numerical estimates for the number of accumulated GW cycles in the frequency band of current and future detectors. Our naive estimation does not take the various detector noises into account, and a more realistic estimation should be performed [79]. Nevertheless, it can be useful to gain insight on the behavior of the PN expansion, which seems to converge well, as we see from Table I. For all the typical compact binaries in Table I, we find that the 4PN and 4.5PN orders amount to about a tenth of a cycle (less than 1 radian). This suggests that systematic errors due to the PN modeling may be dominated by statistical errors and negligible for LISA. However, this should be confirmed by detailed investigations along the lines of [80].

Detector	LIGO/Virgo		ET		LISA	
Masses (M_{\odot})	1.4×1.4	10×10	1.4×1.4	500×500	$10^5 \times 10^5$	$10^7 \times 10^7$
PN order	cumulative number of cycles					
Newtonian	2562.599	95.502	744 401.36	37.90	28095.39	9.534
1PN	143.453	17.879	4 4 3 3.85	9.60	618.31	3.386
1.5PN	-94.817	-20.797	-1005.78	-12.63	-265.70	-5.181
2PN	5.811	2.124	23.94	1.44	11.35	0.677
2.5PN	-8.105	-4.604	-17.01	-3.42	-12.47	-1.821
3PN	1.858	1.731	2.69	1.43	2.59	0.876
3.5PN	-0.627	-0.689	-0.93	-0.59	-0.91	-0.383
4PN	-0.107	-0.064	-0.12	-0.04	-0.12	-0.013
4.5PN	0.098	0.118	0.14	0.10	0.14	0.065

TABLE I. Contribution of each PN order to the total number of accumulated cycles inside the detector's frequency band, for typical (but non-spinning) quasi-circular compact binaries observed by current and future detectors. We have approximated the frequency bands of LIGO/Virgo, Einstein Telescope (ET) and LISA with step functions, respectively between $[30 \text{ Hz}, 10^3 \text{ Hz}]$, $[1 \text{ Hz}, 10^4 \text{ Hz}]$ and $[10^{-4} \text{ Hz}, 10^{-1} \text{ Hz}]$. When the merger occurs within the frequency band of the detector, the exit frequency is taken to be the Schwarzschild ISCO, $f_{\rm ISCO} = c^3/(6^{3/2}\pi Gm)$. The contributions due to the non-linearities of GR (*e.g.*, tails) increase with the PN order and are detailed in [24].

Besides the chirp described by the results (6)-(8), it is also important to compute the wave amplitude, in view of the data analysis of LISA [81-83] and high-accuracy comparisons with numerical relativity (see *e.g.* [84-87]). We decompose the waveform, at leading order in the distance R to the source, onto a basis of spin-weighted spherical harmonics (following the conventions of [88, 89])

$$h_{+} - \mathrm{i}h_{\times} = \frac{8Gm\nu x}{Rc^{2}} \sqrt{\frac{\pi}{5}} \sum_{\ell=2}^{+\infty} \sum_{\mathrm{m}=-\ell}^{\ell} H_{\ell\mathrm{m}} e^{-\mathrm{i}\mathrm{m}\psi} Y_{-2}^{\ell\mathrm{m}}, \qquad (10)$$

where the half-phase variable is given by (8). All $H_{\ell m}$ modes are currently known at 3.5PN order for spinning, non-precessing, quasi-circular orbits [88–92]. Although we were able to derive the phase with 4.5PN accuracy, the same precision for the modes is yet out of reach, since, even though the 4.5PN radiation-reaction terms in the equations of motion are known [93, 94], neither the source quadrupole moment nor the non-linear contributions to the GW propagation are fully controlled at 4.5PN order (only the contributions that enter the 4.5PN flux for circular orbits are known). We thus report the extension of the dominant quadrupole mode (ℓ , m) = (2, 2) for non-spinning, quasi-circular orbits up to 4PN order:

$$\begin{aligned} H_{22} &= 1 + \left(-\frac{107}{42} + \frac{55}{42} \nu \right) x + 2\pi x^{3/2} + \left(-\frac{2173}{1512} - \frac{1069}{216} \nu + \frac{2047}{1512} \nu^2 \right) x^2 + \left[-\frac{107\pi}{21} + \left(\frac{34\pi}{21} - 24 \,\mathrm{i} \right) \nu \right] x^{5/2} \\ &+ \left[\frac{27027409}{646800} - \frac{856}{105} \gamma_{\rm E} + \frac{428 \,\mathrm{i} \pi}{105} + \frac{2\pi^2}{3} + \left(-\frac{278185}{33264} + \frac{41\pi^2}{96} \right) \nu - \frac{20261}{2772} \nu^2 + \frac{114635}{99792} \nu^3 - \frac{428}{105} \ln(16x) \right] x^3 \\ &+ \left[-\frac{2173\pi}{756} + \left(-\frac{2495\pi}{378} + \frac{14333 \,\mathrm{i}}{162} \right) \nu + \left(\frac{40\pi}{27} - \frac{4066 \,\mathrm{i}}{945} \right) \nu^2 \right] x^{7/2} \\ &+ \left[-\frac{846557506853}{12713500800} + \frac{45796}{2205} \gamma_{\rm E} - \frac{22898}{2205} \,\mathrm{i} \pi - \frac{107}{63} \pi^2 + \frac{22898}{2205} \ln(16x) \right. \\ &+ \left(-\frac{336005827477}{4237833600} + \frac{15284}{441} \gamma_{\rm E} - \frac{219314}{2205} \,\mathrm{i} \pi - \frac{9755}{32256} \pi^2 + \frac{7642}{441} \ln(16x) \right) \nu \\ &+ \left(\frac{256450291}{7413120} - \frac{1025}{1008} \pi^2 \right) \nu^2 - \frac{81579187}{15567552} \nu^3 + \frac{26251249}{31135104} \nu^4 \right] x^4 + \mathcal{O}(x^{9/2}) \,. \end{aligned}$$

Satisfyingly, this result is in perfect agreement with linear black-hole perturbation theory in the limit when $\nu \to 0$; see App. B of [58]. Again, it would be interesting to compare the PN prediction (11) with second-order (numerical or analytical) BH perturbation theory.

We thank Adam Pound for discussions on BH perturbation and GSF results, Bala Iyer for his longstanding support and interest in this project, and Alessandra Buonanno for suggestions. F.L. is grateful to the Institut d'Astrophysique de Paris for its hospitality during this project. He received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No 817791). G.F. thanks IIT Madras for a Visiting Faculty Fellow position under the IoE program during the completion of this work.

- [†] faye@iap.fr
- [‡] quentin.henry@aei.mpg.de
- § francois.larrouturou@desy.de
- ¶ david.trestini@obspm.fr
- [1] A. Einstein, Über gravitationswellen, Sitzber. Preuss. Akad. Wiss. Berlin 1, 154 (1918).
- [2] L. Landau and E. Lifshitz, The classical theory of fields (Pergamon, Oxford, 1971).
- [3] P. Peters and J. Mathews, Gravitational radiation from point masses in a keplerian orbit, Phys. Rev. 131, 435 (1963).
- [4] C. Cutler, T. Apostolatos, L. Bildsten, L. Finn, E. Flanagan, D. Kennefick, D. Markovic, A. Ori, E. Poisson, G. Sussman, and K. Thorne, The last three minutes: Issues in gravitational-wave measurements of coalescing compact binaries, Phys. Rev. Lett. 70, 2984 (1993).
- [5] C. Cutler and E. Flanagan, Gravitational waves from merging compact binaries: How accurately can one extract the binary's parameters from the inspiral waveform?, Phys. Rev. D 49, 2658 (1994).
- [6] L. Blanchet and T. Damour, Radiative gravitational fields in general relativity. i. general structure of the field outside the source, Phil. Trans. Roy. Soc. Lond. A 320, 379 (1986).
- [7] L. Blanchet, Radiative gravitational fields in general relativity. ii. asymptotic behaviour at future null infinity, Proc. Roy. Soc. Lond. A 409, 383 (1987).
- [8] L. Blanchet and T. Damour, Tail-transported temporal correlations in the dynamics of a gravitating system, Phys. Rev. D 37, 1410 (1988).
- [9] L. Blanchet and T. Damour, Hereditary effects in gravitational radiation, Phys. Rev. D 46, 4304 (1992).
- [10] L. Blanchet, On the multipole expansion of the gravitational field, Class. Quant. Grav. 15, 1971 (1998), gr-qc/9801101.
- [11] L. Blanchet, T. Damour, and B. R. Iyer, Gravitational waves from inspiralling compact binaries: Energy loss and wave form to second post-Newtonian order, Phys. Rev. D 51, 5360 (1995), gr-qc/9501029.

^{*} luc.blanchet@iap.fr

- [12] L. Blanchet, T. Damour, B. R. Iyer, C. M. Will, and A. G. Wiseman, Gravitational radiation damping of compact binary systems to second post-Newtonian order, Phys. Rev. Lett. 74, 3515 (1995), gr-qc/9501027.
- [13] L. Blanchet, B. R. Iyer, C. M. Will, and A. G. Wiseman, Gravitational wave forms from inspiralling compact binaries to second-post-Newtonian order, Class. Quant. Grav. 13, 575 (1996), gr-qc/9602024.
- [14] L. Blanchet, Energy losses by gravitational radiation in inspiralling compact binaries to five halves post-Newtonian order, Phys. Rev. D 54, 1417 (1996), erratum Phys. Rev. D 71, 129904(E) (2005), gr-qc/9603048.
- [15] L. Blanchet, Gravitational-wave tails of tails, Class. Quant. Grav. 15, 113 (1998), gr-qc/9710038.
- [16] L. Blanchet, B. R. Iyer, and B. Joguet, Gravitational waves from inspiralling compact binaries: Energy flux to third post-Newtonian order, Phys. Rev. D 65, 064005 (2002), erratum Phys. Rev. D, 71:129903(E), 2005, gr-qc/0105098.
- [17] L. Blanchet, G. Faye, B. R. Iyer, and B. Joguet, Gravitational-wave inspiral of compact binary systems to 7/2 post-Newtonian order, Phys. Rev. D 65, 061501(R) (2002), erratum Phys. Rev. D, 71:129902(E), 2005, gr-qc/0105099.
- [18] L. Blanchet and B. R. Iyer, Hadamard regularization of the third post-Newtonian gravitational wave generation of two point masses, Phys. Rev. D 71, 024004 (2004), gr-qc/0409094.
- [19] L. Blanchet, T. Damour, G. Esposito-Farèse, and B. R. Iyer, Gravitational radiation from inspiralling compact binaries completed at the third post-Newtonian order, Phys. Rev. Lett. 93, 091101 (2004), gr-qc/0406012.
- [20] L. Blanchet, T. Damour, G. Esposito-Farèse, and B. R. Iyer, Dimensional regularization of the third post-Newtonian gravitational wave generation of two point masses, Phys. Rev. D 71, 124004 (2005), gr-qc/0503044.
- [21] L. Blanchet, Gravitational radiation from post-Newtonian sources and inspiralling compact binaries, Living Rev. Relativ. 17, 2 (2014), arXiv:1310.1528 [gr-qc].
- [22] G. Cho, R. A. Porto, and Z. Yang, Gravitational radiation from inspiralling compact objects: Spin effects to the fourth post-newtonian order, Physical Review D 106, L101501 (2022).
- [23] The ancillary file GW_4PN_supplementary_material.m collects the results associated with the gravitational wave phase at 4.5PN and the $(\ell, m) = (2, 2)$ mode at 4PN.
- [24] L. Blanchet, G. Faye, Q. Henry, F. Larrouturou, and D. Trestini, Gravitational-wave flux and quadrupole modes from quasicircular nonspinning compact binaries to the fourth post-Newtonian order, Phys. Rev. D 108, 064041 (2023), arXiv:2304.11186 [gr-qc].
- [25] L. Blanchet and B. S. Sathyaprakash, Detecting the tail effect in gravitational wave experiments, Phys. Rev. Lett. 74, 1067 (1995).
- [26] K. Arun, B. Iyer, M. Qusailah, and B. Sathyaprakash, Testing post-Newtonian theory with gravitational wave observations, Class. Quant. Gravity 23, L37 (2006), gr-qc/0604018.
- [27] B. Abbott *et al.* (LIGO Scientific Collaboration and VIRGO Collaboration), Tests of general relativity with GW150914, Phys. Rev. Lett. **116**, 221101 (2016), arXiv:1602.03841 [gr-qc].
- [28] B. P. Abbott et al. (LIGO Scientific, Virgo), Tests of General Relativity with GW170817, Phys. Rev. Lett. 123, 011102 (2019), arXiv:1811.00364 [gr-qc].
- [29] A. Sesana, Prospects for Multiband Gravitational-Wave Astronomy after GW150914, Phys. Rev. Lett. 116, 231102 (2016), arXiv:1602.06951 [gr-qc].
- [30] P. Peters, Gravitational radiation and the motion of two point masses, Phys. Rev. 136, B1224 (1964).
- [31] P. Jaranowski and G. Schäfer, Towards the fourth post-Newtonian Hamiltonian for two-point-mass systems, Phys. Rev. D 86, 061503(R) (2012), arXiv:1207.5448 [gr-qc].
- [32] P. Jaranowski and G. Schäfer, Dimensional regularization of local singularities in the 4th post-Newtonian two-point-mass hamiltonian, Phys. Rev. D 87, 081503(R) (2013), arXiv:1303.3225 [gr-qc].
- [33] T. Damour, P. Jaranowski, and G. Schäfer, Non-local-in-time action for the fourth post-Newtonian conservative dynamics of two-body systems, Phys. Rev. D 89, 064058 (2014), arXiv:1401.4548 [gr-qc].
- [34] T. Damour, P. Jaranowski, and G. Schäfer, On the conservative dynamics of two-body systems at the fourth post-Newtonian approximation of general relativity, Phys. Rev. D **93**, 084014 (2016), arXiv:1601.01283 [gr-qc].
- [35] D. Bini and T. Damour, Analytical determination of the two-body gravitational interaction potential at the fourth post-Newtonian approximation, Phys. Rev. D 87, 121501(R) (2013), arXiv:1305.4884 [gr-qc].
- [36] L. Bernard, L. Blanchet, A. Bohé, G. Faye, and S. Marsat, Fokker action of non-spinning compact binaries at the fourth post-Newtonian approximation, Phys. Rev. D 93, 084037 (2016), arXiv:1512.02876 [gr-qc].
- [37] L. Bernard, L. Blanchet, A. Bohé, G. Faye, and S. Marsat, Energy and periastron advance of compact binaries on circular orbits at the fourth post-Newtonian order, Phys. Rev. D 95, 044026 (2017), arXiv:1610.07934 [gr-qc].
- [38] L. Bernard, L. Blanchet, A. Bohé, G. Faye, and S. Marsat, Dimensional regularization of the ir divergences in the Fokker action of point-particle binaries at the fourth post-Newtonian order, Phys. Rev. D 96, 104043 (2017), arXiv:1706.08480 [gr-qc].
- [39] L. Bernard, L. Blanchet, G. Faye, and T. Marchand, Center-of-mass equations of motion and conserved integrals of compact binary systems at the fourth post-Newtonian order, Phys. Rev. D 97, 044037 (2018), arXiv:1711.00283 [gr-qc].
- [40] S. Foffa and R. Sturani, The dynamics of the gravitational two-body problem in the post-Newtonian approximation at quadratic order in the Newton's constant, Phys. Rev. D 87, 064011 (2013), arXiv:1206.7087 [gr-qc].
- [41] S. Foffa and R. Sturani, Tail terms in gravitational radiation reaction via effective field theory, Phys. Rev. D 87, 044056 (2012), arXiv:1111.5488 [gr-qc].
- [42] C. R. Galley, A. K. Leibovich, R. A. Porto, and A. Ross, Tail effect in gravitational radiation reaction: Time nonlocality and renormalization group evolution, Phys. Rev. D 93, 124010 (2016), arXiv:1511.07379 [gr-qc].
- [43] S. Foffa, P. Mastrolia, R. Sturani, and C. Sturm, Effective field theory approach to the gravitational two-body dynamics at fourth post-Newtonian order and quintic in the newton constant, Phys. Rev. D 95, 104009 (2017), arXiv:1612.00482 [gr-qc].
- [44] R. A. Porto and I. Z. Rothstein, Apparent ambiguities in the post-Newtonian expansion for binary systems,

Phys. Rev. D 96, 024062 (2017), arXiv:1703.06433 [gr-qc].

- [45] S. Foffa and R. Sturani, Conservative dynamics of binary systems to fourth post-Newtonian order in the EFT approach i: Regularized Lagrangian, Phys. Rev. D 100, 024047 (2019), arXiv:1903.05113 [gr-qc].
- [46] S. Foffa, R. Porto, I. Rothstein, and R. Sturani, Conservative dynamics of binary systems to fourth post-Newtonian order in the EFT approach ii: Renormalized Lagrangian, Phys. Rev. D 100, 024048 (2019), arXiv:1903.05118 [gr-qc].
- [47] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, Fourth post-Newtonian hamiltonian dynamics of two-body systems from an effective field theory approach, 2003.01692 [hep-th] (2020), arXiv preprint.
- [48] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, The fifth-order post-Newtonian hamiltonian dynamics of twobody systems from an effective field theory approach: Potential contributions, Nuclear Physics B 983, 115900 (2022), arXiv:2110.13822 [gr-qc].
- [49] D. Bini, T. Damour, and A. Geralico, Sixth post-Newtonian local-in-time dynamics of binary systems, Phys. Rev. D 102, 024061 (2020), arXiv:2004.05407 [gr-qc].
- [50] T. Marchand, Q. Henry, F. Larrouturou, S. Marsat, G. Faye, and L. Blanchet, The mass quadrupole moment of compact binary systems at the fourth post-Newtonian order, Class. Quant. Grav. 37, 215006 (2020), arXiv:2003.13672 [gr-qc].
- [51] F. Larrouturou, Q. Henry, L. Blanchet, and G. Faye, The quadrupole moment of compact binaries to the fourth post-Newtonian order: I. Non-locality in time and infra-red divergencies, Class. Quant. Grav. 39, 115007 (2022), arXiv:2110.02240 [gr-qc].
- [52] F. Larrouturou, L. Blanchet, Q. Henry, and G. Faye, The quadrupole moment of compact binaries to the fourth post-Newtonian order: II. Dimensional regularization and renormalization, Class. Quant. Grav. 39, 115008 (2022), arXiv:2110.02243 [gr-qc].
- [53] L. Blanchet, G. Faye, and F. Larrouturou, The quadrupole moment of compact binaries to the fourth post-Newtonian order: from source to canonical moment, Class. Quant. Grav. **39**, 195003 (2022), arXiv:2204.11293 [gr-qc].
- [54] Q. Henry, G. Faye, and L. Blanchet, The current-type quadrupole moment and gravitational-wave mode $(\ell, m) = (2, 1)$ of compact binary systems at the third post-Newtonian order, Class. Quant. Grav. **38**, 185004 (2021), arXiv:2105.10876 [gr-qc].
- [55] D. Trestini, F. Larrouturou, and L. Blanchet, The quadrupole moment of compact binaries to the fourth post-Newtonian order: relating the harmonic and radiative metrics, Class. Quant. Grav. 40, 055006 (2023), arXiv:2209.02719 [gr-qc].
- [56] D. Trestini and L. Blanchet, Gravitational-wave tails of memory, Phys. Rev. D 107, 104048 (2023), arXiv:2301.09395 [gr-qc].
- [57] M. Sasaki, Post-Newtonian expansion of the ingoing-wave regge-wheeler function, Prog. Theor. Phys. 92, 17 (1994).
- [58] H. Tagoshi and M. Sasaki, Post-Newtonian expansion of gravitational-waves from a particle in circular orbit around a schwarzschild black-hole, Prog. Theor. Phys. 92, 745 (1994), gr-qc/9405062.
- [59] T. Tanaka, H. Tagoshi, and M. Sasaki, Gravitational waves by a particle in circular orbit around a schwarzschild black hole: 5.5 Post-Newtonian formula, Prog. Theor. Phys. 96, 1087 (1996), gr-qc/9701050.
- [60] R. Fujita, Gravitational radiation for extreme mass ratio inspirals to the 14th post-Newtonian order, Prog. Theor. Phys. 127, 583 (2012), arXiv:1104.5615 [gr-qc].
- [61] R. Fujita, Gravitational waves from a particle in circular orbits around a schwarzschild black hole to the 22nd post-Newtonian order, Prog. Theor. Phys. 128, 971 (2012), arXiv:1211.5535 [gr-qc].
- [62] B. Wardell, A. Pound, N. Warburton, J. Miller, L. Durkan, and A. Le Tiec, Gravitational Waveforms for Compact Binaries from Second-Order Self-Force Theory, Phys. Rev. Lett. 130, 241402 (2023), arXiv:2112.12265 [gr-qc].
- [63] A. Albertini, A. Nagar, A. Pound, N. Warburton, B. Wardell, L. Durkan, and J. Miller, Comparing second-order gravitational self-force, numerical relativity, and effective one body waveforms from inspiralling, quasicircular, and nonspinning black hole binaries, Phys. Rev. D 106, 084061 (2022), arXiv:2208.01049 [gr-qc].
- [64] A. Albertini, A. Nagar, A. Pound, N. Warburton, B. Wardell, L. Durkan, and J. Miller, Comparing second-order gravitational self-force and effective one body waveforms from inspiralling, quasicircular and nonspinning black hole binaries. II. The large-mass-ratio case, Phys. Rev. D 106, 084062 (2022), arXiv:2208.02055 [gr-qc].
- [65] E. Poisson and M. Sasaki, Gravitational radiation from a particle in circular orbit around a black hole. v. black-hole absorption and tail corrections, Phys. Rev. D 51, 5753 (1995), gr-qc/9412027.
- [66] H. Tagoshi, S. Mano, and E. Takasugi, Post-Newtonian expansion of gravitational waves from a particle in circular orbits around a rotating black hole, Prog. Theor. Phys. 98, 829 (1997), gr-qc/9711072.
- [67] K. Alvi, Energy and angular momentum flow into a black hole in a binary, Phys. Rev. D 64, 104020 (2001), arXiv:gr-qc/0107080.
- [68] R. A. Porto, Absorption effects due to spin in the worldline approach to black hole dynamics, Phys. Rev. D 77, 064026 (2008), arXiv:0710.5150 [hep-th].
- [69] K. Chatziioannou, E. Poisson, and N. Yunes, Tidal heating and torquing of a kerr black hole to next-to-leading order in the tidal coupling, Phys. Rev. D 87, 044022 (2013), arXiv:1211.1686 [gr-qc].
- [70] M. V. S. Saketh, J. Steinhoff, J. Vines, and A. Buonanno, Modeling horizon absorption in spinning binary black holes using effective worldline theory, Phys. Rev. D 107, 084006 (2023), arXiv:2212.13095 [gr-qc].
- [71] T. Damour, B. R. Iyer, and B. S. Sathyaprakash, Improved filters for gravitational waves from inspiralling compact binaries, Phys. Rev. D 57, 885 (1998), arXiv:gr-qc/9708034.
- [72] A. Buonanno and T. Damour, Effective one-body approach to general relativistic two-body dynamics, Phys. Rev. D 59, 084006 (1999), gr-qc/9811091.
- [73] A. Buonanno, B. Iyer, Y. Pan, E. Ochsner, and B. Sathyaprakash, Comparison of post-Newtonian templates for compact binary inspiral signals in gravitational-wave detectors, Phys. Rev. D 80, 084043 (2009), arXiv:0907.0700 [gr-qc].

- [74] T. Damour and A. Nagar, The effective one-body description of the two-body problem, in *Mass and motion in general relativity*, edited by L. Blanchet, A. Spallicci, and B. Whiting (Springer, 2011) p. 211.
- [75] A. Wiseman, Coalescing binary-systems of compact objects to (post)^{5/2}-Newtonian order. iv. the gravitational-wave tail, Phys. Rev. D 48, 4757 (1993).
- [76] L. Blanchet and G. Schäfer, Gravitational wave tails and binary star systems, Class. Quant. Grav. 10, 2699 (1993).
- [77] W. Tichy, E. E. Flanagan, and E. Poisson, Can the postNewtonian gravitational wave form of an inspiraling binary be improved by solving the energy balance equation numerically?, Phys. Rev. D 61, 104015 (2000), arXiv:gr-qc/9912075.
- [78] G. Pratten, S. Husa, C. Garcia-Quiros, M. Colleoni, A. Ramos-Buades, H. Estelles, and R. Jaume, Setting the cornerstone for a family of models for gravitational waves from compact binaries: The dominant harmonic for nonprecessing quasicircular black holes, Phys. Rev. D 102, 064001 (2020), arXiv:2001.11412 [gr-qc].
- [79] K. Arun, B. Iyer, B. Sathyaprakash, and P. Sundararajan, Parameter estimation of inspiralling compact binaries using 3.5 post-Newtonian gravitational wave phasing: The non-spinning case, Phys. Rev. D 71, 084008 (2005), gr-qc/0411146.
- [80] C. B. Owen, C.-J. Haster, S. Perkins, N. J. Cornish, and N. Yunes, Waveform accuracy and systematic uncertainties in current gravitational wave observations, Phys. Rev. D 108, 044018 (2023), arXiv:2301.11941 [gr-qc].
- [81] R. W. Hellings and T. A. Moore, The Information content of gravitational wave harmonics in compact binary inspiral, Class. Quant. Grav. 20, S181 (2003), arXiv:gr-qc/0207102.
- [82] T. A. Moore and R. W. Hellings, The Angular resolution of space based gravitational wave detectors, AIP Conf. Proc. 523, 255 (2000), arXiv:gr-qc/9910116.
- [83] K. G. Arun, B. R. Iyer, B. S. Sathyaprakash, S. Sinha, and C. Van Den Broeck, Higher signal harmonics, LISA's angular resolution and dark energy, Phys. Rev. D 76, 104016 (2007), [Erratum: Phys.Rev.D 76, 129903 (2007)], arXiv:0707.3920 [astro-ph].
- [84] P. Grandclement and J. Novak, Spectral methods for numerical relativity, Living Rev. Rel. 12, 1 (2009), arXiv:0706.2286 [gr-qc].
- [85] M. Boyle, D. A. Brown, L. E. Kidder, A. H. Mroue, H. P. Pfeiffer, M. A. Scheel, G. B. Cook, and S. A. Teukolsky, Highaccuracy comparison of numerical relativity simulations with post-Newtonian expansions, Phys. Rev. D 76, 124038 (2007), arXiv:0710.0158 [gr-qc].
- [86] S. Bernuzzi, M. Thierfelder, and B. Bruegmann, Accuracy of numerical relativity waveforms from binary neutron star mergers and their comparison with post-Newtonian waveforms, Phys. Rev. D 85, 104030 (2012), arXiv:1109.3611 [gr-qc].
- [87] S. Borhanian, K. G. Arun, H. P. Pfeiffer, and B. S. Sathyaprakash, Comparison of post-Newtonian mode amplitudes with numerical relativity simulations of binary black holes, Class. Quant. Grav. 37, 065006 (2020), arXiv:1901.08516 [gr-qc].
- [88] L. Blanchet, G. Faye, B. R. Iyer, and S. Sinha, The third post-Newtonian gravitational wave polarisations and associated spherical harmonic modes for inspiralling compact binaries in quasi-circular orbits, Class. Quant. Grav. 25, 165003 (2008), arXiv:0802.1249 [gr-qc].
- [89] G. Faye, S. Marsat, L. Blanchet, and B. R. Iyer, The third and a half post-Newtonian gravitational wave quadrupole mode for quasi-circular inspiralling compact binaries, Class. Quant. Grav. 29, 175004 (2012), arXiv:1204.1043 [gr-qc].
- [90] K. Arun, L. Blanchet, B. R. Iyer, and M. S. Qusailah, The 2.5PN gravitational wave polarisations from inspiralling compact binaries in circular orbits, Class. Quant. Grav. 21, 3771 (2004), erratum Class. Quant. Grav., 22:3115, 2005, gr-qc/0404085.
- [91] Q. Henry, S. Marsat, and M. Khalil, Spin contributions to the gravitational-waveform modes for spin-aligned binaries at the 3.5PN order, Phys. Rev. D 106, 124018 (2022), arXiv:2209.00374 [gr-qc].
- [92] Q. Henry, Complete gravitational-waveform amplitude modes for quasicircular compact binaries to the 3.5PN order, Phys. Rev. D 107, 044057 (2023), arXiv:2210.15602 [gr-qc].
- [93] A. Gopakumar, B. R. Iyer, and S. Iyer, Second post-Newtonian gravitational radiation reaction for two-body systems: Nonspinning bodies, Phys. Rev. D 55, 6030 (1997), gr-qc/9703075.
- [94] A. K. Leibovich, B. A. Pardo, and Z. Yang, Radiation reaction for nonspinning bodies at 4.5PN in the effective field theory approach, Phys. Rev. D 108, 024017 (2023), arXiv:2302.11016 [gr-qc].