

CLASSIFICATION, α -INNER DERIVATIONS AND α -CENTROIDS OF FINITE-DIMENSIONAL COMPLEX HOM-TRIALGEBRAS

BOUZID MOSBAHI*

University of Sfax, Faculty of Sciences of Sfax, BP 1171, 3000 Sfax, Tunisia.

AHMED ZAHARI†

Université de Haute Alsace,
IRIMAS-Département de Mathématiques,
18, rue des Frères Lumière F-68093 Mulhouse, France.

IMED BASDOURI‡

University of Gafsa, Faculty of Sciences of Gafsa, 2112 Gafsa, Tunisia.

Abstract. In the current research work, our basic objective is to investigate the structure of Hom-associative trialgebras. Next, we build up one important class of Hom-associative trialgebras and provide properties of right, left and middle operations in Hom-associative trialgebras. Furthermore, we describe the classification of n -dimensional Hom-associative trialgebras for $n \leq 3$. Additionally, the properties of the Inner-derivations and centroids are identified and discussed. Eventually the Inner-derivations and centroids are computed.

AMS Subject Classification: 17A30, 17A32, 16D20, 16W25, 17B63 .

Keywords: Hom-associative trialgebra, Classification, Inner-derivation, Centroid.

1 Introduction

Notably, a Hom-associative trialgebra $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ consists of a vector space, two multiplications and a linear self map. It may be perceived as a deformation of an associative algebra, where the associativity condition is twisted by a linear map α , such that when $\alpha = id$, the Hom-associative trialgebras degenerate to exactly triassociative algebras. We aim in this paper to examine the structure of Hom-associative trialgebras. Let \mathcal{A} be an n -dimensional \mathbb{K} -linear space and $\{e_1, e_2, \dots, e_n\}$ be a basis of \mathcal{A} . A Hom-associative trialgebra structure on \mathcal{A} with products μ and λ is determined by $3n^3$ structure constants γ_{ij}^k and δ_{ij}^k , where $e_i \dashv e_j = \sum_{k=1}^n \gamma_{ij}^k e_k$, $e_i \vdash e_j = \sum_{k=1}^n \delta_{ij}^k e_k$ and $e_i \perp e_j = \sum_{k=1}^n \phi_{ij}^k e_k$ and by α which is given by n^2 structure constants a_{ij} , where $\alpha(e_i) = \sum_{j=1}^n a_{ji} e_j$. Requiring the algebra structure to be a Hom-associative trialgebra gives rise to sub-variety $\mathcal{H}_{\mathcal{T}}$ of

*e-mail address: mosbahibouزيد@yahoo.fr

†e-mail address: zaharymaths@gmail.com

‡e-mail address: basdourimed@yahoo.fr

$K^{3n^3+n^2}$. Basic changes in \mathcal{A} yield the natural transport of structure action of $GL_n(k)$ on $\mathcal{H}_{\mathcal{T}}$. As a matter of facts, isomorphism classes of n -dimensional Hom-associative trialgebras are one-to-one correspondence with the orbits of the action of $GL_n(k)$ on $\mathcal{H}_{\mathcal{T}}$.

Classification problems of the Hom-associative trialgebras using the algebraic and geometric technique have drawn much interest in the derivations and centroids of Hom-associative trialgebras. The associative trialgebras introduced by Loday [1] with a motivation to provide dual dialgebras, have been further investigated with regard to several areas in mathematics and physics. The classification of Hom-associative algebras was set forward elaborated by [3] and A. Zahari and I. Bakayoko who addressed the classification of BiHom-associative dialgebras [2].

This paper involves around examining the Inner-derivations and centroids of finite dimensional associative trialgebras. The algebra of Inner-derivations and centroids are highly needed and extremely useful in terms of algebraic and geometric classification problems of algebras.

The current paper is organized as follows. In the first section, we identify the topic alongside with some previously obtained results. The chief objective of this paper is to specify and classify Inner-derivations as well as centroids of Hom-associative trialgebras. In section 2, we tackle the structure of Hom-associative trialgebras. In section 3, we handle the algebraic varieties of Hom-associative trialgebras, and we depict classifications, up to isomorphism, of two-dimensional and three-dimensional Hom-associative trialgebras. We focus upon the classification of the Inner-derivations. Eventually, in Section 4, we present the classification of the centroids. In this case, the concept of derivations and centroids is notably inspired and whetted significant scientific concern from that of finite-dimensional algebras. The algebra of centroids plays an outstanding role in the classification problems as well as in different applications of algebras. As far as our work is concerned, we elaborated classification results of two and three-dimensional Hom-associative trialgebras. All considered algebras and vectors spaces are supposed to be over a field \mathbb{K} of characteristic zero.

2 Hom-associative trialgebras

Definition 2.1. A Hom-associative trialgebra is a 5-tuple $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ consisting of a linear space \mathcal{A} linear maps $\dashv, \vdash, \perp : \mathcal{A} \times \mathcal{A} \longrightarrow \mathcal{A}$ and $\alpha : \mathcal{A} \longrightarrow \mathcal{A}$ satisfying, for all $x, y, z \in \mathcal{A}$, the following conditions :

$$(x \dashv y) \dashv \alpha(z) = \alpha(x) \dashv (y \dashv z), \quad (2.1)$$

$$(x \dashv y) \dashv \alpha(z) = \alpha(x) \dashv (y \vdash z) = \alpha(x) \dashv (y \perp z), \quad (2.2)$$

$$(x \vdash y) \dashv \alpha(z) = \alpha(x) \vdash (y \dashv z), \quad (2.3)$$

$$(x \dashv y) \vdash \alpha(z) = \alpha(x) \vdash (y \vdash z) = (x \perp y) \vdash \alpha(z), \quad (2.4)$$

$$(x \vdash y) \vdash \alpha(z) = \alpha(x) \vdash (y \vdash z), \quad (2.5)$$

$$(x \perp y) \dashv \alpha(z) = \alpha(x) \perp (y \dashv z), \quad (2.6)$$

$$(x \dashv y) \perp \alpha(z) = \alpha(x) \perp (y \vdash z), \quad (2.7)$$

$$(x \vdash y) \perp \alpha(z) = \alpha(x) \vdash (y \perp z), \quad (2.8)$$

$$(x \perp y) \perp \alpha(z) = \alpha(x) \perp (y \perp z), \quad (2.9)$$

for all $x, y, z \in \mathcal{A}$.

Remark 2.2. In addition, α is an endomorphism with respect to \dashv, \vdash and \perp . Then, \mathcal{A} is said to be a

multiplicative Hom-associative trialgebra

$$\begin{aligned}\alpha(x \dashv y) &= \alpha(x) \dashv \alpha(y) \quad ; \quad \alpha(x \vdash y) = \alpha(x) \vdash \alpha(y), \\ \alpha(x \perp y) &= \alpha(x) \perp \alpha(y) \quad ,\end{aligned}$$

for all $x, y \in \mathcal{A}$.

Definition 2.3. A morphism of Hom-associative trialgebra is a linear map

$$f : (\mathcal{A}, \dashv, \vdash, \perp, \alpha) \rightarrow (\mathcal{A}', \dashv', \vdash', \perp', \alpha')$$

such that

$$\alpha' \circ f = f \circ \alpha,$$

and

$$f(x \dashv y) = f(x) \dashv' f(y), \quad f(x \vdash y) = f(x) \vdash' f(y), \quad f(x \perp y) = f(x) \perp' f(y)$$

for all $x, y \in \mathcal{A}$.

Remark 2.4. A bijective homomorphism is an isomorphism of \mathcal{A}_1 and \mathcal{A}_2 .

Proposition 2.5. Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-associative trialgebra. Therefore, $(\mathcal{A}, \dashv, \perp, *, \alpha)$ is a Hom-associative trialgebra, where for any $x, y, z \in \mathcal{A}$, $x * y = x \vdash y + x \perp y$.

Proof. We prove only one axiom, as others are proven similarly. For any $x, y, z \in \mathcal{A}$,

$$\begin{aligned}(x * y) \dashv \alpha(z) &= (x \vdash y + x \perp y) \dashv \alpha(z) \\ &= (x \vdash y) \dashv \alpha(z) + (x \perp y) \dashv \alpha(z) \\ &= \alpha(x) \vdash (y \dashv z) + \alpha(x) \perp (y \dashv z) = \alpha(x) * (y \dashv z),\end{aligned}$$

which ends the proof. □

Proposition 2.6. Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-associative trialgebra and $x * y = x \vdash y + x \dashv y + x \perp y$. From this perspective, $(\mathcal{A}, *, \alpha)$ is a Hom-associative algebra.

Proof. For any $x, y, z \in \mathcal{A}$,

$$\begin{aligned}(x * y) * \alpha(z) &= \alpha(x) * (y * z) \\ &= (x \vdash y + x \dashv y + x \perp y) * \alpha(z) - \alpha(x) * (y \vdash z + y \dashv z + y \perp z) \\ &= (x \vdash y) * \alpha(z) + (x \dashv y) * \alpha(z) + (x \perp y) * \alpha(z) - \alpha(x) * (y \vdash z) - \alpha(x) * (y \dashv z) - \alpha(x) * (y \perp z) \\ &= (x \vdash y) \vdash \alpha(z) + (x \vdash y) \dashv \alpha(z) + (x \vdash y) \perp \alpha(z) + (x \dashv y) \vdash \alpha(z) + (x \dashv y) \dashv \alpha(z) + (x \dashv y) \perp \alpha(z) \\ &\quad + (x \perp y) \vdash \alpha(z) + (x \perp y) \dashv \alpha(z) + (x \perp y) \perp \alpha(z) - \alpha(x) \vdash (y \vdash z) - \alpha(x) \dashv (y \vdash z) - \alpha(x) \perp (y \vdash z) \\ &\quad - \alpha(x) \vdash (y \dashv z) - \alpha(x) \dashv (y \dashv z) - \alpha(x) \perp (y \dashv z) - \alpha(x) \vdash (y \perp z) - \alpha(x) \dashv (y \perp z) - \alpha(x) \perp (y \perp z).\end{aligned}$$

This corroborates that $(\mathcal{A}, *, \alpha)$ is a Hom-associative algebra. □

Corollary 2.7. Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-associative trialgebra. Then, \mathcal{A} is a Hom-associative algebra with respect to the multiplicative $*$: $\mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$

$$x * y = x \dashv y + x \vdash y - x \perp y \quad \text{for any } x, y \in \mathcal{A}.$$

Proposition 2.8. Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-associative trialgebra. Then $(\mathcal{A}, \dashv, \vdash', \alpha)$ is a Hom-associative trialgebra, where $x \vdash' y = x \vdash y + x \dashv y$, for $x, y \in \mathcal{A}$.

Proof. The proof is straightforward by calculation by using Definition 2.1. \square

Proposition 2.9. *Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-associative trialgebra. Then $(\mathcal{A}, \dashv^{op}, \vdash^{op}, \perp^{op}, \alpha)$ is also a Hom-associative trialgebra, with $x \dashv^{op} y = y \vdash x$, $x \vdash^{op} y = y \dashv x$, $x \perp^{op} y = y \perp x$, for any $x, y \in \mathcal{A}$.*

Proof. It comes immediately from Definition 2.1. \square

Definition 2.10. A Hom-Leibniz Poisson algebra is the triplet $(\mathcal{A}, \cdot, [\cdot, \cdot], \alpha)$ consisting of a linear space \mathcal{T} , two linear maps $\cdot, [\cdot, \cdot] : \mathcal{T} \times \mathcal{T} \longrightarrow \mathcal{T}$ and a linear map $\alpha : \mathcal{T} \longrightarrow \mathcal{T}$ satisfying the following statements

- i) $(\mathcal{T}, \cdot, \alpha)$ is a Hom-associative trialbra,
- ii) $(\mathcal{T}, [\cdot, \cdot], \alpha)$ is a Hom-Leibniz algebra ,
- iii) $[x \cdot y, \alpha(z)] = \alpha(x) \cdot [y, z] + [x, z] \cdot \alpha(y)$ for all $x, y, z \in \mathcal{T}$

Proposition 2.11. *Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-associative trialgebra. Define now binary operations by*

$$\begin{aligned} x * y &= x \dashv y - y \vdash x \\ [x, y] &= x \perp y - y \perp x. \end{aligned}$$

*Then $(\mathcal{A}, *, [\cdot, \cdot], \alpha)$ becomes a Hom-associative trialgebra.*

Proof. By definition, for any $x, y, z \in \mathcal{A}$, we have

$$\begin{aligned} [x, y] * \alpha(z) &= [x, y] \dashv \alpha(z) - \alpha(z) \vdash [x, y] \\ &= (x \vdash y - y \perp x) \dashv \alpha(z) - \alpha(z) \vdash (x \perp y - y \perp x). \end{aligned}$$

and

$$\begin{aligned} [x * z, \alpha(y)] + [\alpha(x), y * z] &= [x \dashv z - z \vdash x, \alpha(y)] + [\alpha(x), y \dashv z - z \vdash y] \\ &= (x \vdash z - z \vdash x) \perp \alpha(y) - \alpha(y) \perp (x \dashv z - z \vdash x) \\ &\quad + \alpha(x) \perp (y \dashv z - z \vdash y) - (y \dashv z - z \vdash y) \perp \alpha(x). \end{aligned}$$

Since

$$\begin{aligned} (x \perp y) \dashv \alpha(z) &= \alpha(x) \perp (y \vdash z), \\ (y \perp x) \dashv \alpha(z) &= \alpha(y) \perp (x \vdash z), \\ \alpha(z) \vdash (x \perp y) &= (z \vdash x) \perp \alpha(y), \\ \alpha(z) \vdash (y \perp x) &= (z \vdash y) \perp \alpha(x), \\ (x \perp z) \perp \alpha(y) &= \alpha(x) \perp (z \vdash y), \\ \alpha(y) \perp (z \vdash x) &= (y \dashv z) \perp \alpha(x), \end{aligned}$$

in a Hom-associative trialgebra, we get $[x, y] * \alpha(z) = [x * z, \alpha(y)] + [\alpha(x), y * z]$.

This ends the proof. \square

Proposition 2.12. *Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-trialgebra. Therefore, $(*, [-, -], \alpha)$ is a non-commutative Hom-Leibniz-Poisson algebra with respect to the operations $x * y = x \perp y$, $[x, y] = x \dashv y - x \vdash y$, for all $x, y, z \in \mathcal{A}$.*

Proof. If following from a straightforward computation. \square

Corollary 2.13. *Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-associative trialgebra. Hence, $(\mathcal{A}, *, [-, -], \alpha)$ is a non-commutative Hom-Leibniz-Poisson algebra with*

$$x * y = x \dashv y + x \vdash y - x \perp y \quad \text{and} \quad [x, y] = x * y - y * x \quad \text{for any } x, y \in \mathcal{A}.$$

2.1 Classification of Finite-dimensional Complex Hom-associative trialgebras

In this section, Hom-associative trialgebras are classified in low dimension. Note that \mathcal{TH}_n^m denotes m^{th} isomorphism class of Hom-associative trialgebra in dimension n .

Let $(\mathcal{A}, +, \vdash, \perp, \alpha)$ be an n -dimensional Hom-trialgebra of \mathcal{A} . For any $i, j \in \mathbb{N}$, $1 \leq i, j \leq n$, let us put

$$e_i + e_j = \sum_{k=1}^n \gamma_{ij}^k e_k, \quad e_i \vdash e_j = \sum_{k=1}^n \delta_{ij}^k e_k, \quad e_i \perp e_j = \sum_{k=1}^n \phi_{ij}^k e_k, \quad \alpha(e_i) = \sum_{j=1}^n a_{ji} e_j.$$

The axioms in Definition 2.1 are, respectively, equivalent to

$$\sum_{p=1}^n \sum_{q=1}^n \gamma_{ij}^p a_{qk} \gamma_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n a_{pi} \gamma_{jk}^q \gamma_{pq}^r, \quad (2.10)$$

$$\sum_{p=1}^n \sum_{q=1}^n \gamma_{ij}^p a_{qk} \gamma_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n a_{pi} \delta_{jk}^q \gamma_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n a_{pi} \phi_{jk}^q \gamma_{pq}^r, \quad (2.11)$$

$$\sum_{p=1}^n \sum_{q=1}^n \delta_{ij}^p a_{qk} \gamma_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n a_{pi} \gamma_{jk}^q \delta_{pq}^r, \quad (2.12)$$

$$\sum_{p=1}^n \sum_{q=1}^n \gamma_{ij}^p a_{qk} \delta_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n a_{pi} \gamma_{jk}^q \delta_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n \phi_{ij}^p a_{qk} \delta_{pq}^r, \quad (2.13)$$

$$\sum_{p=1}^n \sum_{q=1}^n \delta_{ij}^p a_{qk} \delta_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n a_{pi} \delta_{jk}^q \delta_{pq}^r, \quad \sum_{p=1}^n \sum_{q=1}^n \phi_{ij}^p a_{qk} \gamma_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n a_{pi} \gamma_{jk}^q \phi_{pq}^r, \quad (2.14)$$

$$\sum_{p=1}^n \sum_{q=1}^n \gamma_{ij}^p a_{qk} \phi_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n a_{pi} \delta_{jk}^q \phi_{pq}^r, \quad \sum_{p=1}^n \sum_{q=1}^n \delta_{ij}^p a_{qk} \phi_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n a_{pi} \phi_{jk}^q \delta_{pq}^r, \quad (2.15)$$

$$\sum_{p=1}^n \sum_{q=1}^n \phi_{ij}^p a_{qk} \phi_{pq}^r = \sum_{p=1}^n \sum_{q=1}^n a_{pi} \phi_{jk}^q \phi_{pq}^r. \quad (2.16)$$

where $(a_{ij})_{1 \leq i, j \leq n}$ refers to the matrix of α and (γ_{ij}^k) , (δ_{ij}^k) and (ξ_{ij}^k) stand for the structure constants of \mathcal{A} .

The axioms in Remark 2.2 are, respectively, equivalent to

$$\sum_{k=1}^n \gamma_{ij}^k a_{qk} = \sum_{k=1}^n \sum_{p=1}^n a_{ki} a_{pj} \gamma_{kp}^q, \quad \sum_{k=1}^n \delta_{ij}^k a_{qk} = \sum_{k=1}^n \sum_{p=1}^n a_{ki} a_{pj} \delta_{kp}^q, \quad \sum_{k=1}^n \xi_{ij}^k a_{qk} = \sum_{k=1}^n \sum_{p=1}^n a_{ki} a_{pj} \xi_{kp}^q.$$

Theorem 2.14. *Any 2-dimensional complex Hom-associative trialgebra is either associative or isomorphic to one of the following pairwise non-isomorphic Hom-associative trialgebras:*

$$\mathcal{TH}_2^1: \quad \begin{array}{lllll} e_1 + e_2 = e_1, & e_2 + e_2 = e_1, & e_2 \vdash e_1 = e_1, & e_1 \perp e_2 = e_1, & e_2 \perp e_2 = e_1, \\ e_2 + e_1 = e_1, & e_1 \vdash e_2 = e_1, & e_2 \vdash e_2 = e_1, & e_2 \perp e_1 = e_1, & \alpha(e_2) = e_1. \end{array}$$

$$\mathcal{TH}_2^2: \quad \begin{array}{llll} e_1 + e_2 = e_1, & e_2 + e_2 = e_1, & e_2 \vdash e_1 = e_1, & e_2 \perp e_2 = e_1, \\ e_2 + e_1 = e_1, & e_1 \vdash e_2 = e_1, & e_2 \vdash e_2 = e_1, & \alpha(e_2) = e_1. \end{array}$$

$$\mathcal{TH}_2^3: \quad \begin{array}{llll} e_1 + e_2 = e_1, & e_2 + e_2 = e_1, & e_2 \vdash e_1 = e_1, & \alpha(e_2) = e_1, \\ e_2 + e_1 = e_1, & e_1 \vdash e_2 = e_1, & e_2 \perp e_2 = e_1, & \end{array}$$

$$\mathcal{TH}_2^4: \quad \begin{array}{llll} e_2 + e_2 = e_1, & e_2 \perp e_2 = e_1, & \alpha(e_2) = e_1 + e_2, \\ e_2 \vdash e_2 = e_1, & \alpha(e_1) = e_1, & \end{array}$$

$$\begin{array}{l}
\mathcal{TH}_2^5 : \quad e_2 \dashv e_1 = e_1, \quad e_2 \vdash e_1 = e_1, \quad e_2 \perp e_1 = e_1 \quad \alpha(e_1) = e_1, \\
\quad \quad \quad e_2 \dashv e_2 = e_1 + e_2, \quad e_2 \vdash e_2 = e_1 + e_2, \quad e_2 \perp e_2 = e_1 + e_2, \quad \alpha(e_2) = e_1 + e_2. \\
\mathcal{TH}_2^6 : \quad e_1 \dashv e_2 = e_1, \quad e_1 \vdash e_2 = e_1, \quad e_1 \perp e_2 = e_1 \quad \alpha(e_1) = e_1, \\
\quad \quad \quad e_2 \dashv e_2 = e_1 + e_2, \quad e_2 \vdash e_2 = e_1 + e_2, \quad e_2 \perp e_2 = e_1 + e_2, \quad \alpha(e_2) = e_1 + e_2. \\
\mathcal{TH}_2^7 : \quad e_1 \dashv e_2 = e_1, \quad e_2 \vdash e_1 = e_1, \quad e_1 \perp e_1 = e_1 \quad e_2 \perp e_1 = e_1 \quad \alpha(e_1) = e_1, \\
\quad \quad \quad e_2 \dashv e_2 = e_2, \quad e_2 \vdash e_2 = e_2, \quad e_1 \perp e_2 = e_1 \quad e_2 \perp e_2 = e_2, \quad \alpha(e_2) = e_2. \\
\mathcal{TH}_2^8 : \quad e_1 \dashv e_2 = e_1, \quad e_2 \vdash e_1 = e_1, \quad e_1 \perp e_2 = e_1, \quad e_2 \perp e_2 = e_2, \quad \alpha(e_1) = e_1, \\
\quad \quad \quad e_2 \dashv e_2 = e_2, \quad e_2 \vdash e_2 = e_2, \quad e_2 \perp e_1 = e_1, \quad \alpha(e_2) = e_2. \\
\mathcal{TH}_2^9 : \quad e_1 \dashv e_1 = -e_1, \quad e_1 \vdash e_1 = -e_1, \quad e_1 \perp e_1 = -e_1, \quad \alpha(e_1) = e_1, \\
\quad \quad \quad e_1 \dashv e_2 = e_2, \quad e_1 \vdash e_2 = e_2, \quad e_1 \perp e_2 = e_1, \quad \alpha(e_2) = -e_2. \\
\quad \quad \quad e_2 \dashv e_1 = e_2, \quad e_2 \vdash e_1 = e_2, \quad e_2 \perp e_2 = e_2, \\
\mathcal{TH}_2^{10} : \quad e_1 \dashv e_1 = -e_1, \quad e_1 \vdash e_2 = e_2, \quad e_2 \perp e_1 = e_2, \\
\quad \quad \quad e_1 \dashv e_2 = e_2, \quad e_1 \perp e_1 = -e_1, \quad \alpha(e_1) = e_1, \\
\quad \quad \quad e_1 \vdash e_1 = -e_1, \quad e_1 \perp e_2 = e_2, \quad \alpha(e_2) = -e_2. \\
\mathcal{TH}_2^{11} : \quad e_1 \dashv e_1 = e_1, \quad e_1 \vdash e_1 = e_1, \quad e_1 \perp e_1 = e_1, \quad \alpha(e_2) = e_2. \\
\quad \quad \quad e_2 \dashv e_2 = e_2, \quad e_2 \vdash e_2 = e_2, \quad e_2 \perp e_2 = e_2, \\
\mathcal{TH}_2^{12} : \quad e_2 \dashv e_2 = e_2, \quad e_1 \perp e_1 = e_1, \quad \alpha(e_1) = e_1. \\
\quad \quad \quad e_2 \vdash e_2 = e_2, \quad e_2 \perp e_2 = e_2, \\
\mathcal{TH}_2^{13} : \quad e_1 \dashv e_2 = e_1, \quad e_2 \dashv e_2 = e_2, \quad e_2 \perp e_1 = e_1, \quad \alpha(e_1) = e_1, \\
\quad \quad \quad e_2 \dashv e_2 = e_2, \quad e_1 \perp e_1 = e_1, \quad e_2 \perp e_2 = e_1 + e_2, \quad \alpha(e_2) = e_2. \\
\quad \quad \quad e_2 \vdash e_1 = e_1, \quad e_1 \perp e_2 = e_1,
\end{array}$$

Proof. Let \mathcal{A} be a two-dimensional vector space. To determine a Hom-associative trialgebra structure on \mathcal{A} , we consider \mathcal{A} with respect to one Hom-associative trialgebra operation. Let $\mathcal{H}_{\mathcal{T}_2} = (\mathcal{A}, \vdash, \alpha)$ be the Hom-algebra

$$e_1 \vdash e_1 = -e_1, \quad e_1 \vdash e_2 = e_2, \quad e_2 \vdash e_1 = e_2, \quad \alpha(e_1) = e_1, \quad \alpha(e_2) = -e_2.$$

The multiplication operations \dashv, \perp in \mathcal{A} , we define as follows:

$$\begin{array}{ll}
e_1 \dashv e_1 = a_1 e_1 + a_2 e_2, & e_1 \perp e_1 = b_1 e_1 + b_2 e_2, \\
e_1 \dashv e_2 = a_3 e_1 + a_4 e_2, & e_1 \perp e_2 = b_3 e_1 + b_4 e_2, \\
e_2 \dashv e_1 = a_5 e_1 + a_6 e_2, & e_2 \perp e_1 = b_5 e_1 + b_6 e_2, \\
e_2 \dashv e_2 = a_7 e_1 + a_8 e_2, & e_2 \perp e_2 = b_7 e_1 + b_8 e_2.
\end{array}$$

Now verifying Hom-associative trialgebra axioms, we get several constraints for the coefficients $a_i, b_i \in \mathbb{R}$ where $1 \leq i \leq 8$.

Applying $(e_1 \dashv e_1) \vdash \alpha(e_1) = \alpha(e_1) \vdash (e_1 \dashv e_1)$, we get $(a_1 e_1 + a_2 e_2) \vdash e_1 = e_1 \vdash (e_1 \dashv e_1)$ and then $e_1 \vdash e_1 = 1$. Therefore $a_1 = -1$.

The verification of $(e_1 \dashv e_1) \dashv \alpha(e_1) = \alpha(e_1) \dashv (e_1 \dashv e_1)$ leads to $e_1 \dashv e_1 = e_1 \dashv (e_1 + a_2 e_2)$ and from this we get $e_1 + a_2 e_2 = e_1$. Hence we obtain $a_2 = 0$.

Consider $(e_1 \dashv e_1) \dashv \alpha(e_1) = \alpha(e_1) \dashv (e_1 \dashv e_1)$. It implies that $e_1 \dashv e_2 = 1$, therefore $a_3 = 1$ and $a_4 = 0$. The next relation to consider is $(e_1 \dashv e_2) \dashv \alpha(e_1) = \alpha(e_1) \dashv (e_2 \dashv e_1)$. It implies that $1 = e_1 \dashv (a_5 e_1 + a_6 e_2)$ and we get $a_5 = 0$ and $a_6 = 1$. To find a_7 and a_8 , we note that $(e_2 \dashv e_2) \dashv \alpha(e_1) = \alpha(e_2) \dashv (e_2 \dashv e_1) \Rightarrow (a_7 e_1 + a_8 e_2) \dashv e_1 = 0 \Rightarrow a_7 e_1 + a_6 a_8 e_2 = 0$. Hence we have $a_7 = 0, a_6 a_8 = 0$. Finally, we apply $(e_2 \dashv e_2) \dashv \alpha(e_2) = \alpha(e_2) \dashv (e_2 \dashv e_2) \Rightarrow a_8 (e_2 \dashv e_2) = 0$, and get $a_8 = 0$.

Applying $(e_1 \perp e_1) \vdash \beta(e_1) = \alpha(e_1) \vdash (e_1 \perp e_1)$, we get $(y_1 e_1 + y_2 e_2) \vdash e_1 = e_1 \vdash (e_1 \perp e_1)$ and then $e_1 \vdash e_1 = 0$. Therefore $y_1 = 0$ and $y_2 = 0$. Consider $(e_1 \perp e_2) \vdash \beta(e_1) = \alpha(e_1) \perp (e_2 \dashv e_1)$. It implies that

$e_1 \perp e_2 = 0$, therefore $y_3 = 0$ and $y_4 = 0$.

The next relation to consider is $(e_1 \perp e_2) \dashv \alpha(e_1) = \alpha(e_2) \perp (e_2 \dashv e_1)$.

It implies that $0 = e_2 \perp (b_5 e_1 + b_6 e_2)$ and we get $b_5 = 0$ and $b_6 = 0$. To find b_7 and b_8 we note that $(e_2 \perp e_2) \dashv \alpha(e_1) = \alpha(e_2) \dashv (e_2 \dashv e_1) \Rightarrow (b_7 e_1 + b_8 e_2) \dashv e_1 = 1 \Rightarrow b_7 e_1 + b_6 b_8 e_2 = 1$. Hence we have $b_7 = 0, b_6 b_8 = 0$. Finally, we apply $(e_2 \perp e_2) \dashv \alpha(e_2) = \alpha(e_2) \dashv (e_2 \dashv e_2) \Rightarrow b_8 (e_2 \dashv e_2) = 1$, and get $b_8 = 1$. The verification of all other cases leads to the obtained constraints. Thus, in this case we come to the Hom-associative trialgebra with the multiplication table:

$$\begin{array}{llll} e_1 \dashv e_1 = -e_1, & e_1 \dashv e_1 = -e_1, & e_1 \perp e_1 = -e_1, & \alpha(e_1) = e_1, \\ e_1 \dashv e_2 = e_2, & e_1 \dashv e_2 = e_2, & e_1 \perp e_2 = e_1, & \alpha(e_2) = -e_2, \\ e_2 \dashv e_1 = e_2, & e_2 \dashv e_1 = e_2, & e_2 \perp e_2 = e_2, & \end{array}$$

Then $\mathcal{H}_{\mathcal{T}2} = (\mathcal{A}, \dashv, \alpha)$ it is isomorphic to $\mathcal{H}_{\mathcal{T}2}^9$. The other Hom-associative trialgebras of the list of Theorem 2.14 can be obtained by minor modification of the observation above. \square

Theorem 2.15. *Any 3-dimensional complex Hom-associative trialgebra is either associative or isomorphic to one of the following pairwise non-isomorphic Hom-associative trialgebras:*

$$\begin{array}{llll} \mathcal{TH}_3^1: & e_2 \dashv e_2 = e_2 + e_3, & e_2 \dashv e_2 = e_2 + e_3, & e_2 \perp e_2 = e_2 + e_3, & e_3 \perp e_3 = e_2 + e_3, \\ & e_2 \dashv e_3 = e_2 + e_3, & e_3 \dashv e_2 = e_2 + e_3, & e_2 \perp e_3 = e_2 + e_3, & \alpha(e_1) = e_1. \\ & e_3 \dashv e_2 = e_2 + e_3, & e_3 \dashv e_3 = e_2 + e_3, & e_3 \perp e_2 = e_2 + e_3, & \\ \mathcal{TH}_3^2: & e_2 \dashv e_2 = e_2 + e_3, & e_2 \dashv e_2 = e_2 + e_3, & e_2 \perp e_3 = e_2 + e_3, & \alpha(e_1) = e_1. \\ & e_3 \dashv e_2 = e_2 + e_3, & e_2 \dashv e_3 = e_2 + e_3, & e_3 \perp e_2 = e_2 + e_3, & \\ & e_3 \dashv e_3 = e_2 + e_3, & e_3 \dashv e_3 = e_2 + e_3, & e_3 \perp e_3 = e_2 + e_3, & \\ \mathcal{TH}_3^3: & e_1 \dashv e_1 = ae_2, & e_1 \dashv e_1 = e_2, & e_1 \perp e_1 = e_2, & \alpha(e_1) = e_1, \\ & e_3 \dashv e_3 = e_3, & e_3 \dashv e_3 = e_2, & e_3 \perp e_3 = e_2, & \alpha(e_2) = e_2. \\ \mathcal{TH}_3^4: & e_1 \dashv e_1 = e_3, & e_1 \dashv e_1 = e_3, & e_1 \perp e_1 = e_3, & \alpha(e_1) = e_1, \\ & e_2 \dashv e_2 = e_2, & e_2 \dashv e_2 = e_2, & e_2 \perp e_2 = e_2, & \alpha(e_3) = e_3. \\ \mathcal{TH}_3^5: & e_1 \dashv e_1 = ae_1, & e_1 \dashv e_1 = e_1, & e_1 \perp e_1 = e_1, & \alpha(e_2) = e_2, \\ & e_2 \dashv e_2 = e_2, & e_2 \dashv e_2 = e_2, & e_2 \perp e_2 = e_2, & \alpha(e_3) = e_3, \\ & e_3 \dashv e_2 = e_3, & e_3 \dashv e_2 = e_3, & e_3 \perp e_2 = e_3, & \\ \mathcal{TH}_3^6: & e_1 \dashv e_1 = e_1 + e_3, & e_1 \dashv e_1 = e_1 + e_3, & e_1 \perp e_1 = e_1 + e_3, & \alpha(e_2) = e_2. \\ & e_1 \dashv e_3 = e_1 + e_3, & e_1 \dashv e_3 = e_1 + e_3, & e_1 \perp e_3 = e_1 + e_3, & \\ & e_3 \dashv e_1 = e_1 + e_3, & e_3 \dashv e_1 = e_1 + e_3, & e_3 \perp e_1 = e_1 + e_3, & \\ \mathcal{TH}_3^7: & e_1 \dashv e_1 = e_1 + e_2, & e_1 \dashv e_1 = e_1 + e_2, & e_2 \dashv e_2 = e_1 + e_2, & \alpha(e_3) = e_3. \\ & e_1 \dashv e_2 = e_1 + e_2, & e_1 \dashv e_2 = e_1 + e_2, & e_2 \perp e_1 = e_1 + e_2, & \\ & e_2 \dashv e_2 = e_1 + e_2, & e_2 \dashv e_1 = e_1 + e_2, & e_2 \perp e_2 = e_1 + e_2, & \\ \mathcal{TH}_3^8: & e_1 \dashv e_2 = e_1 + e_2, & e_2 \dashv e_1 = e_1 + e_2, & e_1 \perp e_2 = e_1 + e_2, & \alpha(e_3) = e_3. \\ & e_2 \dashv e_1 = e_1 + e_2, & e_2 \dashv e_2 = e_1 + e_2, & e_2 \perp e_2 = e_1 + e_2, & \\ & e_2 \dashv e_2 = e_1 + e_2, & e_1 \perp e_1 = e_1 + e_2, & & \\ \mathcal{TH}_3^9: & e_2 \dashv e_2 = ae_2 - be_3, & e_2 \perp e_2 = be_2 + e_3, & \alpha(e_2) = e_1 + e_2, & \\ & e_2 \dashv e_2 = e_2 + de_3, & \alpha(e_1) = e_1, & \alpha(e_3) = e_2 + e_3. & \\ \mathcal{TH}_3^{10}: & e_1 \dashv e_2 = e_1, & e_2 \dashv e_1 = e_1, & e_2 \perp e_2 = e_1, & \alpha(e_2) = e_1, \\ & e_2 \dashv e_1 = e_1, & e_2 \dashv e_2 = e_1, & e_2 \perp e_3 = e_1, & \alpha(e_3) = e_3. \\ & e_2 \dashv e_2 = e_1, & e_2 \dashv e_3 = e_1, & e_3 \perp e_2 = e_1, & \\ \mathcal{TH}_3^{11}: & e_2 \dashv e_1 = e_1, & e_1 \dashv e_2 = e_1, & e_2 \perp e_2 = e_1, & \alpha(e_2) = e_1, \\ & e_2 \dashv e_2 = e_1, & e_2 \dashv e_1 = e_1, & e_2 \perp e_2 = e_1, & \alpha(e_3) = e_3. \\ & e_3 \dashv e_2 = e_1, & e_3 \dashv e_2 = e_1, & e_3 \perp e_2 = e_1, & \end{array}$$

$$\begin{array}{l}
\mathcal{TH}_3^{12} : \begin{array}{lll} e_2 \vdash e_1 = e_1 + e_3, & e_1 \vdash e_2 = e_1 + e_3, & e_2 \perp e_2 = e_1 + e_3, \\ e_2 \vdash e_2 = e_1 + e_3, & e_2 \vdash e_1 = e_1 + e_3, & e_2 \perp e_3 = e_1 + e_3, \\ e_3 \vdash e_3 = e_1 + e_3, & e_2 \vdash e_3 = e_1 + e_3, & e_3 \perp e_3 = e_1 + e_3, \end{array} \quad \alpha(e_2) = e_1. \\
\mathcal{TH}_3^{13} : \begin{array}{lll} e_3 \vdash e_2 = e_1 + e_3, & e_2 \vdash e_3 = e_1 + e_3, & e_2 \perp e_2 = e_1 + e_3, \\ e_3 \vdash e_3 = e_1 + e_3, & e_3 \vdash e_2 = e_1 + e_3, & e_3 \perp e_2 = e_1 + e_3, \\ e_1 \vdash e_2 = e_1 + e_3, & e_3 \vdash e_3 = e_1 + e_3, & e_3 \perp e_3 = e_1 + e_3, \end{array} \quad \alpha(e_2) = e_1. \\
\mathcal{TH}_3^{14} : \begin{array}{lll} e_2 \vdash e_3 = e_1 + e_3, & e_2 \vdash e_2 = e_1 + e_3, & e_3 \perp e_2 = e_1 + e_3, \\ e_3 \vdash e_2 = e_1 + e_3, & e_3 \vdash e_2 = e_1 + e_3, & e_3 \perp e_3 = e_1 + e_3, \\ e_3 \vdash e_3 = e_1 + e_3, & e_3 \vdash e_3 = e_1 + e_3, & \end{array} \quad \alpha(e_2) = e_1. \\
\mathcal{TH}_3^{15} : \begin{array}{lll} e_1 \vdash e_2 = e_1 + e_3, & e_2 \vdash e_3 = e_1 + e_3, & e_2 \perp e_1 = e_3, \\ e_3 \vdash e_2 = e_1 + e_3, & e_3 \vdash e_2 = e_1 + e_3, & e_2 \perp e_3 = e_3, \\ e_3 \vdash e_3 = e_1 + e_3, & e_3 \vdash e_3 = e_1 + e_3, & e_3 \perp e_3 = e_3, \end{array} \quad \alpha(e_2) = e_1. \\
\mathcal{TH}_3^{16} : \begin{array}{lll} e_2 \vdash e_1 = e_3, & e_1 \vdash e_2 = e_1, & e_3 \perp e_2 = e_3, \\ e_2 \vdash e_2 = e_3, & e_3 \vdash e_2 = e_1 + e_3, & e_3 \perp e_3 = e_3, \\ e_3 \vdash e_3 = e_1 + e_3, & e_3 \vdash e_3 = e_1, & \end{array} \quad \alpha(e_2) = e_1. \\
\mathcal{TH}_3^{17} : \begin{array}{lll} e_1 \vdash e_2 = e_1 + e_3, & e_3 \vdash e_3 = e_1, & e_3 \perp e_2 = e_1, \\ e_2 \vdash e_1 = e_1 + e_3, & e_1 \perp e_2 = e_3, & e_3 \perp e_3 = e_3, \\ e_1 \vdash e_2 = e_1, & e_2 \perp e_1 = e_3, & \end{array} \quad \alpha(e_2) = e_1. \\
\mathcal{TH}_3^{18} : \begin{array}{lll} e_2 \vdash e_3 = e_1 + e_3, & e_2 \vdash e_1 = e_3, & e_1 \perp e_2 = e_1 + e_3, \\ e_3 \vdash e_2 = e_1 + e_3, & e_2 \vdash e_2 = e_1 + e_3, & e_2 \perp e_3 = e_1, \\ e_3 \vdash e_3 = e_1 + e_3, & e_3 \vdash e_3 = e_1 + e_3, & e_3 \perp e_2 = e_1 + e_3, \end{array} \quad \alpha(e_2) = e_1. \\
\mathcal{TH}_3^{19} : \begin{array}{lll} e_2 \vdash e_3 = e_2, & e_2 \vdash e_2 = e_2 + e_3, & e_2 \perp e_3 = e_3, \\ e_3 \vdash e_2 = e_2, & e_3 \vdash e_3 = e_2 + e_3, & e_3 \perp e_2 = e_3, \\ e_3 \vdash e_3 = e_3, & e_2 \perp e_2 = e_2 + e_3, & e_3 \perp e_3 = e_3, \end{array} \quad \alpha(e_1) = e_1. \\
\mathcal{TH}_3^{20} : \begin{array}{lll} e_1 \vdash e_3 = e_1, & e_3 \vdash e_1 = e_1, & e_1 \perp e_3 = e_1, \\ e_2 \vdash e_3 = e_1, & e_3 \vdash e_2 = e_1, & e_2 \perp e_3 = e_1, \\ e_3 \vdash e_3 = e_1, & e_3 \vdash e_3 = e_1, & e_3 \perp e_3 = e_1, \end{array} \quad \begin{array}{l} \alpha(e_2) = e_1, \\ \alpha(e_3) = e_2. \end{array} \\
\mathcal{TH}_3^{21} : \begin{array}{lll} e_2 \vdash e_3 = e_1, & e_3 \vdash e_1 = e_1, & e_2 \perp e_3 = e_1, \\ e_3 \vdash e_2 = e_1, & e_3 \vdash e_3 = e_1, & e_3 \perp e_1 = e_1, \\ e_3 \vdash e_3 = e_1, & e_1 \perp e_3 = e_1, & e_3 \perp e_2 = e_1, \end{array} \quad \begin{array}{l} \alpha(e_2) = e_1, \\ \alpha(e_3) = e_2. \end{array}
\end{array}$$

Proof. Let \mathcal{A} be a three-dimensional vector space. To determine a Hom-associative trialgebra structure on \mathcal{A} , we consider \mathcal{A} with respect to one Hom-associative trialgebra operation. Let $\mathcal{H}_{\mathcal{T}_3} = (\mathcal{A}, \vdash, \alpha)$ be the Hom-algebra

$$e_3 \vdash e_1 = e_1, \quad e_3 \vdash e_2 = e_1, \quad e_3 \vdash e_3 = e_1, \quad \alpha(e_1) = e_1, \quad \alpha(e_3) = e_2.$$

The multiplication operations \vdash, \perp in \mathcal{A} . We use the same method of the Proof of the Theorem 2.15. Then $\mathcal{H}_{\mathcal{T}_3} = (\mathcal{A}, \vdash, \alpha)$ it is isomorphic to $\mathcal{H}_{\mathcal{T}_3}^{20}$. The other Hom-associative trialgebra of the list of Theorem 2.15 can be obtained by minor modification of the observation. \square

3 α -Inner-derivations of Finite-dimensional Hom-associative trialgebras.

This section sets forward a detailed description of α -Inner-derivations of Hom-associative trialgebras in dimensions two and three over the field \mathbb{K} .

Definition 3.1. An α -Inner-derivation of the Hom-associative trialgebra \mathcal{A} is a linear transformation $I : \mathcal{A} \rightarrow \mathcal{A}$ satisfying

$$\alpha \circ I = I \circ \alpha \quad , \quad (3.1)$$

$$ad_z(x) = I(x) \dashv \alpha(z) - \alpha(x) \dashv I(z), \quad (3.2)$$

$$ad_z(x) = I(x) \vdash \alpha(z) - \alpha(x) \vdash I(z), \quad (3.3)$$

$$ad_z(x) = I(x) \perp \alpha(z) - \alpha(x) \perp I(z), \quad (3.4)$$

for all $x, y \in \mathcal{A}$.

Let $(\mathcal{A}, \dashv, \vdash, \alpha)$ be a Hom-associative trialgebra over \mathbb{K} . For $z \in \mathcal{A}$, we have

$$ad_z(X) = X \dashv z - z \vdash X, \quad (\forall, X \in \mathcal{A}).$$

We can, therefore, prove that ad_z is a α -derivation of $(\mathcal{A}, \dashv, \vdash, \alpha)$. For any $X, Y \in \mathcal{A}$, we get

$$ad_z(X \dashv Y) = (X \dashv Y) \dashv \alpha(z) - \alpha(z) \vdash (X \dashv Y)$$

and

$$\begin{aligned} ad_z(X) \dashv \alpha(Y) + \alpha(X) \dashv ad_z(Y) &= (X \dashv z - z \vdash X) \dashv \alpha(Y) + \alpha(X) \dashv (Y \dashv z - z \vdash Y) \\ &= \alpha(X) \dashv (z \dashv Y) - (z \vdash X) \dashv \alpha(Y) + \alpha(X) \dashv (Y \dashv z) - \alpha(X) \dashv (z \dashv Y) \\ &= \alpha(X) \dashv (Y \dashv z) - (z \vdash X) \dashv \alpha(Y) = \alpha(X) \dashv (Y \perp z) - (z \vdash X) \dashv \alpha(Y). \end{aligned}$$

Hence,

$$ad_z(X \dashv Y) = ad_z(X) \dashv \alpha(Y) + \alpha(X) \dashv ad_z(Y).$$

On the other side,

$$ad_z(X \vdash Y) = (X \vdash Y) \dashv \alpha(z) + \alpha(z) \vdash (X \vdash Y) = (X \vdash Y) \dashv (z) - \alpha(z) \vdash (X \perp Y)$$

and

$$\begin{aligned} ad_z(X) \vdash \alpha(Y) + \alpha(X) \vdash ad_z(Y) &= (X \dashv z - z \vdash X) \vdash \alpha(Y) + \alpha(X) \vdash (Y \dashv z - z \vdash Y) \\ &= (X \vdash z) \vdash \alpha(Y) - (z \vdash X) \vdash \alpha(Y) + \alpha(X) \vdash (Y \dashv z) - (X \vdash z) \vdash \alpha(Y) \\ &= \alpha(X) \vdash (Y \dashv z) - (z \vdash X) \vdash \alpha(Y). \end{aligned}$$

Thus, it follows that

$$ad_z(X \vdash Y) = ad_z(X) \vdash \alpha(Y) + \alpha(X) \vdash ad_z(Y).$$

Let $\{e_1, e_2, e_3, \dots, e_n\}$ be a basis of an n -dimensional Hom-associative trialgebra \mathcal{A} . The product of basis is denoted by

$$I(e_p) = \sum_{q=1}^n I_{qp} e_q.$$

We have

$$\sum_{p=1}^n \mathcal{I}_{pk} a_{qp} = \sum_{p=1}^n a_{pk} \mathcal{I}_{qp} \quad , \quad (3.5)$$

$$\sum_{k=1}^n \gamma_{ij}^p \mathcal{I}_{rp} = \sum_{p=1}^n \sum_{q=1}^n \mathcal{I}_{pi} a_{qj} \gamma_{pq}^r - \sum_{p=1}^n \sum_{q=1}^n a_{pi} \mathcal{I}_{qj} \gamma_{pq}^r, \quad (3.6)$$

$$\sum_{k=1}^n \delta_{ij}^p \mathcal{I}_{rp} = \sum_{p=1}^n \sum_{q=1}^n \mathcal{I}_{pi} a_{qj} \delta_{pq}^r - \sum_{p=1}^n \sum_{q=1}^n a_{pi} \mathcal{I}_{qj} \delta_{pq}^r, \quad (3.7)$$

$$\sum_{k=1}^n \phi_{ij}^p \mathcal{I}_{rp} = \sum_{p=1}^n \sum_{q=1}^n \mathcal{I}_{pi} a_{qj} \phi_{pq}^r - \sum_{p=1}^n \sum_{q=1}^n a_{pi} \mathcal{I}_{qj} \phi_{pq}^r. \quad (3.8)$$

Theorem 3.2. *The α -Inner-derivations of 2-dimensional Hom-associative trialgebras have the following form :*

IC	Der(\mathcal{I})	Dim(\mathcal{I})	IC	Der(\mathcal{I})	Dim(\mathcal{I})
\mathcal{TH}_2^1	$\begin{pmatrix} 0 & 0 \\ \mathcal{I}_{21} & 0 \end{pmatrix}$	1	\mathcal{TH}_2^2	$\begin{pmatrix} 0 & 0 \\ \mathcal{I}_{21} & 0 \end{pmatrix}$	1
\mathcal{TH}_2^3	$\begin{pmatrix} 0 & 0 \\ \mathcal{I}_{21} & 0 \end{pmatrix}$	1	\mathcal{TH}_2^4	$\begin{pmatrix} 0 & 0 \\ \mathcal{I}_{21} & \mathcal{I}_{22} \end{pmatrix}$	1
\mathcal{TH}_2^6	$\begin{pmatrix} 0 & 0 \\ \mathcal{I}_{21} & 0 \end{pmatrix}$	1			

Proof. Resting upon Theorem 3.2, we provide the proof only for one case to illustrate the used approach. The other cases can be handled similarly with or without modification(s). Let us consider \mathcal{TH}_2^4 . Applying the systems of equations (3.5), (3.6), (3.7) and (3.8), we get $\mathcal{I}_{11} = \mathcal{I}_{12} = 0$. Thus, the Inner-derivations of \mathcal{TH}_2^4 are expressed as follows

$\mathcal{I}_1 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, $\mathcal{I}_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ is the basis of $Der(\mathcal{I})$ and $DimDer(\mathcal{I}) = 2$. The Inner-derivations of the remaining parts of the two-dimensional associative trialgebras can be tackled in a similar manner as depicted above. \square

Theorem 3.3. *The α -Inner-derivations of 3-dimensional Hom-associative trialgebras have the following form :*

IC	Der(\mathcal{I})	Dim(\mathcal{I})	IC	Der(\mathcal{I})	Dim(\mathcal{I})
\mathcal{TH}_3^1	$\begin{pmatrix} \mathcal{I}_{11} & 0 & 0 \\ 0 & \mathcal{I}_{22} & \mathcal{I}_{23} \\ 0 & -\mathcal{I}_{22} & -\mathcal{I}_{23} \end{pmatrix}$	3	\mathcal{TH}_3^2	$\begin{pmatrix} \mathcal{I}_{11} & 0 & 0 \\ 0 & \mathcal{I}_{22} & \mathcal{I}_{23} \\ 0 & -\mathcal{I}_{22} & -\mathcal{I}_{23} \end{pmatrix}$	3
\mathcal{TH}_3^3	$\begin{pmatrix} \mathcal{I}_{11} & \mathcal{I}_{21} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \mathcal{I}_{33} \end{pmatrix}$	3	\mathcal{TH}_3^4	$\begin{pmatrix} \mathcal{I}_{11} & 0 & \mathcal{I}_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	2
\mathcal{TH}_3^5	$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \mathcal{I}_{23} \\ 0 & 0 & \mathcal{I}_{33} \end{pmatrix}$	2	\mathcal{TH}_3^6	$\begin{pmatrix} \mathcal{I}_{11} & 0 & \mathcal{I}_{13} \\ 0 & \mathcal{I}_{22} & 0 \\ -\mathcal{I}_{11} & 0 & -\mathcal{I}_{13} \end{pmatrix}$	3
\mathcal{TH}_3^7	$\begin{pmatrix} \mathcal{I}_{11} & \mathcal{I}_{12} & 0 \\ -\mathcal{I}_{11} & -\mathcal{I}_{12} & 0 \\ 0 & 0 & \mathcal{I}_{33} \end{pmatrix}$	3	\mathcal{TH}_3^8	$\begin{pmatrix} \mathcal{I}_{11} & \mathcal{I}_{12} & 0 \\ -\mathcal{I}_{11} & -\mathcal{I}_{12} & 0 \\ 0 & 0 & \mathcal{I}_{33} \end{pmatrix}$	3

IC	$Der(\mathcal{I})$	$Dim(\mathcal{I})$	IC	$Der(\mathcal{I})$	$Dim(\mathcal{I})$
\mathcal{TH}_3^{10}	$\begin{pmatrix} 0 & 0 & 0 \\ I_{21} & 0 & 0 \\ 0 & 0 & I_{33} \end{pmatrix}$	2	\mathcal{TH}_3^{11}	$\begin{pmatrix} 0 & 0 & 0 \\ I_{21} & 0 & 0 \\ 0 & 0 & I_{33} \end{pmatrix}$	2
\mathcal{TH}_3^{12}	$\begin{pmatrix} 0 & 0 & 0 \\ I_{21} & I_{23} & 0 \\ 0 & 0 & 0 \end{pmatrix}$	2	\mathcal{TH}_3^{13}	$\begin{pmatrix} 0 & 0 & 0 \\ I_{21} & I_{23} & 0 \\ 0 & 0 & 0 \end{pmatrix}$	2
\mathcal{TH}_3^{14}	$\begin{pmatrix} 0 & 0 & 0 \\ I_{21} & I_{23} & 0 \\ 0 & 0 & 0 \end{pmatrix}$	3	\mathcal{TH}_3^{15}	$\begin{pmatrix} 0 & 0 & 0 \\ I_{21} & I_{23} & 0 \\ 0 & 0 & 0 \end{pmatrix}$	2
\mathcal{TH}_3^{16}	$\begin{pmatrix} 0 & 0 & 0 \\ I_{21} & I_{23} & 0 \\ 0 & 0 & 0 \end{pmatrix}$	2	\mathcal{TH}_3^{17}	$\begin{pmatrix} 0 & 0 & 0 \\ I_{21} & I_{23} & 0 \\ 0 & 0 & 0 \end{pmatrix}$	2
\mathcal{TH}_3^{18}	$\begin{pmatrix} 0 & 0 & 0 \\ I_{21} & I_{23} & 0 \\ 0 & 0 & 0 \end{pmatrix}$	2	\mathcal{TH}_3^{19}	$\begin{pmatrix} I_{21} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	1

Proof. Departing from Theorem 3.3, we provide the proof only for one case to illustrate the used approach, the other cases can be addressed similarly with or without modification(s). Let's consider $Trias_3^1$. Applying the systems of equations (3.5), (3.6), (3.7) and (3.8), we get $I_{21} = I_{31} = I_{21} = I_{31} = 0$. Hence, the derivations of $Trias_3^1$ are indicated as follows

$$I_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad I_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 0 \end{pmatrix} \quad \text{and} \quad I_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & -1 \end{pmatrix}$$

is the basis of $Der(\mathcal{I})$ and $DimDer(\mathcal{I}) = 3$. The centroids of the remaining parts of dimension three associative trialgebras can be handled in a similar manner as illustrated above. \square

Corollary 3.4.

- The dimensions of the α -Inner-derivations of 2-dimensional Hom-associative trialgebras range between zero and one.
- The dimensions of the α -Inner-derivations of 3-dimensional Hom-associative trialgebras range between zero and three.

4 Centroids of low-dimensional Hom-associative trialgebras.

4.1 Properties of centroids Hom-associative trialgebras.

In this section, we draw the following results on properties of centroids of Hom-associative trialgebras \mathcal{A} .

Definition 4.1. Let $(\mathcal{A}, +, \cdot, \alpha)$ be a Hom-associative trialgebra. A linear map $\psi : \mathcal{A} \rightarrow \mathcal{A}$ is called an element of (α) -centroids on \mathcal{A} if, for all $x, y \in \mathcal{A}$,

$$\alpha \circ \psi = \psi \circ \alpha, \quad (4.1)$$

$$\psi(x) + \alpha(y) = \psi(x) + \psi(y) = \alpha(x) + \psi(y), \quad (4.2)$$

$$\psi(x) \cdot \alpha(y) = \psi(x) \cdot \psi(y) = \alpha(x) \cdot \psi(y), \quad (4.3)$$

$$\psi(x) \perp \alpha(y) = \psi(x) \perp \psi(y) = \alpha(x) \perp \psi(y). \quad (4.4)$$

The set of all elements of (α) -**centroid** of \mathcal{A} is denoted $Cent_{(\alpha)}(\mathcal{A})$. The **centroid** of \mathcal{A} is denoted $Cent(\mathcal{A})$.

Definition 4.2. Let \mathcal{H} be a nonempty subset of \mathcal{A} . The subset

$$Z_{\mathcal{A}}(\mathcal{H}) = \{x \in \mathcal{H} | \alpha(x) \bullet \mathcal{H} = \mathcal{H} \bullet \alpha(x) = 0\}, \quad (4.5)$$

is said to be centralizer of \mathcal{H} in \mathcal{A} , where the \bullet is \dashv and \vdash , respectively.

Definition 4.3. Let $\psi \in End(\mathcal{A})$. If $\psi(\mathcal{A}) \subseteq Z(\mathcal{A})$ and $\psi(\mathcal{A}^2) = 0$, then ψ is called a central derivation. The set of all central derivations of \mathcal{A} is denoted by $C(\mathcal{A})$.

Proposition 4.4. Consider $(\mathcal{A}, \dashv, \vdash, \alpha)$ a Hom-associative trialgebra. Then,

- i) $\Gamma(\mathcal{A})Der(\mathcal{T}) \subseteq Der(\mathcal{A})$.
- ii) $[\Gamma(\mathcal{A}), Dr(\mathcal{A})] \subseteq \Gamma(\mathcal{A})$.
- iii) $[\Gamma(\mathcal{A}), \Gamma(\mathcal{A})](\mathcal{A}) \subseteq \Gamma(\mathcal{A})$ and $[\Gamma(\mathcal{A}), \Gamma(\mathcal{A})](\mathcal{A}^2) = 0$.

Proof. The proof of parts i) – iii) is straightforward with reference to definitions of derivation and centroid. \square

Proposition 4.5. Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-associative trialgebra and $\varphi \in Cent(\mathcal{A})$, $d \in Der(\mathcal{A})$. Then, $\varphi \circ d$ is an α -derivation of \mathcal{A} .

Proof. Indeed, if $x, y \in \mathcal{A}$, then

$$\begin{aligned} (\varphi \circ d)(x \bullet y) &= \varphi(d(x) \bullet \alpha(y) + \alpha(x) \bullet d(y)) \\ &= \varphi(d(x) \bullet y) + \varphi(\alpha(x) \bullet d(y)) = (\varphi \circ d)(x) \bullet \alpha(y) + \alpha(x) \bullet (\varphi \circ d)(y), \end{aligned}$$

where \bullet is \dashv, \vdash and \perp , respectively. \square

Proposition 4.6. Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-associative trialgebra over a field \mathbb{F} . Hence, $C(\mathcal{A}) = Cent(\mathcal{A}) \cap Der(\mathcal{A})$.

Proof. If $\psi \in Cent(\mathcal{A}) \cap Der(\mathcal{A})$, then by definition of $Cent(\mathcal{A})$ and $Der(\mathcal{A})$, we get

$\psi(x \bullet y) = \psi(x) \bullet \alpha(y) + \alpha(x) \bullet \psi(y)$ and $\psi(x \bullet y) = \psi(x) \circ \alpha(y) = \alpha(x) \circ \psi(y)$ for $x, y \in \mathcal{A}$. This yields $\psi(\mathcal{A}\mathcal{A}) = 0$ and $\psi(\mathcal{A}) \subseteq Z(\mathcal{A})$, i.e., $Cent(\mathcal{A}) \cap Der(\mathcal{A}) \subseteq Cent(\mathcal{A})$. The opposite is obvious since $C(\mathcal{A})$ is in both $Cent(\mathcal{A})$ and $Der(\mathcal{A})$, where \bullet is \dashv, \vdash and \perp , respectively. \square

Proposition 4.7. Let $(\mathcal{A}, \dashv, \vdash, \perp, \alpha)$ be a Hom-associative trialgebra. Therefore, for any $d \in Der(\mathcal{A})$ and $\varphi \in Cent(\mathcal{A})$, we have

- (i) The composition $d \circ \varphi$ is in $Cent(\mathcal{A})$, if and only if $\varphi \circ d$ is a central α -derivation of \mathcal{A} .
- (ii) The composition $d \circ \varphi$ is a α -derivation of \mathcal{A} , if and only if $[d, \varphi]$ is a central α -derivation of \mathcal{A} .

Proof. i) For any $\varphi \in Cent(\mathcal{A})$, $d \in Der(\mathcal{A})$, $\forall x, y \in \mathcal{A}$, we have

$$\begin{aligned} d \circ \varphi(x \bullet y) &= d \circ \varphi(x) \bullet y = d \circ \varphi(x) \bullet y + \varphi(x) \bullet d(y) \\ &= d \circ \varphi(x) \bullet y + \varphi \circ d(x \bullet y) - \varphi \circ d(x) \bullet y. \end{aligned}$$

Thus, $(d \circ \varphi - \varphi \circ d)(x \bullet y) = (d \bullet \varphi - \varphi \circ d)(x) \bullet y$.

ii) Let $d \circ \varphi \in \text{Der}(\mathcal{A})$. Using $[d, \varphi] \in \text{Cent}(\mathcal{A})$, we get

$$[d, \varphi](x \bullet y) = ([d, \varphi](x)) \bullet \alpha(y) = \alpha(x) \bullet ([d, \varphi](y)) \quad (4.6)$$

On the other side, $[d, \varphi]d \circ \varphi - \varphi \circ d$ and $d \circ \varphi, \varphi \circ d \in \text{Der}(\mathcal{A})$. Therefore,

$$[d, \varphi](x \bullet y) = (d(\varphi \circ x)) \bullet \alpha(y) + \alpha(x) \bullet (d \circ \varphi(y)) - (\varphi \circ d(x)) \bullet \alpha(y) - \alpha(x) \bullet (\varphi \circ d(y)). \quad (4.7)$$

Relying upon (4.6) and (4.7), we get $\alpha(x) \bullet ([d, \varphi](y)) = ([d, \varphi](x)) \bullet \alpha(y) = 0$.

At this stage of analysis, let $[d, \varphi]$ be a central α -derivation of \mathcal{A} . Then,

$$\begin{aligned} d \circ \varphi(x \bullet y) &= [d \circ \varphi](x \bullet y) + (\varphi \circ d)(x \bullet y) \\ &= \varphi(\circ d(x) \bullet \alpha(y)) + \varphi(\alpha(x) \bullet d(y)) \\ &= (\varphi \circ d)(x) \bullet \alpha(y) + \alpha(x) \bullet (\varphi \circ d)(y), \end{aligned}$$

where \bullet indicates the products \dashv , \vdash and \perp , respectively. \square

This section provides pertinent details on α -derivation of Hom-associative trialgebras in dimension two and three over the field \mathbb{K} . Let $\{e_1, e_2, e_3, \dots, e_n\}$ be a basis of an n -dimensional Hom-associative trialgebra \mathcal{A} . The product of basis

$$\psi(e_p) = \sum_{q=1}^n c_{qp} e_q.$$

$$\sum_{p=1}^n c_{pi} a_{qp} = \sum_{p=1}^n a_{pi} c_{qp}, \quad (4.8)$$

$$\sum_{k=1}^n \gamma_{ij}^k c_{qk} = \sum_{k=1}^n \sum_{p=1}^n c_{ki} a_{pj} \gamma_{kp}^q \quad ; \quad \sum_{k=1}^n \gamma_{ij}^k c_{qk} = \sum_{k=1}^n \sum_{p=1}^n a_{ki} c_{pj} \gamma_{kp}^q, \quad (4.9)$$

$$\sum_{k=1}^n \delta_{ij}^k c_{qk} = \sum_{k=1}^n \sum_{p=1}^n c_{ki} a_{pj} \delta_{kp}^q \quad ; \quad \sum_{k=1}^n \delta_{ij}^k c_{qk} = \sum_{k=1}^n \sum_{p=1}^n a_{ki} c_{pj} \delta_{kp}^q, \quad (4.10)$$

$$\sum_{k=1}^n \phi_{ij}^k c_{qk} = \sum_{k=1}^n \sum_{p=1}^n c_{ki} a_{pj} \phi_{kp}^q \quad ; \quad \sum_{k=1}^n \phi_{ij}^k c_{qk} = \sum_{k=1}^n \sum_{p=1}^n a_{ki} c_{pj} \phi_{kp}^q. \quad (4.11)$$

Theorem 4.8. *The centroids of 2-dimensional complex Hom-associative trialgebras are depicted as follows :*

IC	$\text{Cent}(\mathcal{A})$	$\text{Dim}(\text{Cent}(\mathcal{A}))$	IC	$\text{Cent}(\mathcal{A})$	$\text{Dim}(\text{Cent}(\mathcal{A}))$
\mathcal{TH}_2^1	$\begin{pmatrix} 0 & 0 \\ c_{21} & 0 \end{pmatrix}$	1	\mathcal{TH}_2^2	$\begin{pmatrix} 0 & 0 \\ c_{21} & 0 \end{pmatrix}$	1
\mathcal{TH}_2^4	$\begin{pmatrix} c_{11} & 0 \\ c_{21} & d_{11} \end{pmatrix}$	2	\mathcal{TH}_2^5	$\begin{pmatrix} c_{11} & 0 \\ c_{21} & c_{11} \end{pmatrix}$	2
\mathcal{TH}_2^6	$\begin{pmatrix} c_{11} & 0 \\ c_{21} & c_{11} \end{pmatrix}$	2	\mathcal{TH}_2^7	$\begin{pmatrix} c_{11} & 0 \\ 0 & c_{11} \end{pmatrix}$	1
\mathcal{TH}_2^8	$\begin{pmatrix} c_{11} & 0 \\ 0 & c_{11} \end{pmatrix}$	1	\mathcal{TH}_2^{10}	$\begin{pmatrix} c_{11} & 0 \\ 0 & 0 \end{pmatrix}$	1
\mathcal{TH}_2^{11}	$\begin{pmatrix} 0 & 0 \\ 0 & c_{22} \end{pmatrix}$	1	\mathcal{TH}_2^{12}	$\begin{pmatrix} c_{11} & 0 \\ 0 & 0 \end{pmatrix}$	1
\mathcal{TH}_2^{13}	$\begin{pmatrix} c_{11} & 0 \\ 0 & c_{11} \end{pmatrix}$	1			

Proof. Departing from Theorem 4.8, we provide the proof only for one case to illustrate the used approach, the other cases can be addressed similarly with or without modification(s). Let's consider \mathcal{TH}_2^4 . Applying the systems of equations (4.8), (4.9), (4.10) and (4.11), we get $c_{12} = 0$ and $c_{22} = c_{11}$. Hence, the centroids of \mathcal{TH}_2^4 are indicated as follows

$c_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, $c_2 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ is the basis of $Der(c)$ and $\text{Dim}Der(c) = 2$. The centroids of the remaining parts of dimension three associative trialgebras can be handled in a similar manner as illustrated above. \square

Theorem 4.9. *The centroids of 3-dimensional complex Hom-associative trialgebras are depicted as follows :*

IC	$Cent(\mathcal{A})$	$Dim(Cent(\mathcal{A}))$	IC	$Cent(\mathcal{A})$	$Dim(Cent(\mathcal{A}))$
\mathcal{TH}_3^1	$\begin{pmatrix} c_{11} & 0 & 0 \\ 0 & c_{22} & c_{23} \\ 0 & -c_{22} & -c_{23} \end{pmatrix}$	3	\mathcal{TH}_3^2	$\begin{pmatrix} c_{11} & 0 & 0 \\ 0 & c_{22} & c_{23} \\ 0 & -c_{22} & -c_{23} \end{pmatrix}$	3
\mathcal{TH}_3^3	$\begin{pmatrix} 0 & c_{12} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	1	\mathcal{TH}_3^4	$\begin{pmatrix} c_{11} & 0 & c_{13} \\ 0 & 0 & 0 \\ 0 & 0 & c_{11} \end{pmatrix}$	2
\mathcal{TH}_3^5	$\begin{pmatrix} 0 & 0 & 0 \\ 0 & c_{22} & \\ 0 & 0 & c_{33} \end{pmatrix}$	2	\mathcal{TH}_3^6	$\begin{pmatrix} c_{11} & 0 & c_{13} \\ 0 & c_{22} & 0 \\ -c_{11} & 0 & -c_{13} \end{pmatrix}$	3
\mathcal{TH}_3^7	$\begin{pmatrix} c_{11} & c_{12} & 0 \\ -c_{11} & -c_{12} & 0 \\ 0 & 0 & c_{33} \end{pmatrix}$	3	\mathcal{TH}_3^8	$\begin{pmatrix} c_{11} & c_{12} & 0 \\ -c_{11} & -c_{12} & 0 \\ 0 & 0 & c_{33} \end{pmatrix}$	3
\mathcal{TH}_3^{10}	$\begin{pmatrix} 0 & 0 & 0 \\ c_{21} & 0 & 0 \\ 0 & 0 & c_{33} \end{pmatrix}$	2	\mathcal{TH}_3^{11}	$\begin{pmatrix} 0 & 0 & 0 \\ c_{21} & 0 & 0 \\ 0 & 0 & c_{33} \end{pmatrix}$	2
\mathcal{TH}_3^{12}	$\begin{pmatrix} 0 & 0 & 0 \\ c_{21} & c_{23} & 0 \\ 0 & 0 & 0 \end{pmatrix}$	2	\mathcal{TH}_3^{13}	$\begin{pmatrix} 0 & 0 & 0 \\ c_{21} & c_{23} & 0 \\ 0 & 0 & 0 \end{pmatrix}$	2
\mathcal{TH}_3^{14}	$\begin{pmatrix} c_{11} & 0 & 0 \\ c_{21} & c_{11} & c_{23} \\ -c_{11} & 0 & 0 \end{pmatrix}$	3	\mathcal{TH}_3^{15}	$\begin{pmatrix} 0 & 0 & 0 \\ c_{21} & 0 & c_{23} \\ 0 & 0 & 0 \end{pmatrix}$	2
\mathcal{TH}_3^{16}	$\begin{pmatrix} 0 & 0 & 0 \\ c_{21} & 0 & c_{23} \\ 0 & 0 & 0 \end{pmatrix}$	2	\mathcal{TH}_3^{17}	$\begin{pmatrix} 0 & 0 & 0 \\ c_{21} & 0 & c_{23} \\ 0 & 0 & 0 \end{pmatrix}$	2
\mathcal{TH}_3^{18}	$\begin{pmatrix} 0 & 0 & 0 \\ c_{21} & 0 & c_{23} \\ 0 & 0 & 0 \end{pmatrix}$	2	\mathcal{TH}_3^{20}	$\begin{pmatrix} 0 & 0 & 0 \\ c_{21} & 0 & 0 \\ c_{31} & c_{21} & 0 \end{pmatrix}$	2
\mathcal{TH}_3^{21}	$\begin{pmatrix} 0 & 0 & 0 \\ c_{21} & 0 & 0 \\ c_{31} & c_{21} & 0 \end{pmatrix}$	2			

Proof. Departing from Theorem 4.9, we provide the proof only for one case to illustrate the used approach, the other cases can be addressed similarly with or without modification(s). Let's consider \mathcal{TH}_3^6 . Applying the systems of equations (4.8), (4.9), (4.10) and (4.11), we get $c_{12} = c_{21} = c_{23} = c_{32} = 0$, $c_{11} = -c_{31}$ and $c_{13} = -c_{33}$. Hence, the centroids of \mathcal{TH}_3^6 are indicated as follows

$c_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}$, $c_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ and $c_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & -1 \end{pmatrix}$ is the basis of $Der(c)$ and

$\dim Der(c) = 3$. The centroids of the remaining parts of dimension three associative trialgebras can be handled in a similar manner as illustrated above. \square

Corollary 4.10.

- The dimensions of the **centroids** of two-dimensional Hom-associative trialgebras range between zero and two.
- The dimensions of the **centroids** of three-dimensional Hom-associative trialgebras range between zero and three.

References

- [1] Loday J.-L., Frabetti A., Chapoton F., Goichot F., *Dialgebras and Related operads*, lecture Notes in LMaths., 1763 Springer, Berlin, 2001.
- [2] A. Zahari and I. Bakayoko, *On BiHom-Associative dialgebras*, Open J. Math. Sci. vol(7), (2023), 96 – 117.
- [3] A. Zahari and A. Makhlouf, Structure and Classification of Hom-Associative Algebras, Acta et commentationes universitatis Tartuensis de mathematica, vol 24 (1) 2020.
- [4] I. Bakayoko and S. Silvestrov *Hom-left-symmetric color dialgebras, Hom-tridendriform color algebras and Yau's twisting generalizations*, Afrika mathematic 32, (2021), 941 – 958.
- [5] L. Philippe, *Construction of Nijenhuis operators and dendriform trialgebras*, arXiv:math/0311132v1 [math.RA] 3 nov 2003.
- [6] Makhlouf, A. and Silvestrov, S.D. *Hom-algebra structures*. J. Gen. Lie Theory Appl. 2(2), 51–64 (2008)
- [7] Bakayoko, I., Diallo, O.W. *Some generalized Hom-algebra structures*. J. Gen. Lie Theory Appl. 9, 226 (2015)
- [8] A. Majumdar, and G. Mukherjee, (2002). Deformation theory of dialgebras. K-theory, 27(1):33-60.
- [9] A. P. Pozhidaev, (2008). Dialgebras and related triple systems. Siberian Mathematical Journal, 49(4):696-708.
- [10] Rakhimov, I.S., Fiidov, M. A., *Centroids of finite dimensional associative dialgebras*, Far East Journal of Mathematical Sciences, 98(4), (2015), 427 – 443.
- [11] Basri, W., Rakhimov, I., Rikhsiboev, I., et al. (2015). Four-dimensional nilpotent dias- sociative algebras. Journal of Generalized Lie Theory and Applications, 9(1):1-7.
- [12] G. Graziani, A. Makhlouf, C. Menini and F. Panaite, *BiHom-Associative Algebras, BiHom-Lie Algebras and BiHom-Bialgebras*, Symmetry, Integrability and Geometry: Methods and Applications SIGMA 11 (2015), 086, 34 pages.

-
- [13] I. Bakayoko and M. Bangoura, *Bimodules and Rota-Baxter relations*, J. Appl. Mech. Eng 4:178, doi:10.4172/2168-9873.1000178, 2015.
- [14] I. Bakayoko and M. Bangoura, *Left-Hom-symmetric and Hom-Poisson dialgebras*, Konuralp Journal of Mathematics, 3 No.2, 42-53, 2015.
- [15] I.Rikhsiboev, I.Rakhimov and W. Basri, (2014). *Diassociative algebras and their derivations*. In Journal of Physics: Conference Series, volume 553, pages 179–199. IOP Publishing.
- [16] I. M.Rikhsiboev, I. Rakhimov, and W. Basri (2010). *Classification of 3-dimensional complex diassociative algebras*. Malaysian Journal of Mathematical Sciences, 4(2):241-254.
- [17] J.L. Loday, M.Ronco, *Trialgebras and families of polytopes*, *Homotopy Theory: Relations with Algebraic Geometry, Group Cohomology, and Algebraic K-theory*. Comte. Math. 346, 369-398 (2004)
- [18] Ahmed Zahari, *Classification of 3-dimensional BiHom-Associative and BiHom-Bialgebras*, arXiv : 1907.0080 v1 [math.RA] 28 Jun 2019.
- [19] X. Li, *Structures of multiplicative Hom-Lie algebras*, Advances in Mathematics (China), 43(6)(2014)817-823.
- [20] W. Basri, I. S. Rakhimov and I. M. Rokhsiboev, *Four-Dimension Nilpotent Diassociative algebras*, J. Generalised Lie Theory Appl. 28. doi : 10.4172/1736 – 4337.1000218.
- [21] X. Chen and W. Han, *Classification of multiplicative simple Hom-Lie algebras*, J. Lie Theory 21(4)(2015)
- [22] A. Zahari, B. Mosbahi and I. Basdouri *Classification, Derivations and Centroids of Low-dimensional Complex BiHom-Trialgebras*, arXiv:2304.06781v1 [math.RA] 13 Apr 2023.
- [23] A. Zahari, B. Mosbahi and I. Basdouri *Classification, Derivations and Centroids of Low-dimensional Real Trialgebras*, arXiv : 2304.04251v1 [math.RA] 9 Apr 2023.
- [24] Ahmed Zahari *Etude et Classification des algèbres Hom-associatives*, thèse, Université de Haute Alsace à Mulhouse, le 24 – 05 – 2017.
- [25] A. Armour, H. Chen and Y. Zhang, *Classification of 4-dimensional superalgebras*, Comm. in Algebra 37(2009), 3697-3728.
- [26] Y. Fregier, A. Gohr and S. Silvestrov, *Unital algebras of Hom-associative type and surjective or injective twistings*, J. Gen. Lie Theory Appl. Vol. 3 (4), (2009) 285–295.
- [27] D. Larsson and S. Silvestrov, *Quasi-hom-Lie algebras, central extensions and 2-cocycle-like identities*, J. Algebra 288 (2005), 321–344.
- [28] Merville D. J, *Centroids of nilpotent Lie algebras*, Comm. Algebra, 20(12), (1992), 3649 – 3682.
- [29] I. S. Rakhimov, Al-Nashri, Al-Hossain, *On Derivations of low-dimensional complex Leibniz algebras*, JR Journal of Algebra, Number Theory and Applications. Vol. 21(1), (2011), pp. 68 – 81.