

# The Minimum Norm of a Projector under Linear Interpolation on a Euclidean Ball

Mikhail Nevskii\*

May 1, 2023

## Abstract

We prove the following proposition. Under linear interpolation on a Euclidean  $n$ -dimensional ball  $B$ , an interpolation projector whose nodes coincide with the vertices of a regular simplex inscribed into the boundary sphere has the minimum  $C$ -norm. This minimum norm  $\theta_n(B) = \max\{\psi(a_n), \psi(a_n + 1)\}$ , where  $\psi(t) = \frac{2\sqrt{n}}{n+1} \left( t(n+1-t) \right)^{1/2} + \left| 1 - \frac{2t}{n+1} \right|$ ,  $0 \leq t \leq n+1$ , and  $a_n = \left\lfloor \frac{n+1}{2} - \frac{\sqrt{n+1}}{2} \right\rfloor$ . For any  $n$ ,  $\sqrt{n} \leq \theta_n(B) \leq \sqrt{n+1}$ . Moreover,  $\theta_n(B) = \sqrt{n}$  only for  $n = 1$  and  $\theta_n(B) = \sqrt{n+1}$  iff  $\sqrt{n+1}$  is an integer.

Keywords:  $n$ -dimensional ball,  $n$ -dimensional simplex, linear interpolation, projector, norm

MSC: 41A05, 52B55, 52C07

## 1 Main definitions

Further  $n \in \mathbb{N}$ . An element  $x \in \mathbb{R}^n$  is written in the form  $x = (x_1, \dots, x_n)$ . Let  $K$  be a convex body in  $\mathbb{R}^n$ , i. e., a compact convex subset of  $\mathbb{R}^n$  with nonempty interior. Denote by  $C(K)$  a space of continuous functions  $f : K \rightarrow \mathbb{R}$  with the uniform norm

$$\|f\|_{C(K)} := \max_{x \in K} |f(x)|.$$

By  $\Pi_1(\mathbb{R}^n)$  we mean a set of polynomials in  $n$  variables of degree  $\leq 1$ , i. e., a set of linear functions on  $\mathbb{R}^n$ . For  $x^{(0)} \in \mathbb{R}^n, R > 0$ , by  $B(x^{(0)}; R)$  we denote the  $n$ -dimensional Euclidean ball given by the inequality  $\|x - x^{(0)}\| \leq R$ . Here

$$\|x\| := \sqrt{\langle x, x \rangle} = \left( \sum_{i=1}^n x_i^2 \right)^{1/2}.$$

---

\*Department of Mathematics, P.G. Demidov Yaroslavl State University, Sovetskaya str., 14, Yaroslavl, 150003, Russia, mnevsk55@yandex.ru, orcid.org/0000-0002-6392-7618

By definition,  $B_n := B(0; 1)$ .

Let  $S$  be a nondegenerate simplex in  $\mathbb{R}^n$  with the vertices  $x^{(j)} = (x_1^{(j)}, \dots, x_n^{(j)})$ ,  $1 \leq j \leq n+1$ . Consider *the vertex matrix* of this simplex:

$$\mathbf{S} := \begin{pmatrix} x_1^{(1)} & \dots & x_n^{(1)} & 1 \\ x_1^{(2)} & \dots & x_n^{(2)} & 1 \\ \vdots & \vdots & \vdots & \vdots \\ x_1^{(n+1)} & \dots & x_n^{(n+1)} & 1 \end{pmatrix}.$$

Assume  $\mathbf{S}^{-1} = (l_{ij})$ . Linear polynomials  $\lambda_j(x) := l_{1j}x_1 + \dots + l_{nj}x_n + l_{n+1,j}$  have the property  $\lambda_j(x^{(k)}) = \delta_j^k$ . We call  $\lambda_j$  *the basic Lagrange polynomials corresponding to  $S$* . For an arbitrary  $x \in \mathbb{R}^n$ ,

$$x = \sum_{j=1}^{n+1} \lambda_j(x)x^{(j)}, \quad \sum_{j=1}^{n+1} \lambda_j(x) = 1.$$

This means that  $\lambda_j(x)$  are *the barycentric coordinates of  $x$*  (see, e. g., [1]– [2]).

We say that the interpolation projector  $P : C(K) \rightarrow \Pi_1(\mathbb{R}^n)$  corresponds to a simplex  $S \subset K$  if the nodes of  $P$  coincide with the vertices of  $S$ . This projector is defined by the equalities  $Pf(x^{(j)}) = f(x^{(j)})$ . The following analogue of the Lagrange interpolation formula holds:

$$Pf(x) = \sum_{j=1}^{n+1} f(x^{(j)}) \lambda_j(x). \quad (1)$$

Denote by  $\|P\|_K$  the norm of  $P$  as an operator from  $C(K)$  in  $C(K)$ . From (1), it follows that

$$\|P\|_K = \max_{x \in K} \sum_{j=1}^{n+1} |\lambda_j(x)|. \quad (2)$$

If  $K$  is a convex polytope, then maximum in (2) also can be taken only over the vertex set of  $K$ . Although, throughout the paper  $K$  is a ball or an ellipsoid.

By  $\theta_n(K)$  we denote the minimal norm of an interpolation projector  $P : C(K) \rightarrow \Pi_1(\mathbb{R}^n)$  whose nodes belong to  $K$ .

Assume  $K = B := B(x^{(0)}; R)$ . It is proved in [3] that

$$\|P\|_B = \max_{f_j = \pm 1} \left[ R \left( \sum_{i=1}^n \left( \sum_{j=1}^{n+1} f_j l_{ij} \right)^2 \right)^{1/2} + \left| \sum_{j=1}^{n+1} f_j \lambda_j(x^{(0)}) \right| \right]. \quad (3)$$

If  $S$  is a regular simplex inscribed into the ball, then  $\|P\|_B$  does not depend neither on the center  $x^{(0)}$ , nor on the radius  $R$  of the ball, nor on the choice of such a simplex. In this case,  $\|P\|_B = \max\{\psi(a_n), \psi(a_n + 1)\}$  (see [3, Theorem 2]). Here

$$\psi(t) := \frac{2\sqrt{n}}{n+1} \left( t(n+1-t) \right)^{1/2} + \left| 1 - \frac{2t}{n+1} \right|, \quad 0 \leq t \leq n+1, \quad (4)$$

$$a_n := \left[ \frac{n+1}{2} - \frac{\sqrt{n+1}}{2} \right]. \quad (5)$$

## 2 The theorem on a simplex and its minimal ellipsoid

Suppose  $S$  is a nondegenerate  $n$ -dimensional simplex,  $E$  is a minimum volume ellipsoid containing  $S$ , and  $m$  is a natural number,  $1 \leq m \leq n$ . To any set of  $m$  vertices of  $S$ , let us put in correspondence a point  $y \in E$  defined as follows. Let  $g$  be the center of gravity of the  $(m-1)$ -dimensional face of  $S$  which contains the chosen vertices, and let  $h$  be the center of gravity of the  $(n-m)$ -dimensional face which contains the rest  $n+1-m$  vertices. Then  $y$  is the intersection point of the straight line  $(gh)$  with the boundary of the ellipsoid in direction from  $g$  to  $h$ .

**Theorem 1.** *For an arbitrary nondegenerate simplex  $S \subset B_n$ , there is such a set of  $m$  vertices for which  $y \in B_n$ .*

*Proof.* Let  $x^{(j)}$  be the vertices of  $S$ . The center of gravity of the simplex and also the center of gravity of its minimal ellipsoid  $E$  are the point  $c = \frac{1}{n+1} \sum x^{(j)}$ . Denote by  $r$  the ratio of the distance between the center of gravity of a regular simplex and the center of gravity of its  $(m-1)$ -dimensional face to the radius of the circumscribed sphere. It is easy to see that

$$r = \frac{1}{m} \sqrt{m - \frac{m(m-1)}{n}}.$$

By  $N$ , we denote the number of  $(m-1)$ -dimensional faces of an  $n$ -dimensional simplex:  $N = \binom{n+1}{m}$ .

Consider any set  $J$  consisting of  $m$  indices  $j$ . The center of gravity of the  $(m-1)$ -dimensional face with vertices  $x^{(j)}, j \in J$ , is the point  $g_J = \frac{1}{m} \sum_{j \in J} x^{(j)}$ . Suppose  $y_J$  coincides with the point  $y$  for this set of vertices. Then

$$y_J = c + \frac{1}{r} (c - g_J),$$

i. e.,  $y_J = (1/r)((r+1)c - g_J)$ . Summing up over all sets  $J$ , we have

$$\begin{aligned} \sum_J \|y_J\|^2 &= \sum_J \langle y_J, y_J \rangle = \\ &= \frac{1}{r^2} \left( N \|c\|^2 (r+1)^2 + \sum_J \|g_J\|^2 - 2(r+1) \langle \sum_J g_J, c \rangle \right) = \\ &= \frac{1}{r^2} \left( N \|c\|^2 (r^2 - 1) + \sum_J \|g_J\|^2 \right). \end{aligned} \quad (6)$$

We took into account the equality  $\sum g_J = Nc$ . Further, we claim that

$$\sum_J \|g_J\|^2 = \frac{1}{mn} \binom{n}{m} \sum_{j=1}^{n+1} \|x^{(j)}\|^2 + \frac{(m-1)(n+1)}{mn} \binom{n+1}{m} \|c\|^2. \quad (7)$$

To obtain (7), let us remark the following. The value  $\sum \|g_J\|^2$  contains, for every  $j$ , exactly  $\binom{n}{m-1}$  numbers  $\|x^{(j)}\|^2$  taken with coefficient  $1/m^2$ , and exactly  $\binom{n-1}{m-2}$  numbers  $2\langle x^{(i)}, x^{(j)} \rangle$ , for each  $i \neq j$ , with the same coefficient  $1/m^2$ . (The latter ones exist for  $m \geq 2$ ; in case  $m = 1$  there are no pairwise products.) The expression  $\|c\|^2$  contains all numbers  $\|x^{(j)}\|^2$ , for any  $j$ , and numbers  $2\langle x^{(i)}, x^{(j)} \rangle$ , for  $i \neq j$ , with multiplier  $1/(n+1)^2$ . The coefficient at  $\|c\|^2$  in the right-hand part of (7) guarantees the equality of expressions with pairwise products  $\langle x^{(i)}, x^{(j)} \rangle$ ; the coefficient at  $\sum \|x^{(j)}\|^2$  is chosen so that the difference of the left-hand part of (7) and the second item in the right-hand part is equal to the first item. Since

$$N(r^2 - 1) = -\frac{(m-1)(n+1)}{mn} \binom{n+1}{m},$$

after replacing  $\sum \|g_J\|^2$  in (6) with the right-hand part of (7), we notice that items with  $\|c\|^2$  are compensated. After these transformations, we get

$$\frac{1}{N} \sum_J \|y_J\|^2 = \frac{1}{n+1} \sum_{j=1}^{n+1} \|x^{(j)}\|^2. \quad (8)$$

The inclusion  $S \subset B_n$  means that  $\|x^{(j)}\| \leq 1$ . Therefore, the mean value of numbers  $\|y_J\|^2$  is also  $\leq 1$ . Consequently, for some set  $J$  we have  $\|y_J\| \leq 1$ , i. e., this point  $y_J$  lies in  $B_n$ . Theorem 1 is proved.  $\square$

As a conjecture, Theorem 1 was formulated by the author in [4]. Also in [4], the statement of Theorem 1 in case  $m = 1$  was proved in another way.

### 3 The value $\theta_n(B_n)$

The main result of the paper concerns the minimum norm of an interpolation projector with the nodes in an  $n$ -dimensional ball.

**Theorem 2.** *Let function  $\psi(t)$  and number  $a_n$  be defined by equalities (4)–(5). For any  $n \in \mathbb{N}$ ,*

$$\theta_n(B_n) = \max\{\psi(a_n), \psi(a_n + 1)\}. \quad (9)$$

*An interpolation projector  $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$  satisfies the equality  $\|P\|_{B_n} = \theta_n(B_n)$  iff the interpolation nodes coincide with the vertices of a regular simplex inscribed into the boundary sphere.*

*Proof.* First let  $P$  be an interpolation projector corresponding to a regular inscribed simplex  $S$  and let  $\lambda_j(x)$  be the basic Lagrange polynomials for this simplex. Denote  $p_n := \|P\|_{B_n}$ . It is proved in [3] that  $p_n = \max\{\psi(a_n), \psi(a_n + 1)\}$ . The points  $y \in B_n$  satisfying

$$\sum_{j=1}^{n+1} |\lambda_j(y)| = p_n \quad (10)$$

were found in [4]. Namely, assume  $k_n$  coincides with that of the numbers  $a_n$  and  $a_n + 1$  which delivers to  $\psi(t)$  the maximum value. Let us fix  $m = k_n$ . Then equality (10) takes place for every point  $y$  which is constructed above Theorem 1 in case  $E = B_n$ . The number of these points  $y$  is  $N = \binom{n+1}{m}$ .

Now suppose  $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$  is an arbitrary projector with the nodes  $x^{(j)} \in B_n$ ,  $S$  is the simplex with the vertices  $x^{(j)}$ , and  $E$  is a minimum volume ellipsoid contained  $S$ . We can also consider  $P$  as a projector acting from  $C(E)$ . Then

$$\|P\|_E = p_n = \sum_{j=1}^{n+1} |\lambda_j(y)|$$

for  $m = k_n$  and for each of the points  $y$  noted at the beginning of the previous section (now with respect to the minimal ellipsoid  $E$ ). By Theorem 1, at least one of these points, say  $y^*$ , belongs to  $B_n$ . Therefore,

$$\|P\|_{B_n} = \max_{x \in B_n} \sum_{j=1}^{n+1} |\lambda_j(x)| \geq \sum_{j=1}^{n+1} |\lambda_j(y^*)| = p_n.$$

Thus,  $\theta_n(B_n) = p_n = \max\{\psi(a_n), \psi(a_n + 1)\} = \psi(k_n)$ .

Let us demonstrate that if  $\|P\|_{B_n} = \theta_n(B_n)$ , then the simplex with vertices at the interpolation nodes is inscribed into  $B_n$  and regular. For this simplex  $S$ , some point  $y \in E$ , constructed for  $m = k_n$ , falls on the boundary sphere — otherwise we have  $\|P\|_{B_n} > p_n = \theta_n(B_n)$ . This is due to the fact that  $\sum |\lambda_j(x)|$  increases monotonously as  $x$  moves in a straight line in direction from  $c$  to  $y$ . Since  $\|y\| = 1$ , the mean value of  $\|y_J\|$  for the rest  $y_J$  is also  $\leq 1$  (see (8)). Hence, another such point also lies in the boundary of the ball. Thus we obtain  $\|y_J\| = 1$  for all sets  $J$  consisting of  $m = k_n$  indices  $j$ . Now (8) yields  $\|x^{(j)}\| = 1$ , i. e., the simplex is inscribed into the ball. Since all the points  $y_J$  belong to the boundary sphere, the function  $\sum |\lambda_j(x)|$  has maximum upon the ball at  $N = \binom{n+1}{m}$  different points. Consequently, the simplex  $S$  is regular. This completes the proof.  $\square$

For  $1 \leq n \leq 4$ , equality (9) and characterization of minimal projectors were obtained in [3] and [4] by another methods.

**Corollary.** *Always  $\sqrt{n} \leq \theta_n(B_n) \leq \sqrt{n+1}$ . Moreover,  $\theta_n(B_n) = \sqrt{n}$  only for  $n = 1$  and  $\theta_n(B_n) = \sqrt{n+1}$  when and only when  $\sqrt{n+1}$  is an integer.*

Really, it was proved in [3] that the above relations are satisfied for values  $p_n = \max\{\psi(a_n), \psi(a_n + 1)\}$ . It remains to take into account Theorem 2.

Note that the numbers  $k_n$  from the proof of Theorem 2 increase with  $n$ , but not strictly monotonously. If  $n \geq 2$ , then  $k_n \leq n/2$ . The numbers  $a_n, k_n$ , and  $\theta_n(B_n) = \psi(k_n)$  are given in Table 1; we use the results from [3].

## 4 Acknowledgement

The author is grateful to Arseniy Akopyan for his help while proving Theorem 1.

Table 1: The numbers  $a_n, k_n$ , and  $\theta_n(B_n) = \psi(k_n)$

| $n$  | $a_n$ | $a_n + 1$ | $\psi(a_n)$                   | $\psi(a_n + 1)$                   | $k_n$ | $\theta_n(B_n) = \psi(k_n)$                      |
|------|-------|-----------|-------------------------------|-----------------------------------|-------|--|
| 1    | 0     | 1         | 1                             | 1                                 | 1     | 1  |
| 2    | 0     | 1         | 1                             | $\frac{5}{3}$                     | 1     | $\frac{5}{3} = 1.6666\dots$                      |
| 3    | 1     | 2         | 2                             | $\sqrt{3}$                        | 1     | 2  |
| 4    | 1     | 2         | $\frac{11}{5}$                | $\frac{1+4\sqrt{6}}{5}$           | 1     | $\frac{11}{5} = 2.2$                             |
| 5    | 1     | 2         | $\frac{7}{3}$                 | $\frac{1+2\sqrt{10}}{3}$          | 2     | $\frac{1+2\sqrt{10}}{3} = 2.4415\dots$           |
| 6    | 2     | 3         | $\frac{3+4\sqrt{15}}{7}$      | $\frac{1+12\sqrt{2}}{7}$          | 2     | $\frac{3+4\sqrt{15}}{7} = 2.6417\dots$           |
| 7    | 2     | 3         | $\frac{1+\sqrt{21}}{2}$       | $\frac{1+\sqrt{105}}{4}$          | 3     | $\frac{1+\sqrt{105}}{4} = 2.8117\dots$           |
| 8    | 3     | 4         | 3                             | $\frac{1+8\sqrt{10}}{9}$          | 3     | 3  |
| 9    | 3     | 4         | $\frac{2+3\sqrt{21}}{5}$      | $\frac{1+6\sqrt{6}}{5}$           | 3     | $\frac{2+3\sqrt{21}}{5} = 3.1495\dots$           |
| 10   | 3     | 4         | $\frac{5+8\sqrt{15}}{11}$     | $\frac{3+4\sqrt{70}}{11}$         | 4     | $\frac{3+4\sqrt{70}}{11} = 3.3151\dots$          |
| 11   | 4     | 5         | $\frac{1+2\sqrt{22}}{3}$      | $\frac{1+\sqrt{385}}{6}$          | 4     | $\frac{1+2\sqrt{22}}{3} = 3.4602\dots$           |
| 12   | 4     | 5         | $\frac{5+24\sqrt{3}}{13}$     | $\frac{3+8\sqrt{30}}{13}$         | 5     | $\frac{3+8\sqrt{30}}{13} = 3.6013\dots$          |
| 13   | 5     | 6         | $\frac{2+3\sqrt{65}}{7}$      | $\frac{1+4\sqrt{39}}{7}$          | 5     | $\frac{2+3\sqrt{65}}{7} = 3.7409\dots$           |
| 14   | 5     | 6         | $\frac{1+4\sqrt{7}}{3}$       | $\frac{1+4\sqrt{21}}{5}$          | 6     | $\frac{1+4\sqrt{21}}{5} = 3.8660\dots$           |
| 15   | 6     | 7         | 4                             | $\frac{1+3\sqrt{105}}{8}$         | 6     | 4  |
| 50   | 21    | 22        | $\frac{3+20\sqrt{35}}{17}$    | $\frac{7+20\sqrt{319}}{51}$       | 22    | $\frac{7+20\sqrt{319}}{51} = 7.1414\dots$        |
| 100  | 45    | 46        | $\frac{11+120\sqrt{70}}{101}$ | $\frac{9+20\sqrt{2530}}{101}$     | 45    | $\frac{11+120\sqrt{70}}{101} = 10.0494\dots$     |
| 1000 | 484   | 485       | $\frac{3+40\sqrt{5170}}{91}$  | $\frac{31+200\sqrt{25026}}{1001}$ | 485   | $\frac{31+200\sqrt{25026}}{1001} = 31.6385\dots$ |

## References

- [1] M. V. NEVSKII. On a property of  $n$ -dimensional simplices, *Mathematical Notes*, **87**:4 (2010), 543–555.
- [2] M. V. NEVSKII. *Geometricheskie Ocenki v Polinomialnoi Interpolyacii* (Geometric Estimates in Polynomial Interpolation), Yaroslavl': Yarosl. Gos. Univ., 2012 (in Russian).
- [3] M. V. NEVSKII AND A. YU. UKHALOV. Linear interpolation on a Euclidean ball in  $\mathbb{R}^n$ , *Modeling and Analysis of Information Systems*, **26**:2 (2019), 279–296.
- [4] M. V. NEVSKII. On properties of a regular simplex inscribed into a ball, *Modeling and Analysis of Information Systems*, **28**:2 (2021), 186–197.