

ANOMALY CANCELLATION FORMULAS AND E_8 BUNDLES

YONG WANG, YUCHEN YANG

ABSTRACT. Using E_8 bundles, we construct some modular forms over $SL(2, \mathbf{Z})$, $\Gamma^0(2)$ and $\Gamma_0(2)$. By these modular forms, we get some new anomaly cancellation formulas of characteristic forms.

1. INTRODUCTION

In 1983, the physicists Alvarez-Gaumé and Witten [1] discovered the "miraculous cancellation" formula for gravitational anomaly which reveals a beautiful relation between the top components of the Hirzebruch \widehat{L} -form and \widehat{A} -form of a 12-dimensional smooth Riemannian manifold. Kefeng Liu [11] established higher dimensional "miraculous cancellation" formulas for $(8k+4)$ -dimensional Riemannian manifolds by developing modular invariance properties of characteristic forms. These formulas could be used to deduce some divisibility results. In [9], [10], [3], some more general cancellation formulas that involve a complex line bundle and their applications were established. In [7], using the Eisenstein series, a more general cancellation formula was derived. In [8], Han, Liu and Zhang showed that both of the Green-Schwarz anomaly factorization formula for the gauge group $E_8 \times E_8$ and the Horava-Witten anomaly factorization formula for the gauge group E_8 could be derived through modular forms of weight 14. This answered a question of J. H. Schwarz. They also established generalizations of these factorization formulas and obtained a new Horava-Witten type factorization formula on 12-dimensional manifolds. In [6], Han, Huang, Liu and Zhang introduced a modular form of weight 14 over $SL(2, \mathbf{Z})$ and a modular form of weight 10 over $SL(2, \mathbf{Z})$ by E_8 bundles and they got some interesting anomaly cancellation formulas on 12-dimensional manifolds. In [5], Chen, Han and Zhang defined an integral modular form of weight $2k$ for a $4k+2$ -dimensional $spin^c$ manifold. In this paper, we twist the Chen-Han-Zhang $SL(2, \mathbf{Z})$ modular form by E_8 bundles and get $SL(2, \mathbf{Z})$ modular forms of weight 14 and 10 for 14 and 10-dimensional $spin^c$ manifolds. In [11], Liu introduced some $\Gamma^0(2)$ and $\Gamma_0(2)$ modular forms and got some interesting anomaly cancellation formulas. In this paper, we also twist the Liu's modular form by E_8 bundles and get $\Gamma^0(2)$ and $\Gamma_0(2)$ modular forms of weight 14 and 10 for a 12-dimensional spin manifold. By these modular forms, we get some new anomaly cancellation formulas of characteristic forms.

2010 *Mathematics Subject Classification:* 58C20; 57R20; 53C80.

Key words and phrases: Modular forms; anomaly cancellation formulas; E_8 bundles .

This paper is organized as follows: In Section 2, we twist the Chen-Han-Zhang $SL(2, \mathbf{Z})$ modular form by E_8 bundles and get $SL(2, \mathbf{Z})$ modular forms of weight 14 and 10 for 14 and 10-dimensional $spin^c$ manifolds and we get some anomaly cancellation formulas of characteristic forms. In Section 3, we twist the Liu's modular form by E_8 bundles and get $\Gamma^0(2)$ and $\Gamma_0(2)$ modular forms of weight 14 and 10 for a 12-dimensional spin manifold. By these modular forms, we get some new anomaly cancellation formulas of characteristic forms.

2. ANOMALY CANCELLATION FORMULAS FOR 14 AND 10-DIMENSIONAL $spin^c$ MANIFOLDS

For the basic representation theory for the affine E_8 , we can see Section 1 in [8]. Here we omit it. For the principle E_8 bundle P_i , $i = 1, 2$, consider the associated bundles $\mathcal{V}_i = \sum_{k=0}^{\infty} (P_i \times_{\rho_k} V_k) q^k \in K(M)[[q]]$. We let $W_i = P_i \times_{\rho_1} V_1$. Let M be a 14-dimensional spinc manifold and L be the complex line bundle associated to the given spinc structure on M . We also consider L as a real vector bundle denoted by L_R .

Denote by $c = c_1(L) = 2\pi\sqrt{-1}u$ the first Chern class of L . Let $\varphi(\tau) = \prod_{n=1}^{\infty} (1 - q^n)$, where $q = e^{2\pi\sqrt{-1}\tau}$ and $\tau \in H$ the upper half plane. We recall the four Jacobi theta functions are defined as follows(cf. [2]):

$$(2.1) \quad \theta(v, \tau) = 2q^{\frac{1}{8}} \sin(\pi v) \prod_{j=1}^{\infty} [(1 - q^j)(1 - e^{2\pi\sqrt{-1}v}q^j)(1 - e^{-2\pi\sqrt{-1}v}q^j)],$$

$$(2.2) \quad \theta_1(v, \tau) = 2q^{\frac{1}{8}} \cos(\pi v) \prod_{j=1}^{\infty} [(1 - q^j)(1 + e^{2\pi\sqrt{-1}v}q^j)(1 + e^{-2\pi\sqrt{-1}v}q^j)],$$

$$(2.3) \quad \theta_2(v, \tau) = \prod_{j=1}^{\infty} [(1 - q^j)(1 - e^{2\pi\sqrt{-1}v}q^{j-\frac{1}{2}})(1 - e^{-2\pi\sqrt{-1}v}q^{j-\frac{1}{2}})],$$

$$(2.4) \quad \theta_3(v, \tau) = \prod_{j=1}^{\infty} [(1 - q^j)(1 + e^{2\pi\sqrt{-1}v}q^{j-\frac{1}{2}})(1 + e^{-2\pi\sqrt{-1}v}q^{j-\frac{1}{2}})],$$

By (2.3)-(2.7) in [8], we have that there are formal two forms y_l^i , $1 \leq l \leq 8$, $i = 1, 2$ such that

$$(2.5) \quad \varphi(\tau)^8 \operatorname{ch}(\mathcal{V}_i) = \frac{1}{2} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right),$$

and

$$(2.6) \quad \sum_{l=1}^8 (2\pi\sqrt{-1}y_l^i)^2 = -\frac{1}{30} c_2(W_i),$$

where $c_2(W_i)$ denotes the second Chern class of W_i . Let $E_2(\tau)$ is the Eisenstein series satisfying:

$$(2.7) \quad E_2(\tau + 1) = E_2(\tau), \quad E_2(-\frac{1}{\tau}) = \tau^2 E_2(\tau) - \frac{6\sqrt{-1}\tau}{\pi},$$

and

$$(2.8) \quad E_2(\tau) = 1 - 24q - 72q^2 + O(q^3).$$

Let $A := p_1(M) - p_1(L_R) + \frac{1}{30}(c_2(W_i) + c_2(W_j))$, where $p_1(M), p_1(L_R)$ denote the first Pontryagin classes of M and L_R . Let

$$(2.9) \quad Q(M, P_i, P_j, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \operatorname{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^m}(\widetilde{L_C}) \right] \cdot \varphi(\tau)^{16} \operatorname{ch}(\mathcal{V}_i) \operatorname{ch}(\mathcal{V}_j) \right\}^{(14)}.$$

Then

$$(2.10)$$

$$Q(M, P_i, P_j, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A} \left(\prod_{j=1}^7 \frac{x_j \theta'(0, \tau)}{\theta(x_j, \tau)} \right) \frac{\sqrt{-1}\theta(u, \tau)}{\theta_1(0, \tau)\theta_2(0, \tau)\theta_3(0, \tau)} \right. \\ \left. \frac{1}{4} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \left(\prod_{l=1}^8 \theta_1(y_l^j, \tau) + \prod_{l=1}^8 \theta_2(y_l^j, \tau) + \prod_{l=1}^8 \theta_3(y_l^j, \tau) \right) \right\}^{(14)},$$

where $\pm 2\pi\sqrt{-1}x_j, 1 \leq j \leq 7$ denote the Chern roots of $T_C M$. One has the following transformation laws of theta functions (cf. [2]):

$$(2.11) \quad \theta(v, \tau + 1) = e^{\frac{\pi\sqrt{-1}}{4}} \theta(v, \tau), \quad \theta(v, -\frac{1}{\tau}) = \frac{1}{\sqrt{-1}} \left(\frac{\tau}{\sqrt{-1}} \right)^{\frac{1}{2}} e^{\pi\sqrt{-1}\tau v^2} \theta(\tau v, \tau);$$

$$(2.12) \quad \theta_1(v, \tau + 1) = e^{\frac{\pi\sqrt{-1}}{4}} \theta_1(v, \tau), \quad \theta_1(v, -\frac{1}{\tau}) = \left(\frac{\tau}{\sqrt{-1}} \right)^{\frac{1}{2}} e^{\pi\sqrt{-1}\tau v^2} \theta_2(\tau v, \tau);$$

$$(2.13) \quad \theta_2(v, \tau + 1) = \theta_3(v, \tau), \quad \theta_2(v, -\frac{1}{\tau}) = \left(\frac{\tau}{\sqrt{-1}} \right)^{\frac{1}{2}} e^{\pi\sqrt{-1}\tau v^2} \theta_1(\tau v, \tau);$$

$$(2.14) \quad \theta_3(v, \tau + 1) = \theta_2(v, \tau), \quad \theta_3(v, -\frac{1}{\tau}) = \left(\frac{\tau}{\sqrt{-1}} \right)^{\frac{1}{2}} e^{\pi\sqrt{-1}\tau v^2} \theta_3(\tau v, \tau),$$

$$(2.15) \quad \theta'(v, \tau + 1) = e^{\frac{\pi\sqrt{-1}}{4}} \theta'(v, \tau), \quad \theta'(0, -\frac{1}{\tau}) = \frac{1}{\sqrt{-1}} \left(\frac{\tau}{\sqrt{-1}} \right)^{\frac{1}{2}} \tau \theta'(0, \tau).$$

Definition 2.1. A modular form over Γ , a subgroup of $SL_2(\mathbf{Z})$, is a holomorphic function $f(\tau)$ on H such that

$$(2.16) \quad f(g\tau) := f\left(\frac{a\tau + b}{c\tau + d}\right) = \chi(g)(c\tau + d)^k f(\tau), \quad \forall g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma,$$

where $\chi : \Gamma \rightarrow \mathbf{C}^*$ is a character of Γ . k is called the weight of f .

By (2.7) and (2.11)-(2.15), we have

Lemma 2.2. $Q(M, P_i, P_j, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 14.

By

$$(2.17) \quad \text{ch}(\mathcal{V}_i) = 1 + (248 - c_2(W_i) + \dots)q + \dots,$$

and (2.8), we have by expanding the q -series:

(2.18)

$$\begin{aligned} & e^{\frac{1}{24}E_2(\tau)A} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^m}(\widetilde{L_C}) \right] \varphi(\tau)^{16} \text{ch}(\mathcal{V}_i) \text{ch}(\mathcal{V}_j) \\ &= (e^{\frac{1}{24}A} - e^{\frac{1}{24}A} Aq + O(q^2)) \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch} \left[(1 + q\widetilde{T_C M} + O(q^2))(1 - q\widetilde{L_C} + O(q^2)) \right] \\ & \quad \cdot (1 - 16q + O(q^2))(1 + \text{ch}(W_i)q + O(q^2))(1 + \text{ch}(W_j)q + O(q^2)) \\ &= e^{\frac{1}{24}A} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) + \left[e^{\frac{1}{24}A} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} - 16 + W_i + W_j) \right. \\ & \quad \left. - e^{\frac{1}{24}A} A \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right] q + O(q^2). \end{aligned}$$

It is well known that modular forms over $SL_2(\mathbf{Z})$ can be expressed as polynomials of the Eisenstein series $E_4(\tau)$ and $E_6(\tau)$, where

$$(2.19) \quad E_4(\tau) = 1 + 240q + 2160q^2 + 6720q^3 + \dots,$$

$$(2.20) \quad E_6(\tau) = 1 - 504q - 16632q^2 - 122976q^3 + \dots.$$

Their weights are 4 and 6 respectively. Since $Q(M, P_i, P_j, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 14, it must be a multiple of

$$(2.21) \quad E_4(\tau)^2 E_6(\tau) = 1 - 24q - 196632q^2 + \dots.$$

So

$$\begin{aligned} (2.22) \quad & \left\{ e^{\frac{1}{24}A} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} - 16 + W_i + W_j) - e^{\frac{1}{24}A} A \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(14)} \\ &= -24 \left\{ e^{\frac{1}{24}A} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(14)}. \end{aligned}$$

By (2.22), we have

Theorem 2.3. *One has the following equality:*

(2.23)

$$\begin{aligned} & \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} + 8 + W_i + W_j) \right\}^{(14)} \\ &= A \left\{ e^{\frac{1}{24}A} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) - \frac{e^{\frac{1}{24}A} - 1}{A} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} + 8 + W_i + W_j) \right\}^{(10)}. \end{aligned}$$

Corollary 2.4. *When $A = 0$, we have*

$$\begin{aligned} (2.24) \quad & \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} - 16 + W_i + W_j) \right\}^{(14)} \\ &= -24 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(14)}. \end{aligned}$$

When M is a 14-dimensional spin c manifold, then $\text{Ind}D^c \otimes (\widetilde{T_C M} - \widetilde{L_C} - 16 + W_i + W_j)_+$ is a multiply of 24.

Let $A_1 := p_1(M) - p_1(L_R) + \frac{1}{30}c_2(W_i)$. Let

$$(2.25) \quad Q(M, P_i, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^m}(\widetilde{L_C}) \right] \cdot \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \right\}^{(14)}.$$

Then

$$(2.26) \quad Q(M, P_i, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A_1} \left(\prod_{j=1}^7 \frac{x_j \theta'(0, \tau)}{\theta(x_j, \tau)} \right) \frac{\sqrt{-1}\theta(u, \tau)}{\theta_1(0, \tau)\theta_2(0, \tau)\theta_3(0, \tau)} \right. \\ \left. \frac{1}{2} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \right\}^{(14)}.$$

Similar to Lemma 2.2, we have

Lemma 2.5. *$Q(M, P_i, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 10.*

Similar to (2.18), we have

$$\begin{aligned} (2.27) \quad & e^{\frac{1}{24}E_2(\tau)A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^m}(\widetilde{L_C}) \right] \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \\ &= e^{\frac{1}{24}A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) + \left[e^{\frac{1}{24}A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} - 8 + W_i) \right. \\ &\quad \left. - e^{\frac{1}{24}A_1} A_1 \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right] q + O(q^2). \end{aligned}$$

Since $Q(M, P_i, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 10, it must be a multiple of

$$(2.28) \quad E_4(\tau)E_6(\tau) = 1 - 264q - 135432q^2 + \dots .$$

So

$$(2.29) \quad \begin{aligned} & \left\{ e^{\frac{1}{24}A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \operatorname{ch}(\widetilde{T_C M} - \widetilde{L_C} - 8 + W_i) - e^{\frac{1}{24}A_1} A_1 \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(14)} \\ &= -264 \left\{ e^{\frac{1}{24}A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(14)}. \end{aligned}$$

Theorem 2.6. *One has the following equality:*

$$(2.30)$$

$$\begin{aligned} & \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \operatorname{ch}(\widetilde{T_C M} - \widetilde{L_C} + 256 + W_i) \right\}^{(14)} \\ &= A_1 \left\{ e^{\frac{1}{24}A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) - \frac{e^{\frac{1}{24}A_1} - 1}{A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \operatorname{ch}(\widetilde{T_C M} - \widetilde{L_C} + 256 + W_i) \right\}^{(10)}. \end{aligned}$$

Corollary 2.7. *When $A_1 = 0$, we have*

$$(2.31) \quad \begin{aligned} & \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \operatorname{ch}(\widetilde{T_C M} - \widetilde{L_C} - 8 + W_i) \right\}^{(14)} \\ &= -264 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(14)}. \end{aligned}$$

When M is a 14-dimensional $spin^c$ manifold, then $\operatorname{Ind}D^c \otimes (\widetilde{T_C M} - \widetilde{L_C} - 8 + W_i)_+$ is a multiply of 264.

If M is a 10-dimensional $spin^c$ manifold, we let

$$(2.32) \quad R(M, P_i, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \operatorname{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^m}(\widetilde{L_C}) \right] \cdot \varphi(\tau)^8 \operatorname{ch}(\mathcal{V}_i) \right\}^{(10)}.$$

Then

$$(2.33) \quad Q(M, P_i, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A_1} \left(\prod_{j=1}^5 \frac{x_j \theta'(0, \tau)}{\theta(x_j, \tau)} \right) \frac{\sqrt{-1}\theta(u, \tau)}{\theta_1(0, \tau)\theta_2(0, \tau)\theta_3(0, \tau)} \right. \\ \left. \frac{1}{2} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \right\}^{(10)}.$$

Similar to Lemma 2.2, we have

Lemma 2.8. *$R(M, P_i, \tau)$ is a modular form over $SL_2(\mathbf{Z})$ with the weight 8.*

By (2.27) and Lemma 2.8, it must be a multiple of

$$(2.34) \quad E_4(\tau)^2 = 1 + 480q + 61920q^2 + \dots.$$

So

$$(2.35) \quad \begin{aligned} & \left\{ e^{\frac{1}{24}A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} - 8 + W_i) - e^{\frac{1}{24}A_1} A_1 \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(10)} \\ &= 480 \left\{ e^{\frac{1}{24}A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(10)}. \end{aligned}$$

Theorem 2.9. *One has the following equality:*

$$(2.36) \quad \begin{aligned} & \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} - 488 + W_i) \right\}^{(10)} \\ &= A_1 \left\{ e^{\frac{1}{24}A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) - \frac{e^{\frac{1}{24}A_1} - 1}{A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} - 488 + W_i) \right\}^{(6)}. \end{aligned}$$

Corollary 2.10. *When $A_1 = 0$, we have*

$$\left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} + W_i) \right\}^{(10)} = 488 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(10)}.$$

When M is a 10-dimensional spin^c manifold, then $\text{Ind}D^c \otimes (\widetilde{T_C M} - \widetilde{L_C} + W_i)_+$ is a multiply of 488.

In the following, we consider the q^2 -terms in the q -expansions of $Q(M, P_i, P_j, \tau)$, $Q(M, P_i, \tau)$, $R(M, P_i, \tau)$. Let $\overline{W}_i = P_i \times_{\rho_2} V_2$. We assume that $A = 0$, then

$$(2.37) \quad \begin{aligned} & e^{\frac{1}{24}E_2(\tau)A} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^m}(\widetilde{L_C}) \right] \varphi(\tau)^{16} \text{ch}(\mathcal{V}_i) \text{ch}(\mathcal{V}_j) \\ &= \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch} [1 + qB_1 + q^2B_2 + O(q^3)] \\ & \cdot (1 - 16q + 104q^2)(1 + \text{ch}(W_i)q + q^2\text{ch}(\overline{W}_i) + O(q^3))(1 + \text{ch}(W_j)q + q^2\text{ch}(\overline{W}_j) + O(q^3)) \\ &= \widehat{A}(TX) \exp\left(\frac{c}{2}\right) + \left[\widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} - 16 + W_i + W_j) \right] q \\ &+ \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) (\text{ch}(\overline{W}_j) + \text{ch}(\overline{W}_i) + \text{ch}(W_i)\text{ch}(W_j) - 16\text{ch}(W_j) - 16\text{ch}(W_i) + 104 \right. \\ & \left. + B_1 \wedge [\text{ch}(W_j) + \text{ch}(W_i) - 16] + B_2) \right\} q^2 + O(q^3). \end{aligned}$$

where

$$(2.38) \quad \begin{aligned} B_1 &= \widetilde{T_C M} - \widetilde{L_C}, \\ B_2 &= \wedge^2 \widetilde{L_C} - \widetilde{L_C} - \widetilde{T_C M} \otimes \widetilde{L_C} + S^2 \widetilde{T_C M} + \widetilde{T_C M}. \end{aligned}$$

By (2.21) and (2.37), we get

Theorem 2.11. *When $A = 0$, we have*

(2.39)

$$\begin{aligned} & \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) (\text{ch}(\overline{W}_j) + \text{ch}(\overline{W}_i) + \text{ch}(W_i)\text{ch}(W_j) - 16\text{ch}(W_j) - 16\text{ch}(W_i) + 104 \right. \\ & \quad \left. + B_1 \wedge [\text{ch}(W_j) + \text{ch}(W_i) - 16] + B_2) \right\}^{(14)} \\ & = -196632 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(14)}. \end{aligned}$$

We assume that $A_1 = 0$, then

$$\begin{aligned} (2.40) \quad & e^{\frac{1}{24}E_2(\tau)A_1} \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^m}(\widetilde{L_C}) \right] \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \\ & = \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch} [1 + qB_1 + q^2B_2 + O(q^3)] \\ & \quad \cdot (1 - 8q + 20q^2)(1 + \text{ch}(W_i)q + q^2\text{ch}(\overline{W}_i) + O(q^3)) \\ & = \widehat{A}(TX) \exp\left(\frac{c}{2}\right) + \left[\widehat{A}(TX) \exp\left(\frac{c}{2}\right) \text{ch}(\widetilde{T_C M} - \widetilde{L_C} - 8 + W_i) \right] q \\ & \quad + \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) (20 + \text{ch}(\overline{W}_i) - 8\text{ch}(W_i) \right. \\ & \quad \left. + B_1 \wedge [\text{ch}(W_i) - 8] + B_2) \right\} q^2 + O(q^3). \end{aligned}$$

By (2.28) and (2.40), we get

Theorem 2.12. *When $A_1 = 0$, we have*

$$\begin{aligned} (2.41) \quad & \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) (20 + \text{ch}(\overline{W}_i) - 8\text{ch}(W_i) + B_1 \wedge [\text{ch}(W_i) - 8] + B_2) \right\}^{(14)} \\ & = -135432 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(14)}. \end{aligned}$$

Similarly, we have:

Theorem 2.13. *When $A_1 = 0$, we have*

$$\begin{aligned} (2.42) \quad & \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) (20 + \text{ch}(\overline{W}_i) - 8\text{ch}(W_i) + B_1 \wedge [\text{ch}(W_i) - 8] + B_2) \right\}^{(10)} \\ & = 61920 \left\{ \widehat{A}(TX) \exp\left(\frac{c}{2}\right) \right\}^{(10)}. \end{aligned}$$

3. ANOMALY CANCELLATION FORMULAS FOR 12-DIMENSIONAL MANIFOLDS

In this section, we let M be a 12-dimensional manifold. Let $\widehat{L}(TM, \nabla^{TM})$ be the Hirzebruch characteristic forms defined by ([13])

$$\widehat{L}(TM, \nabla^{TM}) = \det^{\frac{1}{2}} \left(\frac{\frac{\sqrt{-1}}{2\pi} R^{TM}}{\tanh(\frac{\sqrt{-1}}{4\pi} R^{TM})} \right).$$

Let $A_2 := \frac{1}{30}(c_2(W_i) + c_2(W_j))$. Let

$$(3.1) \quad Q_1(M, P_i, P_j, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A_2} \widehat{L}(TM, \nabla^{TM}) \operatorname{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{q^m}(\widetilde{T_C M}) \right] \cdot \varphi(\tau)^{16} \operatorname{ch}(\mathcal{V}_i) \operatorname{ch}(\mathcal{V}_j) \right\}^{(12)},$$

(3.2)

$$Q_2(M, P_i, P_j, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A_2} \widehat{A}(TM, \nabla^{TM}) \operatorname{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^{m-\frac{1}{2}}}(\widetilde{T_C M}) \right] \cdot \varphi(\tau)^{16} \operatorname{ch}(\mathcal{V}_i) \operatorname{ch}(\mathcal{V}_j) \right\}^{(12)},$$

Then

(3.3)

$$Q_1(M, P_i, P_j, \tau) = 2^6 \left\{ e^{\frac{1}{24}E_2(\tau)A_2} \left(\prod_{j=1}^6 \frac{x_j \theta'(0, \tau) \theta_1(x_j, \tau)}{\theta(x_j, \tau) \theta_1(0, \tau)} \right) \frac{1}{4} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \left(\prod_{l=1}^8 \theta_1(y_l^j, \tau) + \prod_{l=1}^8 \theta_2(y_l^j, \tau) + \prod_{l=1}^8 \theta_3(y_l^j, \tau) \right) \right\}^{(12)},$$

and

(3.4)

$$Q_2(M, P_i, P_j, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A_2} \left(\prod_{j=1}^6 \frac{x_j \theta'(0, \tau) \theta_2(x_j, \tau)}{\theta(x_j, \tau) \theta_2(0, \tau)} \right) \frac{1}{4} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \left(\prod_{l=1}^8 \theta_1(y_l^j, \tau) + \prod_{l=1}^8 \theta_2(y_l^j, \tau) + \prod_{l=1}^8 \theta_3(y_l^j, \tau) \right) \right\}^{(12)},$$

Let

$$\Gamma_0(2) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbf{Z}) \mid c \equiv 0 \pmod{2} \right\},$$

$$\Gamma^0(2) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbf{Z}) \mid b \equiv 0 \pmod{2} \right\},$$

be the two modular subgroups of $SL_2(\mathbf{Z})$. Writing $\theta_j = \theta_j(0, \tau)$, $1 \leq j \leq 3$, we introduce four explicit modular forms ([11]),

$$(3.5) \quad \begin{aligned} \delta_1(\tau) &= \frac{1}{8}(\theta_2^4 + \theta_3^4), \quad \varepsilon_1(\tau) = \frac{1}{16}\theta_2^4\theta_3^4, \\ \delta_2(\tau) &= -\frac{1}{8}(\theta_1^4 + \theta_3^4), \quad \varepsilon_2(\tau) = \frac{1}{16}\theta_1^4\theta_3^4. \end{aligned}$$

They have the following Fourier expansions in $q^{\frac{1}{2}}$:

$$(3.6) \quad \begin{aligned} \delta_1(\tau) &= \frac{1}{4} + 6q + 6q^2 + \cdots, \quad \varepsilon_1(\tau) = \frac{1}{16} - q + 7q^2 + \cdots, \\ 8\delta_2(\tau) &= -1 - 24q^{\frac{1}{2}} - 24q - 96q^{\frac{3}{2}} + \cdots, \quad \varepsilon_2(\tau) = q^{\frac{1}{2}} + 8q + 28q^{\frac{3}{2}} + \cdots. \end{aligned}$$

They also satisfy the transformation laws,

$$(3.7) \quad \delta_2\left(-\frac{1}{\tau}\right) = \tau^2 \delta_1(\tau), \quad \varepsilon_2\left(-\frac{1}{\tau}\right) = \tau^4 \varepsilon_1(\tau),$$

Lemma 3.1. ([11]) $\delta_1(\tau)$ (resp. $\varepsilon_1(\tau)$) is a modular form of weight 2 (resp. 4) over $\Gamma_0(2)$, $\delta_2(\tau)$ (resp. $\varepsilon_2(\tau)$) is a modular form of weight 2 (resp. 4) over $\Gamma^0(2)$ and moreover $\mathcal{M}_{\mathbf{R}}(\Gamma^0(2)) = \mathbf{R}[\delta_2(\tau), \varepsilon_2(\tau)]$.

By (2.11)-(2.15) and (2.7), we have

Lemma 3.2. $Q_1(M, P_i, P_j, \tau)$ is a modular form of weight 14 over $\Gamma_0(2)$, while $Q_2(M, P_i, P_j, \tau)$ is a modular form of weight 14 over $\Gamma^0(2)$. Moreover, the following identity holds,

$$(3.8) \quad Q_1(M, P_i, P_j, -\frac{1}{\tau}) = 2^6 \tau^{14} Q_2(M, P_i, P_j, \tau).$$

Theorem 3.3. We have the following equality:

$$(3.9) \quad \begin{aligned} 32 \left\{ e^{\frac{1}{24}A_2} \widehat{L}(TM) \right\}^{(12)} &= \left\{ e^{\frac{1}{24}A_2} \widehat{A}(TM) \text{ch}[2240 + 309\widetilde{T_C M} + 24(\wedge^2 \widetilde{T_C M} + W_i + W_j) \right. \\ &\quad \left. + \widetilde{T_C M} \otimes \widetilde{T_C M} + \widetilde{T_C M} \otimes (W_i + W_j) + \wedge^3 \widetilde{T_C M}] - \frac{1}{24} e^{\frac{1}{24}A_2} A_2 \widehat{A}(TX) \text{ch}(576 + 24\widetilde{T_C M}) \right\}^{(12)}. \end{aligned}$$

Proof. By Lemmas 3.1 and 3.2, we have

$$(3.10) \quad Q_2(M, P_i, P_j, \tau) = h_0(8\delta_2)^7 + h_1(8\delta_2)^5 \varepsilon_2 + h_2(8\delta_2)^3 \varepsilon_2^2 + h_3(8\delta_2) \varepsilon_2^3,$$

$$(3.11) \quad Q_1(M, P_i, P_j, \tau) = 2^6 [h_0(8\delta_1)^7 + h_1(8\delta_1)^5 \varepsilon_1 + h_2(8\delta_1)^3 \varepsilon_1^2 + h_3(8\delta_1) \varepsilon_1^3],$$

where each h_r , $0 \leq r \leq 3$, is a real multiple of the volume form at x .

(3.12)

$$\begin{aligned}
& e^{\frac{1}{24}E_2(\tau)A_2} \widehat{A}(TM, \nabla^{TM}) \operatorname{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^{m-\frac{1}{2}}} (\widetilde{T_C M}) \right] \cdot \varphi(\tau)^{16} \operatorname{ch}(\mathcal{V}_i) \operatorname{ch}(\mathcal{V}_j) \\
& = (e^{\frac{1}{24}A_2} - e^{\frac{1}{24}A_2} A_2 q + O(q^2)) \widehat{A}(TM) \operatorname{ch} \left[(1 + q \widetilde{T_C M} + O(q^2)) \right. \\
& \quad \left. \otimes (1 - q^{\frac{1}{2}} \widetilde{T_C M} + q \wedge^2 \widetilde{T_C M} - q^{\frac{3}{2}} \wedge^3 \widetilde{T_C M} + O(q^2)) \otimes (1 - q^{\frac{3}{2}} \widetilde{T_C M}) \right] \\
& \quad \cdot (1 - 16q + O(q^2))(1 + \operatorname{ch}(W_i)q + O(q^2))(1 + \operatorname{ch}(W_j)q + O(q^2)) \\
& = e^{\frac{1}{24}A_2} \widehat{A}(TM) - q^{\frac{1}{2}} e^{\frac{1}{24}A_2} \widehat{A}(TX) \operatorname{ch}(\widetilde{T_C M}) \\
& \quad + q[e^{\frac{1}{24}A_2} \widehat{A}(TM) \operatorname{ch}(\widetilde{T_C M} + \wedge^2 \widetilde{T_C M} + W_i + W_j - 16) - e^{\frac{1}{24}A_2} A_2 \widehat{A}(TM)] \\
& \quad + q^{\frac{3}{2}} \left[-e^{\frac{1}{24}A_2} \widehat{A}(TX) \operatorname{ch}(\widetilde{T_C M} \otimes \widetilde{T_C M} + \widetilde{T_C M} \otimes (W_i + W_j - 16) + \widetilde{T_C M} + \wedge^3 \widetilde{T_C M}) \right. \\
& \quad \left. + e^{\frac{1}{24}A_2} A_2 \widehat{A}(TM) \operatorname{ch}(\widetilde{T_C M}) \right] + O(q^2).
\end{aligned}$$

By (3.10), (3.12) and (3.6), comparing the the constant term, $q^{\frac{1}{2}}, q, q^{\frac{3}{2}}$ terms in (3.10), we get

(3.13)

$$\begin{aligned}
h_0 &= - \left\{ e^{\frac{1}{24}A_2} \widehat{A}(TM) \right\}^{(12)} \\
h_1 &= \left\{ e^{\frac{1}{24}A_2} \widehat{A}(TM) \operatorname{ch}(\widetilde{T_C M} + 168) \right\}^{(12)} \\
h_2 &= \left\{ e^{\frac{1}{24}A_2} \widehat{A}(TM) \operatorname{ch}(-9224 - 129 \widetilde{T_C M} - \wedge^2 \widetilde{T_C M} - W_i - W_j) + e^{\frac{1}{24}A_2} A_2 \widehat{A}(TM) \right\}^{(12)} \\
h_3 &= \left\{ e^{\frac{1}{24}A_2} \widehat{A}(TM) \operatorname{ch}(\widetilde{T_C M} \otimes \widetilde{T_C M} + \widetilde{T_C M} \otimes (W_i + W_j - 16) + \widetilde{T_C M} + \wedge^3 \widetilde{T_C M} \right. \\
&\quad \left. + 508704 - 6868(\widetilde{T_C M} + 168) + 88 \times 9224 + 88 \times 129 \widetilde{T_C M} \right. \\
&\quad \left. + 88 \wedge^2 \widetilde{T_C M} + 88W_i + 88W_j) - e^{\frac{1}{24}A_2} A_2 \widehat{A}(TM) \operatorname{ch}(\widetilde{T_C M} + 88) \right\}^{(12)}.
\end{aligned}$$

By (3.13), (3.11) and (3.6), comparing the the constant term in (3.11), we get Theorem 3.3. \square

Corollary 3.4. *We have the following equality when $A_2 = 0$:*

$$\begin{aligned}
(3.14) \quad 32 \left\{ \widehat{L}(TM) \right\}^{(12)} &= \left\{ \widehat{A}(TM) \operatorname{ch}[2240 + 309 \widetilde{T_C M} + 24(\wedge^2 \widetilde{T_C M} + W_i + W_j) \right. \\
&\quad \left. + \widetilde{T_C M} \otimes \widetilde{T_C M} + \widetilde{T_C M} \otimes (W_i + W_j) + \wedge^3 \widetilde{T_C M}] \right\}^{(12)}.
\end{aligned}$$

When M is spin, then the characteristic number of the right hand of (3.14) is a multiple of 32.

Let $A_3 := \frac{1}{30}c_2(W_i)$. Let

$$(3.15) \quad Q_1(M, P_i, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A_3} \widehat{L}(TM, \nabla^{TM}) \text{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{q^m}(\widetilde{T_C M}) \right] \cdot \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \right\}^{(12)},$$

$$(3.16) \quad Q_2(M, P_i, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A_3} \widehat{A}(TM, \nabla^{TM}) \text{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^{m-\frac{1}{2}}}(\widetilde{T_C M}) \right] \cdot \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \right\}^{(12)}.$$

Then

$$(3.17) \quad Q_1(M, P_i, \tau) = 2^6 \left\{ e^{\frac{1}{24}E_2(\tau)A_3} \left(\prod_{j=1}^6 \frac{x_j \theta'(0, \tau) \theta_1(x_j, \tau)}{\theta(x_j, \tau) \theta_1(0, \tau)} \right) \frac{1}{2} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \right\}^{(12)},$$

and

$$(3.18) \quad Q_2(M, P_i, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)A_3} \left(\prod_{j=1}^6 \frac{x_j \theta'(0, \tau) \theta_2(x_j, \tau)}{\theta(x_j, \tau) \theta_2(0, \tau)} \right) \frac{1}{2} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \right\}^{(12)}.$$

We have

Lemma 3.5. $Q_1(M, P_i, \tau)$ is a modular form of weight 10 over $\Gamma_0(2)$, while $Q_2(M, P_i, \tau)$ is a modular form of weight 10 over $\Gamma^0(2)$. Moreover, the following identity holds,

$$(3.19) \quad Q_1(M, P_i, -\frac{1}{\tau}) = 2^6 \tau^{10} Q_2(M, P_i, \tau).$$

Theorem 3.6. We have the following equality:

$$(3.20) \quad \left\{ e^{\frac{1}{24}A_3} \widehat{L}(TM) \right\}^{(12)} = -\frac{1}{2} \left\{ e^{\frac{1}{24}A_3} \widehat{A}(TM) \text{ch}[17\widetilde{T_C M} + \wedge^2 \widetilde{T_C M} + W_i + 128] \right\}^{(12)} + \frac{1}{60} \left\{ c_2(W_i) e^{\frac{1}{24}A_3} \widehat{A}(TX) \right\}^{(12)}.$$

Proof. By Lemmas 3.1 and 3.5, we have

$$(3.21) \quad Q_2(M, P_i, \tau) = h'_0(8\delta_2)^5 + h'_1(8\delta_2)^3 \varepsilon_2 + h'_2(8\delta_2) \varepsilon_2^2,$$

$$(3.22) \quad Q_1(M, P_i, \tau) = 2^6 [h'_0(8\delta_1)^5 + h'_1(8\delta_1)^3 \varepsilon_1 + h'_2(8\delta_1) \varepsilon_1^2],$$

where each h'_r , $0 \leq r \leq 2$, is a real multiple of the volume form at x .

$$\begin{aligned}
(3.23) \quad & e^{\frac{1}{24}E_2(\tau)A_3} \widehat{A}(TM, \nabla^{TM}) \text{ch} \left[\bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_C M}) \otimes \bigotimes_{m=1}^{\infty} \wedge_{-q^{m-\frac{1}{2}}} (\widetilde{T_C M}) \right] \cdot \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \\
& = (e^{\frac{1}{24}A_3} - e^{\frac{1}{24}A_3} A_3 q + O(q^2)) \widehat{A}(TM) \text{ch} \left[(1 + q \widetilde{T_C M} + O(q^2)) \right. \\
& \quad \otimes (1 - q^{\frac{1}{2}} \widetilde{T_C M} + q \wedge^2 \widetilde{T_C M} + O(q^{\frac{3}{2}})) \\
& \quad \cdot (1 - 8q + O(q^2))(1 + \text{ch}(W_i)q + O(q^2)) \\
& = e^{\frac{1}{24}A_3} \widehat{A}(TM) - q^{\frac{1}{2}} e^{\frac{1}{24}A_3} \widehat{A}(TM) \text{ch}(\widetilde{T_C M}) \\
& \quad + q [e^{\frac{1}{24}A_3} \widehat{A}(TM) \text{ch}(\widetilde{T_C M} + \wedge^2 \widetilde{T_C M} + W_i - 8) - e^{\frac{1}{24}A_3} A_3 \widehat{A}(TM)] + O(q^{\frac{3}{2}})
\end{aligned}$$

By (3.21), (3.23) and (3.6), comparing the the constant term, $q^{\frac{1}{2}}, q$ terms in (3.21), we get

$$\begin{aligned}
(3.24) \quad h'_0 & = - \left\{ e^{\frac{1}{24}A_3} \widehat{A}(TM) \right\}^{(12)} \\
h'_1 & = \left\{ e^{\frac{1}{24}A_3} \widehat{A}(TM) \text{ch}(\widetilde{T_C M} + 120) \right\}^{(12)} \\
h'_2 & = \left\{ -e^{\frac{1}{24}A_3} \widehat{A}(TM) \text{ch}(3712 + 81 \widetilde{T_C M} + \wedge^2 \widetilde{T_C M} + W_i) + e^{\frac{1}{24}A_3} A_3 \widehat{A}(TM) \right\}^{(12)}
\end{aligned}$$

By (3.22), (3.24) and (3.6), comparing the the constant term in (3.22), we get Theorem 3.6. \square

Corollary 3.7. *We have the following equality when $c_2(W_i) = 0$:*

$$- 2 \left\{ \widehat{L}(TM) \right\}^{(12)} = \left\{ \widehat{A}(TM) \text{ch}[17 \widetilde{T_C M} + \wedge^2 \widetilde{T_C M} + W_i + 128] \right\}^{(12)}.$$

When M is a spin manifold, then $\text{Ind}(D \otimes (17 \widetilde{T_C M} + \wedge^2 \widetilde{T_C M} + W_i + 128)_+)$ is a multiple of 2.

Comparing the coefficient of q in (3.22), by (3.24), we have

Theorem 3.8. *We have the following equality:*

$$\begin{aligned}
(3.25) \quad & \left\{ e^{\frac{1}{24}A_3} \widehat{L}(TM) [-A_3 + 2 \text{ch}(\widetilde{T_C M}) - 8 + \text{ch}(W_i)] \right\}^{(12)} \\
& = \left\{ e^{\frac{1}{24}A_3} \widehat{A}(TM) \text{ch}[2116 \widetilde{T_C M} + 4 \wedge^2 \widetilde{T_C M} + 4W_i - 15872] \right\}^{(12)} \\
& \quad - \frac{2}{15} \left\{ c_2(W_i) e^{\frac{1}{24}A_3} \widehat{A}(TX) \right\}^{(12)}.
\end{aligned}$$

Corollary 3.9. *We have the following equality when $c_2(W_i) = 0$:*

$$(3.26) \quad \begin{aligned} & \left\{ \widehat{L}(TM)[2\text{ch}(\widetilde{T_C M}) - 8 + \text{ch}(W_i)] \right\}^{(12)} \\ &= \left\{ \widehat{A}(TM)\text{ch}[2116\widetilde{T_C M} + 4 \wedge^2 \widetilde{T_C M} + 4W_i - 15872] \right\}^{(12)}. \end{aligned}$$

Remark: In [4], Chen and Han constructed some modular forms on odd dimensional manifolds. We can construct some new modular forms on odd dimensional manifolds by twist the Chen-Han modular forms by E_8 bundles.

4. ACKNOWLEDGEMENTS

The author was supported in part by NSFC No.11771070. The first author is indebted to Prof. F. Han for very helpful discussions. The authors also thank the referee for his (or her) careful reading and helpful comments.

REFERENCES

- [1] L. Alvarez-Gaumé, E. Witten, Gravitaional anomalies, Nucl. Phys. B234 (1983), 269-330.
- [2] K. Chandrasekharan, *Elliptic Functions*, Springer-Verlag, 1985.
- [3] Q. Chen, F. Han, Modular invariance and twisted anomaly cancellations of characteristic numbers, Trans. Amer. Math. Soc. 361 (2009), 1463-1493.
- [4] Q. Chen, F. Han, Elliptic genera, transgression and loop space Chern-Simons form, Comm. Anal. Geom. 17 (2009), no. 1, 73-106.
- [5] Q. Chen, F. Han, W. Zhang, Generalized Witten genus and vanishing theorems. J. Differential Geom. 88(2011), no. 1, 1-40.
- [6] F. Han, R. Huang, K. Liu and W. Zhang, Cubic forms, anomaly cancellation and modularity. Adv. Math. 394(2022), Paper No. 108023, 46pp.
- [7] F. Han, K. Liu and W. Zhang, Modular forms and generalized anomaly cancellation formulas, J. Geom. Phys. 62 (2012) 1038-1053.
- [8] F. Han, K. Liu and W. Zhang, Anomaly cancellation and modularity, II: the $E_8 \times E_8$ case. Sci. China Math. 60(2017), no. 6, 985-994.
- [9] F. Han, W. Zhang, Spin^c-manifold and elliptic genera, C. R. Acad. Sci. Paris Serie I., 336 (2003), 1011-1014.
- [10] F. Han, W. Zhang, Modular invariance, characteristic numbers and eta Invariants, J. Diff. Geom. 67 (2004), 257-288.
- [11] K. Liu, Modular invariance and characteristic numbers. Commu. Math. Phys. 174 (1995), 29-42.
- [12] K. Liu, On elliptic genera and theta-functions. Topology 35 (1996), no. 3, 617-640.
- [13] W. Zhang, *Lectures on Chern-weil Theory and Witten Deformations*. Nankai Tracks in Mathematics Vol. 4, World Scientific, Singapore, 2001.

1.Yong Wang, Corresponding author, School of Mathematics and Statistics, Northeast Normal University, Changchun Jilin, 130024, China

E-mail: wangy581@nenu.edu.cn

2.Yuchen Yang, School of Mathematics and Statistics , Northeast Normal University,
Changchun, Jilin 130024, China

E-mail: *yangyc580@nenu.edu.cn*