

Correction to: Powers of the vertex cover ideals (Collect. Math. 65 (2014) 169–181).

Dancheng Lu^{1*} and Zexin Wang^{1†}

¹School of Mathematical Science, Soochow University, No. 1 Shizhi Street, Suzhou, 215021, Jiangsu, P.R.China.

*Corresponding author(s). E-mail(s): ludancheng@suda.edu.cn;
Contributing authors: zexinwang6@outlook.com;

†These authors contributed equally to this work.

Abstract

We point out an essential gap in the proof of one of main results in [3] and then give a corrected proof for it.

Keywords: Cohen-Macaulay, Cactus graph, Weakly polymatroidal, Vertex cover ideal, Power

In [3] F. Mohammadi succeeded in classifying all the Cohen-Macaulay cactus graphs. Based on this result, he proved in [3, Theorem 4.3] that if G is a Cohen-Macaulay cactus graph then all the powers of the vertex cover ideal $J(G)$ of G are weakly polymatroidal. In this note we illustrate by an example that there exists an essential gap in the proof of [3, Theorem 4.3] and then present a corrected proof for this result.

We adopt the notation from the original paper [3], which the reader should have at hand when reading this note. First of all, We explain why some parts of the original proof of [3, Theorem 4.3] is not correct.

There were several cases to consider in the proof. In case 4, it is assumed that $z = y_{i3}$, then one deduces that $\{y_{i1}, y_{i2}, y_{i5}\} \subseteq \text{supp}(g_j)$ for some $1 \leq j \leq k$. Let us see the following example.

Example 1. Let G be a 5-cycle whose vertices are labeled y_1, y_4, y_2, y_3, y_5 in order. Then $f = y_1 y_2 y_3^2 y_4 y_5 = (y_1 y_2 y_3)(y_3 y_4 y_5)$ and $g = y_1 y_2 y_3 y_4^2 y_5 = (y_2 y_4 y_5)(y_1 y_3 y_4)$ are minimal monomial generators of $J(G)^2$. We may write $g_1 = y_2 y_4 y_5$ and $g_2 = y_1 y_3 y_4$. Note that $\deg_{y_i} f = \deg_{y_i} g$ for $i = 1, 2$ and $\deg_{y_3} f > \deg_{y_3} g$. According to the

Next, we denote by T the induced subgraph of G on the subset $V(G) \setminus \{y_{i3}, y_{i4}, y_{i5}\}$, and for $a = 1, 2$, we set

$$T_a = \{z \in V(T) \setminus \{y_{i1}, y_{i2}\} \mid \text{there exists a path in } T \text{ connecting } z \text{ with } y_{ia}\}.$$

Since G is a cactus graph, we have the subsets $T_1, \{y_{i1}, y_{i2}\}$ and T_2 are pairwise disjoint and $V(T) = T_1 \cup \{y_{i1}, y_{i2}\} \cup T_2$.

Case $z = y_{i3}$: If $g_{125} \neq 0$, there exists $s \in \{1, \dots, k\}$ such that $\{y_{i1}, y_{i2}, y_{i5}\} \subseteq \text{supp}(g_s)$. Since the subset $(\text{supp}(g_s) \setminus \{y_{i5}\}) \cup \{y_{i3}\}$ is again a minimal vertex cover of G , we have $y_{i3}g/y_{i5} \in J(G)^k$ and so we are done. Note that $g_{125} \neq 0$ if $k = 1$, we assume now that $g_{125} = 0$ and $k \geq 2$.

Since $\deg_{y_{ia}} f = \deg_{y_{ia}} g$ for $a = 1, 2$ and $\deg_{y_{i3}} f > \deg_{y_{i3}} g$, we have the following formulas:

- $f_{134} + f_{123} + f_{125} = g_{134} + g_{123} + g_{125};$ ❶
- $f_{245} + f_{123} + f_{125} = g_{245} + g_{123} + g_{125};$ ❷
- $f_{134} + f_{123} + f_{345} > g_{134} + g_{123} + g_{345};$ ❸
- $f_{134} + f_{245} + f_{123} + f_{125} + f_{345} = g_{134} + g_{245} + g_{123} + g_{125} + g_{345} = k.$ ❹

Subtracting ❶ from ❹ and subtracting ❷ from ❹, we obtain

- $f_{245} + f_{345} = g_{245} + g_{345};$ ❺
- $f_{134} + f_{345} = g_{134} + g_{345}.$ ❻

On the other hand, by combining ❸ and ❶ with the assumption $g_{125} = 0$, we have

$$f_{134} + f_{123} + f_{345} > g_{134} + g_{123} + g_{345} = f_{134} + f_{123} + f_{125} + g_{345},$$

and so $f_{345} > f_{125} + g_{345} \geq g_{345}$. From this together with ❺ and ❻, it follows that $f_{134} < g_{134}$ and $f_{245} < g_{245}$. This particularly implies $g_{134} > 0$ and $g_{245} > 0$, and thus, there exist $p, q \in \{1, \dots, k\}$ with $p \neq q$ such that $\{y_{i2}, y_{i4}, y_{i5}\} \subseteq g_p$ and $\{y_{i1}, y_{i3}, y_{i4}\} \subseteq g_q$. Say $p = 1$ and $q = 2$. To prove $y_{i3}g/y_{i4}$ belongs to $J(G)^k$, we only need to prove that $y_{i3}g_1g_2/y_{i4}$ belongs to $J(G)^2$.

We may write g_1, g_2 as follows:

$$g_1 = y_{i2}y_{i4}y_{i5}u_1u_2 \quad \text{and} \quad g_2 = y_{i1}y_{i3}y_{i4}v_1v_2,$$

where $\text{supp}(u_i) \cup \text{supp}(v_i) \subseteq T_i$ for $i = 1, 2$. Then there are the following decomposition:

$$y_{i3}g_1g_2/y_{i4} = (y_{i3}y_{i4}y_{i5}u_1u_2)(y_{i1}y_{i2}y_{i3}v_1v_2) = (y_{i3}y_{i4}y_{i5}u_1v_2)(y_{i1}y_{i2}y_{i3}v_1u_2).$$

We next check that both $\text{supp}(y_{i3}y_{i4}y_{i5}u_1v_2)$ and $\text{supp}(y_{i1}y_{i2}y_{i3}v_1u_2)$ are vertex covers of G . To this end, we denote $V_1 = \text{supp}(y_{i3}y_{i4}y_{i5}u_1v_2)$ and $V_2 = \text{supp}(y_{i1}y_{i2}y_{i3}v_1u_2)$. Let $e = \{z_1, z_2\}$ be an edge of G .

If $e \subseteq V(G_i)$, then it is clear that $V_1 \cap e \neq \emptyset$ and $V_2 \cap e \neq \emptyset$.

If $z_1 = y_{i1}$ and $z_2 \in T_1$ then $z_2 \in \text{supp}(u_1) \cap e$. This implies $V_1 \cap e \neq \emptyset$ and $V_2 \cap e \neq \emptyset$; Similarly, we have if $z_1 = y_{i2}$ and $z_2 \in T_2$ then $V_1 \cap e \neq \emptyset$ and $V_2 \cap e \neq \emptyset$.

Finally suppose that $e \cap V(G_i) = \emptyset$. Then either $e \subseteq E(T_1)$ or $e \subseteq E(T_2)$. In the case that $e \subseteq E(T_1)$, we have $\text{supp}(u_1) \cap e \neq \emptyset$ and $\text{supp}(v_1) \cap e \neq \emptyset$ and so $V_1 \cap e \neq \emptyset$ and $V_2 \cap e \neq \emptyset$. The case that $e \subseteq E(T_2)$ can be proved similarly.

Thus, both V_1 and V_2 are vertex covers of G indeed and so $y_{i3}g_1g_2/y_{i4}$ belongs to $J(G)^2$, as required.

Case $z = y_{i4}$ or y_{i5} : Since the case where $z = y_{i5}$ is impossible actually, we only need to consider the case where $z = y_{i4}$. Moreover we may assume that $k \geq 2$ as in the case $z = y_{i3}$. Because $\deg_{y_{ia}} f = \deg_{y_{ia}} g$ for $a = 1, 2, 3$ and $\deg_{y_{i4}} f > \deg_{y_{i4}} g$, we have the following formulas:

- $f_{134} + f_{123} + f_{125} = g_{134} + g_{123} + g_{125};$ ❶
- $f_{245} + f_{123} + f_{125} = g_{245} + g_{123} + g_{125};$ ❷
- $f_{134} + f_{123} + f_{345} = g_{134} + g_{123} + g_{345};$ ❸
- $f_{134} + f_{245} + f_{345} > g_{134} + g_{245} + g_{345};$ ❹
- $f_{134} + f_{245} + f_{123} + f_{125} + f_{345} = g_{134} + g_{245} + g_{123} + g_{125} + g_{345} = k.$ ❺

Subtracting ❶, ❷ and ❸ from ❺ respectively, we have

- $f_{245} + f_{345} = g_{245} + g_{345};$ ❻
- $f_{134} + f_{345} = g_{134} + g_{345};$ ❼
- $f_{245} + f_{125} = g_{245} + g_{125}.$ ❽

We also obtain $f_{245} > g_{245}$ by substituting ❼ into ❹. From this together with ❻ and ❽, it follows that $f_{345} < g_{345}$ and $f_{125} < g_{125}$. In particular, $g_{345} > 0$ and $g_{125} > 0$. Consequently, there exist $p, q \in \{1, \dots, k\}$ such that $\{y_{i3}, y_{i4}, y_{i5}\} \subseteq g_p$ and $\{y_{i1}, y_{i2}, y_{i5}\} \subseteq g_q$. Say $p = 1$ and $q = 2$. It suffices to prove that the monomial $y_{i4}g_1g_2/y_{i5}$ belongs to $J(G)^2$.

For this, we decompose g_1 and g_2 as follows:

$$g_1 = y_{i3}y_{i4}y_{i5}u_1u_2 \quad \text{and} \quad g_2 = y_{i1}y_{i2}y_{i5}v_1v_2,$$

where $\text{supp}(u_i) \cup \text{supp}(v_i) \subseteq T_i$ for $i = 1, 2$. This implies

$$y_{i4}g_1g_2/y_{i5} = (y_{i3}y_{i4}y_{i5}u_1u_2)(y_{i1}y_{i2}y_{i4}v_1v_2) = (y_{i2}y_{i4}y_{i5}u_1v_2)(y_{i1}y_{i3}y_{i4}v_1u_2).$$

We can check that both $\text{supp}(y_{i2}y_{i4}y_{i5}u_1v_2)$ and $\text{supp}(y_{i1}y_{i3}y_{i4}v_1u_2)$ are vertex covers of G , as in the case where $z = y_{i3}$. Hence $y_{i4}g_1g_2/y_{i5}$ belongs to $J(G)^2$. This finishes the proof. □

Remark 1. The method used in our proof of this note depends heavily on the condition that every edge of G belongs to at most cycle of G . The gap of the original proof in [3] was found when we tried to use the idea in the proof to show that $J(G)^k$ is weakly polymatroidal for all $k \geq 1$ if G is a Cohen-Macaulay graph of large girth, whose structure was described explicitly in [1]. But we failed in this regard and we can only prove in [2, Proposition 3.6] that $J(G)$ is weakly polymatroidal for such graphs.

Declarations

- Funding: This work was supported by National Natural Science Foundation of China (Grant No. 11971338)
- The authors have no competing interests to declare that are relevant to the content of this article.
- We would like to express our sincere thanks to the referee, whose comments improved the presentation of the paper greatly.

References

- [1] Hoang, D.T., Minh, N.C., Trung, T.N., Cohen-Macaulay graphs with large girth. *J. Algebra Appl.* 14 (2015), 1550112
- [2] Lu, D., Wang, Z., On powers of cover ideals of graphs. to appear *Osaka Journal of mathematics*
- [3] Mohammadi, F., Powers of the vertex cover ideals, *Collect. Math.* 65, 169–181 (2014)