THE GRUNSKY NORM OF UNIVALENT FUNCTIONS AND ABELIAN HOLOMORPHIC DIFFERENTIALS

SAMUEL L. KRUSHKAL

ABSTRACT. We establish that the Grunsky norm of any normalized univalent function on the disk is completely determined by the squares of holomorphic abelian differentials (in contrast to the Teichmüller norm, which relates to all integrable holomorphic quadratic differentials).

This result has important interesting applications. In particular, it provides an explicit representation of Fredholm eigevalues of all quasiconformal curves.

2020 Mathematics Subject Classification: Primary: 30C55, 30C62, 30F45; Secondary 30F60, 32F45, 32G15, 46G20

Key words and phrases: Univalent function, Grunsky norm, quasiconformal extension, universal Teichmüller space, Teichmüller norm, hyperbolic metric, generalized Gaussian curvature, Minda's maximum principle

1. PRELIMINARIES

1.1. The Grunsky norm and its dominante. The classical Grunsky theorem of 1939 implies the necessary and sufficient conditions for univalence of holomorphic functions in a finitely connected domain on the Riemann sphere $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ in terms of an infinite system of the coefficient inequalities. In particular, for the canonical disk $\mathbb{D}^* = \{z \in \widehat{\mathbb{C}} : |z| > 1\}$ this theorem yields that a holomorphic function $f(z) = z + \text{const} + O(z^{-1})$ in a neighborhood of $z = \infty$ can be extended to a univalent holomorphic function on \mathbb{D}^* if and only if the Taylor coefficients α_{mn} of the function

$$\log \frac{f(z) - f(\zeta)}{z - \zeta} = -\sum_{m,n=1}^{\infty} \alpha_{mn} z^{-m} \zeta^{-n}, \quad (z,\zeta) \in (\mathbb{D}^*)^2,$$
(1)

called the **Grunsky coefficients** of f(z), satisfy the inequality

$$\left|\sum_{m,n=1}^{\infty} \sqrt{mn} \alpha_{mn} x_m x_n\right| \le 1,\tag{2}$$

for any sequence $\mathbf{x} = (x_n)$ from the unit sphere $S(l^2)$ of the Hilbert space l^2 with norm $\|\mathbf{x}\| = (\sum_{1}^{\infty} |x_n|^2)^{1/2}$; here the principal branch of the logarithmic function is chosen (cf. [6]).

Date: May 5, 2023 (GruAbel(2).tex).

The quantity

$$\varkappa(f) = \sup\left\{ \left| \sum_{m,n=1}^{\infty} \sqrt{mn} \, \alpha_{mn} x_m x_n \right| : \ \mathbf{x} = (x_n) \in S(l^2) \right\} \le 1$$

is called the **Grunsky norm** of f.

The univalent functions $f(z) = z + b_0 + b_1 z^{-1} + ...$ in \mathbb{D}^* admitting quasiconformal extensions across the unit circle $\mathbb{S}^1 = \partial \mathbb{D}^*$ onto the disk $\mathbb{D} = \{|z| < 1\}$ form the class Σ_Q . To have their uniqueness for a given Beltrami coefficient $\mu(z) = \partial_{\overline{z}} f/\partial_z f$ in \mathbb{D} , compactness in the topology of locally uniform convergence on \mathbb{C} , etc., we add the third normalization condition

f(0) = 0.

All such $f \in \Sigma_Q$ are zero free in \mathbb{D}^* , hence their inversions $F_f(z) = 1/f(1/z) = z + a_2 z^2 + \ldots$ are holomorphic and univalent in the disk \mathbb{D} with $F_f(\infty) = \infty$. The functions f and F_f have the same Grunsky coefficients, and $\varkappa(F_f) = \varkappa(f)$.

Note also that the norm $\varkappa(f)$ is defined for all $f \in \Sigma_Q$ and does not depend on the additional normalization at 0.

For the functions with k-quasiconformal extensions (k < 1), we have instead of (2) a stronger bound

$$\left|\sum_{m,n=1}^{\infty} \sqrt{mn} \alpha_{mn} x_m x_n\right| \le k \quad \text{for any} \ \mathbf{x} = (x_n) \in S(l^2), \tag{3}$$

established first in [15] (see also [12]).

Note that the Grunsky matrix operator $\mathcal{G}(f) = (\sqrt{mn} \ \alpha_{mn}(f))_{m,n=1}^{\infty}$ acts as a linear operator $l^2 \to l^2$ contracting the norms of elements $\mathbf{x} \in l^2$; the norm of this operator equals $\varkappa(f)$.

The method of Grunsky inequalities was generalized in several directions, with the corresponding generalization of Grunsky norm. In this paper continuing [13], we shall deal only with the canonical case of disk \mathbb{D}^* .

1.2. Some known results. The Grunsky norm $\varkappa(f)$ is dominated by the **Teichmüller norm** k(f), which is equal to the infimum of dilatations $k(w^{\mu}) = ||\mu||_{\infty}$ of quasiconformal extensions of f to $\widehat{\mathbb{C}}$. Here w^{μ} denotes a homeomorphic solution to the Beltrami equation $\partial_{\overline{z}}w = \mu \partial_z w$ on \mathbb{C} extending f; accordingly, μ is called the **Beltrami coefficient** (or complex dilatation) of w.

For most functions f, we have the strong inequality $\varkappa(f) < k(f)$ (moreover, the functions satisfying this inequality form a dense subset of Σ_Q), while the functions with the equal norms play a crucial role in many applications.

On the other hand, the important result of Pommerenke and Zhuravlev states that if a function $f \in \Sigma$ satisfies the inequality $\varkappa(f) < k$ with some constant k < 1, then f has a quasiconformal extension to $\widehat{\mathbb{C}}$ with a dilatation $k_1 = k_1(k) \ge k$ [20], [23]; [14], pp.82-84.

Each coefficient $\alpha_{mn}(f)$ in (4) is represented as a polynomial of a finite number of the initial coefficients b_1, b_2, \ldots, b_s of f; hence it depends holomorphically on Beltrami coefficients of quasiconformal extensions of f as well as on the **Schwarzian derivatives**

$$S_f(z) = \left(\frac{f''(z)}{f'(z)}\right)' - \frac{1}{2} \left(\frac{f''(z)}{f'(z)}\right)^2, \quad z \in \mathbb{D}^*.$$

These derivatives range over a bounded domain in the complex Banach space $\mathbf{B}(\mathbb{D}^*)$ of hyperbolically bounded holomorphic functions $\varphi \in \mathbb{D}^*$ with norm

$$\|\varphi\|_{\mathbf{B}} = \sup_{\mathbb{D}^*} (|z|^2 - 1)^2 |\varphi(z)|.$$

This domain models the **universal Teichmüller space** \mathbf{T} (the space of complex structures on the disk) in holomorphic Bers' embedding of \mathbf{T} . The complex geometry of this space provides a background of many results related to this paper.

The Beltrami coefficients of quasiconformal extensions w^{μ} of functions $f(z) \in \Sigma_Q$ range over the unit ball

Belt
$$(\mathbb{D})_1 = \{ \mu \in L_{\infty}(\mathbb{C}) : \ \mu(z) | \mathbb{D}^* = 0, \ \|\mu\|_{\infty} < 1 \},\$$

and the well-known criterion for extremality (the Hamilton-Krushkal-Reich-Strebel theorem) implies that a Betrami coefficient $\mu_0 \in \text{Belt}(\mathbb{D})_1$ is extremal if an only if

$$\|\mu_0\|_{\infty} = \sup_{\|\psi\|_{A_1(\mathbb{D})}=1} |\langle \mu, \psi \rangle_{\mathbb{D}}|,$$

where

$$\langle \mu, \psi \rangle_{\mathbb{D}} = \iint_{\mathbb{D}} \mu(z)\psi(z)dxdy \quad (z = x + iy),$$

and

 $A_1(\mathbb{D}) = \{ \psi \in L_1(D) : \psi \text{ holomorphic in } \mathbb{D} \}.$

The same condition is necessary and sufficient for the infinitesimal extremality of μ_0 at the origin of the space **T** in the direction $t\phi_{\mathbf{T}}(\mu_0)$, where $\phi_{\mathbf{T}}$ is the defining (factorizing) holomorphic projection $\text{Belt}(\mathbb{D})_1 \to \mathbf{T}$; see, e.g., [3], [4], [7].

The Grunsky norm is connected with the subset of squares of the abelian holomorphic differentials $\omega(z)dz$ on \mathbb{D} :

$$A_1^2(\mathbb{D}) = \{ \psi = \omega^2 \in A_1(\mathbb{D}) : \omega \text{ holomorphic in } \mathbb{D} \}$$

formed by integrable squares of squares of the abelian holomorphic differentials $\omega(z)dz$. Due to [8] (and to the more general result in [12]), the equality $\varkappa(f^{\mu}) = k(f^{\mu})$ is valid if and only if

$$\|\mu\|_{\infty} = \sup_{\psi \in A_1^2(\mathbb{D}), \|\psi\|_{A_1} = 1} |\langle \mu, \psi \rangle_{\mathbb{D}}|;$$

in addition, if μ is of Teichmüller form, i.e., $\mu = k|\psi|/\psi$ with $\psi \in A_1(\mathbb{D})$, then necessarily $\psi \in A_1^2$.

Note also that, due to [8], the elements of $A_1^2(\mathbb{D})$ are represented in the form

$$\psi(z) = \omega(z)^2 = \frac{1}{\pi} \sum_{m+n=2}^{\infty} \sqrt{mn} x_m x_n z^{m+n-2},$$

with $\|\mathbf{x}\|_{l^2} = \|\omega\|_{L_2}$; here $\mathbf{x} = (x_n)$.

1.3. A lower bound for Grunsky norm. The following essential estimate established in [13] is one of the crucial steps in the proof of our main theorem.

Lemma 1. The Grunsky norm $\varkappa(f)$ of every function

$$f(z) = z + b_0 + b_m z^{-m} + b_{m+1} z^{-(m+1)} + \dots \in \Sigma_Q, \quad m \ge 1$$

satisfies the inequality

$$\varkappa(f) \ge \alpha_{\mathbb{D}}(f) := \sup_{\psi \in A_1^2(\mathbb{D}), \|\psi\|_{A_1} = 1} |\langle \mu_0, \psi \rangle_{\mathbb{D}}| = \sup_{\omega \in A_2(\mathbb{D}), \|\omega\|_2 = 1} |\langle \mu_0, \omega^2 \rangle_{\mathbb{D}}|, \tag{4}$$

where μ_0 is an extremal Beltrami coefficient among quasiconformal extensions f^{μ} of f onto \mathbb{D} .

For functions f with equal Grunsky and Teichmüller norms, we have in (4) the equality, and vice versa (see, [8], [12], [17]), but until now there was unknown whether there exist the functions $f \in \Sigma_Q$ with $\varkappa(f) < k(f)$ and $\varkappa(f) = \alpha_{\mathbb{D}}(f)$.

The proof of this lower bound for $\varkappa(f)$ is based on geometric extensions of the Ahlfors Schwarz lemma to metrics of integrable generalized negative curvatures given in [10], [21].

We shall also use the infinitesimal version of the inequality (4). It will be presented later.

1.4. Conformal metrics of negative generalized Gaussian curvature. We shall use some known results on subharmonic conformal metrics $ds = \lambda(t)|dt|$ on the disk \mathbb{D} (with $\lambda(t) \geq 0$) of negative generalized Gaussian curvature.

Recall that the **generalized Gaussian curvature** κ_{λ} of an upper semicontinuous Finsler metric $ds = \lambda |dt|$ in a domain $\Omega \subset \mathbb{C}$ is defined by

$$\kappa_{\lambda}(t) = -\frac{\Delta \log \lambda(t)}{\lambda(t)^2},\tag{5}$$

where Δ is the generalized Laplacian

$$\Delta\lambda(t) = 4\liminf_{r\to 0} \frac{1}{r^2} \left\{ \frac{1}{2\pi} \int_0^{2\pi} \lambda(t + re^{i\theta}) d\theta - \lambda(t) \right\}$$

(provided that $-\infty \leq \lambda(t) < \infty$). Similar to C^2 functions, for which Δ coincides with the usual Laplacian, one obtains that λ is subharmonic on Ω if and only if $\Delta\lambda(t) \geq 0$; hence, at the points t_0 of local maximum of λ with $\lambda(t_0) > -\infty$, we have $\Delta\lambda(t_0) \leq 0$.

The sectional *holomorphic curvature* of a Finsler metric on a complex Banach manifold X is defined in a similar way as the supremum of the curvatures (5) over appropriate collections of holomorphic maps from the disk into X for a given tangent direction in the image.

As is well-known [1], [9], the holomorphic curvature of the Kobayashi-Teichmüller metric $\mathcal{K}_{\mathbf{T}}(x, v)$ of universal Teichmüller space \mathbf{T} equals -4 at all points (x, v) of the tangent bundle $\mathcal{T}(\mathbf{T})$ over \mathbf{T} . Instead, the holomorphic curvature of metric λ_{\varkappa} generated on \mathbb{D} by the Grunsky Finsler structure satisfies the inequality $\Delta \log \lambda \geq 4\lambda^2$, where Δ is again the generalized Laplacian (see [11]). So, $\lambda_{\varkappa}(t) \leq -4$ on any holomorphic disk in the space \mathbf{T} .

This is a special case of metrics whose curvature satisfies the inequality $\Delta \log \lambda \geq K \lambda^2$ with constant K > 0. All such metrics are subharmonic.

We shall also apply the upper semicontinuous metrics λ satisfying the inequality

$$\Delta \log \lambda \ge K\lambda, \quad K = \text{const} > 0 \tag{6}$$

(then $u = \log \lambda$ can be negative). For such metrics, we have the following Minda's maximum principle given by

Lemma 2. If a function $u: D \to [-\infty, +\infty)$ is upper semicontinuous in a domain $D \subset \mathbb{C}$ and its generalized Laplacian satisfies the inequality $\Delta u(z) \geq Ku(z)$ with some positive constant K at any point $z \in D$, where $u(z) > -\infty$, and if

$$\limsup_{z \to \zeta} u(z) \le 0 \quad \text{for all } \zeta \in \partial D,$$

then either u(z) < 0 for all $z \in D$ or $u(z) \equiv 0$ on D.

The proof of this lemma related to the Ahlfors-Schwarz lemma is given in [19]; its variations see in [2].

2. MAIN THEOREM AND ITS APPLICATIONS

2.1. Main theorem. The following theorem implies a complete answer on relation between the Grunsky norm and abelian holomorphic differentials.

Theorem 1. For any univalent function we have the equality

$$\varkappa(f) = \alpha_{\mathbb{D}}(f). \tag{7}$$

It suffices to consider the functions $f \in \Sigma_Q$, because for functions not admitting quasiconformal extension, both sides of (7) are equal to 1.

2.2. Applications. We mention two important applications of Theorem 1.

1. Plurisubharmonicity. Letting $\alpha_{\mathbb{D}}(S_{f^{\mu}}) = \alpha_{\mathbb{D}}(f^{\mu})$, one can regard this quantity as a function on the universal Teichmüller space **T** (as well as on the ball Belt(\mathbb{D})₁. The following corollary to Theorem 1 solves the question posed in [10].

Corollary 1. The function $\alpha_{\mathbb{D}}(S_{f^{\mu}})$ is plurisubharmonic on the space **T** and on the ball $\operatorname{Belt}(\mathbb{D})_1$.

This follows from the equality (7), because its left-hand side is plurisubharmonic on \mathbf{T} . In fact, the indicated plurisubharmonicity is a consequence of the relation (10) and of holomorphy of functions (9) defined below. Another proof is given in [12].

2. Application to Fredholm eigenvalues of Jordan curves. The Fredholm eigenvalues ρ_n of an oriented smooth closed Jordan curve $L \subset \widehat{\mathbb{C}}$ are the eigenvalues of its double-layer potential, or equivalently, of the integral equation

$$u(z) + \frac{\rho}{\pi} \int_{L} u(\zeta) \frac{\partial}{\partial n_{\zeta}} \log \frac{1}{|\zeta - z|} ds_{\zeta} = h(z),$$

where n_{ζ} denotes the outer normal and ds_{ζ} is the length element at $\zeta \in L$. These values have crucial applications in solving many problems in various fields of mathematics.

It is important often to know the least positive eigenvalue $\rho_L = \rho_1$. This value is naturally connected with conformal and quasiconformal maps related to L and can be defined for any oriented closed Jordan curve L by

$$\frac{1}{\rho_L} = \sup \frac{|\mathcal{D}_G(u) - \mathcal{D}_{G^*}(u)|}{\mathcal{D}_G(u) + \mathcal{D}_{G^*}(u)},$$

where G and G^* are, respectively, the interior and exterior of L; \mathcal{D} denotes the Dirichlet integral, and the supremum is taken over all functions u continuous on $\widehat{\mathbb{C}}$ and harmonic on $G \cup G^*$. In particular, $\rho_L = \infty$ only for the circle.

Note that both sides of the last equality remain invariant under the action of the Moebius group $PSL(2, \widehat{\mathbb{C}})$.

The indicated first eigenvalue is intrinsically connected with the Grunsky coefficients of the exterior conformal mapping function $f_L^*: \mathbb{D}^* \to D^*$; this is qualitatively expressed by the remarkable Kühnau-Schiffer theorem, which states that the value ρ_L is reciprocal to the Grunsky norm $\varkappa(f_L^*)$ (see [16], [22]).

Together with this result, Theorem 1 implies explicitly the first Fredholm eigenvalue ρ_L of every quasiconformal curve $L \subset \widehat{\mathbb{C}}$. Namely, we have

Corollary 2. For any quasicircle $L \subset \widehat{\mathbb{C}}$,

$$\frac{1}{\rho_L} = \sup_{\psi \in A_1^2(D), \|\psi\|_{A_1} = 1} |\langle \mu_0, \psi \rangle_{\mathbb{D}}|,$$
(8)

where μ_0 is either of extremal Beltrami coefficients of the appropriately normalized exterior conformal mapping function f_L^* on which the Teichmüller norm of f_L^* is attained.

Moreover, letting in (8) $\mu_0 = t |\psi|/\psi$ with |t| < 1 and $\psi \in A_1$, one obtains a dense subset of a possible eigenvalues ρ_L .

Conversely, given a quasicircle $L \subset \widehat{\mathbb{C}}$, take a quasiconformal extension f^{μ} of the exterior conformal map f_L^* . Its Beltrami coefficient μ determines a linear functional $\mu(\psi) = \langle \mu, \psi \rangle_{\mathbb{D}}$ on the linear span \widetilde{A} of the subset $\{\psi \in A_1^2(\mathbb{D}) : \|\psi\|_{A_1} = 1\}$ of the unit sphere in $A_1(\mathbb{D})$. Its Hahn-Banach extension onto $L_1(\mathbb{D})$ provides an extremal Beltrami coefficient μ_0 in \mathbb{D} for f_L^* satisfying (8).

3. Teichmüller maps with $\varkappa(f) = k(f)$. We also mention the following useful property of Teichmüller extremal maps.

Corollary 3. If the equivalence class of f is a Strebel point and $\varkappa(f) = k(f)$, then μ_0 is necessarily of the form $\mu_0(z) = \|\mu_0\|_{\infty} |\psi_0(z)| / \psi_0(z)$ with $\psi_0 \in A_1^2(\mathbb{D})$, and the relation (8) is completed by

$$\varkappa(f^{\mu_0}) = k(f^{\mu_0}) = q_L = \frac{1}{\rho_L} = \sup_{\psi \in A_1^2(D), \|\psi\|_{A_1} = 1} |\langle \mu_0, \psi \rangle_{\mathbb{D}}|,$$

where $L = f^{\mu_0}(\mathbb{S}^1)$, and q_L denotes its quasireflection coefficient. (equal to the minimum of dilatations $\partial_z W/\partial -\overline{z}W$ of the orientation reversing quasiconformal homeomorphisms of the sphere $\widehat{\mathbb{C}}$ which preserve point-wise the quasicircle $L \subset \widehat{\mathbb{C}}$ and interchange its interior and exterior domains.

This was established by the author earlier even for univalent functions in the arbitrary quasidisks (see, e.g., [12] (and in a special case by Kühnau [17]), but it follows from (7) immediately.

Recall that quasireflection coefficient of an (oriented) quasicircle $L' \subset \widehat{\mathbb{C}}$ is the minimumal dilatation $\partial_z W/\partial -\overline{z}W$ of the orientation reversing quasiconformal homeomorphisms of the sphere $\widehat{\mathbb{C}}$, which preserve point-wise the quasicircle $L \subset \widehat{\mathbb{C}}$ and interchange its interior and exterior domains.

Theorem 1 also has other important consequences and applications. These results will be presented elsewhere.

3. PROOF OF THEOREM 1

1⁰. First observe that the Grunsky coefficients $\alpha_{mn}(f^{\mu})$ of functions $f^{\mu} \in \Sigma_Q$ generate for each $\mathbf{x} = (x_n) \in l^2$ with $\|\mathbf{x}\| = 1$ the holomorphic maps

$$h_{\mathbf{x}}(f^{\mu}) = \sum_{m,n=1}^{\infty} \alpha_{mn}(f^{\mu}) x_m x_n : \text{ Belt}(D)_1 \to \mathbb{D},$$
(9)

and

$$\sup_{\mathbf{x}} |h_{\mathbf{x}}(f^{\mu})| = \varkappa_{D^*}(f^{\mu}).$$
(10)

The holomorphy of these functions follows from the holomorphy of coefficients α_{mn} with respect to Beltrami coefficients $\mu \in Belt(D)_1$ mentioned above, by applying the estimate

$$\left|\sum_{m=j}^{M}\sum_{n=l}^{N} \beta_{mn} x_m x_n\right|^2 \le \sum_{m=j}^{M} |x_m|^2 \sum_{n=l}^{N} |x_n|^2 \tag{11}$$

which holds for any finite M, N and $1 \le j \le M$, $1 \le l \le N$ (see [20], p. 61).

Similar arguments imply that the maps (9) regarded as functions of points $\varphi^{\mu} = S_{f^{\mu}}$ in the universal Teichmüller space **T** are holomorphic on **T**. This holomorphy provides, together with the equality (10), that the Grunsky norm \varkappa_{D^*} regarded as a function of the Schwarzians S_f is logarithmically plurisubharmonic on the space **T**. In addition, as it is established in [12], the functions $\varkappa_{D^*}(S_f)$ is continuous; moreover, it is Lipschitz continuous on this space. The Teichmüller norm has the similar properties.

Now, given a function $f \in \Sigma_Q$, take its extremal extension f^{μ} (i.e., such that $k(f) = \|\mu\|_{\infty}$) and consider its extremal disk

$$\mathbb{D}(\mu) = \{t\mu/\|\mu\|_{\infty} : |t| < 1\} \subset \operatorname{Belt}(\mathbb{D})_1.$$

Put $\mu^* = \mu / \|\mu\|_{\infty}$.

Using the functions (9), we pull back the hyperbolic metric $\lambda_{\mathbb{D}}(t)|dt| = |dt|/(1-|t|^2)$ of the disk \mathbb{D} , which implies on this disk the conformal metrics $\lambda_{h_x}(t)|dt|$ with

$$\lambda_{h_{\mathbf{x}}}(t) = |h'_{\mathbf{x}}(t)|/(1 - |h_{\mathbf{x}}(t)|^2).$$

All these metric have at their noncrical points the Gaussian curvature -4. We take the upper envelope of these metrics

$$\lambda_{\varkappa}(t) = \sup\{\lambda_{h_{\mathbf{x}}}(t) : \mathbf{x} \in S(l^2)\}$$

followed by its upper semicontinuous regularization, which determines a logarithmically subharmonic metric $\lambda_{\varkappa}(t)$ on the unit disk. It is circularly symmetric, i.e., $\lambda_{\varkappa}(t) = \lambda_{\varkappa}(|t|)$. This metric being generated by the Grunsky coefficients of f actually is the differential (infinitesimal) form of the norm $\varkappa(f)$.

On a standard way, one obtains that λ_{\varkappa} has at any its noncritical point t_0 a supporting subharmonic metric λ_0 of Gaussian curvature at most -4, and hence, $\kappa_{\lambda_{\varkappa}} \leq -4$; the details see, e.g., in [9].

It is shown in [13], that if the initial function $f \in \Sigma_Q$ has in \mathbb{D}^* the expansion

$$f(z) = z + b_n z^n + b_{n+1} z^{-n-1} + \dots, \quad b_n \neq 0, \quad n \ge 1.$$

then the corresponding metric λ_{\varkappa} satisfies the relation

$$\lim_{r \to 0} \frac{\lambda_{\varkappa}(r)}{r^n} = \sup_{\mathbf{x} \in S(l^2)} \frac{|dh_{\mathbf{x}}(0)(t\mu_0/\|\mu_0\|_{\infty}|)}{|t|} \ge \sup_{\psi \in A_1^2(\mathbb{D}), \|\psi\|_{A_1} = 1} \left| \iint_{\mathbb{D}} \frac{\mu_0(z)}{\|\mu_0\|_{\infty}} \psi(z) dx dy \right|, \quad (12)$$

which is an infinitesimal version of (4).

2⁰. To find an upper bound for $\varkappa(f)$ (smaller that k(f)), we apply to functions $h_{\mathbf{x}}(f^{t\mu^*})$ a special case of Golusin's improvement [5] of the classical Schwarz lemma given by the following

Lemma 3. A holomorphic function

$$g(t) = c_m t^m + c_{m+1} t^{m+1} + \dots : \mathbb{D} \to \mathbb{D} \quad (c_m \neq 0, \quad m \ge 1),$$

in \mathbb{D} is estimated by

$$|g(t)| \le |t|^m \frac{|t| + |c_m|}{1 + |c_m||t|},\tag{13}$$

with equality only for the function

$$g_m(t) = t^m \frac{t + c_m}{1 + \overline{c}_m t} = c_m t^m + \dots$$
(14)

The corresponding differential metrics are related by

$$g^* \lambda_{\mathbb{D}}(r) \le \lambda_j(r) := \frac{j'(r)}{1 - j(r)^2}$$
(15)

where $r = |t| < 1, \lambda_{\mathbb{D}}(r) = 1/(1-r^2)$ is the hyperbolic metric of the unit disk of curvature -4, and

$$j(r) = r^m \frac{r + |c_m|}{1 + r|c_m|}.$$

Note that the pull-backed metric $\lambda_j(r)$ has the constant curvature -4 in the punctured disk $\{0 < |t| < 1\}$ and in view of (10) any function (9) is dominated by (14) with $|c_m| \ge \alpha_{\mathbb{D}}(f)$.

3⁰. To calculate explicitly the quantity $\alpha(f)$ given by the right hand side of (4), one can use the variational formula for the functions $f^{\mu}(z) = z + b_0 + b_1 z^{-1} + \ldots$ from Σ^0 with extensions satisfying $f^{\mu}(0) = 0$ (cf [13]). Namely, for small $\|\mu\|_{\infty}$,

$$f^{\mu}(z) = z - \frac{1}{\pi} \iint_{\mathbb{D}} \mu(w) \left(\frac{1}{\zeta - z} - \frac{1}{\zeta} \right) d\xi d\eta + O(\|\mu^2\|_{\infty}), \quad \zeta = \xi + i\eta,$$

where the ratio $O(\|\mu^2\|_{\infty}^2)/\|\mu^2\|_{\infty}^2$ is uniformly bounded on compact sets of \mathbb{C} . Then

$$b_n = \frac{1}{\pi} \iint_{\mathbb{D}} \mu(\zeta) \zeta^{n-1} d\xi d\eta + O(\|\mu^2\|_{\infty}), \quad n = 1, 2, \dots,$$

and

$$\alpha_{mn}(\mu) = -\pi^{-1} \iint_{\mathbb{D}} \mu(z) z^{m+n-2} dx dy + O(\|\mu\|_{\infty}^2), \quad \|\mu\|_{\infty} \to 0.$$

Using the normalized coefficient $\mu^* = \mu/\|\mu\|_{\infty}$ and passing accordingly to coefficients $t\mu^*$ with |t| < 1, one derives that the differential at zero of the corresponding map $h_{\mathbf{x}}(t\mu^*)$ with

 $\mathbf{x} = (x_n) \in S(l^2)$ along the extremal disk $\{t\mu^* \in \text{Belt}(\mathbb{D})_1\}$: $|t| < 1\}$ (and its image in the space **T**) is given by

$$dh_{\mathbf{x}}(0)(t\mu^*) = -\frac{t}{\pi} \iint_{\mathbb{D}} \mu^*(z) \sum_{m+n=2}^{\infty} \sqrt{mn} \ x_m x_n z^{m+n-2} dx dy$$

On the other hand, as was established in [7], the elements of $A_1^2(\mathbb{D})$ are represented in the form

$$\psi(z) = \omega(z)^2 = \frac{1}{\pi} \sum_{m+n=2}^{\infty} \sqrt{mn} x_m x_n z^{m+n-2} \quad \text{with} \quad \|\mathbf{x}\|_{l^2} = \|\omega\|_{L_2}.$$

Now, letting |t| < 1 and applying the relation (4) and Lemma 3 to $h_{\mathbf{x}}(t\mu^*)$, one obtains

$$|h_{\mathbf{x}}(t\mu^*)| \le |t| \frac{|t| + |\langle \mu^*, \psi \rangle_{\mathbb{D}}|}{1 + |\langle \mu^*, \psi \rangle_{\mathbb{D}}||t|}$$

and, passing to the supremum over $\mathbf{x} \in S(l^2)$,

$$\varkappa(f^{t\mu^*}) \le |t| \frac{|t| + \alpha(\mu^*)}{1 + \alpha(\mu^*)|t|},\tag{16}$$

with

$$\alpha(\mu^*) = \sup_{\psi \in A_1^2(\mathbb{D}), \|\psi\|_{A_1} = 1} |\langle \mu^*, \psi \rangle_{\mathbb{D}}| =: \alpha.$$

The relations (4), (12), (16) result in

$$\alpha|t| \le \varkappa(f^{t\mu^*}) \le |t| \frac{|t| + \alpha}{1 + \alpha|t|}.$$
(17)

4⁰. Now we may prove the theorem. We first establish the relation between the differential (infinitesimal) metric λ_{\varkappa} corresponding to $\varkappa(f)$ and its dominant of type (15) defined in accordance to Lemma 3.

Noting that the right hand side fraction in (13) is increasing with respect to $|c_m|$ on [0, 1], we take the functions

$$g_a(t) = t^m (t+a)/(1+\overline{a}t)$$

with $a > \alpha$ and define by (15) the corresponding metrics

$$\lambda_a(r) = j'_a(r)/(1 - j_a(r)^2)$$
(18)

(all of curvature -4). We compare these metrics with metric λ_{\varkappa} generated by the Grunsky coefficients of f^{μ} , as was described in the first step.

Comparison of λ_{\varkappa} with any metric (18) (which is maximal for a given value of $c_m = a$) provides by (12) the strict inequality

$$\lambda_{\varkappa}(r) < \lambda_a(r), \quad r < 1,$$

(for any $a > \alpha$). In the limit as $a \to \alpha$, one obtains

$$\lambda_{\varkappa}(r) \le \lambda_{\alpha}(r) = \alpha + (1 - \alpha^2)r + \dots \quad r < 1.$$
⁽¹⁹⁾

On the other hand, as was establishes in [13], we have at the origin t = 0 the equality

$$\lambda_{\varkappa}(0) = \lambda_{\alpha}(0)$$

where

$$\widetilde{\lambda}_{\alpha}(r) = \alpha/(1 - \alpha^2 r^2) = \alpha + O(r^2), \quad r \to 0,$$

is a supporting conformal metric for λ_{\varkappa} at the origin with constant Gaussian curvature -4. Therefore, for small r,

$$\alpha \le \lambda_{\varkappa}(r) \le \lambda_{\alpha}(r) \le \alpha + (1 - \alpha^2)r + O(r^2).$$
⁽²⁰⁾

Now choose a sufficiently small neighborhood U_0 of the origin t = 0 and put

$$M = \{ \sup \lambda_{\alpha}(t) : t \in U_0 \}.$$

Then in this neighborhood, we have $\lambda_{\alpha}(t) + \lambda_{\varkappa}(t) \leq 2M$. Consider the function

$$u = \log \frac{\lambda_{\varkappa}}{\lambda_{\alpha}}.$$

Then (cf. [19], [2]) for $t \in U_0$,

$$\Delta u(t) = \Delta \log \lambda_{\varkappa}(t) - \Delta \log \lambda_{\alpha}(t) \ge 4(\lambda_{\varkappa}^2 - \lambda_{\alpha}^2) \ge 8M(\lambda_{\varkappa} - \lambda_{\alpha}).$$

The elementary estimate

$$M \log(t/s) \ge t - s$$
 for $0 < s \le t < M$

(with equality only for t = s) implies that

$$M\log\frac{\lambda_{\alpha}(t)}{\lambda_{\varkappa}(t)} \ge \lambda_{\alpha}(t) - \lambda_{\varkappa}(t),$$

and hence,

$$\Delta u(t) \ge 8M^2 u(t).$$

It follows that the function u satisfies on U_0 the inequality (6) with $K = 8M^2$.

Applying to u Lemma 2 and noting that (20) yields

$$u(0) = \lim_{t \to 0} \log \frac{\lambda_{\varkappa}(t)}{\lambda_{\alpha}(t)} = 0$$

one derives by this lemma that both metrics λ_{\varkappa} and λ_{α} must be equal on U_0 , and in the similar way their equality on the entire disk \mathbb{D} .

4⁰. Finally, we apply the following reconstruction lemma for the Grunsky norm proven in [9], which provides that this norm is the integrated form of λ_{\varkappa} along the Teichmüller extremal disks.

Lemma 4. On any Teichmüller extremal disk $\mathbb{D}(\mu_0) = \{t\mu_0/||\mu_0||_{\infty} : |t| < 1\} \subset \text{Belt}(\mathbb{D})_1$, we have the equality

$$\tanh^{-1}[\varkappa(f^{r\mu_0/\|\mu_0\|_{\infty}})] = \int_0^r \lambda_\varkappa(t) dt.$$
(21)

Integrating the metrics λ_{\varkappa} and λ_{α} along the indicated extremal disks, one obtains from (21) and the right equality in (4) the required equality (7), completing the proof of the theorem.

4. EXAMPLE

The simplest example illustrating Theorem 1 is given by the map

$$f_{3,t}(z) = z \left(1 + \frac{t}{z^3}\right)^{2/3}, \quad 0 \le |t| < 1,$$
(22)

whose extremal extension to \mathbb{D} is

$$\widehat{f}_{3,t}(z) = z \left[1 + t \left(\frac{|z|}{z} \right)^3 \right]^{2/3}$$

with Beltrami coefficient $\mu_3(z) := \mu_{\hat{f}_{3,t}}(z) = t|z|/z$. In polar coordinates, $\mu_3(re^{i\theta}) = te^{-i\theta}$. This map has threefold rotational symmetry

$$f(e^{2n\pi i/3}z) = e^{2n\pi i/3}f(z), \quad n = 0, 1, 2$$

Note that $f_{3,t}(z) = f_t(z^{3/2})^{2/3}$, where

$$f_t(z) = \begin{cases} z + t/z, & |z| \ge 1, \\ z + t\overline{z}, & |z| < 1. \end{cases}$$

For this function, we have (see [10])

$$\sup_{\psi \in A_1^2, \|\psi\|_{A_1}=1} |\langle \mu_3^*, \psi \rangle_{\mathbb{D}}| = \max_{\psi_3} |\langle \mu_3^*, \psi_3 \rangle_{\mathbb{D}}| = \frac{2\sqrt{2}}{3},$$
(23)

where ψ_3 are the squares of nonconstant linear functions

$$\psi_3(z) = \omega_1(z)^2 := (a_0 + a_1 z)^2$$

with $a_0 \neq 0$, $a_1 \neq 0$, and $\|\psi_3\|_{A_1} = 1$.

The map (22) was the first explicit example of functions $f \in \Sigma_Q$ with $\varkappa(f) < k(f)$. This was established in 1981 independently by Kühnau in [16], applying the technique of Fredholm eigenvalues, and by the author in a different way (this also follows from the general result mentioned above). The quantity (23) provides the value of reciprocal to eigenvalue $\rho_{f_{3,t}}(\mathbb{S}^1)$.

More generally, all functions $f_m(z) = f_t(z^{m/2})^{2/m}$ with odd m = 2p - 1 have similar features.

References

- M. Abate and G. Patrizio, *Isometries of the Teichmüller metric*, Ann. Scuola Super. Pisa Cl. Sci.(4) 26 (1998), 437-452.
- [2] S. Dineen, The Schwarz Lemma, Clarendon Press, Oxford, 1989.
- [3] C.J. Earle, I. Kra and S.L. Krushkal, Holomorphic motions and Teichmüller spaces, Trans. Amer. Math. Soc. 944 (1994), 927-948.
- [4] F.P. Gardiner and N. Lakic, *Quasiconformal Teichmüller Theory*, Amer. Math. Soc., Providence, RI, 2000.
- [5] G.M. Goluzin, Geometric Theory of Functions of Complex Variables, Transl. of Math. Monographs, vol. 26, Amer. Math. Soc., Providence, RI, 1969.
- [6] H. Grunsky, Koeffizientenbedingungen f
 ür schlicht abbildende meromorphe Funktionen, Math. Z. 45 (1939), 29-61.
- [7] S.L. Krushkal, Quasiconformal Mappings and Riemann Surfaces, Wiley, New York, 1979.
- [8] S. L. Krushkal, Grunsky coefficient inequalities, Carathéodory metric and extremal quasiconformal mappings, Comment. Math. Helv. 64 (1989), 650-660.
- [9] S.L. Krushkal, Plurisubharmonic features of the Teichmüller metric, Publications de l'Institut Mathématique-Beograd, Nouvelle série 75(89) (2004), 119-138.
- [10] S.L. Krushkal, Strengthened Moser's conjecture, geometry of Grunsky coefficients and Fredholm eigenvalues, Central European J. Math. 5(3) (2007), 551-580.

- [11] S.L. Krushkal, Proof of the Zalcman conjecture for initial coefficients, Georgian Math. J. 17 (2010), 663-681; Corrigendum 19 (2012), 777.
- [12] S.L. Krushkal, Strengthened Grunsky and Milin inequalities, Contemp. Mathematics 667 (2016), 159-179.
- [13] S.L. Krushkal, On Grunsky norm of univalent functions, J. Math. Sci. (2023), to appear.
- [14] S.L. Krushkal and R. Kühnau, Quasikonforme Abbildungen neue Methoden und Anwendungen, Teubner-Texte zur Math., Bd. 54, Teubner, Leipzig, 1983.
- [15] R. Kühnau, Verzerrungssätze und Koeffizientenbedingungen vom Grunskyschen Typ für quasikonforme Abbildungen, Math. Nachr. 48 (1971), 77-105.
- [16] R. Kühnau, Quasikonforme Fortsetzbarkeit, Fredholmsche Eigenwerte und Grunskysche Koeffizientenbedingungen, Ann. Acad. Sci. Fenn. Ser. AI. Math. 7 (1982), 383-391.
- [17] R. Kühnau, Wann sind die Grunskyschen Koeffizientenbedingungen hinreichend für Qquasikonforme Fortsetzbarkeit?, Comment. Math. Helv. 61 (1986), 290-307.
- [18] O. Lehto, Univalent Functions and Teichmüller Spaces, Springer-Verlag, New York, 1987.
- [19] D. Minda, The strong form of Ahlfors' lemma, Rocky Mountain J. Math., 17 (1987), 457-461.
- [20] Chr. Pommerenke, Univalent Functions, Vandenhoeck & Ruprecht, Göttingen, 1975.
- [21] H.L. Royden, The Ahlfors-Schwarz lemma: the case of equality, J. Anal. Math. 46 (1986), 261-270.
- [22] M. Schiffer, Fredholm eigenvalues and Grunsky matrices, Ann. Polon. Math. 39 (1981), 149-164.
- [23] I.V. Zhuravlev, Univalent functions and Teichmüller spaces, Inst. of Mathematics, Novosibirsk, preprint, 1979, 23 pp. (Russian).

Department of Mathematics, Bar-Ilan University, 5290002 Ramat-Gan, Israel and Department of Mathematics, University of Virginia, Charlottesville, VA 22904-4137, USA