

A Kannappan-cosine functional equation on semigroups

Jafar Ahmed¹, Ajebbar Omar², Elqorachi Elhoucien¹

Abstract. In this paper we determine the complex-valued solutions of the Kannappan cosine functional law $g(xyz_0) = g(x)g(y) - f(x)f(y)$, $x, y \in S$, where S is a semigroup and z_0 is a fixed element in S .

1. Introduction

The addition law for cosine is

$$\cos(x + y) = \cos(x)\cos(y) - \sin(x)\sin(y), \quad x, y \in \mathbb{R}.$$

It is at the origin of the following functional equation on any semigroup S :

$$g(xy) = g(x)g(y) - f(x)f(y), \quad x, y \in S \tag{1.1}$$

for the unknown functions $f, g : S \rightarrow \mathbb{C}$, which is called cosine addition law. In Aczèl's monograph [1, Section 3.2.3] we find the early results about (1.1) and its continuous real valued solutions for $S = \mathbb{R}$.

The functional equation (1.1) has been solved on groups by Poulsen and Stetkær ([8]), on semigroups generated by their squares by Ajebbar and Elqorachi ([2]), and recently by Ebanks [6] on semigroups

In [12, Theorem 3.1] Stetkær solved the following functional equation

$$g(xy) = g(x)g(y) - f(y)f(x) + \alpha f(xy), \quad x, y \in S, \tag{1.2}$$

where α is a fixed constant in \mathbb{C} . He expressed the solutions in terms of multiplicative functions and the solution of the special case of sine addition law. In [11, Proposition 16] he solved the functional equation

$$f(xyz_0) = f(x)f(y), \quad x, y \in S, \tag{1.3}$$

2020 Mathematics Subject Classification: Primary 39B52, Secondary 39B32.

Keywords: Functional equations, Semigroups, Multiplicative function, Additive function, Cosine-Sine equation.

on semigroups, and where z_0 is a fixed element in S . We shall use these results in our computations.

In this paper we deal with the following Kannappan-cosine addition Law

$$g(xyz_0) = g(x)g(y) - f(x)f(y), \quad x, y \in S \quad (1.4)$$

on a semigroup S . The functional equation (1.4) is called Kannappan functional equation because it brings up a fixed element z_0 in S as in the paper of Kannappan [10].

In the special case where $\{f, g\}$ is linearly dependent and $g \neq 0$ we get that there exists a constant $\lambda \in \mathbb{C}$ such that the function $(1 - \lambda^2)g$ satisfies the functional equation (1.3).

If S is a monoid with identity element e , and $f(e) = 0$ and $g(e) \neq 0$, or $g(e) = 0$ and $f(e) \neq 0$, the last functional equation is the cosine addition law which was solved recently on general semigroups by Ebanks [6].

Now, if $\alpha := f(e) \neq 0$ and $\beta := g(e) \neq 0$ we get that the pair $\left(\frac{g}{\beta}, \frac{f}{\beta}\right)$ satisfies the following functional equation

$$\frac{g}{\beta}(xy) = \frac{g}{\beta}(x)\frac{g}{\beta}(y) - \frac{f}{\beta}(x)\frac{f}{\beta}(y) + \frac{\alpha}{\beta}\frac{f}{\beta}(xy),$$

which is of the form (1.2), and then explicit formulas for f and g on groups exist in the literature (see for example [14, Corollary 3.2.]).

The natural general setting of the functional equation (1.4) is for S to be a semigroup, because the very formulation of (1.4) requires only an associative composition in S , not an identity element and inverses. Thus we study in the present paper Kannappan-cosine functional equation (1.4) on semigroups S , generalizing previous works in which S is a group. So, the result of the present paper is a naturel continuation of results contained in the literature.

The purpose of the present paper is to show how the relations between (1.4) and (1.2)-(1.3) on monoids extends to the much wider framework, in which S is a semigroup. We find explicit formulas for the solutions, expressing them in terms of homomorphisms and additives maps from the semigroup S into \mathbb{C} (Theorem 4.1). The continuous solutions on topological semigroups are also found.

2. Set up, notations and terminology

Throughout this paper S is a semigroup (a set with an associative composition) and z_0 is a fixed element in S .

If S is topological, we denote by $\mathcal{C}(S)$ the algebra of continuous functions from S to field of complex numbers \mathbb{C} .

Let $f : S \rightarrow \mathbb{C}$ be a function. We say that f is central if $f(xy) = f(yx)$ for all $x, y \in S$, and that f is abelian if $f(x_1x_2, \dots, x_n) = f(x_{\sigma(1)}x_{\sigma(2)}, \dots, x_{\sigma(n)})$ for all $x_1, x_2, \dots, x_n \in S$, all permutations σ of n elements and all $n \in \mathbb{N}$. A map $A : S \rightarrow \mathbb{C}$ is said to be additive if $A(xy) = A(x) + A(y)$, for all $x, y \in S$

and a map $\chi : S \rightarrow \mathbb{C}$ is multiplicative if $\chi(xy) = \chi(x)\chi(y)$, for all $x, y \in S$.

If $\chi \neq 0$, then the nullspace $I_\chi := \{x \in S \mid \chi(x) = 0\}$ is either empty or a proper subset of S and I_χ is a two sided ideal in S if not empty and $S \setminus I_\chi$ is a subsemigroup of S .

Note that additive and multiplicative functions are abelian.

For any subset $T \subseteq S$ let $T^2 := \{xy \mid x, y \in T\}$ and for any fixed element z_0 in S we let $T^2z_0 := \{xyz_0 \mid x, y \in T\}$.

To express solutions of our functional equations studied in this paper we will use the set $P_\chi := \{p \in I_\chi \setminus I_\chi^2 \mid up, pv, upv \in I_\chi \setminus I_\chi^2 \text{ for all } u, v \in S \setminus I_\chi\}$. For more details about P_χ we refer the reader to [5], [6] and [4].

3. Preliminaries

In this section we give useful results to solve the functional equation (1.3).

Lemma 3.1. *Let S be a semigroup, $n \in \mathbb{N}$, and $\chi, \chi_1, \chi_2, \dots, \chi_n : S \rightarrow \mathbb{C}$ be different non-zero multiplicative functions. Then*

- (a) $\{\chi_1, \chi_2, \dots, \chi_n\}$ is linearly independent.
- (b) If $A : S \setminus I_\chi \rightarrow \mathbb{C}$ is a non-zero additive function, then the set $\{\chi A, \chi\}$ is linearly independent on $S \setminus I_\chi$.

Proof. (a): See [13, Theorem 3.18]. (b): See [3, Lemma 4.4]. □

The proposition below gives the solutions of the functional equation

$$f(xyz_0) = \chi(z_0)f(x)\chi(y) + \chi(z_0)f(y)\chi(x), \quad x, y \in S, \tag{3.1}$$

Proposition 3.2. *Let S be a semigroup, and $\chi : S \rightarrow \mathbb{C}$ be a multiplicative function such that $\chi(z_0) \neq 0$. If $f : S \rightarrow \mathbb{C}$ is a solution of (3.1), then*

$$f(x) = \begin{cases} \chi(x)(A(x) + A(z_0)) & \text{for } x \in S \setminus I_\chi \\ \rho(x) & \text{for } x \in P_\chi \\ 0 & \text{for } x \in I_\chi \setminus P_\chi, \end{cases} \tag{3.2}$$

where $A : S \setminus I_\chi \rightarrow \mathbb{C}$ is additive and $\rho : P_\chi \rightarrow \mathbb{C}$ is the restriction of f to P_χ .

In addition, f is abelian and satisfies the following conditions:

- (I) $f(xy) = f(yx) = 0$ for all $x \in I_\chi \setminus P_\chi$ and $y \in S \setminus I_\chi$.
- (II) If $x \in \{up, pv, upv\}$ with $p \in P_\chi$ and $u, v \in S \setminus I_\chi$, then $x \in P_\chi$ and we have respectively $\rho(x) = \rho(p)\chi(u)$, $\rho(x) = \rho(p)\chi(v)$ or $\rho(x) = \rho(p)\chi(uv)$.

Conversely, the function f of the form (3.2) define a solution of (3.1).

Moreover, if S is a topological semigroup and $f \in \mathcal{C}(S)$, then $\chi \in \mathcal{C}(S)$, $A \in \mathcal{C}(S \setminus I_\chi)$ and $\rho \in \mathcal{C}(P_\chi)$.

Proof. See [7, Proposition 4.3]. □

To shorten the way to the solutions of functional equation (1.3), we prove the following Lemma that contains some key properties.

Lemma 3.3. *Let S be a semigroup and let $f, g : S \mapsto \mathbb{C}$ the solutions of the functional equation (1.3) with $g \neq 0$. Then*

(i) *If $f(z_0) = 0$ then*

(1)

$$g(z_0^2)g(xy) = g(z_0)[g(x)g(y) - f(x)f(y)] + f(z_0^2)f(xy), \quad x, y \in S. \quad (3.3)$$

(2)

$$g(z_0^2)^2 = g(z_0)^3 + f(z_0^2)^2.$$

(3) *If f and g are linearly independent then $g(z_0) \neq 0$.*

(ii) *If $f(z_0) \neq 0$, then there exist $\mu \in \mathbb{C}$ such that*

$$f(xyz_0) = f(x)g(y) + f(y)g(x) + \mu f(x)f(y), \quad x, y \in S. \quad (3.4)$$

Proof. (i) Suppose that $f(z_0) = 0$.

(1) Making the substitution (xy, z_0^2) and (xyz_0, z_0) in (1.3) we get respectively $g(xyz_0^3) = g(xy)g(z_0^2) - f(xy)f(z_0^2)$, and $g(xyz_0^3) = g(xyz_0)g(z_0) - f(xyz_0)f(z_0) = g(z_0)g(x)g(y) - g(z_0)f(x)f(y)$. Comparing these expressions, we deduce that $g(xy)g(z_0^2) - f(z_0^2)f(xy) = g(z_0)g(x)g(y) - g(z_0)f(y)f(x)$. This proves the identity desired.

(2) Follows directly by putting $x = y = z_0$ in the equation (3.3).

(3) By contradiction we suppose that $g(z_0) = 0$. Then by using (1.3) we get $g(xyz_0^2) = g(x)g(yz_0) - f(x)f(yz_0) = g(xy)g(z_0) - f(xy)f(z_0) = 0$ since $f(z_0) = g(z_0) = 0$. Then we deduce that

$$g(x)g(yz_0) = f(x)f(yz_0), \quad x, y \in S. \quad (3.5)$$

If $g(yz_0) = 0$ for all $y \in S$ then $0 = g(xyz_0) = g(x)g(y) - f(x)f(y)$, $x, y \in S$. So, $g(x)g(y) = f(x)f(y)$, $x, y \in S$. Hence $f = g$ or $f = -g$, which contradicts the fact that f and g are linearly independent. So $g \neq 0$ on Sz_0 then from (3.5) we get that $g = c_1f$ with $c_1 =: f(az_0)/g(az_0)$ for some $a \in S$ such that $g(az_0) \neq 0$. This is also a contradiction, since f and g are linearly independent. So we conclude that $g(z_0) \neq 0$.

(ii) Suppose that $f(z_0) \neq 0$. Making the substitution (x, yz_0^2) and (xyz_0, z_0) in (1.3) we get respectively $g(xyz_0^3) = g(x)g(yz_0^2) - f(x)f(yz_0^2) = g(z_0)g(x)g(y) - g(x)f(z_0)f(y) - f(x)f(yz_0^2)$ and $g(xyz_0^3) = g(xyz_0)g(z_0) - f(xyz_0)f(z_0) = g(z_0)g(x)g(y) - g(z_0)f(x)f(y) - f(xyz_0)f(z_0)$. Then, by the associativity of the operation in S we get

$$f(z_0)[f(xyz_0) - f(x)g(y) - f(y)g(x)] = f(x)[f(yz_0^2) - f(y)g(z_0) - f(z_0)g(y)]. \quad (3.6)$$

Since $f(z_0) \neq 0$, dividing (3.6) by $f(z_0)$ we get that

$$f(xyz_0) = f(x)g(y) + f(y)g(x) + f(x)\psi(y), \quad (3.7)$$

where $\psi(y) := f(z_0)^{-1}[f(yz_0^2) - f(y)g(z_0) - f(z_0)g(y)]$. Substituting (3.7) come back into (3.6), we find $f(z_0)[f(x)\psi(y)] = f(x)[f(y)\psi(z_0)]$, which implies that $\psi(y) = \mu f(y)$ with $\mu := \frac{\psi(z_0)}{f(z_0)}$. Therefore (3.7) becomes $f(xyz_0) = f(x)g(y) + f(y)g(x) + \mu f(x)f(y)$. This finishes the proof of Lemma 3.3. \square

4. Main results

Now, we are ready to describe the solutions of the functional equation (1.3). Let $\Psi_{A\chi,\rho} : S \rightarrow \mathbb{C}$ denote a function f of the form of [4, Theorem 3.1 (B)], where $\chi : S \rightarrow \mathbb{C}$ is a non-zero multiplicative function, $A : S \setminus I_\chi \rightarrow \mathbb{C}$ is additive, $\rho : P_\chi \rightarrow \mathbb{C}$ is the restriction of f , and conditions (i) and (ii) hold.

Theorem 4.1. *The solutions $f, g : S \rightarrow \mathbb{C}$ of the functional equation (1.3) are the following pairs of functions.*

(1) $f = g = 0$.

(2) $S \neq S^2z_0$ and we have

$$f = \pm g \text{ and } g(x) = \begin{cases} g_{z_0}(x) & \text{for } x \in S \setminus S^2z_0 \\ 0 & \text{for } x \in S^2z_0, \end{cases}$$

where $g_{z_0} : S \setminus S^2z_0 \rightarrow \mathbb{C}$ is an arbitrary non-zero function.

(3) There exist a constant $d \in \mathbb{C} \setminus \{\pm 1\}$ and a multiplicative function χ on S with $\chi(z_0) \neq 0$, such that

$$f = \frac{d\chi(z_0)}{1-d^2}\chi \text{ and } g = \frac{\chi(z_0)}{1-d^2}\chi.$$

(4) There exist a constant $c \in \mathbb{C}^* \setminus \{\pm i\}$ and two different multiplicative functions χ_1 and χ_2 on S , with $\chi_1(z_0) \neq 0$, $\chi_2(z_0) \neq 0$ and $\chi_1(z_0)\chi_2(z_0) \neq 0$ such that

$$f = -\frac{\chi_1(z_0)\chi_1 - \chi_2(z_0)\chi_2}{i(c^{-1} + c)} \text{ and } g = \frac{c^{-1}\chi_1(z_0)\chi_1 + c\chi_2(z_0)\chi_2}{c^{-1} + c}.$$

(5) There exist constants $q, \gamma \in \mathbb{C}^*$ and two different non-zero multiplicative functions χ_1 and χ_2 on S , with $\chi_1(z_0) = \frac{q^2 - (1 + \xi)^2}{2\gamma q}$, $\chi_2(z_0) = -\frac{q^2 - (1 - \xi)^2}{2\gamma q}$

and $\xi := \pm\sqrt{1 + q^2}$ such that

$$f = \frac{\chi_1 + \chi_2}{2\gamma} + \xi \frac{\chi_1 - \chi_2}{2\gamma} \text{ and } g = q \frac{\chi_1 - \chi_2}{2\gamma}.$$

(6) There exist constants $q \in \mathbb{C} \setminus \{\pm\alpha\}$, $\gamma \in \mathbb{C}^* \setminus \{\pm\alpha\}$ and $\delta \in \mathbb{C} \setminus \{\pm 1\}$, and two different non-zero multiplicative functions χ_1 and χ_2 on S , with $\chi_1(z_0) = \frac{(1 + \delta)^2 - (\alpha + q)^2}{2\gamma(1 + \delta)}$, $\chi_2(z_0) = \frac{(1 - \delta)^2 - (\alpha - q)^2}{2\gamma(1 - \delta)}$ and $\delta := \pm\sqrt{1 + q^2 - \alpha^2}$ such that

$$f = \alpha \frac{\chi_1 + \chi_2}{2\gamma} + q \frac{\chi_1 - \chi_2}{2\gamma} \text{ and } g = \frac{\chi_1 + \chi_2}{2\gamma} + (1 + \delta) \frac{\chi_1 - \chi_2}{2\gamma}.$$

(7) There exist a constant $\beta \in \mathbb{C}^*$, a non-zero multiplicative function χ on S , an additive function $A : S \setminus I_\chi \rightarrow \mathbb{C}$ and a function $\rho : P_\chi \rightarrow \mathbb{C}$ with $\chi(z_0) = \frac{1}{\beta}$ and $A(z_0) = 0$ such that

$$f = \frac{1}{\beta}\Psi_{A\chi,\rho} \text{ and } g = \frac{1}{\beta}(\chi \pm \Psi_{A\chi,\rho}).$$

(8) There exist a multiplicative function χ on S with $\chi(z_0) \neq 0$, an additive function $A : S \setminus I_\chi \rightarrow \mathbb{C}$ and a function $\rho : P_\chi \rightarrow \mathbb{C}$ such that

$$f = A(z_0)\chi + \Psi_{A\chi,\rho} \text{ and } g = (\chi(z_0) \pm A(z_0))\chi + \Psi_{A\chi,\rho}.$$

Moreover if S is a topological semigroup and $f \in \mathcal{C}(S)$ then $g \in \mathcal{C}(S)$ in cases (1), (2) and (4)-(8), and in (3) if $d \neq 0$.

Proof. If $g = 0$, then (1.3) reduces to $f(x)f(y) = 0$ for all $x, y \in S$. This implies that $f = 0$, so we get the first part of solutions. From now we may assume that $g \neq 0$.

If f and g are linearly dependent, then there exist $d \in \mathbb{C}$ such that $f = dg$. Substituting this into (1.3) we get the following equation $g(xyz_0) = (1 - d^2)g(x)g(y)$ $x, y \in S$. If $d^2 = 1$, then $g(xyz_0) = 0$ for all $x, y \in S$. Therefore $S \neq S^2z_0$ because $g \neq 0$. So we are in solution family (2) with g_{z_0} is an arbitrary non-zero function. If $d^2 \neq 1$, then by [11, Proposition 16] there exists a multiplicative function χ on S such that $\chi(z_0)\chi := (1 - d^2)g$ and $\chi(z_0) \neq 0$. Then we deduce that $g = \frac{\chi(z_0)}{1 - d^2}\chi$ and $f = dg = \frac{d\chi(z_0)}{1 - d^2}\chi$, so we have the solution family (3).

For the rest of the proof we assume that f and g are linearly independent. We split the proof into two cases according to whether $f(z_0) = 0$ or $f(z_0) \neq 0$.

Case 1. Suppose $f(z_0) = 0$. Then by Lemma 3.3 (i)-(3) and (i)-(1) we have respectively $g(z_0) \neq 0$ and

$$g(z_0^2)g(xy) = g(z_0)g(x)g(y) - g(z_0)f(x)f(y) + f(z_0^2)f(xy), \quad x, y \in S. \quad (4.1)$$

Subcase 1.1. Assume that $g(z_0^2) = 0$. Then by Lemma 3.3 (i)-(2) and (i)-(3) we get $f(z_0^2) \neq 0$ since f and g are linearly independent and then (4.1) can be rewritten as $f(xy) = \gamma f(x)f(y) - \gamma g(x)g(y)$, $x, y \in S$, where $\gamma := \frac{g(z_0)}{f(z_0^2)} \neq 0$.

Consequently the pair $(\gamma f, \gamma g)$ satisfies the cosine addition formula (1.4). So, according to [12, Theorem 6.1] and taking into account that f and g are linearly independent, we know that there are only the following possibilities.

(1.1.i) There exist a constant $q \in \mathbb{C}^*$ and two different non-zero multiplicative functions χ_1 and χ_2 on S such that $\gamma g = q \frac{\chi_1 - \chi_2}{2}$ and $\gamma f = \frac{\chi_1 + \chi_2}{2} \pm \left(\sqrt{1 + q^2} \right) \frac{\chi_1 - \chi_2}{2}$, which gives $f = \frac{\chi_1 + \chi_2}{2\gamma} \pm \left(\sqrt{1 + q^2} \right) \frac{\chi_1 - \chi_2}{2\gamma}$ and $g = q \frac{\chi_1 - \chi_2}{2\gamma}$. By putting $\xi := \pm \sqrt{1 + q^2}$ and using (1.3) we get

$$\begin{aligned} & \frac{1}{4\gamma^2} \left(q^2 - (1 + \xi)^2 \right) \chi_1(xy) + \frac{1}{4\gamma^2} \left(q^2 - (1 - \xi)^2 \right) \chi_2(xy) \\ &= \frac{q}{2\gamma} \chi_1(z_0)\chi_1(xy) - \frac{q}{2\gamma} \chi_2(z_0)\chi_2(xy), \end{aligned}$$

which implies by Lemma 3.1 (i) that $\frac{q}{2\gamma}\chi_1(z_0) = \frac{1}{4\gamma^2} (q^2 - (1 + \xi)^2)$ and $\frac{q}{2\gamma}\chi_2(z_0) = -\frac{1}{4\gamma^2} (q^2 - (1 - \xi)^2)$, since χ_1 and χ_2 are different and non-zero. Then we deduce that

$$\chi_1(z_0) = \frac{1}{2\gamma q} (q^2 - (1 + \xi)^2) \quad \text{and} \quad \chi_2(z_0) = -\frac{1}{2\gamma q} (q^2 - (1 - \xi)^2).$$

So we are in part (5).

(1.1.ii) There exist a non-zero multiplicative function χ on S and an additive function A on $S \setminus I_\chi$ such that $\gamma g = \Psi_{A\chi, \rho}$ and $\gamma f = \chi \pm \Psi_{A\chi, \rho}$.

If $z_0 \in I_\chi \setminus P_\chi$ we have $\gamma g(z_0) = \Psi_{A\chi, \rho}(z_0) = 0$ by definition of $\Psi_{A\chi, \rho}$. If $z_0 \in P_\chi$ we have $\chi(z_0) = 0$ and $|\gamma g(z_0)| = |\rho(z_0)| = |\chi(z_0) \pm \rho(z_0)| = |\gamma f(z_0)| = 0$. So, If $z_0 \in I_\chi$ we get that $\gamma g(z_0) = 0$ which is a contradiction because $g(z_0) \neq 0$ and $\gamma = \frac{g(z_0)}{f(z_0^2)}$.

Hence, $z_0 \in S \setminus I_\chi$ and we have $\chi(z_0) \neq 0$. Since $f(z_0) = 0$ by assumption we get that $f(z_0) = \frac{1}{\gamma} [\chi(z_0) \pm A(z_0)\chi(z_0)] = 0$, which implies that $A(z_0) = -1$. Now for all $x, y \in S \setminus I_\chi$, we have $xy z_0 \in S \setminus I_\chi$, then by using (1.3) we get that $\left(\frac{1}{\gamma} - \chi(z_0)\right) \chi(xy) + \left(\frac{1}{\gamma} + \chi(z_0)\right) \chi(xy)A(xy) = 0$, which implies according to Lemma 3.1(i), that $\frac{1}{\gamma} - \chi(z_0) = 0$ and $\frac{1}{\gamma} + \chi(z_0) = 0$, since $A \neq 0$.

Therefore, $\chi(z_0) = \frac{1}{\gamma} = -\frac{1}{\gamma}$, which is a contradiction because $\frac{1}{\gamma} \neq 0$ by assumption. So we do not have a solution corresponding this this possibility. Subcase 1.2: Suppose that $g(z_0^2) \neq 0$, then the functional equation (4.1) can be rewritten as follows $\beta g(xy) = \beta^2 g(x)g(y) - \beta^2 f(x)f(y) + \alpha\beta f(xy)$, $x, y \in S$ with $\beta := \frac{g(z_0)}{g(z_0^2)} \neq 0$ and $\alpha := \frac{f(z_0^2)}{g(z_0^2)}$. This shows that the pair $(\beta g, \beta f)$ satisfies the functional equation (1.2). So according to [12, Theorem 3.1], and taking into account that f and g are linearly independent, there are only the following possibilities.

(1.2.i) There exist a constant $q \in \mathbb{C} \setminus \{\pm\alpha\}$ and two different non-zero multiplicative functions χ_1 and χ_2 on S such that $\beta f = \alpha \frac{\chi_1 + \chi_2}{2} + q \frac{\chi_1 - \chi_2}{2}$

and $\beta g = \frac{\chi_1 + \chi_2}{2} \pm \sqrt{1 + q^2 - \alpha^2} \frac{\chi_1 - \chi_2}{2}$. Introducing $\delta := \pm \sqrt{1 + q^2 - \alpha^2}$

we find that $f = \alpha \frac{\chi_1 + \chi_2}{2\beta} + q \frac{\chi_1 - \chi_2}{2\beta}$ and $g = \frac{\chi_1 + \chi_2}{2\beta} + \delta \frac{\chi_1 - \chi_2}{2\beta}$. By using

(1.3) we get $\frac{1}{4\beta^2} \left((1 + \delta)^2 - (\alpha + q)^2 \right) \chi_1(xy) + \frac{1}{4\beta^2} \left((1 - \delta)^2 - (\alpha + q)^2 \right) \chi_2(xy)$

$= \frac{1}{2\beta} (1 + \delta) \chi_1(z_0) \chi_1(xy) + \frac{1}{2\beta} (1 - \delta) \chi_2(z_0) \chi_2(xy)$, which implies by Lemma

3.1(i) that $\frac{1}{2\beta} (1 + \delta) \chi_1(z_0) = \frac{1}{4\beta^2} \left((1 + \delta)^2 - (\alpha + q)^2 \right)$

and

$\frac{1}{2\beta}(1-\delta)\chi_2(z_0) = \frac{1}{4\beta^2}\left((1-\delta)^2 - (\alpha+q)^2\right)$, since χ_1 and χ_2 are different non-zero multiplicative functions. Notice that $\delta \neq \pm 1$ because $q \neq \pm\alpha$. Therefore we deduce that $\chi_1(z_0) = \frac{(1+\delta)^2 - (\alpha+q)^2}{2\beta(1+\delta)}$ and $\chi_2(z_0) = \frac{(1-\delta)^2 - (\alpha+q)^2}{2\beta(1-\delta)}$. Hence, by writing γ instead of β we get part (6).

(1.2.ii) $\alpha \neq 0$ and there exist two different non-zero multiplicative functions χ_1 and χ_2 on S such that $\beta f = \alpha\chi_1$ and $\beta g = \chi_2$. By using (1.3) again we get $\frac{1}{\beta}\left(\chi_2(z_0) - \frac{1}{\beta}\right)\chi_2(xy) + \left(\frac{\alpha^2}{\beta^2}\right)\chi_1(xy) = 0$, which gives $\chi_2(z_0) = \frac{1}{\beta}$ and $\alpha = 0$, since χ_1 and χ_2 are different. So this possibility is excluded because $\alpha \neq 0$.

(1.2.iii) There exist a non-zero multiplicative function χ on S and an additive function A on $S \setminus I_\chi$ such that $\beta f = \alpha\chi + \Psi_{A\chi,\rho}$ and $\beta g = \chi \pm \Psi_{A\chi,\rho}$, which gives $f = \frac{1}{\beta}(\alpha\chi + \Psi_{A\chi,\rho})$ and $g = \frac{1}{\beta}(\chi \pm \Psi_{A\chi,\rho})$.

If $g = \frac{1}{\beta}(\chi + \Psi_{A\chi,\rho})$ then $z_0 \notin I_\chi \setminus P_\chi$. Indeed if not we have $\chi(z_0) = 0$ and $\Psi_{A\chi,\rho}(z_0) = 0$. Then $\beta g(z_0) = \chi(z_0) + \Psi_{A\chi,\rho}(z_0) = 0$. This contradicts the fact that $g(z_0) \neq 0$.

On the other hand $z_0 \notin P_\chi$. Indeed if not we have $\chi(z_0) = 0$. Then $\beta g(z_0) = \Psi_{A\chi,\rho}(z_0) = \beta f(z_0) = 0$, which is a contradiction because $g(z_0) \neq 0$. So, $z_0 \in S \setminus I_\chi$ and then $\chi(z_0) \neq 0$. Since $f(z_0) = 0$ we get that $f(z_0) = \frac{\chi(z_0)}{\beta}[\alpha + A(z_0)] = 0$, which implies that $A(z_0) = -\alpha$. Now let $x, y \in S \setminus I_\chi$ be arbitrary. We have $xy z_0 \in S \setminus I_\chi$. By using (1.3) we get that

$$\left(\frac{1-\alpha^2}{\beta^2} + \frac{\alpha-1}{\beta}\chi(z_0)\right)\chi(xy) + \left(\frac{1-\alpha}{\beta^2} - \frac{1}{\beta}\chi(z_0)\right)\chi(xy)A(xy) = 0. \quad (4.2)$$

If $A = 0$ then $\rho \neq 0$ because $\Psi_{A\chi,\rho} \neq 0$, $\alpha = 0$, and $\chi(z_0) = \frac{1}{\beta}$ by (4.2). This is a special case of solution part (7).

If $A \neq 0$ then by Lemma 3.1 (ii) we get from (4.2) that $\frac{1-\alpha^2}{\beta^2} + \frac{\alpha-1}{\beta}\chi(z_0) = 0$ and $\frac{1-\alpha}{\beta^2} - \frac{1}{\beta}\chi(z_0) = 0$. As $\alpha \neq 1$, because $\chi(z_0) \neq 0$, we deduce that $\chi(z_0) = \frac{1-\alpha}{\beta}$ and $\chi(z_0) = \frac{1+\alpha}{\beta}$. So, we obtain that $\alpha = 0$ and $\chi(z_0) = \frac{1}{\beta}$, and the form of f reduces to $f = \frac{1}{\beta}\Psi_{A\chi,\rho}$. So we are in part (7).

If $g = \frac{1}{\beta}(\chi - \Psi_{A\chi,\rho})$, by using a similar computation as above we show that we are also in part (7).

Case 2. Suppose $f(z_0) \neq 0$. By using system (1.3) and (3.4), we deduce by

an elementary computation that for any $\lambda \in \mathbb{C}$

$$(g - \lambda f)(xyz_0) = (g - \lambda f)(x)(g - \lambda f)(y) - (\lambda^2 + \mu\lambda + 1)f(x)f(y), \quad x, y \in S. \tag{4.3}$$

Let λ_1 and λ_2 the two roots of the equation $\lambda^2 + \mu\lambda + 1 = 0$, then $\lambda_1\lambda_2 = 1$ which gives $\lambda_1 \neq 0$ and $\lambda_2 \neq 0$. According to [11, Proposition 16] we deduce from (4.3) that $g - \lambda_1 f := \chi_1(z_0)\chi_1$ and $g - \lambda_2 f := \chi_2(z_0)\chi_2$, where χ_1 and χ_2 are two multiplicative functions such that $\chi_1(z_0) \neq 0$ and $\chi_2(z_0) \neq 0$, because f and g are linearly independent.

If $\lambda_1 \neq \lambda_2$, then $\chi_1 \neq \chi_2$ and we get that $g = \frac{\lambda_2\chi_1(z_0)\chi_1 - \lambda_1\chi_2(z_0)\chi_2}{\lambda_2 - \lambda_1}$ and $f = \frac{\chi_1(z_0)\chi_1 - \chi_2(z_0)\chi_2}{\lambda_2 - \lambda_1}$. By putting $\lambda_1 = ic$ we get the solution categories (4).

If $\lambda_1 = \lambda_2 =: \lambda$, then $g - \lambda f =: \chi(z_0)\chi$ where χ is a multiplicative function on S such that $\chi(z_0) \neq 0$, because f and g are linearly independent. Hence

$$g = \chi(z_0)\chi + \lambda f. \tag{4.4}$$

Substituting this (3.4) an elementary computation shows that

$$f(xyz_0) = \chi(z_0)f(x)\chi(y) + \chi(z_0)f(y)\chi(x) + (2\lambda + \mu)f(x)f(y),$$

for all $x, y \in S$.

Moreover $\lambda = 1$ or $\lambda = -1$ because $\lambda_1\lambda_2 = 1$. Hence, $(\lambda, \mu) = (1, -2)$ or $(\lambda, \mu) = (-1, 2)$ since $\lambda^2 + \mu\lambda + 1 = 0$ and $\lambda \in \{-1, 1\}$. So, the functional equation above reduces to

$$f(xyz_0) = \chi(z_0)f(x)\chi(y) + \chi(z_0)f(y)\chi(x),$$

for all $x, y \in S$. Thus the pair (f, χ) satisfies the functional equation (3.1). Hence, in view of Proposition 3.2, we get $f = A(z_0)\chi + \Psi_{A\chi, \rho}$. Then, by using (4.4), we derive that $g = \chi(z_0)\chi + \lambda f = (\chi(z_0) + \lambda A(z_0))\chi + \lambda^2\Psi_{A\chi, \rho} = (\chi(z_0) \pm A(z_0))\chi + \Psi_{A\chi, \rho}$. This is part (8).

Conversely it is easy to check that the formulas of f and g listed in Theorem 4.1 define solutions of (1.3).

Finally, suppose that S is a topological semigroup. The continuity of the solutions of the forms (1)-(6) follows directly from [13, Theorem 3.18], and for the ones of the forms (7) and (8) it is parallel to the proof used in [6, Theorem 2.1] for categories (7)-(8). This completes the proof of Theorem 4.1. □

References

- [1] J. Aczél, *Lectures on functional equations and their applications*, Mathematics in Sciences and Engineering, vol. 19. Academic Press, New York, xx+510 pp(1966).
- [2] O. Ajebbar, E. Elqorachi, *Solutions and stability of trigonometric functional equations on an amenable group with an involutive automorphism*, Communications Korean Math. Soc., 34 (1), 55-82, (2019).

- [3] O. Ajebbar, E. Elqorachi, *The Cosine-Sine functional equation on a semigroup with an involutive automorphism*, Aequationes Math., 91, 1115-1146, (2017).
- [4] B. Ebanks, *Around the Sine Addition Law and d'Alembert's Equation on Semigroups*. Results in Math., 77, Article number: 11, (2022).
- [5] B. Ebanks, *The sine addition and subtraction formulas on semigroups*, Acta Math. Hungar., 167 (2), 533-555, (2021).
- [6] B. Ebanks, *The cosine and sine addition and subtraction formulas on semigroups*, Acta Math. Hungar., 165 (2), 337-354, (2021).
- [7] A. Jafar, O. Ajebbar, E. Elqorachi, *A Kannappan-sine addition law on semigroups*, <https://doi.org/10.48550/arXiv.2304.14146>.
- [8] T. A. Poulsen and H. Stetkær, *The trigonometric subtraction and addition formulas*, Aequationes Math., 59, 84-92 (2000).
- [9] B. Ebanks, *A Levi-Civita functional equation Monoids, Two Ways*. Silesianae, vol.36, no.2, 2022, pp.151-166. <https://doi.org/10.2478/amsil-2022-0009>.
- [10] Pl. Kannappan, *A functional equation for the cosine*, Canad. Math. Bull., 2, 495-498, (1968).
- [11] H. Stetkær, *Kannappan's functional equation on semigroups with involution*. Semigroup Forum, 94 (1), 17-30, (2017).
- [12] H. Stetkær, *The cosine addition law with an additional term*, Aequa. Math., 90 (6), 1147-1168, (2016).
- [13] H. Stetkær, : *Functional Equations on Groups*. World Scientific Publishing Company, Singapore (2013).
- [14] S. Kabbaj, M. Tial, D. Zeglami, *The Integral Cosine Addition and Sine Subtraction Laws*, Results in Math., 73, Article number: 97, (2018).

Jafar Ahmed¹, Ajebbar Omar², Elqorachi Elhoucien¹

¹ Ibn Zohr University, Faculty of sciences, Department of mathematics, Agadir, Morocco

² Sultan Moulay Slimane University, Multidisciplinary Faculty, Department of mathematics and computer science, Beni Mellal, Morocco

e-mail: hamadabenali2@gmail.com, omar-ajb@hotmail.com, elqorachi@hotmail.com