On certain paradoxical properties of Einstein photon fields and Dirac electron-positron fields

Beilinson A. A.*

* Department of Theoretical Physics Russian Peoples' Friendship University Russia, 117198, Moscow, Miklukho-Maklaya st., 6 alalbeyl@gmail.com

The article shows that the photons and electrons in the states with «deinterlaced» spins, which are isomorphic to their usual wave states, turned out to be not only nonlocal states on «Feynman paths», but also having a power to penetrate infinitely distant in arbitrarily small time.

Key words and phrases: generalized quantum mechanics, generalized paths integral, Feynman paths, quantum field of a photon.

Introduction

This refers to a number of reliable experimental data, according to which the speed of propagation of signals generated by «entangled» photons and electrons exceeds many times the speed of light, see for example [9].

The article shows that in the scope of generalized solutions of equations for an Einstein photon e.-m. field and Dirac electron-positron field these physical quantum fields along with existance of nonlocal states of corresponding relativistic quantum particles and their path integrals, and mentioned paradoxical properties, also can be successfully described, see [2–4].

The article shows that these phenomena arises due to unitarily equivalent states of relativistic quantum particles associated with cut-off their spin interactions. A similar phenomenon in relativistic quantum particle physics has actually already been considered using known Foldy–Wouthuysen variables, see [5].

For simplicity onedimensional fields will be considered. The speed of light assuming equal 1; More detailed the case of e.-m. field is considered.

The work uses notation and terminology corresponding to monographs «Generalized functions», issues 1, 2 and 4 by I. M. Gel'fand, N. Ya. Vilenkin and G. E. Shilov (see [6-8]).

Two record possibilities for generalized retarded Green functional of e.-m. field and Dirac field

Recall that Maxwell equations of e.-m. field in Majorana variables [1]

$$M_t(x) = E_t(x) + iH_t(x), \quad \bar{M}_t(x) = E_t(x) - iH_t(x),$$

assuming this field is fundamentally quantum (see [1, 4]), are becomes

$$i\frac{\partial}{\partial t}M_t(x) = (S,\hat{p})M_t(x), \quad -i\frac{\partial}{\partial t}\bar{M}_t(x) = (S,\hat{p})\bar{M}_t(x),$$

where $\hat{p} = \frac{1}{i} \nabla$, and S is spin operators of a photon $M_t(x)$, see [1]); antiphoton $bar M_t(x)$ is considered similarly.

In momentum representation these equations are as follows:

$$i\frac{\partial}{\partial t}\tilde{M}_t(p) = (S,p)\tilde{M}_t(p), \quad -i\frac{\partial}{\partial t}\bar{\tilde{M}}_t(p) = (S,p)\bar{\tilde{M}}_t(p).$$

We will consider such motion of a photon along the z axis, in which the coordinates x, y is any. To this e.-m. field, it is easy to see, corresponding to it the three of the momentum representations of it wave functions

$$\tilde{M}_t(p_z) = \begin{pmatrix} \tilde{M}_t^{(1)}(p_z) \\ \tilde{M}_t^{(2)}(p_z) \end{pmatrix}, \quad \tilde{M}_t^{(3)}(p_z)$$

satisfying the equations

$$i\frac{\partial}{\partial t}\tilde{M}_t(p_z) = \begin{pmatrix} 0, & -ip_z\\ ip_z, & 0 \end{pmatrix}\tilde{M}_t(p_z), \quad i\frac{\partial}{\partial t}\tilde{M}_t^{(3)}(p_z) = 0,$$

whose retarded Green function in momentum representation is

$$\tilde{M}_t(p_z) = \exp\left(-it\begin{pmatrix}0, & -ip_z\\ip_z, & 0\end{pmatrix}\right) = \exp\left(-i\begin{pmatrix}0, & -itp_z\\itp_z, & 0\end{pmatrix}\right), \quad \tilde{M}_t^{(3)}(p_z) = 1.$$

(we will not consider last equation).

Bearing in mind the constructing of the coordinate representation the generalized Green function of this field as functional on columns of bump functions $\varphi(z) \in K$ (as Fourier preimage of solution in the momentum representation, see [7]), we will first consider the latter as a functional

$$\int \bar{\tilde{M}}_t(p_z) \begin{pmatrix} \psi_1(p_z) \\ \psi_2(p_z) \end{pmatrix} dp_z$$

on columns of analytic test functions $\psi(p) \in \mathbb{Z}$ (see [4]).

Remark that the momentum representation of solution contains a Hermitian matrix in the exponent, which can therefore be reduced to a diagonal form by unitary transform $\hat{Q}(p_z)$. Therefore we have

$$\tilde{M}_t(p_z) = \exp\left(-i\begin{pmatrix}0, & -itp_z\\itp_z, & 0\end{pmatrix}\right) = \tilde{Q}^+(p_z)\begin{pmatrix}\exp(-it|p_z|), & 0\\0, & \exp(it|p_z|)\end{pmatrix}\tilde{Q}(p_z),$$

where

$$\tilde{Q}(p_z) = \frac{1}{\sqrt{2}} \begin{pmatrix} -i \operatorname{sgn} p_z, & 1\\ i \operatorname{sgn} p_z, & 1 \end{pmatrix}.$$

Note that arised diagonal generalized Green function can be called the Green function of e.-m. field in Foldy–Wouthuysen variables (see [5]).

Therefore, it is easy to see,

$$\tilde{M}_t(p_z) = \frac{1}{2} \begin{pmatrix} \exp(-it|p_z|) + \exp(it|p_z|), & \operatorname{sgn}(p_z) \cdot 2i\sin(t|p_z|) \\ -\operatorname{sgn}(p_z) \cdot 2i\sin(t|p_z|), & \exp(it|p_z|) + \exp(-it|p_z|) \end{pmatrix} = \begin{pmatrix} \cos(tp_z), & \sin(tp_z) \\ -\sin(tp_z), & \cos(tp_z) \end{pmatrix}.$$

But the Fourier preimages of numerical functionals $\exp(it|p_z|)$ and $\operatorname{sgn} p_z$ on Z are the quantum Cauchy functional (see [3])

$$C_{it}(z) = \frac{1}{2} \left(\delta(t-z) + \delta(t+z) \right) + \frac{i}{2\pi} \left(\frac{1}{t-z} + \frac{1}{t+z} \right)$$

and, accordingly, the functional $-\frac{i}{\pi z}$ (see [6], p. 360 formula 19) on bump functions $\varphi(z) \in K$. Hence it is easy to see that we have coordinate representation the generalized Green function

$$\int \bar{M}_t \begin{pmatrix} \varphi_1 \\ \varphi_2 \end{pmatrix} dz$$

of Maxwell-Majorana equations, where

$$M_t(z) = \frac{1}{2} \begin{pmatrix} \delta(t-z) + \delta(t+z), & i\left(\delta(t-z) - \delta(t+z)\right) \\ -i\left(\delta(t-z) - \delta(t+z)\right), & \delta(t-z) + \delta(t+z) \end{pmatrix}.$$

Remark that conducted diagonalization operation of the retarded Green function of e.-m. field have physical interpretation, since diagonalization of matrix underlying of this operation is interpreted as cut-off the spin interaction between components of the considered generalized e.-m. field of a photon and appearance of «new» field in the result with the Green function momentum representation

$$\tilde{m}_t(p_z) = \begin{pmatrix} \exp(-it|p_z|), & 0\\ 0, & \exp(it|p_z|) \end{pmatrix} = \exp\left(it \begin{pmatrix} -|p_z|, & 0\\ 0, & |p_z| \end{pmatrix}\right),$$

with «deinterlaced» components. Here the unitary operator $\tilde{Q}(p_z)$ realizes both «deinterlacing» and «interlacing» of the filed components, and their Fourier preimage as functionals on bump functions

$$Q(z) = \frac{1}{\sqrt{2}} \begin{pmatrix} \frac{-1}{\pi z}, & \delta(z) \\ \frac{1}{\pi z}, & \delta(z) \end{pmatrix}, \quad Q^+(z) * Q(z) = \begin{pmatrix} \delta(z), & 0 \\ 0, & \delta(z) \end{pmatrix}.$$

realizes «interlacing» and «deinterlacing» in the coordinate representation.

Here the Green function coordinate representation as functional on bump functions, it is easy to see, is

$$M_t(z) = Q^+(z) * m_t(z) * Q(z)$$

Compare one more $M_t(z)$ and $m_t(z)$. These generalized retarded Green functions of e.-m. field of a photon are unitarily equivalent and isomorphic; despite of that they are represented of one initial state development, this is radially different branch of a photon quantum state development: $M_t(z)$ is wave one propagating with fundamental velocity, and other one is $m_t(z)$ filling whole coordinate space in arbitrarily small time and, as has been shown, has nonlocal generalized states on «Feynman paths» (see [2, 4].

Thus the isomorphism of «deinterlaced» and wave states indicate the existance of integral over «Feynman paths» also in case of wave brach solution $M_t(z)$.

Let us consider case of Dirac electron-positron field, the retarded Green function of which is delivered by Pauli–Jordan formula that also is functionals on bump functions. As has been shown, the diagonalization of the Hamiltonian of this functional reducing to known Foldy–Wouthuysen variables (see [5]) also is interpreted as «deinterlacing» of the wave functional components when disabling spin interaction and reduces to the Green functional of Dirac electron (see [3])

$$\begin{pmatrix} C_{it}^m(z), & 0\\ 0, & C_{it}^m(z) \end{pmatrix}, \text{ where } C_{it}^m(z) = \frac{1}{2} \left(\delta(t-z) + \delta(t+z) \right) + \frac{1}{\pi} \cdot \frac{-tmK_1 \left(im\sqrt{t^2 - z^2} \right)}{\sqrt{t^2 - z^2}}$$

Thus also in case of Dirac field arising the unitarily equivalent isomorphic form of the field solutions that has lost its usual wave properties and acquired a property instant propogation in whole sapce. It can be shown that, together with this property, this form acquired exactly the same properties of nonlocality as the photon field. Here the corresponding generalized quantum measures are turned out to be absolutely continuous.

It should be remarked that all the above constructions, it is easy to see, are Lorentz-covariant.

References

- 1. Axiezer A. I., Beresteckij V. B. Quantum electrodynamics / Ed. by L. P. Rusakova. 4th edition. M. : Nauka, 1981.
- Beilinson A. A. The Cauchy process in imaginary time and the path integrals in relativistic quantum mechanics of Dirac's electrons and Einstein's photons // International Scientific Journal. — Vol. 4 of Journal of Mathematics. — AU : CreateSpace, 2018. — Aug. — P. 103–116.
- 3. Beilinson A. A. The quantum mechanics of the free Dirac electrons and Einstein photons, and the Cauchy distribution // International Scientific Journal. Vol. 4 of Journal of Mathematics. AU : CreateSpace, 2018. Aug. P. 36–55.
- 4. Beilinson A. A. The generalized quantum mechanics of Einstein photon and Casimir force. ArXiv. 2023. 2302.10176.
- Foldy L. L., A. Wouthuysen S. On the Dirac theory of spin 1/2 particles and its non-relativistic limit // Phys. Rev. — 1950. — Vol. 78.
- 6. Gel'fand I. M., Shilov G. E. Generalized functions: Properties and Operations / Ed. by and trans. from rus. E. Saletan. N. Y. and L. : Academic Press, 1964. Vol. 1st of Generalized functions.
- 7. Gel'fand I. M., Shilov G. E. Spaces of Fundamental and Generalized Functions / Ed. by C. P. Peltzer and trans. from rus. M. D. Friedman, A. Feinstein. N. Y. and L. : Academic Press, 1968. Vol. 2nd of Generalized functions.
- Gel'fand I. M., Vilenkin N. Ya. Application of Harmonic Analysis / Ed. by trans. from rus. A. Feinstein. N. Y. and L. : Academic Press, 1964. — Vol. 4th of Generalized functions.
- 9. Nobel prize in physics. -2022.