

A SHARP INEQUALITY FOR TRACE-FREE MATRICES WITH APPLICATIONS TO HYPERSURFACES

JEFFREY S. CASE AND AARON J. TYRRELL

ABSTRACT. We derive a sharp inequality relating the second and fourth elementary symmetric functions of the eigenvalues of a trace-free matrix and give two applications. First, we give a new proof of the classification of conformally flat hypersurfaces in spaceforms. Second, we construct a functional which characterizes rotational hypersurfaces and catenoids.

1. INTRODUCTION

Let $i: N^n \rightarrow (M^{n+1}, g)$, $n \geq 4$, be an immersed hypersurface in a locally conformally flat Riemannian manifold. This note is motivated by two interesting rigidity results for N in terms of its principal curvatures. First, Nishikawa and Maeda showed [5] that (N^n, i^*g) is locally conformally flat if and only if at each point it has a principal curvature of multiplicity at least $n - 1$. Second, do Carmo and Dajczer showed [1] that, under the stronger assumption that (M^{n+1}, g) is complete, simply connected, and of constant sectional curvature $c \in \mathbb{R}$, the hypersurface N is contained in a so-called rotation hypersurface [1, Definition 2.2] if at each point it has a principal curvature of multiplicity exactly $n - 1$. Furthermore, if N is minimal, then it is contained in a catenoid if and only if at each point it has a principal curvature of multiplicity at least $n - 1$.

The purpose of this short note is to point out that the condition on the principal curvatures can be recast in terms of a sharp inequality relating the squared-norm $|\mathring{A}^2|^2$ of the square of the trace-free part of the second fundamental form \mathring{A} and the square of the squared-norm $|\mathring{A}|^2$ of \mathring{A} . Our main result is a purely algebraic statement about trace-free linear maps on finite-dimensional vector spaces:

Theorem 1.1. *Let V be an n -dimensional inner product space, $n \geq 4$, and let $\mathring{A} \in \text{End}(V)$ be trace-free. Then*

$$|\mathring{A}^2|^2 \leq \frac{n^2 - 3n + 3}{n(n-1)} |\mathring{A}|^4$$

with equality if and only if \mathring{A} has an eigenspace of dimension at least $n - 1$.

The main idea of the proof of Theorem 1.1 is that the Newton inequality relating the first, second, and third elementary symmetric functions of the eigenvalues of a matrix A realizes equality if and only if A is proportional to the identity or has rank 1. Applying this to $\mathring{A} + \lambda I$ for I the identity map yields the sharp inequality

$$\left(\text{tr } \mathring{A}^3\right)^2 \leq \frac{(n-2)^2}{n(n-1)} |\mathring{A}|^6$$

2020 *Mathematics Subject Classification.* Primary 53C42; Secondary 26D05, 53A07, 53C24.
Key words and phrases. rotation hypersurface; catenoid; rigidity.

for any trace-free $\mathring{A} \in \text{End}(V^n)$; this is sometimes called the λ -method [4]. Repeating this argument for the Newton inequality relating the second, third, and fourth elementary symmetric functions yields Theorem 1.1.

We conclude with a number of applications of Theorem 1.1 to the rigidity of locally conformally flat hypersurfaces, of rotation hypersurfaces, and of catenoids.

Our first application is a new proof of the characterization [5] of locally conformally flat hypersurfaces in locally conformally flat manifolds in terms of their principal curvatures.

Corollary 1.2. *Let $i: N^n \rightarrow (M^{n+1}, g)$, $n \geq 4$, be an immersed hypersurface in a locally conformally flat manifold. The induced metric i^*g on N is locally conformally flat if and only if for each $p \in N$, the shape operator $A_p: T_pN \rightarrow T_pN$ has an eigenspace of dimension at least $n - 1$.*

Our second application is a rigidity result for catenoids in simply connected spaceforms amongst all minimal hypersurfaces.

Corollary 1.3. *Let $i: N^n \rightarrow M^{n+1}(c)$, $n \geq 4$, be a minimal hypersurface in the complete, simply connected Riemannian $(n + 1)$ -manifold of constant sectional curvature $c \in \mathbb{R}$. Then the shape operator $A: TN \rightarrow TN$ satisfies*

$$(1.1) \quad |A^2|^2 \leq \frac{n^2 - 3n + 3}{n(n-1)} |A|^4.$$

Moreover, equality holds if and only if $i(N)$ is contained in a catenoid.

The above results motivate the introduction of two energy functionals for hypersurfaces of a Riemannian manifold. First, given an immersion $i: N^n \rightarrow (M^{n+1}, g)$, we define the *rotational energy* of N by

$$E_{rot}[N] := \int_N \left(\frac{n^2 - 3n + 3}{n(n-1)} |\mathring{A}|^4 - |\mathring{A}^2|^2 \right) dA,$$

assuming it is finite. Our terminology is motivated by the characterization [1] of rotation hypersurfaces in simply connected spaceforms.

Corollary 1.4. *Let $i: N^n \rightarrow M^{n+1}(c)$, $n \geq 4$, be a nowhere umbilic immersed hypersurface in the complete, simply connected, Riemannian $(n + 1)$ -manifold of constant sectional curvature $c \in \mathbb{R}$. Then $E_{rot}[N] \geq 0$ with equality if and only if N is contained in a rotation hypersurface.*

This result is sharper for minimal immersions.

Corollary 1.5. *Let $i: N^n \rightarrow M^{n+1}(c)$, $n \geq 4$, be a minimal immersed hypersurface in the complete, simply connected, Riemannian $(n + 1)$ -manifold of constant sectional curvature $c \in \mathbb{R}$. Then $E_{rot}[N] \geq 0$ with equality if and only if N is contained in a catenoid.*

Second, given an immersion $i: N^n \rightarrow (M^{n+1}, g)$, we define the *conformal rotational energy* of N by

$$E_{rot}^{conf}[N] := \int_N |\mathring{A}|^{n-4} \left(\frac{n^2 - 3n + 3}{n(n-1)} |\mathring{A}|^4 - |\mathring{A}^2|^2 \right) dA,$$

assuming it is finite. Note that $E_{rot}^{conf} = E_{rot}$ for four-dimensional hypersurfaces. The main point of this definition is that E_{rot}^{conf} is conformally invariant, and hence

gives a conformally invariant characterization of immersed submanifolds for which the shape operator has an eigenspace of dimension at least $n - 1$.

Corollary 1.6. *Let $i: N^n \rightarrow (M^{n+1}, g)$, $n \geq 4$, be an immersed hypersurface in a Riemannian manifold. Then $E_{rot}^{conf}[N] \geq 0$ with equality if and only if for each $p \in N$, the shape operator $A_p: T_p N \rightarrow T_p N$ has an eigenspace of dimension at least $n - 1$.*

This note is organized as follows. In Section 2 we give an elementary proof of Theorem 1.1. In Section 3 we prove our geometric applications of Theorem 1.1.

2. ELEMENTARY SYMMETRIC FUNCTIONS

Let $A \in \text{End}(V^n)$ be a linear map on an n -dimensional vector space. Denote by

$$(2.1) \quad \sigma_k(A) := \sum_{i_1 < \dots < i_k} \lambda_{i_1} \cdots \lambda_{i_k}$$

the k -th elementary symmetric function of the eigenvalues $\{\lambda_1, \dots, \lambda_n\}$ of A , with the convention $\sigma_0(A) = 1$. Note that $\sigma_1(A) = \text{tr } A$. Define $p_k(A)$, $k \leq n$, by

$$\sigma_k(A) = \binom{n}{k} p_k(A);$$

this renormalization is such that $p_k(\lambda I) = \lambda^k$, where I is the identity map. The main tool in the proof of Theorem 1.1 is the sharp version of Newton's inequalities:

Lemma 2.1 ([6, Section 2]). *Let $A \in \text{End}(V^n)$ and let $k \leq n - 1$ be a positive integer. Then*

$$p_k^2(A) \geq p_{k-1}(A)p_{k+1}(A)$$

with equality if and only if A is proportional to the identity or $\dim \ker A \geq n - k + 1$.

Applying the λ -method to Lemma 2.1 with $k = 2$ and $k = 3$ yields useful inequalities, the second of which is recorded in Theorem 1.1. We record the resulting inequalities separately due to their different dependence on $n = \dim V$.

Our first inequality relates $p_2(\mathring{A})$ and $p_3(\mathring{A})$ for a trace-free linear map \mathring{A} on a vector space of dimension at least 3.

Proposition 2.2. *Let $\mathring{A} \in \text{End}(V^n)$, $n \geq 3$, be such that $p_1(\mathring{A}) = 0$. Then*

$$p_3^2(\mathring{A}) + 4p_2^3(\mathring{A}) \leq 0.$$

Moreover, equality holds if and only if \mathring{A} has an eigenspace of dimension at least $n - 1$.

Proof. It is well-known, and follows easily from Equation (2.1), that

$$(2.2) \quad p_k(A + \lambda I) = \sum_{j=0}^k \binom{k}{j} \lambda^j p_{k-j}(A)$$

for all $A \in \text{End}(V^n)$ and all $\lambda \in \mathbb{R}$.

Now let $\mathring{A} \in \text{End}(V^n)$ be such that $p_1(\mathring{A}) = 0$. Lemma 2.1 implies that $p_2(\mathring{A}) \leq 0$ with equality if and only if $\mathring{A} = 0$. Since the conclusion is trivially true if $\mathring{A} = 0$, we may assume that $p_2(\mathring{A}) < 0$. For notational simplicity, denote $p_k := p_k(\mathring{A})$.

Let $\lambda \in \mathbb{R}$. Applying Lemma 2.1 to $\mathring{A} + \lambda I$ and simplifying via Equation (2.2) yields

$$(2.3) \quad \begin{aligned} 0 &\leq (p_2 + \lambda^2)^2 - \lambda(p_3 + 3\lambda p_2 + \lambda^3) \\ &= p_2^2 - \lambda p_3 - \lambda^2 p_2, \end{aligned}$$

with equality if and only if \mathring{A} has an eigenspace of dimension at least $n-1$. In fact, since $p_1 = 0$ and $p_2 < 0$, the dimension must equal $n-1$ in the case of equality in Inequality (2.3). Multiplying Inequality (2.3) by $4p_2 < 0$ yields

$$\begin{aligned} 0 &\geq 4p_2^3 - 4\lambda p_2 p_3 - 4\lambda^2 p_2^2 \\ &= -(2\lambda p_2 + p_3)^2 + 4p_2^3 + p_3^2. \end{aligned}$$

Therefore $p_2^3 + 4p_3^2 \leq 0$ with equality if and only if \mathring{A} has an eigenspace of dimension $n-1$. \square

Our second inequality relates $p_2(\mathring{A})$ and $p_4(\mathring{A})$ for a trace-free linear map \mathring{A} on a vector space of dimension at least 4.

Proposition 2.3. *Let $\mathring{A} \in \text{End}(V^n)$, $n \geq 4$, be such that $p_1(\mathring{A}) = 0$. Then*

$$p_4(\mathring{A}) + 3p_2^2(\mathring{A}) \geq 0.$$

Moreover, equality holds if and only if \mathring{A} has an eigenspace of dimension at least $n-1$.

Proof. Let $\mathring{A} \in \text{End}(V^n)$ be such that $p_1(\mathring{A}) = 0$. For notational simplicity, denote $p_k := p_k(\mathring{A})$. Lemma 2.1 implies that $p_2(\mathring{A}) \leq 0$ with equality if and only if $\mathring{A} = 0$. Since the conclusion is trivially true if $\mathring{A} = 0$, we may assume that $p_2 < 0$.

Now, applying Lemma 2.1 and Equation (2.2) to $\mathring{A} + \lambda I$ yields

$$\begin{aligned} 0 &\leq (p_3 + 3\lambda p_2 + \lambda^3)^2 - (p_2 + \lambda^2)(p_4 + 4\lambda p_3 + 6\lambda^2 p_2 + \lambda^4) \\ &= p_3^2 - p_2 p_4 + 2\lambda p_2 p_3 + \lambda^2(3p_2^2 - p_4) - 2\lambda^3 p_3 - \lambda^4 p_2 \end{aligned}$$

for all $\lambda \in \mathbb{R}$. Multiplying by $16p_2^3$ and denoting $s := 2\lambda p_2 + p_3$ yields

$$\begin{aligned} 0 &\geq 16p_2^3 p_3^2 - 16p_2^4 p_4 + 32\lambda p_2^4 p_3 + 16\lambda^2 p_2^2(3p_2^3 - p_2 p_4) - 32\lambda^3 p_2^3 p_3 - 16\lambda^4 p_2^4 \\ &= (3p_3^2 - 4p_2 p_4)(p_2^3 + 4p_2^3) + 8s p_3(p_2 p_4 - p_2^3 - p_3^2) + s^2(12p_2^3 - 4p_2 p_4 + 6p_3^2) - s^4. \end{aligned}$$

Setting $s = 0$ yields

$$(2.4) \quad (3p_3^2 - 4p_2 p_4)(p_2^3 + 4p_2^3) \leq 0.$$

Proposition 2.2 implies that $p_2^3 + 4p_3^2 \leq 0$ with equality if and only if, up to a multiplicative constant and a choice of basis,

$$\mathring{A} = \begin{pmatrix} 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & \cdots & 0 & 1-n \end{pmatrix}.$$

If $p_2^3 + 4p_3^2 = 0$, then a straightforward computation yields $p_k = 1 - k$ for all $k \in \mathbb{N}$, and hence $p_4 + 3p_2^2 = 0$. If instead $p_2^3 + 4p_3^2 < 0$, then Inequality (2.4) implies that $3p_3^2 - 4p_2 p_4 \geq 0$. Therefore

$$0 \leq 3p_3^2 - 4p_2 p_4 < -12p_2^3 - 4p_2 p_4 = -4p_2(p_4 + 3p_2^2).$$

Since $p_2 < 0$, we conclude that $p_4 + 3p_2^2 > 0$. \square

Expressing the nonnegative quantity $p_4 + 3p_2^2$ in terms of $|A|^4$ and $|A^2|^2$ yields Theorem 1.1.

Proof of Theorem 1.1. Let $\mathring{A} \in \text{End}(V^n)$ be such that $p_1(\mathring{A}) = 0$. Direct computation gives

$$\begin{aligned}\sigma_2(\mathring{A}) &= -\frac{1}{2}|\mathring{A}|^2, \\ \sigma_4(\mathring{A}) &= \frac{1}{8}|\mathring{A}|^4 - \frac{1}{4}|\mathring{A}^2|^2.\end{aligned}$$

Therefore

$$0 \leq \binom{n}{4} (p_4(\mathring{A}) + 3p_2^2(\mathring{A})) = -\frac{1}{4} \left(|\mathring{A}^2|^2 - \frac{n^2 - 3n + 3}{n(n-1)} |\mathring{A}|^4 \right).$$

The conclusion readily follows from Proposition 2.3. \square

3. GEOMETRIC APPLICATIONS

We conclude by proving our geometric applications of Theorem 1.1.

Proof of Corollary 1.2. Let $i: N^n \rightarrow (M^{n+1}, g)$ be a Riemannian immersion into a locally conformally flat manifold. Denote by \overline{W} the Weyl tensor of i^*g . Denote by

$$F := \frac{1}{n-2} (\mathring{A}^2 - Gi^*g)$$

the Fialkow tensor $F \in \Gamma(S^2T^*N)$, where $G := \text{tr}_{i^*g} F = \frac{1}{2(n-1)} |\mathring{A}|^2$ is its trace (cf. [2, Equation (13.9)]). The Gauss–Codazzi equations (cf. [2, Equation (22.13)]) imply that

$$\overline{W} = \frac{1}{2} \mathring{A} \wedge \mathring{A} + F \wedge g,$$

where $S \wedge T$ denotes the Kulkarni–Nomizu product

$$(S \wedge T)_{abcd} := S_{ac}T_{bd} + S_{bd}T_{ac} - S_{ad}T_{bc} - S_{bc}T_{ad}.$$

Direct computation gives (cf. [3])

$$\begin{aligned}|\mathring{A} \wedge \mathring{A}|^2 &= 8|\mathring{A}|^4 - 8|\mathring{A}^2|^2, \\ \langle \mathring{A} \wedge \mathring{A}, F \wedge g \rangle &= -8\langle \mathring{A}^2, F \rangle, \\ |F \wedge g|^2 &= 4\langle \mathring{A}^2, F \rangle, \\ \langle \mathring{A}^2, F \rangle &= \frac{1}{n-2} |\mathring{A}^2|^2 - \frac{1}{2(n-1)(n-2)} |\mathring{A}|^4.\end{aligned}$$

Therefore

$$|\overline{W}|^2 = \frac{2(n^2 - 3n + 3)}{(n-1)(n-2)} |\mathring{A}|^4 - \frac{2n}{n-2} |\mathring{A}^2|^2.$$

The conclusion readily follows from Theorem 1.1. \square

Proof of Corollary 1.3. Let $i: N^n \rightarrow M^{n+1}(c)$ be a minimal hypersurface. By minimality, the shape operator is trace-free; i.e. $A = \mathring{A}$. Theorem 1.1 implies that A satisfies Inequality (1.1) with equality if and only if A has an eigenvalue of multiplicity at least $n-1$. The final conclusion follows from a result [1, Corollary 4.4] of do Carmo and Dajczer. \square

Proof of Corollary 1.4. Let $i: N^n \rightarrow M^{n+1}(c)$ be a nowhere umbilic immersed hypersurface. Theorem 1.1 implies that $E_{rot}[N] \geq 0$ with equality if and only if at each point $p \in M$ the shape operator A_p has an eigenspace of dimension exactly $n - 1$. The final conclusion follows from a result [1, Theorem 4.2] of do Carmo and Dajczer. \square

Proof of Corollary 1.5. Let $i: N^n \rightarrow M^{n+1}(c)$ be a minimal immersed hypersurface. Corollary 1.3 implies that $E_{rot}[N] \geq 0$ with equality if and only if $i(N)$ is contained in a catenoid. \square

Proof of Corollary 1.6. Let $i: N^n \rightarrow (M^{n+1}, g)$ be an immersed hypersurface. Theorem 1.1 implies that $E_{rot}^{conf}[N] \geq 0$ with equality if and only if

$$|\mathring{A}|^{n-4} \left(\frac{n^2 - 3n + 3}{n(n-1)} |\mathring{A}|^4 - |\mathring{A}^2|^2 \right) = 0.$$

It follows that for each $p \in N$, either $\mathring{A}_p = 0$ or

$$|\mathring{A}_p^2|^2 = \frac{n^2 - 3n + 3}{n(n-1)} |\mathring{A}_p|^4.$$

In the former case, \mathring{A}_p has an eigenspace of dimension n . In the latter case, Theorem 1.1 implies that \mathring{A}_p has an eigenspace of dimension at least $n - 1$. \square

ACKNOWLEDGEMENTS

We thank Eudes Leite de Lima for pointing out a mistake in an early version of this paper. JSC was partially supported by the Simons Foundation (Grant #524601).

REFERENCES

- [1] M. do Carmo and M. Dajczer, *Rotation hypersurfaces in spaces of constant curvature*, Trans. Amer. Math. Soc. **277** (1983), no. 2, 685–709. MR694383
- [2] A. Fialkow, *Conformal differential geometry of a subspace*, Trans. Amer. Math. Soc. **56** (1944), 309–433. MR11023
- [3] M.-L. Labbi, *Double forms, curvature structures and the (p, q) -curvatures*, Trans. Amer. Math. Soc. **357** (2005), no. 10, 3971–3992. MR2159696
- [4] D. S. Mitrinović, *Analytic inequalities*, Die Grundlehren der mathematischen Wissenschaften, Band 165, Springer-Verlag, New York-Berlin, 1970. In cooperation with P. M. Vasić. MR0274686
- [5] S. Nishikawa and Y. Maeda, *Conformally flat hypersurfaces in a conformally flat Riemannian manifold*, Tohoku Math. J. (2) **26** (1974), 159–168. MR338967
- [6] S. Rosset, *Normalized symmetric functions, Newton's inequalities and a new set of stronger inequalities*, Amer. Math. Monthly **96** (1989), no. 9, 815–819. MR1033349

109 McALLISTER BUILDING, PENN STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA
 Email address: jscase@psu.edu

18A DEPARTMENT OF MATHEMATICS AND STATISTICS, TEXAS TECH UNIVERSITY, LUBBOCK, TX 79409, USA
 Email address: aatyrrel@ttu.edu