

FIRST DERIVATIVE OF AUTOMORPHIC FUNCTION OF TRIANGLE GROUP

MD. SHAFIUL ALAM, BIJAN KRISHNA SAHA, AND CHINMAYEE PODDER

ABSTRACT. For a triangle group G , the G -automorphic function is the inverse of Schwarz triangle function. In this paper, we compute the first derivative of the G -automorphic function for the triangle group G in terms of the Gaussian hypergeometric function.

1. INTRODUCTION

If $\alpha, \beta, \gamma \in \mathbb{C}$ for $\gamma \neq 0, -1, -2, \dots$, and $j \in \mathbb{N} \cup \{0\}$, then the function ${}_2F_1(\alpha, \beta; \gamma; \xi)$ defined by

$${}_2F_1(\alpha, \beta; \gamma; \xi) = \sum_{n=0}^{\infty} \frac{(\alpha, j)(\beta, j)}{(\gamma, j)j!} \xi^j, \quad |\xi| < 1,$$

where

$$(\alpha, j) = \begin{cases} 1 & \text{if } j = 0 \text{ and } \alpha \neq 0 \\ \alpha(\alpha + 1) \cdots (\alpha + j - 1) & \text{if } j \geq 1 \end{cases},$$

is known as the Gaussian hypergeometric function. By the Euler's integral representation formula, one can extend ${}_2F_1(\alpha, \beta; \gamma; \xi)$ analytically to $\mathbb{C} \setminus [1, +\infty)$ (cf. [3, Chapter II], [15, Chapter XIV]).

For the group $\mathrm{SL}_2(\mathbb{R})$ given by

$$\mathrm{SL}_2(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{R}, \quad ad - bc = 1 \right\},$$

the group $\mathrm{PSL}_2(\mathbb{R})$ is defined as $\mathrm{PSL}_2(\mathbb{R}) = \mathrm{SL}_2(\mathbb{R}) / \{I_2, -I_2\}$, where I_2 is the 2×2 identity matrix (see [14, Chapter VII]). Let \mathcal{H} denote the complex upper half-plane $\{w \in \mathbb{C} : \mathrm{Im} w > 0\}$. The action of $\mathrm{PSL}_2(\mathbb{R})$ on \mathcal{H} is given by

$$w \mapsto \begin{pmatrix} a & b \\ c & d \end{pmatrix} w = \frac{aw + b}{cw + d}, \quad \text{for } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PSL}_2(\mathbb{R}), \quad w \in \mathcal{H}.$$

Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PSL}_2(\mathbb{R})$. The element A is called parabolic if $a + d = \pm 2$, elliptic if $(a + d) \in \mathbb{R}$ and $|a + d| < 2$, hyperbolic if $(a + d) \in \mathbb{R}$ and $|a + d| > 2$, loxodromic if

2020 *Mathematics Subject Classification.* 11F03; 33C05.

Key words and phrases. Automorphic function, Triangle group, Hypergeometric function.

$(a + d) \in \mathbb{C}$ (see Theorem 15, Section 10 of [10]). Let $G \subset \text{PSL}_2(\mathbb{R})$ be a triangle group with signature $(0; n_1, n_2, n_3)$ acting on \mathcal{H} . Consider a parabolic element $A_1 \in G$ and an elliptic element $A_2 \in G$. If $A_1(x) = x$, where $x \in \mathbb{R} \cup \{\infty\}$, then the point x is called a cusp of the group G . If $A_2(w) = w$, where $w \in \mathcal{H}$, then we call the point w an elliptic point of G . Since the action of G on \mathcal{H} is properly discontinuous, we obtain the quotient Riemann surface $G \backslash \mathcal{H}$ (see [5] and [11]). In [1] and [2], a special type of triangle group known as the Hecke group is investigated to study modular equations.

For a hyperbolic triangle $[w_1, w_2, w_3]$ with angles $\frac{\pi}{n_1}, \frac{\pi}{n_2}, \frac{\pi}{n_3}$, let f denote the Schwarz triangle function. The G -automorphic function $\xi : \mathcal{H} \rightarrow G \backslash \mathcal{H}^*$, where \mathcal{H}^* is the union of \mathcal{H} and the set of cusps of G , is the inverse of f and

$$\xi\left(\frac{aw + b}{cw + d}\right) = \xi(w)$$

for $w \in \mathcal{H}$ and $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G$. The function $\xi(w)$ can be normalized in such a way that the orders of the ramifications are n_1, n_2, n_3 over $0, 1, \infty$, respectively (see [8]). Suppose that n_1, n_2 and n_3 are the orders of the elliptic points w_1, w_2 and w_3 , respectively, such that

$$\xi(w_1) = 0, \quad \xi(w_2) = 1, \quad \xi(w_3) = \infty.$$

Then, $\xi(w)$ satisfies the following hypergeometric differential equation

$$(1.1) \quad \xi(1 - \xi) \frac{d^2 g}{d\xi^2} + \{\gamma - (\alpha + \beta + 1)\xi\} \frac{dg}{d\xi} - \alpha\beta g = 0.$$

Let $g_1(\xi)$ and $g_2(\xi)$ be linearly independent solutions of (1.1), then $f(\xi) = \frac{g_2(\xi)}{g_1(\xi)}$ is the Schwarz triangle function. The Schwarzian derivative, denoted by $\{f, \xi\}$, of $f(\xi)$ is given by

$$\{f, \xi\} = \left(\frac{f''(\xi)}{f'(\xi)}\right)' - \frac{1}{2} \left(\frac{f''(\xi)}{f'(\xi)}\right)^2.$$

The function $f(z)$ satisfies the Schwarzian equation given by

$$(1.2) \quad \{f, \xi\} = \frac{1 - p_1^2}{2\xi^2} + \frac{1 - p_2^2}{2(1 - \xi)^2} + \frac{1 - p_1^2 - p_2^2 + p_3^2}{2\xi(1 - \xi)},$$

where p_1, p_2, p_3 are known as accessory parameters and $p_1 = \frac{1}{n_1}, p_2 = \frac{1}{n_2}, p_3 = \frac{1}{n_3}$ (cf. Chap. V, Sect. 7 of [12]). These facts are presented in the following theorem.

Theorem A ([4, Theorem 6.2]). *For $\gamma \neq 1$, let*

$$\alpha' = \alpha - \gamma + 1, \quad \beta' = \beta - \gamma + 1, \quad \gamma' = 2 - \gamma,$$

then the functions

$$g_1(\xi) = {}_2F_1(\alpha, \beta; \gamma; \xi) \quad \text{and} \quad g_2(\xi) = \xi^{1-\gamma} {}_2F_1(\alpha', \beta'; \gamma'; \xi)$$

are linearly independent solutions of (1.1). Let $w = f(\xi) = \frac{g_2(\xi)}{g_1(\xi)}$, then $\mathcal{H} = \{\xi \in \mathbb{C} : \text{Im } \xi > 0\}$ is mapped by $f(\xi)$ conformally to the hyperbolic (non-Euclidean) triangle $[w_1, w_2, w_3]$ in the w -plane, where

$$\begin{aligned} w_1 &= f(0) = 0, \\ w_2 &= f(1) = \frac{\Gamma(2-\gamma)\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)}{\Gamma(\gamma)\Gamma(1-\alpha)\Gamma(1-\beta)}, \\ w_3 &= f(\infty) = e^{\pi i(1-\gamma)} \frac{\Gamma(\alpha)\Gamma(\gamma-\beta)\Gamma(2-\beta)}{\Gamma(\gamma)\Gamma(\alpha-\gamma+1)\Gamma(1-\beta)}. \end{aligned}$$

The interior angles at the vertices w_1, w_2 and w_3 are, respectively, $(1-\gamma)\pi$, $(\gamma-\alpha-\beta)\pi$ and $(\beta-\alpha)\pi$.

For $w_1, w_2 \in \mathcal{H}$, let us denote the hyperbolic (non-Euclidean) distance between w_1 and w_2 by $d_h(w_1, w_2)$. Since at the vertices w_1, w_2 and w_3 of the hyperbolic triangle $[w_1, w_2, w_3]$, the angles are $\frac{\pi}{n_1}$, $\frac{\pi}{n_2}$ and $\frac{\pi}{n_3}$, respectively, we have

$$(1.3) \quad \tanh^2 \frac{d_h(w_1, w_2)}{2} = \frac{\cos \left\{ \frac{1}{2} \left(\frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3} \right) \pi \right\} \cos \left\{ \frac{1}{2} \left(\frac{1}{n_1} + \frac{1}{n_2} - \frac{1}{n_3} \right) \pi \right\}}{\cos \left\{ \frac{1}{2} \left(\frac{1}{n_2} + \frac{1}{n_3} - \frac{1}{n_1} \right) \pi \right\} \cos \left\{ \frac{1}{2} \left(\frac{1}{n_3} + \frac{1}{n_1} - \frac{1}{n_2} \right) \pi \right\}}$$

(see Equation (75.1) of [7], where $a_i = d_h(w_1, w_2)$, $\alpha_i = \frac{\pi}{n_3}$, $\alpha_j = \frac{\pi}{n_1}$, $\alpha_k = \frac{\pi}{n_2}$). Also, by Equation (85.3) of [7], we have

$$(1.4) \quad \tanh^2 \frac{d_h(w_1, w_2)}{2} = \left| \frac{w_2 - w_1}{w_2 - \bar{w}_1} \right|.$$

In the following section, we present the main results and their proofs.

2. MAIN RESULTS AND PROOFS

Theorem 2.1. *Suppose that G is a triangle group with signature $(0; n_1, n_2, n_3)$. Let*

$$\xi : \mathcal{H} \rightarrow G \backslash \mathcal{H}^*$$

be the G -automorphic function and let $w_1, w_2, w_3 \in \mathcal{H}$ be the elliptic points of orders n_1, n_2 and n_3 , respectively, such that

$$\xi(w_1) = 0, \quad \xi(w_2) = 1, \quad \xi(w_3) = \infty.$$

Then,

$$(2.1) \quad \frac{d\xi}{dw} = C \xi^{(1-\frac{1}{n_1})} (1-\xi)^{(1-\frac{1}{n_2})} \left({}_2F_1(\alpha, \beta; \gamma; \xi) - K \xi^{\frac{1}{n_1}} {}_2F_1(\alpha', \beta'; \gamma'; \xi) \right)^2,$$

where $C \in \mathbb{C} \setminus \{0\}$, $\alpha, \beta \notin \mathbb{Z}$, $\alpha' = \alpha - \gamma + 1$, $\beta' = \beta - \gamma + 1$, $\gamma' = 2 - \gamma$ and

$$K = \left(\frac{\Gamma(1-\alpha)\Gamma(1-\beta)\Gamma(\alpha')\Gamma(\beta')}{\Gamma(\alpha)\Gamma(\beta)\Gamma(1-\alpha')\Gamma(1-\beta')} \right)^{\frac{1}{2}} \frac{\Gamma(\gamma)}{\Gamma(\gamma')}.$$

We need the following lemma in the proof of Theorem 2.1.

Lemma 2.2. *Let $w_1, w_2 \in \mathcal{H}$ be the elliptic points of orders n_1 and n_2 , respectively. Then the hyperbolic (non-Euclidean) distance, $d_h(w_1, w_2)$, between w_1 and w_2 is given by*

$$\tanh^2 \frac{d_h(w_1, w_2)}{2} = \frac{\sin \pi \alpha \sin \pi \beta}{\sin \pi \alpha' \sin \pi \beta'}.$$

Proof. We have

$$(2.2) \quad \frac{1}{n_1} = 1 - \gamma, \quad \frac{1}{n_2} = \gamma - \alpha - \beta, \quad \frac{1}{n_3} = \beta - \alpha$$

and

$$(2.3) \quad \alpha' = \alpha - \gamma + 1, \quad \beta' = \beta - \gamma + 1, \quad \gamma' = 2 - \gamma.$$

From (2.2) and (2.3), it follows that

$$(2.4) \quad \alpha = \frac{1}{2} \left(1 - \frac{1}{n_1} - \frac{1}{n_2} - \frac{1}{n_3} \right), \quad \beta = \frac{1}{2} \left(1 - \frac{1}{n_1} - \frac{1}{n_2} + \frac{1}{n_3} \right), \quad \gamma = 1 - \frac{1}{n_1}$$

and

$$(2.5) \quad \alpha' = \frac{1}{2} \left(1 + \frac{1}{n_1} - \frac{1}{n_2} - \frac{1}{n_3} \right), \quad \beta' = \frac{1}{2} \left(1 + \frac{1}{n_1} - \frac{1}{n_2} + \frac{1}{n_3} \right), \quad \gamma' = 1 + \frac{1}{n_1}.$$

Thus,

$$\begin{aligned} \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3} &= 1 - 2\alpha, \\ \frac{1}{n_1} + \frac{1}{n_2} - \frac{1}{n_3} &= 1 - 2\beta, \\ -\frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3} &= 1 - 2\alpha', \\ -\frac{1}{n_1} + \frac{1}{n_2} - \frac{1}{n_3} &= 1 - 2\beta'. \end{aligned}$$

From (1.3), we have

$$\begin{aligned} \tanh^2 \frac{d_h(w_1, w_2)}{2} &= \frac{\cos \left\{ \frac{1}{2}(1 - 2\alpha)\pi \right\} \cos \left\{ \frac{1}{2}(1 - 2\beta)\pi \right\}}{\cos \left\{ \frac{1}{2}(1 - 2\alpha')\pi \right\} \cos \left\{ -\frac{1}{2}(1 - 2\beta')\pi \right\}} \\ &= \frac{\sin \pi \alpha \sin \pi \beta}{\sin \pi \alpha' \sin \pi \beta'}. \end{aligned}$$

□

Recall that the linearly independent solutions of (1.1) are $g_1(\xi) = {}_2F_1(\alpha, \beta; \gamma; \xi)$ and $g_2(\xi) = \xi^{1-\gamma} {}_2F_1(\alpha', \beta'; \gamma'; \xi)$ for $0 < \gamma < 1$. The quotient $w = f(\xi) = \frac{g_2(\xi)}{g_1(\xi)}$ maps conformally \mathcal{H} to the interior of $[w_1, w_2, w_3]$. Let $\partial[w_1, w_2, w_3]$ and $\partial\mathcal{H}$ denote the boundary of the triangle $[w_1, w_2, w_3]$ and the boundary of \mathcal{H} , respectively. Then $f(\xi)$ forms a homeomorphism between $\partial\mathcal{H} = \mathbb{R} \cup \{\infty\}$ and $\partial[w_1, w_2, w_3]$. The orders of w_1 , w_2 and w_3 are n_1 , n_2 and n_3 , respectively, where n_1, n_2, n_3 are integers greater than 2 or infinity. Since

$$\frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3} < 1,$$

the inverse $\xi(w)$ of the Schwarz triangle function $f(\xi)$ is single-valued (see p. 416 in [13]). Also, $\xi(w_1) = 0$, $\xi(w_2) = 1$ and $\xi(w_3) = \infty$.

Consider the elliptic point $w_1 \in \mathcal{H}$ of order n_1 and let $A \in G$ be the generator of the stabilizer subgroup for w_1 , then

$$(2.6) \quad \frac{Aw - w_1}{Aw - \bar{w}_1} = e^{\frac{2\pi i}{n_1} \frac{w - w_1}{w - \bar{w}_1}}$$

(see (9) of Section 48 in [10]). Also, for $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G$ and $w = \frac{g_2}{g_1}$, we have

$$(2.7) \quad w \mapsto \begin{pmatrix} a & b \\ c & d \end{pmatrix} w = \frac{aw + b}{cw + d} = \frac{ag_2 + bg_1}{cg_2 + dg_1}.$$

The two sides (hyperbolic lines) of $[w_1, w_2, w_3]$ through w_1 are mapped to straight lines through the origin by the Möbius transformation

$$w \rightarrow \frac{w - w_1}{w - \bar{w}_1}.$$

Proof of Theorem 2.1. The points on $G \setminus \mathcal{H}$ are the G -orbits of w near the elliptic point w_1 such that

$$(2.8) \quad \frac{w - w_1}{w - \bar{w}_1} = K \frac{w_2}{w_1} = K \frac{\xi^{1-\gamma} {}_2F_1(\alpha', \beta'; \gamma'; \xi)}{{}_2F_1(\alpha, \beta; \gamma; \xi)}$$

for a nonzero constant K (see (48) of [9]). By assumptions, $\xi(w_2) = 1$ and

$$\begin{aligned} \gamma' - \alpha' - \beta' &= \gamma - \alpha - \beta, \\ \gamma' - \beta' &= 1 - \beta, \\ \gamma' - \alpha' &= 1 - \alpha. \end{aligned}$$

Thus,

$${}_2F_1(\alpha, \beta; \gamma; \xi(w_2)) = \frac{\Gamma(\gamma)\Gamma(\gamma - \alpha - \beta)}{\Gamma(\gamma - \alpha)\Gamma(\gamma - \beta)}$$

(see Section 14.11 of [3]) and

$$\begin{aligned} {}_2F_1(\alpha', \beta'; \gamma'; \xi(w_2)) &= \frac{\Gamma(\gamma')\Gamma(\gamma' - \alpha' - \beta')}{\Gamma(\gamma' - \alpha')\Gamma(\gamma' - \beta')} \\ &= \frac{\Gamma(\gamma')\Gamma(\gamma - \alpha - \beta)}{\Gamma(1 - \alpha)\Gamma(1 - \beta)}. \end{aligned}$$

Therefore, when $w \rightarrow w_2$, from (2.8) we have

$$\frac{w_2 - w_1}{w_2 - \bar{w}_1} = K \frac{\Gamma(\gamma')\Gamma(\gamma - \alpha)\Gamma(\gamma - \beta)}{\Gamma(\gamma)\Gamma(1 - \alpha)\Gamma(1 - \beta)}$$

or,

$$(2.9) \quad K = \frac{w_2 - w_1}{w_2 - \bar{w}_1} \frac{\Gamma(\gamma)\Gamma(1 - \alpha)\Gamma(1 - \beta)}{\Gamma(\gamma')\Gamma(1 - \alpha')\Gamma(1 - \beta')}.$$

Since $\sin \pi\alpha = \frac{\pi}{\Gamma(\alpha)\Gamma(1 - \alpha)}$ for $\alpha \notin \mathbb{Z}$, by Lemma 2.2 we have

$$(2.10) \quad \tanh^2 \frac{d_h(w_1, w_2)}{2} = \frac{\sin \pi\alpha \sin \pi\beta}{\sin \pi\alpha' \sin \pi\beta'} = \frac{\Gamma(\alpha')\Gamma(1 - \alpha')\Gamma(\beta')\Gamma(1 - \beta')}{\Gamma(\alpha)\Gamma(1 - \alpha)\Gamma(\beta)\Gamma(1 - \beta)}.$$

From (1.4), it follows that

$$(2.11) \quad \tanh^2 \frac{d_h(w_1, w_2)}{2} = \left(\frac{w_2 - w_1}{w_2 - \bar{w}_1} \right)^2.$$

Thus, from (2.9), (2.10) and (2.11), we have

$$K = \left(\frac{\Gamma(1 - \alpha)\Gamma(1 - \beta)\Gamma(\alpha')\Gamma(\beta')}{\Gamma(\alpha)\Gamma(\beta)\Gamma(1 - \alpha')\Gamma(1 - \beta')} \right)^{\frac{1}{2}} \frac{\Gamma(\gamma)}{\Gamma(\gamma')}.$$

Now, (2.8) implies

$$(2.12) \quad w = \frac{w_1 g_1 - K \bar{w}_1 g_2}{g_1 - K g_2}.$$

Therefore,

$$(2.13) \quad \frac{dw}{d\xi} = K(w_1 - \bar{w}_1) \frac{g_1 \frac{dg_2}{d\xi} - \frac{dg_1}{d\xi} g_2}{(g_1 - K g_2)^2}.$$

Let W denote the Wronskian of g_1 and g_2 given by

$$W = g_1 \frac{dg_2}{d\xi} - \frac{dg_1}{d\xi} g_2.$$

Then,

$$(2.14) \quad \frac{dw}{d\xi} = K(w_1 - \bar{w}_1) \frac{W}{(g_1 - K g_2)^2}.$$

From (1.1), we have

$$(2.15) \quad \frac{d^2g}{d\xi^2} + P(\xi)\frac{dg}{d\xi} + Q(\xi)g = 0,$$

where

$$P(\xi) = \frac{\gamma - (\alpha + \beta + 1)\xi}{\xi(1 - \xi)}$$

and

$$Q(\xi) = \frac{\alpha\beta}{\xi(\xi - 1)}.$$

Since the linearly independent solutions of (2.15) are $g_1(\xi)$ and $g_2(\xi)$, we have by Abel's theorem (see Theorem 3.3.2 of [6])

$$(2.16) \quad W(\xi) = W(\xi_0) \exp\left(-\int_{\xi_0}^{\xi} P(\xi)d\xi\right),$$

where $W(\xi_0) \neq 0$. For some constant $C_0 \in \mathbb{C} \setminus \{0\}$, it follows from (2.16) that

$$(2.17) \quad W(\xi) = C_0\xi^{-\gamma}(1 - \xi)^{\gamma-\alpha-\beta-1}.$$

Therefore, from (2.14) and (2.17) we have

$$(2.18) \quad \frac{dw}{d\xi} = C_0K(w_1 - \bar{w}_1)\frac{\xi^{-\gamma}(1 - \xi)^{\gamma-\alpha-\beta-1}}{(g_1 - Kg_2)^2},$$

which implies

$$\frac{d\xi(w)}{dw} = C\xi^{(1-\frac{1}{n_1})}(1 - \xi)^{(1-\frac{1}{n_2})}\left({}_2F_1(\alpha, \beta; \gamma; \xi) - K\xi^{\frac{1}{n_1}}{}_2F_1(\alpha', \beta'; \gamma'; \xi)\right)^2,$$

where

$$(2.19) \quad C = \frac{1}{C_0K(w_1 - \bar{w}_1)}.$$

□

Corollary 2.2.1. *The constants C and K in Theorem 2.1 are inversely proportional to each other.*

Proof. From (2.19), it follows that $C \propto \frac{1}{K}$. □

REFERENCES

- [1] Md. S. Alam, *On Ramanujan's modular equations and Hecke groups*, Preprint (2021), arXiv:2105.13152.
- [2] Md. S. Alam and T. Sugawa, *Geometric deduction of the solutions to modular equations*, The Ramanujan Journal **59(2)**(2022), 459–477.
- [3] H. Bateman, *Higher Transcendental Functions*, Vol. I, McGraw-Hill, New York, 1953.
- [4] P. Bayer and A. Travesa, *Uniformizing functions for certain Shimura curves, in the case $D = 6$* , Acta Arithmetica, **126**(2007), 315–339.
- [5] A. F. Beardon, *The Geometry of Discrete Groups*, Graduate Texts in Mathematics **91**, Springer-Verlag, New York, 1983.
- [6] W. E. Boyce, R. C. DiPrima and D. B. Meade, *Elementary differential equations and boundary value problems*, John Wiley & Sons, 2021.
- [7] C. Carathéodory, *Theory of Functions of a Complex Variable*, Chelsea, New York, 1954.
- [8] P. B. Cohen, C. Itzykson and J. Wolfart, *Fuchsian triangle groups and Grothendieck dessins. Variations on a theme of Belyi*, Communications in Mathematical Physics, **163(3)**(1994), 605–627.
- [9] N. D. Elkies, *Shimura curve computations, in Algorithmic number theory (Portland, OR, 1998)*, Lecture Notes in Computer Science, vol. 1423 (Springer, Berlin, 1998), 1–47.
- [10] L. R. Ford, *Automorphic Functions*, McGraw-Hill, New York, 1929.
- [11] S. Katok, *Fuchsian Groups*, The University of Chicago Press, Chicago and London, 1992.
- [12] Z. Nehari, *Conformal Mapping*, McGraw-Hill, New York, 1952.
- [13] G. Sansone and J. Gerretsen, *Lectures on the Theory of Functions of a Complex Variable: II. Geometric Theory*, Wolters-Noordhoff Publ., Groningen, 1969.
- [14] J.-P. Serre. *A Course in Arithmetic*, Graduate Texts in Mathematics 7, Springer-Verlag, New York, 1973.
- [15] E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, 4th Edition, Cambridge University Press, Cambridge, 1927.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BARISHAL, BARISHAL-8254, BANGLADESH
Email address: msalam@bu.ac.bd, shafiulmt@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BARISHAL, BARISHAL-8254, BANGLADESH
Email address: bijandumath@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BARISHAL, BARISHAL-8254, BANGLADESH
Email address: chinmayeepodder1@gmail.com