

# Symphonic map from ellipsoid to ellipsoid

Xiang-Zhi Cao

May 9, 2023

## Abstract

In this paper, we proved the existence of Symphonic map from ellipsoid to ellipsoid. We also geive give Hopf construction of Symphonic map from ellipsoid to ellipsoid.

## 1 Introductory

It is well known that a map  $f : (M, g) \rightarrow S^n(1)$  is harmonic map if and only if

$$\Delta_g f + |\nabla f|^2 f = 0.$$

When  $M$  is  $S^m$ , we can construct harmonic map  $f : (S^m(1), g) \rightarrow S^n(1)$  by restricting the map between Euclidean space, whose component is harmonic homogeneous polynomials.

In 1960s, Toda [25] gived complete details of these calculations of the homotopy group of spheres. In 1970s, Smith [22] considered harmonic map between spheres using join method. Peng and Tang[17] construct harmonic representation of homotopy group of spheres. Tang [16, 24] consider harmonic representation of the homotopy group  $\pi_{16p+j}(S^{16p+j})$  under some conditons on  $j$ . Ratto [4, 18, 21] used ODE method to construct harmonic map between sphere or ellipsoid. Ding [2] obtained the existence of symmetric harmonic map between sphere under some conditons.

We can parametrize  $x \in S^{m_1+m_2-1}$  uniquely by  $(x_1 \sin s, x_2 \cos s)$  with  $x_1 \in S^{m_1-1}, x_2 \in S^{m_2-1}, s \in [0, \pi/2]$ . Given a map  $f : S^{m_1-1} \times S^{m_2-1} \rightarrow S^{n-1}$ , the Hopf construction is given by  $h_f : S^{m_1+m_2-1} \rightarrow S^n$ , which is

$$h_f(x_1 \sin s, x_2 \cos s) := (f(x_1, x_2) \sin \alpha(s), \cos \alpha(s))$$

Let  $\lambda_1$  and  $\lambda_2$  be the eigenvalues of  $f$ .

Smith [23] proved that a harmonic Hopf construction exists in the case that  $m_1 = m_2 = 2, \lambda_1 = \lambda_2$ . Ratto [20] considered hopf construction in the case that  $4\lambda_1 < (m_1 - 2)^2, 4\lambda_2 < (m_2 - 2)^2$  or  $m_1 = m_2 = 2, \lambda_1 = \lambda_2$ . Ratto[19] proved that no harmonic Hopf construction can exist if  $\lambda_1 \leq (m_1 - 1)\lambda_2, m_1 \geq 3, m_2 = 2$ . Gastel [6] consider Hopf construction exists if  $\lambda_1 > (m_1 - 1)\lambda_2, m_1 \geq 3, m_2 = 2$ . Ding [3] proved that there is

always a harmonic Hopf construction in the case  $m_1, m_2 \geq 3$ . Later, Gastel [5] gives simplified the proof of Dings result in [3]. Later, Ding, Fan and Li [1] obtained the existence of Hopf construction if  $m_1 = 2, m_2 > 2, \lambda_1 \geq 1, \lambda_2 > \lambda_1(m_2 - 1)$ .

In 2011, Nakauchi and Takenaka [15] introduced the symphonic map, which is the critical point of

$$E(u) = \int_M \frac{|u^*h|^2}{2} dv_g.$$

In 2014, Nakauchi [14] used symphonic joining method to construct symphonic map between spheres. Inspired by the work [14] and the history of harmonic map, in this paper, our main aims are to construct symphonic map between ellipsoid and gives hopf construction of symphonic map.

One can refer to [7, 10, 9, 8, 11, 13, 12] for the studies for symphonic map.

## 2 preliminary

We denote  $\mathbb{S}^n$  the unit sphere of dimension  $n$ , we denote  $\mathbb{S}^n(r)$  the sphere of dimension  $n$  with radius  $r$ . The ellipsoid  $Q^{m_1+m_2+1}(a, b)$  is parametrized by

$$z = a \cos t \cdot x + b \sin t \cdot y$$

for  $x \in \mathbb{S}^{m_1}, y \in \mathbb{S}^{m_2}$  and  $0 \leq t \leq \pi/2$ . The induced Riemannian metric on  $Q^{m_1+m_2+1}(a, b)$  is:

$$g = (a^2 \cos^2 t) g_{\mathbb{S}^{m_2}} + (b^2 \sin^2 t) g_{\mathbb{S}^{m_1}} + h^2(t) dt^2$$

where

$$h(s) = [b^2 \sin^2 t + a^2 \cos^2 t]^{1/2}$$

Its volume density is

$$v_g = a^{m_1} b^{m_2} (\cos^{m_1} t \sin^{m_2} t) h(t) \cdot v_{\mathbb{S}^{m_1}} \cdot v_{\mathbb{S}^{m_2}}$$

where  $v_{\mathbb{S}^{m_1}}, v_{\mathbb{S}^{m_2}}$  are volume densities of the indicated Euclidean spheres.

We shall refer to  $(Q^{m_1+m_2+1}(a, b), g)$  as an ellipsoidal join of  $\mathbb{S}^{m_1}, \mathbb{S}^{m_2}$ . We observe that  $\text{SO}(m_1 + 1) \times \text{SO}(m_2 + 1)$  is a group of isometries of  $Q^{p+q+1}(a, b)$  and that  $Q^{m_1+m_2+1}(a, b)$  and  $Q^{m_2+m_1+1}(b, a)$  are isometric.

An eigenmap  $u : \mathbb{S}^p \rightarrow \mathbb{S}^r$  is one whose components (as a map into  $\mathbb{R}^{r+1}$ ) are harmonic  $k$ -homogeneous polynomials; its associated eigenvalue is  $\lambda_u = k(k + p - 1)$ . It is easy to calculate that  $|du(x)|^2 = \lambda_u$  for all  $x \in \mathbb{S}^p$ ; and that  $u$  is a harmonic map.

**Lemma 2.1** ((cf. [4, 21])). *Let  $\varphi$  be the harmonic map from Riemannian manifold  $(M, g)$  into ellipsoid  $Q^n(c, d)$ . we denote by  $\Phi = i \circ \varphi$  the composition of  $\varphi$  with the canonical embedding  $i$  of  $Q^n(c, d)$  into  $\mathbb{R}^{n+1}$ . Then  $\varphi$  is harmonic iff*

$$\Delta_g \Phi = \left( |P^{-1/2} d\Phi|^2 / |P^{-1} \Phi|^2 \right) P^{-1} \Phi \quad (2.1)$$

where

$$P = \begin{pmatrix} c^2 & & & & & 0 \\ & \ddots & & & & \\ & & c^2 & & & \\ & & & d^2 & & \\ & & & & \ddots & \\ 0 & & & & & d^2 \end{pmatrix}$$

If we write  $\Phi = (\Phi_1, \Phi_2)$ , the components being the projections on the factors of the ambient space  $\mathbb{R}^{n+1}$  following the join construction of  $Q^n(c, d)$ , then eq. (2.1) becomes the system (cf. [4, 21])

$$\begin{cases} \Delta\Phi_1 = (\Lambda/c^2)\Phi_1 \\ \Delta\Phi_2 = (\Lambda/d^2)\Phi_2 \end{cases}$$

with

$$\Lambda = \frac{|d\Phi_1|^2/c^2 + |d\Phi_2|^2/d^2}{|\Phi_1|^2/c^4 + |\Phi_2|^2/d^4}$$

Such harmonic maps are real analytic.

Given two maps  $f_1 : \mathbb{S}^{m_1} \rightarrow \mathbb{S}^{n_1}$  and  $f_2 : \mathbb{S}^{m_2} \rightarrow \mathbb{S}^{n_2}$ , we consider their join  $f_1 * f_2$ , which is a map between ellipsoids

$$\phi = f_1 * f_2 : Q^{m_1+m_2+1}(a, b) \rightarrow Q^{n_1+n_2+1}(c, d)$$

indeed, for any continuous function  $\varphi : [0, \pi/2] \rightarrow [0, \pi/2]$  with  $\varphi(0) = 0, \varphi(\pi/2) = \pi/2$ , we can define

$$\phi(z) = c \cos \varphi(t) \cdot f_1(x) + d \sin \varphi(t) \cdot f_2(y) \quad (2.2)$$

for  $x \in \mathbb{S}^{m_1}, y \in \mathbb{S}^{m_2}$ , and  $0 \leq s \leq \pi/2$ . We assume  $m_1, m_2 \geq 1$ .

**Lemma 2.2.** *Let  $\Phi = i \circ \phi$  the composition of  $\phi$  with the canonical embedding  $i$  of  $Q^n(c, d)$  into  $\mathbb{R}^{n+1}$ . Then  $\phi$  is symphonic map iff*

$$\Delta^{\Phi^*h}\Phi = \left( |P^{-1/2}\Phi^*h|^2 / |P^{-1}\Phi|^2 \right) P^{-1}\Phi \quad (2.3)$$

where we borrowed the notations in [14],

$$\Delta^{f^*h}f^\alpha = \sum_{i,j=1}^m \frac{\partial}{\partial x^j} \left( (f^*h)^{ij} \frac{\partial f^\alpha}{\partial x^i} \right) \quad (2.4)$$

and

$$(f^*h)^{ij} = \sum_{k,\ell} g^{ik} g^{j\ell} (f^*h)(e_k, e_\ell).$$

If we write  $\Phi = (\Phi_1, \Phi_2)$ , the components being the projections on the factors of the ambient space  $\mathbb{R}^{n+1}$  following the join construction of  $Q^n(c, d)$ , then (2.3) becomes the system (cf. [4, 21])

$$\begin{cases} \Delta^{\Phi_1^* h} \Phi_1 = (\Lambda/c^2) \Phi_1 \\ \Delta^{\Phi_2^* h} \Phi_2 = (\Lambda/d^2) \Phi_2 \end{cases}$$

with

$$\Lambda = \frac{|\Phi_1^* h|^2 / c^2 + |\Phi_2^* h|^2 / d^2}{|\Phi_1|^2 / c^4 + |\Phi_2|^2 / d^4}$$

*Proof.* We follow the proof of [14, Lemma 1]. Let  $A$  be the second fundamental form of  $Q^n(c, d)$  in  $\mathbb{R}^{n+1}$

$$\Delta^{\Phi^* h} \Phi = A(\Phi^* h, \Phi^* h)$$

and

$$A(X, Y)|_{x,y} = \frac{|X|^2 / c^2 + |Y|^2 / d^2}{|x|^2 / c^4 + |y|^2 / d^4} \begin{pmatrix} x & y \\ c & d \end{pmatrix}$$

The Lemma follows from this. □

**Theorem 2.1.** *Let  $f_1 : \mathbb{S}^{m_1}(r_1) \rightarrow \mathbb{S}^{n_1}$ ,  $f_2 : \mathbb{S}^{m_2}(r_2) \rightarrow \mathbb{S}^{n_2}$  be two symphonic maps. Then the following two conditions are equivalent:*

- (1) *The join  $f_1 * f_2$  given by (2.2) is a symphonic map.*
- (2)  *$\varphi(t)$  satisfies the ODE*

$$\begin{aligned} & \frac{k(\alpha)^2}{h^2} (\varphi'(t)^3)' - \frac{k(\alpha)^2}{h^2} \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} - 2 \frac{h'(t)}{h(t)} \right) \varphi'(t)^3 + \frac{2kk'}{h^2} \varphi'(t)^4 \\ & + \left( \frac{\cos^2 \varphi(t)}{\cos^4 t} \|f_1^* g_{S^{n_1}}\|^2 - \frac{\sin^2 \varphi(t)}{\sin^4 t} \|f_2^* g_{S^{n_2}}\|^2 \right) \sin \varphi(t) \cos \varphi(t) = 0 \end{aligned} \tag{2.5}$$

**Remark 2.1.** *This equation has extra terms compared with the equation of Theorem 1 in [14].*

*Proof.* Using the notation in (2.4), we get

$$\begin{aligned}
& \Delta^{(f_1 * f_2)^* g_{Q(c,d)}} \\
&= \frac{1}{\sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})}} \frac{\partial}{\partial x_i} \left\{ \sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})} \frac{1}{a^2 \cos^2 t} (g_{\mathbb{S}^{m_1}})^{ia} \frac{1}{a^2 \cos^2 t} (g_{\mathbb{S}^{m_1}})^{jb} c^2 \cos^2 \varphi(t) \right. \\
& \quad \left. (f_1^* g_{\mathbb{S}^{n_1}})_{ab} \frac{\partial}{\partial x^j} \right\} \\
&+ \frac{1}{\sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})}} \frac{\partial}{\partial y^k} \left\{ \sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})} \frac{1}{b^2 \sin^2 t} (g_{\mathbb{S}^{m_2}})^{kc} \frac{1}{b^2 \sin^2 t} (g_{\mathbb{S}^{m_2}})^{ld} c^2 \sin^2 \varphi(t) \right. \\
& \quad \left. (f_1^* g_{\mathbb{S}^{n_1}})_{cd} \frac{\partial}{\partial y^l} \right\} \\
&+ \frac{1}{\sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})}} \frac{\partial}{\partial t} \left\{ \sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})} \frac{1}{h(t)^2} \varphi'(t)^2 k^2(\varphi) \frac{\partial}{\partial t} \right\}
\end{aligned}$$

Thus, we get

$$\begin{aligned}
\Delta^{(f_1 * f_2)^* g_{Q(c,d)}} &= \frac{c^2 \cos^2 \varphi(t)}{a^4 \cos^4 t} \frac{1}{\sqrt{\det(g_{\mathbb{S}^{m_1}(r_2)})}} \sum_{k,l} \frac{\partial}{\partial x^i} \left\{ \sqrt{\det(g_{\mathbb{S}^{m_1}(r_2)})} (f_2^* g_{\mathbb{S}^{n_1}(R_2)})^{ij} \frac{\partial}{\partial y^j} \right\} \\
&+ \frac{d^2 \sin^2 \varphi(t)}{b^4 \sin^4 t} \frac{1}{\sqrt{\det(g_{\mathbb{S}^{m_2}(r_2)})}} \sum_{k,l} \frac{\partial}{\partial y^k} \left\{ \sqrt{\det(g_{\mathbb{S}^{m_2}(r_2)})} (f_2^* g_{\mathbb{S}^{n_2}(R_2)})^{kl} \frac{\partial}{\partial y^l} \right\} \\
&+ \frac{1}{\left(\frac{\cos t}{r_1}\right)^{m_1} \left(\frac{\sin t}{r_2}\right)^{m_2}} \frac{\partial}{\partial t} \left\{ \left(\frac{\cos t}{r_1}\right)^{m_1} \left(\frac{\sin t}{r_2}\right)^{m_2} \frac{1}{h^2(t)} \varphi'(t)^2 k^2(\varphi) \frac{\partial}{\partial t} \right\} \\
&= \frac{c^2 \cos^2 \varphi(t)}{a^4 \cos^4 t} \Delta_{f_1^* g_{\mathbb{S}^{n_1}}} + \frac{d^2 \sin^2 \varphi(t)}{b^4 \sin^4 t} \Delta_{f_2^* g_{\mathbb{S}^{n_2}}} \\
&+ \frac{\partial}{\partial t} \left( \frac{1}{h^2(t)} \varphi'(t)^2 k^2(\varphi) \frac{\partial}{\partial t} \right) - \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} \right) \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t}.
\end{aligned} \tag{2.6}$$

On the other hand,

$$\begin{aligned}
& \left\| ((f_1 * f_2))^* g_{\mathbb{S}^{n_1+n_2+1}(1)} \right\|^2 \\
&= \sum_{i,j,k,\ell} (g_{Q^{m_1+m_2+1}(a,b)})^{ik} (g_{Q^{m_1+m_2+1}(a,b)})^{j\ell} ((f_1 * f_2)^* g_{\mathbb{S}^{n_1+n_2+1}(1)})_{ij} ((f_1 * f_2)^* g_{\mathbb{S}^{n_1+n_2+1}(1)})_{k\ell} \\
&= \sum_{a,b,c,d} \left( \frac{1}{a^2 \cos^2 t} (g_{\mathbb{S}^{m_1}(r_1)})^{ac} \right) \left( \frac{1}{a^2 \cos^2 t} (g_{\mathbb{S}^{m_1}(r_1)})^{bd} \right) \\
&\quad \times \left( c^2 \cos^2 \varphi(t) (f_1^* g_{\mathbb{S}^{n_1}(1)})_{ab} \right) \left( c^2 \cos^2 \varphi(t) (f_1^* g_{\mathbb{S}^{n_1}(1)})_{cd} \right) \\
&\quad + \sum_{p,q,r,s} \left( \frac{1}{b^2 \sin^2 t} (g_{\mathbb{S}^{m_2}(r_2)})^{pr} \right) \left( \frac{1}{b^2 \sin^2 t} (g_{\mathbb{S}^{m_2}(r_2)})^{qs} \right) \\
&\quad \times \left( d^2 \sin^2 \varphi(t) (f_2^* g_{\mathbb{S}^{n_2}(1)})_{pq} \right) \left( d^2 \sin^2 \varphi(t) (f_2^* g_{\mathbb{S}^{n_2}(1)})_{rs} \right) + \frac{k^4(\varphi(t))}{h^4(t)} \varphi'(t)^4 \\
&= \left( \frac{b^4}{a^4} \right) \left( \frac{\cos^2 \varphi(t)}{\cos^2 t} \right)^2 \|f_1^* g_{\mathbb{S}^{n_1}}\|^2 + \left( \frac{d^4}{b^4} \right) \left( \frac{\sin^2 \varphi(t)}{\sin^2 t} \right)^2 \|f_2^* g_{\mathbb{S}^{n_2}}\|^2 + \frac{k^4(\varphi(t))}{h^4(t)} \varphi'(t)^4.
\end{aligned} \tag{2.7}$$

Since  $f_1 * f_2$  is symphonic map, we have (cf. Theorem 1 in [14] )

$$\begin{aligned}
\Delta^{((f_1 * f_2))^* g_{\mathbb{S}^{n_1+n_2+1}(1)}} (\cos \varphi(t) f_1(x)) + \frac{\Lambda}{c^2} \cos \varphi(t) f_1(x) &= 0 \\
\Delta^{((f_1 * f_2))^* g_{\mathbb{S}^{n_1+n_2+1}(1)}} (\sin \varphi(t) f_2(x)) + \frac{\Lambda}{d^2} \sin \varphi(t) f_2(x) &= 0
\end{aligned} \tag{2.8}$$

By (2.6) and (2.8), we get

$$\begin{aligned}
& \frac{c^2 \cos^2 \varphi(t)}{a^4 \cos^4 t} \Delta^{f^* g_{\mathbb{S}^n}}|_{\mathbb{S}^{m_1}} \cos(\varphi(t) f_1) + \frac{d^2 \sin^2 \varphi(t)}{b^4 \sin^4 t} \Delta^{f^* g_{\mathbb{S}^n}}|_{\mathbb{S}^{m_2}} (\cos \varphi(t) f_1) \\
&+ \frac{\partial}{\partial t} \left( \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t} \right) (\cos \varphi(t) f_1) - \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} \right) \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t} (\cos \varphi(t) f_2) \\
&+ \frac{\Lambda}{c^2} \cos \varphi(t) f_1 = 0.
\end{aligned} \tag{2.9}$$

and

$$\begin{aligned}
& \frac{c^2 \cos^2 \varphi(t)}{a^4 \cos^4 t} \Delta^{f^* g_{\mathbb{S}^n}}|_{\mathbb{S}^{m_1}} \sin(\varphi(t) f_2) + \frac{d^2 \sin^2 \varphi(t)}{b^4 \sin^4 t} \Delta^{f^* g_{\mathbb{S}^n}}|_{\mathbb{S}^{m_2}} (\sin \varphi(t) f_2) \\
&+ \frac{\partial}{\partial t} \left( \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t} \right) (\sin \varphi(t) f_2) - \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} \right) \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t} (\sin \varphi(t) f_2) \\
&+ \frac{\Lambda}{d^2} \sin \varphi(t) f_2 = 0.
\end{aligned} \tag{2.10}$$

Then, by the same trick in [14], we can see that (2.9)  $\times c^2 \sin(\varphi(t))$  - (2.10)  $\times d^2 \cos(\varphi(t)) = 0$  is equivalent to

$$\begin{aligned} & \frac{k(\alpha)^2}{h^2} (\varphi'(t)^3)' - \frac{k(\alpha)^2}{h^2} \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} - 2 \frac{h'(t)}{h(t)} \right) \varphi'(t)^3 + \frac{2kk'}{h^2} \varphi'(t)^4 \\ & + \left( \frac{c^4 \cos^2 \varphi(t)}{a^4 \cos^4 t} \|f_1^* g_{S^n}\|^2 - \frac{d^4 \sin^2 \varphi(t)}{b^4 \sin^4 t} \|f_2^* g_{S^n}\|^2 \right) \sin \varphi(t) \cos \varphi(t) = 0 \end{aligned} \quad (2.11)$$

and that  $c^2 \cos(\varphi(t)) \times (2.9) + (2.10) \times d^2 \sin(\varphi(t)) = 0$  is equivalent to (2.7).  $\square$

On the other hand, we have

$$\|d(f_1 * f_2)\|^2 = \frac{c^2 \cos^2 \varphi(t)}{a^2 \cos^2(t)} |df_1|^2 + \frac{d^2 \sin^2 \varphi(t)}{b^2 \sin^2(t)} |df_2|^2 + \frac{k^2(\varphi(t))}{h^2(t)} \varphi'(t)^2$$

### 3 The existence of symphonic join

In order to apply the variation method to obtain the existence of symphonic map, we need to consider the Hilbert space

$$X = \left\{ \alpha \in W^{1,4}([0, \pi/2], \mathbb{R}) : \|\alpha\|^4 = \int_0^{\pi/2} (\dot{\alpha}^4 + \alpha^4) v ds < \infty \right\}$$

where  $v$  is the function on  $[0, \frac{\pi}{2}]$  and the functional

$$J(\varphi) = \int_0^{\frac{\pi}{2}} \left( \frac{k^4(\varphi(t))}{h^4(t)} \varphi'(t)^4 + a_1 \frac{\cos^4 \varphi(t)}{\cos^4 t} + a_2 \frac{\sin^4 \varphi(t)}{\sin^4 t} \right) \cos^{m_1} t \sin^{m_2} t dt$$

where we used the following notations

$$a_1 = \frac{c^4}{a^4} \frac{\mathcal{F}(f_1)}{\text{Vol}(\mathbb{S}^{m_1}(r_1))}, \quad a_2 = \frac{d^4}{b^4} \frac{\mathcal{F}(f_2)}{\text{Vol}(\mathbb{S}^{m_2}(r_2))}.$$

$$h = [b^2 \sin^2(t) + a^2 \cos^2(t)]^{1/2}, \quad k(\varphi) = [d^2 \sin^2 \varphi + c^2 \cos^2 \varphi]^{1/2}, \quad \mathcal{F}(f_1) = \int_M |f_1^* g_{S^{n_1}}|^2, \\ \mathcal{F}(f_2) = \int_M |f_2^* g_{S^{n_2}}|^2$$

Similar to Lemma 2.8 in [4], we have

**Lemma 3.1.** *There is a constant  $C$  such that for all  $\alpha \in X$*

$$\left. \begin{aligned} & \int_0^{\pi/2} \alpha^4 \sin^{p-4} \cos^q ds \\ & \int_0^{\pi/2} \alpha^4 \sin^p \cos^{q-4} ds \end{aligned} \right\} \leq C \int_0^{\pi/2} [\dot{\alpha}^4 + \alpha^4] \sin^p \cos^q ds.$$

If either  $p = 1$  or  $q = 1$ , we extend the definition of  $J$ , allowing it to assume the value  $+\infty$

*Proof.* This follows from Sobolev inequality for the Riemannian manifolds  $([0, \frac{\pi}{2}], \sin^p \cos^{q-4} ds)$  and  $([0, \frac{\pi}{2}], \sin^{p-4} \cos^q ds)$ .  $\square$

Thus, for  $p, q > 1$ , the functional  $J$  is defined and smooth on  $X$ . Since  $h$  is bounded above and below by positive constants.

**Lemma 3.2.** *There exists a minimizer  $\hat{\varphi}$  of  $J(\varphi)$  in  $W^{1,4}((0, \frac{\pi}{2}))$  with the conditions*

$$\hat{\varphi}(0) = 0, \quad \hat{\varphi}\left(\frac{\pi}{2}\right) = \frac{\pi}{2} \text{ and } 0 \leq \hat{\varphi} \leq \frac{\pi}{2}.$$

Then  $\hat{\varphi}(t)$  satisfies the Euler-Lagrange equation (2.5)

**Remark 3.1.** *Unlike the second order ODE for harmonic map, here we deal with one order ODE. Thus, it is much simpler.*

*Proof.* We modify the proof of [14].

$$\mathcal{A} = \left\{ \psi \in W^{1,4} \left( \left(0, \frac{\pi}{2}\right) \right) \mid \varphi(0) = 0, \varphi\left(\frac{\pi}{2}\right) = \frac{\pi}{2}, 0 \leq \varphi(t) \leq \frac{\pi}{2} \right\}.$$

Take a minimizing sequence  $\varphi_i$  in  $\mathcal{A}$ , we get a weak limit  $\varphi_\infty$ . Since the functional  $J$  is lower semicontinuous,  $\varphi_\infty$  is a minimizer in  $\mathcal{A}$ . By Ding's method (cf. Lemma 2.1 in [2]), we get a weak limit  $\varphi_\infty$  which is also a minimizer in

$$\tilde{\mathcal{A}} = \left\{ \varphi \in L^{1,4} \left( \left(0, \frac{\pi}{2}\right) \right) \mid \varphi(0) = 0, \varphi\left(\frac{\pi}{2}\right) = \frac{\pi}{2} \right\}.$$

and therefore  $\varphi_\infty$  solves (2.5), set  $\hat{\varphi} = \varphi_\infty$  and we finished the proof.  $\square$

By the Sobolev embedding from  $L^{1,4}((0, \frac{\pi}{2}))$  into  $C^0([0, \frac{\pi}{2}])$ , we have  $\hat{\varphi}(t) \in C^0([0, \frac{\pi}{2}])$ .

**Lemma 3.3.**  $\hat{\varphi} \in C^\infty((0, \frac{\pi}{2}))$

*Proof.* Let  $\psi(t) = k^2(\varphi)(\varphi'(t))^3$ . By Theorem 2.1, we have

$$\psi'(t) + A(t)\psi(t) + B(t) = 0$$

where

$$A(t) = \frac{k(\alpha)^2}{h^2} \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} - 2 \frac{h'(t)}{h(t)} \right)$$

$$B(t) = -\sin \hat{\varphi}(t) \cos \hat{\varphi}(t) \left( \frac{d^4 \sin^2 \hat{\varphi}(t)}{b^4 \sin^4 t} \|f_1^* g_{g^{n_1(1)}}\|^2 - \frac{c^4 \cos^2 \hat{\varphi}(t)}{a^4 \cos^4 t} \|f_2^* g_{s^{2_2(t)}}\|^2 \right).$$



Fix any  $t_0 \in (0, \frac{\pi}{2})$ . For any  $t \in (0, \frac{\pi}{2})$ , as in [14], we have

$$\begin{aligned} |\psi(t) - \psi(t_0)| &= \left| \int_{r_0}^t \psi(t) dt \right| \\ &\leq \left\{ \int_{t_0}^t |A(t)|^4 dt \right\}^{\frac{1}{4}} \left\{ \int_{t_0}^t |\hat{\varphi}'(t)|^4 dt \right\}^{\frac{3}{4}} + \left| \int_{t_0}^t |B(t)| dt \right| \\ &\longrightarrow 0 \text{ as } t \longrightarrow t_0. \end{aligned}$$

since  $\hat{\varphi} \in W^{1,4}(0, \frac{\pi}{2})$ , and  $\int_{t_0-\varepsilon}^{t_0+\varepsilon} |A(t)|^4 dt < \infty$  and  $\int_{t_0-\varepsilon}^{t_0+\varepsilon} |B(t)|^4 dt < \infty$ . Thus we have  $\psi \in C^0((0, \frac{\pi}{2}))$ . We use a bootstrap argument for the equality

$$\psi'(t) = -A(t)\psi(t) - B(t)$$

implies that  $\hat{\varphi}(t)$  is of class  $C^\infty$ . □

The above lemmas imply the existence of the join of maps. Just as in [14], we get

**Theorem 3.1.** *For any two symphonic maps*

$$\begin{aligned} f_1 : S^{m_1}(r_1) &\longrightarrow S^{m_1}(1) \\ f_2 : S^{m_2}(r_2) &\longrightarrow S^{m_2}(1) \end{aligned}$$

there exists a symphonic join

$$f_1 * f_2 : Q^{m_1+m_2+1}(a, b) \longrightarrow Q^{n_1+n_2+1}(a, b)$$

ie., a symphonic map which is a join of maps  $f_1$  and  $f_2$ .

## 4 Hopf construction of symphonic map

Let  $X = \{(x_1, x_2, \dots, x_{n+1}) \mid \frac{x_1^2+x_2^2+\dots+x_n^2}{c^2} + \frac{x_{n+1}^2}{d^2} = 1\}$ . Topologically, the *Hopf* construction on a map  $f : S^{m_1} \times S^{m_2} \rightarrow S^{n-1}$  is the map  $h_f : Q^{m_1+m_2+1}(a, b) \rightarrow X$  given by

$$h_f(ax_1 \cos t, bx_2 \sin t) := (cf(x_1, x_2) \cos \varphi(t), d \sin \varphi(t)) \quad (4.1)$$

where  $x \in \mathbb{S}^{m_1+m_2-1}$  can be parametrized as  $(x_1 \sin s, x_2 \cos s)$  with  $x_1 \in \mathbb{S}^{m_1}, x_2 \in \mathbb{S}^{m_2}, t \in [0, \pi/2]$ . The metric tensor on ellipsoid  $Q^{m_1+m_2+1}(a, b)$  is

$$g = (a^2 \cos^2 t) g_{\mathbb{S}^{m_1}} + (b^2 \sin^2 t) g_{\mathbb{S}^{m_2}} + h^2(t) dt^2$$

The metric tensor on ellipsoid  $X$  is

$$g = (c^2 \cos^2 s) g_{\mathbb{S}^{n-1}} + (d^2 \sin^2 s) g_{\mathbb{S}^1} + k^2(s) ds^2$$

Here,

$$h(t) = [b^2 \sin^2 t + a^2 \cos^2 t]^{1/2}, k(s) = [c^2 \sin^2 s + d^2 \cos^2 s]^{1/2}$$

**Lemma 4.1.** *Given a map  $f : S^{m_1}(1) \times S^{m_2}(1) \rightarrow S^{n-1}(1)$ , the associated Hopf map  $h_f : Q^{m_1+m_2+1}(a, b) \rightarrow X$  given by (4.1).  $f_1(\cdot) = f(\cdot, y)$ ,  $f(\cdot) = f_1(x, \cdot)$ , where  $x \in \mathbb{S}^{m_1}(1)$ ,  $y \in \mathbb{S}^{m_2}(1)$ . Assume that  $f_1$  and  $f_2$  are both symphonic map.*

*Then the following two conditions are equivalent:*

- (1) *The join  $h_f$  is a symphonic map.*
- (2)  *$\varphi(t)$  satisfies that*

$$\begin{aligned} & \frac{k^2(\varphi)}{h^2}(\varphi'(t)^3)' - \frac{k^2(\varphi)}{h^2} \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} - 2 \frac{h'(t)}{h(t)} \right) \varphi'(t)^3 + \frac{2kk'}{h^2} \varphi'(t)^4 \\ & + \left( \frac{c^4 \cos^2 \varphi(t)}{a^4 \cos^4 t} \|f^* g_{\mathbb{S}^n}\|^2 + \frac{c^4 \sin^2 \varphi(t)}{b^4 \sin^4 t} \|f^* g_{\mathbb{S}^n}\|^2 \right) \sin \varphi(t) \cos \varphi(t) = 0 \end{aligned} \quad (4.2)$$

*Proof.* Using the notation in (2.4), we get

$$\begin{aligned} & \Delta^{f^* g_{\mathbb{S}^{n-1}}} \\ & = \frac{1}{\sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})}} \frac{\partial}{\partial x_i} \left\{ \sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})} \frac{1}{a^2 \cos^2 t} (g_{\mathbb{S}^{m_1}})^{ia} \frac{1}{a^2 \cos^2 t} (g_{\mathbb{S}^{m_1}})^{jb} c^2 \cos^2 \varphi(t) \right. \\ & \quad \left. (f^* g_{\mathbb{S}^{n_1}})_{ab} \frac{\partial}{\partial x^j} \right\} \\ & + \frac{1}{\sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})}} \frac{\partial}{\partial y^j} \left\{ \sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})} \frac{1}{b^2 \sin^2 t} (g_{\mathbb{S}^{m_2}})^{kc} \frac{1}{b^2 \sin^2 t} (g_{\mathbb{S}^{m_2}})^{ld} c^2 \cos^2 \varphi(t) \right. \\ & \quad \left. (f^* g_{\mathbb{S}^{n_1}})_{cd} \frac{\partial}{\partial y^j} \right\} \\ & + \frac{1}{\sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})}} \frac{\partial}{\partial t} \left\{ \sqrt{\det(g_{Q^{m_1+m_2+1}(a,b)})} \frac{1}{h(t)^2} \varphi'(t)^2 \frac{\partial}{\partial t} \right\} \end{aligned} \quad (4.3)$$

Thus,

$$\begin{aligned}
\Delta^{f^* g_{\mathbb{S}^{n-1}}} &= \frac{c^2 \cos^2 \varphi(t)}{a^4 \cos^4 t} \frac{1}{\sqrt{\det(g_{\mathbb{S}^{m_1}(r_2)})}} \sum_{k,\ell} \frac{\partial}{\partial x^i} \left\{ \sqrt{\det(g_{\mathbb{S}^{m_1}(r_2)})} (f^* g_{\mathbb{S}^n(R_2)})^{ij} \frac{\partial}{\partial x^j} \right\} \\
&+ \frac{c^2 \cos^2 \varphi(t)}{b^4 \sin^4 t} \frac{1}{\sqrt{\det(g_{\mathbb{S}^{m_2}(r_2)})}} \sum_{k,\ell} \frac{\partial}{\partial y^k} \left\{ \sqrt{\det(g_{\mathbb{S}^{m_2}(r_2)})} (f^* g_{\mathbb{S}^n(R_2)})^{k\ell} \frac{\partial}{\partial y^\ell} \right\} \\
&+ \frac{1}{\left(\frac{\cos t}{r_1}\right)^{m_1} \left(\frac{\sin t}{r_2}\right)^{m_2}} \frac{\partial}{\partial t} \left\{ \left(\frac{\cos t}{r_1}\right)^{m_1} \left(\frac{\sin t}{r_2}\right)^{m_2} \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t} \right\} \\
&= \frac{c^2 \cos^2 \varphi(t)}{a^4 \cos^4 t} \Delta^{f^* g_{\mathbb{S}^n}|_{S^{m_1}}} + \frac{c^2 \cos^2 \varphi(t)}{b^4 \sin^4 t} \Delta^{f^* g_{\mathbb{S}^n}|_{S^{m_2}}} \\
&+ \frac{\partial}{\partial t} \left( \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t} \right) - \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} \right) \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t}.
\end{aligned} \tag{4.4}$$

So,

$$\begin{aligned}
&\|f^* g_{\mathbb{S}^{n-1}}\|^2 \\
&= \sum_{i,j,k,\ell} (g_{Q^{m_1+m_2+1}(a,b)})^{ik} (g_{Q^{m_1+m_2+1}(a,b)})^{j\ell} \times (f^* g_{\mathbb{S}^n(1)})_{ij} (f^* g_{\mathbb{S}^n(1)})_{k\ell} \\
&= \sum_{a,b,c,d} \left( \frac{1}{a^2 \cos^2 t} (g_{\mathbb{S}^{m_1}(r_1)})^{ac} \right) \left( \frac{1}{a^2 \cos^2 t} (g_{\mathbb{S}^{m_1}(r_1)})^{bd} \right) \\
&\quad \times \left( c^2 \cos^2 \varphi(t) (f_1^* g_{\mathbb{S}^{n-1}})_{ab} \right) \left( c^2 \cos^2 \varphi(t) (f_1^* g_{\mathbb{S}^{n-1}(1)})_{cd} \right) \\
&+ \sum_{p,q,r,s} \left( \frac{1}{b^2 \sin^2 t} (g_{\mathbb{S}^{m_2}(r_2)})^{pr} \right) \left( \frac{1}{b^2 \sin^2 t} (g_{\mathbb{S}^{m_2}(r_2)})^{qs} \right) \\
&\quad \times \left( c^2 \cos^2 \varphi(t) (f_2^* g_{\mathbb{S}^{n-1}})_{pq} \right) \left( c^2 \cos^2 \varphi(t) (f_2^* g_{\mathbb{S}^{n-1}})_{rs} \right) + \frac{k^2(\varphi(t))}{h^4(t)} \varphi'(t)^4 \\
&= \left( \frac{c^2}{a^2 \cos^2 t} \right)^2 \cos^4 \varphi(t) \|f_1^* g_{\mathbb{S}^{n-1}}\|^2 + \left( \frac{c^2}{b^2 \sin^2 t} \right)^2 \cos^4 \varphi(t) \|f_2^* g_{\mathbb{S}^{n-1}}\|^2 + \frac{k^2(\varphi(t))}{h^4(t)} \varphi'(t)
\end{aligned} \tag{4.5}$$

Since the Hopf map is symphonic map, we get

$$\Delta^{f^* g_{\mathbb{S}^n(1)}} (\cos \varphi(t) f(x)) + \frac{\Lambda}{c^2} \cos \varphi(t) f(x) = 0 \tag{4.6}$$

$$\Delta^{f^* g_{\mathbb{S}^n(1)}} (\sin \varphi(t)) + \frac{\Lambda}{d^2} \sin \varphi(t) = 0 \tag{4.7}$$

Then,

$$\begin{aligned} & \left[ \frac{c^2 \cos^2 \varphi(t)}{a^4 \cos^4 t} \Delta^{f^* g_{S^{m_1}}} \cos \varphi(t) f + \frac{c^2 \cos^2 \varphi(t)}{b^4 \sin^4 t} \Delta^{f^* g_{S^{m_2}}} \cos \varphi(t) f \right. \\ & \left. + \frac{\partial}{\partial t} \left( \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t} \right) (\cos \varphi(t) f) - \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} \right) \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t} (\cos \varphi(t) f) \right] \\ & + \frac{\Lambda}{c^2} \cos \varphi(t) f = 0. \end{aligned} \quad (4.8)$$

and

$$\begin{aligned} & \left[ \frac{\partial}{\partial t} \left( \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t} \right) (\sin \varphi(t) f) - \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} \right) \frac{1}{h^2(t)} \varphi'(t)^2 \frac{\partial}{\partial t} (\sin \varphi(t) f) \right] \\ & + \frac{\Lambda}{d^2} \sin \varphi(t) = 0. \end{aligned} \quad (4.9)$$

Since  $f$  is symphonic map,  $c^2 \sin(\varphi(t)) \times (4.8) - d^2 \cos(\varphi(t)) \times (4.9) = 0$  is equivalent to

$$\begin{aligned} & \frac{k^2(\varphi)}{h^2} (\varphi'(t)^3)' - \frac{k^2(\varphi)}{h^2} \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} - 2 \frac{h'(t)}{h(t)} \right) \varphi'(t)^3 + \frac{2kk'}{h^2} \varphi'(t)^4 \\ & + \left( \frac{c^4 \cos^2 \varphi(t)}{a^4 \cos^4 t} \|f_1^* g_{S^n}\|^2 + \frac{c^4 \cos^2 \varphi(t)}{b^4 \sin^4 t} \|f_2^* g_{S^n}\|^2 \right) \sin \varphi(t) \cos \varphi(t) = 0 \end{aligned} \quad (4.10)$$

Moreover, we also see that  $c^2 \cos \varphi \times (4.8) + d^2 \sin \varphi \times (4.9) = 0$  is just (4.5).  $\square$

By a direct computation, we have

$$\|dh_f\|^2 = \frac{c^2 \cos^2 \varphi(t)}{a^2 \cos^2(t)} |df_1|^2 + \frac{c^2 \cos^2 \varphi(t)}{b^2 \sin^2(t)} |df_2|^2 + \frac{\kappa^2(\varphi)}{h^2(t)} \varphi'(t)^2$$

where  $f_1(\cdot) = f_1(\cdot, y)$ ,  $f_2(\cdot) = f_1(x, \cdot)$ , where  $x \in \mathbb{S}^{m_1}$ ,  $y \in \mathbb{S}^{m_2}$ .

$$J(\varphi) = \int_0^{\frac{\pi}{2}} \left( \frac{k^4(\varphi(t))}{h^4(t)} \varphi'(t)^4 + a_1 \frac{\sin^4 \varphi(t)}{\cos^4 t} + a_2 \frac{\sin^4 \varphi(t)}{\sin^4 t} \right) \cos^{m_1} t \sin^{m_2} t dt$$

$$a_1 = \frac{\mathcal{F}(c^4 f_1)}{a^4 \text{Vol}(\mathbb{S}^{m_1}(r_1))}, \quad a_2 = \frac{c^4 \mathcal{F}(f_2)}{b^4 \text{Vol}(\mathbb{S}^{m_2}(r_2))}.$$

where  $\text{Vol}(A)$  denotes the volume of  $A$ ,  $\mathcal{F}(f_1) = \int_M |f_1^* g_{S^n}|^2$ ,  $\mathcal{F}(f_2) = \int_M |f_2^* g_{S^n}|^2$

**Lemma 4.2.** *There exists a minimizer  $\hat{\varphi}$  of  $J(\varphi)$  in  $L^{1,4}((0, \frac{\pi}{2}))$  with the conditions*

$$\hat{\varphi}(0) = 0, \quad \hat{\varphi}\left(\frac{\pi}{2}\right) = \frac{\pi}{2} \text{ and } 0 \leq \hat{\varphi}(t) \leq \frac{\pi}{2}.$$

where  $L^{1,4}((0, \frac{\pi}{2}))$  denotes the Sobolev space of  $L^4$ -functions with derivatives in  $L^4$ . Furthermore,  $\hat{\varphi}(t)$  satisfies the Euler-Lagrange (eq. (4.10))

**Remark 4.1.** *Unlike the second order ODE for harmonic map, here we deal with one order ODE. Thus, it is much simpler.*

*Proof.* The proof is the same as Lemma 3.2. □

By the Sobolev embedding from  $L^{1,4}((0, \frac{\pi}{2}))$  into  $C^0([0, \frac{\pi}{2}])$ , we have  $\widehat{\varphi}(t) \in C^0([0, \frac{\pi}{2}])$ .

**Lemma 4.3.**  $\widehat{\varphi} \in C^\infty((0, \frac{\pi}{2}))$

*Proof.* Let  $\psi(t) = k(\alpha)^2 (\widehat{\varphi}'(t))^3$ . By Lemma 2 we have

$$\psi'(t) + A(t)\psi(t) + B(t) = 0$$

where

$$A(t) = \frac{k(\alpha)^2}{h^2(t)} \left( m_1 \frac{\sin t}{\cos t} - m_2 \frac{\cos t}{\sin t} - 2 \frac{h'(t)}{h(t)} \right)$$

$$B(t) = -h^2(t) \sin \widehat{\varphi}(t) \cos \widehat{\varphi}(t) \left( \frac{c^4 \cos^2 \widehat{\varphi}(t)}{b^4 \sin^4 t} \|f_1^* g_{g^n(1)}\|^2 + \frac{c^4 \cos^2 \widehat{\varphi}(t)}{a^4 \cos^4 t} \|f_2^* g_{S^{n_2}}\|^2 \right).$$

The rest proof is the same as [Lemma 3.3](#). □

**Theorem 4.1.** *For map  $f : S^{m_1} \times S^{m_2} \rightarrow S^{n-1}$ , its associated hopf map defined in [\(4.1\)](#) exists and is a symhonic map.*

## References

- [1] DING, W., FAN, H., AND LI, J. Harmonic Hopf constructions between spheres. II. *Calc. Var. Partial Differential Equations* 16, 3 (2003), 273–282. [2](#)
- [2] DING, W. Y. Symmetric harmonic maps between spheres. *Comm. Math. Phys.* 118, 4 (1988), 641–649. [1](#), [8](#)
- [3] DING, W. Y. Harmonic Hopf constructions between spheres. *Internat. J. Math.* 5, 6 (1994), 849–860. [1](#), [2](#)
- [4] EELLS, J., AND RATTO, A. Harmonic maps between spheres and ellipsoids. *Internat. J. Math.* 1, 1 (1990), 1–27. [1](#), [2](#), [3](#), [4](#), [7](#)
- [5] GASTEL, A. Singularities of first kind in the harmonic map and Yang-Mills heat flows. *Math. Z.* 242, 1 (2002), 47–62. [2](#)
- [6] GASTEL, A. On the harmonic Hopf construction. *Proc. Amer. Math. Soc.* 132, 8 (2004), 607–615. [1](#)

- [7] KAWAI, S., AND NAKAUCHI, N. Results of Liouville type for symphonic maps. *Differ. Geom. Appl.* 65 (2019), 147–159. 2
- [8] MISAWA, M., AND NAKAUCHI, N. A Hölder continuity of symphonic maps into the spheres. *Calc. Var. Partial Differential Equations* 55, 1 (2016), Art. 16, 20. 2
- [9] MISAWA, M., AND NAKAUCHI, N. Global existence for the heat flow of symphonic maps into spheres. *Adv. Differential Equations* 23, 9-10 (2018), 693–724. 2
- [10] MISAWA, M., AND NAKAUCHI, N. On the finite-time blow-up of symphonic map flows. *Differ. Integral Equ.* 36, 1-2 (2023), 93–131. 2
- [11] NAKAUCHI, N. Stress energy tensor for symphonic maps. *Boll. Unione Mat. Ital.* 12, 3 (2019), 431–440. 2
- [12] NAKAUCHI, N. Rotationally symmetric symphonic maps. *Ann. Global Anal. Geom.* 62, 1 (2022), 83–92. 2
- [13] NAKAUCHI, N. Two results for symphonic maps under assumptions on  $m$ -symphonic energy. *Results Math.* 77, 6 (2022), Paper No. 216, 23. 2
- [14] NAKAUCHI, N., AND TAKAKUWA, S. Symphonic join of maps between the spheres. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods* 108 (2014), 87–98. 2, 3, 4, 6, 7, 8, 9
- [15] NAKAUCHI, N., AND TAKENAKA, Y. A variational problem for pullback metrics. *Ric. Mat.* 60, 2 (2011), 219–235. 2
- [16] PENG, C., AND TANG, Z. Harmonic maps from spheres to spheres. *Topology* 37, 1 (1998), 39–43. 1
- [17] PENG, C. K., AND TANG, Z. Z. On representing homotopy classes of spheres by harmonic maps. *Topology* 36, 4 (1997), 867–879. 1
- [18] PETTINATI, V., AND RATTO, A. Existence and nonexistence results for harmonic maps between spheres. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* 17, 2 (1990), 273–282. 1
- [19] RATTO, A. Harmonic maps from deformed spheres to spheres. *Amer. J. Math.* 111, 2 (1989), 225–238. 1
- [20] RATTO, A. Harmonic maps of spheres and the Hopf construction. *Topology* 28, 3 (1989), 379–388. 1
- [21] RATTO, A. On the Dirichlet problem for harmonic maps into spheres or ellipsoids and equivariant harmonic maps from warped products. *Bull. Soc. Math. Belg. Sér. B* 41, 2 (1989), 145–168. 1, 2, 3, 4

- [22] SMITH, R. T. Harmonic mappings of spheres. *Bull. Amer. Math. Soc.* 78 (1972), 593–596. [1](#)
- [23] SMITH, R. T. Harmonic mappings of spheres. *Amer. J. Math.* 97 (1975), 364–385. [1](#)
- [24] TANG, Z. Harmonic Hopf constructions and isoparametric gradient maps. *Differential Geom. Appl.* 25, 5 (2007), 461–465. [1](#)
- [25] TODA, H. *Composition methods in homotopy groups of spheres*. Annals of Mathematics Studies, No. 49. Princeton University Press, Princeton, N.J., 1962. [1](#)