Polynomials with exponents in compact convex sets and associated weighted extremal functions -Fundamental results

Benedikt Steinar Magnússon, Álfheiður Edda Sigurðardóttir, Ragnar Sigurðsson and Bergur Snorrason

Abstract

This paper is a collection of fundamental results for the study of polynomial rings $\mathcal{P}^{S}(\mathbb{C}^{n})$ where the *m*-th degree polynomials have exponents restricted to mS, where $S \subseteq \mathbb{R}^{n}_{+}$ is compact, convex and $0 \in S$. We study the relationship between $\mathcal{P}^{S}(\mathbb{C}^{n})$ and the class $\mathcal{L}^{S}(\mathbb{C}^{n})$ of global plurisubharmonic functions where the growth is determined by the logarithmic supporting function of S. We present properties of their respective weighted extremal functions $\Phi^{S}_{K,q}$ and $V^{S}_{K,q}$ in connection with properties of S. Our ambition is to give detailed proofs with minimal assumptions of all results, thus creating a self-contained exposition.

AMS Subject Classification: 32U35. Secondary 32A08, 32A15, 32U15, 32W05.

1 Introduction

Approximation theory deals with problems of determining whether a given function in some prescribed function space can be approximated by functions in a certain subspace. The theorems of Runge and Mergelyan are prototypes of results from approximation theory. The Runge theorem states that every function f holomorphic in some neighborhood of a simply connected compact subset K of \mathbb{C} can be approximated in the uniform norm $\|\cdot\|_K$ on K by polynomials and the Mergelyan theorem states that it is enough to assume that f is continuous on K and holomorphic in the interior of K.

Quantitative approximation theory deals with problems of relating properties of the given function f to the error in approximation which is usually measured as the distance from f to a certain finite dimensional subspaces of the approximating subspace. The Bernstein-Walsh theorem is a prototype of a result from quantitative approximation theory. It states that a holomorphic function f defined in some neighborhood of a compact simply connected $K \subset \mathbb{C}$ extends as a holomorphic function to the sublevel set $\{z \in \mathbb{C}; g_{\mathbb{C}\setminus K}(z, \infty) < \log R\}$ if and only if $\overline{\lim_{m \to +\infty} d_m}(f, K)^{1/m} \leq 1/R$, where $R > 1, g_{\mathbb{C}\setminus K}(\cdot, \infty)$ is the Green function of $\mathbb{C}\setminus K$ with logarithmic pole at $\infty, d_m(f, K)$ is the distance from f to the space $\mathcal{P}_m(\mathbb{C})$ of all polynomials of degree $\leq m$ with respect to the uniform norm on K, and it is assumed that the domain $\mathbb{C}\setminus K$ is regular for the Dirichlet problem in the sense that $g_{\mathbb{C}\setminus K}(\cdot,\infty)$ vanishes at the boundary of K.

The Runge theorem is generalized to several variables where simple connectedness of K generalizes as polynomial convexity. This generalization is usually called the Oka-Weil theorem. There only exist fragmentary, but interesting, results generalizing the Mergelyan theorem to several complex variables. See Levenberg [20].

The origin of the subject of the present paper is the generalization by Siciak [30] of the Bernstein-Walsh theorem. There he introduced the extremal function $\Phi_K = \lim_{m \to \infty} \Phi_{K,m}$, where $\Phi_{K,m} = \sup\{|p|^{1/m}; p \in \mathcal{P}_m(\mathbb{C}^n), \|p\|_K \leq 1\}$ and $\mathcal{P}_m(\mathbb{C}^n)$ is the space of all polynomials of degree $\leq m$. In his work log Φ_K plays the role of $g_{\mathbb{C}\setminus K}(\cdot,\infty)$ and the regularity condition at ∂K is that $\Phi_K^*(z) = \lim_{\zeta \to z} \Phi_K(\zeta) \leq 1$ for every $z \in \partial K$. Siciak's paper is a seminal work on the understanding of the interrelation between quantitative approximation theory in several complex variables and pluripotential theory. He studied this subject further in many of his works, for example [31, 32, 33, 34]. See also Bloom's Appendix B in the monograph by Saff and Totik [28].

We will systematically work with subspaces of the polynomial space $\mathcal{P}(\mathbb{C}^n)$ in n variables. For every non-empty bounded subset S of \mathbb{R}^n_+ we let $\mathcal{P}^S_m(\mathbb{C}^n)$ denote the space of all polynomials p which can be written on the form $p(z) = \sum_{\alpha \in (mS) \cap \mathbb{N}^n} a_{\alpha} z^{\alpha}, z \in \mathbb{C}^n$, and let $\mathcal{P}^S(\mathbb{C}^n) = \bigcup_{m \in \mathbb{N}} \mathcal{P}^S_m(\mathbb{C}^n)$. The standard simplex $\Sigma = ch\{0, e_1, \ldots, e_n\}$, where ch A denotes the convex hull of a set A and (e_1, \ldots, e_n) is the standard basis in \mathbb{R}^n , yields the standard grading of polynomials of degree $\leq m$, that is $\mathcal{P}^{\Sigma}_m(\mathbb{C}^n) = \mathcal{P}_m(\mathbb{C}^n)$. If S is a compact convex set in \mathbb{R}^n_+ with $0 \in S$ then the space $\mathcal{P}^S(\mathbb{C}^n)$ forms a polynomial ring where the degree of a polynomial p is the minimal $m \in \mathbb{N}$ such that $p \in \mathcal{P}^S_m(\mathbb{C}^n)$.

The polynomial spaces $\mathcal{P}_m^S(\mathbb{C}^n)$ appear in Shiffman-Zelditch [29] with S as an integral polytope and later in Bayraktar [4] as sparse polynomials. These polynomial classes and their relation to pluripotential theory have been studied by Bloom, Levenberg and their collaborators [1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 21, 25]. Unfortunately, in some of these papers there are false results stated that we have not seen corrected. We point them out and correct them as far as we can.

In Section 2 we define the Siciak function with respect to S, E, and q for every function $q: E \to \mathbb{R} \cup \{+\infty\}$ on a subset E of \mathbb{C}^n by $\Phi^S_{E,q} = \overline{\lim}_{m \to \infty} \Phi^S_{E,q,m}$, where

$$\Phi_{E,q,m}^{S} = \sup\{|p|^{1/m}; p \in \mathcal{P}_{m}^{S}(\mathbb{C}^{n}), \|pe^{-mq}\|_{E} \le 1\}, \qquad m = 1, 2, 3, \dots$$

We drop S in the superscript if $S = \Sigma$ and q in the subscript if q = 0. The grading of the polynomial classes $\mathcal{P}_m^S(\mathbb{C}^n)$ implies that for every $z \in \mathbb{C}^n$ the sequence $(-\log(\Phi_{E,q,m}^S(z))^m)_{m\in\mathbb{N}^*}$ is subadditive and a lemma by Fekete implies that

$$\Phi_{E,q}^S = \lim_{m \to \infty} \Phi_{E,q,m}^S = \sup_{m \in \mathbb{N}^*} \Phi_{E,q,m}^S$$

without any restriction on S or q. This was first proved by Siciak [30, Theorem 6.1] for $S = \Sigma$. For the reader's convenience we prove the Fekete Lemma 2.3, because ingredients from the proof are needed in the proof of Proposition 2.2, where it is shown that the convergence is uniform on a compact subset X, if q is bounded below on Eand $\Phi_{E,q}^S$ is continuous on X. This statement on uniform convergence appears in many arguments in pluripotential theory. See for example Bloom and Shiffman [12, Lemma 3.2], Bayraktar [1, 4, Theorem 2.10] and Bayraktar, Hussung, Levenberg and Perera [8, Theorem 1.1]. The statements on uniform convergence in these papers all follow directly from Proposition 2.2 under the conditions that S is compact and convex with $0 \in S$ and q is bounded below.

In Section 3 we introduce the logarithmic supporting function H_S of S and the Lelong class with respect to S. The function H_S is defined on \mathbb{C}^{*n} as the supporting function φ_S of S, $\varphi_S(\xi) = \sup_{s \in S} \langle s, \xi \rangle$, $\xi \in \mathbb{R}^n$, in logarithmic coordinates extended to $\mathbb{C}^n \setminus \mathbb{C}^{*n}$ as an upper semicontinuous function and the Lelong class $\mathcal{L}^S(\mathbb{C}^n)$ with respect to S is defined as the set of all $u \in \mathcal{PSH}(\mathbb{C}^n)$ satisfying $u \leq c_u + H_S$, where c_u is a constant only depending on u and $\mathcal{PSH}(X)$ denotes the set of plurisubharmonic functions on an open set $X \subset \mathbb{C}^n$. We have $H_{\Sigma}(z) = \log^+ ||z||_{\infty}$, which implies that $\mathcal{L}^{\Sigma}(\mathbb{C}^n)$ is equal to the Lelong class $\mathcal{L}(\mathbb{C}^n)$.

What values H_S takes at points on the coordinate hyperplanes $\mathbb{C}^n \setminus \mathbb{C}^{*n}$ is opaque from its definition itself, but Proposition 3.3 provides an explicit formula. Additionally, this formula shows that the zero set $\mathcal{N}(H_S)$ of H_S may very well be unbounded. We prove in Proposition 3.4 that H_S extends from \mathbb{C}^{*n} as a continuous plurisubharmonic function on \mathbb{C}^n .

The usual grading of the polynomials can be characterized by growth, and so is also the case for $\mathcal{P}^{S}(\mathbb{C}^{n})$. It is clear that if $p \in \mathcal{P}_{m}^{S}(\mathbb{C}^{n})$ then $|p| \leq C_{p}e^{H_{S}}$ for some constant $C_{p} > 0$. In Theorem 3.6 we prove a Liouville type theorem which states that if $f \in \mathcal{O}(\mathbb{C}^{n})$ satisfies a growth estimate $|f(z)| \leq C(1+|z|)^{a}e^{mH_{S}(z)}$ for some C > 0and some a > 0 strictly less than the distance from mS to $\mathbb{N}^{n} \setminus mS$ in L^{1} -norm, then $f \in \mathcal{P}_{m}^{S}(\mathbb{C}^{n})$.

In Section 4 we define the Siciak-Zakharyuta function

$$V_{E,q}^S = \sup\{u \, ; \, u \in \mathcal{L}^S(\mathbb{C}^n), \, u|_E \le q\}$$

with respect to S and any $q: E \to \mathbb{R} \cup \{+\infty\}$ on a subset E of \mathbb{C}^n . We drop S in the superscript if $S = \Sigma$ and q in the subscript if q = 0. By Klimek [19, Example 5.1.1] $V_K(z) = \log^+(||z - a||/r)$ for any norm $|| \cdot ||$ and $K = \{z \in \mathbb{C}^n; ||z - a|| \le r\}, r > 0$, the closed ball in $|| \cdot ||$ with center a and radius r. These classical examples can not be generalized for V_E^S for the simple reason that the Lelong class $\mathcal{L}^S(\mathbb{C}^n)$ with respect to S does not need to be translation invariant. We prove in Proposition 4.3 that $V_E^S = H_S$ for any subset E of \mathbb{C}^n such that $\mathbb{T}^n \subseteq E \subseteq \mathcal{N}(H_S)$, where \mathbb{T} denotes the unit circle in \mathbb{C} and $\mathcal{N}(H_S)$ the zero set of H_S . This is a generalization of Bayraktar [4, Example 2.3].

We introduce *admissible weights* in Definition 4.4, where we follow Bloom [28, Appendix B] with the natural generalization for the Lelong classes with respect to S given in Bayraktar, Bloom, and Levenberg [6]. In Proposition 4.5 we prove that for every compact convex $S \subset \mathbb{R}^n_+$ with $0 \in S$ and admissible weight q on a closed subset E the upper regularization $V_{E,q}^{S*}$ of $V_{E,q}^S$ is in $\mathcal{L}^S_+(\mathbb{C}^n)$, the set of $u \in \mathcal{L}^S(\mathbb{C}^n)$ such that $H_S - c_u \leq u$ for some constant c_u .

It is a fundamental problem to characterize those S and admissible weights q for which $V_{E,q}^S = \log \Phi_{E,q}^S$. The classical Siciak-Zakharyuta theorem states that $V_K =$ log Φ_K , for every compact subset K of \mathbb{C}^n , see [19, Theorem 5.1.7]. It was first proved by Zakharyuta [38] and generalized to $V_{K,q} = \log \Phi_{K,q}$ for continuous q on a compact K by Siciak [31, Theorem 4.12]. Magnússon, Sigurðardóttir, and Sigurðsson [22] prove that for every compact and convex $S \subset \mathbb{R}^n_+$ with $0 \in S$ and every admissible weight qon a closed $E \subset \mathbb{C}^n$, the equation $V_{E,q}^S = \log \Phi_{E,q}^S$ holds on \mathbb{C}^{*n} if and only if $S \cap \mathbb{Q}^n$ is dense in S. In the case when $V_{E,q}^S$ is lower semicontinuous equality holds on \mathbb{C}^n . In Proposition 4.7 we show that if $S \cap \mathbb{Q}^n$ is not dense in S, then $V_{E,q}^S \neq \log \Phi_{E,q}^S$ for any admissible weight q on a closed set $E \subseteq \mathbb{C}^n$. Magnússon, Sigurðsson, and Snorrason [23] prove a generalization of the Bernstein-Walsh-Siciak theorem to the weighted setting with approximation by polynomials from the polynomial ring $\mathcal{P}^S(\mathbb{C}^n)$.

In Section 5 we study regularity of $V_{E,q}^S$. We introduce a regularization operator R_{δ} that preserves the class $\mathcal{L}^S(\mathbb{C}^n)$ and has the property that $R_{\delta}u \in \mathcal{C}^{\infty}(\mathbb{C}^{*n}) \cap \mathcal{L}^S(\mathbb{C}^n)$ for every $u \in \mathcal{L}^S(\mathbb{C}^n)$ and $R_{\delta}u \searrow u$ as $\delta \searrow 0$. This property enables us to prove in Proposition 5.4 that $V_{K,q}^S$ is lower semicontinuous on \mathbb{C}^{*n} for every S and every admissible weight q on a compact set K. It is crucial to know if the upper regularization $V_{E,q}^{S*}$ of $V_{E,q}^S$ satisfies $V_{E,q}^{S*} \leq q$ on K, for then $V_{E,q}^{S*} = V_{E,q}^S$ and $V_{E,q}^S$ is continuous at every point where it is lower semicontinuous.

It is a natural question to ask under which conditions on S the class $\mathcal{L}^{S}(\mathbb{C}^{n})$ is preserved under the standard method for regularization of plurisubharmonic functions, that is convolution $u \mapsto u * \psi$, where $\psi \in \mathcal{C}_{0}^{\infty}(\mathbb{C}^{n})$ with $\psi \geq 0$, and $\int_{\mathbb{C}^{n}} \psi \, d\lambda = 1$. In Theorem 5.8 we prove that $\mathcal{L}^{S}(\mathbb{C}^{n})$ is preserved under convolution if and only if S is a lower set, which says that the cube $C_{s} = [0, s_{1}] \times \cdots \times [0, s_{n}]$ is contained in S for every $s \in S$. This is a complete answer to a question raised by Bayraktar, Hussung, Levenberg, and Perera [8, Section 2].

In Section 6 we consider an equilibrium measure $\mu_{E,q}^S = \left(dd^c V_{E,q}^{S*}\right)^n$ for $V_{E,q}^{S*}$. We prove in Theorem 6.1 that its support is located where $V_{K,q}^{S*} \ge q$ and in Theorem 6.2 we prove that its total mass is $\mu_{E,q}^S(\mathbb{C}^n) = (2\pi)^n n! \operatorname{vol}(S)$ where vol denotes the euclidean volume. See also Bayraktar [4, Proposition 2.7] and Rashkovskii [26, Section 3].

In Section 7 we return to the problem of characterizing the classes $\mathcal{P}_m^S(\mathbb{C}^n)$ by growth properties we started in Theorem 3.6. Our main result, Theorem 7.2, is that every entire function f in the weighted L^2 -space $L^2(\mathbb{C}^n, \psi)$ consisting of all measurable functions which are square integrable with respect to the Lebesgue measure λ on \mathbb{C}^n with weight $e^{-\psi}$ and $\psi = 2mH_S + a\log(1+|\cdot|^2)$ is a polynomial in $\mathcal{P}_m^{\widehat{S}_{\Gamma}}(\mathbb{C}^n)$, where \widehat{S}_{Γ} is the Γ -hull of S defined by $\widehat{S}_{\Gamma} = \{x \in \mathbb{R}^n_+; \langle x, \xi \rangle \leq \varphi_S(\xi), \forall \xi \in \Gamma\}$ for a certain cone $\Gamma \subset \mathbb{R}^n$. Bayraktar, Hussung, Levenberg, and Perera [8, Proposition 4.3] claim that $f \in \mathcal{P}_m^S(\mathbb{C}^n)$ for any polytope S and a sufficiently small. Example 7.4 shows that their claim is false even if S is a polytope containing a neighborhood of 0 in \mathbb{R}^n_+ . The Siciak-Zakharyuta type theorem [8, Theorem 1.1] does not have a sound proof as it is based on their Proposition 4.3. As far as we can see the proof is only valid if S is a lower set.

In Section 8 we continue the discussion in Perera [25] and show that for any polynomial map $f: \mathbb{C}^n \to \mathbb{C}^{\ell}$ and any compact convex $0 \in S \subseteq \mathbb{C}^n$, there is a canonical minimal choice of S' such that $f^*: \mathcal{P}_m^S(\mathbb{C}^n) \to \mathcal{P}_m^{S'}(\mathbb{C}^{\ell})$ is well-defined for all $m \in \mathbb{N}$ and $f^*: \mathcal{L}^S(\mathbb{C}^n) \to \mathcal{L}^{S'}(\mathbb{C}^{\ell})$ is well-defined. In the case when $\ell = n$ we show when such pullbacks are bijective.

Snorrason [36] generalized the Siciak product formula, Siciak [30, 31] and Klimek [19, Theorem 5.1.8]. He shows in Corollary 1.3, that the generalization of Bos and Levenberg [13] of Siciak's product formula only holds for lower sets. This shows that both Levenberg and Perera [21, Proposition 1.3], and Nguyen Quang Dieu and Tang Van Long [24, Theorem 1.3] do not hold. The error in both the papers is the same, the authors implicitly assume that $\varphi_S(\xi) = \varphi_S(\xi^+)$ holds for every $\xi \in \mathbb{R}^n$, where $\xi_j^+ = \max\{0, \xi_j\}$, but in Theorem 5.8 we prove that this identity holds if and only if S is a lower set. In [36, Section 5] Snorrason shows that the sublevel sets of V_K^S are not convex in general. This contradicts Nguyen Quang Dieu and Tang Van Long [24, Theorem 1.2] where they claim that for every convex body S and every compact convex $K \subset \mathbb{C}^n$ the sublevel sets of V_K^S are convex. All these mistakes showed us the importance of a careful study of the values of H_S near points on the union of the coordinate hyperplanes as we have done in Propositions 3.3 and 3.4.

Acknowledgment

The results of this paper are a part of a research project, *Holomorphic Approximations* and *Pluripotential Theory*, project grant no. 207236-051, supported by the Icelandic Research Fund. We would like to thank the Fund for its support and the Mathematics Division, Science Institute, University of Iceland, for hosting the project. Parts of the paper were written while the second and third authors were visitors at Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, Sweden. They would like to thank the members of the institute for their hospitality and generosity. We thank the referees for their constructive and helpful comments which have improved the quality and clarity of the paper.

2 Weighted polynomial classes and Siciak functions

Let S be a bounded subset of $\mathbb{R}^n_+ = \{x \in \mathbb{R}^n; x_j \ge 0 \text{ for all } j = 1, \ldots, n\}$. For every $m \in \mathbb{N}$ we associate to S the space $\mathcal{P}^S_m(\mathbb{C}^n)$ of all polynomials in n complex variables of the form

(2.1)
$$p(z) = \sum_{\alpha \in (mS) \cap \mathbb{N}^n} a_{\alpha} z^{\alpha}, \qquad z \in \mathbb{C}^n,$$

with the standard multi-index notation and let $\mathcal{P}^{S}(\mathbb{C}^{n}) = \bigcup_{m \in \mathbb{N}} \mathcal{P}_{m}^{S}(\mathbb{C}^{n})$. If S is the standard simplex $\Sigma = ch\{0, e_{1}, \ldots, e_{n}\}$, then the space $\mathcal{P}_{m}^{\Sigma}(\mathbb{C}^{n})$ consists of all polynomials of degree $\leq m$, which we denote by $\mathcal{P}_{m}(\mathbb{C}^{n})$. We let $\mathcal{P}(\mathbb{C}^{n}) = \bigcup_{m \in \mathbb{N}} \mathcal{P}_{m}(\mathbb{C}^{n})$ denote the space of all polynomials in n complex variables.

Assume now that S is a compact convex subset of \mathbb{R}^n_+ with $0 \in S$. If $\alpha \in jS$ and $\beta \in kS$ for some $j, k \in \mathbb{N}^*$, say $\alpha = ja$ and $\beta = kb$ with $a, b \in S$, then convexity of S gives $\alpha + \beta = (j+k)((1-\lambda)a + \lambda b) \in (j+k)S$, where $\lambda = k/(j+k) \in [0,1]$. Thus, $z^{\alpha}z^{\beta} \in \mathcal{P}^S_{j+k}(\mathbb{C}^n)$ and by taking linear combinations of products of monomials we get

(2.2)
$$\mathcal{P}_{j}^{S}(\mathbb{C}^{n})\mathcal{P}_{k}^{S}(\mathbb{C}^{n}) \subseteq \mathcal{P}_{j+k}^{S}(\mathbb{C}^{n}).$$

This tells us that $\mathcal{P}^{S}(\mathbb{C}^{n})$ is a subring of $\mathcal{P}(\mathbb{C}^{n})$. For every $p \in \mathcal{P}^{S}(\mathbb{C}^{n})$ we define the *S*-degree deg^S(p) of p as the infimum over m for which $p \in \mathcal{P}_{m}^{S}(\mathbb{C}^{n})$. We have

(2.3)
$$\deg^{S}(p_1 + p_2) \le \max\{\deg^{S}(p_1), \deg^{S}(p_2)\},\$$

(2.4)
$$\deg^{S}(p_1p_2) \le \deg^{S}(p_1) + \deg^{S}(p_2)$$

Equality does not hold in general in either of these inequalities.

Definition 2.1 Let $S \subset \mathbb{R}^n_+$ be a compact convex set, $0 \in S$, and $q: E \to \mathbb{R} \cup \{+\infty\}$ be a function on $E \subset \mathbb{C}^n$. For $m \in \mathbb{N}^* = \{1, 2, 3, ...\}$ the *m*-th Siciak extremal function with respect to S, E, and q is defined by

$$\Phi_{E,q,m}^{S}(z) = \sup\{|p(z)|^{1/m} \, ; \, p \in \mathcal{P}_{m}^{S}(\mathbb{C}^{n}), \|pe^{-mq}\|_{E} \le 1\}, \qquad z \in \mathbb{C}^{n},$$

the Siciak extremal function with respect to S, E, and q is defined by

$$\Phi^{S}_{E,q}(z) = \overline{\lim}_{m \to \infty} \Phi^{S}_{E,q,m}(z), \qquad z \in \mathbb{C}^{n}.$$

We drop S in the superscripts if $S = \Sigma$ and q in the subscripts if q = 0. Note that the family $\{p \in \mathcal{P}_m^S(\mathbb{C}^n); \|pe^{-mq}\|_E \leq 1\}$ is never empty since it always contains the zero polynomial. Furthermore, we define $\Phi_{E,q,0}^S = 1$.

Observe that our definition of $\Phi_{E,q,m}^S$ deviates from the original definition of Siciak [30], which is $(\Phi_{E,q,m}^S)^m$ in our notation. We have that $\Phi_{E,q}^S, \Phi_{E,q,m}^S$ are lower semicontinuous on \mathbb{C}^n for $m = 1, 2, 3, \ldots$, for all these functions are suprema of continuous functions.

If q is bounded below, say by the real number q_0 , then the constant polynomial $p(z) = e^{mq_0}$ is in $\mathcal{P}_m^S(\mathbb{C}^n)$ and $\|pe^{-mq}\|_E = \|e^{-m(q-q_0)}\|_E \leq 1$. Hence, it follows that $\Phi_{E,q,m}^S(z) \geq e^{q_0}$ for every $z \in \mathbb{C}^n$ and $m \in \mathbb{N}^*$.

Proposition 2.2 Let $S \subset \mathbb{R}^n_+$ be a compact convex set with $0 \in S$, $E \subset \mathbb{C}^n$, and $q: E \to \mathbb{R} \cup \{+\infty\}$ be a function. Then for j, k = 1, 2, 3, ...

(2.5)
$$\left(\Phi_{E,q,j}^S(z)\right)^j \left(\Phi_{E,q,k}^S(z)\right)^k \le \left(\Phi_{E,q,j+k}^S(z)\right)^{j+k}, \qquad z \in \mathbb{C}^n,$$

and

(2.6)
$$\Phi_{E,q}^S(z) = \lim_{m \to \infty} \Phi_{E,q,m}^S(z) = \sup_{m \ge 1} \Phi_{E,q,m}^S(z), \qquad z \in \mathbb{C}^n$$

If q is bounded below and $\Phi_{E,q}^S$ is continuous on some compact subset X of \mathbb{C}^n , then the convergence is uniform on X.

We need ingredients from the proof from Tsuji [37, Lemma after Theorem III.25 on page 73]:

Lemma 2.3 (Fekete lemma) Let $(a_m)_{m \in \mathbb{N}^*}$ be a subadditive real sequence, that is $a_{j+k} \leq a_j + a_k$ for $j, k = 1, 2, 3, \ldots$ Then $\lim_{m \to \infty} a_m/m = \inf_{m \geq 1} a_m/m$.

Proof: Denote the infimum of a_m/m by α and take $\beta \in \mathbb{R}$ such that $-\infty \leq \alpha < \beta$. Then there exists $j \in \mathbb{N}^*$ such that $a_j/j < \beta$. Every m > j can be written as m = sj + r for some $s, r \in \mathbb{N}$ with $s \geq 1$ and $0 \leq r < j$. By assumption, we have $a_m \leq a_{sj} + a_r \leq sa_j + a_r$, so

(2.7)
$$\frac{a_m}{m} \le \frac{sa_j + a_r}{m} < \beta \left(1 - r/m \right) + \frac{\max_{\ell < j} a_\ell}{m},$$

which implies $\overline{\lim}_{m\to\infty} a_m/m \leq \beta$. Since β is arbitrary we have

$$\overline{\lim_{n \to \infty}} a_m / m \le \inf_{m \ge 1} a_m / m \le \lim_{m \to \infty} a_m / m.$$

Hence, the limit (possibly $-\infty$) exists and the equality holds.

Proof of Proposition 2.2: Take $p_j \in \mathcal{P}_j^S(\mathbb{C}^n)$ with $\|p_j e^{-jq}\|_E \leq 1$ and $p_k \in \mathcal{P}_k^S(\mathbb{C}^n)$ with $\|p_k e^{-kq}\|_E \leq 1$. Then $\|p_j p_k e^{-(j+k)q}\|_E \leq 1$ and (2.2) implies $p_j p_k \in \mathcal{P}_{j+k}^S(\mathbb{C}^n)$. Hence, $|p_j(z)p_k(z)| \leq (\Phi_{E,q,j+k}^S(z))^{j+k}$. By taking supremum over p_j and p_k (2.5) follows. By (2.5) the sequence defined by $a_m = -\log(\Phi_{E,q,m}^S(z))^m$ is subadditive for every z, so (2.6) follows from Lemma 2.3.

Assume now that $q \ge q_0$ for some $q_0 \in \mathbb{R}$ and that $\Phi_{E,q}^S$ is continuous on X. By the discussion after Definition 2.1 we have $\Phi_{E,q,m}^S \ge e^{q_0}$. Since $\Phi_{E,q}^S = \sup_{m\ge 1} \Phi_{E,q,m}^S$, it follows by a simple compactness argument that it is sufficient to show that for every $z_0 \in X$ and every $\varepsilon > 0$ there exists $\delta > 0$ and $k \in \mathbb{N}^*$ both depending on z_0 and ε such that

(2.8)
$$\Phi_{E,q}^S(z) - \Phi_{E,q,m}^S(z) < \varepsilon, \qquad z \in B(z_0, \delta) \cap X, \ m \ge k.$$

Let $c = \sup_X \Phi_{E,q}^S$ and choose $\gamma > 0$ such that $c(1 - e^{-\gamma}) < \frac{1}{4}\varepsilon$ and $j \in \mathbb{N}^*$ so large that $\Phi_{E,q}^S(z_0) - \Phi_{E,q,j}^S(z_0) < \frac{1}{4}\varepsilon$. Since $\Phi_{E,q}^S$ is continuous on X and $\Phi_{E,q,j}^S$ is lower semicontinuous on \mathbb{C}^n , there exists $\delta > 0$ such that for all $z \in B(z_0, \delta) \cap X$ we have

$$\Phi_{E,q}^S(z) - \Phi_{E,q}^S(z_0) < \frac{1}{4}\varepsilon \quad \text{and} \quad \Phi_{E,q,j}^S(z_0) - \Phi_{E,q,j}^S(z) < \frac{1}{4}\varepsilon.$$

The three estimates imply

(2.9)
$$\Phi_{E,q}^S(z) - \Phi_{E,q,j}^S(z) < \frac{3}{4}\varepsilon, \qquad z \in B(z_0,\delta) \cap X,$$

and (2.8) follows from (2.9) if we can prove that there exists k > j such that

(2.10)
$$\Phi_{E,q}^{S}(z) - \Phi_{E,q,m}^{S}(z) \le \Phi_{E,q}^{S}(z) - \Phi_{E,q,j}^{S}(z) + \frac{1}{4}\varepsilon, \qquad z \in X, \ m \ge k.$$

For every $z \in \mathbb{C}^n$ the sequence $a_m = -\log \left(\Phi_{E,q,m}^S(z)\right)^m$ is subadditive. By (2.7) we have for every m > j written as m = sj + r with $s \in \mathbb{N}^*$ and $r \in \mathbb{N}$ with $0 \le r < j$ that

$$-\log \Phi_{E,q,m}^{S}(z) \le \frac{-sj\log \Phi_{E,q,j}^{S}(z) - r\log \Phi_{E,q,r}^{S}(z)}{m}, \qquad z \in \mathbb{C}^{n}.$$

We have sj/m = 1 - r/m, $-\log \Phi^S_{E,q,j}(z) \ge -\log c$ for every $z \in X$, and that $\log \Phi^S_{E,q,r}(z) \ge q_0$ for every $z \in \mathbb{C}^n$, so

$$\log \Phi_{E,q,m}^{S}(z) \ge \log \Phi_{E,q,j}^{S}(z) + \frac{-r \log \Phi_{E,q,j}^{S}(z) + r \log \Phi_{E,q,r}^{S}(z)}{m}$$
$$\ge \log \Phi_{E,q,j}^{S}(z) - \frac{j \log(c/e^{q_0})}{m}, \qquad z \in X.$$

We choose k > j so large that $j \log(c/e^{q_0})/k < \gamma$. Then $\Phi^S_{E,q,m}(z) \ge \Phi^S_{E,q,j}(z)e^{-\gamma}$ and

$$\Phi_{E,q}^{S}(z) - \Phi_{E,q,m}^{S}(z) \le \Phi_{E,q}^{S}(z) - \Phi_{E,q,j}^{S}(z) + \Phi_{E,q,j}^{S}(z) (1 - e^{-\gamma})$$

for every $z \in X$ and $m \geq k$. Since $\Phi_{E,q,j}^S(z) \leq c$ for every $z \in X$ and $c(1 - e^{-\gamma}) < \frac{1}{4}\varepsilon$ the estimate (2.10) holds.

3 The Lelong class with respect to a convex set

Let us begin by setting some notation for the sequel. We denote by $\mathcal{H}(X)$, $\mathcal{SH}(X)$, $\mathcal{PSH}(X)$, $\mathcal{O}(X)$, $\mathcal{LSC}(X)$, and $\mathcal{USC}(X)$ the classes of harmonic and subharmonic functions on a domain X in \mathbb{C} , plurisubharmonic and holomorphic functions on a complex manifold X, and lower and upper semicontinuous functions on a topological space X, respectively. We define the coordinate-wise logarithm of the modulus, exponential function, and positive part, by

Log:
$$\mathbb{C}^{*n} \to \mathbb{R}^n$$
, Log $(z) = (\log |z_1|, \dots, \log |z_n|)$, $z \in \mathbb{C}^{*n}$,
Exp: $\mathbb{R}^n \to \mathbb{R}^n_+$, Exp $(\xi) = e^{\xi} = (e^{\xi_1}, \dots, e^{\xi_n})$, $\xi \in \mathbb{R}^n$,
 $^+$: $\mathbb{R}^n \to \mathbb{R}^n_+$, $\xi^+ = (\xi_1^+, \dots, \xi_n^+)$, $\xi_j^+ = \max\{\xi_j, 0\}$ $\xi \in \mathbb{R}^n$.

We let \mathbb{D} denote the unit disc and \mathbb{T} the unit circle in \mathbb{C} . The Lelong class $\mathcal{L}(\mathbb{C}^n)$ is the set of all $u \in \mathcal{PSH}(\mathbb{C}^n)$ such that for some constant c_u depending on u we have

$$u(z) \le c_u + \log^+ ||z||_{\infty}, \qquad z \in \mathbb{C}^n.$$

It is clear that $\log^+ \|\cdot\|_{\infty}$ can be replaced by $\log^+ \|\cdot\|$ or $\log(1+\|\cdot\|)$ for any norm $\|\cdot\|$ on \mathbb{C}^n .

Definition 3.1 For every compact subset of \mathbb{R}^n_+ with $0 \in S$ we define the supporting function of S as

$$\varphi(x) = \sup_{s \in S} \langle s, x \rangle, \qquad x \in \mathbb{R}^n,$$

and the logarithmic supporting function of S as the function $H_S \colon \mathbb{C}^n \to \mathbb{R}_+$ defined on \mathbb{C}^{*n} by

$$H_S(z) = (\varphi_S \circ \operatorname{Log})(z) = \max_{s \in S} \left(s_1 \log |z_1| + \dots + s_n \log |z_n| \right), \qquad z \in \mathbb{C}^{*n},$$

and extended to $\mathbb{C}^n \setminus \mathbb{C}^{*n}$ by

$$H_S(z) = \overline{\lim_{\mathbb{C}^{*n} \ni w \to z}} H_S(w), \qquad z \in \mathbb{C}^n \setminus \mathbb{C}^{*n}.$$

The real number $\sigma_S = \varphi_S(\mathbf{1})$, where $\mathbf{1} = (1, \ldots, 1) \in \mathbb{R}^n_+$, is called the *logarithmic type* of H_S .

Since φ_S is positively homogeneous of degree 1 and convex, that is $\varphi_S(t\xi) = t\varphi_S(\xi)$ and $\varphi_S(\xi + \eta) \leq \varphi_S(\xi) + \varphi_S(\eta)$ for every $t \in \mathbb{R}_+$ and $\xi, \eta \in \mathbb{R}^n$, we have

(3.1)
$$H_S(z) = \frac{1}{\lambda} H_S(|z_1|^{\lambda}, \dots, |z_n|^{\lambda}), \qquad \lambda \in \mathbb{R}^*_+, \ z \in \mathbb{C}^{*n},$$

(3.2)
$$H_S(z_1w_1,\ldots,z_nw_n) \le H_S(z) + H_S(w), \qquad z, w \in \mathbb{C}^{*n}$$

Observe that $\varphi_S(-1) = 0$ and that for $z \in \mathbb{C}^{*n}$, $\lambda \in \mathbb{C}^*$

(3.3)
$$H_S(\lambda z) \le H_S(z) + H_S(|\lambda|\mathbf{1}) = H_S(z) + \sigma_S \log^+ |\lambda|.$$

If we write $z = ||z||_{\infty} w$ with $||w||_{\infty} = 1$, then this formula implies $H_S/\sigma_S \in \mathcal{L}(\mathbb{C}^n)$,

(3.4)
$$H_S(z) \le \sigma_S \log^+ \|z\|_{\infty}, \qquad z \in \mathbb{C}^n$$

Directly from the definition we see that $H_S \in \mathcal{PSH}(\mathbb{C}^{*n}) \cap \mathcal{C}(\mathbb{C}^{*n})$. Since $\mathbb{C}^n \setminus \mathbb{C}^{*n}$ is pluripolar and H_S is locally bounded above we have $H_S \in \mathcal{PSH}(\mathbb{C}^n)$.

Proposition 3.2 Let $S \subset \mathbb{R}^n_+$ be compact convex and with $0 \in S$. Then for every $z \in \mathbb{C}^{*n}$ and $w \in \mathbb{C}^n$ we have

$$H_S(z+w) \le H_S(z) + \varphi_S(|w_1|/|z_1|, \dots, |w_n|/|z_n|),$$

and in particular, for every $w \in \overline{\mathbb{D}}^n$ and $\delta \in]0,1[$ we have

$$H_S(\mathbf{1} + \delta w) \le \delta \sigma_S.$$

Furthermore,

 $H_S(z+w) \le H_S(z) + \varphi_S(\log^+(|w_1|/|z_1|), \dots, \log^+(|w_n|/|z_n|)) + (\log 2)\sigma_S.$

Proof: By plurisubharmonicity of H_S and convexity of the functions given by $w \mapsto \varphi_S(|w_1|/|z_1|, \ldots, |w_n|/|z_n|)$ we may assume that $w, z + w \in \mathbb{C}^{*n}$. For some $t \in S$ we have

$$H_S(z+w) = \log(|z_1+w_1|^{t_1}\cdots|z_n+w_n|^{t_n}) \le \langle t, \operatorname{Log} z \rangle + \sum_{j=1}^n t_j \log\left(1+\frac{|w_j|}{|z_j|}\right).$$

Since $\log(1+x) \leq x$, for $x \geq 0$ the first estimate follows. Since $1 + \delta w \in \mathbb{C}^{*n}$ for $w \in \overline{\mathbb{D}}^n$, $\delta \in]0, 1[$, and $H_S(1) = 0$, the second estimate follows. For the third estimate we use the fact that $\log(1+x) \leq \log 2 + \log^+ x$ for $x \geq 0$.

The zero set $\mathcal{N}(H_S)$ of H_S can be understood in terms of the zero set of φ_S which is a cone. Since $\mathcal{N}(H_S) \cap \mathbb{C}^{*n} = \mathrm{Log}^{-1}(\mathcal{N}(\varphi_S))$ and $\mathbb{R}^n_- \subseteq \mathcal{N}(\varphi_S)$, the closed unit polydisc $\overline{\mathbb{D}}^n$ is contained in $\mathcal{N}(H_S)$. Furthermore, $\mathcal{N}(H_S)$ is equal to $\overline{\mathbb{D}}^n$ if and only if $\mathbb{R}_+S = \mathbb{R}^n_+$. We have a complete description of the values of H_S at every point in $\mathbb{C}^n \setminus \mathbb{C}^{*n}$, the union of the coordinate hyperplanes. **Proposition 3.3** Let S be a compact convex subset of \mathbb{R}^n_+ with $0 \in S$. For every $a \neq 0$ in some coordinate hyperplane we have

$$H_S(a) = H_{S_J}(a_{j_1}, \ldots, a_{j_\ell}),$$

where $J \subset \{1, \ldots, n\}$ consists of the indices $j_1 < \cdots < j_\ell$ of the non-zero coordinates $a_{j_1}, \ldots, a_{j_\ell}$ of a and $S_J \subseteq \mathbb{R}^\ell$ consists of all $t \in \mathbb{R}^\ell$ such that if $s \in \mathbb{R}^n_+$ is defined by $s_{j_k} = t_k$ for $j_k \in J$ and $s_j = 0$ for $j \notin J$, then $s \in S$.

Proof: After renumbering the variables we may assume that $J = \{1, \ldots, \ell\}$ and $a_{\ell+1} = \cdots = a_n = 0$. We write $z = (z', z'') \in \mathbb{C}^n$, where $z' \in \mathbb{C}^\ell$ and $z'' \in \mathbb{C}^{n-\ell}$. Then $a' \in \mathbb{C}^{*\ell}$, so H_{S_J} is continuous at a' and we have

$$H_{S_J}(a') = \lim_{\mathbb{C}^{*\ell} \ni w' \to a'} H_{S_J}(w') = \lim_{\mathbb{C}^{*m} \ni w' \to a'} \sup_{t \in S_J} \langle t, \log w' \rangle$$
$$= \lim_{\mathbb{C}^{*m} \ni w' \to a'} \sup_{s = (t,0) \in S} \langle s, (\log w', 0, \dots, 0) \rangle$$
$$\leq \lim_{\mathbb{C}^{*n} \ni w \to a} \sup_{s \in S} \langle s, \log w \rangle = H_S(a).$$

In order to prove the converse inequality we take a convergent sequence in \mathbb{C}^{*n} , $w_j = (w_{j,1}, \ldots, w_{j,n}) \to a$ such that $\lim_{j\to\infty} H_S(w_j) = H_S(a)$. There exists $s_j \in S$ such that $H_S(w_j) = \langle s_j, \log w_j \rangle$ for every j. Since $H_S \geq 0$ and $\log |w_{j,\kappa}| \to -\infty$ as $j \to \infty$ for $\kappa = \ell + 1, \ldots, n$, it follows that $s_{j,\kappa} \to 0$ as $j \to \infty$ for $\kappa = \ell + 1, \ldots, n$. By compactness of S there exists a subsequence s_{j_k} converging to $(t,0) \in S$. We have $t \in S_J$ and conclude that

$$H_S(a) = \lim_{k \to \infty} H_S(w_{j_k}) = \lim_{k \to \infty} \langle s_{j_k}, \operatorname{Log} w_{j_k} \rangle = \langle t, \operatorname{Log} a' \rangle \le H_{S_J}(a').$$

With a proof similar to Rashkovskii [27, Proposition 2.2] we are able to show that $H_S \in \mathcal{C}(\mathbb{C}^n)$:

Proposition 3.4 Let S be a compact convex subset of \mathbb{R}^n_+ with $0 \in S$. Then H_S is plurisubharmonic and continuous on \mathbb{C}^n .

Proof: Let $0 \neq a \in \mathbb{C}^n \setminus \mathbb{C}^{*n}$. Since $H_S \in \mathcal{PSH}(\mathbb{C}^n) \cap \mathcal{C}(\mathbb{C}^{*n})$, it suffices to prove that H_S is lower semicontinuous at a. After renumbering the variables we may assume $a_{\ell+1} = \cdots = a_n = 0$ and $a_j \neq 0$ for $j \leq \ell$. We also write $z = (z', z'') \in \mathbb{C}^n$, where $z' \in \mathbb{C}^\ell$ and $z'' \in \mathbb{C}^{n-\ell}$. Let $z_j \in \mathbb{C}^n$ be such that $z_j \to a$. Since H_S is rotationally invariant in each variable it takes the constant value $H_S(z_j)$ on the distinguished boundary of the $n-\ell$ dimensional polydisc $\{(z', \zeta''); |\zeta_k''| \leq |z_{j,k}''|, k = 1, \ldots, n-\ell\}$, so by the maximum principle $H_S(z'_j, 0) \leq H_S(z_j)$. By Proposition 3.3 we have that $H_S(z'_j, 0) = H_{S_J}(z'_j)$, for $J = \{1, \ldots, \ell\}$. By the continuity of H_{S_J} at a' we have

$$\lim_{j \to \infty} H_S(z_j) \ge \lim_{j \to \infty} H_S(z'_j, 0) = \lim_{j \to \infty} H_{S_J}(z'_j) = H_{S_J}(a') = H_S(a).$$

This proves the lower semicontinuity of H_S at a.

Definition 3.5 For every compact convex subset S of \mathbb{R}^n_+ with $0 \in S$ we define the *S*-Lelong class $\mathcal{L}^S(\mathbb{C}^n)$ as the set of all $u \in \mathcal{PSH}(\mathbb{C}^n)$ such that

$$u(z) \le c_u + H_S(z), \qquad z \in \mathbb{C}^n,$$

for some constant c_u depending on u, and define $\mathcal{L}^S_+(\mathbb{C}^n)$ as the subclass of functions u, that have the same asymptotic behavior at infinity as the function H_S , that is

$$(3.5) -c_u + H_S(z) \le u(z) \le c_u + H_S(z), z \in \mathbb{C}^n.$$

The Liouville theorem tells us that an entire function $f \in \mathcal{O}(\mathbb{C}^n)$ which satisfies a growth estimate $|f(z)| \leq C(1+|z|)^{a+m}$, $z \in \mathbb{C}^n$, for some $m \in \mathbb{N}$ and $a \in [0,1[$, is a polynomial of degree $\leq m$, that is $f \in \mathcal{P}_m(\mathbb{C}^n)$. The following is a Liouville type theorem for the polynomial classes $\mathcal{P}_m^S(\mathbb{C}^n)$:

Theorem 3.6 Let d_m denote the distance between mS and $\mathbb{N}^n \setminus mS$ in the L^1 -norm. Then for every $f \in \mathcal{O}(\mathbb{C}^n)$ the following are equivalent:

- (i) $f \in \mathcal{P}_m^S(\mathbb{C}^n)$.
- (ii) $\log |f|^{1/m} \in \mathcal{L}^S(\mathbb{C}^n).$
- (iii) there exists $a \in [0, d_m[$ such that $|f|e^{-mH_S a\log^+ \|\cdot\|_{\infty}} \in L^{\infty}(\mathbb{C}^n).$
- (iv) there exists $a \in [0, d_m]$ and a constant C > 0 such that

$$|f(z)| \le C(1+|z|)^a e^{mH_S(z)}, \qquad z \in \mathbb{C}^n.$$

Proof: (i) \Rightarrow (ii): If $\alpha \in mS$, then $|z^{\alpha}| \leq e^{mH_S(z)}$, so for $f \in \mathcal{P}_m^S(\mathbb{C}^n)$, $f(z) = \sum_{\alpha \in (mS) \cap \mathbb{N}^n} a_{\alpha} z^{\alpha}$, we have $\log |f(z)|^{1/m} \leq c_f/m + H_S(z)$ with $c_f = \log \sum_{\alpha \in (mS) \cap \mathbb{N}^n} |a_{\alpha}|$. (ii) \Rightarrow (iii): We have $|f|e^{-mH_S-a\log^+} \|\cdot\|_{\infty} \leq |f|e^{-mH_S} \in L^{\infty}(\mathbb{C}^n)$.

(iii) \Rightarrow (i): Observe first that $m\varphi_S(\xi) + a \|\xi^+\|_{\infty} = \varphi_{mS+a\Sigma}(\xi)$ for every $\xi \in \mathbb{R}^n$. Let $f(z) = \sum_{\alpha \in \mathbb{N}^n} a_{\alpha} z^{\alpha}$ be the power series expansion of f at 0. We need to show that $a_{\alpha} = 0$ for all $\alpha \in \mathbb{N}^n \setminus mS$. Since $a < d_m$ we have $(mS + a\Sigma) \cap \mathbb{N}^n = (mS) \cap \mathbb{N}^n$, so $\alpha \in \mathbb{N}^n \setminus (mS + a\Sigma)$. Hence, there exists $\xi \in \mathbb{R}^n$ such that $\langle \alpha, \xi \rangle > \varphi_{mS+a\Sigma}(\xi)$. We let C_t denote the polycircle with center 0 and polyradius $(e^{t\xi_1}, \ldots, e^{t\xi_n})$ and observe that by the Cauchy formula for derivatives we have

$$a_{\alpha} = \frac{1}{(2\pi i)^n} \int_{C_t} \frac{f(\zeta)}{\zeta^{\alpha}} \frac{d\zeta_1 \cdots d\zeta_n}{\zeta_1 \cdots \zeta_n}.$$

For $\zeta = (e^{t\xi_1 + i\theta_1}, \dots, e^{t\xi_n + i\theta_n})$ on C_t we have

$$|f(\zeta)|/|\zeta^{\alpha}| \le C e^{-t(\langle \alpha,\xi\rangle - \varphi_{mS+a\Sigma}(\xi))},$$

so the right-hand side tends to 0 as $t \to +\infty$, and we conclude that $a_{\alpha} = 0$. (iii) \Leftrightarrow (iv): Follows from the equivalence of the euclidean norm $|\cdot|$ and $||\cdot||_{\infty}$. \Box

Recall from Klimek [19, p. 87] that a real valued $u \in \mathcal{PSH}(X)$ on an open subset X of \mathbb{C}^n is said to be *maximal*, if for every relatively compact open subset G of X and every $v \in \mathcal{USC}(\overline{G}) \cap \mathcal{PSH}(G)$ satisfying $v \leq u$ on ∂G we have $v \leq u$ on G. For the reader's convenience we prove the following well known result.

Lemma 3.7 Let X be an open subset of \mathbb{C}^n and $u \in \mathcal{PSH}(X)$ be real valued. Assume that for every relatively compact open subset G of X there exists a family $(g_z)_{z\in G}$ of holomorphic maps $g_z \colon D_z \to \mathbb{C}^n$ defined on open subsets D_z of \mathbb{C} , such that $K_z = g_z^{-1}(\overline{G})$ is compact, $z = g_z(\tau_z)$ for some $\tau_z \in K_z$, and $u \circ g_z$ is harmonic on D_z . Then u is maximal on X.

Proof: By Klimek [19, Proposition 3.1.1] we may take $v \in \mathcal{PSH}(X)$ in the definition of maximality. We assume that $v \leq u$ on ∂G and need to prove that $v(z) \leq u(z)$ for every $z \in G$. The set $K_z = g_z^{-1}(\overline{G})$ is compact, the function $s_z = v \circ g_z \in \mathcal{SH}(D_z)$ is less than or equal to $h_z = u \circ g_z \in \mathcal{H}(D_z)$ on the boundary of K_z . By the maximum principle for harmonic functions $v(z) = s_z(\tau_z) \leq h_z(\tau_z) = u(z)$.

For every $z \in \mathbb{C}^{n*}$ we define a parametric curve $f_z \colon \mathbb{C} \to \mathbb{C}^n$, $f_z = (f_{z,1}, \ldots, f_{z,n})$, by

(3.6)
$$f_{z,j}(\tau) = \begin{cases} e^{-i\tau \log |z_j|} (z_j/|z_j|), & z_j \neq 0, \\ 0, & z_j = 0, \end{cases} \quad \tau \in \mathbb{C}.$$

We have $f_z(i) = z$ and $||f_z(\tau)||_{\infty} = 1$ for every $\tau \in \mathbb{R}$. If $||z||_{\infty} > 1$, then for j with $|z_j| = ||z||_{\infty}$ we have

(3.7)
$$f'_{z,j}(\tau) = -ie^{-i\tau \log |z_j|} (\log |z_j|) (z_j/|z_j|) \neq 0. \quad \tau \in \mathbb{C}.$$

Hence f_z parametrizes an open Riemann surface in \mathbb{C}^n through the point z. Furthermore, we have $H_S(f_z(\tau)) = \operatorname{Im} \tau H_S(z)$ for $\operatorname{Im} \tau \geq 0$. The function $\mathbb{C} \ni \tau \mapsto H_S(f_z(\tau))$ is subharmonic, harmonic in the upper half plane, equal to 0 on the real axis, and takes the value $H_S(z)$ at *i*. By Lemma 3.7 with f_z in the role of g_z and H_S in the role of *u* we get:

Proposition 3.8 Let S be a compact convex subset of \mathbb{R}^n_+ with $0 \in S$. Then H_S is maximal on $\mathbb{C}^n \setminus \partial \mathcal{N}(H_S)$, where $\mathcal{N}(H_S)$ is the zero set of H_S .

4 Weighted Siciak-Zakharyuta functions

Definition 4.1 Let $S \subset \mathbb{R}^n_+$ be a compact convex set, $0 \in S$, and $q: E \to \mathbb{R} \cup \{+\infty\}$ be a function on $E \subset \mathbb{C}^n$. The Siciak-Zakharyuta function with respect to S, E, and q is defined by

$$V_{E,q}^S(z) = \sup\{u(z) \, ; \, u \in \mathcal{L}^S(\mathbb{C}^n), \, u|_E \le q\}, \qquad z \in \mathbb{C}^n.$$

We drop S in superscripts if $S = \Sigma$ and q in subscripts if q = 0.

From Theorem 3.6 it follows that $\log \Phi_{E,q}^S \leq V_{E,q}^S$ for every compact convex $S \subset \mathbb{R}^n_+$ and every $q: E \to \mathbb{R} \cup \{+\infty\}$ on $E \subset \mathbb{C}^n$. We will need a variant of the Phragmén-Lindelöf principle, see [18, Lemma 2.1]. **Lemma 4.2** Let v be subharmonic in the upper half plane $\mathbb{C}_+ = \{z \in \mathbb{C} ; \operatorname{Im} z > 0\}$ such that for some real constants C and A we have $v(z) \leq C + A|z|$ for all $z \in \mathbb{C}_+$, and $\overline{\lim_{\mathbb{C}_+ \ni z \to x}} v(z) \leq 0$ for all $x \in \mathbb{R}$. Then $v(z) \leq A \operatorname{Im} z$ for all $z \in \mathbb{C}_+$.

By Klimek [19, Example 5.1.1] $V_K(z) = \log^+(||z - a||/r)$ if $|| \cdot ||$ is any norm and $K = \{z \in \mathbb{C}^n; ||z - a|| \le r\}, r > 0$, is the closed ball in this norm with center a and radius r. The polynomial classes $\mathcal{P}_m^S(\mathbb{C}^n)$ are in general not translation invariant, so we can not expect to have a generalization of this example. The following is proved in special cases by Bos and Levenberg [13]:

Proposition 4.3 Let S be a compact convex subset of \mathbb{R}^n_+ with $0 \in S$ and let E be a subset of \mathbb{C}^n such that $\mathbb{T}^n \subseteq E \subseteq \mathcal{N}(H_S)$. Then $V_E^S = H_S$.

Proof: Since $H_S \in \mathcal{L}^S(\mathbb{C}^n)$ and $H_S|_E = 0$ we have $H_S \leq V_E^S$, so it is sufficient to prove that if $u \in \mathcal{L}^S(\mathbb{C}^n)$ with $u|_E \leq 0$ we have $u(z) \leq H_S(z)$ for every $z \in \mathbb{C}^{*n}$ such that $H_S(z) > 0$. Define f_z by (3.6) and $v \in \mathcal{SH}(\mathbb{C})$ by $v(\tau) = u(f_z(\tau)), \tau \in \mathbb{C}$. Since $u \in \mathcal{L}^S(\mathbb{C}^n)$ we have $v(\tau) = u(f_z(\tau)) \leq c_u + H_S(f_z(\tau)) = c_u + \operatorname{Im} \tau H_S(z), \operatorname{Im} \tau \geq 0$, and since $f_z(\mathbb{R}) \subseteq \overline{\mathbb{D}}^n$ we have $v \leq 0$ on \mathbb{R} . Lemma 4.2 gives $u(z) = v(i) \leq H_S(z)$. \Box

Since $\mathbb{T}^n \subset \mathcal{N}(H_S)$ for every S the maximum principle implies that $\overline{\mathbb{D}}^n \subset \mathcal{N}(H_S)$, where \mathbb{D} denotes the unit disc in \mathbb{C} .

Definition 4.4 Let $0 \in S \subset \mathbb{R}^n_+$ be compact and convex and $q: E \to \mathbb{R} \cup \{+\infty\}$ be a function on $E \subset \mathbb{C}^n$. We say that q is an *admissible weight with respect to S on E* if

- (i) q is lower semicontinuous,
- (ii) the set $\{z \in E; q(z) < +\infty\}$ is non-pluripolar, and
- (iii) if E is unbounded, then $\lim_{E \ni z, |z| \to \infty} (H_S(z) q(z)) = -\infty.$

This definition is taken from Bloom [28, Appendix B: Definition 2.1]. Some authors use the term *admissible external field* for q rather than *weight* in this situation and then they refer to e^{-q} as a weight. Observe that if q is an admissible weight, then E is nonpluripolar and that if E is unbounded then q = 0 is not an admissible weight.

Proposition 4.5 Let S be a compact convex subset of \mathbb{R}^n_+ with $0 \in S$ and q be an admissible weight with respect to S on a compact subset K of \mathbb{C}^n . Then $V^{S*}_{K,q} \in \mathcal{L}^S_+(\mathbb{C}^n)$.

Proof: The upper regularization $V_{E,q}^{S*}$ of $V_{E,q}^S$ is plurisubharmonic in \mathbb{C}^n if q is an admissible weight with respect to S on E. Since $V_{K,q}^S \leq q$ on K, q is admissible on K and $\{z \in K; q(z) < +\infty\} \subseteq \{z \in \mathbb{C}^n; V_{K,q}^S(z) < +\infty\}$, the set on the right is non-pluripolar. By Klimek [19, Proposition 5.2.1] it follows that the family $\mathcal{U} = \{u \in \mathcal{L}^S(\mathbb{C}^n); u|_K \leq q\} \subset \sigma_S \mathcal{L}(\mathbb{C}^n)$ is locally uniformly bounded above, where $\sigma_S = \varphi_S(\mathbf{1})$ is the logarithmic type of H_S . Let c > 0 be such that $u|_{\overline{\mathbb{D}}^n} \leq c$ for all $u \in \mathcal{U}$. Then by Proposition 4.3, we have $V_{K,q}^S \leq V_{\overline{\mathbb{D}}^n}^S + c = H_S + c$. Hence, $V_{K,q}^{S*} \in \mathcal{L}^S(\mathbb{C}^n)$. If $c = \max_{w \in K} H_S(w) - \min_{w \in K} q(w)$ then $H_S - c \leq V_{K,q}^S$, so $V_{K,q}^{S*} \in \mathcal{L}_+^S(\mathbb{C}^n)$.

Admissible weights on unbounded closed sets yield the same Siciak and Siciak-Zakharyuta functions as some compact subsets. Admissible weights do not obstruct the growth of Siciak-Zakharyuta functions. See Bloom [28, Appendix B].

Proposition 4.6 Let S be a compact convex subset of \mathbb{R}^n_+ with $0 \in S$ and q an admissible weight on a closed subset E of \mathbb{C}^n , and set $E_R = E \cap \overline{B}(0, R)$ for every R > 0. Then $V_{E,q}^S = V_{E_R,q}^S$ and $\Phi_{E,q}^S = \Phi_{E_R,q}^S$ for R sufficiently large. Furthermore, $V_{E,q}^{S*} \in \mathcal{L}^S_+(\mathbb{C}^n)$.

Proof: Since $E_R \subseteq E$ for every R > 0 we have $\Phi_{E,q}^S \leq \Phi_{E_R,q}^S$ and $V_{E,q}^S \leq V_{E_R,q}^S$, so we need to prove the reverse inequalities. By Definition 4.4, E is non-pluripolar and by [19, Corollary 4.7.7] a countable union of pluripolar sets is pluripolar. It follows that E_{R_0} is non-pluripolar for some $R_0 > 0$ and consequently E_R is pluripolar for every $R \geq R_0$. By Proposition 4.5, $V_{E_R,q}^{S*} \in \mathcal{L}^S(\mathbb{C}^n)$ and it follows that for some constant c > 0

(4.1)
$$V_{E_{R},q}^{S*}(z) \le c + H_S(z), \qquad z \in \mathbb{C}^n.$$

Condition (iii) in Definition 4.4 implies that we can choose $R_1 \ge R_0$ such that

(4.2)
$$q(z) - H_S(z) \ge c, \qquad z \in E \setminus E_{R_1}.$$

Now we take $u \in \mathcal{L}^{S}(\mathbb{C}^{n})$ and assume that $u \leq q$ on $E_{R_{1}}$. By (4.1) we have $u \leq c+H_{S}(z)$ for all $z \in \mathbb{C}^{n}$ and by (4.2) we have $u \leq q$ on $E \setminus E_{R_{1}}$. Hence, $u \leq V_{E,q}^{S}$.

Let $p \in \mathcal{P}_m^S(\mathbb{C}^n)$ for some $m \in \mathbb{N}$ be such that $\|pe^{-mq}\|_{E_{R_1}} \leq 1$. By Theorem 3.6 we have $u = \log |p|^{1/m} \in \mathcal{L}^S(\mathbb{C}^n)$ and $u \leq q$ on E_{R_1} . Again by (4.1) and (4.2) we have $u \leq q$ on $E \setminus E_{R_1}$ as well. Then $\|pe^{-mq}\|_E \leq 1$ and $|p|^{1/m} \leq \Phi_{E,q}^S$. The last statement follows directly from Proposition 4.5.

For a general compact convex S with $0 \in S$, let $S' = \overline{S \cap \mathbb{Q}^n}$ be the closure of the set of rational points in S. Then $\mathcal{P}_m^S(\mathbb{C}^n) = \mathcal{P}_m^{S'}(\mathbb{C}^n)$ for every $m \in \mathbb{N}^*$ and consequently $\log \Phi_{E,q}^S = \log \Phi_{E,q}^{S'} \leq V_{E,q}^{S'} \leq V_{E,q}^S$. Observe that S = S' if S is a convex body but $S \neq S'$ for example if S is a line segment in \mathbb{R}^2 with irrational slope.

Proposition 4.7 Let $S \subset \mathbb{R}^n_+$ be compact and convex with $0 \in S$. If $S \cap \mathbb{Q}^n$ is not dense in S, then for every admissible weight q on a closed $E \subset \mathbb{C}^n$ we have $\log \Phi^S_{E,q} \neq V^S_{E,q}$.

Proof: Since $S' = \overline{S \cap \mathbb{Q}^n} \subsetneq S$, there exists a $\xi \in \mathbb{R}^n$ with $\varphi_{S'}(\xi) < \varphi_S(\xi)$. By Proposition 4.5 there exist constants c and c' such that

$$H_S(z) - c \le V_{E,q}^S$$
 and $V_{E,q}^{S'}(z) \le H_{S'}(z) + c'.$

For r > 0 sufficiently large we have $\varphi_{S'}(r\xi) + c' < \varphi_S(r\xi) - c$, so for $z = (e^{r\xi_1}, \ldots, e^{r\xi_n})$

$$\log \Phi_{E,q}^{S}(z) = \log \Phi_{E,q}^{S'}(z) \le V_{E,q}^{S'}(z) \le H_{S'}(z) + c' < H_{S}(z) - c \le V_{E,q}^{S}(z).$$

The next result regards the Siciak-Zakharyuta functions with respect to S, E and q when we have decreasing sequences of sets $S_j \searrow S$ or increasing sequences of weights $q_j \nearrow q$.

Proposition 4.8 Let S_j , $j \in \mathbb{N}$ and S be compact convex subsets of \mathbb{R}^n_+ with $0 \in S$ and $S_j \searrow S$, q be an admissible weight on a compact subset K of \mathbb{C}^n , $(K_j)_{j\in\mathbb{N}}$ be a decreasing sequence of compact sets with $\bigcap_j K_j = K$, and $(q_j)_{j\in\mathbb{N}}$ be a sequence in $\mathcal{LSC}(K_j)$ such that $q_j \nearrow q$. Then:

- (i) $\mathcal{L}^{S}(\mathbb{C}^{n}) = \bigcap_{j \in \mathbb{N}} \mathcal{L}^{S_{j}}(\mathbb{C}^{n}).$
- (ii) If $V_{K,q}^{S_j*} \leq q$ on K for some j, then $V_{K,q}^{S_j} \searrow V_{K,q}^S$ as $j \to \infty$.
- (iii) Every q_j is an admissible weight on K_j , $V_{K_j,q_j}^S \nearrow V_{K,q}^S$ and $\Phi_{K_j,q_j}^S \nearrow \Phi_{K,q}^S$ as $j \to \infty$.

Proof: (i) Obviously $\mathcal{L}^{S}(\mathbb{C}^{n}) \subseteq \bigcap_{j} \mathcal{L}^{S_{j}}(\mathbb{C}^{n})$. Let $u \in \bigcap_{j} \mathcal{L}^{S_{j}}(\mathbb{C}^{n})$ and set $c_{u} = \sup_{\overline{\mathbb{D}}^{n}} u$. Then by Proposition 4.3 we have $u - c_{u} \leq V_{\overline{\mathbb{D}}^{n}}^{S_{j}} = H_{S_{j}}$ for every j. We have $H_{S_{j}} \searrow H_{S}$, so $u \leq c_{u} + H_{S}$ and $u \in \mathcal{L}^{S}(\mathbb{C}^{n})$.

(ii) We have $V_{K,q}^S \leq V_{K,q}^{S_j}$ and since $V_{K,q}^{S_j*} \leq q$ the equation $V_{K,q}^{S_j*} = V_{K,q}^{S_j}$ holds. By Proposition 4.5 we have $V_{K,q}^{S_j} \in \mathcal{L}^{S_j}(\mathbb{C}^n)$. Since the sequence is decreasing we have $V = \lim_{j \to \infty} V_{K,q}^{S_j*} \in \bigcap_j \mathcal{L}^{S_j}(\mathbb{C}^n) = \mathcal{L}^S(\mathbb{C}^n)$ and $V \leq q$ on K. Hence,

$$V_{K,q}^S \le \lim_{j \to \infty} V_{K,q}^{S_j} \le V \le V_{K,q}^S.$$

(iii) Since $F = \{z \in K; q(z) < +\infty\} \subseteq F_j = \{z \in K_j; q_j(z) < +\infty\}$ for every j and F is non-pluripolar, the set F_j in non-pluripolar and every q_j is an admissible weight on K_j . Since the sequence $(V_{K_j,q_j}^S)_{j\in\mathbb{N}}$ is increasing and bounded above by $V_{K,q}^S$ we need to show that $V_{K,q}^S \leq V = \lim_{j\to\infty} V_{K_j,q_j}^S$. For that purpose we take $u \in \mathcal{L}^S(\mathbb{C}^n)$ with $u \leq q$ on K and $\varepsilon > 0$. For every $z_0 \in K$ there exists j_{ε} such that $u(z_0) - \varepsilon < q_{j_{\varepsilon}}(z_0)$. Since $u \in \mathcal{USC}(\mathbb{C}^n)$ and $q_{j_{\varepsilon}} \in \mathcal{LSC}(K_j)$ it follows that we have $u(z) - \varepsilon < q_{j_{\varepsilon}}(z)$ for all $z \in K_j \cap U_0$ for some neighborhood U_0 of z_0 in \mathbb{C}^n . A simple compactness argument gives that there exists an open neighborhood U of K such that $u - \varepsilon < q_j$ on $K_j \subset U$ for $j \geq j_{\varepsilon}$, possibly with j_{ε} replaced by a larger number. Hence, $u - \varepsilon \leq V$, and since ε is arbitrary we conclude that $u \leq V$. By taking supremum over u we get $V_{K,q}^S \leq V$. The same argument for $\log |p|^{1/m}$, $p \in \mathcal{P}_m^S(\mathbb{C}^n)$, in the role of u implies that $\Phi_{K,q}^S \leq \lim_{j\to\infty} \Phi_{K_j,q_j}^S$ by Proposition 2.2.

5 Regularity of Siciak-Zakharyuta functions

In this section we study regularity of the functions $\Phi_{E,q}^S$ and $V_{E,q}^S$, where we assume that S is a convex subset of \mathbb{R}^n_+ with $0 \in S$ and that $q: E \to \mathbb{R} \cup \{+\infty\}$ is a function on a subset E of \mathbb{C}^n . Since the Siciak function $\Phi_{E,q}^S$ is the supremum of a subclass of $\mathcal{C}(\mathbb{C}^n)$ we have $\Phi_{E,q}^S \in \mathcal{LSC}(\mathbb{C}^n)$. If q is an admissible weight on a closed set E, then by Proposition 4.6 we have $V_{E,q}^{S*} \in \mathcal{L}^S_+(\mathbb{C}^n)$.

by Proposition 4.6 we have $V_{E,q}^{S*} \in \mathcal{L}_{+}^{S}(\mathbb{C}^{n})$. If $V_{E,q}^{S*} \leq q$ on E then $V_{E,q}^{S} = V_{E,q}^{S*}$ and we conclude that $V_{E,q}^{S}$ is continuous at every point where it is lower semicontinuous. In this section we prove that $V_{E,q}^{S} \in \mathcal{LSC}(\mathbb{C}^{*n})$ for every admissible weight on a compact set K. For that purpose we need to discuss regularization of plurisubharmonic functions, but we begin with local \mathcal{L} -regularity of sets.

Definition 5.1 A subset E of \mathbb{C}^n is said to be \mathcal{L} -regular at a point $a \in \overline{E}$ if V_E is continuous at a and E is said to be *locally* \mathcal{L} -regular at a if $E \cap U$ is \mathcal{L} -regular at a for every open neighborhood U of a. We say that E is (*locally*) \mathcal{L} -regular if E is (locally) \mathcal{L} -regular at every $a \in \overline{E}$.

The function V_E is continuous at every interior point of E. Moreover, V_E is continuous at $a \in \overline{E}$ if and only if $V_E^*(a) = 0$. Thus, it is sufficient to check the condition for local \mathcal{L} -regularity at boundary points only.

Lemma 5.2 Let *E* be a closed subset of \mathbb{C}^n , $a \in E$ and assume that there exists a norm with closed unit ball \mathbb{B} such that for some $\delta > 0$ and $b \in E$ we have $a \in B = b + \delta \mathbb{B} \subset E$. Then *E* is locally \mathcal{L} -regular at *a*.

Proof: Let $\|\cdot\|$ denote the norm and U be open with $a \in U$. Choose $\tau > 0$ so small that $C = c + \tau \delta \mathbb{B} \subset U$, where $c = (1 - \tau)a + \tau b$. We have $C \subset B \subset E$, so

$$0 \le V_E^*(a) \le V_C^*(a) = \log^+(\|a - c\|/\tau\delta) = \log^+(\|a - b\|/\delta) = 0.$$

Observe that the lemma implies that every set of the form $E = A + \delta \mathbb{B}$ is locally \mathcal{L} -regular, where $A \subset \mathbb{C}^n$ is closed and \mathbb{B} is the closed unit ball with respect to some norm. The following result is a generalization of Siciak [31, Proposition 2.16].

Proposition 5.3 For every continuous function q on a locally \mathcal{L} -regular closed subset E of \mathbb{C}^n we have $V_{E,q}^{S*} \leq q$ and consequently $V_{E,q}^S = V_{E,q}^{S*}$.

Proof: Let $a \in E$ and take $\varepsilon > 0$. Since q is continuous on E there exists an open neighborhood U of a such that $q(z) \leq q(a) + \varepsilon$ for every $z \in E \cap U$. Since E is locally \mathcal{L} -regular we have $V_{E \cap U}^*(a) = 0$ and

$$V_{E,q}^{S*}(a) \le V_{E\cap U,q(a)+\varepsilon}^{S*}(a) \le \sigma_S V_{E\cap U}^*(a) + q(a) + \varepsilon = q(a) + \varepsilon.$$

Since a and ε are arbitrary the inequality holds.

Convolution is a standard tool for approximating functions $u \in \mathcal{PSH}(\mathbb{C}^n)$. We define a convolution operator $L^1_{\text{loc}}(\mathbb{C}^n) \to L^1_{\text{loc}}(\mathbb{C}^n)$, $u \mapsto u * \mu$, for a given positive Borel measure μ with compact support,

(5.1)
$$u * \mu(z) = \int_{\mathbb{C}^n} u(z-w) \, d\mu(w), \qquad z \in \mathbb{C}^n.$$

If $u \in \mathcal{PSH}(\mathbb{C}^n)$ then $u * \mu \in \mathcal{PSH}(\mathbb{C}^n)$ and if μ is a probability measure then $u \mapsto u * \mu$ preserves $\mathcal{L}(\mathbb{C}^n)$. In the case that μ is presented by a \mathcal{C}^{∞} function, $\mu = \psi \, d\lambda$, then $u * \mu = u * \psi$ is a \mathcal{C}^{∞} function. If we define $\psi_{\delta} \in \mathcal{C}_0^{\infty}(\mathbb{C}^n)$ by $\psi_{\delta}(z) = \delta^{-2n}\psi(z/\delta)$, $\delta > 0$, with $\psi \in \mathcal{C}_0^{\infty}(\mathbb{C}^n)$ radially symmetric, $\psi \ge 0$, and $\int_{\mathbb{C}^n} \psi \, d\lambda = 1$, then $u * \psi_{\delta} \searrow u$ as $\delta \searrow 0$. See Klimek [19] or Hörmander [15, 16]. Siciak [31, Proposition 2.12] used convolution to prove that $V_{E,q} \in \mathcal{LSC}(\mathbb{C}^n)$ for compact E and $q \in \mathcal{LSC}(E)$. In general, the class $\mathcal{L}^S(\mathbb{C}^n)$ is not preserved under convolution (cf. Theorem 5.8).

In order to preserve a particular subclass of $\mathcal{PSH}(\mathbb{C}^n)$ under regularization, special methods are sometimes needed. For example homogeneity is preserved under

(5.2)
$$R_{\delta}u(z) = \int_{G} u(Az)\psi_{\delta}(A) \, d\mu(A), \qquad z \in \mathbb{C}^{n}$$

where G is some group of $n \times n$ matrices with real or complex entries and μ is a positive measure on the matrix space $\mathbb{R}^{n \times n}$ or $\mathbb{C}^{n \times n}$. The smoothing kernel ψ_{δ} is chosen so that it converges to the Dirac measure δ_I at the identity I as $\delta \to 0$. This method only gives a \mathcal{C}^{∞} function on $\mathbb{C}^n \setminus \{0\}$ when we integrate over the group of *complex* invertible matrices, and it gives a \mathcal{C}^{∞} function on $\mathbb{C}^n \setminus \mathbb{C}\mathbb{R}^n$, where $\mathbb{C}\mathbb{R}^n = \{\lambda x; \lambda \in \mathbb{C}, x \in \mathbb{R}^n\}$, when we integrate over the group of *real* invertible matrices. See Sigurðsson [35] and Hörmander and Sigurðsson [18].

In order to preserve the classes $\mathcal{L}^{S}(\mathbb{C}^{n})$ for a compact convex $S \subset \mathbb{R}^{n}_{+}$ with $0 \in S$ it is natural to choose G as the group of invertible diagonal matrices. This group can be identified with \mathbb{C}^{*n} with coordinate wise multiplication, so it is natural to choose μ as the Lebesgue measure λ on \mathbb{C}^{n} .

In the following text we allow us a slight abuse of notation by identifying a vector denoted by a lower case letter with a diagonal matrix denoted by the corresponding upper case letter. Thus, we identify the vector $a \in \mathbb{C}^n$ with the diagonal matrix A with diagonal a and in particular the vector $\mathbf{1}$ with the identity matrix I. We define

(5.3)
$$R_{\delta}u(z) = \int_{\mathbb{C}^n} u(Az)\psi_{\delta}(A) \, d\lambda(A) = \int_{\mathbb{C}^n} u((I+\delta B)z)\psi(B) \, d\lambda(B)$$
$$= \int_{\mathbb{C}^n} u((1+\delta w_1)z_1, \dots, (1+\delta w_n)z_n)\psi(w) \, d\lambda(w), \qquad z \in \mathbb{C}^n,$$

where we choose a function $0 \leq \psi \in \mathcal{C}_0^{\infty}(\mathbb{C}^n)$, rotationally symmetric in each variable with $\int_{\mathbb{C}^n} \psi d\lambda = 1$, and set $\psi_{\delta}(z) = \delta^{-2n} \psi((z-1)/\delta)$. By the Fubini-Tonelli theorem $R_{\delta} \colon L^1_{\text{loc}}(\mathbb{C}^n) \to L^1_{\text{loc}}(\mathbb{C}^n)$ and $R_{\delta}u \to u$ in the L^1_{loc} topology as $\delta \to 0$ and with local uniform convergence for $u \in \mathcal{C}(\mathbb{C}^n)$. Furthermore, $R_{\delta} \colon \mathcal{PSH}(\mathbb{C}^n) \to \mathcal{PSH}(\mathbb{C}^n)$. This implies that if $u \in \mathcal{L}^S(\mathbb{C}^n)$ then $R_{\delta}u \in \mathcal{L}^S(\mathbb{C}^n)$, more precisely, if $u \leq c_u + H_S$, Proposition 3.2 implies

$$R_{\delta}u(z) \leq \int_{\mathbb{C}^n} \left(c_u + H_S(\mathbf{1} + \delta w) + H_S(z) \right) \psi(w) \, d\lambda(w)$$

$$\leq c_u + C\sigma_S \delta + H_S(z), \qquad z \in \mathbb{C}^n,$$

where $C = \sup_{w \in \operatorname{supp} \psi} ||w||_{\infty}$. The linear map $\mathbb{C}^n \to \mathbb{C}^n$, $A \mapsto Az$, has the Jacobi determinant $|z_1 \cdots z_n|^2$, so for every $z \in \mathbb{C}^{*n}$ and corresponding matrix Z with z on the diagonal we have

$$R_{\delta}u(z) = \int_{\mathbb{C}^n} u(w)\psi_{\delta}(Z^{-1}w) |z_1\cdots z_n|^{-2} d\lambda(w)$$
$$= \int_{\mathbb{C}^n} u(w)\psi_{\delta}(w_1/z_1,\dots,w_n/z_n) |z_1\cdots z_n|^{-2} d\lambda(w).$$

By applying the Lebesgue dominated convergence theorem we may differentiate with respect to z_j under the integral sign infinitely often, so this shows that for every $u \in L^1_{\text{loc}}(\mathbb{C}^n)$ we have $R_{\delta}u \in \mathcal{C}^{\infty}(\mathbb{C}^{*n})$.

Proposition 5.4 Let S be a compact convex subset of \mathbb{R}^n_+ with $0 \in S$ and q be an admissible weight on a compact subset K of \mathbb{C}^n . Then

- (i) $V_{K,q}^S \in \mathcal{LSC}(\mathbb{C}^{*n})$, and if $\mathcal{L}^S(\mathbb{C}^n)$ is preserved under convolution (see Theorem 5.8) then $V_{K,q}^S \in \mathcal{LSC}(\mathbb{C}^n)$.
- (ii) If $V_{K,q}^{S*} \leq q$ on K then $V_{K,q}^S \in \mathcal{L}^S(\mathbb{C}^n) \cap \mathcal{C}(\mathbb{C}^{*n})$ and if $\log \Phi_{K,q}^S = V_{K,q}^S$ then $V_{K,q}^S \in \mathcal{C}(\mathbb{C}^n)$.

Proof: (i) It is sufficient to show that there exists an increasing sequence $(u_j)_{j\in\mathbb{N}}$ in $\mathcal{L}^S(\mathbb{C}^n)\cap\mathcal{C}(\mathbb{C}^{*n})$ such that $u_j\leq q$ on K for every j and such that $\lim_{j\to\infty} u_j(z) = V_{K,q}^S(z)$ for every $z\in\mathbb{C}^{*n}$. (See Klimek [19, Section 2.3].) We take $u\in\mathcal{L}^S(\mathbb{C}^n), \varepsilon>0$, and let R_δ be the regularization operator (5.3). Since $R_\delta u\searrow u$ and $q\in\mathcal{LSC}(K)$ there exists δ_ε such that $u-\varepsilon\leq R_\delta u-\varepsilon< q$ on K for every $\delta\leq\delta_\varepsilon$. These estimates tell us that there exists a family $\mathcal{F}\subseteq\mathcal{L}^S(\mathbb{C}^n)\cap\mathcal{C}^\infty(\mathbb{C}^{*n})$ such that $V_{K,q}^S=\sup\mathcal{F}$. By the Choquet lemma $V_{K,q}^S=\sup\mathcal{G}$ for some countable subfamily \mathcal{G} . By arranging the elements of \mathcal{G} into a sequence $(v_j)_{j\in\mathbb{N}}$ and then setting $u_j=\max_{k\leq j}v_k$, we have $u_j\nearrow V_{K,q}^S$ as $j\to\infty$. Hence, $V_{K,q}^S\in\mathcal{LSC}(\mathbb{C}^{*n})$. This is a modification of the proof of [31, Proposition 2.12] where the regularization operator is given by (5.1), and that is the second statement. (ii) We have $V_{K,q}^{S*}\geq V_{K,q}^S$ and by the definition of $V_{K,q}^S$ we have $V_{K,q}^{S*}\leq V_{K,q}^S$. Hence, $V_{K,q}^S=V_{K,q}^{S*}\in\mathcal{USC}(\mathbb{C}^n)$ and from (i) it follows that $V_{K,q}^S\in\mathcal{C}(\mathbb{C}^{*n})$. The last statement follows from the fact that $\Phi_{E,q}^S\in\mathcal{LSC}(\mathbb{C}^n)$.

Our next task is to characterize the $\mathcal{L}^{S}(\mathbb{C}^{n})$ classes that are invariant under the convolution operator $u \mapsto u * \psi$ given by (5.1). For that purpose we need to review a few facts from convexity theory and define the hull of a convex set in \mathbb{R}^{n}_{+} with respect to a cone. Recall that a point x in a convex set S is said to be an *extreme point of* S if the only representation of x as a convex combination x = (1 - t)a + tb with $a, b \in S$ and $t \in]0, 1[$ is the case x = a = b. We let ext S denote the set of all extreme points in the convex set S. Note that by the Minkowski theorem [17, Theorem 2.1.9] every non-empty compact convex set S is the closed convex hull of its extreme points and

(5.4)
$$\varphi_S(\xi) = \max_{x \in \text{ext}\,S} \langle x, \xi \rangle, \qquad \xi \in \mathbb{R}^n.$$

Every affine hyperplane $\{x \in \mathbb{R}^n ; \langle x, \xi \rangle = \varphi_S(\xi)\}$ with $\xi \in \mathbb{R}^{n*}$ is called a supporting hyperplane of the set S. For every $s \in \partial S$ the set $N_s^S = \{\xi \in \mathbb{R}^n ; \langle s, \xi \rangle = \varphi_S(\xi)\}$ is a convex cone which is called the normal cone of S at the point s. For every ξ the upper bound in (5.4) is attained at some $s \in \text{ext } S$, so $\mathbb{R}^n = \bigcup_{s \in \text{ext } S} N_s^S$.

Every normal cone N_s^S is an intersection of closed half spaces in \mathbb{R}^n with the origin in the boundary hyperplanes, that is $N_s^S = \bigcap_{t \in \partial S} \{\xi \in \mathbb{R}^n ; \langle s - t, \xi \rangle \ge 0\}$. Observe that ∂S can be replaced by the set of extreme points ext S. In the case S is a convex polytope and s is an extreme point, then N_s^S is an intersection of finitely many half spaces in \mathbb{R}^n with boundary hyperplanes containing the origin. See Figure 2(b).

If $\xi \in \mathbb{R}^{n*}$ and $\xi_0 \in \mathbb{R}$, then $|\langle x, \xi \rangle + \xi_0|/|\xi|$ is the distance from the point $x \in \mathbb{R}^n$ to the hyperplane $\{y \in \mathbb{R}^n; \langle y, \xi \rangle + \xi_0 = 0\}$. For supporting hyperplanes with normal ξ we have $\xi_0 = -\varphi_S(\xi)$. Hence, $|\varphi_S(\xi)|/|\xi|$ is the euclidean distance from the origin in \mathbb{R}^n to the supporting hyperplane $\{x \in \mathbb{R}^n; \langle x, \xi \rangle = \varphi_S(\xi)\}$.

Recall that a subset Γ of \mathbb{R}^n is said to be a cone if $t\xi \in \Gamma$ for every $\xi \in \Gamma$ and $t \in \mathbb{R}_+$. The dual cone $\Gamma^\circ = \{x \in \mathbb{R}^n ; \langle x, \xi \rangle \ge 0 \ \forall \xi \in \Gamma\}$ of Γ is a closed convex cone and if $\Gamma \neq \mathbb{R}^n$ is closed and convex, then $\Gamma^{\circ\circ} = \Gamma$.

Definition 5.5 For every cone $\Gamma \subset (\mathbb{R}^n \setminus \mathbb{R}^n_-) \cup \{0\}$ with $\Gamma \neq \{0\}$ and every subset S of \mathbb{R}^n_+ we define the Γ -hull of S by

$$\widehat{S}_{\Gamma} = \{ x \in \mathbb{R}^n_+ ; \, \langle x, \xi \rangle \le \varphi_S(\xi) \,\,\forall \xi \in \Gamma \},\$$

and we say that S is Γ -convex if $S = \widehat{S}_{\Gamma}$.

Note that if $\Gamma_1 \subseteq \Gamma_2$ then $\widehat{S}_{\Gamma_2} \subseteq \widehat{S}_{\Gamma_1}$. For every compact and convex S we have $S = \{x \in \mathbb{R}^n ; \langle x, \xi \rangle \leq \varphi_S(\xi) \ \forall \xi \in \mathbb{R}^n\} = \widehat{S}_{\mathbb{R}^n}$, which implies that $S \subseteq \widehat{S}_{\Gamma}$ for every cone $\Gamma \subseteq \mathbb{R}^n$.

Proposition 5.6 Let S be a compact convex subset of \mathbb{R}^n_+ with $0 \in S$ and Γ be a proper closed convex cone containing at least one point in \mathbb{R}^{*n}_+ . Then

$$\widehat{S}_{\Gamma} = (S - \Gamma^{\circ}) \cap \mathbb{R}^{n}_{+},$$

and $\varphi_{\widehat{S}_{\Gamma}}(\xi) = \varphi_{S}(\xi)$ holds for every $\xi \in \Gamma$. If for every $\xi \in \mathbb{R}^{n}$ and every extreme point x of \widehat{S}_{Γ} there exists $\eta \in \Gamma$ such that $\langle x, \xi \rangle \leq \langle x, \eta \rangle$ and $\varphi_{S}(\eta) = \varphi_{S}(\xi)$, then S is Γ -convex.

Proof: Take $a = s - t \in S' = (S - \Gamma^{\circ}) \cap \mathbb{R}^{n}_{+}$ with $s \in S$ and $t \in \Gamma^{\circ}$. For every $\xi \in \Gamma$ we have $\langle t, \xi \rangle \geq 0$ which implies $\langle a, \xi \rangle = \langle s, \xi \rangle - \langle t, \xi \rangle \leq \varphi_{S}(\xi)$ and $a \in \widehat{S}_{\Gamma}$.

Conversely, we take $a \notin S'$ and prove that $a \notin \widehat{S}_{\Gamma}$. Without restriction, we may assume that $a \in \mathbb{R}^n_+$. Since $S - \Gamma^\circ$ is convex the Hahn-Banach theorem implies that $\{a\}$ and $S - \Gamma^\circ$ can be separated by an affine hyperplane. Hence, there exists $\xi \in \mathbb{R}^n \setminus \{0\}$ and $c \in \mathbb{R}$ such that $\langle a, \xi \rangle > c$ and $\langle x, \xi \rangle \leq c$ for every $x \in S - \Gamma^\circ$. By replacing c by $\sup_{x \in S - \Gamma^\circ} \langle x, \xi \rangle$ we may assume that there exists $s \in S$ and $t \in \Gamma^\circ$ such that $\langle s - t, \xi \rangle = c$. Now we need to prove that $\xi \in \Gamma = \Gamma^{\circ\circ}$ by showing that $\langle y, \xi \rangle \geq 0$ for every $y \in \Gamma^\circ$. Since Γ° is a convex cone, we have $t + y \in \Gamma^\circ$ and $c - \langle y, \xi \rangle =$ $\langle s - t - y, \xi \rangle \leq c$. Hence, $\langle y, \xi \rangle \geq 0$. This implies that $\langle a, \xi \rangle > c \geq \varphi_S(\xi)$ and we conclude that $a \notin \widehat{S}_{\Gamma}$.

Let $\xi \in \Gamma$ and take $a = s - t \in \widehat{S}_{\Gamma}$, such that $s \in S$, $t \in \Gamma^{\circ}$, and $\varphi_{\widehat{S}_{\Gamma}}(\xi) = \langle s - t, \xi \rangle$. Since $\langle s, \xi \rangle \leq \varphi_{S}(\xi) \leq \varphi_{\widehat{S}_{\Gamma}}(\xi) = \langle s, \xi \rangle - \langle t, \xi \rangle$ and $\langle t, \xi \rangle \geq 0$, we conclude that t = 0and $\varphi_{S}(\xi) = \varphi_{\widehat{S}_{\Gamma}}(\xi)$. If for every $\xi \in \mathbb{R}^{n}$ and every extreme point x of \widehat{S}_{Γ} there exists $\eta \in \Gamma$ such that $\langle x, \xi \rangle \leq \langle x, \eta \rangle$ and $\varphi_{S}(\eta) = \varphi_{S}(\xi)$, then $\widehat{S}_{\Gamma} = \widehat{S}_{\mathbb{R}^{n}} = S$. **Definition 5.7** We say that a subset S of \mathbb{R}^n_+ is a *lower set* if for every $s \in S$ the cube $C_s = [0, s_1] \times \cdots \times [0, s_n]$ is a subset of S and we call

$$\widehat{S}_{\mathbb{R}^n_+} = \bigcup_{s \in S} C_s = (S - \mathbb{R}^n_+) \cap \mathbb{R}^n_+$$

the lower hull of S.

We have a characterization of lower sets:

Theorem 5.8 Let $S \subset \mathbb{R}^n_+$ be compact and convex with $0 \in S$ and set $\sigma_S = \varphi_S(\mathbf{1})$. Then the following are equivalent:

(i) $S \subset \mathbb{R}^n_+$ is a lower set,

(ii)
$$S = \widehat{S}_{\mathbb{R}^n_+}$$

- (iii) $\varphi_S(\xi) = \varphi_S(\xi^+)$ for every $\xi \in \mathbb{R}^n$,
- (iv) $H_S(z-w) \leq \sigma_S ||w||_{\infty} + H_S(z)$ for every $z, w \in \mathbb{C}^n$,
- (v) $\mathcal{L}^{S}(\mathbb{C}^{n})$ is translation invariant, and
- (vi) if $u \in \mathcal{L}^{S}(\mathbb{C}^{n})$ then $u * \psi \in \mathcal{L}^{S}(\mathbb{C}^{n})$ for every $\psi \in \mathcal{C}_{0}^{\infty}(\mathbb{C}^{n})$ with $\psi \geq 0$ and $\int_{\mathbb{C}^{n}} \psi \, d\lambda = 1$.

Proof: (i) \Leftrightarrow (ii) \Leftrightarrow (iii): Observe that the extreme points of C_s are $t = (t_1, \ldots, t_n)$ with $t_j = 0$ or $t_j = s_j$, so $\varphi_{C_s}(\xi) = \sup_{t \in \text{ext } C_s} \langle t, \xi \rangle = \langle s, \xi^+ \rangle$ for every $\xi \in \mathbb{R}^n$. Hence, for every lower set S we have $\varphi_S(\xi) = \sup_{s \in S} \langle s, \xi^+ \rangle = \varphi_S(\xi^+)$ for $\xi \in \mathbb{R}^n$. This equivalence follows from Proposition 5.6 with $\eta = \xi^+$.

(iii) \Rightarrow (iv): Let $w \in \mathbb{C}^n$, $z \in \mathbb{C}^{*n}$ and assume that $z - w \in \mathbb{C}^{*n}$. Since $\varphi_S(\xi) - \varphi_S(\eta) \le \varphi_S(\xi - \eta)$ and $\varphi(\xi) \le \sigma_S ||\xi||_{\infty}$ for every $\xi, \eta \in \mathbb{R}^n$ and $|\log^+ x - \log^+ y| \le |x - y|$ for every $x, y \in \mathbb{R}_+$, (iii) implies that

$$H_{S}(z-w) - H_{S}(z) = \varphi_{S}(\operatorname{Log}^{+}(z-w)) - \varphi_{S}(\operatorname{Log}^{+}(z)) \\ \leq \sigma_{S} \max_{j} |\log^{+}|z_{j} - w_{j}| - \log^{+}|z_{j}|| \leq \sigma_{S} \max_{j} ||z_{j} - w_{j}| - |z_{j}|| \leq \sigma_{S} ||w||_{\infty}.$$

By continuity of the function H_S the inequality follows.

(iv) \Rightarrow (v): If $u \leq c_u + H_S$, then $u(\cdot - w) \leq c_u + \sigma_S ||w||_{\infty} + H_S$ for every $w \in \mathbb{C}^n$. (v) \Rightarrow (vi): It is sufficient to show that $H_S * \psi \in \mathcal{L}^S(\mathbb{C}^n)$. We observe that for every $\gamma \in]0,1[$ the Riemann sum $A_{\gamma} = \sum_{\alpha \in \mathbb{Z}^{2n}} \psi(\gamma \alpha) \gamma^{-2n}$ tends to $1 = \int_{\mathbb{C}^n} \psi \, d\lambda$ as $\gamma \to 0$. This implies that the function $u_{\gamma} : \mathbb{C}^n \to \mathbb{R}$ defined by the Riemann sum

$$u_{\gamma}(z) = \sum_{\alpha \in \mathbb{Z}^{2n}} H_S(z - \gamma \alpha) \psi(\gamma \alpha) \gamma^{-2n} / A_{\gamma}$$

tends to $H_S * \psi$ as $\gamma \to 0$. By (v) the function u_{γ} is in $\mathcal{L}^S(\mathbb{C}^n)$ for it is a convex combination of functions in $\mathcal{L}^S(\mathbb{C}^n)$ and we have $u_{\gamma} - c_{\gamma} \leq V_{\mathbb{D}^n}^S = H_S$, where $c_{\gamma} =$ $\sup_{\overline{\mathbb{D}}^n} u_{\gamma}$. Furthermore, since $H_S \in \mathcal{C}(\mathbb{C}^n)$ the convergence is locally uniform, so by Proposition 4.3 we have $H_S * \psi(z) \leq \sup_{\overline{\mathbb{D}}^n} H_S * \psi + H_S(z)$ for $z \in \mathbb{C}^n$ and conclude that $H_S * \psi \in \mathcal{L}^S(\mathbb{C}^n)$.

(vi) \Rightarrow (iii): We begin by taking $\psi(\zeta) = \chi(\zeta_1) \cdots \chi(\zeta_n)$, where $0 \leq \chi \in \mathcal{C}_0^{\infty}(\mathbb{C})$ is rotationally invariant, supp $\chi \subset \overline{\mathbb{D}}$, and $\int_{\mathbb{C}} \chi d\lambda = 1$ and observe that with this choice of ψ we have $(f * \psi)(z) = \sum_{j=1}^n (f_j * \chi)(z_j)$ for any locally integrable function of the form $f(z) = \sum_{j=1}^n f_j(z_j)$.

Let $\eta \in \mathbb{R}^{*n}$ have at least one strictly negative coordinate, enumerate the coordinates so that $\eta_j < 0$ for $j = 1, \ldots, \ell$ and $\eta_j > 0$ for $j = \ell + 1, \ldots, n$, and take $s \in S$ such that $\varphi_S(\eta) = \langle s, \eta \rangle$. Then for every t > 0 we have

$$\begin{split} H_{S} * \psi(e^{t\eta}) - H_{S}(e^{t\eta}) &= \int_{\mathbb{C}^{n}} \varphi_{S}(\operatorname{Log}(e^{t\eta} - \zeta)) \psi(\zeta) \, d\lambda(\zeta) - \langle s, t\eta \rangle \\ &\geq \int_{\mathbb{C}^{n}} \langle s, \operatorname{Log}(e^{t\eta} - \zeta) - t\eta \rangle \psi(\zeta) \, d\lambda(\zeta) \\ &= \sum_{j=1}^{n} s_{j} \int_{\mathbb{D}} \left(\log |e^{t\eta_{j}} - \zeta_{j}| - t\eta_{j} \right) \chi(\zeta_{j}) \, d\lambda(\zeta_{j}) \\ &= -t \langle s', \eta' \rangle + \sum_{j=1}^{\ell} s_{j} \int_{\mathbb{D}} \log |e^{t\eta_{j}} - \zeta_{j}| \chi(\zeta_{j}) \, d\lambda(\zeta_{j}) \\ &+ \sum_{j=\ell+1}^{n} s_{j} \int_{\mathbb{D}} \log |1 - e^{-t\eta_{j}} \zeta_{j}| \chi(\zeta_{j}) \, d\lambda(\zeta_{j}). \end{split}$$

We let $v = \log |\cdot| \in SH(\mathbb{C}) \cap H(\mathbb{C}^*)$. Then for $j = 1, \ldots, \ell$

$$\int_{\mathbb{D}} \log |e^{t\eta_j} - \zeta_j| \, \chi(\zeta_j) \, d\lambda(\zeta_j) = (v * \chi)(e^{t\eta_j}) \to v * \chi(0), \qquad t \to +\infty.$$

Since $v \in \mathcal{H}(\mathbb{C}^*)$, χ is rotationally invariant, and $\operatorname{supp} \chi \subset \overline{\mathbb{D}}$, the mean value property gives for $j = \ell + 1, \ldots, n$

$$\int_{\mathbb{D}} \log |1 - e^{-t\eta_j} \zeta_j| \, \chi(\zeta_j) \, d\lambda(\zeta_j) = \log 1 = 0.$$

Hence,

(5.5)
$$H_S * \psi(e^{t\eta}) - H_S(e^{t\eta}) \ge -t\langle s', \eta' \rangle + \sum_{j=1}^{\ell} s_j(v * \chi)(e^{t\eta_j}).$$

Assume that (iii) does not hold and take $\xi_0 \in \mathbb{R}^n$ such that $\varphi_S(\xi_0) < \varphi_S(\xi_0^+)$. Then at least one coordinate of ξ_0 is strictly negative. By continuity we can choose $\xi_0 \in \mathbb{R}^{*n}$, and we can renumber the coordinates so that $\xi_{0,j} < 0$ for $j = 1, \ldots, \ell$ and $\xi_{0,j} > 0$ for $j = \ell + 1, \ldots, n$. By continuity there exists an open neighborhood U of ξ_0^+ such that $\varphi_S(\xi_0) < \varphi_S(\eta)$ for every $\eta \in U$. We fix $\eta = (\eta', \xi_0'') \in U$ with $\eta_j < 0$ for $j = 1, \ldots, \ell$. There exists a point $s = (s', s'') \in \partial S$, such that $\langle s, \eta \rangle = \langle s', \eta' \rangle + \langle s'', \eta'' \rangle = \varphi_S(\eta)$. We have $\langle s', \eta' \rangle \leq 0$. Equality is excluded for it would imply s' = 0' and $\varphi_S(\eta) = \langle s'', \xi_0'' \rangle =$ $\langle (0', s''), \xi_0 \rangle = \langle s, \xi_0 \rangle \leq \varphi_S(\xi_0)$, contradicting the choice of η . Hence, $\langle s', \eta' \rangle < 0$ and the estimate (5.5) implies that $H_S * \psi - H_S$ is unbounded, contradicting (vi).

6 Monge-Ampère masses

The equilibrium measure for a bounded non-pluripolar set $E \subset \mathbb{C}^n$ is the Monge-Ampère operator of V_E , defined as $\mu_E = (dd^c V_E^*)^n$ where $d^c = i(\overline{\partial} - \partial)$ and $(dd^c V_E^*)^n = dd^c V_E^* \wedge \cdots \wedge dd^c V_E^*$ is defined in terms of currents. Similarly denote the Monge-Ampère measure of $V_{E,q}^{S*}$ by

$$\mu_{E,q}^S = \left(dd^c V_{E,q}^{S*} \right)^n.$$

Theorem 6.1 Let $S \subset \mathbb{R}^n_+$ be compact and convex with $0 \in S$, $E \subset \mathbb{C}^n$ be closed, and q be an admissible weight on E. Then

- (i) supp $\mu_{E,q}^S \subseteq \{z \in E; V_{E,q}^{S*}(z) \ge q(z)\}$, and
- (ii) $\{z \in E; V_{E,q}^{S*}(z) > q(z)\}$ is pluripolar.

Proof: (i) We need to prove that $V_{E,q}^{S*}$ is maximal in $U = (\mathbb{C}^n \setminus E) \cup \{z \in E; V_{E,q}^{S*}(z) < q(z)\}$, which is open. Take $a \in U$. If $a \in \mathbb{C}^n \setminus E$ then we take r > 0 such that $B(a,r) \in \mathbb{C}^n \setminus E$. If on the other hand $a \in E$ and $V_{E,q}^{S*}(a) < q(a)$ then by upper semicontinuity of $V_{E,q}^{S*}$ and lower semicontinuity of q there exists r > 0 such that

$$\sup_{\zeta \in B(a,r)} V_{E,q}^{S*}(\zeta) < \inf_{\zeta \in E \cap B(a,r)} q(\zeta).$$

We need to prove that the restriction of $\mu_{E,q}^S$ to some B(a,r) is the zero measure. We let $V \in \mathcal{PSH}(\mathbb{C}^n)$ be the function given by

$$V(z) = \begin{cases} V_{E,q}^{S*}(z), & z \in \mathbb{C}^n \setminus B(a,r), \\ u(z), & z \in B(a,r), \end{cases}$$

where u is the Perron-Bremermann function on B(a, r) with boundary values $V_{E,a}^{S*}$, i.e.,

$$u(z) = \sup\{v(z); v \in \mathcal{PSH}(B(a,r)), v^* \leq V_{E,q}^{S*} \text{ on } \partial B(a,r)\}.$$

Then $V_{E,q}^{S*} \leq V$ and since V only deviates from $V_{E,q}^{S*}$ on a compact set we have $V \in \mathcal{L}^{S}(\mathbb{C}^{n})$ by Proposition 4.6. Furthermore, in the case $V_{E,q}^{S*}(a) < q(a)$ the maximum principle implies

$$V(z) \le \sup_{\zeta \in B(a,r)} V_{E,q}^{S*}(\zeta) < \inf_{\zeta \in B(a,r)} q(\zeta) \le q(z), \qquad z \in B(a,r),$$

and it implies that $V \leq q$ on E. Hence, $V = V_{E,q}^{S*}$ and by Klimek [19, Theorem 4.4.1], $\mu_{E,q}^S = \left(dd^c V_{E,q}^{S*} \right)^n = 0$ on B(a, r).

(ii) Since $V_{E,q}^S \leq q$ on E we have $\{z \in E; q(z) < +\infty\} \subseteq \{z \in \mathbb{C}^n; V_{E,q}^S < +\infty\}$ Since q is admissible, the left-hand side is non-pluripolar, and then so is the right-hand side. Since $\mathcal{L}^S(\mathbb{C}^n)/\varphi_S(\mathbf{1}) \subset \mathcal{L}(\mathbb{C}^n)$ it follows from Klimek [19, Proposition 5.2.1 and Theorems 5.2.4 and 4.7.6] the set $\{z \in E; q < V_{E,q}^{S*}(z)\} \subseteq \{z \in \mathbb{C}^n; V_{E,q}^S(z) < V_{E,q}^{S*}(z)\}$ is pluripolar. The complex Monge-Ampère mass of H_S can be described in terms of the real Monge-Ampère mass of φ_S . Let $U_{\ell} = H_S * \psi_{\delta_{\ell}} \searrow H_S$, where $0 < \delta_{\ell} \searrow 0$ and $\psi_{\delta_{\ell}}(\zeta) = \delta_{\ell}^{-2n} \psi(\zeta/\delta_{\ell}), \ \psi(\zeta) = \chi(\zeta_1) \cdots \chi(\zeta_n)$, where $0 \leq \chi \in \mathcal{C}_0^{\infty}(\mathbb{C})$ is rotationally invariant, $\operatorname{supp} \chi \subset \overline{\mathbb{D}}$, and $\int_{\mathbb{C}} \chi d\lambda = 1$. Then $U_{\ell} \in \mathcal{C}^{\infty}(\mathbb{C}^n) \cap \mathcal{PSH}(\mathbb{C}^n)$ and $U_{\ell}(z) = U_{\ell}(|z_1|, \ldots, |z_n|)$ holds. We set $v_{\ell}(\xi) = U_{\ell}(e^{\xi_1}, \ldots, e^{\xi_n})$. Since

$$\frac{\partial^2 U_\ell(z)}{\partial z_j \partial \overline{z}_k} = \frac{1}{4z_j \overline{z}_k} \cdot \frac{\partial^2 v_\ell(\xi)}{\partial \xi_j \partial \xi_k}, \qquad z \in \mathbb{C}^{*n},$$

it follows that

$$\det\left(\frac{\partial^2 U_{\ell}(z)}{\partial z_j \partial \overline{z}_k}\right) = \frac{1}{4^n |z_1 \cdots z_n|^2} \det\left(\frac{\partial^2 v_{\ell}(\xi)}{\partial \xi_j \partial \xi_k}\right) \Big|_{\xi = \operatorname{Log} z}$$

With $z_j = e^{\xi_j + i\theta_j}$, $\theta_j \in [0, 2\pi]$ we have

$$\left(\frac{i}{2}\right)^n dz_1 \wedge d\overline{z}_1 \wedge \dots \wedge dz_n \wedge d\overline{z}_n = |z_1 \cdots z_n|^2 d\xi d\theta$$

and on \mathbb{C}^{*n} we have

$$(dd^{c}U_{\ell})^{n} = n! \det\left(\frac{\partial^{2}v_{\ell}(\xi)}{\partial\xi_{j}\partial\xi_{k}}\right) d\xi d\theta.$$

The real Monge-Ampère measure of v, denoted $\mathcal{MA}_{\mathbb{R}}(v)$, is defined for \mathcal{C}^2 function by

$$\mathcal{MA}_{\mathbb{R}}(v) = \det\left(\frac{\partial^2 v(\xi)}{\partial \xi_j \partial \xi_k}\right) d\xi,$$

and extended to convex functions by locally uniform limits. This is done in Figalli [14, Proposition 2.6].

Letting $\ell \to \infty$, we have for every Borel set $E \subset \mathbb{R}^n$

$$\int_{\text{Log}^{-1}(E)} (dd^c H_S)^n = n! \left(\mathcal{MA}_{\mathbb{R}}(\varphi_S) \otimes d\theta \right) (E \times [0, 2\pi]^n) = (2\pi)^n n! \mathcal{MA}_{\mathbb{R}}(\varphi_S)(E).$$

In particular

(6.1)
$$\int_{\mathbb{T}^n} (dd^c H_S)^n = (2\pi)^n n! \mathcal{MA}_{\mathbb{R}}(\varphi_S)(\{0\}).$$

This will be useful for our next result. Its proof is borrowed from Rashkovskii [27, Theorem 3.4].

Theorem 6.2 Let $S \subset \mathbb{R}^n_+$ be compact and convex with $0 \in S$, $E \subset \mathbb{C}^n$ be closed, and q be an admissible weight on E. Then

$$\mu_{E,q}^S(\mathbb{C}^n) = (2\pi)^n n! \operatorname{vol}(S),$$

where vol denotes the euclidean volume.

Proof: By Proposition 4.5, $V_{K,q}^S - H_S$ is bounded. The comparison principle, Klimek [19, Theorem 3.7.1], then implies that

$$\mu_{K,q}^S(\mathbb{C}^n) = \int_{\mathbb{C}^n} (dd^c H_S)^n = \mu_{\mathbb{T}^n,0}^S(\mathbb{C}^n).$$

By Theorem 6.1 (i), $\mu_{\mathbb{T}^n,0}^S(\mathbb{C}^n) = \int_{\mathbb{T}^n} (dd^c H_S)^n$. We already established in (6.1) that

$$\int_{\mathbb{T}^n} (dd^c H_S)^n = (2\pi)^n n! \mathcal{MA}_{\mathbb{R}}(\varphi_S)(\{0\})$$

By Blocki [9], see also Figalli [14], we have

$$\mathcal{MA}_{\mathbb{R}}(\varphi_S)(\{0\}) = \operatorname{vol}(\{s \in \mathbb{R}^n ; \langle s, \xi \rangle \le \varphi_S(\xi), \ \forall \xi \in \mathbb{R}^n\}) = \operatorname{vol}(S).$$

7 Characterization of polynomials by L^2 -estimates

In this section we study characterization of the polynomial spaces $\mathcal{P}_m^S(\mathbb{C}^n)$ with weighted L^2 -norms of entire functions. Recall that the Liouville type Theorem 3.6 states that if $f \in \mathcal{O}(\mathbb{C}^n)$ and $|f|e^{-mH_S-a\log^+ \|\cdot\|_{\infty}} \in L^{\infty}(\mathbb{C}^n)$ for some $a \in [0, d_m[$, where d_m is the distance between mS and $\mathbb{N}^n \setminus mS$ in the L^1 -norm, then $f \in \mathcal{P}_m^S(\mathbb{C}^n)$. We take a measurable function $\psi \colon \mathbb{C}^n \to \mathbb{R}$ and let $L^2(\mathbb{C}^n, \psi)$ denote the space of all measurable $f \colon \mathbb{C}^n \to \mathbb{C}$ such that

(7.1)
$$||f||_{\psi}^2 = \int_{\mathbb{C}^n} |f|^2 e^{-\psi} d\lambda < +\infty.$$

Proposition 7.1 Let $f \in L^2(\mathbb{C}^n, \psi) \cap \mathcal{O}(\mathbb{C}^n)$ for some measurable $\psi \colon \mathbb{C}^n \to \overline{\mathbb{R}}$ with power series expansion $f(z) = \sum_{\alpha \in \mathbb{N}^n} a_\alpha z^\alpha$ at the origin. Then for every polyannulus $A_{\sigma,\tau} = \{\zeta \in \mathbb{C}^n ; e^{\sigma_j} \leq |\zeta_j| < e^{\tau_j}\}$ in \mathbb{C}^n , where $\sigma, \tau \in \mathbb{R}^n, \sigma_j < \tau_j$ for $j = 1, \ldots, n$, with volume $v(A_{\sigma,\tau}) = \pi^n \prod_{j=1}^n (e^{2\tau_j} - e^{2\sigma_j})$, we have

(7.2)
$$|a_{\alpha}| \leq \frac{\|f\|_{\psi}}{v(A_{\sigma,\tau})} \left(\int_{A_{\sigma,\tau}} \frac{e^{\psi(\zeta)}}{|\zeta^{\alpha}|^2} d\lambda(\zeta) \right)^{1/2}.$$

Furthermore, if ψ is rotationally invariant in each variable ζ_j , then for $K_{\sigma,\tau} = \prod_{j=1}^n [\sigma_j, \tau_j] \subset \mathbb{R}^n$ and $\chi(\xi) = \frac{1}{2}\psi(e^{\xi_1}, \ldots, e^{\xi_n})$ we have

(7.3)
$$|a_{\alpha}| \leq \frac{\|f\|_{\psi}}{\prod_{j=1}^{n} (1 - e^{-2(\tau_j - \sigma_j)})} e^{-\langle \mathbf{1}, \tau \rangle} \left(\int_{K_{\sigma, \tau}} e^{2(\chi(\xi) - \langle \alpha, \xi \rangle)} d\lambda(\xi) \right)^{1/2}.$$

Proof: By the Cauchy formula for derivatives

$$a_{\alpha} = \frac{1}{(2\pi i)^n} \int_{C_r} \frac{f(\zeta)}{\zeta^{\alpha}} \cdot \frac{d\zeta_1 \cdots d\zeta_n}{\zeta_1 \cdots \zeta_n} = \frac{1}{(2\pi)^n} \int_{[-\pi,\pi]^n} \frac{f(r_1 e^{i\theta_1}, \dots, r_n e^{i\theta_n})}{r^{\alpha} e^{i\langle \alpha, \theta \rangle}} \, d\theta_1 \cdots d\theta_n,$$

where $C_r = \{z \in \mathbb{C}^n; |z_j| = r_j\}$, is any polycircle with center 0 and polyradius $r \in \mathbb{R}^{*n}_+$. We parametrize C_r by $[-\pi, \pi]^n \ni \theta \mapsto (r_1 e^{i\theta_1}, \ldots, r_n e^{i\theta_n})$, multiply the integral by $r_1 \cdots r_n dr_1 \cdots dr_n$, integrate with respect to r_j over $[e^{\sigma_j}, e^{\tau_j}]$, note that $\int_{[e^{\sigma_j}, e^{\tau_j}]} r_j dr_j = \frac{1}{2}(e^{2\tau_j} - e^{2\sigma_j})$, set $L_{\sigma,\tau} = \prod_{j=1}^n ([e^{\sigma_j}, e^{\tau_j}] \times [-\pi, \pi])$, and get

$$a_{\alpha} = \frac{1}{v(A_{\sigma,\tau})} \int_{L_{\sigma,\tau}} \frac{f(r_1 e^{i\theta_1}, \dots, r_n e^{i\theta_n})}{r^{\alpha} e^{i\langle\alpha,\theta\rangle}} (r_1 dr_1 d\theta_1) \cdots (r_n dr_n d\theta_n)$$

$$= \frac{1}{v(A_{\sigma,\tau})} \int_{A_{\sigma,\tau}} \frac{f(\zeta)}{\zeta^{\alpha}} d\lambda(\zeta) = \frac{1}{v(A_{\sigma,\tau})} \int_{A_{\sigma,\tau}} f(\zeta) e^{-\psi(\zeta)/2} \cdot \frac{e^{\psi(\zeta)/2}}{\zeta^{\alpha}} d\lambda(\zeta).$$

Now (7.2) follows from the Cauchy-Schwarz inequality and (7.3) by integrating over the angular variables and also using the fact that $v(A_{\sigma,\tau}) = \pi^n \prod_{j=1}^n (e^{2\tau_j} - e^{2\sigma_j})$. \Box

Theorem 7.2 Let S be a compact convex subset of \mathbb{R}^n_+ with $0 \in S$, $m \in \mathbb{N}^*$, and $d_m = d(mS, \mathbb{N}^n \setminus mS)$ denote the euclidean distance between mS and $\mathbb{N}^n \setminus mS$. If $f \in \mathcal{O}(\mathbb{C}^n)$ and for some $a \in [0, d_m]$

(7.4)
$$\int_{\mathbb{C}^n} |f|^2 (1+|\zeta|^2)^{-a} e^{-2mH_S} \, d\lambda < +\infty,$$

then $f \in \mathcal{P}_m^{\widehat{S}_{\Gamma}}(\mathbb{C}^n)$, where \widehat{S}_{Γ} is the Γ -hull of S and $\Gamma = \Gamma_a$ consists of all ξ such that the angle between the vectors $\mathbf{1} = (1, \ldots, 1)$ and ξ is $\leq \arccos(-(d_m - a)/\sqrt{n})$.



Figure 1: The cone Γ has opening angle $\theta = \arccos(-(d_m - a)/\sqrt{n})$

Observe that the largest possible d_m is 1 and smallest possible a is 0, which implies that the largest possible opening angle of the cone Γ is $\operatorname{arccos}(-1/\sqrt{n})$ in the case $d_m = 1$ and $\Gamma = \mathbb{R}^n \setminus \mathbb{R}^{*n}_-$. If a_0 is the infimum of a such that (7.4) holds, then $\Gamma_{a_0} = \bigcup_{a > a_0} \Gamma_a$. Therefore f is a polynomial with exponents in $\widehat{mS}_{\Gamma(a_0)} = \bigcap_{a > a_0} \widehat{mS}_{\Gamma(a)}$.

We are interested in conditions on cones $\Lambda \subseteq \Gamma$ which guarantee that $\widehat{S}_{\Lambda} = S$:

Corollary 7.3 The function f in Theorem 7.2 is in $\mathcal{P}_m^S(\mathbb{C}^n)$ in the cases:

(i) $S = \widehat{S}_{\Lambda}$ for some cone Λ contained in $\{\xi \in \mathbb{R}^n ; \langle \mathbf{1}, \xi \rangle \ge 0\}$.

(ii) S is a lower set, that is $S = \widehat{S}_{\mathbb{R}^n_+}$.

(iii) $(mS) \cap \mathbb{N}^n = (m\widehat{S}_{\Gamma}) \cap \mathbb{N}^n$.

Proof of Theorem 7.2: Let $f(z) = \sum_{\alpha \in \mathbb{N}^n} a_\alpha z^\alpha$ be the Taylor expansion of f at the origin. We need to show that $a_\alpha = 0$ for every $\alpha \in \mathbb{N}^n \setminus m\widehat{S}_{\Gamma}$. Since $\alpha \notin m\widehat{S}_{\Gamma}$, there exists $\tau \in \Gamma$ such that $|\tau| = 1$ and $\langle \alpha, \tau \rangle > m\varphi_S(\tau)$. By rotating τ we may assume that τ is an interior point of Γ . Since the angle between τ and $\mathbf{1}$ is $\leq \arccos(-(d_m - a)/\sqrt{n})$ we have $-\langle \mathbf{1}, \tau \rangle < d_m - a$. We choose $\varepsilon > 0$ such that $d_m - a - \varepsilon > 0$, and $-\langle \mathbf{1}, \tau \rangle < d_m - a - \varepsilon$. Recall that $\langle \alpha, \tau \rangle - m\varphi_S(\tau)$ is the euclidean distance from α to the supporting hyperplane $\{x; \langle x, \tau \rangle = m\varphi_S(\tau)\}$, so by assumption $m\varphi_S(\tau) - \langle \alpha, \tau \rangle \leq -d_m$. Hence

$$-\langle \mathbf{1}, \tau \rangle + m\varphi_S(\tau) - \langle \alpha, \tau \rangle < -a - \varepsilon_s$$

We take $\sigma \in \mathbb{R}^n \setminus \{0\}$ with $\sigma_j < \tau_j$ for $j = 1, \ldots, n$ such that

$$-\langle \mathbf{1}, \tau \rangle + m\varphi_S(\xi) - \langle \alpha, \xi \rangle < -(a+\varepsilon)|\xi|, \qquad \xi \in K_{\sigma,\tau} = \prod_{j=1}^n [\sigma_j, \tau_j].$$

By homogeneity we get

$$-t\langle \mathbf{1}, \tau \rangle + m\varphi_S(\xi) - \langle \alpha, \xi \rangle < -(a+\varepsilon)|\xi|, \qquad t > 0, \ \xi \in tK_{\sigma,\tau}.$$

Let $\xi_j = \log |\zeta_j|$ and observe that $(1 + |\zeta|^2)^a \leq (n+1)^a \max\{1, \|\zeta\|_{\infty}^{2a}\}$. From this inequality and (7.4) it follows that $f \in L^2(\mathbb{C}^n, \psi)$, where

$$\frac{1}{2}\psi(\zeta) = a\log \|\zeta\|_{\infty} + mH_S(\zeta) = a\|\xi\|_{\infty} + m\varphi_S(\xi).$$

We set $\chi(\xi) = \frac{1}{2}\psi(e^{\xi_1},\ldots,e^{\xi_n})$. Then

$$-t\langle \mathbf{1}, \tau \rangle + \chi(\xi) - \langle \alpha, \xi \rangle < -\varepsilon |\xi|, \qquad t > 0, \ \xi \in tK_{\sigma,\tau}.$$

The estimate (7.3) with $tK_{\sigma,\tau}$ in the role of $K_{\sigma,\tau}$ gives

$$|a_{\alpha}| \leq \frac{\|f\|_{\psi}}{\prod_{j=1}^{n} (1 - e^{-2(\tau_{j} - \sigma_{j})t})} e^{-t\langle \mathbf{1}, \tau \rangle} \left(\int_{tK_{\sigma,\tau}} e^{2(\chi(\xi) - \langle \alpha, \xi \rangle)} d\lambda(\xi) \right)^{1/2} \\ \leq \frac{\|f\|_{\psi}}{\prod_{j=1}^{n} (1 - e^{-2(\tau_{j} - \sigma_{j})t})} e^{-\varepsilon |\sigma| t} t^{n/2} v(K_{\sigma,\tau})^{1/2} \to 0, \quad t \to +\infty,$$

and we conclude that $a_{\alpha} = 0$.

The Γ -hull \widehat{S}_{Γ} can not be replaced by S in Theorem 7.2:

Example 7.4 Let $m \ge 4$ and $S \subseteq \mathbb{R}^2_+$ be the quadrilateral

$$S = ch\{(0,0), (a,0), (b,1-b), (0,1)\}.$$



Figure 2: (a) The set S. (b) The normal cones N_s^S of the extreme points of S.

where 0 < a < 1/m, 0 < a < b < 1, m(1-b) < 1, and (b-a)/(1-b) > m-2-am. Then $(1,0), (2,0), \ldots, (m-3,0) \notin mS$, but the calculations below show that $\|p_k\|_{2mH_S} < +\infty$ for $p_k(z) = z^{(k,0)} = z_1^k$ with $k = 1, \ldots, m-3$.

for $p_k(z) = z^{(k,0)} = z_1^k$ with $k = 1, \ldots, m-3$. Since the map $(\mathbb{R} \times] - \pi, \pi[)^2 \to \mathbb{C}^2$, $(\xi_1, \theta_1, \xi_2, \theta_2) \mapsto (e^{\xi_1 + i\theta_1}, e^{\xi_2 + i\theta_2})$ has the Jacobi determinant $e^{2\xi_1 + 2\xi_2}$, we have

$$\begin{aligned} \|p_k\|_{2mH_S}^2 &= \int_{\mathbb{C}^2} |z_1|^{2k} e^{-2mH_S(z)} \, d\lambda(z) = 4\pi^2 \int_{\mathbb{R}^2} e^{2(k+1)\xi_1 + 2\xi_2 - 2m\varphi_S(\xi)} \, d\xi_1 d\xi_2, \\ \varphi_S(\xi) &= \max_{x \in \text{ext}(S)} \langle x, \xi \rangle = \begin{cases} 0, & \xi \in N_{(0,0)}^S, \\ a\xi_1, & \xi \in N_{(a,0)}^S, \\ b\xi_1 + (1-b)\xi_2, & \xi \in N_{(b,(1-b))}^S, \\ \xi_2, & \xi \in N_{(0,1)}^S. \end{cases} \end{aligned}$$

We split the integral over \mathbb{R}^2 into the sum of the integrals over the normal cones at the extreme points of S, which we calculate as

$$\int_{N_{(a,0)}^{S}} e^{2(k+1)\xi_{1}+2\xi_{2}} d\xi = \int_{-\infty}^{0} e^{2(k+1)\xi_{1}} d\xi_{1} \int_{-\infty}^{0} e^{2\xi_{2}} d\xi_{2} = \frac{1}{4(k+1)},$$

$$\int_{N_{(a,0)}^{S}} e^{2(k+1)\xi_{1}+2\xi_{2}-2ma\xi_{1}} d\xi = \int_{0}^{\infty} e^{2(k+1-ma)\xi_{1}} \int_{-\infty}^{-\xi_{1}(b-a)/(1-b)} e^{2\xi_{2}} d\xi_{2} d\xi_{1}$$

$$= \frac{1}{4((b-a)/(1-b)+ma-1-k)},$$

 $\int_{N^S_{(b,1-b)}} e^{2(k+1)\xi_1 + 2\xi_2 - 2m(b\xi_1 + (1-b)\xi_2)} d\xi$

$$= \int_0^\infty e^{2(k+1-mb)\xi_1} \int_{-\xi_1(b-a)/(1-b)}^{\xi_1} e^{2(1-m(1-b))\xi_2} d\xi_2 d\xi_1$$

= $\frac{1}{4(1-m(1-b))} \left(\frac{1}{m-2-k} + \frac{1}{(b-a)/(1-b)+ma-1-k)} \right),$

$$\int_{N_{(0,1)}^{S}} e^{2(k+1)\xi_{1}+2\xi_{2}-2m\xi_{2}} d\xi = \int_{0}^{\infty} e^{2(1-m)\xi_{2}} \int_{-\infty}^{\xi_{2}} e^{2(k+1)\xi_{1}} d\xi_{1} d\xi_{2}$$
$$= \frac{1}{4(k+1)(m-2-k)}.$$

This shows that (7.4) is satisfied with p_k in the role of f, but $p_k \notin \mathcal{P}_m^S(\mathbb{C}^n)$.

8 Pullbacks of polynomial classes by polynomial maps

Let $S \subset \mathbb{R}^n_+$ be compact and convex with $0 \in S$ and $f = (f_1, \ldots, f_n) \colon \mathbb{C}^\ell \to \mathbb{C}^n$ be a polynomial map. If $f_j(z) = \sum_{\alpha \in \mathbb{N}^\ell} a_{j,\alpha} z^{\alpha}$, $I_j = \{\alpha \in \mathbb{N}^\ell : a_{j,\alpha} \neq 0\}$, and $S_j \subset \mathbb{R}^\ell_+$ be the convex hull of I_j . Then $f_j \in \mathcal{P}_1^{S_j}(\mathbb{C}^\ell)$ and for every $p(w) = \sum_{\beta \in (mS) \cap \mathbb{N}^n} b_\beta w^\beta$ in $\mathcal{P}_m^S(\mathbb{C}^n)$ we have

$$(f^*p)(z) = \sum_{\beta \in (mS) \cap \mathbb{N}^n} b_\beta f_1(z)^{\beta_1} \cdots f_n(z)^{\beta_n}$$

By Theorem 3.6 we have $|f_j(z)| \leq e^{c_j + H_{S_j}(z)}$ which implies that for every $z \in \mathbb{C}^{*\ell}$

$$\begin{split} |(f^*p)(z)| &\leq \sum_{\beta \in (mS) \cap \mathbb{N}^n} |b_\beta| e^{\langle c, \beta \rangle + \beta_1 H_{S_1}(z) + \dots + \beta_n H_{S_n}(z)} \\ &\leq \left(\sum_{\beta \in (mS) \cap \mathbb{N}^n} |b_\beta| e^{\langle c, \beta \rangle}\right) e^{m\varphi_S(\varphi_{S_1}(\operatorname{Log} z), \dots, \varphi_{S_n}(\operatorname{Log} z))}. \end{split}$$

We have for every $\xi \in \mathbb{R}^{\ell}$ that

$$\varphi_S(\varphi_{S_1}(\xi), \dots, \varphi_{S_n}(\xi))$$

$$= \sup\{\langle x_1s_1 + \dots + x_ns_n, \xi\rangle; x \in S, s_j \in S_j, j = 1, \dots, n\}$$

$$= \sup_{x \in S} \varphi_{x_1S_1 + \dots + x_nS_n}(\xi) = \varphi_{S'}(\xi),$$

where $S' = \bigcup_{x \in S} x_1 S_1 + \dots + x_n S_n$. The set $S' \subset \mathbb{R}^{\ell}_+$ is compact and convex.

Proposition 8.1 Assume that $\overline{S \cap \mathbb{Q}^n} = S$. Then with the notation above S' is the smallest compact convex subset T of \mathbb{R}^{ℓ}_+ with $0 \in T$ for which $f^*(\mathcal{P}^S_m(\mathbb{C}^n)) \subseteq \mathcal{P}^T_m(\mathbb{C}^{\ell})$ for all $m \in \mathbb{N}$.

Proof: Assume that $T \subsetneq S'$ such that $f^*(\mathcal{P}^S_m(\mathbb{C}^n)) \subseteq \mathcal{P}^T_m(\mathbb{C}^\ell)$ for every $m \in \mathbb{N}$. Then there exists $\xi \in \mathbb{R}^\ell$ such that $\varphi_T(\xi) < \varphi_{S'}(\xi)$ and since $S \cap \mathbb{Q}^n$ is dense in S we can choose $r \in S \cap \mathbb{Q}^n$ such that

(8.1)
$$\varphi_T(\xi) < r_1 \langle \alpha_1, \xi \rangle + \dots + r_n \langle \alpha_n, \xi \rangle \le \varphi_{S'}(\xi).$$

Now we fix $m \in \mathbb{N}$ such that $\gamma = mr \in \mathbb{N}^n$, define $p \in \mathcal{P}^S_{\mu}(\mathbb{C}^n)$ by $p(w) = w^{\gamma}$. By Theorem 3.6, there exists a constant c_p such that

(8.2)
$$|f^*p(z)| \le e^{c_p + mH_T(z)}, \qquad z \in \mathbb{C}^{\ell}.$$

Since S_j is a convex polytope in \mathbb{R}^{ℓ}_+ , the set $\bigcap_{j=1}^n \bigcup_{\alpha \in \operatorname{ext} S_j} \mathring{N}^{S_j}_{\alpha}$ is dense in \mathbb{R}^{ℓ} , where $\mathring{N}^{S_j}_{\alpha}$ is the interior of $N^{S_j}_{\alpha}$. Hence ξ may be chosen from $\mathring{N}^{S_j}_{\alpha_j}$ where $\alpha_j \in \operatorname{ext} S_j$ and $\varphi_{S_j}(\xi) = \langle \alpha_j, \xi \rangle > \langle \alpha, \xi \rangle$ for every $\alpha \in I_j \setminus \{\alpha_j\}$. All the α_j are from \mathbb{N}^{ℓ} by the definition of S_j . We define the sequence $(\zeta_k)_{k \in \mathbb{N}}$ in \mathbb{C}^{ℓ} by $\zeta_k = (e^{k\xi_1}, \ldots, e^{k\xi_\ell})$ and we will show that (8.1) implies that the sequence $(f^*p(\zeta_k)e^{-mH_T(\zeta_k)})_{k \in \mathbb{N}}$ is unbounded, contradicting the estimate (8.2). First we observe that $\zeta_k^{\alpha_j} = e^{k\langle \alpha_j, \xi \rangle}$ and then that (8.1) implies

(8.3)
$$(\zeta_k^{\alpha_1}, \dots, \zeta_k^{\alpha_n})^{\gamma} e^{-mH_T(\zeta_k)} = e^{km(r_1\langle \alpha_1, \xi \rangle + \dots + r_n\langle \alpha_n, \xi \rangle - \varphi_T(\xi))} \to +\infty$$

when $k \to +\infty$. Next we observe that

(8.4)
$$f_j(\zeta_k)/\zeta_k^{\alpha_j} = a_{j,\alpha_j} + \sum_{\alpha \in I_j \setminus \{\alpha_j\}} a_{j,\alpha} e^{-k(\langle \alpha_j, \xi \rangle - \langle \alpha, \xi \rangle)} \to a_{j,\alpha_j} \neq 0$$

when $k \to +\infty$, and

(8.5)
$$f^*p(\zeta_k)/(\zeta_k^{\alpha_1},\ldots,\zeta_k^{\alpha_n})^{\gamma} = (f_1(\zeta_k)/\zeta_k^{\alpha_1})^{\gamma_1}\cdots(f_n(\zeta_k)/\zeta_k^{\alpha_n})^{\gamma_n}.$$

By combining (8.3), (8.4), and (8.5), we see that $(f^*p(\zeta_k)e^{-mH_T(\zeta_k)})_{k\in\mathbb{N}}$ is unbounded.

Assume now that f is a proper map and that q is a given admissible weight function on a compact set $K \subseteq \mathbb{C}^n$. Then f^*q is lower semicontinuous and

$$\{z \in f^{-1}(K); f^*q(z) < +\infty\} = f^{-1}(\{w \in K; q(w) < +\infty\})$$

Since inverse images of non-pluripolar sets by proper maps are non-pluripolar it follows that f^*q is an admissible weight on $f^{-1}(K)$. Furthermore, we have

$$||f^*pe^{-mf^*q}||_{f^{-1}(K)} = ||pe^{-mq}||_K.$$

From Proposition 8.1 we conclude that $f^*(\Phi^S_{K,q,m}) \leq \Phi^{S'}_{f^{-1}(K),f^*q,m}$, for every $m \in \mathbb{N}^*$, consequently $f^*(\Phi^S_{K,q}) \leq \Phi^{S'}_{f^{-1}(K),f^*q}$ and equality holds if $f^* \colon \mathcal{P}^S_m(\mathbb{C}^n) \to \mathcal{P}^{S'}_m(\mathbb{C}^\ell)$ is surjective.

Next we look at the pullback of Lelong classes. Let $u \in \mathcal{L}^{S}(\mathbb{C}^{n})$, say $u \leq c_{u} + H_{S}$. Then for every $z \in \mathbb{C}^{\ell}$ with $\text{Log} f(z) \in \mathbb{C}^{*n}$ we have

$$(f^*u)(z) \le c_u + H_S(f(z)) = c_u + \varphi_S(\log |f_1(z)|, \dots, \log |f_1(z)|)$$

$$\le c_u + \varphi_S(c_1 + \varphi_{S_1}(\operatorname{Log} z), \dots, c_n + \varphi_{S_n}(\operatorname{Log} z))$$

$$\le c_u + \varphi_S(c) + \varphi_S(\varphi_{S_1}(\operatorname{Log} z), \dots, \varphi_{S_n}(\operatorname{Log} z))$$

$$= c_u + \varphi_S(c) + H_{S'}(z)$$

and conclude that $f^*u \in \mathcal{L}^{S'}(\mathbb{C}^{\ell})$. If $u \leq q$ on K, then $f^*u \leq f^*q$ on $f^{-1}(K)$, we have $f^*(V_{K,q}^S) \leq V_{f^{-1}(K),f^*q}^{S'}$ and that equality holds if $f^* \colon \mathcal{L}^S(\mathbb{C}^n) \to \mathcal{L}^{S'}(\mathbb{C}^{\ell})$ is surjective. When $\ell = n$ we have the following weighted transformation rule, which generalizes

When $\ell = n$ we have the following weighted transformation rule, which generalizes Klimek [19, Theorem 5.3.1] and Perera [25, Theorem 1]:

Proposition 8.2 If $f : \mathbb{C}^n \to \mathbb{C}^n$ is proper, the following are equivalent:

- (i) The difference $f^*H_S H_{S'}$ is bounded, that is $f^*H_S \in \mathcal{L}^{S'}_+(\mathbb{C}^n)$.
- (ii) For every compact set $K \subset \mathbb{C}^n$ and every admissible weight q on K we have $f^*V^S_{K,q} = V^{S'}_{f^{-1}(K), f^*q}$.

Proof: (i) \Rightarrow (ii): Let $u \in \mathcal{L}^{S'}(\mathbb{C}^n)$ with $u|_{f^{-1}(K)} \leq f^*q$. By Klimek [19, Theorem 2.9.26] the function $v(z) = \max\{u(w) : w \in f^{-1}(z)\}$ is plurisubharmonic on \mathbb{C}^n . Let c be a constant such that $f^*H_S - H_{S'} \geq -c$. Then $v|_K \leq q$ and

$$v(z) \le \max_{w \in f^{-1}(z)} H_{S'}(w) + c_u \le \max_{w \in f^{-1}(z)} f^* H_S(w) + c + c_u = H_S(z) + c + c_u.$$

It follows that $u \leq f^* v \leq f^* V^S_{K,q}$. (ii) \Rightarrow (i): Let $c = \max_{w \in f^{-1}(\overline{\mathbb{D}}^n)} H_{S'}(w)$ and $c' = \max_{w \in f(\overline{\mathbb{D}}^n)} H_S(w)$. Then

$$H_{S'} - c \le V_{f^{-1}(\overline{\mathbb{D}}^n)}^{S'} = f^* V_{\overline{\mathbb{D}}^n}^S = f^* H_S$$

and

(8.6)
$$H_{S'} = V_{\mathbb{D}^n}^{S'} \ge V_{f^{-1}(f(\overline{\mathbb{D}^n}))}^{S'} = f^* V_{f(\overline{\mathbb{D}^n})}^S \ge f^* H_S - c'.$$

References

- T. BAYRAKTAR, Zero distribution of random sparse polynomials, arXiv:1503.00630v4, (2015), pp. 1–28.
- [2] —, Equidistribution of zeros of random holomorphic sections, Indiana Univ. Math. J., 65 (2016), pp. 1759–1793.
- [3] —, Asymptotic normality of linear statistics of zeros of random polynomials, Proc. Amer. Math. Soc., 145 (2017), pp. 2917–2929.
- [4] , Zero distribution of random sparse polynomials, Michigan Math. J., 66 (2017), pp. 389–419.
- [5] , Mass equidistribution for random polynomials, Potential Anal., 53 (2020), pp. 1403– 1421.
- [6] T. BAYRAKTAR, T. BLOOM, AND N. LEVENBERG, Pluripotential theory and convex bodies, Mat. Sb., 209 (2018), pp. 67–101.
- [7] T. BAYRAKTAR, T. BLOOM, N. LEVENBERG, AND C. H. LU, *Pluripotential theory and convex bodies: large deviation principle*, Ark. Mat., 57 (2019), pp. 247–283.
- [8] T. BAYRAKTAR, S. HUSSUNG, N. LEVENBERG, AND M. PERERA, *Pluripotential theory and convex bodies: a Siciak-Zaharjuta theorem*, Comput. Methods Funct. Theory, 20 (2020), pp. 571–590.

- [9] Z. BLOCKI, A note on maximal plurisubharmonic functions, Uzbek. Mat. Zh., (2009), pp. 28–32.
- [10] T. BLOOM AND N. LEVENBERG, Weighted pluripotential theory in \mathbb{C}^N , Amer. J. Math., 125 (2003), pp. 57–103.
- [11] —, Random polynomials and pluripotential-theoretic extremal functions, Potential Anal., 42 (2015), pp. 311–334.
- [12] T. BLOOM AND B. SHIFFMAN, Zeros of random polynomials on C^m, Math. Res. Lett., 14 (2007), pp. 469–479.
- [13] L. BOS AND N. LEVENBERG, Bernstein-Walsh theory associated to convex bodies and applications to multivariate approximation theory, Comput. Methods Funct. Theory, 18 (2018), pp. 361–388.
- [14] A. FIGALLI, The Monge-Ampère equation and its applications, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2017.
- [15] L. HÖRMANDER, The analysis of linear partial differential operators I and II, Springer Verlag, New York Berlin Heidelberg, 1983.
- [16] —, An introduction to complex analysis in several variables, North-Holland Publishing Co., Amsterdam, 3rd ed., 1990.
- [17] —, Notions of convexity, vol. 127 of Progress in Mathematics, Birkhäuser Boston Inc., Boston, MA, 1994.
- [18] L. HÖRMANDER AND R. SIGURDSSON, Growth properties of plurisubharmonic functions related to Fourier-Laplace transforms, J. Geom. Anal., 8 (1998), pp. 251–311.
- [19] M. KLIMEK, *Pluripotential theory*, vol. 6 of London Mathematical Society Monographs. New Series, The Clarendon Press Oxford University Press, New York, 1991.
- [20] N. LEVENBERG, Approximation in \mathbb{C}^N , Surv. Approx. Theory, 2 (2006), pp. 92–140.
- [21] N. LEVENBERG AND M. PERERA, A global domination principle for P-pluripotential theory, in Complex analysis and spectral theory, vol. 743 of Contemp. Math., Amer. Math. Soc., Providence, RI, 2020, pp. 11–19.
- [22] B. S. MAGNÚSSON, Á. E. SIGURÐARDÓTTIR, AND R. SIGURÐSSON, Polynomials with exponents in compact convex sets and associated weighted extremal functions: The Siciak-Zakharyuta theorem, Complex Analysis and its Synergies, 10 (2024).
- [23] B. S. MAGNÚSSON, R. SIGURÐSSON, AND B. SNORRASON, Polynomials with exponents in compact convex sets and associated weighted extremal functions – The Bernstein-Walsh-Siciak theorem, arXiv:2306.02486, (2023).
- [24] Q. D. NGUYEN AND T. V. LONG, Product property of global P-extremal functions, Math. Scand., 127 (2021), pp. 509–520.
- [25] M. PERERA, A transformation rule associated with P-extremal functions and holomorphic mappings., J.Anal., 31 (2023), pp. 2097–2113.

- [26] A. RASHKOVSKII, Newton numbers and residual measures of plurisubharmonic functions, Ann. Polon. Math., 75 (2000), pp. 213–231.
- [27] —, Indicators for plurisubharmonic functions of logarithmic growth, Indiana Univ. Math. J., 50 (2001), pp. 1433–1446.
- [28] E. B. SAFF AND V. TOTIK, Logarithmic potentials with external fields, vol. 316 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Springer-Verlag, Berlin, 1997. Appendix B by Thomas Bloom.
- [29] B. SHIFFMAN AND S. ZELDITCH, Random polynomials with prescribed Newton polytope, J. Amer. Math. Soc., 17 (2004), pp. 49–108.
- [30] J. SICIAK, On some extremal functions and their applications in the theory of analytic functions of several complex variables, Trans. Amer. Math. Soc., 105 (1962), pp. 322–357.
- [31] —, Extremal plurisubharmonic functions in \mathbf{C}^n , Ann. Polon. Math., 39 (1981), pp. 175–211.
- [32] —, Extremal plurisubharmonic functions and capacities in \mathbb{C}^n , Sophia Kokyuroku in Mathematics, 14 (1982).
- [33] ____, On series of homogeneous polynomials and their partial sums, Ann. Polon. Math., 51 (1990), pp. 289–302.
- [34] —, A remark on Tchebysheff polynomials in \mathbb{C}^N , Univ. Iagel. Acta Math., (1997), pp. 37–45.
- [35] R. SIGURDSSON, Growth properties of analytic and plurisubharmonic functions of finite order, Math. Scand., 59 (1986), pp. 235–304.
- [36] B. SNORRASON, Polynomials with exponents in compact convex sets and associated weighted extremal functions – Generalized product property, to appear in Mathematica Scandinavica, (2024).
- [37] M. TSUJI, Potential theory in modern function theory, Chelsea Publishing Co., New York, 1975. Reprinting of the 1959 original.
- [38] V. P. ZAKHARYUTA, Extremal plurisubharmonic functions, orthogonal polynomials, and the Bernštein-Walsh theorem for functions of several complex variables, Ann. Polon. Math., 33 (1976/77), pp. 137–148.