Energy balance for fractional anti-Zener and Zener models in terms of relaxation modulus and creep compliance

Slađan Jelić¹ Dušan Zorica^{2 3 *}

May 11, 2023

Abstract

Relaxation modulus and creep compliance corresponding to fractional anti-Zener and Zener models are calculated and restrictions on model parameters narrowing thermodynamical constraints are posed in order to ensure relaxation modulus and creep compliance to be completely monotone and Bernstein function respectively, that a priori guarantee the positivity of stored energy and dissipated power per unit volume, derived in time domain by considering the power per unit volume. Both relaxation modulus and creep compliance for model parameters obeying thermodynamical constraints, proved that can also be oscillatory functions with decreasing amplitude. Model used in numerical examples of relaxation modulus and creep compliance is also analyzed for the asymptotic behavior near the initial time instant and for large time. **Key words:** thermodynamically consistent fractional anti-Zener and Zener models, energy balance properties in time domain, stored energy and dissipated power per unit volume, relaxation modulus and creep compliance, completely monotonic and Bernstein functions

1 Introduction

Fractional anti-Zener and Zener models, that are the subject of the energy balance analysis in time domain, are formulated in [33] by considering the rheological schemes corresponding to the classical anti-Zener and Zener models having the classical spring replaced by the fractional element whose stress-strain relation is assumed in terms of fractional integral, as well as having the classical dash-pot replaced by the Scott-Blair element, i.e., element whose stress-strain relation is assumed in terms of fractional derivative. Note, the classical anti-Zener model is also referred to as the Jeffrey model, as discussed in [52] reviewing classical and fractional models of viscoelasticity. Some of the first to use the Scott-Blair element in posing the constitutive equations using classical rheological schemes are [51], while some of the most widely used fractional order models of viscoelastic body derived using the rheological schemes containing Scott-Blair element are reviewed in [36, 50]. Rheological scheme of the Burgers model with the classical dash-pot replaced by the Scott-Blair element is adopted in [40], in order to fractionalize the classical Burgers model. Infinite number of springs and dash-pots in the rheological scheme is used in [18, 23, 24] in order to formulate the Bessel model, while the setting of fractal rheological models is used in [29].

The analysis of thermodynamical consistency of constitutive models in the steady state regime, i.e., by requesting non-negativity of the storage and loss modulus for any frequency, is given in [10]. This method is used in [33] for formulation of thermodynamical restrictions on model parameters appearing in the fractional anti-Zener and Zener models, as well as in [40] and [12] for the fractional Burgers model, where the fractionalization is performed by replacing the classical dash-pot element with the fractional one in the former, as well as by replacing the ordinary derivatives with the fractional ones in the classical Burgers model in the latter one. Constitutive models of viscoelastic body having the orders of fractional derivatives not exceeding the first order are examined for thermodynamical consistency in [5]. Considering time domain, the dissipation inequality is also used in order to pose thermodynamical restrictions on model parameters, see for example [3, 4], while the energy balance properties of the fractional wave equations are examined in [53]. Dissipation of energy in viscoelastic media of Bessel type, the wave equation thermodynamical consistency and the requirements guaranteeing the complete monotonicity, as well as the analysis of the Zener wave equation are presented in [19, 22, 31, 39]. The extensive account on thermodynamics of hereditary materials can be found in [2].

 $^{^1}$ Department of Physics, Faculty of Sciences, University of Novi Sad, Trg D. Obradovića 4, 21000 Novi Sad, Serbia, sladjan.jelic@df.uns.ac.rs

² Department of Physics, Faculty of Sciences, University of Novi Sad, Trg D. Obradovića 4, 21000 Novi Sad, Serbia, dusan.zorica@df.uns.ac.rs

 $^{^3}$ Mathematical Institute, Serbian Academy of Arts and Sciences, Kneza Mihaila 36, 11000 Belgrade, Serbia, dusan_zorica@mi.sanu.ac.rs

^{*} Corresponding author

A priori energy balance properties of fractional anti-Zener and Zener models, that are formulated in [33] and analyzed for thermodynamical consistency in the steady-state regime, are discussed in time domain and it is proved that the power per unit volume consists of two terms: the time derivative of the energy per unit volume, stored in the viscoelastic body and dissipated power per unit volume. The stored energy per unit volume, expressed through strain, consists of the instantaneous term, that is of the same form as the potential energy of elastic body in which the relaxation modulus plays the role of the Young modulus, as well as of the hereditary type term, where the time derivative of the relaxation modulus represents the memory kernel of the difference of current strain and strain in the previous time instant. Dissipated power per unit volume, expressed in terms of strain, also consists of the instantaneous and hereditary type terms, respectively containing the first and second time derivative of the relaxation modulus. The positivity of stored energy and dissipated power per unit volume is guaranteed if the relaxation modulus is a positive monotonically decreasing and convex function, or moreover a completely monotone function. Note, the relaxation modulus is a function representing the time evolution of stress in a stress relaxation experiment, occurring as a consequence of the sudden and later constant strain, i.e., strain prescribed as the Heaviside step function.

The stored energy per unit volume can also be expressed through stress, when there is only the hereditary type term, having the time derivative of the creep compliance as the memory kernel multiplying the time evolution of stress, while the dissipated power per unit volume consists of the instantaneous and hereditary type terms, respectively containing the first and second time derivative of the creep compliance. Again, the positivity of stored energy and dissipated power per unit volume is guaranteed if the creep compliance is a positive monotonically increasing and concave function, or moreover a Bernstein function. Note, the creep compliance is a function representing the time evolution of strain in a creep experiment, occurring as a consequence of the sudden and later constant stress, i.e., stress prescribed as the Heaviside step function.

The important property of relaxation modulus being completely monotonic and creep compliance being a Bernstein function is discussed in [36]. On the other hand, the complete monotonicity of relaxation modus in the case of distributed-order fractional Zener model is studied in [11], while [37] deals with the fractional Burgers model in creep and stress relaxation tests. The relaxation modulus and creep compliance corresponding to the eight thermodynamically consistent fractional Burgers models formulated in [40] are examined in [41], where the thermodynamical requirements are narrowed in order to ensure the complete monotonicity of the relaxation modulus and to ensure that creep compliance is a Bernstein function. The role of Fox function is underlined in [25, 26] in seeking the responses of fractional constitutive models if stress or strain is assumed to be given. The experimental data, obtained in creep and stress relaxation experiments on biological tissues, is used for curve fitting in [21, 28]. In [38], the creep compliances in the steady state regime are reviewed in the case of integer-order models of viscoelasticity. Considering the inertial effects of the one-dimensional viscoelastic body, the displacement and stress in stress relaxation and creep tests are analyzed in [8, 9] when material is described by the distributed-order fractional model, while in [32] the thermodynamically consistent fractional Burgers models are used.

The qualitative characteristics of relaxation modulus and creep compliance play a crucial role in energy balance properties of the viscoelastic body, and therefore their explicit form is calculated and used in order to pose the conditions guaranteeing that the relaxation modulus is a completely monotone function and that creep compliance is a Bernstein function. Previously mentioned conditions prove to narrow the thermodynamical restrictions on model parameters appearing in the fractional anti-Zener and Zener models, that are obtained in [33]. The thermodynamical conditions actually allow for the relaxation modulus and creep compliance to be nonmonotonic and even oscillatory functions with exponentially decreasing amplitude, which does not necessarily violate the positivity of stored energy and dissipated power per unit volume.

Fractional anti-Zener and Zener constitutive equations, listed in Appendix A, take the following form

$$\tilde{\sigma}\left(s\right) = s^{\xi} \frac{\phi_{\varepsilon}\left(s\right)}{\phi_{\sigma}\left(s\right)} \tilde{\varepsilon}\left(s\right) \tag{1}$$

after the Laplace transform, defined as

$$\hat{f}(s) = \mathcal{L}[f(t)](s) = \int_0^\infty f(t) e^{-st} dt$$
, for $\operatorname{Re} s > 0$,

is applied, where ϕ_{ε} and ϕ_{σ} are model dependent constitutive functions in the Laplace domain, that are, along with the model dependent constitutive parameter ξ , listed in Table 1.

	sləbom əirtəmmyS						sləbom əirtəmmyzA								
Function ϕ_{σ}	$a_1+a_2s^{lpha+eta}$		$a_1 + a_2 s^{\lambda} + a_3 s^{\alpha+\gamma}$		$a_1 + a_2 s^{\frac{1+\alpha+\gamma}{2}} + a_3 s^{1+\alpha+\gamma}$			$a_1 + a_2 s^{\lambda} + a_3 s^{\kappa}$		(8 + ~)6 · · 8 + ~ ·	$a_1 + a_2 s^{\alpha+\beta} + a_3 s^{2(\alpha+\beta)}$		$a_1+a_2s^{lpha+eta}$		
Function ϕ_{ε}	8+2 	$b_1 + b_2 s^{lpha + eta}$		$b_1 + b_2 s^\lambda + b_3 s^{lpha + \gamma}$		$b_1 + b_2 s^{\frac{1+\alpha+\gamma}{2}} + b_3 s^{1+\alpha+\gamma}$		$b_1+b_2s^{lpha+eta}$			$b_1 + b_2 s^{\lambda} + b_3 s^{\kappa}$		$b_1 + b_2 s^{\alpha+\beta} + b_3 s^{2(\alpha+\beta)}$		
Model	ID.ID	ID.DD+	IID.IID	IDD.IDD	IID.IDD	I+ID.I+ID	IDD+.IDD+	I+ID.IDD+	IID.ID	IDD.DD+	I+ID.ID	IDD+.DD+	ID.IDD	ID.DDD+	ID.IDD ⁺
0	~	σ	<i>u</i> ×	× C	0	0	ı	1 - 1	7	α –	β +	σ	0	α –	1
rder ξ	$\pi - \chi$	$\vdash \mu < 1$	$\lambda - l$	$\eta - \chi$	$\eta - \chi$	$\eta - \chi$	$\lambda - l$	$(\lambda - \lambda)$	$\lambda - \delta$	$\vdash \mu < 1$	$+ \nu < 1$	$+\mu < 1$	$\eta - \chi$	$\vdash \mu < 1$	$\gamma - \beta$
Order λ	I	I	lpha-eta	$\alpha+\beta<1$	$\mu+\nu<1$	I	I	I	$(\alpha+\beta)-(\nu+\gamma)$	$\alpha+\beta<1$	I	1	$\mu+\nu<1$	$\eta - \eta$	I
Order κ	Ι	Ι	-	-	lpha-eta	Ι	I	Ι	$\alpha+\beta<1$	$\alpha+\mu<1$	Η	I	$\alpha+\beta<1$	$\alpha + \nu - (\mu - \beta)$	I

Table 1: Summary of constitutive functions ϕ_{σ} and ϕ_{ε} , along with the order ξ , corresponding to the thermodynamically consistent fractional anti-Zener and Zener models. The notation \ast means that the orders $\alpha + \beta$ and $\alpha + \gamma$ belong either to the interval (0, 1) or interval (1, 2).

Constitutive model in the Laplace domain (1) is transformed into the relaxation modulus in the Laplace domain

$$\tilde{\sigma}_{sr}\left(s\right) = \frac{1}{s^{1-\xi}} \frac{\phi_{\varepsilon}\left(s\right)}{\phi_{\sigma}\left(s\right)},\tag{2}$$

when the strain is prescribed in the form of Heaviside step function, i.e., as $\varepsilon = H$, while if the stress is prescribed in the form of Heaviside step function, i.e., as $\sigma = H$, then the constitutive model (1) becomes the creep compliance in the Laplace domain, expressed as

$$\tilde{\varepsilon}_{cr}\left(s\right) = \frac{1}{s^{1+\xi}} \frac{\phi_{\sigma}\left(s\right)}{\phi_{\varepsilon}\left(s\right)}.$$
(3)

The use of constitutive models in modeling wave propagation is extensive, where some of the first studies of viscoelastic materials of fractional order is performed in [16, 17]. Damped oscillations and wave propagation in viscoelastic materials modeled by the Zener, modified Zener, and modified Maxwell models is conducted in [44, 45, 46, 47, 48]. Waves in fractional Zener and distributed type media are investigated in [13, 34] and in [14, 35], while in [43] wave propagation in the fractional Burgers type material is considered. Multidimensional generalizations of the fractional Zener model and its application in wave propagation is studied in [1, 20, 42]. Fractional order constitutive equations of viscoelastic media, along with the wave propagation, dispersion, and attenuation processes are reviewed in [6, 7, 30, 36, 49], while [15] surveys the propagation of acoustic waves in complex media.

2 Models' dissipativity properties in the time domain

In order to investigate energy balance properties of the one-dimensional viscoelastic body, the constitutive equation in the Laplace domain (1) is rewritten either as

$$\tilde{\sigma}(s) = s\tilde{\sigma}_{sr}(s)\tilde{\varepsilon}(s), \quad \text{or as} \quad \tilde{\varepsilon}(s) = s\tilde{\varepsilon}_{cr}(s)\tilde{\sigma}(s),$$
(4)

using the relaxation modulus (2) and creep compliance (3) in the Laplace domain, so that, after performing the inverse Laplace transform in (4), the constitutive equation in time domain reads either

$$\sigma(t) = \frac{\mathrm{d}}{\mathrm{d}t} \left(\sigma_{sr}(t) * \varepsilon(t) \right), \quad \text{or} \quad \varepsilon(t) = \dot{\varepsilon}_{cr}(t) * \sigma(t), \tag{5}$$

since $(\sigma_{sr}(t) * \varepsilon(t))|_{t=0} = 0$ and $\varepsilon_{cr}^{(g)} = \varepsilon_{cr}(0) = 0$. The former holds true if the relaxation modulus tends to infinity as a power-type function for small time, i.e., if $\sigma_{sr}(t) \sim Kt^{-\alpha}$, as $t \to 0$ with $\alpha \in (0, 1)$, as well as if strain is bounded at zero by ε_0 , since then $\sigma_{sr}(t) * \varepsilon(t) = \int_0^t \sigma_{sr}(t') \varepsilon(t-t') dt' \sim K\varepsilon_0 \int_0^t (t')^{-\alpha} dt' = K\varepsilon_0 \frac{t^{1-\alpha}}{1-\alpha} \to 0$ as $t \to 0$, while the latter holds true, since the glass compliance $\varepsilon_{cr}^{(g)}$ is zero in $\mathcal{L}^{-1}[s\tilde{\varepsilon}_{cr}(s)] = \dot{\varepsilon}_{cr}(t) + \varepsilon_{cr}^{(g)}\delta(t)$ for all considered models.

The power per unit volume, expressed by

$$P(t) = \sigma(t)\dot{\varepsilon}(t), \qquad (6)$$

can be written either in terms of strain as

$$P(t) = \frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{1}{2} \sigma_{sr}(t) \varepsilon^{2}(t) + \frac{1}{2} \int_{0}^{t} \left(-\dot{\sigma}_{sr}(t-t') \right) \left(\varepsilon(t) - \varepsilon(t') \right)^{2} \mathrm{d}t' \right) + \frac{1}{2} \left(-\dot{\sigma}_{sr}(t) \right) \varepsilon^{2}(t) + \frac{1}{2} \int_{0}^{t} \ddot{\sigma}_{sr}(t-t') \left(\varepsilon(t) - \varepsilon(t') \right)^{2} \mathrm{d}t',$$
(7)

using $(5)_1$, or in terms of stress

$$P(t) = \frac{d}{dt} \left(\frac{1}{2} \int_0^t \dot{\varepsilon}_{cr} (t - t') \,\sigma^2(t') \,dt' \right) + \frac{1}{2} \dot{\varepsilon}_{cr} (t) \,\sigma^2(t) + \frac{1}{2} \int_0^t \left(-\ddot{\varepsilon}_{cr} (t) (t - t') \right) \left(\sigma(t) - \sigma(t') \right)^2 dt',$$
(8)

using $(5)_2$, where the terms

$$W(t) = \frac{1}{2}\sigma_{sr}(t)\varepsilon^{2}(t) + \frac{1}{2}\int_{0}^{t} \left(-\dot{\sigma}_{sr}(t-t')\right)\left(\varepsilon(t) - \varepsilon(t')\right)^{2} \mathrm{d}t' > 0 \quad \text{and} \tag{9}$$

$$W(t) = \frac{1}{2} \int_0^t \dot{\varepsilon}_{cr} (t - t') \,\sigma^2(t') \,\mathrm{d}t' > 0, \tag{10}$$

occurring in (7) and (8) respectively, can be interpreted as the energy per unit volume stored in the viscoelastic body, taking into account the memory of strain, respectively stress, weighted by the derivative of relaxation modulus, respectively creep compliance, carrying the information about material properties, while the first term in (9) resembles to the potential energy of the elastic body with Young's modulus replaced by the relaxation modulus. On the other hand, the terms

$$\mathcal{P}(t) = \frac{1}{2} \left(-\dot{\sigma}_{sr}(t) \right) \varepsilon^2(t) + \frac{1}{2} \int_0^t \ddot{\sigma}_{sr}(t-t') \left(\varepsilon(t) - \varepsilon(t') \right)^2 \mathrm{d}t' > 0 \quad \text{and} \tag{11}$$

$$\mathcal{P}(t) = \frac{1}{2}\dot{\varepsilon}_{cr}(t)\,\sigma^2(t) + \frac{1}{2}\int_0^t \left(-\ddot{\varepsilon}_{cr}(t)\,(t-t')\right)\left(\sigma\left(t\right) - \sigma\left(t'\right)\right)^2\mathrm{d}t' > 0,\tag{12}$$

occurring in (7) and (8) respectively, can be interpreted as the dissipated power per unit volume having two types of contributions: the first one being instantaneous and depending on a material dependent positive and decreasing function $-\dot{\sigma}_{sr}$, respectively $\dot{\varepsilon}_{cr}$, and the second contribution being of hereditary type with material dependent kernel. The positivity of energy and dissipativity of power is guaranteed by the properties of the relaxation modulus and creep compliance, namely by requesting the relaxation modulus to be completely monotonic function, i.e.,

$$\sigma_{sr}\left(t\right) \geqslant 0 \quad \text{and} \quad \left(-1\right)^{k} \frac{\mathrm{d}^{k}}{\mathrm{d}t^{k}} \dot{\sigma}_{sr}\left(t\right) \leqslant 0, \quad \text{for} \quad t > 0, \quad k \in \mathbb{N}_{0},$$

as well as by requesting the creep compliance to be Bernstein's function, i.e., a positive function with completely monotonic first derivative.

Introducing the energy and dissipated power per unit volume, the power per unit volume can be rewritten as

$$P(t) = \frac{\mathrm{d}}{\mathrm{d}t}W(t) + \mathcal{P}(t),$$

where the first term corresponds to the elastic type properties of viscoelastic body, while the second term corresponds to its properties of viscous type.

The expression (7) for the power per unit volume follows from

$$P(t) = \frac{\mathrm{d}}{\mathrm{d}t} \left(\sigma_{sr}(t) * \varepsilon(t) \right) \dot{\varepsilon}(t) ,$$

obtained by (6) and $(5)_1$, transforming into

$$P(t) = \frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\mathrm{d}}{\mathrm{d}t} \left(\sigma_{sr} \left(t \right) * \varepsilon \left(t \right) \right) \varepsilon \left(t \right) \right) - \frac{\mathrm{d}}{\mathrm{d}t} \left(\dot{\sigma}_{sr} \left(t \right) * \varepsilon \left(t \right) + \sigma_{sr}^{(g)} \varepsilon \left(t \right) \right) \varepsilon \left(t \right)$$
$$= \frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\mathrm{d}}{\mathrm{d}t} \left(\sigma_{sr} \left(t \right) * \varepsilon \left(t \right) \right) \varepsilon \left(t \right) \right) - \frac{\mathrm{d}}{\mathrm{d}t} \left(\dot{\sigma}_{sr} \left(t \right) * \varepsilon \left(t \right) \right) \varepsilon \left(t \right) - \frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{1}{2} \sigma_{sr}^{(g)} \varepsilon^{2} \left(t \right) \right),$$
(13)

using the derivative of a product of two functions and a derivative of a convolution

$$\frac{\mathrm{d}}{\mathrm{d}t}\left(f\left(t\right)\ast g\left(t\right)\right) = \dot{f}\left(t\right)\ast g\left(t\right) + f\left(0\right)g\left(t\right),$$

since the first two terms in (13) become

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} \left(\sigma_{sr}\left(t\right) \ast \varepsilon\left(t\right)\right) \varepsilon\left(t\right) \\ &= \frac{1}{2}\sigma_{sr}\left(t\right)\varepsilon^{2}\left(t\right) + \frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\left(\sigma_{sr}\left(t\right) \ast \varepsilon^{2}\left(t\right)\right) - \frac{1}{2}\int_{0}^{t} \dot{\sigma}_{sr}\left(t-t'\right)\left(\varepsilon\left(t\right) - \varepsilon\left(t'\right)\right)^{2} \mathrm{d}t' \\ &= \frac{1}{2}\sigma_{sr}\left(t\right)\varepsilon^{2}\left(t\right) + \frac{1}{2}\dot{\sigma}_{sr}\left(t\right) \ast \varepsilon^{2}\left(t\right) + \frac{1}{2}\sigma_{sr}^{\left(g\right)}\varepsilon^{2}\left(t\right) - \frac{1}{2}\int_{0}^{t} \dot{\sigma}_{sr}\left(t-t'\right)\left(\varepsilon\left(t\right) - \varepsilon\left(t'\right)\right)^{2} \mathrm{d}t' \end{aligned}$$

and

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\dot{\sigma}_{sr} \left(t \right) \ast \varepsilon \left(t \right) \right) \varepsilon \left(t \right)$$

= $\frac{1}{2} \dot{\sigma}_{sr} \left(t \right) \varepsilon^{2} \left(t \right) + \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \left(\dot{\sigma}_{sr} \left(t \right) \ast \varepsilon^{2} \left(t \right) \right) - \frac{1}{2} \int_{0}^{t} \ddot{\sigma}_{sr} \left(t - t' \right) \left(\varepsilon \left(t \right) - \varepsilon \left(t' \right) \right)^{2} \mathrm{d}t',$

according to

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(k\left(t\right) * u\left(t\right)\right) u\left(t\right) = \frac{1}{2}k\left(t\right)u^{2}\left(t\right) + \frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t} \left(k\left(t\right) * u^{2}\left(t\right)\right) - \frac{1}{2}\int_{0}^{t}\dot{k}\left(t - t'\right)\left(u\left(t\right) - u\left(t'\right)\right)^{2}\mathrm{d}t'$$
(14)

and a derivative of a convolution.

On the other hand, the expression (8) for the power per unit volume follows from

$$P(t) = \frac{\mathrm{d}}{\mathrm{d}t} \left(\dot{\varepsilon}_{cr} \left(t \right) * \sigma \left(t \right) \right) \sigma \left(t \right),$$

obtained by (6) and $(5)_2$, transforming into

$$P(t) = \frac{1}{2}\dot{\varepsilon}_{cr}(t)\,\sigma^{2}(t) + \frac{1}{2}\frac{d}{dt}\left(\dot{\varepsilon}_{cr}(t)*\sigma^{2}(t)\right) - \frac{1}{2}\int_{0}^{t}\ddot{\varepsilon}_{cr}(t-t')\left(\sigma(t) - \sigma(t')\right)^{2}dt',$$

according to (14).

3 Stress relaxation and creep

Relaxation modulus and creep compliance play a crucial role in energy balance properties of viscoelastic body, since being a completely monotonic function and a Bernstein function respectively, they guarantee the positivity of the energy stored in viscoelastic body, see (9) and (10), as well as the positivity of the dissipated power, see (11) and (12). Therefore, the relaxation modulus and creep compliance, given by (2) and (3) in the Laplace domain, are calculated in the time domain and their properties are investigated, yielding that the relaxation modulus and creep compliance are completely monotonic and Bernstein functions respectively, if the model parameters satisfy narrowed thermodynamical restrictions in addition to request that the relaxation modulus and creep compliance in the Laplace domain (2) and (3) do not have poles, while the character of relaxation modulus and creep compliance in time domain changes if their counterparts in the Laplace domain have poles. Namely, in the case of negative real pole there exists an additional term decaying exponentially in time, while if there exists a pair of complex conjugated poles with negative real part, then there is an additional term displaying damped oscillatory character.

3.1Relaxation modulus and creep compliance

3.1.1**Relaxation** modulus

The relaxation modulus is obtained as

$$\sigma_{sr}(t) = \sigma_{sr}^{(\text{NP})}(t) + \begin{cases} 0, & \text{if } \tilde{\sigma}_{sr} \text{ has no poles,} \\ \sigma_{sr}^{(\text{RP})}(t), & \text{if } \tilde{\sigma}_{sr} \text{ has a negative real pole,} \\ \sigma_{sr}^{(\text{CCP})}(t), & \text{if } \tilde{\sigma}_{sr} \text{ has a pair of complex conjugated poles,} \end{cases}$$
(15)

by inverting the Laplace transform in the relaxation modulus in the Laplace domain $\tilde{\sigma}_{sr}$, given by (2), using the definition and integration in the complex plane along the Bromwich contour, with the functions $\sigma_{sr}^{(NP)}$, $\sigma_{sr}^{(RP)}$, and $\sigma_{sr}^{(\text{CCP})}$ respectively taking the forms

$$\sigma_{sr}^{(\rm NP)}(t) = \frac{1}{\pi} \int_0^\infty \frac{1}{\rho^{1-\xi}} \frac{\left|\phi_{\varepsilon}\left(\rho e^{i\pi}\right)\right|}{\left|\phi_{\sigma}\left(\rho e^{i\pi}\right)\right|} \sin\left(\arg\phi_{\varepsilon}\left(\rho e^{i\pi}\right) - \arg\phi_{\sigma}\left(\rho e^{i\pi}\right) + \xi\pi\right) e^{-\rho t} d\rho,\tag{16}$$

$$\sigma_{sr}^{(\text{RP})}(t) = -\frac{1}{\rho_{\text{RP}}^{1-\xi}} \frac{|\phi_{\varepsilon}(s_{\text{RP}})|}{|\phi_{\sigma}'(s_{\text{RP}})|} \cos\left(\arg\phi_{\varepsilon}(s_{\text{RP}}) - \arg\phi_{\sigma}'(s_{\text{RP}}) + \xi\pi\right) e^{-\rho_{\text{RP}}t},\tag{17}$$

$$\sigma_{sr}^{(\text{CCP})}(t) = 2 \frac{1}{\rho_{\text{CCP}}^{1-\xi}} \frac{|\phi_{\varepsilon}(s_{\text{CCP}})|}{|\phi_{\sigma}'(s_{\text{CCP}})|} e^{-|\text{Re}\,s_{\text{CCP}}|t} \cos\left(\text{Im}\,s_{\text{CCP}}t + \arg\phi_{\varepsilon}(s_{\text{CCP}}) - \arg\phi_{\sigma}'(s_{\text{CCP}}) - (1-\xi)\,\varphi_{\text{CCP}}\right), \quad (18)$$

where $\phi'_{\sigma}(s) = \frac{d}{ds}\phi_{\sigma}(s)$ and where poles of function $\tilde{\sigma}_{sr}$: $s_{\rm RP} = \rho_{\rm RP} e^{i\pi}$ and $s_{\rm CCP} = \rho_{\rm CCP} e^{i\varphi_{\rm CCP}}$ are respectively a negative real zero of function ϕ_{σ} and its complex zero having negative real part. The equivalent form of function $\sigma_{sr}^{(\rm NP)}$, given by (16), is

$$\sigma_{sr}^{(\text{NP})}(t) = \frac{1}{\pi} \int_0^\infty \frac{1}{\rho^{1-\xi}} \frac{K(\rho)}{\left|\phi_\sigma\left(\rho e^{i\pi}\right)\right|^2} e^{-\rho t} d\rho,$$
(19)

with

$$K(\rho) = \frac{1}{2i} \left(e^{i\xi\pi} \phi_{\varepsilon} \left(\rho e^{i\pi} \right) \bar{\phi}_{\sigma} \left(\rho e^{i\pi} \right) - e^{-i\xi\pi} \bar{\phi}_{\varepsilon} \left(\rho e^{i\pi} \right) \phi_{\sigma} \left(\rho e^{i\pi} \right) \right)$$

$$= \left| \phi_{\varepsilon} \left(\rho e^{i\pi} \right) \right| \left| \phi_{\sigma} \left(\rho e^{i\pi} \right) \right| \sin \left(\arg \phi_{\varepsilon} \left(\rho e^{i\pi} \right) - \arg \phi_{\sigma} \left(\rho e^{i\pi} \right) + \xi\pi \right)$$

$$= \cos \left(\xi\pi \right) \left(\operatorname{Im}\phi_{\varepsilon} \left(\rho e^{i\pi} \right) \operatorname{Re} \phi_{\sigma} \left(\rho e^{i\pi} \right) - \operatorname{Re} \phi_{\varepsilon} \left(\rho e^{i\pi} \right) \operatorname{Im} \phi_{\sigma} \left(\rho e^{i\pi} \right) \right)$$

$$+ \sin \left(\xi\pi \right) \left(\operatorname{Re}\phi_{\varepsilon} \left(\rho e^{i\pi} \right) \operatorname{Re} \phi_{\sigma} \left(\rho e^{i\pi} \right) + \operatorname{Im} \phi_{\varepsilon} \left(\rho e^{i\pi} \right) \operatorname{Im} \phi_{\sigma} \left(\rho e^{i\pi} \right) \right), \qquad (20)$$

being convenient for examining whether the function $\sigma_{sr}^{(NP)}$ is completely monotonic, that is guaranteed by requiring

$$K(\rho) \ge 0, \quad \text{i.e.,} \quad \sin\left(\arg\phi_{\varepsilon}\left(\rho e^{i\pi}\right) - \arg\phi_{\sigma}\left(\rho e^{i\pi}\right) + \xi\pi\right) \ge 0 \quad \text{for} \quad \rho \ge 0, \tag{21}$$

while if the condition (21) is not satisfied, then function $\sigma_{sr}^{(\text{NP})}$ can be non-monotonic. Function $\sigma_{sr}^{(\text{RP})}$ is either a positive exponentially decreasing function tending to zero or a negative exponentially increasing function also tending to zero, with time constant ρ_{RP} , obtained as a negative real pole of function $\tilde{\sigma}_{sr}$, see (17), while function $\sigma_{sr}^{(\text{CCP})}$ is an oscillatory function of an exponentially decreasing amplitude, having angular frequency defined by the imaginary part of a complex pole of function $\tilde{\sigma}_{sr}$, i.e., by $\text{Im } s_{\text{CCP}}$, and damping parameter defined by the real part of a complex pole of function $\tilde{\sigma}_{sr}$, i.e., by $|\text{Re } s_{\text{CCP}}|$, see (18).

If the function ϕ_{σ} , appearing in the relaxation modulus in the Laplace domain (2), consists of two terms, as it is the case for models ID.ID, ID.DD⁺, ID.IDD, ID.DDD⁺, and ID.IDD⁺, see Table 1, then the relaxation modulus can be written in terms of the two-parameter Mittag-Leffler function $e_{\xi,\zeta,\lambda}$ as

$$\sigma_{sr}(t) = \frac{b_1}{a_2} e_{\alpha+\beta,1-\xi+\alpha+\beta,\frac{a_1}{a_2}}(t) + \frac{b_2}{a_2} e_{\alpha+\beta,1-\xi+\alpha+\beta-\lambda,\frac{a_1}{a_2}}(t) + \frac{b_3}{a_2} e_{\alpha+\beta,1-\xi+\alpha+\beta-\kappa,\frac{a_1}{a_2}}(t),$$
(22)

where one takes $b_3 = 0$ for models ID.ID and ID.DD⁺, with the notation

$$e_{\xi,\zeta,\lambda}\left(t\right) = t^{\zeta-1} E_{\xi,\zeta}\left(-\lambda t^{\xi}\right), \quad \text{where} \quad E_{\xi,\zeta}\left(z\right) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma\left(\xi n + \zeta\right)}, \tag{23}$$

see [27]. Namely, the relaxation modulus in the Laplace domain (2) for the mentioned models can be written as

$$\tilde{\sigma}_{sr}\left(s\right) = \frac{1}{s^{1-\xi}} \frac{b_1 + b_2 s^{\lambda} + b_3 s^{\kappa}}{a_1 + a_2 s^{\alpha+\beta}},$$

see Table 1, transforming into

$$\tilde{\sigma}_{sr}\left(s\right) = \frac{b_1}{a_2} \frac{s^{-(1-\xi)}}{s^{\alpha+\beta} + \frac{a_1}{a_2}} + \frac{b_2}{a_2} \frac{s^{-(1-\xi-\lambda)}}{s^{\alpha+\beta} + \frac{a_1}{a_2}} + \frac{b_3}{a_2} \frac{s^{-(1-\xi-\kappa)}}{s^{\alpha+\beta} + \frac{a_1}{a_2}},$$

that yields the relaxation modulus in the form (22) by the Laplace transform of the two-parameter Mittag-Leffler function

$$e_{\xi,\zeta,\lambda}\left(t\right) = \mathcal{L}^{-1}\left[\frac{s^{\xi-\zeta}}{s^{\xi}+\lambda}\right]\left(t\right).$$
(24)

The integral representation of the relaxation modulus, already given by (19) in the case when $\tilde{\sigma}_{sr}$ has no poles, using the relaxation modulus (22) expressed in terms of the two-parameter Mittag-Leffler function (23), according to its integral representation

$$e_{\xi,\zeta,\lambda}\left(t\right) = \frac{1}{\pi} \int_0^\infty \frac{\lambda \sin\left(\left(\zeta - \xi\right)\pi\right) + \rho^{\xi} \sin\left(\zeta\pi\right)}{\left|\rho^{\xi} \mathrm{e}^{\mathrm{i}\xi\pi} + \lambda\right|^2} \rho^{\xi-\zeta} \mathrm{e}^{-\rho t} \mathrm{d}\rho,\tag{25}$$

see [27], provided that $\xi \in (0, 1)$ and $\zeta < 1 + \xi$, is reobtained in the form

$$\sigma_{sr}(t) = \frac{1}{a_2\pi} \int_0^\infty \frac{1}{\rho^{1-\xi}} \left(\frac{a_1}{a_2} \frac{b_1 \sin\left(\xi\pi\right) + b_2 \rho^\lambda \sin\left(\left(\xi + \lambda\right)\pi\right) + b_3 \rho^\kappa \sin\left(\left(\xi + \kappa\right)\pi\right)}{\left|\rho^{\alpha+\beta} \mathrm{e}^{\mathrm{i}(\alpha+\beta)\pi} + \frac{a_1}{a_2}\right|^2} + \rho^{\alpha+\beta} \frac{b_1 \sin\left(\left(\xi - \alpha - \beta\right)\pi\right) + b_2 \rho^\lambda \sin\left(\left(\xi + \lambda - \alpha - \beta\right)\pi\right) + b_3 \rho^\kappa \sin\left(\left(\xi + \kappa - \alpha - \beta\right)\pi\right)}{\left|\rho^{\alpha+\beta} \mathrm{e}^{\mathrm{i}(\alpha+\beta)\pi} + \frac{a_1}{a_2}\right|^2} \right) \mathrm{e}^{-\rho t} \mathrm{d}\rho,$$

$$(26)$$

so that its complete monotonicity, provided that $\alpha + \beta \in (0,1)$, is ensured by requiring the function

$$K(\rho) = a_1 b_1 \sin(\xi \pi) + a_1 b_2 \rho^{\lambda} \sin((\xi + \lambda) \pi) + a_1 b_3 \rho^{\kappa} \sin((\xi + \kappa) \pi) + a_2 b_1 \rho^{\alpha + \beta} \sin((\xi - \alpha - \beta) \pi) + a_2 b_2 \rho^{\alpha + \beta + \lambda} \sin((\xi + \lambda - \alpha - \beta) \pi) + a_2 b_3 \rho^{\alpha + \beta + \kappa} \sin((\xi + \kappa - \alpha - \beta) \pi)$$

to be non-negative for $\rho \ge 0$. Note, the forms of the relaxation modulus (19) and (26) coincide.

3.1.2 Creep compliance

The creep compliance is obtained as

$$\varepsilon_{cr}(t) = \varepsilon_{cr}^{(\text{NP})}(t) + \begin{cases} 0, & \text{if } \tilde{\varepsilon}_{cr} \text{ has no poles,} \\ \varepsilon_{cr}^{(\text{RP})}(t), & \text{if } \tilde{\varepsilon}_{cr} \text{ has a negative real pole,} \\ \varepsilon_{cr}^{(\text{CCP})}(t), & \text{if } \tilde{\varepsilon}_{cr} \text{ has a pair of complex conjugated poles,} \end{cases}$$
(27)

with the functions $\varepsilon_{cr}^{(\rm NP)}$, $\varepsilon_{cr}^{(\rm RP)}$, and $\varepsilon_{cr}^{(\rm CCP)}$ respectively taking the forms

$$\varepsilon_{cr}^{(\rm NP)}(t) = \frac{1}{\pi} \int_0^\infty \frac{1}{\rho^{1+\xi}} \frac{\left|\phi_\sigma\left(\rho e^{i\pi}\right)\right|}{\left|\phi_\varepsilon\left(\rho e^{i\pi}\right)\right|} \sin\left(\arg\phi_\varepsilon\left(\rho e^{i\pi}\right) - \arg\phi_\sigma\left(\rho e^{i\pi}\right) + \xi\pi\right) \left(1 - e^{-\rho t}\right) d\rho,\tag{28}$$

$$\varepsilon_{cr}^{(\mathrm{RP})}(t) = -\frac{1}{\rho_{\mathrm{RP}}^{1+\xi}} \frac{|\phi_{\sigma}(s_{\mathrm{RP}})|}{|\phi_{\varepsilon}'(s_{\mathrm{RP}})|} \cos\left(\arg\phi_{\varepsilon}'(s_{\mathrm{RP}}) - \arg\phi_{\sigma}(s_{\mathrm{RP}}) + \xi\pi\right) \left(1 - \mathrm{e}^{-\rho_{\mathrm{RP}}t}\right),\tag{29}$$
$$\varepsilon_{cr}^{(\mathrm{CCP})}(t) = 2\frac{1}{\rho_{\mathrm{RP}}^{1+\xi}} \frac{|\phi_{\sigma}(s_{\mathrm{CCP}})|}{|\phi_{\varepsilon}'(s_{\mathrm{CCP}})|}$$

$$\times \left(e^{-|\operatorname{Re} s_{\operatorname{CCP}}|t} \cos\left(\operatorname{Im} s_{\operatorname{CCP}}t - \arg \phi_{\varepsilon}'(s_{\operatorname{CCP}}) + \arg \phi_{\sigma}(s_{\operatorname{CCP}}) - (1+\xi)\varphi_{\operatorname{CCP}} \right) - \cos\left(\arg \phi_{\varepsilon}'(s_{\operatorname{CCP}}) - \arg \phi_{\sigma}(s_{\operatorname{CCP}}) + (1+\xi)\varphi_{\operatorname{CCP}} \right) \right),$$
(30)

where $\phi'_{\varepsilon}(s) = \frac{\mathrm{d}}{\mathrm{d}s}\phi_{\varepsilon}(s)$ and where $s_{\mathrm{RP}} = \rho_{\mathrm{RP}} e^{\mathrm{i}\pi}$ and $s_{\mathrm{CCP}} = \rho_{\mathrm{CCP}} e^{\mathrm{i}\varphi_{\mathrm{CCP}}}$ are respectively a negative real zero of function ϕ_{ε} and its complex zero having negative real part.

The equivalent form of function $\varepsilon_{cr}^{(NP)}$, given by (28), is

$$\varepsilon_{cr}^{(\text{NP})}(t) = \frac{1}{\pi} \int_0^\infty \frac{1}{\rho^{1+\xi}} \frac{K(\rho)}{\left|\phi_{\varepsilon}\left(\rho e^{i\pi}\right)\right|^2} \left(1 - e^{-\rho t}\right) d\rho,\tag{31}$$

with function K defined by (20), so that the conditions (21), which guaranteed that the relaxation modulus $\sigma_{sr}^{(\text{NP})}$, see (19), is completely monotonic, also guarantee that the creep compliance $\varepsilon_{cr}^{(\text{NP})}$ is a Bernstein function. On the other hand, if the condition (21) is not satisfied, then function $\varepsilon_{cr}^{(\text{NP})}$ can be non-monotonic. Function $\varepsilon_{cr}^{(\text{RP})}$ is a function starting from zero, that either exponentially increases to a positive horizontal asymptote, or exponentially decreases to a negative horizontal asymptote, having the time constant ρ_{RP} , obtained as a negative real pole of function $\tilde{\varepsilon}_{cr}$, see (29), while function $\varepsilon_{cr}^{(\text{CCP})}$ is a vertically shifted oscillatory function of an exponentially decreasing amplitude, having angular frequency defined by the imaginary part of a complex pole of function $\tilde{\varepsilon}_{cr}$, i.e., by Im s_{CCP} , and damping parameter defined by the real part of a complex pole of function $\tilde{\varepsilon}_{cr}$, i.e., by [Re s_{CCP}], see (30).

If the function ϕ_{ε} , appearing in the creep compliance in the Laplace domain (3), consists of two terms, as it is the case for models ID.ID, ID.DD⁺, IID.ID, IDD.DD⁺, I⁺ID.ID, and IDD⁺.DD⁺, see Table 1, then the creep compliance can be written in terms of the two-parameter Mittag-Leffler function $e_{\xi,\zeta,\lambda}$, given by (23), as

$$\varepsilon_{cr}(t) = \frac{a_1}{b_2} e_{\alpha+\beta,1+\xi+\alpha+\beta,\frac{b_1}{b_2}}(t) + \frac{a_2}{b_2} e_{\alpha+\beta,1+\xi+\alpha+\beta-\lambda,\frac{b_1}{b_2}}(t) + \frac{a_3}{b_2} e_{\alpha+\beta,1+\xi+\alpha+\beta-\kappa,\frac{b_1}{b_2}}(t), \qquad (32)$$

where one takes $a_3 = 0$ for models ID.ID and ID.DD⁺. Namely, the creep compliance in the Laplace domain (3) for the mentioned models can be written as

$$\tilde{\varepsilon}_{cr}\left(s\right) = \frac{1}{s^{1+\xi}} \frac{a_1 + a_2 s^{\lambda} + a_3 s^{\kappa}}{b_1 + b_2 s^{\alpha+\beta}},$$

see Table 1, transforming into

$$\tilde{\varepsilon}_{cr}\left(s\right) = \frac{a_1}{b_2} \frac{s^{-(1+\xi)}}{s^{\alpha+\beta} + \frac{b_1}{b_2}} + \frac{a_2}{b_2} \frac{s^{-(1+\xi-\lambda)}}{s^{\alpha+\beta} + \frac{b_1}{b_2}} + \frac{a_3}{b_2} \frac{s^{-(1+\xi-\kappa)}}{s^{\alpha+\beta} + \frac{b_1}{b_2}},$$

that yields the creep compliance in the form (32) by the Laplace transform of the two-parameter Mittag-Leffler function, given by (24).

Note, the creep compliance written in terms of the two-parameter Mittag-Leffler function (32), does not admit the integral representation in the form (25), since the condition $\zeta < 1 + \xi$ is not satisfied for the first term (32). However, the derivative of creep compliance (32), obtained as

$$\dot{\varepsilon}_{cr}\left(t\right) = \frac{a_1}{b_2} e_{\alpha+\beta,\xi+\alpha+\beta,\frac{b_1}{b_2}}\left(t\right) + \frac{a_2}{b_2} e_{\alpha+\beta,\xi+\alpha+\beta-\lambda,\frac{b_1}{b_2}}\left(t\right) + \frac{a_3}{b_2} e_{\alpha+\beta,\xi+\alpha+\beta-\kappa,\frac{b_1}{b_2}}\left(t\right),$$

according to the property $\frac{d}{dt}e_{\xi,\zeta,\lambda}(t) = e_{\xi,\zeta-1,\lambda}(t)$ of the two-parameter Mittag-Leffler function, admits the integral representation in the form

$$\dot{\varepsilon}_{cr}(t) = \frac{1}{b_2 \pi} \int_0^\infty \frac{1}{\rho^{\xi}} \left(\frac{b_1}{b_2} \frac{a_1 \sin\left(\xi\pi\right) + a_2 \rho^{\lambda} \sin\left(\left(\xi - \lambda\right)\pi\right) + a_3 \rho^{\kappa} \sin\left(\left(\xi - \kappa\right)\pi\right)}{\left|\rho^{\alpha + \beta} \mathrm{e}^{\mathrm{i}(\alpha + \beta)\pi} + \frac{b_1}{b_2}\right|^2} + \rho^{\alpha + \beta} \frac{a_1 \sin\left(\left(\xi + \alpha + \beta\right)\pi\right) + a_2 \rho^{\lambda} \sin\left(\left(\xi - \lambda + \alpha + \beta\right)\pi\right) + a_3 \rho^{\kappa} \sin\left(\left(\xi - \kappa + \alpha + \beta\right)\pi\right)}{\left|\rho^{\alpha + \beta} \mathrm{e}^{\mathrm{i}(\alpha + \beta)\pi} + \frac{b_1}{b_2}\right|^2} \right) \mathrm{e}^{-\rho t} \mathrm{d}\rho,$$

$$(33)$$

according to the expression (25), provided that $\alpha + \beta \in (0, 1)$, so that by requiring the function

$$K(\rho) = a_1 b_1 \sin(\xi \pi) + a_2 b_1 \rho^{\lambda} \sin((\xi - \lambda) \pi) + a_3 b_1 \rho^{\kappa} \sin((\xi - \kappa) \pi) + a_1 b_2 \sin((\xi + \alpha + \beta) \pi) + a_2 b_2 \rho^{\lambda} \sin((\xi - \lambda + \alpha + \beta) \pi) + a_3 b_2 \rho^{\kappa} \sin((\xi - \kappa + \alpha + \beta) \pi)$$

to be non-negative, it is ensured that function $\dot{\varepsilon}_{cr}$, see (25), is completely monotonic and therefore the creep compliance is a Bernstein function. Note, the creep compliance in the form (31) coincides with the integral of (33), since ε_{cr} (0) = 0.

3.2 Calculation of relaxation modulus and creep compliance

3.2.1 Relaxation modulus calculation

The relaxation modulus is obtained in the form (15), containing functions given by (16), (17), and (18), by inverting the relaxation modulus in the Laplace domain (2) according to the definition of inverse Laplace transform

$$\sigma_{sr}(t) = \mathcal{L}^{-1}\left[\tilde{\sigma}_{sr}(s)\right](t) = \frac{1}{2\pi i} \int_{p_0 - i\infty}^{p_0 + i\infty} \tilde{\sigma}_{sr}(s) e^{st} ds$$

More precisely, the relaxation modulus σ_{sr} in the forms $\sigma_{sr} = \sigma_{sr}^{(NP)}$ and $\sigma_{sr} = \sigma_{sr}^{(NP)} + \sigma_{sr}^{(RP)}$ is obtained by the Cauchy integral theorem

$$\oint_{\Gamma^{(\mathrm{I},\mathrm{II})}} \tilde{\sigma}_{sr}\left(s\right) \mathrm{e}^{st} \mathrm{d}s = 0, \tag{34}$$

since function $\tilde{\sigma}_{sr}$ either does not have poles and then integration is performed along the contour $\Gamma^{(I)}$, depicted in Figure 1, or has a negative real pole lying outside of the contour $\Gamma^{(II)}$, depicted in Figure 2, while the relaxation modulus σ_{sr} in the form $\sigma_{sr} = \sigma_{sr}^{(NP)} + \sigma_{sr}^{(CCP)}$ is obtained by the Cauchy residue theorem

$$\oint_{\Gamma^{(\mathrm{I})}} \tilde{\sigma}_{sr}\left(s\right) \mathrm{e}^{st} \mathrm{d}s = 2\pi \mathrm{i}\left(\operatorname{Res}\left(\tilde{\sigma}_{sr}\left(s\right) \mathrm{e}^{st}, s_{\mathrm{CCP}}\right) + \operatorname{Res}\left(\tilde{\sigma}_{sr}\left(s\right) \mathrm{e}^{st}, \bar{s}_{\mathrm{CCP}}\right)\right),\tag{35}$$

since function $\tilde{\sigma}_{sr}$ has s_{CCP} and its complex conjugate \bar{s}_{CCP} as poles lying within the contour $\Gamma^{(I)}$, depicted in Figure 1.



Table 2: Parametrization of integration contour $\Gamma^{(I)}$.

Figure 1: Integration contour $\Gamma^{(I)}$.

Ims

Table 3: Parametrization of integration contour $\Gamma^{(II)}$.

Figure 2: Integration contour $\Gamma^{(II)}$.

In the case when relaxation modulus in the Laplace domain (2) does not have poles, the Cauchy integral theorem (34), with the contour $\Gamma^{(I)}$ depicted in Figure 1 and by taking into account integrals having non-zero contributions, yields

$$\int_{\Gamma_0} \tilde{\sigma}_{sr}(s) e^{st} ds + \int_{\Gamma_3} \tilde{\sigma}_{sr}(s) e^{st} ds + \int_{\Gamma_5} \tilde{\sigma}_{sr}(s) e^{st} ds = 0,$$
(36)

that, according to the parameterization of contours Γ_3 and Γ_5 , given in Table 2, in the limit when $r \to 0$ and $R \to \infty$ becomes

$$2\pi \mathrm{i}\,\sigma_{sr}\left(t\right) + \int_{\infty}^{0} \frac{1}{\rho^{1-\xi}\mathrm{e}^{\mathrm{i}(1-\xi)\pi}} \frac{\phi_{\varepsilon}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)}{\phi_{\sigma}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)} \mathrm{e}^{\rho t\mathrm{e}^{\mathrm{i}\pi}} \mathrm{e}^{\mathrm{i}\pi} \mathrm{d}\rho + \int_{0}^{\infty} \frac{1}{\rho^{1-\xi}\mathrm{e}^{-\mathrm{i}(1-\xi)\pi}} \frac{\phi_{\varepsilon}\left(\rho\mathrm{e}^{-\mathrm{i}\pi}\right)}{\phi_{\sigma}\left(\rho\mathrm{e}^{-\mathrm{i}\pi}\right)} \mathrm{e}^{\rho t\mathrm{e}^{-\mathrm{i}\pi}} \mathrm{e}^{-\mathrm{i}\pi} \mathrm{d}\rho = 0, \quad (37)$$

transforming into

$$\begin{split} \sigma_{sr}\left(t\right) &= \sigma_{sr}^{(\mathrm{NP})}\left(t\right) \\ &= \frac{1}{2\pi\mathrm{i}} \int_{0}^{\infty} \frac{1}{\rho^{1-\xi}} \frac{\mathrm{e}^{\mathrm{i}(1-\xi)\pi} \bar{\phi}_{\varepsilon}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right) \phi_{\sigma}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right) - \mathrm{e}^{-\mathrm{i}(1-\xi)\pi} \phi_{\varepsilon}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right) \bar{\phi}_{\sigma}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right)}{|\phi_{\sigma}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right)|^{2}} \mathrm{e}^{-\rho t} \mathrm{d}\rho, \quad \mathrm{i.e.}, \\ \sigma_{sr}\left(t\right) &= \sigma_{sr}^{(\mathrm{NP})}\left(t\right) \\ &= \frac{1}{2\pi\mathrm{i}} \int_{0}^{\infty} \frac{1}{\rho^{1-\xi}} \frac{\left|\phi_{\varepsilon}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right)\right|}{|\phi_{\sigma}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right)|} \left(\mathrm{e}^{\mathrm{i}(1-\xi)\pi - \mathrm{arg}\,\phi_{\varepsilon}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right) + \mathrm{arg}\,\phi_{\sigma}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right)} - \mathrm{e}^{-\mathrm{i}(1-\xi)\pi + \mathrm{arg}\,\phi_{\varepsilon}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right) - \mathrm{arg}\,\phi_{\sigma}\left(\rho \mathrm{e}^{\mathrm{i}\pi}\right)}\right) \mathrm{e}^{-\rho t} \mathrm{d}\rho, \end{split}$$

and becoming of the form (16) or (19) having the function K, given by (20), taken into account.

In addition to the integrals along contours $\Gamma_{3a} \cup \Gamma_{3b}$ and $\Gamma_{5a} \cup \Gamma_{5b}$, that are parts of contour $\Gamma^{(\text{II})}$ from Figure 2, there are additional integrals along contours Γ_8 and Γ_9 having non-zero contributions in the case when relaxation modulus in the Laplace domain (2) has a negative real pole $s_{\text{RP}} = \rho_{\text{RP}} e^{i\pi}$, so that the Cauchy integral theorem (34) yields

$$\int_{\Gamma_0} \tilde{\sigma}_{sr}\left(s\right) \mathrm{e}^{st} \mathrm{d}s + \int_{\Gamma_{3a} \cup \Gamma_{3b}} \tilde{\sigma}_{sr}\left(s\right) \mathrm{e}^{st} \mathrm{d}s + \int_{\Gamma_{5a} \cup \Gamma_{5b}} \tilde{\sigma}_{sr}\left(s\right) \mathrm{e}^{st} \mathrm{d}s + \int_{\Gamma_8} \tilde{\sigma}_{sr}\left(s\right) \mathrm{e}^{st} \mathrm{d}s + \int_{\Gamma_9} \tilde{\sigma}_{sr}\left(s\right) \mathrm{e}^{st} \mathrm{d}s = 0,$$

with the first three terms being already defined by (37) and with the remaining terms, containing integrals along contours Γ_8 and Γ_9 parameterized as in Table 3, transforming the previous expression into

$$2\pi i \sigma_{sr}(t) + 2\pi i \sigma_{sr}^{(NP)}(t) + \int_{\pi}^{0} \frac{1}{(s_{RP} + re^{i\varphi})^{1-\xi}} \frac{\phi_{\varepsilon} \left(s_{RP} + re^{i\varphi}\right)}{\phi_{\sigma} \left(s_{RP} + re^{i\varphi}\right)} e^{\left(s_{RP} + re^{i\varphi}\right)t} ire^{i\varphi} d\varphi + \int_{0}^{-\pi} \frac{1}{(\bar{s}_{RP} + re^{i\varphi})^{1-\xi}} \frac{\phi_{\varepsilon} \left(\bar{s}_{RP} + re^{i\varphi}\right)}{\phi_{\sigma} \left(\bar{s}_{RP} + re^{i\varphi}\right)} e^{\left(\bar{s}_{RP} + re^{i\varphi}\right)t} ire^{i\varphi} d\varphi = 0,$$

in the limit when $r \to 0$ and $R \to \infty$, so that

$$\sigma_{sr}\left(t\right) = \sigma_{sr}^{\left(\mathrm{NP}\right)}\left(t\right) + \sigma_{sr}^{\left(\mathrm{RP}\right)}\left(t\right) + \sigma_$$

where

$$\sigma_{sr}^{(\mathrm{RP})}(t) = \frac{1}{2\pi\mathrm{i}} \left(-\mathrm{i}\pi \frac{1}{s_{\mathrm{RP}}^{1-\xi}} \frac{\phi_{\varepsilon}\left(s_{\mathrm{RP}}\right)}{\phi_{\sigma}'\left(s_{\mathrm{RP}}\right)} \mathrm{e}^{s_{\mathrm{RP}}t} - \mathrm{i}\pi \frac{1}{\bar{s}_{\mathrm{RP}}^{1-\xi}} \frac{\phi_{\varepsilon}\left(\bar{s}_{\mathrm{RP}}\right)}{\phi_{\sigma}'\left(\bar{s}_{\mathrm{RP}}\right)} \mathrm{e}^{\bar{s}_{\mathrm{RP}}t} \right)$$
$$= -\frac{1}{2} \frac{1}{\rho_{\mathrm{RP}}^{1-\xi}} \frac{\mathrm{e}^{-\mathrm{i}(1-\xi)\pi}\phi_{\varepsilon}\left(s_{\mathrm{RP}}\right)}{\phi_{\varepsilon}'\left(s_{\mathrm{RP}}\right)} \frac{\bar{\phi}_{\sigma}'\left(s_{\mathrm{RP}}\right)}{\phi_{\sigma}'\left(s_{\mathrm{RP}}\right)} \mathrm{e}^{-\rho_{\mathrm{RP}}t}} \mathrm{e}^{-\rho_{\mathrm{RP}}t}$$

becoming of the form given by (17). The integral along contour Γ_8 is calculated as

$$\begin{split} \lim_{r \to 0} \int_{\Gamma_8} \tilde{\sigma}_{sr} \left(s \right) \mathrm{e}^{st} \mathrm{d}s &= \lim_{r \to 0} \int_{\pi}^0 \frac{1}{\left(s_{\mathrm{RP}} + r \mathrm{e}^{\mathrm{i}\varphi} \right)^{1-\xi}} \frac{\phi_{\varepsilon} \left(s_{\mathrm{RP}} + r \mathrm{e}^{\mathrm{i}\varphi} \right)}{\phi_{\sigma} \left(s_{\mathrm{RP}} + r \mathrm{e}^{\mathrm{i}\varphi} \right)} \mathrm{e}^{\left(s_{\mathrm{RP}} + r \mathrm{e}^{\mathrm{i}\varphi} \right) t} \mathrm{i} r \mathrm{e}^{\mathrm{i}\varphi} \mathrm{d}\varphi \\ &= \lim_{r \to 0} \int_{\pi}^0 \frac{1}{\left(s_{\mathrm{RP}} + r \mathrm{e}^{\mathrm{i}\varphi} \right)^{1-\xi}} \frac{\phi_{\varepsilon} \left(s_{\mathrm{RP}} + r \mathrm{e}^{\mathrm{i}\varphi} \right)}{\phi_{\sigma} \left(s_{\mathrm{RP}} \right) + \phi_{\sigma}' \left(s \right) \left(s - s_{\mathrm{RP}} \right) \right|_{s=s_{\mathrm{RP}} + r \mathrm{e}^{\mathrm{i}\varphi}} \mathrm{e}^{\left(s_{\mathrm{RP}} + r \mathrm{e}^{\mathrm{i}\varphi} \right) t} \mathrm{i} r \mathrm{e}^{\mathrm{i}\varphi} \mathrm{d}\varphi \\ &= -\mathrm{i}\pi \frac{1}{s_{\mathrm{RP}}^{1-\xi}} \frac{\phi_{\varepsilon} \left(s_{\mathrm{RP}} \right)}{\phi_{\sigma}' \left(s_{\mathrm{RP}} \right)} \mathrm{e}^{s_{\mathrm{RP}} t}, \end{split}$$

by expanding function ϕ_{σ} into the series and by taking into account $\phi_{\sigma}(s_{\rm RP}) = 0$, while the similar calculation yields the integral along contour Γ_9 in the form

$$\lim_{r \to 0} \int_{\Gamma_9} \tilde{\sigma}_{sr}(s) e^{st} ds = -i\pi \frac{1}{\bar{s}_{\rm RP}^{1-\xi}} \frac{\phi_{\varepsilon}(\bar{s}_{\rm RP})}{\phi_{\sigma}'(\bar{s}_{\rm RP})} e^{\bar{s}_{\rm RP}t}.$$

On the right-hand side of equation (36), the term originating from the residues appears when the function $\tilde{\sigma}_{sr}$, given by (2), has a pair of complex conjugated poles $s_{\rm RP}$ and $\bar{s}_{\rm RP}$, since the Cauchy residues theorem (35), with the contour $\Gamma^{(I)}$ depicted in Figure 1, where the poles are located within the area bounded by the contour, by taking into account integrals having non-zero contributions, yields

$$\int_{\Gamma_0} \tilde{\sigma}_{sr}(s) e^{st} ds + \int_{\Gamma_3} \tilde{\sigma}_{sr}(s) e^{st} ds + \int_{\Gamma_5} \tilde{\sigma}_{sr}(s) e^{st} ds = 2\pi i \left(\operatorname{Res}\left(\tilde{\sigma}_{sr}(s) e^{st}, s_{\text{CCP}} \right) + \operatorname{Res}\left(\tilde{\sigma}_{sr}(s) e^{st}, \bar{s}_{\text{CCP}} \right) \right),$$

that becomes

$$\sigma_{sr}\left(t\right) - \sigma_{sr}^{\left(\mathrm{NP}\right)}\left(t\right) = \sigma_{sr}^{\left(\mathrm{CCP}\right)}\left(t\right),$$

where

$$\begin{split} \sigma_{sr}^{(\mathrm{CCP})}\left(t\right) &= \frac{1}{\rho_{\mathrm{CCP}}^{1-\xi}\mathrm{e}^{\mathrm{i}(1-\xi)\varphi_{\mathrm{CCP}}}} \frac{\phi_{\varepsilon}\left(s_{\mathrm{CCP}}\right)}{\phi_{\sigma}'\left(s_{\mathrm{CCP}}\right)} \mathrm{e}^{\mathrm{Re}\,s_{\mathrm{CCP}}t + \mathrm{i}\,\mathrm{Im}\,s_{\mathrm{CCP}}t} \\ &+ \frac{1}{\rho_{\mathrm{CCP}}^{1-\xi}\mathrm{e}^{-\mathrm{i}(1-\xi)\varphi_{\mathrm{CCP}}}} \frac{\phi_{\varepsilon}\left(\bar{s}_{\mathrm{CCP}}\right)}{\phi_{\sigma}'\left(\bar{s}_{\mathrm{CCP}}\right)} \mathrm{e}^{\mathrm{Re}\,\bar{s}_{\mathrm{CCP}}t + \mathrm{i}\,\mathrm{Im}\,\bar{s}_{\mathrm{CCP}}t} \\ &= \frac{1}{\rho_{\mathrm{CCP}}^{1-\xi}} \mathrm{e}^{\mathrm{Re}\,s_{\mathrm{CCP}}t} \left(\frac{\phi_{\varepsilon}\left(s_{\mathrm{CCP}}\right)}{\phi_{\sigma}'\left(s_{\mathrm{CCP}}\right)} \frac{\mathrm{e}^{\mathrm{i}\,\mathrm{Im}\,s_{\mathrm{CCP}}t}}{\mathrm{e}^{\mathrm{i}(1-\xi)\varphi_{\mathrm{CCP}}}} + \frac{\phi_{\varepsilon}\left(\bar{s}_{\mathrm{CCP}}\right)}{\phi_{\sigma}'\left(\bar{s}_{\mathrm{CCP}}\right)} \frac{\mathrm{e}^{\mathrm{i}\,\mathrm{Im}\,\bar{s}_{\mathrm{CCP}}t}}{\mathrm{e}^{-\mathrm{i}(1-\xi)\varphi_{\mathrm{CCP}}}}\right) \\ &= \frac{1}{\rho_{\mathrm{CCP}}^{1-\xi}} \mathrm{e}^{-|\mathrm{Re}\,s_{\mathrm{CCP}}|t} \\ &\times \frac{\mathrm{e}^{\mathrm{i}(\mathrm{Im}\,s_{\mathrm{CCP}}t - (1-\xi)\varphi_{\mathrm{CCP}})\phi_{\varepsilon}\left(s_{\mathrm{CCP}}\right)}{\left|\phi_{\sigma}'\left(s_{\mathrm{CCP}}\right)\right|^{2}} + \mathrm{e}^{-\mathrm{i}(\mathrm{Im}\,s_{\mathrm{CCP}}t - (1-\xi)\varphi_{\mathrm{CCP}})}\overline{\phi}_{\varepsilon}\left(s_{\mathrm{CCP}}\right)}, \end{split}$$

becoming of the form (18).

It is left to be shown that the integrals along contours Γ_1 , Γ_2 , Γ_4 , Γ_6 , and Γ_7 , as parts of contours $\Gamma^{(I)}$ and $\Gamma^{(II)}$ from Figures 1 and 2, have zero contributions in both Cauchy integral and residues theorems, see (34) and (35), in the limit when $r \to 0$ and $R \to \infty$.

The integral along the contour Γ_1 reads

$$I_{\Gamma_1} = \int_{p_0}^0 \frac{1}{(p+\mathrm{i}R)^{1-\xi}} \frac{\phi_\varepsilon (p+\mathrm{i}R)}{\phi_\sigma (p+\mathrm{i}R)} \mathrm{e}^{(p+\mathrm{i}R)t} \mathrm{d}p,$$

according to the contour parametrization given in Tables 2 and 3, so that it has zero contribution, since its absolute value yields

$$\begin{split} |I_{\Gamma_1}| &\leqslant \int_0^{p_0} \frac{1}{|p+\mathrm{i}R|^{1-\xi}} \frac{|\phi_{\varepsilon}(p+\mathrm{i}R)|}{|\phi_{\sigma}(p+\mathrm{i}R)|} \mathrm{e}^{pt} \mathrm{d}p \\ &\leqslant \int_0^{p_0} \frac{1}{R^{1-\xi} \left|1-\mathrm{i}\frac{p}{R}\right|^{1-\xi}} \frac{|\phi_{\varepsilon}(p+\mathrm{i}R)|}{|\phi_{\sigma}(p+\mathrm{i}R)|} \mathrm{e}^{pt} \mathrm{d}p \\ &\leqslant \int_0^{p_0} \frac{1}{R^{1-\xi}} \frac{|\phi_{\varepsilon}(p+\mathrm{i}R)|}{|\phi_{\sigma}(p+\mathrm{i}R)|} \mathrm{e}^{pt} \mathrm{d}p \\ &\leqslant \int_0^{p_0} \frac{1}{R^{1-\xi-\zeta_R}} \mathrm{e}^{pt} \mathrm{d}p \to 0, \quad \text{for} \quad R \to \infty, \end{split}$$

if the functions ϕ_{ε} and ϕ_{σ} , that are power-type functions, see Table 1, satisfy $\frac{|\phi_{\varepsilon}(p+iR)|}{|\phi_{\sigma}(p+iR)|} \sim R^{\zeta_R}$, where $\zeta_R < 1-\xi$ when $R \to \infty$ for $p \in [0, p_0]$. Using the similar argumentation, one can prove that the integral along contour Γ_7 has zero contribution as well.

The integral along contour Γ_2 , parametrized as in Tables 2 and 3, takes the form

$$I_{\Gamma_{2}} = \int_{\frac{\pi}{2}}^{\pi} \frac{1}{R^{1-\xi} \mathrm{e}^{\mathrm{i}(1-\xi)\varphi}} \frac{\phi_{\varepsilon}\left(R\mathrm{e}^{\mathrm{i}\varphi}\right)}{\phi_{\sigma}\left(R\mathrm{e}^{\mathrm{i}\varphi}\right)} \mathrm{e}^{Rt\mathrm{e}^{\mathrm{i}\varphi}} \mathrm{i}R\mathrm{e}^{\mathrm{i}\varphi} \mathrm{d}\varphi$$

and has zero contribution, that can be proved by considering its absolute value

$$|I_{\Gamma_2}| \leqslant \int_{\frac{\pi}{2}}^{\pi} R^{\xi} \frac{\left|\phi_{\varepsilon}\left(R e^{i\varphi}\right)\right|}{\left|\phi_{\sigma}\left(R e^{i\varphi}\right)\right|} e^{Rt \cos\varphi} d\varphi \to 0, \quad \text{for} \quad R \to \infty,$$

since the integrand term $e^{Rt\cos\varphi} \to 0$ for $\varphi \in \left[\frac{\pi}{2}, \pi\right]$ when $R \to \infty$ and since functions ϕ_{ε} and ϕ_{σ} are of power-type, see Table 1. The integral along contour Γ_6 also has zero contribution, that is proved by the similar argumentation.

If the power-type functions ϕ_{ε} and ϕ_{σ} , see Table 1, are such that $\frac{|\phi_{\varepsilon}(re^{i\varphi})|}{|\phi_{\sigma}(re^{i\varphi})|} \sim \frac{1}{r^{\zeta_r}}$, where $\zeta_r < \xi$ when $r \to 0$ for $\varphi \in [-\pi, \pi]$, it can be proved that the integral along contour Γ_4 has zero contribution, since, according to the parametrization given in Tables 2 and 3, the integral is

$$I_{\Gamma_4} = \int_{\pi}^{-\pi} \frac{1}{r^{1-\xi} \mathrm{e}^{\mathrm{i}(1-\xi)\varphi}} \frac{\phi_{\varepsilon} \left(r \mathrm{e}^{\mathrm{i}\varphi}\right)}{\phi_{\sigma} \left(r \mathrm{e}^{\mathrm{i}\varphi}\right)} \mathrm{e}^{r t \mathrm{e}^{\mathrm{i}\varphi}} \mathrm{i} r \mathrm{e}^{\mathrm{i}\varphi} \mathrm{d}\varphi,$$

so that

$$\begin{aligned} |I_{\Gamma_4}| &\leqslant \int_{-\pi}^{\pi} r^{\xi} \frac{\left|\phi_{\varepsilon}\left(r \mathrm{e}^{\mathrm{i}\varphi}\right)\right|}{\left|\phi_{\sigma}\left(r \mathrm{e}^{\mathrm{i}\varphi}\right)\right|} \mathrm{e}^{rt \mathrm{cos}\varphi} \mathrm{d}\varphi \\ &\leqslant \int_{-\pi}^{\pi} r^{\xi - \zeta_r} \mathrm{d}\varphi \to 0, \quad \mathrm{for} \quad r \to 0. \end{aligned}$$

3.2.2 Creep compliance calculation

The creep compliance in the form (27), containing functions given by (28), (29), and (30), is obtained according to

$$\varepsilon_{cr}\left(t\right) = \int_{0}^{t} \epsilon_{cr}\left(t'\right) \mathrm{d}t',\tag{38}$$

that is a consequence of considering the function

$$\tilde{\epsilon}_{cr}\left(s\right) = s\tilde{\varepsilon}_{cr}\left(s\right) = \frac{1}{s^{\xi}}\frac{\phi_{\sigma}\left(s\right)}{\phi_{\varepsilon}\left(s\right)},\tag{39}$$

having the inverse Laplace transform obtained as

$$\epsilon_{cr}\left(t\right) = \frac{\mathrm{d}}{\mathrm{d}t}\varepsilon_{cr}\left(t\right) + \varepsilon_{cr}^{\left(g\right)}\delta\left(t\right) = \frac{\mathrm{d}}{\mathrm{d}t}\varepsilon_{cr}\left(t\right),$$

with the glass compliance $\varepsilon_{cr}^{(g)} = \lim_{t\to 0} \varepsilon_{cr}(t) = \lim_{s\to\infty} s\tilde{\varepsilon}_{cr}(s) = \lim_{s\to\infty} \frac{1}{s^{\xi}} \frac{\phi_{\sigma}(s)}{\phi_{\varepsilon}(s)} = 0$ for all considered models, rather than the creep compliance in Laplace domain (3). The function ϵ_{cr} is calculated using the definition of the inverse Laplace transform applied to function $\tilde{\epsilon}_{cr}$, see (39), and integration in the complex plane along the Bromwich contour, i.e., by

$$\epsilon_{cr}\left(t\right) = \frac{1}{2\pi i} \int_{p_0 - i\infty}^{p_0 + i\infty} \tilde{\epsilon}_{cr}\left(s\right) e^{st} ds.$$
(40)

More precisely, the creep compliance ε_{cr} in the forms $\varepsilon_{cr} = \varepsilon_{cr}^{(NP)}$ and $\varepsilon_{cr} = \varepsilon_{cr}^{(NP)} + \varepsilon_{cr}^{(RP)}$ is obtained according to (38), with the function ϵ_{cr} calculated by the Cauchy integral theorem

$$\oint_{\Gamma^{(\mathrm{I},\mathrm{II})}} \tilde{\epsilon}_{cr}\left(s\right) \mathrm{e}^{st} \mathrm{d}s = 0,\tag{41}$$

where the integration is performed either along the contour $\Gamma^{(I)}$, depicted in Figure 1, if the function $\tilde{\epsilon}_{cr}$ does not have poles, or along the contour $\Gamma^{(II)}$, depicted in Figure 2, if the function $\tilde{\epsilon}_{cr}$ has a negative real pole

that lies outside of contour $\Gamma^{(\text{II})}$, while the creep compliance ε_{cr} in the form $\varepsilon_{cr} = \varepsilon_{cr}^{(\text{NP})} + \varepsilon_{cr}^{(\text{CCP})}$ is obtained according to (38), with the function ϵ_{cr} calculated by the Cauchy residue theorem

$$\oint_{\Gamma^{(\mathrm{I})}} \tilde{\epsilon}_{cr}\left(s\right) \mathrm{e}^{st} \mathrm{d}s = 2\pi \mathrm{i}\left(\operatorname{Res}\left(\tilde{\epsilon}_{cr}\left(s\right) \mathrm{e}^{st}, s_{\mathrm{CCP}}\right) + \operatorname{Res}\left(\tilde{\epsilon}_{cr}\left(s\right) \mathrm{e}^{st}, \bar{s}_{\mathrm{CCP}}\right)\right),\tag{42}$$

since function $\tilde{\epsilon}_{cr}$ has s_{CCP} and its complex conjugate \bar{s}_{CCP} as poles lying within the contour $\Gamma^{(I)}$, depicted in Figure 1.

Namely, in the case when function $\tilde{\epsilon}_{cr}$, given by (39), does not have poles, the Cauchy integral theorem (41), with the contour $\Gamma^{(I)}$ depicted in Figure 1, taking into account integrals that have non-zero contributions, yields

$$\int_{\Gamma_0} \tilde{\epsilon}_{cr}(s) e^{st} ds + \int_{\Gamma_3} \tilde{\epsilon}_{cr}(s) e^{st} ds + \int_{\Gamma_5} \tilde{\epsilon}_{cr}(s) e^{st} ds = 0,$$

where the integrals along contours Γ_0 , Γ_3 , and Γ_5 , parameterized as in Table 2, in the limit when $r \to 0$ and $R \to \infty$ become

$$2\pi i \epsilon_{cr} (t) + \int_{\infty}^{0} \frac{1}{\rho^{\xi} e^{i\xi\pi}} \frac{\phi_{\sigma} (\rho e^{i\pi})}{\phi_{\varepsilon} (\rho e^{i\pi})} e^{\rho t e^{i\pi}} e^{i\pi} d\rho + \int_{0}^{\infty} \frac{1}{\rho^{\xi} e^{-i\xi\pi}} \frac{\phi_{\sigma} (\rho e^{-i\pi})}{\phi_{\varepsilon} (\rho e^{-i\pi})} e^{\rho t e^{-i\pi}} e^{-i\pi} d\rho = 0$$
(43)

transforming into

$$\begin{split} \epsilon_{cr}\left(t\right) &= \epsilon_{cr}^{(\mathrm{NP})}\left(t\right) \\ &= \frac{1}{2\pi\mathrm{i}} \int_{0}^{\infty} \frac{1}{\rho^{\xi}} \frac{\mathrm{e}^{\mathrm{i}\xi\pi} \phi_{\varepsilon}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right) \bar{\phi}_{\sigma}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right) - \mathrm{e}^{-\mathrm{i}\xi\pi} \bar{\phi}_{\varepsilon}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right) \phi_{\sigma}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)}{\left|\phi_{\varepsilon}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)\right|^{2}} \mathrm{e}^{-\rho t} \mathrm{d}\rho \\ &= \frac{1}{\pi} \int_{0}^{\infty} \frac{1}{\rho^{\xi}} \frac{K\left(\rho\right)}{\left|\phi_{\varepsilon}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)\right|^{2}} \mathrm{e}^{-\rho t} \mathrm{d}\rho, \quad \mathrm{i.e.}, \\ \epsilon_{cr}\left(t\right) &= \epsilon_{cr}^{(\mathrm{NP})}\left(t\right) \\ &= \frac{1}{2\pi\mathrm{i}} \int_{0}^{\infty} \frac{1}{\rho^{\xi}} \frac{\left|\phi_{\sigma}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)\right|}{\left|\phi_{\varepsilon}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)\right|} \left(\mathrm{e}^{\mathrm{i}\left(\xi\pi + \mathrm{arg}\,\phi_{\varepsilon}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right) - \mathrm{arg}\,\phi_{\sigma}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)\right)} - \mathrm{e}^{-\mathrm{i}\left(\xi\pi + \mathrm{arg}\,\phi_{\varepsilon}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right) - \mathrm{arg}\,\phi_{\sigma}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)\right)} \right| \mathrm{e}^{-\rho t} \mathrm{d}\rho \\ &= \frac{1}{\pi} \int_{0}^{\infty} \frac{1}{\rho^{\xi}} \frac{\left|\phi_{\sigma}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)\right|}{\left|\phi_{\varepsilon}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right)\right|} \sin\left(\mathrm{arg}\,\phi_{\varepsilon}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right) - \mathrm{arg}\,\phi_{\sigma}\left(\rho\mathrm{e}^{\mathrm{i}\pi}\right) + \xi\pi\right)\mathrm{e}^{-\rho t} \mathrm{d}\rho, \end{split}$$

that, according to (38) and according to the form (20) of function K, become creep compliance $\varepsilon_{cr}^{(NP)}$ in the equivalent forms (28) and (31).

On the other hand, if the function $\tilde{\epsilon}_{cr}$, given by (39), has a negative real pole $s_{\rm RP} = \rho_{\rm RP} e^{i\pi}$, then the Cauchy integral theorem (41) is considered for contour $\Gamma^{(\rm II)}$ from Figure 2, so that, in addition to the integrals along contours $\Gamma_{3a} \cup \Gamma_{3b}$ and $\Gamma_{5a} \cup \Gamma_{5b}$, there are also integrals along contours Γ_8 and Γ_9 , that have non-zero contributions, implying

$$\int_{\Gamma_0} \tilde{\epsilon}_{cr}(s) e^{st} ds + \int_{\Gamma_{3a} \cup \Gamma_{3b}} \tilde{\epsilon}_{cr}(s) e^{st} ds + \int_{\Gamma_{5a} \cup \Gamma_{5b}} \tilde{\epsilon}_{cr}(s) e^{st} ds + \int_{\Gamma_8} \tilde{\epsilon}_{cr}(s) e^{st} ds + \int_{\Gamma_9} \tilde{\epsilon}_{cr}(s) e^{st} ds = 0,$$

with the first three terms being already defined by (43) and with the remaining terms calculated along the contours Γ_8 and Γ_9 , parameterized as in Table 3, transforming the previous expression in the limit when $r \to 0$ and $R \to \infty$ into

$$2\pi \mathbf{i} \,\epsilon_{cr}\left(t\right) + 2\pi \mathbf{i} \,\epsilon_{cr}^{(\mathrm{NP})}\left(t\right) + \int_{\pi}^{0} \frac{1}{(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi})^{\xi}} \frac{\phi_{\sigma}\left(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)}{\phi_{\varepsilon}\left(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)} \mathbf{e}^{\left(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)t} \mathrm{i} r\mathrm{e}^{\mathrm{i}\varphi} \mathrm{d}\varphi + \int_{0}^{-\pi} \frac{1}{\left(\bar{s}_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)^{\xi}} \frac{\phi_{\sigma}\left(\bar{s}_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)}{\phi_{\varepsilon}\left(\bar{s}_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)} \mathbf{e}^{\left(\bar{s}_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)t} \mathrm{i} r\mathrm{e}^{\mathrm{i}\varphi} \mathrm{d}\varphi = 0,$$

that becomes

$$\begin{aligned} \epsilon_{cr}\left(t\right) &= \epsilon_{cr}^{\left(\mathrm{NP}\right)}\left(t\right) + \frac{1}{2\pi\mathrm{i}} \left(-\mathrm{i}\pi \frac{1}{s_{\mathrm{RP}}^{\xi}} \frac{\phi_{\sigma}\left(s_{\mathrm{RP}}\right)}{\phi_{\varepsilon}'\left(s_{\mathrm{RP}}\right)} \mathrm{e}^{s_{\mathrm{RP}}t} - \mathrm{i}\pi \frac{1}{\bar{s}_{\mathrm{RP}}^{\xi}} \frac{\phi_{\sigma}\left(\bar{s}_{\mathrm{RP}}\right)}{\phi_{\varepsilon}'\left(\bar{s}_{\mathrm{RP}}\right)} \mathrm{e}^{\bar{s}_{\mathrm{RP}}t}\right) \\ &= \epsilon_{cr}^{\left(\mathrm{NP}\right)}\left(t\right) - \frac{1}{2} \frac{1}{\rho_{\mathrm{RP}}^{\xi}} \frac{\mathrm{e}^{-\mathrm{i}\xi\pi} \bar{\phi}_{\varepsilon}'\left(s_{\mathrm{RP}}\right) \phi_{\sigma}\left(s_{\mathrm{RP}}\right) + \mathrm{e}^{\mathrm{i}\xi\pi} \phi_{\varepsilon}'\left(s_{\mathrm{RP}}\right) \bar{\phi}_{\sigma}\left(s_{\mathrm{RP}}\right)}{\left|\phi_{\varepsilon}'\left(s_{\mathrm{RP}}\right)\right|^{2}} \mathrm{e}^{-\rho_{\mathrm{RP}}t} \end{aligned}$$

where the last term transforms into

$$\epsilon_{cr}^{(\text{RP})}(t) = -\frac{1}{\rho_{\text{RP}}^{\xi}} \frac{|\phi_{\sigma}(s_{\text{RP}})|}{|\phi_{\varepsilon}'(s_{\text{RP}})|} \cos\left(\arg\phi_{\varepsilon}'(s_{\text{RP}}) - \arg\phi_{\sigma}(s_{\text{RP}}) + \xi\pi\right) e^{-\rho_{\text{RP}}t},$$

yielding the function $\varepsilon_{cr}^{(\text{RP})}$ in the form (29), according to (38). The integral along contour Γ_8 is calculated as

$$\begin{split} \lim_{r \to 0} \int_{\Gamma_{8}} \tilde{\epsilon}_{cr}\left(s\right) \mathrm{e}^{st} \mathrm{d}s &= \lim_{r \to 0} \int_{\pi}^{0} \frac{1}{\left(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)^{\xi}} \frac{\phi_{\sigma}\left(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)}{\phi_{\varepsilon}\left(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)} \mathrm{e}^{\left(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)t} \mathrm{i}r\mathrm{e}^{\mathrm{i}\varphi} \mathrm{d}\varphi \\ &= \lim_{r \to 0} \int_{\pi}^{0} \frac{1}{\left(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)^{\xi}} \frac{\phi_{\sigma}\left(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)}{\phi_{\varepsilon}\left(s_{\mathrm{RP}}\right) + \phi_{\varepsilon}'\left(s\right)\left(s - s_{\mathrm{RP}}\right)\Big|_{s=s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}} + \dots} \mathrm{e}^{\left(s_{\mathrm{RP}} + r\mathrm{e}^{\mathrm{i}\varphi}\right)t} \mathrm{i}r\mathrm{e}^{\mathrm{i}\varphi} \mathrm{d}\varphi \\ &= -\mathrm{i}\pi \frac{1}{s_{\mathrm{RP}}^{\xi}} \frac{\phi_{\sigma}\left(s_{\mathrm{RP}}\right)}{\phi_{\varepsilon}'\left(s_{\mathrm{RP}}\right)} \mathrm{e}^{s_{\mathrm{RP}}t}, \end{split}$$

by expanding function ϕ_{ε} into the series and by taking into account $\phi_{\varepsilon}(s_{\text{RP}}) = 0$, while the similar calculation yields the integral along contour Γ_9 in the form

$$\lim_{r \to 0} \int_{\Gamma_9} \tilde{\epsilon}_{cr}(s) \, \mathrm{e}^{st} \mathrm{d}s = -\mathrm{i}\pi \frac{1}{\bar{s}_{\mathrm{RP}}^{\xi}} \frac{\phi_{\sigma}(\bar{s}_{\mathrm{RP}})}{\phi_{\varepsilon}'(\bar{s}_{\mathrm{RP}})} \mathrm{e}^{\bar{s}_{\mathrm{RP}}t}.$$

In addition, when the function $\tilde{\epsilon}_{cr}$, see (39), has a pair of complex conjugated poles s_{RP} and \bar{s}_{RP} , located within the area bounded by the contour $\Gamma^{(I)}$ depicted in Figure 1, the Cauchy residues theorem (42), by taking into account integrals having non-zero contributions, yields

$$\int_{\Gamma_0} \tilde{\epsilon}_{cr}(s) e^{st} ds + \int_{\Gamma_3} \tilde{\epsilon}_{cr}(s) e^{st} ds + \int_{\Gamma_5} \tilde{\epsilon}_{cr}(s) e^{st} ds = 2\pi i \left(\operatorname{Res}\left(\tilde{\epsilon}_{cr}(s) e^{st}, s_{CCP} \right) + \operatorname{Res}\left(\tilde{\epsilon}_{cr}(s) e^{st}, \bar{s}_{CCP} \right) \right),$$

that, in the limit when $r \to 0$ and $R \to \infty$ and with the terms on the left-hand-side of the previous expression already defined by (43), becomes

$$\epsilon_{cr}\left(t\right) - \epsilon_{cr}^{(\text{NP})}\left(t\right) = \epsilon_{cr}^{(\text{CCP})}\left(t\right),$$

with

$$\begin{split} \epsilon_{cr}^{(\mathrm{CCP})}\left(t\right) &= \frac{1}{\rho_{\mathrm{CcP}}^{\xi} \mathrm{e}^{\mathrm{i}\xi\varphi_{\mathrm{CCP}}}} \frac{\phi_{\sigma}\left(s_{\mathrm{CCP}}\right)}{\phi_{\varepsilon}'\left(s_{\mathrm{CCP}}\right)} \mathrm{e}^{\mathrm{Re}\,s_{\mathrm{CCP}}t + \mathrm{i}\,\mathrm{Im}\,s_{\mathrm{CCP}}t} \\ &+ \frac{1}{\rho_{\mathrm{CcP}}^{\xi} \mathrm{e}^{-\mathrm{i}\xi\varphi_{\mathrm{CCP}}}} \frac{\phi_{\sigma}\left(\bar{s}_{\mathrm{CCP}}\right)}{\phi_{\varepsilon}'\left(\bar{s}_{\mathrm{CCP}}\right)} \mathrm{e}^{\mathrm{Re}\,\bar{s}_{\mathrm{CCP}}t + \mathrm{i}\,\mathrm{Im}\,\bar{s}_{\mathrm{CCP}}t} \\ &= \frac{1}{\rho_{\mathrm{CCP}}^{\xi}} \mathrm{e}^{\mathrm{Re}\,s_{\mathrm{CCP}}t} \left(\frac{1}{\mathrm{e}^{\mathrm{i}\xi\varphi_{\mathrm{CCP}}}} \frac{\phi_{\sigma}\left(s_{\mathrm{CCP}}\right)}{\phi_{\varepsilon}'\left(s_{\mathrm{CCP}}\right)} \mathrm{e}^{\mathrm{i}\,\mathrm{Im}\,s_{\mathrm{CCP}}t} + \frac{1}{\mathrm{e}^{-\mathrm{i}\xi\varphi_{\mathrm{CCP}}}} \frac{\phi_{\sigma}\left(\bar{s}_{\mathrm{CCP}}\right)}{\phi_{\varepsilon}'\left(\bar{s}_{\mathrm{CCP}}\right)} \mathrm{e}^{\mathrm{i}\,\mathrm{Im}\,\bar{s}_{\mathrm{CCP}}t} \right) \\ &= \frac{1}{\rho_{\mathrm{CCP}}^{\xi}} \mathrm{e}^{-|\mathrm{Re}\,s_{\mathrm{CCP}}|t} \\ &\times \frac{\mathrm{e}^{\mathrm{i}(\mathrm{Im}\,s_{\mathrm{CCP}}t - \xi\varphi_{\mathrm{CCP}})} \bar{\phi}_{\varepsilon}'\left(s_{\mathrm{CCP}}\right) \phi_{\sigma}\left(s_{\mathrm{CCP}}\right) + \mathrm{e}^{-\mathrm{i}(\mathrm{Im}\,s_{\mathrm{CCP}}t - \xi\varphi_{\mathrm{CCP}})} \phi_{\varepsilon}'\left(s_{\mathrm{CCP}}\right)}{\left|\phi_{\varepsilon}'\left(s_{\mathrm{CCP}}\right)\right|^{2}} \\ &= 2\frac{1}{\rho_{\mathrm{CCP}}^{\xi}} \frac{\left|\phi_{\sigma}\left(s_{\mathrm{CCP}}\right)\right|}{\left|\phi_{\varepsilon}'\left(s_{\mathrm{CCP}}\right|\right|}} \mathrm{e}^{-|\mathrm{Re}\,s_{\mathrm{CCP}}|t}\cos\left(\mathrm{Im}\,s_{\mathrm{CCP}}t - \mathrm{arg}\,\phi_{\varepsilon}'\left(s_{\mathrm{CCP}}\right) + \mathrm{arg}\,\phi_{\sigma}\left(s_{\mathrm{CCP}}\right) - \xi\varphi_{\mathrm{CCP}}\right)} \right) \end{split}$$

implying, according to (38),

$$\varepsilon_{cr}\left(t\right) = \varepsilon_{cr}^{\left(\mathrm{NP}\right)}\left(t\right) + \varepsilon_{cr}^{\left(\mathrm{CCP}\right)}\left(t\right),$$

,

with

$$\begin{split} \varepsilon_{cr}^{(\text{CCP})}\left(t\right) &= 2\frac{1}{\rho_{\text{CCP}}^{1+\xi}} \frac{|\phi_{\sigma}\left(s_{\text{CCP}}\right)|}{|\phi_{\varepsilon}'\left(s_{\text{CCP}}\right)|} \\ &\times \left(e^{-|\text{Re}\,s_{\text{CCP}}|t}\cos\left(\text{Im}\,s_{\text{CCP}}t - \arg\phi_{\varepsilon}'\left(s_{\text{CCP}}\right) + \arg\phi_{\sigma}\left(s_{\text{CCP}}\right) - \left(1+\xi\right)\varphi_{\text{CCP}}\right) \\ &- \cos\left(\arg\phi_{\varepsilon}'\left(s_{\text{CCP}}\right) - \arg\phi_{\sigma}\left(s_{\text{CCP}}\right) + \left(1+\xi\right)\varphi_{\text{CCP}}\right)\right), \end{split}$$

where the integral

$$\int_0^t e^{\mu t'} \cos\left(\omega t' + \phi\right) dt' = \frac{1}{\mu^2 + \omega^2} \left(e^{\mu t} \left(\omega \sin\left(\omega t + \phi\right) + \mu \cos\left(\omega t + \phi\right)\right) - \left(\omega \sin\phi + \mu \cos\phi\right) \right)$$
$$= \frac{1}{\sqrt{\mu^2 + \omega^2}} \left(e^{\mu t} \cos\left(\omega t + \phi - \varphi\right) - \cos\left(\phi - \varphi\right) \right)$$

is used with substitutions $\mu = \operatorname{Re} s_{CCP}$, $\omega = \operatorname{Im} s_{CCP}$, $\phi = -\arg \phi'_{\varepsilon}(s_{CCP}) + \arg \phi_{\sigma}(s_{CCP}) - \xi \varphi_{CCP}$, and $\varphi = \arctan \frac{\omega}{\mu} = \varphi_{CCP}$.

The integrals along the contours Γ_1 (Γ_7), Γ_2 (Γ_6), and Γ_4 tend to zero in the Cauchy integral and residue theorems (41) and (42) in the limit when $r \to 0$ and $R \to \infty$. Namely, if the power-type functions ϕ_{ε} and ϕ_{σ} , see Table 1, satisfy conditions

$$\begin{split} & \frac{|\phi_{\varepsilon}\left(p+\mathrm{i}R\right)|}{|\phi_{\sigma}\left(p+\mathrm{i}R\right)|} \sim R^{\zeta_{R}}, \quad \mathrm{when} \quad R \to \infty \quad \mathrm{for} \quad p \in [0, p_{0}] \,, \\ & \frac{|\phi_{\varepsilon}\left(r\mathrm{e}^{\mathrm{i}\varphi}\right)|}{|\phi_{\sigma}\left(r\mathrm{e}^{\mathrm{i}\varphi}\right)|} \sim \frac{1}{r^{\zeta_{r}}}, \quad \mathrm{when} \quad r \to 0 \quad \mathrm{for} \quad \varphi \in [-\pi, \pi] \,, \end{split}$$

that are already posed in Section 3.2.1 in order to ensure zero contributions of the integrals of function $\tilde{\sigma}_{sr}(s) e^{st}$ along the contours $\Gamma_1(\Gamma_7)$, $\Gamma_2(\Gamma_6)$, and Γ_4 , then one has

$$\begin{split} & \frac{|\phi_{\sigma}\left(p+\mathrm{i}R\right)|}{|\phi_{\varepsilon}\left(p+\mathrm{i}R\right)|} \sim \frac{1}{R^{\zeta_{R}}}, \quad \mathrm{when} \quad R \to \infty \quad \mathrm{for} \quad p \in [0, p_{0}] \,, \\ & \frac{|\phi_{\sigma}\left(r\mathrm{e}^{\mathrm{i}\varphi}\right)|}{|\phi_{\varepsilon}\left(r\mathrm{e}^{\mathrm{i}\varphi}\right)|} \sim r^{\zeta_{r}}, \quad \mathrm{when} \quad r \to 0 \quad \mathrm{for} \quad \varphi \in [-\pi, \pi] \,, \end{split}$$

and due to the similar forms of the relaxation modulus in the Laplace domain and function $\tilde{\epsilon}_{cr}$, see (2) and (39), one has the zero contributions of integrals along the contours Γ_1 (Γ_7) and Γ_4 if $\zeta_R > -\xi$ and $\zeta_r > -(1-\xi)$, yielding

$$-\xi < \zeta_R < 1-\xi$$
 and $-(1-\xi) < \zeta_r < \xi$

with the previously posed conditions $\zeta_R < 1 - \xi$ and $\zeta_r < \xi$, while the zero contributions of integrals along the contours Γ_2 (Γ_6) is simply guaranteed by the fact that ϕ_{ε} and ϕ_{σ} are power-type functions.

4 Narrowing the thermodynamical requirements

4.1 Symmetric models

4.1.1 Model ID.ID

Thermodynamical, see (168) and (169), along with the narrowed thermodynamical restrictions on the parameters for model ID.ID, given by the expression

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\beta}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{I}_{t}^{\mu}+a_{2\,0}\mathbf{D}_{t}^{\alpha+\beta-\mu}\right)\varepsilon\left(t\right),$$

are of the following form

$$0 \leqslant \alpha + \beta - \mu \leqslant 1, \quad \mu \leqslant \alpha, \quad \beta + \mu \leqslant 1, \tag{44}$$

$$-\frac{a_1}{a_2}\frac{\cos\frac{(2\alpha+\beta-\mu)\pi}{2}}{\cos\frac{(\beta+\mu)\pi}{2}} \leqslant \frac{b_1}{b_2} \leqslant \frac{a_1}{a_2}\frac{\sin\frac{(2\alpha+\beta-\mu)\pi}{2}}{\sin\frac{(\beta+\mu)\pi}{2}}\frac{\cos\frac{(2\alpha+\beta-\mu)\pi}{2}}{\cos\frac{(\beta+\mu)\pi}{2}} \leqslant \frac{a_1}{a_2}\frac{\sin\frac{(2\alpha+\beta-\mu)\pi}{2}}{\sin\frac{(\beta+\mu)\pi}{2}},\tag{45}$$

where the inequality $(45)_3$ narrows the thermodynamical restriction (169) if $\alpha \leq 2\alpha + \beta - \mu < 1$, and it is obtained by requesting the function K, given in the general form by (20) and reducing to the expression

$$K(\rho) = a_1 b_1 \sin\left(\left(\alpha - \mu\right)\pi\right) + a_1 b_2 \rho^{\alpha+\beta} \sin\left(\left(2\alpha + \beta - \mu\right)\pi\right) - a_2 b_1 \rho^{\alpha+\beta} \sin\left(\left(\beta + \mu\right)\pi\right) + a_2 b_2 \rho^{2(\alpha+\beta)} \sin\left(\left(\alpha - \mu\right)\pi\right)$$
(46)

for the model ID.ID, to have a non-negative value. The second term in (46), that can be either positive for $2\alpha + \beta - \mu \in [\alpha, 1)$, or non-positive for $2\alpha + \beta - \mu \in [1, 1 + \alpha)$, see (44)₁, combined with the third non-positive term in (46) yields narrowed thermodynamical restriction (45)₃ if $2\alpha + \beta - \mu \in [\alpha, 1)$, since $\beta + \mu \leq 2\alpha + \beta - \mu < 1$ reducing to $\mu \leq \alpha$, see (44)₂, implies $\frac{\cos \frac{(2\alpha + \beta - \mu)\pi}{2}}{\cos \frac{(\beta + \mu)\pi}{2}} \leq 1$, while if $2\alpha + \beta - \mu \in [1, 1 + \alpha)$, then one has the second and third term in (46) non-positive and therefore non-negativity of function K cannot be guaranteed, see (46).

In conclusion, the non-negativity of function K, corresponding to the model ID.ID and given by (46), is ensured if $2\alpha + \beta - \mu \in [\alpha, 1)$ by requesting (45) in addition to (44) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function, while if $2\alpha + \beta - \mu \in [1, 1 + \alpha)$, then the mentioned properties cannot be ensured.

4.1.2 Model $ID.DD^+$

Thermodynamical, see (170) and (171), along with the narrowed thermodynamical restrictions on the parameters for model ID.DD⁺, given by the expression

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\beta}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{D}_{t}^{\mu}+b_{2\,0}\mathbf{D}_{t}^{\alpha+\beta+\mu}\right)\varepsilon\left(t\right),$$

are of the following form

$$1 \leqslant \alpha + \beta + \mu \leqslant 2, \quad \beta \leqslant \mu \leqslant 1 - \alpha, \tag{47}$$

$$\frac{a_1}{a_2} \frac{\left|\cos\frac{(2\alpha+\beta+\mu)\pi}{2}\right|}{\cos\frac{(\mu-\beta)\pi}{2}} \leqslant \frac{a_1}{a_2} \frac{\left|\cos\frac{(2\alpha+\beta+\mu)\pi}{2}\right|}{\cos\frac{(\mu-\beta)\pi}{2}} \frac{\sin\frac{(2\alpha+\beta+\mu)\pi}{2}}{\sin\frac{(\mu-\beta)\pi}{2}} \leqslant \frac{b_1}{b_2},\tag{48}$$

where the inequality $(48)_1$ narrows the thermodynamical restriction (171) and it is obtained by requesting the function K, given in the general form by (20) and reducing to the expression

$$K(\rho) = a_1 b_1 \sin((\alpha + \mu) \pi) + a_1 b_2 \rho^{\alpha + \beta} \sin((2\alpha + \beta + \mu) \pi) + a_2 b_1 \rho^{\alpha + \beta} \sin((\mu - \beta) \pi) + a_2 b_2 \rho^{2(\alpha + \beta)} \sin((\alpha + \mu) \pi),$$
(49)

for the model ID.DD⁺, to have a non-negative value. The second term in (49), that is non-positive since (47)₁ implies $2\alpha + \beta + \mu \ge 1 + \alpha$ and (47)₂ implies $2\alpha + \beta + \mu = (\alpha + \beta) + (\alpha + \mu) \le 2$ and therefore $2\alpha + \beta + \mu \in [1 + \alpha, 2]$, combined with the third non-negative term in (49) yields narrowed thermodynamical restriction (48)₁, since $\frac{(\mu - \beta)\pi}{2} \le \pi - \frac{(2\alpha + \beta + \mu)\pi}{2} < \frac{\pi}{2}$ reducing to $\alpha + \mu \le 1$, see (47)₂, implies $\frac{\sin \frac{(2\alpha + \beta + \mu)\pi}{2}}{\sin \frac{(\mu - \beta)\pi}{2}} \ge 1$.

In conclusion, the non-negativity of function K, corresponding to the model ID.DD⁺ and given by (49), is ensured by requesting (48) in addition to (47), implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function.

4.1.3 Model IID.IID

Function K, given by (20), in the case of model IID.IID, having the constitutive equation given in the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{I}_{t}^{\beta}+a_{3\,0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{I}_{t}^{\alpha+\gamma-\eta}+b_{2\,0}\mathbf{I}_{t}^{\beta+\gamma-\eta}+b_{3\,0}\mathbf{D}_{t}^{\eta}\right)\varepsilon\left(t\right),$$

and with corresponding thermodynamical, see (172) - (174), as well as with the narrowed thermodynamical requirements

$$\beta < \alpha, \quad \gamma \leqslant \eta, \quad 0 \leqslant \beta + \gamma - \eta \leqslant \alpha + 2\gamma - \eta \leqslant 1, \quad \alpha + \gamma \leqslant \beta + \eta, \tag{50}$$

$$-\frac{b_3}{b_1}\frac{\cos\frac{(\alpha+\eta)\pi}{2}}{\cos\frac{(\alpha+2\gamma-\eta)\pi}{2}} \leqslant \frac{a_3}{a_1} \leqslant \frac{b_3}{b_1}\frac{\sin\frac{(\alpha+\eta)\pi}{2}}{\sin\frac{(\alpha+2\gamma-\eta)\pi}{2}}\frac{\cos\frac{(\alpha+\eta)\pi}{2}}{\cos\frac{(\alpha+2\gamma-\eta)\pi}{2}} \leqslant \frac{b_3}{b_1}\frac{\sin\frac{(\alpha+\eta)\pi}{2}}{\sin\frac{(\alpha+2\gamma-\eta)\pi}{2}},\tag{51}$$

$$-\frac{b_3}{b_2} \frac{\cos\frac{(\beta+\eta)\pi}{2}}{\cos\frac{(\beta+2\gamma-\eta)\pi}{2}} \leqslant \frac{a_3}{a_2} \leqslant \frac{b_3}{b_2} \frac{\sin\frac{(\beta+\eta)\pi}{2}}{\sin\frac{(\beta+2\gamma-\eta)\pi}{2}} \frac{\cos\frac{(\beta+\eta)\pi}{2}}{\cos\frac{(\beta+2\gamma-\eta)\pi}{2}} \leqslant \frac{b_3}{b_2} \frac{\sin\frac{(\beta+\eta)\pi}{2}}{\sin\frac{(\beta+2\gamma-\eta)\pi}{2}},\tag{52}$$

valid if $\beta + \eta < \alpha + \eta < 1$, becomes

$$K(\rho) = a_{1}b_{1}\sin((\eta - \gamma)\pi) + a_{1}b_{2}\rho^{\alpha-\beta}\sin((\alpha + \eta - \beta - \gamma)\pi) + a_{1}b_{3}\rho^{\alpha+\gamma}\sin((\alpha + \eta)\pi) + a_{2}b_{1}\rho^{\alpha-\beta}\sin((\beta + \eta - \alpha - \gamma)\pi) + a_{2}b_{2}\rho^{2(\alpha-\beta)}\sin((\eta - \gamma)\pi) + a_{2}b_{3}\rho^{2\alpha-\beta+\gamma}\sin((\beta + \eta)\pi) - a_{3}b_{1}\rho^{\alpha+\gamma}\sin((\alpha + 2\gamma - \eta)\pi) - a_{3}b_{2}\rho^{2\alpha-\beta+\gamma}\sin((\beta + 2\gamma - \eta)\pi) + a_{3}b_{3}\rho^{2(\alpha+\gamma)}\sin((\eta - \gamma)\pi),$$
(53)

so that the non-negativity requirement of function K implies

$$a_1 b_2 \sin\left(\left(\alpha + \eta - \beta - \gamma\right)\pi\right) + a_2 b_1 \sin\left(\left(\beta + \eta - \alpha - \gamma\right)\pi\right) \ge 0,\tag{54}$$

 $a_1 b_3 \sin\left(\left(\alpha + \eta\right)\pi\right) - a_3 b_1 \sin\left(\left(\alpha + 2\gamma - \eta\right)\pi\right) \ge 0,\tag{55}$

$$a_2 b_3 \sin\left(\left(\beta + \eta\right)\pi\right) - a_3 b_2 \sin\left(\left(\beta + 2\gamma - \eta\right)\pi\right) \ge 0.$$
(56)

According to the thermodynamical requirements $(50)_{1,2}$, one finds that $\alpha + \eta - \beta - \gamma = (\alpha - \beta) + (\eta - \gamma) \ge 0$, while the thermodynamical requirement $\beta + \gamma - \eta \ge 0$, given by $(50)_3$, can be modified into $\alpha + \eta - \beta - \gamma \le \alpha \le 1$, and therefore the first term in (54) has a non-negative value. The argument $(\beta + \eta - \alpha - \gamma)\pi = ((\eta - \gamma) - (\alpha - \beta))\pi$ of the second term in (54), firstly has a value that is less than a value of the argument $(\alpha + \eta - \beta - \gamma)\pi = ((\eta - \gamma) + (\alpha - \beta))\pi$ of the first term in (54), more precisely $\beta + \eta - \alpha - \gamma = (\eta - \gamma) - (\alpha - \beta) \leqslant \alpha + \eta - \beta - \gamma = (\eta - \gamma) + (\alpha - \beta) \leqslant \alpha \leqslant 1$, and, secondly it has a non-negative value, i.e., $\beta + \eta - \alpha - \gamma = (\beta + \eta) - (\alpha + \gamma) \ge 0$, according to the thermodynamical requirement (50)₄. Therefore, the inequality (54) is trivially satisfied.

According to $(50)_3$, the sine in the second term in (55) has a non-negative value, while the first term has either a positive value if $\alpha + \eta < 1$, or a non-positive value if $\alpha + \eta \ge 1$, so that if $\alpha + \eta < 1$, then the requirement (55) is transformed into

$$\frac{a_3}{a_1} \leqslant \frac{b_3}{b_1} \frac{\sin \frac{(\alpha+\eta)\pi}{2}}{\sin \frac{(\alpha+2\gamma-\eta)\pi}{2}} \frac{\cos \frac{(\alpha+\eta)\pi}{2}}{\cos \frac{(\alpha+2\gamma-\eta)\pi}{2}},$$

representing the narrowed thermodynamical restriction $(51)_3$, since $\alpha + 2\gamma - \eta \leq \alpha + \eta < 1$, reducing to $\gamma \leq \eta$, see $(50)_2$, implies $\frac{\cos \frac{(\alpha+\eta)\pi}{2}}{\cos \frac{(\alpha+2\gamma-\eta)\pi}{2}} \leq 1$. Similarly, according to $(50)_{1,3}$, the sine in the second term in (56) has a non-negative value, since $0 \leq \beta + 2\gamma - \eta \leq \alpha + 2\gamma - \eta \leq 1$, while the first term has either a positive value if $\beta + \eta < 1$, or a non-positive value if $\beta + \eta \geq 1$, so that if $\beta + \eta < 1$, then the requirement (56) is transformed into

$$\frac{a_3}{a_2} \leqslant \frac{b_3}{b_2} \frac{\sin\frac{(\beta+\eta)\pi}{2}}{\sin\frac{(\beta+2\gamma-\eta)\pi}{2}} \frac{\cos\frac{(\beta+\eta)\pi}{2}}{\cos\frac{(\beta+2\gamma-\eta)\pi}{2}}$$

representing the narrowed thermodynamical restriction $(52)_3$, since $\beta + 2\gamma - \eta \leq \beta + \eta < 1$ reducing to $\gamma \leq \eta$, see $(50)_2$, implies $\frac{\cos \frac{(\beta+\eta)\pi}{2}}{\cos \frac{(\beta+2\gamma-\eta)\pi}{2}} \leq 1$.

In conclusion, the non-negativity of function K, corresponding to the model IID.IID and given by (53), is ensured if $\beta + \eta < \alpha + \eta < 1$, see (50)₁, by requesting (51) and (52) in addition to (50) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function, while if either $\alpha + \eta \ge 1$ or $\alpha + \eta > \beta + \eta \ge 1$, then the mentioned properties cannot be ensured, since either (55), or both (55) and (56) cannot be satisfied.

4.1.4 Model IDD.IDD

Function K, given by (20), in the case of the model IDD.IDD, having the constitutive equation given in the form $\left(\frac{1}{2} - \frac{1}{2} + \frac{1}{2} +$

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha} + a_{2\,0}\mathbf{D}_{t}^{\beta} + a_{3\,0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right) = \left(b_{1\,0}\mathbf{I}_{t}^{\mu} + b_{2\,0}\mathbf{D}_{t}^{\alpha+\beta-\mu} + b_{3\,0}\mathbf{D}_{t}^{\alpha+\gamma-\mu}\right)\varepsilon\left(t\right)$$

and with corresponding thermodynamical, see (175) - (177), as well as with the narrowed thermodynamical requirements

$$0 \leqslant \alpha + \gamma - \mu \leqslant 1, \quad \beta < \gamma, \quad \mu \leqslant \alpha, \quad \gamma + \mu \leqslant \alpha + \beta, \quad \gamma + \mu \leqslant 1, \tag{57}$$

$$\frac{b_2}{b_1} \frac{\cos\frac{(2\alpha+\beta-\mu)\pi}{2}}{\cos\frac{(\beta+\mu)\pi}{2}} \leqslant \frac{a_2}{a_1} \leqslant \frac{b_2}{b_1} \frac{\sin\frac{(2\alpha+\beta-\mu)\pi}{2}}{\sin\frac{(\beta+\mu)\pi}{2}} \frac{\cos\frac{(2\alpha+\beta-\mu)\pi}{2}}{\cos\frac{(\beta+\mu)\pi}{2}} \leqslant \frac{b_2}{b_1} \frac{\sin\frac{(2\alpha+\beta-\mu)\pi}{2}}{\sin\frac{(\beta+\mu)\pi}{2}},\tag{58}$$

$$-\frac{b_3}{b_1}\frac{\cos\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\cos\frac{(\gamma+\mu)\pi}{2}} \leqslant \frac{a_3}{a_1} \leqslant \frac{b_3}{b_1}\frac{\sin\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\sin\frac{(\gamma+\mu)\pi}{2}}\frac{\cos\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\cos\frac{(\gamma+\mu)\pi}{2}} \leqslant \frac{b_3}{b_1}\frac{\sin\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\sin\frac{(\gamma+\mu)\pi}{2}},\tag{59}$$

valid if $2\alpha + \beta - \mu < 2\alpha + \gamma - \mu < 1$, becomes

$$K(\rho) = a_{1}b_{1}\sin((\alpha - \mu)\pi) + a_{1}b_{2}\rho^{\alpha+\beta}\sin((2\alpha + \beta - \mu)\pi) + a_{1}b_{3}\rho^{\alpha+\gamma}\sin((2\alpha + \gamma - \mu)\pi) - a_{2}b_{1}\rho^{\alpha+\beta}\sin((\beta + \mu)\pi) + a_{2}b_{2}\rho^{2(\alpha+\beta)}\sin((\alpha - \mu)\pi) + a_{2}b_{3}\rho^{2\alpha+\beta+\gamma}\sin((\alpha + \gamma - \beta - \mu)\pi) - a_{3}b_{1}\rho^{\alpha+\gamma}\sin((\gamma + \mu)\pi) + a_{3}b_{2}\rho^{2\alpha+\beta+\gamma}\sin((\alpha + \beta - \gamma - \mu)\pi) + a_{3}b_{3}\rho^{2(\alpha+\gamma)}\sin((\alpha - \mu)\pi),$$
(60)

so that the non-negativity requirement of function K implies

$$a_2 b_3 \sin\left(\left(\alpha + \gamma - \beta - \mu\right)\pi\right) + a_3 b_2 \sin\left(\left(\alpha + \beta - \gamma - \mu\right)\pi\right) \ge 0,\tag{61}$$

$$a_1 b_2 \sin\left(\left(2\alpha + \beta - \mu\right)\pi\right) - a_2 b_1 \sin\left(\left(\beta + \mu\right)\pi\right) \ge 0,\tag{62}$$

$$a_1 b_3 \sin\left(\left(2\alpha + \gamma - \mu\right)\pi\right) - a_3 b_1 \sin\left(\left(\gamma + \mu\right)\pi\right) \ge 0.$$
(63)

The first term in (61) has a non-negative value since $\alpha + \gamma - \beta - \mu = (\alpha - \mu) + (\gamma - \beta) \ge 0$, according to the thermodynamical requirements $(57)_{2,3}$, as well as $\alpha + \gamma - \beta - \mu = (\alpha + \gamma - \mu) - \beta \le 1 - \beta$, according to the thermodynamical requirement $(57)_1$. The second term in (61) also has a non-negative value, since $\alpha + \beta - \gamma - \mu = (\alpha + \beta) - (\gamma + \mu) \ge 0$, according to the thermodynamical requirement $(57)_4$, as well as $\alpha + \beta - \gamma - \mu = \alpha - (\gamma + \mu - \beta) \le \alpha \le 1$, since $(\gamma - \beta) + \mu \ge 0$, according to $(57)_2$. Therefore, the inequality (61) is trivially satisfied. According to $(57)_{2,5}$, yielding $0 \le \beta + \mu < \gamma + \mu \le 1$, the sine in the second term in (62) as well as in (63) has a non-negative value, while the first term has either a positive value if $2\alpha + \beta - \mu < 1$, or a non-positive value if $2\alpha + \beta - \mu < 1$, so that if $2\alpha + \beta - \mu < 1$, then the requirement (62) is transformed into

$$\frac{a_2}{a_1} \leqslant \frac{b_2}{b_1} \frac{\sin \frac{(2\alpha + \beta - \mu)\pi}{2}}{\sin \frac{(\beta + \mu)\pi}{2}} \frac{\cos \frac{(2\alpha + \beta - \mu)\pi}{2}}{\cos \frac{(\beta + \mu)\pi}{2}},$$

representing the narrowed thermodynamical restriction $(58)_3$, since $\beta + \mu \leq 2\alpha + \beta - \mu < 1$, reducing to $\mu \leq \alpha$, see $(57)_3$, implies $\frac{\cos \frac{(2\alpha+\beta-\mu)\pi}{2}}{\cos \frac{(\beta+\mu)\pi}{2}} \leq 1$. Similarly, the second term in (63), as already mentioned, has a non-negative value, while the first term has either a positive value if $2\alpha + \gamma - \mu < 1$ or a non-positive value if $2\alpha + \gamma - \mu \geq 1$, so that if $2\alpha + \gamma - \mu < 1$, then the requirement (63) is transformed into

$$\frac{a_3}{a_1} \leqslant \frac{b_3}{b_1} \frac{\sin \frac{(2\alpha + \gamma - \mu)\pi}{2}}{\sin \frac{(\gamma + \mu)\pi}{2}} \frac{\cos \frac{(2\alpha + \gamma - \mu)\pi}{2}}{\cos \frac{(\gamma + \mu)\pi}{2}},$$

representing the narrowed thermodynamical restriction (59)₃, since $\gamma + \mu \leq 2\alpha + \gamma - \mu < 1$ reducing to $\mu \leq \alpha$, see (57)₃, implies $\frac{\cos \frac{(2\alpha + \gamma - \mu)\pi}{2}}{\cos \frac{(\gamma + \mu)\pi}{2}} \leq 1$.

In conclusion, the non-negativity of function K, corresponding to the model IDD.IDD and given by (60), is ensured if $2\alpha + \beta - \mu < 2\alpha + \gamma - \mu < 1$, see (57)₂, by requesting (58) and (59) in addition to (57) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function, while if either $2\alpha + \gamma - \mu \ge 1$ or $2\alpha + \gamma - \mu \ge 2\alpha + \beta - \mu \ge 1$, then the mentioned properties cannot be ensured, since either (63), or both (62) and (63) cannot be satisfied.

4.1.5 Model IID.IDD

Function K, given by (20), in the case of the model IID.IDD, having the constitutive equation given in the form

$$\left(a_{1\ 0}\mathbf{I}_{t}^{\alpha} + a_{2\ 0}\mathbf{I}_{t}^{\beta} + a_{3\ 0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right) = \left(b_{1\ 0}\mathbf{I}_{t}^{\mu} + b_{2\ 0}\mathbf{D}_{t}^{\nu} + b_{3\ 0}\mathbf{D}_{t}^{\alpha+\gamma-\mu}\right)\varepsilon\left(t\right)$$

and with corresponding thermodynamical, see (178) and (179), as well as with the narrowed thermodynamical requirements

$$\mu \leq \beta < \alpha, \quad \gamma \leq \nu, \quad \alpha + \beta + \gamma \leq 1 + \mu, \quad \mu + \nu - \gamma < \alpha \leq 1 - \nu, \tag{64}$$

$$0 \leqslant \left\{ \begin{array}{c} \alpha - \beta - \gamma - \mu \\ \alpha - 2\mu - \nu \end{array} \right\} \leqslant 2\alpha - \beta - 2\mu - \nu \leqslant \left\{ \begin{array}{c} 2\alpha - \beta - \mu \\ 2\alpha + \gamma - 2\mu - \nu \end{array} \right\} \leqslant 2\alpha + \gamma - \mu < 1, \tag{65}$$

$$-\frac{b_3}{b_1}\frac{\cos\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\cos\frac{(\gamma+\mu)\pi}{2}} \leqslant \frac{a_3}{a_1} \leqslant \frac{b_3}{b_1}\frac{\sin\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\sin\frac{(\gamma+\mu)\pi}{2}}\frac{\cos\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\cos\frac{(\gamma+\mu)\pi}{2}} \leqslant \frac{b_3}{b_1}\frac{\sin\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\sin\frac{(\gamma+\mu)\pi}{2}},\tag{66}$$

becomes

$$K(\rho) = a_{1}b_{1}\sin((\alpha - \mu)\pi) + a_{1}b_{2}\rho^{\alpha - \beta}\sin((2\alpha - \beta - \mu)\pi) + a_{1}b_{3}\rho^{\alpha + \gamma}\sin((2\alpha + \gamma - \mu)\pi) + a_{2}b_{1}\rho^{\mu + \nu}\sin((\alpha - 2\mu - \nu)\pi) + a_{2}b_{2}\rho^{\alpha - \beta + \mu + \nu}\sin((2\alpha - \beta - 2\mu - \nu)\pi) + a_{2}b_{3}\rho^{\alpha + \gamma + \mu + \nu}\sin((2\alpha + \gamma - 2\mu - \nu)\pi) - a_{3}b_{1}\rho^{\alpha + \gamma}\sin((\mu + \gamma)\pi) + a_{3}b_{2}\rho^{2\alpha - \beta + \gamma}\sin((\alpha - \beta - \gamma - \mu)\pi) + a_{3}b_{3}\rho^{2(\alpha + \gamma)}\sin((\alpha - \mu)\pi),$$
(67)

so that the non-negativity requirement of function K implies

$$a_1 b_3 \sin\left(\left(2\alpha + \gamma - \mu\right)\pi\right) - a_3 b_1 \sin\left(\left(\gamma + \mu\right)\pi\right) \ge 0,\tag{68}$$

$$\sin\left(\left(2\alpha - \beta - \mu\right)\pi\right) \ge 0, \quad \sin\left(\left(\alpha - 2\mu - \nu\right)\pi\right) \ge 0,\tag{69}$$

$$\sin\left(\left(2\alpha - \beta - 2\mu - \nu\right)\pi\right) \ge 0, \quad \sin\left(\left(2\alpha + \gamma - 2\mu - \nu\right)\pi\right) \ge 0, \quad \sin\left(\left(\alpha - \beta - \gamma - \mu\right)\pi\right) \ge 0. \tag{70}$$

The notation used in the narrowed thermodynamical requirement $(65)_{1,2}$ means that both $\alpha - \beta - \gamma - \mu$ and $\alpha - 2\mu - \nu$ are non-negative and not greater than $2\alpha - \beta - 2\mu - \nu$, while the relation between $\alpha - \beta - \gamma - \mu$ and $\alpha - 2\mu - \nu$ cannot be established and the same interpretation holds for the notation used in $(65)_{3,4}$.

The first term in (68) has either a positive value if $2\alpha + \gamma - \mu < 1$, or a non-positive value if $2\alpha + \gamma - \mu \ge 1$, since $2\alpha + \gamma - \mu = (\alpha + \gamma) + (\alpha - \mu) \in (0, 2)$, due to $\alpha + \gamma \le \alpha + \nu \le 1$, according to the thermodynamical requirements (64)_{1,2,4}, while the second term in (68) has a non-positive value, since $\gamma + \mu \le \alpha + \nu \le 1$, according to the thermodynamical requirements (64)_{1,2,4}, so that if $2\alpha + \gamma - \mu < 1$, then the requirement (68) is transformed into $(2\alpha \pm \alpha \pm \mu)\pi$ $(2\alpha \pm \mu)\pi$

$$\frac{a_3}{a_1} \leqslant \frac{b_3}{b_1} \frac{\sin \frac{(2\alpha+\gamma-\mu)\pi}{2}}{\sin \frac{(\gamma+\mu)\pi}{2}} \frac{\cos \frac{(2\alpha+\gamma-\mu)\pi}{2}}{\cos \frac{(\gamma+\mu)\pi}{2}},$$

representing the narrowed thermodynamical restriction (66)₃, since $\gamma + \mu \leq 2\alpha + \gamma - \mu < 1$ reducing to $\mu \leq \alpha$, see (64)₁, implies $\frac{\cos \frac{(2\alpha+\gamma-\mu)\pi}{2}}{\cos \frac{(\gamma+\mu)\pi}{2}} \leq 1.$

The chain of inequalities

$$\left\{\begin{array}{c} \alpha - \beta - \gamma - \mu \\ \alpha - 2\mu - \nu \end{array}\right\} \leqslant 2\alpha - \beta - 2\mu - \nu \leqslant \left\{\begin{array}{c} 2\alpha - \beta - \mu \\ 2\alpha + \gamma - 2\mu - \nu \end{array}\right\} \leqslant 2\alpha + \gamma - \mu,$$

holds since inequality $\alpha - \beta - \gamma - \mu \leq 2\alpha - \beta - 2\mu - \nu$ reduces to the thermodynamical requirement (64)₄, inequality $\alpha - 2\mu - \nu \leq 2\alpha - \beta - 2\mu - \nu$ reduces to the requirement (64)₁, while all other inequalities are trivially satisfied. On the other hand, neither can be stated that $\alpha - 2\mu - \nu \leq \alpha - \beta - \gamma - \mu$, nor that $2\alpha + \gamma - 2\mu - \nu \leq 2\alpha - \beta - \mu$, since both inequalities reduce to $\beta + \gamma \leq \mu + \nu$, while the thermodynamical requirements $(64)_{1,2}$ imply $\beta + \gamma \leq \alpha + \nu$.

In addition to the request $2\alpha + \gamma - \mu < 1$, that guarantees non-negativity of (68), in order to satisfy inequalities (69) and (70), one requests that $\alpha - \beta - \gamma - \mu \ge 0$ as well as that $\alpha - 2\mu - \nu \ge 0$, since both $\alpha - \beta - \gamma - \mu = \alpha - \beta - (\gamma + \mu) \ge -1 + \alpha - \beta \ge -1 \text{ and } \alpha - 2\mu - \nu = \alpha - \mu - (\mu + \nu) \ge -1 + \alpha - \mu \ge -1,$ according to the thermodynamical requirement $(64)_{1,2,4}$.

In conclusion, the non-negativity of function K, corresponding to the model IID.IDD and given by (67), is ensured if the request (65) on the orders of fractional integrals and derivatives, along with the thermodynamical restriction (64), is satisfied in addition to the request (66) on the model parameters, implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function, while if the request (65) is violated, then the mentioned properties cannot be ensured, since some or all of the conditions in (68), (69), and (70) may not satisfied.

4.1.6 Model $I^{+}ID.I^{+}ID$

Function K, given by (20), in the case of model $I^{+}ID.I^{+}ID$, having the constitutive equation given in the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{1+\alpha} + a_{2\,0}\mathbf{I}_{t}^{\frac{1+\alpha-\gamma}{2}} + a_{3\,0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right) = \left(b_{1\,0}\mathbf{I}_{t}^{1+\mu} + b_{2\,0}\mathbf{I}_{t}^{\frac{1+\mu-(\alpha+\gamma-\mu)}{2}} + b_{3\,0}\mathbf{D}_{t}^{\alpha+\gamma-\mu}\right)\varepsilon\left(t\right),$$

and with corresponding thermodynamical, see (180) - (183), as well as with the narrowed thermodynamical requirements

$$\mu \leqslant \alpha, \quad \alpha + \gamma + 2 \left(\alpha - \mu \right) = 3\alpha + \gamma - 2\mu \leqslant 1, \tag{71}$$

$$\frac{a_2}{a_1} \leqslant \frac{b_2}{b_1} \frac{\cos\frac{(1-3\alpha-\gamma+2\mu)\pi}{4}}{\cos\frac{(1+\alpha-\gamma-2\mu)\pi}{4}} \frac{\sin\frac{(1-3\alpha-\gamma+2\mu)\pi}{4}}{\sin\frac{(1+\alpha-\gamma-2\mu)\pi}{4}} \leqslant \frac{b_2}{b_1} \frac{\cos\frac{(1-3\alpha-\gamma+2\mu)\pi}{4}}{\cos\frac{(1+\alpha-\gamma-2\mu)\pi}{4}},\tag{72}$$

$$\frac{a_3}{a_2} \leqslant \frac{b_3}{b_2} \frac{\sin\frac{(1+3\alpha+\gamma-2\mu)\pi}{4}}{\sin\frac{(1-\alpha+\gamma+2\mu)\pi}{4}} \frac{\cos\frac{(1+3\alpha+\gamma-2\mu)\pi}{4}}{\cos\frac{(1-\alpha+\gamma+2\mu)\pi}{4}} \leqslant \frac{b_3}{b_2} \frac{\sin\frac{(1+3\alpha+\gamma-2\mu)\pi}{4}}{\sin\frac{(1-\alpha+\gamma+2\mu)\pi}{4}},\tag{73}$$

$$a_3b_1\cos\frac{(\gamma+\mu)\pi}{2} \le a_2b_2\sin\frac{(\alpha-\mu)\pi}{2} + a_1b_3\cos\frac{(2\alpha+\gamma-\mu)\pi}{2},$$
 (74)

$$a_1 b_3 \sin \frac{(2\alpha + \gamma - \mu)\pi}{2} \leqslant a_2 b_2 \cos \frac{(\alpha - \mu)\pi}{2} - a_3 b_1 \sin \frac{(\gamma + \mu)\pi}{2},$$
 (75)

$$a_{1}b_{3}\sin\frac{(2\alpha+\gamma-\mu)\pi}{2} \leqslant a_{2}b_{2}\cos\frac{(\alpha-\mu)\pi}{2}\frac{\sin\frac{(\alpha-\mu)\pi}{2}}{\cos\frac{(2\alpha+\gamma-\mu)\pi}{2}} + a_{3}b_{1}\sin\frac{(\gamma+\mu)\pi}{2}\frac{\cos\frac{(\gamma+\mu)\pi}{2}}{\cos\frac{(2\alpha+\gamma-\mu)\pi}{2}},$$
 (76)

becomes

$$K(\rho) = a_1 b_1 \sin((\alpha - \mu)\pi) + a_1 b_2 \rho^{\frac{1+\alpha+\gamma}{2}} \sin\frac{(1 + 3\alpha + \gamma - 2\mu)\pi}{2} + a_1 b_3 \rho^{1+\alpha+\gamma} \sin((1 + 2\alpha + \gamma - \mu)\pi) - a_2 b_1 \rho^{\frac{1+\alpha+\gamma}{2}} \sin\frac{(1 - \alpha + \gamma + 2\mu)\pi}{2} + a_2 b_2 \rho^{1+\alpha+\gamma} \sin((\alpha - \mu)\pi) + a_2 b_3 \rho^{3\frac{1+\alpha+\gamma}{2}} \sin\frac{(1 + 3\alpha + \gamma - 2\mu)\pi}{2} - a_3 b_1 \rho^{1+\alpha+\gamma} \sin((1 + \gamma + \mu)\pi) - a_3 b_2 \rho^{3\frac{1+\alpha+\gamma}{2}} \sin\frac{(1 - \alpha + \gamma + 2\mu)\pi}{2} + a_3 b_3 \rho^{2(1+\alpha+\gamma)} \sin((\alpha - \mu)\pi),$$
(77)

so that the non-negativity requirement of function K implies

$$a_1 b_2 \sin \frac{(1+3\alpha+\gamma-2\mu)\pi}{2} - a_2 b_1 \sin \frac{(1-\alpha+\gamma+2\mu)\pi}{2} \ge 0,$$
(78)

$$a_{2}b_{3}\sin\frac{(1+3\alpha+\gamma-2\mu)\pi}{2} - a_{3}b_{2}\sin\frac{(1-\alpha+\gamma+2\mu)\pi}{2} \ge 0,$$
(79)

$$-a_{1}b_{3}\sin((2\alpha + \gamma - \mu)\pi) + a_{2}b_{2}\sin((\alpha - \mu)\pi) + a_{3}b_{1}\sin((\gamma + \mu)\pi) \ge 0.$$
(80)

The sines appearing in the first term in (78) and (79) have a non-negative value, since $3\alpha + \gamma - 2\mu \in (0, 1)$, i.e., $1+3\alpha+\gamma-2\mu \in (1, 2)$, according to the thermodynamical requirement $(71)_2$, as well as the sines appearing in the second term in (78) and in (79), since $1-\alpha+\gamma+2\mu \in (0, 2)$, because of $-\alpha+\gamma+2\mu \in (-1, 1)$, that is valid since $-\alpha+\gamma+2\mu = (\alpha+\gamma)-2(\alpha-\mu) \leq 1-4(\alpha-\mu) \leq 1$ according to the thermodynamical restrictions (71), as well as since the thermodynamical requirement $(71)_2$ transforms into $-\alpha+\gamma+2\mu = 2(\alpha+\gamma) = -(3\alpha+\gamma-2\mu) \geq -1$ implying $-\alpha+\gamma+2\mu \geq -1+2(\alpha+\gamma) \geq -1$, so that $1-\alpha+\gamma+2\mu \geq 2(\alpha+\gamma) \geq 0$. Therefore, the requirements (78) and (79) respectively transform into

$$\frac{a_2}{a_1} \leqslant \frac{b_2}{b_1} \frac{\cos\frac{(1-3\alpha-\gamma+2\mu)\pi}{4}}{\cos\frac{(1+\alpha-\gamma-2\mu)\pi}{4}} \frac{\sin\frac{(1-3\alpha-\gamma+2\mu)\pi}{4}}{\sin\frac{(1+\alpha-\gamma-2\mu)\pi}{4}} \quad \text{and} \quad \frac{a_3}{a_2} \leqslant \frac{b_3}{b_2} \frac{\sin\frac{(1+3\alpha+\gamma-2\mu)\pi}{4}}{\sin\frac{(1-\alpha+\gamma+2\mu)\pi}{4}} \frac{\cos\frac{(1+3\alpha+\gamma-2\mu)\pi}{4}}{\cos\frac{(1-\alpha+\gamma+2\mu)\pi}{4}}, \tag{81}$$

where the requirement (78) is rewritten as

$$a_1 b_2 \sin \frac{(1 - 3\alpha - \gamma + 2\mu)\pi}{2} - a_2 b_1 \sin \frac{(1 + \alpha - \gamma - 2\mu)\pi}{2} \ge 0,$$

since $\sin\left(\frac{\pi}{2}+\phi\right) = \sin\left(\frac{\pi}{2}-\phi\right)$ and then transformed into $(81)_1$ in order to be combined with the thermodynamical restriction $(181)_1$. Expressions given by (81) represent the narrowed thermodynamical restrictions $(72)_2$ and $(73)_2$, since $\frac{\sin\left(\frac{1-3\alpha-\gamma+2\mu\right)\pi}{4}\right)}{\sin\left(\frac{1+\alpha-\gamma-2\mu\right)\pi}{4}} \leq 1$ and $\frac{\cos\left(\frac{1+3\alpha+\gamma-2\mu\right)\pi}{4}}{\cos\left(\frac{1-\alpha+\gamma+2\mu\right)\pi}{4}} \leq 1$, due to $1-3\alpha-\gamma+2\mu \leq 1+\alpha-\gamma-2\mu$ and $1-\alpha+\gamma+2\mu \leq 1+3\alpha+\gamma-2\mu$, both reducing to $\mu \leq \alpha$, see $(71)_1$.

The requirement $(71)_2$, reading $3\alpha + \gamma - 2\mu = 2\alpha + \gamma - \mu + (\alpha - \mu) \leq 1$, i.e., $2\alpha + \gamma - \mu \leq 1 - (\alpha - \mu) \leq 1$, combined with the requirement $(71)_1$ guarantees a non-negative value of the sine in the first term in (80), while the requirement $(71)_1$ ensures non-negativity of the sine in the second term in (80), and finally non-negativity of the sine in the third term in (80) is guaranteed by the requirement $(71)_2$, so that the requirement (80) is transformed into the narrowed thermodynamical restriction (76) if the right-hand-side of (76) is less than the right-hand-side of (75), since $\frac{\sin \frac{(\alpha - \mu)\pi}{2}}{\cos \frac{(2\alpha + \gamma - \mu)\pi}{2}} = \frac{\cos \frac{(1 - (\alpha - \mu))\pi}{\cos \frac{(2\alpha + \gamma - \mu)\pi}{2}} \leq 1$ and $\frac{\cos \frac{(\gamma + \mu)\pi}{2}}{\cos \frac{(2\alpha + \gamma - \mu)\pi}{2}} \geq 1$, due to $2\alpha + \gamma - \mu \leq 1 - (\alpha - \mu) < 1$ and $\gamma + \mu \leq 2\alpha + \gamma - \mu < 1$, respectively reducing to $2\alpha + \gamma - \mu \leq 1$ and $\mu \leq \alpha$, see (71).

In conclusion, the non-negativity of function K, corresponding to the model $I^{+}ID.I^{+}ID$ and given by (77), is ensured by requesting (72), (73), and possibly (76) in addition to (71), (74), and (75) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function.

4.1.7 Model $IDD^+.IDD^+$

Function K, given by (20), in the case of the model IDD^+ . IDD^+ , having the constitutive equation given in the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha} + a_{2\,0}\mathbf{D}_{t}^{\frac{1+\gamma-\alpha}{2}} + a_{3\,0}\mathbf{D}_{t}^{1+\gamma}\right)\sigma\left(t\right) = \left(b_{1\,0}\mathbf{I}_{t}^{\alpha+\gamma-\eta} + b_{2\,0}\mathbf{D}_{t}^{\frac{1+\eta-(\alpha+\gamma-\eta)}{2}} + b_{3\,0}\mathbf{D}_{t}^{1+\eta}\right)\varepsilon\left(t\right),$$

and with corresponding thermodynamical, see (184) - (187), as well as with the narrowed thermodynamical requirements

$$0 \leqslant \alpha + \gamma - \eta \leqslant 1, \quad \gamma \leqslant \eta, \quad \alpha + \eta + (\eta - \gamma) = \alpha - \gamma + 2\eta \leqslant 1, \tag{82}$$

$$\frac{a_2}{a_1} \leqslant \frac{b_2}{b_1} \frac{\sin \frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\sin \frac{(1+\alpha+3\gamma-2\eta)\pi}{4}} \frac{\cos \frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\cos \frac{(1+\alpha+3\gamma-2\eta)\pi}{4}} \leqslant \frac{b_2}{b_1} \frac{\sin \frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\sin \frac{(1+\alpha+3\gamma-2\eta)\pi}{4}},\tag{83}$$

$$\frac{a_3}{a_2} \leqslant \frac{b_3}{b_2} \frac{\cos\frac{(1-\alpha+\gamma-2\eta)\pi}{4}}{\cos\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}} \frac{\sin\frac{(1-\alpha+\gamma-2\eta)\pi}{4}}{\sin\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}} \leqslant \frac{b_3}{b_2} \frac{\cos\frac{(1-\alpha+\gamma-2\eta)\pi}{4}}{\cos\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}},\tag{84}$$

$$a_{3}b_{1}\cos\frac{(\alpha+2\gamma-\eta)\pi}{2} - a_{2}b_{2}\sin\frac{(\eta-\gamma)\pi}{2} \leqslant a_{1}b_{3}\cos\frac{(\alpha+\eta)\pi}{2},\tag{85}$$

$$a_1 b_3 \sin \frac{(\alpha + \eta) \pi}{2} \leqslant a_2 b_2 \cos \frac{(\eta - \gamma) \pi}{2} - a_3 b_1 \sin \frac{(\alpha + 2\gamma - \eta) \pi}{2}, \tag{86}$$

$$a_1 b_3 \sin \frac{(\alpha+\eta)\pi}{2} \leqslant a_2 b_2 \cos \frac{(\eta-\gamma)\pi}{2} \frac{\sin \frac{(\eta-\gamma)\pi}{2}}{\cos \frac{(\alpha+\eta)\pi}{2}} + a_3 b_1 \sin \frac{(\alpha+2\gamma-\eta)\pi}{2} \frac{\cos \frac{(\alpha+2\gamma-\eta)\pi}{2}}{\cos \frac{(\alpha+\eta)\pi}{2}},\tag{87}$$

becomes

$$K(\rho) = a_{1}b_{1}\sin((\eta - \gamma)\pi) + a_{1}b_{2}\rho^{\frac{1+\alpha+\gamma}{2}}\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{2} - a_{1}b_{3}\rho^{1+\alpha+\gamma}\sin((\alpha+\eta)\pi) - a_{2}b_{1}\rho^{\frac{1+\alpha+\gamma}{2}}\sin\frac{(1+\alpha+3\gamma-2\eta)\pi}{2} + a_{2}b_{2}\rho^{1+\alpha+\gamma}\sin((\eta-\gamma)\pi) + a_{2}b_{3}\rho^{3\frac{1+\alpha+\gamma}{2}}\sin\frac{(1+\alpha+2\eta-\gamma)\pi}{2} + a_{3}b_{1}\rho^{1+\alpha+\gamma}\sin((\alpha+2\gamma-\eta)\pi) - a_{3}b_{2}\rho^{3\frac{1+\alpha+\gamma}{2}}\sin\frac{(1+\alpha+3\gamma-2\eta)\pi}{2} + a_{3}b_{3}\rho^{2(1+\alpha+\gamma)}\sin((\eta-\gamma)\pi),$$
(88)

so that the non-negativity requirement of function K implies

$$a_{1}b_{2}\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{2} - a_{2}b_{1}\sin\frac{(1+\alpha+3\gamma-2\eta)\pi}{2} \ge 0$$
(89)

$$a_{2}b_{3}\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{2} - a_{3}b_{2}\sin\frac{(1+\alpha+3\gamma-2\eta)\pi}{2} \ge 0$$
(90)

$$-a_{1}b_{3}\sin((\alpha + \eta)\pi) + a_{2}b_{2}\sin((\eta - \gamma)\pi) + a_{3}b_{1}\sin((\alpha + 2\gamma - \eta)\pi) \ge 0.$$
(91)

According to the thermodynamical requirement $(82)_3$, the first term in (89), as well as in (90) has a non-negative value, since $1+\alpha-\gamma+2\eta \in (1,2)$, while the second term in (89), as well as in (90) is non-positive, since one has $1+\alpha+3\gamma-2\eta \in (0,2)$, due to $1+\alpha+3\gamma-2\eta = 1-(\alpha-\gamma+2\eta)+2(\alpha+\gamma) \ge 2(\alpha+\gamma) \ge 0$, where the restriction $(82)_3$ is used in the form $-(\alpha-\gamma+2\eta) \ge -1$, as well as $1+\alpha+3\gamma-2\eta = 1+(\alpha-\gamma+2\eta)-4(\eta-\gamma) \le 2-4(\eta-\gamma) \le 2$, with the restriction $(82)_3$ used again, so that the requirements (89) and (90) are transformed into

$$\frac{a_2}{a_1} \leqslant \frac{b_2}{b_1} \frac{\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\sin\frac{(1+\alpha+3\gamma-2\eta)\pi}{4}} \frac{\cos\frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\cos\frac{(1+\alpha+3\gamma-2\eta)\pi}{4}} \quad \text{and} \quad \frac{a_3}{a_2} \leqslant \frac{b_3}{b_2} \frac{\cos\frac{(1-\alpha+\gamma-2\eta)\pi}{4}}{\cos\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}} \frac{\sin\frac{(1-\alpha+\gamma-2\eta)\pi}{4}}{\sin\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}}, \tag{92}$$

and represent narrowed thermodynamical requirements (83) and (84), where the requirement (90) becomes

$$a_2b_3\sin\frac{(1-\alpha+\gamma-2\eta)\pi}{2} - a_3b_2\sin\frac{(1-\alpha-3\gamma+2\eta)\pi}{2} \ge 0,$$

since $\sin\left(\frac{\pi}{2}+\phi\right) = \sin\left(\frac{\pi}{2}-\phi\right)$ and then transformed into $(92)_2$ in order to be combined with the thermodynamical restriction (185)₂. Expressions given by (92) represent the narrowed thermodynamical restrictions $(83)_2$ and $(84)_2$, since $\frac{\cos\left(\frac{(1+\alpha-\gamma+2\eta)\pi}{4}\right)}{\cos\left(\frac{(1+\alpha+3\gamma-2\eta)\pi}{4}\right)} \leq 1$ and $\frac{\sin\left(\frac{(1-\alpha+\gamma-2\eta)\pi}{4}\right)}{\sin\left(\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}\right)} \leq 1$, due to $1+\alpha+3\gamma-2\eta \leq 1+\alpha-\gamma+2\eta$ and $1-\alpha+\gamma-2\eta \leq 1-\alpha-3\gamma+2\eta$, both reducing to $\gamma \leq \eta$, see $(82)_2$.

The requirements (82) ensure non-negative values of sines in the first and third term in (91), since $\alpha + \eta \leq 1 - (\eta - \gamma) \leq 1$ and $\alpha + 2\gamma - \eta = (\alpha - \gamma + 2\eta) - 3(\eta - \gamma) \leq 1$ as well as $\alpha + 2\gamma - \eta = (\alpha + \gamma - \eta) + \gamma \geq \gamma$, while the second term in (91) has a non-negative value according to (82)₂, so that the requirement (91) is transformed into the narrowed thermodynamical restriction (87) if the right-hand-side of (87) is less than the right-hand-side of (86), since $\frac{\sin \frac{(\eta - \gamma)\pi}{2}}{\cos \frac{(\alpha + \eta)\pi}{2}} = \frac{\cos \frac{(1 - (\eta - \gamma))\pi}{2}}{\cos \frac{(\alpha + \eta)\pi}{2}} \leq 1$ and $\frac{\cos \frac{(\alpha + 2\gamma - \eta)\pi}{2}}{\cos \frac{(\alpha + \eta)\pi}{2}} \geq 1$, due to $\alpha + \eta \leq 1 - (\eta - \gamma) < 1$ and $\alpha + 2\gamma - \eta \leq \alpha + \eta < 1$, respectively reducing to $\alpha - \gamma + 2\eta \leq 1$ and $\gamma \leq \eta$, see (82)_{3,2}.

In conclusion, the non-negativity of function K, corresponding to the model IDD^+ . IDD^+ and given by (88), is ensured by requesting (83), (84), and possibly (87) in addition to (82), (85), and (86) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function.

4.1.8 Model $I^{+}ID.IDD^{+}$

Function K, given by (20), in the case of the model $I^{+}ID.IDD^{+}$, having the constitutive equation given in the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{1+\alpha} + a_{2\,0}\mathbf{I}_{t}^{\frac{1+\alpha-\gamma}{2}} + a_{3\,0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right) = \left(b_{1\,0}\mathbf{I}_{t}^{\alpha+\gamma-\eta} + b_{2\,0}\mathbf{D}_{t}^{\frac{1+\eta-(\alpha+\gamma-\eta)}{2}} + b_{3\,0}\mathbf{D}_{t}^{1+\eta}\right)\varepsilon\left(t\right),$$

and with corresponding thermodynamical, see (188) - (191), as well as with the narrowed thermodynamical requirements

$$\eta \leqslant \gamma, \quad \alpha + \gamma + 2\left(\gamma - \eta\right) = \alpha + 3\gamma - 2\eta \leqslant 1, \tag{93}$$

$$\frac{a_1}{b_1} \frac{\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\cos\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}} \leqslant \frac{a_1}{b_1} \frac{\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\cos\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}} \frac{\cos\frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\sin\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}} \leqslant \frac{a_2}{b_2},\tag{94}$$

$$\frac{a_2}{b_2} \leqslant \frac{a_3}{b_3} \frac{\cos\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}}{\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{4}} \frac{\sin\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}}{\cos\frac{(1+\alpha-\gamma+2\eta)\pi}{4}} \leqslant \frac{a_3}{b_3} \frac{\cos\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}}{\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{4}},\tag{95}$$

$$a_1 b_3 \cos \frac{(\alpha+\eta)\pi}{2} - a_2 b_2 \sin \frac{(\gamma-\eta)\pi}{2} \leqslant a_3 b_1 \cos \frac{(\alpha+2\gamma-\eta)\pi}{2}, \tag{96}$$

$$a_3b_1\sin\frac{(\alpha+2\gamma-\eta)\pi}{2} \leqslant a_2b_2\cos\frac{(\gamma-\eta)\pi}{2} - a_1b_3\sin\frac{(\alpha+\eta)\pi}{2},\tag{97}$$

$$a_{3}b_{1}\sin\frac{(\alpha+2\gamma-\eta)\pi}{2} \leqslant a_{2}b_{2}\cos\frac{(\gamma-\eta)\pi}{2}\frac{\sin\frac{(\gamma-\eta)\pi}{2}}{\cos\frac{(\alpha+2\gamma-\eta)\pi}{2}} + a_{1}b_{3}\sin\frac{(\alpha+\eta)\pi}{2}\frac{\cos\frac{(\alpha+\eta)\pi}{2}}{\cos\frac{(\alpha+2\gamma-\eta)\pi}{2}}, \tag{98}$$

becomes

$$K(\rho) = a_{1}b_{1}\sin((\gamma - \eta)\pi) - a_{1}b_{2}\rho^{\frac{1+\alpha+\gamma}{2}}\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{2} + a_{1}b_{3}\rho^{1+\alpha+\gamma}\sin((\alpha + \eta)\pi) + a_{2}b_{1}\rho^{\frac{1+\alpha+\gamma}{2}}\sin\frac{(1-\alpha-3\gamma+2\eta)\pi}{2} + a_{2}b_{2}\rho^{1+\alpha+\gamma}\sin((\gamma - \eta)\pi) - a_{2}b_{3}\rho^{3\frac{1+\alpha+\gamma}{2}}\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{2} - a_{3}b_{1}\rho^{1+\alpha+\gamma}\sin((\alpha + 2\gamma - \eta)\pi) + a_{3}b_{2}\rho^{3\frac{1+\alpha+\gamma}{2}}\sin\frac{(1-\alpha-3\gamma+2\eta)\pi}{2} + a_{3}b_{3}\rho^{2(1+\alpha+\gamma)}\sin((\gamma - \eta)\pi),$$
(99)

so that the non-negativity requirement of function K implies

$$-a_{1}b_{2}\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{2} + a_{2}b_{1}\sin\frac{(1-\alpha-3\gamma+2\eta)\pi}{2} \ge 0,$$
(100)

$$-a_{2}b_{3}\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{2} + a_{3}b_{2}\sin\frac{(1-\alpha-3\gamma+2\eta)\pi}{2} \ge 0,$$
(101)

$$a_{1}b_{3}\sin((\alpha+\eta)\pi) + a_{2}b_{2}\sin((\gamma-\eta)\pi) - a_{3}b_{1}\sin((\alpha+2\gamma-\eta)\pi) \ge 0.$$
(102)

According to the thermodynamical requirements (93), the sine in the first term in (100), as well as in (101), has a non-negative value, since $1 + \alpha - \gamma + 2\eta \ge (1 - (\gamma - \eta)) + \alpha + \eta \ge 0$ and $1 + \alpha - \gamma + 2\eta = 1 + (\alpha + 3\gamma - 2\eta) - 4(\gamma - \eta) \le 2 - 4(\gamma - \eta) \le 2$, while the argument $\frac{(1 - \alpha - 3\gamma + 2\eta)\pi}{2}$ in second term in (100) and (101) is in the interval $(0, \frac{\pi}{2})$, since $\alpha + 3\gamma - 2\eta \in (0, 1)$, according to (93)₂, so that the requirements (100) and (101) are transformed into

$$\frac{a_1}{b_1} \frac{\sin \frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\cos \frac{(1-\alpha-3\gamma+2\eta)\pi}{4}} \frac{\cos \frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\sin \frac{(1-\alpha-3\gamma+2\eta)\pi}{4}} \leqslant \frac{a_2}{b_2} \leqslant \frac{a_3}{b_3} \frac{\cos \frac{(1-\alpha-3\gamma+2\eta)\pi}{4}}{\sin \frac{(1+\alpha-\gamma+2\eta)\pi}{4}} \frac{\sin \frac{(1-\alpha-3\gamma+2\eta)\pi}{4}}{\cos \frac{(1+\alpha-\gamma+2\eta)\pi}{4}},$$

and represent narrowed thermodynamical requirements $(94)_1$ and $(95)_2$, when combined with the thermodynamical restriction (189), since $\frac{\cos\left(1+\alpha-\gamma+2\eta\right)\pi}{\sin\left(\frac{1-\alpha-3\gamma+2\eta\right)\pi}{4}\right)} = \frac{\cos\left(1+\alpha-\gamma+2\eta\right)\pi}{\cos\left(\frac{1+\alpha+3\gamma-2\eta\right)\pi}{4}\right)} \ge 1$ and $\frac{\sin\left(\frac{1-\alpha-3\gamma+2\eta\right)\pi}{4}\right)}{\cos\left(\frac{1+\alpha-\gamma+2\eta\right)\pi}{4}} = \frac{\cos\left(1+\alpha+3\gamma-2\eta\right)\pi}{\cos\left(\frac{1+\alpha-\gamma+2\eta\right)\pi}{4}\right)} \le 1$, due to $1 + \alpha - \gamma + 2\eta \le 1 + \alpha + 3\gamma - 2\eta$ reducing to $\eta \le \gamma$, see $(94)_1$.

The requirements (93) ensure non-negative values of sines in the first and third term in (102), since $0 \leq \alpha + \eta \leq \alpha + \gamma + 2(\gamma - \eta) \leq 1$ and $0 \leq \alpha + 2\gamma - \eta = (\alpha + 3\gamma - 2\eta) - (\gamma - \eta) \leq 1$, while sine in the second term in (102) has a non-negative value according to the requirement (93)₁, so that the requirement (102) is transformed into the narrowed thermodynamical restriction (98) if the right-hand-side of (98) is less than the right-hand-side of (97), since $\frac{\sin \frac{(\gamma - \eta)\pi}{2}}{\cos \frac{(\alpha + 2\gamma - \eta)\pi}{2}} = \frac{\cos \frac{(1 - (\gamma - \eta))\pi}{2}}{\cos \frac{(\alpha + 2\gamma - \eta)\pi}{2}} \leq 1$ and $\frac{\cos \frac{(\alpha + 2\eta)\pi}{2}}{\cos \frac{(\alpha + 2\gamma - \eta)\pi}{2}} \geq 1$, due to $\alpha + 2\gamma - \eta \leq 1 - (\gamma - \eta) < 1$ and $\alpha + \eta \leq \alpha + 2\gamma - \eta < 1$, respectively reducing to $\alpha + 3\gamma - 2\eta \leq 1$ and $\eta \leq \gamma$, see (93).

In conclusion, the non-negativity of function K, corresponding to the model $I^{+}ID.IDD^{+}$ and given by (99), is ensured by requesting (94), (95) and possibly (98) in addition to (93), (96) and (97) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function.

4.2 Asymmetric models

4.2.1 Model IID.ID

Function K, given by (20), in the case of model IID.ID, having the constitutive equation given in the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha+\beta-\gamma}+a_{2\,0}\mathbf{I}_{t}^{\nu}+a_{3\,0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{I}_{t}^{\alpha}+b_{2\,0}\mathbf{D}_{t}^{\beta}\right)\varepsilon\left(t\right)$$

and with corresponding thermodynamical, see (192) and (193), as well as with the narrowed thermodynamical requirements

$$0 \leqslant \alpha \leqslant \nu < \alpha + \beta - \gamma \leqslant 1, \quad \beta + \nu \leqslant 1, \tag{103}$$

$$\frac{b_1}{b_2} \leqslant \frac{a_1}{a_3} \frac{\sin\frac{(\alpha+2\beta-\gamma)\pi}{2}}{\sin\frac{(\alpha+\gamma)\pi}{2}} \frac{\cos\frac{(\alpha+2\beta-\gamma)\pi}{2}}{\cos\frac{(\alpha+\gamma)\pi}{2}} \leqslant \frac{a_1}{a_3} \frac{\sin\frac{(\alpha+2\beta-\gamma)\pi}{2}}{\sin\frac{(\alpha+\gamma)\pi}{2}},\tag{104}$$

valid if $\alpha + 2\beta - \gamma < 1$, becomes

$$K(\rho) = a_1 b_1 \sin\left(\left(\beta - \gamma\right)\pi\right) + a_1 b_2 \rho^{\alpha+\beta} \sin\left(\left(\alpha + 2\beta - \gamma\right)\pi\right) + a_2 b_1 \rho^{\alpha+\beta-\gamma-\nu} \sin\left(\left(\nu - \alpha\right)\pi\right) + a_2 b_2 \rho^{2(\alpha+\beta)-\gamma-\nu} \sin\left(\left(\beta + \nu\right)\pi\right) - a_3 b_1 \rho^{\alpha+\beta} \sin\left(\left(\alpha + \gamma\right)\pi\right) + a_3 b_2 \rho^{2(\alpha+\beta)} \sin\left(\left(\beta - \gamma\right)\pi\right),$$
(105)

so that the non-negativity requirement of function K implies

$$a_1 b_2 \sin\left(\left(\alpha + 2\beta - \gamma\right)\pi\right) - a_3 b_1 \sin\left(\left(\alpha + \gamma\right)\pi\right) \ge 0.$$
(106)

The first term in (106) has either a positive value if $\alpha + 2\beta - \gamma < 1$, or a non-positive value if $\alpha + 2\beta - \gamma \ge 1$, since $\alpha + 2\beta - \gamma = (\alpha + \beta - \gamma) + \beta \in (0, 2)$, according to the thermodynamical requirement (103)₁, while the second term in (106) has a non-positive value, since the thermodynamical requirement (103)₁ gives $\alpha + \gamma \le \nu + \gamma < \alpha + \beta$ implying $\gamma < \beta$, that combined with $\alpha \le \nu$, see (103)₁, yields $\alpha + \gamma \le \beta + \nu \le 1$ according to the thermodynamical requirement (103)₂, so that the non-negativity of requirement (106), as well as of the function K, given by (105), can be guaranteed only if $\alpha + 2\beta - \gamma < 1$, and than the requirement (106) is transformed into

$$\frac{b_1}{b_2} \leqslant \frac{a_1}{a_3} \frac{\sin \frac{(\alpha + 2\beta - \gamma)\pi}{2}}{\sin \frac{(\alpha + \gamma)\pi}{2}} \frac{\cos \frac{(\alpha + 2\beta - \gamma)\pi}{2}}{\cos \frac{(\alpha + \gamma)\pi}{2}},$$

representing the narrowed thermodynamical restriction $(104)_2$, since $\alpha + \gamma \leq \alpha + 2\beta - \gamma < 1$ reducing to $\gamma < \beta$, see $(103)_1$, implies $\frac{\cos \frac{(\alpha+2\beta-\gamma)\pi}{2}}{\cos \frac{(\alpha+\gamma)\pi}{2}} \leq 1$.

In conclusion, the non-negativity of function K, corresponding to the model IID.ID and given by (105), is ensured if $\alpha + 2\beta - \gamma < 1$ by requesting (104), in addition to (103) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function, while if $\alpha + 2\beta - \gamma \ge 1$, then the mentioned properties cannot be ensured, since (106) cannot be satisfied.

4.2.2 Model $IDD.DD^+$

Function K, given by (20), in the case of model IDD.DD^+ , having the constitutive equation given in the form

$$\left(a_{1\ 0}\mathbf{I}_{t}^{\alpha} + a_{2\ 0}\mathbf{D}_{t}^{\beta} + a_{3\ 0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right) = \left(b_{1\ 0}\mathbf{D}_{t}^{\mu} + b_{2\ 0}\mathbf{D}_{t}^{\alpha+\beta+\mu}\right)\varepsilon\left(t\right),$$

and with corresponding thermodynamical, see (194) and (195), as well as with the narrowed thermodynamical requirements

$$1 \leqslant \alpha + \beta + \mu \leqslant 2, \quad \beta < \gamma \leqslant \mu \leqslant 1 - \alpha, \tag{107}$$

$$\frac{a_1}{a_2} \frac{\left| \cos \frac{(2\alpha+\beta+\mu)\pi}{2} \right|}{\cos \frac{(\mu-\beta)\pi}{2}} \leqslant \frac{a_1}{a_2} \frac{\left| \cos \frac{(2\alpha+\beta+\mu)\pi}{2} \right|}{\cos \frac{(\mu-\beta)\pi}{2}} \frac{\sin \frac{(2\alpha+\beta+\mu)\pi}{2}}{\sin \frac{(\mu-\beta)\pi}{2}} \leqslant \frac{b_1}{b_2}, \tag{108}$$

becomes

$$K(\rho) = a_1 b_1 \sin((\alpha + \mu)\pi) + a_1 b_2 \rho^{\alpha + \beta} \sin((2\alpha + \beta + \mu)\pi) + a_2 b_1 \rho^{\alpha + \beta} \sin((\mu - \beta)\pi) + a_2 b_2 \rho^{2(\alpha + \beta)} \sin((\alpha + \mu)\pi) + a_3 b_2 \rho^{2\alpha + \beta + \mu} \sin((\alpha + \beta)\pi),$$
(109)

so that the non-negativity requirement of function K implies

$$a_1 b_2 \sin((2\alpha + \beta + \mu)\pi) + a_2 b_1 \sin((\mu - \beta)\pi) > 0.$$
(110)

According to the thermodynamical requirement $(107)_2$, one finds $2\alpha + \beta + \mu = (\alpha + \beta) + (\alpha + \mu) \leq 2$, as well as according to the thermodynamical requirement $(107)_1$, one has $2\alpha + \beta + \mu = (\alpha + \beta + \mu) + \alpha \geq 1$, so that the first term in (110) has a non-positive value, while the second term in (110) has a non-negative value, according the thermodynamical requirement $(107)_2$. Therefore, the requirement (110) is transformed into

$$\frac{a_1}{a_2} \frac{\left| \cos \frac{(2\alpha+\beta+\mu)\pi}{2} \right|}{\cos \frac{(\mu-\beta)\pi}{2}} \frac{\sin \frac{(2\alpha+\beta+\mu)\pi}{2}}{\sin \frac{(\mu-\beta)\pi}{2}} \leqslant \frac{b_1}{b_2},$$

representing the narrowed thermodynamical restriction $(108)_1$, since $\pi - \frac{(2\alpha + \beta + \mu)\pi}{2} \ge \frac{(\mu - \beta)\pi}{2}$, reducing to $\alpha + \mu \le 1$, see $(107)_2$, implies $\frac{\sin \frac{(2\alpha + \beta + \mu)\pi}{2}}{\sin \frac{(\mu - \beta)\pi}{2}} \ge 1$.

In conclusion, the non-negativity of function K, corresponding to the model IDD.DD⁺ and given by (109), is ensured by requesting (108) in addition to (107) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function.

4.2.3 Model $I^+ID.ID$

Function K, given by (20), in the case of the model I^{\dagger} ID.ID, having the constitutive equation given in the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha+\beta+\nu} + a_{2\,0}\mathbf{I}_{t}^{\nu} + a_{3\,0}\mathbf{D}_{t}^{\alpha+\beta-\nu}\right)\sigma\left(t\right) = \left(b_{1\,0}\mathbf{I}_{t}^{\alpha} + b_{2\,0}\mathbf{D}_{t}^{\beta}\right)\varepsilon\left(t\right),\tag{111}$$

and with corresponding thermodynamical, see (196) and (197), as well as with the narrowed thermodynamical requirements

$$0 \leqslant \alpha + \beta - \nu \leqslant 1, \quad 1 \leqslant \alpha + \beta + \nu \leqslant 2, \quad \alpha \leqslant \nu \leqslant 1 - \beta, \tag{112}$$

$$\frac{a_1}{a_2} \frac{\left| \cos \frac{(\alpha+2\beta+\nu)\pi}{2} \right|}{\cos \frac{(\nu-\alpha)\pi}{2}} \leqslant \frac{a_1}{a_2} \frac{\left| \cos \frac{(\alpha+2\beta+\nu)\pi}{2} \right|}{\cos \frac{(\nu-\alpha)\pi}{2}} \frac{\sin \frac{(\alpha+2\beta+\nu)\pi}{2}}{\sin \frac{(\nu-\alpha)\pi}{2}} \leqslant \frac{b_1}{b_2}, \tag{113}$$

$$\frac{b_1}{b_2} \leqslant \frac{a_2}{a_3} \frac{\sin\frac{(\beta+\nu)\pi}{2}}{\sin\frac{(2\alpha+\beta-\nu)\pi}{2}} \frac{\cos\frac{(\beta+\nu)\pi}{2}}{\cos\frac{(2\alpha+\beta-\nu)\pi}{2}} \leqslant \frac{a_2}{a_3} \frac{\sin\frac{(\beta+\nu)\pi}{2}}{\sin\frac{(2\alpha+\beta-\nu)\pi}{2}},\tag{114}$$

becomes

$$K(\rho) = a_1 b_1 \sin((\beta + \nu) \pi) + a_1 b_2 \rho^{\alpha + \beta} \sin((\alpha + 2\beta + \nu) \pi) + a_2 b_1 \rho^{\alpha + \beta} \sin((\nu - \alpha) \pi) + a_2 b_2 \rho^{2(\alpha + \beta)} \sin((\beta + \nu) \pi) - a_3 b_1 \rho^{2(\alpha + \beta)} \sin((2\alpha + \beta - \nu) \pi) + a_3 b_2 \rho^{3(\alpha + \beta)} \sin((\nu - \alpha) \pi),$$
(115)

so that the non-negativity requirement of function K implies

$$a_1 b_2 \sin((\alpha + 2\beta + \nu)\pi) + a_2 b_1 \sin((\nu - \alpha)\pi) \ge 0, \tag{116}$$

$$a_2 b_2 \sin\left(\left(\beta + \nu\right)\pi\right) - a_3 b_1 \sin\left(\left(2\alpha + \beta - \nu\right)\pi\right) \ge 0. \tag{117}$$

According to the thermodynamical requirement $(112)_3$, one finds $\alpha + 2\beta + \nu = (\alpha + \beta) + (\beta + \nu) \leq 2$, while according to the thermodynamical requirement $(112)_2$ one finds $\alpha + 2\beta + \nu = (\alpha + \beta + \nu) + \beta \geq 1$, so that the first term in (116) has a non-positive value, while the second term in (116) has a non-negative value, according to the thermodynamical requirement $(112)_3$. Therefore, the inequality (116) is transformed into

$$\frac{a_1}{a_2} \frac{\left| \cos \frac{(\alpha+2\beta+\nu)\pi}{2} \right|}{\cos \frac{(\nu-\alpha)\pi}{2}} \frac{\sin \frac{(\alpha+2\beta+\nu)\pi}{2}}{\sin \frac{(\nu-\alpha)\pi}{2}} \leqslant \frac{b_1}{b_2}$$

representing the narrowed thermodynamical restriction $(113)_1$, since $\pi - \frac{(\alpha+2\beta+\nu)\pi}{2} \ge \frac{(\nu-\alpha)\pi}{2}$, reducing to $\beta + \nu \le 1$, see $(112)_3$, implies $\frac{\sin\frac{(\alpha+2\beta+\nu)\pi}{2}}{\sin\frac{(\nu-\alpha)\pi}{2}} \ge 1$.

According to the thermodynamical requirement $(112)_3$, the first term in (117) has a non-negative value, while according to the thermodynamical requirement $(112)_1$, one finds $2\alpha + \beta - \nu = (\alpha + \beta - \nu) + \alpha \ge 0$, as well as according to the requirement $(112)_3$, one has $2\alpha + \beta - \nu = (\alpha + \beta) - (\nu - \alpha) \le 1$, and therefore the second term in (117) has a non-positive value, so that the inequality (117) is transformed into

$$\frac{b_1}{b_2} \leqslant \frac{a_2}{a_3} \frac{\sin \frac{(\beta+\nu)\pi}{2}}{\sin \frac{(2\alpha+\beta-\nu)\pi}{2}} \frac{\cos \frac{(\beta+\nu)\pi}{2}}{\cos \frac{(2\alpha+\beta-\nu)\pi}{2}},$$

representing the narrowed thermodynamical restriction $(114)_2$, since $2\alpha + \beta - \nu \leq \beta + \nu < 1$, reducing to $\alpha \leq \nu$, see $(112)_3$, implies $\frac{\cos \frac{(\beta + \nu)\pi}{2}}{\cos \frac{(2\alpha + \beta - \nu)\pi}{2}} \leq 1$.

In conclusion, the non-negativity of function K, corresponding to the model I^+ ID.ID and given by (115), is ensured by requesting (113) and (114) in addition to (112) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function.

4.2.4 Model IDD^+ .DD⁺

Function K, given by (20), in the case of model IDD^+ , DD^+ , having the constitutive equation given in the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\beta}+a_{3\,0}\mathbf{D}_{t}^{\alpha+2\beta}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{D}_{t}^{\mu}+b_{2\,0}\mathbf{D}_{t}^{\alpha+\beta+\mu}\right)\varepsilon\left(t\right),$$

and with corresponding thermodynamical, see (198) and (199), as well as with the narrowed thermodynamical requirements

$$1 \leqslant \alpha + 2\beta \leqslant 2, \quad 1 \leqslant \alpha + \beta + \mu \leqslant 2, \quad \beta \leqslant \mu \leqslant 1 - \alpha, \tag{118}$$

$$\frac{a_1}{a_2} \frac{\left|\cos\frac{(2\alpha+\beta+\mu)\pi}{2}\right|}{\cos\frac{(\mu-\beta)\pi}{2}} \leqslant \frac{a_1}{a_2} \frac{\left|\cos\frac{(2\alpha+\beta+\mu)\pi}{2}\right|}{\cos\frac{(\mu-\beta)\pi}{2}} \frac{\sin\frac{(2\alpha+\beta+\mu)\pi}{2}}{\sin\frac{(\mu-\beta)\pi}{2}} \leqslant \frac{b_1}{b_2},\tag{119}$$

$$\frac{b_1}{b_2} \leqslant \frac{a_2}{a_3} \frac{\sin\frac{(\alpha+\mu)\pi}{2}}{\sin\frac{(\alpha+2\beta-\mu)\pi}{2}} \frac{\cos\frac{(\alpha+\mu)\pi}{2}}{\cos\frac{(\alpha+2\beta-\mu)\pi}{2}} \leqslant \frac{a_2}{a_3} \frac{\sin\frac{(\alpha+\mu)\pi}{2}}{\sin\frac{(\alpha+2\beta-\mu)\pi}{2}},\tag{120}$$

valid if $2\alpha + \beta - \mu < 1$, becomes

$$K(\rho) = a_1 b_1 \sin((\alpha + \mu) \pi) + a_1 b_2 \rho^{\alpha + \beta} \sin((2\alpha + \beta + \mu) \pi) + a_2 b_1 \rho^{\alpha + \beta} \sin((\mu - \beta) \pi) + a_2 b_2 \rho^{2(\alpha + \beta)} \sin((\alpha + \mu) \pi) - a_3 b_1 \rho^{2(\alpha + \beta)} \sin((\alpha + 2\beta - \mu) \pi) + a_3 b_2 \rho^{3(\alpha + \beta)} \sin((\mu - \beta) \pi),$$
(121)

so that the non-negativity requirement of function K implies

$$a_{1}b_{2}\sin((2\alpha + \beta + \mu)\pi) + a_{2}b_{1}\sin((\mu - \beta)\pi) \ge 0,$$
(122)

$$a_2 b_2 \sin\left(\left(\alpha + \mu\right)\pi\right) - a_3 b_1 \sin\left(\left(\alpha + 2\beta - \mu\right)\pi\right) \ge 0.$$
(123)

According to the thermodynamical requirement $(118)_3$, one finds $\alpha + 2\beta + \mu = (\alpha + \beta) + (\beta + \mu) \leq 2$, while according to the thermodynamical requirement $(118)_2$, one finds $\alpha + 2\beta + \mu = (\alpha + \beta + \mu) + \beta \geq 1$, so that the first term in (122) has a non-positive value, while the second term in (122) has a non-negative value according to the thermodynamical requirement $(118)_3$. Therefore, the inequality (122) is transformed into

$$\frac{a_1}{a_2} \frac{\left| \cos \frac{(2\alpha + \beta + \mu)\pi}{2} \right|}{\cos \frac{(\mu - \beta)\pi}{2}} \frac{\sin \frac{(2\alpha + \beta + \mu)\pi}{2}}{\sin \frac{(\mu - \beta)\pi}{2}} \leqslant \frac{b_1}{b_2}$$

representing the narrowed thermodynamical restriction $(119)_1$, since $\pi - \frac{(2\alpha + \beta + \mu)\pi}{2} \ge \frac{(\mu - \beta)\pi}{2}$, reducing to $\alpha + \mu \le 1$, see $(118)_3$, implies $\frac{\sin \frac{(2\alpha + \beta + \mu)\pi}{2}}{\sin \frac{(\mu - \beta)\pi}{2}} \ge 1$.

According to the thermodynamical requirement $(118)_3$, the first term in (123) has a non-negative value, while according to the thermodynamical requirement $(118)_1$, one finds $\alpha + 2\beta - \mu = (\alpha + 2\beta) - \mu \ge 0$, as well as according to the requirement $(118)_3$, one finds $\alpha + 2\beta - \mu = (\alpha + \beta) - (\mu - \beta) \le 1$, and therefore the second term in (123) has a non-positive value, so that the inequality (123) is transformed into

$$\frac{b_1}{b_2} \leqslant \frac{a_2}{a_3} \frac{\sin\frac{(\alpha+\mu)\pi}{2}}{\sin\frac{(\alpha+2\beta-\mu)\pi}{2}} \frac{\cos\frac{(\alpha+\mu)\pi}{2}}{\cos\frac{(\alpha+2\beta-\mu)\pi}{2}}$$

representing the narrowed thermodynamical restriction $(120)_2$, since $\alpha + 2\beta - \mu \leq \alpha + \mu < 1$, reducing to $\beta \leq \mu$, see $(118)_3$, implies $\frac{\cos \frac{(\alpha + \mu)\pi}{2}}{\cos \frac{(\alpha + 2\beta - \mu)\pi}{2}} \leq 1$.

In conclusion, the non-negativity of function K, corresponding to the model IDD^+ .DD⁺ and given by (121), is ensured by requesting (119) and (120) in addition to (118), implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function.

4.2.5 Model ID.IDD

Function K, given by (20), in the case of the model ID.IDD, having the constitutive equation given in the form

$$(a_{1\,0}\mathbf{I}_{t}^{\mu} + a_{2\,0}\mathbf{D}_{t}^{\nu})\,\sigma(t) = \left(b_{1\,0}\mathbf{I}_{t}^{\alpha} + b_{2\,0}\mathbf{D}_{t}^{\beta} + b_{3\,0}\mathbf{D}_{t}^{\mu+\nu-\alpha}\right)\varepsilon(t)$$

and with corresponding thermodynamical, see (200) and (201), as well as with the narrowed thermodynamical requirements

$$0 \leqslant \nu \leqslant \beta < \mu + \nu - \alpha \leqslant 1, \quad \alpha \leqslant \mu \leqslant 1 - \beta, \tag{124}$$

$$-\frac{a_1}{a_2} \frac{\cos\frac{(2\alpha+\beta-\mu)\pi}{2}}{\cos\frac{(\beta+\mu)\pi}{2}} \leqslant \frac{b_1}{b_3} \leqslant \frac{a_1}{a_2} \frac{\sin\frac{(2\mu+\nu-\alpha)\pi}{2}}{\sin\frac{(\alpha+\nu)\pi}{2}} \frac{\cos\frac{(2\mu+\nu-\alpha)\pi}{2}}{\cos\frac{(\alpha+\nu)\pi}{2}} \leqslant \frac{a_1}{a_2} \frac{\sin\frac{(2\alpha+\beta-\mu)\pi}{2}}{\sin\frac{(\beta+\mu)\pi}{2}},$$
(125)

valid if $2\mu + \nu - \alpha < 1$, becomes

$$K(\rho) = a_1 b_1 \sin(((\mu - \alpha)\pi) + a_1 b_2 \rho^{\alpha + \beta} \sin((\beta + \mu)\pi) + a_1 b_3 \rho^{\mu + \nu} \sin((2\mu + \nu - \alpha)\pi) - a_2 b_1 \rho^{\mu + \nu} \sin((\alpha + \nu)\pi) + a_2 b_2 \rho^{\alpha + \beta + \mu + \nu} \sin((\beta - \nu)\pi) + a_2 b_3 \rho^{2(\mu + \nu)} \sin((\mu - \alpha)\pi), \quad (126)$$

so that the non-negativity requirement of function K implies

 $a_1 b_3 \sin((2\mu + \nu - \alpha)\pi) - a_2 b_1 \sin((\alpha + \nu)\pi) \ge 0.$ (127)

The first term in (127) has either a positive value if $2\mu + \nu - \alpha < 1$, or non-positive value if $2\mu + \nu - \alpha \ge 1$, since $2\mu + \nu - \alpha = (\mu + \nu - \alpha) + \mu \in (0, 2)$ according to the thermodynamical requirement (124)₁, while the second term in (127) has a non-positive value, since $\alpha + \nu \le \alpha + \beta \le 1$, according to the thermodynamical requirements (124). Therefore, the inequality (127) can be satisfied only if $2\mu + \nu - \alpha < 1$ and it is transformed into

$$\frac{b_1}{b_3} \leqslant \frac{a_1}{a_2} \frac{\sin \frac{(2\mu+\nu-\alpha)\pi}{2}}{\sin \frac{(\alpha+\nu)\pi}{2}} \frac{\cos \frac{(2\mu+\nu-\alpha)\pi}{2}}{\cos \frac{(\alpha+\nu)\pi}{2}},$$

representing the narrowed thermodynamical restriction (125)₃, since $\alpha + \nu \leq 2\mu + \nu - \alpha < 1$, reducing to $\alpha \leq \mu$, see (124)₂, implies $\frac{\cos \frac{(2\mu+\nu-\alpha)\pi}{2}}{\cos \frac{(\alpha+\nu)\pi}{2}} \leq 1$.

In conclusion, the non-negativity of function K, corresponding to the model ID.IDD and given by (126), is ensured if $2\mu + \nu - \alpha < 1$ by requesting (125) in addition to (124) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function, while if $2\mu + \nu - \alpha \ge 1$, then the mentioned properties cannot be ensured, since (127) cannot be satisfied.

4.2.6 Model ID.DDD^+

Function K, given by (20), in the case of the model ID.DDD^+ , having the constitutive equation given in the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\beta}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{D}_{t}^{\mu}+b_{2\,0}\mathbf{D}_{t}^{\nu}+b_{3\,0}\mathbf{D}_{t}^{\alpha+\beta+\nu}\right)\varepsilon\left(t\right),$$

and with corresponding thermodynamical, see (202) and (203), as well as with the narrowed thermodynamical requirements

$$1 \leqslant \alpha + \beta + \nu \leqslant 2, \quad \beta \leqslant \mu < \nu \leqslant 1 - \alpha, \tag{128}$$

$$\frac{a_1}{a_2} \frac{\left|\cos\frac{(2\alpha+\beta+\nu)\pi}{2}\right|}{\cos\frac{(\nu-\beta)\pi}{2}} \leqslant \frac{a_1}{a_2} \frac{\left|\cos\frac{(2\alpha+\beta+\nu)\pi}{2}\right|}{\cos\frac{(\nu-\beta)\pi}{2}} \frac{\sin\frac{(2\alpha+\beta+\nu)\pi}{2}}{\sin\frac{(\nu-\beta)\pi}{2}} \leqslant \frac{b_2}{b_3},\tag{129}$$

becomes

$$K(\rho) = a_1 b_1 \sin((\alpha + \mu) \pi) + a_1 b_2 \rho^{\nu - \mu} \sin((\alpha + \nu) \pi) + a_1 b_3 \rho^{\alpha + \beta + \nu - \mu} \sin((2\alpha + \beta + \nu) \pi) + a_2 b_1 \rho^{\alpha + \beta} \sin((\mu - \beta) \pi) + a_2 b_2 \rho^{\alpha + \beta + \nu - \mu} \sin((\nu - \beta) \pi) + a_2 b_3 \rho^{2\alpha + 2\beta + \nu - \mu} \sin((\alpha + \nu) \pi), \quad (130)$$

so that the non-negativity requirement of function K implies

$$a_1 b_3 \sin\left(\left(2\alpha + \beta + \nu\right)\pi\right) + a_2 b_2 \sin\left(\left(\nu - \beta\right)\pi\right) \ge 0.$$
(131)

The first term in (131) has a non-positive value, since $2\alpha + \beta + \nu = (\alpha + \beta + \nu) + \alpha \ge 1$, according to the thermodynamical requirements $(128)_1$, as well as $2\alpha + \beta + \nu = (\alpha + \beta) + (\alpha + \nu) \le 2$ according to the thermodynamical requirements $(128)_2$, while the second term in (131) has a non-negative value, according to the thermodynamical requirement $(128)_2$. Therefore, the inequality (131) is transformed into

$$\frac{a_1}{a_2} \frac{\left|\cos\frac{(2\alpha+\beta+\nu)\pi}{2}\right|}{\cos\frac{(\nu-\beta)\pi}{2}} \frac{\sin\frac{(2\alpha+\beta+\nu)\pi}{2}}{\sin\frac{(\nu-\beta)\pi}{2}} \leqslant \frac{b_2}{b_3},$$

representing the narrowed thermodynamical restriction $(129)_1$, since $\frac{(\nu-\beta)\pi}{2} \leq \pi - \frac{(2\alpha+\beta+\nu)\pi}{2}$ reducing to $\alpha+\nu \leq 1$, see $(128)_2$, implies $\frac{\sin\frac{(2\alpha+\beta+\nu)\pi}{2}}{\sin\frac{(\nu-\beta)\pi}{2}} \geq 1$.

In conclusion, the non-negativity of function K, corresponding to the model ID.DDD⁺ and given by (130), is ensured by requesting (129) in addition to (128) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function.

4.2.7 Model ID.IDD^+

Function K, given by (20), in the case of the model ID.IDD^+ , having the constitutive equation given in the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\beta}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{I}_{t}^{\alpha+\beta-\nu}+b_{2\,0}\mathbf{D}_{t}^{\nu}+b_{3\,0}\mathbf{D}_{t}^{\alpha+\beta+\nu}\right)\varepsilon\left(t\right),$$

and with corresponding thermodynamical, see (204) and (205), as well as with the narrowed thermodynamical requirements

$$0 \leqslant \alpha + \beta - \nu \leqslant 1, \quad 1 \leqslant \alpha + \beta + \nu \leqslant 2, \quad \beta \leqslant \nu \leqslant 1 - \alpha, \tag{132}$$

$$\frac{b_1}{b_2} \frac{\sin\frac{(\alpha+2\beta-\nu)\pi}{2}}{\sin\frac{(\alpha+\nu)\pi}{2}} \leqslant \frac{b_1}{b_2} \frac{\sin\frac{(\alpha+2\beta-\nu)\pi}{2}}{\sin\frac{(\alpha+\nu)\pi}{2}} \frac{\cos\frac{(\alpha+2\beta-\nu)\pi}{2}}{\cos\frac{(\alpha+\nu)\pi}{2}} \leqslant \frac{a_1}{a_2},\tag{133}$$

$$\frac{a_1}{a_2} \leqslant \frac{b_2}{b_3} \frac{\cos\frac{(\nu-\beta)\pi}{2}}{\left|\cos\frac{(2\alpha+\beta+\nu)\pi}{2}\right|} \frac{\sin\frac{(\nu-\beta)\pi}{2}}{\sin\frac{(2\alpha+\beta+\nu)\pi}{2}} \leqslant \frac{b_2}{b_3} \frac{\cos\frac{(\nu-\beta)\pi}{2}}{\left|\cos\frac{(2\alpha+\beta+\nu)\pi}{2}\right|},\tag{134}$$

becomes

$$K(\rho) = a_1 b_1 \sin((\nu - \beta)\pi) + a_1 b_2 \rho^{\alpha + \beta} \sin((\alpha + \nu)\pi) + a_1 b_3 \rho^{2(\alpha + \beta)} \sin((2\alpha + \beta + \nu)\pi) - a_2 b_1 \rho^{\alpha + \beta} \sin((\alpha + 2\beta - \nu)\pi) + a_2 b_2 \rho^{2(\alpha + \beta)} \sin((\nu - \beta)\pi) + a_2 b_3 \rho^{3(\alpha + \beta)} \sin((\alpha + \nu)\pi), \quad (135)$$

so that the non-negativity requirement of function K implies

$$a_{1}b_{2}\sin((\alpha+\nu)\pi) - a_{2}b_{1}\sin((\alpha+2\beta-\nu)\pi) \ge 0,$$
(136)

$$a_1 b_3 \sin((2\alpha + \beta + \nu)\pi) + a_2 b_2 \sin((\nu - \beta)\pi) \ge 0.$$
(137)

According to the thermodynamical requirement $(132)_3$, the first term in (136) has a non-negative value, while according to the thermodynamical requirement $(132)_1$, one finds $\alpha + 2\beta - \nu = (\alpha + \beta - \nu) + \beta \ge 0$, as well as according to the requirement $(132)_3$, one has $\alpha + 2\beta - \nu = (\alpha + \beta) - (\nu - \beta) \le 1$, and therefore the second term in (136) has a non-positive value, so that the requirement (136) is transformed into

$$\frac{b_1}{b_2} \frac{\sin \frac{(\alpha+2\beta-\nu)\pi}{2}}{\sin \frac{(\alpha+\nu)\pi}{2}} \frac{\cos \frac{(\alpha+2\beta-\nu)\pi}{2}}{\cos \frac{(\alpha+\nu)\pi}{2}} \leqslant \frac{a_1}{a_2}$$

representing the narrowed thermodynamical restriction (133)₁, since $\alpha + 2\beta - \nu \leq \alpha + \nu < 1$ reducing to $\beta \leq \nu$, see (132)₃, implies $\frac{\cos \frac{(\alpha+2\beta-\nu)\pi}{2}}{\cos \frac{(\alpha+\nu)\pi}{2}} \ge 1$.

The first term in (137) has a non-positive value, since $2\alpha + \beta + \nu = (\alpha + \beta + \nu) + \alpha \ge 1$, according to the thermodynamical requirement (132)₂, as well as $2\alpha + \beta + \nu = (\alpha + \beta) + (\alpha + \nu) \le 2$ according to the thermodynamical requirement (132)₃, while the second term in (137) has a non-negative value, according to the thermodynamical requirement (132)₃. Therefore, the inequality (137) is transformed into

$$\frac{a_1}{a_2} \leqslant \frac{b_2}{b_3} \frac{\cos\frac{(\nu-\beta)\pi}{2}}{\left|\cos\frac{(2\alpha+\beta+\nu)\pi}{2}\right|} \frac{\sin\frac{(\nu-\beta)\pi}{2}}{\sin\frac{(2\alpha+\beta+\nu)\pi}{2}}$$

representing the narrowed thermodynamical restriction $(134)_2$, since $\frac{(\nu-\beta)\pi}{2} \leqslant \pi - \frac{(2\alpha+\beta+\nu)\pi}{2}$ reducing to $\alpha+\nu \leqslant 1$, see $(132)_3$, implies $\frac{\sin\frac{(\nu-\beta)\pi}{2}}{\sin\frac{(2\alpha+\beta+\nu)\pi}{2}} \leqslant 1$.

In conclusion, the non-negativity of function K, corresponding to the model ID.IDD⁺ and given by (135), is ensured by requesting (133) and (134) in addition to (132) implying the corresponding relaxation modulus to be a completely monotonic function and creep compliance a Bernstein function.

5 Numerical examples

Thermodynamical and narrowed thermodynamical restrictions on the parameters for the model I⁺ID.ID, taking the form

$$\left(a_{1\ 0}\mathbf{I}_{t}^{\alpha+\beta+\nu} + a_{2\ 0}\mathbf{I}_{t}^{\nu} + a_{3\ 0}\mathbf{D}_{t}^{\alpha+\beta-\nu}\right)\sigma(t) = \left(b_{1\ 0}\mathbf{I}_{t}^{\alpha} + b_{2\ 0}\mathbf{D}_{t}^{\beta}\right)\varepsilon(t),$$
(138)

are given as

$$0 \leqslant \alpha + \beta - \nu \leqslant 1, \quad 1 \leqslant \alpha + \beta + \nu \leqslant 2, \quad \alpha \leqslant \nu \leqslant 1 - \beta, \tag{139}$$

$$\frac{a_1}{a_2} \frac{\left|\cos\frac{(\alpha+2\beta+\nu)\pi}{2}\right|}{\cos\frac{(\nu-\alpha)\pi}{2}} \leqslant \frac{a_1}{a_2} \frac{\sin\frac{(\alpha+2\beta+\nu)\pi}{2}}{\sin\frac{(\nu-\alpha)\pi}{2}} \frac{\left|\cos\frac{(\alpha+2\beta+\nu)\pi}{2}\right|}{\cos\frac{(\nu-\alpha)\pi}{2}} \leqslant \frac{b_1}{b_2},\tag{140}$$

$$\frac{b_1}{b_2} \leqslant \frac{a_2}{a_3} \frac{\sin\frac{(\beta+\nu)\pi}{2}}{\sin\frac{(2\alpha+\beta-\nu)\pi}{2}} \frac{\cos\frac{(\beta+\nu)\pi}{2}}{\cos\frac{(2\alpha+\beta-\nu)\pi}{2}} \leqslant \frac{a_2}{a_3} \frac{\sin\frac{(\beta+\nu)\pi}{2}}{\sin\frac{(2\alpha+\beta-\nu)\pi}{2}},\tag{141}$$

see also Section 4.2.3 and Appendix A.2.

5.1 Stress relaxation for the model I⁺ID.ID

The relaxation modulus for the model I⁺ID.ID, given by (138), takes the form

$$\tilde{\sigma}_{sr}(s) = \frac{1}{s^{1-(\beta+\nu)}} \frac{\phi_{\varepsilon}(s)}{\phi_{\sigma}(s)} = \frac{1}{s^{1-(\beta+\nu)}} \frac{b_1 + b_2 s^{\alpha+\beta}}{a_1 + a_2 s^{\alpha+\beta} + a_3 s^{2(\alpha+\beta)}}$$
(142)

in the Laplace domain, after applying the Laplace transform to the constitutive equation (138), where displacement ε is prescribed as a Heaviside step function, see also (2) and Table 1, so that, after applying the inverse Laplace transform to (142), according to (15), one obtains the relaxation modulus for the model I⁺ID.ID in the time domain as

$$\sigma_{sr}(t) = \sigma_{sr}^{(\text{NP})}(t) + \begin{cases} 0, & \text{if } \tilde{\sigma}_{sr} \text{ has no poles,} \\ \sigma_{sr}^{(\text{RP})}(t), & \text{if } \tilde{\sigma}_{sr} \text{ has a negative real pole,} \\ \sigma_{sr}^{(\text{CCP})}(t), & \text{if } \tilde{\sigma}_{sr} \text{ has a pair of complex conjugated poles,} \end{cases}$$
(143)

where functions $\sigma_{sr}^{(\rm NP)},\,\sigma_{sr}^{(\rm RP)},$ and $\sigma_{sr}^{(\rm CCP)}$ are given by

$$\begin{split} \sigma_{sr}^{(\mathrm{NP})}\left(t\right) &= \frac{1}{\pi} \int_{0}^{\infty} \frac{1}{\rho^{1-(\beta+\nu)}} \frac{\left|b_{1}+b_{2}\rho^{\alpha+\beta}\mathrm{e}^{\mathrm{i}(\alpha+\beta)\pi}\right|}{\left|a_{1}+a_{2}\rho^{\alpha+\beta}\mathrm{e}^{\mathrm{i}(\alpha+\beta)\pi}+a_{3}\rho^{2(\alpha+\beta)}\mathrm{e}^{2\mathrm{i}(\alpha+\beta)\pi}\right|} \\ &\times \sin\left(\left(\beta+\nu\right)\pi + \arctan\frac{b_{2}\rho^{\alpha+\beta}\sin\left(\left(\alpha+\beta\right)\pi\right)}{b_{1}+b_{2}\rho^{\alpha+\beta}\cos\left(\left(\alpha+\beta\right)\pi\right)} \\ &-\arctan\frac{a_{2}\rho^{\alpha+\beta}\sin\left(\left(\alpha+\beta\right)\pi\right)+a_{3}\rho^{2(\alpha+\beta)}\sin\left(2\left(\alpha+\beta\right)\pi\right)}{a_{1}+a_{2}\rho^{\alpha+\beta}\cos\left(\left(\alpha+\beta\right)\pi\right)+a_{3}\rho^{2(\alpha+\beta)}\cos\left(2\left(\alpha+\beta\right)\pi\right)}\right)} e^{-\rho t}\mathrm{d}\rho, \end{split}$$
(144)
$$\sigma_{sr}^{(\mathrm{NP})}\left(t\right) &= \frac{1}{\pi} \int_{0}^{\infty} \frac{1}{\rho^{1-(\beta+\nu)}} \frac{K\left(\rho\right)}{\left|a_{1}+a_{2}\rho^{\alpha+\beta}\mathrm{e}^{\mathrm{i}(\alpha+\beta)\pi}+a_{3}\rho^{2(\alpha+\beta)}\mathrm{e}^{2\mathrm{i}(\alpha+\beta)\pi}\right|^{2}} e^{-\rho t}\mathrm{d}\rho, \end{aligned} \\ \sigma_{sr}^{(\mathrm{RP})}\left(t\right) &= -\rho_{\mathrm{RP}}^{\beta-\mu} \frac{\left|b_{1}+b_{2}\rho_{\mathrm{RP}}^{\alpha+\beta}\mathrm{e}^{\mathrm{i}(\alpha+\beta)\pi}\right|}{\left(\alpha+\beta\right)\left|a_{2}+2a_{3}\rho_{\mathrm{RP}}^{\alpha+\beta}\mathrm{e}^{\mathrm{i}(\alpha+\beta)\pi}\right|} \\ &\times \cos\left(\left(\beta+\nu\right)\pi + \arctan\frac{b_{2}\rho_{\mathrm{RP}}^{\alpha+\beta}\sin\left(2\left(\alpha+\beta\right)\pi\right)}{b_{1}+b_{2}\rho_{\mathrm{RP}}^{\alpha+\beta}\cos\left(\left(\alpha+\beta\right)\pi\right)}\right)} \\ &-\arctan\frac{a_{2}\sin\left(\left(\alpha+\beta\right)\pi\right)+2a_{3}\rho_{\mathrm{RP}}^{\alpha+\beta}\sin\left(2\left(\alpha+\beta\right)\pi\right)}{\left(\alpha+\beta\right)\left|a_{2}+2a_{3}\rho_{\mathrm{CP}}^{\alpha+\beta}\mathrm{e}^{\mathrm{i}(\alpha+\beta)\varphi_{\mathrm{CP}}}\right|} \\ &\times \mathrm{e}^{-|\mathrm{Re}\operatorname{sccr}|t}\cos\left(\mathrm{Im}\operatorname{sccp}t-\left(1-\left(\beta+\nu\right)\right)\varphi_{\mathrm{CCP}}+\arctan\frac{b_{2}\rho_{\mathrm{CCP}}^{\alpha+\beta}\sin\left(\left(\alpha+\beta\right)\varphi_{\mathrm{CCP}}\right)}{b_{1}+b_{2}\rho_{\mathrm{CCP}}^{\alpha+\beta}\cos\left(\left(1-2\left(\alpha+\beta\right)\right)\varphi_{\mathrm{CCP}}\right)}\right) \right)$$
(146)

according to (16), (19), (17), and (18) respectively, with

$$\phi_{\varepsilon}(s) = b_1 + b_2 s^{\alpha+\beta} \quad \text{and} \quad \phi_{\sigma}(s) = a_1 + a_2 s^{\alpha+\beta} + a_3 s^{2(\alpha+\beta)}, \tag{147}$$

where the function K, defined by (20), is represented by

$$K(\rho) = a_1 b_1 \sin((\beta + \nu)\pi) + \rho^{\alpha + \beta} (a_1 b_2 \sin((\alpha + 2\beta + \nu)\pi) + a_2 b_1 \sin((\nu - \alpha)\pi))$$

$$+ \rho^{2(\alpha+\beta)} \left(a_2 b_2 \sin \left((\beta+\nu) \,\pi \right) - a_3 b_1 \sin \left((2\alpha+\beta-\nu) \,\pi \right) \right) + \rho^{3(\alpha+\beta)} a_3 b_2 \sin \left((\nu-\alpha) \,\pi \right), \tag{148}$$

and with poles of the relaxation modulus in Laplace domain $\rho_{\rm RP} = -\frac{2(\alpha+\beta)}{a_3} \sqrt{\frac{a_1}{a_3}}$, if $\tan((\alpha+\beta)\pi) = -\sqrt{\frac{4a_1a_3}{a_2^2}-1}$ and $s_{\rm CCP} = \rho_{\rm CCP} e^{i\varphi_{\rm CCP}}$, with $\rho_{\rm CCP} = \frac{2(\alpha+\beta)}{\sqrt{\frac{a_1}{a_3}}}$ and $\varphi_{\rm CCP} = \left(1 - \frac{1}{\pi} \arctan\sqrt{\frac{4a_1a_3}{a_2^2}-1}\right) \frac{\pi}{\alpha+\beta}$, if $\tan((\alpha+\beta)\pi) > -\sqrt{\frac{4a_1a_3}{a_2^2}-1}$, obtained in Section 6.3 as zeros of function ϕ_{σ} , given by (147)₂, by (166) and (167).

Considering the model I⁺ID.ID, the asymptotic behavior of the corresponding relaxation modulus in the Laplace domain, according to (142), takes the form

$$\begin{split} \tilde{\sigma}_{sr}\left(s\right) &= \frac{b_2}{a_3} \frac{1}{s^{1-(\nu-\alpha)}} \frac{1 + \frac{b_1}{b_2} \frac{1}{s^{\alpha+\beta}}}{1 + \frac{a_2}{a_3} \frac{1}{s^{\alpha+\beta}} + \frac{a_1}{a_3} \frac{1}{s^{2(\alpha+\beta)}}} \\ &= \frac{b_2}{a_3} \frac{1}{s^{1-(\nu-\alpha)}} \left(1 + \frac{b_1}{b_2} \frac{1}{s^{\alpha+\beta}}\right) \left(1 + \frac{a_2}{a_3} \frac{1}{s^{\alpha+\beta}} + \frac{a_1}{a_3} \frac{1}{s^{2(\alpha+\beta)}}\right)^{-1} \\ &= \frac{b_2}{a_3} \frac{1}{s^{1-(\nu-\alpha)}} \left(1 + \frac{b_1}{b_2} \frac{1}{s^{\alpha+\beta}}\right) \left(1 - \frac{a_2}{a_3} \frac{1}{s^{\alpha+\beta}} + \left(-\frac{a_1}{a_3} + \left(\frac{a_2}{a_3}\right)^2\right) \frac{1}{s^{2(\alpha+\beta)}} + O\left(s^{-3(\alpha+\beta)}\right)\right) \\ &= \frac{b_2}{a_3} \frac{1}{s^{1-(\nu-\alpha)}} \left(1 + \left(\frac{b_1}{b_2} - \frac{a_2}{a_3}\right) \frac{1}{s^{\alpha+\beta}} + \left(\left(\frac{a_2}{a_3}\right)^2 - \frac{a_1}{a_3} - \frac{a_2}{a_3} \frac{b_1}{b_2}\right) \frac{1}{s^{2(\alpha+\beta)}} + O\left(s^{-3(\alpha+\beta)}\right)\right) \\ &= \frac{b_2}{a_3} \frac{1}{s^{1-(\nu-\alpha)}} + \frac{b_2}{a_3} \left(\frac{b_1}{b_2} - \frac{a_2}{a_3}\right) \frac{1}{s^{1+2\alpha+\beta-\nu}} \\ &+ \frac{b_2}{a_3} \left(\left(\frac{a_2}{a_3}\right)^2 - \frac{a_1}{a_3} - \frac{a_2}{a_3} \frac{b_1}{b_2}\right) \frac{1}{s^{1+3\alpha+2\beta-\nu}} + O\left(s^{-(1+4\alpha+3\beta-\nu)}\right), \quad \text{when} \quad s \to \infty, \end{split}$$

that implies the short time asymptotics of the relaxation modulus as

$$\sigma_{sr}(t) = \frac{b_2}{a_3} \frac{t^{-(\nu-\alpha)}}{\Gamma(1-(\nu-\alpha))} + \frac{b_2}{a_3} \left(\frac{b_1}{b_2} - \frac{a_2}{a_3}\right) \frac{t^{2\alpha+\beta-\nu}}{\Gamma(1+2\alpha+\beta-\nu)} + \frac{b_2}{a_3} \left(\left(\frac{a_2}{a_3}\right)^2 - \frac{a_1}{a_3} - \frac{a_2}{a_3}\frac{b_1}{b_2}\right) \frac{t^{3\alpha+2\beta-\nu}}{\Gamma(1+3\alpha+2\beta-\nu)} + O\left(t^{-(4\alpha+3\beta-\nu)}\right), \quad \text{when} \quad t \to 0,$$
(149)

according to the theorem that if $\tilde{f}(s) \sim \tilde{g}(s)$ as $s \to \infty$, then $f(t) \sim g(t)$ as $t \to 0$, while the large time asymptotics of the relaxation modulus

$$\sigma_{sr}\left(t\right) = \frac{b_1}{a_1} \frac{t^{-(\beta+\nu)}}{\Gamma\left(1-(\beta+\nu)\right)} + O\left(t^{-(1-\delta)}\right), \quad \text{when} \quad t \to \infty,$$
(150)

follows from the asymptotics of relaxation modulus in the Laplace domain (142), obtained as

$$\begin{split} \tilde{\sigma}_{sr}\left(s\right) &= \frac{b_{1}}{a_{1}} \frac{1}{s^{1-(\beta+\nu)}} \frac{1 + \frac{b_{2}}{b_{1}} s^{\alpha+\beta}}{1 + \frac{a_{2}}{a_{1}} s^{\alpha+\beta} + \frac{a_{3}}{a_{1}} s^{2(\alpha+\beta)}} \\ &= \frac{b_{1}}{a_{1}} \frac{1}{s^{1-(\beta+\nu)}} \left(1 + \frac{b_{2}}{b_{1}} s^{\alpha+\beta}\right) \left(1 + \frac{a_{2}}{a_{1}} s^{\alpha+\beta} + \frac{a_{3}}{a_{1}} s^{2(\alpha+\beta)}\right)^{-1} \\ &= \frac{b_{1}}{a_{1}} \frac{1}{s^{1-(\beta+\nu)}} \left(1 + \frac{b_{2}}{b_{1}} s^{\alpha+\beta}\right) \left(1 - \frac{a_{2}}{a_{1}} s^{\alpha+\beta} + O\left(s^{2(\alpha+\beta)}\right)\right) \\ &= \frac{b_{1}}{a_{1}} \frac{1}{s^{1-(\beta+\nu)}} \left(1 + \left(\frac{b_{2}}{b_{1}} - \frac{a_{2}}{a_{1}}\right) s^{\alpha+\beta} + O\left(s^{2(\alpha+\beta)}\right)\right) \\ &= \frac{b_{1}}{a_{1}} \frac{1}{s^{1-(\beta+\nu)}} + O\left(s^{-\delta}\right), \quad \text{when} \quad s \to 0, \end{split}$$

with $0 < \delta < 1 - (\beta + \nu)$, since $\alpha + 2\beta + \nu > 1$, according to the theorem that if $\tilde{f}(s) \sim \tilde{g}(s)$ as $s \to 0$, then $f(t) \sim g(t)$ as $t \to \infty$.

Figure 3 presents time evolution of the relaxation modulus for different values of model parameters given in Table 4, so that in the case when model parameters are such that the relaxation modulus in Laplace domain does not have poles and additionally satisfy narrowed thermodynamical requirements (139) - (141), the relaxation modulus from Figure 3a, obtained according to (143) with function $\sigma_{sr}^{(NP)}$ given by (144), is completely monotonic, while the complete monotonicity of the relaxation modulus from Figure 3b, obtained according to

(143) with function $\sigma_{sr}^{(\text{RP})}$ given by (145), is not guaranteed, although its form is seemingly such, since model parameters do not satisfy narrowed thermodynamical requirements (139) - (141). Figures 3c and 3d, obtained according to (143) with function $\sigma_{sr}^{(CCP)}$ given by (146), present oscillatory relaxation modulus having amplitude exponentially decreasing in time, displaying the significant damping and the negative stress for a certain time interval, however stress being positive and decreasing to zero for large time. Short and long time behavior of the relaxation modulus, as obvious from all graphs from Figure 3, is governed by the asymptotic expressions (149) and (150).

Case when $\tilde{\sigma}_{sr}$ has	α	β	ν	a_1	a_2	a_3	b_1	b_2
no poles				0.05	1.5	0.45	0.7	0.95
a negative real pole	0.35	0.55	0.4	11	28.4029	20.27	7	9.5
a pair of complex conjugated poles				11	15	20.27	7	9.5

Table 4: Model parameters used for numerical examples.



2

(a) Case when $\tilde{\sigma}_{sr}$ has no poles. Relaxation modulus is completely monotonic.

0.6

0.5

0.4

0.3

0.2

0.1





Figure 3: Time evolution of the relaxation modulus.

5.2Creep compliance for the model I⁺ID.ID

The creep compliance for the model I^+ ID.ID, given by (111), takes the form

$$\tilde{\varepsilon}_{cr}\left(s\right) = \frac{1}{s^{1+\beta+\nu}} \frac{\phi_{\sigma}\left(s\right)}{\phi_{\varepsilon}\left(s\right)} = \frac{1}{s^{1+\beta+\nu}} \frac{a_1 + a_2 s^{\alpha+\beta} + a_3 s^{2(\alpha+\beta)}}{b_1 + b_2 s^{\alpha+\beta}} \tag{151}$$

in the Laplace domain, after applying the Laplace transform to the constitutive equation (138), where stress σ is prescribed as the Heaviside step function, see also (3) and Table 1, so that, after applying the inverse Laplace transform to (151), according to (27), one obtains the creep compliance for the model $I^+ID.ID$ in the time domain as

$$\varepsilon_{cr}(t) = \varepsilon_{cr}^{(\text{NP})}(t) = \frac{1}{\pi} \int_0^\infty \frac{1}{\rho^{1+\beta+\nu}} \frac{\left|a_1 + a_2\rho^{\alpha+\beta} e^{i(\alpha+\beta)\pi} + a_3\rho^{2(\alpha+\beta)} e^{2i(\alpha+\beta)\pi}\right|}{\left|b_1 + b_2\rho^{\alpha+\beta} e^{i(\alpha+\beta)\pi}\right|} \\ \times \sin\left(\left(\beta+\nu\right)\pi + \arctan\frac{b_2\rho^{\alpha+\beta}\sin\left(\left(\alpha+\beta\right)\pi\right)}{b_1 + b_2\rho^{\alpha+\beta}\cos\left(\left(\alpha+\beta\right)\pi\right)} - \arctan\frac{a_2\rho^{\alpha+\beta}\sin\left(\left(\alpha+\beta\right)\pi\right) + a_3\rho^{2(\alpha+\beta)}\sin\left(2\left(\alpha+\beta\right)\pi\right)}{a_1 + a_2\rho^{\alpha+\beta}\cos\left(\left(\alpha+\beta\right)\pi\right) + a_3\rho^{2(\alpha+\beta)}\cos\left(2\left(\alpha+\beta\right)\pi\right)}\right) \left(1 - e^{-\rho t}\right) d\rho, \quad (152)$$

or equivalently as

$$\varepsilon_{cr}\left(t\right) = \varepsilon_{cr}^{(\mathrm{NP})}\left(t\right) = \frac{1}{\pi} \int_{0}^{\infty} \frac{1}{\rho^{1+\beta+\nu}} \frac{K\left(\rho\right)}{\left|b_{1}+b_{2}\rho^{\alpha+\beta}\mathrm{e}^{\mathrm{i}\left(\alpha+\beta\right)\pi}\right|^{2}} \left(1-\mathrm{e}^{-\rho t}\right) \mathrm{d}\rho,\tag{153}$$

where the equivalent forms of function $\varepsilon_{cr}^{(\text{NP})}$ are given by (28) and (31) and where function K is given by (148), since function ϕ_{ε} , given by (147)₁, has no zeros in the complex plane, implying that the creep compliance in Laplace domain admits no poles.

Rather than using any of the expressions (152) and (153), for the purpose of better numerical convergence, one rewrites the creep compliance in Laplace domain (151) as

$$\tilde{\epsilon}_{cr}(s) = s\tilde{\varepsilon}_{cr}(s) = \frac{1}{s^{\beta+\nu}} \left(\frac{a_1}{b_2} \frac{1}{s^{\alpha+\beta} + \frac{b_1}{b_2}} + \frac{a_3}{b_2} s^{\alpha+\beta} \frac{s^{\alpha+\beta} + \frac{a_2}{a_3}}{s^{\alpha+\beta} + \frac{b_1}{b_2}} \right)$$
$$= \frac{a_3}{b_2} \frac{1}{s^{\nu-\alpha}} + \frac{a_3}{b_2} \left(\frac{a_2}{a_3} - \frac{b_1}{b_2} \right) \frac{1}{s^{\nu-\alpha}} \frac{1}{s^{\alpha+\beta} + \frac{b_1}{b_2}} + \frac{a_1}{b_2} \frac{1}{s^{\beta+\nu}} \frac{1}{s^{\alpha+\beta} + \frac{b_1}{b_2}}$$

yielding the creep compliance in time domain in the form

$$\varepsilon_{cr}(t) = \int_{0}^{t} \epsilon_{cr}(t') dt'$$

$$= \frac{a_{3}}{b_{2}} \frac{t^{-(1-(\nu-\alpha))}}{\Gamma(\nu-\alpha)} + \frac{1}{\pi} \frac{a_{3}}{b_{2}} \left(\frac{a_{2}}{a_{3}} - \frac{b_{1}}{b_{2}}\right) \int_{0}^{\infty} \frac{1}{\rho^{1+\nu-\alpha}} \frac{\rho^{\alpha+\beta} \sin\left((\beta+\nu)\pi\right) + \frac{b_{1}}{b_{2}} \sin\left((\nu-\alpha)\pi\right)}{\left|\rho^{\alpha+\beta} e^{i(\alpha+\beta)\pi} + \frac{b_{1}}{b_{2}}\right|^{2}} \left(1 - e^{-\rho t}\right) d\rho$$

$$+ \frac{1}{\pi} \frac{a_{1}}{b_{2}} \int_{0}^{\infty} \frac{1}{\rho^{1+\beta+\nu}} \frac{\rho^{\alpha+\beta} \sin\left((\alpha+2\beta+\nu)\pi\right) + \frac{b_{1}}{b_{2}} \sin\left((\beta+\nu)\pi\right)}{\left|\rho^{\alpha+\beta} e^{i(\alpha+\beta)\pi} + \frac{b_{1}}{b_{2}}\right|^{2}} \left(1 - e^{-\rho t}\right) d\rho, \tag{154}$$

since the definition of inverse Laplace transform (40), used along with the Cauchy integral theorem (41) where the contour is depicted in Figure 1, implies

$$\mathcal{L}^{-1}\left[\frac{1}{s^{\xi}}\frac{1}{s^{\alpha+\beta}+\frac{b_1}{b_2}}\right](t) = \frac{1}{\pi}\int_0^{\infty}\frac{1}{\rho^{\xi}}\frac{\rho^{\alpha+\beta}\sin\left(\left(\alpha+\beta+\xi\right)\pi\right)+\frac{b_1}{b_2}\sin\left(\xi\pi\right)}{\left|\rho^{\alpha+\beta}\mathrm{e}^{\mathrm{i}\left(\alpha+\beta\right)\pi}+\frac{b_1}{b_2}\right|^2}\mathrm{e}^{-\rho t}\mathrm{d}\rho.$$

Equivalently, the creep compliance is also derived in terms of two-parameter Mittag-Leffler function, see (32), that in the case of model I⁺ID.ID yields

$$\varepsilon_{cr}\left(t\right) = \frac{a_1}{b_2} e_{\alpha+\beta,1+\alpha+2\beta+\nu,\frac{b_1}{b_2}}\left(t\right) + \frac{a_2}{b_2} e_{\alpha+\beta,1+\beta+\nu,\frac{b_1}{b_2}}\left(t\right) + \frac{a_3}{b_2} e_{\alpha+\beta,1+\nu-\alpha,\frac{b_1}{b_2}}\left(t\right).$$
(155)

Considering the model $I^+ID.ID$, the asymptotic behavior of the corresponding creep compliance in Laplace domain, according to (151), takes the form

$$\begin{split} \tilde{\varepsilon}_{cr}\left(s\right) &= \frac{a_3}{b_2} \frac{1}{s^{1+\nu-\alpha}} \frac{1 + \frac{a_2}{a_3} \frac{1}{s^{\alpha+\beta}} + \frac{a_1}{a_3} \frac{1}{s^{2(\alpha+\beta)}}}{1 + \frac{b_1}{b_2} \frac{1}{s^{\alpha+\beta}}} \\ &= \frac{a_3}{b_2} \frac{1}{s^{1+\nu-\alpha}} \left(1 + \frac{a_2}{a_3} \frac{1}{s^{\alpha+\beta}} + \frac{a_1}{a_3} \frac{1}{s^{2(\alpha+\beta)}}\right) \left(1 + \frac{b_1}{b_2} \frac{1}{s^{\alpha+\beta}}\right)^{-1} \\ &= \frac{a_3}{b_2} \frac{1}{s^{1+\nu-\alpha}} \left(1 + \frac{a_2}{a_3} \frac{1}{s^{\alpha+\beta}} + \frac{a_1}{a_3} \frac{1}{s^{2(\alpha+\beta)}}\right) \left(1 - \frac{b_1}{b_2} \frac{1}{s^{\alpha+\beta}} + \left(\frac{b_1}{b_2}\right)^2 \frac{1}{s^{2(\alpha+\beta)}} + O\left(s^{-3(\alpha+\beta)}\right)\right) \end{split}$$

$$= \frac{a_3}{b_2} \frac{1}{s^{1+\nu-\alpha}} \left(1 + \left(\frac{a_2}{a_3} - \frac{b_1}{b_2}\right) \frac{1}{s^{\alpha+\beta}} + \left(\frac{a_1}{a_3} - \frac{a_2}{a_3}\frac{b_1}{b_2} + \left(\frac{b_1}{b_2}\right)^2\right) \frac{1}{s^{2(\alpha+\beta)}} + O\left(s^{-3(\alpha+\beta)}\right) \right) = \frac{a_3}{b_2} \frac{1}{s^{1+\nu-\alpha}} + \frac{a_3}{b_2} \left(\frac{a_2}{a_3} - \frac{b_1}{b_2}\right) \frac{1}{s^{1+\beta+\nu}} + \frac{a_3}{b_2} \left(\frac{a_1}{a_3} - \frac{a_2}{a_3}\frac{b_1}{b_2} + \left(\frac{b_1}{b_2}\right)^2\right) \frac{1}{s^{1+\alpha+2\beta+\nu}} + O\left(s^{-(1+2\alpha+3\beta+\nu)}\right), \quad \text{when} \quad s \to \infty,$$

that implies the short time asymptotics of the creep compliance as

$$\varepsilon_{cr}(t) = \frac{a_3}{b_2} \frac{t^{\nu-\alpha}}{\Gamma(1+\nu-\alpha)} + \frac{a_3}{b_2} \left(\frac{a_2}{a_3} - \frac{b_1}{b_2}\right) \frac{t^{\beta+\nu}}{\Gamma(1+\beta+\nu)} + \frac{a_3}{b_2} \left(\frac{a_1}{a_3} - \frac{a_2}{a_3} \frac{b_1}{b_2} + \left(\frac{b_1}{b_2}\right)^2\right) \frac{t^{\alpha+2\beta+\nu}}{\Gamma(1+\alpha+2\beta+\nu)} + O\left(t^{2\alpha+3\beta+\nu}\right), \quad \text{when} \quad t \to 0,$$
(156)

while the large time asymptotics of the creep compliance

$$\varepsilon_{cr}(t) = \frac{a_1}{b_1} \frac{t^{\beta+\nu}}{\Gamma(1+\beta+\nu)} + \frac{a_1}{b_1} \left(\frac{a_2}{a_1} - \frac{b_2}{b_1}\right) \frac{t^{\nu-\alpha}}{\Gamma(1+\nu-\alpha)} + \frac{a_1}{b_1} \left(\frac{a_3}{a_1} - \frac{a_2}{a_1}\frac{b_2}{b_1} + \left(\frac{b_2}{b_1}\right)^2\right) \frac{t^{-2\alpha-\beta+\nu}}{\Gamma(1-2\alpha-\beta+\nu)} + O\left(t^{-\xi}\right), \quad \text{when} \quad t \to \infty,$$
(157)

follows from the asymptotics of creep compliance in Laplace domain, obtained as

$$\begin{split} \tilde{\varepsilon}_{cr}\left(s\right) &= \frac{a_{1}}{b_{1}} \frac{1}{s^{1+\beta+\nu}} \frac{1 + \frac{a_{2}}{a_{1}} s^{\alpha+\beta} + \frac{a_{3}}{a_{2}} s^{2(\alpha+\beta)}}{1 + \frac{b_{2}}{b_{1}} s^{\alpha+\beta}} \\ &= \frac{a_{1}}{b_{1}} \frac{1}{s^{1+\beta+\nu}} \left(1 + \frac{a_{2}}{a_{1}} s^{\alpha+\beta} + \frac{a_{3}}{a_{1}} s^{2(\alpha+\beta)}\right) \left(1 + \frac{b_{2}}{b_{1}} s^{\alpha+\beta}\right)^{-1} \\ &= \frac{a_{1}}{b_{1}} \frac{1}{s^{1+\beta+\nu}} \left(1 + \frac{a_{2}}{a_{1}} s^{\alpha+\beta} + \frac{a_{3}}{a_{1}} s^{2(\alpha+\beta)}\right) \left(1 - \frac{b_{2}}{b_{1}} s^{\alpha+\beta} + \left(\frac{b_{2}}{b_{1}}\right)^{2} s^{2(\alpha+\beta)} + O\left(s^{3(\alpha+\beta)}\right)\right) \\ &= \frac{a_{1}}{b_{1}} \frac{1}{s^{1+\beta+\nu}} \left(1 + \left(\frac{a_{2}}{a_{1}} - \frac{b_{2}}{b_{1}}\right) s^{\alpha+\beta} + \left(\frac{a_{3}}{a_{1}} - \frac{a_{2}}{a_{1}} \frac{b_{2}}{b_{1}} + \left(\frac{b_{2}}{b_{1}}\right)^{2}\right) s^{2(\alpha+\beta)} + O\left(s^{3(\alpha+\beta)}\right) \right) \\ &= \frac{a_{1}}{b_{1}} \frac{1}{s^{1+\beta+\nu}} + \frac{a_{1}}{b_{1}} \left(\frac{a_{2}}{a_{1}} - \frac{b_{2}}{b_{1}}\right) \frac{1}{s^{1+\nu-\alpha}} \\ &+ \frac{a_{1}}{b_{1}} \left(\frac{a_{3}}{a_{1}} - \frac{a_{2}}{a_{1}} \frac{b_{2}}{b_{1}} + \left(\frac{b_{2}}{b_{1}}\right)^{2}\right) \frac{1}{s^{1-2\alpha-\beta+\nu}} + O\left(s^{-(1-\xi)}\right), \quad \text{when} \quad s \to 0, \end{split}$$

with $\xi \in (0, 1)$.

Figure 4 presents time evolution of the creep compliance for different values of model parameters, so that the creep compliance from Figure 4a, obtained according to (154) for the same set of parameters as the relaxation modulus from Figure 3a, is a Bernstein function, while the creep compliance from Figure 4b, obtained for the set of parameters in Table 4 guaranteeing that function $\tilde{\sigma}_{sr}$ has a pair of complex conjugated poles, is obviously not a Bernstein function, since it is a convex, rather than a concave function. Creep compliance curves from Figure 4, obtained according to the integral representation (154), perfectly overlap with the creep compliance curves from short and long time asymptotic expressions (156) and (157) generate curves that show satisfactory overlapping with the creep compliance curves.

6 Position and number of zeros of functions ϕ_{σ} and ϕ_{ε}

Functions ϕ_{σ} and ϕ_{ε} in the case of fractional anti-Zener and Zener models are of three different forms

$$\phi\left(s\right) = \begin{cases} as^{\xi} + b, \\ as^{1+\xi} + b, \end{cases} \quad \phi\left(s\right) = \begin{cases} as^{\xi} + bs^{\zeta} + c, \\ as^{1+\xi} + bs^{\zeta} + c, \end{cases} \quad \text{and} \quad \phi\left(s\right) = as^{2\xi} + bs^{\xi} + c, \end{cases}$$

see Table 1, where a, b, c > 0 and $\xi, \zeta \in (0, 1)$. Functions containing powers of complex variable s that are in the interval (0, 1), as it is well-known, do not admit zeros in the main Riemann branch $\arg s \in (-\pi, \pi)$, while functions having the power of leading term in the interval (1, 2) may admit no zeros, a negative real zero, and a pair of complex conjugated zeros with negative real part, as it is proved in the sequel.



Figure 4: Time evolution of the creep compliance.

6.1 Two-term function

Functions ϕ_{σ} and ϕ_{ε} in the case of model ID.ID takes the form

$$\phi(s) = as^{1+\xi} + b, \quad s \in \mathbb{C}, \quad \arg s \in (-\pi, \pi),$$

see Table 1, where a, b > 0 and $\xi \in (0, 1)$. The function ϕ proves to admit a pair of complex conjugated zeros with negative real part determined by

$$s_{\rm CCP} = \left(\frac{b}{a}\right)^{\frac{1}{1+\xi}} e^{i\frac{\pi}{1+\xi}} \quad \text{and} \quad \bar{s}_{\rm CCP} = \left(\frac{b}{a}\right)^{\frac{1}{1+\xi}} e^{-i\frac{\pi}{1+\xi}}.$$

The real and imaginary part of function ϕ , by substituting $s = \rho e^{i\varphi}$, become

$$\operatorname{Re} \phi\left(\rho,\varphi\right) = a\rho^{1+\xi} \cos\left(\left(1+\xi\right)\varphi\right) + b \quad \text{and} \quad \operatorname{Im} \phi\left(\rho,\varphi\right) = a\rho^{1+\xi} \sin\left(\left(1+\xi\right)\varphi\right),$$

so that $\operatorname{Im} \phi(\rho, \varphi) = 0$ implies $\varphi_k = \frac{k\pi}{1+\xi}$. By requesting $\varphi_k \in (-\pi, \pi)$, one finds $-(1+\xi) \leq k \leq (1+\xi)$, implying $k \in \{-1, 0, 1\}$, transforming the real part of function ϕ into

$$\operatorname{Re} \phi \left(\rho, \varphi_k \right) = a \rho^{1+\xi} \cos \left(k \pi \right) + b,$$

that is zero for $k \in \{-1,1\}$, implying $\rho = \left(\frac{b}{a}\right)^{\frac{1}{1+\xi}}$, while for k = 0 one has $\operatorname{Re} \phi(\rho, \varphi_0) = a\rho^{1+\xi} + b > 0$. Therefore, the zeros of function ϕ are $s_1 = \left(\frac{b}{a}\right)^{\frac{1}{1+\xi}} \operatorname{e}^{i\frac{\pi}{1+\xi}}$ and $s_{-1} = \left(\frac{b}{a}\right)^{\frac{1}{1+\xi}} \operatorname{e}^{-i\frac{\pi}{1+\xi}}$.

6.2 Three-term function

Functions ϕ_{σ} and ϕ_{ε} in the case of models IID.IID and IDD.IDD may take the form

$$\phi(s) = as^{1+\xi} + bs^{\zeta} + c, \quad s \in \mathbb{C}, \quad \arg s \in (-\pi, \pi)$$

see Table 1, where a, b, c > 0 and $\xi, \zeta \in (0, 1)$. The function ϕ proves to admit: a pair of complex conjugated zeros with negative real part if $\operatorname{Re} \phi(\rho^*, \pi) > 0$, a negative real zero $s = -\rho^*$ if $\operatorname{Re} \phi(\rho^*, \pi) = 0$, no zeros if

Re $\phi(\rho^*, \pi) < 0$, where Re ϕ is given by (158)₁ and where $\rho^* = \left(\frac{b}{a}\frac{\sin(\zeta\pi)}{\sin(\zeta\pi)}\right)^{\frac{1}{1+\xi-\zeta}}$

The real and imaginary part of function ϕ , by substituting $s = \rho e^{i\varphi}$, become

$$\operatorname{Re}\phi\left(\rho,\varphi\right) = a\rho^{1+\xi}\cos\left(\left(1+\xi\right)\varphi\right) + b\rho^{\zeta}\cos\left(\zeta\varphi\right) + c \quad \text{and} \quad \operatorname{Im}\phi\left(\rho,\varphi\right) = a\rho^{1+\xi}\sin\left(\left(1+\xi\right)\varphi\right) + b\rho^{\zeta}\sin\left(\zeta\varphi\right).$$
(158)

According to $(158)_2$, one has $\operatorname{Im} \phi(\rho, -\varphi) = -\operatorname{Im} \phi(\rho, \varphi)$ implying that $s_0 = \rho_0 e^{i\varphi_0}$ and its complex conjugate $\bar{s}_0 = \rho_0 e^{-i\varphi_0}$ are both zeros of function ϕ , so that it is sufficient to to consider the upper complex half-plane only. Clearly, there are no zeros of function ϕ lying in the right complex half-plane, since $\operatorname{Im} \phi(\rho, \varphi) > 0$ for $\varphi \in (0, \frac{\pi}{2})$ and moreover there are no positive zeros of function ϕ as well, since $\operatorname{Re} \phi(\rho, 0) > 0$. Therefore, zeros of function ϕ may only lie in the left complex half-plane.

In order to show that function ϕ admits a single zero in the left complex quarter-plane, the argument principle and contour γ , shown in Figure 5, is used, stating that if the variable s changes along the contour γ closed in the complex plane, then the number of zeros N of function ϕ in the domain encircled by contour γ is given by $\Delta \arg \phi(s) = 2\pi N$, provided that function ϕ does not have poles in the mentioned domain.



Figure 5: Contour γ .

The argument of complex numbers s belonging to the contour γ_1 has a fixed value $\varphi = \frac{\pi}{2}$, while their modulus changes so that $\rho \in (r, R)$, see the parameterization given in Table 5, so that one has

$$\operatorname{Im} \phi\left(\rho, \frac{\pi}{2}\right) = a\rho^{1+\xi} \cos\frac{\xi\pi}{2} + b\rho^{\zeta} \sin\frac{\zeta\pi}{2} > 0,$$

by (158), which also implies that function ϕ does not have purely imaginary zeros. The asymptotics of real and imaginary parts of function ϕ , given by (158), taking the form

$$\begin{aligned} &\operatorname{Re} \phi \left(r, \frac{\pi}{2} \right) \sim c \quad \text{and} \quad \operatorname{Im} \phi \left(r, \frac{\pi}{2} \right) \sim b r^{\zeta} \sin \frac{\zeta \pi}{2} \to 0^{+}, \quad \text{for} \quad r \to 0, \quad \text{as well as} \\ &\operatorname{Re} \phi \left(R, \frac{\pi}{2} \right) \sim -a R^{1+\xi} \sin \frac{\xi \pi}{2} \to -\infty \quad \text{and} \quad \operatorname{Im} \phi \left(R, \frac{\pi}{2} \right) \sim a R^{1+\xi} \cos \frac{\xi \pi}{2} \to \infty, \quad \text{for} \quad R \to \infty, \end{aligned}$$

so that $\left|\phi\left(R,\frac{\pi}{2}\right)\right| \sim aR^{1+\xi} \to \infty$ and $\tan \arg \phi\left(R,\frac{\pi}{2}\right) \sim -\tan \frac{\xi\pi}{2}$, implying $\arg \phi\left(R,\frac{\pi}{2}\right) \sim \pi - \frac{\xi\pi}{2} \in \left(\frac{\pi}{2},\pi\right)$. Therefore, as *s* changes along the contour γ_1 , the argument of function ϕ changes from zero to $\pi - \frac{\xi\pi}{2}$.

Complex number s belonging to the contour γ_2 has large but fixed modulus $\rho = R$, while its argument changes along the contour γ_2 taking the values $\varphi \in \left[\frac{\pi}{2}, \pi\right]$, see Table 5, so that the asymptotics of (158) as $R \to \infty$ gives

$$\operatorname{Re}\phi\left(R,\varphi\right)\sim aR^{1+\xi}\cos\left(\left(1+\xi\right)\varphi\right)\quad\text{and}\quad\operatorname{Im}\phi\left(R,\varphi\right)\sim aR^{1+\xi}\sin\left(\left(1+\xi\right)\varphi\right),$$

so that $|\phi(R,\varphi)| \sim aR^{1+\xi} \to \infty$ and $\arg \phi(R,\varphi) \sim (1+\xi)\varphi$, implying $\arg \phi(R,\varphi) \in \left[\frac{(1+\xi)\pi}{2}, (1+\xi)\pi\right]$. Therefore, as *s* changes along contour γ_2 , the argument of function ϕ starts from the second quadrant, since $\arg \phi\left(R, \frac{\pi}{2}\right) = \frac{(1+\xi)\pi}{2}$ and ends either in the third, or possibly the fourth quadrant, since $\arg \phi(R,\pi) = (1+\xi)\pi$, however not crossing through the first quadrant, since the real and imaginary parts of function ϕ cannot be simultaneously positive for $\varphi \in \left[\frac{\pi}{2}, \pi\right]$, since $\frac{\pi}{2} \leq (1+\xi)\varphi \leq 2\pi$.

The argument of complex numbers s lying on the contour γ_3 has a fixed value $\varphi = \pi$, while their modulus changes in the interval $\rho \in (r, R)$, yielding

$$\operatorname{Re}\phi\left(\rho,\pi\right) = -a\rho^{1+\xi}\cos\left(\xi\pi\right) + b\rho^{\zeta}\cos\left(\zeta\pi\right) + c \quad \text{and} \quad \operatorname{Im}\phi\left(\rho,\pi\right) = -a\rho^{1+\xi}\sin\left(\xi\pi\right) + b\rho^{\zeta}\sin\left(\zeta\pi\right),$$

according to (158), so that the asymptotics of real and imaginary parts of function ϕ is obtained in the form

$$\operatorname{Re}\phi\left(R,\pi\right)\sim-aR^{1+\xi}\cos\left(\xi\pi\right)\rightarrow-\infty\quad\text{and}\quad\operatorname{Im}\phi\left(R,\pi\right)\sim-aR^{1+\xi}\sin\left(\xi\pi\right)\rightarrow-\infty,\quad\text{for}\quad R\rightarrow\infty,$$

as well as

$$\operatorname{Re}\phi\left(r,\pi\right)\sim c\quad \text{and}\quad \operatorname{Im}\phi\left(r,\pi\right)\sim br^{\zeta}\sin\left(\zeta\pi\right)\rightarrow0^{+}\quad \text{for}\quad r\rightarrow0,$$

implying that $\arg \phi(R,\pi) \sim (1+\xi)\pi$ as $R \to \infty$, as well as $\tan \arg \phi(r,\pi) \sim 0$ as $r \to 0$. Obviously, $\operatorname{Im} \phi(\rho,\pi)$ changes sign as ρ changes from $R \to \infty$ to $r \to 0$ and it has a single zero at $\rho^* = \left(\frac{b}{a}\frac{\sin(\zeta\pi)}{\sin(\xi\pi)}\right)^{\frac{1}{1+\xi-\zeta}}$, since $\rho^{1+\xi}$ is a convex and ρ^{ζ} is a concave function, both monotonically increasing from zero to infinity. Therefore, the sign of $\operatorname{Re} \phi(\rho^*,\pi)$ determines whether the argument of function ϕ , starting from $\arg \phi(R,\pi) = (1+\xi)\pi$, as s changes along contour γ_3 , ends at $\arg \phi(r,\pi) = 0$ or at $\arg \phi(r,\pi) = 2\pi$, so that the former is achieved if $\operatorname{Re} \phi(\rho^*,\pi) < 0$ and the latter is achieved if $\operatorname{Re} \phi(\rho^*,\pi) > 0$. Moreover, if $\operatorname{Re} \phi(\rho^*,\pi) = 0$, then function ϕ admits a negative real zero $s = -\rho^* = -\left(\frac{b}{a}\frac{\sin(\zeta\pi)}{\sin(\xi\pi)}\right)^{\frac{1}{1+\xi-\zeta}}$, since ρ^* solves $\operatorname{Im} \phi(\rho,\pi) = 0$.

The modulus of complex number s belonging to the contour γ_4 has a fixed but small value $\rho = r$, while its argument changes in the interval $\varphi \in \left[\frac{\pi}{2}, \pi\right]$, so that the asymptotics of real and imaginary parts of function ϕ , see (158), become

$$\operatorname{Re}\phi(r,\varphi) \sim c$$
 and $\operatorname{Im}\phi(r,\varphi) \sim br^{\zeta}\sin(\zeta\varphi) \to 0^+$ for $r \to 0$

implying that $\arg \phi$ does not really change.

Summing up, the change of argument of function ϕ , as s changes along the contour γ , is either $\Delta \arg \phi(s) = 2\pi$ if $\operatorname{Re} \phi(\rho^*, \pi) > 0$, or $\Delta \arg \phi(s) = 0$ if $\operatorname{Re} \phi(\rho^*, \pi) < 0$, implying that function ϕ either has one zero, or has no zeros in the upper left complex quarter-plane. If $\operatorname{Re} \phi(\rho^*, \pi) = 0$, then function ϕ admits a negative real zero.

6.3 Function quadratic in s^{ξ}

Function ϕ_{σ} in the case of models I⁺ID.ID and IDD⁺.DD⁺, as well as the function ϕ_{ε} in the case of model ID.IDD⁺, so as both functions ϕ_{σ} and ϕ_{ε} in the case of models I⁺ID.I⁺ID, IDD⁺.IDD⁺, and I⁺ID.IDD⁺, take the same form

$$b(s) = as^{2\xi} + bs^{\xi} + c, \quad s \in \mathbb{C}, \quad \arg s \in (-\pi, \pi),$$
(159)

see Table 1, where a, b, c > 0 and $\xi \in (0, 1)$. If $\xi \in (0, \frac{1}{2}]$, then all powers of variable s belong to interval (0, 1) and, as it is well-known, the function ϕ does not have zeros, while if $\xi \in (\frac{1}{2}, 1)$, then the zeros of function ϕ are

$$s^{\xi} = -\frac{b}{2a} \left(1 \pm \sqrt{1 - \frac{4ac}{b^2}} \right).$$

Assuming that the discriminant of equation quadratic in s^{ξ} is non-negative, i.e., that $\frac{2\sqrt{ac}}{b} \leq 1$, one has

$$s^{\xi} = -\frac{b}{2a} \left(1 \pm \sqrt{1 - \frac{4ac}{b^2}} \right) < 0 \quad \text{i.e.,} \quad \rho^{\xi} \operatorname{e}^{\mathrm{i}\xi\varphi} = \frac{b}{2a} \left(1 \pm \sqrt{1 - \frac{4ac}{b^2}} \right) \operatorname{e}^{\mathrm{i}(2k+1)\pi}, \quad k \in \mathbb{Z},$$

implying

$$\rho = \sqrt[\epsilon]{\frac{b}{2a} \left(1 \pm \sqrt{1 - \frac{4ac}{b^2}} \right)} \quad \text{and} \quad \varphi = \frac{(2k+1)\pi}{\xi}.$$
(160)

The complex number s with modulus ρ and argument φ , given by (160), is not a zero of function ϕ , given by (159), since the requirement $\varphi \in (-\pi, \pi)$ cannot be satisfied for any $k \in \mathbb{Z}$, due to $-\pi < \frac{(2k+1)\pi}{\xi} < \pi$ implying $-1 < -\xi < 2k + 1 < \xi < 1$, i.e., -1 < k < 0.

On the other hand, if the discriminant of equation quadratic in s^{ξ} is negative, i.e., if $\frac{2\sqrt{ac}}{b} > 1$, then one has

$$s^{\xi} = -\frac{b}{2a} \left(1 \pm i\sqrt{\frac{4ac}{b^2} - 1} \right) = -\lambda \pm i\eta = z_{\pm} \quad \text{with} \quad \lambda = \frac{b}{2a} > 0 \quad \text{and} \quad \eta = \frac{b}{2a}\sqrt{\frac{4ac}{b^2} - 1} > 0, \tag{161}$$

so that

$$s^{\xi} = \rho^{\xi} e^{i\xi\varphi_{\pm}} = |z_{\pm}| e^{i(\arg z_{\pm} + 2k\pi)} \quad \text{with} \quad |z_{\pm}| = \sqrt{\lambda^2 + \eta^2} \quad \text{and} \quad \arg z_{\pm} = \pm \left(1 - \frac{1}{\pi}\arctan\frac{\eta}{\lambda}\right)\pi, \quad (162)$$

implying

s

$$_{\pm} = \rho e^{i\varphi_{\pm}} \quad \text{with} \quad \rho = \sqrt[\xi]{|z_{\pm}|} \quad \text{and} \quad \varphi_{\pm} = \frac{\arg z_{\pm} + 2k\pi}{\xi}.$$
(163)

In order for φ_+ to be an argument of zero of the function ϕ , given by (159), one requires $\varphi_+ = \frac{\arg z_+ + 2k\pi}{\xi} = \frac{2k+1-\delta}{\xi}\pi \in (0,\pi)$, with $\delta = \frac{1}{\pi}\arctan\frac{\eta}{\lambda} \in (0,\frac{1}{2})$, so that one obtains $0 < (2k+1-\delta)\pi < \xi\pi$, transforming into $0 < \delta < 2k+1 < \delta + \xi < \frac{3}{2}$, i.e., $-\frac{1}{2} < k < \frac{1}{4}$, yielding k = 0. Similar argumentation in the case of φ_- also implies that k = 0. Therefore, the zeros $s_{\pm} = \rho e^{i\varphi_{\pm}}$ of function ϕ , given by (159), are such that

$$\rho = \sqrt[2\epsilon]{\lambda^2 + \eta^2} \quad \text{and} \quad \varphi_{\pm} = \pm \frac{1 - \frac{1}{\pi} \arctan \frac{\eta}{\lambda}}{\xi} \pi_{\pm}$$

with λ and η defined by (161), according to (162) and (163). Actually, the function ϕ admits a pair of complex conjugated zeros

$$s_{\rm CCP} = \sqrt[2\xi]{\lambda^2 + \eta^2} e^{i\frac{1 - \frac{1}{\pi} \arctan \frac{\eta}{\lambda}}{\xi}\pi} \quad \text{and} \quad \bar{s}_{\rm CCP} = \sqrt[2\xi]{\lambda^2 + \eta^2} e^{-i\frac{1 - \frac{1}{\pi} \arctan \frac{\eta}{\lambda}}{\xi}\pi}, \tag{164}$$

having negative real part, if $\tan(\xi\pi) > -\frac{\eta}{\lambda}$, since $\varphi_+ = \frac{1-\frac{1}{\pi}\arctan\frac{\eta}{\lambda}}{\xi}\pi < \pi$ implies $1-\frac{1}{\pi}\arctan\frac{\eta}{\lambda} < \xi$, reducing to $\tan(\xi\pi) > -\frac{\eta}{\lambda}$, while one already knows that $\arg z_+ = \pi - \arctan\frac{\eta}{\lambda} > \frac{\pi}{2}$, see (162)₃, and therefore $\varphi_+ = \frac{\arg z_+}{\xi} > \frac{\pi}{2}$, see (163), due to $\xi < 1$. Note, if $\tan(\xi\pi) < -\frac{\eta}{\lambda}$, then $\varphi_+ > \pi$ and function ϕ , given by (159) does not have zeros for $\arg s \in (-\pi, \pi)$.

In the special case when $\varphi_{\pm} = \pm \frac{1 - \frac{1}{\pi} \arctan \frac{\eta}{\lambda}}{\xi} \pi = \pm \pi$, i.e., when $\frac{1 - \frac{1}{\pi} \arctan \frac{\eta}{\lambda}}{\xi} = 1$ reducing to $\tan(\xi \pi) = -\frac{\eta}{\lambda}$, one has that the function ϕ , given by (159), admits a negative real zero

$$s_{\rm RP} = -\sqrt[2\xi]{\lambda^2 + \eta^2},\tag{165}$$

according to (164), with the order ξ obtained as a solution to $\tan(\xi\pi) = -\frac{\eta}{\lambda}$.

In conclusion, the function ϕ , given by (159), does not have zeros if $\xi \in (0, \frac{1}{2}]$, or if $\xi \in (\frac{1}{2}, 1)$ and $\frac{2\sqrt{ac}}{b} \leq 1$, or if $\xi \in (\frac{1}{2}, 1)$, $\frac{2\sqrt{ac}}{b} > 1$, and $\tan(\xi\pi) < -\sqrt{\frac{4ac}{b^2} - 1}$, while if $\xi \in (\frac{1}{2}, 1)$ and $\frac{2\sqrt{ac}}{b} > 1$, then it has either a negative real zero

$$s_{\rm RP} = -\sqrt[2\varepsilon]{\frac{c}{a}},\tag{166}$$

according to (165) and (161), if $\tan(\xi \pi) = -\sqrt{\frac{4ac}{b^2} - 1}$, that actually determines the order ξ , or it has a pair of complex conjugated zeros

$$s_{\rm CCP} = \sqrt[2\xi]{\frac{c}{a}} e^{i\frac{1-\frac{1}{\pi}\arctan\sqrt{\frac{4ac}{b^2}-1}}{\xi}\pi} \quad \text{and} \quad \bar{s}_{\rm CCP} = \sqrt[2\xi]{\frac{c}{a}} e^{-i\frac{1-\frac{1}{\pi}\arctan\sqrt{\frac{4ac}{b^2}-1}}{\xi}\pi}, \tag{167}$$

according to (164) and (161), if $\tan(\xi \pi) > -\sqrt{\frac{4ac}{b^2} - 1}$.

7 Conclusion

The analysis of the energy balance of one-dimensional viscoelastic body modeled by the fractional anti-Zener and Zener models, conducted in Section 2, yields two expressions for the power per unit volume of the viscoelastic body, where the first one is expressed through the strain (7) and the second one is expressed through the stress (8), both of them containing information about the elastic properties of material, represented by the energy per unit volume stored in the viscoelastic body (9) and (10), as well as about the viscous properties of the material, represented by the dissipated power per unit volume (11) and (12).

The positivity of stored energy and dissipated power per unit volume, both of them consisting of the instantaneous and hereditary terms and depending either on the relaxation modulus and its derivatives, or on the creep compliance and its derivatives, is guaranteed if the relaxation modulus, respectively creep compliance, is a completely monotone function, respectively a Bernstein function, and therefore the Laplace transform method is used in Section 3 in order to obtain the explicit forms of the relaxation modulus σ_{sr} , expressed by (15) in terms of functions $\sigma_{sr}^{(\text{NP})}$, $\sigma_{sr}^{(\text{RP})}$, and $\sigma_{sr}^{(\text{CCP})}$, given by (16), (17), and (18), as well as of the creep compliance ε_{cr} , expressed by (27) in terms of functions $\varepsilon_{cr}^{(\text{NP})}$, $\varepsilon_{cr}^{(\text{RP})}$, and $\varepsilon_{cr}^{(\text{CCP})}$, given by (28), (29), and (30), allowing for the relaxation modulus and creep compliance to be non-monotonic and even oscillatory functions with exponentially decreasing amplitude. These qualitative properties are due to the existence of poles of the relaxation modulus is proved to be a completely monotone function, as well as the creep compliance proved to be a Bernstein function, if additional restrictions on model parameters, derived in Section 4 and narrowing the thermodynamical restrictions on the parameters of fractional anti-Zener and Zener models, listed in Appendix A, are posed.

In Section 5, the time evolution of relaxation modulus and creep compliance for model I⁺ID.ID is examined numerically for the set of parameters from Table 4 and presented in Figures 3 and 4 along with the short and long time asymptotics, displaying a good agreement. In the case of model I⁺ID.ID, the relaxation modulus can be a completely monotone function, as obvious Figure 3a, as well as a function resembling to a completely monotone function, because of the choice of model parameters not guaranteeing the mentioned property, see Figure 3b, and moreover an oscillatory function having exponentially decreasing amplitude, see Figures 3c and 3d. On the other hand, the creep compliance can be at most a non-monotonic function, with the possibility to be a Bernstein function, see Figure 4a, and a function that is positive, monotonically increasing and a convex rather than a concave function, see Figure 4b. The creep compliance curves, obtained according to the integral representation (154) because of convergence reasons rather than (152), are compared with the ones obtained by the Mittag-Leffler representation (155) and showed a good agreement.

Declaration of Competing Interest

The authors declare that they have no conflict of interest.

Data availability

No data was used for the research described in the article.

Acknowledgements

The work is supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia under grant 451-03-47/2023-01/200125 (SJ and DZ).

A Fractional anti-Zener and Zener models

A.1 Symmetric models

 ${\bf Model \ ID.ID}$ takes the form

$$\left(a_{1\ 0}\mathbf{I}_{t}^{\alpha}+a_{2\ 0}\mathbf{D}_{t}^{\beta}\right)\sigma(t)=\left(b_{1\ 0}\mathbf{I}_{t}^{\mu}+a_{2\ 0}\mathbf{D}_{t}^{\alpha+\beta-\mu}\right)\varepsilon(t)\,,$$

with thermodynamical restrictions

$$0 \leqslant \alpha + \beta - \mu \leqslant 1, \quad \mu \leqslant \alpha, \quad \beta + \mu \leqslant 1, \tag{168}$$

$$-\frac{a_1}{a_2} \frac{\cos\frac{(2\alpha+\beta-\mu)\pi}{2}}{\cos\frac{(\beta+\mu)\pi}{2}} \leqslant \frac{b_1}{b_2} \leqslant \frac{a_1}{a_2} \frac{\sin\frac{(2\alpha+\beta-\mu)\pi}{2}}{\sin\frac{(\beta+\mu)\pi}{2}}.$$
(169)

Model ID.DD⁺ takes the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\beta}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{D}_{t}^{\mu}+b_{2\,0}\mathbf{D}_{t}^{\alpha+\beta+\mu}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$1 \leqslant \alpha + \beta + \mu \leqslant 2, \quad \beta \leqslant \mu \leqslant 1 - \alpha, \tag{170}$$

$$\frac{a_1}{a_2} \frac{\left| \cos \frac{(2\alpha + \beta + \mu)\pi}{2} \right|}{\cos \frac{(\mu - \beta)\pi}{2}} \leqslant \frac{b_1}{b_2}.$$
(171)

Model IID.IID takes the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{I}_{t}^{\beta}+a_{3\,0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{I}_{t}^{\alpha+\gamma-\eta}+b_{2\,0}\mathbf{I}_{t}^{\beta+\gamma-\eta}+b_{3\,0}\mathbf{D}_{t}^{\eta}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$\beta < \alpha, \quad \gamma \leqslant \eta, \quad 0 \leqslant \beta + \gamma - \eta \leqslant \alpha + 2\gamma - \eta \leqslant 1, \quad \alpha + \gamma \leqslant \beta + \eta, \tag{172}$$

$$-\frac{b_3}{b_1} \frac{\cos\frac{(\alpha+\gamma)\pi}{2}}{\cos\frac{(\alpha+2\gamma-\eta)\pi}{2}} \leqslant \frac{a_3}{a_1} \leqslant \frac{b_3}{b_1} \frac{\sin\frac{(\alpha+\gamma)\pi}{2}}{\sin\frac{(\alpha+2\gamma-\eta)\pi}{2}},\tag{173}$$

$$-\frac{b_3}{b_2} \frac{\cos\frac{(\beta+\eta)\pi}{2}}{\cos\frac{(\beta+2\gamma-\eta)\pi}{2}} \leqslant \frac{a_3}{a_2} \leqslant \frac{b_3}{b_2} \frac{\sin\frac{(\beta+\eta)\pi}{2}}{\sin\frac{(\beta+2\gamma-\eta)\pi}{2}}.$$
(174)

Model IDD.IDD takes the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\beta}+a_{3\,0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{I}_{t}^{\mu}+b_{2\,0}\mathbf{D}_{t}^{\alpha+\beta-\mu}+b_{3\,0}\mathbf{D}_{t}^{\alpha+\gamma-\mu}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$0 \leqslant \alpha + \gamma - \mu \leqslant 1, \quad \beta < \gamma, \quad \mu \leqslant \alpha, \quad \gamma + \mu \leqslant \alpha + \beta, \quad \gamma + \mu \leqslant 1, \tag{175}$$

$$-\frac{b_2}{b_1}\frac{\cos\frac{(\beta+\mu)\pi}{2}}{\cos\frac{(\beta+\mu)\pi}{2}} \leqslant \frac{a_2}{a_1} \leqslant \frac{b_2}{b_1}\frac{\sin\frac{(\beta+\mu)\pi}{2}}{\sin\frac{(\beta+\mu)\pi}{2}},\tag{176}$$

$$-\frac{b_3}{b_1} \frac{\cos\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\cos\frac{(\gamma+\mu)\pi}{2}} \leqslant \frac{a_3}{a_1} \leqslant \frac{b_3}{b_1} \frac{\sin\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\sin\frac{(\gamma+\mu)\pi}{2}}.$$
(177)

Model IID.IDD takes the form

$$\left(a_{1\ 0}\mathbf{I}_{t}^{\alpha} + a_{2\ 0}\mathbf{I}_{t}^{\beta} + a_{3\ 0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right) = \left(b_{1\ 0}\mathbf{I}_{t}^{\mu} + b_{2\ 0}\mathbf{D}_{t}^{\nu} + b_{3\ 0}\mathbf{D}_{t}^{\alpha+\gamma-\mu}\right)\varepsilon\left(t\right)$$

with thermodynamical restrictions

$$\mu \leqslant \beta < \alpha, \quad \gamma \leqslant \nu, \quad \alpha + \beta + \gamma \leqslant 1 + \mu, \quad \mu + \nu - \gamma < \alpha \leqslant 1 - \nu, \tag{178}$$

$$-\frac{b_3}{b_1} \frac{\cos\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\cos\frac{(\gamma+\mu)\pi}{2}} \leqslant \frac{a_3}{a_1} \leqslant \frac{b_3}{b_1} \frac{\sin\frac{(2\alpha+\gamma-\mu)\pi}{2}}{\sin\frac{(\gamma+\mu)\pi}{2}}.$$
(179)

Model $I^{+}ID.I^{+}ID$ takes the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{1+\alpha} + a_{2\,0}\mathbf{I}_{t}^{\frac{1+\alpha-\gamma}{2}} + a_{3\,0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right) = \left(b_{1\,0}\mathbf{I}_{t}^{1+\mu} + b_{2\,0}\mathbf{I}_{t}^{\frac{1+\mu-(\alpha+\gamma-\mu)}{2}} + b_{3\,0}\mathbf{D}_{t}^{\alpha+\gamma-\mu}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$\mu \leq \alpha, \quad \alpha + \gamma + 2 \left(\alpha - \mu \right) = 3\alpha + \gamma - 2\mu \leq 1, \tag{180}$$

$$\frac{a_2}{a_1} \leqslant \frac{b_2}{b_1} \frac{\cos\frac{(1-3\alpha-\gamma+2\mu)\pi}{4}}{\cos\frac{(1+\alpha-\gamma-2\mu)\pi}{4}}, \quad \frac{a_3}{a_2} \leqslant \frac{b_3}{b_2} \frac{\sin\frac{(1+3\alpha+\gamma-2\mu)\pi}{4}}{\sin\frac{(1-\alpha+\gamma+2\mu)\pi}{4}}, \tag{181}$$

$$a_{3}b_{1}\cos\frac{(\gamma+\mu)\pi}{2} - a_{2}b_{2}\sin\frac{(\alpha-\mu)\pi}{2} \leqslant a_{1}b_{3}\cos\frac{(2\alpha+\gamma-\mu)\pi}{2},$$
(182)

$$a_1 b_3 \sin \frac{(2\alpha + \gamma - \mu)\pi}{2} \leq a_2 b_2 \cos \frac{(\alpha - \mu)\pi}{2} - a_3 b_1 \sin \frac{(\gamma + \mu)\pi}{2}.$$
 (183)

Model IDD^+.IDD⁺ takes the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\frac{1+\gamma-\alpha}{2}}+a_{3\,0}\mathbf{D}_{t}^{1+\gamma}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{I}_{t}^{\alpha+\gamma-\eta}+b_{2\,0}\mathbf{D}_{t}^{\frac{1+\eta-(\alpha+\gamma-\eta)}{2}}+b_{3\,0}\mathbf{D}_{t}^{1+\eta}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$0 \leqslant \alpha + \gamma - \eta \leqslant 1, \quad \gamma \leqslant \eta, \quad \alpha + \eta + (\eta - \gamma) = \alpha - \gamma + 2\eta \leqslant 1, \tag{184}$$

$$\frac{a_2}{a_1} \leqslant \frac{b_2}{b_1} \frac{\sin\frac{(1+\alpha-\gamma+2\eta)\pi}{4}}{\sin\frac{(1+\alpha+3\gamma-2\eta)\pi}{4}}, \quad \frac{a_3}{a_2} \leqslant \frac{b_3}{b_2} \frac{\cos\frac{(1-\alpha+\gamma-2\eta)\pi}{4}}{\cos\frac{(1-\alpha-3\gamma+2\eta)\pi}{4}}, \tag{185}$$

$$a_3b_1\cos\frac{(\alpha+2\gamma-\eta)\pi}{2} - a_2b_2\sin\frac{(\eta-\gamma)\pi}{2} \leqslant a_1b_3\cos\frac{(\alpha+\eta)\pi}{2},\tag{186}$$

$$a_1 b_3 \sin \frac{(\alpha + \eta) \pi}{2} \leqslant a_2 b_2 \cos \frac{(\eta - \gamma) \pi}{2} - a_3 b_1 \sin \frac{(\alpha + 2\gamma - \eta) \pi}{2}.$$
(187)

Model $I^{+}ID.IDD^{+}$ takes the form

$$\left(a_{1\ 0}\mathbf{I}_{t}^{1+\alpha} + a_{2\ 0}\mathbf{I}_{t}^{\frac{1+\alpha-\gamma}{2}} + a_{3\ 0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right) = \left(b_{1\ 0}\mathbf{I}_{t}^{\alpha+\gamma-\eta} + b_{2\ 0}\mathbf{D}_{t}^{\frac{1+\eta-(\alpha+\gamma-\eta)}{2}} + b_{3\ 0}\mathbf{D}_{t}^{1+\eta}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$\eta \leq \gamma, \quad \alpha + \gamma + 2(\gamma - \eta) = \alpha + 3\gamma - 2\eta \leq 1,$$
(188)

$$\eta \leqslant \gamma, \quad \alpha + \gamma + 2 \left(\gamma - \eta\right) = \alpha + 3\gamma - 2\eta \leqslant 1, \tag{188}$$

$$\frac{a_1}{b_1} \frac{\sin \frac{(1 + \alpha - \gamma + 2\eta)\pi}{4}}{\cos \frac{(1 - \alpha - 3\gamma + 2\eta)\pi}{4}} \leqslant \frac{a_2}{b_2} \leqslant \frac{a_3}{b_3} \frac{\cos \frac{(1 - \alpha - 3\gamma + 2\eta)\pi}{4}}{\sin \frac{(1 + \alpha - \gamma + 2\eta)\pi}{4}}, \tag{189}$$

$$a_1 b_3 \cos \frac{(\alpha + \eta)\pi}{2} - a_2 b_2 \sin \frac{(\gamma - \eta)\pi}{2} \leqslant a_3 b_1 \cos \frac{(\alpha + 2\gamma - \eta)\pi}{2}, \tag{190}$$

$$a_3b_1\sin\frac{(\alpha+2\gamma-\eta)\pi}{2} \leqslant a_2b_2\cos\frac{(\gamma-\eta)\pi}{2} - a_1b_3\sin\frac{(\alpha+\eta)\pi}{2}.$$
(191)

A.2 Asymmetric models

Model IID.ID takes the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha+\beta-\gamma}+a_{2\,0}\mathbf{I}_{t}^{\nu}+a_{3\,0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{I}_{t}^{\alpha}+b_{2\,0}\mathbf{D}_{t}^{\beta}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$0 \leqslant \alpha \leqslant \nu < \alpha + \beta - \gamma \leqslant 1, \quad \beta + \nu \leqslant 1, \tag{192}$$

$$-\frac{a_1}{a_3} \frac{\cos\frac{(\alpha+2\beta-\gamma)\pi}{2}}{\cos\frac{(\alpha+\gamma)\pi}{2}} \leqslant \frac{b_1}{b_2} \leqslant \frac{a_1}{a_3} \frac{\sin\frac{(\alpha+2\beta-\gamma)\pi}{2}}{\sin\frac{(\alpha+\gamma)\pi}{2}}.$$
(193)

Model IDD.DD $^+$ takes the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\beta}+a_{3\,0}\mathbf{D}_{t}^{\gamma}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{D}_{t}^{\mu}+b_{2\,0}\mathbf{D}_{t}^{\alpha+\beta+\mu}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$1 \leqslant \alpha + \beta + \mu \leqslant 2, \quad \beta < \gamma \leqslant \mu \leqslant 1 - \alpha, \tag{194}$$

$$\frac{a_1}{a_2} \frac{\left|\cos\frac{(2\alpha+\beta+\mu)\pi}{2}\right|}{\cos\frac{(\mu-\beta)\pi}{2}} \leqslant \frac{b_1}{b_2}.$$
(195)

Model $I^+ID.ID$ takes the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha+\beta+\nu}+a_{2\,0}\mathbf{I}_{t}^{\nu}+a_{3\,0}\mathbf{D}_{t}^{\alpha+\beta-\nu}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{I}_{t}^{\alpha}+b_{2\,0}\mathbf{D}_{t}^{\beta}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$0 \leqslant \alpha + \beta - \nu \leqslant 1, \quad 1 \leqslant \alpha + \beta + \nu \leqslant 2, \quad \alpha \leqslant \nu \leqslant 1 - \beta,$$
(196)

$$\frac{a_1}{a_2} \frac{\left| \cos \frac{(\alpha + 2\beta + \nu)\pi}{2} \right|}{\cos \frac{(\nu - \alpha)\pi}{2}} \leqslant \frac{b_1}{b_2} \leqslant \frac{a_2}{a_3} \frac{\sin \frac{(\beta + \nu)\pi}{2}}{\sin \frac{(2\alpha + \beta - \nu)\pi}{2}}.$$
(197)

Model \mathbf{IDD}^+ **.** \mathbf{DD}^+ takes the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\beta}+a_{3\,0}\mathbf{D}_{t}^{\alpha+2\beta}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{D}_{t}^{\mu}+b_{2\,0}\mathbf{D}_{t}^{\alpha+\beta+\mu}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$1 \leqslant \alpha + 2\beta \leqslant 2, \quad 1 \leqslant \alpha + \beta + \mu \leqslant 2, \quad \beta \leqslant \mu \leqslant 1 - \alpha, \tag{198}$$

$$\frac{a_1}{a_2} \frac{\left| \cos \frac{(2\alpha + \beta + \mu)\pi}{2} \right|}{\cos \frac{(\mu - \beta)\pi}{2}} \leqslant \frac{b_1}{b_2} \leqslant \frac{a_2}{a_3} \frac{\sin \frac{(\alpha + \mu)\pi}{2}}{\sin \frac{(\alpha + 2\beta - \mu)\pi}{2}}.$$
(199)

Model ID.IDD takes the form

$$(a_{1\,0}\mathbf{I}_{t}^{\mu} + a_{2\,0}\mathbf{D}_{t}^{\nu})\,\sigma\left(t\right) = \left(b_{1\,0}\mathbf{I}_{t}^{\alpha} + b_{2\,0}\mathbf{D}_{t}^{\beta} + b_{3\,0}\mathbf{D}_{t}^{\mu+\nu-\alpha}\right)\varepsilon\left(t\right)$$

with thermodynamical restrictions

$$0 \leqslant \nu \leqslant \beta < \mu + \nu - \alpha \leqslant 1, \quad \alpha \leqslant \mu \leqslant 1 - \beta, \tag{200}$$

$$-\frac{a_1}{a_2} \frac{\cos\frac{(2\mu+\nu-\alpha)\pi}{2}}{\cos\frac{(\alpha+\nu)\pi}{2}} \leqslant \frac{b_1}{b_3} \leqslant \frac{a_1}{a_2} \frac{\sin\frac{(2\mu+\nu-\alpha)\pi}{2}}{\sin\frac{(\alpha+\nu)\pi}{2}}.$$
(201)

Model ID.DDD⁺ takes the form

$$\left(a_{1\,0}\mathbf{I}_{t}^{\alpha}+a_{2\,0}\mathbf{D}_{t}^{\beta}\right)\sigma\left(t\right)=\left(b_{1\,0}\mathbf{D}_{t}^{\mu}+b_{2\,0}\mathbf{D}_{t}^{\nu}+b_{3\,0}\mathbf{D}_{t}^{\alpha+\beta+\nu}\right)\varepsilon\left(t\right),$$

with thermodynamical restrictions

$$1 \leqslant \alpha + \beta + \nu \leqslant 2, \quad \beta \leqslant \mu < \nu \leqslant 1 - \alpha, \tag{202}$$

$$\frac{a_1}{a_2} \frac{\left|\cos\frac{(2\alpha+\beta+\nu)\pi}{2}\right|}{\cos\frac{(\nu-\beta)\pi}{2}} \leqslant \frac{b_2}{b_3}.$$
(203)

Model ID.IDD $^+$ takes the form

$$\left(a_{1 0} \mathbf{I}_{t}^{\alpha} + a_{2 0} \mathbf{D}_{t}^{\beta} \right) \sigma \left(t \right) = \left(b_{1 0} \mathbf{I}_{t}^{\alpha+\beta-\nu} + b_{2 0} \mathbf{D}_{t}^{\nu} + b_{3 0} \mathbf{D}_{t}^{\alpha+\beta+\nu} \right) \varepsilon \left(t \right),$$

with thermodynamical restrictions

$$0 \leqslant \alpha + \beta - \nu \leqslant 1, \quad 1 \leqslant \alpha + \beta + \nu \leqslant 2, \quad \beta \leqslant \nu \leqslant 1 - \alpha, \tag{204}$$

$$\frac{b_1}{b_2} \frac{\sin\frac{(\alpha+\nu)\pi}{2}}{\sin\frac{(\alpha+\nu)\pi}{2}} \leqslant \frac{a_1}{a_2} \leqslant \frac{b_2}{b_3} \frac{\cos\frac{(\nu-\beta)\pi}{2}}{\left|\cos\frac{(2\alpha+\beta+\nu)\pi}{2}\right|}.$$
(205)

References

- M. Ait Ichou, H. El Amri, and A. Ezziani. Mathematical modeling of wave propagation in viscoelastic media with the fractional Zener model. *Mathematical Modeling and Computing*, 8:601–615, 2021.
- [2] G. Amendola, M. Fabrizio, and J. M. Golden. Thermodynamics of Materials with Memory: Theory and Applications. Springer Science+Business Media LLC, New York, 2012.
- [3] T. M. Atanackovic. A modified Zener model of a viscoelastic body. Continuum Mechanics and Thermodynamics, 14:137–148, 2002.
- [4] T. M. Atanackovic, M. Janev, and S. Pilipovic. On the thermodynamical restrictions in isothermal deformations of fractional Burgers model. *Philosophical Transactions of the Royal Society A*, 378:20190278–1–13, 2020.
- [5] T. M. Atanackovic, S. Konjik, Lj. Oparnica, and D. Zorica. Thermodynamical restrictions and wave propagation for a class of fractional order viscoelastic rods. *Abstract and Applied Analysis*, 2011:ID975694– 1–32, 2011.
- [6] T. M. Atanackovic, S. Pilipovic, B. Stankovic, and D. Zorica. Fractional Calculus with Applications in Mechanics: Vibrations and Diffusion Processes. Wiley-ISTE, London, 2014.
- [7] T. M. Atanackovic, S. Pilipovic, B. Stankovic, and D. Zorica. Fractional Calculus with Applications in Mechanics: Wave Propagation, Impact and Variational Principles. Wiley-ISTE, London, 2014.
- [8] T. M. Atanackovic, S. Pilipovic, and D. Zorica. Distributed-order fractional wave equation on a finite domain: creep and forced oscillations of a rod. *Continuum Mechanics and Thermodynamics*, 23:305–318, 2011.
- [9] T. M. Atanackovic, S. Pilipovic, and D. Zorica. Distributed-order fractional wave equation on a finite domain. Stress relaxation in a rod. *International Journal of Engineering Science*, 49:175–190, 2011.
- [10] R. L. Bagley and P. J. Torvik. On the fractional calculus model of viscoelastic behavior. Journal of Rheology, 30:133–155, 1986.
- [11] E. Bazhlekova and I. Bazhlekov. Complete monotonicity of the relaxation moduli of distributed-order fractional Zener model. In Proceedings of the 44th International Conference on Applications of Mathematics in Engineering and Economics, AIP Conference Proceedings 2048, pages 050008–1–8, 2018.
- [12] E. Bazhlekova and K. Tsocheva. Fractional Burgers' model: thermodynamic constraints and completely monotonic relaxation function. Comptes rendus de l'Académie bulgare des Sciences, 69:825–834, 2016.
- [13] F. Broucke and Lj. Oparnica. Micro-local and qualitative analysis of the fractional Zener wave equation. Journal of Differential Equations, 321:217–257, 2022.
- [14] F. Broucke and Lj. Oparnica. Distributed-order time-fractional wave equations. Zeitschrift f
 ür angewandte Mathematik und Physik, 74:19–1–25, 2023.
- [15] W. Cai, W. Chen, J. Fang, and S. Holm. A survey on fractional derivative modeling of power-law frequencydependent viscous dissipative and scattering attenuation in acoustic wave propagation. *Applied Mechanics Reviews*, 70:1–12, 2018.
- [16] M. Caputo and F. Mainardi. Linear models of dissipation in anelastic solids. La Rivista del Nuovo Cimento, 1:161–198, 1971.
- [17] M. Caputo and F. Mainardi. A new dissipation model based on memory mechanism. Pure and Applied Geophysics, 91:134–147, 1971.
- [18] I. Colombaro, A. Giusti, and F. Mainardi. A class of linear viscoelastic models based on Bessel functions. *Meccanica*, 52:825–832, 2017.
- [19] I. Colombaro, A. Giusti, and A. Mentrelli. Energy dissipation in viscoelastic Bessel media. Acta Mechanica, pages 1–10, 2023.
- [20] A. G. Cunha-Filho, Y. Briend, A. M. G. de Lima, and M. V. Donadon. A new and efficient constitutive model based on fractional time derivatives for transient analyses of viscoelastic systems. *Mechanical Systems* and Signal Processing, 146:107042–1–12, 2021.

- [21] N. Demirci and E. Tönük. Non-integer viscoelastic constitutive law to model soft biological tissues to in-vivo indentation. Acta of Bioengineering and Biomechanics, 16:14–21, 2014.
- [22] S. von Ende, A. Lion, and R. Lammering. On the thermodynamically consistent fractional wave equation for viscoelastic solids. Acta Mechanica, 221:1–10, 2011.
- [23] A. Giusti. On infinite order differential operators in fractional viscoelasticity. Fractional Calculus and Applied Analysis, 20:854–867, 2017.
- [24] A. Giusti and F. Mainardi. A dynamic viscoelastic analogy for fluid-filled elastic tubes. *Meccanica*, 51:2321– 2330, 2016.
- [25] W. Glöckle and T. Nonnenmacher. Fractional integral operators and Fox functions in the theory of viscoelasticity. *Macromolecules*, 24:6426–6434, 1991.
- [26] W. Glöckle and T. Nonnenmacher. Fox function representation of non-Debye relaxation processes. Journal of Statistical Physics, 71:741–757, 1993.
- [27] R. Gorenflo and F. Mainardi. Fractional Calculus: Integral and Differential Equations of Fractional Order. In: Fractals and Fractional Calculus in Continuum Mechanics (eds A. Carpinteri, F. Mainardi), volume 378 of CISM Courses and Lecture Notes. Springer Verlag, Wien and New York, 1997.
- [28] N. M. Grahovac and M. M. Žigić. Modelling of the hamstring muscle group by use of fractional derivatives. Computers and Mathematics with Applications, 59:1695–1700, 2010.
- [29] N. Heymans and J-C. Bauwens. Fractal rheological models and fractional differential equations for viscoelastic behavior. *Rheologica Acta*, 33:210–219, 1994.
- [30] S. Holm. Waves with Power-Law Attenuation. Springer Nature Switzerland AG, Cham, 2019.
- [31] S. Holm and M. B. Holm. Restrictions on wave equations for passive media. Journal of the Acoustical Society of America, 142:1888–1896, 2017.
- [32] S. Jelić and D. Zorica. Fractional Burgers wave equation on a finite domain. Chaos, Solitons, and Fractals, 154:111632–1–26, 2022.
- [33] S. Jelić and D. Zorica. Fractionalization of anti-Zener and Zener models via rheological analogy. Acta Mechanica, pages 1–42, 2022.
- [34] S. Konjik, Lj. Oparnica, and D. Zorica. Waves in fractional Zener type viscoelastic media. Journal of Mathematical Analysis and Applications, 365:259–268, 2010.
- [35] S. Konjik, Lj. Oparnica, and D. Zorica. Distributed-order fractional constitutive stress-strain relation in wave propagation modeling. Zeitschrift f
 ür angewandte Mathematik und Physik, 70:51–1–21, 2019.
- [36] F. Mainardi. Fractional Calculus and Waves in Linear Viscoelasticity. Imperial College Press, London, 2010.
- [37] F. Mainardi and G. Spada. Creep, relaxation and viscosity properties for basic fractional models in rheology. European Physical Journal Special Topics, 193:133–160, 2011.
- [38] N. Makris. The frequency response function of the creep compliance. *Meccanica*, 54:19–31, 2019.
- [39] S. P. Näsholm and S. Holm. On a fractional Zener elastic wave equation. Fractional Calculus and Applied Analysis, 16:26–50, 2013.
- [40] A. S. Okuka and D. Zorica. Formulation of thermodynamically consistent fractional Burgers models. Acta Mechanica, 229:3557–3570, 2018.
- [41] A. S. Okuka and D. Zorica. Fractional Burgers models in creep and stress relaxation tests. Applied Mathematical Modelling, 77:1894–1935, 2020.
- [42] Lj. Oparnica and E. Süli. Well-posedness of the fractional Zener wave equation for heterogenous viscoelastic materials. *Fractional Calculus and Applied Analysis*, 23:126–166, 2020.
- [43] Lj. Oparnica, D. Zorica, and A. S. Okuka. Fractional Burgers wave equation. Acta Mechanica, 230:4321– 4340, 2019.

- [44] Yu. A. Rossikhin and M. V. Shitikova. Analysis of dynamic behavior of viscoelastic rods whose rheological models contain fractional derivatives of two different orders. *Zeitschrift für angewandte Mathematik und Mechanik*, 81:363–376, 2001.
- [45] Yu. A. Rossikhin and M. V. Shitikova. Analysis of rheological equations involving more than one fractional parameters by the use of the simplest mechanical systems based on these equations. *Mechanics of Time-Dependent Materials*, 5:131–175, 2001.
- [46] Yu. A. Rossikhin and M. V. Shitikova. A new method for solving dynamic problems of fractional derivative viscoelasticity. *International Journal of Engineering Science*, 39:149–176, 2001.
- [47] Yu. A. Rossikhin and M. V. Shitikova. Analysis of the viscoelastic rod dynamics via models involving fractional derivatives or operators of two different orders. *Shock and Vibration Digest*, 36:3–26, 2004.
- [48] Yu. A. Rossikhin and M. V. Shitikova. Free damped vibrations of a viscoelastic oscillator based on Rabotnov's model. *Mechanics of Time-Dependent Materials*, 12:129–149, 2008.
- [49] Yu. A. Rossikhin and M. V. Shitikova. Application of fractional calculus for dynamic problems of solid mechanics: Novel trends and recent results. *Applied Mechanics Reviews*, 63:010801–1–52, 2010.
- [50] H. Schiessel, Chr. Friedrich, and A. Blumen. Applications to Problems in Polymer Physics and Rheology. In: Applications of Fractional Calculus in Physics (ed. R. Hilfer). World Scientific, Singapore, 2000.
- [51] H. Schiessel, R. Metzler, A. Blumen, and T. Nonnemacher. Generalized viscoelastic models: their fractional equations with solutions. *Journal of Physics A: Mathematical and General*, 28:6567–6584, 1995.
- [52] M. V. Shitikova. Fractional operator viscoelastic models in dynamic problems of mechanics of solids: A review. *Mechanics of Solids*, 57:1–33, 2022.
- [53] D. Zorica and Lj. Oparnica. Energy dissipation for hereditary and energy conservation for non-local fractional wave equations. *Philosophical Transactions of the Royal Society A*, 378:20190295–1–24, 2020.