SUPERVISED LEARNING WITH PROBABILISTIC MORPHISMS AND KERNEL MEAN EMBEDDINGS

HÔNG VÂN LÊ

Abstract. In this paper I propose a generative model of supervised learning that unifies two approaches to supervised learning, using a concept of a correct loss function. Addressing two measurability problems, which have been ignored in statistical learning theory, I propose to use convergence in outer probability to characterize the consistency of a learning algorithm. Building upon these results, I extend a result due to Cucker-Smale, which addresses the learnability of a regression model, to the setting of a conditional probability estimation problem. Additionally, I present a variant of Vapnik-Stefanuyk's regularization method for solving stochastic ill-posed problems, and using it to prove the generalizability of overparameterized supervised learning models.

CONTENTS

2010 Mathematics Subject Classification. Primary: 46N30, Secondary: 60B10, 62G05, 18N99.

Date: November 16, 2023.

Key words and phrases. Markov kernel, supervised learning model, loss function, generalization ability, inner measure, kernel mean embedding, ε -minimizer.

Research of HVL was supported by GAČR-project 22-00091S and the Czech Academy of Sciences (RVO: 67985840).

2 H. V. LÊ

1. INTRODUCTION

1.1. The concept of a correct loss function in supervised learning **theory.** Let \mathcal{X} and \mathcal{Y} be measurable spaces. In supervised learning, we are given a data set of labeled items:

$$
S_n = \{(x_1, y_1), \ldots, (x_n, y_n)\} \in ((\mathcal{X} \times \mathcal{Y})^n, \mu^n),
$$

where μ is an (unknown) probability measure that governs the distribution of i.i.d. labeled items $(x_n, y_n) \in \mathcal{X} \times \mathcal{Y}$. The goal of a learner in this scenario is to find the best approximation f_{S_n} of the stochastic relation between the input $x \in \mathcal{X}$ and its label $y \in \mathcal{Y}$, formalized as the conditional probability measure $[\mu_{\mathcal{Y}|\mathcal{X}}]$ for a probability measure μ on $(\mathcal{X} \times \mathcal{Y})$ with respect to the projection $\Pi_{\mathcal{X}} : \mathcal{X} \times \mathcal{Y} \to \mathcal{X}$. We refer to $[\mu_{\mathcal{Y}|\mathcal{X}}]$ as the supervisor operator [\[Vapnik98,](#page-50-0) p. 35-36, p. 48]. It is important to note that if $\mathcal X$ consists of a single point, this problem is equivalent to estimating and approximating probability measures on Y.

The concept of a best approximation requires a specification of a hypothesis space H of possible predictors as well a the notion of a *correct* loss function that measures the deviation of a possible predictor from the supervisor operator. The loss function concept in statistical analysis was introduced by Wald in his statistical decision theory [\[Wald50\]](#page-50-1), which can be traced back to Laplace's theory of errors, further developed by Gauss and many others [\[Sheynin77\]](#page-49-0). The importance of a correct choice of a loss function has been discussed by Chentsov. He noted that well-known examples of statistical point estimation problems, when properly formulated, admit asymptotically optimal decision rules, and the simplicity of the results proved there is largely due to the correct choice of a loss function [\[Chentsov72,](#page-48-1) p. 391].

In this article, we focus on characterizing the deviation of a predictor h in a hypothesis class H from the supervisor operator $[\mu_{\mathcal{V}|\mathcal{X}}]$, using the notion of a correct loss function in the underlying generative model of supervised learning. Our approach utilizes the concept of a probabilistic morphism, which is a categorical name for a Markov kernel and can also be interpreted

as a regular conditional probability. If $\mathcal Y$ is a separable metrizable topological space with the Borel σ -algebra $\mathcal{B}(\mathcal{Y})$, we propose an alternative characterization of a regular conditional probability measure μ_{ν} as a minimizer of mean square error on the space of all probabilistic morphisms from $\mathcal X$ to $\mathcal Y$, employing kernel mean embeddings.

1.2. Previous works. 1. In his book "Statistical Learning Theory" [\[Vapnik98\]](#page-50-0) Vapnik discussed two approaches to the learning problem: the first approach evaluates the quality of the chosen function using a risk function, while the second approach involves estimating stochastic dependencies. According to Vapnik, by estimating stochastic dependencies, problems such as pattern recognition and regression estimation can be solved effectively [\[Vapnik98,](#page-50-0) p. 19. Vapnik connected the estimation of the supervisor operator $[\mu_{\nu}^{\dagger}]$, which corresponds to estimating stochastic dependencies in supervised learning, to classical aspects of supervised learning theory (such as pattern recognition and regression estimation problems) through the Bayes decision rule [\[Vapnik98,](#page-50-0) p. 37]. It is important to note that Vapnik considered conditional probability measures $[\mu_{\nu}{}_{|\mathcal{X}}]$ but did not explicitly consider their regular versions, denoted here as $\mu_{\mathcal{Y}|\mathcal{X}}$. However, in all cases where \mathcal{X}, \mathcal{Y} are Borel subsets of \mathbb{R}^n , a regular conditional probability measure $\mu_{\mathcal{Y}|\mathcal{X}}$ exists for any Borel probability measure μ on $\mathcal{X} \times \mathcal{Y}$ with respect to the projection $\Pi_{\mathcal{X}} : \mathcal{X} \times \mathcal{Y} \to \mathcal{X}$. For further details, refer to Subsection [2.1.](#page-5-0) Vapnik addressed the problem of conditional probability estimation in multiclassification supervised learning as well as conditional density estimation in supervised learning when $\mathcal{X} \subset \mathbb{R}^n$ and $\mathcal{Y} \subset \mathbb{R}$. He proposed a class of solutions for these problems [\[Vapnik98,](#page-50-0) p. 36-39, 333-334, 337-338]. See also Examples [3.4\(](#page-18-0)3 $\&$ 4) and Remark [6.4](#page-38-1) for additional information.

2. The significance of conditional probability in classical supervised learning is evident in Bayes' decision rule, which is discussed in detail in [\[DGL96,](#page-48-2) Chapter 2]. In [\[VI2019\]](#page-50-2) Vapnik-Izmailov considered the machine learning problem as a problem of estimating the conditional probability function rather than problem of of finding function that minimizes a given loss function. In their work, Talwai, Shameli, and Simchi-Levi [\[TSS2022\]](#page-50-3) treated the estimation of conditional probability as a supervised learning model, utilizing kernel mean embedding. Their work builds upon the results of Park-Muandet [\[PM20\]](#page-49-1), where the authors formalized Grünerwälder and coauthors' proposal to use mean square loss for estimating conditional mean embedding [\[GLGB12\]](#page-49-2). For precise formulations of their results related to the present paper, refer to Remark [4.9](#page-31-3) (2 $\&$ 3) and Remark [5.2.](#page-32-0)

1.3. Main contributions. In this article, we present several contributions to statistical learning theory:

1. Introduction of the Generative Model: We introduce the concept of a generative model of supervised learning, which serves as a comprehensive framework encompassing all models in supervised learning theory, as well as models of probability measure estimation. In particular, by incorporating the concept of a correct loss function, we unify two approaches to statistical learning mentioned above in the Generative Model.

2. Characterization of Regular Conditional Probability Measures: We provide characterizations of regular conditional probability measures $\mu_{\mathcal{V}|\mathcal{X}}$, which are essential for understanding the underlying probabilistic relations in supervised learning. These characterizations allow us to establish a solid foundation for our subsequent analyses and investigations (Theorems [2.22,](#page-16-1) [4.6\)](#page-29-0).

3. Identification of Correct Loss Functions: By leveraging our characterizations of regular conditional probability measures, we identify and analyze various examples of correct loss functions. These include well-known loss functions such as the mean square error, the 0-1 loss function, the log-likelihood function and the Kullback-Leibler divergence widely used in statistics and statistical learning theory. We also highlight other natural loss functions employed in mathematical statistics and statistical learning theory (Examples [3.4,](#page-18-0) Examples [4.8\)](#page-30-0).

4. Quantifying the Consistency of a Learning Algorithm using Inner Measure: We propose the use of inner measure, equivalently, convergence in outer probability, to quantify the consistency of a learning algorithm. We also provide a sufficient condition for the consistency of a learning algorithm. These findings shed light on the potential of a learning algorithm to perform well on unseen data (Definition [3.9,](#page-22-0) Theorem [3.13,](#page-23-0) Lemma [3.15,](#page-24-1) Remarks [3.10,](#page-22-1) [3.16\)](#page-24-2). In addition, relaxing the convergence in probability in the classical theory of statistical learning by convergence in outer probability in our theory made many previous results in statistical learning theory rigorous (Remarks [3.10,](#page-22-1) [3.16,](#page-24-2) [5.2,](#page-32-0) [6.4.](#page-38-1))

5. Generalization of Cucker-Smale's Result: Building upon our characterizations of regular conditional probability measures, we present a generalization of Cucker-Smale's result (Theorem [5.1,](#page-31-2) Remark [5.2\)](#page-32-0).

6. Variant of Vapnik-Stefanyuk's Regularization Method: We introduce a variant of Vapnik-Stefanyuk's regularization method for solving stochastic ill-posed problems. By incorporating inner measure, we demonstrate its effectiveness in proving the learnability of conditional probability estimation problems (Theorem [6.1,](#page-36-0) Remark [6.4,](#page-38-1) Theorem [6.5,](#page-38-0) Corollary [6.13.](#page-44-0) Example [6.14\)](#page-46-0). As a result, we provide examples of learnable "overparameterized" discriminative models of supervised learning (Corollary [6.13\(](#page-44-0)2)).

1.4. Organization of this article. This article is organized into several sections as follows.

Section [2:](#page-4-0) Bounded s-Probabilistic Morphisms. In this section, we introduce the concept of a bounded s-probabilistic morphism, which extends the notion of a probabilistic morphism. Bounded s-probabilistic morphisms are useful for our investigation of variations of probabilistic morphisms. We characterize a regular conditional probability measure $\mu_{\mathcal{V}|\mathcal{X}}$ among bounded

s-probabilistic morphisms as a solution to a linear operator equation (Theorem [2.22\)](#page-16-1). Additionally, this section includes various technical results that will be utilized in later parts of the paper.

Section [3:](#page-17-0) Generative Model of Supervised Learning. In Section [3,](#page-17-0) we utilize probabilistic morphisms to introduce the concept of a generative model of supervised learning and the notion of a correct loss function. Drawing on the results from Section [2,](#page-4-0) we illustrate these concepts using examples such as the Fisher-Wald setting of density estimation and Vapnik's setting of conditional probability estimation. We propose the concept of consistency of a learning algorithm using inner measure (Definition [3.9\)](#page-22-0), which is equivalent to the concept of convergence in outer probability (Remark [3.10\)](#page-22-1), and provide a sufficient condition for the consistency of a learning algorithm (Theorem [3.13,](#page-23-0) Lemma [3.15\)](#page-24-1). Furthermore, we discuss the relationship between these concepts and classical notions of consistency (Remarks [3.10,](#page-22-1) [3.16\)](#page-24-2). We also introduce the concept of a C-empirical risk minimizing (C-ERM) algorithm, which guarantees the existence of such algorithms for any statistical learning model (Definition [3.11,](#page-22-2) Remark [3.12\)](#page-23-1).

Section [4:](#page-24-0) Regular Conditional Probability Measures via Kernel Mean Embedding. Assuming that $\mathcal Y$ is a separable metrizable topological space, this section builds upon the results of Section [2](#page-4-0) to characterize regular conditional probability measures μ_{ν} as the minimizers of mean square error (Theorem [4.6,](#page-29-0) Corollary [4.7\)](#page-30-1). We discuss related findings by Park-Muandet [\[PM20\]](#page-49-1), Grünerwalder et al. [\[GLGB12\]](#page-49-2), and Talwai-Shami-Simchi-Levi [\[TSS2022\]](#page-50-3) in the context of our analysis (Remark [4.9](#page-31-3) $(2 \& 3)$).

Section [5:](#page-31-0) Generalization of Cucker-Smale's Result. Drawing on the previous sections, we present a generalization of a result originally established by Cucker-Smale (Theorem [5.1\)](#page-31-2). We also discuss related findings and implications in Remark [5.2.](#page-32-0)

Section [6](#page-35-0) : Variant of Stefanyuk-Vapnik's Result and Applications. In this section, we present a variant of Vapnik's result that incorporates inner measure (Theorem [6.1\)](#page-36-0). We explore its applications in conditional probability estimation problems, discussing the implications in Remark [6.2,](#page-36-1) Theorem [6.5,](#page-38-0) Corollary [6.13](#page-44-0) and Example [6.14.](#page-46-0)

Last Section [7:](#page-47-0) Discussion of Results. In the final section of the paper, we summarize the main findings, highlight their significance, and offer potential future research directions.

2. Bounded s-probabilistic morphisms

In this section we introduce the concept of a *bounded s-probabilistic mor*phism and compare it with the concept of a probabilistic morphism and with the concept of an s-finite kernel (Definitions [2.5,](#page-8-1) Remark [2.6\)](#page-8-2). We then study the properties of bounded s-probabilistic morphisms (Definition [2.16,](#page-13-0) Lemmas [2.12,](#page-11-0) [2.14,](#page-12-0) [2.21\)](#page-15-0). Finally, we provide a characterization of regular conditional measures using bounded s-probabilistic morphisms (Theorem [2.22\)](#page-16-1).

2.1. Notation, conventions, and preliminaries.

• Given a measurable space \mathcal{X} , let $\Sigma_{\mathcal{X}}$ denote the σ -algebra of \mathcal{X} . We use $\mathcal{S}(\mathcal{X}), \mathcal{M}(\mathcal{X})$ and $\mathcal{P}(\mathcal{X})$ to represent the space of all finite signed measures, the space of all finite (nonnegative) measures, and the space of all probability measures on \mathcal{X} , respectively. The total variation norm on $\mathcal{S}(\mathcal{Y})$ is denoted as $\|\cdot\|_{TV}$.

• For a given $x \in \mathcal{X}$, δ_x represents the Dirac measure concentrated at x.

• For a data set $S = (x_1, \ldots, x_n) \in \mathcal{X}^n$, μ_S denotes the empirical probability measure defined as

$$
\mu_S := \frac{1}{n} \sum_{i=1}^n \delta_{x_i} \in \mathcal{P}(\mathcal{X}).
$$

• $\mathcal{P}_{emp}(\mathcal{X})$ represents the set of all empirical probability measures on \mathcal{X} .

• When X is a topological space, we consider the Borel σ -algebra $\mathcal{B}(\mathcal{X})$ unless stated otherwise.

• For a measurable mapping $f : \mathcal{X} \to \mathcal{Y}$ and $\mu \in \mathcal{S}(\mathcal{X})$, $f_*\mu$ denotes the pushforwarded measure on \mathcal{Y} , defined as $f_*\mu(B) := \mu(f^{-1}(B))$ for any $B \in \Sigma_{\mathcal{Y}}$.

• The product $\mathcal{X}\times\mathcal{Y}$ of measurable spaces $(\mathcal{X},\Sigma_{\mathcal{X}})$ and $(\mathcal{Y},\Sigma_{\mathcal{Y}})$ is assumed to be endowed with the σ -algebra $\Sigma_{\mathcal{X}} \otimes \Sigma_{\mathcal{Y}}$ unless otherwise stated. For a probability measure $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y}), \mu_{\mathcal{X}} := (\Pi_{\mathcal{X}})_*\mu$ represents the marginal probability measure of μ on \mathcal{X} .

• In this paper we consider only measurable spaces \mathcal{X}, \mathcal{Y} such that any $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ has a regular conditional probability measure $\mu_{\mathcal{Y}|\mathcal{X}}$ for μ relative to $\Pi_{\mathcal{X}}$. If \mathcal{Y} is a Souslin measurable space (a measurable space isomorphic to a measurable space associated with a Souslin metrizable space [\[DM1978,](#page-48-3) Definition 16, p.46-III], [\[JLT21,](#page-49-3) Remark 1]), then this regular conditional probability measure $\mu_{\mathcal{V}}|_{\mathcal{X}}$ always exists, [\[Bogachev07,](#page-48-4) Corollary 10.4.15, p. 366, vol. 2], [\[LFR04,](#page-49-4) Theorem 3.1 (6)].

• Meas $(\mathcal{X}, \mathcal{Y})$ represents the space of all measurable mappings from X to Y. If \mathcal{X}, \mathcal{Y} are topological spaces, $C(\mathcal{X}, \mathcal{Y})$ denotes the space of all continuous mappings from $\mathcal X$ to $\mathcal Y$.

• Id_X represents the identity mapping on X.

• For $A \in \Sigma_{\mathcal{X}}$, 1_A denotes the characteristic function of A.

• $\mathcal{F}_s(\mathcal{X})$ denotes the vector space of simple (step) functions on X and $\mathcal{F}_{b}(\mathcal{X})$ denotes the vector space of measurable bounded functions on X. $\mathcal{F}_b(\mathcal{X})$ is a Banach space equipped with the sup-norm $\|\cdot\|_{\infty}$.

• We endow $\mathcal{S}(\mathcal{X})$ with the σ -algebra Σ_w , which is the smallest σ -algebra such that for any $f \in \mathcal{F}_s(\mathcal{X})$, the map

$$
I_f: \mathcal{S}(\mathcal{X}) \to \mathbb{R}, \mu \mapsto \int_{\mathcal{X}} f \, d\mu,
$$

is measurable. The restriction of Σ_w to $\mathcal{P}(\mathcal{X})$ and $\mathcal{M}(\mathcal{X})$ is also denoted by Σ_w [\[Lawere62\]](#page-49-5), [\[JLT21\]](#page-49-3).

• Let $C_b(\mathcal{X})$ be the space of all bounded continuous functions on a topological space X. We denote by τ_w the weakest topology on $\mathcal{S}(\mathcal{X})$ such that for any $f \in C_b(\mathcal{X})$ the map $I_f : (\mathcal{S}(\mathcal{X}), \tau_w) \to \mathbb{R}$ is continuous. We also denote by τ_w the restriction of τ_w to $\mathcal{M}(\mathcal{X})$ and $\mathcal{P}(\mathcal{X})$. If X is separable and metrizable then $(\mathcal{P}(\mathcal{X}), \tau_w)$ is separable and metrizable [\[Bogachev18,](#page-48-5) Theorem 3.1.4, p. 104], [\[Parthasarathy67,](#page-49-6) Theorem 6.2, p.43], and the Borel σ-algebra $\mathcal{B}(\tau_w)$ on $\mathcal{P}(\mathcal{X})$ generated by τ_w coincides with Σ_w [\[GH89,](#page-49-7) Theorem 2.3].

Lemma 2.1. For any $h \in \mathcal{F}_b(\mathcal{X})$ the evaluation mapping $I_h : (\mathcal{S}(\mathcal{X}), \Sigma_w) \to$ $\mathbb{R}, \mu \mapsto \int_{\mathcal{X}} h d\mu$, is a measurable mapping. Consequently, Σ_w is the smallest σ-algebra such that $I_h : (\mathcal{S}(\mathcal{X}), \Sigma_w) \to \mathbb{R}$ is measurable for any $h \in \mathcal{F}_b(\mathcal{X})$.

Proof. Let $h \in \mathcal{F}_b(\mathcal{X})$. Then there exist sequences of simple functions $\{h_n^{\pm}, n \in \mathbb{N}^+\}$ such that

$$
h_n^-(x) \le h(x) \le h_n^+(x) \text{ for all } x \in \mathcal{X} \text{ and } ||h_n^- - h_n^+||_{\infty} \le \frac{1}{n},
$$

see e.g. [\[Chentsov72,](#page-48-1) p. 66]. It follows that for any $\mu \in \mathcal{S}(\mathcal{X})$ we have

(2.1)
$$
\lim_{n \to \infty} \int_{\mathcal{X}} h_n^{\pm} d\mu = \int_{\mathcal{X}} h d\mu.
$$

In other words, the sequence of functions $I_{h_n^{\pm}} : \mathcal{S}(\mathcal{Y}) \to \mathbb{R}$ converges to I_h pointwise. Hence, I_h is also measurable. The last assertion follows since $\mathcal{F}_s(\mathcal{X}) \subset \mathcal{F}_b(\mathcal{X}).$

Lemma 2.2. Assume that X is a complete separable metric space. Let $\mathcal{A} := \{A_i | i \in \mathbb{N}^+\}$ be the collection of closed balls of rational radius centered at a countable dense subset in X. Then for any $A \in \Sigma_{\mathcal{X}}$ and $\mu \in \mathcal{S}(\mathcal{X})$ we have

(2.2)
$$
|\mu|(A) = \sup\{|\mu|(A \cap \bigcup_{i=1}^{N} A_i) : N \in \mathbb{N}^+, A_i \in \mathcal{A}\}.
$$

Proof. Since X is a complete metric space, μ is a Radon measure, see e.g. [\[Bogachev07,](#page-48-4) Theorem 7.4.3, p. 85, vol. 2], i.e. for any $\varepsilon > 0$ there exists a compact set $K_{\varepsilon} \subset A$ such that

$$
(2.3) \t\t |\mu|(A \setminus K_{\varepsilon}) \leq \varepsilon.
$$

Let $\{A_1, \ldots, A_n \in \mathcal{A}\}\$ be a finite cover of K_{ε} . Then it follows from (2.3)

$$
(2.4) \qquad |\mu|(A) \leq \sup\{|\mu|(A \cap \bigcup_{i=1}^{N} A_i) : N \in \mathbb{N}^+, A_i \in \mathcal{A}\} + \varepsilon.
$$

Taking into account

$$
|\mu|(A) \ge \sup\{|\mu|(A \cap \bigcup_{i=1}^{N} A_i) : N \in \mathbb{N}^+, A_i \in \mathcal{A}\},\
$$

this completes the proof of Lemma [2.2.](#page-6-1) \Box

8 H. V. LÊ

Corollary 2.3. Let $\mathcal{F}_\mathcal{A}$ be the countable family consisting of all finite disjoint unions of elements in the countable algebra $G(\mathcal{A})$ generated by \mathcal{A} . Then we have

(2.5)
$$
|\mu|(A) = \sup \{ \sum_{i=1}^{N} |\mu(A \cap B_i)| : \dot{\cup}_{i=1}^{N} B_i \in \mathcal{F}_A, B_i \in G(\mathcal{A}) \}.
$$

Proof. Since $\dot{\cup}_{i=1}^{N} B_i$ is a disjoint union, we have

(2.6)
$$
|\mu|(A) \ge \sup \{ \sum_{i=1}^{N} |\mu(A \cap B_i)| : \dot{\cup}_{i=1}^{N} B_i \in \mathcal{F}_A, B_i \in G(\mathcal{A}) \}.
$$

Now let $\mathcal{X} = \mathcal{X}^+ \cup \mathcal{X}^-$ be the Hahn decomposition of X for μ and $\mu =$ $\mu^+ - \mu^-$ be the Jordan-Hahn decomposition of μ . Then

(2.7)
$$
|\mu|(A) = |\mu(A^+)| + |\mu(A^-)|, \text{ where } A^{\pm} := \mathcal{X}^{\pm} \cap A.
$$

Applying Lemma [2.2](#page-6-1) to μ , A^{\pm} we obtain from [\(2.7\)](#page-7-0)

(2.8)
$$
|\mu|(A) \leq \sup \{ \sum_{i=1}^{N} |\mu(A \cap B_i)| : \bigcup_{i=1}^{N} B_i \in \mathcal{F}_A, B_i \in G(\mathcal{A}) \}.
$$

Taking into account [\(2.6\)](#page-7-1), this completes the proof of Corollary [2.3.](#page-7-2) \Box

Proposition 2.4 (Measurability of the Jordan-Hahn Decomposition and Total Variation). Assume that X is a Polish space. The map $v : (\mathcal{S}(\mathcal{X}), \Sigma_w)$ \rightarrow $(\mathcal{M}(\mathcal{X}), \Sigma_w), \mu \mapsto |\mu|$, is a measurable function. Consequently, the Jordan-Hahn decomposition $S(\mathcal{X}) \to \mathcal{M}(\mathcal{X}) \times \mathcal{M}(\mathcal{X}), \mu \mapsto (\mu^+, \mu^-),$ is also measurable mapping. Additionally, the map $S(\mathcal{X}) \to \mathbb{R}, \mu \mapsto ||\mu||_{TV}$, is measurable.

Proof. By the definition of Σ_w , to prove that the map $v : (\mathcal{S}(\mathcal{X}), \Sigma_w) \to$ $\mathcal{M}(\mathcal{X}), \mu \mapsto |\mu|$, is measurable, it suffices to show that for any $A \in \Sigma_{\mathcal{X}}$ the composition: $I_{1_A} \circ v : \mathcal{S}(\mathcal{X}) \to \mathbb{R}_{\geq 0}$ is a measurable function. Let $\mathcal{F}_{\mathcal{A}} := \{\mathcal{F}_1, \ldots, \mathcal{F}_n, \ldots\}$. By Corollary [2.3,](#page-7-2) we have

(2.9)
$$
I_{1_A} \circ \mathbf{v}(\mu) = \sup_i f_i^A(\mu),
$$

where

$$
f_i^A(\mu) = \sum_{B_{ij} \in \mathcal{F}_i} |\mu(A \cap B_{ij})| : \dot{\cup}_{j=1}^n B_{ij} = \mathcal{F}_i \in \mathcal{F}_A, B_i \in G(\mathcal{A}).
$$

Since \mathcal{F}_i is fixed, taking into account that the function $\mathbb{R} \to \mathbb{R}_{\geq 0}, x \mapsto |x|$, is measurable, the function $f_i^A : \mathcal{S}(\mathcal{X}) \to \mathbb{R}, \mu \mapsto f_i^A(\mu)$, is also measurable for all *i*. Since for any $-\infty < a < \infty$ we have

$$
(\sup_{i} f_{i}^{A})^{-1} ((-\infty, b]) = \bigcap_{k=1}^{\infty} \bigcup_{i=1}^{\infty} (f_{i}^{A})^{-1} (-\infty, b + \frac{1}{k}],
$$

taking into account [\(2.9\)](#page-7-3) and the fact that $\mathcal{B}(\mathbb{R})$ is generated by the sets ${(-\infty, b], b \in \mathbb{R}}$, we conclude that the function $I_{1_A} \circ \mathbf{v} : \mathcal{S}(\mathcal{X}) \to \mathbb{R}$ is a measurable. This completes the proof of the first aseertion of Proposition [2.4.](#page-7-4)

The second and third assertion of Proposition [2.4](#page-7-4) follow immediately. \Box

2.2. Bounded s-probabilistic morphisms and their joints. Follow-ing Lawvere [\[Lawere62\]](#page-49-5), we consider the σ -algebra Σ_w on $\mathcal{P}(\mathcal{Y})$ and treat Markov kernels from X to Y as measurable mappings from X to $\mathcal{P}(y)$.

Definition 2.5. cf. [\[JLT21,](#page-49-3) Definition 1], cf. [\[Lawere62\]](#page-49-5). Let \mathcal{X}, \mathcal{Y} be measurable space. A bounded s-probabilistic morphism $T : \mathcal{X} \rightarrow \mathcal{Y}$ is a measurable mapping \overline{T} : $\mathcal{X} \to (\mathcal{S}(\mathcal{Y}), \Sigma_w)$ such that the mapping \overline{T} : $\mathcal{X} \to$ $\mathcal{S}(\mathcal{Y})_{TV}$ is bounded, i.e. there exists a constant $C > 0$ such that $\|\overline{T}(x)\|_{TV} \leq$ C for all $x \in \mathcal{X}$. In this case we shall say that $\overline{T} : \mathcal{X} \to \mathcal{S}(\mathcal{Y})_{TV}$ is a bounded *measurable map.* A bounded s-probabilistic morphism $T : \mathcal{X} \rightarrow \mathcal{Y}$ shall be called a probabilistic morphism, if $\overline{T}(\mathcal{X}) \subset \mathcal{P}(\mathcal{Y})$.

Remark 2.6. (1) The space of all bounded s-probabilistic morphisms is a normed vector space with the sup-norm $||T||_{\infty} := \sup_{x \in \mathcal{X}} ||\overline{T}(x)||_{TV}$.

(2) A kernel k from X to Y is function $k : \mathcal{X} \times \Sigma_{\mathcal{Y}} \to [0,\infty]$ such that for any $x \in \mathcal{X}$ the function $k(x, \cdot) : \Sigma_{\mathcal{Y}} \to [0, \infty]$ is a measure and for any $B \in \Sigma_{\mathcal{Y}}$ the function $k(\cdot, B) : \mathcal{X} \to [0, \infty]$ is measurable. A kernel k is finite, if there is a finite $r \in [0,\infty)$ such that $k(x,\mathcal{Y}) < r$ for all $x \in$ X . Clearly any finite kernel k generates a bounded measurable mapping $k : \mathcal{X} \to (\mathcal{M}(\mathcal{X}), \Sigma_w), x \mapsto k(x, \cdot)$, and hence a bounded s-probabilistic morphism. Proposition [2.4](#page-7-4) implies that if $\mathcal Y$ is a Polish space, then any bounded measurable mapping $\overline{T} : \mathcal{X} \to \mathcal{S}(\mathcal{Y})$ can be written as $T = T^+ - T^-$, where T^{\pm} : $\mathcal{X} \to \mathcal{M}(\mathcal{Y})$ are finite kernels.

(3) An s-finite kernel is a map $k : \mathcal{X} \times \Sigma_{\mathcal{Y}} \to [0, \infty]$ such that there exists a sequence k_1, \ldots, k_n of finite kernels such that $k = \sum_{i=1}^{\infty} k_i$ [\[Staton17,](#page-49-8) Definition 2]. The space of all s-finite kernels is a cone. According to Staton, the concept of an s-finite kernel has been introduced by Kallenberg [\[Kallenberg2014\]](#page-49-9) and by Last-Penrose [\[LP16\]](#page-49-10).

Let $\textbf{sbProbm}(\mathcal{X}, \mathcal{Y})$ denote the vector space of all bounded s-probabilistic morphisms from $\mathcal X$ to $\mathcal Y$ and by $\textbf{Probm}(\mathcal X, \mathcal Y)$ the set of all probabilistic morphisms from $\mathcal X$ to $\mathcal Y$.

For a bounded s-probabilistic morphism $T : \mathcal{X} \rightarrow \mathcal{Y}$, we denote the associated measurable mapping as \overline{T} : $\mathcal{X} \to \mathcal{S}(\mathcal{Y})$. Conversely, for a bounded measurable mapping $f : \mathcal{X} \to \mathcal{S}(\mathcal{Y})$ we denote the generated bounded sprobabilistic morphism as $f : \mathcal{X} \rightarrow \mathcal{Y}$.

Examples 2.7. (i) Any regular conditional probability measure $\mu_{\mathcal{V}|\mathcal{X}}$ for $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ with respect to the projection $\Pi_{\mathcal{X}} : \mathcal{X} \times \mathcal{Y} \to \mathcal{X}$ assigns a measurable mapping $\mu_{\mathcal{Y}|\mathcal{X}} : \mathcal{X} \to \mathcal{P}(\mathcal{Y})$ by the formula $\mu_{\mathcal{Y}|\mathcal{X}}(x)(A) =$ $\mu_{\mathcal{V}|\mathcal{X}}(A|x)$ for $x \in \mathcal{X}$ and $A \in \Sigma_{\mathcal{Y}}$. We shall also call the map $\mu_{\mathcal{Y}|\mathcal{X}}$: $\mathcal{X} \to \mathcal{P}(\mathcal{Y})$ a regular conditional probability measure for μ . The associated probabilistic morphism is denoted by $\mu_{\mathcal{Y}|\mathcal{X}} : \mathcal{X} \rightarrow \mathcal{Y}$.

10 H. V. LEˆ

(ii) The map $\delta : \mathcal{X} \to (\mathcal{P}(\mathcal{X}), \Sigma_w), x \mapsto \delta(x) := \delta_x$, is measurable [\[Lawere62\]](#page-49-5), [\[Giry82,](#page-49-11) §1.2]. If $\mathcal X$ is a topological space, then the map δ : $\mathcal{X} \to (\mathcal{P}(\mathcal{X}), \tau_w)$ is continuous, since the composition $I_f \circ \delta : \mathcal{X} \to \mathbb{R}$ is continuous for any $f \in C_b(\mathcal{X})$. We regard a measurable mapping $\kappa : \mathcal{X} \to \mathcal{Y}$ as a deterministic probabilistic morphism defined by $\overline{\kappa} := \delta \circ \kappa : \mathcal{X} \to \mathcal{P}(\mathcal{Y})$. In particular, the identity mapping $\mathrm{Id}_{\mathcal{X}} : \mathcal{X} \to \mathcal{X}$ of a measurable space \mathcal{X} is a probabilistic morphism generated by $\delta : \mathcal{X} \to \mathcal{P}(\mathcal{X})$, so $\delta = \text{Id}_{\mathcal{X}}$. Graphically speaking, any straight arrow (a measurable mapping) $\kappa : \mathcal{X} \to \mathcal{Y}$ between measurable spaces can be seen as a curved arrow (a probabilistic morphism).

Given a bounded s-probabilistic morphism, following Chentsov [\[Chentsov72,](#page-48-1) Lemma 5.9, p. 72, we define a linear map $S_*(T): S(\mathcal{X}) \to S(\mathcal{Y})$ as follows

(2.10)
$$
S_*(T)(\mu)(B) := \int_{\mathcal{X}} \overline{T}(x)(B) d\mu(x).
$$

for any $\mu \in \mathcal{S}(\mathcal{X})$ and $B \in \Sigma_{\mathcal{Y}}$. Following [\[Chentsov72,](#page-48-1) (5.1), p. 66], we define a linear map $T^* : \mathcal{F}_b(\mathcal{Y}) \to \mathbb{R}^{\mathcal{X}}$ by letting

(2.11)
$$
T^*(f)(x) := \int_{\mathcal{Y}} f d\overline{T}(x) \text{ for } x \in \mathcal{X}.
$$

Lemma 2.8. The map T^* is a bounded linear map between Banach spaces $\mathcal{F}_b(\mathcal{Y})_\infty$ and $\mathcal{F}_b(\mathcal{X})_\infty$.

Proof. We shall use the Chentsov argument in his proof for the case that $T \in$ **Probm** (X, Y) [\[Chentsov72,](#page-48-1) Corollary of Lemma 5.1, p. 66]. First we shall show that $T^*(\mathcal{F}_b(\mathcal{Y})) \subset \mathcal{F}_b(\mathcal{X})$. Note that $T^*(1_B)(\cdot) = \overline{T}(\cdot)(B) \in \mathcal{F}_b(\mathcal{X})$, since $\overline{T}(x)(B) \leq ||T(x)||_{TV}$. Now let $h \in \mathcal{F}_b(\mathcal{Y})$. Then there exist sequence of simple functions $\{h_n^{\pm}, n \in \mathbb{N}^+\}$ on $\mathcal Y$ such that

(2.12)
$$
h_n^-(y) \le h(y) \le h_n^+(y)
$$
 for all $y \in \mathcal{Y}$ and $||h_n - h_n^+||_{\infty} \le \frac{1}{n}$.

Let $x \in \mathcal{X}$. Then

$$
T^*h(x) = \int_{\mathcal{Y}} h \, d\overline{T}(x)^+ - \int_{\mathcal{Y}} h \, \overline{T}(x)^-.
$$

For $x \in \mathcal{X}$ and $f \in \mathcal{F}_b(\mathcal{Y})$, we shall use the following shorthand notation

$$
(T^{\pm})^*(f)(x) := \int_{\mathcal{Y}} f d\overline{T}(x)^{\pm}.
$$

By [\(2.12\)](#page-9-0), for any $x \in \mathcal{X}$ and any $n \in \mathbb{N}^+$, we obtain

$$
(2.13) \qquad (T^{\pm})^*(h_n^{-})(x) \le (T^{\pm})^*(h)(x) \le (T^{\pm})^*(h_n^{+})(x).
$$

From [\(2.12\)](#page-9-0) and [\(2.13\)](#page-9-1), it follows that for any $x \in \mathcal{X}$

$$
\lim_{n \to \infty} (T^{\pm})^*(h_n^{-})(x) = (T^{\pm})^*h(x) = \lim_{n \to \infty} (T^{\pm})^*(h_n^{+})(x).
$$

Hence, for any $x \in \mathcal{X}$ we have $\lim_{n \to \infty} T^*h(x) = \lim_{n \to \infty} T^*(h_n^{\pm})$. It follows that $T^*h \in \mathcal{F}_b(\mathcal{X})$, since we have shown that $T^*(h_n^{\pm}) \in \mathcal{F}_b(\mathcal{X})$.

Finally, we note that $||T^*f||_{\infty} \le ||T||_{\infty} \cdot ||f||_{\infty}$. Hence, T^* is a bounded linear map. This completes the proof of Lemma [2.8.](#page-9-2)

 \Box

For $\overline{T} \in \text{Meas}(\mathcal{X}, (\mathcal{S}(\mathcal{Y}), \Sigma_w)), x \in \mathcal{X}, \text{ and } y \in \mathcal{Y}, \text{ we shall write}$ $d\overline{T}(y|x)$ instead of $d\overline{T}(x)(y)$, which has been denoted as $T(dy|x)$ by Chentsov [\[Chentsov72\]](#page-48-1).

Definition 2.9. Given' $T_1 \in \text{sbProbm}(\mathcal{X}, \mathcal{Y}), T_2 \in \text{sbProbm}(\mathcal{Y}, \mathcal{Z}),$ the composition $\overline{T_2 \circ T_1} : \mathcal{X} \to \mathcal{S}(\mathcal{Z})$ is defined as follows

(2.14)
$$
\overline{T_2 \circ T_1}(x, C) = \int_{\mathcal{Y}} \overline{T}_2(y, C) d\overline{T}_1(y|x)
$$

for any $x \in \mathcal{X}, C \in \Sigma_{\mathcal{Z}}$.

Lemma 2.10. (1) The composition $T_2 \circ T_1$ of two bounded s-probabilistic morphisms is a bounded s-probabilistic morphism.

(2) Furthermore, the composition is associative, i.e. $(T_3 \circ T_2) \circ T_1 =$ $T_3 \circ (T_2 \circ T_1).$

(3) For any $T_i \in \mathbf{sbfProbm}(\mathcal{X}_i, \mathcal{X}_{i+1}), i \in [1, 2],$ we have

(2.15)
$$
\overline{T_2 \circ T_1} = S_*(T_2) \circ \overline{T_1}.
$$

Lemma [2.10](#page-10-0) for probabilistic morphisms are well-known [\[Giry82\]](#page-49-11), see also [\[Chentsov72,](#page-48-1) Lemmas 5.4-5.6, p. 68-69].

Proof of Lemma [2.10.](#page-10-0) (1) First we note that $\overline{T_2 \circ T_1}$ is bounded map from $\mathcal{X} \to \mathcal{S}(\mathcal{Z})_{TV}$, since \overline{T}_1 and \overline{T}_2 are bounded mappings. To show that $T_2 \circ T_1 : \mathcal{X} \to (\mathcal{S}(\mathcal{Z}), \Sigma_w)$ is a measurable map, it suffices to show that for any $C \in \Sigma_{\mathcal{Z}}$ the function

$$
I_{1_C}: \mathcal{X} \to \mathbb{R}, x \mapsto I_{1_C}(\overline{T_2 \circ T_1}(x))
$$

is measurable. Since

$$
I_{1_C}(\overline{T_2 \circ T_1}(x)) = \int_{\mathcal{Y}} \overline{T}_2(y, C) d\overline{T}_1(y|x) = (T_1)^*(\overline{T}_2(\cdot|C))(x),
$$

by Lemma [2.8,](#page-9-2) the function $I_{1_C}(\overline{T_2 \circ T_1})$ belongs to $\mathcal{F}_b(\mathcal{X})$. This proves the first assertion of Lemma [2.10.](#page-10-0)

(2) Let $T_1 \in \mathbf{s}$ **bProbm** $(\mathcal{X}, \mathcal{Y}), T_2 \in \mathbf{s}$ **bProbm** $(\mathcal{Y}, \mathcal{Z})$ and $T_3 \in \mathbf{s}$ **bProbm** $(\mathcal{Z}, \mathcal{V})$. To prove Lemma [2.10](#page-10-0) (2), we have to show that for any $x \in \mathcal{X}$ and $D \in \Sigma_{\mathcal{V}}$ we have:

$$
(2.16)
$$

$$
\int_{\mathcal{Y}} \int_{\mathcal{Z}} \overline{T_3}(z)(D) d\overline{T_2}(z|y) d\overline{T_1}(y|x) = \int_{\mathcal{Z}} \int_{\mathcal{Y}} \overline{T_3}(z)(D) d\overline{T_2}(z|y) d\overline{T_1}(y|x).
$$

In order to prove [\(2.16\)](#page-10-1), we shall approximate the bounded measurable function $T_3(\cdot)(D)$ on Z by step functions $h_n \in \mathcal{F}_s(\mathcal{Z})$. This reduces to 12 H. V. LÊ

verify [\(2.16\)](#page-10-1) for the case that $\overline{T_3}(\cdot)(D) = 1_C$ for some $C \in \Sigma_{\mathcal{Z}}$. In this case straightforward computations yield

$$
\int_{\mathcal{Y}} \int_{\mathcal{Z}} 1_C \overline{T_2}(y, dz) \overline{T_1}(x, dy) = \int_{\mathcal{Y}} \overline{T_2}(y) (C) \overline{T_1}(x, dy) = \int_{\mathcal{Z}} \int_{\mathcal{Y}} 1_C \overline{T_2}(y, dz) \overline{T_1}(x, dy).
$$

This completes the proof of Lemma [2.10\(](#page-10-0)2).

(3) To prove the last assertion of Lemma [2.10,](#page-10-0) we notice that $\overline{T_2 \circ T_1}(x)(C) =$ $S_*(T_2) \circ \overline{T_1}(x)(C)$ for any $x \in \mathcal{X}_1$ and $C \in \Sigma_{\mathcal{X}_3}$ by comparing [\(2.14\)](#page-10-2) with (2.10). This completes the proof of Lemma 2.10. [\(2.10\)](#page-9-3). This completes the proof of Lemma [2.10.](#page-10-0)

Remark 2.11. (1) If T is a probabilistic morphism, then the restriction $M_*(T)$ of $S_*(T)$ to $\mathcal{M}(\mathcal{X})$ and the restriction $P_*(T)$ of $S_*(T)$ to $\mathcal{P}(\mathcal{X})$ maps $\mathcal{M}(\mathcal{X})$ to $\mathcal{M}(\mathcal{Y})$ and $\mathcal{P}(\mathcal{X})$ to $\mathcal{P}(\mathcal{Y})$, respectively [\[Chentsov72,](#page-48-1) Lemma 5.9, p. 72].

(2) If a probabilistic morphism is deterministic, i.e. it is generated by a measurable mapping $\overline{\kappa} = \delta \circ \kappa : \mathcal{X} \to \mathcal{P}(\mathcal{Y})$, then $S_*(\kappa) : \mathcal{S}(\mathcal{X}) \to \mathcal{S}(\mathcal{Y})$ is the push-forward operator $\kappa_* : \mathcal{S}(\mathcal{X}) \to \mathcal{S}(\mathcal{Y})$, i.e.

(2.17)
$$
\mathcal{S}_{*}(\kappa)(\mu)(B) = \mu(\kappa^{-1}(B)) \text{ for } \mu \in \mathcal{S}(\mathcal{X}) \text{ and } B \in \Sigma_{\mathcal{Y}}.
$$

(3) A bounded measurable mapping \overline{T} : $\mathcal{X} \to (\mathcal{S}(\mathcal{Y}), \Sigma_w)$ generates two homomorphisms: $\overline{T}_* : \mathcal{S}(\mathcal{X}) \to \mathcal{S}(\mathcal{S}(\mathcal{Y}), \Sigma_w)$ and $\overline{T}_* : \mathcal{S}(\mathcal{X}) \to \mathcal{S}(\mathcal{Y})$.

In view of Remark [2.11,](#page-11-1) we will use the shorthand notation T_* for $S_*(T)$, $M_*(T)$ and $P_*(T)$ when referring to a s-probabilistic morphism T, as long as there is no possiblity of confusion or misunderstanding.

Lemma 2.12. Assume that $T \in \mathbf{Probm}(\mathcal{X}, \mathcal{Y})$.

(1) Then $T_* : (\mathcal{S}(\mathcal{X}), \Sigma_w) \to (\mathcal{S}(\mathcal{Y}), \Sigma_w)$ is a measurable mapping. In particular, the map $T_* : (\mathcal{P}(\mathcal{X}), \Sigma_w) \to (\mathcal{P}(\mathcal{Y}), \Sigma_w)$ is measurable.

(2) If $\overline{T} \in C(\mathcal{X},(\mathcal{P}(\mathcal{Y}), \tau_w))$ then $T_* : (\mathcal{S}(\mathcal{X}), \tau_w) \to (\mathcal{S}(\mathcal{Y}), \tau_w)$ is a continuous mapping. In particular, the map $T_* : (\mathcal{M}(\mathcal{X}), \tau_w) \to (\mathcal{M}(\mathcal{Y}), \tau_w)$ is continuous.

Proof. (1) To prove the first assertion of Lemma [2.12\(](#page-11-0)1), it suffices to show that for any $f \in \mathcal{F}_b(\mathcal{Y})$ the composition

$$
I_f \circ T_* : (\mathcal{S}(\mathcal{X}), \Sigma_{\omega}) \to \mathbb{R}, \mu \mapsto I_f \circ T_*(\mu),
$$

is a measurable mapping. Note that

$$
I_f \circ T_*(\mu) = \int_{\mathcal{Y}} f dT_* \mu \stackrel{(2.10)}{=} \int_{\mathcal{X}} \int_{\mathcal{Y}} f d\overline{T}(x) d\mu(x) \stackrel{(2.11)}{=} \int_{\mathcal{X}} T^*(f) d\mu.
$$

Hence

$$
(2.18) \t\t I_f \circ T_* = I_{T^*f}.
$$

Using Lemma [2.1](#page-6-2) we conclude the first assertion of Lemma [2.12\(](#page-11-0)1) from [\(2.18\)](#page-11-2). The second assertion of Lemma [2.12\(](#page-11-0)1) follows immediately.

(2) The proof of the second assertion is similar to the the proof of the first one, noting that if \overline{T} : $\mathcal{X} \to (\mathcal{P}(\mathcal{Y}), \tau_w)$ is continuous, then for any $f \in C_b(\mathcal{Y})$ the function

$$
\mathcal{X} \to \mathbb{R}, x \mapsto T^*f(x) = \int_{\mathcal{Y}} f d\overline{T}(x) = (f\overline{T}(x))(\mathcal{Y}),
$$

is continuous. Hence the composition

$$
I_f \circ T_* : (\mathcal{S}(\mathcal{X}), \tau_w) \to \mathbb{R}, \, \mu \mapsto I_f \circ T_*(\mu) = I_{T^*f}(\mu),
$$

is continuous. This completes the proof of Lemma [2.12.](#page-11-0) \Box

Remark 2.13. The assertion "in particular" of Lemma [2.12](#page-11-0) (1) has been stated by Lawvere [\[Lawere62\]](#page-49-5) and proved by Giry [\[Giry82\]](#page-49-11).

Now we are going to define the *joint of two bounded s-probabilistic mor*phisms with the same source. First we need the following.

Lemma 2.14. (1) The multiplication mapping

$$
\mathfrak{m}:\left(\mathcal{S}(\mathcal{X}),\Sigma_w\right)\times\left(\mathcal{S}(\mathcal{Y}),\Sigma_w\right)\rightarrow\mathcal{S}(\mathcal{X}\times\mathcal{Y},\Sigma_w),\,(\mu,\nu)\mapsto\mu\cdot\nu
$$

is measurable. If X and Y are topological spaces, then \mathfrak{m} is (τ_w, τ_w) -continuous. (2) The diagonal mapping

diag:
$$
(\mathcal{S}(\mathcal{X}), \Sigma_w) \to (\mathcal{S}(\mathcal{X} \times \mathcal{X}), \Sigma_w), \mu \mapsto \mu^2
$$

is measurable. If X is a topological space, then diag is (τ_w, τ_w) -continuous. (3) For any $n \in \mathbb{N}^+$ the addition $\mathfrak{a}^n : (\mathcal{S}(\mathcal{X})^n, \otimes^n \Sigma_w) \to (\mathcal{S}(\mathcal{X}), \Sigma_w)$,

 $(\mu_1, \ldots, \mu_n) \mapsto \sum_{i=1}^n \mu_i$, is measurable. Consequently the n-Dirac map

$$
\delta^n: \mathcal{X}^n \to (\mathcal{S}(\mathcal{X}), \Sigma_w), \, S \mapsto \mu_S,
$$

is also measurable. If X is a topological space then δ^n is τ_w -continuous.

Proof. (1) To prove that the map $\mathfrak m$ is measurable, it suffices to show that for any $A \in \Sigma_{\mathcal{X}}$ and $B \in \Sigma_{\mathcal{Y}}$ the map $I_{1_{A \otimes B}} : (\mathcal{S}(\mathcal{X}), \Sigma_{w}) \times (\mathcal{S}(\mathcal{Y}), \Sigma_{w}) \to$ $\mathbb{R}, (\mu, \nu) \mapsto \mu(A)\nu(B)$, is measurable. The map $I_{1_{A\otimes B}}$ is measurable, since it can be written as the composition of measurable mappings

(2.19)
$$
(\mathcal{S}(\mathcal{X}), \Sigma_w) \times (\mathcal{S}(\mathcal{Y}), \Sigma_w) \stackrel{(1_A,1_B)}{\longrightarrow} \mathbb{R} \times \mathbb{R} \longrightarrow^{\mathfrak{m}_{\mathbb{R}}}
$$

where $\mathfrak{m}_{\mathbb{R}}(x, y) = x \cdot y$.

Similarly, we prove that \mathfrak{m} is (τ_w, τ_w) -continuous, if X and Y are topological spaces, since all the mappings in [\(2.19\)](#page-12-1) are continuous.

(2) To prove the second assertion, we write diag $= \mathfrak{m} \circ \text{Diag}$, where

Diag:
$$
(\mathcal{S}(\mathcal{X}), \Sigma_w) \to (\mathcal{S}(\mathcal{X}), \Sigma_w) \times (\mathcal{S}(\mathcal{X}), \Sigma_w), \mu \mapsto (\mu, \mu).
$$

Taking into account the measurability of the map Diag and of the map m, we conclude that diag is a measurable map.

If X and Y are topological spaces, then Diag is a continuous map. Taking into account Lemma [2.14\(](#page-12-0)1), we prove the continuity of diag.

(3) First we prove the case $n = 2$. To prove the measurability of the map \mathfrak{a}^2 it suffices to show that for any $f \in \mathcal{F}_s(\mathcal{X})$ the composition $I_f \circ \mathfrak{a}$: $(\mathcal{S}(\mathcal{X}) \times \mathcal{S}(\mathcal{X}), \Sigma_w \otimes \Sigma_w) \to \mathbb{R}$ is measurable. Using the formula

$$
I_f\circ \mathfrak{a}(\mu,\nu)=I_{1_{\mathcal{X}}}(f\mu)+I_{1_{\mathcal{X}}}(f\nu),
$$

we reduce the proof of the measurability of $I_f \circ \mathfrak{a}$ to proving the measurability of the map $\mathfrak{a}^2 : \mathbb{R} \times \mathbb{R} \to \mathbb{R}, (x, y) \mapsto (x + y)$, which is well-known.

In the same way, we prove the continuity of the map \mathfrak{a}^2 , if X is a topological space.

For $n \geq 2$ we use the formula $\mathfrak{a}^n(\mu_1, \ldots, \mu_n) = \mathfrak{a}^2(\mathfrak{a}^{n-1}(\mu_1, \ldots, \mu_{n_1}), \mu_n)$ and taking into account the validity of the assertion for $n = 2$ we prove the first part of assertion (3).

Similarly, the proof of continuity of δ^n if X is topological space can be reduced to the case $n = 2$, which is well-known.

This completes the proof of Lemma [2.14.](#page-12-0)

$$
\qquad \qquad \Box
$$

Remark 2.15. For the general abstract story behind the formation of the map m, defined in Lemma [2.14,](#page-12-0) see Kock [\[Kock11\]](#page-49-12), and Fritz-Perrone-Rezagholi [\[FPR2021\]](#page-48-6) for a similar result.

From now on we shall drop Σ_w , as long as there is no possibility of confusion or misunderstanding.

Definition 2.16. (1) Given two bounded s-probabilistic morphisms T_i : $\mathcal{X} \rightsquigarrow \mathcal{Y}_i$ i = 1,2, the joint of T_1 and T_2 is the bounded s-probabilistic morphisms $T_1 \cdot T_2 : \mathcal{X} \to \mathcal{Y}_1 \times \mathcal{Y}_2$ whose generating mapping $\overline{T_1 \cdot T_2} : \mathcal{X} \to$ $\mathcal{S}(\mathcal{Y}_1 \times \mathcal{Y}_2)$ is given by:

$$
\overline{T_1 \cdot T_2}(x) = \mathfrak{m}(\overline{T}_1(x), \overline{T}_2(x)) : \mathcal{X} \to \mathcal{S}(\mathcal{Y}_1 \times \mathcal{Y}_2).
$$

(2) Given a bounded s-probabilistic morphism $T : \mathcal{X} \rightarrow \mathcal{Y}$ we denote the joint of Id_X with T by $\Gamma_T : \mathcal{X} \to \mathcal{X} \times \mathcal{Y}$ and call it the graph of T.

It follows from Definition [2.16\(](#page-13-0)2) that for any $T \in s**b**$ Probm $(\mathcal{X}, \mathcal{Y})$, $A \in \Sigma_{\mathcal{X}}$ and $B \in \Sigma_{\mathcal{Y}}$, we have

$$
(2.20)\ \ (\Gamma_T)_*\mu_X(A \times B) = \int_{\mathcal{X}} \overline{\Gamma_T}(x) (A \times B) \, d\mu_X(x) = \int_A \overline{T}(x) (B) \, d\mu_X(x).
$$

Hence, for any $f \in \mathcal{F}_b(\mathcal{X} \times \mathcal{Y})$, we have

(2.21)
$$
\int_{\mathcal{X}\times\mathcal{Y}} f(x,y) d(\Gamma_T)_*\mu_{\mathcal{X}}(x,y) = \int_{\mathcal{X}} \int_{\mathcal{Y}} f(x,y) d\overline{T}(y|x)\mu_{\mathcal{X}}(x).
$$

Remark 2.17. (1) The notion of a graph of a probabilistic morphism f has been appeared first in Jost-Lê-Tran [\[JLT21\]](#page-49-3), the arXiv version, but without a definition. The first definition of this concept has been given in Fritz-Gonda-Perrone-Rischel's paper [\[FGPR2020\]](#page-48-7), where they call the graph of f the input-copy version or bloom of f.

(2) If $\kappa : \mathcal{X} \to \mathcal{Y}$ is a measurable mapping, then $\overline{\Gamma_{\kappa}}(x, y) = \delta_x \delta_{\kappa(x)} =$ $\delta_{(x,\kappa(x))}$. Thus Γ_{κ} is the graph of κ , i.e. $\Gamma_{\kappa}(x,y)=(x,\kappa(y)).$

We shall show the (τ_w, τ_w) -continuity of the map $(\Gamma_T)_*$ under certain conditions.

Proposition 2.18. Assume that X is a topological space, Y is a compact metrizable topological space, and \overline{T} : $\mathcal{X} \to (\mathcal{P}(\mathcal{Y}), \tau_w)$ is a continuous mapping. Then \overline{T} is a Markov kernel, i.e. \overline{T} : $\mathcal{X} \to (\mathcal{P}(\mathcal{Y}), \Sigma_w)$ is a measurable mapping. Furthermore the map

$$
(\Gamma_T)_*: (\mathcal{S}(\mathcal{X}), \tau_w) \to (\mathcal{S}(\mathcal{X} \times \mathcal{Y}), \tau_w), \mu \mapsto (\Gamma_T)_* \mu,
$$

is a continuous map.

Proof. Assume the condition of Proposition [2.18.](#page-14-0) Then \overline{T} is a Markov kernel, since Y is a separable metrizable topological space and $\mathcal{B}(\tau_w) = \Sigma_w$ by [\[GH89,](#page-49-7) Theorem 2.3].

Now let us prove the (τ_w, τ_w) -continuity of $(\Gamma_T)_*$. Since $(\Gamma_T)_*$ is a linear map, it suffices to show that

$$
\tau_{w} \text{-} \lim_{n \to \infty} (\Gamma_T)_* \mu_n = 0 \in (\mathcal{S}(\mathcal{X} \times \mathcal{Y}), \tau_w) \text{ if } \tau_{w} \text{-} \lim_{n \to \infty} \mu_n = 0 \in (\mathcal{S}(\mathcal{X}), \tau_w).
$$

Thus it suffices to show that if τ_w - lim_{n→∞} $\mu_n = 0$ then for any $f \in C_b(\mathcal{X} \times \mathcal{Y})$ we have

(2.22)
$$
\lim_{n \to \infty} \int_{\mathcal{X} \times \mathcal{Y}} f d(\Gamma_T)_* \mu_n = 0.
$$

Lemma 2.19. Assume the conditions of Proposition [2.18.](#page-14-0) Then for any $f \in C_b(\mathcal{X} \times \mathcal{Y})$ the function $F_{T,f} : \mathcal{X} \to \mathbb{R}$

$$
F_{T,f}(x) := \int_{\mathcal{Y}} f(x,y) d\overline{T}(y|x),
$$

belongs to $C_b(\mathcal{X})$.

Proof. Since f is bounded, $F_{T,f} : \mathcal{X} \to \mathbb{R}$ is bounded. To show that $F_{T,f}$ is continuous, it suffices to show that

(2.23)
$$
\lim_{x' \to x} F_{T,f}(x') = F_{T,f}(x).
$$

We write

$$
F_{T,f}(x') - F_{T,f}(x) = \int_{\mathcal{Y}} \left(f(x, \cdot) - f(x', \cdot) \right) dT(x) + \int_{\mathcal{Y}} f(x', \cdot) d(T(x) - T(x')).
$$

Since $\mathcal Y$ is compact,

$$
\lim_{x' \to x} ||f(x', \cdot) - f(x, \cdot)||_{\infty} = 0.
$$

It follows that

$$
\lim_{x' \to x} \int_{\mathcal{Y}} \left(f(x, \cdot) - f(x', \cdot) \right) dT(x) = 0.
$$

Since $T: \mathcal{X} \to (\mathcal{P}(\mathcal{Y}), \tau_w)$ is continuous

$$
\lim_{x' \to x} \int_{\mathcal{Y}} f(x', \cdot) d(T(x) - T(x')) = 0,
$$

Thus (2.23) holds. This completes the proof of Lemma [2.19.](#page-14-2)

16 H. V. LEˆ

Completion of the proof of Proposition [2.18](#page-14-0). Let $f \in C_b(\mathcal{X} \times \mathcal{Y})$. Then

$$
\int_{\mathcal{X}\times\mathcal{Y}} f d(\Gamma_T)_*\mu_n = \int_{\mathcal{X}} F_{T,f} d\mu_n.
$$

By Lemma [2.19,](#page-14-2) $F_{T,f} \in C_b(\mathcal{X})$. Hence Equation [\(2.22\)](#page-14-3) holds, if $\lim_{n\to\infty} \mu_n = 0 \in (S(\mathcal{X}) \subset \mathcal{X})$. This completes the proof of Proposition 2.18 $0 \in (\mathcal{S}(\mathcal{X}), \tau_w)$. This completes the proof of Proposition [2.18.](#page-14-0)

Next we present a concept of almost surely equality of two bounded sprobabilistic morphisms, which is consistent with the concept of almost equality in Markov category proposed by Fritz [\[Fritz2019,](#page-48-8) Example 13.3, §13].

Definition 2.20. Let $\mu \in \mathcal{P}(\mathcal{X})$. Two bounded measurable mappings T, T' : $\mathcal{X} \to \mathcal{S}(\mathcal{Y})$ will be called *equal* μ *-a.e.* (with the shorthand notation $T = T'$ μ -a.e.), if for any $B \in \Sigma_{\mathcal{Y}}$

$$
\mu\{x \in \mathcal{X} : T(x)(B) \neq T'(x)(B)\} = 0.
$$

Lemma 2.21. (1) Assume that $\mu_{\mathcal{X}} \in \mathcal{P}(\mathcal{X})$ and $T_1, T_2 \in \textbf{sbProbm}(\mathcal{X}, \mathcal{Y})$. Then $(\Gamma_{T_1})_* \mu_X = (\Gamma_{T_2})_* \mu_X$ if and only if $T_1 = T_2 \mu_X$ -a.e.

(2) Assume that $\mu_X \in \mathcal{S}(\mathcal{X})$ and $T_1, T_2 \in \mathbf{Probm}(\mathcal{X}, \mathcal{Y})$. Then

- $(\Gamma_{T_1})_*\mu_{\mathcal{X}} = (\Gamma_{T_2})_*\mu_{\mathcal{X}}$ if and only if $T_1 = T_2$ $|\mu_{\mathcal{X}}|$ -a.e.
	- (3) We have $\Gamma_{T_1+T_2} = \Gamma_{T_1} + \Gamma_{T_2}$.

(4) For any $T \in {\bf sbProblem}(\mathcal{X}, \mathcal{Y})$ we have

$$
(2.24) \t\t T = \Pi_{\mathcal{Y}} \circ \Gamma_T.
$$

Proof. (1) Assume the condition of Lemma [2.21.](#page-15-0) To prove the first assertion of Lemma [2.21](#page-15-0) it suffices to show that $(\Gamma_T)_*\mu_{\mathcal{X}}=0$ if and only if $\overline{T}=0$ $\mu_{\mathcal{X}}$ -a.e.

The 'iff" assertion follows immediately from [\(2.10\)](#page-9-3), so let us prove the "only if" assertion. Since for any $B \in \Sigma_{\mathcal{Y}}$ the function $\overline{T}(\cdot)(B) : \mathcal{X} \to \mathbb{R}$ is measurable, B_{χ}^{+} $\chi^+_{\mathcal{X}} := \{x \in \mathcal{X} | T(x)(B) > 0\}$ is a measurable subset of \mathcal{X} . By the assumption, we have

$$
(\Gamma_T)_*\mu_{\mathcal{X}}(B^+_{\mathcal{X}} \times B) = \int_{B^+_{\mathcal{X}}} \overline{T}(x)(B) d\mu_{\mathcal{X}}(x) = 0.
$$

Since $\overline{T}(x)(B) > 0$ for all $x \in B_{\mathcal{X}}^{+}$ χ^+ , it follows that $\mu_{\mathcal{X}}(B^+_{\mathcal{X}})$ χ^+ = 0. Similarly we prove $\mu_{\mathcal{X}}(B_{\mathcal{X}}^{-})$ $\overline{\mathcal{X}}$) = 0, where $B_{\mathcal{X}}^ \overline{\chi} := \{x \in \mathcal{X} | T(x)(B) < 0\}.$ This proves the first assertion.

(2) The second assertion of Lemma [2.21](#page-15-0) follows from the first one, taking into account the Jordan-Hahn decomposition of $\mu_{\mathcal{X}}$ and Remark [2.11](#page-11-1) (1).

(3) The third assertion of Lemma [2.21](#page-15-0) is straightforward.

(4) We verify immediately from Definition [2.16](#page-13-0) (2) that

(2.25)
$$
\overline{T} = (\Pi_{\mathcal{Y}})_* \circ \overline{\Gamma_T}.
$$

Combining [\(2.25\)](#page-15-1) with [\(2.15\)](#page-10-3), we complete the proof of the last assertion of Lemma [2.21.](#page-15-0) \Box

Lemma [2.21](#page-15-0) motivates the following notation. Given $\mu \in \mathcal{M}(\mathcal{X})$ we denote by $\text{Meas}(\mathcal{X}, \mathcal{S}(\mathcal{Y}); \mu)$ the quotient space $\text{Meas}(\mathcal{X}, \mathcal{S}(\mathcal{Y}))$ under μ -a.e. equality and by $[T]_u$, **Meas** $(\mathcal{X}, \mathcal{P}(\mathcal{Y}); \mu)$ the image of $T \in \mathbf{Meas}(\mathcal{X}, \mathcal{S}(\mathcal{Y}))$ and of $\text{Meas}(\mathcal{X}, \mathcal{P}(\mathcal{Y}))$, respectively, in the quotient space.

2.3. A characterization of regular conditional probability measures. For $p \ge 1$ and a σ -finite measure μ on X we set [\[Le2022,](#page-49-13) §3.3], cf. [\[AJLS17,](#page-48-9) (3.51), p. 144]

$$
\mathcal{S}^{1/p}(\mathcal{X}, \mu) := \{ \nu \in \mathcal{S}(\mathcal{X}) | \nu \ll \mu \text{ and } \frac{d\nu}{d\mu} \in L^p(\mu) \}.
$$

The natural identification $S^{1/p}(\mathcal{X}, \mu) = L^p(\mu)$ defines a p-norm on $S^{1/p}(\mathcal{X}, \mu)$ by setting

$$
||f\mu||_p = ||f||_{L^p(\mu)}.
$$

Then $S^{1/p}(\mathcal{X},\mu)$ endowed with the p-norm is a Banach space, which we denote by $S^{1/p}(\mathcal{X}, \mu)_p$. For $\mu_1 \ll \mu_2$ the linear inclusion

$$
(2.26) \qquad \mathcal{S}^{1/p}(\mathcal{X}, \mu_1) \to \mathcal{S}^{1/p}(\mathcal{X}, \mu_2), \ f\mu_1 \mapsto f(\frac{d\mu_1}{d\mu_2})^{1/p}\mu_2 \in \mathcal{S}^{1/p}(\mathcal{X}, \mu_2)
$$

preserves the p-norm.

For a σ -finite measure μ on $\mathcal X$ we set $\mathcal P^{1/p}(\mu) := \mathcal P(\mathcal X) \cap \mathcal S^{1/p}(\mathcal X, \mu)$.

Theorem 2.22. (1) A bounded measurable mapping \overline{T} : $\mathcal{X} \rightarrow \mathcal{S}(\mathcal{Y})$ is a regular conditional probability measure for $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ with respect to the projection $\Pi_{\mathcal{X}}$ if and only if

$$
(2.27) \t\t (\Gamma_T)_*\mu_X = \mu.
$$

(2) If \overline{T} , \overline{T}' : $\mathcal{X} \to \mathcal{P}(\mathcal{Y})$ are regular conditional probability measures for $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ with respect to the projection $\Pi_{\mathcal{X}}$ then $\overline{T} = \overline{T}' \mu_{\mathcal{X}}$ -a.e.

(3) Assume that $T \in \mathbf{Probm}(\mathcal{X}, \mathcal{Y})$ and ρ_0 is a σ -finite measure on $\mathcal Y$ such that $T(x) \ll \rho_0$ for all $x \in \mathcal{X}$. Let ν_0 be a σ -finite measure on X and $\nu \in \mathcal{P}^{1/p}(\mathcal{X}, \nu_0)$ for some $p \geq 1$. Then $(\Gamma_T)_* \nu \in \mathcal{S}^{1/p}(\mathcal{X} \times \mathcal{Y}, \nu_0 \rho_0)$.

Proof. (1) Assume that $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y}), T \in \mathbf{sProbm}(\mathcal{X}, \mathcal{Y})$ and [\(2.27\)](#page-16-2) holds. Then by (2.25) , T must be a probabilistic morphism. Now recall that for any $A \in \Sigma_{\mathcal{X}}$, $B \in \Sigma_{\mathcal{Y}}$ we have

$$
(\Gamma_T)_*\mu_{\mathcal{X}}(A \times B) = \int_{\mathcal{X}} \overline{\Gamma_T}(x) (A \times B) \, d\mu_{\mathcal{X}}(x) = \int_A \overline{T}(x) (B) \, d\mu_{\mathcal{X}}(x),
$$

This implies that T is a regular conditional measure for μ with respect to the projection $\Pi_{\mathcal{X}}$. This proves the "if" assertion.

Conversely, if \overline{T} : $\mathcal{X} \to \mathcal{S}(\mathcal{Y})$ is a regular conditional probability measure for $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ then $\overline{T}(\mathcal{X}) \subset \mathcal{P}(\mathcal{X})$ and [\(2.20\)](#page-13-1) holds. Taking into account (2.10) , we conclude that (2.27) holds. This completes the proof of assertion (1) of Theorem [2.22.](#page-16-1)

(2) The second assertion follows from the first one and Lemma [2.21\(](#page-15-0)1).

(3) Let $T \in \text{Probm}(\mathcal{X}, \mathcal{Y})$ and $T(x) \ll \rho_0$ for all $x \in \mathcal{X}$. We write $T(x) = g(\cdot|x)\rho_0$ where $g(\cdot|x)$ is the density of $T(x)$ with respect to ρ_0 . Taking into account the disintegration formula for $f \in \mathcal{F}_s(\mathcal{X} \times \mathcal{Y}),$

(2.28)
$$
\int_{\mathcal{X}\times\mathcal{Y}} f(x,y) d(\Gamma_T)_*\nu(x,y) = \int_{\mathcal{X}\times\mathcal{Y}} f(x,y)g(y|x) d\rho_0(y)d\nu(x),
$$

it follows that $(\Gamma_T)_*\nu = g\nu \rho_0$. Hence

$$
(2.29) \t\t (\Gamma_T)_*\nu \ll \nu_0 \rho_0.
$$

Since $\nu \in \mathcal{P}^{1/p}(\mathcal{X}, \nu)$, by [\[AJLS17,](#page-48-9) Corollary 5.1, p. 260], we have

(2.30)
$$
(\Gamma_T)_*\nu \in \mathcal{S}^{1/p}(\mathcal{X}\times\mathcal{Y}, (\Gamma_T)_*\nu).
$$

Taking into account [\(2.28\)](#page-17-2), we obtain immediately the last assertion of The-orem [2.22.](#page-16-1)

Remark 2.23. In the setting of Markov categories, Equation [\(2.27\)](#page-16-2) for $T \in \mathbf{Probm}(\mathcal{X}, \mathcal{Y})$ is exactly the definition of conditional [\[CJ19,](#page-48-10) §3].

3. Generative models of supervised learning

In this section, we introduce the concept of a generative model of supervised learning (Definition [3.1\)](#page-17-3), which encompasses various models used for multi-classification, regression tasks, conditional probability estimation and probability measure estimation. We also present the concept of a correct loss function (Definition [3.3\)](#page-18-1), which includes many commonly used loss functions in classical statistics and statistical learning theory (Examples [3.4\)](#page-18-0). Additionally, we discuss important properties of inner and outer measures. Finally, we explore general statistical learning models (Definition [3.8\)](#page-21-1), their learnability (Definition [3.9,](#page-22-0) Remark [3.10\)](#page-22-1), and provide a sufficient condition for the learnability of a statistical learning model (Theorem [3.13,](#page-23-0) Lemma [3.15\)](#page-24-1) by introducing the concept of a C-ERM algorithm (Definition [3.11,](#page-22-2) Remark [3.12\)](#page-23-1).

3.1. Generative models of supervised learning.

Definition 3.1. A generative model of supervised learning is defined as a quintuple $(\mathcal{X}, \mathcal{Y}, \mathcal{H}, R, \mathcal{P}_{\mathcal{X}\times\mathcal{Y}})$, where X and Y are measurable spaces, H is a family of measurable mappings $h : \mathcal{X} \to \mathcal{P}(\mathcal{Y}), \mathcal{P}_{\mathcal{X} \times \mathcal{Y}} \subset \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ contains all possible probability measures that govern the distributions of labeled pairs (x, y) , and $R: \mathcal{H} \times (\mathcal{P}_{\mathcal{X} \times \mathcal{Y}} \cup \mathcal{P}_{emp}(\mathcal{X} \times \mathcal{Y})) \to \mathbb{R} \cup \{+\infty\}$ is a risk/loss function. It should satisfy the condition that for any $\mu \in \mathcal{P}_{\mathcal{X}\times\mathcal{Y}} \cup \mathcal{P}_{emp}(\mathcal{X}\times\mathcal{Y})$ we have $\inf_{h\in\mathcal{H}} R(h,\mu) \neq \pm \infty$. If $R(h,\mu) = \mathbb{E}_{\mu}(L(h))$ where $L : \mathcal{X} \times \mathcal{Y} \times \mathcal{H} \rightarrow$ $\mathbb{R}\{\infty\}$ is an instantaneous loss function then we can represent the model as $(X, Y, H, L, \mathcal{P}_{X \times Y})$. A classical generative model of supervised learning is a special case of a generative model $(\mathcal{X}, \mathcal{Y}, \mathcal{H}, L, \mathcal{P}_{\mathcal{X}\times\mathcal{Y}})$, where \mathcal{H} is a family of dominated regular conditional probability measures, i.e., there exists a σ-finite measure $ρ_0$ on Y such that $h(x) \ll ρ_0$ for all $x \in \mathcal{X}$, $h \in \mathcal{H}$.

Remark 3.2. (1) If X consists of a single point $\{pt\}$, then $\mathcal{P}_{\{pt\}\times\mathcal{Y}}$ can be identified with a statistical model $\mathcal{P}_{\mathcal{Y}} \subset \mathcal{P}(\mathcal{Y})$. In this case it is natural to assume that the set $\{h({\tau}\) | h \in \mathcal{H}\}\$ is identified with $\mathcal{P}_\mathcal{V}$ and our model $({\{pt\}},\mathcal{Y},\mathcal{H}\cong \mathcal{P}_{\mathcal{Y}},R,\mathcal{P}_{\mathcal{Y}})$ of supervised learning is a model of density estimation, or more precisely, a model of probability measure estimation in unsupervised learning.

(2) If R is generated by an instantaneous loss function $L : \mathcal{X} \times \mathcal{Y} \times$ $\mathcal{H} \to \mathbb{R} \cup \{+\infty\}$, then for any $S = (x_1, y_1, \dots, x_n, y_n) \in (\mathcal{X} \times \mathcal{Y})^n$ we have $R_{\mu_S}^L(h) = \frac{1}{n} \sum_{i=1}^n L(x_i, y_i, h).$

(3) Generative models of supervised learning encompass discriminative models of supervised learning where the hypothesis space H consists of measurable mappings. For classical multi-classification and regression tasks, as well as models for learning conditional density functions and conditional probability considered by Vapnik (see Examples $3.4(3 \& 4)$ below), generative models provide a comprehensive framework.

Definition 3.3. A loss function $R: \mathcal{H} \times (\mathcal{P}_{\mathcal{X} \times \mathcal{Y}} \cup \mathcal{P}_{\text{emp}}(\mathcal{X} \times \mathcal{Y})) \rightarrow \mathbb{R} \cup \{+\infty\}$ will be called $\mathcal{P}_{\mathcal{X}\times\mathcal{Y}}$ -correct, if there exists a set $\widetilde{\mathcal{H}} \subset \mathbf{Meas}(\mathcal{X}, \mathcal{P}(\mathcal{Y}))$ such that the following three conditions hold

(1) $\mathcal{H} \subset \widetilde{\mathcal{H}}$.

(2) For any $\mu \in \mathcal{P}_{\mathcal{X}\times\mathcal{Y}}$ there exists $h \in \mathcal{H}$ such that h is a regular conditional measure for μ relative to the projection $\Pi_{\mathcal{X}}$, i.e.. $[h]_{\mu_{\mathcal{X}}} = [\mu_{\mathcal{Y}|\mathcal{X}}]$.

(3) R is the restriction of a loss function \widetilde{R} : $\widetilde{\mathcal{H}} \times (\mathcal{P}_{\mathcal{X} \times \mathcal{Y}} \cup \mathcal{P}_{emp}(\mathcal{X} \times \mathcal{Y})) \rightarrow$ $\mathbb{R} \cup \{+\infty\}$ such that for any $\mu \in \mathcal{P}_{\mathcal{X}\times\mathcal{Y}}$

$$
\arg\min_{h\in\widetilde{\mathcal{H}}} \widetilde{R}(h,\mu) = \{h\in\widetilde{\mathcal{H}}|\,[h]_{\mu_{\mathcal{X}}} = [\mu_{\mathcal{Y}|\mathcal{X}}]\}.
$$

A loss function $R : \mathcal{H} \times (\mathcal{P}_{\mathcal{X} \times \mathcal{Y}} \cup \mathcal{P}_{emp}(\mathcal{X} \times \mathcal{Y})) \rightarrow \mathbb{R} \cup \{+\infty\}$ will be called *correct*, if R is the restriction of a $\mathcal{P}(\mathcal{X} \times \mathcal{Y})$ -correct loss function $\widetilde{R}: \mathcal{H} \times \mathcal{P}(\mathcal{X} \times \mathcal{Y}) \to \mathbb{R} \cup \{+\infty\}.$

Given a σ -finite measure μ on $\mathcal X$ we denote by $\mathcal L^1(\mathcal X, \mu)$ the set of all μ integrable functions on X and by $L^1(\mathcal{X}, \mu)$ the set of $\stackrel{\mu}{\sim}$ equivalence classes in $\mathcal{L}^1(\mathcal{X}, \mu)$ where $f \stackrel{\mu}{\sim} g$ iff $f - g = 0$ μ -a.e.

Examples 3.4. (1) Let $\mathcal{X} = \{pt\}$, $\mathcal{Y} = \mathbb{R}$, and $\mathcal{H} = \mathcal{P}(\mathcal{Y}, dy)$ the set of all probability measures *f dy* where *dy* is the Lebesgue measure on $\mathcal{Y} = \mathbb{R}$ and $f \in \mathcal{L}^1(\mathbb{R}, dy)$. Let $\mathcal{P}_\mathcal{Y} = \mathcal{P}(\mathcal{Y}, dy)$. We define the minus log-likehood instantaneous loss function $L : \mathcal{H} \times \mathbb{R} \to \mathbb{R}$ by setting

$$
L(f, y) = -\log f(y),
$$

which leads to the loss function $R^L : \mathcal{H} \times \mathcal{P}(\mathbb{R}, dy) \to \mathbb{R}$, and hence, for any $\mu \in \mathcal{P}(\mathbb{R}, dy)$, to the expected loss function

$$
R^L_\mu: \mathcal{H} \to \mathbb{R}, f dy \mapsto -\int_{\mathbb{R}} \log f d\mu.
$$

Now we write $\mu = pdy$ where $p \in \mathcal{L}^1(\mathbb{R}, dy)$. By the Bretagnolle-Huber inequality, given in the following form [\[Vapnik98,](#page-50-0) (1.11), p.30]

$$
\int_{\mathbb{R}} |f(y) - p(y)| dy \le 2\sqrt{1 - \exp(R^L_\mu(\mu) - R^L_\mu(fdy))},
$$

the function $R^L : \mathcal{H} \times \mathcal{P}(\mathbb{R}, dy) \to \mathbb{R}$ is a $\mathcal{P}(\mathbb{R}, dy)$ -correct loss function.

(2) Let $(X, Y, H, \mathcal{P}_{X \times Y})$ be an arbitrary geometric model of supervised learning, i.e. a quadruple underlying a supervised learning model $(\mathcal{X}, \mathcal{Y}, \mathcal{H}, L, \mathcal{P}_{\mathcal{X}\times\mathcal{Y}})$. We defines a loss function: $R^{(k)}$: $\mathcal{H} \times \mathcal{P}_{\mathcal{X} \times \mathcal{Y}} \to \mathbb{R}_{\geq 0}$ as follows

(3.1)
$$
R^{(k)}(h,\mu) := \|(\Gamma_h)_*\mu_X - \mu\|_{TV}^k.
$$

By Theorem [2.22](#page-16-1) (1), R is a correct loss function for any $k \in \mathbb{N}^+$.

Now assume that $\mathfrak{M} : \mathcal{P}(\mathcal{X} \times \mathcal{Y}) \to E$ is an embedding from $\mathcal{P}(\mathcal{X} \times \mathcal{Y})$ to a metric vector space (E, d) . By Theorem [2.22\(](#page-16-1)1), the function

(3.2)
$$
R^d: \mathcal{H} \times \mathcal{P}_{\mathcal{X} \times \mathcal{Y}} \to \mathbb{R}_{\geq 0}, (h,\mu) \mapsto d\Big(\mathfrak{M}\big((\Gamma_h)_*\mu_{\mathcal{X}}\big), \mathfrak{M}(\mu)\Big),
$$

is a correct loss function, since \mathfrak{M} is injective.

(3) Assume that $\mathcal{Y} = {\omega_1, \ldots, \omega_n}$ is a finite sample space consisting of *n* elements, and $\mathcal{X} \subset \mathbb{R}^n$. Let $h \in \mathbf{Probm}(\mathcal{X}, \mathcal{Y})$. In this case the equation $(\Gamma_h)_*\mu_{\mathcal{X}} = \mu$ is equivalent to the following equation:

(3.3)
$$
\int_{-\infty}^{a} \overline{h}(x)(\omega_i) dF_{\mu\chi}(x) = F_{\mu}(\omega_i, a) \text{ for } i \in [1, n] \text{ and } a \in \mathbb{R}^n
$$

where $F_{\mu\chi}$ and F_{μ} are cummulative distribution function of μ_{χ} and of μ , respectively, and the integral in the LHS of [\(3.3\)](#page-19-0) is the Lebesgue-Stieltjets integral. Note that the LHS and RHS of (3.3) is well-defined, if μ , and hence $\mu_{\mathcal{X}}$, are empirical measures. That is Vapnik's equation (1.20) in [\[Vapnik98,](#page-50-0) p. 36] for conditional probability. Assuming further that F_{μ} belongs to a metric space (E, d) , we obtain the following correct loss function:

(3.4)
$$
R: \text{Probm}(\mathcal{X}, \mathcal{Y}) \times \mathcal{P}(\mathcal{X} \times \mathcal{Y}) \to \mathbb{R}_{\geq 0}, R(h, \mu) = d(F_{(\Gamma_h)_* \mu_{\mathcal{X}}}, F_{\mu}).
$$

(4) Assume that $\mathcal{X} = \mathbb{R}^m$ and $\mathcal{Y} = \mathbb{R}^n$. Let dy denote the Lebesgue measure on \mathbb{R}^n and $\mathcal{H} \subset \mathbf{Meas}(\mathbb{R}^m, \mathcal{P}(\mathbb{R}^n, dy))$. For $h \in \mathcal{H}$ we write

$$
h(x) := \bar{h}(\cdot|x)dy \in \mathcal{P}(\mathbb{R}^n),
$$

where the density function $\bar{h}(\cdot|x)$ belongs to $\mathcal{L}^1(\mathbb{R}^n, dy)$ for all $x \in \mathbb{R}^m$. By Tonelli's theorem, $\bar{h} \in \mathcal{L}^1(\mathbb{R}^m \times \mathbb{R}^n, dxdy)$, where dx is the Lebesgue measure on \mathbb{R}^m s. Let $\mu \in \mathcal{P}(\mathbb{R}^m \times \mathbb{R}^n, dxdy)$. By [\(2.30\)](#page-17-4), $\mu_{\mathcal{X}} \in \mathcal{P}(\mathbb{R}^m, dx)$. By Theorem $2.22(5)$, we have

(3.5)
$$
(\Gamma_h)_*(\mu_{\mathcal{X}}) \in \mathcal{S}(\mathbb{R}^m \times \mathbb{R}^n, dxdy).
$$

Now we shall rewrite Equation [\(2.27\)](#page-16-2), using cummulative distribution functions as in Example [3.4\(](#page-18-0)3), so that the loss function $R_{\mu}: \mathcal{H} \to \mathbb{R}$ is also well-defined for any $\mu \in \mathcal{P}_{emp}(\mathcal{X} \times \mathcal{Y})$. By Theorem [2.22](#page-16-1) (1), h is a

$$
20\quad
$$

conditional density for $\mu \in \mathcal{P}(\mathbb{R}^m \times \mathbb{R}^n)$, if and only if for any $(a, b) \in$ $\mathbb{R}^m \times \mathbb{R}^n$ we have

(3.6)
$$
(\Gamma_h)_*\mu_X((-\infty, a) \times (-\infty, b)) = F_\mu(a, b).
$$

Equivalently,

(3.7)
$$
\int_{-\infty}^{b} \int_{-\infty}^{a} \bar{h} dF_{\mu\chi} dy = F_{\mu}(a, b).
$$

This is Equation (1.21) in [\[Vapnik98,](#page-50-0) p. 37]. Assuming further that F_{μ} belongs to a metric space (E, d) , we obtain the following $P(\mathbb{R}^m \times \mathbb{R}^n, dxdy)$. correct loss function, cf. [\(3.4\)](#page-19-1),

$$
R: \mathbf{Probm}(\mathcal{X}, \mathcal{Y}) \times \left(\mathcal{P}(\mathbb{R}^m \times \mathbb{R}^n, dxdy) \cup \mathcal{P}_{emp}(\mathbb{R}^m \times \mathbb{R}^n) \right) \to \mathbb{R}_{\geq 0},
$$

(3.8)
$$
R(h, \mu) = d(F_{(\Gamma_h)_*\mu_\mathcal{X}}, F_\mu).
$$

(5) Let $d : \mathcal{P}(\mathcal{X} \times \mathcal{Y}) \times \mathcal{P}(\mathcal{X}, \mathcal{Y}) \to \mathbb{R}_{\geq 0} \cup \{\infty\}$ be an arbitrary divergence on $\mathcal{P}(\mathcal{X} \times \mathcal{Y})$. For instance, d is the Kullback-Leibler divergence, and if X and $\mathcal Y$ are separable, metrizable topological spaces then d can be any metric, which generates the weak*-topology τ_w on $\mathcal{P}(\mathcal{X} \times \mathcal{Y})$. Then

(3.9)
$$
R^d : \text{Problem}(\mathcal{X}, \mathcal{Y}) \times \mathcal{P}(\mathcal{X} \times \mathcal{Y}) \to \mathbb{R}_{\geq 0}, R(h, \mu) = d((\Gamma_h)_* \mu_{\mathcal{X}}, \mu)
$$

is a correct loss function. This is a general form of correct loss functions defined in Equations [3.1,](#page-19-2) [3.2,](#page-19-3) [3.4,](#page-19-1) [\(3.8\)](#page-20-1).

3.2. Inner and outer measure: preliminaries. In this subsection we collect necessary properties of inner and outer measure which we shall need in this article.

• Given a (nonnegative) measure μ on \mathcal{X} , we denote by μ^* the outer measure defined by μ and by μ_* the inner measure defined by μ , i.e. for any $S \subset \mathcal{X}$ we have [\[Bogachev07,](#page-48-4) p. 16, 56, 57, vol. 1]:

$$
\mu^*(S) = \inf \{ \mu(A) : S \subset A, A \in \Sigma_{\mathcal{X}} \}
$$

$$
\mu_*(S) = \sup \{ \mu(A) : S \supset A, A \in \Sigma_{\mathcal{X}} \}.
$$

Then we have [\[Bogachev07,](#page-48-4) p. 23, vol. 1]:

(3.10)
$$
\mu^*(S) + \mu_*(\mathcal{X} \setminus S) = \mu(\mathcal{X}).
$$

• Monotonicity of inner and outer measure [\[Bogachev07,](#page-48-4) p. 17, p. 70, vol. 1]:

$$
\mu_*(S_1) \le \mu_*(S_2)
$$
 and $\mu^*(S_1) \le \mu^*(S_2)$ if $S_1 \subset S_2$.

• Countable subadditivity of outer measure [\[Bogachev07,](#page-48-4) (1.5.1), p. 17, vol. 1]:

$$
\mu^*(\cup_{n=1}^{\infty} X_n) \le \sum_{n=1}^{\infty} \mu^*(S_n).
$$

Proposition 3.5 (Continuity from below of outer measure). [\[Bogachev07,](#page-48-4) Proposition 1.5.12, p. 23, vol. 1. Let μ be a (nonnegative) measure on X. Suppose that the set S_n are such that $S_n \subset S_{n+1}$ for all $n \in \mathbb{N}$. Then one has

(3.11)
$$
\mu^* \left(\bigcup_{n=1}^{\infty} S_n \right) = \lim_{n \to \infty} \mu^*(S_n).
$$

• For every decreasing sequence $S_1 \supset S_2 \ldots \supset S_n$ such that $\mu_*(S_1) < \infty$ we have [\[Bogachev07,](#page-48-4) p. 70, vol. 1]

(3.12)
$$
\mu_*\left(\bigcap_{n=1}^{\infty} S_n\right) = \lim_{n \to \infty} \mu_*(S_n).
$$

Proposition 3.6. [\[Bogachev07,](#page-48-4) Proposition 1.5.11, p. 22] If $A \in \Sigma_{\mathcal{X}}$ then

(3.13)
$$
\mu^*(S \cap A) + \mu^*(S \setminus A) = \mu^*(S) \text{ for all } S \subset \mathcal{X}.
$$

Corollary 3.7. For any $S \subset \mathcal{X}$ and $A \in \Sigma_{\mathcal{X}}$ we have

$$
(3.14) \qquad \qquad \mu_*(S \cap A) \ge \mu_*(S) - \mu(\mathcal{X} \setminus A).
$$

Proof. Using (3.10) , the validity of inequality (3.14) is equivalent to validity of the following

$$
\mu^*(\mathcal{X} \setminus (S \cap A)) \leq \mu^*(\mathcal{X} \setminus S) + \mu(\mathcal{X} \setminus A)
$$

$$
\iff \mu^*\big((\mathcal{X} \setminus S) \cup (\mathcal{X} \setminus A)\big) \leq \mu^*(\mathcal{X} \setminus S) + \mu(\mathcal{X} \setminus A),
$$

which holds because of the subadditivity of outer measure.

3.3. Consistency of a learning algorithm. The concept of a generative model of supervised learning is a particular case of the concept of a statistical learning model defined below.

Definition 3.8. A *statistical learning model* consists of a quadruple $(\mathcal{Z}, \mathcal{H}, R, \mathcal{P}_{\mathcal{Z}})$ where $\mathcal Z$ is a measurable space, $\mathcal H$ is a decision space containing all possible decisions we have to find based on a sequence of observables $(z_1, \dots, z_n) \in$ \mathcal{Z}^n , $\mathcal{P}_{\mathcal{Z}} \subset \mathcal{P}(\mathcal{Z})$ is a statistical model that contains all possible probability measures μ on $\mathcal Z$ that govern the distribution of an i.i.d. sample of data $z_1, \ldots, z_n \in \mathcal{Z}, R: \mathcal{H} \times (\mathcal{P}_{\mathcal{Z}} \cup \mathcal{P}_{emp}(\mathcal{Z})) \to \mathbb{R} \cup \{+\infty\}$ is a loss function such that $\inf_{h \in \mathcal{H}} R(h,\mu) \neq \pm \infty$ for any $\mu \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y}} \cup \mathcal{P}_{emp}(\mathcal{X} \times \mathcal{Y})$. We say that R is generated by an instantaneous loss function $L : \mathcal{Z} \times \mathcal{H} \to \mathbb{R}$ if $R(h,\mu) = R_{\mu}^{L}(h) := \mathbb{E}_{\mu} L(z,h)$. In this case we shall write R^{L} instead of R. A learning algorithm is a map

$$
A:\bigcup_{n=1}^{\infty}\mathcal{Z}^n\to\mathcal{H}.
$$

Given a statistical learning model $(\mathcal{Z}, \mathcal{H}, R, \mathcal{P}_\mathcal{Z})$ and $\mu \in \mathcal{P}_\mathcal{Z}$, we set (3.15) $R_{\mu, H} := \inf_{h \in \mathcal{H}} R_{\mu}(h).$

For $h \in \mathcal{H}$ we denote its *estimation error* as follows:

(3.16)
$$
\mathcal{E}_{\mathcal{H},R,\mu}(h) := R_{\mu}(h) - R_{\mu,\mathcal{H}}.
$$

If $R = R^L$ we shall write $\mathcal{E}_{\mathcal{H}, L, \mu}$ instead of $\mathcal{E}_{\mathcal{H}, R^L, \mu}$.

Definition 3.9. A statistical learning model $(\mathcal{Z}, \mathcal{H}, R, \mathcal{P}_\mathcal{Z})$ will be said to have a *generalization ability* or it will be called *learnable*, if there exists a uniformly consistent learning algorithm

$$
A:\bigcup_{n=1}^{\infty}\mathcal{Z}^n\to\mathcal{H},
$$

i.e. for any $(\varepsilon, \delta) \in (0, 1)^2$ there exists a number $m_A(\varepsilon, \delta)$ such that for any $m \geq m_A(\varepsilon, \delta)$ and any $\mu \in \mathcal{P}_\mathcal{Z}$ we have

$$
(3.17) \qquad (\mu^n)_*\{S \in \mathcal{Z}^n, \mathcal{E}_{\mathcal{H},R,\mu}(A(S)) \le \varepsilon\} \ge 1 - \delta.
$$

In this case A will be called uniformly consistent.

Remark 3.10. The current definition of a uniform consistency, also called a generalizability, of a learning algorithm A in literature is almost identical to our definition but the inner measure $(\mu^n)_*$ is replaced by μ^n . Equivalently, in Definition [3.9,](#page-22-0) we relax the convergence in probability in the classical requirement by the convergence in outer probability for the sequence of functions $\mathcal{E}_{\mathcal{H},R,\mu} \circ A$. Note that convergence is outer probability has been used in empirical processes [\[Dudley2014,](#page-48-11) §. 3.3], [\[VW1996,](#page-50-4) §1.9]. The classical requirement poses the following condition on a learning algorithm A: for any $\mu \in \mathcal{P}_{\mathcal{Z}}$ and any $n \in \mathbb{N}^+$ the function $\mathcal{E}_{\mathcal{H},R,\mu} \circ A : \mathcal{Z}^n \to \mathbb{R}$ is μ^n -measurable. Equivalently, for any $\mu \in \mathcal{P}_\mathcal{Z}$ and any $n \in \mathbb{N}^+$ the function $R_\mu \circ A : \mathcal{Z}^n \to \mathbb{R}$ is μ^n -measurable. Many algorithms in statistics and statistic learning theory may not be measurable, see e.g. [\[BP1973\]](#page-48-12).

Given a sequence of data $S \in \mathcal{Z}^n$ we define the empirical risk

$$
R_S: \mathcal{H} \to \mathbb{R}, h \mapsto R_{\mu_S}(h).
$$

Definition 3.11. Given a statistical model model $(\mathcal{Z}, \mathcal{H}, R, \mathcal{P}_{\mathcal{Z}})$ and a sequence $C = (c_1 \geq \ldots \geq c_n \geq \ldots : c_i \geq 0)$, a learning algorithm

$$
A:\bigcup_{n\in\mathbb{N}}\mathcal{Z}^n\to\mathcal{H}
$$

will be called a *C*-empirical risk minimizing algorithm, abbreviated as *C*-ERM algorithm, if for any $n \in \mathbb{N}$ and any $S \in \mathcal{Z}^n$, we have

$$
\widehat{R}_S(A(S)) - \inf_{h \in \mathcal{H}} R_S(h) \le c_n.
$$

If $c_i = 0$ for all i we write $C = 0$,

Clearly, a 0-ERM algorithm is an ERM algorithm.

24 H. V. LEˆ

Remark 3.12. The concept of a C-ERM algorithm is motivated by the fact that, given a sequence $C = (c_1 \geq c_2 \geq \ldots, : c_i > 0)$, a C-ERM algorithm always exists and an ERM-algorithm (a 0-ERM algorithm) may not exist. Furthermore, with fine tuning, a gradient flow could yield a C-ERM algorithm. Examples of C-ERM are solutions of regularized ERM with parameter c_n for $S \in \mathbb{Z}^n$, i.e.

$$
A(S) \in \arg\min_{h \in \mathcal{H}} (\hat{R}_S(h) + c_n W(h))
$$

if a solution of this equation exists and if we know that $W(h) \in [0, C]$ for all $h \in \mathcal{H}$ where $C < \infty$.

Theorem 3.13. Let $(\mathcal{Z}, \mathcal{H}, R, \mathcal{P}_{\mathcal{Z}})$ be a statistical learning model such that for any n the function

(3.18)
$$
\hat{R}(\cdot,h): \mathcal{Z}^n \to \mathbb{R}, S \mapsto \hat{R}_S(h),
$$

is measurable for any $h \in \mathcal{H}$. Assume that there exists a function $m_{\mathcal{H},R,\mathcal{P}_\mathcal{Z}}$: $(0,1)^2 \to \mathbb{R}_+$ such that for any $(\varepsilon,\delta) \in (0,1)^2$, any $n \geq m_{\mathcal{H},R,\mathcal{P}_\mathcal{Z}}(\varepsilon,\delta)$ and any $\mu \in \mathcal{P}_{\mathcal{Z}}$ we have

(3.19)
$$
(\mu^n)_* \{ S \in \mathcal{Z}^n : \sup_{h \in \mathcal{H}} |\hat{R}_S(h) - R_\mu(h)| \le \varepsilon \} \ge 1 - \delta.
$$

Given any sequence $C = (c_1, \ldots c_m \ldots c_i \geq 0)$, and a C-ERM algorithm $A: \bigcup_{n\in\mathbb{N}^+} \mathcal{Z}^n \to \mathcal{H}$, for any $m \geq m_{\mathcal{H},R,\mathcal{P}_\mathcal{Z}}(\varepsilon,\delta)$ any $\mu \in \mathcal{P}_\mathcal{Z}$ we have

$$
(3.20) \qquad (\mu^m)_* \{ S \in \mathcal{Z}^m : \mathcal{E}_{\mathcal{H},L,\mu}(A(S)) \le 2\varepsilon + c_m \} \ge 1 - 2\delta.
$$

Consequently, A is a uniformly consistent algorithm, if $\lim_{n\to\infty} c_n = 0$.

Proof. Assume the condition [\(3.19\)](#page-23-2) of Theorem [3.13.](#page-23-0) Let $\mu \in \mathcal{P}_\mathcal{Z}$ and $(\varepsilon, \delta) \in (0, 1)$. Then for any $m \geq m_{\mathcal{H}, R, \mathcal{P}_{\mathcal{Z}}}(\varepsilon, \delta)$ we have

$$
(3.21) \qquad (\mu^m)_* \{ S \in \mathcal{Z}^m : R_\mu(A(S)) \le \hat{R}_S(A(S)) + \varepsilon \} \ge 1 - \delta.
$$

Given $\theta > 0$ let $h_{\theta} \in \mathcal{H}$ be such that

$$
(3.22) \t\t R_{\mu}(h_{\theta}) - R_{\mu,\mathcal{H}} \leq \theta.
$$

By [\(3.19\)](#page-23-2), for any $m \geq m_{\mathcal{H},R,\mathcal{P}_{\mathcal{Z}}}(\varepsilon,\delta)$, taking into account the condition [\(3.18\)](#page-23-3), we have

(3.23)
$$
\mu^m\{S \in \mathcal{Z}^m : \hat{R}_S(h_\theta) \le R_\mu(h_\theta) + \varepsilon\} \ge 1 - \delta.
$$

Since A is a C -ERM, we have

(3.24)
$$
\forall S \in \mathcal{Z}^m : \hat{R}_S(A(S)) \leq \hat{R}_S(h_\theta) + \theta + c_m.
$$

Taking into account $(3.21), (3.23), (3.24),$ $(3.21), (3.23), (3.24),$ $(3.21), (3.23), (3.24),$ $(3.21), (3.23), (3.24),$ $(3.21), (3.23), (3.24),$ we obtain

$$
(\mu^m)_* \{ S \in \mathcal{Z}^m : R_{\mu}(A(S)) \le \hat{R}_S(A(S)) + \varepsilon \le \hat{R}_S(h_{\theta}) + \varepsilon + \theta + c_m \le
$$

\n
$$
R_{\mu}(h_{\theta}) + 2\varepsilon + \theta + c_m \} \ge
$$

\n(by (3.14))
\n
$$
1 - \delta - \mu^m (\{ S \in \mathcal{Z}^m : \hat{R}_S(h_{\theta}) > R_{\mu}(h_{\theta}) + \varepsilon \})
$$

\n(3.25)
\n
$$
\ge 1 - 2\delta.
$$

Letting θ go to zero and taking into account [\(3.12\)](#page-21-3), we obtain [\(3.20\)](#page-23-7) from (3.25) immediately.

Definition 3.14. The function $m_{\mathcal{H},R,\mathcal{P}_\mathcal{Z}}:(0,1)^2 \to \mathbb{R}$ defined by the requirement that $m_{\mathcal{H},R,\mathcal{P}_\mathcal{Z}}(\varepsilon,\delta)$ is the least number for which [\(3.19\)](#page-23-2) holds for all $\mu \in \mathcal{P}(\mathcal{Z})$ is called the sample complexity of a statistical learning model $(\mathcal{Z}, \mathcal{H}, R, \mathcal{P}_{\mathcal{Z}})$. If $R = R^L$ we shall write $m_{\mathcal{H}, L, \mathcal{P}_{\mathcal{Z}}}$ instead of $m_{\mathcal{H}, R, \mathcal{P}_{\mathcal{Z}}}$. If $\mathcal{P}_{\mathcal{Z}} = \mathcal{P}(\mathcal{Z})$ then we shall use the shorthand notation $m_{\mathcal{H},R}$ for $m_{\mathcal{H},R,\mathcal{P}(\mathcal{Z})}$.

The following Lemma provides a sufficient condition for the validity of the assumption [\(3.18\)](#page-23-3) in Theorem [3.13.](#page-23-0)

Lemma 3.15. Assume that for any $h \in \mathcal{H}$ the function $L(\cdot, h) : \mathcal{Z} \to \mathbb{R}$ is measurable. Then for any $n \geq 1$ the function

$$
R^L(\cdot,h): \mathcal{Z}^n \to \mathbb{R}, S \mapsto \hat{R}^L_S(h),
$$

is measurable.

Proof. For $k \in [1, n]$ denote by $\Pi_k : \mathcal{X}^n \to \mathcal{X}$ the projection on the k-th factor of \mathcal{X}^n . Then $\hat{R}^L(\cdot, h) : \mathcal{X}^n \to \mathbb{R}$ can be written as $\hat{R}^L =$ 1 $\frac{1}{n}\sum_{i=1}^{n} L(\cdot, h) \circ \Pi_i$. Since $L(\cdot, h)$ and Π_i are measurable, $\hat{R}^L(\cdot, h)$ is measurable.

Remark 3.16. Theorem [3.13](#page-23-0) is a generalization of Cucker-Smale's result [\[CS01,](#page-48-13) Lemma 2], though the coefficient -2 of δ in [\(3.20\)](#page-23-7) is missing in their formulas. Cucker-Smale also did not discuss the measurability of ERMalgorithms that we discussed in Remark [3.10.](#page-22-1) There are many generalizations of [\[CS01,](#page-48-13) Lemma 2] in textbooks on machine learning, where authors did not consider inner/outer measures and take uncountable intersection of measurable sets in [\(3.19\)](#page-23-2).

4. Regular conditional measures via kernel mean embedding

In this section we assume that $K: \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ is a measurable positive definite symmetric (PDS) kernel on a measurable space \mathcal{Y} . For $y \in \mathcal{Y}$ let K_y be the function on $\mathcal Y$ defined by

$$
K_y(y') = K(y, y') \text{ for } y' \in \mathcal{Y}.
$$

We denote by $\mathcal{H}(K)$ the associated RKHS [\[Aronszajn50\]](#page-48-14), see also [\[BT04\]](#page-48-15), [\[SC08\]](#page-49-14), i.e.

$$
\mathcal{H}(K) = \overline{\operatorname{span}}\{K_y, y \in \mathcal{Y}\},\
$$

where the closure is taken with respect to the $\mathcal{H}(K)$ -norm defined by

$$
\langle K_y, K_{y'} \rangle_{\mathcal{H}(K)} = K(y, y').
$$

Then for any $f \in \mathcal{H}(K)$ we have

(4.1)
$$
f(y) = \langle f, K_y \rangle_{\mathcal{H}(K)}.
$$

First, we summarize known results concerning kernel mean embeddings $\mathfrak{M}_K : \mathcal{S}(\mathcal{Y}) \to \mathcal{H}(K)$ and their direct consequences, which we shall need 26 H. V. LÊ

in our paper. Using results concerning probabilistic morphisms obtained in section [2,](#page-4-0) we study measurability and continuity of kernel mean embeddings. Then we give a characterization of regular conditional probability measure as a minimizer of mean square error (Theorem [4.6\)](#page-29-0). We show that the constructed mean square error is a correct loss function (Corollary [4.7\)](#page-30-1)). Finally, we present examples of correct loss functions obtained using this characterization, among them there are the 0-1 loss function and the mean square error (Examples [4.8\)](#page-30-0).

4.1. Kernel mean embeddings: preliminaries. By the Bochner theo-rem [\[Bochner33\]](#page-48-16), see also [\[Yosida95,](#page-50-5) Theorem 1, p. 133], $\int_{\mathcal{Y}} \sqrt{K(y,y)} \, d\mu(y) <$ ∞ for $\mu \in \mathcal{P}(\mathcal{Y})$ if and only if the kernel mean embedding $\mathfrak{M}_K(\mu)$ of μ via the Bochner integral is well-defined [\[BT04\]](#page-48-15), where

(4.2)
$$
\mathfrak{M}_K(\mu) = \int_{\mathcal{Y}} K_y d\mu(y) \in \mathcal{H}(K).
$$

By [\(4.1\)](#page-24-3), if $\mathfrak{M}_K(\mu)$ is well-defined, for any $f \in \mathcal{H}(K)$ we have

(4.3)
$$
\int_{\mathcal{X}} f(x) d\mu(x) = \langle \mathfrak{M}_K(\mu), f \rangle_{\mathcal{H}(K)}.
$$

• For a Banach space V, τ_s denotes the strong topology on V and the induced topology on its subsets, and τ_W denotes the weak topology on V and the induced topology on its subsets.

• For a locally compact Hausdorff topological space $\mathcal Y$ we denote by $C_0(\mathcal Y)$ the set of all continuous functions on which vanish at infinity.

Proposition 4.1. [\[Sriperumbudur16,](#page-49-15) Theorem 3.2] Assume that \mathcal{Y} is a Polish space that is locally compact Hausdorff. Assume that there exists a continuous bounded kernel $K : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ such that the following conditions hold.

(1) $K_y \in C_0(\mathcal{Y})$ for all $y \in \mathcal{Y}$.

(2) For any $\mu \in \mathcal{S}(\mathcal{Y})$ we have

(4.4)
$$
\int_{\mathcal{Y}} \int_{\mathcal{Y}} K(y, y') d\mu(y) d\mu(y') > 0 \text{ for all } \mu \in \mathcal{S}(\mathcal{Y}) \setminus \{0\}.
$$

In other words, $\mathfrak{M}_K : \mathcal{S}(\mathcal{Y}) \to \mathcal{H}(K)$ is injective. (3) K satisfies the following property (P)

 $\forall y \in \mathcal{Y}, \forall \varepsilon > 0, \exists \text{ open } U_{y,\varepsilon} \subset \mathcal{Y} \text{ such that } ||K_y - K_{y'}||_{\mathcal{H}(K)} < \varepsilon, \forall y' \in U_{y,\varepsilon}.$

Then the induced topology $\mathfrak{M}_{K}^{*}(\tau_{s})$ is the weak* topology τ_{w} on $\mathcal{P}(\mathcal{Y})$. In particular, the kernel mean embedding $\mathfrak{M}_K : (\mathcal{P}(\mathcal{Y}), \tau_w) \to (\mathcal{H}(\mathcal{Y}), \tau_s), \mu \mapsto$ $\mathfrak{M}_K(\mu)$, is continuous.

Proposition 4.2. [\[LMST15,](#page-49-16) Theorem 1] Let $K : \mathcal{Y} \times \mathcal{Y} \rightarrow \mathbb{R}$ be a measurable kernel such that $\mathfrak{M}_K : \mathcal{S}(\mathcal{Y}) \to \mathcal{H}(K)$ is well-defined. Assume that $||f||_{\infty} \leq 1$

for all $f \in \mathcal{H}(K)$ with $||f||_{\mathcal{H}(K)} \leq 1$. Then for any $\varepsilon \in (0,1)$ we have

$$
\mu\Big\{S_n \in \mathcal{Y}^n : \|\mathfrak{M}_K(\mu_{S_n}) - \mathfrak{M}_K(\mu)\|_{\mathcal{H}(K)} \le 2\sqrt{\frac{\int_{\mathcal{Y}} K(y, y) d\mu(y)}{n}}
$$

(4.5)
$$
+\sqrt{\frac{2\log\frac{1}{\varepsilon}}{n}} \Big\} \ge 1 - \varepsilon.
$$

• Examples of PDS kernels, which satisfy the conditions of Propositions [4.1](#page-25-1) and [4.2,](#page-25-2) are Gaussian kernels $K : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}, (x, y) \mapsto \exp(-\sigma ||x - y||)$ $y\|_2^2$, where $\sigma > 0$, Laplacian kernels $(x, y) \mapsto -\sigma \|x - y\|_1$, where $\sigma > 0$, and Matérn kernel

$$
K(x,y) = \frac{c^{2r-d}}{\Gamma(r-d/2)2^{r-1-d/2}} \left(\frac{\|x-y\|_2}{c}\right)^{r-d/2} B_{-d/2-r}(c\|x-y\|_2),
$$

where $r > d/2, c > 0$, B_a is the third modified Bessel function of the third kind of order a, and Γ is the Gamma function, see e.g. [\[MFSS17,](#page-49-17) §2.1] for a more complete list of examples.

• Assume that $\mathcal{X} \subset \mathbb{R}^n$ with the induced metric structure is a separable metric space. Since $i : \mathcal{X} \to \mathbb{R}^n$ is a continuous mapping, by Lemma [2.12,](#page-11-0) the push-forward map $i_* : \mathcal{S}(\mathcal{X}) \to \mathcal{S}(\mathbb{R}^n)$ is (τ_w, τ_w) -continuous and hence measurable mapping. Clearly i_* is a linear injective embedding.

Lemma 4.3. Assume that \mathcal{Y} is a Polish subspace of \mathbb{R}^n and $K : \mathbb{R}^n \times \mathbb{R}^n \to$ $\mathbb R$ is a continuous PDS kernel that satisfies the condition of Proposition [4.1.](#page-25-1) Let K be the restriction of K to $\mathcal{Y} \times \mathcal{Y}$. Then

(1) \tilde{K} is a PDS kernel that also satisfies the condition of Proposition [4.1.](#page-25-1)

(2) Denote by $i: \mathcal{Y} \to \mathbb{R}^n$ the canonical embedding. For any $\mu, \nu \in \mathcal{S}(\mathcal{Y})$ we have

(4.6)
$$
\langle \mathfrak{M}_{\tilde{K}}(\mu), \mathfrak{M}_{\tilde{K}}(\nu) \rangle_{\mathcal{H}(\tilde{K})} = \langle \mathfrak{M}_{K}(i_{*}\mu), \mathfrak{M}_{K}(i_{*}\nu) \rangle_{\mathcal{H}(K)}.
$$

Proof. (1) The first assertion of Lemma [4.3](#page-26-0) can be verified easily.

(2) Let $\mu, \nu \in \mathcal{S}(\mathcal{Y})$. We compute

$$
\langle \mathfrak{M}_{\tilde{K}}(\mu), \mathfrak{M}_{\tilde{K}}(\nu) \rangle_{\mathcal{H}(\tilde{K})} = \int_{\mathcal{Y}} \int_{\mathcal{Y}} \tilde{K}(y, y') d\mu(y) d\nu(y')
$$

=
$$
\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} K(y, y') di_*\mu(y) di_*\nu(y') = \langle \mathfrak{M}_K(i_*(\mu)), \mathfrak{M}_K(i_*(\nu)) \rangle_{\mathcal{H}(K)}.
$$

This completes the proof of Lemma [4.3.](#page-26-0)

If K is a measurable kernel on \mathbb{R}^n such that the kernel mean embedding $\mathfrak{M}_K : \mathcal{S}(\mathbb{R}^n) \to \mathcal{H}(K)$ is well-defined, for any Polish subspace \mathcal{Y} in \mathbb{R}^n we shall denote the pullback inner product on $\mathcal{S}(\mathcal{Y})$ by $\langle \cdot, \cdot \rangle_{\tilde{K}}$.

4.2. Measurability and continuity of kernel mean embeddings. Let $\mathcal{H}(K)$ be the RKHS associated to a measurable SPD kernel $K: \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$. Denote by $\mathcal{B}a(\tau_W)$ the smallest σ -algebra on $\mathcal{H}(K)$ such that any continuous linear function on $\mathcal{H}(K)$ is $\mathcal{B}a(\tau_W)$ -measurable, and by $\mathcal{B}(\mathcal{H}(K))$ the Borel σ-algebra of $\mathcal{H}(K)$. If $\mathcal H$ is separable, $\mathcal{B}a(\tau_W) = \mathcal{B}(\mathcal{H})$ [\[BT04,](#page-48-15) Theorem 88, p. 194. It is known that if $\mathcal Y$ is a separable topological space and K is a continuous PDS kernel on $\mathcal Y$ then $\mathcal H(K)$ is separable [\[BT04,](#page-48-15) Theorems 15, 17, p. 33-34] and [\[SC08,](#page-49-14) Lemma 4.33, p. 130].

Lemma 4.4. Assume that $K: \mathcal{Y} \times \mathcal{Y} \rightarrow \mathbb{R}$ is a bounded measurable PDS kernel. Write $\mathcal{H} := \mathcal{H}(K)$.

(1) Then the kernel mean embedding $\mathfrak{M}_K : (\mathcal{S}(\mathcal{Y}), \Sigma_w) \to (\mathcal{H}, \mathcal{B}a(\tau_W))$ is measurable. If H is separable, then $\mathfrak{M}_K : (\mathcal{S}(\mathcal{Y}), \Sigma_w) \to (\mathcal{H}, \mathcal{B}(\mathcal{H}))$ is measurable.

(2) Assume further that $\mathcal Y$ is a topological space and K is separately continuous, i.e. $K_y \in C(\mathcal{Y}, \mathbb{R})$ for all $y \in \mathcal{Y}$. Then $\mathfrak{M}_K : (\mathcal{S}(\mathcal{Y}), \tau_w) \to (\mathcal{H}, \tau_W)$ is continuous, where τ_W is the weak topology on \mathcal{H} .

(3) Assume further that $\mathcal Y$ is a topological space, K is continuous. Then the norm $\| \cdot \|_{\tilde{K}} : (\mathcal{S}(\mathcal{Y}), \tau_w) \to \mathbb{R}, \mu \mapsto \| \mu \|_{\tilde{K}}$ is continuous.

Proof. (1) To prove that $\mathfrak{M}_K : (\mathcal{S}(\mathcal{Y}), \Sigma_w) \to (\mathcal{H}, \mathcal{B}a(\tau_W))$ is measurable, taking into account Lemma [2.1,](#page-6-2) it suffices to show that for any $f \in \mathcal{H}(K)$ the composition $\langle f, \mathfrak{M}_K \rangle : (\mathcal{S}(\mathcal{Y}), \Sigma_w) \to \mathbb{R}, \mu \mapsto \langle \mathfrak{M}_K(\mu), f \rangle_{\mathcal{H}(K)}$, is a bounded measurable map. We have

$$
\langle f, \mathfrak{M}_K(\mu) \rangle_{\mathcal{H}(K)} = \int_{\mathcal{Y}} \langle f, K_y \rangle_{\mathcal{H}(K)} d\mu(y) = \int_{\mathcal{Y}} f(y) d\mu(y).
$$

We always can assume that there exists a sequence of functions K_{y_1}, \ldots, K_{y_n} converging point-wise to f by [\[BT04,](#page-48-15) Corollary 1, p. 10]. Since K is bounded, f is bounded. Since K_{y_n} is measurable, f is measurable. Since K is bounded, f is bounded. Hence

$$
\langle f,\mathfrak{M}_K \rangle(\mu) = I_f(\mu)
$$

where $f \in \mathcal{F}_b(\mathcal{Y})$. This completes the first assertion of Lemma [4.4](#page-27-1) (1).

The second assertion of Lemma [4.4](#page-27-1) (1) follows immediately by [\[BT04,](#page-48-15) Theorem 88, p. 194].

(2) Assume further that $\mathcal Y$ is a topological space and K is separately continuous. Then by [\[SC08,](#page-49-14) Lemma 4.28, p. 128], any element in $\mathcal{H} = \mathcal{H}(K)$ is bounded and continuous. Hence for any $f \in \mathcal{H}$ we have

$$
\langle f, \mathfrak{M}_K \rangle : (\mathcal{S}(\mathcal{Y}), \tau_w) \to \mathbb{R}, \, \mu \mapsto \int_{\mathcal{Y}} \langle f, K_y \rangle d\mu(y) = \int_{\mathcal{Y}} f(y) d\mu(y)
$$

i.e. $\langle f, \mathfrak{M}_K \rangle(\mu) = I_f(\mu)$. Taking into account the boundedness and continuity of f , this completes the second assertion of Lemma [4.4.](#page-27-1)

(3) Note that the map : $(\mathcal{S}(\mathcal{Y}), \tau_w) \to \mathbb{R}, \mu \mapsto ||\mu||_{\tilde{K}}$, is the composition of the map diag : $(S(\mathcal{Y}), \tau_w) \to (S(\mathcal{Y} \times \mathcal{Y}), \tau_w), \mu \mapsto \mu^2$, and the evaluation map $I_K: (\mathcal{S}(\mathcal{Y} \times \mathcal{Y}), \tau_w) \to \mathbb{R}, \nu \mapsto \int_{\mathcal{Y} \times \mathcal{Y}} K d\nu$. By Lemma [2.14](#page-12-0) (2), the

map diag is continuous. Since K is bounded and continuous, the map I_K is continuous. Hence $\|\cdot\|_{\tilde{K}}$ is a continuous map. This completes the proof of Lemma 4.4. Lemma [4.4.](#page-27-1)

4.3. Instantaneous correct loss functions via kernel mean embeddings.

Lemma 4.5. Assume that $\mathcal Y$ is a separable metrizable topological space. Let $K : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ be a bounded SPD continuous kernel and $h \in \text{Meas}(\mathcal{X}, (\mathcal{S}(\mathcal{Y}), \Sigma_w)).$ Write $\mathcal{H} = \mathcal{H}(K)$.

(1) Then the map \mathfrak{M}_K is well-defined on $\mathcal{S}(\mathcal{Y})$ and the function

(4.7)
$$
L_h^K: (\mathcal{X} \times \mathcal{Y}) \to \mathbb{R}_{\geq 0} \colon (x, y) \mapsto ||\mathfrak{M}_K(h(x)) - K_y||^2_{\mathcal{H}},
$$

is measurable.

(2) If
$$
h \in \text{Meas}(\mathcal{X}, (\mathcal{P}(\mathcal{Y}), \Sigma_w))
$$
 then the function L_h^K is bounded.

Proof of Lemma [4.5.](#page-28-1) (1) The first assertion of Lemma [4.5](#page-28-1) (1) is a consequence of the Bochner theorem.

To prove the second assertion of Lemma [4.5](#page-28-1) (1), we write

$$
(4.8)\ \|\mathfrak{M}_{K}(h(x)) - K_{y}\|_{\mathcal{H}}^{2} = \|\mathfrak{M}_{K}(h(x))\|_{\mathcal{H}}^{2} + \|K_{y}\|_{\mathcal{H}}^{2} - 2\langle\mathfrak{M}_{K}(h(x)), K_{y}\rangle_{\mathcal{H}}.
$$

We shall show that the first summand in the RHS of (4.8) , $\rho_1 : \mathcal{X} \times \mathcal{Y} \rightarrow$ $\mathbb{R}, (x, y) \mapsto \|\mathfrak{M}_K(h(x))\|_{\mathcal{H}}^2$ is a measurable bounded function. We write ρ_1 as the composition of the following maps

$$
(\mathcal{X}\times\mathcal{Y})\stackrel{\Pi_{\mathcal{X}}}{\rightarrow}\mathcal{X}\stackrel{h}{\rightarrow}(\mathcal{S}(\mathcal{Y}),\Sigma_w)\stackrel{\text{diag}}{\rightarrow}(\mathcal{S}(\mathcal{Y}\times\mathcal{Y}),\Sigma_w)\stackrel{\widehat{M}_K}{\rightarrow}\mathbb{R},
$$

where

$$
\widehat{M}_K : (\mathcal{S}(\mathcal{Y} \times \mathcal{Y}), \Sigma_w) \to \mathbb{R}, \nu \mapsto \int_{\mathcal{Y}} \int_{\mathcal{Y}} K(y, y') d\nu(y, y'),
$$

and diag is defined in Lemma [2.14](#page-12-0) (2). Since $K: \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ is a bounded and continuous function, and $\mathcal{Y} \times \mathcal{Y}$ is separable and metrizable, the function \tilde{M}_K continuous in the weak-topology τ_w on $\mathcal{S}(\mathcal{Y}\times\mathcal{Y})$, and hence measurable with respect to the σ -algebra Σ_w on $\mathcal{S}(\mathcal{Y} \times \mathcal{Y})$. Since $\Pi_{\mathcal{X}}$ and h are measurable maps, taking into account Lemma [2.14,](#page-12-0) we conclude that ρ_1 is a measurable function.

Next, we observe that the second summand in the RHS of [\(4.8\)](#page-28-2),

$$
\rho_2: \mathcal{X} \times \mathcal{Y}, (x, y) \mapsto ||K_y||_{\mathcal{H}}^2 = K(y, y),
$$

is a measurable and bounded function, since the function $K: \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ is measurable and bounded, and the mappings $\Pi_{\mathcal{X}}$ and $\Delta_{\mathcal{Y}} : \mathcal{Y} \to \mathcal{Y} \times \mathcal{Y}, y \mapsto$ (y, y) , are measurable.

Let $p: \mathcal{H} \times \mathcal{H} \to \mathbb{R}, (h, h') \mapsto \langle h, h' \rangle_{\mathcal{H}}$ be the pairing map. Now we shall prove that the function

$$
\rho_3: \mathcal{X} \times \mathcal{Y} \stackrel{(h,\mathrm{Id}_{\mathcal{Y}})}{\rightarrow} \mathcal{S}(\mathcal{Y}) \times \mathcal{Y} \stackrel{(\mathfrak{M}_{K},\hat{K})}{\rightarrow} (\mathcal{H} \times \mathcal{H}, \mathcal{B}(\mathcal{H}) \otimes \mathcal{B}(\mathcal{H})) \stackrel{p}{\rightarrow} \mathbb{R},
$$

is measurable.

By Lemma [4.4,](#page-27-1) the map $\mathfrak{M}_K : (\mathcal{S}(\mathcal{Y}), \Sigma_w) \to (\mathcal{H}, \mathcal{B}a(\tau_W))$ is measurable. Since Y is a separable topological space and K is a continuous PDS kernel, H is separable, and hence $\mathcal{B}a(\tau_W) = \mathcal{B}(\mathcal{H})$ by [\[BT04,](#page-48-15) Theorem 88, p. 194. Hence $\mathfrak{M}_K : (\mathcal{S}(\mathcal{Y}), \Sigma_w) \to (\mathcal{H}, \mathcal{B}(\mathcal{H}))$ is measurable. Since $\hat{K}: \mathcal{Y} \to (\mathcal{H}, \mathcal{B}(\mathcal{H}))$, $y \mapsto K_y$, is the composition of the measurable mappings $\delta : \mathcal{Y} \to \mathcal{S}(\mathcal{Y})$, see Lemma [2.14\(](#page-12-0)3), and \mathfrak{M}_K , the map \hat{K} is measurable. We note the pairing map $p : (\mathcal{H} \times \mathcal{H}, \mathcal{B}(\mathcal{H}) \otimes \mathcal{B}(\mathcal{H})) \to \mathbb{R}$ is measurable,

since $p : (\mathcal{H} \times \mathcal{H}, \tau_s \otimes \tau_s) \to \mathbb{R}$ is continuous.

Taking into account [\(4.8\)](#page-28-2), we complete the proof of Lemma [4.5](#page-28-1) (1).

(2) Let $C = \sup_{y \in \mathcal{Y}} K(y, y)$. If $h \in \text{Meas}(\mathcal{X}, (\mathcal{P}(\mathcal{Y}), \Sigma_w))$ then we have $L_h^K(x, y) \leq 4C$, This completes the proof of Lemma [4.5.](#page-28-1)

Theorem 4.6. Assume that \mathcal{Y} is a separable metrizable topological space, $K: \mathcal{Y} \times \mathcal{Y} \rightarrow \mathbb{R}$ is a bounded continuous PDS kernel.

(1) Then the kernel mean embedding \mathfrak{M}_K is well-defined on $\mathcal{S}(\mathcal{Y})$.

(2) For any $h \in \text{Meas}(\mathcal{X}, \mathcal{S}(\mathcal{Y}))$ the function

$$
L_h^K: (\mathcal{X} \times \mathcal{Y}) \to \mathbb{R}_{\geq 0}, (x, y) \mapsto ||h(x) - \delta_y||_K^2,
$$

is measurable.

(3) We define an instantaneous quadratic loss function as follows

$$
L^K: \mathbf{Meas}(\mathcal{X}, \mathcal{P}(\mathcal{Y})) \times \mathcal{X} \times \mathcal{Y} \to \mathbb{R}_{\geq 0}, (h, x, y) \mapsto L_h^K(x, y).
$$

For any $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ the expected loss function $R_{\mu}^{L^K}$: $\mathbf{Meas}(\mathcal{X}, \mathcal{P}(\mathcal{Y})) \rightarrow$ $\mathbb{R}.$

$$
R_{\mu}^{L^K}(h) = \int_{\mathcal{X} \times \mathcal{Y}} L_h^K(x, y) d\mu(x, y)
$$

takes a finite value at any $h \in \text{Meas}(\mathcal{X}, \mathcal{P}(\mathcal{Y})).$

(4) A regular conditional probability measure $\mu_{\mathcal{Y}|\mathcal{X}}$ for $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ is a minimizer of $R_{\mu}^{L^{K}}$.

(5) If the kernel embedding $\mathfrak{M}_K : \mathcal{S}(\mathcal{Y}) \to \mathcal{H}(K)$ is injective, then any minimizer of R^L_μ : **Meas** $(\mathcal{X}, \mathcal{P}(\mathcal{Y})) \to \mathbb{R}$, where $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$, is a regular conditional measure $\mu_{\mathcal{Y}|\mathcal{X}}$.

Proof of Theorem [4.6.](#page-29-0) (1) The first assertion of Theorem [4.6](#page-29-0) follows from Lemma [4.5.](#page-28-1)

(2) The second assertion is a reformulation of the second assertion of Lemma $4.5(1)$.

(3) The third assertion of Theorem [4.6](#page-29-0) follows from Lemma [4.5\(](#page-28-1)2).

(4) We write

 \ddotsc

(4.9)
$$
L_{h}^{K}(x, y) = ||h(x) - \mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x)) + \mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x) - \delta_{y}||_{\tilde{K}}^{2}
$$

$$
= ||h(x) - \mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x)||_{\tilde{K}}^{2} + ||\mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x) - \delta_{y}||_{\tilde{K}}^{2}
$$

$$
+ 2\langle (h(x) - \mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x), \mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x) - \delta_{y} \rangle_{\tilde{K}}.
$$

Next, using the following disintegration formula for $f \in L^1(\mathcal{X} \times \mathcal{Y}, \mu)$, where \mathcal{X}, \mathcal{Y} are measurable spaces and $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ admits a regular conditional probability $\mu_{\mathcal{Y}|\mathcal{X}}$,

(4.10)
$$
\int_{\mathcal{X}\times\mathcal{Y}} f(x,y)d\mu(x,y) = \int_{\mathcal{X}} \int_{\mathcal{Y}} f(x,y) d\mu_{\mathcal{Y}|\mathcal{X}}(y|x) d\mu_{\mathcal{X}},
$$

we obtain

$$
\int_{\mathcal{X}\times\mathcal{Y}} \langle h(x) - \mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x)), \mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x) - \delta_{y} \rangle_{\tilde{K}} d\mu(x, y)
$$
\n
$$
= \int_{\mathcal{X}} \left\langle h(x) - \mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x), \int_{\mathcal{Y}} (\mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x) - \delta_{y}) d\mu_{\mathcal{Y}|\mathcal{X}}(y|x) \right\rangle_{\tilde{K}} d(\pi_{\mathcal{X}})_{*} \mu(x) = 0,
$$

since by (4.2) we have

(4.11)
$$
\int_{\mathcal{Y}} (\mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x) - \delta_{y}) d\mu_{\mathcal{Y}|\mathcal{X}}(y|x) = 0.
$$

Hence, by [\(4.9\)](#page-29-1), we obtain

$$
(4.12) \ \ R_{\mu}^{L^K}(h) = \int_{\mathcal{X}\times\mathcal{Y}} \|h(x) - \mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x))\|_{\tilde{K}}^2 + \|\mu_{\mathcal{Y}|\mathcal{X}}(\cdot|x) - \delta_{y}\|_{\tilde{K}}^2 d\mu(x, y).
$$

Theorem [4.6](#page-29-0) (4) follows immediately from [\(4.12\)](#page-30-2).

(5) The last assertion of Theorem [4.6](#page-29-0) follows immediately from [\(4.12\)](#page-30-2). \Box

Corollary 4.7. Assume X is a measurable space and $K: \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ is a continuous bounded characteristic PDS kernel on a separable metrizable topological space Y. Then the instantaneous loss function

(4.13)
$$
L^K: \mathcal{X} \times \mathcal{Y} \times \mathbf{Meas}(\mathcal{X}, \mathcal{P}(\mathcal{Y})): (x, y, h) \mapsto L_h^K(x, y),
$$

is a correct loss function.

Examples 4.8. (1) Let $\mathcal{Y} \subset \mathbb{R}^n$ and $K : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ be defined by $K(y, y') =$ $\langle y, y' \rangle$. The restriction of L^K to $\mathcal{X} \times \mathcal{Y} \times \mathbf{Meas}(\mathcal{X}, \mathcal{Y})$ is the quadratic loss function $L_2(x, y, f) = |f(x) - y|^2$, and the minimizer of $R_{\mu}^{L_2}$ in **Meas** $(\mathcal{X}, \mathcal{Y})$ is the regression function $f(x) = E_\mu(\cdot|x)$.

(2) Let $\mathcal{Y} = \{0,1\}$. Let us embed \mathcal{Y} in \mathbb{R} by setting $\psi(i) = i \in \mathbb{R}$ for $i = 0, 1$. Then we set for $i, j \in \{0, 1\}$

$$
K(i,j) = \langle \psi(i), \psi(j) \rangle.
$$

Then $L^K : \mathcal{X} \times \{0,1\} \times \textbf{Probm}(\mathcal{X}, \mathcal{Y}) \to \mathbb{R}$ is

(4.14)
$$
L_h(x, y) = |\psi(\overline{h}(x)) - \psi(y)|^2
$$

and its restriction to $\mathcal{X} \times \{0,1\} \times \mathbf{Meas}(\mathcal{X}, \mathcal{Y})$ is the 0-1 loss function.

(3) Let us generalize Example [4.8](#page-30-0) (2) and assume that $\mathcal{Y} = \{y_1, \dots, y_n\}.$ Let $K: \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$, $K(y, y') = 1/\sqrt{2\delta_y(y')}$. Then $\mathcal{H}(\mathcal{Y}) = \mathbb{R}^n = \mathcal{S}(\mathcal{Y})$ and $K_y = \delta_y$ for $y \in \mathcal{Y}$. Then for any $h \in \mathbf{Meas}(\mathcal{X}, \mathcal{Y})$, we have

(4.15)
$$
L_h(x,y) = \frac{1}{2} ||\delta_{h(x)} - \delta_y||^2.
$$

(4) Let $\mathcal Y$ be a Polish subset in $\mathbb R^n$ an $K : \mathbb R^n \times \mathbb R^n \to \mathbb R$ be a continuous bounded PDS kernel that satisfies the conditions of Propositions [4.1](#page-25-1) and [4.2.](#page-25-2) Then the restriction of K to $\mathcal Y$ also satisfies all the conditions in Theorem [4.6.](#page-29-0)

Remark 4.9. (1) The instantaneous loss function in [\(4.14\)](#page-30-3) was proposed to the author by Frederic Protin in 2021.

(2) Park-Muandet considered the kernel embedding $\mathfrak{M}_{K}(\mu_{\mathcal{V}|\mathcal{X}})$ of a regular conditional probability measure $\mu_{\mathcal{V}|\mathcal{X}}$ in their paper [\[PM20,](#page-49-1) Definition 3.1], assuming that $\mathcal{H}(K)$ is separable. They proved that $\mathfrak{M}_K(\mu_{\mathcal{Y}|\mathcal{X}}): \mathcal{X} \to$ $\mathcal{H}(K)$ is measurable with respect to the Borel σ -algebra $\mathcal{B}(\mathcal{H})$ [\[PM20,](#page-49-1) Theorem 4.1, moreover it minimizes the loss function L^K_μ defined on $L^2(\mathcal{X}, \mathcal{H}(K), \mu_\mathcal{X})$ [\[PM20,](#page-49-1) Theorem 4.2]. Their results generalize the results due to Grünerwalder et al. in [\[GLGB12\]](#page-49-2), where the authors considered the problem of estimating the conditional expectation $\mathbb{E}_{\mu}(h(Y)|X=x)$, where h belongs to a $\mathcal{H}(K)$ valued RKHS \mathcal{H}_{Γ} , using the mean square error R_{μ}^{L} we considered in Lemma [4.5,](#page-28-1) under many strong assumptions. They also noted that such a \mathcal{H}_{Γ} belongs to the space of continuous functions from $\mathcal X$ to $\mathcal H(K)$.

(3) In [\[TSS2022\]](#page-50-3) Talwai-Shami-Simchi-Levi considered the problem of estimating conditional distribution $[\mu_{\mathcal{Y}|\mathcal{X}}]$ by representing them as a operator $C_{\mathcal{Y}|\mathcal{X}} : \mathcal{H}_{\mathcal{X}} \to \mathcal{H}_{\mathcal{Y}}$, where $\mathcal{H}_{\mathcal{X}}$ and $\mathcal{H}_{\mathcal{Y}}$ are RKHSs associated with PSD kernels on X and Y respectively.

5. A generalization of Cucker-Smale's result

In this section, we keep the notation used in the previous sections.

• For a Hilbert space H and a topological space \mathcal{X} , we denote by $C_b(\mathcal{X}, \mathcal{H})$ the space of all continuous bounded mappings from X to H endowed with the sup-norm

$$
||f||_{\infty} = \sup_{x \in \mathcal{X}} ||f(x)||_{\mathcal{H}}.
$$

• For any precompact metric space \mathcal{H} , and $s > 0$, we denote the s-covering number of H by $\mathcal{N}(\mathcal{H},s)$, i.e.

 $\mathcal{N}(\mathcal{H}, s) := \min\{l \in \mathbb{N} | \exists l \text{ balls centered in } \mathcal{H} \text{ of radius } s \text{ covering } \mathcal{H}\} < \infty.$

In Subsection [5.1,](#page-31-1) we state our theorem (Theorem [5.1\)](#page-31-2) and discuss the relation with previous results. In Subsection [5.2,](#page-33-0) we give a proof of Theorem [5.1.](#page-31-2)

5.1. Statement of the result.

Theorem 5.1. Let X be a topological space and \mathcal{Y} a Polish subset in \mathbb{R}^m . Let K be the restriction to $\mathcal Y$ of a continuous bounded SPD kernel on $\mathbb R^m$, which satisfies the conditions of Proposition [4.1.](#page-25-1) Assume that $\mathcal{H} \subset C(\mathcal{X}, \mathcal{P}(\mathcal{Y}), \tau_w)$.

(1) Then $\mathcal{H} \subset C_b(\mathcal{X}, \mathcal{S}(\mathcal{Y})_{\tilde{K}})$.

(2) Let $A^K: \bigcup_{n=1}^{\infty} (\mathcal{X} \times \mathcal{Y})^n \to \mathcal{H}, S_n \mapsto A_{S_n}^K$, be a C-ERM algorithm for the supervised learning model $(\mathcal{X}, \mathcal{Y}, \mathcal{H}, L^K, \mathcal{P}(\mathcal{X} \times \mathcal{Y}))$, where L^K is defined

in [\(4.13\)](#page-30-4), i.e.

$$
L^K(x, y, h) = ||h(x) - \delta_y||_{\tilde{K}}.
$$

If H is a pre-compact subset in $C_b(\mathcal{X}, \mathcal{S}(\mathcal{Y})_{\tilde{K}})_{\infty}$ then for any $\varepsilon > 0, \delta > 0$ there exists $m(\varepsilon, \delta)$ such that for any $m \ge m(\varepsilon, \delta)$ and any $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ we have

(5.1)

$$
(\mu^m)_*\{S_m \in (\mathcal{X} \times \mathcal{Y})^m | \mathcal{E}_{\mathcal{H}, L^K, \mu}(A_{S_m}^K) < 2\varepsilon + c_m\} \ge 1 - 2\mathcal{N}(\mathcal{H}, \frac{\varepsilon}{8C_K})2 \exp\left(-\frac{m\varepsilon^2}{4C_K^2}\right)
$$

where

$$
C_K := \sup_{y \in \mathcal{Y}} \sqrt{|K(y, y)|} < \infty.
$$

Hence a C-ERM algorithm for the supervised learning model $(\mathcal{X}, \mathcal{Y}, \mathcal{H}, L^K, \mathcal{P}(\mathcal{X} \times$ (Y)) is a uniformly consistent learning algorithm, if $\lim_{n\to\infty} c_n = 0$.

In particular, if

$$
A_{S_n}^K \in \arg\min_{h \in \mathcal{H}} (R_{\mu_{S_n}}^{L^K}(h) + c_n \|h\|_{\infty}),
$$

then A^K is a C-ERM algorithm and (5.2)

$$
(\mu^m)_*\{S_m\in (\mathcal{X}\times \mathcal{Y})^m|\,\mathcal{E}_{\mathcal{H},L^K,\mu}(A_{S_m}^K)<4\varepsilon+c_mC_K\}\geq 1-2\mathcal{N}(\mathcal{H},\frac{\varepsilon}{8C_K})2\exp(-\frac{m\varepsilon^2}{4C_K^2}),
$$

Remark 5.2. (1) Theorem [5.1\(](#page-31-2)2) is a generalization of Cucker-Smale's result [\[CS01,](#page-48-13) Theorem C], where Cucker-Smale also assumed implicitly that their ERM algorithm must satisfy certain measurability, see Remark [3.16.](#page-24-2) Cucker-Smale also assumed the boundedness of predictors h in hypothesis classes $\mathcal{H} \subset \text{Meas}(\mathcal{X}, \mathbb{R}^n)$. In our setting of conditional probability estimation, the boundedness of Markov kernels $h \in C(X, \mathcal{P}(\mathcal{Y})_{K_2})$ is automatically satisfied (Theorem [5.1](#page-31-2) (1)).

 (2) Grünerwalder et al. [\[GLGB12\]](#page-49-2), based on results due to Capponento and de Vito [\[CD07\]](#page-48-17), proved the learnability of their statistical learning model for conditional mean embeddings under a strong assumptions on statistical model $\mathcal{P}_{\mathcal{X}\times\mathcal{Y}}$, on the finiteness dimension of a RKHS $\mathcal{H}(K)$ associated to a PSD kernel K on $\mathcal Y$ and moreover the hypothesis space $\mathcal H_{\Gamma}$, which is a vector valued RKHS, must contain a minimizer $\mu_{\mathcal{Y}|\mathcal{X}}$ of the mean square error. Park-Muandet proved the universal consistency of a regularized ERM algorithm of their statistical learning model $(\mathcal{X}, \mathcal{Y}, \mathcal{H}_{\Gamma}, L^{K}, \mathcal{P}(\mathcal{X} \times \mathcal{Y}))$, where a hypothesis space \mathcal{H}_{Γ} is a vector-valued RKHS associated with a C_0 -universal kernel $l_{\mathcal{X}\mathcal{V}} : \mathcal{X} \times \mathcal{X} \to \mathcal{H}(K_{\mathcal{V}}), \mathcal{H}(K_{\mathcal{V}})$ is a separable RKHS associated with a measurable bounded SPD kernel $K_{\mathcal{Y}}$ on \mathcal{Y} and \mathcal{X} also admits a bounded measurable kernel $K_{\mathcal{X}}$ such that $\mathcal{H}(K_{\mathcal{X}})$ is a separable Hilbert space [\[PM20,](#page-49-1) Theorem 4.4]. They provided a convergence rate of their learning algorithm under the condition that \mathcal{H}_{Γ} contains a minimizer $\mu_{\mathcal{Y}|\mathcal{X}}$ of the mean square error [\[PM20,](#page-49-1) Theorem 4.5]. In [\[TSS2022\]](#page-50-3) Talwai-Shameli and Simchi-Levi improved the result by Grünerwalder et al. [\[GLGB12\]](#page-49-2) by dropping many

technical assumptions in [\[GLGB12\]](#page-49-2) and applying the theory of interpolation spaces for RKHS [\[FS20\]](#page-49-18). Note that a minimizer of the loss function in \mathcal{H}_{Γ} in the papers considered above may not correspond to a Markov kernel, i.e. to a measurable map \overline{T} : $\mathcal{X} \to \mathcal{P}(\mathcal{Y})$.

5.2. Proof of Theorem [5.1.](#page-31-2) (1) Assertion (1) of Theorem [5.1](#page-31-2) stating that $\mathcal{H} \subset C_b(\mathcal{X}, \mathcal{S}(\mathcal{Y})_{\tilde{K}})$ follows from Proposition [4.1](#page-25-1) and Lemma [4.3.](#page-26-0)

(2) To prove assertion (2) of Theorem [5.1,](#page-31-2) we shall apply Theorem [3.13,](#page-23-0) namely we shall give an upper bound for the sample complexity of the supervised learning model $(\mathcal{X}, \mathcal{Y}, \mathcal{H}, L^K, \mathcal{P}(\mathcal{X} \times \mathcal{Y}))$ in Proposition [5.6](#page-34-0) below, using the same strategy in the proof of Cucker-Smale's theorem [\[CS01,](#page-48-13) Theorem C, though Cucker-Smale considered $\mathcal{Y} = \mathbb{R}^n$, $n < \infty$ and assumed that $\mathcal{H} \subset C(\mathcal{X}, \mathbb{R}^n)$ is compact [\[CS01,](#page-48-13) Remark 14], they also did not use inner/outer measure.

First we shall prove Lemmas [5.3,](#page-33-1) [5.4,](#page-33-2) [5.5,](#page-34-1) then we shall prove Proposition [5.6](#page-34-0) and complete the proof of Theorem [5.1](#page-31-2) (2).

Lemma 5.3. Assume the condition of Theorem [5.1.](#page-31-2) Then for any $h \in \mathcal{H}$ the following claims hold.

(1) The function $L_h^K : (\mathcal{X} \times \mathcal{Y}) \to \mathbb{R}, (x, y) \to L^K(x, y, h)$ is measurable. (2) For any $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ we have

$$
(5.3) \quad \mu^m \{ S \in (\mathcal{X} \times \mathcal{Y})^m : \left| R_{\mu}^{L_K}(h) - \widehat{R_S^{L_K}}(h) \right| \le \varepsilon \} \ge 1 - 2 \exp\left(-\frac{m\varepsilon^2}{4C_K^2}\right) \}
$$

Proof. (1) The first assertion of Lemma [5.3](#page-33-1) follows from Lemma [4.5.](#page-28-1)

(2) We note that

(5.4)
$$
\|\mu\|_{\tilde{K}} \leq C_K \text{ for all } \mu \in \mathcal{P}(\mathcal{Y}).
$$

Using the first assertion of Lemma [5.3](#page-33-1) and (5.4) , we derive (5.3) from the Hoeffding inequality, stating that for any measurable mapping $\xi : \mathcal{Z} \rightarrow$ $[a, b] \subset \mathbb{R}, \mu \in \mathcal{P}(\mathcal{Z})$, we have [\[BLM13,](#page-48-18) Theorem 2.8, p. 34] (5.5)

$$
\mu^m\Big\{(z_1,\ldots,z_m)\in\mathcal{Z}^m\big|\frac{1}{m}\sum_{i=1}^m\xi(z_i)-\mathbb{E}_{\mu}(\xi)\big|\geq\varepsilon\Big\}\leq 2\exp(-\frac{2m\varepsilon^2}{(b-a)^2}),
$$

noting that [\(5.4\)](#page-33-3) implies

(5.6)
$$
0 \le L_h^K(x, y) = ||h(x) - \delta_y||_K^2 \le 4C_K^2,
$$

and then plugging $\xi := L_h^K : \mathcal{X} \times \mathcal{Y} \to [0, 4C_K^2]$ into [\(5.5\)](#page-33-5).

Lemma 5.4. For any $f, g \in \mathcal{H}$, $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ and $S_n \in (\mathcal{X} \times \mathcal{Y})^n$ we have

$$
(5.7) \qquad |(R_{\mu}^{L^K}(f) - R_{S_n}^{L^K}(f)) - (R_{\mu}^{L^K}(g) - R_{S_n}^{L^K}(g))| \leq 8C_K \|f - g\|_{\infty}.
$$

Proof. Let $f, g \in C(X, \mathcal{P}(Y))$. Using [\(5.4\)](#page-33-3), we obtain

$$
\left| \|f(x)\|_{\tilde{K}}^2 - \|g(x)\|_{\tilde{K}}^2 \right| = \left| \langle f(x) - g(x)|f(x) \rangle_{\tilde{K}} + \langle g(x) - f(x)| - g(x) \rangle_{\tilde{K}} \right|
$$

$$
\leq |||f(x) - g(x)||_{\tilde{K}}| \cdot (||f(x)||_{\tilde{K}} + ||g(x)||_{\tilde{K}}) \leq 2C_K ||f(x) - g(x)||_{\tilde{K}}.
$$

the also have

We

$$
|2\langle f(x) - g(x)|\delta_y\rangle_{\tilde{K}}| \leq 2C_K ||f(x) - g(x)||_{\tilde{K}}.
$$

Hence,

$$
|R_{\mu}^{L^K}(f) - R_{\mu}^{L^K}(g)| = \left| \int_{\mathcal{X} \times \mathcal{Y}} ||f(x)||_{\tilde{K}}^2 - ||g(x)||_{\tilde{K}}^2 - 2\langle f(x) - g(x)|\delta_y \rangle_{\tilde{K}} d\mu(x, y) \right|
$$

(5.8)
$$
\leq 4C_K ||f - g||_{\infty}.
$$

In particular, for any $S_n = (x_1, y_1, \dots, x_n, y_n) \in (\mathcal{X} \times \mathcal{Y})^n$ we have

$$
\left| R_{S_n}^{L^K}(f) - R_{S_n}^{L^K}(g) \right| = \frac{1}{n} \sum_{i=1}^n \left(\|f(x_i)\|_{\tilde{K}}^2 - \|g(x_i)\|_{\tilde{K}}^2 \right)
$$

(5.9)
$$
-2\langle f(x_i) - g(x_i)|\delta_{y_i} \rangle_{\tilde{K}} \le 4C_K \|f - g\|_{\infty}.
$$

Clearly (5.7) follows from (5.8) and (5.9) .

Lemma 5.5. cf. [\[CS01,](#page-48-13) Lemma 1] Let $\mathcal{H} = D_1 \cup ... \cup D_l$, where D_i are open balls centered in $h_i \in \mathcal{H}$, $i \in [1, l]$, and $\varepsilon > 0$. Then

$$
(\mu^m)^* \{ S \in (\mathcal{X} \times \mathcal{Y})^m : \sup_{h \in \mathcal{H}} |R_{\mu}^{L^K}(h) - \widehat{R}_{S}^{L^K}(h)| > \varepsilon \}
$$

$$
(5.10) \qquad \leq \sum_{j=1}^{N} (\mu^m)^* \{ S \in (\mathcal{X} \times \mathcal{Y})^m : \sup_{h \in D_j} |R_{\mu}^{L^K}(h) - \widehat{R}_{S}^{L^K}(h)| > \varepsilon \}
$$

Proof. Lemma [5.5](#page-34-1) follows from the equivalence

$$
\sup_{h\in\mathcal{H}}|R_{\mu}^{L^K}(h)-\widehat{R}_{S}^{L^K}(h)|\geq \varepsilon\}\Longleftrightarrow \exists j\in[1,l]:\sup_{h\in D_j}|R_{\mu}^{L^K}(h)-\widehat{R}_{S}^{L^K}(h)|\geq \varepsilon,
$$

taking into account the countable subadditivity of outer measure. \Box

Proposition 5.6. Assume the condition of Theorem [5.1.](#page-31-2) Then for any $\varepsilon > 0$, $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ and $h \in \mathcal{H}$ we have

$$
(\mu^m)_* \{ S \in (\mathcal{X} \times \mathcal{Y})^m : \sup_{h \in \mathcal{H}} |R_{\mu}^{L_K}(h) - R_S^{L_K}(h)| \le 2\varepsilon \} \ge
$$

(5.11)

$$
1 - \mathcal{N}(\mathcal{H}, \frac{\varepsilon}{8C_K^2}) 2 \exp(-\frac{m\varepsilon^2}{4C_K^2}) \}.
$$

Proof. Let $l = \mathcal{N}(\mathcal{H}, \frac{\varepsilon}{8C})$ $\frac{\varepsilon}{8C_K}$ and consider h_1, \ldots, h_l such that the collection of balls D_j centered at h_j with radius $\frac{\varepsilon}{8C_K}$ covers H. By Lemma [5.4,](#page-33-2) for any $S \in (\mathcal{X} \times \mathcal{Y})^m$ and $f \in D_j$ we have

$$
|(R_{\mu}^{L^K}(f) - \hat{R}_{S}^{L^K}(f)) - (R_{\mu}^{L^K}(f_j) - \hat{R}_{S}^{L^K}(f_j))| \leq 8C_K \frac{\varepsilon}{8C_K} = \varepsilon.
$$

It follows that

$$
\sup_{f \in D_j} |R_{\mu}^{L^K}(f) - \widehat{R}_{S}^{L^K}(f)| \ge 2\varepsilon \Longrightarrow |R_{\mu}^{L^K}(f_j) - \widehat{R}_{S}^{L^K}(f_j)| \ge \varepsilon.
$$

Hence, taking into account Lemmas [4.5](#page-28-1) and [3.15,](#page-24-1) for any $j \in [1, l]$ we obtain:

$$
(\mu^m)^* \{ S \in (\mathcal{X} \times \mathcal{Y})^m : \sup_{f \in D_j} |R_{\mu}^{L^K}(f) - \widehat{R}_{S}^{L^K}(f)| \ge 2\varepsilon \} \le
$$

$$
\mu^m \{ S \in (\mathcal{X} \times \mathcal{Y})^m : |R_{\mu}^{L^K}(f_j) - \widehat{R}_{S}^{L^K}(f_j)| \ge \varepsilon \}
$$

$$
(5.12)
$$
(by Lemma 5.3) $\le 2 \exp(-\frac{m\varepsilon^2}{4CK^2}).$

Taking into account Lemma [5.5](#page-34-1) and [\(3.10\)](#page-20-2), we obtain from [\(5.12\)](#page-35-2) Proposi-tion [5.6.](#page-34-0) \Box

Proof of Theorem [5.1\(](#page-31-2)2). Using Theorem [3.13,](#page-23-0) we obtain the second assertion of Theorem $5.1(2)$ from Proposition [5.6.](#page-34-0)

The last assertion of Theorem [5.1\(](#page-31-2)2) follows immediately. \Box

6. A variant of Vapnik-Stefanyuk's theorem and applications

6.1. A variant of Vapnik-Stefanyuk's regularization method. In [\[Vapnik98,](#page-50-0) Chapter 7] Vapnik explained methods due to Stefanyuk-Vapnik [\[VS1978\]](#page-50-6) to solve the following operator equation

$$
(6.1)\t\t\t Af = F
$$

defined by a continuous operator A which maps in a one-to one manner the elements f of a metric space E_1 into the elements of a metric space E_2 assuming that a solution $f \in E_1$ of [\(6.1\)](#page-35-3) exists and is unique.

We consider the situation when A belongs to a space A and instead of Equation [\(6.1\)](#page-35-3) we are given a sequence $\{F_{S_l} \in E_2, l \in \mathbb{N}^+\}$, a sequence ${A}_{S_l} \in \mathcal{A}, l \in \mathbb{N}^+$, where S_l belongs to a probability space (\mathcal{X}_l, μ_l) and A_{S_l}, F_{S_l} are defined by a family of maps $\mathcal{X}_l \to E_2, S_l \mapsto \mathcal{F}_{S_l}$, and $\mathcal{X}_l \to$ $\mathcal{A}, S_l \mapsto A_{S_l}.$

Let $W: E_1 \to \mathbb{R}_{\geq 0}$ be a lower semi-continuous function that satisfies the following property (W).

(W) The set $\mathcal{M}_c = W^{-1}([0, c])$ for $c \ge 0$ are all compact.

Given A_{S_l}, F_{S_l} , and $\gamma_l > 0$, let us define a regularized risk function $R_{\gamma_l}^*(\cdot, F_{S_l}, A_{S_l}) : E_1 \to \mathbb{R}$ by

(6.2)
$$
R_{\gamma_l}^*(\hat{f}, F_{S_l}, A_{S_l}) = \rho_{E_2}^2(A_{S_l}\hat{f}, F_{S_l}) + \gamma_l W(\hat{f}).
$$

We shall say that $f_{S_l} \in E_1$ is an ε_l -minimizer of $R^*_{\gamma_l}$ if

(6.3)
$$
R_{\gamma_l}^*(f_{S_l}, F_{S_l}, A_{S_l}) \le R_{\gamma_l}(\hat{f}, F_{S_l}, A_{S_l}) + \varepsilon_l \text{ for all } \hat{f} \in D.
$$

We shall also use the shorthand notation A_l for A_{S_l} , F_l for F_{S_l} , f_l for f_{S_l} , ρ_2 for ρ_{E_2} , ρ_1 for ρ_{E_1} . For any $\varepsilon_l > 0$, an ε_l -minimizer of $R^*_{\gamma_l}$ exists. We will measure the closedness of operator A and operator A_l by the distance

(6.4)
$$
||A_l - A|| = \sup_{\hat{f} \in E_1} \frac{||A_l \hat{f} - A\hat{f}||_{E_2}}{W^{1/2}(\hat{f})}.
$$

Theorem 6.1. *cf.* [\[Vapnik98,](#page-50-0) Theorem 7.3, p. 299] Let f_{S_l} be a γ_l^2 minimizer of $R_{\gamma_l}^*$ in [\(6.2\)](#page-35-4) and f the solution of [\(6.1\)](#page-35-3). For any $\varepsilon > 0$ and any constant $C_1, C_2 > 0$ there exists a value $\gamma_0 > 0$ such that for any $\gamma_l \leq \gamma_0$

$$
(\mu_l)^* \{ S_l \in \mathcal{X}_l : \rho_1(f_{S_l}, f) > \varepsilon \} \leq (\mu_l)^* \{ S_l \in \mathcal{X}_l : \rho_2(F_{S_l}, F) > C_1 \sqrt{\gamma_l} \} + (\mu_l)^* \{ S_l \in \mathcal{X}_l : ||A_{S_l} - A|| > C_2 \sqrt{\gamma_l} \}
$$

holds true.

Remark 6.2. Note that Theorem [6.1](#page-36-0) is a slight generalization of [\[Vapnik98,](#page-50-0) Theorem 7.3, p. 299], where Vapnik considered the case that f_l is a minimizer of $R_{\gamma_l}^*$. Our proof of Theorem [6.1](#page-36-0) follows the arguments in Vapnik's proof of [\[Vapnik98,](#page-50-0) Theorem 7.3], carefully estimating all " ε , δ " for outer measure, instead of measure as in Vapnik's proof, which requires also measurablity of sets involved.

Proof of Theorem [6.1.](#page-36-0) Since f_l is a γ_l^2 -minimizer of $R_{\gamma_l}^*(\cdot, F_l, A_l)$, by [\(6.3\)](#page-35-5) we have

(6.6)
$$
\gamma_l W(f_l) \leq R_{\gamma_l}^*(f_l, F_l, A_l) \leq R_{\gamma_l}^*(f, F_l, A_l) + \gamma_l^2
$$

(6.7)
$$
= \rho_2^2(A_l f, F_l) + \gamma_l W(f) + \gamma_l^2
$$

where f is the desired solution of (6.1) . From (6.6) we find

(6.8)
$$
W(f_l) \le W(f) + \frac{\rho_2^2(A_l f, F_l)}{\gamma_l} + \gamma_l.
$$

Since according to the triangle inequality, we have

(6.9)
$$
\rho_2(A_l f, F_l) \leq \rho_2(A_l f, F) + \rho_2(F, F_l)
$$

$$
\leq \|A_l - A\| W^{1/2}(f) + \rho_2(F, F_l),
$$

we obtain

(6.10)
$$
W(f_l) \leq W(f) + \frac{1}{\gamma_l} \Big(\|A_l - A\| W^{1/2}(f) + \rho_2(F, F_l) \Big)^2 + \gamma_l.
$$

Since f_l is a γ_l^2 -minimizer of $R_{\gamma_l}^*(\cdot, F_l, A_l)$, we have

(6.11)
$$
\rho_2^2(A_l f_l, F_l) \leq R_{\gamma_l}^*(f_l, F_l, A_l) + \gamma_l^2.
$$

From [\(6.11\)](#page-36-3), [\(6.6\)](#page-36-2), and [\(6.9\)](#page-36-4), we obtain

$$
(6.12) \qquad \rho_2^2(A_l f_l, F_l) \le \gamma_l W(f) + \left(\|A_l - A\| W^{1/2}(f) + \rho_2(F, F_l) \right) + \gamma_l^2.
$$

From this, using [\(6.4\)](#page-35-6) and [\(6.9\)](#page-36-4), we derive

$$
\rho_2(Af_l, F) \leq \rho_2(Af_l, A_l f_l) + \rho_2(A_l f_l, F_l) + \rho_2(F_l, F)
$$

\n
$$
\leq W^{1/2}(f_l) \|A_l - A\|
$$

\n
$$
+ \left(\gamma_l W(f) + (\|A_l - A\| W^{1/2}(f) + \rho_2(F, F_l))^2 + \gamma_l^2\right)^{1/2}
$$

\n
$$
+ \rho_2(F_l, F) = \sqrt{\gamma_l} \left(\frac{\rho_2(F_l, F)}{\sqrt{\gamma_l}} + W^{1/2}(f) \frac{\|A_l - A\|}{\sqrt{\gamma_l}}\right)
$$

\n(6.13)
$$
+ \sqrt{\gamma_l} \left(W(f) + \left(\frac{\rho_2(F_l, F)}{\sqrt{\gamma_l}} + W^{1/2}(f) \frac{\|A_l - A\|}{\sqrt{\gamma_l}}\right)^2 + \gamma_l^{3/2}\right)^{1/2}
$$

Given $C_1, C_2 > 0$ we set

(6.14)
$$
C_l := \{ S_l \in \mathcal{X}_l : \frac{\rho_2(F, F_l)}{\sqrt{\gamma_l}} \le C_1 \text{ and } \frac{\|A_l - A\|}{\sqrt{\gamma_l}} \le C_2 \}.
$$

Now assume that

$$
(6.15) \t S_l \in \mathcal{C}_l.
$$

From [\(6.10\)](#page-36-5) we have

(6.16)
$$
W(f_l) \le W(f) + (C_1 + C_2 W^{1/2}(f))^2 + \gamma_l = d + \gamma_l < \infty
$$

where

(6.17)
$$
d = W(f) + (C_1 + C_2 W^{1/2}(f))^2 > 0.
$$

From (6.13) , using (6.14) and (6.16) , we obtain

$$
\rho_2(Af_l, F) \le \sqrt{\gamma_l} (C_1 + W^{1/2}(f) C_2) \n+ \sqrt{\gamma_l} (d + \gamma_l^{3/2})^{1/2} \n\le \sqrt{\gamma_l} (\sqrt{d} + \sqrt{d + \gamma_l^{3/2}}) \n\le 2\sqrt{\gamma_l} \sqrt{d + \gamma_l^{3/2}}.
$$
\n(6.18)

Now we pose the following condition on γ_l :

(6.19) $\gamma_l \le \min\{d, d^{2/3}\}.$

Taking into account [\(6.17\)](#page-37-3) and the properties of the functional $W : D \to$ $\mathbb{R}_{\geq 0}$, inequality [\(6.16\)](#page-37-2) implies that, under the condition [\(6.19\)](#page-37-4) on γ_l , both f and f_l belong to the compactum $W^{-1}([0, 2d])$.

Lemma 6.3. [\[Vapnik98,](#page-50-0) Lemma in p. 53]. Let E_1, E_2 be metric spaces. If $A: E_1 \rightarrow E_2$ is a one-to one continuous operator defined on a compact set $M \subset E_1$, then the inverse operator A^{-1} is continuous on the set $N = A(M)$.

By Lemma [6.3,](#page-37-5) for any $\varepsilon > 0$ there exists $\delta(\varepsilon) > 0$ such that

(6.20)
$$
\rho_2(Af_l, Af) \leq \delta(\varepsilon) \Longrightarrow \rho_1(f, f_l) \leq \varepsilon.
$$

Now we set

(6.21)
$$
\gamma_0 = \min\{d, d^{2/3}, \left(\frac{\delta(\varepsilon)^2}{8d}\right)\} > 0.
$$

Equations [\(6.19\)](#page-37-4), [\(6.21\)](#page-38-2), and [\(6.18\)](#page-37-6) imply that for $\gamma_l \leq \gamma_0$ we have

$$
(6.22) \t\t \rho_1(f, f_l) \le \varepsilon.
$$

Hence, the condition [\(6.21\)](#page-38-2) for γ_0 implies that, for all $\gamma_l \leq \gamma_0$, by [\(6.14\)](#page-37-1) and [\(6.15\)](#page-37-7) and taking into account the subadditivity of outer measure, we have

$$
(\mu_l)^* \big\{ S_l : \rho_1(f_l, f) > \varepsilon \big\} \leq (\mu_l)^* \big\{ S_l : \frac{\rho_2(F_l, F)}{\sqrt{\gamma_l}} > C_1 \big\} + (\mu_l)^* \big\{ S_l : \frac{\|A_l - A\|}{\sqrt{\gamma_l}} > C_2 \big\}.
$$

Note that $\gamma_0 = \gamma(C_1, C_2, W(f), A^{-1}, \varepsilon)$, where C_1, C_2, ε are arbitrary fixed constant. This completes the proof of Theorem [6.1.](#page-36-0) \Box

Remark 6.4. In [\[Vapnik98,](#page-50-0) §7.10] Vapnik applied Stefanyuk-Vapnik's theorem [\[Vapnik98,](#page-50-0) Theorem 7.3, p. 299] for estimation of smooth conditional probability densities on a line. He also presented a method of estimation of a conditional probability $\mu_{\mathcal{V}|\mathcal{X}}$ where \mathcal{Y} is finite and $\mathcal{X} = \mathbb{R}$ [\[Vapnik98,](#page-50-0) §7.12]. In these cases Vapnik assumed that the hypothesis spaces are finite dimensional and explicit minimum of the expected risk function can be found. In [\[VI2019\]](#page-50-2) Vapnik and Izmailov consider estimation of a conditional probability $\mu_{\mathcal{Y}|\mathcal{X}}$ where \mathcal{Y} is finite and $\mathcal{X} \subset \mathbb{R}^n$, using cumulative functions. We shall show an application of Theorem [6.1](#page-36-0) below for proving the consistency of a model for estimating conditional probability where \mathcal{X}, \mathcal{Y} are Polish subsets in \mathbb{R}^m .

For metric spaces F_1, F_2 , we denote by $C_{Lip}(F_1, F_2)$ the space of all Lipschitz continuous mappings from F_1 to F_2 . By Lemma [2.14\(](#page-12-0)1), $\Gamma_h : \mathcal{X} \to$ $\mathcal{M}(\mathcal{X}\times\mathcal{Y})_{\tilde{K}_1}$ is continuous, since $h\in C(\mathcal{X},\mathcal{M}(\mathcal{Y})_{\tilde{K}_2})$. Since Γ_h is continuous, the function $x \mapsto ||\Gamma_h(x)||_{\tilde{K}_1}$ is continuous, and it is bounded, since $\mathcal X$ is compact. Now we define the distance d_M on $C(\mathcal{X}, \mathcal{M}(\mathcal{Y})_{\tilde{K}_2})$ by

(6.23)
$$
d_M(f, f') = \sup_{x \in \mathcal{X}} (\|(f - f')(x)\|_{\tilde{K}_2} + \|\Gamma_f(x) - \Gamma_{f'}(x)\|_{\tilde{K}_1}).
$$

In other words, the metric d_M is induced by the norm $\|\cdot\|_M$ defined on the space $C(\mathcal{X}, \mathcal{S}(\mathcal{Y})_{\tilde{K}_2})$ as follows:

$$
||f||_M = \sup_{x \in \mathcal{X}} (||f(x)||_{\tilde{K}_2} + ||\Gamma_f(x)||_{\tilde{K}_1}).
$$

The space $C(\mathcal{X}, \mathcal{M}(\mathcal{Y})_{\tilde{K}_2})$ endowed with the metric d_M shall be denoted by $C(\mathcal{X},\mathcal{M}(\mathcal{Y})_{\tilde{K}_2})_M.$

Theorem 6.5. Let \mathcal{X} be a compact subset in $\mathbb{R}^n \times \{0\} \subset \mathbb{R}^{n+m}$ and \mathcal{Y} a compact subset in $\{0\} \times \mathbb{R}^m$. Let $K_2 : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ be the restriction of a continuous bounded SPD kernel K on $\mathbb{R}^{m+n} \times \mathbb{R}^{m+n}$, which satisfies the conditions of Propositions [4.1](#page-25-1) and [4.2.](#page-25-2) Denote by $\mathcal{P}_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}, vol_{\mathcal{X}})$ the set of all probability measures $\mu \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ such that:

(i) sppt $\mu_{\mathcal{X}} = \mathcal{X}$, where $\mu_{\mathcal{X}} = (\Pi_{\mathcal{X}})_*\mu$; (ii) there exists a regular conditional measure $\mu_{\mathcal{Y}|\mathcal{X}} \in C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})$ for μ with respect to the projection $\Pi_{\mathcal{X}} : \mathcal{X} \times \mathcal{Y} \to \mathcal{X}$. Let $K_1: \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$ be the restriction of the kernel K to $(\mathcal{X} \times \mathcal{Y})$. We

define a loss function (6.24) $R^{K_1}: C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}) \times \mathcal{P}_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}, vol_{\mathcal{X}}) \to \mathbb{R}_{\geq 0}, (h, \mu) \mapsto \|(\Gamma_h)_*\mu_{\mathcal{X}} - \mu\|_{\tilde{K}_1}.$

Then for any $\mu \in \mathcal{P}_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}, vol_{\mathcal{X}})$, there exists a consistent C-ERM algorithm A for the supervised learning model $(\mathcal{X}, \mathcal{Y}, C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{K_2}), R^{K_1},$ $\mathcal{P}_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}, vol_{\mathcal{X}}))$. Moreover, for any $\varepsilon, \delta > 0$ there exists $N(\varepsilon, \delta)$ such that for any $n \geq N(\varepsilon, \delta)$ we have

(6.25)
$$
(\mu^n)^* \{ S_n \in (\mathcal{X} \times \mathcal{Y})^n : d_M(A(S_n), \mu_{\mathcal{Y}|\mathcal{X}}) > \varepsilon \} \le \delta,
$$

where $\mu_{\mathcal{Y}|\mathcal{X}} \in C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}$ is the unique regular conditional probability measure for μ with respect to the projection $\mathcal{P}_{\mathcal{X}} : \mathcal{X} \times \mathcal{Y} \to \mathcal{X}$.

A proof of Theorem [6.5](#page-38-0) shall be given in the next subsection, in particular the uniqueness of $\mu_{\mathcal{V}|\mathcal{X}}$ in [\(6.25\)](#page-39-1) shall be proved in Lemma [6.7.](#page-40-0) We shall provide the uniform consistency of a learning algorithm A for a supervised learning model $(\mathcal{X}, \mathcal{Y}, C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{K_2}), R^{K_1}, \mathcal{P}_{\mathcal{X}\times\mathcal{Y}})$, where $\mathcal{P}_{\mathcal{X}\times\mathcal{Y}} \subset$ $\mathcal{P}_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}, vol_{\mathcal{X}})$, in Corollary [6.13](#page-44-0) at the end of the next subsection.

6.2. Proof of Theorem [6.5.](#page-38-0) Assume the condition of Theorem [6.5.](#page-38-0) We shall apply Theorem [6.1](#page-36-0) to prove Theorem [6.5.](#page-38-0) First we shall prove Lemmas [6.6,](#page-39-2) [6.7](#page-40-0) and Propositions [6.9,](#page-40-1) [6.11](#page-42-0) to ensure that the conditions of Theorem [6.1](#page-36-0) are satisfied. We also prove Lemma [6.8,](#page-40-2) which shall guide us to define the lower semi-continuous function $W : C(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}) \to \mathbb{R}$ in Proposition [6.11,](#page-42-0) which is the most delicate technical part of the proof. Lemma [6.8](#page-40-2) is also important in the proof of Corollary [6.13.](#page-44-0)

For $h \in C(\mathcal{X}, \mathcal{M}(\mathcal{Y})_{\tilde{K}_2})$, we use the notation Γ_h instead of

$$
\overline{\Gamma_{\underline{h}}}:\mathcal{X}\to \mathcal{M}(\mathcal{X}\times \mathcal{Y}), x\mapsto \overline{\Gamma_{\underline{h}}}(x)=\delta_x\cdot h(x).
$$

Lemma 6.6. For any $\mu \in \mathcal{S}(\mathcal{X})$, the map

$$
\widehat{\mu}: C(\mathcal{X}, \mathcal{M}(\mathcal{Y})_{\tilde{K}_2})_M \to \mathcal{S}(\mathcal{X} \times \mathcal{Y})_{\tilde{K}_1}, h \mapsto (\Gamma_{\underline{h}})_*\mu
$$

is continuous. If $\text{sppt}\,\mu = \mathcal{X}$, then μ maps $C_b(\mathcal{X},\mathcal{M}(\mathcal{Y})_{\tilde{K}_2})$ 1-1 onto its image.

Proof. (1) Let $h, h' \in C(X, \mathcal{M}(\mathcal{Y})_{\tilde{K}_2})_M$ be bounded by a number α , i.e.,

$$
||h||_M \leq \alpha \& ||h'||_M \leq \alpha
$$

for some positive number $\alpha < \infty$. Let $\mu \in \mathcal{S}(\mathcal{X})$. By [\(4.6\)](#page-26-1) we have

$$
\begin{split} |||(\Gamma_{\underline{h}})_*\mu - (\Gamma_{\underline{h'}})_*\mu||^2_{\tilde{K}_1}| &\leq |\langle (\Gamma_{\underline{h}})_*\mu - (\Gamma_{\underline{h'}})_*\mu|(\Gamma_{\underline{h}})_*\mu\rangle_{\tilde{K}_1}| \\ &+ |\langle (\Gamma_{\underline{h'}})_*\mu|(\Gamma_{\underline{h}})_*\mu - (\Gamma_{\underline{h'}})_*\mu\rangle_{\tilde{K}_1}| \\ & = |\int_{\mathcal{X}\times\mathcal{X}} \langle \Gamma_h(x) - \Gamma_{h'}(x), \Gamma_h(x')\rangle_{\tilde{K}_1} d\mu(x) d\mu(x')| \\ &+ |\int_{\mathcal{X}\times\mathcal{X}} \langle \Gamma_{h'}(x), \Gamma_{h'}(x') - \Gamma_h(x')\rangle_{\tilde{K}_1} d\mu(x) d\mu(x')| \\ &\leq 2\alpha \cdot d_M(h, h') \cdot \mu(\mathcal{X})^2. \end{split}
$$

Inequality [\(6.26\)](#page-40-3) implies that $\hat{\mu}$ is a continuous map. This completes the proof of the first assertion of Lemma [6.6.](#page-39-2)

(2) The second assertion of Lemma [6.6](#page-39-2) follows from Lemma [2.21\(](#page-15-0)1). This completes the proof of Lemma [6.6.](#page-39-2) \Box

Lemma 6.7. Assume that $\mu \in \mathcal{P}_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}, vol_{\mathcal{X}})$. There exists a unique regular conditional probability measure $\mu_{\mathcal{Y}|\mathcal{X}} \in C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})$ for μ with respect to the projection $\Pi_{\mathcal{X}} : \mathcal{X} \times \mathcal{Y} \to \mathcal{X}$.

Proof. The existence of a regular conditional probability measure $\mu_{\mathcal{V}|\mathcal{X}} \in$ $C(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})$ for μ follows from the condition (ii) of Theorem [6.5.](#page-38-0) The uniqueness follows from Theorem [2.22\(](#page-16-1)2), taking into account the condition (i) of Theorem [6.5.](#page-38-0)

The following Lemma is a variant of Proposition [2.18.](#page-14-0)

Lemma 6.8. Assume the condition of Theorem [6.5.](#page-38-0) Let $T \in C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})$. Let K_3 denote the restriction of K to X. Then the map

$$
(\Gamma_T)_*: \mathcal{P}(\mathcal{X})_{\tilde{K}_3} \to \mathcal{S}(\mathcal{X} \times \mathcal{Y})_{\tilde{K}_1}
$$

is continuous.

Proof. By Proposition [2.18,](#page-14-0) the map

$$
(\Gamma_T)_*: (\mathcal{P}(\mathcal{X}), \tau_w) \to (\mathcal{P}(\mathcal{X} \times \mathcal{Y}), \tau_w)
$$

is continuous. Since the weak* topology τ_w on $\mathcal{P}(\mathcal{X})$ and on $\mathcal{P}(\mathcal{X} \times \mathcal{Y})$ are generated by the metric \tilde{K}_3 and \tilde{K}_1 , respectively, it follows that

$$
(\Gamma_T)_*: \mathcal{P}(\mathcal{X})_{\tilde{K}_3} \to (\mathcal{P}(\mathcal{X} \times \mathcal{Y})_{\tilde{K}_2})
$$

is continuous. Taking into account the continuity of the inclusion $(\mathcal{P}(\mathcal{X} \times \mathcal{Y})_{\tilde{\mathcal{E}}} \to \mathcal{S}(\mathcal{X} \times \mathcal{Y})_{\tilde{\mathcal{E}}}$, we conclude Lemma 6.8. $(\mathcal{Y})_{\tilde{K}_2} \to \mathcal{S}(\mathcal{X} \times \mathcal{Y})_{\tilde{K}_1}$, we conclude Lemma [6.8.](#page-40-2)

Proposition 6.9. Let $\mu \in \mathcal{P}_{epm}(\mathcal{X} \times \mathcal{Y})$. Then there exists a map $f \in$ $C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})$ such that

$$
(\Gamma_f)_*\mu_{\mathcal{X}}=\mu.
$$

Proof. Let $\mu := \sum_{i=1}^{N} \sum_{j=1}^{M} a_{ij} \delta_{x_i} \delta_{y_j} \in \mathcal{P}_{emp}(\mathcal{X} \times \mathcal{Y})$ where $a_{ij} \in \mathbb{Q}^+, x_i \in \mathcal{X}$ and $y_j \in \mathcal{Y}$. Then

(6.27)
$$
\mu_{\mathcal{X}} = \sum_{i=1}^{N} \sum_{j=1}^{M} a_{ij} \delta_{x_i}.
$$

Lemma 6.10. If $f \in C_{Lip}(\mathcal{X}, \mathcal{S}(\mathcal{Y})_{\tilde{K}_2})$ satisfies

(6.28)
$$
f(x_i) = \frac{\sum_{j=1}^{M} a_{ij} \delta_{y_j}}{\sum_{j=1}^{M} a_{ij}} \text{ for any } i \in [1, N],
$$

then

$$
(\Gamma_f)_*\mu_{\mathcal{X}}=\mu.
$$

Proof. Lemma [6.10](#page-41-0) follows directly from the equation [\(2.10\)](#page-9-3).

Continuation of the proof of Proposition [6.9.](#page-40-1) For $i \in [1, N]$ we set

$$
Y_i = \frac{\sum_{j=1}^{M} a_{ij} \delta_{y_j}}{\sum_{j=1}^{M} a_{ij}}.
$$

Let $V = \text{span}(Y_1, \ldots, Y_N)$ be the linear span of Y_1, \ldots, Y_N . First, we shall find a Lipschitz continuous map f from \mathbb{R}^n to V such that f satisfies the Equation [\(6.28\)](#page-41-1), equivalently:

$$
f(x_i) = Y_i.
$$

Then the restriction of f to $\mathcal X$ is the required Lipschitz continuous map in Proposition [6.9.](#page-40-1)

Let \mathbb{R} be a straight line in \mathbb{R}^n such that the projection $\Pi_1 : \mathbb{R}^n \to \mathbb{R}$ maps $\{x_1 \ldots, x_N \in \mathcal{X}\}\$ 1-1 onto its image in R. Since the restriction $\Pi_1|_{\mathcal{X}} : \mathcal{X} \to \mathbb{R}$ of Π_1 to X is a Lipschitz continuous map, to construct the required Lipschitz continuous map $f : \mathbb{R}^n \to V$, it suffices assume that $n = 1$. Now we shall construct a polynomial mapping f from $\mathbb R$ to V, which satisfies the interpolation equation [\(6.28\)](#page-41-1).

Given a tuple of N points (Y_1, \ldots, Y_N) in the vector space V of dimension $d \leq N-1$ and a tuple of N points $x_0 = 0 < x_1 < \ldots < x_{N-1} \in \mathbb{R}$, we shall construct a polynomial mapping $f : \mathbb{R} \to V$ such that

$$
(6.29) \t\t f(x_i) = Y_i.
$$

The coordinates of Y_i are denoted by y_i^j i^j . Our map f consists of polynomial functions $f^1, \ldots, f^d : \mathbb{R} \to \mathbb{R}$ such that

(6.30)
$$
f^{j}(x_{i}) = y_{i}^{j} \text{ for } j \in [1, d].
$$

We set

$$
f^i(x) := \lambda_i^0 + \lambda_i^1 x + \lambda_i^2 x(x - x_1) + \cdots + \lambda_i^n x(x - x_1) \cdots (x - x_n),
$$

where the coefficients λ_i^k are defined inductively: $\lambda_i^0 = y_i^0$, $\lambda_i^1 = y_i^1 - \lambda_i^0$, etc... Clearly the constructed polynomial mapping $f = (f^1, \ldots, f^d)$ satisfies (6.30) and hence (6.29) . This completes the proof of Proposition [6.9.](#page-40-1)

For any subset S of the metric space $C(\mathcal{X}, \mathcal{M}(\mathcal{Y})_{\tilde{K}_2})_M$ we denote by S_M the metric space S endowed with the induced metric d_M .

Proof of Theorem [6.5.](#page-38-0) Let $E_1 = C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})_M$, $\mathcal{A} = \mathcal{P}(\mathcal{X})$, $E_2 =$ $\mathcal{S}(\mathcal{X}\times\mathcal{Y})_{\tilde{K}_1}$ and $\mu \in \mathcal{P}(\mathcal{X}, \mathcal{P}(\mathcal{Y}_{\tilde{K}_2}), vol_{\mathcal{X}}) \subset E_2$. By Lemma [6.6,](#page-39-2) for any $\mu \in A$, the operator $\hat{\mu}: C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})_M \to E_2$ is a continuous operator. We shall apply Theorem [6.1](#page-36-0) to prove the existence of approximate solutions $f_{S_l} \in E_1$ for solving the following equation of a regular conditional probability measure f for μ relative to the projection $\Pi_{\mathcal{X}} : \mathcal{X} \times \mathcal{Y} \to \mathcal{X}$:

(6.31)
$$
(\Gamma_f)_*((\Pi_{\mathcal{X}})_*(\mu)) = \mu.
$$

By Theorem [4.6,](#page-29-0) f is the minimizer of the loss function R^{K_1} .

Let us define a function $W: C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})_M \to \mathbb{R}_{\geq 0}$ as follows

(6.32)
$$
W(f) := (\|f\|_M + L(f) + \|\Gamma_{\underline{f}}\|_{(\tilde{K}_3, \tilde{K}_1)})^2
$$

where $L(f)$ is the Lipschitz constant of f and

(6.33)
$$
\|\Gamma_{\underline{f}}\|_{\tilde{K}_3,\tilde{K}_1} := \sup_{A,B \in \mathcal{P}(\mathcal{X})} \frac{\|(\Gamma_{\underline{f}})_*(A-B)\|_{\tilde{K}_1}}{\|A-B\|_{\tilde{K}_3}}
$$

By Lemma [6.8,](#page-40-2) $\|\Gamma_{\underline{f}}\|_{\tilde{K}_3,\tilde{K}_1} < \infty$ for $f \in C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})_M$.

Proposition 6.11. (1) $W : C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})_M \to \mathbb{R}_{\geq 0}$ is a lower semicontinuous function.

.

(2) Furthermore, for any $c \geq 0$, the set $W^{-1}[0,c]$ is a compact set in $C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})_M$.

Proof. (1) First, we shall show that W is a lower semi-continuous function. Since the function $C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})_M \to \mathbb{R}_{\geq 0}, f \mapsto L(f)$, is a lower semicontinuous function, it suffices to show that the function $C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})_M \to$ $\mathbb{R}_{>0}, f \mapsto \|\Gamma_{\underline{f}}\|_{\tilde{K}_3, \tilde{K}_2}$, is lower semi-continuous. For $f \in C_L(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})_M$, let $A_{\varepsilon}, B_{\varepsilon} \in \mathcal{P}(\mathcal{X})$ be chosen such that

(6.34)
$$
\|\Gamma_{\underline{f}}\|_{\tilde{K}_3,\tilde{K}_1} \leq \frac{\|(\Gamma_{\underline{f}})_*(A_{\varepsilon} - B_{\varepsilon})\|_{\tilde{K}_1}}{\|A_{\varepsilon} - B_{\varepsilon}\|_{\tilde{K}_3}} + \varepsilon.
$$

Write $C_{\varepsilon} = A_{\varepsilon} - B_{\varepsilon} \in \mathcal{S}(\mathcal{X})$. Let $f' \in C_L(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})_M$ such that

(6.35)
$$
||f||_{\infty} \le \alpha \text{ and } ||f'||_{\infty} < \alpha
$$

for some positive $\alpha < \infty$.

Lemma 6.12. Under the assumption [\(6.35\)](#page-42-1) we have

(6.36)
$$
\|(\Gamma_{\underline{f}})_*C_{\varepsilon}\|_{\tilde{K}_1} \leq 2C_K\alpha \text{ and } \|(\Gamma_{\underline{f'}})_*C_{\varepsilon}\|_{\tilde{K}_1} \leq 2C_K\alpha.
$$

Proof. Lemma [6.12](#page-42-2) follows immediately from [\(6.26\)](#page-40-3) taking into account that $\|C_{\varepsilon}\|_{TV} \leq 2.$

Completion of the proof of Proposition [6.11](#page-42-0)(1). Let

$$
C_K = \max_{z \in \mathcal{X} \times \mathcal{Y}} K(z, z).
$$

Then we have (cf. the proof of Lemma [6.6](#page-39-2) and the proof of Lemma [5.4\)](#page-33-2)

$$
\begin{split} |||(\Gamma_{\underline{f'}})_*(C_{\varepsilon})||^2_{\tilde{K}_1} - ||(\Gamma_{\underline{f}})_*(C_{\varepsilon})||^2_{\tilde{K}_1}| &\leq |\langle (\Gamma_{\underline{f'}})_*(C_{\varepsilon}) - (\Gamma_{\underline{f}})_*(C_{\varepsilon})|(\Gamma_{\underline{f'}})_*(C_{\varepsilon})\rangle_{\tilde{K}_1}| \\ &+ |\langle (\Gamma_{\underline{f}})_*(C_{\varepsilon}) - (\Gamma_{\underline{f'}})_*(C_{\varepsilon})|(\Gamma_{\underline{f}})_*(C_{\varepsilon})\rangle_{\tilde{K}_1}| \\ (6.37) &\leq 4\alpha \cdot C_k ||(\Gamma_{\underline{f-f'}})_*(C_{\varepsilon})||_{\tilde{K}_1} \leq 4\alpha \cdot C_K ||f - f'||_{\infty} \cdot 2, \end{split}
$$

since $||C_{\varepsilon}||_{TV} \leq 2$. Since

(6.38)
$$
\|\Gamma_{\underline{f'}}\|_{\tilde{K}_3,\tilde{K}_1} \ge \Big(\frac{\|(\Gamma_{\underline{f'}})_*(C_{\varepsilon})\|_{\tilde{K}_1}}{\|C_{\varepsilon}\|_{\tilde{K}_3}}\Big),
$$

taking into account (6.34) and (6.38) , we conclude from (6.37) that

(6.39)
$$
\lim_{f' \to f} \|\Gamma_{\underline{f'}}\|_{\tilde{K}_3, \tilde{K}_1} \ge \|\Gamma_f\|_{\tilde{K}_3, K_1}.
$$

Equation [6.39](#page-43-2) implies that the function $f \mapsto ||\Gamma_f||_{\tilde{K}_3, \tilde{K}_2}$ is lower semicontinuous, and hence W is a lower semi-continuous function. This completes the proof of the first assertion of Proposition [6.11.](#page-42-0)

(2) Since $\mathcal Y$ is a compact subset of \mathbb{R}^m , by Sturm's result [\[Sturm06,](#page-50-7) Proposition 2.10 (iii)], the set $\mathcal{P}(\mathcal{Y})$ is compact in the weak*-topology τ_w . Since $\mathcal X$ is compact, by Arzelà-Ascoli theorem [\[Kelley75,](#page-49-19) Theorem 18, Chapter 7, p. 234], for any $c \geq 0$ the set $W^{-1}[0, c]$ is compact in E_1 . This completes the proof of Proposition [6.11.](#page-42-0)

By Lemmas [6.6,](#page-39-2) [6.7](#page-40-0) and Propositions [6.9,](#page-40-1) [6.11,](#page-42-0) all the requirements of Theorem [6.1](#page-36-0) are satisfied.

Let $\mu_{\mathcal{X},l} = (\pi_{\mathcal{X}})_*\mu_{S_l} \in \mathcal{P}(\mathcal{X})$ where $\mu_{S_l} \in \mathcal{P}_{emp}(\mathcal{X}\times\mathcal{Y})$. For $\mu \in \mathcal{P}(\mathcal{X}\times\mathcal{Y})$, we have

$$
\|\mu_{\mathcal{X},l} - \mu_{\mathcal{X}}\| \stackrel{(6.4)}{=} \sup_{f \in E_1} \frac{\|(\Gamma_{\underline{f}})_*(\mu_{\mathcal{X},l} - \mu)\|_{\tilde{K}_1}}{W^{1/2}(f)}
$$

(6.40)
$$
\leq \sup_{f \in E_1} \frac{\|(\Gamma_{\underline{f}})_*(\mu_{\mathcal{X},l} - \mu_{\mathcal{X}})\|_{\tilde{K}_1}}{\|\Gamma_f\|_{\tilde{K}_3,\tilde{K}_2}} \stackrel{(6.33)}{\leq} \|\mu_{\mathcal{X},l} - \mu_{\mathcal{X}}\|_{\tilde{K}_1}.
$$

Applying Theorem [6.1](#page-36-0) to our case with $(\mathcal{X}_l, \mu_l) = ((\mathcal{X} \times \mathcal{Y})^l, \mu^l)$, and letting f_{S_l} to be a γ_l^2 -minimizer of the regularized risk function

$$
R_{\gamma_l}^{K_1}(f, \mu_{S_l}) = R^{K_1}(f, \mu_{S_l}) + \gamma_l W(f),
$$

taking into account [\(6.40\)](#page-43-3), we obtain the following estimation

$$
(\mu^l)^* \{ S_l : d_M(f_{S_l}, \mu_{\mathcal{Y}|\mathcal{X}}) > \varepsilon \} \le (\mu^l)^* \{ S_l : \frac{\|\mu_l - \mu\|_{\tilde{K}_1}}{\sqrt{\gamma_l}} \ge C_1 \}
$$

(6.41)

$$
+ (\mu^l)^* \{ S_l : \frac{\|\mu_{\mathcal{X},l} - \mu_{\mathcal{X}}\|_{\tilde{K}_3}}{\sqrt{\gamma_l}} \ge C_2 \}.
$$

Taking into account Proposition [4.2](#page-25-2) and Lemma [6.6,](#page-39-2) we conclude Equation [\(6.25\)](#page-39-1) in Theorem [6.5](#page-38-0) from [\(6.41\)](#page-44-1). The consistency of the C-ERM algorithm follows from Equations [\(6.25\)](#page-39-1) and [\(6.26\)](#page-40-3).

Corollary 6.13. Assume that $\mathcal{P}_{\mathcal{X}\times\mathcal{Y}} \subset \mathcal{P}_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}, vol_{\mathcal{X}})$ satisfies the following condition (L).

(L) The function $(\mathcal{P}_{\mathcal{X}\times\mathcal{Y}})_{\tilde{K}_1} \to \mathbb{R}, \mu \mapsto L(\mu_{\mathcal{Y}|\mathcal{X}})$, where $\mu_{\mathcal{Y}|\mathcal{X}} \in C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})$, takes a value on a finite interval $[a, b] \subset \mathbb{R}$.

(1) Then the algorithm A defined in the proof of Theorem [6.5](#page-38-0) is uniformly consistent for the supervised learning model $(\mathcal{X}, \mathcal{Y}, C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}), \mathcal{P}_{\mathcal{X}\times\mathcal{Y}}).$

(2) Assume further that the PDS kernel $K : \mathbb{R}^{n+m} \times \mathbb{R}^{n+m} \to \mathbb{R}$ in Theorem [6.5](#page-38-0) is continuously (C^1-) differentiable and H is a subspace of the space $C_{Lip}(\mathcal{X}, \mathcal{Y})$. Then the supervised learning model $(\mathcal{X}, \mathcal{Y}, \mathcal{H}, R^{K_1}, \mathcal{P}_{\mathcal{X}\times\mathcal{Y}})$ has a generalization ability.

Proof. (1) To prove the first assertion of Corollary [6.13,](#page-44-0) taking into account the condition (L) and Equation [6.26,](#page-40-3) it suffices to show that the coefficient $\gamma_0 = \gamma(C_1, C_2, W(\mu_{\mathcal{Y}|\mathcal{X}}), A^{-1}(\mu), \varepsilon)$ defined in [\(6.19\)](#page-37-4) in the proof of Theorem [6.1,](#page-36-0) adapted to the proof of Theorem [6.5,](#page-38-0) can be chosen independently of $\mu \in \mathcal{P}_{\mathcal{X}\times\mathcal{Y}}$, where $A^{-1}(\mu) = \mu_{\mathcal{Y}|\mathcal{X}}$. By the condition of Corollary [6.13,](#page-44-0) $W(f)$, and hence the constant d in Equation [\(6.17\)](#page-37-3), and therefore the compactum $W^{-1}[0, 2d]$ can be chosen independently from μ . Let $\mathcal{A}^{\circ} := \{\mu \in \mathcal{P}(\mathcal{X}) :$ sppt $\mu = \mathcal{X}$. Since X is compact, by Sturm's result [\[Sturm06,](#page-50-7) Proposition] 2.10 (iii)], $\mathcal{P}(\mathcal{X})$ is compact is the weak*-topology and hence \mathcal{A}° is compact in the weak*-topology. Now we define the operator

$$
U: \mathcal{P}(\mathcal{X})_{\tilde{K}_3} \times W^{-1}[0, 2d] \to \mathcal{P}_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}, vol_{\mathcal{X}}), (\mu, h) \mapsto \Gamma_{\underline{h}}(\mu).
$$

We shall show that U is a continuous map. Let $\mu, \mu' \in \mathcal{A}^{\circ}$ and $h, h' \in$ $W^{-1}[0, 2d]$. Then we have

$$
\begin{aligned} \|(\Gamma_{\underline{h}'})_* \mu' - \Gamma_{\underline{h}})_* \mu \|_{\tilde{K}_1} &\le \|(\Gamma_{\underline{h}'})_* \mu' - (\Gamma_{\underline{h}})_* \mu' \|_{\tilde{K}_1} \\ &+ \|(\Gamma_{\underline{h}})_* \mu' - \Gamma_{\underline{h}} \mu \|_{\tilde{K}_1} \\ (6.42) &\le 2C_K \cdot d_M(h, h') + (\Gamma_{\underline{h}})_* (\mu' - \mu). \end{aligned}
$$

By Lemma [6.8,](#page-40-2) $(\Gamma_h)_*(\mu'-\mu)$ converges to zero when μ' converges to μ . Taking into account (6.42) , we conclude that U is a continuous map. By Theorem [2.22](#page-16-1) and taking into account the condition (i) of Theorem [6.5,](#page-38-0) U is a 1-1 map. By Lemma [6.3,](#page-37-5) the map

$$
U^{-1} : (U(\mathcal{A}^{\circ} \times W^{-1}[0, 2d])) \to \mathcal{A}^{\circ} \times W^{-1}[0, d], \mu \mapsto (\mu_{\mathcal{X}}, \mu_{\mathcal{Y}|\mathcal{X}}),
$$

46 H. V. LEˆ

is continuous. Since $U(\mathcal{A}^{\circ} \times W^{-1}[0, 2d])$ is compact, U^{-1} is uniformly continuous. It follows that for any $\varepsilon > 0$ there exists $\delta(\varepsilon) > 0$ such that

(6.43)
$$
\|\mu - \mu'\|_{\tilde{K}_1} \le \delta(\varepsilon) \Longrightarrow d_M(\mu_{\mathcal{Y}|\mathcal{X}}, \mu_{\mathcal{Y}|\mathcal{X}}') \le \varepsilon.
$$

Since $\mathcal{P}_{\mathcal{X}\times\mathcal{Y}} \subset U(\mathcal{A}^{\circ}\times W^{-1}[0,2d]),$ [\(6.43\)](#page-45-0) implies that $\delta(\varepsilon)$ in Equation [6.20](#page-37-8) and hence $\gamma_0 = \gamma(C_1, C_2, W(\mu_{\mathcal{Y}|\mathcal{X}}), A^{-1}\mu, \varepsilon)$ can be chosen independently from μ . This completes the proof of Corollary [6.13\(](#page-44-0)1).

(2) Assume the condition of Corollary [6.13\(](#page-44-0)2). Since the PDS kernel K is C¹-differentiable, the map $\delta: \mathcal{Y} \to \mathcal{P}(\mathcal{Y})_{\tilde{K}_2}, y \mapsto \delta_y$, is a Lipschitz map. Hence for any $h \in \mathcal{H}$ the composition $\delta \circ h$ belongs to $C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})$. To prove Corollary [6.13\(](#page-44-0)2), it suffices to find an algorithm $\bar{A}: \cup_n (\mathcal{X} \times \mathcal{Y})^n \to \mathcal{H}$ with the following property. For any $\varepsilon, \delta > 0$ there exists $N(\varepsilon, \delta)$ such that for any $n \geq N(\varepsilon, \delta)$ and for any $\mu \in \mathcal{P}_{\mathcal{X} \times \mathcal{Y}}$ we have

$$
(6.44) \qquad (\mu^n)^* \{ S_n \in (\mathcal{X} \times \mathcal{Y})^n : R_\mu^{K_1}(\bar{A}(S_n)) - \inf_{h \in \mathcal{H}} R_\mu^{K_1}(h) \ge \varepsilon \} \le \delta.
$$

Let $A: \cup_n (\mathcal{X} \times \mathcal{Y})^n \to C_{Lip}(\mathcal{X}, \mathcal{P}(\mathcal{Y})_{\tilde{K}_2})$ be the uniformly consistent learning algorithm in the proof of Theorem [6.5.](#page-38-0) By [\(6.25\)](#page-39-1), for any $\varepsilon, \delta > 0$ there exists $N_A(\varepsilon,\delta)$ such that for any $n \geq N_A(\varepsilon,\delta)$ and for any $\mu \in \mathcal{P}_{\mathcal{X}\times\mathcal{Y}}$ we have

$$
(6.45) \qquad (\mu^n)^* \{ S_n \in (\mathcal{X} \times \mathcal{Y})^n : \| (\Gamma_{A(S_n)})_*(\mu_{S_n})_{\mathcal{X}} - \mu \|_{\tilde{K}_1} \ge \frac{\varepsilon}{3} \} \le \delta.
$$

Let $\bar{A}(S_n) \in \mathcal{H}$ be any element such that

$$
\| (\Gamma_{\bar{A}(S_n)})_*(\mu_{S_n})_{\mathcal{X}} - (\Gamma_{A(S_n)})_*(\mu_{S_n})_{\mathcal{X}} \|_{\tilde{K}_1}
$$

(6.46)
$$
\leq \inf_{h \in \mathcal{H}} \| (\Gamma_h)_*(\mu_{S_n})_{\mathcal{X}} - (\Gamma_{A(S_n)})_*(\mu_{S_n})_{\mathcal{X}} \|_{\tilde{K}_1} + \alpha_n
$$

where α_n is a sufficiently small number to be defined later. Then we have

$$
R_{\mu}^{K_{1}}(\bar{A}(S_{n})) - \inf_{h \in \mathcal{H}} R_{\mu}^{K_{1}}(h)
$$

\n
$$
\leq ||(\Gamma_{\bar{A}(S_{n})})_{*}(\mu_{S_{n}})_{\mathcal{X}} - (\Gamma_{A(S_{n})})_{*}(\mu_{S_{n}})_{\mathcal{X}}||_{\tilde{K}_{1}} + ||(\Gamma_{A(S_{n})})_{*}(\mu_{S_{n}})_{\mathcal{X}} - \mu||_{\tilde{K}_{1}}
$$

\n
$$
- \inf_{h \in \mathcal{H}} R_{\mu}^{K_{1}}(h)
$$

\n
$$
\leq ||(\Gamma_{\bar{A}(S_{n})})_{*}(\mu_{S_{n}})_{\mathcal{X}} - (\Gamma_{A(S_{n})})_{*}(\mu_{S_{n}})_{\mathcal{X}}||_{\tilde{K}_{1}} + ||(\Gamma_{A(S_{n})})_{*}(\mu_{S_{n}})_{\mathcal{X}} - \mu||_{\tilde{K}_{1}}
$$

\n
$$
- \inf_{h \in \mathcal{H}} ||(\Gamma_{h})_{*}(\mu_{S_{n}})_{\mathcal{X}} - (\Gamma_{A(S_{n})})_{*}(\mu_{S_{n}})_{\mathcal{X}}||_{\tilde{K}_{1}} + ||(\Gamma_{A(S_{n})})_{*}(\mu_{S_{n}})_{\mathcal{X}} - \mu||_{\tilde{K}_{1}}
$$

\n(6.46)
\n
$$
\leq 2 ||(\Gamma_{A(S_{n})})_{*} \mu_{\mathcal{X}} - \mu||_{\tilde{K}_{1}} + \alpha_{n}.
$$

Taking into account [\(6.45\)](#page-45-2), we obtain from Equation [6.47](#page-45-3) the following

$$
(6.48) \quad (\mu^n)^* \{ S_n \in (\mathcal{X} \times \mathcal{Y})^n : \| (\Gamma_{\bar{A}(S_n)})_*(\mu_{S_n})_{\mathcal{X}} - \mu \|_{\tilde{K}_1} \ge \frac{2\varepsilon}{3} + \alpha_n \} \le \delta.
$$

By letting $\alpha_n \leq \frac{\varepsilon}{3}$ $\frac{\varepsilon}{3}$, we obtain [\(6.44\)](#page-45-4) and, hence, the last assertion of Corollary [6.13](#page-44-0) immediately from [\(6.48\)](#page-45-5).

Example 6.14. Let $\mathcal{X} = [0,1]^n \subset \mathbb{R}^n \times \{0\} \subset \mathbb{R}^{n+m}$ and $\mathcal{Y} = [0,1]^m \subset \mathbb{R}^n$ $\{0\}\times\mathbb{R}^m\subset\mathbb{R}^{n+m}$. Denote by dx the restriction of the Lebesgue measure on \mathbb{R}^n to X and by dy the restriction of the Lebesgue measure on \mathbb{R}^m to Y. Let $\mathcal{P}_{\mathcal{X}\times\mathcal{Y}}$ consist of probability measures $\mu_f := fdxdy$, where $f \in C^1(\mathcal{X}\times\mathcal{Y})$ and moreover $f(x, y) > 0$ for all $x, y \in X \times Y$. We shall show that if the PDS kernel $K : \mathbb{R}^{n+m} \times \mathbb{R}^{n+m} \to \mathbb{R}$ in Theorem [6.5](#page-38-0) is continuously (C^1-) differentiable, then $P_{\chi \times \chi}$ satisfies the condition of Corollary [6.13.](#page-44-0) First we note that

$$
\mu_X = q dx
$$
 where $q(x) = \int_X f(x, y) dy$.

Since $f(x, y) > 0$, we have $q(x) > 0$ for all $x \in \mathcal{X}$. Therefore sppt $\mu_{\mathcal{X}} = \mathcal{X}$. Hence the condition (i) in Theorem [6.5](#page-38-0) on μ_f is satisfied.

Next we note that $q \in C^1(\mathcal{X})$, since $f \in C^1(\mathcal{X} \times \mathcal{Y})$ and $\mathcal{X} \times \mathcal{Y}$ is compact [\[Jost05,](#page-49-20) Theorem 16.11, p. 213]. We observe that

$$
(\mu_f)_{\mathcal{Y}|\mathcal{X}}(x) = g(\cdot|x) dy, \text{ where } g(y|x) = \frac{f(x,y)}{q(x)}.
$$

Note that $\mu_f : \mathcal{X} \to \mathcal{P}(\mathcal{Y}), x \mapsto (\mu_f)_{\mathcal{Y}|\mathcal{X}}(x)$, is a continuous map with respect to the strong topology on $\mathcal{P}(y)$, and hence μ_f is a measurable map. Let $K_2: \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ and $K_1: (\mathcal{X} \times \mathcal{Y}) \times (\mathcal{X} \times \mathcal{Y}) \to \mathbb{R}$ be continuous bounded SPD kernels defined as in Theorem [6.5.](#page-38-0) We shall show that both the maps

$$
\mathfrak{M}_{K_2} \circ \mu_f: \mathcal{X} \to \mathcal{H}(K_2), x \mapsto \mathfrak{M}_{K_2} \circ \mu_f(x),
$$

$$
\mathfrak{M}_{K_1} \circ \Gamma_{\mu_f}: \mathcal{X} \to \mathcal{H}(K), x \mapsto \mathfrak{M}_{K_1} \circ \Gamma_{\mu_f}(x),
$$

are C^1 -differentiable.

Let us first prove that $\mathfrak{M}_{K_2} \circ \mu_f$ is C^1 -differentiable. Let x_t be a smooth curve $(-\varepsilon, \varepsilon) \to \mathcal{X}, t \mapsto x_t$. Then

(6.49)
$$
\mathfrak{M}_{K_2}(\mu_f(x_t)) = \int_{\mathcal{Y}} (K_2)_y \, d\mu_f(y|x_t) = \int_{\mathcal{Y}} (K_2)_y g(y|x_t) \, dy
$$

where the RHS of (6.49) is understood as the Bochner integral. Since K_2 : $\mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ is bounded and continuous, and $g(\cdot|\cdot)$ is bounded and C^1 differentiable in x, we can perform differentiation under integral. Thus, for $v = \frac{dx_t}{dt}_{|t=0} \in T_{x_0} \mathcal{X}$ we have

$$
d_{x_0}(\mathfrak{M}_{K_2} \circ \mu_f) v = \int_{\mathcal{Y}} (K_2)_y \frac{dg(y|x_t)}{dt}_{|t=0} dy.
$$

Hence, $\mathfrak{M}_{K_2} \circ \mu_f$ is C^1 -differentiable. Similarly, we prove that $\mathfrak{M}_{K_1} \circ \Gamma_{\mu_f}$ is C^1 -differentiable, using also the differentiability of K_1 and of $g(\cdot|\cdot)$ in variable x . Hence, these maps are Lipschitz continuous with Lipschitz constants belonging to an bounded interval $[a', b'] \subset \mathbb{R}$. Consequently, μ_f satisfies the condition of Corollary [6.13.](#page-44-0)

Note that the Gaussian kernels on \mathbb{R}^n are C^1 -differentiable but the Laplacian kernels are not C^1 -differentiable.

48 H. V. LEˆ

7. Discussion of results

(1) In this paper we have demonstrated the usefulness of the new concept of a generative model of supervised learning, which naturally incorporates the concept of a correct loss function. The generative model provides a unified framework for density estimation problems, supervised learning problems, and many problems in Bayesian statistics.

(2) We have developed the theory of probabilistic morphisms and applied the obtained results to characterize regular conditional probability measures, which are key tools for defining a correct loss function in a generative model of supervised learning. Probabilistic morphisms also offer tools and insights for proving the generalizability of generative models in supervised learning.

(3) The usefulness of the inner measure, equivalently, the convergence in outer probability, for defining and proving the generalization ability of statistical learning models has been demonstrated.

(4) We present examples of uniformly consistent algorithms for overparameterized discriminative models of supervised learning (Corollary [6.13](#page-44-0) (2)).

(5) To improve Theorem [6.5](#page-38-0) and Corollary [6.13,](#page-44-0) we can relax the condition (i) in Theorem [6.5.](#page-38-0) One possible approach could be a combination of the methods developed in this paper and techniques of vector-valued reproducing kernel Hilbert spaces (RKHSs), as developed in [\[CDT06\]](#page-48-19), [\[CDTU10\]](#page-48-20). Corollary [6.13](#page-44-0) and Examples [6.14](#page-46-0) suggest that the generalizability of a generative model of supervised learning $(\mathcal{X}, \mathcal{Y}, \mathcal{H}, L, \mathcal{P}_{\mathcal{X}\times\mathcal{Y}})$ depends on the geometry of H and $\mathcal{P}_{\mathcal{X}\times\mathcal{Y}}$ as well as on the choice of positive definite symmetric (PDS) kernels.

(6) Another possible research direction could involve combining the methods developed in this paper with those presented in [\[LV2009\]](#page-49-21), [\[Sturm06\]](#page-50-7).

Acknowledgement

The author wishes to thank Frederic Protin for proposing her the loss function in [\(4.14\)](#page-30-3), which led her to use kernel mean embedding for a characterization of regular conditional probability measures (Theorem [4.6\)](#page-29-0). She would like to thank Tobias Fritz for helpful comments on results in Section [2.](#page-4-0) A part of this paper has been prepared while the author was Visiting Professor of the Kyoto University from July till October 2022. She is grateful to Kaoru Ono and the Research Institute of Mathematical Sciences for their hospitality and excellent working conditions. The author also thanks Wilderich Tuschmann and Xia Kelin for inviting her to the Online Conference "Applied Geometry for Data Sciences" in Chongquing, China, July 2022, where she had an opportunity to report some results of this paper. Finally she expresses her indebtedness to Alexei Tuzhilin for stimulating discussions on measure theory for a long time and his suggestion to use outer measure about five years ago.

REFERENCES

- [AJLS17] N, Ay, J. Jost, H.V. Lê, L. Schwachhöfer, Information geometry, Springer Nature: Cham, Switzerland, 2017.
- [Aronszajn50] N. Aronszajn, Theory of reproducing kernels. Trans. Amer. Math. Soc. 68 (1950),337-404.
- [BH78] J. Bretagnolle, C. Huber, Estimation des densités : Risque minimax, Séminaire de Probabilit´es XII, Lecture notes in Mathematics, Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 342-363 (1978).
- [BLM13] S. Boucheron, G. Lugosi, P. Massart, Concentration Inequalities, A Nonasymptotic Theory of Independence, Oxford University Press, 2013.
- [BP1973] L. D. Brown, R. Purves, Measurable Selections of Extrema, Ann. Statist. 1(1973), 902-912 .
- [BT04] A. Berlinet and C. Thomas-Agnan, Reproducing Kernel Hilbert Spaces in Probability and Statistics. Kluwer Academic Publishers, 2004.
- [Bochner33] S. Bochner, Integation von Funktionen, deren Wert die Elemente eines Vektorraumes sind, Fund. Math. 20 (1933), 262-276.
- [Bogachev07] V. I. Bogachev, Measure Theory, I, II, Springer, 2007.
- [Bogachev18] V.I. Bogachev, Weak convergence of measures, Mathematical Surveys and Monographs, vol. 234, Amer. Math. Soc.: Providence, RI, USA, 2018.
- [CD07] A. Capponentto and E. De Vito, Optimal rates for the regularized least-square algorithms, Foundations of Computational Mathematics, 7(2007), 331-368.
- [CDT06] C. Carmeli, E. De Vito, and A. Toigo, Vector Valued Reproducing Kernel Hilbert Spaces of Integrable Functions and Mercer Theorem. Analysis and Applications, 4(2006), 377-408.
- [CDTU10] C. Carmeli, E. De Vito, A. Toigo, and V. Umanitá. Vector valued reproducing kernel hilbert spaces and universality, Analysis and Applications, 8(2010), 19-61.
- [CJ19] K. Cho and B. Jacobs, Disintegration and Bayesian inversion via string diagrams, Mathematical Structures in Computer Science, Volume 29 , Issue 7 (2019), 938 - 971.
- [Chentsov72] N. Chentsov, Statistical decision rules and optimal inference, Nauka: Moscow, Russia, 1972, English translation in: Translation of Math. Monograph vol. 53, Amer. Math. Soc.: Providence, RI, USA, 1982.
- [CS01] F. Cucker and S. Smale, On mathematical foundations of learning. Bulletin of AMS, 39 (2002), 1-49.
- [DGL96] L. Devroye, L. Györfi and G. Lugosi, A Probabilistic Theory of Pattern Recognition. Springer 1996.
- [DM1978] C. Dellacherie and P.-A. Meyer, Probabilities and potential, North Holland, 1978.
- [Dudley2014] R. M. Dudley, Uniform Central Limit Theorems, Second Edition, Cambridge, 2014.
- [FGPR2020] T. Fritz, T. Gonda, P. Perrone, E. F. Rischel, Representable Markov Categories and Comparison of Statistical Experiments in Categorical Probability, Theoretical Computer Science, Volume 961, 15 June 2023, 113896, [arXiv:2010.07416.](http://arxiv.org/abs/2010.07416)
- [FGSS08] K. Fukumizu, A. Gretton, X. Sun, and B. Schölkopf. Kernel measures of conditional dependence. In Advances in neural information processing systems 20, pages 489-496, Red Hook, NY, USA, 9 2008. Curran Associates, Inc.
- [Fritz2019] T. Fritz, A synthetic approach to Markov kernel, conditional independence and theorem of sufficient statistics. Adv. Math. 370, 107239 (2020), [arXiv:1908.07021.](http://arxiv.org/abs/1908.07021)
- [FPR2021] T. Fritz, P. Perrone and S. Rezagholi, Probability, valuations, hyperspace: Three monads on Top and the support as a morphism, Mathematical Structures in Computer Science 31(8), 850-897 (2021), [arXiv:1910.03752.](http://arxiv.org/abs/1910.03752)
- [FR19] T. Fritz, E. F. Rischel, Infinite products and zero-one laws in categorical probability. Compositionality 2, 3 (2020), DOI: 10.32408/compositionality-2-3, [arXiv:1912.02769.](http://arxiv.org/abs/1912.02769)
- [FS20] S. Fisher and I. Steinwart, Sobolev Norm Learning rate for regularied Least-Squares algorithms, J. of Machine Learning Research, 21(2020), 1-38.
- [GH89] M. Gaudard, and D. Hadwin, Sigma-algebras on spaces of probability measures, Scandinavian Journal of Statistics 16, 2 (1989), 169–175.
- [Giry82] M. Giry, A categorical approach to probability theory, In: B. Banaschewski, editor, Categorical Aspects of Topology and Analysis, Lecture Notes in Mathematics 915, 68- 85, Springer, 1982.
- [GLGB12] G. Grünewälder, G. Lever, A. Gretton, L. Baldassarre, S. Patterson, and M. Pontil. Conditional mean embeddings as regressors. In Proceedings of the 29th International Conference on Machine Learning (ICML), 2012.
- [JLT21] J. Jost, H. V. Lê, and T. D. Tran, Probabilistic morphisms and Bayesian nonparametrics, [Eur. Phys. J. Plus 136, 441 \(2021\),](https://doi.org/10.1140/epjp/s13360-021-01427-7) [arXiv:1905.11448.](http://arxiv.org/abs/1905.11448)
- [Jost05] J. Jost, Postmodern Analysis, Springer 2005.
- [Kelley75] J. L. Kelley, General Topology, Springer 1975.
- [Kallenberg2014] O. Kallenberg, Stationary and invariant densities and disintegration kernels, Probab. Theory Relat. Fields, 160(2014), 567-592.
- [Kock11] A. Kock, Commutative monads as a theory of distributions, [arXiv:1108.5952.](http://arxiv.org/abs/1108.5952)
- [Lawere62] W. F. Lawvere, The category of probabilistic mappings. Available at https://ncatlab.org/nlab/files/lawvereprobability1962.pdf., 1962.
- [Le2022] H. V. Lê, Natural differentiable structures on statistical models and the Fisher metric, [Information Geometry, 2022.](https://doi.org/10.1007/s41884-022-00090-w)
- [LFR04] D. Leao Jr., M. Fragoso and P. Ruffino, Regular conditional probability, disintegration of probability and Radon spaces. Proyecciones vol. 23 (2004)Nr. 1, Universidad Catolica Norte, Antofagasta, Chile, 15-29.
- [LMST15] D. Lopez-Paz, K. Muandet, B. Schölkopf, and I. Tolstikhin. Towards a learning theory of cause-effect inference. In Proceedings of the 32nd International Conference on Machine Learning (ICML2015), 2015.
- [LP16] G. Last and M. Penrose, Lectures on Poisson process, CUP, 2016.
- [LV2009] J. Lott and C. Villani, Ricci curvature for metric-measure spaces via optimal tranport, Annals of Mathematics, 169 (2009), 903-991.
- [MFSS17] K. Muandet, K. Fukumizu, B. Sriperumbudur and B. Schölkopf, Kernel Mean Embedding of Distributions: A Review and Beyond. Foundations and Trends in Machine Learning: Vol. 10: No. 1-2, pp 1-141 (2017), [arXiv:1605.09522.](http://arxiv.org/abs/1605.09522)
- [Parthasarathy67] K. R. Parthasarathy, Probability Measures on Metric Spaces. Academic Press, New York, 1967.
- [PM20] J. Park and K. Muandet, A measure-theoretic approach to kernel conditional mean embeddings, 34th Conference on Neural Information Processing Systems (NeurIPS 2020), Vancouver, Canada, [arXiv:2002.03689.](http://arxiv.org/abs/2002.03689)
- [SC08] I. Steinwart and A. Christmann, Support Vector Machines, Information Sciences and Statistics, Springer, 2008.
- [Sheynin77] O. B. Sheynin, Laplace's Theory of Errors, Archive for History of Exact Sciences , 31.V.1977, Vol. 17, No. 1 (31.V.1977), 1-61, Springer.
- [SFL11] B.K. Sriperumbudur, K. Fukumizu and G.R.G. Lanckriet, Universality, characteristic kernels and RKHS embedding of measures, J. Mach. Learn. Res. 12 2389-2410(2011).
- [Sriperumbudur16] B. Sriperumbudur, On the optimal estimation of probability measures in weak and strong topologies. Bernoulli, 22(3):1839-1893, 08, 2016.
- [Staton17] S. Staton, Commutative semantics for probabilistic programming. In European Symposium on Programming, pages 855-879, Springer, 2017.
- [Sturm06] K.-L. Sturm, On the geometry of metric measure spaces. I, Acta Math., 196 (2006), 65-131.
- [TSS2022] P. Talwai, A. Shameli, D. Simchi-Levi, Sobolev Norm Learning Rates for Conditional Mean Embeddings, Proceedings of the 25th International Conference on Artificial Intelligence and Statistics (AISTATS) 2022, Valencia, Spain. PMLR: volume 151.
- [Vapnik98] V. Vapnik, Statistical Learning Theory. John Willey & Sons, 1998.
- [VS1978] V. Vapnik and A. Stefanyuk, Nonparametric methods for estimating probability densities. Automation and Remote Control, 8, 38-52.
- [VW1996] A. W. Van der Vaart and J. A. Wellner, Weak convergence and Empirical Processes, Springer, 1996.
- [VI2019] V. Vapnik and R. Izmailov, Rethinking statistical learning theory: learning using statistical invariants, Machine Learning (2019) 108:381-423.
- [Wald50] A. Wald, Statistical Decision Functions. Wiley, 1950.
- [Yosida95] K. Yosida, Functional Analysis. Springer, 1995.

Institute of Mathematics of the Czech Academy of Sciences, Zitna 25, 11567 PRAHA 1, CZECH REPUBLIC

Email address: hvle@math.cas.cz