

A NOTE ON LOCALLY CONFORMALLY KÄHLER STRUCTURES AND SMALL DEFORMATIONS OF HOPF MANIFOLDS

KEIZO HASEGAWA

ABSTRACT. A Hopf manifold is a compact complex manifold of which the universal covering is $\mathbb{C}^n \setminus \{0\}$. In this note we show that any Hopf manifold admits a locally conformally Kähler structure (shortly lcK structure), by constructing a complex analytic family around a Hopf manifold of diagonal type, which admits a lcK potential, and applying a well known fact (due to Ornea and Verbitsky) that the property of lcK potential is preserved under a complex analytic family over a sufficiently small parameter space.

1. INTRODUCTION

A *generalized Hopf manifold* or simply a *Hopf manifold* M of complex dimension n is a compact complex manifold of which the universal covering is $W := \mathbb{C}^n \setminus \{0\}$, where n is a positive integer ($n \geq 2$). Let G be the covering transformation group of M . Then G acts properly discontinuously without fixed points as a holomorphic automorphism group on W . We can see [7] that G contains an infinite cyclic subgroup Z of finite index, which defines a contraction map on W (see Sec. 2 for definition). A *primary Hopf manifold* M is a compact complex manifold of which the covering transformation group is an infinite cyclic subgroup Z , that is, $M = W/Z$. It turns out that a Hopf manifold in general has a primary Hopf manifold as a finite normal covering. Note that in the literature a primary Hopf manifold is often referred to as a Hopf manifold. The generalized definition above is due to Kodaira [4].

It is well known that a Hopf manifold is the first example of a non-Kähler compact complex manifold, that is, it does not admit a Kähler structure compatible with its complex structure. And this is also the case for a generalized Hopf manifold; that is, any compact complex manifold whose universal covering is W admit no Kähler structures. This is easily seen, for instance, due to the fact that the diffeomorphism type of a primary Hopf manifold is $S^1 \times S^{2n-1}$. On the other hands, a Hopf manifold is also known to give an example of a *locally conformally Kähler manifold* (or shortly *lcK manifold*) (see Sec. 2 for definition), which is a generalization of Kähler manifold. A problem whether any (generalized) Hopf manifold admits a lcK structure is affirmative, as we see in this note. A (primary) Hopf manifold W/Z , where Z is the covering transformation group generated by a contraction (see Sec. 2 for definition), is called of *linear type* if Z is generated by a linear contraction (non-singular linear map on \mathbb{C}^n); and it is called of *diagonal type* if Z is generated by a diagonalizable linear contraction. It is known that a linear Hopf manifold admit a lcK structure. There is recently a preprint [12] where the authors show

that a non-linear Hopf manifold can be embedded into a linear Hopf manifold, and thus admit a lcK structure. There is a more direct and succinct way to show this result. In fact, the author (of this note) already made a notice at some workshops (including the workshop “*Geometric Structures on Complex Manifolds*”, Steklov Institute of Mathematics, 2011, Moscow) that there is a complex analytic family over \mathbb{C} deforming a non-linear Hopf manifold into a linear Hopf manifold, and another complex analytic family over \mathbb{C} deforming a linear Hopf manifold to a Hopf manifold of diagonal type such that all fibers over $\mathbb{C} \setminus \{0\}$ are biholomorphic. Since a Hopf manifold of diagonal type is of Vaisman type and thus admit a lcK potential (see Sec. 2 for definition), applying the result (due to Ornea and Verbitsky) that the property of lcK potential is preserved under a complex analytic family over a sufficiently small parameter space, we can conclude that any Hopf manifold admits a lcK structure.

In this note, we briefly review basic results relating to lcK structures on Hopf manifolds with concise proofs to them (for the seek of completeness), leading to a proof for the theorem that any Hopf manifold admits a lcK structure. Note that the construction of lcK structures on Hopf manifolds is non-trivial and historic as seen in a series of papers [5, 2, 9, 11].

2. LOCALLY CONFORMALLY KÄHLER STRUCTURE

Definition 1. A *locally conformally Kähler structure* (or shortly *lcK structure*) on a differentiable manifold M is a Hermitian structure (h, J) on M with its associated fundamental form Ω satisfying $d\Omega = \theta \wedge \Omega$ for some closed 1-form θ (which is called *Lee form*). A lcK manifold M is of *Vaisman type* if its Lee form θ is parallel w.r.t. the Levi-Civita connection of h ; or equivalently, the Lee field $\xi = h^{-1}\theta$ is parallel.

Note that A lcK structure Ω is *locally conformally Kähler*, in the sense that there is a open covering $\{U_i\}$ of M such that $\Omega_i = e^{-\sigma_i} \Omega$ is Kähler form on U_i for some functions σ_i , that is, $d\Omega_i = 0$. The condition $d\Omega = \theta \wedge \Omega$ is equivalent to the existence of a global close 1-form θ such that $\theta|_{U_i} = d\sigma_i$.

Example 1. A complex projective space $\mathbb{C}P^1$ is a quotient manifold of $W = \mathbb{C}^2 - \{O\}$ by the action of \mathbb{C}^* ,

$$\phi_\lambda : (z_1, z_2) \rightarrow (\lambda z_1, \lambda z_2) \quad (\lambda \in \mathbb{C}^*).$$

On the other hand, a Hopf surface S is a quotient manifold of W by the action of \mathbb{Z}

$$\psi^k : (z_1, z_2) \rightarrow (\mu^k z_1, \mu^k z_2) \quad (k \in \mathbb{Z}),$$

for some $\mu \in \mathbb{C}^*$ ($|\mu| > 1$).

Since $\Gamma = \{\mu^k \mid k \in \mathbb{Z}\}$ is a discrete subgroup of \mathbb{C}^* and \mathbb{C}^*/Γ is a complex torus $T_{\mathbb{C}}^1$, we see S is a $T_{\mathbb{C}}^1$ bundle over $\mathbb{C}P^1$.

Consider a $(1, 1)$ -form on W ,

$$\omega = -\sqrt{-1} (dz_1 \wedge d\bar{z}_1 + dz_2 \wedge d\bar{z}_2),$$

and put

$$\Omega = \frac{1}{|z_1|^2 + |z_2|^2} \omega,$$

then Ω defines a real 2-form on S . Ω is not closed, but satisfies

$$d\Omega = \theta \wedge \Omega,$$

with

$$\theta = -\frac{1}{|z_1|^2 + |z_2|^2} (z_1 d\bar{z}_1 + z_2 d\bar{z}_2 + \bar{z}_1 dz_1 + \bar{z}_2 dz_2).$$

For $\psi(X) = \theta(JX)$, we have

$$\psi = \sqrt{-1} \frac{1}{|z_1|^2 + |z_2|^2} (z_1 d\bar{z}_1 + z_2 d\bar{z}_2 - \bar{z}_1 dz_1 - \bar{z}_2 dz_2),$$

and if we defines $\bar{\omega} = d\psi$, then

$$\bar{\omega} = \frac{-\sqrt{-1}}{(|z_1|^2 + |z_2|^2)^2} (|z_2|^2 dz_1 \wedge d\bar{z}_1 + |z_1|^2 dz_2 \wedge d\bar{z}_2 - \bar{z}_1 z_2 dz_1 \wedge d\bar{z}_2 - \bar{z}_2 z_1 dz_2 \wedge d\bar{z}_1)$$

which is so called Fubini-Study form. In the affine coordinates $z = \frac{z_2}{z_1}$, $\bar{\omega}$ is expressed as

$$\bar{\omega} = \frac{-\sqrt{-1}}{(1 + |z|^2)^2} dz \wedge d\bar{z}$$

We can express $\mathbb{C}P^1$ and Hopf surface S as homogeneous complex manifolds. Since $G = \mathrm{SL}_2(\mathbb{C})$ acts on $\mathbb{C}P^1$ transitively, we have

$$\mathbb{C}P^1 = G/B,$$

where B is a Borel subgroup of G :

$$B = \left\{ \begin{pmatrix} \alpha & \gamma \\ 0 & \beta \end{pmatrix} \middle| \alpha, \beta \in \mathbb{C}^*, \alpha\beta = 1, \gamma \in \mathbb{C} \right\}$$

Consider a subgroup B_μ of B

$$B_\mu = \left\{ \begin{pmatrix} \mu^t & \gamma \\ 0 & \mu^{-t} \end{pmatrix} \middle| \mu, \gamma \in \mathbb{C}, |\mu| > 1, t \in \mathbb{Z} \right\}.$$

Then we have $S = G/B_\mu$, and B/B_μ is a complex torus $T_{\mathbb{C}}^1$. S is a holomorphic fiber bundle over $\mathbb{C}P^1$ with fiber $T_{\mathbb{C}}^1$.

Note that this structure is generalized for a compact homogeneous lcK manifold: it has a structure of a holomorphic fiber bundle over a complex flag manifold with fiber $T_{\mathbb{C}}^1$ [8, 1].

It is known that lcK structure on M may be defined as a Kähler structure $\tilde{\omega}$ on the universal covering \tilde{M} on which the the fundamental group Γ acts homothetically; that is, for every $\gamma \in \Gamma$, $\gamma^* \tilde{\omega} = \rho(\gamma) \tilde{\omega}$ holds for some positive constant $\rho(\gamma)$.

Definition 2. Let M be a lcK manifold. Suppose that the universal covering \tilde{M} admits a *Kähler potential* Φ , which is a real positive function on \tilde{M} such that $\tilde{\omega} = -\sqrt{-1} \partial \bar{\partial} \Phi$ defines a Kähler structure on \tilde{M} . If the fundamental group Γ acts homothetically on Φ , then we call Φ a *lcK potential* for M . $\tilde{\omega}$ clearly defines a lcK structure on M .

Example 2. A diagonal Hopf surfaces $S = W/\Gamma_\mu$, where Γ_μ is generated by a contraction $\psi : (z_1, z_2) \rightarrow (\mu z_1, \mu z_2)$ with $|\mu| > 1$ on W , admits a lcK potential

$$\Phi(z_1, z_2) = |z_1|^2 + |z_2|^2.$$

We have a Kähler structure $\tilde{\omega} = -\sqrt{-1} (dz_1 \wedge d\bar{z}_1 + dz_2 \wedge d\bar{z}_2)$ on W for which $\tilde{\omega} = -\sqrt{-1} \partial \bar{\partial} \Phi$ holds.

3. STRUCTURE OF HOPF MANIFOLDS

We recall some results [7] on a structure theorem of the covering transformation groups of (generalized) Hopf manifolds and their diffeomorphism types. An analytic automorphism ϕ of W may be considered as an analytic automorphism of \mathbb{C}^n which fixes the origin (by Hartogs' theorem). ϕ is a *contraction* if the sequence $\{\phi^n\}$ converges uniformly to 0 on any compact neighborhood of the origin, or equivalently for any relatively compact neighborhoods U, V of the origin, there exists $N \in \mathbb{N}$ such that $\phi^n(\overline{V}) \subset U$ holds for any $n \geq N$.

Let $M = W/G$ be a Hopf manifold, where G is the covering transformation group of M consisting of analytic automorphisms over \mathbb{C}^n which fixes the origin $\mathbf{0}$. Then G acts on W properly discontinuously and fixed point free. We can express G as

$$G = H \rtimes Z,$$

where Z is an infinite cyclic group generated by a contraction ϕ on W , and H is a finite normal subgroup of G [7]. There exists $m \in \mathbb{N}$ such that for $Z' = \langle \phi^m \rangle$, $\phi^m = \phi^m$, $G' = H \times Z'$ (direct product) is a normal subgroup of finite index in G ; and thus $M = W/G$ has a finite normal covering $M' = W/G'$, which is diffeomorphic to $S^{2n-1}/H \times S^1$, where H is a finite unitary group acting freely on S^{2n-1} , as shown as a consequence of the deformation arguments we describe in the following.

Definition 3. A complex analytic family $\pi : \mathcal{M} \rightarrow U$ is a proper holomorphic surbmersion from a complex manifold \mathcal{M} into a domain $U \subset \mathbb{C}^k$. Two complex analytic families $\pi : \mathcal{M} \rightarrow U$ and $\pi' : \mathcal{M}' \rightarrow U$ over the same domain U are isomorphic if there is an holomorphic isomorphism $\psi : \mathcal{M} \rightarrow \mathcal{M}'$ commuting with π and π' . A complex analytic family $\pi : \mathcal{M} \rightarrow U$ is said to be complete at $p \in U$ if any complex analytic family $\pi' : \mathcal{M}' \rightarrow U'$ ($p \in U'$) with $\pi^{-1}(p) = \pi'^{-1}(p)$, there exists an open neighborhood $V \subset U'$ ($p \in V$) and a holomorphic map $q : V \rightarrow U$ such that the induced family $\pi' : q^{-1}\mathcal{M}' \rightarrow V$ by q is isomorphic to $\pi'|_V$; and it is said to be complete if it is complete at any point $p \in U$.

For an analytic automorphism ϕ over \mathbb{C}^n which fixes the origin $\mathbf{0}$, we denote the linear part of ϕ (i.e. the Jacobian matrix $d\phi(\mathbf{0})$) by $L(\phi)$. It is shown [7] that for the covering transformation group G of a Hopf manifold W/G , $L(G)$, as an automorphism group over W is properly discontinuously without fixed point free, defining a compact complex manifold $W/L(G)$ which is itself a Hopf manifold. This is more or less trivial for an infinite cyclic group Z ; and for a general case, it resorts to the Cartan's Uniqueness theorem.

We can construct a complex analytic family $\pi : \mathcal{M} \rightarrow \mathbb{C}$ which transforms $M = \pi^{-1}(1) = W/G$ into $\pi^{-1}(0) = W/L(G)$. In fact, let $T_t, (t \neq 0)$ be an analytic automorphism over W defined by

$$T_t(z_1, z_2, \dots, z_n) = (tz_1, tz_2, \dots, tz_n),$$

and set $g_t = T_t g T_t^{-1}$, $G(t) = \{g_t \mid g \in G\}$ and $G(0) = L(G)$. We define for $g \in G$ an analytic automorphism \tilde{g} over $W \times \mathbb{C}$ as

$$\tilde{g} : (z, t) \rightarrow (g_t(z), t),$$

where $z = (z_1, z_2, \dots, z_n) \in W$ and $t \in \mathbb{C}$. Then $\tilde{G} = \{\tilde{g} \mid g \in G\}$ is properly discontinuous and fixed-point-free as an analytic automorphism group over $W \times \mathbb{C}$ [7]. Let $\mathcal{M} = W \times \mathbb{C}/\tilde{G}$ and $\pi : \mathcal{M} \rightarrow \mathbb{C}$ be the canonical map induced from

the $Pr_2 : W \times \mathbb{C} \rightarrow \mathbb{C}$. Then $\pi : \mathcal{M} \rightarrow \mathbb{C}$ defines a complex analytic family with $\pi^{-1}(1) = W/G$ and $\pi^{-1}(0) = W/L(G)$.

We can further deform a Hopf manifold $W/L(G)$ to $W/L_0(G)$ with $L_0(G) = L_0(H) \times L_0(Z)$, where $L_0(Z)$ is the contraction generated by a diagonal matrices $d(\alpha_1, \alpha_2, \dots, \alpha_n)$ with eigenvalues of $\alpha_1, \alpha_2, \dots, \alpha_n$ ($0 < |\alpha_i| < 1, i = 1, 2, \dots, n$) of $L(\phi)$ and $L_0(H) \subset U(n)$. In fact, we can assume that $L(\phi)$ is of Jordan form $J(\alpha, n)$. Let $T_t, (t \neq 0)$ be an analytic automorphism over W defined by

$$T_t(z_1, z_2, \dots, z_n) = (t^{n-1}z_1, t^{n-2}z_2, \dots, tz_{n-1}, z_n),$$

and set $g_t = T_t g T_t^{-1}$, $L(G)(t) = \{g_t \mid g \in L(G)\}$. Note that $L(H)(t)$ may be assumed to stay in $U(n)$ [7]. As for the case of the complex analytic family $\{W/G(t)\}, \{W/LG(t)\}$ defines a complex analytic family $\pi' : \mathcal{M}' \rightarrow \mathbb{C}$ which transforms $\mathcal{M}' = \pi'^{-1}(1) = W/L(G)$ into $\pi'^{-1}(0) = W/L_0(G)$.

Example 3. The complex analytic family constructed above may be considered as a generalization of the following complex family of Hopf surfaces due to Kodaira [4]. For analytic automorphisms over $W = \mathbb{C}^2$,

$$g_t : (z_1, z_2) \rightarrow (\alpha z_1 + tz_2, \alpha z_2), 0 < |\alpha| < 1, t \in \mathbb{C},$$

we set $Z_t = \langle g_t \rangle$ and $M_t = W/Z_t$. Then $\{M_t\}$ defines a complex analytic family, where $M_t (t \neq 0)$ are all biholomorphic while M_0 is not biholomorphic to $M_t (t \neq 0)$.

Remark 1. The complete complex analytic families (so-called *Kuranishi families*) for Hopf surfaces was determined by J. Wehler [13]; for Hopf manifolds of diagonal type by C. Borcea [3]; and for any Hopf manifolds by A. Haefliger [6]. The result that any deformations of Hopf manifolds are Hopf manifolds (whose universal covering is $W = \mathbb{C}^n \setminus \{0\}$) is non-trivial and a consequence of the above results.

We recall that any Hopf manifold of diagonal type admits a Vaisman structure.

Example 4. A Hopf manifold of diagonal type with eigen values $\lambda_i, i = 1, 2, \dots, n$ admits a locally homogeneous Vaisman structure if all the eigen values λ_i have the same absolute value (which is less than 1) [8]. For simplicity, we consider here the case of Hopf surfaces. We construct a locally homogeneous l.c.K. manifolds $\Gamma \backslash G/H$ with a discrete subgroups Γ of $G = \mathbb{R} \times U(2)$ and $H = U(1)$. Let $\Gamma_{p,q} (p, q \neq 0)$ be a discrete subgroup of G :

$$\Gamma_{p,q} = \left\{ \left(k, \begin{pmatrix} e^{\sqrt{-1}pk} & 0 \\ 0 & e^{\sqrt{-1}qk} \end{pmatrix} \right) \in \mathbb{R} \times U(2) \mid k \in \mathbb{Z} \right\}.$$

Then $\Gamma_{p,q} \backslash G/H$ is biholomorphic to a Hopf surface $S_{p,q} = W/\Gamma_{\lambda_1, \lambda_2}$, where $\Gamma_{\lambda_1, \lambda_2}$ is the cyclic group of automorphisms on W generated by

$$\phi : (z_1, z_2) \longrightarrow (\lambda_1 z_1, \lambda_2 z_2)$$

with $\lambda_1 = e^{-r+\sqrt{-1}p}, \lambda_2 = e^{-r+\sqrt{-1}q}, r > 0$. In fact, we have a homogeneous complex structure J_r on G/H for which the biholomorphism $\Phi_r : G/H = \mathbb{R} \times S^3 \rightarrow W$ defined by $(t, z_1, z_2) \longrightarrow (e^{-rt}z_1, e^{-rt}z_2)$ induces a biholomorphism between $\Gamma_{p,q} \backslash G/H$ and $S_{p,q}$. Note that in case $p = q$, $S_{p,q}$ defines a homogeneous Vaisman Hopf surface $S^1 \times SU(2)$ [8]. For the standard homogeneous Sasaki structure ψ (see Example 1) on S^3 and $\theta = dt$ on \mathbb{R} , we have a locally homogeneous Vaisman structure Ω on $S_{p,q}$:

$$\Omega = -\theta \wedge \psi + d\psi.$$

For a Hopf manifold of diagonal type with eigen values $\lambda_i, i = 1, 2, \dots, n$, where $\lambda_i = e^{-r_i + \sqrt{-1}p_i}$ may have different absolute values $|\lambda_i| = e^{-r_i}, r_i > 0$, we consider the biholomorphism $\Phi : \mathbb{R} \times S^{2n-1} \rightarrow W$ defined by

$$\Phi : (t, z_1, z_2, \dots, z_n) \longrightarrow (e^{-r_1 t} z_1, e^{-r_2 t} z_2, \dots, e^{-r_n t} z_n).$$

For simplicity we consider here again the case of Hopf surfaces S_{λ_1, λ_2} . In place of the standard Sasaki structure ψ , we take a *weighted Sasaki form*:

$$\psi = \sqrt{-1} \frac{1}{r_1 |z_1|^2 + r_2 |z_2|^2} (z_1 d\bar{z}_1 + z_2 d\bar{z}_2 - \bar{z}_1 dz_1 - \bar{z}_2 dz_2).$$

The discrete group Γ_{p_1, p_2} ($p_1, p_2 \neq 0$) acts on $\mathbb{R} \times S^3$ preserving the weighted Sasaki structure ψ , and Φ induces a biholomorphism from $\Gamma_{p_1, p_2} \backslash \mathbb{R} \times S^{2n-1}$ to S_{λ_1, λ_2} . For the Lee form $\theta = dt$ on \mathbb{R} , we have a Vaisman structure Ω on S_{λ_1, λ_2} :

$$\Omega = -\theta \wedge \psi + d\psi.$$

4. MAIN THEOREM

We now give a proof for the main theorem, based on the above arguments.

Main Theorem. *Any Hopf manifold admits a locally conformally Kähler structure.*

Proof. We have seen that any Hopf manifold has a primary Hopf manifold as a finite normal covering; and thus it is sufficient to show the assertion for a primary Hopf manifold. We have just seen that a Hopf manifold of diagonal type has a Vaisman structure. As is known [10], a compact Vaisman manifold M admits a Kähler potential Φ on the universal covering \widetilde{M} such that the fundamental group Γ acts homothetically on Φ . As observed in [10], for a complex analytic family $\pi : \mathcal{M} \rightarrow U$, where U is an open neighborhood of the origin 0 in \mathbb{C}^n , with $M = \pi^{-1}(0)$, if M admits a Kähler potential Φ , then $M_t = \pi^{-1}(t)$ admit also a Kähler potential for a sufficiently small $t \in U$. This is seen as follows: $\tilde{\omega} = -\sqrt{-1} \partial \bar{\partial} \Phi$ defines a Kähler structure on \widetilde{M} . For $t \in U$, let ∂_t and $\bar{\partial}_t$ be partial derivatives with respect to the complex structures J_t on M_t and also on \widetilde{M}_t . For a sufficiently small $t \in U$, $\tilde{\omega} = -\sqrt{-1} \partial_t \bar{\partial}_t \Phi$ is positive definite for a coordinate neighborhood O_p of each point $p \in M$ and thus also for \widetilde{O}_p of \widetilde{M} . Since M is compact, we can take a finite number of such evenly covered O_p of M . Hence, $\tilde{\omega} = -\sqrt{-1} \partial_t \bar{\partial}_t \Phi$ defines a Kähler potential for a sufficiently small $t \in U$. We have seen that any primary Hopf manifold can be deformed through a complex analytic family into a linear primary Hopf manifold; and any linear primary Hopf manifold can be also deformed into a Hopf manifold of diagonal type. It should be emphasized that the complex analytic family $\pi : \mathcal{M} \rightarrow M$ we have constructed is such that $\pi^{-1}(t), t \neq 0$ are all biholomorphic; and thus for a given Hopf manifold M , M is biholomorphic to M_t for arbitrarily small $t \in U$. We have therefore completed the proof for the theorem. \square

REFERENCES

- [1] D.V. Alekseevsky, K. Hasegawa, and Y. Kamishima, *Homogeneous Sasaki and Vaisman manifolds of unimodular Lie groups*, Nagoya Math. J. **243** (2021), 83-96.
- [2] F. A. Belgun, *On the metric structure of non-Kähler complex surfaces*, Math. Ann. **317** (2000), 1-40.

- [3] C. Borcea, *Some remarks on deformations of Hopf manifolds*. Rev. Roum. Math. pures et appl. **26** (1981), 1287-1294.
- [4] K. Kodaira, *On the structure of compact complex analytic surfaces*, II, III, Amer. J. Math. **88** (1966), 682-721; 90 (1968), 55-83.
- [5] P. Gauduchon and L. Ornea, *Locally conformally Kähler metrics on Hopf surfaces*, Ann. Inst. Fourier. **48** (1998), 1107-1127.
- [6] A. Haefliger, *Deformations of transversely holomorphic flows on spheres and deformations of Hopf manifolds*, Compositio Math., **55** (1985), no. 2241-251.
- [7] K. Hasegawa, *Deformations and diffeomorphism types of generalized Hopf manifolds*, Illinois J. Math. **37** (1993), no. 7, 643-651.
- [8] K. Hasegawa. and Y. Kamishima, *Compact homogeneous locally conformally Kähler manifolds*, Osaka J. Math. **53**, 683-703 (2016).
- [9] Y. Kamishima and L. Ornea, *Geometric flow on compact locally conformally Kähler manifolds*, Tohoku Math. J. **57** (2005), no. 2, 201-221.
- [10] L. Ornea and M. Verbitsky, *Locally conformal Kähler manifolds with potential*, Math. Ann. **348** (2010), no. 1, 25-33.
- [11] L. Ornea and M. Verbitsky, *Locally conformal Kähler metrics obtained from pseudoconvex shells*, Proc. AMS, **144** (2016), no. 1, 325-335.
- [12] L. Ornea and M. Verbitsky, *Non-linear Hopf manifolds are locally conformally Kähler*, arXiv.2202.12398.
- [13] J. Wehler, *Versal deformation of Hopf surfaces*, J. für die reine und ang. Math. **328** (1981), 22-32.

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN

Email address: hasegawa@math.sci.osaka-u.ac.jp

FACULTY OF EDUCATION, NIIGATA UNIVERSITY, IKARASHI-NINO-CHO, NISHI-KU, NIIGATA 950-2181, JAPAN

Email address: hasegawa@ed.niigata-u.ac.jp