Perspective on complexity measures targetting read-once branching programs

Yaqiao Li^{*}, Pierre McKenzie[†]

May 22, 2023

Abstract

A model of computation for which reasonable yet still incomplete lower bounds are known is the read-once branching program. Here variants of complexity measures successful in the study of read-once branching programs are defined and studied. Some new or simpler proofs of known bounds are uncovered. Branching program resources and the new measures are compared extensively. The new variants are developed in part in the hope of tackling read-k branching programs for the tree evaluation problem [8]. Other computation problems are studied as well. In particular, a common view of a function studied by Gál [11] and a function studied by Bollig and Wegener [3] leads to the general combinatorics of blocking sets. Technical combinatorial results of independent interest are obtained. New leads towards further progress are discussed. An exponential lower bound for non-deterministic read-k branching programs for the GEN function [17] is also derived, independently from the new measures.

Contents

1	Intr	roduction	2
2	Mea	asures, branching programs and common functions	3
	2.1	Notation	3
	2.2	Measures from counting subfunctions: S and \widehat{S}	4
	2.3	Rectangles	4
	2.4	Measures from covering: C and \widehat{C}	5
	2.5	Measures from partitioning: P and \hat{P}	5
	2.6	max-min Communication complexity	6
	2.7	Branching programs	6
	2.8	Roster of functions	7
3	The	e measures vs read-once branching program sizes	9
	3.1	\widehat{P} is a lower bound for BP ₁	9
	3.2	S^* (almost) characterizes OBDD	10
	3.3	Relations and separations	11
	3.4	Lower bound measures for read- k BPs \ldots	15

*Concordia University, yaqiao.li@concordia.ca

[†]Université de Montréal, mckenzie@iro.umontreal.ca

4	\widehat{S} is small for TEP 4.1 S on TEP	17 19
5	The read-once non-deterministic BP lower bound of satisfiable Tseitin formulas via \hat{C} : a short proof	20
6	$ \widehat{S} \text{ and } \widehat{C} \text{ on Gál-type functions, and related blocking sets} $	 21 21 22 24 25
7	A read-k non-deterministic BP lower bound for GEN	27
8	Discussion and open problems	28
\mathbf{A}	The proof for Theorem 34	31

1 Introduction

Proving lower bounds on the resources needed to perform a computation often relies on confronting the computation model with complexity measures capturing its combinatorics. For example, partitions into rectangles and ranks of matrices offer viewpoints on the two-party communication complexity of a boolean function. As a rule of thumb, the more viewpoints are available, the more successful our study of the model is.

Polynomial size branching programs (bps) have long been known to capture logarithmic space. Much effort was devoted to their study and lower bounds on restricted bps abound (see [30]). But even in the context of as severe a restriction as *read-once*, or even *ordered read-once*, no lower bound method applies naturally to all hard functions in L, in NL, or even in P for that matter, despite the belief that P-hard functions require exponential size *unrestricted* bps. Read-once bps and their variants remain themselves to this day an object of study in connection with derandomization (e.g. [10, 6]) and proof complexity (e.g. [12, 26]).

In this work we first define two types of complexity measures inspired by known lower bounds for read-once bps. The first type (Section 2.2) derives from counting subfunctions of the function being computed. The second type (Sections 2.4, 2.5) exploits the variable partition model in communication complexity and leads to a framework tersely described as max-min communication complexity (Section 2.6).

The new measures are then compared with each other and with read-once deterministic and nondeterministic BP sizes (Figure 1 and Table 1). Several separations there follow from known (or adaptations of known) upper and lower bounds for the functions defined and grouped in Section 2.8. But a perspective on the combinatorics of read-limited bps emerges and in subsequent sections we pick up on some of the threads that arise.

In Section 4 we consider the tree evaluation problem, proposed in [8] as a candidate to separate L from larger classes. A read-once BP size lower bound for TEP is known [16]. In fact, our initial motivation for considering subfunctions counting lower bound measures for read-once bps was to use them to give an alternative proof for a read-once BP size lower bound for TEP, in hoping that

the alternative proof can be generalized to the read-k case. However, we show that a weak form of subfunctions counting (measure \hat{S}) will not suffice for this purpose. Our stronger measure S implies lower bounds on ordered read-once BP size in general (Corollary 19). We provide an incomplete report on S(TEP).

In Section 5, we use our weak covering measure (measure \widehat{C}) to give an alternative and simpler proof of a lower bound on the size of Tseitin formulas, used in [13] as a tool to obtain nondeterministic read-once BP lower bounds on the satisfiability problem for such formulas.

In Section 6 we cast two functions known to require large read-once bps, namely GAL defined in [11] from projective geometry and BW defined in [3] from representing numbers in a prime basis, as problem instances in a common regular $K_{2,2}$ -free bipartite graph. GAL and BW are known to have small CNF size and DNF size respectively, yet no function with small weight (i.e., small DNF and small CNF sizes *combined*) is known to require large read-once bps. We observe that weight and *ordered* read-once BP size are provably unrelated. Then we extend the GAL lower bounds to the bipartite graph setting by means of our measures \hat{S} and \hat{C} . Our analysis suggests the need to better understand the properties of blocking sets in \mathbb{F}_p^2 , raising several questions of a purely combinatorial nature. We make some observations in that direction (such as Lemma 46 describing an elegant property of intersecting points in \mathbb{F}_p^2).

In Section 7 we record an exponential nondeterministic read-k BP lower bound for the Pcomplete GEN function. This is obtained independently from our measures, by merely exhibiting a read-once reduction from the BRS function defined in [4].

In Section 8 we take stock and highlight several open questions arising from this work, some of which are of independent interest from the viewpoint of combinatorics alone.

2 Measures, branching programs and common functions

This section introduces notation, defines our measures, defines max-min complexity, recalls the definitions of branching programs and finally collects the definitions of several known functions whose complexities are at stake in the paper.

2.1 Notation

The set $\{1, \ldots, k\} \subseteq \mathbb{N}$ is denoted [k]. The operations \bigvee and \bigwedge denote the Boolean OR and AND on $\{0, 1\}$. The operations + and \cdot are the usual sum and product on \mathbb{N} or \mathbb{R} . When $f, g: D \to E$ and **op** is a binary operation on E, f **op** g is the function defined by (f op g)(x) = f(x) op g(x).

Let $f : \{0,1\}^n \to \mathbb{N}$. Let $A = \{i_1, \ldots, i_\ell\} \subseteq [n], \overline{A} = \{j_1, \ldots, j_{n-\ell}\} \subseteq [n], i_1 < \cdots < i_\ell$ and $j_1 < \cdots < j_{n-\ell}$. We define $\pi_A \in \text{Sym}(n)$ as the permutation

and write f_A for the $2^{\ell} \times 2^{(n-\ell)}$ matrix defined by

$$f_A[(x_{\pi_A(1)},\ldots,x_{\pi_A(\ell)}),(x_{\pi_A(\ell+1)},\ldots,x_{\pi_A(n)})] = f(x_1,\ldots,x_n).$$

Any $\alpha \in \{0,1\}^A$ then identifies a row of f_A and prescribes the subfunction $f_{A,\alpha}: \{0,1\}^{\overline{A}} \to \mathbb{N}$ of f given by $f_{A,\alpha}(y) = f_A(\alpha, y)$.

2.2 Measures from counting subfunctions: S and \widehat{S}

Definition 1. Let $f : \{0,1\}^n \to \mathbb{N}$ and $A \subseteq [n]$. Let $\operatorname{nrows}(f_A)$ denote the number of distinct rows of the matrix f_A and $\operatorname{mult}(f_A)$ the number of occurrences of a most frequent row. Define

$$S(f) = \max_{1 \le k \le n} \min_{A \subseteq [n], |A| = k} \operatorname{nrows}(f_A), \tag{1}$$

$$\widehat{S}(f) = \max_{1 \le k \le n} \min_{A \subseteq [n], |A| = k} \frac{2^{\kappa}}{\mathsf{mult}(f_A)}.$$
(2)

Consider the multiset of subfunctions $\{f_{A,\alpha} : \alpha \in \{0,1\}^A\}$. This is the same as the multiset of rows of the matrix f_A . Then, $\operatorname{nrows}(f_A)$ is the number of distinct subfunctions, and $\operatorname{mult}(f_A)$ is the *multiplicity* of this multiset. For example, the notion of an *m*-mixed Boolean function (see [20]) is captured via $\operatorname{mult}(f_A)$ as follows: f is *m*-mixed if $\operatorname{mult}(f_A) = 1$ for every $A \subseteq [n]$ of size |A| = m.

Lemma 2. If f is m-mixed, then $\widehat{S}(f) \ge 2^m$.

Remark 3. The measures S(f) and $\widehat{S}(f)$ can be generalized to non-Boolean functions $f: D^n \to R$, by leaving (1) unchanged and replacing in (2) the numerator 2^k with $|D|^k$.

A lower bound on OBDD size (Section 3.2) will involve a min-max "dual" to the S measure. For $\sigma \in \text{Sym}(n)$ and $1 \leq k \leq n$, let $\sigma([k])$ denote the set $\{\sigma(1), \ldots, \sigma(k)\}$ of size k. With this notation, $S(f) = \max_{1 \leq k \leq n} \min_{|A|=k,A \subseteq [n]} \operatorname{nrows}(f_A) = \max_{1 \leq k \leq n} \min_{\sigma \in \text{Sym}(n)} \operatorname{nrows}(f_{\sigma([k])})$.

Definition 4. Define the dual form of S, denoted by S^* , as

$$S^*(f) = \min_{\sigma \in \operatorname{Sym}(n)} \max_{1 \le k \le n} \operatorname{nrows}(f_{\sigma([k])}).$$
(3)

By definition, $S^*(f) \ge S(f)$.

2.3 Rectangles

The measures to be defined in Sections 2.4 and 2.5 build on the notion of a rectangle, standard for functions with a Boolean range but in need of a definition for functions with range \mathbb{N} .

Definition 5. A matrix $M \in \mathbb{N}^{k \times \ell}$ is a rectangle if $M(i, j) = g_i \cdot h_j$ for some $g \in \{0, 1\}^k$ and $h \in \mathbb{N}^{\ell}$.

Note the provenance of g in Definition 5 (drawing g from \mathbb{N}^k would make sense as well but we only have need for Boolean gs in our applications). Note further that when M is Boolean, a rectangle as defined is a rectangle in the standard sense [21, Def 1.12].

Definition 6. Let $f : \{0,1\}^n \to \mathbb{N}$, $A \subseteq [n]$ and $k \in [n]$. Then f is declared an A-rectangle if the matrix f_A is a rectangle, and declared a k-rectangle if it is an A-rectangle for some A of size k.

Example 7. Let n be an integer multiple of 4. Consider the equality function $g : \{0,1\}^n \to \{0,1\}$ defined by $g(x_1,\ldots,x_n) = 1$ if and only if $x_i = x_{n/2+i}$ for every $1 \le i \le n/2$. With $A = \{1,\ldots,n/2\}, g_A$ is simply the identity matrix, hence is not an A-rectangle. Consider $A' = \{1,\ldots,n/4,n/2+1,\ldots,3n/4\}$. It is easy to see that for every $a \in \{0,1\}^{A'}, b \in \{0,1\}^{\overline{A'}}, g(a,b) = g_1(a) \land g_2(b) = g_1(a) \cdot g_2(b)$, where g_1 and g_2 are both the equality function on n/2 bits. Hence g is an A'-rectangle. As |A'| = n/2, g is an n/2-rectangle.

2.4 Measures from covering: C and \widehat{C}

The measures defined in this section relate to nondeterministic communication complexity, hence apply solely to functions with a Boolean range.

Definition 8. Let $f : \{0,1\}^n \to \{0,1\}, A \subseteq [n] \text{ and } k \in [n].$ Define

$$C(f,A) = \min\left\{r : f \text{ is the } \bigvee \text{ of } r \text{ functions that are } A \text{-rectangles}\right\}$$
(4)

$$C(f,k) = \min\left\{r: f \text{ is the } \bigvee \text{ of } r \text{ functions that are } k\text{-rectangles}\right\}$$
(5)

$$C(f) = \max_{k \in [n]} \min_{A \subseteq [n], |A| = k} C(f, A)$$

$$\tag{6}$$

$$\widehat{C}(f) = \max_{k \in [n]} C(f, k).$$
(7)

We will refer to C(f) as the cover number of f and to $\widehat{C}(f) \leq C(f)$ as its weak cover number. Justification for naming C(f) in this way comes from observing that when $A \subseteq [n]$, C(f, A) is the measure $C^1(f_A)$ defined in [21, Def 2.1] as the minimal number of 1-monochromatic rectangles needed to cover the ones in the matrix f_A . Of course $\widehat{C}(f)$ is a new measure to be studied. We note that the appellation "cover number of f" in [21] applies in the fixed partition communication model and differs from our C(f) here.

2.5 Measures from partitioning: P and \hat{P}

Here we allow functions with a non-Boolean range. We say that $f: D \to \mathbb{N}$ and $g: D \to \mathbb{N}$ are orthogonal if $f \cdot g$ is the zero function.

Definition 9. Let $f : \{0,1\}^n \to \mathbb{N}, A \subseteq [n] \text{ and } k \in [n].$

$$P^+(f,A) = \min\{r: f \text{ is the integer sum of } r \text{ pairwise orthogonal } A \text{-rectangles}\}$$
(8)

 $P^{+}(f,k) = \min \{r : f \text{ is the integer sum of } r \text{ pairwise orthogonal } k \text{-rectangles} \}$ (9)

$$P^+(f) = \max_{k \in [n]} \min_{A \subseteq [n], |A| = k} P^+(f, A)$$
 (10)

$$\widehat{P^+}(f) = \max_{k \in [n]} P^+(f,k) \tag{11}$$

and further when $Im(f) \subseteq \{0, 1\} \subseteq \mathbb{N}$,

$$P(f) = \max_{k \in [n]} \min_{A \subseteq [n], |A| = k} (P^+(f, A) + P^+(\neg f, A))$$
(12)

$$\widehat{P}(f) = \max\left\{\widehat{P^+}(f), \widehat{P^+}(\neg f)\right\}.$$
(13)

Again here we will refer to $P^+(f)$ and to $\widehat{P^+}(f) \leq P^+(f)$ (and to P(f) and to $\widehat{P}(f) \leq P(f)$ when applicable) respectively as the partition number and the weak partition number of f. Justification for this naming comes from the partition number $C^D(M)$ [21, Def 2.1] of a Boolean matrix M, defined as the minimum number of monochromatic rectangles needed to partition M. When $\operatorname{Im}(f) \subseteq \{0,1\}, P^+(f,A)$ refers to partitioning the ones of the Boolean matrix f_A , so $P^+(f,A) + P^+(\neg f, A) = C^D(f_A)$.

Remark 10. The partition number P(f) is investigated under the name "rectangle complexity" in [24], where errors are allowed in representing f using rectangles.

2.6 max-min Communication complexity

Let M be a Boolean matrix. Let ccm(M) denote its deterministic communication complexity (i.e., measure D in [21, Def 1.2]) and nccm(M) denote the nondeterministic communication complexity of M (i.e. measure N^1 in [21, Def 2.3]).

Definition 11. Let $f : \{0,1\}^n \to \{0,1\}$. Define the max-min communication complexity of f, denoted by CC(f), to be

$$CC(f) = \max_{1 \le k \le n} \min_{A \subseteq [n], |A| = k} \mathsf{ccm}(f_A)$$
(14)

and the max-min nondeterministic communication complexity of f, denoted by NCC(f), to be

$$NCC(f) = \max_{1 \le k \le n} \min_{A \subseteq [n], |A| = k} \mathsf{nccm}(f_A).$$
(15)

Remark 12. The quantity $\min_{A \subseteq [n], |A| = n/2} \operatorname{ccm}(f_A)$ corresponds to the notion $D^{best}(f)$ in [21]. By definition,

$$NCC(f) = \log C(f).$$
(16)

Note that CC(f) seamlessly generalizes to non-Boolean functions $f : D^n \to \{0, 1\}$. Reassuringly CC(f) and NCC(f) in the Boolean case behave as we expect:

Lemma 13. Let $f: \{0,1\}^n \to \{0,1\}$. Then

$$NCC(f) \le CC(f) \le n/2 + 1, \tag{17}$$

$$\log P(f) \le \operatorname{CC}(f) \le O(\log^2 P(f)).$$
(18)

Proof. The left of (17) holds since nccm \leq ccm. The right of (17) holds because ccm $(f_A) \leq 1 + \min\{|A|, n - |A|\}$. As to (18), recall that $P^+(f, A) + P^+(\neg f, A) = C^D(f_A)$, which is at most $2^{\text{ccm}(f_A)}$ by [21, Prop 2.2], so applying max_k min_A yields $P(f) \leq 2^{\text{CC}(f)}$. Finally, any Boolean matrix of the form f_A satisfies ccm $(f_A) \leq c \cdot \log^2 P^+(f, A)$ for some constant c (see [22, Ex 1.1]). Hence ccm $(f_A) \leq c \cdot \log^2 (P^+(f, A) + P^+(\neg f, A))$, so that applying max_k min_A yields CC $(f) \leq c \cdot \log^2 P(f)$.

2.7 Branching programs

A deterministic (binary) branching program (BP for short) is a directed acyclic graph (DAG for short) with a unique source node and two sink nodes (one sink node labelled by 1, while the other sink node labelled by 0). Each non-sink node has outdegree 2, the node is labelled by a variable x_i for some $i \in [n]$, one of the two out-edges of the node is labelled by $x_i = 0$ and the other is labelled by $x_i = 1$. Every $x \in \{0,1\}^n$ defines a unique source-to-sink path in a BP. A BP computes a Boolean function $f : \{0,1\}^n \to \{0,1\}$ if the unique source-to-sink path for x ends at the sink with label f(x). The size of a BP is defined to be the number of nodes. Let BP(f) denote the smallest deterministic BP size computing f correctly. A BP is called syntactic *read-k* if on every source-to-sink path, each variable x_i appears at most k times. Let BP_k(f) denote the minimal size among all read-k BPs that compute f.

A nondeterministic BP (NBP for short) is a DAG with a unique source node and a unique sink node (the sink node is labelled by 1). Each non-sink node has outdegree at most 2. The

non-sink nodes have no labels. Each edge is either labelled by $x_i = 0$, or by $x_i = 1$, or has no label. An edge with no label will pass through all inputs. An NBP computes a Boolean function $f : \{0,1\}^n \to \{0,1\}$ if for every input x such that f(x) = 1, there exists at least one source-to-sink consistent path with x. The size of an NBP is defined to be the number of labelled edges. An NBP is called syntactic *read-k* if on every source-to-sink path, each variable x_i appears at most k times. Let NBP_k(f) denote the minimal size among all read-k NBPs that compute f.

An OBDD (ordered binary decision diagram, a.k.a., oblivious read-once branching program) is a read-once BP with the following property: there is a permutation π on [n], so that variables on every source-to-sink path follow the order π . Specifically, if x_j appears after x_i in a source-to-sink path, then $\pi^{-1}(j) > \pi^{-1}(i)$. Let OBDD(f) denote the minimal size among all OBDDs that compute f.

By definition, for every Boolean function f, $OBDD(f) \ge BP_1(f) \ge NBP_1(f)$.

2.8 Roster of functions

Here we define all the functions that will be used.

- The equality function $EQ_n : \{0,1\}^n \times \{0,1\}^n \to \{0,1\}$ is defined by $EQ_n(x,y) = 1$ if and only if x = y.
- The shifted equality function $\text{SEQ}_n : \{0,1\}^n \times \{0,1\}^n \times [n] \to \{0,1\}$, is defined by $\text{SEQ}_n(x,y,i) = 1$ if and only if $x_j = y_{t_j}$ for all $j = 1, \ldots, n$, where $t_j \in [n]$ and $t_j \equiv j + i 1 \mod n$. That is, $\text{SEQ}_n(x, y, i)$ computes the equality function on x and y, where bits of y are shifted by i 1 to the left.
- The parity function $\text{PARITY}_n : \{0,1\}^n \to \{0,1\}$ is defined by $\text{PARITY}_n(x) = 1$ if and only if the number of 1s in x is odd.
- The exact half clique function $\text{CLIQUE}_{n,n/2} : \{0,1\}^{n(n-1)/2} \to \{0,1\}$. Given an input $x \in \{0,1\}^{n(n-1)/2}$ as a graph on n vertices, $\text{CLIQUE}_{n,n/2}(x) = 1$ if and only if the graph x contains exactly a clique of size $\lceil n/2 \rceil$ and $\lfloor n/2 \rfloor$ isolated vertices.
- The pointer function $\pi_n : \{0,1\}^n \to \{0,1\}$. To define the pointer function we firstly define the $(\operatorname{OR} \circ \operatorname{AND})_{m^2} : \{0,1\}^{m^2} \to \{0,1\}$ as $(\operatorname{OR} \circ \operatorname{AND})_{m^2}(y) = \bigvee_{i=1}^m \bigwedge_{j=1}^m y_{(i-1)m+j}$. Now write $n = \log n \times \frac{n}{\log n}$ and partition the input *n*-bits into $\log n$ blocks $x = (x_1, \dots, x_{\log n})$ where each $x_i \in \{0,1\}^{n/\log n}$ for $i = 1, \dots, \log n$. Let $z_i = (\operatorname{OR} \circ \operatorname{AND})_{n/\log n}(x_i)$. Let $0 \le z \le n-1$ denote the unique integer represented by the $\log n$ bits $z_1 \cdots z_{\log n}$. Then, define $\pi_n(x) = x_{z+1}$, i.e., the (z+1)-th bit in x.
- The Tree Evaluation Problem (TEP). Let (TEP, 1): $[k] \to [k]$ be defined as (TEP, 1)(x) = x. For $h \ge 2$, let (TEP, h): $[k]^{n_h} \to [k]$ denote the Tree Evaluation Problem (see detail in [8]) of height h, where $n_h = 2n_{h-1} + k^2 = (2^{h-1} - 1)k^2 + 2^{h-1}$ denotes the input size, and $n_1 = 1$. Specifically, the input is a complete binary tree of height¹ h, in which every leaf is given an integer in [k], and every internal node is given a matrix in $[k]^{k \times k}$. One can naturally evaluate the binary tree in a bottom-up fashion and the output of the root node is defined as the output of (TEP, h).

¹Here the binary tree with a root and two leaves is deemed of height 2.

- The BRS function. Let $d \in \mathbb{N}$, $n = 2^d$. The BRS function $BRS_n : \{0,1\}^{2n} \times \{0,1\}^{2n} \to \{0,1\}$ is defined in [4] as follows. For every $x \in \{0,1\}^{2n}$, we write it as $x = (\ldots, x_{a1}, x_{a2}, \ldots)$, where $a \in \{0,1\}^d$ and $x_{a1}, x_{a2} \in \{0,1\}$. For every $y \in \{0,1\}^{2n}$, we write it as $y = (\ldots, x_{b1}, y_{b2}, \ldots)$, where $b \in \{0,1\}^d$ and $y_{b1}, y_{b2} \in \{0,1\}$. Define $BRS_n(x,y) = 1$ if and only if $\sum_{a,b \in \{0,1\}^d} (-1)^{\langle a,b \rangle} (x_{a1} + x_{a2})(y_{b1} + y_{b2}) \equiv 0 \mod 3$, where $\langle a, b \rangle$ is the usual inner product mod 2, but the rest operations are performed mod 3. More details of the definition is in [4]. The BRS function is called as bilinear Sylvester function in [30, Theorem 10.3.10].
- The indirect storage access function $\text{ISA}_n : [n] \times \{0,1\}^n \to \{0,1\}$ defined as follows, for $i \in [n]$ and $x \in \{0,1\}^n$, $\text{ISA}_n(i,x) = x_p$ where p is the integer represented by the length $\log n$ binary string $x_i x_{i+1} \cdots x_{i+\log n-1}$ where the addition of indices is mod n.
- The iterated NAND_n function. Define NAND₂ : $\{0,1\} \times \{0,1\} \rightarrow \{0,1\}$ by NAND₂ $(x,y) = \neg x \vee \neg y$. For $n = 2^h$, the iterated function NAND_n : $\{0,1\}^n \rightarrow \{0,1\}$ is computed by the balanced read-once formula of height h in which every gate is NAND₂.
- The satisfiable Tseitin formulas. Let G = (V, E) be a graph, |V| = n, |E| = m, and $c : V \to \{0, 1\}$ be a labelling function. The pair (G, c) defines a Tseitin formula as a Boolean function on $\{0, 1\}^E$, denoted by $TS_{G,c}$,

$$\mathrm{TS}_{G,c}: \{0,1\}^E \to \{0,1\}, \quad x = (\cdots, x_e, \cdots) \mapsto \bigwedge_{v \in V} \left(\left(\sum_{e \text{ is incident to } v} x_e \right) \equiv c(v) \mod 2 \right).$$
(19)

The Tseitin formula $TS_{G,c}$ is said to be *satisfiable* if the Boolean function $TS_{G,c}$ is not identically 0.

- The generalized Gál's function and Bollig-Wegener function. Let $G = (A \cup B, E)$ be a bipartite graph where A and B are the two parts of vertices. For $S \subseteq A$, let $Nb(S) \subseteq B$ denote the set of neighbors of S in B. The generalized Gál's function on G, denoted by $GAL_G : \{0,1\}^A \to \{0,1\}$, is defined as $GAL_G(x) = 1$ if and only if Nb(x) = B. The Bollig-Wegener function on G, denoted by $BW_G : \{0,1\}^A \times \{0,1\}^B \to \{0,1\}$, is defined as $BW_G(x,y) = 1$ if and only if $Nb(x) \cap y \neq \emptyset$.
- The GEN problem. For $m \ge 2$, let $n = \{(i, j) : i, j \in [m-1], i \le j\}$, then $n = \frac{m(m-1)}{2}$. Every $X \in [m]^n$ defines an upper triangular matrix which can be thought of as a (commutative) multiplication table: $X_{ij} = k$ means i * j = j * i = k, where $1 \le i \le j \le m-1$ and $1 \le k \le m$. Define GEN_n : $[m]^n \to \{0, 1\}$ as follows: GEN_n(X) = 1 if and only if $m \in \langle 1 \rangle$, where $\langle 1 \rangle$ denotes the set of elements generated starting from 1 * 1 and using the multiplication table X. Note that if interpreted as a Boolean function, then GEN_n is defined on the domain $\{0, 1\}^{n \log m} = \{0, 1\}^{\Theta(n \log n)}$.

Lastly, for every Boolean function $f : \{0,1\}^n \to \{0,1\}$, let $\neg f$ denote the negation of f, i.e., $(\neg f)(x) = 1$ if and only if f(x) = 0.

3 The measures vs read-once branching program sizes

It is known that \widehat{S} and \widehat{C} are lower bounds for BP₁ and NBP₁, as proved in [25]² and [4] respectively. The argument in [25], [4, Corollary 2] in fact proved NBP₁(f) $\geq \widehat{C}(f, n/2)$, but it is easy to see that NBP₁(f) $\geq \widehat{C}(f)$ holds.

Proposition 14 ([25, 4]). Let $f : \{0, 1\}^n \to \{0, 1\}$. Then, $BP_1(f) \ge \widehat{S}(f)$, $NBP_1(f) \ge \widehat{C}(f)$.

In this section we will prove more lower bounds, study relations among measures and provide examples (many are derived from the existing literature) that separate the measures from each other and from read-once BP sizes in all possible cases. An interesting feature is that communication complexity often plays a role.

3.1 \widehat{P} is a lower bound for BP₁

Theorem 15. For every integer-valued function $f : \{0,1\}^n \to \mathbb{N}$, $BP_1(f) \ge \widehat{P^+}(f)$. When $f : \{0,1\}^n \to \{0,1\}$ is a Boolean function, $BP_1(f) \ge \widehat{P}(f)$.

Proof. We adapt the proof strategy and ideas introduced in [25, 4]. Let B be a deterministic read-once BP computing f. We think of B = (V(B), E(B)) as a directed acyclic graph (DAG). Given a non-sink node $v \in V(B)$, let x(v) denote the variable queried at node v. Given two nodes $u, v \in V(B)$, let $u \leq v$ denote the relation that either there is a directed path from u to v or u = v. Define

$$X(u,v) = \{x(w) : u \le w \le v\}.$$

Let $s, t_0, t_1 \in V(B)$ denote the source node and the two sink nodes (where t_0 and t_1 are the sink nodes with output 0 and 1, respectively). Note that $X(s, t_0) = X(s, t_1)$. Without loss of generality we assume $X(s, t_0) = X(s, t_1) = [n]$. Fix a parameter $1 \leq k \leq n$. Consider a mapping

$$\varphi_k: \{0,1\}^n \to E(B), \quad x \mapsto e = (u,v),$$

where e = (u, v) is the edge in the computation path of x in B that satisfies

$$|X(s,u)| \le k < |X(s,v)|.$$

Observe that such edge (u, v) is unique, hence φ_k is well-defined. As usual, we think of a subset $A \subseteq [n]$ as the corresponding subset of variables. Choose a subset $A_e \subseteq [n]$ such that

$$X(s,u) \subseteq A_e \subseteq X(s,v), \quad |A_e| = k.$$

Given two non-sink nodes $u, v \in V(B)$, let $f_{u,v} : X(u,v) \setminus \{x(v)\} \to \{0,1\}$ denote the function computed by the sub-BP in B with u to be the source node, and $f_{u,v}(y) = 1$ if and only if the input y starts at u and reaches v. Note that $f_{u,v}$ is well-defined because the BP B is read-once. Given a non-sink node $v \in V(B)$, let $f_v : X(v,t_0) \cup X(v,t_1) \to \{0,1\}$ denote the function computed by the sub-BP in B with v to be the source node. The definitions of $f_{u,v}$ and f_v can be extended to be defined on larger domains by ignoring the irrelevant variables. Let $\mu(e) \in \{0,1\}$ denote the value

²In fact, [25] proved a stronger lower bound than \widehat{S} , but we focus on \widehat{S} in this paper for its naturalness.

marked on edge e, i.e., the edge e = (u, v) tests whether $x(u) = \mu(e)$. Let $\operatorname{Im}(\varphi_k)$ denote the image of φ_k . With these notations, we define a function f_e for every $e = (u, v) \in \operatorname{Im}(\varphi_k)$,

$$f_e: \{0,1\}^n \to \mathbb{N}, \quad x \mapsto f_{s,u}(x|_{A_e \setminus \{x(u)\}}) \cdot \left(x(u) = = \mu(e)\right) \cdot f_v(x|_{\overline{A_e}}). \tag{20}$$

In other word, the function f_e checks whether an input reaches the node u and passes through the edge e = (u, v) and is then computed by f_v . Hence,

$$f_e(x) = \begin{cases} f(x), & x \in \varphi_k^{-1}(e), \\ 0, & x \notin \varphi_k^{-1}(e). \end{cases}$$
(21)

Equation (21) implies that

$$f(x) = \sum_{e \in \operatorname{Im}(\varphi_k)} f_e \quad \text{and} \quad \langle f_e, f_{e'} \rangle = 0 \ \forall \ e, e' \in \operatorname{Im}(\varphi_k), e \neq e'.$$
(22)

If we denote $f_{s,u,v}(x|_{A_e}) = f_{s,u}(x|_{A_e-x(u)}) \cdot (x(u) == \mu(e))$, then,

$$f_e = f_{s,u,v} \cdot f_v,$$

where $f_{s,u,v}$ is a Boolean-valued function defined on variables in A_e and f_v is an (non-negative) integer-valued function defined on variables in $\overline{A_e}$. By definition in Section 2.5, f_e is an k-rectangle since $|A_e| = k$. Hence, by (22) and Definition 9,

$$|\operatorname{Im}(\varphi_k)| \ge \widehat{P^+}(f,k).$$

Since every node in *B* has degree at most 2, $|V(B)| \ge |\operatorname{Im}(\varphi_k)|/2$. By observing that the edges in $\operatorname{Im}(\varphi_k)$ do not lie in the same path, the factor 2 can be removed (see [25, Theorem 2.4]), we omit this technical detail. As $1 \le k \le n$ can be arbitrary, $\operatorname{BP}_1(f) \ge \max_k \widehat{P^+}(f,k) = \widehat{P^+}(f)$. The lower bound for a Boolean function f follows by noting that $\operatorname{BP}_1(f) = \operatorname{BP}_1(\neg f)$.

Remark 16. The proofs for Proposition 14 in [25, 4] both follow a similar strategy as shown above. Adopting the method in [4] one could also generalize the measure $\widehat{P^+}(f)$ appropriately so that it becomes a lower bound for $BP_k(f)$, see Section 3.4.

3.2 S^* (almost) characterizes OBDD

Theorem 17. Let $f : \{0,1\}^n \to \{0,1\}$. Then, $S^*(f) \leq \text{OBDD}(f) \leq 1 + n \cdot S^*(f)$.

Proof. The claim is true if f is a constant function, in which case $OBDD(f) = S^*(f) = 1$. Assume now f is not a constant function.

The lower bound: By changing the names of the variables if necessary, assume that the size OBDD(f) is achieved with respect to the order of variables x_1, \ldots, x_n . Let $B_i = \{x_1, \ldots, x_i\}$ for $1 \leq i \leq n$. For $A \subseteq \{x_1, \ldots, x_n\}$, let R(A) = the set of distinct rows of the matrix f_A , i.e., it is the set of all distinct subfunctions $f_{A,\alpha}$ where $\alpha \in \{0, 1\}^A$. Consider $R(B_i)$. Let $U(B_i) = \{g \in R(B_i) : g|_{x_{i+1}=0} \neq g|_{x_{i+1}=1}\}$, i.e., it is the subset of subfunctions in $R(B_i)$ that essentially depend on x_{i+1} . Let $W(B_i) = R(B_i) \setminus U(B_i)$, i.e., the subset of subfunctions in $R(B_i)$ that do not essentially depend on x_{i+1} . Let $u(B_i) = |U(B_i)|$ and $w(B_i) = |W(B_i)|$. So,

$$S(f_{B_i}) = |R(B_i)| = u(B_i) + w(B_i).$$
(23)

By [30, Theorem 3.1.4],

OBDD
$$(f) = 1 + \sum_{i=1}^{n} u(B_i).$$
 (24)

We claim $w(B_i) \leq S(f_{B_{i+1}})$. Since we assume f is not a constant function, $S(f_{B_n}) = u(B_n) = 2$. This and (23) together imply that $S(f_{B_i}) \leq \sum_{j=i}^n u(B_j) \leq \text{OBDD}(f)$ holds for every $1 \leq i \leq n$. Hence, $S^*(f) = \min_{\sigma} \max_{1 \leq i \leq n} S(f_{\sigma([i])}) \leq \max_{1 \leq i \leq n} S(f_{B_i}) \leq \text{OBDD}(f)$.

We proceed to show the claim. Consider a mapping

$$\phi: W(B_i) \to R(B_{i+1}), \quad g \mapsto \phi(g)$$

where $\phi(g) = g|_{x_{i+1}=0}$. It suffices to show ϕ is injective. Indeed, if $g, g' \in W(B_i)$ and $g \neq g'$, then there exists $(x_{i+1}, x_{i+2}, \ldots, x_n)$ such that $g(x_{i+1}, x_{i+2}, \ldots, x_n) \neq g'(x_{i+1}, x_{i+2}, \ldots, x_n)$. Since $g, g' \in W(B_i)$, one has

$$\phi(g)(x_{i+2},\dots,x_n) = g(0,x_{i+2},\dots,x_n) = g(x_{i+1},x_{i+2},\dots,x_n)$$

$$\neq g'(x_{i+1},x_{i+2},\dots,x_n)$$

$$= g'(0,x_{i+2},\dots,x_n) = \phi(g')(x_{i+2},\dots,x_n).$$

That is, $\phi(g) \neq \phi(g')$ as desired.

The upper bound: suppose $S^*(f)$ is minimized with respect to the order of variables x_1, \ldots, x_n , denote this order by π . Let π -OBDD(f) denote the least OBDD size of f when variables are queried with respect to the order π . Use the notation in the lower bound proof, by (23) and (24),

$$OBDD(f) \le \pi - OBDD(f) = 1 + \sum_{i=1}^{n} S(f_{B_i}) \le 1 + n \max_{1 \le i \le n} S(f_{B_i}) = 1 + n \max_{1 \le i \le n} S(f_{\pi([i])}) = 1 + n \cdot S^*(f),$$

where the last step follows by the assumption that $S^*(f)$ is minimized at π .

Remark 18. The measure $S^*(f)$ as a lower bound for OBDD(f) has essentially been applied in the literature, e.g., [23, Proposition 1] as well as in [30] etc, however, to the best knowledge of the authors there was no formal proof for this fact.

Corollary 19. $OBDD(f) \ge S(f)$.

3.3 Relations and separations

In this section we focus exclusively on Boolean functions $f: \{0,1\}^n \to \{0,1\}$.

Theorem 20. Let $f : \{0,1\}^n \to \{0,1\}$ be a Boolean function.

(1)
$$\widehat{S}(f) \leq S(f), \ \widehat{C}(f) \leq C(f), \ \widehat{P}(f) \leq P(f), \ \widehat{C}(f) \leq \widehat{P}(f), \ C(f) \leq P(f),$$

(2)
$$P(f)/2 \le S(f) \le 2^{C(f)}$$
,

(3)
$$S(f) = O(2^n/n), P(f) \le 2^{n/2+1} = O(2^{n/2}).$$

- (4) for every $\delta > 1$, most f satisfies $\widehat{S}(f) \ge 2^n/n^{\delta}$,
- (5) most f satisfies $\widehat{C}(f) \ge \Omega(2^{n/2}/\log n)$.

Proof. (1). These follow directly from definitions.

(2). It suffices to show that for every subset $A \subseteq [n]$, $P(f_A)/2 \leq S(f_A) \leq 2^{C(f_A)}$. The first inequality follows from $P^+(f_A) \leq S(f_A)$ and $P^+(\neg f_A) \leq S(\neg f_A)$ and noting that $S(f_A) = S(\neg f_A)$. For the second, by the definition of C(f, A), suppose C(f, A) = r, then

$$f_A = g_1 \vee \cdots \vee g_r,$$

where each matrix g_i has rank 1. For every $1 \le i \le r$, let X_i denote the unique nonzero row in the matrix g_i . Then, the above equation implies that each row of the matrix f_A can be written in the form $\bigvee_{i \in S} X_i$ for some subset $S \subseteq [r]$. Hence, f_A contains at most $2^r = 2^{C(f,A)}$ distinct rows. (3). By definition,

$$S(f) = \max_{1 \le k \le n} \min_{A \subseteq [n], |A| = k} \operatorname{nrows}(f_A) \le \max_{1 \le k \le n} \min\{2^k, 2^{2^{n-k}}\} = O(2^n/n).$$

By (17) and (18), $P(f) \le 2^{CC(f)} \le 2^{n/2+1}$.

(4). Observe that for a random Boolean matrix $M_{a \times b} \in \{0, 1\}^{a \times b}$ of a rows and b columns, Pr[M contains at least 2 identical rows] $\leq \frac{a^2}{2^b}$. Now consider $\widehat{S}(f)$. Consider a random function $f: \{0,1\}^n \to \{0,1\}$. Then, for every $A \subseteq [n]$ of size |A| = k, the matrix f_A is a random Boolean matrix of $a = 2^k$ rows and $b = 2^{n-k}$ columns. Since there are $\binom{n}{k}$ ways of choosing subsets $A \subseteq [n]$,

Pr[There exists A of size |A| = k, s.t. f_A contains at least 2 identical rows] $\leq \binom{n}{k} \cdot \frac{2^{2k}}{2^{2^{n-k}}}$.

Let $\delta > 1$. Choose $k = n - \delta \log n$. Then, the above probability is upper bounded by $2^{(2+\delta)n}/2^{n^{\delta}}$. Hence,

$$\begin{aligned} &\Pr\left[\max_{|A|=n-\delta\log n} \mathsf{mult}(f_A) = 1\right] \\ &= \Pr[\text{For every } A \text{ of size } |A| = n - \delta\log n, \text{ rows in } f_A \text{ are all distinct}] \\ &\geq 1 - \frac{2^{(2+\delta)n}}{2^{n^{\delta}}}. \end{aligned}$$

Equivalently, for a random function f, with probability at least $1 - 2^{(2+\delta)n}/2^{n^{\delta}}$, $\widehat{S}(f) \ge 2^{n-\delta \log n}$.

(5). Let $r = \widehat{C}(f)$, then $\widehat{C}(f, n/2) \leq r$ (if n is odd one chooses (n+1)/2). By definition of $\widehat{C}(f, n/2),$

$$f = f_1 \vee \cdots \vee f_r$$

where each f_i is an n/2-rectangle. That is, $f_i = f_{i1} \wedge f_{i2}$ where $f_{i1} : \{0,1\}^{n/2} \to \{0,1\}$ and $f_{i2}: \{0,1\}^{n/2} \to \{0,1\}$ are two Boolean functions each is defined on n/2 bits. This shows the formula size of f is at most $O(r \cdot 2^{n/2})$. It is well known that most Boolean functions have formula size $\Omega(2^n/\log n)$ (see, e.g., [28]). Hence, $r \ge \Omega(2^{n/2}/\log n)$.

Corollary 21. Let $f: \{0,1\}^n \to \{0,1\}$ be a Boolean function. Then,

$$\log \log S(f) \le \log C(f) = \operatorname{NCC}(f) \le \log P(f) \le \operatorname{CC}(f) \le 1 + \log S(f).$$

Proof. Inequalities except the last one follow from Theorem 20 and (18). The last inequality can be easily proved by the definition of communication complexity (see [21]). We omit the detail. \Box We now show the inequality $C(f) \ge \log S(f)$ is tight up to a polynomial (the rest inequalities in Theorem 20 are easily seen to be tight).

Viewing SEQ_n as a Boolean function on $2n + \log n$ bits, the max-min deterministic and nondeterministic communication complexity of SEQ_n are given below.

Proposition 22. $CC(SEQ_n) = CC(\neg SEQ_n) = \Theta(n), NCC(SEQ_n) = \Theta(n), NCC(\neg SEQ_n) \le 1 + 2 \log n = O(\log n).$

Proof. [21, Example 7.9] shows that $\min_{A \subseteq [n], |A| = n + \frac{\log n}{2}} \operatorname{ccm}((\operatorname{SEQ}_n)_A) = \Theta(n)$, hence $\operatorname{CC}(\operatorname{SEQ}_n) = \Theta(n)$. The proof for [21, Example 7.9] uses the fact that $\operatorname{ccm}(\operatorname{EQ}_n) \ge n$. Since $\operatorname{nccm}(\operatorname{EQ}_n) \ge n$ also holds (see [21]), it is not hard to see that the proof for [21, Example 7.9] can be adapted to show $\operatorname{NCC}(\operatorname{SEQ}_n) = \Theta(n)$. We omit the detail. Since $\operatorname{nccm}(\neg \operatorname{EQ}_n) \le 1 + \log n$ (see [21]), then $\operatorname{NCC}(\neg \operatorname{SEQ}_n) \le 1 + 2\log n$.

In viewing of Proposition 14 and Theorem 15, it is natural to ask whether S, P and C are also lower bounds for BP₁ or NBP₁. Perhaps not surprisingly, none of them is.

 $\textbf{Proposition 23. } \mathrm{BP}_1(\mathrm{SEQ}_n) = O(n^2), \ S(\mathrm{SEQ}_n) = 2^{\Theta(n)}, \ P(\mathrm{SEQ}_n) \geq C(\mathrm{SEQ}_n) = 2^{\Theta(n)}.$

Proof. Consider a BP that firstly reads the index i, using $O(2^{\log n}) = O(n)$ number of nodes, then for each index i, it continues to compute an equality function using O(n) nodes. This BP is read-once. Hence, $BP_1(SEQ_n) = O(n^2)$. Lower bounds for S and C follow from Corollary 21 and Proposition 22.

Proposition 24. $S(\neg SEQ_n) = 2^{\Theta(n)}, P(\neg SEQ_n) = 2^{\Omega(\sqrt{n})}, C(\neg SEQ_n) = O(n^2).$

Proof. Apply Corollary 21 and Proposition 22 and Equation (18) and (16).

Theorem 20 shows that $\widehat{S}(f)$, $\widehat{P}(f)$ and $\widehat{C}(f)$ are good lower bounds for BP₁(f) and NBP₁(f), respectively, in the following sense: since BP₁(f) $\leq O(2^n/n)$ and NBP₁(f) $\leq O(2^{n/2})$ (see, e.g., [1, Lemma 3.6]), property (4) and (5) of Theorem 20 show that \widehat{S} , \widehat{P} and \widehat{C} provide almost tight exponential lower bounds for BP₁(f) and NBP₁(f), respectively, for most functions. Similarly, S(f) is a good lower bound for OBDD(f). Nonetheless, they do not characterize the corresponding BP sizes. For example, BP₁(PARITY_n) = $\Theta(n)$, but $S(\text{PARITY}_n) = 2$ which implies that $\widehat{S}(\text{PARITY}_n)$, $\widehat{P}(\text{PARITY}_n)$ and $\widehat{C}(\text{PARITY}_n)$ all equal to 2 via Theorem 20. In fact, the gaps can be exponential.

Proposition 25 ([25, 4]).
$$\widehat{C}(\text{CLIQUE}_{n,n/2}) = 2^{\Theta(n)}, \ \widehat{S}(\text{CLIQUE}_{n,n/2}) \le 2^{49} = O(1).$$

Proof. The lower bound for \widehat{C} is shown in [4, Theorem 3]. [25] mentioned that "the uniform weighting provably cannot work", using our notation, this is equivalent to say that $\widehat{S}(\text{CLIQUE}_{n,n/2}) = n^{O(1)}$. We proceed to give an explicit bound.

Let N = n(n-1)/2. For every $1 \le k \le N$, we choose a subset $A \subseteq [N]$ so that $\mathsf{mult}((\mathsf{CLIQUE}_{n,n/2})_A)$ is large, as follows. Let p_k be the largest integer so that $T_k = p_k(p_k - 1)/2 \le k$. Choose p_k vertices from n vertices, and choose all the T_k edges on these p_k vertices to be in A, and choose the remaining $k - T_k$ edges, if any, arbitrarily. By our choice, $k - T_k \le p_k \le 3\sqrt{k}$.

Let $(\alpha, \beta) \in \{0, 1\}^A \times \{0, 1\}^{\overline{A}} = \{0, 1\}^N$, let α_{p_k} denote the subgraph on the p_k vertices given by α . We say the subgraph α_{p_k} is a *q*-clique if it contains exactly a clique of size *q* and isolated $p_k - q$ vertices, where $0 \le q \le p_k$. Observe that $\text{CLIQUE}_{n,n/2}(\alpha, \beta) = 1$ implies that α_{p_k} must be

a q-clique for some $0 \le q \le p_k$. Alternatively, if for every $0 \le q \le p_k$, the subgraph α_{p_k} is not a q-clique, then the subfunction (CLIQUE_{n,n/2})_{A, $\alpha} = 0$. Since the number of q-cliques on p_k vertices is $\binom{p_k}{q}$, we get the number of choices of α such that (CLIQUE_{n,n/2})_{A, $\alpha} = 0$ is at least}}

$$\left(2^{T_k} - \sum_{q=0}^{p_k} \binom{p_k}{q}\right) \cdot 2^{k-T_k} = (2^{T_k} - 2^{p_k}) \cdot 2^{k-T_k} \ge 2^k - 2^{6\sqrt{k}}.$$

This shows $\mathsf{mult}((\mathrm{CLIQUE}_{n,n/2})_A) \geq 2^k - 2^{6\sqrt{k}}$. Hence, $2^k/\mathsf{mult}((\mathrm{CLIQUE}_{n,n/2})_A) \leq 2$ when $k \geq 49$.

The proof for $\widehat{S}(\text{CLIQUE}_{n,n/2}) = O(1)$ in Proposition 25 lies in the fact that there are many constant 0 subfunctions in $\text{CLIQUE}_{n,n/2}$. This may seem too special. In Section 4 we provide another separation example of BP₁ from \widehat{S} via the function TEP. Note that obviously $\widehat{S}(f) = \widehat{S}(\neg f)$ and $\widehat{P}(f) = \widehat{P}(\neg f)$ for every Boolean function f, but this is not true for \widehat{C} .

Proposition 26. $\widehat{C}(\neg \operatorname{CLIQUE}_{n,n/2}) \le \operatorname{NBP}_1(\neg \operatorname{CLIQUE}_{n,n/2}) = O(n^4), \operatorname{NCC}(\neg \operatorname{CLIQUE}_{n,n/2}) = O(\log n), \ \widehat{P}(\neg \operatorname{CLIQUE}_{n,n/2}) = 2^{\Theta(n)}.$

Proof. The first upper bound follows from Proposition 14 and $\text{NBP}_1(\neg \text{CLIQUE}_{n,n/2}) = O(n^4)$ showed in [4, Theorem 5].

For the second upper bound, for any subset $A \subseteq [n(n-1)/2]$, consider $\operatorname{nccm}(\neg \operatorname{CLIQUE}_{n,n/2})_A$. By the proof of [4, Theorem 5], a graph G is not an exact $\lceil n/2 \rceil$ -clique if and only if at least one of the following is true:

- (a) there are two edges (v_1, v_2) and (v_3, v_4) in G such that $v_1 \neq v_3$ and (v_1, v_3) is not an edge;
- (b) there exists at least one vertex v whose degree differs from both 0 and $\lceil n/2 \rceil$;
- (c) G is empty.

Alice and Bob can verify whether G is empty using O(1) communication. For case (a), the prover can give the four vertices v_1, v_2, v_3, v_4 as a proof, which has length $O(\log n)$, and Alice and Bob need to use only O(1) communication to verify it. For case (b), the prover can give the name of v as a proof, which has length $O(\log n)$, and Alice and Bob need to use $O(\log n)$ communication to verify it. Hence, $\operatorname{nccm}(\neg \operatorname{CLIQUE}_{n,n/2})_A = O(\log n)$, independent of the choice of A. Hence, $\operatorname{NCC}(\neg \operatorname{CLIQUE}_{n,n/2}) = O(\log n)$.

Lastly,
$$\widehat{P}(\neg \text{CLIQUE}_{n,n/2}) = \widehat{P}(\text{CLIQUE}_{n,n/2}) \ge \widehat{C}(\text{CLIQUE}_{n,n/2}) = 2^{\Theta(n)}.$$

Proposition 27.
$$\widehat{S}(\pi_n) \ge 2^{\sqrt{n/\log n} - 1}, \ P(\pi_n) \le 16n, \ CC(\pi_n) \le \log n + 4, \ NBP_1(\pi_n) = O(n).$$

Proof. [18, Theorem 16.8] showed that π_n is $(\sqrt{n/\log n} - 1)$ -mixed, hence $\widehat{S}(\pi_n) \ge 2\sqrt{n/\log n} - 1$. [18, Theorem 16.8] also showed NBP₁(π_n) = O(n). For the rest two, (18) implies it suffices to show $CC(\pi_n) \le \log n + 4$. Indeed, for every $1 \le k \le n$, choose A = [k]. It is easy to see that $ccm((\pi_n)_A) \le \log n + 4$.

Lastly, we consider separation of \widehat{C} from NBP₁.

Proposition 28. The following hold.

- (1) There exists a Boolean function $f : \{0,1\}^n \to \{0,1\}$ that depends on all its input variables, such that $\text{NBP}_k(f) = 2^{\Omega(\frac{n}{4^k k^3})}$ and $S(f) \leq 16$.
- (2) There exists a Boolean function $f : \{0,1\}^n \to \{0,1\}$ that depends on all its input variables, such that the circuit size of f is $\Omega(2^{n/2}/n)$ and $S(f) \leq 16$.

Proof. (1) Assume n is even (it will be clear that odd n can be handled similarly). Write $z \in \{0, 1\}^n$ as z = (x, y) where $x, y \in \{0, 1\}^{n/2}$. Define $f(z) = g(x) \wedge \text{PARITY}_{n/2}(y)$. Trivially, f depends on all its variables as long as g depends on every variable in x. Observe that $\text{NBP}_k(f) \ge \text{NBP}_k(g)$. On the other hand, it is easy to see that $\text{CC}(f) \le 2$, hence $S(f) \le 16$ by Corollary 21. Set $g = \text{BRS}_{n/2}$ gives the result.

(2) This can be proved similarly as (1).

We summarize the relations and separations in Figure 1 and Table 1, respectively. Note that all measures in Table 1 are lower bounds for OBDD(f). The measures OBDD(f) and $2^{C(f)}$ in Figure 1 can be shown to be incomparable by the method in Proposition 28.



Figure 1: The relation among measures for Boolean functions $f : \{0,1\}^n \to \{0,1\}$. A directed edge from a to b indicates $a \leq b$.

Remark 29. Proposition 23 and Proposition 28 show that there are exponential separations between circuit size (or branching program size, or formula size) of f and S(f) in both directions. In fact, simple functions can provide separations such as $SEQ \in AC^0$. As mentioned in the introduction of [25], Uhlig showed that if the average number of subfunctions is a constant, then the circuit size is linear. The proof for (2) in Proposition 28 shows that this is not true for S(f), S(f) can be exponentially smaller than the average number of subfunctions.

3.4 Lower bound measures for read-*k* BPs

Although we will not elaborate on these, in this subsection we define lower bound measures for BP_k and NBP_k . We say d (not necessarily disjoint) subsets $A_1, \ldots, A_d \subseteq [n]$ is a read-k partition

Table 1: Comparison of measures. The notation \ll (resp. \gg) indicates an exponential separation in the direction Row \ll (resp. \gg) Column. All measures in the table are lower bounds for OBDD.

	NBP_1	BP_1	\widehat{C}	C	\widehat{P}	Р	\widehat{S}	S	
NBP1	=	\leq	$\geq: [4, \text{Thm 1}]$	≪: Prop 23	≪: Prop 26	≪: Prop 23	≪: Prop 27	≪: Prop 23	
TIDI 1		$\ll: [18, \text{Thm 16.8}]$	$\gg:$ Prop 28	$\gg:$ Prop 28	$\gg: Prop 28$	$\gg:$ Prop 28	»: Prop 28	$\gg: Prop 28$	
BD.		_	$\geq: [4, \text{Thm 1}]$	≪: Prop 23	\geq : Thm 15	≪: Prop 23	$\geq: [25, \text{Thm } 2.1]$	≪: Prop 23	
$\mathbf{D}1$		_	$\gg:$ Prop 28	$\gg:$ Prop 28	$\gg:$ Prop 27	$\gg:$ Prop 27	$\gg:$ Prop 25	$\gg:$ Prop 28	
â			_	$\leq:$ Thm 20	\leq : Thm 20	\leq : Thm 20	≪: Prop 27	\leq : Thm 20	
C			=	$\ll:$ Prop 23	$\ll:$ Prop 26	$\ll:$ Prop 23	$\gg:$ Prop 25	$\ll:$ Prop 23	
C					≪: Prop 26	$\leq:$ Thm 20	≪: Prop 27	\leq : Thm 20	
C					_	$\gg:$ Prop 23	$\ll:$ Prop 24	$\gg:$ Prop 25	$\ll:$ Prop 24
\hat{D}					_	$\leq:$ Thm 20	≪: Prop 27	\leq : Thm 20	
Γ						—	$\ll:$ Prop 23	$\gg:$ Prop 25	$\ll:$ Prop 23
D						_	≪: Prop 27	\leq : Thm 20	
1						_	$\gg:$ Prop 25	$\ll:$ Prop 27	
ĉ							_	\leq : Thm 20	
5							—	$\ll:$ Prop 25	

of size ℓ for [n], if they satisfy the following conditions: (1) $\cup_{i=1}^{d} A_i = [n]$; (2) $|A_i| = \ell$ for every $i \leq d-1$, and $|A_d| \leq \ell$; and (3) every element $t \in [n]$ appears in at most $k A_i$'s. A Boolean function $g: \{0,1\}^n \to \{0,1\}$ is said to be a (k,ℓ) -hyperrectangle³ if there exists a read-k partition of size ℓ for [n], say (A_1,\ldots,A_d) , such that for every $x \in \{0,1\}^n$,

$$g(x) = \bigwedge_{i=1}^{a} g_i(x_i), \text{ where } x_i = x|_{A_i}, \ g_i : \{0,1\}^{A_i} \to \{0,1\} \text{ is a Boolean function.}$$
 (25)

Let $f: \{0,1\}^n \to \{0,1\}$ be a Boolean function. We define $\widehat{\mathcal{C}}_k(f,\ell)$ to be the minimal integer r such that $f = \bigvee_{j=1}^r f_j$ where each $f_j: \{0,1\}^n \to \{0,1\}$ is a (k,ℓ) -hyperrectangle. Finally, define⁴ $\widehat{\mathcal{C}}_k(f) = \left(\max_{1 \le \ell \le n} \widehat{\mathcal{C}}_k(f,\ell)^\ell\right)^{\frac{1}{kn}}$.

We proceed to define the counterpart for deterministic case. An integer-valued function $g : \{0,1\}^n \to \mathbb{N}$ is said to be a (k,ℓ) -hyperrectangle if a similar condition as (25) holds, except that the \wedge is replaced by the integer multiplication, and the last function $g_d : \{0,1\}^{A_d} \to \mathbb{N}$ is integer-valued. Given $f : \{0,1\}^n \to \mathbb{N}$ to be an integer-valued function. Similar to Definition 9, define $\widehat{\mathcal{P}}_k^+(f,\ell)$ to be the minimal integer r such that f can be decomposed into r orthogonal (k,ℓ) -hyperrectangles. Then, we define $\widehat{\mathcal{P}}_k^+(f) = \left(\max_{1 \le \ell \le n} \widehat{\mathcal{P}}_k^+(f,\ell)^\ell\right)^{\frac{1}{kn}}$. When $f : \{0,1\}^n \to \{0,1\}$ is a Boolean function, we define $\widehat{\mathcal{P}}_k(f) = \max\{\widehat{\mathcal{P}}_k^+(f), \widehat{\mathcal{P}}_k^+(\neg f)\}$.

Proposition 30. Let $f : \{0,1\}^n \to \{0,1\}$ be a Boolean function. Then,

$$\operatorname{BP}_k(f) \ge \widehat{\mathcal{P}_k}(f), \quad \operatorname{NBP}_k(f) \ge \frac{1}{2}\sqrt{\widehat{\mathcal{C}_k}(f)}.$$

³This is called $(k, n/\ell)$ -rectangle in [30]. If we view d as the dimension of a hyperrectangle, we remark that a Boolean function g can be viewed as a (k, ℓ) -hyperrectangle in different dimensions. For example, consider (2, n/4)hyperrectangles. One can have the following two distinct read-2 partitions of size n/4 for [n] as follows: (1) an equi-partition into four disjoint subsets, so d = 4; (2) equi-partition of [n] into 8 disjoint subsets, say B_1, \ldots, B_8 , then set $A_i = B_i \cup B_{i+1}$ for $i = 1, \ldots, 7$, and set $A_8 = B_8 \cup B_1$, so d = 8. Though (k, ℓ) -hyperrectangles can be of different dimensions, the dimension satisfies $d \leq kn/\ell$.

⁴Here the exponent ℓ/kn in defining $\widehat{\mathcal{C}}_k$ is the inverse of the maximal possible dimension of the (k, ℓ) -hyperrectangles, see the previous footnote.

If $f: \{0,1\}^n \to \mathbb{N}$ is integer-valued, then $\mathrm{BP}_k(f) \ge \widehat{\mathcal{P}_k^+}(f)$.

The lower bound for NBP_k by $\widehat{\mathcal{C}}_k(f)$ is proved in [4] (see also [30]), and it is shown that $\widehat{\mathcal{C}}_k(BRS_n) = 2^{\Omega(\frac{n}{k^3 4^k})}$. The lower bounds for BP_k can be proved by a small modification of the proof for Theorem 15, using the method in [4]. We omit the detail.

4 \widehat{S} is small for TEP

Iwama et al. [16] showed that $BP_1(TEP, h) \ge k^h$ as long as $h \le (\frac{1}{3} - \frac{1}{2\log k})k$. The approach in [16] is to directly exploit properties satisfied by a read-once BP for (TEP, h), and seems hard to be generalized to the read-k case. It would be desirable to give an alternative proof that is amenable for a possible generalization to read-k. Unfortunately, below we show that \hat{S} does not work: $\hat{S}(\text{TEP}, h)$ is small. In this section we will use the definition of \hat{S} on k-ary functions, i.e., |D| = k in Remark 3 (do not confuse |D| = k with the k in $|D|^k$ there).

We use the following notation. Let (M, L, R) denote a partition of the input variables for (TEP, h) where M denotes the matrix at the root of the binary tree, L and R denote the left and right child, respectively. Note that both L and R correspond to inputs for (TEP, h - 1). Recall n_h denotes the input size for (TEP, h). For a subset $A \subseteq [n_h]$, we think of A as a subset of input variables for (TEP, h) and write $A = (A_M, A_L, A_R)$, where $A_M = A \cap M$, $A_L = A \cap L$ and $A_R = A \cap R$. For notational simplicity, when the parameter h is clear from the context, we use TEP_A to denote the matrix $(\text{TEP}, h)_A$, and for $\alpha \in [k]^A$, we use TEP_α to denote the subfunction $(\text{TEP}, h)_{A,\alpha}$. We use $M_{ij} \in A_M$ to mean that A_M contains the variable at entry (i, j) of the root matrix M.

Lemma 31. Let $h \geq 2$, $A \subseteq [n_h]$. Suppose $A = (\emptyset, A_L, A_R)$. Let $\alpha = (\alpha_L, \alpha_R) \in [k]^A = [k]^{A_L} \times [k]^{A_R}$, $\alpha' = (\alpha'_L, \alpha'_R) \in [k]^A = [k]^{A_L} \times [k]^{A_R}$. Then,

$$(\text{TEP}, h)_{A,\alpha} = (\text{TEP}, h)_{A,\alpha'}$$
$$\iff \left((\text{TEP}, h-1)_{A_L,\alpha_L}, (\text{TEP}, h-1)_{A_R,\alpha_R} \right) = \left((\text{TEP}, h-1)_{A_L,\alpha'_L}, (\text{TEP}, h-1)_{A_R,\alpha'_R} \right). \quad (26)$$

Proof. The direction \Leftarrow . Obvious.

The direction \implies . Assume for the sake of a contradiction the implication is not true. Without loss of generality we may assume $(\text{TEP}, h - 1)_{A_L,\alpha_L} \neq (\text{TEP}, h - 1)_{A_L,\alpha'_L}$. We will show

$$(\text{TEP}, h-1)_{A_L, \alpha_L} \neq (\text{TEP}, h-1)_{A_L, \alpha'_L} \Longrightarrow (\text{TEP}, h)_{A, \alpha} \neq (\text{TEP}, h)_{A, \alpha'}$$

Indeed, let $\beta_L \in [k]^{L-A_L}$ be such that $(\text{TEP}, h-1)_{A_L,\alpha_L}(\beta_L) = i \neq i' = (\text{TEP}, h-1)_{A_L,\alpha'_L}(\gamma_L)$. Then, set $\beta_M \in [k]^M$ to be such that every entry in the *i*-th row equals to *i*, and every entry in the *i'*-th row equals to *i'*. Choose $\beta_R \in [k]^{R-A_R}$ arbitrarily. Set $\beta = (\beta_M, \beta_L, \beta_R)$. Then, $(\text{TEP}, h)_{A,\alpha}(\beta) = i \neq i' = (\text{TEP}, h)_{A,\alpha'}(\beta)$.

Lemma 32. For every $h \ge 2$, for every $\frac{n_h}{2} + 2^{h-2} \le \ell \le n_h$, there exists a subset $A \subseteq [n_h]$ of size $|A| = \ell$ such that for every fixed $i \in [k]$,

$$\Pr_{\alpha}[\operatorname{TEP}_{\alpha} = i] \ge \frac{1}{2^{2^{h-1}-1} \cdot k},$$

where α is chosen from $[k]^A$ uniformly at random. Here $\text{TEP}_{\alpha} = i$ means the subfunction $(\text{TEP}, h)_{A,\alpha}$ identically equals to *i*.

Proof. Suppose k is even. Consider firstly the case h = 2. Then $|A| \ge \frac{k^2+2}{2} + 1 = \frac{k^2}{2} + 2$. Choose A such that $|A_M| = |A| - 2 \ge k^2/2$, $|A_R| = |A_L| = 1$. In particular, both of the two leaves belong to A. Let $\alpha = (\alpha_M, \alpha_L, \alpha_R) \in [k]^{A_M} \times [k]^{A_L} \times [k]^{A_R}$. Let $Q = \{(r, s) \in [k] \times [k] : M_{rs} \in A_M\}$. Let α be chosen from $[k]^A$ uniformly at random. Then, for every fixed $i \in [k]$,

$$\Pr_{\alpha}[\text{TEP}_{\alpha} = i] = \sum_{(r,s)\in Q} \Pr_{\alpha_{L}}[\alpha_{L} = r] \times \Pr_{\alpha_{R}}[\alpha_{R} = s] \times \Pr_{\alpha_{M}}[M_{rs} = i] = \frac{|Q|}{k^{3}} = \frac{|A_{M}|}{k^{3}} \ge \frac{1}{2k}.$$
 (27)

We use induction to prove the general case. Specifically, we choose A such that,

- (1) $|A_M| \ge k^2/2;$
- (2) choose A_L so that all the leave variables in L are chosen, and for every matrix node in L, A_L contains at least a half of its entries;
- (3) choose A_R similarly as A_L .

The above choice for A is feasible because $|A| \ge \frac{n_h}{2} + 2^{h-2}$. Observe that the choice of A_L and A_R allows induction on height h-1. Hence, by the same calculation as (27) and by induction on height h-1, one has that for every fixed $i \in [k]$,

$$\Pr_{\alpha}[\text{TEP}_{\alpha} = i] \ge |A_M| \times \frac{1}{2^{2^{h-2}-1} \cdot k} \times \frac{1}{2^{2^{h-2}-1} \cdot k} \times \frac{1}{k} \ge \frac{1}{2^{2^{h-1}-1} \cdot k}$$

Finally, it is easy to see that the same argument also applies when k is odd. We omit the details. \Box

Theorem 33. For $k \ge 2$ and $1 \le h \le \log k$, one has

$$k \le \widehat{S}(\text{TEP}, h) \le \frac{2^{2^{h-1}+1}}{3} \cdot k.$$

In particular, $\widehat{S}(\text{TEP}, \log \log k) \le k^2/3$.

Proof. By definition, $\widehat{S}(\text{TEP}, h) = \max_{1 \le \ell \le n_h} \min_{|A| = \ell} \frac{k^{|A|}}{\mathsf{mult}((\text{TEP}, h)_A)}$.

The lower bound. Choosing $\ell = n_h$ and $A = [n_h]$ implies that $\mathsf{mult}(\mathsf{TEP}_A) = k^{n_h - 1}$, hence $\widehat{S}(\mathsf{TEP}, h) \ge k$.

The upper bound. We will use induction on h. The base case when h = 1 is clear: $\widehat{S}(\text{TEP}, 1) = k \leq 4k/3$. Assume now $h \geq 2$ and the claim is true for h - 1. By definition of \widehat{S} , we will show that for each $1 \leq \ell \leq n_h$, there exists a subset $A \subseteq [n_h]$ such that $|A| = \ell$ and $\text{mult}(\text{TEP}_A) \geq \phi(h)k^{|A|-1} = \phi(h)k^{\ell-1}$, where $\phi(h) = 3/(2^{2^{h-1}+1})$. We consider three cases.

• Case 1: $\ell \leq n_{h-1}$. Choose A such that $A_M = \emptyset$, $|A_L| = \ell$ and $A_R = \emptyset$, i.e., $A = A_L$. Apply Lemma 53 in the case when $A_R = \emptyset$, one has that for every $\alpha, \alpha' \in [k]^A = [k]^{A_L}$,

$$(\text{TEP}, h)_{A,\alpha} = (\text{TEP}, h)_{A,\alpha'} \iff (\text{TEP}, h-1)_{A_L,\alpha} = (\text{TEP}, h-1)_{A_L,\alpha}$$

Hence, $m((\text{TEP}, h)_A) = m((\text{TEP}, h - 1)_{A_L})$. By induction, there exists A_L such that $m((\text{TEP}, h - 1)_{A_L}) \ge \phi(h - 1)k^{\ell - 1}$. Hence, $m((\text{TEP}, h)_A) \ge \phi(h - 1)k^{\ell - 1} \ge \phi(h)k^{\ell - 1}$.

- Case 2: $n_{h-1} < \ell \le n_{h-1} + k(k-1)$. Choose A to be of size ℓ such that,
 - (1) A_M is a subset of the entries in the first k-1 rows of M;
 - (2) $|A_L| = n_{h-1};$
 - (3) $A_R = \emptyset$.

Clearly, the above choice is feasible. Let

 $\mathcal{F} = \{ \alpha = (\alpha_M, \alpha_L, \alpha_R) \in [k]^{A_M} \times [k]^{A_L} \times [k]^{A_R} : (\text{TEP}, h - 1)_{A_L, \alpha_L} = k \}.$

Since $|A_L| = n_{h-1}$, we have $\Pr_{\alpha_L}[(\text{TEP}, h-1)_{A_L,\alpha_L} = k] = 1/k$. Hence, $|\mathcal{F}| = k^{|A|-1}$. The choice of A implies that $(\text{TEP}, h)_{A,\alpha} = (\text{TEP}, h)_{A,\beta}$ whenever $\alpha, \beta \in \mathcal{F}$. Hence, $m((\text{TEP}, h)_A) \geq |\mathcal{F}| = k^{|A|-1} > \phi(h)k^{|A|-1}$.

- Case 3: $\ell > n_{h-1} + k(k-1)$. Choose A to be of size ℓ such that,
 - (1) $k(k-1) 2^{h-2} < |A_M| \le k^2;$ (2) $|A_L| \ge \frac{n_{h-1}}{2} + 2^{h-3};$ (3) $|A_R| \ge \frac{n_{h-1}}{2} + 2^{h-3}.$

Clearly, the above choice is feasible. Moreover, $|A_M| \ge k^2 - k - 2^{h-2} \ge k^2 - 5k/4$ as $h \le \log k$. Specification on A_L an A_R will be given by Lemma 32. Indeed, by Lemma 32, there exist choices of A_L and A_R such that for every $(i, j) \in [k] \times [k]$,

$$\begin{aligned} \Pr_{\alpha_L}[(\text{TEP}, h-1)_{A_L, \alpha_L} &= i] \geq \frac{1}{2^{2^{h-2}-1} \cdot k}, \quad \Pr_{\alpha_R}[(\text{TEP}, h-1)_{A_R, \alpha_R} &= j] \geq \frac{1}{2^{2^{h-2}-1} \cdot k}. \end{aligned} (28) \\ \text{Let } Q &= \{(i, j) \in [k] \times [k] : M_{ij} \in A_M \}. \text{ Similar to } (27), \text{ apply } (28), \\ \Pr_{\alpha}[(\text{TEP}, h)_{A, \alpha} &= 1] \\ &= \sum_{(i, j) \in Q} \Pr_{\alpha_L}[(\text{TEP}, h-1)_{A_L, \alpha_L} &= i] \times \Pr_{\alpha_R}[(\text{TEP}, h-1)_{A_R, \alpha_R} &= j] \times \Pr_{\alpha_M}[M_{ij} = 1] \\ &\geq \frac{|A_M|}{2^{2^{h-1}-2} \cdot k^3} \geq \frac{1-\frac{5}{4k}}{2^{2^{h-1}-2}} \cdot \frac{1}{k} \geq \frac{3}{2^{2^{h-1}+1}} \cdot \frac{1}{k} \\ \text{Hence, mult}((\text{TEP}, h)_A) \geq \frac{3}{2^{2^{h-1}+1}} \cdot k^{|A|-1} &= \phi(h) \cdot k^{|A|-1}. \end{aligned}$$

In all three cases we have verified the existence of the desired A, the proof is completed.

4.1 *S* **on** TEP

Let $S((\text{TEP}, h), \ell) = \min_{A \subseteq [n_h], |A| = \ell} \operatorname{nrows}((\text{TEP}, h)_A)$. Then, $S(\text{TEP}, h) = \max_{1 \le \ell \le n_h} S((\text{TEP}, h), \ell)$. **Theorem 34.** (1) $S(\text{TEP}, 2) = k^2$ and is achieved at either $\ell = k + 1$ or $\ell = k + 2$, *i.e.*, $S((\text{TEP}, 2), k + 1) = S((\text{TEP}, 2), k + 2) = k^2$;

- (2) $S(\text{TEP}, 3) \ge k^3/16$ for $k \ge 4$;
- (3) $S(\text{TEP}, h) \le k^h k^{h-2}(k-2) < k^h \text{ for } h \ge 3.$

Theorem 34-(1) shows that the value of S(f) is not necessarily achieved at |A| = n/2. The complete proof is lengthy, we provide it in Appendix A.

5 The read-once non-deterministic BP lower bound of satisfiable Tseitin formulas via \hat{C} : a short proof

In a line of recent works [15, 12, 13], BP (OBDD and NBP₁) sizes of satisfiable Tseitin formulas are studied for applications in proof complexity. Let $\kappa(H)$ denote the number of connected components of graph H. Define $\kappa_G(\ell) = \max_{H \leq G, |E(H)| = \ell} \kappa(H)$, where $H \leq G$ means H is a subgraph of G. [12, 13] proved that if $\kappa(G) = 1$, then NBP₁(TS_{G,c}) $\geq \max_{1 \leq \ell \leq m} 2^{n-\kappa_G(\ell)-\kappa_G(m-\ell)+1}$, and used this as a starting point to prove NBP₁ lower bounds for satisfiable Tseitin formulas. Below we give a short proof using \widehat{C} for arbitrary $\kappa(G)$. We need the following fact.

Fact ([12, Lemma 2]). $|TS_{G,c}^{-1}(1)| = 2^{m-n+\kappa(G)}$.

Theorem 35. Let $TS_{G,c}$ be a satisfiable Tseitin formula. Then,

$$\widehat{C}(\mathrm{TS}_{G,c}) \ge \max_{1 \le \ell \le m} 2^{n - \kappa_G(\ell) - \kappa_G(m-\ell) + \kappa(G)}.$$

Proof. Fix an $1 \leq \ell \leq m$, consider an arbitrary $A \subseteq [m]$ of size $|A| = \ell$, let B = [m] - A. We view $A \subseteq E$, so $A \cup B = E$. Consider an A-rectangle $(p,q) : \{0,1\}^A \times \{0,1\}^B \to \{0,1\}$, that is, $(p,q)(a,b) = p(a) \wedge q(b)$ for every $(a,b) \in \{0,1\}^A \times \{0,1\}^B = \{0,1\}^E$. Suppose further the rectangle satisfies (p,q)(a,b) = 1 implies $\operatorname{TS}_{G,c}(a,b) = 1$ for every (a,b). Let $P = p^{-1}(1) \subseteq \{0,1\}^A$ and $Q = q^{-1}(1) \subseteq \{0,1\}^B$. Then, it suffices to show $|P| \times |Q|$ is "small".

Let $v \in V$. Let E(v) denote the set of edges in E that are incident to v, let $E(A, v) = E(v) \cap A$ and $E(B, v) = E(v) \cap B$. Then, $E(v) = E(A, v) \cup E(B, v)$. Given $b \in \{0, 1\}^B$, by an abuse of notation, we define a labelling function

$$b: V \to \{0,1\}, \quad v \mapsto \begin{cases} 0, & E(B,v) = \emptyset, \\ \sum_{e \in E(B,v)} b_e \mod 2, & \text{otherwise.} \end{cases}$$

Then, for $x = (a, b) \in \{0, 1\}^A \times \{0, 1\}^B = \{0, 1\}^E$, we have

$$\sum_{e \in E(v)} x_e = \sum_{e \in E(A,v)} a_e + \sum_{e \in E(B,v)} b_e \equiv b(v) + \sum_{e \in E(A,v)} a_e \mod 2.$$
(29)

Now, fix an arbitrary $b^* \in Q$. By (29), for every $a \in P$, we have

$$(p,q)(a,b^*) = 1 \Longrightarrow \mathrm{TS}_{G,c}(a,b^*) = 1 \Longrightarrow \forall \ v \in V, \ b^*(v) + \sum_{e \in E(A,v)} a_e \equiv c(v) \mod 2$$
$$\Longrightarrow \forall \ v \in V, \ \sum_{e \in E(A,v)} a_e \equiv c(v) - b^*(v) \mod 2.$$

Let $H_A = (V, A)$ be the subgraph of G given by edges in A. The above and the **Fact** imply that

$$\forall \ a \in P, \ \mathrm{TS}_{H_A, c-b^*}(a) = 1 \Longrightarrow |P| \le |\mathrm{TS}_{H_A, c-b^*}^{-1}(1)| = 2^{|A| - n + \kappa(H_A)} \le 2^{\ell - n + \kappa_G(\ell)}.$$

By symmetry, $|Q| \leq 2^{m-\ell-n+\kappa_G(m-\ell)}$. Hence, $|P| \times |Q| \leq 2^{m-2n+\kappa_G(\ell)+\kappa_G(m-\ell)}$. The claim follows by applying the **Fact** again to $TS_{G,c}$.

6 \widehat{S} and \widehat{C} on Gál-type functions, and related blocking sets

Gál's original function [11] was defined using projective planes, while a closely related function the Bollig-Wegener function [3] was defined from representing numbers in a prime basis. In Section 2.8 we have defined both functions with respect to a given bipartite graph. This definition makes the connection between the two readily apparent. Indeed, by definition in Section 2.8, for every $x \subseteq A$,

$$\operatorname{GAL}_G(x) = 1 \Longleftrightarrow \operatorname{BW}_G(x, y) = 1, \forall \ \emptyset \neq y \subseteq B \Longleftrightarrow \operatorname{BW}_G(x, \{b\}) = 1, \forall \ b \in B.$$
(30)

Furthermore, our definition also naturally allows a possible further approach to tackle the BP₁ vs weight problem, as we will see shortly. Below we discuss \hat{S} and \hat{C} on generalized Gál's function and Bollig-Wegener function, and related mathematical problems.

6.1 The point-line incidence graph over finite fields

Here we specify a bipartite graph to instantiate the GAL and BW functions we will study. Let \mathbb{F}_q be a finite field of order q, we use the notation \mathbb{F}_p to denote the case when p is a prime and hence \mathbb{F}_p is a prime field. We associate every $(i, j) \in \mathbb{F}_q^2$ a unique *non-vertical* line $\ell_{(i,j)} \subseteq \mathbb{F}_q^2$ given by

$$\ell_{(i,j)} = \{ (t, i+jt) : t \in \mathbb{F}_q \}.$$
(31)

Note that the "direction" of the line $\ell_{(i,j)}$ is (1,j), hence is "non-vertical", whereas we think of the direction (0,1) as the vertical direction. Let $A = B = \mathbb{F}_q^2$. Consider the bipartite graph $G_q(A \cup B, E)$ defined by the point-line incidence relation. Specifically, there is an edge between $a \in A$ and $b \in B$ if and only if $a \in \ell_b$. In other words, we think of A as the set of q^2 points of \mathbb{F}_q^2 , and B as the set of q^2 non-vertical lines in \mathbb{F}_q^2 . It is easy to verify the following.

Lemma 36. The bipartite graph G_q is q-regular and $K_{2,2}$ -free.

Definition 37. A subset of points $S \subseteq \mathbb{F}_q^2$ is called a blocking set if and only if it intersects every non-vertical line, it is called a minimal blocking set if no subset of S is a blocking set.

Remark 38. The usual definition of blocking sets in an affine plane requires to intersect all lines, not just the non-vertical lines, see e.g., [5]. Blocking sets under this definition in both affine and projective spaces have been widely studied in the past several decades, see [14, 2], most studies focus on classifying minimal blocking sets. Our definition is slightly different, we will also ask a different question (Question 1) in Section 6.2.

Using the bipartite graph $G_q(A \cup B, E)$, a subset $S \subseteq A$ is a blocking set if and only if Nb(S) = B. It is also not hard to show that $S \subseteq \mathbb{F}_p^2$ is a blocking set if and only if the polynomial $P_S(x, y) \in \mathbb{F}_p[x, y]$

$$P_{S}(x,y) = \prod_{(a,b)\in S} (x+ay-b)$$
(32)

is identically 0 on \mathbb{F}_p^2 .

We will consider GAL_{G_q} and BW_{G_q} . In this case, $\operatorname{GAL}_{G_q}(x) = 1$ if and only if x is a blocking set in \mathbb{F}_q^2 . Gál in [11] defined her function similar to the one given above, but in the projective plane $\operatorname{PG}(2,q)$ (see the detail in [11]), instead of the affine plane \mathbb{F}_q^2 . Bollig and Wegener defined their function in [3] in a different language, and remarked that their function BW "is somehow similar to the construction in [11]". By phrasing both functions over G_q , this connection is shown in (30).

6.2 \widehat{S} and \widehat{C}

Gál in [11] showed that her function defined using the projective plane PG(2, q) is q-mixed, hence has exponential BP₁ complexity. Below we generalize Gál's result to arbitrary bipartite graphs satisfying appropriate conditions, by a similar argument as [11].



Figure 2: The illustration for Theorem 39.

Theorem 39. Let $G = (A \cup B, E)$ be a bipartite graph that is r-regular and $K_{2,s}$ -free for some $s \ge 2$. Then, $\widehat{S}(\text{GAL}_G) \ge 2^{\lfloor (r-1)/(s-1) \rfloor}$.

Proof. By Lemma 2, it suffices to show that GAL_G is $\lfloor (r-1)/(s-1) \rfloor$ -mixed. Let $S \subseteq A$ be an arbitrary subset of size $\lfloor (r-1)/(s-1) \rfloor$, let $\alpha \neq \beta$ be two subsets of S and assume $a \in \alpha \setminus \beta$, see Figure 2. We need to show that there exists a subset $\gamma \subseteq A \setminus S$ such that $\operatorname{Nb}(\alpha \cup \gamma) = B$ but $\operatorname{Nb}(\beta \cup \gamma) \neq B$.

Since G is $K_{2,s}$ -free and r-regular, and $|S| = \lfloor (r-1)/(s-1) \rfloor$,

$$|\operatorname{Nb}(\beta) \cap \operatorname{Nb}(a)| \le (s-1)|\beta| < (s-1)|S| \le r-1 < |\operatorname{Nb}(a)| \Longrightarrow \exists b \in \operatorname{Nb}(a) \setminus \operatorname{Nb}(\beta).$$

Take $\gamma = A \setminus (S \cup \operatorname{Nb}(b))$. Then, $b \notin (\operatorname{Nb}(\beta) \cup \operatorname{Nb}(\gamma))$, i.e., $\operatorname{Nb}(\beta \cup \gamma) \neq B$. To show $\operatorname{Nb}(\alpha \cup \gamma) = B$, consider an arbitrary $b' \notin \operatorname{Nb}(a)$. Again, since G is $K_{2,s}$ -free and r-regular and $|S| = \lfloor (r-1)/(s-1) \rfloor$,

$$|(S \cup \operatorname{Nb}(b)) \cap \operatorname{Nb}(b')| \le |(S \setminus \{a\}) \cap \operatorname{Nb}(b')| + |\operatorname{Nb}(b) \cap \operatorname{Nb}(b')|$$
$$\le (s-1)|S \setminus \{a\}| + (s-1) \le r-1 < |\operatorname{Nb}(b')|.$$

Hence, there exists $t \in \gamma \cap Nb(b')$. This together with $Nb(a) \subseteq Nb(\alpha)$ shows $Nb(\alpha \cup \gamma) = B$. \Box

Corollary 40. $BP_1(GAL_{G_q}) \ge 2^{q-1}$.

Proof. Apply Proposition 14, Lemma 36 and Theorem 39.

Similar to Gál's BP₁ lower bound [11], Bollig and Wegener in [3] also showed that BP₁(BW_{G_q}) has an exponential lower bound. In [3] they pointed out that BW_{G_q} is not mixed, here we strengthen this by showing that in fact $\hat{S}(BW_{G_q})$ is small.

Theorem 41.
$$\widehat{S}(BW_{G_a}) = O(1).$$

Proof. Let $k_t = t(q+1)$. We first show for $1 \le t \le q$, there exist $S \subseteq A \cup B$ with size $|S| = k_t$ such that $\mathsf{mult}((\mathsf{BW}_{G_q})_S)$ is large. Let $V = \{x_1, \ldots, x_q\} \subseteq A$ be a subset of q points that form a vertical line. Then, $Nb(x_i) \cap Nb(x_j) = \emptyset$ and V is a minimal blocking set. Let $T = \{x_1, \ldots, x_t\} \subseteq V$.



Figure 3: The illustration for Theorem 41.

Choose $S = T \cup Nb(T)$, so $|S| = t(q+1) = k_t$, see Figure 41. Let $H(S, E(S)) \leq G_q(A \cup B, E)$ denote the induced subgraph by S. Let $(\alpha, \beta) \in \{0, 1\}^T \times \{0, 1\}^{Nb(T)}$. With this notation, one has

$$(\mathrm{BW}_{G_q})_{S,(\alpha,\beta)} = 1 \Longleftrightarrow \mathrm{BW}_H(\alpha,\beta) = 1.$$
(33)

Let P_t denote the probability that $BW_H(\alpha, \beta) = 0$ where (α, β) is chosen uniformly at random, i.e., the probability that there are no edges between α and β in H.

$$P_t = \sum_{i=0}^{t} {t \choose i} 2^{-t} 2^{-qi} = (2^{-1} + 2^{-(q+1)})^t \le 5/8 \Longrightarrow \Pr[BW_H(\alpha, \beta) = 1] \ge 3/8$$

By (33), this implies $\mathsf{mult}((\mathsf{BW}_{G_q})_S) \ge 2^{k_t} \cdot 3/8$ as desired. With slight modification, the other values of $1 \le k \le 2q^2$ can all be handled, we omit the details.

Next we discuss $\widehat{C}(\text{GAL}_{G_q})$, it is best to look at this problem within a background. Using the terminology from [19], define the *weight* of $f : \{0, 1\}^n \to \{0, 1\}$ to be

$$w(f) = \text{DNFSize}(f) + \text{CNFSize}(f), \tag{34}$$

where DNFSize(f) and CNFSize(f) denote the DNF and CNF sizes of f, respectively.

Open Problem ([3, 19, 30]). Does there exist a Boolean function f such that $BP_1(f)$ is superpolynomially larger than w(f)?

This problem has been open for more than two decades. If we replace BP_1 by OBDD, we could deduce separations in both sides. Let DTSize(f) denote the decision tree size of f.

Proposition 42.
$$w(\text{NAND}_n) \ge \exp\left(\Omega(\sqrt{n/\log n})\right)$$
 but $\text{OBDD}(\text{NAND}_n) = O(n)$
 $w(\text{ISA}_n) \le \text{DTSize}(\text{ISA}_n) = O(n^2)$ but $\text{OBDD}(\text{ISA}_n) = \exp(\Omega(n/\log n)).$

Proof. In [9] it shows (see also a proof from [18]) DTSize $(f) \leq \exp(O((\log n) \cdot \log^2 w(f)))$. By [19], DTSize $(\text{NAND}_n) \geq 2^{\Omega(n)}$. These two together implies the lower bound for $w(\text{NAND}_n)$. Theorem 4.3.3 from [30] shows that OBDD(ISA_n) $\geq \exp(\Omega(n/\log n))$. The upper bounds are easy to see. \Box

It is easy to see that $\text{CNFSize}(\text{GAL}_{G_q}) = O(q^3)$ and $\text{DNFSize}(\text{BW}_{G_q}) = O(q^3)$. Hence, it is interesting to ask what is $\text{DNFSize}(\text{GAL}_{G_q})$? It is not hard to see that $\text{DNFSize}(f) \ge \widehat{C}(f)$. Hence, for the purpose of a lower bound, it suffices to show $\widehat{C}(\text{GAL}_{G_q}, n/2)$ is large, where $n = q^2$. This raises naturally the following question. Question 1. Let $M \subseteq \mathbb{F}_q^2$ be an arbitrary subset of size $q^2/2$. Is the number of blocking sets inside M always exponentially (or superpolynomially) smaller than $2^{q^2/2}$?

Indeed, it is not hard to see that if there is an $M \subseteq \mathbb{F}_q^2$ of size $n/2 = q^2/2$ with t blocking sets inside, then GAL_{G_q} contains a 1-monochromatic n/2-rectangle of size $t2^{p^2/2}$. If one wishes to show $\widehat{C}(\operatorname{GAL}_{G_q}, n/2)$ is large by showing that every 1-monochromatic n/2-rectangle is small, then answering Question 1 is crucial. The following simple fact is a contrast to Question 1.

Proposition 43. As $q \to \infty$, almost all subsets in \mathbb{F}_q^2 are blocking sets.

Proof. By definition, every non-blocking-set is a subset of $\mathbb{F}_q^2 \setminus \ell$ for some (non-vertical) line ℓ . As there are q^2 such lines, the number of non-blocking-set $\leq q^2 \cdot 2^{q^2-q}$. This implies the number of blocking sets $\geq 2^{q^2}(1-q^2/2^q) \to 2^{q^2}$, as $q \to \infty$.

One potential way towards answering Question 1 might be the following. Question 2. Classify all minimal blocking sets in \mathbb{F}_q^2 .

6.3 Some results on minimal blocking sets in \mathbb{F}_n^2

Here we give some results for Question 2. Structures of minimal blocking sets (MBS for short) might also be of independent interest. For simplicity throughout this section we work with \mathbb{F}_p^2 where p is a prime. When we say lines we mean non-vertical lines unless specified otherwise. We say two lines are parallel if they are two distinct lines with the same direction.

Theorem 44. MBS in \mathbb{F}_p^2 have size at least p. Furthermore,

- (1) There are exactly p MBS of size p, each of which is a vertical line.
- (2) There are no MBS of size p + 1.
- (3) Let ℓ and ℓ' be two intersecting lines. Let $x \in \ell \setminus \ell'$. Let $y \in \ell'$ be the unique point in y such that the line determined by x and y is vertical. Let ℓ_y be the unique line parallel to ℓ and passing through y. Let ℓ'_x be the unique line parallel to ℓ' and passing through x. Define $\phi(x) = \ell_y \cap \ell'_x$. Then, $\ell \cup \ell' \cup \{\phi(x)\} \setminus \{x, y\}$ is an MBS of size 2p - 2, see Figure 4-(i).
- (4) Every pair of two intersecting lines is an MBS of size 2p 1.
- (5) Let ℓ be a line. Let a_1, \ldots, a_{p-1} be p-1 points not in ℓ , such that no two of them are in the same vertical line, and such that every line parallel to ℓ passes through some a_i . Then, $\ell \cup \{a_1, \ldots, a_{p-1}\}$ is an MBS of size 2p-1, see Figure 4-(ii).
- (6) Let ℓ and ℓ' be two intersecting lines, let $x \in \ell$, let $y \in \ell'$ be such that the line determined by xand y is not vertical. Let ℓ_y and ℓ'_x be the two lines that are parallel to ℓ and ℓ' , respectively. Let $a \neq x, y$ be a point in the line determined by x and y. Let $b \in \ell_y$, $b \neq y$ and $b \neq \ell_y \cap \ell'_x$. Let $c \in \ell'_x$, $c \neq x$ and $c \neq \ell_y \cap \ell'_x$. Then, $\ell \cup \ell' \cup \{a, b, c\} \setminus \{x, y\}$ is an MBS of size 2p, see Figure 4-(iii).
- (7) Let ℓ_1 be a vertical line. Let ℓ_2 and ℓ_3 be two parallel lines intersecting with ℓ_1 at a and b, respectively. Then, $\ell_1 \cup \ell_2 \cup \ell_3 \setminus \{a, b\}$ is an MBS of size 3p 4, see Figure 4-(iv).



Figure 4: The illustration for Theorem 44.

Most of the statements in Theorem 44 are not hard to prove. For example, the MBS has size at least p can be deduced via a degree argument by the polynomial characterization (32) (of course, it can also be proved directly). We find the proof for (5) is particularly interesting, and the lemmas used to prove it might be of independent interest. So we give its proof below.

6.3.1 A lemma of intersecting points in \mathbb{F}_p^2

Lemma 45. Let p be a prime, let $x_1, \ldots, x_{p-1} \in \{1, \ldots, p-1\}$ be distinct. Then, $\prod_{1 \le i < j \le p-1} (ix_j - jx_i) \equiv 0 \mod p$.

Proof. For $1 \le i \le p-1$, let $y_i = x_i/i \in \{1, \ldots, p-1\}$. Writing $ix_j - jx_i = ij(y_j - y_i)$, one observes that in \mathbb{F}_p , $\prod_{1\le i < j\le p-1}(ix_j - jx_i) = 0$ is equivalent to $\prod_{1\le i < j\le p-1}(y_j - y_i) = 0$. So it suffices to show that $y_i = y_j$ for some $i \ne j$. Assume otherwise, i.e., $y_1, \ldots, y_{p-1} \in \{1, \ldots, p-1\}$ are all distinct. Then, on one hand, by Wilson's theorem, $\prod_{i=1}^{p-1} y_i = (p-1)! \equiv -1 \mod p$. On the other hand, since x_i are all distinct, $\prod_{i=1}^{p-1} y_i = \prod_{i=1}^{p-1} \frac{x_i}{i} = \frac{\prod_{i=1}^{p-1} x_i}{\prod_{i=1}^{p-1} i} = \frac{(p-1)!}{(p-1)!} = 1 \mod p$, contradicting to the previous equation.

Below in Lemma 46 lines mean arbitrary lines, i.e., either vertical or non-vertical.

Lemma 46. Let ℓ_1, \ldots, ℓ_p be p parallel lines in \mathbb{F}_p^2 , let ℓ'_1, \ldots, ℓ'_p be p parallels in a different direction. For $1 \leq i \leq p$, let $a_i = \ell_i \cap \ell'_i$. Then, for every point a_i , there exists $a_j, a_k, j, k \neq i$ and $j \neq k$, such that these three points a_i, a_j, a_k are colinear.

Proof. Note that there are p+1 different directions, i.e., (0,1) and (1,d) for $d \in \mathbb{F}_p$.

Consider the case where the two directions are (1, d) and (1, d'), where $d \neq d'$, for lines ℓ_1, \ldots, ℓ_p and ℓ'_1, \ldots, ℓ'_p , respectively. By (31), the lines ℓ_i can be parametrized as

 $\ell_i = \{(t, m_i + dt) : t \in \mathbb{F}_p\}, \quad i = 1, \dots, p, \quad \text{where } m_1, \dots, m_p \in \mathbb{F}_p \text{ are all distinct.}$ (35)

Similarly, lines ℓ'_i can be parametrized as

 $\ell'_{i} = \{(t, n_{i} + d't) : t \in \mathbb{F}_{p}\}, \quad i = 1, \dots, p, \quad \text{where } n_{1}, \dots, n_{p} \in \mathbb{F}_{p} \text{ are all distinct.}$ (36)

By (35) and (36), we have

$$a_i = (t_i, m_i + dt_i), \quad \text{where } t_i = \frac{m_i - n_i}{d' - d}, \quad i = 1, \dots, p.$$
 (37)

Note that since $d' \neq d$, $t_i \in \mathbb{F}_p$ is well-defined.

Without loss of generality, we may assume the point that is fixed is a_1 , and we wish to show there exist $2 \leq j < k \leq p$ such that a_1, a_j, a_k are in the same line. This is equivalent to showing the p-1 directions a_2-a_1, \ldots, a_p-a_1 are not all distinct. By (37), $a_i-a_1 = (t_i-t_1, m_i-m_1+d(t_i-t_1))$. The directions $a_j - a_1$ and $a_k - a_1$ are the same if and only if

$$\det(A_{jk}) = 0, \quad \text{where matrix } A_{jk} = \begin{pmatrix} t_j - t_1 & m_j - m_1 + d(t_j - t_1) \\ t_k - t_1 & m_k - m_1 + d(t_k - t_1) \end{pmatrix}, \quad 2 \le j < k \le p.$$

In the above, $\det(A_{jk}) = 0$ means $\det(A_{jk}) \equiv 0 \mod p$, and this notation is used for what follows. By simple linear algebra and (37), we have

$$\det(A_{jk}) = 0 \iff \det(A'_{jk}) = 0, \quad \text{where matrix } A'_{jk} = \begin{pmatrix} n_j - n_1 & m_j - m_1 \\ n_k - n_1 & m_k - m_1 \end{pmatrix}, \quad 2 \le j < k \le p.$$

Hence, it suffices to show $\det(A'_{jk}) = 0$, for some $2 \le j < k \le p$. Observe further that since m_i are all distinct, one has for every $2 \le i \le p$, $m_i - m_1 \in \{1, \ldots, p-1\}$ and these p-1 numbers are all distinct. Similarly, $n_i - n_1 \in \{1, \ldots, p-1\}$ are all distinct. Hence, without loss of generality, we may assume $m_i - m_1 = i - 1$ and let $n_i - n_1 = x_i \in \{1, \ldots, p-1\}$ for $2 \le i \le p$. In this notation, the problem is then reduced to show

$$\det(A'_{jk}) \equiv 0 \mod p, \quad \text{for some } 2 \le j < k \le p, \tag{38}$$

where

$$A'_{jk} = \begin{pmatrix} x_j & j-1 \\ x_k & k-1 \end{pmatrix}, \text{ and } x_2, \dots, x_p \in \{1, \dots, p-1\} \text{ are distinct.}$$

Above we have reduced the problem to (38) when the two directions are (1, d) and (1, d'). The case when one of the two directions is (0, 1) can be handled in the same manner and can be reduced to (38) too, we omit the detail.

Finally, observe that (38) is equivalent to Lemma 45. This completes the proof.

Proof of Theorem 44-(5). Let $A = \{a_1, \ldots, a_{p-1}\}$. Observe that the set $S = \ell \cup A$ is a blocking set. Indeed, let ℓ' be a non-vertical line, then it either intersect ℓ or is parallel to ℓ , in the latter case it passes through some a_i . Hence, $\ell' \cap S \neq \emptyset$ always hold.

To show S is an MBS, we say a point $x \in S$ is *essential* if there is a non-vertical line ℓ^* passing through x and ℓ^* is disjoint from $S \setminus \{x\}$. It suffices to show every point in S is essential.

- Every a_i is essential: since there are p-1 lines parallel to ℓ , each parallel line of ℓ passes through exactly one point a_i .
- As no two of a_i are in the same vertical line, for $1 \leq i \leq p-1$, let ℓ_i denote the unique vertical line passing through a_i , and let $b_i = \ell \cap \ell_i$. Every $b_i \in \ell$ is also essential. To see this, let \mathcal{L} be the set of the p-1 lines passing through b_i that are not ℓ . Let $\ell' \in \mathcal{L}$, then $\ell' \cap (S \setminus \{b_i\}) = \ell' \cap (A \setminus \{a_i\})$. And the intersection $\ell' \cap (A \setminus \{a_i\})$ are disjoint from each for distinct $\ell' \in \mathcal{L}$. As $A \setminus \{a_i\} = p-2$, but \mathcal{L} contains p-1 lines, hence there must be a line $\ell^* \in \mathcal{L}$ such that $\ell^* \cap (A \setminus \{a_i\}) = \emptyset$ as desired.
- Since there are p-1 points b_i , there is one unique point $b \in \ell$ left. We show b is also essential. Observe that the set of p points $\{a_1, \ldots, a_{p-1}, b\}$ satisfies the condition of Lemma 46. Indeed, they are the p intersecting points of the two sets of parallel lines: the set of p vertical lines and the set of p lines in the direction of ℓ . Apply Lemma 46, there are a_i, a_j such that b, a_i, a_j are collinear. Repeating a similar argument as the previous case finishes the proof.

7 A read-k non-deterministic BP lower bound for GEN

In this section we derive an exponential lower bound for NBP_k for the GEN function via a direct reduction from the BRS function.

Definition 47. A Boolean function⁵ sequence $f = (f_n)_{n\geq 1}$ is said to be a projection of Boolean function $g = (g_n)_{n\geq 1}$, written $f \leq_{p(n)} g$, if $f(x_1, \ldots, x_n) = g_{p(n)}(y_1, \ldots, y_{p(n)})$ for some polynomial p and $y_j \in \{0, 1, x_1, \overline{x_1}, \ldots, x_n, \overline{x_n}\}$. If for every $i = 1, \ldots, n$, the number of j for which $y_j \in \{x_i, \overline{x_i}\}$ is at most t, then we denote the projection by $f \leq_{p(n)}^t g$.

Lemma 48. If $f \leq_{O(n^s)}^t g$, then $\text{NBP}_k(g_n) \geq \text{NBP}_{kt}(f_{\Omega(n^{1/s})})$.

Proof. Let P be a nondeterministic BP for g. By the definition of the projection $f \leq_{O(n^s)}^t g$, replacing the variables in P using variables for f (or constants) gives a nondeterministic BP Q for f. Hence, $|Q| \leq |P|$. Furthermore, Q will be read-(kt) if P is read-k. This implies $\text{NBP}_{kt}(f_n) \leq \text{NBP}_k(g_{O(n^s)})$. Rewriting this inequality gives the lemma.

Remark 49. By the construction in Proposition 28 one can show that, however, even $f \leq_{O(n^s)}^1 g$ (i.e., so-called read-once projection) does not imply a similar inequality as in the Observation for the measures \hat{S}, S, \hat{C} and C. That is, for example, one can have $f \leq_{O(n^s)}^1 g$ and $\hat{S}(f)$ is exponentially large but $\hat{S}(g)$ is a constant.

Theorem 50. Let $f = (f_n)_{n\geq 1}$ be a sequence of Boolean functions so that f_n has a De Morgan circuit of size $S(n) \geq n$ with the following property: no two gates have the same pair of gates as their two inputs. Then, $f \leq_{O(S(n)^2 \log S(n))}^{1} \text{GEN}$.

Proof. Fix a circuit C_n for f_n with the desired property, we define the projection from GEN as follows. Consider a set M that contains the following elements:

- n+1 elements $\$_0, \$_1, \ldots, \$_n;$
- 2 elements $(X_i, 0), (X_i, 1)$ for every $1 \le i \le n$;
- 2 elements (g, 0), (g, 1) for every gate g in the circuit C_n .

Let m = |M|, then m = n + 1 + 2n + 2S(n) = O(S(n)). To compare this with the usual definition of GEN (see definition of GEN_q in Section 2.8), we think of $\$_0$ as the element 1, and (g, 1) as the *m*-th element in M, where g is the output gate of C_n . Let $q = \frac{m(m-1)}{2} = O(S(n)^2)$. We show that f_n is a projection of GEN_q : $M^q \to \{0, 1\}$. Specifically, given an input *n*-bit string $(x_1, \ldots, x_n) \in \{0, 1\}^n$ for f_n , define the corresponding input $Y \in M^q$ for GEN_q as follows.

- (1) $\$_0 * \$_i = \$_{i+1}$ for every $0 \le i \le n-1$;
- (2) $\$_i * \$_i = (X_i, x_i)$ for every $1 \le i \le n$;
- (3) If g is an \wedge gate in C_n with two input gates h, l, then define $(h, a) * (l, b) = (g, a \wedge b)$ for every $a, b \in \{0, 1\}$, Define similarly if g is an \vee gate.
- (4) If g is a \neg gate with input gate h, then define (h, 0) * (h, 0) = (g, 1) and (h, 1) * (h, 1) = (g, 0).

⁵The domain is $\{0,1\}^n$ and the range is $\{0,1\}$.

Define $\delta * \gamma = \$_0$ for every $\delta, \gamma \in M$ where the operation $\delta * \gamma$ has not been defined above. Let the obtained variable for GEN_q be $Y \in M^q$. One can directly check that $f_n(x_1, \ldots, x_n) = \operatorname{GEN}_q(Y)$. Furthermore, by (2), each variable x_i only appears once in Y, this proves the theorem.

Corollary 51. $\text{NBP}_k(\text{GEN}_n) = 2^{\Omega(\frac{\sqrt{n}}{4^k k^3 \sqrt{\log n}})}.$

Proof. It is not hard to construct a size O(n) De Morgan circuit for BRS_n satisfying the property in Theorem 50. Hence, BRS $\leq_{O(n^2 \log n)}^{1}$ GEN. In [4] it is shown that $\text{NBP}_k(\text{BRS}_n) = 2^{\Omega(\frac{n}{4^k k^3})}$. Lemma 48 implies the desired lower bound for GEN.

8 Discussion and open problems

Although exponential lower bounds for deterministic read-once BPs have been proved more than three decades ago [31, 29], the read-once BP models still offer challenges such as (a) finding a function having small DNF and small CNF, yet having no small deterministic read-once BP, and (b) finding a Boolean function having an exponential lower bound for *semantic* nondeterministic read-once BP (we did not discuss this topic in our paper, see reference [7]). Besides, read-once BPs are also important for their connections to derandomization and proof complexity. Below we discuss questions naturally inspired from our work.

- (1) What is $\widehat{P}(\text{TEP}, h)$? By Theorem 15, an exponential lower bound for $\widehat{P}(\text{TEP}, h)$ would give an alternative proof, besides [16], for (TEP, h) having exponential deterministic BP₁ lower bound. The reason to pursue an alternative proof is the hope that it might be generalized to read-k BP. Indeed, $\widehat{\mathcal{P}}_k$ as a natural generalization of \widehat{P} is a lower bound for BP_k as discussed in Section 3.4. Similarly, what is $\widehat{C}(\text{TEP}, h)$? As well, we have not been able to solve S(TEP, h), despite the fact that the only immediate implication of an S lower bound is a lower bound on the size of OBDDs, a very weak BP model.
- (2) What is CC(TEP, h)? An easy upper bound is $CC(TEP, h) = O(h \log k)$ following from Theorem 34. Referring to Figure 1, note that a lower bound for CC(TEP, h) does not imply lower bounds for $\hat{P}(TEP, h)$ or $\hat{C}(TEP, h)$, or vice versa. Nonetheless, studying the communication complexity of TEP seems to be an interesting question itself.
- (3) In view of Theorem 39, does there exist an *r*-regular and $K_{2,s}$ -free bipartite graph G, such that $\widehat{S}(\text{GAL}_G)$ is large but $\widehat{C}(\text{GAL}_G)$ is small, for appropriate parameters r and s? This means that for the purpose of tackling the BP₁ vs weight problem there is no reason to restrict oneself to the graph G_q . Of course, determining $\widehat{C}(\text{GAL}_{G_q})$ is still an interesting problem for NBP₁(GAL_{G_q}).
- (4) The **Question 1** and **Question 2** asked in Section 6.2. Question 2 is a typical question in studying the standard blocking sets in either projective or affine spaces. In contrast, the number of blocking sets satisfying certain conditions (i.e., Question 1) receives little attention (see [14, 27]). Our work provides a strong motivation for it. Yet, to fully classify blocking sets in \mathbb{F}_p^2 in order to help answer Question 1 seems an daunting task. Is it possible to solve Question 1 without answering Question 2, and in general, is it possible to determine $\widehat{C}(\text{GAL}_{G_q})$ without answering Question 2? This again points to the direction that, if one is only interested in complexity applications, perhaps it will be useful to try other graphs besides G_q . Having

said that, studying the number and structures of (minimal) blocking sets in \mathbb{F}_p^2 are interesting mathematical problems on their own, as we try to demonstrate in Section 6.3.

(5) By Proposition 28 and Figure 1, none of the measures (such as S, \hat{C} , etc) defined in the paper lie in-between BP₁ and circuit size. In view of derandomization, it could be beneficial to have a sequence of measures, say m_1, m_2, \ldots , such that BP₁ $\geq m_1 \geq m_2 \geq \ldots \geq$ circuit size. This in theory might allow a progressive way of adapting techniques for derandomizing space (i.e., BP₁), where things are better understood, to derandomizing time (i.e., circuit size). For example, a typical question could be to construct pseudo random generators for Boolean functions f satisfying $m_i(f) \leq O(n^2)$.

Acknowledgement

The authors acknowledge support from the Natural Sciences and Engineering Research Council of Canada, P.M. as holder of discovery grant RGPIN-04500 and Y.L. as postdoctoral collaborator. Y.L. acknowledges the support from University of Montréal where this work was done when Y.L. was a postdoctoral researcher there.

References

- Paul Beame, Nathan Grosshans, Pierre McKenzie, and Luc Segoufin. Nondeterminism and an abstract formulation of Nečiporuk's lower bound method. ACM Trans. Comput. Theory, 9(1):Art. 5, 34, 2016.
- [2] A Blokhuis. Combinatorial problems in finite geometry and lacunary polynomials. In Proceedings International Congress of Mathematicians (ICM 2002, Beijing, China, August 20-28, 2002), Volume III: Invited lectures, pages 537–545. Higher Education Press, 2002.
- [3] Beate Bollig and Ingo Wegener. A very simple function that requires exponential size read-once branching programs. *Information Processing Letters*, 66(2):53–57, 1998.
- [4] Allan Borodin, Alexander Razborov, and Roman Smolensky. On lower bounds for read-k-times branching programs. *Computational Complexity*, 3(1):1–18, 1993.
- [5] Andries E Brouwer and Alexander Schrijver. The blocking number of an affine space. Journal of Combinatorial Theory, Series A, 24(2):251–253, 1978.
- [6] Gil Cohen, Dean Doron, Oren Renard, Ori Sberlo, and Amnon Ta-Shma. Error reduction for weighted prgs against read once branching programs. In Valentine Kabanets, editor, 36th Computational Complexity Conference, CCC 2021, July 20-23, 2021, Toronto, Ontario, Canada (Virtual Conference), volume 200 of LIPIcs, pages 22:1–22:17. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [7] Stephen Cook, Jeff Edmonds, Venkatesh Medabalimi, and Toniann Pitassi. Lower bounds for nondeterministic semantic read-once branching programs. In 43rd International Colloquium on Automata, Languages, and Programming (ICALP 2016). Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2016.

- [8] Stephen Cook, Pierre McKenzie, Dustin Wehr, Mark Braverman, and Rahul Santhanam. Pebbles and branching programs for tree evaluation. ACM Transactions on Computation Theory (TOCT), 3(2):1–43, 2012.
- [9] Andrzej Ehrenfeucht and David Haussler. Learning decision trees from random examples. Information and Computation, 82(3):231–246, 1989.
- [10] Michael A Forbes and Zander Kelley. Pseudorandom generators for read-once branching programs, in any order. In 2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS), pages 946–955. IEEE, 2018.
- [11] Anna Gál. A simple function that requires exponential size read-once branching programs. Information Processing Letters, 62(1):13–16, 1997.
- [12] Ludmila Glinskih and Dmitry Itsykson. Satisfiable tseitin formulas are hard for nondeterministic read-once branching programs. In 42nd International Symposium on Mathematical Foundations of Computer Science (MFCS 2017). Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2017.
- [13] Ludmila Glinskih and Dmitry Itsykson. On tseitin formulas, read-once branching programs and treewidth. *Theory of Computing Systems*, pages 1–21, 2020.
- [14] JWP Hirschfeld. Projective geometries over finite fields. Oxford mathematical monographs. Oxford University Press New York, 1998.
- [15] Dmitry Itsykson, Alexander Knop, Andrei Romashchenko, and Dmitry Sokolov. On obddbased algorithms and proof systems that dynamically change order of variables. *The Journal* of Symbolic Logic, pages 1–41, 2020.
- [16] Kazuo Iwama and Atsuki Nagao. Read-once branching programs for tree evaluation problems. ACM Transactions on Computation Theory (TOCT), 11(1):1–12, 2018.
- [17] Neil D. Jones and William T. Laaser. Complete problems for deterministic polynomial time. *Theor. Comput. Sci.*, 3(1):105–117, 1976.
- [18] Stasys Jukna. Boolean function complexity: advances and frontiers, volume 27. Springer Science & Business Media, 2012.
- [19] Stasys Jukna, A Razborov, P Savicky, and Ingo Wegener. On p versus np ∩ co-np for decision trees and read-once branching programs. *Computational Complexity*, 8(4):357–370, 1999.
- [20] Stasys P Jukna. Entropy of contact circuits and lower bounds on their complexity. Theoretical Computer Science, 57(1):113–129, 1988.
- [21] Eyal Kushilevitz and Noam Nisan. Communication complexity, 1997.
- [22] Anup Rao and Amir Yehudayoff. *Communication Complexity: and Applications*. Cambridge University Press, 2020.
- [23] Igor Razgon. On obdds for cnfs of bounded treewidth. In Proceedings of the Fourteenth International Conference on Principles of Knowledge Representation and Reasoning, pages 92–100, 2014.

- [24] Martin Sauerhoff. Approximation of boolean functions by combinatorial rectangles. Theoretical computer science, 301(1-3):45–78, 2003.
- [25] Janos Simon and Mario Szegedy. A new lower bound theorem for read-only-once branching programs and its applications. In Advances in Computational Complexity Theory, pages 183– 193, 1990.
- [26] Anastasia Sofronova and Dmitry Sokolov. Branching programs with bounded repetitions and flow formulas. In Valentine Kabanets, editor, 36th Computational Complexity Conference, CCC 2021, July 20-23, 2021, Toronto, Ontario, Canada (Virtual Conference), volume 200 of LIPIcs, pages 17:1–17:25. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [27] Tamás Szőnyi, András Gács, and Zsuzsa Weiner. On the spectrum of minimal blocking sets in pg (2,q). Journal of Geometry, 76(1-2):256–281, 2003.
- [28] Ingo Wegener. The complexity of Boolean functions. BG Teubner, 1987.
- [29] Ingo Wegener. On the complexity of branching programs and decision trees for clique functions. Journal of the ACM (JACM), 35(2):461–471, 1988.
- [30] Ingo Wegener. Branching programs and binary decision diagrams: theory and applications. SIAM, 2000.
- [31] Stanislav Žák. An exponential lower bound for one-time-only branching programs. In International Symposium on Mathematical Foundations of Computer Science, pages 562–566. Springer, 1984.

A The proof for Theorem 34

In this appendix, we provide calculation and proof for Theorem 34. We will prove each part of Theorem 34 separately, as in Theorem 57, Theorem 58 and Theorem 59, in the following.

We use the following notation. Let (M, L, R) denote a partition of the input variables for (TEP, h) where M corresponds to the root matrix, L and R correspond to the left and right child, respectively. Note that both L and R correspond to inputs for (TEP, h - 1). Recall n_h denotes the input size for (TEP, h). For a subset $A \subseteq [n_h]$, we think of A as a subset of input variables for (TEP, h) and write $A = (A_M, A_L, A_R)$, where $A_M = A \cap M$, $A_L = A \cap L$ and $A_R = A \cap R$. For notational simplicity, when the parameter h is clear from the context, we use TEP_A to denote the matrix $(\text{TEP}, h)_A$, and for $\alpha \in [k]^A$, we use TEP_α to denote the subfunction $(\text{TEP}, h)_{A,\alpha}$.

By an abuse of notation, sometimes we use $M_{ij} \in A_M$ or $(i, j) \in A_M$, depending on which one is more convenient in the context, to mean that A_M contains the variable at entry (i, j) of the root matrix M.

Let $A \subseteq [n_h]$. Let $\alpha, \alpha' \in [k]^A$. We say $\alpha \sim \alpha'$ if $(\text{TEP}, h)_{A,\alpha} = (\text{TEP}, h)_{A,\alpha'}$. Obviously, \sim is an equivalence relation. We use the notation $\langle \alpha \rangle$ to denote the equivalence class represented by α .

A function is said to be *non-constant* if it evaluates to at least two distinct values. A subfunction $(\text{TEP}, h)_{A,\alpha}$ is a *full-range* function if for every $r \in [k]$ there exists $\beta \in [k]^{[n_h]\setminus A}$ satisfying $(\text{TEP}, h)_{A,\alpha}(\beta) = r$.

Lemma 52. Let $h \ge 1$. For any $A \subseteq [n_h]$ of size $|A| \le n_h - 1$, there exists $\alpha^* \in [k]^A$ such that $(\text{TEP}, h)_{A,\alpha^*}$ is a full-range function.

Proof. We use induction on h. The base case h = 1 is clear. Assume the lemma is true for h - 1. Consider (TEP, h). Let $A = (A_M, A_L, A_R)$ satisfy $|A| \le n_h - 1$.

- $A_L \neq L$, i.e., $|A_L| \leq n_{h-1} 1$. By induction hypothesis on h-1, there exists $\alpha_L^* \in [k]^{A_L}$ such that $(\text{TEP}, h-1)_{A_L, \alpha_L^*}$ is a full-range function. Fix $\alpha_R^* \in [k]^{A_R}$ and $\beta_R \in [k]^{R-A_R}$ arbitrarily. Suppose $(\text{TEP}, h-1)_{A_R, \alpha_R^*}(\beta_R) = j$ for some $j \in [k]$. Fix $\alpha_M^* \in [k]^{A_M}$ and $\beta_M \in [k]^{M-A_M}$ such that the *j*-th column of the matrix M is the vector $(1, 2, ..., k)^T$. For every $r \in [k]$, there exists $\beta_L \in [k]^{L-A_L}$ such that $(\text{TEP}, h-1)_{A_L, \alpha_L^*}(\beta_L) = r$. Set $\alpha^* = (\alpha_M^*, \alpha_L^*, \alpha_R^*)$ and $\beta = (\beta_M, \beta_L, \beta_R)$. Then, $(\text{TEP}, h)_{A, \alpha^*}(\beta) = r$.
- $A_R \neq R$. This is symmetric to the previous case.
- $A_L = L$ and $A_R = R$ but $A_M \neq M$. Choose α_L^* and α_R^* such that $((\text{TEP}, h-1)(\alpha_L^*), (\text{TEP}, h-1)(\alpha_R^*)) = (i, j) \notin A_M$. Set α_M^* arbitrarily. Let $\alpha^* = (\alpha_M^*, \alpha_L^*, \alpha_R^*)$. It is easy to see that $(\text{TEP}, h)_{A,\alpha^*}$ is a full-range function.

Lemma 53. Let $h \geq 2$, $A \subseteq [n_h]$. Suppose $A = (\emptyset, A_L, A_R)$. Let $\alpha = (\alpha_L, \alpha_R) \in [k]^A = [k]^{A_L} \times [k]^{A_R}$, $\alpha' = (\alpha'_L, \alpha'_R) \in [k]^A = [k]^{A_L} \times [k]^{A_R}$. Then,

$$(\text{TEP}, h)_{A,\alpha} = (\text{TEP}, h)_{A,\alpha'}$$
$$\iff \left((\text{TEP}, h-1)_{A_L,\alpha_L}, (\text{TEP}, h-1)_{A_R,\alpha_R} \right) = \left((\text{TEP}, h-1)_{A_L,\alpha'_L}, (\text{TEP}, h-1)_{A_R,\alpha'_R} \right).$$
(39)

Proof. The direction \Leftarrow . Obvious.

The direction \implies . Assume for the sake of a contradiction the implication is not true. Without loss of generality we may assume $(\text{TEP}, h - 1)_{A_L,\alpha_L} \neq (\text{TEP}, h - 1)_{A_L,\alpha'_I}$. We will show

 $(\text{TEP}, h-1)_{A_L, \alpha_L} \neq (\text{TEP}, h-1)_{A_L, \alpha'_L} \Longrightarrow (\text{TEP}, h)_{A, \alpha} \neq (\text{TEP}, h)_{A, \alpha'}.$

Indeed, let $\beta_L \in [k]^{L-A_L}$ be such that $(\text{TEP}, h-1)_{A_L,\alpha_L}(\beta_L) = i \neq i' = (\text{TEP}, h-1)_{A_L,\alpha'_L}(\gamma_L)$. Then, set $\beta_M \in [k]^M$ to be such that every entry in the *i*-th row equals to *i*, and every entry in the *i'*-th row equals to *i'*. Choose $\beta_R \in [k]^{R-A_R}$ arbitrarily. Set $\beta = (\beta_M, \beta_L, \beta_R)$. Then, $(\text{TEP}, h)_{A,\alpha}(\beta) = i \neq i' = (\text{TEP}, h)_{A,\alpha'}(\beta)$.

Lemma 54. Let $h \ge 2$. Let $A = (A_M, A_L, A_R) \subseteq [n_h]$. If $|A_L|, |A_R| \le n_{h-1}-1$, then $S((\text{TEP}, h)_A) \ge k^{|A_M|}$.

Proof. Since $|A_L| \leq n_{h-1} - 1$, Lemma 52 implies that there exists $\alpha_L^* \in [k]^{A_L}$ such that $(\text{TEP}, h - 1)_{A_L,\alpha_L^*}$ is a full-range function. Similarly, let $\alpha_R^* \in [k]^{A_R}$ be such that $(\text{TEP}, h - 1)_{A_R,\alpha_R^*}$ is a full-range function. Consider the set

$$\Omega = \{ \alpha = (\alpha_M, \alpha_L^*, \alpha_R^*) : \alpha_M \in [k]^{A_M} \}.$$

We claim that

$$\alpha, \alpha' \in \Omega, \alpha \neq \alpha' \Longrightarrow (\text{TEP}, h)_{A, \alpha} \neq (\text{TEP}, h)_{A, \alpha'}$$

This implies $S((\text{TEP}, h)_A) \ge |\Omega| = k^{|A_M|}$.

To show the claim, suppose $\alpha_M(i,j) \neq \alpha'_M(i,j)$ for some $(i,j) \in A_M$. By the choice of α_L^* and α_R^* , there exist $(\beta_L, \beta_R) \in [k]^{L-A_L} \times [k]^{R-A_R}$ such that $(\text{TEP}, h-1)_{A_L,\alpha_L^*}(\beta_L) = i$ and $(\text{TEP}, h-1)_{A_L,\alpha_R^*}(\beta_R) = j$. Let $\beta = (\beta_M, \beta_L, \beta_R)$ where $\beta_M \in [k]^{M-A_M}$ is chosen arbitrarily. Then

$$(\text{TEP}, h)_{A,\alpha}(\beta) = \alpha_M(i, j) \neq \alpha'_M(i, j) = (\text{TEP}, h)_{A,\alpha'}(\beta)$$

as claimed.

Lemma 55. Let $h \ge 2$. Let $A = (A_M, L, A_R)$, i.e., $A_L = L$. Suppose $|A_R| \le n_{h-1} - 1$. For $i \in [k]$, let r_i denote the number of entries of A_M in row i.

- (1) If $r_i = k$ for some *i*, then $S((\text{TEP}, h)_A) \ge k^k$.
- (2) If $r_i < k$ for every $i \in [k]$, then $S((\text{TEP}, h)_A) \ge \sum_{i=1}^k k^{r_i}$.

Proof. For $i \in [k]$, let $\langle i \rangle$ denote $\langle \alpha_L \rangle$ for which $(\text{TEP}, h - 1)(\alpha_L) = i$. Lemma 52 implies the existence of $\alpha_R^* \in [k]^{A_R}$ such that $(\text{TEP}, h - 1)_{A_R, \alpha_R^*}$ is a full-range function.

Claim: There are exactly k^{r_i} distinct subfunctions $(\text{TEP}, h)_{A,\alpha}$ for which α is of the form $\alpha = (\alpha_M, \langle i \rangle, \alpha_R^*) \in [k]^A$.

Proof of the Claim: Let $\alpha' = (\alpha'_M, \langle i \rangle, \alpha^*_R) \in [k]^A$. Suppose α_M and α'_M differ on row *i*, i.e., $\alpha_M(i, j) \neq \alpha'_M(i, j)$ for some *j* where $(i, j) \in A_M$. It suffices to show (TEP, $h)_{A,\alpha} \neq$ (TEP, $h)_{A,\alpha'}$. Indeed, since (TEP, $h - 1)_{A_R,\alpha^*_R}$ is a full-range function, there exists $\beta_R \in [k]^{R-A_R}$ such that $(\text{TEP}, h - 1)_{A_R,\alpha^*_R}(\beta_R) = j$. Let $\beta = (\beta_M, \beta_R) \in [k]^{A-A_M} \times [k]^{R-A_R}$ where β_M is chosen arbitrarily. Then,

$$(\text{TEP}, h)_{A,\alpha}(\beta) = \alpha_M(i, j) \neq \alpha'_M(i, j) = (\text{TEP}, h)_{A,\alpha'}(\beta).$$

Hence, $(\text{TEP}, h)_{A,\alpha} \neq (\text{TEP}, h)_{A,\alpha'}$. Since the entries in A_M that are not in row *i* are irrelevant for the subfunction $(\text{TEP}, h)_{A,\alpha}$ for which α is of the form defined before, the number of such subfunctions is equal to k^t .

We proceed to prove the Lemma.

- (1) This follows directly from the Claim.
- (2) By the Claim, it suffices to show subfunctions given by $\alpha = (\alpha_M, \langle i \rangle, \alpha_R^*)$ are distinct for distinct $i \in [k]$.

Let $i, i' \in [k]$ and $i \neq i'$. Consider $\langle i \rangle$ and $\langle i' \rangle$. Let $\alpha = (\alpha_M, \langle i \rangle, \alpha_R^*)$ and $\alpha' = (\alpha'_M, \langle i' \rangle, \alpha_R^*)$. We show (TEP, h)_{$A,\alpha'} <math>\neq$ (TEP, h)_{A,α'}. Since $r_i < k$, there exists $j \in [k]$ such that $(i, j) \notin A_M$. Choose β_R as before such that (TEP, h - 1)_{A_R,α_R^*} $(\beta_R) = j$.</sub>

- Case 1: $(i', j) \notin A_M$. Choose $\beta_M \in [k]^{M-A_M}$ such that $\beta_M(i, j) = 1$ and $\beta_M(i', j) = 2$. Set $\beta = (\beta_M, \beta_R)$. Then, (TEP, $h)_{A,\alpha}(\beta) = 1$ but (TEP, $h)_{A,\alpha'}(\beta) = 2$.
- Case 2: $(i', j) \in A_M$. Choose $\beta_M \in [k]^{M-A_M}$ such that $\beta_M(i, j) \neq \alpha'_M(i', j)$. Set $\beta = (\beta_M, \beta_R)$. Then, (TEP, $h)_{A,\alpha}(\beta) = \beta_M(i, j) \neq \alpha'_M(i', j) = (\text{TEP}, h)_{A,\alpha'}(\beta)$.

In both cases, we have $(\text{TEP}, h)_{A,\alpha} \neq (\text{TEP}, h)_{A,\alpha'}$ as desired.

Lemma 56. Let $h \ge 2$. Let $A = (A_M, A_L, A_R)$. If $|A_M| \le k$, then $S((\text{TEP}, h)_A) \ge S((\text{TEP}, h - 1)_{A_L}) \cdot S((\text{TEP}, h - 1)_{A_R})$. Furthermore, if $|A_M| = 0$, then the equality holds.

Proof. The "Furthermore" part follows from Lemma 53.

To show the inequality, choose $\alpha_M^* \in [k]^{A_M}$ such that α_M^* assigns distinct values for entries in A_M . This is possible because $|A_M| \leq k$. Let $(\alpha_L, \alpha_R) \in [k]^{A_L} \times [k]^{A_R}$. It suffices to show that each different pair $(\langle \alpha_L \rangle, \langle \alpha_R \rangle)$ gives rise to a different subfunction (TEP, $h)_{A,\alpha}$ where $\alpha = (\alpha_M^*, \alpha_L, \alpha_R)$. Note that the number of distinct pairs $(\langle \alpha_L \rangle, \langle \alpha_R \rangle)$ is exactly the desired lower bound.

To verify the claim, consider two distinct pairs

$$(\langle \alpha_L \rangle, \langle \alpha_R \rangle) \neq (\langle \alpha'_L \rangle, \langle \alpha'_R \rangle).$$
(40)

Let $\alpha = (\alpha_M^*, \langle \alpha_L \rangle, \langle \alpha_R \rangle)$ and $\alpha' = (\alpha_M^*, \langle \alpha'_L \rangle, \langle \alpha'_R \rangle)$. The assumption (40) implies that there exists $(\beta_L, \beta_R) \in [k]^{L-A_L} \times [k]^{R-A_R}$ such that

$$\left((\text{TEP}, h-1)_{A_L,\alpha_L}(\beta_L), (\text{TEP}, h-1)_{A_R,\alpha_R}(\beta_R) \right)$$

= $(i,j) \neq (i',j') = \left((\text{TEP}, h-1)_{A_L,\alpha'_L}(\beta_L), (\text{TEP}, h-1)_{A_R,\alpha'_R}(\beta_R) \right).$

By the choice of α_M^* , it is easy to see that there exists $\beta_M \in [k]^{M-A_M}$ such that $(\text{TEP}, h)_{A,\alpha}(\beta) \neq (\text{TEP}, h)_{A,\alpha'}(\beta)$ for $\beta = (\beta_M, \beta_L, \beta_R)$.

Let $S((\text{TEP}, h), \ell) = \min_{A \subseteq [n_h], |A| = \ell} S((\text{TEP}, h)_A)$. Then, $S(\text{TEP}, h) = \max_{1 \le \ell \le n_h} S((\text{TEP}, h), \ell)$.

Theorem 57. Let $1 \le \ell \le n_2$ be an integer. Let $c \in [0,1)$. Define $V_c = k+2$ and $W_c = ck^2$.

(1) For $1 \le \ell \le k+1$, $S((\text{TEP}, 2), \ell) = S((\text{TEP}, 2), n_2 - (k-1)(\ell-1))$.

(2) Let $c \in [0,1)$. If $V_c \leq \ell \leq W_c$, then $S((\text{TEP},2),\ell) \geq (1-c)k^2$. In particular,

$$S(\text{TEP}, 2) = \max_{1 \le \ell \le k+1} S((\text{TEP}, 2), \ell) = \max_{k+2 \le \ell \le n_2} S((\text{TEP}, 2), \ell) = k^2,$$

and is achieved at either $\ell = k + 1$ or $\ell = k + 2$.

Proof. Recall $n_2 = k^2 + 2$. Assume the calculation of $S((\text{TEP}, 2)_A)$ in Table 2 is correct. For every $1 \le \ell \le n_2$, in Table 3 we give a minimizer $A^* \subseteq [n_2]$ of size $|A^*| = \ell$ such that $S((\text{TEP}, 2)_{A^*}) = S((\text{TEP}, 2), \ell)$. The minimizer is obtained from Table 2. The theorem follows by a simple calculation using Table 3.

We now prove the calculation of $S((\text{TEP}, 2)_A)$ in Table 2 is correct. Let $|A| = \ell$. There are four cases.

- $A = (A_M, \emptyset, \emptyset)$. The upper bound is trivial, the lower bound follows from Lemma 54.
- $A = (A_M, x, \emptyset).$

Consider the case $|A_M| = \ell - 1 > (k - 1)k$ first. This implies that A_M must contain all the k entries for some row. Then, Lemma 55 implies $S((\text{TEP}, 2)_A) \ge k^k$

Now assume $|A_M| = \ell - 1 \le (k-1)k$. Observe that this implies $qk^{p+1} + (k-q)k^p \le k^k$ where $\ell - 1 = pk + q$.

We consider two cases depending on the choice of A_M .

- Case 1: A_M contains all the k entries of some row. By Lemma 55, $S((\text{TEP}, 2)_A) \ge k^k$.
- Case 2: A_M contains at most k-1 entries from each row. Then, Lemma 55 implies that to minimize $S((\text{TEP}, 2)_A)$ the A_M should be chosen greedily according to the columns, as described in Table 2. For this choice of A_M , Lemma 55 implies $S((\text{TEP}, 2)_A) = qk^{p+1} + (k-q)k^p \leq k^k$.

The above two cases imply that the optimal choice for A_M is as described in Table 2.

- $A = (\emptyset, x, y)$. Obvious.
- $A = (A_M, x, y)$. Let $\alpha = (\alpha_M, i, j) \in [k]^A$.

- Case 1: $(i, j) \in A_M$. In this case, for all $\gamma \in [k]^{\overline{A}}$, one has

$$(\text{TEP}, 2)_{A,\alpha}(\gamma) = \alpha_M(i, j), \tag{41}$$

i.e., $(\text{TEP}, 2)_{A,\alpha}$ is a constant function that is identically equal to $\alpha_M(i, j)$. Trivially, there are k distinct constant functions. Note that $A_M \neq \emptyset$ because $\ell \geq 3$, Hence, by (41), each constant function can be achieved by some $\alpha \in [k]^A$.

- Case 2: $(i, j) \notin A_M$. Let $\gamma \in [k]^{\overline{A}}$. Then, $(\text{TEP}, 2)_{A,\alpha}(\gamma) = \gamma(i, j)$. Let $\alpha' = (\alpha'_M, i', j') \in [k]^A$ such that $(i', j') \notin A_M$. Then, it is easy to see that $(\text{TEP}, 2)_{A,\alpha} = (\text{TEP}, 2)_{A,\alpha'}$ is equivalent to (i, j) = (i', j'). In other words, every pair $(i, j) \notin A_M$ defines a distinct subfunction, and no more. Hence, there are $k^2 - |A_M|$ such subfunctions.

To summarize the two cases, we get that $S((\text{TEP}, 2)_A) = k + (k^2 - |A_M|) = k^2 + k + 2 - \ell$.

Note that the case $A = (A_M, \emptyset, y)$ is symmetric to the case $A = (A_M, x, \emptyset)$. Hence, we have verified all possible cases.

Table 2: The calculation for $S((\text{TEP}, 2)_A)$, categorized according to the pattern of $A = (A_M, A_L, A_R)$. The variables x and y denote the left and right leaf, respectively. For each pattern, only the one that minimizes $S((\text{TEP}, 2)_A)$ under this pattern is given.

	$(A_M, \emptyset, \emptyset)$	(A_M, x, \emptyset)	(A_M, x, \emptyset)	(\emptyset, x, y)	(A_M, x, y)
$ A = \ell$	$ A_M = \ell$	$ A_M + 1 = \ell$ $\ell - 1 \le (k - 1)k$ $\ell - 1 = pk + q, 0 \le q < k$	$ A_M + 1 = \ell$ $\ell - 1 > (k - 1)k$	$\ell=2$	$ A_M + 2 = \ell$ $\ell \ge 3$
Choice of A_M	arbitrary	The first p columns plus the first q entries in the $(p + 1)$ -th column	n/a	irrelevant	arbitrary
$S((\text{TEP}, 2)_A) = ?$	k^ℓ	$qk^{p+1} + (k-q)k^p$	$\geq k^k$	k^2	$k^2 + k + 2 - \ell$

Table 3: The minimizer A^* for $\min_{|A|=\ell} S((\text{TEP}, 2)_A)$.

	$1 \leq \ell \leq k+1$	$k+2 \le \ell \le n_2$	
$A^{*} = ?$	(A_M, x, \emptyset)	(A_M, x, y)	
$S((\text{TEP},2)_{A^*}) =?$	$(\ell - 1)(k - 1) + k$ = $k^2 - (k - 1)(k + 1 - \ell)$	$k^{2} + k + 2 - \ell = k^{2} - (\ell - (k + 2))$	

Theorem 58. Let $1 \le \ell \le n_3$ be an integer. Let $c \in [0,1)$. Define $V_c = k+2$ and $W_c = ck^2$.

- (1) $S(\text{TEP},3) \le k(k^2 k + 2) \le k^3$. In particular, $S(\text{TEP},3) < k^3$ for $k \ge 3$.
- (2) Let $k \ge 2/(1-c)$. If $n_2 + V_c + 3k/2 \le \ell \le n_2 + W_c$, then $S((\text{TEP}, 3), \ell) \ge (1-c)k^3/8$. In particular, $S(\text{TEP}, 3) \ge (1-c)k^3/8$.

Proof. (1) To show the upper bound, it suffices to show for every $1 \le \ell \le n_3$, there exists $A \subseteq [n_3]$ of size $|A| = \ell$ such that $S((\text{TEP}, 3)_A) \le k(k^2 - k + 2)$. Let $A = (A_M, A_L, A_R)$. We consider the following cases.

- (i) $2n_2 < \ell \leq n_3$. Choose $A = (A_M, L, R)$. Observe that this choice reduces the problem to that height h = 2 case corresponding to the pattern (A_M, x, y) in Table 2. Hence, by Table 2, $S((\text{TEP}, 3)_A) \leq k^2 + k 1$.
- (ii) $n_2 + k^2 \leq \ell \leq 2n_2$. Equivalently, $\ell = n_2 + k^2$ or $n_2 + k^2 + 1$ or $n_2 + k^2 + 2$. Choose $A = (\emptyset, L, A_R)$. Then, $|A_R| = k^2$ or $k^2 + 1$ or $k^2 + 2$. By Lemma 56 and Table 3,

$$S((\text{TEP}, 3)_A) = S((\text{TEP}, 2)_L) \cdot S((\text{TEP}, 2)_{A_B}) \le k(k+2),$$

where the upper bound is achieved at $|A_R| = k^2$.

(iii) $n_2 + 2k \le \ell < n_2 + k^2$. Choose $A = (\emptyset, L, A_R)$. Then, $2k \le |A_R| < k^2$. By Lemma 56 and Table 3

$$S((\text{TEP},3)_A) = S((\text{TEP},2)_L) \cdot S((\text{TEP},2)_{A_B}) \le k(k^2 - k + 2),$$

where the upper bound is achieved at $|A_R| = 2k$.

- (iv) $n_2 < \ell \le n_2 + 2k 1$. Choose $A = (A_M, L, \emptyset)$. Observe that this choice reduces the problem to height h = 2 case corresponding to the pattern (A_M, x, \emptyset) in Table 2. Also $1 \le |A_M| = \ell n_2 \le 2k 1$. By Table 2, there exists A_M , such that $S((\text{TEP}, 3)_A) \le (k 1)k^2 + k = k(k^2 k + 1)$, where the upper bound is achieved at $|A_M| = 2k 1$.
- (v) $1 \leq \ell \leq n_2$. Choose $A = (\emptyset, A_L, \emptyset)$. By Lemma 56 and Theorem 57, $S((\text{TEP}, 3)_A) = S((\text{TEP}, 2)_{A_L}) \leq k^2$.

To sum up, in all cases there exists A such that $S((\text{TEP}, 3)_A) \leq k(k^2 - k + 2)$ as desired.

(2) Let $A = (A_M, A_L, A_R) \subseteq [n_3]$ be such that $|A_M| + |A_R| + |A_R| = \ell$ where $n_2 + V_c + 3k/2 \leq \ell \leq n_2 + W_c$. We show $S((\text{TEP}, 3)_A) \geq \Omega(k^3)$.

(i) $|A_M| \leq k$. Let $|A_M| = \ell'$. Then, $0 \leq \ell' \leq k$. Let $a = |A_L|$ and $b = |A_R|$. Hence,

$$a+b = \ell - |A_M| \ge n_2 + V_c + 3k/2 - \ell' \ge n_2 + V_c + k/2.$$

Since $a, b \leq n_2$, one has $\min\{a, b\} \geq V_c + k/2 \geq V_c$. Hence, by Table 3, $S((\text{TEP}, 2), a) = k^2 + k + 2 - a$ and $S((\text{TEP}, 2), b) = k^2 + k + 2 - b$. Since $a + b = \ell - |A_M| \leq n_2 + W_c$, by Lemma 56,

$$S((\text{TEP}, 3)_A) \ge S((\text{TEP}, 2)_{A_L}) \cdot S((\text{TEP}, 2)_{A_R})$$
$$\ge S((\text{TEP}, 2), a) \cdot S((\text{TEP}, 2), b)$$
$$\ge k \cdot (1 - c)k^2 = (1 - c)k^3$$

minimized at $a = n_2, b = W_c$.

- (ii) $|A_M| \ge k+1$ and $|A_L|, |A_R| \le n_2 1$. By Lemma 54, $S((\text{TEP}, 3)_A) \ge k^{|A_M|} \ge k^{k+1}$.
- (iii) $|A_M| \ge k+1$ and $|A_L| = n_2$.

For every $i \in [k]$, let r_i denote the number of entries of A_M in row i.

• $r_i \ge 3$ for some *i*. By Lemma 55, $S((\text{TEP}, 3)_A) \ge k^3$.

• $r_i \leq 2$ for every $i \in [k]$, and $|A_M| \geq 3k/2$. For t = 0, 1, 2, let $p_t = |\{r_i : r_i = t\}|$. Then,

$$p_0 + p_1 + p_2 = k$$
, $p_1 + 2p_2 = |A_M| \ge 3k/2$.

Hence, $k + p_2 \ge (p_1 + p_2) + p_2 \ge 3k/2$, this implies $p_2 \ge k/2$. By Lemma 55,

$$S((\text{TEP}, 3)_A) \ge \sum_{i=1}^k k^{r_i} \ge p_2 k^2 \ge k^3/2$$

• $r_i \leq 2$ for every $i \in [k]$, and $k+1 \leq |A_M| < 3k/2$. Hence, $V_c \leq |A_R| = \ell - n_2 - |A_M| \leq W_c$. Use the notation p_0, p_1, p_2 from above. Let

$$G(A_M) = \{ i \in [k] : r_i = 0, 1 \}.$$

Then, $|G(A_M)| = p_0 + p_1 \ge k/4$. Since $A_L = L$, we use the notation $\langle i \rangle$ to denote $\langle \alpha_L \rangle$ for which (TEP, L)(α_L) = i. Consider the set of pairs

 $\{(\langle i \rangle, \langle \alpha_R \rangle) : i \in G(A_M), (\text{TEP}, 2)_{A_R, \alpha_R} \text{ is a non-constant function}\}.$

Then, we claim that each such different pair defines a distinct subfunction $(\text{TEP}, 3)_{A,\alpha}$ for $\alpha = (\alpha_M^*, \langle i \rangle, \langle \alpha_R \rangle)$ where α_M^* can be appropriately chosen. Indeed, since $i \in G(A_M)$ and $(\text{TEP}, 2)_{A_R,\alpha_R}$ is a non-constant function, there exists $\beta_R \in [k]^{R-A_R}$ such that $(i, (\text{TEP}, 2)_{A_R,\alpha_R}(\beta_R)) \notin A_M$. Then, it is not hard to see that each such pair would define a distinct subfunction. By Theorem 57, the number of such pairs is

$$|G(A_M)| \cdot \left(S((\text{TEP}, 2)_{A_R}) - k\right) \ge k/4 \cdot \left((1 - c)k^2 - k\right) \ge (1 - c)k^3/8,$$

as long as $k \ge 2/(1-c)$.

To sum up, if $k \ge 2/(1-c)$, then in all cases we have shown $S((\text{TEP},3),\ell) \ge (1-c)k^3/8$ as desired.

Theorem 59. For every $h \ge 3$,

$$S((\text{TEP}, h), \ell) \leq \begin{cases} k^{h-1}, & 1 \leq \ell \leq \sum_{i=2}^{h-1} n_i, \\ k^{h-2}(k^2 - k + 2), & \sum_{i=2}^{h-1} n_i < \ell < \sum_{i=2}^{h-1} n_i + k^2, \\ k^{h-2}(k+2), & \sum_{i=2}^{h-1} n_i + k^2 \leq \ell \leq 2n_{h-1}, \\ k^2 + k - 1, & 2n_{h-1} < \ell \leq n_h. \end{cases}$$

In particular, $S(\text{TEP}, h) \le k^h - k^{h-2}(k-2) < k^h$ when $k \ge 3$.

Proof. We use induction on h. The base case h = 3 has been proved in Theorem 58. Assume now $h \ge 4$ and the claim is true for h - 1. Let $1 \le \ell \le n_h$. Consider $S((\text{TEP}, h), \ell)$.

(i) $1 \le \ell \le \sum_{i=2}^{h-1} n_i$. When $1 \le \ell \le n_{h-1}$, choose $A = (\emptyset, A_L, \emptyset)$. Apply the induction hypothesis and Lemma 56,

$$S((\text{TEP}, h)_A) = S((\text{TEP}, h - 1)_{A_L}) \le k^{h-1}.$$

When $n_{h-1} < \ell \leq \sum_{i=2}^{h-1} n_i$, choose $A = (\emptyset, L, A_R)$. Then, $1 \leq |A_R| \leq \sum_{i=2}^{h-2} n_i$. Apply the induction hypothesis and Lemma 56,

$$S((\text{TEP}, h)_A) = S((\text{TEP}, h - 1)_L) \cdot S((\text{TEP}, h - 1)_{A_R}) \le k \cdot k^{h-2} = k^{h-1}.$$

(ii) $\sum_{i=2}^{h-1} n_i < \ell < \sum_{i=2}^{h-1} n_i + k^2$. Choose $A = (\emptyset, L, A_R)$. Then, $\sum_{i=2}^{h-2} n_i < |A_R| < \sum_{i=2}^{h-2} n_i + k^2$. Apply the induction hypothesis and Lemma 56,

$$S((\text{TEP}, h)_A) = S((\text{TEP}, h-1)_L) \cdot S((\text{TEP}, h-1)_{A_R}) \le k \cdot k^{h-3}(k^2 - k + 2) = k^{h-2}(k^2 - k + 2).$$

(iii) $\sum_{i=2}^{h-1} n_i + k^2 \leq \ell \leq 2n_{h-1}$. Choose $A = (\emptyset, L, A_R)$. Then, $\sum_{i=2}^{h-2} n_i + k^2 \leq |A_R| \leq n_{h-1}$. Apply the induction hypothesis and Lemma 56,

$$S((\text{TEP}, h)_A) = S((\text{TEP}, h - 1)_L) \cdot S((\text{TEP}, h - 1)_{A_R})$$

$$\leq k \cdot \max\{k^{h-3}(k+2), k^2 + k + 1\} = k^{h-2}(k+2),$$

where we used $h \ge 4$ in the last step.

(iv) $2n_{h-1} < \ell \le n_h$. This can be proved in the same way as (i) in the proof of Theorem 58. \Box