GLOBAL SOLUTIONS OF THE TWO-DIMENSIONAL **RIEMANN PROBLEM WITH FOUR-SHOCK INTERACTIONS** FOR THE EULER EQUATIONS FOR POTENTIAL FLOW

GUI-QIANG G. CHEN, ALEX CLIFFE, FEIMIN HUANG, SONG LIU, AND QIN WANG

ABSTRACT. We present a rigorous approach and related techniques to construct global solutions of the two-dimensional (2-D) Riemann problem with four-shock interactions for the Euler equations for potential flow. With the introduction of three critical angles: the vacuum critical angle from the compatibility conditions, the detachment angle, and the sonic angle, we clarify all configurations of the Riemann solutions for the interactions of two-forward and two-backward shocks, including the subsonic-subsonic reflection configuration that has not emerged in previous results. To achieve this, we first identify the three critical angles that determine the configurations, whose existence and uniqueness follow from our rigorous proof of the strict monotonicity of the steady detachment and sonic angles for 2-D steady potential flow with respect to the Mach number of the upstream state. Then we reformulate the 2-D Riemann problem into the shock reflection-diffraction problem with respect to a symmetric line, along with two independent incident angles and two sonic boundaries varying with the choice of incident angles. With these, the problem can be further reformulated as a free boundary problem for a second-order quasilinear equation of mixed elliptic-hyperbolic type. The difficulties arise from the degenerate ellipticity of the nonlinear equation near the sonic boundaries, the nonlinearity of the free boundary condition, the singularity of the solution near the corners of the domain, and the geometric properties of the free boundary. To solve the problem, we need to analyze the solutions for a quasilinear degenerate elliptic equation by the maximum principle of the mixed-boundary value problem, the theory of the oblique derivative boundary value problem, the uniform a priori estimates, and the iteration method. To the best of our knowledge, this is the first rigorous result for the 2-D Riemann problem with four-shock interactions for the Euler equations. The approach and techniques developed for the Riemann problem for four-wave interactions should be useful for solving other 2-D Riemann problems for more general Euler equations and related nonlinear hyperbolic systems of conservation laws.

CONTENTS

2

1. Introduction	2
2. The Riemann Problem and Main Theorems for Four-Shock Interactions	5
2.1. Mathematical formulation	5
2.2. Regular reflection configurations	10
2.3. Admissible solutions and main theorems	15
3. Uniform Estimates for Admissible Solutions	17
3.1. Monotonicity and ellipticity	17
3.2. Interior Hölder estimates away from the sonic boundaries	24
3.3. Weighted Hölder estimates near the sonic boundaries	25
3.4. Compactness of the set of admissible solutions	35
4. Iteration Method and Existence of Admissible Solutions	35
4.1. Mapping between the elliptic and standard domains	36
4.2. Definition and properties of the iteration set	43
4.3. Definition of the iteration map	53
4.4. Proof of Theorem 2.1: Existence of admissible solutions	56

Date: May 25, 2023.

²⁰²⁰ Mathematics Subject Classification. 35M30, 35M12, 35R35, 35Q31, 76N10, 35L65, 35L70, 35B65, 35D30, 35J70. 35J66.

Key words and phrases. Riemann problem, shock interaction, Euler equations, potential flow, free boundary, mixed elliptic-hyperbolic type, a priori estimates, degenerate elliptic equation, regularity, convexity.

5. Optimal Regularity of Solutions and Convexity of Free Boundaries	60
5.1. Proof of Theorem 2.2: Optimal regularity of solutions	60
5.2. Proof of Theorem 2.3: Convexity of free boundaries and transonic shocks	62
Appendix A. Proof of Lemma 2.5 and Related Properties of Solutions	63
A.1. Proof of Lemma 2.5: Monotonicity of critical angles for 2-D steady potential flow	63
A.2. Monotonicity properties with respect to the incident angles	64
Appendix B. Some Known Results Needed for the Proofs	67
B.1. Well-posedness of the iteration boundary value problem	67
B.2. Regularized distance function	71
B.3. Leray-Schauder degree theorem	71
B.4. Regularity theorem	71
B.5. General framework for the convexity of transonic shocks	72
Acknowledgements	73
References	74

1. INTRODUCTION

We are concerned with the two-dimensional (2-D) Riemann problem for the Euler equations for potential flow, consisting of the conservation law of mass and the Bernoulli law:

(1.1)
$$\begin{cases} \partial_t \rho + \operatorname{div}_{\mathbf{x}}(\rho \nabla_{\mathbf{x}} \Phi) = 0, \\ \partial_t \Phi + \frac{1}{2} |\nabla_{\mathbf{x}} \Phi|^2 + h(\rho) = B, \end{cases}$$

where $(t, \mathbf{x}) \in (0, \infty) \times \mathbb{R}^2$, $\rho = \rho(t, \mathbf{x})$ is the density, $\Phi = \Phi(t, \mathbf{x})$ is the velocity potential (*i.e.*, $\nabla_{\mathbf{x}} \Phi = (u, v)$ is the velocity), $h(\rho) \coloneqq \int_1^{\rho} \frac{p'(s)}{s} \, ds$ is the enthalpy with the polytropic pressure law $p(\rho) = A\rho^{\gamma}$ for $\gamma \ge 1$ and A > 0, and the Bernoulli constant *B* is determined by the far field flow. Without loss of generality, we may fix $A = \gamma^{-1}$ by scaling and denote the sonic speed by $c(\rho) \coloneqq \sqrt{p'(\rho)} = \rho^{\frac{\gamma-1}{2}}$.

In this paper, we focus on the 2-D Riemann problem with four-shock interactions and corresponding Riemann initial given by

(1.2)
$$(\rho, \Phi)(0, \mathbf{x}) = (\rho_i, (u_i, v_i) \cdot \mathbf{x}) \quad \text{for } \mathbf{x} \in \Lambda_i,$$

for suitable constant states (i) with values $U_i := (\rho_i, u_i, v_i)$, and corresponding scale-invariant domains $\Lambda_i \subseteq \mathbb{R}^2$ for i = 1, 2, 3, 4. We choose these domains to be four sectors of \mathbb{R}^2 , defined by

(1.3)
$$\Lambda_1 \coloneqq \left\{ \mathbf{x} \in \mathbb{R}^2 : -\theta_{14} < \theta < \theta_{12} \right\}, \qquad \Lambda_2 \coloneqq \left\{ \mathbf{x} \in \mathbb{R}^2 : \theta_{12} < \theta < \pi - \theta_{32} \right\},$$
$$\Lambda_3 \coloneqq \left\{ \mathbf{x} \in \mathbb{R}^2 : \pi - \theta_{32} < \theta < \pi + \theta_{34} \right\}, \qquad \Lambda_4 \coloneqq \left\{ \mathbf{x} \in \mathbb{R}^2 : \pi + \theta_{34} < \theta < 2\pi - \theta_{14} \right\},$$

where θ is the polar angle of point $\mathbf{x} \in \mathbb{R}^2$, and the four parameters $\theta_{12}, \theta_{32}, \theta_{34}, \theta_{14} \in (0, \frac{\pi}{2})$. The initial data and shock discontinuities are depicted in Fig. 1.1.

The Euler equations for potential flow (1.1) are the oldest, yet still prominent, paradigm of fluid dynamic equations for inviscid fluids, which have been widely used in aerodynamics. System (1.1) has further been extended to the full Euler equations for more general inviscid compressible fluids when the effect of vortex sheets and the derivation of vorticity become significant. One of the main features of the Euler equations is the formation of shocks in finite time, no matter how smooth the initial data are. The shock is one of the most fundamental nonlinear waves in compressible fluid flow, and the analysis of shocks dates back to the 19th century; see *e.g.* [20, 27, 43, 44, 47]. In 1860, Riemann [44] first considered a special initial value problem for the one-dimensional (1-D) Euler system, with two constant states separated at the origin—now known as the Riemann problem. For general 1-D strictly hyperbolic systems of conservation laws, Lax [29] showed that the Riemann solutions are the combination of three fundamental waves: the rarefaction waves, the shocks, and the contact discontinuities. In 1965, Glimm [24] employed the Riemann solutions as building blocks of the Glimm scheme to establish the global existence of solutions in BV (bounded variation) of 1-D strictly hyperbolic systems of conservation laws for the initial data of small BV,



FIG. 1.1. The four sectors of the Riemann initial data.

for which the Riemann solutions play a fundamental role. For the uniqueness and stability of BV solutions, we refer to [3, 4, 5, 40, 41] and the references cited therein. For a systematic theory of 1-D hyperbolic systems of conservation laws, we refer to [3, 19, 26, 30, 42] and the references cited therein.

Multi-dimensional (M-D) Riemann problems for the Euler system are vastly more complicated; see [8, 19, 26, 30, 31] and the references cited therein. For the 2-D four-quadrant Riemann problem with four initial constant states given in the four quadrants so that each jump between two neighbouring quadrants projects exactly one planar fundamental wave, Zhang-Zheng [49] predicted that there are a total of 16 genuinely different configurations of the Riemann solutions for polytropic gases. In Chang-Chen-Yang [6, 7], it is first observed that, when two initially parallel slip lines are present, it makes a distinguished difference whether the vorticity waves generated have the same or opposite sign, which, along with Lax-Liu [32], leads to the classification with a total of 19 genuinely different configurations of the Riemann solutions for the compressible Euler equations for polytropic gases, via characteristic analysis; also see [25, 28, 33, 45] and the references cited therein. However, there have been few rigorous global results for the 2-D fourquadrant Riemann problem for the Euler equations, except for the result obtained by Li-Zheng in [36], in which the four-rarefaction wave case was considered. On the other hand, the 2-D Riemann problem has been solved for simplified models such as the pressure gradient system [14], and the local structure of wave interactions in the Riemann problem has been analyzed in [34, 35, 37]. For the 2-D Riemann problem for Chaplygin gases, we refer to [17, 46] and the references cited therein.

The purpose of this paper is to solve rigorously the 2-D Riemann problem with four-shock interactions globally for the Euler equations for potential flow (1.1). The 2-D Riemann initial data (1.2) under consideration in this paper satisfy that

 $\begin{cases} \text{one forward shock } \overrightarrow{S}_{1j} \text{ is generated between states (1) and } (j), \\ \text{one backward shock } \overleftarrow{S}_{3j} \text{ is generated between states (3) and } (j), \end{cases}$

for j = 2, 4, and that the angular bisectors at the vertices of Λ_1 and Λ_3 in Fig. 1.1 coincide. Without loss of generality, we choose $\theta_{12} = \theta_{14}$ and $\theta_{32} = \theta_{34}$, and then the x_1 -axis may be considered as a rigid wall. The above Riemann problem corresponds to one of the conjectures for the Euler equations proposed in Zhang-Zheng [49].

The problem considered has a close relation to the shock reflection-diffraction problem, whose research dates back to 1878, in which Ernst Mach revealed the complexity of the shock reflection-diffraction diffraction configurations that include two fundamental modes: the regular reflection-diffraction and the Mach reflection-diffraction. The shock reflection phenomena occur when a moving planar shock wave impinges on a rigid wall; see Courant-Friedrichs [18] and Chen-Feldman [10]. In the recent three decades, there have been many theoretical results on shock reflection/diffraction problems. Indeed, Elling-Liu [22] considered the self-similar solutions of the Prandtl-Meyer problem for supersonic potential flow onto a solid wedge for a class of physical parameters. Chen-Feldman [9]

proved the first global existence of solutions of the shock reflection-diffraction problem for potential flow when the incident angles are small; later, they succeeded to establish the global existence of solutions for the incident angles all the way up to the detachment angle in [10]. Furthermore, Chen-Feldman-Xiang [12] proved the convexity of transonic shocks, which leads to the uniqueness and stability of the admissible solution [13]. More recently, Bae-Chen-Feldman [2] obtained the existence and the optimal regularity of admissible solutions of the Prandtl-Meyer reflection problem for the general case up to the detachment angle. For the supersonic flow passing through a wing or curved wedge, see also [15, 16, 11].

For the 2-D Riemann problem with four-shock interactions under consideration, we can reformulate the problem into a shock reflection-diffraction problem with respect to a symmetric line. Compared with the previous results, such as the shock reflection-diffraction problem [10] and the Prandtl-Meyer reflection problem [2], we now have two independent incident angles in the four-shock reflection problem. Correspondingly, we have two sonic boundaries varying with the choice of incident angles, which leads to more complicated configurations. After introducing the self-similar coordinates, the original problem can be reformulated as a free boundary problem for a second-order quasilinear partial differential equation (PDE) of mixed elliptic-hyperbolic type. The difficulties come from the degenerate ellipticity of the nonlinear PDE near the sonic boundaries, the nonlinearity of the free boundary conditions, the singularity of the solution near the corners of the domain, and the geometric properties of the free boundary. To solve our problem, we need to analyze the solutions for a quasilinear degenerate elliptic equation by the maximum principle of mixed-boundary value problems and the theory of the oblique derivative boundary value problems [23, 38, 39], as well as the uniform estimates and the iteration method from [2, 10, 21].

An important part of our analysis includes the determination of the three critical angles: the vacuum critical angle, the detachment angle, and the sonic angle. Their existence and uniqueness follow from Lemma 2.5, whereby we prove the strict monotonicity of the steady detachment and sonic angles for 2-D steady potential flow with respect to the Mach number of the upstream state. Using these critical angles, we classify all configurations of the solutions of the Riemann problem. In particular, the subsonic-subsonic reflection configuration has never emerged in the previous works. Lemma 2.5 should be useful in the analysis of other shock reflection-diffraction problems.

To the best of our knowledge, this is the first result on the shock reflection-diffraction of the Riemann problem for potential flow with piecewise constant initial data in four sectors. Furthermore, the mathematical methods developed in solving the free boundary problem for nonlinear PDEs such as the uniform *a priori* estimate on the ellipticity of the equation, the anisotropic scaling method, and the sophisticated construction of the iteration set and iteration map will provide insights for the analysis of more general nonlinear problems. In particular, the Euler equations for potential flow are the core part of the full Euler equations, coupled with the incompressible-type Euler equations, in a local domain, the techniques and approaches developed for the Riemann problem for four-wave interactions will be useful to guide the analysis of the structure of Riemann solutions, shock reflection/diffraction problems, and numerical simulations of the full Euler equations.

The organisation of this paper is as follows: In Section 2, we present the mathematical formulation of the Riemann problem with four-shock interactions and the definition of admissible solutions of the free boundary problem, and state three main results. In Section 3, we obtain the directional monotonicity, the strict ellipticity, and the uniform weighted Hölder estimates of admissible solutions. In Section 4, we prove the existence of admissible solutions by constructing a suitable iteration set and iteration map via applying the Leray-Schauder degree theory. Finally, in Section 5, we give the proof of the optimal regularity of admissible solutions, as well as the proof of the convexity of free boundaries and transonic shocks. In Appendix A, we give the details for some calculations involving the potential flow equation and, in Appendix B, we present some known results that are needed throughout the proofs.

2. The Riemann Problem and Main Theorems for Four-Shock Interactions

In this section, we first present the mathematical formulation of the Riemann problem with four-shock interactions as a free boundary problem, along with the definition of admissible solutions of the free boundary problem, and then state three main theorems.

2.1. Mathematical formulation. The Riemann problem (1.1)-(1.2) is invariant under the selfsimilar scaling:

$$(t, \mathbf{x}) \mapsto (\lambda t, \lambda \mathbf{x}), \quad (\rho, \Phi)(t, \mathbf{x}) \mapsto (\rho, \frac{1}{\lambda} \Phi)(\lambda t, \lambda \mathbf{x}) \qquad \text{for any } \lambda > 0.$$

This leads us to consider the self-similar solution $(\rho, \phi)(\boldsymbol{\xi}) := (\rho, \frac{\Phi}{t})(t, \mathbf{x})$ in the self-similar coordinates $\boldsymbol{\xi} = (\xi, \eta) \coloneqq \frac{\mathbf{x}}{t} \in \mathbb{R}^2$. Moreover, by introducing the pseudo-potential function $\varphi(\boldsymbol{\xi}) \coloneqq \phi(\boldsymbol{\xi}) - \frac{1}{2} |\boldsymbol{\xi}|^2$, we obtain the pseudo-steady potential flow equation for $\varphi(\boldsymbol{\xi})$ in the form:

(2.1)
$$\operatorname{div}(\rho(|D\varphi|,\varphi)D\varphi) + 2\rho(|D\varphi|,\varphi) = 0,$$

where div and D represent the divergence and gradient operators with respect to the self-similar coordinates $\boldsymbol{\xi} \in \mathbb{R}^2$, and the density is determined from the Bernoulli law as

(2.2)
$$\rho(|D\varphi|,\varphi) = \begin{cases} \left(1 + (\gamma - 1)(B - \frac{1}{2}|D\varphi|^2 - \varphi)\right)^{\frac{1}{\gamma - 1}} & \text{if } \gamma > 1,\\ \exp(B - \frac{1}{2}|D\varphi|^2 - \varphi) & \text{if } \gamma = 1. \end{cases}$$

Equation (2.1) with (2.2) is a second-order nonlinear equation of mixed elliptic-hyperbolic type. It is elliptic if and only if $|D\varphi| < c(|D\varphi|, \varphi)$ (pseudo-subsonic), while it is hyperbolic if and only if $|D\varphi| > c(|D\varphi|, \varphi)$ (pseudo-supersonic).

We seek a weak solution for the pseudo-steady potential flow equation (2.1)-(2.2).

Definition 2.1. A function $\varphi \in W^{1,\infty}_{\text{loc}}(\mathbb{R}^2)$ is called a weak solution of equation (2.1)–(2.2) in \mathbb{R}^2 if φ satisfies the following properties:

- (i) $B \frac{1}{2} |D\varphi|^2 \varphi \ge h(0^+)$ a.e. in \mathbb{R}^2 ; (ii) $\int_{\mathbb{R}^2} \left(\rho(|D\varphi|, \varphi) D\varphi \cdot D\zeta 2\rho(|D\varphi|, \varphi)\zeta \right) d\boldsymbol{\xi} = 0$ for every $\zeta \in C_c^{\infty}(\mathbb{R}^2)$.

Let S be a C^1 -curve across which $|D\varphi|$ is discontinuous. Then φ is a weak solution if and only if φ satisfies the Rankine–Hugoniot conditions on S:

(2.3)
$$[\varphi]_S = [\rho(|D\varphi|, \varphi)D\varphi \cdot \boldsymbol{\nu}_S]_S = 0,$$

where $[F]_S$ represents the jump of a quantity F across S, and ν_S is any unit normal vector on S. By Definition 2.1(i) and (2.3), $D\varphi \cdot \nu_S$ is either positive or negative across the discontinuity S. We say that ν_S points from upstream to downstream if $D\varphi \cdot \nu_S > 0$, and from downstream to upstream if $D\varphi \cdot \boldsymbol{\nu}_S < 0$. The discontinuity S is called a *shock* if it further satisfies the physical entropy condition: the corresponding density $\rho(|D\varphi|,\varphi)$ increases across S from upstream to downstream, or equivalently, $|D\varphi \cdot \nu_S|$ decreases across S from upstream to downstream.

In the self-similar coordinates $\boldsymbol{\xi} = (\xi, \eta)$, the initial condition (1.2) becomes the following asymptotic boundary condition at infinity: For any $\theta \in [0, 2\pi)$ and i = 1, 2, 3, 4,

$$\lim_{r \to \infty} \|\varphi - \varphi_i\|_{C^{0,1}(R_\theta \setminus B_r(\mathbf{0}))} = 0$$

whenever ray $R_{\theta} \coloneqq \{r(\cos\theta, \sin\theta) : r > 0\}$ is contained in sector Λ_i as defined in (1.3), where the pseudo-potential function φ_i corresponding to the uniform state (i) is given by

(2.4)
$$\varphi_i(\boldsymbol{\xi}) \coloneqq -\frac{1}{2}|\boldsymbol{\xi}|^2 + (u_i, v_i) \cdot \boldsymbol{\xi} + k_i$$

and constant k_i is uniquely determined to satisfy (2.2) with $(\rho, \varphi) = (\rho_i, \varphi_i)$ for i = 1, 2, 3, 4. We may simply fix $k_2 = 0$ in (2.4) since the Bernoulli constant can be shifted to $B - k_2$.

Since system (1.1) is of Galilean invariance, it is convenient to normalize the initial data to satisfy $u_2 = v_1 = 0$. From (2.2) and (2.4) with $k_2 = 0$, the Bernoulli constant B is then fixed as

(2.5)
$$B = \frac{1}{2}v_2^2 + h(\rho_2).$$



FIG. 2.1. Four-shock interaction with symmetric incident shocks.

In this paper, we assume that the initial data (1.2) are chosen such that, in the far field region, any two neighboring states are connected by exactly one planar shock discontinuity. Furthermore, we restrict our attention to the case:

(2.6)
$$\max\{\rho_1, \rho_3\} < \min\{\rho_2, \rho_4\},\$$

so that the initial data consist precisely of two forward shocks $\{\vec{S}_{12}, \vec{S}_{14}\}$ and two backward shocks $\{\vec{S}_{32}, \vec{S}_{34}\}$, which travel away from the origin in pairs; see Case 5.2 in [6]. Moreover, to facilitate our analysis, we focus on the symmetric case:

(2.7)
$$\theta_{12} = \theta_{14} \eqqcolon \theta_1, \quad \theta_{32} = \theta_{34} \eqqcolon \theta_2 \qquad \text{with } \theta_1, \theta_2 \in (0, \frac{\pi}{2}).$$

We refer to parameters $\boldsymbol{\theta} \coloneqq (\theta_1, \theta_2)$ as the incident angles of the shock discontinuities; see Fig. 2.1.

2.1.1. Compatibility conditions. It is clear that the initial data and parameters $\boldsymbol{\theta}$ cannot be specified arbitrarily. In the following lemma, we give the necessary and sufficient conditions for the initial data $U_i = (\rho_i, u_i, v_i)$ and parameters $\boldsymbol{\theta}$ to generate two forward shocks $\{\vec{S}_{12}, \vec{S}_{14}\}$ and two backward shocks $\{\vec{S}_{32}, \vec{S}_{34}\}$.

Lemma 2.2. Fix $\gamma \geq 1$ and $\rho_2 > 0$. There exist both a constant $v_{\min} \in [-\infty, 0)$ depending on (γ, ρ_2) and a critical angle $\theta^{\operatorname{cr}} \in (0, \frac{\pi}{2}]$ depending on (γ, ρ_2, v_2) such that, whenever $v_2 \in (v_{\min}, 0)$ and $\boldsymbol{\theta} \in (0, \theta^{\operatorname{cr}})^2$, there exists a unique choice of constant states $U_i = (\rho_i, u_i, v_i)$, for i = 1, 3, 4, satisfying the Rankine–Hugoniot conditions (2.3) between any two neighbouring states and the entropy condition (2.6) so that two forward shocks $\{\vec{S}_{12}, \vec{S}_{14}\}$ and two backward shocks $\{\vec{S}_{32}, \vec{S}_{34}\}$ are generated.

Proof. Throughout the proof, we take i = 1, 3, and j = 2, 4. In the self-similar coordinates, we denote the shock discontinuity line between state (i) and state (j) by

(2.8)
$$S_{ij} \coloneqq \{\boldsymbol{\xi} \in \mathbb{R}^2 : \varphi_i(\boldsymbol{\xi}) = \varphi_j(\boldsymbol{\xi})\} = \{\boldsymbol{\xi} \in \mathbb{R}^2 : \eta = (-1)^{\frac{i+j+1}{2}} \xi \tan \theta_{ij} + a_{ij}\},$$

where the expressions of φ_i and φ_j are given in (2.4), $\theta_{ij} \in (0, \frac{\pi}{2})$ is the angle between the ξ -axis and the shock discontinuity line S_{ij} , and a_{ij} is a constant determined below. We fix the unit normal vector on S_{ij} to be

$$\boldsymbol{\nu}_{ij} = \left((-1)^{\frac{i+1}{2}} \sin \theta_{ij}, (-1)^{\frac{j+2}{2}} \cos \theta_{ij} \right),$$

which points towards the downstream state according to the entropy condition (2.6), so that

(2.9)
$$D\varphi_i(\boldsymbol{\xi}) \cdot \boldsymbol{\nu}_{ij} > D\varphi_j(\boldsymbol{\xi}) \cdot \boldsymbol{\nu}_{ij} > 0 \qquad \text{for } \boldsymbol{\xi} \in S_{ij}$$

A straightforward calculation by using (2.2)-(2.4) and (2.8)-(2.9) gives

(2.10)
$$(u_i, v_i) - (u_j, v_j) = \ell_{ij} \boldsymbol{\nu}_{ij}, \quad a_{ij} = (-1)^{\frac{i+j+1}{2}} \left(v_i - u_i \tan \theta_{ij} + \frac{(-1)^{\frac{i-1}{2}} \rho_j \ell_{ij}}{(\rho_i - \rho_j) \cos \theta_{ij}} \right),$$

where $\ell_{ij} \coloneqq \ell(\rho_i, \rho_j)$ is defined as

(2.11)
$$\ell(\rho_i, \rho_j) \coloneqq \sqrt{\frac{2(\rho_i - \rho_j)(h(\rho_i) - h(\rho_j))}{\rho_i + \rho_j}}$$

It is direct to check that ℓ is symmetric and satisfies the strict monotonicity: For any $\bar{\rho} > 0$, (2.12) $\rho \mapsto \ell(\bar{\rho}, \rho)$ is strictly increasing on $\rho \in (\bar{\rho}, \infty)$ and strictly decreasing on $\rho \in (0, \bar{\rho})$. From (2.7) and (2.10), we deduce that

$$(\ell_{12} - \ell_{14})\sin\theta_1 = (\ell_{34} - \ell_{32})\sin\theta_2,$$

from which $\rho_4 = \rho_2$ must hold by virtue of the entropy condition (2.6) and (2.12). Thus, we obtain the following relations:

$$\ell_{12}\cos\theta_1 = \ell_{32}\cos\theta_2, \qquad \begin{cases} u_2 = u_4 = u_1 + \ell_{12}\sin\theta_1 = u_3 - \ell_{32}\sin\theta_2, \\ v_1 = v_3 = v_2 + \ell_{12}\cos\theta_1 = v_4 - \ell_{32}\cos\theta_2. \end{cases}$$

Recall that $u_2 = v_1 = 0$ are fixed, so that

(2.13)
$$U_1 = (\rho_1, -\ell_{12}\sin\theta_1, 0), \quad U_2 = (\rho_2, 0, v_2), \quad U_3 = (\rho_3, \ell_{32}\sin\theta_2, 0), \quad U_4 = (\rho_2, 0, -v_2),$$

whenever $\rho_1, \rho_3 \in (0, \rho_2)$ satisfy

(2.14)
$$\ell_{12}\cos\theta_1 = \ell_{32}\cos\theta_2 = -v_2 \,.$$

We refer to (2.14) as the compatibility conditions for the initial data. We define

(2.15)
$$v_{\min} \coloneqq -\ell(0^+, \rho_2) \in [-\infty, 0), \qquad \theta^{\operatorname{cr}} \coloneqq \arccos\left(\frac{-v_2}{\ell(0^+, \rho_2)}\right) \in (0, \frac{\pi}{2}]$$

Then, by (2.12), the necessary and sufficient conditions for the existence of ρ_1 and ρ_3 uniquely solving (2.14) subject to (2.6) are

$$v_2 \in (v_{\min}, 0), \qquad \boldsymbol{\theta} \in (0, \theta^{\operatorname{cr}})^2.$$

The constant states U_i , i = 1, 3, 4, are then uniquely determined by (2.13)–(2.14) depending only on $(\gamma, \rho_2, v_2, \theta)$.

We call θ^{cr} the vacuum critical angle, because $\rho_1 \to 0^+$ as $\theta_1 \to \theta^{cr-}$ and $\rho_3 \to 0^+$ as $\theta_2 \to \theta^{cr-}$, according to (2.14). Note that, in potential flow, the incident shock with vacuum upstream state is mathematically consistent, which is in contrast to the non-potential Euler flow for which the maximal density ratio across a shock is bounded.

It follows from Lemma 2.2 that $\rho_2 = \rho_4$. Without loss of generality, we fix $\rho_2 = \rho_4 = 1$ via the scale-invariance of equation (2.1) as follows

$$\boldsymbol{\xi} \mapsto c_2 \boldsymbol{\xi}, \qquad (\rho, \varphi, \rho_2, v_2) \mapsto \left(\frac{\rho}{\rho_2}, \frac{\varphi}{c_2^2}, 1, \frac{v_2}{c_2}\right)$$

Then the entropy condition (2.6) becomes $\max\{\rho_1, \rho_3\} < 1$, and $v_{\min} = -\ell(0^+, 1)$ depends only on $\gamma \ge 1$. For simplicity, we occasionally use the abbreviation $\ell(\cdot) := \ell(\cdot, 1)$.

2.1.2. The asymptotic boundary value problem in the upper half-plane. Equation (1.1) and the initial data in (1.2) given by (2.13) are invariant under reflection with respect to the x_1 -axis. Thus, we look for solutions of (1.1)–(1.2) in the symmetric form:

$$\Phi(t, x_1, x_2) = \Phi(t, x_1, -x_2)$$
 for all $(t, x_1, x_2) \in (0, \infty) \times \mathbb{R}^2$.

which is equivalent to

$$\varphi(\xi,\eta) = \varphi(\xi,-\eta) \quad \text{for all } (\xi,\eta) \in \mathbb{R}^2$$

For this reason, it suffices to consider the restriction of solutions to the upper half-plane

$$\mathbb{R}^2_+ \coloneqq \left\{ \boldsymbol{\xi} \in \mathbb{R}^2 : \eta > 0 \right\},\,$$

so long as we impose the slip boundary condition

$$Darphi \cdot (0,1) = 0$$
 on $L_{ ext{sym}} \coloneqq \left\{ oldsymbol{\xi} \in \mathbb{R}^2 \, : \, \eta = 0
ight\}.$

This condition means that L_{sym} can be regarded as a solid wall.

Hereafter, we use symbols S_{12} and S_{32} to represent only the closure of the rays of the incident shocks S_{12} and S_{32} lying in the upper half-plane. We denote by $P_0^1 = (\xi^{P_0^1}, \eta^{P_0^1})$ the point of intersection between S_{12} and L_{sym} , and by $P_0^2 = (\xi^{P_0^2}, \eta^{P_0^2})$ the point of intersection between S_{32} and L_{sym} . We expect the incident shocks S_{12} and S_{32} to undergo shock reflection-diffraction at points P_0^1 and P_0^2 respectively; see Fig. 2.1. From (2.8), (2.10), and (2.13), we have

(2.16)
$$P_0^1 = (-\ell(\rho_1)\sin\theta_1 + \frac{1}{1-\rho_1}\frac{\ell(\rho_1)}{\sin\theta_1}, 0), \qquad P_0^2 = (\ell(\rho_3)\sin\theta_2 - \frac{1}{1-\rho_3}\frac{\ell(\rho_3)}{\sin\theta_2}, 0).$$

Using (2.4), (2.13)-(2.14), and (2.16), the pseudo-potentials of states (1)-(3) are given by

(2.17)
$$\varphi_1 = -\frac{1}{2}|\boldsymbol{\xi}|^2 + v_2(\boldsymbol{\xi} - \boldsymbol{\xi}^{P_0^1})\tan\theta_1, \ \varphi_2 = -\frac{1}{2}|\boldsymbol{\xi}|^2 + v_2\eta, \ \varphi_3 = -\frac{1}{2}|\boldsymbol{\xi}|^2 - v_2(\boldsymbol{\xi} - \boldsymbol{\xi}^{P_0^2})\tan\theta_2.$$

In particular, φ_1 is independent of θ_2 , while φ_3 is independent of θ_1 . The following lemma concerning the pseudo-Mach number of state (2) at the reflection points P_0^1 and P_0^2 is fundamental to our later analysis of the critical angles in §2.2.1.

Lemma 2.3. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For any incident angle $\theta_1 \in (0, \theta^{cr})$, the pseudo-Mach number of state (2) at the reflection point P_0^1 is given by

(2.18)
$$M_2^{(\theta_1)} \coloneqq |D\varphi_2(P_0^1)| = \frac{\ell(\rho_1)}{1-\rho_1} (\rho_1^2 + \cot^2 \theta_1)^{\frac{1}{2}},$$

with $\rho_1 = \rho_1(\theta_1; v_2) \in (0, 1)$ given by (2.14). Moreover, the pseudo-Mach number $M_2^{(\theta_1)}$ is a strictly decreasing function of the incident angle θ_1 and satisfies that $M_2^{(\theta_1)} \to \infty$ as $\theta_1 \to 0^+$.

Proof. Formula (2.18) follows directly from (2.14) and (2.16)–(2.17).

For the second property, the limit is clear since (2.14) implies that $\ell(\rho_1) \to -v_2 > 0$ as $\theta_1 \to 0^+$, whilst $\cot^2 \theta_1 \to \infty$ as $\theta_1 \to 0^+$. To determine the strict monotonicity of $M_2^{(\theta_1)}$, we differentiate (2.14) and (2.18) with respect to θ_1 and re-arrange it to obtain

(2.19)
$$\frac{(1-\rho_1)^2 \cot \theta_1}{2(\rho_1 + \cot^2 \theta_1)(\ell(\rho_1))^2} \frac{\mathrm{d}(M_2^{(\theta_1)})^2}{\mathrm{d}\theta_1} = \rho_1 + \frac{1}{1-\rho_1} \frac{\ell(\rho_1)}{\ell'(\rho_1)} - \cot^2 \theta_1 \,.$$

Furthermore, from the definition of $\ell(\rho)$ in (2.11), we directly compute

$$(1-\rho)\frac{\ell'(\rho)}{\ell(\rho)} = -\frac{(\ell(\rho))^2 + (1-\rho)^2 h'(\rho)}{(\ell(\rho))^2 (1+\rho)}$$

Substituting the above expression into the right-hand side of (2.19) gives

RHS of (2.19) =
$$\frac{\rho_1(1-\rho_1)^2 h'(\rho_1) - (\ell(\rho_1))^2}{(\ell(\rho_1))^2 + (1-\rho_1)^2 h'(\rho_1)} - \cot^2 \theta_1.$$

Denote the numerator of the first term above by

$$f(\rho) \coloneqq \rho(1-\rho)^2 h'(\rho) - (\ell(\rho))^2 \equiv \frac{1-\rho}{1+\rho} \tilde{f}(\rho)$$

where $\tilde{f}(\rho) \coloneqq (\gamma + 1)h(\rho) - (\gamma - 1)\rho^2 h(\rho) + (1 - \rho^2)$. Observe that $\tilde{f}(1) = 0$ and $\tilde{f}'(\rho) = (\gamma + 1)(1 - \rho^2)h'(\rho) > 0$ for any $\rho \in (0, 1)$.

Thus, $\tilde{f}(\rho) < 0$ for all $\rho \in (0,1)$. It directly follows that the RHS of (2.19) is negative. We conclude that $M_2^{(\theta_1)}$ is a strictly decreasing function of $\theta_1 \in (0, \theta^{cr})$.

It follows immediately from Lemma 2.3 that $\xi^{P_0^1} > 0$ is a strictly decreasing function of $\theta_1 \in (0, \theta^{cr})$ with $\xi^{P_0^1} \to \infty$ as $\theta_1 \to 0^+$, since $M_2^{(\theta_1)} = |(-\xi^{P_0^1}, v_2)|$. Similarly, $\xi^{P_0^2} < 0$ is a strictly increasing function of $\theta_2 \in (0, \theta^{cr})$ with $\xi^{P_0^2} \to -\infty$ as $\theta_2 \to 0^+$.

We define the background solution $\bar{\varphi} \in C^{0,1}_{\text{loc}}(\mathbb{R}^2_+)$ as

(2.20)
$$\bar{\varphi}(\boldsymbol{\xi}) \coloneqq \min\left\{\varphi_1(\boldsymbol{\xi}), \varphi_2(\boldsymbol{\xi}), \varphi_3(\boldsymbol{\xi})\right\},$$

and define the three open domains

(2.21)
$$\Omega_i \coloneqq \inf \left\{ \boldsymbol{\xi} \in \mathbb{R}^2_+ : \varphi_i(\boldsymbol{\xi}) = \bar{\varphi}(\boldsymbol{\xi}) \right\} \quad \text{for } i = 1, 2, 3,$$

where int G denotes the interior of a set $G \subseteq \mathbb{R}^2$. Note that Ω_1 and Ω_3 are the wedge-shaped domains with wedge angles θ_1 and θ_2 respectively, and they become empty sets in the limits: $\theta_1 \to 0^+$ and $\theta_2 \to 0^+$, respectively.

In light of the above discussion, we seek solutions of the following asymptotic boundary value problem in the self-similar coordinates $\boldsymbol{\xi} = (\xi, \eta)$ in the upper half-plane \mathbb{R}^2_+ .

Problem 2.4 (Asymptotic boundary value problem). Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For any $\boldsymbol{\theta} \in (0, \theta^{cr})^2$, determine the existence of a weak solution $\varphi \in W^{1,\infty}_{\text{loc}}(\mathbb{R}^2_+)$ of equations (2.1)–(2.2) in \mathbb{R}^2_+ satisfying the following conditions:

(i) The asymptotic boundary condition at infinity:

$$\lim_{r \to \infty} \|\varphi - \bar{\varphi}\|_{C^{0,1}(R_{\theta} \setminus B_r(\mathbf{0}))} = 0 \qquad \text{for any } \theta \in (0,\pi);$$

(ii) The slip boundary condition on the symmetric boundary:

$$D\varphi \cdot (0,1) = 0$$
 on $L_{\text{sym}} = \left\{ \boldsymbol{\xi} \in \mathbb{R}^2 : \eta = 0 \right\}$.

2.1.3. Normal reflection configurations. It is meaningful to extend the range of parameters in Problem 2.4 to allow $\boldsymbol{\theta} \in [0, \theta^{cr})^2$. We study the reflection for the case: $\boldsymbol{\theta} = \mathbf{0}$, which we call the normal reflection configuration. It is clear that setting $\varphi = \bar{\varphi} \equiv \varphi_2$ would satisfy Problem 2.4(i) at infinity. However, since φ_2 does not satisfy Problem 2.4(ii), a boundary layer must be present. For this reason, we introduce a uniform downstream state (0), determined by a pseudo-potential φ_0 satisfying Problem 2.4(ii), such that a straight reflected shock S_0 is formed between states (2) and (0). In fact, the only possible straight reflected shock that satisfies 2.4(i) is a normal shock $S_0 \coloneqq \{\boldsymbol{\xi} \in \mathbb{R}^2_+ : \eta = \eta_0\}$ for some $\eta_0 > 0$, which is parallel to L_{sym} . The constant density of state (0) given by $\rho_0 \coloneqq \rho(|D\varphi_0|, \varphi_0)$ should satisfy the entropy condition $\rho_0 > 1$. We demonstrate that state (0) described above is uniquely determined by (γ, v_2) .

The pseudo-potential φ_0 has form (2.4) with i = 0, for suitable constants (u_0, v_0, k_0) to be determined. It follows from Problem 2.4(ii) that $v_0 = 0$, whilst $u_0 = 0$ follows from the Rankine–Hugoniot conditions (2.3) between states (2) and (0) because S_0 is parallel to L_{sym} . The value of $k_0 = v_2\eta_0$ is then obtained by using the continuity of $\varphi = \varphi_0 = \varphi_2$ on S_0 .

It remains to determine constant $\eta_0 > 0$, which fixes the location of S_0 . Combining the Bernoulli law (2.2) with the Rankine–Hugoniot conditions (2.3), we find the necessary condition (2.22) $\ell(\rho_0) = -v_2$.

It is clear that (2.22) admits a unique solution $\rho_0 \in (1, \infty)$ depending only on (γ, v_2) . Indeed, $\ell(\cdot)$ is strictly increasing on $(1, \infty)$ and $\ell(1) = 0$. Constant η_0 is then uniquely determined by the Rankine–Hugoniot conditions (2.3) to be

(2.23)
$$\eta_0 \coloneqq -\frac{v_2}{\rho_0 - 1} > 0$$
 where $\rho_0 \in (1, \infty)$ satisfies (2.22).

Thus, the pseudo-potential of state (0) has been uniquely determined above by (γ, v_2) as

$$arphi_0(oldsymbol{\xi})\coloneqq -rac{1}{2}|oldsymbol{\xi}|^2+v_2\eta_0\,.$$

It is direct to verify that

(2.24)
$$\varphi_{\text{norm}}(\boldsymbol{\xi}) \coloneqq \min\{\varphi_0(\boldsymbol{\xi}), \varphi_2(\boldsymbol{\xi})\} = \begin{cases} \varphi_2(\boldsymbol{\xi}) & \text{for } \eta > \eta_0, \\ \varphi_0(\boldsymbol{\xi}) & \text{for } 0 < \eta < \eta_0, \end{cases}$$

is a solution to Problem 2.4 when $\theta = 0$; see Fig. 2.2(a).

Briefly, we mention the case: $\boldsymbol{\theta} \in (\{0\} \times (0, \theta^{cr})) \cup ((0, \theta^{cr}) \times \{0\})$, which is called the unilateral normal reflection; a normal reflection on one side with respect to the symmetric boundary L_{sym} occurs by the same argument as the normal reflection case $\boldsymbol{\theta} = \mathbf{0}$ above, while the other side undergoes a regular reflection at the reflection point, of which details will be discussed in §2.2 below; see the case: $\boldsymbol{\theta} \in (0, \theta^{cr}) \times \{0\}$ in Fig. 2.2(b). Moreover, the unilateral normal reflections are essentially the same as the Prandtl-Meyer reflection configurations considered in [2].



(a) Normal reflection when $\theta_1 = \theta_2 = 0$. (b) Unilateral normal reflection when $\theta_1 > \theta_2 = 0$.

FIG. 2.2. Structure of solutions of Problem 2.4 involving normal shocks.

2.2. **Regular reflection configurations.** In this section, we introduce the detachment and sonic angles and describe the structure of three genuinely different configurations of the four-shock interactions.

2.2.1. Detachment and sonic angles. For $\boldsymbol{\theta} \in (0, \theta^{cr})^2$, the incident shocks S_{12} and S_{32} intersect the symmetric line L_{sym} at points P_0^1 and P_0^2 respectively, with coordinates given by (2.16). As discussed above, the symmetric line can be regarded as a rigid wall and, similarly to the previous works [10], we expect that, if the incident angle θ_1 is less than a detachment angle θ^d , there exists a regular reflection of S_{12} at point P_0^1 , *i.e.*, there exist both a uniform state (5) determined by a pseudo-potential φ_5 and a straight reflected shock S_{25} passing through P_0^1 such that S_{25} separates the uniform upstream state (2) from the uniform downstream state (5). Similarly, we expect that there exists a regular reflection of S_{32} at point P_0^2 if θ_2 is less than θ^d ; that is, there exist both a uniform state (6) determined by a pseudo-potential φ_6 and a straight reflected shock S_{26} passing through P_0^2 . The expected structure of shock regular reflection on the upper half-plane can be seen in Fig. 2.1. We now determine the detachment angle θ^d and the sonic angle θ^s .

The regular reflection of S_{12} at point P_0^1 can be reduced to the study of an algebraic system. By the argument in [10, §7.1], to determine the uniform state (5), it suffices to apply the Rankine– Hugoniot conditions at the reflection point P_0^1 only. The pseudo-potential φ_5 has form (2.4) with i = 5, and the straight reflected shock S_{25} must have the form:

$$S_{25} \coloneqq \left\{ \boldsymbol{\xi} \in \overline{\mathbb{R}^2_+} : \varphi_5(\boldsymbol{\xi}) = \varphi_2(\boldsymbol{\xi}) \right\} = \left\{ \boldsymbol{\xi} \in \overline{\mathbb{R}^2_+} : \eta = \boldsymbol{\xi} \tan \theta_{25} + a_{25} \right\}$$

for some constants $(u_5, v_5, k_5, \theta_{25}, a_{25})$ depending on (γ, v_2, θ_1) to be determined, where θ_{25} is the angle between the reflected shock S_{25} and the positive ξ -axis, and a_{25} is the η -intercept of the reflected shock S_{25} . Applying the Rankine–Hugoniot conditions (2.3) between states (2) and (5) at P_0^1 , we have

(2.25)
$$\varphi_5(P_0^1) = \varphi_2(P_0^1), \quad D\varphi_5(P_0^1) \cdot \boldsymbol{\tau}_{25} = D\varphi_2(P_0^1) \cdot \boldsymbol{\tau}_{25}, \quad \rho_5 D\varphi_5(P_0^1) \cdot \boldsymbol{\nu}_{25} = D\varphi_2(P_0^1) \cdot \boldsymbol{\nu}_{25},$$

where $\boldsymbol{\nu}_{25} \coloneqq \frac{D(\varphi_2 - \varphi_5)(P_0^1)}{|D(\varphi_2 - \varphi_5)(P_0^1)|}$, $\boldsymbol{\tau}_{25} \coloneqq \boldsymbol{\nu}^{\perp}$, and $\rho_5 \coloneqq \rho(|D\varphi_5|, \varphi_5)$ is given by (2.2). Furthermore, the pseudo-potential φ_5 should satisfy the entropy condition:

(2.26) $\rho_5 > 1$, or equivalently, $D\varphi_2(P_0^1) \cdot \boldsymbol{\nu}_{25} > D\varphi_5(P_0^1) \cdot \boldsymbol{\nu}_{25} > 0$;

and the slip boundary condition:

$$(2.27) D\varphi_5 \cdot (0,1) = 0 on L_{\text{sym}}$$

A necessary condition for the regular reflection of S_{12} at P_0^1 is clearly that state (2) must be pseudo-supersonic at P_0^1 . From Lemma 2.3, there exists a unique $\theta^+ \in (0, \theta^{\rm cr}]$ such that $M_2^{(\theta_1)} > 1$ if and only if $\theta_1 \in (0, \theta^+)$. However, note that $M_2^{(\theta_1)}|_{\theta_1=\theta^+}$ is not necessarily 1 if $\theta^+ = \theta^{\rm cr}$.

Introduce the steady detachment angle $\theta_{\text{stdy}}^{d}(\rho_{\infty}, u_{\infty})$ for the steady potential flow with constant supersonic upstream state $U_{\infty} = (\rho_{\infty}, u_{\infty}, 0)$, as defined in [10, §7.1]. Also, for $\theta_1 \in (0, \theta^{\text{cr}})$, denote by $\hat{\theta}_{25}(\theta_1)$ the acute angle between the pseudo-velocity $D\varphi_2(P_0^1) = (-\xi^{P_0^1}, v_2)$ and the symmetric line L_{sym} :

(2.28)
$$\hat{\theta}_{25}(\theta_1) \coloneqq \arccos\left(\frac{\xi^{P_0^1}}{|D\varphi_2(P_0^1)|}\right) = \arcsin\left(\frac{-v_2}{|D\varphi_2(P_0^1)|}\right) \in (0, \frac{\pi}{2}).$$

Similarly to [10, §7.4], whenever $\theta_1 \in (0, \theta^+)$, the existence and multiplicity of solutions of the algebraic system (2.25)–(2.27) are determined as follows:

- (a) If $\hat{\theta}_{25}(\theta_1) < \theta_{\text{stdy}}^{\text{d}}(1, |D\varphi_2(P_0^1)|)$, there are two solutions;
- (b) If $\hat{\theta}_{25}(\theta_1) = \theta_{\text{stdy}}^{\text{d}}(1, |D\varphi_2(P_0^1)|)$, there is one solution;
- (c) If $\hat{\theta}_{25}(\theta_1) > \theta_{\text{stdy}}^{\text{d}}(1, |D\varphi_2(P_0^1)|)$, there are no solutions.

We demonstrate that there exists at most one value $\theta^{d} \in (0, \theta^{+})$ such that $\theta_{1} = \theta^{d}$ satisfies the equality in (b) above. Indeed, using (2.28) and Lemma 2.3, it is clear that $\hat{\theta}_{25}(\theta_{1})$ is a strictly increasing function of $\theta_{1} \in (0, \theta^{cr})$, whilst, for $\theta^{d}_{stdy}(1, |D\varphi_{2}(P_{0}^{1})|)$, we state below a general fact about 2-D steady potential flow, from which it is clear that $\theta^{d}_{stdy}(1, |D\varphi_{2}(P_{0}^{1})|)$ is strictly decreasing with $\theta_{1} \in (0, \theta^{+})$. The proof of this statement is given in Appendix A.1.

Lemma 2.5. Let $\theta_{\text{stdy}}^{\text{d}}$ and $\theta_{\text{stdy}}^{\text{s}}$ denote the steady detachment and sonic angles for the steady potential flow with constant supersonic upstream state $U_{\infty} = (\rho_{\infty}, u_{\infty}, 0)$. Then $\theta_{\text{stdy}}^{\text{d}}$ and $\theta_{\text{stdy}}^{\text{s}}$ are smooth, strictly increasing functions of the upstream Mach number $M_{\infty} \coloneqq \frac{u_{\infty}}{c(\rho_{\infty})} > 1$. Moreover, the following limits hold:

$$\lim_{M_{\infty} \to 1^{+}} (\theta_{\text{stdy}}^{\text{d}}, \theta_{\text{stdy}}^{\text{s}}) = (0, 0), \qquad \lim_{M_{\infty} \to \infty} (\theta_{\text{stdy}}^{\text{d}}, \theta_{\text{stdy}}^{\text{s}}) = (\frac{\pi}{2}, \arctan\sqrt{\frac{2}{\gamma - 1}}).$$

With the above discussion, the following proposition gives the necessary and sufficient criterion on the incident angle θ_1 for the existence of a regular reflection of S_{12} at point P_0^1 .

Proposition 2.6 (Local reflection theory). Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. There exists a unique $\theta^{d} = \theta^{d}(\gamma, v_2) \in (0, \theta^{cr}]$, called the detachment angle, such that, for $\theta_1 \in (\theta^{d}, \theta^{cr})$, there are no states (5) of form (2.4) satisfying (2.25)–(2.27) and, for each $\theta_1 \in (0, \theta^{d})$, there are exactly two states (5) of form (2.4) satisfying (2.25)–(2.27): the weak reflection state φ_5^{sg} , distinguished by $1 < \rho_5^{sg} < \rho_5^{sg}$, where $\rho_5^{sk} \coloneqq \rho(|D\varphi_5^{sk}|, \varphi_5^{sk})$ and $\rho_5^{sg} \coloneqq \rho(|D\varphi_5^{sg}|, \varphi_5^{sg})$ are the constant densities of the weak and strong reflection states (5) respectively. Furthermore, denoting by $O_5^{sk} \coloneqq (u_5^{sk}, v_5^{sk})$ and $O_5^{sg} \coloneqq (u_5^{sg}, v_5^{sg})$ the sonic centres, and by c_5^{sk} and c_5^{sg} the sonic speeds of the weak and strong reflection states (5) respectively, then

(i) $(\rho_5^{\text{wk}}, O_5^{\text{wk}}) \in C([0, \theta^d]) \cap C^{\infty}([0, \theta^d])$ and $(\rho_5^{\text{sg}}, O_5^{\text{sg}}) \in C((0, \theta^d]) \cap C^{\infty}((0, \theta^d))$. In particular, the following limits exist:

$$\lim_{\theta_1 \to 0^+} (\rho_5^{\text{wk}}, O_5^{\text{wk}}, u_5^{\text{wk}} \xi^{P_0^1}) = (\rho_0, \mathbf{0}, -v_2 \eta_0), \qquad \lim_{\theta_1 \to \theta^{d-}} (\rho_5^{\text{wk}}, \rho_5^{\text{sg}}, O_5^{\text{wk}}, O_5^{\text{sg}}),$$

where ρ_0 and η_0 are the density and shock location of the normal reflection state (0) which are given in §2.1.3.

(ii) For any $\theta_1 \in (0, \theta^d)$,

(

(2.29)
$$|D\varphi_2(P_0^1)| > |D\varphi_2(P_0^1|_{\theta_1 = \theta^d})| > 1,$$

(2.30)
$$0 < u_5^{\text{wk}} < u_5^{\text{sg}} < \xi^{P_0^1}, \quad v_5^{\text{wk}} = v_5^{\text{sg}} = 0.$$

(iii) There exists a unique $\theta^{s} = \theta^{s}(\gamma, v_{2}) \in (0, \theta^{d}]$, called the sonic angle, such that

$$\begin{aligned} |D\varphi_5^{\text{wk}}(P_0^1)| &> c_5^{\text{wk}} \qquad \text{for all } \theta_1 \in (0, \theta^{\text{s}}) \,, \\ |D\varphi_5^{\text{wk}}(P_0^1)| &< c_5^{\text{wk}} \qquad \text{for all } \theta_1 \in (\theta^{\text{s}}, \theta^{\text{d}}] \,. \end{aligned}$$

Furthermore, if $\theta^{s} < \theta^{cr}$, then $\theta^{s} < \theta^{d}$ and $|D\varphi_{5}^{wk}(P_{0}^{1})| = c_{5}^{wk}$ when $\theta_{1} = \theta^{s}$.

Proof. The proof follows similarly to [10, Theorem 7.1.1], once θ^{d} and θ^{s} are determined. Thus, it remains to show the existence and uniqueness of θ^{d} and θ^{s} .

As discussed above, $\hat{\theta}_{25}(\theta_1)$ is strictly increasing with $\theta_1 \in (0, \theta^{cr})$, whilst $\theta_{stdy}^d(1, |D\varphi_2(P_0^1)|)$ is strictly decreasing with $\theta_1 \in (0, \theta^+)$. Furthermore, Lemmas 2.3 and 2.5 show that

$$\lim_{\theta_1 \to 0^+} \hat{\theta}_{25}(\theta_1) = 0 < \frac{\pi}{2} = \lim_{\theta_1 \to 0^+} \theta^{\mathrm{d}}_{\mathrm{stdy}}(1, |D\varphi_2(P_0^1)|)$$

Then it follows that there exists at most one solution $\vartheta^{d} \in (0, \theta^{+})$ of the equation:

(2.31)
$$\hat{\theta}_{25}(\vartheta^{\mathrm{d}}) = \theta^{\mathrm{d}}_{\mathrm{stdy}}(1, |D\varphi_2(P_0^1|_{\theta_1 = \vartheta^{\mathrm{d}}})|)$$

We define the detachment angle $\theta^{d} \in (0, \theta^{+}]$ by

$$\theta^{d} \coloneqq \min \left\{ \theta^{+}, \inf \left\{ \vartheta^{d} \in (0, \theta^{+}) : \vartheta^{d} \text{ satisfies } (2.31) \right\} \right\}.$$

In this sense, θ^{d} may refer to either the conventional detachment angle ϑ^{d} that satisfies equation (2.31) when such a solution exists, or angle θ^{+} otherwise.

The existence and properties of the sonic angle θ^{s} contained in statement (iii) of Proposition 2.6 follow from a similar argument as above. In particular, symbol θ^{s} may refer to either the conventional sonic angle $\vartheta^{s} \in (0, \theta^{d}]$ which should satisfy

(2.32)
$$\theta_{25}(\vartheta^{\mathrm{s}}) = \theta_{\mathrm{stdy}}^{\mathrm{s}}(1, |D\varphi(P_0^1|_{\theta_1 = \vartheta^{\mathrm{s}}})|);$$

or we simply define $\theta^{s} = \theta^{d}$ if no such solution exists. That is, we define

$$\theta^{\mathbf{s}} \coloneqq \min \left\{ \theta^{\mathbf{d}}, \inf \{ \vartheta^{\mathbf{s}} \in (0, \theta^{\mathbf{d}}) : \vartheta^{\mathbf{s}} \text{ satisfies } (2.32) \} \right\}.$$

Remark 2.1. We have the following remarks about Proposition 2.6:

(i) Fix $\gamma \geq 1$. For any $v_2 \in (v_{\min}, 0)$, let θ^{cr} be the critical angle given by (2.15), and let $v_2^s, v_2^d \in (v_{\min}, 0)$ be the constants given by Lemma A.1. Then

$$\begin{aligned} \theta^{\mathrm{s}} &= \theta^{\mathrm{d}} = \theta^{\mathrm{cr}} & \Longleftrightarrow \quad v_2 \in (v_{\min}, v_2^{\mathrm{s}}] \,, \\ \theta^{\mathrm{s}} &< \theta^{\mathrm{d}} = \theta^{\mathrm{cr}} & \Longleftrightarrow \quad v_2 \in (v_2^{\mathrm{s}}, v_2^{\mathrm{d}}] \,, \\ \theta^{\mathrm{s}} &< \theta^{\mathrm{d}} < \theta^{\mathrm{cr}} & \Longleftrightarrow \quad v_2 \in (v_2^{\mathrm{d}}, 0) \,. \end{aligned}$$

- (ii) By the symmetry of the problem, the conclusions of Proposition 2.6 immediately apply to the regular reflection of S₃₂ at P₀² after exchanging (θ₁, P₀¹, (5)) with (θ₂, P₀², (6)), and reversing the inequalities in (2.30). In particular, the same detachment angle θ^d(γ, v₂) and sonic angle θ^s(γ, v₂) apply to the incident angle θ₂ and state (6).
- (iii) The main difference between Proposition 2.6 and [10, Theorem 7.1.1] is that we assert the uniqueness of solutions to (2.31) and (2.32), although we cannot in general assert the existence of such solutions due to the restriction $\theta_1 \in (0, \theta^{cr})$; whereas [10, Theorem 7.1.1] asserts the existence, but not the uniqueness of such solutions.

Definition 2.7 (Admissible parameters and states (5) and (6)). Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. Let $\theta^{s} \coloneqq \theta^{s}(\gamma, v_2)$ and $\theta^{d} \coloneqq \theta^{d}(\gamma, v_2)$ be given by Proposition 2.6. Denote the set of admissible parameters

$$\Theta \coloneqq [0, \theta^{\mathrm{d}})^2 \setminus \{\mathbf{0}\} \subseteq [0, \theta^{\mathrm{cr}}]^2$$
.

Hereafter, for $\boldsymbol{\theta} \in \overline{\Theta}$, states (5) and (6) refer to the weak reflection states (5) and (6) from Proposition 2.6. For j = 5, 6, we write $(\rho_j, c_j, u_j, v_j) \coloneqq (\rho_j^{\text{wk}}, c_j^{\text{wk}}, u_j^{\text{wk}}, v_j^{\text{wk}})$, $O_j \coloneqq (u_j, v_j)$, and $\varphi_j \coloneqq \varphi_j^{\text{wk}}$. The pseudo-potential of state (j) is given by

(2.33)
$$\varphi_j(\boldsymbol{\xi}) = -\frac{1}{2}|\boldsymbol{\xi}|^2 + (u_j, v_j) \cdot \boldsymbol{\xi} + v_2 a_{2j},$$

where $a_{2j} \coloneqq -\frac{1}{v_2} \xi^{P_0^{j-4}} u_j$ for $\theta_{j-4} > 0$, and $a_{2j} \coloneqq \eta_0$ when $\theta_{j-4} = 0$ (cf. Proposition 2.6(i)). Define the reflected shocks S_{25} and S_{26} by

$$S_{2j} \coloneqq \left\{ \boldsymbol{\xi} \in \overline{\mathbb{R}^2_+} : \varphi_j(\boldsymbol{\xi}) = \varphi_2(\boldsymbol{\xi}) \right\} = \left\{ \boldsymbol{\xi} \in \overline{\mathbb{R}^2_+} : \eta = \xi \tan \theta_{2j} + a_{2j} \right\} \quad \text{for } j = 5, 6,$$

where θ_{25} and θ_{26} are the angles between the reflected shocks S_{25} and S_{26} and the positive ξ -axis, given respectively by

(2.34)
$$\theta_{25} \coloneqq \arctan\left(\frac{u_5}{v_2}\right) \in \left(\frac{\pi}{2}, \pi\right], \qquad \theta_{26} \coloneqq \arctan\left(\frac{u_6}{v_2}\right) \in \left[0, \frac{\pi}{2}\right).$$

The unit tangent vectors to S_{25} and S_{26} are given respectively by

(2.35)
$$\boldsymbol{e}_{S_{25}} \coloneqq (\cos \theta_{25}, \sin \theta_{25}), \qquad \boldsymbol{e}_{S_{26}} \coloneqq (\cos \theta_{26}, \sin \theta_{26}).$$

It follows from Proposition 2.6(i) and Definition 2.7 that $\varphi_j = \varphi_0$, and $\theta_{2j} = (6-j)\pi$ when $\theta_{j-4} = 0$ for j = 5, 6.

Lemma 2.8 (Further properties of states (5) and (6)). Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. Let ρ_5 and ρ_6 be the constant densities of states (5) and (6) given by Proposition 2.6, and let θ_{25} and θ_{26} be the reflected shock angles given by (2.34). Then, for j = 5, 6, there exists a constant $\delta_{5,6} > 0$ depending only on (γ, v_2) such that, for any $\boldsymbol{\theta} \in \overline{\Theta}$,

$$(2.36) 1 + \delta_{5,6} \le \rho_j \le \delta_{5,6}^{-1}$$

(2.37)
$$\delta_{5,6} \le |\theta_{2j} - \frac{\pi}{2}|.$$

Moreover, for any $\bar{\theta} \in (0, \theta^d)$, there exists $\delta_{5,6}^{(\bar{\theta})} > 0$ depending only on $(\gamma, v_2, \bar{\theta})$ such that

(2.38)
$$\delta_{5,6}^{(\theta)} < |\theta_{2j} - (6-j)\pi| \qquad \text{whenever } \boldsymbol{\theta} \in \overline{\Theta} \cap \{\theta_{j-4} > \overline{\theta}\}.$$

Proof. We prove the results only for state (5), since the proof for state (6) is similar.

Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. By Proposition 2.6(i), we see that $\rho_5 \in C(\overline{\Theta})$, so that the uniform upper bound in (2.36) is clear. For the uniform lower bound, the entropy condition (2.26) gives $\rho_5 > 1$. Furthermore, the Rankine–Hugoniot conditions between states (2) and (5) imply that

(2.39)
$$\ell(\rho_5) = |(u_2, v_2) - (u_5, v_5)| = \sqrt{v_2^2 + u_5^2} \ge |v_2| > 0 \quad \text{for any } \boldsymbol{\theta} \in \overline{\Theta},$$

with $\ell(\cdot)$ given by (2.11). In particular, $\ell(\cdot)$ is an increasing function on $[1, \infty)$ with $\ell(1) = 0$, and hence (2.39) implies the existence of a small constant $\delta_{5,6}^{(1)} > 0$, depending only on (γ, v_2) , such that $\rho_5 > 1 + \delta_{5,6}^{(1)}$ for any $\boldsymbol{\theta} \in \overline{\Theta}$.

Next, we prove (2.37). From (2.34) and the geometric properties of the shock polar curve for 2-D steady potential flow, we have

(2.40)
$$\frac{\pi}{2} + \hat{\theta}_{25} < \theta_{25} \le \pi \qquad \text{for any } \boldsymbol{\theta} \in \overline{\Theta}$$

with $\hat{\theta}_{25}$ given by (2.28). From Proposition 2.6(i) and (2.34), we see that $\theta_{25} \to \pi^-$ as $\theta_1 \to 0^+$. By continuity, there exists a small constant $\delta_{5,6}^{(2)} > 0$ depending only on (γ, v_2) such that $\theta_{25} > \frac{3\pi}{4}$ whenever $\theta_1 \in [0, \delta_{5,6}^{(2)})$. On the other hand, by (2.28) and Lemma 2.3, there exists a constant $\delta_{5,6}^{(3)} > 0$ depending only on (γ, v_2) such that $\hat{\theta}_{25} > \delta_{5,6}^{(3)}$ whenever $\theta_1 \in [\delta_{5,6}^{(2)}, \theta^d]$. Together with (2.40), we conclude that $\theta_{25} > \min\{\frac{3\pi}{4}, \frac{\pi}{2} + \delta_{5,6}^{(3)}\}$ for all $\theta_1 \in [0, \theta^d]$, which leads to (2.37).

Finally, bound (2.38) follows directly from (2.40) and Lemma A.2, which states that θ_{25} is a strictly decreasing function of $\theta_1 \in [0, \theta^d]$.

2.2.2. Configurations for four-shock interactions. The four-shock interaction configurations are split into three genuinely different cases, depending on parameters $\boldsymbol{\theta} \in \Theta$ and the critical angles θ^{s} and θ^{d} . We briefly describe the expected structure of each configuration and introduce some important notation.

Case I. Supersonic-supersonic reflection configuration. When $\boldsymbol{\theta} \in (0, \theta^{s})^{2}$, the supersonic-supersonic reflection configuration is expected; see Fig. 2.3. In this case, both states (5) and (6) are pseudo-supersonic at the reflection points P_{0}^{1} and P_{0}^{2} , respectively.

Case II. Supersonic-subsonic reflection configuration. When $\boldsymbol{\theta} \in [\theta^{s}, \theta^{d}) \times (0, \theta^{s})$, or $\boldsymbol{\theta} \in (0, \theta^{s}) \times [\theta^{s}, \theta^{d})$, the supersonic-subsonic reflection configuration is expected; see Fig. 2.4 for the case: $\boldsymbol{\theta} \in [\theta^{s}, \theta^{d}) \times (0, \theta^{s})$. In this case, either state (5) is pseudo-sonic/pseudo-subsonic at P_{0}^{1} and state (6) is pseudo-supersonic at P_{0}^{2} , or vice versa.

Case III. Subsonic-subsonic reflection configuration. When $\theta \in [\theta^{s}, \theta^{d})^{2}$, the subsonic-subsonic reflection configuration is expected; see Fig. 2.5. In this case, both states (5) and (6) are pseudo-subsonic at the reflection points P_{0}^{1} and P_{0}^{2} , respectively.



FIG. 2.3. Case I. The supersonic-supersonic reflection configuration.



FIG. 2.4. **Case II.** The supersonic-subsonic reflection configuration in the case: $\boldsymbol{\theta} \in [\theta^{s}, \theta^{d}) \times (0, \theta^{s}), i.e., \text{ state } (5) \text{ is pseudo-sonic/pseudo-subsonic at } P_{0}^{1}.$



FIG. 2.5. Case III. The subsonic-subsonic reflection configuration.

It follows from Remark 2.1(i) that the configurations of **Case II** and **Case III** are not possible unless $v_2 \in (v_2^s, 0)$, for constant v_2^s from Lemma A.1.

Definition 2.9 (The sonic boundaries and intersection points). Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For any $\boldsymbol{\theta} \in \overline{\Theta}$, let Q_5 and Q_6 be the orthogonal projections of O_5 and O_6 onto S_{25} and S_{26} , respectively. When $\theta_1 \in (0, \theta^{\rm s})$, the right sonic boundary and intersection points are given by

$$\Gamma^{5}_{\text{sonic}} \coloneqq \left\{ \boldsymbol{\xi} \in \partial B_{c_{5}}(O_{5}) \cap \overline{\mathbb{R}^{2}_{+}} : \boldsymbol{\xi}^{Q_{5}} \leq \boldsymbol{\xi} \leq \boldsymbol{\xi}^{P^{1}_{0}}, \, \varphi_{5}(\boldsymbol{\xi}) \leq \varphi_{2}(\boldsymbol{\xi}) \right\}, \\ \{P_{1}\} \coloneqq \Gamma^{5}_{\text{sonic}} \cap L_{\text{sym}}, \quad \{P_{2}\} \coloneqq \Gamma^{5}_{\text{sonic}} \cap S_{25},$$

whilst, for $\theta_1 \in [\theta^s, \theta^d)$,

$$\Gamma_{\text{sonic}}^5 \equiv \{P_1\} \equiv \{P_2\} \coloneqq \{P_0^1\}$$

Similarly, when $\theta_2 \in (0, \theta^s)$, the left sonic boundary and intersection points are given by

$$\Gamma_{\text{sonic}}^6 \coloneqq \left\{ \boldsymbol{\xi} \in \partial B_{c_6}(O_6) \cap \mathbb{R}^2_+ : \boldsymbol{\xi}^{P_0^4} \leq \boldsymbol{\xi} \leq \boldsymbol{\xi}^{Q_6}, \, \varphi_6(\boldsymbol{\xi}) \leq \varphi_2(\boldsymbol{\xi}) \right\}, \\ \{P_3\} \coloneqq \Gamma_{\text{sonic}}^6 \cap S_{26}, \quad \{P_4\} \coloneqq \Gamma_{\text{sonic}}^6 \cap L_{\text{sym}},$$

whilst, for $\theta_2 \in [\theta^s, \theta^d)$,

$$\Gamma_{\text{sonic}}^6 \equiv \{P_3\} \equiv \{P_4\} \coloneqq \{P_0^2\}.$$

The open regions are given by

$$\begin{split} \Omega_5 &\coloneqq \left\{ \boldsymbol{\xi} \in \mathbb{R}^2_+ \setminus \overline{B_{c_5}(O_5)} \,:\, \varphi_5(\boldsymbol{\xi}) < \varphi_2(\boldsymbol{\xi}), \, \boldsymbol{\xi} > \boldsymbol{\xi}^{P_2} \right\}, \\ \Omega_6 &\coloneqq \left\{ \boldsymbol{\xi} \in \mathbb{R}^2_+ \setminus \overline{B_{c_6}(O_6)} \,:\, \varphi_6(\boldsymbol{\xi}) < \varphi_2(\boldsymbol{\xi}), \, \boldsymbol{\xi} < \boldsymbol{\xi}^{P_3} \right\}. \end{split}$$

Finally, define the symmetric boundary

$$\Gamma_{\text{sym}} \coloneqq \left\{ (\xi, 0) \in L_{\text{sym}} : \xi^{P_4} < \xi < \xi^{P_1} \right\},\,$$

and the line segments

$$S_{25}^{\text{seg}} \coloneqq S_{25} \cap \left\{ \xi^{P_2} \le \xi \le \xi^{P_0^1} \right\}, \qquad S_{26}^{\text{seg}} \coloneqq S_{26} \cap \left\{ \xi^{P_0^2} \le \xi \le \xi^{P_3} \right\}.$$

Note that curves Γ_{sonic}^5 , Γ_{sonic}^6 , and $\overline{\Gamma_{\text{sym}}}$ as defined in Definition 2.9 above are fixed by $(\gamma, v_2, \boldsymbol{\theta})$ and do not share any common points except their endpoints $\{P_1, P_4\}$. Also, note that $\Omega_j = \emptyset$ if $\theta_{j-4} \in [\theta^{\text{s}}, \theta^{\text{d}})$ for j = 5, 6. For a suitable simple open curve segment Γ_{shock} (yet to be determined) with endpoints P_2 and P_3 , we use Ω to denote the open bounded domain enclosed by the simple closed curve $\Gamma_{\text{shock}} \cup \Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sym}} \cup \Gamma_{\text{sonic}}^6$; see Figs. 2.3–2.5.

2.3. Admissible solutions and main theorems. We reformulate the asymptotic boundary value problem, Problem 2.4, into a free boundary problem. Our goal is to find a suitable curved free boundary Γ_{shock} between points P_2 and P_3 , and a solution φ of Problem 2.4 which satisfies (2.1)–(2.2) in the open region Ω (as depicted in Figs. 2.3–2.5) and the corresponding boundary conditions on $\partial \Omega := \Gamma_{\text{shock}} \cup \Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sym}} \cup \Gamma_{\text{sonic}}^6$. Furthermore, equation (2.1) should be elliptic in Ω , and the free boundary Γ_{shock} should be a transonic shock that separates the hyperbolic and elliptic regions of the solution.

Problem 2.10 (Free boundary problem). Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For any $\boldsymbol{\theta} \in \overline{\Theta}$, find a free boundary (curved reflected-diffracted shock) Γ_{shock} in region $\mathbb{R}^2_+ \cap \{\xi^{P_3} < \xi < \xi^{P_2}\}$ and a function φ defined in region Ω , as shown in Figs. 2.3–2.5, such that φ satisfies:

- (i) Equation (2.1) in Ω so that (2.1) is elliptic inside Ω ;
- (ii) $\varphi = \varphi_2$ and $\rho D \varphi \cdot \boldsymbol{\nu} = \rho_2 D \varphi_2 \cdot \boldsymbol{\nu}$ on the free boundary Γ_{shock} ;
- (iii) $\varphi = \varphi_j$ and $D\varphi = D\varphi_j$ on Γ_{sonic}^j for j = 5, 6;
- (iv) $D\varphi \cdot (0,1) = 0$ on Γ_{sym} ,

where $\boldsymbol{\nu}$ is the unit normal vector on Γ_{shock} pointing into Ω .

Definition 2.11 (Admissible solutions). Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For any $\theta \in \overline{\Theta}$, a function $\varphi \in W^{1,\infty}_{\text{loc}}(\mathbb{R}^2_+)$ is called an admissible solution to the four-shock interaction problem if φ is a solution of Problem 2.10 and satisfies the following properties:

(i) There exists a relatively open curve segment Γ_{shock} (without self-intersection) with endpoints P₂ = (ξ^{P₂}, η^{P₂}) and P₃ = (ξ^{P₃}, η^{P₃}) given by Definition 2.9 such that Γ_{shock} is C² in its relative interior, Γ^{ext}_{shock} := S^{seg}₂₅ ∪ Γ_{shock} ∪ S^{seg}₂₆ is a C¹-curve including at P₂ and P₃, with S^{seg}₂₅ and S^{seg}₂₆ given in Definition 2.9,

$$\Gamma_{ ext{shock}} \subseteq \left\{ oldsymbol{\xi} \in \mathbb{R}^2_+ \setminus \overline{B_{c_2}(O_2)} \, : \, oldsymbol{\xi}^{P_3} < oldsymbol{\xi} < oldsymbol{\xi}^{P_2}
ight\},$$

with $\partial B_{c_2}(O_2)$ as the sonic circle of state (2), and curves $\overline{\Gamma_{\text{shock}}}$ and $\Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sym}} \cup \Gamma_{\text{sonic}}^6$ share no common points except their endpoints $\{P_2, P_3\}$ so that $\Gamma_{\text{shock}} \cup \Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sym}} \cup \Gamma_{\text{sonic}}^6$ is a closed curve without self-intersection.

(ii) For the open bounded domain $\Omega \subseteq \mathbb{R}^2_+$ enclosed by $\Gamma_{\text{shock}} \cup \Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sym}} \cup \Gamma_{\text{sonic}}^6$, solution $\varphi \text{ is in } C^3(\Omega) \cap C^2(\overline{\Omega} \setminus (\Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6)) \cap C^1(\overline{\Omega}) \text{ and satisfies } \varphi \in C^{0,1}_{\text{loc}}(\mathbb{R}^2_+) \cap C^1_{\text{loc}}(\overline{\mathbb{R}^2_+} \setminus \overline{\Gamma_{\text{shock}}^{\text{ext}} \cup S_{12} \cup S_{32}}) \text{ and}$

(2.41)
$$\varphi = \begin{cases} \max\{\varphi_5, \varphi_6\} & \text{ in } \Omega_5 \cup \Omega_6, \\ \bar{\varphi} & \text{ in } \mathbb{R}^2_+ \setminus (\Omega \cup \Omega_5 \cup \Omega_6), \end{cases}$$

where Ω_5 and Ω_6 are given in Definition 2.9, and $\bar{\varphi}$ is given by (2.20).

(iii) Equation (2.1) is strictly elliptic in $\overline{\Omega} \setminus (\Gamma^5_{\text{sonic}} \cup \Gamma^6_{\text{sonic}})$, i.e., solution φ satisfies

$$|D\varphi(\boldsymbol{\xi})| < c(|D\varphi(\boldsymbol{\xi})|, \varphi(\boldsymbol{\xi})) \qquad for \ all \ \boldsymbol{\xi} \in \overline{\Omega} \setminus (\Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6) \ .$$

- (iv) In Ω , solution φ satisfies
- (2.42) $\max\{\varphi_5, \varphi_6\} \le \varphi \le \varphi_2,$
- (2.43) $D(\varphi_2 \varphi) \cdot (\cos \theta, \sin \theta) \le 0 \quad \text{for all } \theta \in [\theta_{26}, \theta_{25}].$

Remark 2.2. From the definition of admissible solutions, we have

- (i) The requirement that φ is a solution of Problem 2.10 includes the condition that $\varphi = \varphi_j$ and $D\varphi = D\varphi_j$ on Γ^j_{sonic} for j = 5, 6, which becomes a one-point boundary condition in the case $\theta_{j-4} \in (\theta^{\text{s}}, \theta^{\text{d}}]$, according to the definition of Γ^j_{sonic} given by Definition 2.9.
- (ii) Let φ be an admissible solution in the sense of Definition 2.11, and let ν be the unit normal vector on Γ_{shock} interior to Ω . Similarly to [2, Lemma 2.26], it can be shown that Γ_{shock} is a transonic shock, since φ satisfies

$$D arphi_2 \cdot oldsymbol{
u} > D arphi \cdot oldsymbol{
u} > 0 \,, \quad 0 < rac{D arphi \cdot oldsymbol{
u}}{c(|D arphi|, arphi)} < 1 < D arphi_2 \cdot oldsymbol{
u} \qquad on \ \Gamma_{
m shock} \,.$$

(iii) It follows from Proposition 2.6(i) that φ_j depends continuously on $\theta_{j-4} \in [0, \theta^d]$ for j = 5, 6. Moreover, for any r > 0,

$$\lim_{\theta_{j-4}\to 0^+} \|\varphi_j - \varphi_0\|_{C^{0,1}(B_r(\mathbf{0})\cap\mathbb{R}^2_+)} = 0 \qquad \text{for } j = 5, 6 \,,$$

i.e., the weak reflection state (j) coincides with the normal reflection state (0) when $\theta_{j-4} = 0$. One can verify directly that the normal reflection solution $\varphi_{\text{norm}} \in C^{0,1}_{\text{loc}}(\mathbb{R}^2_+)$ as defined in (2.24) is an admissible solution corresponding to $\boldsymbol{\theta} = \mathbf{0}$ in the sense of Definition 2.11.

The first main theorem of this paper is to provide the global existence of such admissible solutions in the sense of Definition 2.11. The proof is given in §4.4.

Theorem 2.1 (Existence of admissible solutions). Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For any $\boldsymbol{\theta} \in \Theta$, there exists an admissible solution φ of the four-shock interaction problem corresponding to parameters $\boldsymbol{\theta}$ in the sense of Definition 2.11. Moreover, the global solution φ converges to the normal reflection solution φ_{norm} in $W^{1,1}_{\text{loc}}(\mathbb{R}^2_+)$ as $\boldsymbol{\theta} \in \Theta$ converges to 0, where φ_{norm} is given by (2.24).

The admissible solutions have additional regularity properties, given in Theorem 2.2 below. Moreover, the transonic shocks, as free boundaries of admissible solutions, satisfy additional convexity properties, given in Theorem 2.3 below. The proofs of Theorems 2.2–2.3 are given in §5.1–§5.2, respectively.

Theorem 2.2 (Regularity of admissible solutions). Fix $\gamma \geq 1$, $v_2 \in (v_{\min}, 0)$, and $\theta \in \Theta$. Let φ be an admissible solution corresponding to parameters θ in the sense of Definition 2.11, with transonic shock Γ_{shock} . Then the following properties hold:

(i) Γ_{shock} is C^{∞} in its relative interior and

Case I. When $\boldsymbol{\theta} \in [0, \theta^{\mathrm{s}})^2$, $\varphi \in C^{\infty}(\overline{\Omega} \setminus (\Gamma_{\mathrm{sonic}}^5 \cup \Gamma_{\mathrm{sonic}}^6)) \cap C^{1,1}(\overline{\Omega})$, and $\overline{S_{25}^{\mathrm{seg}} \cup \Gamma_{\mathrm{shock}} \cup S_{26}^{\mathrm{seg}}}$ is a $C^{2,\alpha}$ -curve for any $\alpha \in (0, 1)$, including at points P_2 and P_3 .

Case II. When $\theta \in [\theta^{s}, \theta^{d}) \times [0, \theta^{s}), \varphi \in C^{\infty}(\overline{\Omega} \setminus (\Gamma_{\text{sonic}}^{6} \cup \{P_{1}\})) \cap C^{1,1}(\overline{\Omega} \setminus \{P_{1}\}) \cap C^{1,\bar{\alpha}}(\overline{\Omega}),$ and $\overline{S_{26}^{\text{seg}} \cup \Gamma_{\text{shock}}}$ is a $C^{1,\bar{\alpha}}$ -curve for some $\bar{\alpha} \in (0,1)$. Furthermore, $\overline{S_{26}^{\text{seg}} \cup \Gamma_{\text{shock}}} \setminus \{P_{1}\}$ is a $C^{2,\alpha}$ -curve for any $\alpha \in (0,1)$, including at P_{3} .

 $\begin{array}{l} \hline When \ \boldsymbol{\theta} \in [0, \theta^{\mathrm{s}}) \times [\theta^{\mathrm{s}}, \theta^{\mathrm{d}}), \ \varphi \in C^{\infty}(\overline{\Omega} \setminus (\Gamma_{\mathrm{sonic}}^{5} \cup \{P_{4}\})) \cap C^{1,1}(\overline{\Omega} \setminus \{P_{4}\}) \cap C^{1,\bar{\alpha}}(\overline{\Omega}), \ and \\ \hline S_{25}^{\mathrm{seg}} \cup \Gamma_{\mathrm{shock}} \ is \ a \ C^{1,\bar{\alpha}}-curve \ for \ some \ \bar{\alpha} \in (0,1). \ Furthermore, \ \overline{S_{25}^{\mathrm{seg}} \cup \Gamma_{\mathrm{shock}}} \setminus \{P_{4}\} \ is \ a \ C^{2,\alpha}-curve \ for \ any \ \alpha \in (0,1), \ including \ at \ P_{2}. \end{array}$

Case III. When $\boldsymbol{\theta} \in [\theta^{s}, \theta^{d})^{2}$, $\varphi \in C^{\infty}(\overline{\Omega} \setminus \{P_{1}, P_{4}\}) \cap C^{1,\overline{\alpha}}(\overline{\Omega})$, and $\overline{\Gamma_{shock}}$ is a $C^{1,\overline{\alpha}}$ -curve for some $\overline{\alpha} \in (0, 1)$. Furthermore, $\overline{\Gamma_{shock}} \setminus \{P_{1}, P_{4}\}$ is a $C^{2,\alpha}$ -curve for any $\alpha \in (0, 1)$.

(ii) Let $\mathcal{U} \coloneqq \{\boldsymbol{\xi} \in \mathbb{R}^2_+ : \max\{\varphi_5(\boldsymbol{\xi}), \varphi_6(\boldsymbol{\xi})\} < \varphi_2(\boldsymbol{\xi})\}$. For any constant $\sigma > 0$ and j = 5, 6, define domain $\mathcal{U}^{\sigma}_{\sigma}$ by

$$\mathcal{U}_{\sigma}^{j} \coloneqq \begin{cases} \mathcal{U} \cap \left\{ \boldsymbol{\xi} \in \mathbb{R}_{+}^{2} : \operatorname{dist}(\boldsymbol{\xi}, \Gamma_{\operatorname{sonic}}^{j}) < \sigma \right\} \cap B_{c_{j}}(O_{j}) & \text{if } \theta_{j-4} \in [0, \theta^{\operatorname{s}}), \\ \emptyset & \text{if } \theta_{j-4} \in [\theta^{\operatorname{s}}, \theta^{\operatorname{d}}). \end{cases}$$

Fix any point $\boldsymbol{\xi}_0 \in (\Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6) \setminus \{P_2, P_3\}$ and denote $d \coloneqq \text{dist}(\boldsymbol{\xi}_0, \Gamma_{\text{shock}})$. Then, for a sufficiently small constant $\varepsilon_0 > 0$ and for any $\alpha \in (0, 1)$, there exists a constant $K < \infty$

depending on $(\gamma, v_2, \varepsilon_0, \alpha, d)$ and $\|\varphi\|_{C^{1,1}(\Omega \cap (\mathcal{U}^5_{\varepsilon_0} \cup \mathcal{U}^6_{\varepsilon_0}))}$ such that

$$\left\|\varphi\right\|_{2,\alpha,\overline{\Omega\cap B_{d/2}(\boldsymbol{\xi}_0)\cap(\mathcal{U}^5_{\varepsilon_0/2}\cup\mathcal{U}^6_{\varepsilon_0/2})}}\leq K\,;$$

(iii) For any $\boldsymbol{\xi}_0 \in (\Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6) \setminus \{P_2, P_3\},\$

$$\lim_{T \to \boldsymbol{\xi}_0, \, \boldsymbol{\xi} \in \Omega} (D_{rr} \varphi - D_{rr} \max\{\varphi_5, \varphi_6\})(\boldsymbol{\xi}) = \frac{1}{\gamma + 1} \,,$$

where $r = |\boldsymbol{\xi} - O_j|$ for $\boldsymbol{\xi}$ near Γ_{sonic}^j for j = 5, 6; (iv) For i = 1, 2, whenever $\theta_i \in [0, \theta^{\text{s}})$, $\lim_{\boldsymbol{\xi} \to P_{i+1}, \boldsymbol{\xi} \in \Omega} D^2 \varphi(\boldsymbol{\xi})$ does not exist.

Theorem 2.3 (Convexity of transonic shocks). Fix $\theta \in \Theta$. For any admissible solution of the four-shock interaction problem in the sense of Definition 2.11, the transonic shock $\Gamma_{\rm shock}$ is uniformly convex on closed subsets of its relative interior in the sense described in Theorems B.4– B.5. Furthermore, for any weak solution of (2.1)–(2.2) in the sense of Definition 2.1, satisfying also all the properties of Definition 2.11 except (2.43), the transonic shock $\Gamma_{\rm shock}$ is a strictly convex graph if and only if condition (2.43) holds.

3. Uniform Estimates for Admissible Solutions

To establish the first main theorem, Theorem 2.1, we apply the Leray-Schauder degree theorem to a suitable iteration map, similar to [2, 10]. To achieve this, in this section, we make several essential uniform estimates of the admissible solutions. In §4, we construct a suitable iteration map between two function spaces with weighted Hölder norms, where any fixed point of the iteration map is an admissible solution in the sense of Definition 2.11. The main results we establish in this section are the following:

- (a) The strict directional monotonicity of $\varphi_2 \varphi$, and the directional monotonicity of $\varphi \varphi_5$ and $\varphi - \varphi_6$;
- (b) The uniform positive lower bound of dist($\Gamma_{\text{shock}}, \partial B_1(O_2)$);
- (c) The uniform estimate of the ellipticity of equation (2.1) in Ω ;
- (d) The uniform estimates of admissible solutions in some suitable weighted Hölder norms.

Our problem shares many similarities with the Prandtl-Meyer shock reflection problem [2], and the von Neumann shock reflection-diffraction problem [10]. Many of the following proofs are similar to those within [2, 10], but have been adapted to the current setting with careful estimates.

3.1. Monotonicity and ellipticity.

3.1.1. Directional monotonicity and uniform bounds. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. Let φ be an admissible solution corresponding to parameters $\theta \in \Theta$ in the sense of Definition 2.11, and let the constant states φ_2 and φ_j , j = 5, 6, be given by (2.17) and (2.33) respectively. We aim to establish the directional monotonicity properties for ϕ that satisfies

(3.1)
$$(c^2 - \varphi_{\xi}^2)\phi_{\xi\xi} - 2\varphi_{\xi}\varphi_{\eta}\phi_{\xi\eta} + (c^2 - \varphi_{\eta}^2)\phi_{\eta\eta} = 0 \quad \text{in } \Omega$$

where ϕ can be taken as either $\varphi_2 - \varphi$ or $\varphi - \varphi_j$, j = 5, 6, and

(3.2)
$$c^{2}(|D\varphi|,\varphi) = 1 + (\gamma - 1)\left(B - \frac{1}{2}|D\varphi|^{2} - \varphi\right),$$

where B is the Bernoulli constant given by (2.5).

Lemma 3.1. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For $\theta \in \Theta$, let φ be an admissible solution in the sense of Definition 2.11 corresponding to parameters θ . Then φ satisfies that, for j = 5, 6,

- (i) $\partial_{\boldsymbol{e}}(\varphi \varphi_j)$ is not constant in Ω for any unit vector $\boldsymbol{e} \in \mathbb{R}^2$;
- (ii) For vector $\boldsymbol{e}_{S_{2i}}$ given by (2.35),

$$\partial_{\boldsymbol{e}_{S_{2i}}}(\varphi_2-\varphi)<0 \quad in \ \overline{\Omega}\setminus\Gamma^j_{\text{sonic}};$$

(iii) $\partial_{\boldsymbol{e}_{S_{2j}}}(\varphi - \varphi_j), -\partial_{\eta}(\varphi - \varphi_j) \ge 0$ in $\overline{\Omega}$.

Proof. The proof of (i) is given by a slight modification of [2, Lemma 3.1]. Indeed, for $\phi := \varphi - \varphi_5$, if $\partial_{\boldsymbol{e}}\phi$ is constant in Ω , then $\partial_{\boldsymbol{e}}\phi = 0$ in Ω since $D\phi = \mathbf{0}$ at P_1 . Furthermore, $\partial_{\boldsymbol{e}}\phi = \boldsymbol{e} \cdot (u_6 - u_5, 0)$ at P_4 so that \boldsymbol{e} must be parallel to (0, 1), since $u_5 - u_6 > 0$ when $\boldsymbol{\theta} \neq \mathbf{0}$. Thus, $\partial_{\eta}\phi \equiv 0$ in Ω and hence $\partial_{\xi\eta}\phi = \partial_{\eta\eta}\phi \equiv 0$ in Ω . By the strict ellipticity of equation (3.1) in Ω , $\partial_{\xi\xi}\phi \equiv 0$ in Ω , which implies that there exist constants (u, v, k) such that $\phi = u\xi + v\eta + k$ in Ω . We note that v = 0 since $\partial_{\eta}\phi \equiv 0$. Applying the boundary conditions: $\phi = 0$ and $D\phi = \mathbf{0}$ at P_1 , we find that u = k = 0, so that $\phi \equiv 0$ in Ω . However, $D\phi(P_4) = D\varphi_6(P_4) - D\varphi_5(P_4) = (u_6 - u_5, 0) \neq \mathbf{0}$, which is a contradiction. The proof is similar for $\phi = \varphi - \varphi_6$. This concludes the proof of (i).

The proofs of (ii) and (iii) are similar to [2, Lemma 3.2] and [2, Lemma 3.6], respectively. \Box

For any $\theta \in \Theta$, define the following fan-shaped set by

$$\operatorname{Cone}(\boldsymbol{e}_{S_{25}}, \boldsymbol{e}_{S_{26}}) \coloneqq \{r(\cos\theta, \sin\theta) : \theta_{26} \le \theta \le \theta_{25}, r \ge 0\}$$

Let $\text{Cone}^{0}(\boldsymbol{e}_{S_{25}}, \boldsymbol{e}_{S_{26}})$ be the interior of $\text{Cone}(\boldsymbol{e}_{S_{25}}, \boldsymbol{e}_{S_{26}})$. It follows from Lemma 3.1 that, if φ is an admissible solution corresponding to $\boldsymbol{\theta} \in \Theta$, then φ satisfies

(3.3)
$$\partial_{\boldsymbol{e}}(\varphi_2 - \varphi) < 0 \quad \text{in } \overline{\Omega} \text{ for any } \boldsymbol{e} \in \operatorname{Cone}^0(\boldsymbol{e}_{S_{25}}, \boldsymbol{e}_{S_{26}})$$

and $-\boldsymbol{\nu}(P) \in \left\{ (\cos\theta, \sin\theta) : \theta_{25} - \frac{\pi}{2} < \theta < \theta_{26} + \frac{\pi}{2} \right\}$ for all $P \in \Gamma_{\text{shock}}$, where $\boldsymbol{\nu}$ is the unit normal vector on Γ_{shock} pointing into the interior of Ω .

Lemma 3.2. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For $\theta \in \Theta$, let φ be an admissible solution in the sense of Definition 2.11 with parameters θ . Then there exists a C^1 -function f_{sh} such that

$$\Gamma_{\text{shock}} = \{ \boldsymbol{\xi} \in \mathbb{R}^2_+ : \eta = f_{\text{sh}}(\xi), \, \xi^{P_3} < \xi < \xi^{P_2} \}$$

and $f'_{\rm sh}(\xi) \in (\tan \theta_{25}, \tan \theta_{26})$ for $\xi \in (\xi^{P_3}, \xi^{P_2})$ with $f'_{\rm sh}(\xi^{P_{j-3}}) = \tan \theta_{2j}$, for j = 5, 6.

Proof. By Lemma 2.8, $e_{\eta} = (0,1) \in \text{Cone}^{0}(e_{S_{25}}, e_{S_{26}})$. From the strict directional monotonicity (3.3), we see that $\partial_{e_{\eta}}(\varphi_{2} - \varphi) < 0$ on $\overline{\Gamma_{\text{shock}}}$, which implies the existence of a C^{1} -function f_{sh} that solves the implicit relation $\varphi_{2}(\xi, f_{\text{sh}}(\xi)) = \varphi(\xi, f_{\text{sh}}(\xi))$. Taking the derivative, we have

$$f_{\rm sh}'(\xi) = -\frac{\partial_{\boldsymbol{e}_{\xi}}(\varphi_2 - \varphi)(\xi, f_{\rm sh}(\xi))}{\partial_{\boldsymbol{e}_{\eta}}(\varphi_2 - \varphi)(\xi, f_{\rm sh}(\xi))} \,.$$

From Definition 2.11(i), we obtain that $f'_{\rm sh}(\xi^{P_2}) = \tan \theta_{25}$ and $f'_{\rm sh}(\xi^{P_3}) = \tan \theta_{26}$. Denoting by $\boldsymbol{\nu}(P)$ the unit normal vector on $\Gamma_{\rm shock}$ at a point $P \in \Gamma_{\rm shock}$, interior to Ω , we have

$$\boldsymbol{\nu}(P) = \frac{D(\varphi_2 - \varphi)(P)}{|D(\varphi_2 - \varphi)(P)|} = \frac{(f'_{\rm sh}(\xi), -1)}{\sqrt{1 + (f'_{\rm sh}(\xi))^2}}.$$

Combining the above with property (ii) in Lemma 3.1, for any $(a_1, a_2) \in [0, \infty)^2$, $(a_1, a_2) \neq (0, 0)$,

$$a_1(f'_{\rm sh}(\xi)\cos\theta_{25} - \sin\theta_{25}) + a_2(f'_{\rm sh}(\xi)\cos\theta_{26} - \sin\theta_{26})$$

= $\sqrt{1 + (f'_{\rm sh}(\xi))^2} \nu(P) \cdot (a_1 e_{S_{25}} + a_2 e_{S_{26}}) < 0$ for any $\xi \in (\xi^{P_3}, \xi^{P_2})$.

Choosing (a_1, a_2) as (1, 0) and (0, 1) in turn, we obtain that $f'_{sh}(\xi) > \tan \theta_{25}$ and $f'_{sh}(\xi) < \tan \theta_{26}$, respectively, where we have used $\theta_{25} \in (\frac{\pi}{2}, \pi]$ and $\theta_{26} \in [0, \frac{\pi}{2})$.

Lemma 3.3. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For any $\boldsymbol{\theta} \in \Theta$, let φ be an admissible solution corresponding to parameters $\boldsymbol{\theta} \in \Theta$ in the sense of Definition 2.11, and let Ω be the pseudo-subsonic region of φ . Then there exists a constant $C_{ub} > 0$ depending only on (γ, v_2) such that the following properties hold:

(3.4)
$$\overline{\Omega} \subseteq B_{C_{\rm ub}}(\mathbf{0}),$$

(3.5)
$$\|\varphi\|_{C^{0,1}(\overline{\Omega})} \le C_{\mathrm{ub}},$$

(3.6)
$$\rho^*(\gamma) \le \rho \le C_{\rm ub} \quad in \ \Omega \,, \qquad 1 < \rho \le C_{\rm ub} \quad on \ \Gamma_{\rm shock} \,,$$

where
$$\rho^*(\gamma) := (\frac{2}{\gamma+1})^{\frac{1}{\gamma-1}}$$
 when $\gamma > 1$, and $\rho^*(1) := e^{-\frac{1}{2}} = \lim_{\gamma \to 1+} \rho^*(\gamma)$

Proof. We divide the proof into three steps.

1. To prove (3.4), we use the expression of Γ_{shock} from Lemma 3.2. In particular, it follows from from Definition 2.11(i) and Lemma 3.2 that $0 < f_{\text{sh}}(\xi) \leq \min\{\xi \tan \theta_{25} + a_{25}, \xi \tan \theta_{26} + a_{26}\}$ whenever $\xi^{P_3} < \xi < \xi^{P_2}$, which implies that

$$\Omega \subseteq \{ \boldsymbol{\xi} : u_6 - c_6 < \boldsymbol{\xi} < u_5 + c_5, \, 0 < \eta \le \max\{a_{25}, a_{26}\} \}$$

From (2.16) and Proposition 2.6(i) and Definition 2.7, we know that (ρ_5, u_5, a_{25}) depend continuously on $\theta_1 \in [0, \theta^d]$, while (ρ_6, u_6, a_{26}) depend continuously on $\theta_2 \in [0, \theta^d]$, from which (3.4) follows directly.

2. From Definition 2.11(iv), we see that $\inf_{\overline{\Omega}} \max\{\varphi_5, \varphi_6\} \leq \varphi \leq \sup_{\overline{\Omega}} \varphi_2$. By (3.4) and the expressions of $(\varphi_2, \varphi_5, \varphi_6)$ given in (2.17) and (2.33), there exists a constant $C_1 > 0$ depending only on (γ, v_2) such that $-C_1 \leq \min_{\overline{\Omega}} \max\{\varphi_5, \varphi_6\} \leq \max_{\overline{\Omega}} \varphi_2 \leq C_1$, which implies that $\max_{\overline{\Omega}} |\varphi| \leq C_1$.

Then the uniform bound (3.5) follows from

$$|D\varphi|^2 < c^2(|D\varphi|,\varphi) = \frac{2}{\gamma+1} \left(1 + (\gamma-1)(B-\varphi)\right),$$

where we have used Definition 2.11(iii) and (3.2).

3. The upper bounds in (3.6) follow directly from the expression of density (2.2) along with (3.5). For the lower bound in Ω , we combine the Bernoulli law (2.2) with Definition 2.11(iv) to obtain

$$\frac{1}{2}|D\varphi|^2 + h(\rho) = \frac{1}{2}|D\varphi_2|^2 + (\varphi_2 - \varphi) + h(\rho_2) \ge \frac{1}{2}|D\varphi_2|^2 \ge 0.$$

Then, from Definition 2.11(iii), we have

$$\frac{1}{\gamma - 1} \left(\frac{1}{2} (\gamma + 1) \rho^{\gamma - 1} - 1 \right) = \frac{1}{2} c^2 + h(\rho) \ge \frac{1}{2} |D\varphi|^2 + h(\rho) \ge 0 \,,$$

so that the lower bound of ρ in Ω is obtained. For the lower bound of ρ on Γ_{shock} , we combine Problem 2.10(ii) and Remark 2.2(iii) to obtain

$$\rho = \frac{D\varphi_2 \cdot \boldsymbol{\nu}}{D\varphi \cdot \boldsymbol{\nu}} > 1 \quad \text{on } \Gamma_{\text{shock}} \,.$$

3.1.2. Uniform positive lower bound of dist($\Gamma_{\text{shock}}, \partial B_1(O_2)$). We now obtain a uniform estimate for the positive lower bound of dist($\Gamma_{\text{shock}}, \partial B_1(O_2)$) for any admissible solution. This allows us to make the uniform estimate for the ellipticity of equation (2.1) within the pseudo-subsonic domain of admissible solutions.

Proposition 3.4. Fix $\gamma \ge 1$ and $v_2 \in (v_{\min}, 0)$. There exists a constant $C_{\text{sh}} > 0$ depending only on (γ, v_2) such that any admissible solution corresponding to parameters $\boldsymbol{\theta} \in \Theta$ satisfies

(3.7)
$$\operatorname{dist}(\Gamma_{\operatorname{shock}}, \partial B_1(O_2)) \ge C_{\operatorname{sh}}^{-1}.$$

Proof. The proof of (3.7) follows the same argument as for [2, Proposition 3.7] except the case that $v_2 \leq -1$. In the following, we focus on the case that $v_2 \leq -1$ and give the proof in three steps.

1. We first rewrite equation (2.1) as

$$\operatorname{div}\mathcal{A}(D\varphi,\varphi) + \mathcal{B}(D\varphi,\varphi) = 0$$

where $\mathcal{A}(\boldsymbol{p}, z) \coloneqq \rho(|\boldsymbol{p}|, z)\boldsymbol{p}$ and $\mathcal{B}(|\boldsymbol{p}|, z) \coloneqq 2\rho(|\boldsymbol{p}|, z)$ with $\boldsymbol{p} = (p_1, p_2) \in \mathbb{R}^2, z \in \mathbb{R}$, and

(3.8)
$$\rho(|\mathbf{p}|, z) = \begin{cases} \left(1 + (\gamma - 1)(\frac{1}{2}v_2^2 - \frac{1}{2}|\mathbf{p}|^2 - z)\right)^{\frac{1}{\gamma - 1}} & \text{if } \gamma > 1, \\ \exp(\frac{1}{2}v_2^2 - \frac{1}{2}|\mathbf{p}|^2 - z) & \text{if } \gamma = 1. \end{cases}$$

We denote the sonic speed as $c(|\mathbf{p}|, z) \coloneqq \rho^{(\gamma-1)/2}$ for $\gamma \ge 1$. For a constant R > 1, define

(3.9)
$$\mathcal{K}_R \coloneqq \left\{ (\boldsymbol{p}, z) \in \mathbb{R}^2 \times \mathbb{R} : |\boldsymbol{p}| + |z| \le R, \, \rho(|\boldsymbol{p}|, z) \ge R^{-1}, \, \frac{|\boldsymbol{p}|^2}{c^2(|\boldsymbol{p}|, z)} \le 1 - R^{-1} \right\}.$$

For each R > 1, there exists a constant $\lambda_R > 0$ depending only on (γ, v_2, R) such that, for any $\boldsymbol{\kappa} = (\kappa_1, \kappa_2) \in \mathbb{R}^2,$

$$\sum_{i,j=1}^{2} \partial_{p_j} \mathcal{A}_i(\boldsymbol{p}, z) \kappa_i \kappa_j \geq \lambda_R |\boldsymbol{\kappa}|^2 \quad \text{for any } (\boldsymbol{p}, z) \in \mathcal{K}_R \,.$$

The next two lemmas follow directly from [2, Lemmas 3.8–3.9].

Lemma 3.5. For R > 2, let \mathcal{K}_R be given by (3.9). Then there exist functions $(\tilde{\mathcal{A}}, \tilde{\mathcal{B}})(\boldsymbol{p}, z)$ in $\mathbb{R}^2 \times \mathbb{R}$ satisfying the following properties:

- (i) For any $(\boldsymbol{p}_0, z_0) \in \mathcal{K}_R$, $(\tilde{\mathcal{A}}, \tilde{\mathcal{B}})(\boldsymbol{p}, z) = (\mathcal{A}, \mathcal{B})(\boldsymbol{p}, z)$ for all $(\boldsymbol{p}, z) \in B_{\varepsilon}((\boldsymbol{p}_0, z_0))$; (ii) For any $(\boldsymbol{p}, z) \in \mathbb{R}^2 \times \mathbb{R}$ and $\boldsymbol{\kappa} = (\kappa_1, \kappa_2) \in \mathbb{R}^2$, $\sum_{i,j=1}^2 \partial_{p_j} \tilde{\mathcal{A}}_i(\boldsymbol{p}, z) \kappa_i \kappa_j \ge \lambda_R |\boldsymbol{\kappa}|^2$; (iii) $|\tilde{\mathcal{B}}(\boldsymbol{p}, z)| \le C_0$ and $|D_{(\boldsymbol{p}, z)}^m(\tilde{\mathcal{A}}, \tilde{\mathcal{B}})(\boldsymbol{p}, z)| \le C_m$ in $\mathbb{R}^2 \times \mathbb{R}$ for each $m = 1, 2, \cdots$,

where the positive constants ε and λ_R depend only on (γ, v_2, R) , and C_m depends on (γ, v_2, R, m) .

Lemma 3.6. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For any given constants $\alpha \in (0, 1)$, $m \in \mathbb{N}$, and r > 0, there exist constants $C, C_m > 0$ depending only on (γ, v_2, α, r) , with C_m depending additionally on m, such that any admissible solution φ corresponding to parameters $\theta \in \Theta$ satisfies the following *estimates*:

(i) For any $B_{4r}(P) \subseteq \Omega$,

$$\begin{aligned} \|\varphi\|_{2,\alpha,\overline{B_{2r}(P)}} &\leq C \,, \qquad \|\varphi\|_{m,\overline{B_r(P)}} \leq C_m \,; \end{aligned}$$
(ii) If $P \in \Gamma_{\text{sym}}$, and $B_{4r}(P) \cap \Omega$ is the half-ball $B_{4r}^+(P) \coloneqq B_{4r}(P) \cap \{\eta > 0\}$, then
 $\|\varphi\|_{2,\alpha,\overline{B_{2r}(P)} \cap \Omega} \leq C \,, \qquad \|\varphi\|_{m,\overline{B_r(P)} \cap \Omega} \leq C_m \,. \end{aligned}$

2. Fix $\varepsilon > 0$ small enough such that $2\xi^{P_3} < \xi^{P_0}$ for all $\theta_2 \in [0,\varepsilon)$, where $\xi^{P_0} < 0$ is the ξ -coordinate of $P_0 \coloneqq \lim_{\theta_2 \to 0^+} P_3$, which depends only on (γ, v_2) . We define

(3.10)
$$d_{\text{sep}} \coloneqq \min\left\{-\frac{\xi^{P_0}}{2}, \inf_{\theta_2 \in [\varepsilon, \theta^d]} |u_6|\right\}$$

It is direct to verify that $d_{sep} > 0$ by using $u_6 = v_2 \tan \theta_{26}$ and property (2.38), and d_{sep} depends only on (γ, v_2) . Then, for any $\boldsymbol{\theta} \in \Theta$, dist $(P_2, P_3) \geq 2d_{sep}$. We also define

(3.11)
$$r_1 \coloneqq \frac{1}{2} \inf_{\theta \in \Theta} \left\{ |D\varphi_5(P_0^1)|, |D\varphi_6(P_0^2)| \right\}.$$

By (2.25) and Lemma 2.8, it is direct to verify that $r_1 > 0$ depends only on (γ, v_2) .

For any $r \in (0, r_1)$ and for i = 1, 2, we define two sets:

$$\mathscr{A} \coloneqq \left\{ \theta_i \in (0, \theta^{\mathrm{d}}] : |P_0^i - P_{i+1}| \ge \frac{r}{20} \right\} \cup \{0\}, \quad \mathscr{B} \coloneqq \left\{ \theta_i \in [0, \theta^{\mathrm{d}}] : |P_{3i-2} - P_{i+1}| \ge \frac{r}{20} \right\}.$$

From the continuous dependence of P_0^i, P_{3i-2} , and P_{i+1} on $\theta_i \in (0, \theta^d]$, we know that \mathscr{A} and \mathscr{B} both are closed sets. Then there exists a constant $C_1 > 0$ such that

$$\operatorname{dist}(P_{i+1}, \Gamma_{\operatorname{sym}}) \geq \frac{2}{C_1} \quad \text{for all } \theta_i \in \mathscr{A} \cup \mathscr{B}.$$

If dist $(P_{i+1}, \Gamma_{sym}) \leq \frac{1}{C_1}$, then $\theta_i \notin \mathscr{A} \cup \mathscr{B}$, which implies that

$$\max\{|P_0^i - P_{i+1}|, |P_{3i-2} - P_{i+1}|\} < \frac{r}{20}$$

We now seek a constant $C_r > 0$ depending only on (γ, v_2, r) such that, for any admissible solution of Problem 2.10 corresponding to parameters $\theta \in \Theta$,

(3.12)
$$\sup_{P \in \Gamma_{\text{shock}} \cap B_{r/2}(P_{3i-2})} \operatorname{dist}(P, \Gamma_{\text{sym}}) > C_r^{-1} \quad \text{if } |P_{3i-2} - P_{i+1}| \le \frac{r}{10}.$$

To obtain the above result, we follow the proof as in [10, Lemma 9.4.2], in which we need to take a limit of parameters $\boldsymbol{\theta}^{(n)}$.

In order to take the limit, we use a compactness lemma below. For concreteness, in the following, we use the notation defined in Definition 2.9, and append superscripts (n) and * to indicate that the corresponding quantities are related to parameters $\theta^{(n)}$ and θ^* , respectively.

Lemma 3.7. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. Let $\{\boldsymbol{\theta}^{(n)}\}_{n \in \mathbb{N}} \subseteq \Theta$ be any sequence of parameters satisfying $\lim_{n\to\infty} \theta^{(n)} = \theta^*$ for some $\theta^* \in \overline{\Theta}$. For each $n \in \mathbb{N}$, let $\varphi^{(n)}$ be an admissible solution corresponding to parameters $\theta^{(n)}$. Then there exists a subsequence $\{\varphi^{(n_j)}\}_{j\in\mathbb{N}}$ converging uniformly on any compact subset of \mathbb{R}^2_+ to a function $\varphi^* \in C^{0,1}_{\text{loc}}(\mathbb{R}^2_+)$. Moreover, φ^* is a weak solution of equation (2.1) in \mathbb{R}^2_+ and satisfies the following properties:

(i) For the corner points P_m , m = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4, and the reflection points P_0^i , i = 1, 2, 3, 4.

$$\lim_{j \to \infty} P_m^{(n_j)} = P_m^*, \qquad \lim_{j \to \infty} P_0^{i,(n_j)} = P_0^{i,*}$$

Note that $\xi^{P_0^{1,*}} = \infty$ in the case that $\theta_1^* = 0$, and $\xi^{P_0^{2,*}} = -\infty$ in the case that $\theta_2^* = 0$. (ii) Let $\eta = f_{2l}^{(n_j)}(\xi)$, l = 5, 6, and $\eta = f_{\rm sh}^{(n_j)}(\xi)$ be the expressions of $S_{2l}^{(n_j)}$ and $\Gamma_{\rm shock}^{(n_j)}$ from

Definition 2.7 and Lemma 3.2, respectively. Extend $f_{\rm sh}^{(n_j)}$ by

$$f_{\rm sh}^{(n_j)}(\xi) = \begin{cases} f_{25}^{(n_j)}(\xi) & \text{for } \xi^{P_2^{(n_j)}} < \xi < \xi^{P_0^{1,(n_j)}}, \\ f_{26}^{(n_j)}(\xi) & \text{for } \xi^{P_0^{2,(n_j)}} < \xi < \xi^{P_3^{(n_j)}}. \end{cases}$$

- Then $\{f_{\rm sh}^{(n_j)}\}_{j\in\mathbb{N}}$ converges uniformly to a function $f_{\rm sh}^* \in C^{0,1}([\xi^{P_0^{2,*}}, \xi^{P_0^{1,*}}])$. (iii) Let smooth functions $\eta = g_{l,\rm so}^{(n_j)}(\xi)$ be the expressions of $\Gamma_{\rm sonic}^{l,(n_j)}$ for l = 5, 6. Then $\{g_{5,\rm so}^{(n_j)}\}_{j\in\mathbb{N}}$ converges pointwise to $g_{5,so}^*$ on $(\xi^{P_2^*}, \xi^{P_1^*})$, and $\{g_{6,so}^{(n_j)}\}_{j\in\mathbb{N}}$ converges pointwise to $g_{6,so}^*$ on $(\xi^{P_4^*},\xi^{P_3^*}).$
- (iv) Denote $\widehat{\Omega^*} := \{ \boldsymbol{\xi} \in [\xi^{P_4^*}, \xi^{P_1^*}] \times \overline{\mathbb{R}_+} : 0 \le \eta \le f_{\mathrm{bd}}^*(\xi) \}, \text{ where } f_{\mathrm{bd}}^*(\cdot) \text{ is given by}$

$$f_{\rm bd}^*(\xi) = \begin{cases} g_{6,\rm so}^*(\xi) & \text{for } \xi^{P_4^*} \le \xi \le \xi^{P_3^*} ,\\ f_{\rm sh}^*(\xi) & \text{for } \xi^{P_3^*} < \xi \le \xi^{P_2} ,\\ g_{5,\rm so}^*(\xi) & \text{for } \xi^{P_2^*} < \xi \le \xi^{P_1^*} . \end{cases}$$

Denote by Ω^* the interior of $\widehat{\Omega^*}$, $\Gamma^*_{\text{shock}} \coloneqq \{(\xi, f^*_{\text{sh}}(\xi)) : \xi \in (\xi^{P_3^*}, \xi^{P_2^*})\}$, and $\Gamma^*_{\text{sym}} \coloneqq \{(\xi, 0) : \xi \in (\xi^{P_4^*}, \xi^{P_1^*})\}$, and denote by $\Gamma^{*,0}_{\text{sym}}$ the relative interior of $\Gamma^*_{\text{sym}} \setminus \Gamma^*_{\text{shock}}$. Then $\varphi^* \in C^{\infty}(\Omega^* \cup \Gamma^{*,0}_{\text{sym}}), \text{ and}$ (iv.1) $\lim_{j \to \infty} \|\varphi^{(n_j)} - \varphi^*\|_{C^2(K)} = 0 \text{ for any } K \subseteq \Omega^* \cup \Gamma^{*,0}_{\text{sym}},$

- (iv.2) $D(\varphi_2 \varphi^*) \cdot (\cos \theta, \sin \theta) \le 0 \text{ for all } \theta \in (\theta_{26}^*, \theta_{25}^*),$
- (iv.3) Equation (2.1) is strictly elliptic in $\Omega^* \cup \Gamma^{*,0}_{sym}$,
- (iv.4) In $\mathbb{R}^2_+ \setminus \Omega^*$, φ^* is equal to the constant states $\bar{\varphi}^*$, φ_5^* , and φ_6^* in their respective domains as given in Definition 2.11(ii), where $\bar{\varphi}^*$ is given by (2.20) with $\theta = \theta^*$. (v) $\Gamma^{*,0}_{\text{sym}} = \{\varphi_2 > \varphi^*\} \cap \Gamma^*_{\text{sym}}$ is dense in Γ^*_{sym} .

Using Lemmas 3.3 and 3.6, properties (i)-(iv) in Lemma 3.7 can be proved by similar arguments as for [2, Corollary 3.10].

For property (v), if it is not true, we can choose $\xi_1, \xi_2 \in (\xi^{P_3^*}, \xi^{P_2^*})$ such that $\{(\xi, 0) : \xi_1 < \xi < \xi_2\} \subseteq \{\varphi_2 \le \varphi^*\} \cap \Gamma^*_{\text{sym}}$. By Definition 2.11(iv), $\varphi^{(n_j)} \le \varphi_2$ in $\Omega^{(n_j)}$ so that $\varphi^* \le \varphi_2$ in $\overline{\Omega^*}$. Combining with property (iv.4), we obtain that $\varphi^*(\boldsymbol{\xi}) = \varphi_2(\boldsymbol{\xi})$ on $\{\boldsymbol{\xi} \in \mathbb{R}^2_+ : \xi_1 < \xi_2\}$. We take any non-negative function $\zeta \in C_{c}^{\infty}(\overline{\mathbb{R}^{2}_{+}})$ satisfying that $\operatorname{supp} \zeta \subseteq \{\boldsymbol{\xi} \in \overline{\mathbb{R}^{2}_{+}} : \xi_{1} < \xi < \xi_{2}\}$ and $\zeta > 0$ on some subset of $\{(\xi, 0) : \xi_{1} < \xi < \xi_{2}\}$ with positive measure. Then

$$\int_{\mathbb{R}^2_+} (\rho(|D\varphi^*|,\varphi^*)D\varphi^* \cdot D\zeta(\boldsymbol{\xi}) - 2\rho(|D\varphi^*|,\varphi^*)\zeta(\boldsymbol{\xi})) \,\mathrm{d}\boldsymbol{\xi}$$
$$= \int_{\mathbb{R}^2_+} (\rho_2 D\varphi_2 \cdot D\zeta(\boldsymbol{\xi}) - 2\rho_2\zeta(\boldsymbol{\xi})) \,\mathrm{d}\boldsymbol{\xi} = -v_2 \int_{\xi_1}^{\xi_2} \zeta(\xi,0) \,\mathrm{d}\boldsymbol{\xi} > 0$$

which contradicts the fact that φ^* is a weak solution of equation (2.1) in \mathbb{R}^2_+ . Then Lemma 3.7(v) is verified.

We prove (3.12) by contradiction. If it is not true, we can choose a sequence of parameters $\{\boldsymbol{\theta}^{(n)}\}_{n\in\mathbb{N}}$ such that

$$\sup_{P \in \Gamma_{\text{shock}}^{(n)} \cap B_{r/2}(P_{3i-2}^{(n)})} \operatorname{dist}(P, \Gamma_{\text{sym}}^{(n)}) \le \frac{1}{n} \quad \text{if } |P_{3i-2}^{(n)} - P_{i+1}^{(n)}| \le \frac{r}{10}$$

Then, by Lemma 3.7, we can further choose a subsequence $\{\theta^{(n_j)}\}_{j\in\mathbb{N}}$ and take a limit to some $\theta^* \in \overline{\Theta}$, which provides a contradiction to Lemma 3.7(v).

From (3.12), we can choose $\hat{Q}_i \in \overline{\Gamma_{\text{shock}} \cap B_{r/2}(P_{3i-2})}$ with $\operatorname{dist}(\hat{Q}_i, \Gamma_{\text{sym}}) \geq C_r^{-1}$, so that $\hat{Q}_i \in \overline{\Gamma_{\text{shock}} \cap B_{3r/4}(P_0^i)}$. For i = 1, 2, we take $Q_i = P_{i+1} \in \overline{\Gamma_{\text{shock}}}$ if $\operatorname{dist}(P_{i+1}, \Gamma_{\text{sym}}) > C_1^{-1}$; otherwise, we take $Q_i = \hat{Q}_i \in \overline{\Gamma_{\text{shock}} \cap B_{3r/4}(P_0^i)}$ as chosen above. Then it follows that

 $\min\{\operatorname{dist}(Q_1, Q_2), \operatorname{dist}(Q_1, \Gamma_{\operatorname{sym}}), \operatorname{dist}(Q_2, \Gamma_{\operatorname{sym}})\} \ge \min\{d_{\operatorname{sep}}, C_1^{-1}, C_r^{-1}\} \rightleftharpoons a.$

Combining with Lemma 3.7 and following a similar proof as for [10, Lemmas 9.4.1 and 15.4.1], we obtain that, for a > 0 defined above, there exists a constant $C_2 > 0$ depending only on (γ, v_2, a) such that, for any admissible solution corresponding to parameters $\boldsymbol{\theta} \in \Theta$,

$$\operatorname{dist}(\Gamma_{\operatorname{shock}}[Q_1, Q_2], \Gamma_{\operatorname{sym}}) \ge C_2^{-1},$$

where $\Gamma_{\text{shock}}[Q_1, Q_2]$ represents the segment on Γ_{shock} between points Q_1 and Q_2 . Therefore,

(3.13)
$$\operatorname{dist}(\Gamma_{\operatorname{shock}} \setminus (B_r(P_0^1) \cup B_r(P_0^2)), \Gamma_{\operatorname{sym}}) > C_2^{-1}$$

3. Since $v_2 \leq -1$, $B_1(O_2) \subseteq \mathbb{R} \times (-\infty, 0)$. By Lemma 2.3 and property (ii) of Proposition 2.6, $d_{\text{ref}} \coloneqq |D\varphi_2(P_0^1|_{\theta_1=\theta^d})| -1 > 0$ depending only on (γ, v_2) satisfies

$$\operatorname{dist}(P_0^1, B_1(O_2)) = |D\varphi_2(P_0^1)| - 1 \ge d_{\operatorname{ref}}.$$

By symmetry, we also see that $\operatorname{dist}(P_0^2, B_1(O_2)) \ge d_{\operatorname{ref}}$. Denote $\overline{r} := \frac{1}{2} \min\{r_1, d_{\operatorname{ref}}\} > 0$ with r_1 defined by (3.11). Then

$$\operatorname{dist}(\Gamma_{\operatorname{shock}} \cap (B_{\bar{r}}(P_0^1) \cup B_{\bar{r}}(P_0^2)), B_1(O_2)) \geq \bar{r}.$$

Meanwhile, from the argument in Step 2, there exists a constant $C_2 > 0$ depending only on (γ, v_2, \bar{r}) such that (3.13) holds with r replaced by \bar{r} . Thus, for any admissible solution corresponding to parameters $\boldsymbol{\theta} \in \Theta$, we have

$$\operatorname{dist}(\Gamma_{\operatorname{shock}}, B_1(O_2)) \ge \min\{\bar{r}, C_2^{-1}\} \rightleftharpoons C_{\operatorname{sh}}^{-1}$$

This completes the proof of (3.7).

3.1.3. Uniform estimate for the ellipticity of equation (3.1). We now show a uniform estimate for the ellipticity of equation (3.1), which may be degenerate near $\Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6$. Fix a function $h^* \in C^{\infty}(\mathbb{R})$ such that

$$h^*(s) = \begin{cases} s & \text{for } s \le \frac{1}{2}, \\ 1 & \text{for } s \ge 1, \end{cases}$$

satisfying $0 \leq (h^*)'(\cdot) \leq 2$ on \mathbb{R} . For each $\boldsymbol{\theta} \in \overline{\Theta}$, define

(3.14)
$$\hat{c}_j \coloneqq \min\{c_j, |D\varphi_j(P_0^{j-4})|\}, \quad q_j \coloneqq \operatorname{dist}(O_j, S_{2j}) \quad \text{for } j = 5, 6.$$

By Lemma 2.8 and [10, Lemma 6.1.2], we see that $\hat{c}_j > q_j$ for every $\boldsymbol{\theta} \in \overline{\Theta}$, so that there exists a constant $\hat{\delta} > 0$ depending only on (γ, v_2) such that

(3.15)
$$\inf_{\boldsymbol{\theta}\in\overline{\Theta}}\min\{\hat{c}_5-q_5,\,\hat{c}_6-q_6\}\geq\hat{\delta}\,.$$

For j = 5, 6, define $g_j^* : \mathbb{R}^2 \to \mathbb{R}_+$ by

$$g_j^*(\boldsymbol{\xi}) \coloneqq \frac{1}{2} (\hat{c}_j - q_j) h^* (\frac{\text{dist}(\boldsymbol{\xi}, \partial B_{\hat{c}_j}(O_j))}{\hat{c}_j - q_j}) + \left(1 - \frac{\hat{c}_j^2}{c_j^2}\right).$$

We choose a function $\chi \in C^{\infty}(\mathbb{R})$ such that

$$\chi(\xi) = \begin{cases} 1 & \text{for } \xi \leq -\frac{d_{\text{sep}}}{4} \\ 0 & \text{for } \xi \geq \frac{d_{\text{sep}}}{4} \end{cases},$$

satisfying $-\frac{4}{d_{sep}} \leq \chi'(\cdot) \leq 0$ on \mathbb{R} , where $d_{sep} > 0$ is given by (3.10) depending only on (γ, v_2) . Finally, define a function $g: \mathbb{R}^2 \to \mathbb{R}_+$ by

(3.16)
$$g(\boldsymbol{\xi}) \coloneqq \chi(\boldsymbol{\xi}) g_6^*(\boldsymbol{\xi}) + (1 - \chi(\boldsymbol{\xi})) g_5^*(\boldsymbol{\xi}) \,.$$

By Definition 2.11 and Lemma 3.3, there exist constants d > 0 and C > 1 depending only on (γ, v_2) such that, if φ is an admissible solution corresponding to $\theta \in \Theta$ with pseudo-subsonic region Ω , then g given by (3.16) satisfies the following properties:

- (a) For ξ ∈ Ω satisfying dist(ξ, Γ^j_{sonic}) < d, C⁻¹dist_j(ξ, Γ^j_{sonic}) ≤ g(ξ) ≤ C dist_j(ξ, Γ^j_{sonic}), where dist_j(ξ, Γ^j_{sonic}) := dist(ξ, Γ^j_{sonic}) + (c_j ĉ_j) for j = 5, 6.
 (b) For each ε > 0, there exists a constant C_ε > 1 depending only on (γ, v₂, ε) such that, if a point ξ ∈ Ω satisfies dist(ξ, Γ⁵_{sonic} ∪ Γ⁶_{sonic}) > ε, then C⁻¹_ε ≤ g(ξ) ≤ C_ε.

In fact, for property (a) above, it suffices to choose $d := \min\{\frac{d_{\text{sep}}}{2}, \hat{\delta}\}$, with constants d_{sep} and $\hat{\delta}$ given by (3.10) and (3.15) respectively. Then, choosing $\varepsilon \coloneqq \frac{d}{2}$ in property (b), we deduce that there exists a constant $C_{\flat} > 1$ depending only on (γ, v_2) such that, for any $\theta \in \Theta$,

(3.17)
$$C_{\flat}^{-1} \operatorname{dist}^{\flat}(\boldsymbol{\xi}, \Gamma_{\operatorname{sonic}}^{5} \cup \Gamma_{\operatorname{sonic}}^{6}) \leq g(\boldsymbol{\xi}) \leq C_{\flat} \operatorname{dist}^{\flat}(\boldsymbol{\xi}, \Gamma_{\operatorname{sonic}}^{5} \cup \Gamma_{\operatorname{sonic}}^{6})$$
 for all $\boldsymbol{\xi} \in \overline{\Omega}$,

where

(3.18)
$$\operatorname{dist}^{\flat}(\boldsymbol{\xi},\,\Gamma_{\operatorname{sonic}}^{5}\cup\Gamma_{\operatorname{sonic}}^{6}) \coloneqq \min\left\{\frac{d}{2},\,\operatorname{dist}_{5}(\boldsymbol{\xi},\,\Gamma_{\operatorname{sonic}}^{5}),\,\operatorname{dist}_{6}(\boldsymbol{\xi},\,\Gamma_{\operatorname{sonic}}^{6})\right\}.$$

Proposition 3.8. There exists a constant $\mu > 0$ such that, if φ is an admissible solution corresponding to $\boldsymbol{\theta} \in \Theta$ and Ω is its pseudo-subsonic region, then the pseudo-Mach number

$$M(\boldsymbol{\xi}) \coloneqq \frac{|D\varphi(\boldsymbol{\xi})|}{c(|D\varphi(\boldsymbol{\xi})|,\varphi(\boldsymbol{\xi}))}$$

satisfies

(3.19)
$$M^2(\boldsymbol{\xi}) \le 1 - \mu g(\boldsymbol{\xi}) \qquad in \ \overline{\Omega} \,,$$

and there exists a constant C > 1 such that

(3.20)
$$C^{-1}\operatorname{dist}^{\flat}(\boldsymbol{\xi},\,\Gamma^{5}_{\operatorname{sonic}}\cup\Gamma^{6}_{\operatorname{sonic}})|\boldsymbol{\kappa}|^{2} \leq \sum_{i,j=1}^{2}\mathcal{A}^{i}_{p_{j}}(D\varphi(\boldsymbol{\xi}),\varphi(\boldsymbol{\xi}))\kappa_{i}\kappa_{j} \leq C|\boldsymbol{\kappa}|^{2}$$

for all $\boldsymbol{\xi} \in \overline{\Omega}$ and $\boldsymbol{\kappa} = (\kappa_1, \kappa_2) \in \mathbb{R}^2$, where $\mathcal{A}(\boldsymbol{p}, z) = \rho(|\boldsymbol{p}|, z)\boldsymbol{p}$, dist^b (\cdot, \cdot) is given by (3.18), and constants μ and C depend only on (γ, v_2) .

Proof. Let φ be any admissible solution corresponding to $\theta \in \Theta$ with pseudo-subsonic region Ω and curved transonic shock $\Gamma_{\rm shock}$.

1. From Lemma 3.3, there exists a constant R > 1 depending only on (γ, v_2) such that

$$\Omega \subseteq B_R(\mathbf{0}), \qquad \|c(|D\varphi|,\varphi)\|_{C^0(\overline{\Omega})} + \|g\|_{C^2(\overline{\Omega})} \le R,$$

where $g(\cdot)$ is given by (3.16). Since $O_5, O_6 \in \Gamma_{\text{sym}}, \partial_\eta g = 0$ on Γ_{sym} . Then, by [10, Theorem 5.3.1], we can choose constants $C_0 > 1$, $\delta \in (0, \frac{3}{4C_0})$, and $\mu_1 \in (0, 1)$ depending only on (γ, v_2) such that, for any $\mu \in (0, \mu_1]$, either inequality $M^2 + \mu g \leq C_0 \delta < 1$ holds in Ω or the maximum of $M^2 + \mu g$ over $\overline{\Omega}$ cannot be attained in $\Omega \cup \Gamma_{\text{sym}}$.

In the first case, (3.19) follows immediately. Thus, we focus on the second case that the maximum of $M^2 + \mu g$ must be attained on $\partial \Omega \setminus \Gamma_{\text{sym}}$.

2. We claim that there exists a constant $\mu_2 \in (0, \mu_1]$ depending only on (γ, v_2) such that, whenever $\mu \in (0, \mu_2]$, $(M^2 + \mu g) \leq 1$ holds in Ω , from which (3.19) follows.

Let the maximum of $M^2 + \mu g$ be attained at $P_{\max} \in \partial \Omega \setminus \Gamma_{\text{sym}}$, and let $(M^2 + \mu g)(P_{\max}) > 1$. Then, using Proposition 3.4, we may proceed as in Steps 2–3 of the proof of [2, Proposition 3.15]

to find $\mu_2 \in (0, \mu_1]$ depending only on (γ, v_2) such that $P_{\max} \notin \Gamma_{\text{shock}}$ when $\mu \in (0, \mu_2]$. Therefore, $P_{\max} \in \Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6$. However, reducing μ_2 depending only on (γ, v_2) if necessary, we have

$$(M^2 + \mu g)(P_{\max}) = \sup_{\boldsymbol{\xi} \in \Gamma^5_{\text{sonic}} \cup \Gamma^6_{\text{sonic}}} (M^2 + \mu g)(\boldsymbol{\xi}) \le 1,$$

which is a contradiction. In the final inequality above, we have used that $g|_{\Gamma_{\text{sonic}}^{j}} = 0$ for $\theta_{j-4} \in [0, \theta^{\text{s}})$, whilst $g|_{\Gamma_{\text{sonic}}^{j}} \leq C_{\flat}(c_{j} - |D\varphi_{j}(P_{0}^{j-4})|)$ for $\theta_{j-4} \in [\theta^{\text{s}}, \theta^{\text{d}})$, for j = 5, 6.

3.2. Interior Hölder estimates away from the sonic boundaries. For any set $U \subseteq \mathbb{R}^2$ and any constant $\varepsilon > 0$, define the ε -neighbourhood of U as

(3.21)
$$\mathcal{N}_{\varepsilon}(U) \coloneqq \{ \boldsymbol{\xi} \in \mathbb{R}^2 : \operatorname{dist}(\boldsymbol{\xi}, U) < \varepsilon \}$$

From Proposition 3.4, there exists a constant $C_{\rm sh} > 0$ depending only on (γ, v_2) such that

dist(
$$\Gamma_{\text{shock}}, \partial B_1(O_2)$$
) $\geq C_{\text{sh}}^{-1}$

for any admissible solution and parameters $\theta \in \Theta$. It follows that

$$|D\varphi_2(\boldsymbol{\xi})|^2 \ge 1 + d_0$$
 for any $\boldsymbol{\xi} \in \mathcal{N}_{rac{1}{2C_{\mathrm{sh}}}}(\Gamma_{\mathrm{shock}})$

for some constant $d_0 > 0$ depending only on (γ, v_2) . Subsequently, we claim that there exist constants $d'_0 \in (0, d_0)$ and $\varepsilon \in (0, \frac{1}{2C_{\rm sh}})$ depending only on (γ, v_2) such that, for any admissible solution φ ,

(3.22)
$$|D\varphi_2(\boldsymbol{\xi})| - |D\varphi(\boldsymbol{\xi})| \ge d'_0 \quad \text{for any } \boldsymbol{\xi} \in \overline{\Omega} \cap \mathcal{N}_{\varepsilon}(\Gamma_{\text{shock}})$$

Indeed, in the case: $\gamma > 1$, from Definition 2.11(iii)–(iv) and the Bernoulli law:

$$\frac{1}{2}|D\varphi|^2 + \varphi + \frac{\rho^{\gamma - 1} - 1}{\gamma - 1} = B = \frac{1}{2}|D\varphi_2|^2 + \varphi_2,$$

we have

$$\begin{aligned} \frac{\gamma+1}{2} \left(\rho^{\gamma-1} - 1 \right) &\geq \left(\rho^{\gamma-1} - 1 \right) + \frac{\gamma-1}{2} \left(|D\varphi|^2 - 1 \right) \\ &= (\gamma-1)(\varphi_2 - \varphi) + \frac{\gamma-1}{2} \left(|D\varphi_2|^2 - 1 \right) \geq \frac{\gamma-1}{2} d_0 \,, \end{aligned}$$

and

$$|D\varphi_2|^2 - |D\varphi|^2 = \frac{2(\rho^{\gamma-1}-1)}{\gamma-1} + (\varphi-\varphi_2) \ge \frac{2d_0}{\gamma+1} + (\varphi-\varphi_2)$$

Since φ is Lipschitz continuous in $\overline{\Omega}$ from Lemma 3.3, and $\varphi = \varphi_2$ on Γ_{shock} , we obtain

$$|D\varphi_2(\boldsymbol{\xi})|^2 - |D\varphi(\boldsymbol{\xi})|^2 \ge rac{d_0}{\gamma+1} \qquad ext{for any } \boldsymbol{\xi} \in \overline{\Omega} \cap \mathcal{N}_{arepsilon}(\Gamma_{ ext{shock}}),$$

with some $\varepsilon \in (0, \frac{1}{2C_{\rm sh}})$ sufficiently small. Then (3.22) follows from the above inequality and the uniform boundedness property in Lemma 3.3. In the case: $\gamma = 1$, it is straightforward to obtain (3.22) by using Definition 2.11(iii): for any admissible solution φ , $|D\varphi|^2 \leq c^2(|D\varphi|, \varphi) \equiv 1$ in $\overline{\Omega}$, since the sonic speed is a constant.

For $\boldsymbol{\xi} = (\xi, \eta) \in \mathbb{R}^2 \setminus \{O_2\}$, define the polar coordinates (r, θ) centred at $O_2 = (u_2, v_2)$ by

$$r(\cos\theta, \sin\theta) \coloneqq (\xi, \eta) - O_2$$
.

It follows from (3.22) that

(3.23)
$$\partial_r(\varphi_2 - \varphi) \le -(|D\varphi_2| - |D\varphi|) \le -d'_0 \quad \text{in } \overline{\Omega} \cap \mathcal{N}_{\varepsilon}(\Gamma_{\text{shock}})$$

Therefore, we may apply the implicit function theorem to express Γ_{shock} under these polar coordinates as

$$\Gamma_{\text{shock}} = \left\{ \boldsymbol{\xi}(r,\theta) : r = f_{O_2,\text{sh}}(\theta), \, \theta_{P_2} < \theta < \theta_{P_3} \right\},\,$$

where $(f_{O_2,\text{sh}}(\theta_{P_i}), \theta_{P_i})$ are the polar coordinates of points P_i , for i = 2, 3. By Lemma 3.3 and (3.23), there exists a constant $C_1 > 0$ depending only on (γ, v_2) such that

$$||f_{O_2, \text{sh}}||_{C^{0,1}([\theta_{P_2}, \theta_{P_3}])} \le C_1$$

From the above, with the help of Lemma 3.5, we deduce the following properties; the proof is similar to that of [2, Lemmas 3.17–3.18].

Lemma 3.9. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. Let φ be an admissible solution corresponding to parameters $\theta \in \Theta$.

(i) There exists a constant $\delta' > 0$ depending only on (γ, v_2) such that

$$\partial_{\boldsymbol{\nu}}\varphi_2 - \delta' > \partial_{\boldsymbol{\nu}}\varphi \ge \delta' \qquad on \ \overline{\Gamma_{\mathrm{shock}}}$$

where $\boldsymbol{\nu} = \frac{D(\varphi_2 - \varphi)}{|D(\varphi_2 - \varphi)|}$ is the unit normal vector on Γ_{shock} pointing into the interior of Ω . (ii) For each d > 0 and $m = 2, 3, \cdots$, there exist positive constants s and C_m depending only

(ii) For each a > 0 and $m = 2, 3, \cdots$, there exist positive constants s and \mathbb{C}_m depending only on (γ, v_2, d) such that, whenever $P = \boldsymbol{\xi}(r_P, \theta_P) \in \Gamma_{\text{shock}}$ satisfies $\operatorname{dist}(P, \Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6) \ge d$,

$$|f_{O_2,\mathrm{sh}}^{(m)}(\theta_P)| + \sup_{B_s(P)\cap\Omega} |D^m\varphi| \le C_m$$

From Lemmas 3.6, 3.8, and 3.9(ii), we have the following *a priori* interior estimates.

Corollary 3.10. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For each d > 0 and $m = 2, 3, \cdots$, there exists a constant $C_{m,d} > 0$ depending only on (γ, v_2, m, d) such that any admissible solution φ corresponding to $\theta \in \Theta$ satisfies

$$\left\|\varphi\right\|_{m,\overline{\Omega\cap\{\mathrm{dist}(\boldsymbol{\xi},\Gamma^5_{\mathrm{sonic}}\cup\Gamma^6_{\mathrm{sonic}})>d\}}}\leq C_{m,d}\,.$$

3.3. Weighted Hölder estimates near the sonic boundaries. From Definition 2.9, the sonic boundary Γ_{sonic}^5 depends continuously on $\theta_1 \in [0, \theta^d)$ and shrinks into a single point $P_0^1 = P_1 = P_2$ when $\theta_1 \in [0, \theta^s)$ tends to θ^s from below (if additionally $v_2 \in [v_2^s, 0)$). Moreover, from Proposition 3.8, the ellipticity of equation (3.1) degenerates on the sonic boundary Γ_{sonic}^5 for $\theta_1 \in [0, \theta^s)$, while equation (3.1) is uniformly elliptic in the case that $\theta_1 \in (\theta^s, \theta^d)$ up to Γ_{sonic}^5 away from Γ_{sonic}^6 .

The above reasons lead us to consider the following four cases separately for the weighted Hölder estimates of the admissible solutions near Γ_{sonic}^5 . Since θ^{s} can be equal to θ^{cr} depending on (γ, v_2) , some of the following cases may not occur (*cf.* Remark 2.1(i)):

- Case 1. $\theta_1 \in [0, \theta^s)$ away from θ^s ; Case 2. $\theta_1 \in [0, \theta^s)$ close to θ^s ;
- Case 3. $\theta_1 \in [\theta^{s}, \theta^{d})$ close to θ^{s} ;
- Case 4. $\theta_1 \in [\theta^s, \theta^d)$ away from θ^s .

Corresponding weighted $C^{2,\alpha}$ -estimates near Γ_{sonic}^6 for the four cases with respect to $\theta_2 \in [0, \theta^d)$ are obtained immediately due to the symmetry of our problem.

For any $\boldsymbol{\theta} \in \Theta$, define an open set $\Omega^5 \subseteq \Omega$ by

$$\Omega^5 \coloneqq \left(\Omega \cap \{\boldsymbol{\xi} \in \mathbb{R}^2_+ : \boldsymbol{\xi} > u_5 + q_5 \sin \theta_{25}\}\right) \setminus \overline{B_{c_5^*}(O_5)},$$

where $q_5 = \text{dist}(O_5, S_{25})$ is given by (3.14), and $c_5^* := \frac{1}{2} (|O_5P_2| + \text{dist}(O_5, S_{25}))$. For $\boldsymbol{\xi} \in \overline{\Omega^5}$, we introduce the modified polar coordinates (x, y) centred at O_5 defined by

(3.24)
$$(x,y) \coloneqq \mathcal{R}(\xi,\eta) \quad \Longleftrightarrow \quad (\xi,\eta) - O_5 = (c_5 - x)(\cos y, \sin y) \quad \text{with } y \in [0, \frac{\pi}{2}].$$

It is clear that $\mathcal{R}(\Omega^5) \subseteq \{(x, y) : 0 < x < c_5 - c_5^*\}$. For the pseudo-potential φ_5 given by (2.33), we have

$$\varphi_5 \circ \mathcal{R}^{-1}(x,y) = -\frac{1}{2}(c_5 - x)^2 + \frac{1}{2}u_5^2 - u_5\xi^{P_0^1} \quad \text{for } (x,y) \in \mathcal{R}(\Omega^5).$$

Note that $u_5\xi^{P_0^1} \to -v_2\eta_0$ as $\theta_1 \to 0^+$, where $\eta_0 > 0$ is defined in §2.1.3.

For any admissible solution φ corresponding to parameters $\theta \in \Theta$, let ψ be given by

$$\psi(x,y) \coloneqq (\varphi - \varphi_5) \circ \mathcal{R}^{-1}(x,y) \quad \text{for } (x,y) \in \mathcal{R}(\Omega^5).$$

Then ψ satisfies a free boundary problem for an elliptic equation:

(i) Equation for ψ in $\mathcal{R}(\Omega^5)$: Direct calculation gives

(3.25)
$$\mathcal{N}(\psi) = (2x - (\gamma + 1)\psi_x + \mathcal{O}_1)\psi_{xx} + \mathcal{O}_2\psi_{xy} + (\frac{1}{c_5} + \mathcal{O}_3)\psi_{yy} - (1 + \mathcal{O}_4)\psi_x + \mathcal{O}_5\psi_y = 0,$$

with $\mathcal{O}_j = \mathcal{O}_j(\psi_x, \psi_y, \psi, x, c_5)$ for $j = 1, \dots, 5$, defined as follows:

$$\mathcal{O}_1(p_x, p_y, z, x, c) = -\frac{x^2}{c} + \frac{\gamma + 1}{2c}(2x - p_x)p_x - \frac{\gamma - 1}{c}\left(z + \frac{p_y^2}{2(c - x)^2}\right),$$

$$\mathcal{O}_2(p_x, p_y, z, x, c) = -\frac{2(p_x + c - x)p_y}{c(c - x)^2},$$

$$(3.26) \quad \mathcal{O}_{3}(p_{x}, p_{y}, z, x, c) = \frac{1}{c(c-x)^{2}} \Big((2c-x)x - (\gamma-1)\big(z + (c-x)p_{x} - \frac{p_{x}^{2}}{2}\big) + \frac{(\gamma-1)p_{y}^{2}}{2(c-x)^{2}} \Big), \\ \mathcal{O}_{4}(p_{x}, p_{y}, z, x, c) = \frac{1}{c-x} \Big(x - \frac{\gamma-1}{c}\big(z + (c-x)p_{x} + \frac{p_{x}^{2}}{2} + \frac{(\gamma+1)p_{y}^{2}}{2(\gamma-1)(c-x)^{2}}\big)\Big), \\ \mathcal{O}_{5}(p_{x}, p_{y}, z, x, c) = -\frac{2(p_{x} + c - x)p_{y}}{c(c-x)^{3}}.$$

(ii) Boundary condition for ψ on $\mathcal{R}(\Gamma_{\text{shock}} \cap \partial \Omega^5)$: Denote the Rankine–Hugoniot condition for φ on Γ_{shock} as

(3.27)
$$g^{\rm sh}(D\varphi,\varphi,\boldsymbol{\xi}) \coloneqq \rho(|D\varphi|,\varphi)D\varphi \cdot \boldsymbol{\nu} - D\varphi_2 \cdot \boldsymbol{\nu} = (\mathcal{A}(D\varphi,\varphi) - D\varphi_2) \cdot \boldsymbol{\nu} = 0$$

where $\mathcal{A}(\boldsymbol{p}, z) = \rho(|\boldsymbol{p}|, z)\boldsymbol{p}$, and $\boldsymbol{\nu} = \frac{D(\varphi_2 - \varphi)}{|D(\varphi_2 - \varphi)|}$ is the unit interior normal vector on Γ_{shock} . For constant $\delta' > 0$ given in Lemma 3.9, we define a function $\zeta \in C^{\infty}(\mathbb{R})$ by

$$\zeta(t) = \begin{cases} t & \text{for } t \ge \frac{3\delta'}{4} ,\\ \frac{\delta'}{2} & \text{for } t < \frac{\delta'}{2} , \end{cases}$$

satisfying $\zeta'(t) \geq 0$ on \mathbb{R} . Also, we define an extension of $g^{\mathrm{sh}}(\boldsymbol{p}, z, \boldsymbol{\xi})$ onto $\mathbb{R}^2 \times \mathbb{R} \times \overline{\Omega^5}$:

(3.28)
$$g_{\text{mod}}^{\text{sh}}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq \left(\tilde{\mathcal{A}}(\boldsymbol{p}, z) - D\varphi_2(\boldsymbol{\xi})\right) \cdot \frac{D\varphi_2(\boldsymbol{\xi}) - \boldsymbol{p}}{\zeta(|D\varphi_2(\boldsymbol{\xi}) - \boldsymbol{p}|)}$$

where $\tilde{\mathcal{A}}(\boldsymbol{p}, z)$ is given by Lemma 3.5, and $g_{\text{mod}}^{\text{sh}}(\boldsymbol{p}, z, \boldsymbol{\xi}) = g^{\text{sh}}(\boldsymbol{p}, z, \boldsymbol{\xi})$ if $|D\varphi_2(\boldsymbol{\xi}) - \boldsymbol{p}| \geq \frac{3\delta_1}{4}$. Then we define

$$M^{(\theta_1)}(\boldsymbol{p}, z, \xi) \coloneqq g_{\text{mod}}^{\text{sh}}(\boldsymbol{p} + D\varphi_5, z + \varphi_5, \xi, \xi \tan \theta_{25} + a_{25} + \frac{z}{v_2})$$

with $(D\varphi_5, \varphi_5)$ evaluated at $\boldsymbol{\xi} = (\xi, \xi \tan \theta_{25} + a_{25} + \frac{z}{v_2})$. Note that, by Proposition 2.6(i) and Definition 2.7, (θ_{25}, a_{25}) depend continuously on $\theta_1 \in [0, \theta^d]$ with the limit: $\lim_{\theta_1 \to 0^+} (\theta_{25}, a_{25}) =$

 (π, η_0) , where $\eta_0 > 0$ is given by (2.23).

We prescribe the free boundary condition for ψ to be

 $\mathcal{B}^{(\theta_1)}(\psi_x,\psi_y,\psi,x,y) = 0 \qquad \text{on } \mathcal{R}(\Gamma_{\text{shock}} \cap \partial \Omega^5) \,,$

where $\mathcal{B}^{(\theta_1)}$ is given by

(3.29)
$$\mathcal{B}^{(\theta_1)}(p_x, p_y, z, x, y) \coloneqq |D\bar{\phi}_{25} - (p_1, p_2)| M^{(\theta_1)}(p_1, p_2, z, \xi) ,$$

with function $\bar{\phi}_{25}(\boldsymbol{\xi}) \coloneqq (\varphi_2 - \varphi_5)(\boldsymbol{\xi}) = v_2(\eta - \xi \tan \theta_{25} - a_{25})$, and

$$\xi = v_2 \tan \theta_{25} + (c_5 - x) \cos y, \qquad \begin{pmatrix} p_1 \\ p_2 \end{pmatrix} = \begin{pmatrix} \cos y & \sin y \\ \sin y & -\cos y \end{pmatrix} \begin{pmatrix} -p_x \\ -\frac{p_y}{c_5 - x} \end{pmatrix}.$$

(iii) Other conditions for ψ : From Problem 2.10(iii)–(iv) and Definition 2.11(iv), ψ satisfies

$$\psi \ge 0$$
 in $\mathcal{R}(\Omega^5)$, $\psi = 0$ on $\mathcal{R}(\Gamma_{\text{sonic}}^5)$, $\psi_y = 0$ on $\mathcal{R}(\Gamma_{\text{sym}} \cap \partial \Omega^5)$.

Definition 3.11 (Parabolic norm). Fix $\alpha \in (0,1)$. Define the α -th parabolic distance between points $\mathbf{x} = (x, y), \ \tilde{\mathbf{x}} = (\tilde{x}, \tilde{y}) \in (0, \infty) \times \mathbb{R}$ to be

$$\delta_{\alpha}^{(\text{par})}(\boldsymbol{x}, \tilde{\boldsymbol{x}}) \coloneqq \left(|x - \tilde{x}|^2 + \max\{x, \tilde{x}\} |y - \tilde{y}|^2 \right)^{\frac{1}{2}}$$

Then, given constants $\sigma > 0$ and $m \in \mathbb{Z}_+$, the parabolic norms are defined as follows:

(i) For any open set $\mathcal{D} \subseteq (0,\infty) \times \mathbb{R}$ and any function $u \in C^2(\mathcal{D})$ in the (x,y)-coordinates, define

$$\begin{split} \|u\|_{m,0,\mathcal{D}}^{(\sigma),(\mathrm{par})} &\coloneqq \sum_{0 \le k+l \le m} \sup_{\boldsymbol{x} \in \mathcal{D}} \left(x^{k+\frac{l}{2}-\sigma} |\partial_x^k \partial_y^l u(\boldsymbol{x})| \right), \\ [u]_{m,\alpha,\mathcal{D}}^{(\sigma),(\mathrm{par})} &\coloneqq \sum_{k+l=m} \sup_{\substack{\boldsymbol{x}, \tilde{\boldsymbol{x}} \in \mathcal{D} \\ \boldsymbol{x} \neq \tilde{\boldsymbol{x}}}} \left(\min\left\{ x^{\alpha+k+\frac{l}{2}-\sigma}, \tilde{x}^{\alpha+k+\frac{l}{2}-\sigma} \right\} \frac{|\partial_x^k \partial_y^l u(\boldsymbol{x}) - \partial_x^k \partial_y^l u(\tilde{\boldsymbol{x}})|}{\delta_{\alpha}^{(\mathrm{par})}(\boldsymbol{x}, \tilde{\boldsymbol{x}})} \right), \\ \|u\|_{m,\alpha,\mathcal{D}}^{(\sigma),(\mathrm{par})} &\coloneqq \|u\|_{m,0,\mathcal{D}}^{(\sigma),(\mathrm{par})} + [u]_{m,\alpha,\mathcal{D}}^{(\sigma),(\mathrm{par})}. \end{split}$$

(ii) For any a > 0, fix an open interval I := (0, a). For a function $f \in C^2(I)$, define

$$\begin{split} \|f\|_{m,0,I}^{(\sigma),(\text{par})} &\coloneqq \sum_{k=0}^{m} \sup_{x \in I} \left(x^{k-\sigma} |\partial_{x}^{k} f(x)| \right), \\ [f]_{m,\alpha,I}^{(\sigma),(\text{par})} &\coloneqq \sup_{\substack{x, \tilde{x} \in I \\ x \neq \tilde{x}}} \left(\min\left\{ x^{\alpha+m-\sigma}, \tilde{x}^{\alpha+m-\sigma} \right\} \frac{|\partial_{x}^{m} f(x) - \partial_{x}^{m} f(\tilde{x})|}{|x - \tilde{x}|^{\alpha}} \right), \\ \|f\|_{m,\alpha,I}^{(\sigma),(\text{par})} &\coloneqq \|f\|_{m,0,I}^{(\sigma),(\text{par})} + [f]_{m,\alpha,I}^{(\sigma),(\text{par})}. \end{split}$$

(iii) In the case $\sigma = 2$, write $\|u\|_{2,\alpha,\mathcal{D}}^{(\text{par})} \coloneqq \|u\|_{2,\alpha,\mathcal{D}}^{(2),(\text{par})}$ and $\|f\|_{2,\alpha,I}^{(\text{par})} \coloneqq \|f\|_{2,\alpha,I}^{(2),(\text{par})}.$

Case 1. Admissible solutions for $\theta_1 < \theta^s$ away from θ^s .

Proposition 3.12. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For some constant $\bar{\varepsilon} > 0$ depending only on (γ, v_2) , any $\sigma \in (0, \theta^s)$, and any $\alpha \in (0, 1)$, there exist both $\varepsilon \in (0, \bar{\varepsilon}]$ depending only on (γ, v_2, σ) and C > 0 depending only on (γ, v_2, α) such that any admissible solution $\varphi = \psi \circ \mathcal{R} + \varphi_5$ corresponding to $\theta \in \Theta \cap \{0 \leq \theta_1 \leq \theta^s - \sigma\}$ satisfies

(3.30)
$$\|\psi\|_{2,\alpha,\mathcal{R}(\Omega_{\varepsilon}^{5})}^{(\text{par})} + \|f_{5,\text{sh}} - f_{5,0}\|_{2,\alpha,(0,\varepsilon)}^{(\text{par})} \le C,$$

where $\Omega_{\varepsilon}^5 \subseteq \Omega$ is a neighbourhood of Γ_{sonic}^5 , and functions $f_{5,\text{sh}}$ and $f_{5,0}$ represent Γ_{shock} and S_{25} respectively to be defined in Step 1 below.

The proof of this proposition is similar to that of [2, Propositions 3.26 and 3.30]. We sketch the main steps of the proof, while omitting most details.

1. The depiction and the properties of the reflected shock S_{25} and the free boundary Γ_{shock} in a small neighbourhood of the sonic boundary Γ_{sonic}^5 . It follows from the geometric properties of $\overline{\Omega^5}$ that there exist positive constants $(\varepsilon_1, \varepsilon_0, \omega_0)$ depending only on (γ, v_2) , with $\varepsilon_0 < \varepsilon_1$ and $\omega_0 \in (0, 1)$, and a unique function $f_{5,0} \in C^{\infty}([-\varepsilon_0, \varepsilon_0])$ satisfying

$$\mathcal{R}\left(S_{25} \cap \mathcal{N}_{\varepsilon_1}(\Gamma_{\text{sonic}}^5)\right) \cap \left\{|x| < \varepsilon_0\right\} = \left\{(x, y) : |x| < \varepsilon_0, \ y = f_{5,0}(x)\right\},\$$

where $\mathcal{N}_{\varepsilon_1}(\Gamma_{\text{sonic}}^5)$ is defined by (3.21), and $2\omega_0 \leq f'_{5,0}(x) \leq \omega_0^{-1}$ for $x \in (-\varepsilon_0, \varepsilon_0)$. Then, for any $\varepsilon \in (0, \varepsilon_1]$, we define a set $\Omega^5_{\varepsilon} \subseteq \Omega$ by

(3.31)
$$\Omega^{5}_{\varepsilon} \coloneqq \Omega \cap \mathcal{N}_{\varepsilon_{1}}(\Gamma^{5}_{\text{sonic}}) \cap \mathcal{R}^{-1}(\{x < \varepsilon\}).$$

Using (3.20) and (3.25), we can show that there exist $\bar{\varepsilon} \in (0, \varepsilon_0]$, $L \ge 1$, $\delta \in (0, \frac{1}{2})$, and $\omega \in (0, \omega_0)$ depending only on (γ, v_2) such that, for all $(x, y) \in \mathcal{R}(\Omega_{\bar{\varepsilon}}^5)$,

(3.32)
$$0 \le \psi(x, y) \le Lx^2, \quad 0 \le \psi_x \le \frac{2-\delta}{1+\gamma} x \le Lx, \quad |\psi_y(x, y)| \le Lx,$$

and there exists a unique function $f_{5,sh} \in C^1([0, \bar{\varepsilon}])$ such that

$$\mathcal{R}(\Omega^{5}_{\bar{\varepsilon}}) = \left\{ (x, y) : 0 < x < \bar{\varepsilon}, \ 0 < y < f_{5, \text{sh}}(x) \right\},$$
$$\mathcal{R}(\Gamma_{\text{shock}} \cap \partial \Omega^{5}_{\bar{\varepsilon}}) = \left\{ (x, y) : 0 < x < \bar{\varepsilon}, \ y = f_{5, \text{sh}}(x) \right\},$$

and $\omega \leq f'_{5,\mathrm{sh}}(x) \leq L$ for any $x \in (0, \bar{\varepsilon})$.

2. Ellipticity for the equation and the free boundary condition. Equation (3.25) for ψ in $\mathcal{R}(\Omega^5)$ can be written as

(3.33)
$$\sum_{i,j=1}^{2} \hat{A}_{ij}^{(\theta_1)}(\psi_x, \psi_y, \psi, x) \partial_i \partial_j \psi + \sum_{i,j=1}^{2} \hat{A}_i^{(\theta_1)} \partial_i \psi = 0,$$

with $(\partial_1, \partial_2) = (\partial_x, \partial_y)$ and $\hat{A}_{12}^{(\theta_1)} = \hat{A}_{21}^{(\theta_1)}$. Then there exist $\varepsilon_5 \in (0, \frac{\overline{\varepsilon}}{4}]$ and $\lambda_5 > 0$ depending only on (γ, v_2) such that, for all $(x, y) \in \mathcal{R}(\overline{\Omega_{4\varepsilon_5}^5})$,

$$\frac{\lambda_5}{2}|\boldsymbol{\kappa}|^2 \leq \sum_{i,j=1}^2 \hat{A}_{ij}^{(\theta_1)}(\psi_x(x,y),\psi_y(x,y),\psi(x,y),x)\frac{\kappa_i\kappa_j}{x^{2-\frac{i+j}{2}}} \leq \frac{2}{\lambda_5}|\boldsymbol{\kappa}|^2 \quad \text{for all } \boldsymbol{\kappa} \in \mathbb{R}^2.$$

Moreover, $\mathcal{B}^{(\theta_1)}$ defined by (3.29) satisfies $\mathcal{B}^{(\theta_1)}(\mathbf{0}, 0, x, y) = 0$ for all $(x, y) \in \mathbb{R}^2$. Let y_{P_2} be the y-coordinate of point P_2 defined in Definition 2.9. For each $m = 2, 3, \cdots$, there exist constants $\delta_{\mathrm{bc}} > 0$ and C > 1 depending only on (γ, v_2, m) such that, whenever $|(p_x, p_y, z, x)| \leq \delta_{\mathrm{bc}}$ and $|y - y_{P_2}| \leq \delta_{\mathrm{bc}}$,

$$\left| D^m \mathcal{B}^{(\theta_1)}(p_x, p_y, z, x, y) \right| \le C$$

Furthermore, there exist constants $\hat{\delta}_{bc} > 0$, C > 1, and $\varepsilon' > 0$ depending only on (γ, v_2) such that, whenever $|(p_x, p_y, z, x)| \leq \hat{\delta}_{bc}$ and $|y - y_{P_2}| \leq \hat{\delta}_{bc}$,

$$\begin{aligned} \partial_a \mathcal{B}^{(\theta_1)}(p_x, p_y, z, x, y) &\leq -C^{-1} & \text{for } \partial_a = \partial_{p_x}, \partial_{p_y}, \text{ or } \partial_z \\ D_{(p_x, p_y)} \mathcal{B}^{(\theta_1)}(p_x, p_y, z, x, y) \cdot \boldsymbol{\nu}_{\mathrm{sh}}^{(x, y)} &\geq C^{-1} & \text{on } \mathcal{R}(\Gamma_{\mathrm{shock}} \cap \partial \Omega_{\varepsilon'}^5) \,, \end{aligned}$$

where the vector field $\boldsymbol{\nu}_{\rm sh}^{(x,y)}$ is the unit normal vector to $\mathcal{R}(\Gamma_{\rm shock})$ expressed in the (x,y)coordinates and oriented towards the interior of $\mathcal{R}(\Omega)$.

3. Extension of the domain for the coefficients to equation (3.33). Let $\varepsilon_0 > 0$ and $L \ge 1$ be the constants defined as above. Then there exist constants $\varepsilon \in (0, \frac{\varepsilon_0}{2}]$ and C > 0 depending only on (γ, v_2) such that, for any admissible solution φ corresponding to parameters $\theta \in \Theta$, function $\psi = (\varphi - \varphi_5) \circ \mathcal{R}^{-1}$ satisfies

(3.34)
$$\sum_{i,j=1}^{2} \hat{A}_{ij}^{(\text{mod})}(\psi_x,\psi_y,\psi,x)\partial_i\partial_j\psi + \sum_{i,j=1}^{2} \hat{A}_i^{(\text{mod})}(\psi_x,\psi_y,\psi,x)\partial_i\psi = 0 \quad \text{in } \mathcal{R}(\Omega_{\varepsilon}^5),$$

with $(\partial_1, \partial_2) \coloneqq (\partial_x, \partial_y)$ and coefficients $(\hat{A}_{ij}^{(\text{mod})}, \hat{A}_i^{(\text{mod})})$ satisfying

$$(\hat{A}_{ij}^{(\text{mod})}, \hat{A}_i^{(\text{mod})}) = (\hat{A}_{ij}^{(\theta_1)}, \hat{A}_i^{(\theta_1)}) \quad \text{in } \{(p_x, p_y, z, x) : |(p_x, p_y)| \le Lx, \, |z| \le Lx^2, \, 0 < x < \varepsilon\}.$$

Moreover, the modified coefficients satisfy

$$\begin{split} \big| (\hat{A}_{11}^{(\mathrm{mod})}, \hat{A}_{12}^{(\mathrm{mod})}, \hat{A}_{2}^{(\mathrm{mod})})(p_x, p_y, z, x) \big| &\leq Cx \qquad \text{in } \mathbb{R}^2 \times \mathbb{R} \times (0, \varepsilon) \,, \\ \big\| (\hat{A}_{22}^{(\mathrm{mod})}, \hat{A}_{1}^{(\mathrm{mod})}) \big\|_{0, \mathbb{R}^2 \times \mathbb{R} \times (0, \varepsilon)} + \big\| D_{(p_x, p_y, z, x)}(\hat{A}_{ij}^{(\mathrm{mod})}, \hat{A}_i^{(\mathrm{mod})}) \big\|_{0, \mathbb{R}^2 \times \mathbb{R} \times (0, \varepsilon)} \leq C \,. \end{split}$$

4. Rescaling method in the local domain to obtain the weighted Hölder estimate. Using the strictly decreasing property of η^{P_2} with respect to $\theta_1 \in [0, \theta^s]$ which is shown in Lemma A.2(ii), we obtain a constant $l_{so} > 0$ depending only on (γ, v_2, σ) such that, for any admissible solution φ corresponding to $\theta \in \Theta \cap \{0 \leq \theta_1 \leq \theta^s - \sigma\}$,

$$y_P = f_{5,\mathrm{sh}}(x_P) \ge l_{\mathrm{so}}$$
 for any $x_P \in [0, \bar{\varepsilon}]$.

Subsequently, we define $\varepsilon_* \coloneqq \min\{\frac{\overline{\varepsilon}}{2}, l_{so}^2\}.$

For fixed $\varepsilon \in (0, \varepsilon_*]$ and any $z_0 = (x_0, y_0) \in \mathcal{R}(\overline{\Omega_{\varepsilon}^5} \setminus \overline{\Gamma_{\text{sonic}}^5})$, define the rescaled coordinates (S, T) by $(x, y) \coloneqq z_0 + \frac{1}{4}(x_0 S, \sqrt{x_0}T)$, which take values in a local domain

$$Q_r^{(z_0)} \coloneqq \left\{ (S,T) \in (-1,1)^2 : z = z_0 + \frac{r}{4} (x_0 S, \sqrt{x_0} T) \in \mathcal{R}(\Omega_{2\varepsilon}^5) \right\} \quad \text{for any } r \in (0,1]$$

Considering property (3.32), we define a rescaled function $\psi^{(z_0)}$ by

$$\psi^{(z_0)}(S,T) \coloneqq \frac{1}{x_0^2} \psi(x_0 + \frac{x_0}{4}S, y_0 + \frac{\sqrt{x_0}}{4}T) \qquad \text{for } (S,T) \in Q_1^{(z_0)}$$

With a suitable rescaling on the coefficients of the modified equation (3.34), we obtain a uniformly elliptic equation for $\psi^{(z_0)}$ under the (S, T)-coordinates.

Note that, for $z_0 \in \mathcal{R}(\overline{\Omega_{\varepsilon}^5} \cap \Gamma_{\text{shock}})$, we can express Γ_{shock} locally under the rescaled coordinates as $\Gamma_{\text{shock}}^{(z_0)} \coloneqq \{(S,T) \in (-1,1)^2 : T = F^{(z_0)}(S)\} \subseteq \partial Q_1^{(z_0)}$ with $F^{(z_0)}(S)$ defined by

$$F^{(z_0)}(S) \coloneqq \frac{4}{\sqrt{x_0}} \left(f_{5,\text{sh}}(x_0 + \frac{x_0}{4}S) - f_{5,\text{sh}}(x_0) \right) \quad \text{for } S \in (-1,1)$$

Then, in the local domain $Q_1^{(z_0)}$, we apply [10, Theorem 4.2.3] for $z_0 \in \mathcal{R}(\Omega_{\varepsilon}^5)$, [10, Theorem 4.2.8] for $z_0 \in \mathcal{R}(\Gamma_{\text{shock}} \cap \partial \Omega_{\varepsilon}^5)$, and [10, Theorem 4.2.10] for $z_0 \in \mathcal{R}(\Gamma_{\text{sym}} \cap \partial \Omega_{\varepsilon}^5)$, respectively, to obtain the uniform bound:

$$\sup_{0 \in \mathcal{R}(\Omega_{\varepsilon}^{5})} \left\| \psi^{(z_{0})} \right\|_{2,\alpha,\overline{Q_{1/4}^{(z_{0})}}} + \sup_{z_{0} \in \mathcal{R}(\Gamma_{\mathrm{shock}} \cap \partial \Omega_{\varepsilon}^{5})} \frac{1}{\sqrt{x_{0}}} \left\| F^{(z_{0})} \right\|_{2,\alpha,[-\frac{1}{4},\frac{1}{4}]} \leq C.$$

Scaling back into the (x, y)-coordinates, we obtain the weighted Hölder estimates (3.30).

Case 2. Admissible solutions for $\theta_1 < \theta^s$ close to θ^s .

Proposition 3.13. Fix $\gamma \geq 1$ and $v_2 \in [v_2^s, 0)$. Let $\bar{\varepsilon} > 0$ be the constant from Proposition 3.12. For any $\alpha \in (0, 1)$, there exist constants $\varepsilon \in (0, \bar{\varepsilon}]$ and $\sigma_1 \in (0, \theta^s)$ depending only on (γ, v_2) , and C > 0 depending only on (γ, v_2, α) such that any admissible solution $\varphi = \psi \circ \mathcal{R} + \varphi_5$ corresponding to $\theta \in \Theta \cap \{\theta^s - \sigma_1 \leq \theta_1 < \theta^s\}$ satisfies

$$\|\psi\|_{2,\alpha,\mathcal{R}(\Omega_{\varepsilon}^{5})}^{(\text{par})} + \|f_{5,\text{sh}} - f_{5,0}\|_{2,\alpha,(0,\varepsilon)}^{(\text{par})} \le C$$

where $\Omega^5_{\varepsilon} \subseteq \Omega$, and functions $f_{5,\text{sh}}$ and $f_{5,0}$ are defined in Step 1 of Case 1 above.

The proof is similar to that of [2, Proposition 3.32]. The admissible solutions for Case 2 share the same properties as in Steps 1-3 of Case 1. We omit the details of the proof and only draw the main differences compared with the proof of Proposition 3.12 here.

1. Weighted Hölder estimate near $\mathcal{R}(P_0^1)$. Because $v_2 \in [v_2^s, 0), y_{P_2} > 0$ tends to zero as $\theta_1 \to \theta^{s-}$. Therefore, for the same rescaled coordinates (S,T) and local domain $Q_r^{(z_0)}$ with $z_0 = (x_0, y_0) \in \mathcal{R}(\Omega_{\varepsilon}^5)$ as given in Step 4 of Case 1 above, it is possible that " $Q_r^{(z_0)}$ does not fit into Ω_{ε}^{5} ", meaning that the image of $Q_r^{(z_0)}$ under transform $(S,T) \mapsto (x,y)$ intersects the boundaries: $\mathcal{R}(\Gamma_{\text{shock}} \cap \partial \Omega_{\varepsilon}^5)$ and $\mathcal{R}(\Gamma_{\text{sym}} \cap \partial \Omega_{\varepsilon}^5)$ simultaneously.

For that reason, we define $\Omega_{fc}^5 := \Omega_{\varepsilon}^5 \cap \mathcal{R}^{-1}(\{x < y_{P_2}^2\})$. It follows that $Q_r^{(z_0)}$ fits into Ω_{fc}^5 for any $z_0 \in \mathcal{R}(\Omega_{fc}^5)$ due to similar considerations as the choice of ε_* above. In this case, we follow the same idea as for Proposition 3.12 to obtain that there exists a constant C > 0 depending only on (γ, v_2, α) such that, for $\boldsymbol{\theta} \in \Theta \cap \{\theta^s - \sigma^* \leq \theta_1 < \theta^s\}$, $\|\psi\|_{2,\alpha,\mathcal{R}(\Omega_{fc}^5)}^{(\text{par})} \leq C$ for some small constant $\sigma^* > 0$ depending only on (γ, v_2) .

2. Weighted Hölder estimate away from $\mathcal{R}(P_0^1)$. Note that there exists sufficiently large k > 1 depending only on (γ, v_2) such that the following geometric relations hold:

$$\{0 < x < 2\bar{\varepsilon}, \ 0 < y < y_{P_2} + \frac{x}{k}\} \subseteq \mathcal{R}(\Omega_{2\bar{\varepsilon}}^5) \subseteq \{0 < x < 2\bar{\varepsilon}, \ 0 < y < y_{P_2} + kx\}.$$

In the rest of the domain, there exist constants $\hat{\varepsilon} \in (0, \frac{\tilde{\varepsilon}}{2}], \sigma' \in (0, \sigma^*]$, and $C^* > 0$ depending only on (γ, v_2) such that, for $\boldsymbol{\theta} \in \Theta \cap \{\theta^s - \sigma' \leq \theta_1 < \theta^s\}$,

(3.35)
$$0 \le \psi(x, y) \le C^* x^4 \quad \text{in } \mathcal{R}(\Omega_{2\hat{\varepsilon}}^5) \cap \{x > \frac{1}{10} y_{P_2}^2\}.$$

Then, for any $z_0 = (x_0, y_0) \in \mathcal{R}(\overline{\Omega_{\hat{\varepsilon}}^5}) \cap \{x > \frac{1}{5}y_{P_2}^2\}$, we introduce the rescaled coordinates (S, T) as $(x, y) =: z_0 + \frac{\sqrt{x_0}}{10k}(x_0S, \sqrt{x_0}T)$, which take values in a local domain $Q_r^{(z_0)}$ defined by

$$Q_r^{(z_0)} \coloneqq \left\{ (S,T) \in (-1,1)^2 : z = z_0 + \frac{r\sqrt{x_0}}{10k} (x_0 S, \sqrt{x_0}T) \in \mathcal{R}(\Omega_{2\hat{\varepsilon}}^5) \right\} \quad \text{for any } r \in (0,1].$$

Thus, the local domain $Q_r^{(z_0)}$ fits into $\Omega_{2\hat{\varepsilon}}^5$, meaning that the image of $Q_r^{(z_0)}$ under transform $(S,T) \mapsto (x,y)$ does not intersect both $\mathcal{R}(\Gamma_{\text{shock}})$ and $\mathcal{R}(\Gamma_{\text{sym}})$ simultaneously. Considering property (3.35), we define a rescaled function $\psi^{(z_0)}$ by

$$\psi^{(z_0)}(S,T) \coloneqq \frac{1}{x_0^4} \psi(x_0 + \frac{x_0^{3/2}}{10k}S, y_0 + \frac{x_0}{10k}T) \quad \text{for } (S,T) \in Q_1^{(z_0)}.$$

The rescaled function $\psi^{(z_0)}$ satisfies a uniformly elliptic equation in the (S, T)-coordinates after a suitable rescaling on the coefficients of the modified equation (3.34). The rest of the proof follows the same idea as in Step 4 of the proof of Proposition 3.12.

Case 3. Admissible solutions for $\theta_1 \geq \theta^s$ close to θ^s .

Proposition 3.14. Fix $\gamma \geq 1$ and $v_2 \in (v_2^s, 0)$. Let $\bar{\varepsilon} > 0$ be the constant from Proposition 3.12. For each $\alpha \in (0, 1)$, there exist constants $\varepsilon \in (0, \bar{\varepsilon}]$ and $\sigma_3 \in (0, \theta^d - \theta^s)$ depending only on (γ, v_2) , and a constant C > 0 depending only on (γ, v_2, α) such that any admissible solution φ corresponding to $\theta \in \Theta \cap \{\theta^s \leq \theta_1 < \theta^s + \sigma_3\}$ satisfies

$$\left\|\varphi - \varphi_5\right\|_{2,\alpha,\overline{\Omega_{\varepsilon}^5}} + \left\|f_{5,\mathrm{sh}} - f_{5,0}\right\|_{2,\alpha,(0,\varepsilon)} \le C,$$

$$|D_{\boldsymbol{\xi}}^{m}(\varphi - \varphi_{5})(P_{0}^{1})| = \frac{\mathrm{d}^{m}}{\mathrm{d}x^{m}} \left(f_{5,\mathrm{sh}} - f_{5,0}\right)(0) = 0 \qquad \text{for } m = 0, 1, 2 \,,$$

where $\Omega^5_{\varepsilon} \subseteq \Omega$, and functions $f_{5,\text{sh}}$ and $f_{5,0}$ are defined in Step 1 below.

The proof is similar to that of [2, Proposition 3.39], so we omit most details here and only emphasize the difference compared with the previous two cases.

1. The depiction and the properties of the reflected shock S_{25} and the free boundary Γ_{shock} in a small neighbourhood of the sonic boundary Γ_{sonic}^5 . Let $\varepsilon_1, \varepsilon_0, \omega_0 > 0$ be chosen as in Step 1 of the proof of Proposition 3.12. For the coordinates: $(x, y) = \mathcal{R}(\boldsymbol{\xi})$ given by (3.24), we define $\hat{x} \coloneqq x - x_{P_2}$. Note that $x_{P_2} = 0$ if $\theta_1 \in [0, \theta^s]$ and $x_{P_2} > 0$ if $\theta_1 \in (\theta^s, \theta^s + \sigma_2]$. Then there exists a unique function $f_{5,0} \in C^{\infty}([-\varepsilon_0, \varepsilon_0])$ satisfying

$$\mathcal{R}\left(S_{25} \cap \mathcal{N}_{\varepsilon_1}(\Gamma^5_{\text{sonic}})\right) \cap \{(x,y) : |\hat{x}| < \varepsilon_0\} = \left\{(x,y) : |\hat{x}| < \varepsilon_0, \, y = f_{5,0}(\hat{x})\right\},$$

and $2\omega_0 \leq f_{5,0}'(\hat{x}) \leq \omega_0^{-1}$ for $\hat{x} \in (-\varepsilon_0, \varepsilon_0)$.

Reducing $\bar{\varepsilon} \in (0, \varepsilon_0]$ further, we can choose $\sigma_2 \in (0, \theta^d - \theta^s)$ depending only on $(\gamma, v_2, \bar{\varepsilon})$ such that, for any admissible solution φ corresponding to $\theta \in \Theta \cap \{0 \leq \theta_1 \leq \theta^s + \sigma_2\}, \Omega_{\bar{\varepsilon}}^5$ defined by (3.31) is nonempty due to the monotonicity of the pseudo-Mach number in Lemma 2.3, from which we can generalize the definition of $\Omega_{\bar{\varepsilon}}^5$ to

(3.37)
$$\Omega^{5}_{\bar{\varepsilon}} \coloneqq \Omega \cap \mathcal{N}_{\varepsilon_{1}}(\Gamma^{5}_{\text{sonic}}) \cap \mathcal{R}^{-1}(\{(x,y) : 0 < \hat{x} < \bar{\varepsilon}\}).$$

There exist $L \geq 1$, $\delta \in (0, \frac{1}{2})$, and $\omega \in (0, \omega_0)$ depending only on (γ, v_2) such that, for all $(x, y) \in \mathcal{R}(\Omega^5_{\overline{\varepsilon}})$,

(3.38)
$$0 \le \psi(x, y) \le Lx^2, \quad 0 \le \psi_x \le \frac{2-\delta}{1+\gamma} x \le Lx, \quad |\psi_y(x, y)| \le Lx,$$

for $\psi := (\varphi - \varphi_5) \circ \mathcal{R}^{-1}$, and there exists a unique function $f_{5,\text{sh}} \in C^1([0,\bar{\varepsilon}])$ such that

$$\begin{aligned} \mathcal{R}(\Omega^{5}_{\bar{\varepsilon}}) &= \left\{ (x,y) \, : \, 0 < \hat{x} < \bar{\varepsilon}, \, 0 < y < f_{5,\mathrm{sh}}(\hat{x}) \right\}, \\ \mathcal{R}(\Gamma_{\mathrm{shock}} \cap \partial \Omega^{5}_{\bar{\varepsilon}}) &= \left\{ (x,y) \, : \, 0 < \hat{x} < \bar{\varepsilon}, \, y = f_{5,\mathrm{sh}}(\hat{x}) \right\} \end{aligned}$$

,

and $\omega \leq f'_{5\,\mathrm{sh}}(\hat{x}) \leq L$ for any $\hat{x} \in (0, \bar{\varepsilon})$.

2. The free boundary condition and the modification on equation (3.33). Similar to [2, Lemma 3.37 and Corollary 3.38], it follows that, for $\sigma_2 \in (0, \theta^d - \theta^s)$ defined in Step 1, there exists

a constant $\mu_0 > 0$ depending only on (γ, v_2, σ_2) such that, for any $\boldsymbol{\theta} \in \Theta \cap \{\boldsymbol{\theta}^s \leq \boldsymbol{\theta}_1 \leq \boldsymbol{\theta}^d - \sigma_2\}$, $g_{\text{mod}}^{\text{sh}}$ defined by (3.28) satisfies the following properties:

$$\partial_a g_{\mathrm{mod}}^{\mathrm{sh}}(D\varphi_5(P_2),\varphi_5(P_2),P_2) \leq -\mu_0,$$

where $\partial_a = \partial_{p_1}, \partial_{p_2}$, or ∂_z . From these properties above, we can show that $\mathcal{B}^{(\theta_1)}$ defined by (3.29) satisfies the same properties as in Step **2** of Case 1. The extension of the domain for the coefficients of equation (3.33) is the same as Step **3** of Case 1.

3. Rescaling method in the local domain to obtain the weighted Hölder estimate. Note that, from the properties of $f_{5,0}$ and $f_{5,\text{sh}}$ in Step 1, there exists k > 1 depending only on (γ, v_2) such that

$$\left\{ 0 < \hat{x} < \bar{\varepsilon}, \, 0 < y < \frac{\hat{x}}{k} \right\}_{-} \subseteq \mathcal{R}(\Omega^5_{\bar{\varepsilon}}) \subseteq \left\{ 0 < \hat{x} < \bar{\varepsilon}, \, 0 < y < k\hat{x} \right\}.$$

Moreover, there exist constants $\varepsilon \in (0, \frac{\varepsilon}{2}], \sigma_3 \in (0, \sigma_2]$, and C > 1 depending only on (γ, v_2) such that, for $\boldsymbol{\theta} \in \Theta \cap \{\theta^{s} \leq \theta_1 \leq \theta^{s} + \sigma_3\},\$

(3.39)
$$x_{P_2} \leq \frac{\varepsilon}{10}, \qquad 0 \leq \psi(x,y) \leq C(x-x_{P_2})^5 \quad \text{in } \mathcal{R}(\Omega_{2\varepsilon}^5).$$

Then, for any $z_0 = (x_0, y_0) \in \mathcal{R}(\overline{\Omega_{\varepsilon}^5} \setminus \{P_2\})$, we introduce the rescaled coordinates (S, T) as $(x, y) =: z_0 + \frac{x_0 - x_{P_2}}{10k}(\sqrt{x_0}S, T)$, taking values in a local domain $Q_r^{(z_0)}$ defined by

$$Q_r^{(z_0)} \coloneqq \left\{ (S,T) \in (-1,1)^2 : z = z_0 + \frac{r(x_0 - x_{P_2})}{10k} (\sqrt{x_0}S,T) \in \mathcal{R}(\Omega_{2\varepsilon}^5) \right\} \quad \text{for any } r \in (0,1].$$

Then the local domain $Q_r^{(z_0)}$ fits into $\Omega_{2\varepsilon}^5$, meaning that the image of $Q_r^{(z_0)}$ under transform $(S,T) \mapsto (x,y)$ does not intersect both $\mathcal{R}(\Gamma_{\text{shock}})$ and $\mathcal{R}(\Gamma_{\text{sym}})$ simultaneously. Considering property (3.39), we define a rescaled function $\psi^{(z_0)}$ by

$$\psi^{(z_0)}(S,T) \coloneqq \frac{1}{(x_0 - x_{P_2})^5} \psi(x_0 + \frac{x_0 - x_{P_2}}{10k} \sqrt{x_0} S, y_0 + \frac{x_0 - x_{P_2}}{10k} T) \quad \text{for } (S,T) \in Q_1^{(z_0)}.$$

The rescaled function $\psi^{(z_0)}$ satisfies a uniformly elliptic equation in the (S, T)-coordinates after a suitable rescaling of the coefficients of the modified equation (3.34). The rest of the proof follows the same idea as Step 4 of Case 1.

Case 4. Admissible solutions for $\theta_1 \geq \theta^s$ away from θ^s .

Let $\sigma_3 > 0$ be from Proposition 3.14. By Proposition 3.8, there exists a constant $\delta \in (0, 1)$ depending only on (γ, v_2) such that any admissible solution φ corresponding to $\theta \in \Theta \cap \{\theta^s + \frac{\sigma_3}{2} \leq \theta_1 < \theta^d\}$ satisfies

(3.40)
$$\frac{|D\varphi|^2}{c^2(|D\varphi|,\varphi)} \le 1 - \delta \qquad \text{in } \overline{\Omega} \cap \{\xi \ge 0\},\$$

where $c(|D\varphi|,\varphi)$ is defined by (3.2). By Lemma 3.3 and (3.40), $(D\varphi(\boldsymbol{\xi}),\varphi(\boldsymbol{\xi})) \in \mathcal{K}_{R_*}$ for some constant $R_* \geq 2$ depending only on (γ, v_2) , where \mathcal{K}_R is defined by (3.9). In particular, there exist $\lambda_* > 0$ and $r_* > 0$, depending only on (γ, v_2) , such that any admissible solution φ corresponding to $\boldsymbol{\theta} \in \Theta \cap \{\theta^s + \frac{\sigma_3}{2} \leq \theta_1 < \theta^d\}$ satisfies

(3.41)
$$\lambda_* |\boldsymbol{\kappa}|^2 \le \sum_{i,j=1}^2 \partial_{p_j} \mathcal{A}_i(D\varphi(\boldsymbol{\xi}), \varphi(\boldsymbol{\xi})) \kappa_i \kappa_j \le \lambda_*^{-1} |\boldsymbol{\kappa}|^2$$

for any $\boldsymbol{\xi} \in \overline{\Omega} \cap B_{2r_*}(P_0^1)$ and any $\boldsymbol{\kappa} = (\kappa_1, \kappa_2) \in \mathbb{R}^2$.

Set $\bar{\phi}(\boldsymbol{\xi}) \coloneqq \varphi_2(\boldsymbol{\xi}) - \varphi(\boldsymbol{\xi})$. Recalling equation (3.1) for function $\phi = \varphi - \varphi_5$, the free boundary condition (3.27) on Γ_{shock} , and the slip boundary condition defined in Problem 2.10(iv) on Γ_{sym} , it follows that $\bar{\phi}$ satisfies the system:

(3.42)
$$\begin{cases} \sum_{i,j=1}^{2} \bar{A}_{ij} (D\bar{\phi}, \bar{\phi}, \boldsymbol{\xi}) \partial_{i} \partial_{j} \bar{\phi} = 0 & \text{ in } \Omega \cap B_{r_{*}} (P_{0}^{1}) \\ \bar{g}^{\text{sym}} (D\bar{\phi}, \bar{\phi}, \boldsymbol{\xi}) = 0 & \text{ on } \Gamma_{\text{sym}} , \\ \bar{g}^{\text{sh}} (D\bar{\phi}, \bar{\phi}, \boldsymbol{\xi}) = 0 & \text{ on } \Gamma_{\text{shock}} , \end{cases}$$

with coefficients \bar{A}_{ij} and the boundary functions \bar{g}^{sym} and \bar{g}^{sh} defined by

$$\bar{A}_{ij}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq c^2(|D\varphi_2 - \boldsymbol{p}|, \varphi_2 - z)\delta_{ij} - (\partial_i\varphi_2 - p_i)(\partial_j\varphi_2 - p_j)$$

$$\bar{g}^{\text{sym}}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq v_2 - p_2 - \eta,$$

(3.43)

$$\bar{g}^{\mathrm{sh}}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq -g^{\mathrm{sh}}(D\varphi_2 - \boldsymbol{p}, \varphi_2 - z, \boldsymbol{\xi}),$$

for $g^{\rm sh}$ given by (3.27), where $(\partial_1, \partial_2) = (\partial_{\xi}, \partial_{\eta})$.

We define a vector \boldsymbol{e} as

$$e \coloneqq rac{D arphi_2(P_0^1)}{|D arphi_2(P_0^1)|} = rac{O_2 - P_0^1}{|O_2 - P_0^1|},$$

and choose e^{\perp} as the clockwise rotation of e by $\frac{\pi}{2}$. Note that e depends only on (γ, v_2, θ_1) and satisfies that, for any $\boldsymbol{\xi} \in \Gamma_{\text{shock}} \cap B_{r_*}(P_0^1)$,

(3.44)
$$\partial_{\boldsymbol{e}}(\varphi_2 - \varphi)(\boldsymbol{\xi}) = -\partial_r(\varphi_2 - \varphi) + \left(\boldsymbol{e} - \frac{D\varphi(\boldsymbol{\xi})}{|D\varphi(\boldsymbol{\xi})|}\right) \cdot D(\varphi_2 - \varphi) \ge \frac{d'_0}{2}$$

where $d'_0 > 0$ is the constant from (3.23), depending only on (γ, v_2) . Then we define the new coordinates (S, T) by

(3.45)
$$\boldsymbol{\xi} \coloneqq P_0^1 + S\boldsymbol{e} + T\boldsymbol{e}^{\perp}$$

Under these (S,T)-coordinates, we can express $S_{25}, \Gamma_{\text{shock}}, \Gamma_{\text{sym}}$, and the domain near P_0^1 as

$$S_{25} \cap B_{r_*}(P_0^1) = \{ \boldsymbol{\xi}(S,T) : S = a_{\text{ref}}T, T > 0, (S,T) \in B_{r_*}(\mathbf{0}) \},\$$

$$\Gamma_{\text{shock}} \cap B_{r_*}(P_0^1) = \{ \boldsymbol{\xi}(S,T) : S = f_{\boldsymbol{e}}(T), T > 0, (S,T) \in B_{r_*}(\mathbf{0}) \},\$$

(3.46)

$$\Gamma_{\text{sym}} \cap B_{r_*}(P_0^1) = \{ \boldsymbol{\xi}(S,T) : S = a_{\text{sym}}T, T > 0, (S,T) \in B_{r_*}(\mathbf{0}) \},$$

$$\Omega \cap B_{r_*}(P_0^1) = \{ \boldsymbol{\xi}(S,T) : a_{\text{ref}}T \le f_{\boldsymbol{e}}(T) < S < a_{\text{sym}}T, T > 0, (S,T) \in B_{r_*}(\mathbf{0}) \},$$

where the positive constants $a_{\text{ref}} = \tan(\theta_{25} - \hat{\theta}_{25} - \frac{\pi}{2})$ and $a_{\text{sym}} = \cot \hat{\theta}_{25}$ depend continuously on $\theta_1 \in (0, \theta^d)$, and there exists a constant C > 0 depending only on (γ, v_2) such that $C^{-1} \leq a_{\text{ref}} < a_{\text{sym}} \leq C$ for all $\theta_1 \in [\theta^s, \theta^d]$. Note that $\hat{\theta}_{25}$ is defined by (2.28), and we have used Proposition 2.6, including (2.40).

Definition 3.15 (Weighted Hölder norm). For any $\sigma \in \mathbb{R}$, $\alpha \in (0,1)$, and $m \in \mathbb{N}$, the weighted Hölder norms are defined as follows:

(i) For any open bounded connected set $U \subseteq \mathbb{R}^2$, let Γ be a closed portion of ∂U . Write

$$\delta_{\boldsymbol{\xi}_1} \coloneqq \operatorname{dist}(\boldsymbol{\xi}_1, \Gamma), \quad \delta_{\boldsymbol{\xi}_1, \boldsymbol{\xi}_2} \coloneqq \min\{\delta_{\boldsymbol{\xi}_1}, \delta_{\boldsymbol{\xi}_2}\} \quad \text{for any } \boldsymbol{\xi}_1, \boldsymbol{\xi}_2 \in U.$$

For any $u \in C^m(U)$, define

$$\begin{split} \|u\|_{m,0,U}^{(\sigma),\Gamma} &\coloneqq \sum_{0 \le |\boldsymbol{\beta}| \le m} \sup_{\boldsymbol{\xi}_1 \in U} \left(\delta_{\boldsymbol{\xi}_1}^{\max\{|\boldsymbol{\beta}| + \sigma, 0\}} \left| D^{\boldsymbol{\beta}} u(\boldsymbol{\xi}_1) \right| \right), \\ [u]_{m,\alpha,U}^{(\sigma),\Gamma} &\coloneqq \sum_{|\boldsymbol{\beta}| = m} \sup_{\substack{\boldsymbol{\xi}_1, \boldsymbol{\xi}_2 \in U\\ \boldsymbol{\xi}_1 \neq \boldsymbol{\xi}_2}} \left(\delta_{\boldsymbol{\xi}_1, \boldsymbol{\xi}_2}^{\max\{m + \alpha + \sigma, 0\}} \frac{|D^{\boldsymbol{\beta}} u(\boldsymbol{\xi}_1) - D^{\boldsymbol{\beta}} u(\boldsymbol{\xi}_2)|}{|\boldsymbol{\xi}_1 - \boldsymbol{\xi}_2|^{\alpha}} \right), \\ \|u\|_{m,\alpha,U}^{(\sigma),\Gamma} &\coloneqq \|u\|_{m,0,U}^{(\sigma),\Gamma} + [u]_{m,\alpha,U}^{(\sigma),\Gamma}, \end{split}$$

where $D^{\boldsymbol{\beta}} \coloneqq \partial_{\boldsymbol{\xi}}^{\beta_1} \partial_{\boldsymbol{\eta}}^{\beta_2}$ for $\boldsymbol{\beta} \coloneqq (\beta_1, \beta_2)$ with $\beta_1, \beta_2 \in \mathbb{N}$ and $|\boldsymbol{\beta}| = \beta_1 + \beta_2$.

(ii) For any open bounded interval $I \subseteq \mathbb{R}$, let $x_0 \in \partial I$ be an endpoint of I. For any $f \in C^m(I)$, define

$$\begin{split} \|f\|_{m,0,I}^{(\sigma),\{x_0\}} &\coloneqq \sum_{k=0}^m \sup_{x \in I} \left(|x - x_0|^{\max\{k + \sigma, 0\}} |f^{(k)}(x)| \right), \\ [f]_{m,\alpha,I}^{(\sigma),\{x_0\}} &\coloneqq \sup_{\substack{x_1, x_2 \in I \\ x_1 \neq x_2}} \left(\min\left\{ |x_1 - x_0|, |x_2 - x_0| \right\}^{\max\{m + \alpha + \sigma, 0\}} \frac{|f^{(m)}(x_1) - f^{(m)}(x_2)|}{|x_1 - x_2|^{\alpha}} \right), \\ \|f\|_{m,\alpha,I}^{(\sigma),\{x_0\}} &\coloneqq \|f\|_{m,0,I}^{(\sigma),\{x_0\}} + [f]_{m,\alpha,I}^{(\sigma),\{x_0\}}. \end{split}$$

Proposition 3.16. Fix $\gamma \geq 1$ and $v_2 \in (v_2^s, 0)$. Let $\sigma_3 > 0$ be the constant from Proposition 3.14. For small constants $\sigma_{\rm s} \in (0, \frac{\sigma_3}{2}]$ and $\sigma_{\rm d} \in (0, \frac{\theta^{\rm d}}{10})$, there exist $r \in (0, r_*)$, $\alpha \in (0, 1)$, and C > 0 depending only on $(\gamma, v_2, \sigma_{\rm s}, \sigma_{\rm d})$ such that any admissible solution φ corresponding to $\theta \in \Theta \cap$ $\{\theta^{s} + \sigma_{s} \leq \theta_{1} \leq \theta^{d} - \sigma_{d}\}$ satisfies the estimates:

(3.47)
$$\|\varphi\|_{2,\alpha,\Omega\cap B_r(P_0^1)}^{(-1-\alpha),\{P_0^1\}} + \|f_e\|_{2,\alpha,(0,r)}^{(-1-\alpha),\{0\}} \le C, \qquad |D^m_{\boldsymbol{\xi}}(\varphi-\varphi_5)(P_0^1)| = 0 \text{ for } m = 0,1.$$

The proof is similar to the proof of [2, Proposition 3.42]. We omit the details of the proof and sketch only the main ideas here.

1. Hodograph transform. Define $\bar{\phi}^{(h)}(S,T) := \bar{\phi}(\boldsymbol{\xi}(S,T))$ for function $\bar{\phi} = \varphi_2 - \varphi$ and the (S,T)-coordinates given by (3.45). By property (3.44), we can apply the hodograph transform $y = (y_1, y_2) = (\overline{\phi}^{(h)}(S, T), T)$ to $\overline{\phi}$ for the original system (3.42). We write $S \rightleftharpoons v(y_1, y_2)$ if it satisfies $\bar{\phi}^{(h)}(S, y_2) = y_1$ for any

$$\boldsymbol{y} \in \mathcal{D}_{r_*}^{(\theta_1)} \coloneqq \left\{ (\bar{\phi}^{(h)}(S,T),T) : \boldsymbol{\xi}(S,T) \in \Omega \cap B_{r_*}(P_0^1) \right\},\$$

from which we obtain

$$\partial_{y_1} v = \frac{1}{\partial_S \bar{\phi}^{(h)}}, \qquad \quad \partial_{y_2} v = -\frac{\partial_T \bar{\phi}^{(h)}}{\partial_S \bar{\phi}^{(h)}}.$$

It follows from Lemma 3.3 and property (3.44) that there exists a constant K > 1 depending only on (γ, v_2) such that

(3.48)
$$\frac{1}{K} \le \partial_{y_1} v \le \frac{2}{d'_0}, \qquad |v| + |D_y v| \le 2K.$$

Under the hodograph transform, system (3.42) is equivalent to

(3.49)
$$\begin{cases} \sum_{i,j=1}^{2} a_{ij}(D_{\boldsymbol{y}}v,v,\boldsymbol{y})\partial_{y_{i}y_{j}}v = 0 & \text{ in } \mathcal{D}_{r_{*}}^{(\theta_{1})}, \\ g_{h}^{\text{sym}}(D_{\boldsymbol{y}}v,v,\boldsymbol{y}) = 0 & \text{ on } \Gamma_{\text{sym}}^{(h)}, \\ g_{h}^{\text{sh}}(D_{\boldsymbol{y}}v,v,\boldsymbol{y}) = 0 & \text{ on } \Gamma_{\text{shock}}^{(h)}, \end{cases}$$

for suitable functions $(a_{ij}, g_h^{\text{sym}}, g_h^{\text{sh}})$, with the boundary sets given by

$$\begin{split} \Gamma_{\mathrm{sym}}^{(h)} &\coloneqq \left\{ (\bar{\phi}^{(h)}(S,T),T) \,:\, \boldsymbol{\xi}(S,T) \in \Gamma_{\mathrm{sym}} \cap B_{r_*}(P_0^1) \right\}, \\ \Gamma_{\mathrm{shock}}^{(h)} &\coloneqq \left\{ (0,T) \,:\, \boldsymbol{\xi}(S,T) \in \Gamma_{\mathrm{shock}} \cap B_{r_*}(P_0^1) \right\}. \end{split}$$

Due to (3.48), we extend the domain of functions $(a_{ij}, g_h^{\text{sym}}, g_h^{\text{sh}})$ to the set:

$$U \coloneqq \left\{ (\boldsymbol{p}, z, \boldsymbol{y}) \in \mathbb{R}^2 \times \mathbb{R} \times \mathcal{D}_{r_*}^{(\theta_1)} : |\boldsymbol{p}| + |z| \le 2K \right\},\$$

and continue to use the same notation as in (3.49) for this modified system.

2. Hölder continuity of g_h^{sym} at the corner. From Lemma 3.3, there exists a constant $\tau_0 > 0$ depending only on (γ, v_2) such that

$$\Gamma_{\text{sym}}^{(h)} \cup \Gamma_{\text{shock}}^{(h)} \subseteq \{ \boldsymbol{y} \in \mathbb{R}^2 : y_2 > \tau_0 |y_1| \}.$$

Moreover, $\Gamma_{\text{shock}}^{(h)}$ is in C^2 up to endpoint $\{\mathbf{0}\}$. From the definition of admissible solutions and the hodograph transform, we obtain that $v \in$ $C^{1}(\overline{\mathcal{D}_{r_{*}}^{(\theta_{1})}}) \cap C^{2}(\mathcal{D}_{r_{*}}^{(\theta_{1})} \cup \Gamma_{\mathrm{shock}}^{(h)}) \cap C^{3}(\mathcal{D}_{r_{*}}^{(\theta_{1})})$ satisfies (3.48). From (3.43) and the notation defined in Step **1**, we can obtain the boundedness of coefficients $(a_{ij}, g_{h}^{\mathrm{sym}}, g_{h}^{\mathrm{sh}})$ in the modified system (3.49). For any $\theta \in \Theta \cap \{\theta^{s} + \sigma_{s} \leq \theta_{1} < \theta^{d}\}$ with $\sigma_{s} \in (0, \frac{\sigma_{3}}{2}]$, we may reduce $\lambda_{*} > 0$ depending only on (γ, v_2, σ_s) in (3.41) to obtain the uniform ellipticity of the modified system (3.49) in $\mathcal{D}_{r_*}^{(\theta_1)}$ by using (3.48) and the hodograph transform. It follows from Lemma 3.3, (3.41), and (3.48) that there exists a constant $\lambda_1 > 0$ depending only on (γ, v_2, σ_s) such that, for any $y \in \Gamma_{\text{shock}}^{(h)}$,

$$\boldsymbol{\nu} \cdot D_{\boldsymbol{p}} g_h^{\mathrm{sh}}(D_{\boldsymbol{y}} v, v, \boldsymbol{y}) = \frac{1}{v_{y_1}} D \bar{\phi} \cdot D_{\boldsymbol{q}} g^{\mathrm{sh}}(D \varphi, \varphi, \boldsymbol{\xi}) \ge \lambda_1 \,,$$

where we have used the expression of $D_{\boldsymbol{q}}g^{\mathrm{sh}}(D\varphi,\varphi,\boldsymbol{\xi})$ on $\Gamma_{\mathrm{shock}}^{(h)}$, and $\boldsymbol{\nu}=(1,0)$ is the unit normal vector to $\Gamma_{\text{shock}}^{(h)}$. From (3.48), $|D_{\boldsymbol{p}}g_h^{\text{sym}}(D_{\boldsymbol{y}}v, v, \boldsymbol{y})| \ge \frac{d'_0}{2} > 0.$

Similar to the proof of [2, Lemma 3.40], one can show that, for small $\sigma_{\rm d} \in (0, \frac{\theta^{\rm d}}{10})$, there exist constants M > 0 and $\bar{r} \in (0, r_*)$ depending only on (γ, v_2, σ_d) such that any admissible solution φ corresponding to $\theta \in \Theta \cap \{\theta^{s} \leq \theta_{1} < \theta^{d} - \sigma_{d}\}$ satisfies

$$\partial_{q_1} g_{\mathrm{mod}}^{\mathrm{sh}}(D\varphi,\varphi,\boldsymbol{\xi}) \leq -M^{-1} \quad \text{for all } \boldsymbol{\xi} \in \Gamma_{\mathrm{shock}} \cap B_{\bar{r}}(P_0^1).$$

Then, for any $\boldsymbol{y} \in \Gamma_{\text{shock}}^{(h)}$, $|\det D_{\boldsymbol{p}}(g_h^{\text{sym}}, g_h^{\text{sh}})| > 0$ holds. It follows from [10, Proposition 4.3.7] that there exist constants $\alpha_1 \in (0, 1)$, C > 0, and $r_0 \in (0, \bar{r}]$ depending only on $(\gamma, v_2, \sigma_s, \sigma_d)$ such that, for any $\boldsymbol{y} \in \overline{\mathcal{D}_{\bar{r}}^{(\theta_1)} \cap B_{r_0}(\mathbf{0})}$,

(3.50)
$$\left|g_h^{\text{sym}}(D_{\boldsymbol{y}}v(\boldsymbol{y}),v(\boldsymbol{y}),\boldsymbol{y}) - g_h^{\text{sym}}(D_{\boldsymbol{y}}v(\boldsymbol{0}),v(\boldsymbol{0}),\boldsymbol{0})\right| \le C|\boldsymbol{y}|^{\alpha_1}.$$

3. Hölder continuity of $v(\mathbf{y})$ at the corner. Notice that both g_h^{sym} and g_h^{sh} are Lipschitz continuous on (p, z, \boldsymbol{y}) . Since $v \in C^1(\overline{\mathcal{D}_{\bar{r}}^{(\theta_1)}})$, inequality (3.50) also holds for any $\boldsymbol{y} \in \Gamma_{\text{shock}}^{(h)}$ after replacing g_h^{sym} by g_h^{sh} . Moreover, functions $(g_h^{\text{sym}}, g_h^{\text{sh}})(\cdot, v(\mathbf{0}), \mathbf{0})$ are bounded in $C^{1,\alpha_1}(\overline{B_{2K}(D_{\boldsymbol{y}}v(\mathbf{0}))})$, and $|\det D_{\boldsymbol{p}}(g_h^{\text{sym}}, g_h^{\text{sh}})(D_{\boldsymbol{y}}v(\mathbf{0}), \mathbf{0})| > 0$ follows from Step 2. Then, from [10, Proposition 4.3.9], there are a constant of the properties of the p there exist $r_1 \in (0, r_0)$ and $C_1 > 0$ depending only on $(\gamma, v_2, \sigma_s, \sigma_d)$ such that

(3.51)
$$|D_{\boldsymbol{y}}v(\boldsymbol{y}) - D_{\boldsymbol{y}}v(\boldsymbol{0})| \le C_1 |\boldsymbol{y}|^{\alpha_1} \quad \text{for any } \boldsymbol{y} \in \Gamma_{\text{shock}}^{(h)} \cap B_{r_1}(\boldsymbol{0}).$$

By the hodograph transform and the geometric properties of the domain near corner P_0^1 , it follows from (3.50) - (3.51) that

(3.52)
$$\begin{aligned} |\partial_{\eta}\varphi(\boldsymbol{\xi}) - \partial_{\eta}\varphi(P_{0}^{1})| &\leq C|\boldsymbol{\xi} - P_{0}^{1}|^{\alpha_{1}} \quad \text{for any } \boldsymbol{\xi} \in \overline{\Omega \cap B_{r_{1}}(P_{0}^{1})}, \\ |D\varphi(\boldsymbol{\xi}) - D\varphi(P_{0}^{1})| &\leq C|\boldsymbol{\xi} - P_{0}^{1}|^{\alpha_{1}} \quad \text{for any } \boldsymbol{\xi} \in \Gamma_{\text{shock}} \cap B_{r_{1}}(P_{0}^{1}). \end{aligned}$$

The open domain $\Omega_{r_1} \coloneqq \Omega \cap B_{r_1}(P_0^1)$ contains two Lipschitz boundaries $\Gamma^1 \coloneqq \Gamma_{\text{shock}} \cap B_{r_1}(P_0^1)$ and $\Gamma^2 := \Gamma_{\text{sym}} \cap B_{r_1}(P_0^1)$ intersecting at corner P_0^1 . We also take

$$B^{(1)}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq (1, 0) \cdot (\boldsymbol{p} - D\bar{\phi}(P_0^1)), \qquad B^{(2)}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq \bar{g}_{\text{sym}}(\boldsymbol{p}, z, \boldsymbol{\xi}).$$

Now we update system (3.42) for $\bar{\phi} \in C^1(\overline{\Omega_{r_1}}) \cap C^2(\Omega_{r_1} \cup \Gamma^1) \cap C^3(\Omega_{r_1})$ into domain Ω_{r_1} , replacing the boundary conditions by

$$B^{(1)}(D\bar{\phi},\bar{\phi},\boldsymbol{\xi}) = \bar{h}(\boldsymbol{\xi}) \text{ on } \Gamma^1, \qquad B^{(2)}(D\bar{\phi},\bar{\phi},\boldsymbol{\xi}) = 0 \text{ on } \Gamma^2,$$

where $\bar{h}(\boldsymbol{\xi}) \coloneqq (1,0) \cdot (D\bar{\phi}(\boldsymbol{\xi}) - D\bar{\phi}(P_0^1))$, which satisfies

$$|\bar{h}(\boldsymbol{\xi}) - \bar{h}(P_0^1)| \le C |\boldsymbol{\xi} - P_0^1|^{\alpha_1} \quad \text{for any } \boldsymbol{\xi} \in \Gamma_{\text{shock}} \cap B_{r_1}(P_0^1) \,.$$

We can also extend the domain of functions $(\bar{A}_{ij}, B^{(1)}, B^{(2)})$ for the updated system to the set:

$$V \coloneqq \{(\boldsymbol{p}, z, \boldsymbol{\xi}) \in \mathbb{R}^2 \times \mathbb{R} \times \overline{\Omega_{r_1}} : |\boldsymbol{p}| + |z| \le 2K_1\}.$$

Using [10, Proposition 4.3.7] again, we obtain that there exist $\alpha \in (0, \alpha_1], r_2 \in (0, r_1]$, and C > 0depending only on $(\gamma, v_2, \sigma_s, \sigma_d)$ such that, for any $\boldsymbol{\xi} \in \Omega \cap B_{r_2}(P_0^1)$,

$$|B^{(1)}(D\bar{\phi}(\boldsymbol{\xi}),\bar{\phi}(\boldsymbol{\xi}),\boldsymbol{\xi}) - B^{(1)}(D\bar{\phi}(P_0^1),\bar{\phi}(P_0^1),P_0^1)| \le C|\boldsymbol{\xi} - P_0^1|^{lpha},$$

which, combined with (3.52), implies that

(3.53)
$$|D\varphi(\boldsymbol{\xi}) - D\varphi(P_0^1)| \le C |\boldsymbol{\xi} - P_0^1|^{\alpha} \quad \text{for any } \boldsymbol{\xi} \in \overline{\Omega \cap B_{r_2}(P_0^1)}.$$

4. Weighted Hölder estimate near corner P_0^1 . From Lemma 3.2, Remark 2.2(i), and (3.53), there exists a constant $\varepsilon_{\rm bd} > 0$ depending only on (γ, v_2) such that, for any $\boldsymbol{\xi}_0 \in \Gamma_{\rm bd} \cap B_{r_3}(P_0^1)$ for some $r_3 \in (0, r_2]$ with either $\Gamma_{bd} \coloneqq \Gamma_{shock}$ or $\Gamma_{bd} \coloneqq \Gamma_{sym}$,

$$\left\{ oldsymbol{\xi} \in \partial \Omega \, : \, |oldsymbol{\xi} - oldsymbol{\xi}_0| < arepsilon_{
m bd} |oldsymbol{\xi}_0 - P_0^1|
ight\} \subseteq \Gamma_{
m bd}$$
 .

We now consider the following problem for $\varphi \in C^1(\overline{\Omega \cap B_{r_3}(P_0^1)}) \cap C^3(\Omega \cap B_{r_3}(P_0^1))$:

$$\begin{cases} \operatorname{div} \tilde{\mathcal{A}}(D\varphi,\varphi) + \tilde{\mathcal{B}}(D\varphi,\varphi) = 0 & \text{ in } \Omega \cap B_{r_3}(P_0^1) ,\\ g_{\mathrm{mod}}^{\mathrm{sh}}(D\varphi,\varphi,\boldsymbol{\xi}) = 0 & \text{ on } \Gamma_{\mathrm{shock}} \cap B_{r_3}(P_0^1) ,\\ g_{\mathrm{sym}}(D\varphi,\varphi,\boldsymbol{\xi}) = 0 & \text{ on } \Gamma_{\mathrm{sym}} \cap B_{r_3}(P_0^1) , \end{cases}$$

where $(\tilde{\mathcal{A}}, \tilde{\mathcal{B}})(\boldsymbol{q}, w)$ are defined on $\mathbb{R}^2 \times \mathbb{R}$ from Lemma 3.5, $g_{\text{mod}}^{\text{sh}}(\boldsymbol{q}, w, \boldsymbol{\xi})$ is defined on $\mathbb{R}^2 \times \mathbb{R} \times \overline{\Omega \cap B_{r_3}(P_0^1)}$ by (3.28), and $g_{\text{sym}}(\boldsymbol{q}, w, \boldsymbol{\xi}) = (0, 1) \cdot \boldsymbol{q}$. Combining the above with property (3.53) and using [10, Proposition 4.3.11(i)], we obtain that there exist $\tilde{\alpha} \in (0, \alpha]$ and C > 0 depending only on $(\gamma, v_2, \sigma_{\text{s}}, \sigma_{\text{d}})$ such that $\varphi \in C^{1, \tilde{\alpha}}(\Omega \cap B_r(P_0^1))$ satisfies $\|\varphi\|_{1, \tilde{\alpha}, \Omega \cap B_r(P_0^1)} \leq C$ and, for Γ_{shock} given by (3.46), $\|f_{\boldsymbol{e}}\|_{1, \tilde{\alpha}, (0, r)} \leq C$. Finally, using [10, Proposition 4.3.11(i)], we obtain (3.47).

3.4. Compactness of the set of admissible solutions. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$. For any $\theta_* \in (0, \theta^d)$, the arguments in §3.3 are also valid for the weighted $C^{2,\alpha}$ -estimates near Γ_{sonic}^6 with respect to $\theta_2 \in [0, \theta^d)$. According to all the *a priori* estimates obtained in Lemma 3.9, Corollary 3.10, and Propositions 3.12–3.14 and 3.16, there exists a constant $\bar{\alpha} \in (0, 1)$ depending only on (γ, v_2, θ_*) such that the set:

$$\left\{ \left\|\varphi\right\|_{1,\bar{\alpha},\overline{\Omega}} + \left\|f_{\mathrm{sh}}\right\|_{1,\bar{\alpha},\left[\xi^{P_{3}},\xi^{P_{2}}\right]} : \begin{array}{c} \varphi \text{ is an admissible solution} \\ \mathrm{corresponding to } \boldsymbol{\theta} \in \Theta \cap [0,\theta_{*}]^{2} \end{array} \right\}$$

is bounded, where $\eta = f_{\rm sh}(\xi)$ is the expression of $\Gamma_{\rm shock}$ as a graph, given by Lemma 3.2. For each admissible solution, the corresponding pseudo-subsonic region Ω is a bounded domain enclosed by $\Gamma_{\rm sonic}^5$, $\Gamma_{\rm shock}$, $\Gamma_{\rm sonic}^6$, and $\Gamma_{\rm sym}$. These four curves intersect only at P_i for i = 1, 2, 3, 4. Note that $\Gamma_{\rm sonic}^5$, O_5 , P_1 , and P_2 depend continuously on $\theta_1 \in [0, \theta^d]$, while $\Gamma_{\rm sonic}^6$, O_6 , P_3 , and P_4 depend continuously on $\theta_2 \in [0, \theta^d]$. Combining the above results with (3.13) and Lemma 3.7, we have the following compactness result; the details of the proof can be found in [10, Proposition 11.6.1].

Lemma 3.17. Fix $\gamma \geq 1$, $v_2 \in (v_{\min}, 0)$, and $\theta_* \in (0, \theta^d)$. Let $\{\boldsymbol{\theta}^{(j)}\}_{j \in \mathbb{N}} \subseteq \Theta \cap [0, \theta_*]^2$ be a sequence of parameters satisfying that $\lim_{j\to\infty} \boldsymbol{\theta}^{(j)} = \boldsymbol{\theta}^{(\infty)}$ for some $\boldsymbol{\theta}^{(\infty)} \in [0, \theta_*]^2$. For each $j \in \mathbb{N}$, let $\varphi^{(j)}$ be an admissible solution corresponding to parameters $\boldsymbol{\theta}^{(j)}$, with pseudo-subsonic region $\Omega^{(j)}$ and curved transonic shock $\Gamma^{(j)}_{\text{shock}}$. Then there exists a subsequence $\{\varphi^{(j_k)}\}_{k\in\mathbb{N}}$ such that the following properties hold:

- (i) $\{\varphi^{(j_k)}\}_{k\in\mathbb{N}}$ converges uniformly on any compact subset of $\overline{\mathbb{R}^2_+}$ to a function $\varphi^{(\infty)} \in C^{0,1}_{\text{loc}}(\overline{\mathbb{R}^2_+})$, and the limit function $\varphi^{(\infty)}$ is an admissible solution corresponding to $\theta^{(\infty)}$;
- (ii) $\Omega^{(j_k)} \to \Omega^{(\infty)}$ in the Hausdorff metric;
- (iii) If $\boldsymbol{\xi}^{(j_k)} \in \overline{\Omega^{(j_k)}}$ for each $k \in \mathbb{N}$, and $\{\boldsymbol{\xi}^{(j_k)}\}_{k \in \mathbb{N}}$ converges to $\boldsymbol{\xi}^{(\infty)} \in \overline{\Omega^{(\infty)}}$, then

$$\varphi^{(j_k)}(\boldsymbol{\xi}^{(j_k)}) \to \varphi^{(\infty)}(\boldsymbol{\xi}^{(\infty)}), \quad D\varphi^{(j_k)}(\boldsymbol{\xi}^{(j_k)}) \to D\varphi^{(\infty)}(\boldsymbol{\xi}^{(\infty)}) \qquad as \ k \to \infty,$$

where, in the case that $\boldsymbol{\xi}^{(j_k)} \in \Gamma_{\text{shock}}^{(j_k)}$, the derivative of $\varphi^{(j_k)}$ at $\boldsymbol{\xi}^{(j_k)}$ is defined as the limit of the derivative from the interior of $\Omega^{(j_k)}$:

$$D\varphi^{(j_k)}(\boldsymbol{\xi}^{(j_k)}) \coloneqq \lim_{\boldsymbol{\xi} \to \boldsymbol{\xi}^{(j_k)}, \, \boldsymbol{\xi} \in \Omega^{(j_k)}} D\varphi^{(j_k)}(\boldsymbol{\xi}) \,,$$

and, similarly, for $\boldsymbol{\xi}^{(\infty)} \in \Gamma_{\text{shock}}^{(\infty)}$,

$$D\varphi^{(\infty)}(\boldsymbol{\xi}^{(\infty)}) \coloneqq \lim_{\boldsymbol{\xi} \to \boldsymbol{\xi}^{(\infty)}, \, \boldsymbol{\xi} \in \Omega^{(\infty)}} D\varphi^{(\infty)}(\boldsymbol{\xi}) \,.$$

4. Iteration Method and Existence of Admissible Solutions

In this section, we carefully construct the iteration set and the iteration map based on the uniform estimates obtained in Section 3. Then we apply the Leray-Schauder degree theorem to show the existence of admissible solutions. The construction of the iteration set follows closely the process described in [2, Chapter 4] with the main difference that both sonic arcs Γ_{sonic}^5 and Γ_{sonic}^6 may degenerate into single points. In this section, we always fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$.

4.1. Mapping between the elliptic and standard domains.

4.1.1. Mapping into the standard domain. Define the standard domain to be the rectangle $Q^{\text{iter}} := (-1,1) \times (0,1)$. Before constructing a map between Ω and Q^{iter} , we introduce some useful notation and basic geometric properties. Let φ_2 be given by (2.17), and let (φ_5, φ_6) be given by (2.33). For any $\delta_0 > 0$, we define $S_{2j}^{\delta_0} := \{ \boldsymbol{\xi} \in \mathbb{R}^2 : (\varphi_2 - \varphi_j)(\boldsymbol{\xi}) = -\delta_0 \}$ and $q_j^{\delta_0} := \text{dist}(O_j, S_{2j}^{\delta_0})$ for j = 5, 6, and also write

$$u_5^{\delta_0} \coloneqq u_5 + q_5^{\delta_0} \sin \theta_{25} \ge 0, \qquad u_6^{\delta_0} \coloneqq u_6 - q_6^{\delta_0} \sin \theta_{26} \le 0.$$

For constant $\hat{\delta} > 0$ from (3.15), we set $\tilde{\delta}_0 \coloneqq \frac{\hat{\delta}}{4}$, which depends only on (γ, v_2) . Then, for \hat{c}_j given by (3.14), a direct calculation gives that, for any $\boldsymbol{\theta} \in \overline{\Theta}$,

$$\hat{c}_j - q_j^{\delta_0} \ge \hat{\delta} - \frac{\delta_0}{\ell(\rho_j)} \ge 2\tilde{\delta}_0$$

after choosing $\delta_0 > 0$ small enough, depending only on (γ, v_2) , and using Lemma 2.8. Therefore, $S_{2j}^{\delta_0}$ and $\partial B_{\hat{c}_j}(O_j)$ intersect at two distinct points for any $\boldsymbol{\theta} \in \overline{\Theta}$.

Definition 4.1 (Extended domain Q^{θ}). Let $\delta_0 > 0$ be chosen as above. Define P'_2 and P'_3 by

$$\{P_2'\} \coloneqq S_{25}^{\delta_0} \cap \partial B_{\hat{c}_5}(O_5) \cap \{\xi \ge u_5^{\delta_0}\}, \qquad \{P_3'\} \coloneqq S_{26}^{\delta_0} \cap \partial B_{\hat{c}_6}(O_6) \cap \{\xi \le u_6^{\delta_0}\}.$$

Denote by $\Gamma_{\text{sonic}}^{5,\delta_0}$ the arc of $\partial B_{\hat{c}_5}(O_5) \cap \{\xi \geq u_5^{\delta_0}\}$ with endpoints $\{P_1, P_2'\}$, and denote by $\Gamma_{\text{sonic}}^{6,\delta_0}$ the arc of $\partial B_{\hat{c}_6}(O_6) \cap \{\xi \leq u_6^{\delta_0}\}$ with endpoints $\{P_4, P_3'\}$. Define the extended domain Q^{θ} as the open bounded region enclosed by $\Gamma_{\text{sonic}}^{5,\delta_0}, \Gamma_{\text{sonic}}^{6,\delta_0}, S_{25}^{\delta_0}, S_{26}^{\delta_0}$, and Γ_{sym} ; see Fig. 4.1.

For any $\boldsymbol{\xi} = (\xi, \eta) \in (\mathbb{R} \setminus [u_6^{\delta_0}, u_5^{\delta_0}]) \times [0, \infty)$, define the (x, y)-coordinates by $(x, y) \coloneqq \mathcal{R}(\boldsymbol{\xi})$ if and only if

(4.1)
$$\boldsymbol{\xi} = \begin{cases} O_6 + (c_6 - x) \left(\cos(\pi - y), \sin(\pi - y) \right) & \text{if } \xi < u_6^{\delta_0}, \\ O_5 + (c_5 - x) \left(\cos y, \sin y \right) & \text{if } \xi > u_5^{\delta_0}, \end{cases}$$

with $y \in (0, \pi)$. For any constant $\epsilon > 0$, we define two sets $\mathcal{D}_{\epsilon}^{5}, \mathcal{D}_{\epsilon}^{6} \subseteq Q^{\theta}$ by (4.2) $\mathcal{D}_{\epsilon}^{5} \coloneqq \{\boldsymbol{\xi} \in Q^{\theta} : \boldsymbol{\xi} > u_{5}^{\delta_{0}}\} \setminus \overline{B_{\hat{c}_{5}-\epsilon}(O_{5})}, \qquad \mathcal{D}_{\epsilon}^{6} \coloneqq \{\boldsymbol{\xi} \in Q^{\theta} : \boldsymbol{\xi} < u_{6}^{\delta_{0}}\} \setminus \overline{B_{\hat{c}_{6}-\epsilon}(O_{6})}.$ Let $\omega_{0} \in (0, \frac{\pi}{2})$ be the solution of $(q^{\delta_{0}} + \tilde{\delta}_{0}) \cos \omega_{0} = q^{\delta_{0}}$, where $q^{\delta_{0}} \coloneqq \max\{q_{5}^{\delta_{0}}, q_{6}^{\delta_{0}}\}$. Then $\tilde{\omega} \leftarrow \frac{1}{2} \inf \{Q_{1}, \dots, \overline{\alpha}, \overline{\alpha}, \overline{\alpha}, \dots, \overline{\alpha}\} > 0$

$$\tilde{\omega}_0 \coloneqq \frac{1}{2} \inf_{\boldsymbol{\theta} \in \overline{\Theta}} \left\{ \theta_{25} - \frac{\pi}{2}, \, \frac{\pi}{2} - \theta_{26}, \, \omega_0 \right\} > 0$$

by (3.14)–(3.15) and Lemma 2.8. A direct computation shows that, for any $\epsilon \in (0, \delta_0)$,

(4.3)
$$\mathcal{D}_{\epsilon}^{5} \subseteq \left\{ \boldsymbol{\xi} : x_{P_{2}} < x < x_{P_{2}} + \epsilon, \ 0 < y < \theta_{25} - \frac{\pi}{2} - \tilde{\omega}_{0} \right\} \cap \left\{ \boldsymbol{\xi} > u_{5}^{\delta_{0}} + \tilde{\delta}_{0} \sin \tilde{\omega}_{0} \right\},$$
$$\mathcal{D}_{\epsilon}^{6} \subseteq \left\{ \boldsymbol{\xi} : x_{P_{3}} < x < x_{P_{3}} + \epsilon, \ 0 < y < \frac{\pi}{2} - \theta_{26} - \tilde{\omega}_{0} \right\} \cap \left\{ \boldsymbol{\xi} < u_{6}^{\delta_{0}} - \tilde{\delta}_{0} \sin \tilde{\omega}_{0} \right\}.$$



FIG. 4.1. The extended domain Q^{θ} in grey for the case: $\theta \in [\theta^{s}, \theta^{d}) \times (0, \theta^{s})$.
In particular, we fix a constant k > 4 sufficiently large, depending only on (γ, v_2) , such that $\max\{\hat{c}_5, \hat{c}_6\} \leq \frac{k\tilde{\delta}_0}{4} \sin \tilde{\omega}_0$ for any $\boldsymbol{\theta} \in \overline{\Theta}$ and

(4.4)
$$\mathcal{D}_{\frac{3\hat{c}_5}{k}}^5 \subseteq \left\{ \xi > u_5^{\delta_0} + \frac{3\hat{c}_5}{k} \right\}, \qquad \mathcal{D}_{\frac{3\hat{c}_6}{k}}^6 \subseteq \left\{ \xi < u_6^{\delta_0} - \frac{3\hat{c}_6}{k} \right\}.$$

For each $\boldsymbol{\theta} \in \overline{\Theta}$, we define an invertible map $\mathcal{G}_1^{\boldsymbol{\theta}} : \overline{Q^{\boldsymbol{\theta}}} \to [-1,1] \times [0,\infty)$, which flattens the extended sonic arcs $\Gamma_{\text{sonic}}^{5,\delta_0}$ and $\Gamma_{\text{sonic}}^{6,\delta_0}$. For each admissible solution φ corresponding to parameters $\boldsymbol{\theta} \in \overline{\Theta}$, we further construct a map $G_{2,\mathfrak{g}_{\text{sh}}}$ from $\mathcal{G}_1^{\boldsymbol{\theta}}(\Omega)$ to the standard domain $\mathcal{Q}^{\text{iter}}$, where Ω is the pseudo-subsonic domain associated with φ . Finally, we present the *a priori* estimates in the standard domain for admissible solutions under map $G_{2,\mathfrak{g}_{\text{sh}}} \circ \mathcal{G}_1^{\boldsymbol{\theta}}$.

Construction of $\mathcal{G}_1^{\boldsymbol{\theta}}$. For $r_0 \in \mathbb{R}$ and $r_d > 0$, we fix a basic cut-off function $\tilde{\zeta}(\cdot; r_0, r_d) \in C^4(\mathbb{R})$ such that $\tilde{\zeta}'(\cdot; r_0, r_d) \in [0, \frac{2}{r_d}]$ and

(4.5)
$$\tilde{\zeta}(r; r_0, r_d) = \begin{cases} 0 & \text{for } r \le r_0, \\ 1 & \text{for } r \ge r_0 + r_d. \end{cases}$$

For j = 5, 6, let the cut-off functions (ζ_j, χ_j) be defined by

$$\zeta_j(r) \coloneqq \tilde{\zeta}(r; (1 - \frac{3}{k})\hat{c}_j, \frac{\hat{c}_j}{k}), \qquad \chi_j(\xi) \coloneqq \tilde{\zeta}((-1)^{j-1}\xi; (-1)^{j-1}u_j^{\delta_0}, \frac{2\hat{c}_j}{k}).$$

Then we define $F_1: \overline{Q^{\theta}} \to \mathbb{R}^2$ by $F_1(\xi, \eta) \coloneqq (h_1(\xi, \eta), \eta)$, where

$$h_1(\boldsymbol{\xi}) \coloneqq \zeta_{\mathrm{d}}(\boldsymbol{\xi}) u_r + (1 - \zeta_{\mathrm{d}}(\boldsymbol{\xi})) \xi,$$

and $\zeta_{\rm d}$ and u_r are given by

$$\begin{aligned} \zeta_{\rm d}(\boldsymbol{\xi}) &\coloneqq \chi_5(\xi)\zeta_5(|\boldsymbol{\xi} - O_5|) + \chi_6(\xi)\zeta_6(|\boldsymbol{\xi} - O_6|) \,, \\ u_r &\coloneqq (u_5 + |\boldsymbol{\xi} - O_5|)\chi_5(\xi) + (u_6 - |\boldsymbol{\xi} - O_6|)\chi_6(\xi) \,. \end{aligned}$$

For each $\theta \in \overline{\Theta}$, we write $s_5 \coloneqq u_5 + \hat{c}_5 > 0$ and $s_6 \coloneqq u_6 - \hat{c}_6 < 0$, which depend continuously on $\theta \in \overline{\Theta}$. Let $\mathcal{N}_{\varepsilon}(\Gamma)$ be defined by (3.21). By construction of map F_1 , we observe that

$$h_1(\boldsymbol{\xi}) = u_j + (-1)^{j-1} \operatorname{dist}(\boldsymbol{\xi}, O_j) \quad \text{for any } \boldsymbol{\xi} \in Q^{\boldsymbol{\theta}} \cap \mathcal{N}_{\frac{2\hat{c}_j}{k}}(\Gamma_{\operatorname{sonic}}^{j,\delta_0}),$$

and $\overline{F_1(Q^{\theta})} \subseteq [s_6, s_5] \times [0, \infty)$. Let the linear map $L_{\theta} : [s_6, s_5] \to [-1, 1]$ be given by

(4.6)
$$L_{\boldsymbol{\theta}}(s') \coloneqq d_L(s'-s_6) - 1 \quad \text{with } d_L \coloneqq \frac{2}{s_5 - s_6},$$

which maps interval $[s_6, s_5]$ to the standard interval [-1, 1]. For k > 4 given by (4.4), define the cut-off functions $\tilde{\chi}_j$ by

$$\tilde{\chi}_j(s') \coloneqq \tilde{\zeta}((-1)^{j-1}s'; (-1)^{j-1}u_j + (1-\frac{1}{k})\hat{c}_j, \frac{\hat{c}_j}{2k}) \quad \text{for } j = 5, 6.$$

We define a function $h_2: \overline{F_1(Q^{\theta})} \to [0,\infty)$ by

$$h_2(s',t'') \coloneqq \tilde{\chi}_5 \arcsin\left(\frac{t''}{s'-u_5}\right) + \tilde{\chi}_6 \arcsin\left(\frac{t''}{u_6-s'}\right) + (1-\tilde{\chi}_5-\tilde{\chi}_6)t'',$$

where $\tilde{\chi}_5$ and $\tilde{\chi}_6$ are evaluated at s'. For maps F_1 and h_2 constructed above, we see that

$$h_2 \circ F_1(\boldsymbol{\xi}) = y \qquad \text{for any } \boldsymbol{\xi} \in Q^{\boldsymbol{\theta}} \cap \left(\mathcal{N}_{\frac{\hat{c}_5}{2k}}(\Gamma_{\text{sonic}}^{5,\delta_0}) \cup \mathcal{N}_{\frac{\hat{c}_6}{2k}}(\Gamma_{\text{sonic}}^{6,\delta_0}) \right),$$

for the (x, y)-coordinates given by (4.1).

Finally, we define map $\overline{\mathcal{G}_1^{\theta}} : \overline{Q^{\theta}} \to [-1, 1] \times [0, \infty)$ by

(4.7)
$$\mathcal{G}_{1}^{\boldsymbol{\theta}}(\boldsymbol{\xi}) \coloneqq (L_{\boldsymbol{\theta}} \circ h_{1}(\boldsymbol{\xi}), h_{2} \circ F_{1}(\boldsymbol{\xi})) \quad \text{for any } \boldsymbol{\xi} \in \overline{Q^{\boldsymbol{\theta}}},$$

and write the coordinates $(s, t') \coloneqq \mathcal{G}_1^{\boldsymbol{\theta}}(\boldsymbol{\xi})$.

It can be verified directly that $\mathcal{G}_1^{\boldsymbol{\theta}}(\Gamma_{\text{sym}}) = \{(s,0) : -1 < s < 1\}$. Moreover, map $\mathcal{G}_1^{\boldsymbol{\theta}} : \overline{Q^{\boldsymbol{\theta}}} \rightarrow [-1,1] \times [0,\infty)$ constructed above satisfies the following properties by using (3.3), (4.5), and Lemma 3.17; the proof is similar to [2, Lemmas 4.3 and 4.5].

Lemma 4.2. There exists a constant $\mu_{\mathcal{G}_1} \in (0,1)$ depending only on (γ, v_2) such that, for each $\theta \in \overline{\Theta}$, map \mathcal{G}_1^{θ} defined by (4.7) satisfies the following:

- (i) $\|\mathcal{G}_1^{\boldsymbol{\theta}}\|_{C^4(\overline{Q^{\boldsymbol{\theta}}})} + \|(\mathcal{G}_1^{\boldsymbol{\theta}})^{-1}\|_{C^4(\overline{\mathcal{G}_1^{\boldsymbol{\theta}}(Q^{\boldsymbol{\theta}}))}} \leq \mu_{\mathcal{G}_1}^{-1}, and |\det(D\mathcal{G}_1^{\boldsymbol{\theta}})| \geq \mu_{\mathcal{G}_1} in \overline{Q^{\boldsymbol{\theta}}};$
- (ii) For j = 5, 6, and for any $(s, t') \in \overline{\mathcal{G}_1^{\theta}(Q^{\theta})}, \quad \partial_{t'} ((\varphi_2 \varphi_j) \circ (\mathcal{G}_1^{\theta})^{-1})(s, t') \leq -\mu_{\mathcal{G}_1};$
- (iii) For any $\theta_* \in (0, \theta^d)$, there exists a constant $\mu_* > 0$ depending only on (γ, v_2, θ_*) such that any admissible solution φ corresponding to $\theta \in \Theta \cap \{\theta_1, \theta_2 \leq \theta_*\}$ satisfies

$$\partial_{t'} ((\varphi_2 - \varphi) \circ (\mathcal{G}_1^{\boldsymbol{\theta}})^{-1})(s, t') \le -\mu_* < 0 \quad \text{for all } (s, t') \in \mathcal{G}_1^{\boldsymbol{\theta}}(\Omega)$$

Using Lemma 4.2(ii), there exists a unique function $f_{\theta} \in C^{0,1}([-1,1])$ such that

$$\mathcal{G}_{1}^{\theta}(\{\boldsymbol{\xi} \in \mathbb{R}^{2}_{+} : \varphi_{2} = \max\{\varphi_{5}, \varphi_{6}\} - \delta_{0}\}) = \{(s, f_{\theta}(s)) : -1 < s < 1\},\$$

which follows from the implicit function theorem, and

(4.8)
$$\mathcal{G}_{1}^{\theta}(Q^{\theta}) = \left\{ (s, t') \in \mathbb{R}^{2}_{+} : -1 < s < 1, \ 0 < t' < f_{\theta}(s) \right\}$$

Using (3.13), Definition 2.11(ii), Lemma 3.9(ii), Corollary 3.10, Propositions 3.12–3.14 and 3.16, and Lemma 4.2, we introduce a function $\mathfrak{g}_{sh} : [-1,1] \to [0,\infty)$ representing Γ_{shock} in the (s,t')-coordinates and state its important properties; the proof is similar to [2, Proposition 4.6].

Proposition 4.3. For each admissible solution φ corresponding to $\theta \in \Theta$, there exists a unique function $\mathfrak{g}_{sh} : [-1,1] \to [0,\infty)$ satisfying the following properties:

(i) In the (s, t')-coordinates:

$$\begin{aligned} \mathcal{G}_{1}^{\boldsymbol{\theta}}(\Omega) &= \left\{ (s,t') : -1 < s < 1, \ 0 < t' < \mathfrak{g}_{\mathrm{sh}}(s) \right\}, \\ \mathcal{G}_{1}^{\boldsymbol{\theta}}(\Gamma_{\mathrm{shock}}) &= \left\{ (s,\mathfrak{g}_{\mathrm{sh}}(s)) : -1 < s < 1, \ t' = \mathfrak{g}_{\mathrm{sh}}(s) \right\} \end{aligned}$$

and, for any interval $I_1 \in (-1,1)$, $\|\mathfrak{g}_{sh}\|_{C^3(\overline{I_1})} \leq C_{I_1}$ for some constant $C_{I_1} > 0$ depending only on (γ, v_2, I_1) .

(ii) Let Q_0^{θ} be the bounded region enclosed by $\Gamma_{\text{sonic}}^5, \Gamma_{\text{sonic}}^6, S_{25}, S_{26}$, and Γ_{sym} , which satisfies $\Omega \subseteq Q_0^{\theta} \subseteq Q^{\theta}$. For \mathcal{D}_{ϵ}^5 and \mathcal{D}_{ϵ}^6 given by (4.2), there exist unique functions \mathfrak{g}_5 and \mathfrak{g}_6 with

 $\mathcal{G}_1^{\boldsymbol{\theta}}(Q_0^{\boldsymbol{\theta}} \cap \mathcal{D}_{\epsilon}^j) = \left\{ (s, t') \in \mathcal{Q}^{\text{iter}} : 1 + (-1)^j s < d_L \epsilon, \ 0 < t' < \mathfrak{g}_j(s) \right\} \qquad \text{for } j = 5, 6.$

Let $\epsilon_0^* > 0$ be the smallest constant ε_0 in Step 1 of the proofs of Propositions 3.12 and 3.14. For $d_L > 0$ given by (4.6), set $\hat{\epsilon}_0^* \coloneqq d_L \epsilon_0^*$. There exists a constant C > 0 depending only on (γ, v_2) such that

$$\|\mathfrak{g}_5\|_{C^3([1-\hat{\epsilon}_0^*,1])} + \|\mathfrak{g}_6\|_{C^3([-1,-1+\hat{\epsilon}_0^*])} \le C.$$

(iii) For each $\theta_* \in (0, \theta^d)$, there exist $\bar{\alpha} \in (0, 1)$ and $C_{\theta_*} > 0$ depending only on (γ, v_2, θ_*) such that, for any admissible solution corresponding to $\boldsymbol{\theta} \in \Theta \cap \{\theta_1, \theta_2 \leq \theta_*\}$,

$$\begin{aligned} \|\mathfrak{g}_{\rm sh}\|_{2,\bar{\alpha},(-1,-1+\hat{\epsilon}_0^*)}^{(-1-\bar{\alpha}),\{-1\}} + \|\mathfrak{g}_{\rm sh}\|_{2,\bar{\alpha},(1-\hat{\epsilon}_0^*,1)}^{(-1-\bar{\alpha}),\{1\}} &\leq C_{\theta_*} ,\\ \frac{\mathrm{d}^m}{\mathrm{d}s^m}(\mathfrak{g}_{\rm sh} - \mathfrak{g}_j)((-1)^{j-1}) = 0 \qquad for \ m = 0,1, \ and \ j = 5,6 \,. \end{aligned}$$

where $\|\cdot\|_{2,\alpha,I}^{(\sigma),\{x_0\}}$ is given by Definition 3.15(ii). Note that the properties above are equivalent to

$$\|\mathfrak{g}_{\rm sh} - \mathfrak{g}_5\|_{2,\bar{\alpha},(1-\hat{\epsilon}_0^*,1)}^{(1+\bar{\alpha}),({\rm par})} + \|\mathfrak{g}_{\rm sh} - \mathfrak{g}_6\|_{2,\bar{\alpha},(-1,-1+\hat{\epsilon}_0^*)}^{(1+\bar{\alpha}),({\rm par})} \le C_{\theta_*}'$$

for some constant $C'_{\theta_*} > 0$ depending only on (γ, v_2, θ_*) , where $\|\cdot\|^{(1+\alpha),(\text{par})}_{2,\alpha,I}$ is given by Definition 3.11(ii) with replacement of x by 1 - |s|.

(iv) For each $\theta_* \in (0, \theta^d)$, there exists a constant $\hat{k} > 1$ depending only on (γ, v_2, θ_*) such that, for any admissible solution φ corresponding to $\theta \in \Theta \cap \{\theta_1, \theta_2 \leq \theta_*\}$ and any $s \in [-1, 1]$,

$$\min\{\mathfrak{g}_{\rm sh}(-1) + \frac{s+1}{\hat{k}}, \,\mathfrak{g}_{\rm sh}(1) + \frac{1-s}{\hat{k}}, \,\frac{1}{\hat{k}}\} \le \mathfrak{g}_{\rm sh}(s) \\ \le \min\{\mathfrak{g}_{\rm sh}(-1) + \hat{k}(s+1), \,\mathfrak{g}_{\rm sh}(1) + \hat{k}(1-s), \, f_{\theta}(s) - \frac{1}{\hat{k}}\}.$$

Remark 4.1. By Propositions 3.12–3.13, for each $\alpha \in (0,1)$, there exist $\hat{\epsilon}_3 > 0$ and $C_{\alpha} > 0$ depending only on (γ, v_2, α) such that, for any admissible solution corresponding to $\boldsymbol{\theta} \in \Theta$,

 $\|\mathfrak{g}_{\rm sh} - \mathfrak{g}_j\|_{2,\alpha,I_j}^{(\rm par)} \le C_\alpha \qquad whenever \ \theta_{j-4} \le \theta^{\rm s} \,,$

for j = 5, 6, where $I_j := \{s \in (-1, 1) : 1 + (-1)^j s < \hat{\epsilon}_3\}$, and $\|\cdot\|_{2,\alpha,I_j}^{(\text{par})}$ is given by Definition 3.11(ii) with replacement of x by 1 - |s|.

By Proposition 3.14, for each $\alpha \in (0,1)$, there exist constants $\hat{\epsilon}_4 > 0$ and $C'_{\alpha} > 0$ depending only on (γ, v_2, α) such that, for any admissible solution corresponding to $\boldsymbol{\theta} \in \Theta$,

$$\left\|\mathfrak{g}_{\rm sh} - \mathfrak{g}_j\right\|_{2,\alpha,\overline{I'_j}} \le C'_{\alpha}, \qquad \frac{\mathrm{d}^m}{\mathrm{d}s^m}(\mathfrak{g}_{\rm sh} - \mathfrak{g}_j)((-1)^{j-1}) = 0 \quad for \ m = 0, 1, 2, j \le 1, 2, j \le 1, 2, \dots, 2,$$

whenever $\theta_{j-4} \in [\theta^{s}, \theta^{s} + \sigma_{3}]$ for j = 5, 6, where $I'_{j} \coloneqq \{s \in (-1, 1) : 1 + (-1)^{j} s < \hat{\epsilon}_{4}\}.$

Construction of $u^{(\varphi,\theta)}$. For each $\theta \in \overline{\Theta} \setminus \{\mathbf{0}\}$, it can be verified directly that S_{25} and S_{26} given by Definition 2.7 intersect at the unique point $P_I := (\xi^{P_I}, \eta^{P_I})$ with

(4.9)
$$\xi^{P_I} \coloneqq \frac{u_5 \xi^{P_0^1} - u_6 \xi^{P_0^2}}{u_5 - u_6}, \qquad \eta^{P_I} \coloneqq \frac{u_5 u_6}{u_5 - u_6} \frac{\xi^{P_0^1} - \xi^{P_0^2}}{v_2}$$

Similarly, $S_{25}^{\delta_0}$ and $S_{26}^{\delta_0}$ intersect at the unique point $P_I^{\delta_0} \coloneqq (\xi^{P_I}, \eta^{P_I^{\delta_0}})$ with $\eta^{P_I^{\delta_0}} \coloneqq \eta^{P_I} - \frac{\delta_0}{v_2} > \eta^{P_I}$. In Lemma A.3, we show the limits:

$$\lim_{\boldsymbol{\theta}\to\mathbf{0},\,\boldsymbol{\theta}\in\overline{\Theta}\setminus\{\mathbf{0}\}}(\xi^{P_I},\eta^{P_I})=(0,\eta_0)\,$$

where η_0 is given by (2.23). Then we define $P_I|_{\boldsymbol{\theta}=\mathbf{0}} \coloneqq (0,\eta_0)$ such that P_I is $C(\overline{\Theta})$. By (4.9) and Lemma A.2(ii), $\xi^{P_I} \ge 0$ for any $\boldsymbol{\theta} \in \overline{\Theta} \cap \{\theta_2 = 0\}$. Therefore, there exists a constant $\sigma_{\text{sep}} > 0$ small enough, depending only on (γ, v_2) , such that

$$\xi^{P_3} \leq -d_{\operatorname{sep}} < -\frac{d_{\operatorname{sep}}}{2} \leq \xi^{P_I} \quad \text{for any } \boldsymbol{\theta} \in \overline{\Theta} \cap \{0 \leq \theta_2 \leq \sigma_{\operatorname{sep}}\}$$

where $d_{sep} > 0$ is given by (3.10). Also, by Lemma A.2(ii), there exists a small constant $\delta_{sep} > 0$ depending only on $(\gamma, v_2, \sigma_{sep})$ such that

$$\eta^{P_3} \le \eta_0 - \delta_{\text{sep}} < \eta_0 \le \min\{a_{25}, a_{26}\} \le \eta^{P_I} \quad \text{for any } \boldsymbol{\theta} \in \overline{\Theta} \cap \{\sigma_{\text{sep}} \le \theta_2 \le \theta^d\}.$$

From the above two inequalities, using $P_3, P_I \in S_{26}$ and Lemma 2.8, we obtain the uniform positive lower bound:

$$\inf_{\boldsymbol{\theta}\in\overline{\Theta}}(\xi^{P_I}-\xi^{P_3})>0\,.$$

Definition 4.4. Let $\tilde{\zeta}(r; r_0, r_d)$ be given by (4.5), and let ξ^{P_I} be given by (4.9). For $\theta \in \overline{\Theta}$, define

$$\chi_{\boldsymbol{\theta}}^*(\xi) \coloneqq \tilde{\zeta}(-\xi; -\xi^{P_I}, \frac{\xi^{P_I} - \xi^{P_3}}{5C_{\text{bg}}})$$

for some constant $C_{\text{bg}} > 1$. The smooth background function $\varphi_{\theta}^{*}(\boldsymbol{\xi})$ is defined by

$$\varphi_{\boldsymbol{\theta}}^{*}(\boldsymbol{\xi}) \coloneqq \varphi_{5}(\boldsymbol{\xi}) \left(1 - \chi_{\boldsymbol{\theta}}^{*}(\boldsymbol{\xi})\right) + \varphi_{6}(\boldsymbol{\xi}) \chi_{\boldsymbol{\theta}}^{*}(\boldsymbol{\xi})$$

For constant $\delta_0 > 0$ from Definition 4.1, constant C_{bg} is fixed sufficiently large, depending only on (γ, v_2) , such that $\max\{\varphi_5, \varphi_6\} - \varphi_{\theta}^* \leq \frac{\delta_0}{2}$.

From the construction above, it can be checked that $\varphi_{\theta}^* \leq \max\{\varphi_5, \varphi_6\}$ in \mathbb{R}^2 and there exists a constant $\bar{k} > 1$ sufficiently large, depending only on (γ, v_2) , such that, for j = 5, 6,

(4.10)
$$\varphi_{\boldsymbol{\theta}}^* = \varphi_{\boldsymbol{\theta}} = \varphi_j \qquad \text{in } \mathcal{D}_{\hat{c}_j/\bar{k}}^j.$$

where \hat{c}_i is given by (3.14). Then

(4.11)
$$\left\{\boldsymbol{\xi} \in \mathbb{R}^2_+ : \boldsymbol{\xi}^{P_3} < \boldsymbol{\xi} < \boldsymbol{\xi}^{P_2}, \, (\varphi_2 - \varphi_{\boldsymbol{\theta}}^*)(\boldsymbol{\xi}) = 0\right\} \subseteq Q^{\boldsymbol{\theta}}.$$

Using Lemma 4.2(ii) and Definition 4.4, we can directly obtain that, for any $\theta \in \overline{\Theta}$, there exists a constant $\overline{\mu} > 0$ depending only on (γ, v_2) such that

(4.12)
$$\partial_{t'} ((\varphi_2 - \varphi_{\boldsymbol{\theta}}^*) \circ (\mathcal{G}_1^{\boldsymbol{\theta}})^{-1})(s, t') \leq -\bar{\mu} \quad \text{for any } (s, t') \in \overline{\mathcal{G}_1^{\boldsymbol{\theta}}(Q^{\boldsymbol{\theta}})}.$$

For any admissible solution φ corresponding to parameters $\boldsymbol{\theta} \in \Theta$, let $\mathfrak{g}_{sh} : [-1,1] \to [0,\infty)$ be the unique function given by Proposition 4.3. Then $G_{2,\mathfrak{g}_{sh}} : \mathcal{G}_1^{\boldsymbol{\theta}}(Q^{\boldsymbol{\theta}}) \to \mathbb{R}^2$ is defined by

(4.13)
$$(s,t) = G_{2,\mathfrak{g}_{\mathrm{sh}}}(s,t') \coloneqq (s,\frac{t'}{\mathfrak{g}_{\mathrm{sh}}(s)}) \quad \text{for any } (s,t') \in \mathcal{G}_1^{\boldsymbol{\theta}}(Q^{\boldsymbol{\theta}})$$

By Proposition 4.3(iv), map $G_{2,\mathfrak{g}_{sh}}$ is well defined and invertible with $G_{2,\mathfrak{g}_{sh}}^{-1}(s,t) = (s,t\mathfrak{g}_{sh}(s))$. Then $G_{2,\mathfrak{g}_{sh}} \circ \mathcal{G}_{1}^{\boldsymbol{\theta}}(\Omega) = \mathcal{Q}^{\text{iter}} = (-1,1) \times (0,1)$ by Proposition 4.3(i). Therefore, $u^{(\varphi,\boldsymbol{\theta})}$, defined as

(4.14)
$$u^{(\varphi,\theta)}(s,t) \coloneqq (\varphi - \varphi_{\theta}^*) \circ (G_{2,\mathfrak{g}_{sh}} \circ \mathcal{G}_1^{\theta})^{-1}(s,t) +$$

is well defined for all $(s,t) \in \mathcal{Q}^{\text{iter}}$.

A priori estimates for $u^{(\varphi,\theta)}$. For $\epsilon_0^* > 0$ from Proposition 4.3(ii), define a constant $\epsilon_0' > 0$ depending only on (γ, v_2) as

(4.15)
$$\epsilon'_0 \coloneqq \min_{\boldsymbol{\theta} \in \overline{\Theta}} d_L \epsilon^*_0 \leq \hat{\epsilon}^*_0,$$

where $d_L > 0$ is given by (4.6). For $r \in (0, 1)$ and for j = 5, 6, we define

$$(4.16) \qquad \mathcal{Q}_r^{\text{int}} \coloneqq \left\{ (s,t) \in \mathcal{Q}^{\text{iter}} : |s| < 1-r \right\}, \qquad \mathcal{Q}_r^j \coloneqq \left\{ (s,t) \in \mathcal{Q}^{\text{iter}} : 1+(-1)^j \, s < r \right\}.$$

Definition 4.5 (Weighted Hölder norms in the standard domain). For $\alpha \in (0,1)$ and any $u \in C^2(\mathcal{Q}^{\text{iter}})$, introduce the norm:

$$\|u\|_{2,\alpha,\mathcal{Q}^{\text{iter}}}^{(*)} \coloneqq \|u\|_{2,\alpha,\overline{\mathcal{Q}_{\epsilon_0^{j/4}}^{\text{int}}}} + \sum_{j=5,6} \left(\|u\|_{2,\alpha,\mathcal{Q}_{\epsilon_0^{j}}^{j}}^{(1+\alpha),(\text{par})} + \|u\|_{1,\alpha,\mathcal{Q}_{\epsilon_0^{j}}^{j}}^{(1+\alpha),(\text{subs})} \right).$$

where $\|\cdot\|_{2,\alpha,U}$ is the standard Hölder norm, and the weighted Hölder norms $\|\cdot\|_{m,\alpha,U}^{(\sigma),(\text{subs})}$ and $\|\cdot\|_{m,\alpha,U}^{(\sigma),(\text{par})}$, for an open set $U \subseteq \mathcal{Q}^{\text{iter}}$, $\sigma > 0$, and $m \in \mathbb{N}$, are given as follows:

(i) For any $\mathbf{s} = (s, t), \ \tilde{\mathbf{s}} = (\tilde{s}, \tilde{t}) \in \mathcal{Q}^{\text{iter}}, \ define$

$$\delta_{\alpha}^{(\text{subs})}(\boldsymbol{s}, \tilde{\boldsymbol{s}}) \coloneqq \left((s - \tilde{s})^2 + (\max\{1 - |\boldsymbol{s}|, 1 - |\tilde{s}|\})^2 (t - \tilde{t})^2 \right)^{\frac{\alpha}{2}}$$

The Hölder norms with subsonic scaling are given by

$$\begin{split} \|u\|_{m,0,U}^{(\sigma),(\mathrm{subs})} &\coloneqq \sum_{0 \le k+l \le m} \sup_{\boldsymbol{s} \in U} \left((1-|\boldsymbol{s}|)^{k-\sigma} |\partial_{\boldsymbol{s}}^{k} \partial_{t}^{l} u(\boldsymbol{s})| \right), \\ [u]_{m,\alpha,U}^{(\sigma),(\mathrm{subs})} &\coloneqq \sum_{k+l=m} \sup_{\substack{\boldsymbol{s}, \tilde{\boldsymbol{s}} \in U\\ \boldsymbol{s} \neq \tilde{\boldsymbol{s}}}} \left(\min\left\{ (1-|\boldsymbol{s}|)^{\alpha+k-\sigma}, (1-|\tilde{\boldsymbol{s}}|)^{\alpha+k-\sigma} \right\} \frac{|\partial_{\boldsymbol{s}}^{k} \partial_{t}^{l} u(\boldsymbol{s}) - \partial_{\boldsymbol{s}}^{k} \partial_{t}^{l} u(\tilde{\boldsymbol{s}})|}{\delta_{\alpha}^{(\mathrm{subs})}(\boldsymbol{s}, \tilde{\boldsymbol{s}})} \right), \\ \|u\|_{m,\alpha,U}^{(\sigma),(\mathrm{subs})} &= \|u\|_{m,0,U}^{(\sigma),(\mathrm{subs})} + [u]_{m,\alpha,U}^{(\sigma),(\mathrm{subs})}. \end{split}$$

(ii) For any $\mathbf{s} = (s, t), \tilde{\mathbf{s}} = (\tilde{s}, \tilde{t}) \in \mathcal{Q}^{\text{iter}}, define$

$$\delta_{\alpha}^{(\text{par})}(\boldsymbol{s}, \tilde{\boldsymbol{s}}) \coloneqq \left((s - \tilde{s})^2 + \max\{1 - |s|, 1 - |\tilde{s}|\}(t - \tilde{t})^2 \right)^{\frac{1}{2}}.$$

The Hölder norms with parabolic scaling are given by

$$\begin{split} \|u\|_{m,0,U}^{(\sigma),(\text{par})} &\coloneqq \sum_{0 \le k+l \le m} \sup_{s \in U} \left((1-|s|)^{k+\frac{l}{2}-\sigma} |\partial_s^k \partial_t^l u(s)| \right), \\ [u]_{m,\alpha,U}^{(\sigma),(\text{par})} &\coloneqq \sum_{k+l=m} \sup_{\substack{s, \tilde{s} \in U \\ s \ne \tilde{s}}} \left(\min\left\{ (1-|s|)^{\alpha+k+\frac{l}{2}-\sigma}, (1-|\tilde{s}|)^{\alpha+k+\frac{l}{2}-\sigma} \right\} \frac{|\partial_s^k \partial_t^l u(s) - \partial_s^k \partial_t^l u(\tilde{s})|}{\delta_{\alpha}^{(\text{par})}(s, \tilde{s})} \right), \\ \|u\|_{m,\alpha,U}^{(\sigma),(\text{par})} &= \|u\|_{m,0,U}^{(\sigma),(\text{par})} + [u]_{m,\alpha,U}^{(\sigma),(\text{par})}. \end{split}$$

Denote by $C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}})$ the set of all functions in $C^2(\mathcal{Q}^{\text{iter}})$ whose $\|\cdot\|^{(*)}_{2,\alpha,\mathcal{Q}^{\text{iter}}}$ -norms are finite.

Note that $C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}})$ is compactly embedded into $C^{2,\tilde{\alpha}}_{(*)}(\mathcal{Q}^{\text{iter}})$ whenever $0 \leq \tilde{\alpha} < \alpha < 1$, which follows from the properties given in [10, Lemma 4.6.3 and Corollary 17.2.7].

Using Corollary 3.10 and Propositions 3.12–3.14 and 3.16, we have the following estimate for admissible solutions φ after mapping the pseudo-subsonic domain Ω to the standard domain Q^{iter} ; the proof is similar to [2, Proposition 4.12].

Proposition 4.6. For each $\theta_* \in (0, \theta^d)$, there exist constants M > 0 and $\bar{\alpha} \in (0, \frac{1}{3}]$ depending only on (γ, v_2, θ_*) such that, for any admissible solution φ corresponding to $\theta \in \Theta \cap \{\theta_1, \theta_2 \leq \theta_*\},\$ $u = u^{(\varphi, \theta)} : \mathcal{Q}^{\text{iter}} \to \mathbb{R}, \text{ defined by (4.14), satisfies } u \in C_{(*)}^{2,\bar{\alpha}}(\mathcal{Q}^{\text{iter}}) \text{ and}$

$$\|u\|_{2,\bar{\alpha},\mathcal{Q}^{\mathrm{iter}}}^{(*)} = \|u\|_{2,\bar{\alpha},\mathcal{Q}^{\mathrm{int}}_{\epsilon'_0/4}} + \sum_{j=5,6} \left(\|u\|_{2,\bar{\alpha},\mathcal{Q}^{j}_{\epsilon'_0}}^{(1+\bar{\alpha}),(\mathrm{par})} + \|u\|_{1,\bar{\alpha},\mathcal{Q}^{j}_{\epsilon'_0}}^{(1+\bar{\alpha}),(\mathrm{subs})} \right) \le M$$

4.1.2. Mapping to approximate admissible solutions. For each $\theta \in \overline{\Theta}$, let the extended domain Q^{θ} be given by Definition 4.1, and let φ_{θ}^* be given by Definition 4.4. For each $s^* \in (-1, 1)$, define $Q^{\boldsymbol{\theta}}(s^*) \coloneqq Q^{\boldsymbol{\theta}} \cap (\mathcal{G}_1^{\boldsymbol{\theta}})^{-1}(\{s = s^*\}).$ Then, for i = 1, 2,

$$\inf_{Q^{\boldsymbol{\theta}}((-1)^{i-1})} (\varphi_2 - \varphi_{\boldsymbol{\theta}}^*) < 0 \le \sup_{Q^{\boldsymbol{\theta}}((-1)^{i-1})} (\varphi_2 - \varphi_{\boldsymbol{\theta}}^*)$$

where the inequality on the right above becomes a strict inequality when $\theta_i \in [0, \theta^s)$ and an equality when $\theta_i \in [\theta^s, \theta^d]$. Moreover, by the choice of $C_{bg} > 0$ in Definition 4.4,

$$\sup_{Q^{\boldsymbol{\theta}}(s)} (\varphi_2 - \varphi_{\boldsymbol{\theta}}^*) - \inf_{Q^{\boldsymbol{\theta}}(s)} (\varphi_2 - \varphi_{\boldsymbol{\theta}}^*) \ge \frac{\delta_0}{2} > 0 \quad \text{for any } s \in (-1, 1).$$

Definition 4.7. Fix $\alpha \in (0,1)$, $\theta_* \in (0,\theta^d)$, and $\theta \in \Theta \cap \{\theta_1, \theta_2 \leq \theta_*\}$. Let $u \in C^{1,\alpha}(\overline{\mathcal{Q}^{\text{iter}}})$ be a function satisfying that

(4.17)
$$\inf_{Q^{\boldsymbol{\theta}}(s)} (\varphi_2 - \varphi_{\boldsymbol{\theta}}^*) < u(s, 1) < \sup_{Q^{\boldsymbol{\theta}}(s)} (\varphi_2 - \varphi_{\boldsymbol{\theta}}^*) \quad \text{for any } s \in (-1, 1).$$

- (i) For each $s \in (-1,1)$, let $\bar{t}' > 0$ be the unique solution of $(\varphi_2 \varphi_{\theta}^*) \circ (\mathcal{G}_1^{\theta})^{-1}(s,\bar{t}') = u(s,1)$,
- which exists by (4.12) and (4.17). Define $\mathfrak{g}_{\mathrm{sh}}^{(u,\theta)}: (-1,1) \to (0,\infty)$ by $\mathfrak{g}_{\mathrm{sh}}^{(u,\theta)}(s) \coloneqq \overline{t'}$. (ii) For \mathcal{G}_1^{θ} and $G_{2,\mathfrak{g}_{\mathrm{sh}}^{(u,\theta)}}$ given by (4.7) and (4.13) respectively, define $\mathfrak{F}_{(u,\theta)}: \mathcal{Q}^{\mathrm{iter}} \to \mathcal{Q}^{\theta}$ by $\mathfrak{F}_{(u,\theta)} \coloneqq (\mathcal{G}_1^{\theta})^{-1} \circ G_{2,\mathfrak{g}_{cb}^{(u,\theta)}}^{-1}.$
- (iii) Define sets $\Omega(u, \theta) \coloneqq \mathfrak{F}_{(u,\theta)}(\mathcal{Q}^{\text{iter}})$ and $\Gamma_{\text{shock}}(u, \theta) \coloneqq \mathfrak{F}_{(u,\theta)}((-1,1) \times \{1\})$. Finally, define function $\varphi^{(u,\theta)}(\boldsymbol{\xi}) : \Omega(u, \theta) \to \mathbb{R}$ by $\varphi^{(u,\theta)}(\boldsymbol{\xi}) \coloneqq (u \circ \mathfrak{F}_{(u,\theta)}^{-1})(\boldsymbol{\xi}) + \varphi_{\boldsymbol{\theta}}^{*}(\boldsymbol{\xi})$ for all $\boldsymbol{\xi} \in \Omega(u, \theta)$.

We call $\varphi^{(u,\theta)}$ above an approximate admissible solution if it satisfies the same boundary conditions on $\Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6$ as the admissible solutions in Definition 2.11. For $\alpha \in (0,1)$ and $\theta_* \in (0, \theta^d)$, define the space of such approximate admissible solutions:

(4.18)
$$\mathfrak{G}^{\theta_*}_{\alpha} \coloneqq \left\{ (u, \boldsymbol{\theta}) \in C^{1, \alpha}(\overline{\mathcal{Q}^{\text{iter}}}) \times [0, \theta_*]^2 : \begin{array}{c} (u, \boldsymbol{\theta}) \text{ satisfies (4.17) and} \\ (u, Du)(\pm 1, \cdot) = (0, \mathbf{0}) \end{array} \right\}.$$

(

By construction, for any $(u, \theta) \in \mathfrak{G}_{\alpha}^{\theta_*}$, map $\mathfrak{F}_{(u,\theta)}$ satisfies that $P_1 = \mathfrak{F}_{(u,\theta)}(1,0)$, $P_2 = \mathfrak{F}_{(u,\theta)}(1,1)$, $P_3 = \mathfrak{F}_{(u,\theta)}(-1,1)$, and $P_4 = \mathfrak{F}_{(u,\theta)}(-1,0)$ for points P_i , i = 1, 2, 3, 4, given by Definition 2.9. Furthermore, $\varphi^{(u,\theta)} = \varphi_2$ on $\Gamma_{\text{shock}}(u,\theta)$. Other properties of set $\mathfrak{G}^{\theta_*}_{\alpha}$ are given below; the proof is similar to [10, Lemmas 12.2.7 and 17.2.13].

Lemma 4.8. Fix $\alpha \in (0,1)$ and $\theta_* \in (0,\theta^d)$. For each $(u, \theta) \in \mathfrak{G}^{\theta_*}_{\alpha}$, $\mathfrak{g}^{(u,\theta)}_{\mathrm{sh}} \in C^{1,\alpha}([-1,1])$ and the following properties hold:

(i) $\Omega(u, \theta) \cup \Gamma_{\text{shock}}(u, \theta) \subseteq Q^{\theta} \subseteq \mathbb{R}^2_+$. Moreover, $\Gamma_{\text{shock}}(u, \theta)$ is a $C^{1,\alpha}$ -curve up to its endpoints P_2 and P_3 , and is tangential to S_{25} and S_{26} at P_2 and P_3 respectively. If $f_{j,0}, j = 5, 6$, are given as in Step 1 of the proofs of Propositions 3.12 and 3.14, then

$$\left| \left(\mathfrak{g}_{sh}^{(u,\theta)}, \frac{\mathrm{d}}{\mathrm{d}s} \mathfrak{g}_{sh}^{(u,\theta)} \right) \right|_{s=(-1)^{j-1}} = \left(f_{j,0}(0), \frac{(-1)^j}{d_L} f_{j,0}'(0) \right) \qquad \text{for } j = 5, 6 \,,$$

where d_L is given by (4.6).

(ii) Let $\mathcal{N}_{\varepsilon}(\Gamma)$ denote the open ε -neighbourhood of a set Γ , as given by (3.21). Let δ_0 be from Definition 4.1, and let the coordinates $(x, y) = \mathcal{R}(\boldsymbol{\xi})$ be given by (4.1). Then there exists a constant $\epsilon_0 > 0$ depending only on (γ, v_2) such that, for j = 5, 6, and any $\epsilon \in (0, \epsilon_0)$,

$$\begin{split} \hat{\Omega}^{j}_{\epsilon} &\coloneqq \mathcal{N}_{\epsilon_{0}}(\Gamma^{j,\delta_{0}}_{\text{sonic}}) \cap \left\{ \boldsymbol{\xi} = \mathcal{R}^{-1}(x,y) \,:\, x_{P_{j-3}} < x < x_{P_{j-3}} + \epsilon \right\} \cap \Omega(u,\boldsymbol{\theta}) \\ &= \left\{ \boldsymbol{\xi} = \mathcal{R}^{-1}(x,y) \,:\, x_{P_{j-3}} < x < x_{P_{j-3}} + \epsilon, \, 0 < y < \hat{f}_{j,\text{sh}}(x) \right\}, \\ \Gamma_{\text{shock}}(u,\boldsymbol{\theta}) \cap \partial \hat{\Omega}^{j}_{\epsilon} = \left\{ \boldsymbol{\xi} = \mathcal{R}^{-1}(x,\hat{f}_{j,\text{sh}}(x)) \,:\, x_{P_{j-3}} < x < x_{P_{j-3}} + \epsilon \right\}, \\ \Gamma^{j}_{\text{sonic}} \cap \partial \hat{\Omega}^{j}_{\epsilon} = \left\{ \boldsymbol{\xi} = \mathcal{R}^{-1}(x_{P_{j-3}},y) \,:\, 0 < y < \hat{f}_{j,\text{sh}}(x_{P_{j-3}}) \right\} = \Gamma^{j}_{\text{sonic}}, \\ \Gamma_{\text{sym}} \cap \partial \hat{\Omega}^{j}_{\epsilon} = \left\{ \boldsymbol{\xi} = \mathcal{R}^{-1}(x,0) \,:\, x_{P_{j-3}} < x < x_{P_{j-3}} + \epsilon \right\}, \end{split}$$

where $\hat{f}_{j,\mathrm{sh}}(x) \coloneqq \mathfrak{g}_{\mathrm{sh}}^{(u,\theta)} \circ L_{\theta}(u_j + (-1)^{j-1}(c_j - x))$ and L_{θ} is given by (4.6). (iii) Let $(u,\theta), (\tilde{u},\tilde{\theta}) \in \mathfrak{G}_{\alpha}^{\theta_*}$ satisfy $||(u,\tilde{u})||_{C^{1,\alpha}(\overline{\mathcal{Q}^{\mathrm{iter}}})} < M$ for some constant M > 0. Then there

exists a constant C > 0 depending only on $(\gamma, v_2, \theta_*, M, \alpha)$ such that

$$\begin{split} \|\mathfrak{g}_{\mathrm{sh}}^{(u,\boldsymbol{\theta})}\|_{C^{1,\alpha}([-1,1])} + \|\mathfrak{F}_{(u,\boldsymbol{\theta})}\|_{C^{1,\alpha}(\overline{\mathcal{Q}^{\mathrm{iter}}})} &\leq C \,, \\ \|\mathfrak{g}_{\mathrm{sh}}^{(u,\boldsymbol{\theta})} - \mathfrak{g}_{\mathrm{sh}}^{(\tilde{u},\tilde{\boldsymbol{\theta}})}\|_{C^{1,\alpha}([-1,1])} + \|\mathfrak{F}_{(u,\boldsymbol{\theta})} - \mathfrak{F}_{(\tilde{u},\tilde{\boldsymbol{\theta}})}\|_{C^{1,\alpha}(\overline{\mathcal{Q}^{\mathrm{iter}}})} \\ &+ \|\varphi^{(u,\boldsymbol{\theta})} \circ \mathfrak{F}_{(u,\boldsymbol{\theta})} - \varphi^{(\tilde{u},\tilde{\boldsymbol{\theta}})} \circ \mathfrak{F}_{(\tilde{u},\tilde{\boldsymbol{\theta}})}\|_{C^{1,\alpha}(\overline{\mathcal{Q}^{\mathrm{iter}}})} + \|\varphi_{\boldsymbol{\theta}}^{*} \circ \mathfrak{F}_{(u,\boldsymbol{\theta})} - \varphi_{\tilde{\boldsymbol{\theta}}}^{*} \circ \mathfrak{F}_{(\tilde{u},\tilde{\boldsymbol{\theta}})}\|_{C^{1,\alpha}(\overline{\mathcal{Q}^{\mathrm{iter}}})} \\ &\leq C \big(\|u - \tilde{u}\|_{C^{1,\alpha}(\overline{\mathcal{Q}^{\mathrm{iter}}})} + |\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}}| \big) \,. \end{split}$$

(iv) Let $\epsilon_0 > 0$ be the constant from (ii), and let $\hat{\epsilon}_0 := d_L \epsilon_0$ with $d_L > 0$ given by (4.6). Assume that, for constants $\alpha \in (0, 1), \sigma \in (1, 2], and M > 0$,

(4.19)
$$\|u\|_{2,\alpha,\mathcal{Q}^{\text{iter}} \cap \{|s| < 1 - \frac{\hat{\epsilon}_0}{10}\}} + \|u\|_{2,\alpha,\mathcal{Q}^{\text{iter}} \cap \{|s| > 1 - \hat{\epsilon}_0\}}^{(\sigma),(\text{par})} \le M.$$

Then there exists a constant C > 0 depending only on $(\gamma, v_2, \theta_*, \alpha, \sigma)$ such that

$$\|\mathfrak{g}_{\rm sh}^{(u,\boldsymbol{\theta})}\|_{2,\alpha,[-1+\frac{\hat{\epsilon}_0}{10},1-\frac{\hat{\epsilon}_0}{10}]} + \|\mathfrak{g}_{\rm sh}^{(u,\boldsymbol{\theta})} - \mathfrak{g}_5\|_{2,\alpha,(1,1-\hat{\epsilon}_0)}^{(\sigma),({\rm par})} + \|\mathfrak{g}_{\rm sh}^{(u,\boldsymbol{\theta})} - \mathfrak{g}_6\|_{2,\alpha,(-1,-1+\hat{\epsilon}_0)}^{(\sigma),({\rm par})} \le CM\,,$$

where \mathfrak{g}_5 and \mathfrak{g}_6 are defined in Proposition 4.3(ii).

Furthermore, for $I_{so} \coloneqq (-1, -1 + \hat{\epsilon}_0) \cup (1 - \hat{\epsilon}_0, 1)$, define $\mathfrak{F}_{(0,\boldsymbol{\theta})} : I_{so} \times (0, \infty) \to \mathbb{R}^2$ by

$$\mathfrak{F}_{(0,\boldsymbol{\theta})}(s,t') \coloneqq \begin{cases} (G_{2,\mathfrak{g}_5} \circ \mathcal{G}_1^{\boldsymbol{\theta}})^{-1}(s,t') & \text{if } s \in I_{\mathrm{so}} \cap \{s > 0\}, \\ (G_{2,\mathfrak{g}_6} \circ \mathcal{G}_1^{\boldsymbol{\theta}})^{-1}(s,t') & \text{if } s \in I_{\mathrm{so}} \cap \{s < 0\}. \end{cases}$$

Then there exists a constant $C_0 > 0$ depending only on (γ, v_2, θ_*) such that

$$\|\mathfrak{F}_{(0,\boldsymbol{\theta})}\|_{C^{3}(\overline{I_{\mathrm{so}}}\times[0,1])} + \|\mathfrak{F}_{(u,\boldsymbol{\theta})}\|_{2,\alpha,\mathcal{Q}^{\mathrm{iter}}\cap\{|s|<1-\frac{\hat{\epsilon}_{0}}{10}\}} + \|\mathfrak{F}_{(u,\boldsymbol{\theta})} - \mathfrak{F}_{(0,\boldsymbol{\theta})}\|_{2,\alpha,I_{\mathrm{so}}\times(0,1)}^{(\sigma),(\mathrm{par})} \leq C_{0}.$$

(v) Let f_{θ} be given by (4.8). For constants M > 0 and $\delta_{sh} > 0$, assume that $(u, \theta) \in \mathfrak{G}_{\alpha}^{\theta_*}$ satisfies (4.19) and, for any $s \in [-1, 1]$,

$$\min\left\{\mathfrak{g}_{\mathrm{sh}}^{(u,\boldsymbol{\theta})}(-1) + \frac{s+1}{M}, \,\mathfrak{g}_{\mathrm{sh}}^{(u,\boldsymbol{\theta})}(1) + \frac{1-s}{M}, \,\delta_{\mathrm{sh}}\right\} \leq \mathfrak{g}_{\mathrm{sh}}^{(u,\boldsymbol{\theta})}(s)$$
$$\leq \min\left\{\mathfrak{g}_{\mathrm{sh}}^{(u,\boldsymbol{\theta})}(-1) + M(s+1), \,\mathfrak{g}_{\mathrm{sh}}^{(u,\boldsymbol{\theta})}(1) + M(1-s), f_{\boldsymbol{\theta}}(s) - \frac{1}{M}\right\}.$$

Then, for any $\epsilon \in (0, \frac{1}{4} \min \{s_5, -s_6\})$, there exists a constant $C_{\epsilon} > 0$ depending only on $(\gamma, v_2, \theta_*, \alpha, \delta_{\rm sh}, \epsilon, M)$ such that

$$\|(\mathfrak{F}_{(u,\theta)}^{-1},\varphi-\varphi_{\theta}^*)\|_{2,\alpha,\Omega(u,\theta)\setminus(\overline{\Omega_{\epsilon}^5\cup\widehat{\Omega_{\epsilon}^6}})} \leq C_{\epsilon}\,.$$

(vi) Let $(u, \theta), (\tilde{u}, \tilde{\theta}) \in \mathfrak{G}_{\alpha}^{\theta_*}$ satisfy $||(u, \tilde{u})||_{C^{1,\alpha}(\overline{\mathcal{Q}^{\text{iter}}})} < M$ for some constant M > 0. For any open set $K \Subset \mathcal{Q}^{\text{iter}}$ with $\delta = \text{dist}(K, \overline{\mathcal{Q}^{\text{iter}}} \cap \{|s| = 1\})$, there exists a constant $C_{\delta} > 0$ depending only on $(\gamma, v_2, \theta_*, \alpha, \sigma, \delta, M)$ such that

$$\begin{split} \|\mathfrak{F}_{(u,\boldsymbol{\theta})} - \mathfrak{F}_{(\tilde{u},\tilde{\boldsymbol{\theta}})}\|_{C^{2,\alpha}(\overline{K})} &\leq C_{\delta} \big(\|(u-\tilde{u})(\cdot,1)\|_{C^{2,\alpha}([-1+\delta,1-\delta])} + |\boldsymbol{\theta}-\boldsymbol{\theta}| \big) \,, \\ \|(\varphi^{(u,\boldsymbol{\theta})} \circ \mathfrak{F}_{(u,\boldsymbol{\theta})} - \varphi^{(\tilde{u},\tilde{\boldsymbol{\theta}})} \circ \mathfrak{F}_{(\tilde{u},\tilde{\boldsymbol{\theta}})}, \, \psi^{(u,\boldsymbol{\theta})} \circ \mathfrak{F}_{(u,\boldsymbol{\theta})} - \psi^{(\tilde{u},\tilde{\boldsymbol{\theta}})} \circ \mathfrak{F}_{(\tilde{u},\tilde{\boldsymbol{\theta}})} \|_{C^{2,\alpha}(\overline{K})} \\ &\leq C_{\delta} \big(\|u-\tilde{u}\|_{C^{2,\alpha}(\overline{K})} + |\boldsymbol{\theta}-\tilde{\boldsymbol{\theta}}| \big) \,, \end{split}$$

where $\psi^{(u,\theta)}$ is given by $\psi^{(u,\theta)} \coloneqq \varphi^{(u,\theta)} - \varphi^*_{\theta}$ for each $(u,\theta) \in \mathfrak{G}^{\theta_*}_{\alpha}$.

4.2. Definition and properties of the iteration set. For any $\theta_* \in (0, \theta^d)$, we now define the iteration set $\mathcal{K} \subseteq C^{1,\alpha}(\overline{\mathcal{Q}^{\text{iter}}}) \times [0, \theta_*]^2$ for some $\alpha \in (0, 1)$, as a set of functions satisfying the properties similar to the admissible solutions. In the definition of \mathcal{K} , we construct a nonlinear boundary value problem in a fixed domain and obtain the well-posedness and *a priori* estimates for the solutions.

4.2.1. *Definition of the iteration set.* Before giving the definition of the iteration set, we collect some useful estimates for admissible solutions.

Lemma 4.9. Fix $\theta_* \in (0, \theta^d)$. For any admissible solution φ corresponding to $\theta \in \overline{\Theta} \cap \{\theta_1, \theta_2 \leq \theta_*\}$, denote $\psi \coloneqq (\varphi - \max\{\varphi_5, \varphi_6\}) \circ \mathcal{R}^{-1}$. Then there exist constants $C_1, \overline{\epsilon} > 0$ depending only on (γ, v_2) , and $C_2^* > 0$ depending on (γ, v_2, θ_*) such that, for any $\epsilon \in (0, \overline{\epsilon}]$,

$$\begin{aligned} \|\varphi - \varphi_5\|_{C^{0,1}(\overline{\Omega})} + \|\varphi - \varphi_6\|_{C^{0,1}(\overline{\Omega})} &\leq C_1 \,, \\ |D_{(x,y)}\psi(x,y)| &\leq C_2^* x \qquad for \, (x,y) \in \mathcal{R}(\overline{\Omega} \cap (\mathcal{D}^5_{\epsilon} \cup \mathcal{D}^6_{\epsilon})) \,. \end{aligned}$$

Proof. The first result follows directly from (2.33), Proposition 2.6, and Lemma 3.3.

Without loss of generality, we assume that $\theta_* \in (\theta^s, \theta^d)$. Then, for sufficiently small $\sigma > 0$ and any $\theta \in [0, \theta^s + \sigma] \times [0, \theta_*]$,

$$|D_{(x,y)}\psi(x,y)| \le C_2 x$$
 for $(x,y) \in \mathcal{R}(\overline{\Omega} \cap \mathcal{D}^5_{\epsilon})$.

whenever $\epsilon \in (0, \bar{\epsilon}]$, where $\bar{\epsilon} > 0$ is the minimum of the constants, $\varepsilon > 0$, in Propositions 3.12–3.14.

From the relations: $\ell(\rho_5) \cos \theta_{25} = v_2$ and $M_2^{(\theta_1)} = |(-\xi^{P_0^1}, v_2)|$, and Lemmas A.2(i) and 2.3,

(4.20)
$$\frac{\mathrm{d}x_{P_0^1}}{\mathrm{d}\theta_1} = \frac{\mathrm{d}}{\mathrm{d}\theta_1} \left(c_5 - (\xi^{P_0^1} - u_5) \right) = \frac{\mathrm{d}}{\mathrm{d}\theta_1} \left(c_5 - \xi^{P_0^1} + v_2 \tan \theta_{25} \right) > 0.$$

Then there exists $C'_2 > 0$ depending only on (γ, v_2, θ_*) such that, for any $\boldsymbol{\theta} \in [\theta^s + \frac{\sigma}{2}, \theta_*] \times [0, \theta_*]$,

$$|D_{(x,y)}\psi(x,y)| \le C_2' x^{\bar{\alpha}} \le C_2' \left(x_{P_0^1}|_{\theta_1 = \theta^s + \frac{\sigma}{2}} \right)^{\bar{\alpha} - 1} x \qquad \text{for } (x,y) \in \mathcal{R}\left(\overline{\Omega} \cap \mathcal{D}^5_\epsilon\right),$$

after possibly reducing $\bar{\epsilon}$ further so that $\bar{\epsilon} < r$, for r > 0 from Proposition 3.16 and $\bar{\alpha} \in (0, \frac{1}{3}]$ from Proposition 4.6. From the symmetry of the four-shock interaction problem, we can repeat the above discussion for $\psi(x, y)$ in $\mathcal{R}(\overline{\Omega} \cap \mathcal{D}_{\epsilon}^{6})$.

Define $u^{(\text{norm})} \in C^3(\overline{\mathcal{Q}^{\text{iter}}})$ by (4.14) with $\boldsymbol{\theta} = \mathbf{0}$ and $\varphi = \varphi_0$ from §2.1.3. Note that $\varphi_{\boldsymbol{\theta}}^* \equiv \varphi_0$ in $Q^{\boldsymbol{\theta}}$ by Definition 4.4 because $\varphi_5 = \varphi_6 = \varphi_0$ when $\boldsymbol{\theta} = \mathbf{0}$, which yields that

$$u^{(\text{norm})} \equiv 0 \quad \text{in } \mathcal{Q}^{\text{iter}}.$$

Let $\bar{\alpha} \in (0, \frac{1}{3}]$ be the constant from Proposition 4.6, $\epsilon_0 > 0$ from Lemma 4.8(ii), and $\bar{\epsilon} > 0$ from Lemma 4.9. For constants $\alpha \in (0, \frac{\bar{\alpha}}{2}]$, δ_1 , δ_2 , δ_3 , $\epsilon \in (0, \frac{1}{2}\min\{\epsilon_0, \bar{\epsilon}\})$, and $N_1 > 1$ to be specified later, we now define the iteration set $\mathcal{K} \subseteq C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2$, where space $C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}})$ is given by Definition 4.5. For simplicity, in the following, we use $(\mathfrak{g}_{\text{sh}}, \varphi, \Omega, \Gamma_{\text{shock}})$ to denote $(\mathfrak{g}_{\text{sh}}^{(u,\theta)}, \varphi^{(u,\theta)}, \Omega(u,\theta), \Gamma_{\text{shock}}(u,\theta))$ that are given by Definition 4.7.

Definition 4.10 (Iteration set). Fix $\theta_* \in (0, \theta^d)$. The iteration set $\mathcal{K} \subseteq C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2$ is the set of all (u, θ) satisfying the following properties:

(i)
$$\|u - u^{(\text{norm})}\|_{2,\alpha,\mathcal{Q}^{\text{iter}}}^{(*)} < \mathscr{K}_1(\max\{\theta_1,\theta_2\}) \text{ for } \mathscr{K}_1 \in C^{0,1}(\mathbb{R}) \text{ given by}$$

 $\begin{cases} \delta_1 & \text{if } \theta \leq \frac{\delta_1}{N_1}, \end{cases}$

$$\mathscr{K}_{1}(\theta) = \begin{cases} \sigma_{1} & \sigma_{1} & \sigma_{2} & \sigma_{1} \\ N_{0} & \text{if } \theta \geq \frac{2\delta_{1}}{N_{1}}, \\ \text{linear} & \text{otherwise}, \end{cases}$$

with $N_0 \coloneqq \max\{10M, 1\}$ for $M = M(\gamma, v_2, \theta_*) > 0$ from Proposition 4.6.

- (ii) $(u, \boldsymbol{\theta}) \in \mathfrak{G}_{\alpha}^{\theta_*}$, where $\mathfrak{G}_{\alpha}^{\theta_*}$ is defined by (4.18).
- (iii) Γ_{shock} satisfies dist $(\Gamma_{\text{shock}}, B_1(O_2)) > N_2^{-1}$, and \mathfrak{g}_{sh} satisfies $\mathfrak{g}_{\text{sh}}(\pm 1) \ge 0$ and

$$\min_{i=1,2} \left\{ \mathfrak{g}_{\rm sh}((-1)^{i-1}) + \frac{1 + (-1)^i s}{N_3}, \frac{1}{N_3} \right\} \le \mathfrak{g}_{\rm sh}(s) \\
\le \min_{i=1,2} \left\{ \mathfrak{g}_{\rm sh}((-1)^{i-1}) + N_3(1 + (-1)^i s), f_{\theta}(s) - \frac{1}{N_3} \right\} \quad \text{for all } s \in (-1,1),$$

with $N_2 \coloneqq 2C_{\rm sh}$ for $C_{\rm sh} = C_{\rm sh}(\gamma, v_2) > 0$ from Proposition 3.4, $N_3 \coloneqq 2\hat{k}$ for $\hat{k} = \hat{k}(\gamma, v_2, \theta_*) > 0$ from Proposition 4.3(iv), and f_{θ} given by (4.8).

(iv) Let coordinates $(x, y) = \mathcal{R}(\boldsymbol{\xi})$ be given by (4.1) and, for r > 0, let \mathcal{D}_r^5 and \mathcal{D}_r^6 be given by (4.2). Then φ and $\psi \coloneqq (\varphi - \max{\{\varphi_5, \varphi_6\}}) \circ \mathcal{R}^{-1}$ satisfy

(4.21)
$$\begin{aligned} \varphi - \max\left\{\varphi_{5}, \varphi_{6}\right\} &> \mathscr{K}_{2}(\max\{\theta_{1}, \theta_{2}\}) \qquad \text{ in } \overline{\Omega} \setminus \left(\mathcal{D}_{\epsilon/10}^{5} \cup \mathcal{D}_{\epsilon/10}^{6}\right), \\ \partial_{\boldsymbol{e}_{S_{2j}}}(\varphi_{2} - \varphi) &< -\mathscr{K}_{2}(\max\{\theta_{1}, \theta_{2}\}) \qquad \text{ in } \overline{\Omega} \setminus \mathcal{D}_{\epsilon/10}^{j} \text{ for } j = 5, 6, \end{aligned}$$

and

$$\begin{aligned} |\partial_x \psi(x,y)| &< \mathscr{K}_3(\theta_{j-4})x & in \ \mathcal{R}\big(\overline{\Omega} \cap \big(\mathcal{D}^j_{\epsilon_0} \setminus \mathcal{D}^j_{\epsilon/10}\big)\big) \ for \ j = 5, 6, \\ |\partial_y \psi(x,y)| &< N_4 x & in \ \mathcal{R}\big(\overline{\Omega} \cap \big(\big(\mathcal{D}^5_{\epsilon_0} \setminus \mathcal{D}^5_{\epsilon/10}\big) \cup \big(\mathcal{D}^6_{\epsilon_0} \setminus \mathcal{D}^6_{\epsilon/10}\big)\big)\big), \\ (4.22) & |D_{(x,y)}\psi| &< N_4 \epsilon & in \ \mathcal{R}\big(\overline{\Omega} \cap \big(\overline{\mathcal{D}^5_{\epsilon}} \cup \overline{\mathcal{D}^6_{\epsilon}}\big)\big), \\ \min \left\{\partial_{\boldsymbol{\nu}}(\varphi_2 - \varphi), \ \partial_{\boldsymbol{\nu}}\varphi\right\} > \mu_1 & on \ \overline{\Gamma_{\text{shock}}}, \\ \|\varphi - \varphi_j\|_{C^{0,1}(\overline{\Omega})} < N_5 & for \ j = 5, 6, \end{aligned}$$

for $\mathbf{e}_{S_{25}}$ and $\mathbf{e}_{S_{26}}$ given by (2.35), and the unit normal vector $\boldsymbol{\nu}$ on Γ_{shock} towards the interior of Ω . In the above conditions, functions $\mathscr{K}_2, \mathscr{K}_3 \in C^{0,1}(\mathbb{R})$ are defined by

$$\mathscr{K}_{2}(\theta) \coloneqq \delta_{2} \min\left\{\theta - \frac{\delta_{1}}{N_{1}^{2}}, \frac{\delta_{1}}{N_{1}^{2}}\right\}, \qquad \mathscr{K}_{3}(\theta) \coloneqq \begin{cases} \frac{2-\mu_{0}}{1+\gamma} & \text{if } 0 \leq \theta \leq \theta^{s} + \frac{\sigma_{2}}{2} \\ N_{4} & \text{if } \theta \geq \theta^{s} + \sigma_{2}, \\ linear & otherwise, \end{cases}$$

for constants μ_1 , σ_2 , μ_0 , ϵ_0 , N_4 , and N_5 chosen as follows:

- Let $\delta' = \delta'(\gamma, v_2) > 0$ be from Lemma 3.9. Choose $\mu_1 \coloneqq \frac{\delta'}{2}$.
- Let $\sigma_2 = \sigma_2(\gamma, v_2) > 0$ and $\delta = \delta(\gamma, v_2) > 0$ be from Step 1 of the proof of Proposition 3.14. Choose $\mu_0 := \frac{\delta}{2}$.
- Let $\epsilon_0 = \epsilon_0(\gamma, v_2) > 0$ be from Lemma 4.8(ii).
- Let $C_1 = C_1(\gamma, v_2) > 0$ and $C_2^* = C_2^*(\gamma, v_2, \theta_*) > 0$ be from Lemma 4.9. Choose $N_4 := 10C_2^*$ and $N_5 := 10C_1$.
- (v) The density function $\rho(|D\varphi|, \varphi)$ defined by (3.8) satisfies

$$\frac{\rho^*(\gamma)}{2} < \rho(|D\varphi|,\varphi) < 2C_{\rm ub} \qquad in \ \overline{\Omega} \setminus (\mathcal{D}^5_{\epsilon/10} \cup \mathcal{D}^6_{\epsilon/10})$$

for constants $\rho^*(\gamma) > 0$ and $C_{ub} = C_{ub}(\gamma, v_2) > 0$ given by (3.6) in Lemma 3.3.

(vi) Let the sound speed $c(|D\varphi|, \varphi)$ as defined in (3.2). Function φ satisfies

$$\frac{|D\varphi(\boldsymbol{\xi})|^2}{{}^2(|D\varphi(\boldsymbol{\xi})|,\varphi(\boldsymbol{\xi}))} < 1 - \tilde{\mu}\operatorname{dist}^{\flat}(\boldsymbol{\xi},\Gamma_{\operatorname{sonic}}^5 \cup \Gamma_{\operatorname{sonic}}^6) \qquad for \ \boldsymbol{\xi} \in \overline{\Omega} \setminus \left(\mathcal{D}_{\epsilon/10}^5 \cup \mathcal{D}_{\epsilon/10}^6\right),$$

where $\tilde{\mu} \coloneqq \frac{\mu_{\text{el}}}{2}$ for $\mu_{\text{el}} = \frac{\mu}{C_{\flat}} > 0$ with $C_{\flat} = C_{\flat}(\gamma, v_2) > 0$ from (3.17), $\mu = \mu(\gamma, v_2) > 0$ is from Proposition 3.8, and $\text{dist}^{\flat}(\cdot, \cdot)$ is defined in (3.18).

(vii) The iteration boundary value problem

 $\begin{cases} \mathcal{N}_{(u,\theta)}(\hat{\phi}) = A_{11}\hat{\phi}_{\xi\xi} + 2A_{12}\hat{\phi}_{\xi\eta} + A_{22}\hat{\phi}_{\eta\eta} = 0 & \text{in } \Omega ,\\ \mathcal{M}_{(u,\theta)}(D\hat{\phi}, \hat{\phi}, \boldsymbol{\xi}) = 0 & \text{on } \Gamma_{\text{shock}} ,\\ \hat{\phi} = \max\{\varphi_5, \varphi_6\} + \frac{1}{2}|\boldsymbol{\xi}|^2 & \text{on } \Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6 ,\\ \partial_n \hat{\phi} = 0 & \text{on } \Gamma_{\text{wedge}} , \end{cases}$

has a unique solution $\hat{\phi} \in C^2(\Omega) \cap C^1(\overline{\Omega})$, where the nonlinear operators $\mathcal{N}_{(u,\theta)}$ and $\mathcal{M}_{(u,\theta)}$ are determined later. Moreover, solution $\hat{\phi}$, under the transformation:

(4.24)
$$\hat{u} \coloneqq (\hat{\phi} - \frac{1}{2} |\boldsymbol{\xi}|^2 - \varphi_{\boldsymbol{\theta}}^*) \circ \mathfrak{F}_{(u,\boldsymbol{\theta})} \qquad in \ \mathcal{Q}^{\text{iter}}$$

satisfies the estimate:

(4.25)
$$\|\hat{u} - u\|_{2,\alpha,\mathcal{Q}^{\text{iter}}}^{(*)} < \delta_3$$

Definition 4.11 (Extended iteration set). Define the extended iteration set \mathcal{K}^{ext} as

$$\mathcal{K}^{\text{ext}} \coloneqq \left\{ (u, \boldsymbol{\theta}) \in C^{2, \alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2 : (u, \boldsymbol{\theta}) \text{ satisfies Definition 4.10(i)-(v)} \right\}.$$

Write $\overline{\mathcal{K}}$ and $\overline{\mathcal{K}^{\text{ext}}}$ for the closures of \mathcal{K} and \mathcal{K}^{ext} in $C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0,\theta_*]^2$ respectively.

To complete Definition 4.10, it remains to construct a suitable iteration boundary value problem (4.23) through the definition of the nonlinear differential operators $\mathcal{N}_{(u,\theta)}$ and $\mathcal{M}_{(u,\theta)}$ for each $(u,\theta) \in \overline{\mathcal{K}^{\text{ext}}}$. The construction of these operators is similar to [2, §4.4].

In the following, we write $\Gamma_{\text{sonic}} \coloneqq \Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6$ and $\mathcal{D}_r \coloneqq \mathcal{D}_r^5 \cup \mathcal{D}_r^6$ for any r > 0 with \mathcal{D}_r^5 and \mathcal{D}_r^6 given by (4.2), respectively. Moreover, for any $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}}$, we define

$$\phi \coloneqq \varphi + \frac{1}{2} |\boldsymbol{\xi}|^2, \qquad \psi \coloneqq (\varphi - \varphi_{\boldsymbol{\theta}}^*) \circ \mathcal{R}^{-1} \equiv (\phi - \phi_{\boldsymbol{\theta}}^*) \circ \mathcal{R}^{-1},$$

where $\varphi = \varphi^{(u,\theta)}$ is given by Definition 4.7, φ_{θ}^* is given by Definition 4.4, $\phi_{\theta}^* \coloneqq \varphi_{\theta}^* + \frac{1}{2} |\boldsymbol{\xi}|^2$, and coordinates $(x, y) = \mathcal{R}(\boldsymbol{\xi})$ are given by (4.1).

Construction of $\mathcal{N}_{(u,\theta)}$ in (4.23). For $(u,\theta) \in \overline{\mathcal{K}^{\text{ext}}}$, the nonlinear operator $\mathcal{N}_{(u,\theta)}$ in (4.23) is defined by

(4.26)
$$\mathcal{N}_{(u,\boldsymbol{\theta})}(\hat{\phi}) \coloneqq \sum_{i,j=1}^{2} A_{ij}(D\hat{\phi},\boldsymbol{\xi})\partial_i\partial_j\hat{\phi} \quad \text{for any } \hat{\phi} \in C^2(\Omega) ,$$

with $(\partial_1, \partial_2) \coloneqq (\partial_{\xi}, \partial_{\eta})$ and coefficients $A_{ij}(\boldsymbol{p}, \boldsymbol{\xi}), i, j = 1, 2$, determined below such that equation $\mathcal{N}_{(u,\boldsymbol{\theta})}(\hat{\phi}) = 0$ is strictly elliptic in $\overline{\Omega} \setminus \Gamma_{\text{sonic}}$ and, if ϕ solves (4.23), then $\mathcal{N}_{(u,\boldsymbol{\theta})}(\phi) = 0$ coincides with (3.1). Coefficients $A_{ij}(\boldsymbol{p}, \boldsymbol{\xi})$ are constructed similarly to [2, §4.4.1], the main difference being that both sonic boundaries may now degenerate into single points. We briefly describe the construction in the following four steps and highlight where the differences arise.

1. Away from Γ_{sonic} . Let $\epsilon_0 > 0$ be the constant from Lemma 4.8(ii), and let $\epsilon_{\text{eq}} \in (0, \frac{\epsilon_0}{2})$ be chosen later. Away from Γ_{sonic} , coefficients $A_{ij}(\boldsymbol{p}, \boldsymbol{\xi})$ are independent of \boldsymbol{p} . Indeed, for i, j = 1, 2, and $\boldsymbol{\xi} \in \Omega \setminus \mathcal{D}_{\epsilon_{\text{eq}}/10}$, we define $A_{ij}^{(1)}(\boldsymbol{\xi})$ by

(4.27)
$$A_{ij}^{(1)}(\boldsymbol{\xi}) \coloneqq A_{ij}^{\text{pot}}(D\phi(\boldsymbol{\xi}), \phi(\boldsymbol{\xi}), \boldsymbol{\xi}),$$

where $A_{ij}^{\text{pot}}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq \delta_{ij}c^2 - (p_i - \boldsymbol{\xi} \cdot \boldsymbol{e}_i)(p_j - \boldsymbol{\xi} \cdot \boldsymbol{e}_j)$ for $c = c(|\boldsymbol{p} - \boldsymbol{\xi}|, z - \frac{1}{2}|\boldsymbol{\xi}|^2)$ given in Definition 4.10(vi), where $\boldsymbol{e}_1 \coloneqq (1,0)$ and $\boldsymbol{e}_2 \coloneqq (0,1)$.

2. Supersonic near Γ_{sonic} . For constant $\mu_0 > 0$ from Definition 4.10(iv), fix an odd function of a single variable $\zeta_1 \in C^3(\mathbb{R})$ such that, for all $s \in \mathbb{R}$,

$$\zeta_1(s) = \begin{cases} s & \text{if } |s| \le \frac{2 - \frac{\mu_0}{5}}{1 + \gamma}, \\ \frac{(2 - \frac{\mu_0}{10}) \operatorname{sgn}(s)}{1 + \gamma} & \text{if } |s| > \frac{2}{1 + \gamma}, \end{cases} \quad 0 \le \zeta_1'(s) \le 10, \quad -\frac{20(1 + \gamma)}{\mu_0} \le \zeta_1''(|s|) \le 0.$$

For any $\hat{\phi} \in C^2(\Omega)$, set

$$\hat{\psi} \coloneqq \left(\hat{\phi} - (\varphi_{\theta}^* + \frac{1}{2} |\boldsymbol{\xi}|^2)\right) \circ \mathcal{R}^{-1} \equiv \left(\hat{\phi} - \phi_{\theta}^*\right) \circ \mathcal{R}^{-1} \quad \text{in } \mathcal{R}(\Omega \cap \mathcal{D}_{2\epsilon_{\text{eq}}}),$$

and define the nonlinear operator $\mathcal{N}_{(u,\theta)}^{\text{polar}}(\hat{\phi})$ in $\Omega \cap \mathcal{D}_{2\epsilon_{\text{eq}}}$ by

$$\mathcal{N}_{(u,\theta)}^{\text{polar}}(\hat{\phi}) \coloneqq \left(\left(2x - (\gamma+1)x\,\zeta_1(\frac{\psi_x}{x}) + \mathcal{O}_1^{\text{mod}} \right) \hat{\psi}_{xx} + \mathcal{O}_2^{\text{mod}} \hat{\psi}_{xy} + \left(\frac{1}{c_{\theta}} + \mathcal{O}_3^{\text{mod}} \right) \hat{\psi}_{yy} - \left(1 + \mathcal{O}_4^{\text{mod}} \right) \hat{\psi}_x + \mathcal{O}_5^{\text{mod}} \hat{\psi}_y \right) \circ \mathcal{R} \,,$$

with the modified remainder terms $\mathcal{O}_j^{\text{mod}}$, $j = 1, \dots, 5$, given by

$$\mathcal{O}_j^{\text{mod}} \coloneqq \mathcal{O}_j\big(x^{3/4}\zeta_1\big(\frac{\hat{\psi}_x}{x^{3/4}}\big), N_4(\gamma+1)x\,\zeta_1\big(\frac{\hat{\psi}_y}{N_4(\gamma+1)x}\big), \psi(x,y), x, c_{\theta}\big)$$

where $\mathcal{O}_j(p_x, p_y, z, x, c)$ are given by (3.26), constant $N_4 > 0$ is from Definition 4.10(iv), and $c_{\theta} \coloneqq c_5(1 - \chi_{\theta}^*) + c_6 \chi_{\theta}^*$ with function χ_{θ}^* from Definition 4.4.

For $(\boldsymbol{p}, \boldsymbol{\xi}) \in \mathbb{R}^2 \times (\Omega \cap \mathcal{D}_{2\epsilon_{eq}})$, coefficients $A_{ij}^{(2)}(\boldsymbol{p}, \boldsymbol{\xi})$, i, j = 1, 2, are defined to be the second-order coefficients of operator $c_{\boldsymbol{\theta}} \mathcal{N}_{(u,\boldsymbol{\theta})}^{\text{polar}}(\hat{\phi})$ in the $\boldsymbol{\xi}$ -coordinates, that is,

$$c_{\boldsymbol{\theta}} \mathcal{N}^{\text{polar}}_{(u,\boldsymbol{\theta})}(\hat{\phi}) \coloneqq \sum_{i,j=1}^{2} A^{(2)}_{ij}(D_{\boldsymbol{\xi}}\hat{\phi}, \boldsymbol{\xi}) \partial_{i} \partial_{j} \hat{\phi} \qquad \text{in } \Omega \cap \mathcal{D}_{2\epsilon_{\text{eq}}}$$

In particular, there are no lower order terms in the above expression, since they vanish identically in $\Omega \cap \mathcal{D}_{2\epsilon_{eq}}$, which can be verified by direct computation.

3. Subsonic cut-off near Γ_{sonic} . For any $\sigma \in (0, 1)$, there exists a constant $C_{\sigma} > 0$ depending only on $(\gamma, v_2, \theta_*, \sigma)$ such that, for any $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}}$, there is a function $v_{\sigma}^{(u,\theta)} \in C^4(\overline{\Omega})$ with

- (a) $\|v_{\sigma}^{(u,\theta)} \phi\|_{C^1(\overline{\Omega})} \leq \sigma^2$ and $\|v_{\sigma}^{(u,\theta)}\|_{C^4(\overline{\Omega})} \leq C_{\sigma};$
- (b) For any sequence $\{(u_k, \boldsymbol{\theta}_k)\} \subset \overline{\mathcal{K}^{\text{ext}}}$ converging to $(u, \boldsymbol{\theta})$ in $C^{1,\alpha}(\overline{\mathcal{Q}^{\text{iter}}}) \times [0, \theta_*]^2$,

$$v_{\sigma}^{(u_k,\boldsymbol{\theta}_k)} \circ \mathfrak{F}_{(u_k,\boldsymbol{\theta}_k)} \to v_{\sigma}^{(u,\boldsymbol{\theta})} \circ \mathfrak{F}_{(u,\boldsymbol{\theta})} \quad \text{in } C^{1,\alpha}(\mathcal{Q}^{\text{iter}}).$$

The proof of this statement can be found in [2, Lemma 4.26].

For any $\sigma > 0$, fix a family of cut-off functions $\varsigma_{\sigma} \in C^{\infty}(\mathbb{R})$ satisfying

(4.28)
$$\varsigma_{\sigma}(t) \coloneqq \varsigma_1\left(\frac{t}{\sigma}\right)$$
 with $\varsigma_1(t) = \begin{cases} 1 & \text{for } t < 1, \\ 0 & \text{for } t > 2, \end{cases}$ and $0 \le \varsigma_1 \le 1$ on \mathbb{R} .

For a constant $\sigma_{cf} \in (0, 1)$ to be specified later, we define the subsonic coefficients:

$$A_{ij}^{\text{subs}}(\boldsymbol{p},\boldsymbol{\xi}) \coloneqq \varsigma_{\sigma_{\text{cf}}} A_{ij}^{\text{pot}}(\boldsymbol{p},\phi(\boldsymbol{\xi}),\boldsymbol{\xi}) + (1-\varsigma_{\sigma_{\text{cf}}}) A_{ij}^{\text{pot}}(Dv_{\sigma_{\text{cf}}}^{(u,\boldsymbol{\theta})}(\boldsymbol{\xi}),\phi(\boldsymbol{\xi}),\boldsymbol{\xi}) \quad \text{for } i,j=1,2\,,$$

with $\varsigma_{\sigma_{\rm cf}} = \varsigma_{\sigma_{\rm cf}}(|\boldsymbol{p} - Dv_{\sigma_{\rm cf}}^{(\boldsymbol{u},\boldsymbol{\theta})}(\boldsymbol{\xi})|)$. In particular, from now on, we choose $\sigma_{\rm cf} \coloneqq \sqrt{\delta_1}$. Fix a cut-off function $\chi_{\rm eq} \in C^{\infty}(\mathbb{R})$ satisfying

$$\chi_{\rm eq}(\theta) = \begin{cases} 1 & \text{for } \theta \le \theta^{\rm s} + \frac{\sigma_3}{4} ,\\ 0 & \text{for } \theta \ge \theta^{\rm s} + \frac{\sigma_3}{2} , \end{cases} \quad \chi_{\rm eq}'(\theta) \le 0 \text{ on } \mathbb{R} .$$

Then, for $(\boldsymbol{p}, \boldsymbol{\xi}) \in \mathbb{R}^2 \times (\Omega \cap \mathcal{D}_{2\epsilon_{eq}})$, coefficients $A_{ij}^{(3)}(\boldsymbol{p}, \boldsymbol{\xi}), i, j = 1, 2$, are defined by

$$A_{ij}^{(3)}(\boldsymbol{p},\boldsymbol{\xi}) \coloneqq \chi_{\rm eq}(\theta_{l-4}) A_{ij}^{(2)}(\boldsymbol{p},\boldsymbol{\xi}) + (1 - \chi_{\rm eq}(\theta_{l-4})) A_{ij}^{\rm subs}(\boldsymbol{p},\boldsymbol{\xi}) \qquad \text{for } \boldsymbol{\xi} \in \Omega \cap \mathcal{D}_{2\epsilon_{\rm eq}}^{l}, \, l = 5, 6 \,.$$

4. Coefficients $A_{ij}(\boldsymbol{p},\boldsymbol{\xi})$. We now complete the definition of coefficients $A_{ij}(\boldsymbol{p},\boldsymbol{\xi})$ for the nonlinear differential operator $\mathcal{N}_{(u,\theta)}$ given by (4.26). Let ϵ_0 be from Lemma 4.8(ii). For each $\epsilon \in (0, \frac{\epsilon_0}{2})$, fix a family of cut-off functions $\zeta_2^{(\epsilon,\theta)} \in C^4(\overline{Q^{\theta}})$ with the following properties:

- (a) Function $\zeta_2^{(\epsilon,\cdot)}(\cdot): (\boldsymbol{\xi},\boldsymbol{\theta}) \mapsto \zeta_2^{(\epsilon,\boldsymbol{\theta})}(\boldsymbol{\xi})$ is continuous on $\cup_{\boldsymbol{\theta}\in[0,\theta_*]^2} Q^{\boldsymbol{\theta}} \times \{\boldsymbol{\theta}\};$
- (b) There exists a constant $C_{\epsilon} > 0$ depending only on (γ, v_2, ϵ) such that $\|\zeta_2^{(\epsilon, \theta)}\|_{C^4(\overline{O^{\theta}})} \leq C_{\epsilon}$;

(c)
$$\zeta_2^{(\epsilon,\boldsymbol{\theta})}(\boldsymbol{\xi}) = \begin{cases} 1 & \text{for } \boldsymbol{\xi} \in \Omega \setminus \mathcal{D}_{\epsilon}, \\ 0 & \text{for } \boldsymbol{\xi} \in \Omega \cap \mathcal{D}_{\epsilon/2} \end{cases}$$

Such a family of cut-off functions is constructed in [2, Definition 4.28].

Coefficients $A_{ij}(\boldsymbol{p},\boldsymbol{\xi}), i, j = 1, 2$, are defined by

(4.29)
$$A_{ij}(\boldsymbol{p},\boldsymbol{\xi}) \coloneqq \zeta_2^{(\epsilon_{eq},\boldsymbol{\theta})}(\boldsymbol{\xi}) A_{ij}^{(1)}(\boldsymbol{\xi}) + (1 - \zeta_2^{(\epsilon_{eq},\boldsymbol{\theta})}(\boldsymbol{\xi})) A_{ij}^{(3)}(\boldsymbol{p},\boldsymbol{\xi}) \quad \text{for } (\boldsymbol{p},\boldsymbol{\xi}) \in \mathbb{R}^2 \times \Omega.$$

We have the following lemma concerning the properties of coefficients A_{ij} ; see [2, Lemma 4.30].

Lemma 4.12. There exist positive constants $\epsilon^{(1)}, \delta_1^{(1)}, \epsilon_{eq} \in (0, \frac{\epsilon_0}{2}), \lambda_0 \in (0, 1), N_{eq} \geq 1$, and C > 0 with $(\epsilon^{(1)}, \delta_1^{(1)}, \lambda_0)$ depending only on (γ, v_2) , (ϵ_{eq}, N_{eq}) depending only on (γ, v_2, θ_*) , and C > 0 depending only on $(\gamma, v_2, \theta_*, \alpha)$ such that, whenever (ϵ, δ_1) from Definition 4.10 are chosen from $(0, \epsilon^{(1)}] \times (0, \delta_1^{(1)}]$, for each $(u, \theta) \in \overline{\mathcal{K}^{ext}}$, coefficients $A_{ij}(\boldsymbol{p}, \boldsymbol{\xi})$ of $\mathcal{N}_{(u,\theta)}$ given by (4.29) satisfy the following properties:

(i) $A_{12}(\boldsymbol{p},\boldsymbol{\xi}) = A_{21}(\boldsymbol{p},\boldsymbol{\xi})$ for all $(\boldsymbol{p},\boldsymbol{\xi}) \in \mathbb{R}^2 \times \Omega$, and $A_{ij}(\boldsymbol{p},\boldsymbol{\xi}) = A_{ij}^{(1)}(\boldsymbol{\xi})$ if $\boldsymbol{\xi} \in \Omega \setminus \mathcal{D}_{\epsilon_{eq}}$. Furthermore, $D_{\boldsymbol{p}}^m A_{ij} \in C^{1,\alpha}(\mathbb{R}^2 \times (\overline{\Omega} \setminus \Gamma_{\text{sonic}}))$ for m = 0, 1, 2,

$$\lambda_0 \operatorname{dist}(\boldsymbol{\xi}, \, \Gamma_{\operatorname{sonic}}) |\boldsymbol{\kappa}|^2 \leq \sum_{i,j=1}^2 A_{ij}(\boldsymbol{p}, \boldsymbol{\xi}) \kappa_i \kappa_j \leq \lambda_0^{-1} |\boldsymbol{\kappa}|^2 \qquad \text{for any } \boldsymbol{\kappa} = (\kappa_1, \kappa_2) \in \mathbb{R}^2 \,,$$

for any $(\boldsymbol{p}, \boldsymbol{\xi}) \in \mathbb{R}^2 \times \Omega$, and

$$\begin{split} \|A_{ij}\|_{L^{\infty}(\mathbb{R}^{2}\times\Omega)} + \|A_{ij}(\boldsymbol{p},\cdot)\|_{C^{3/4}(\overline{\Omega})} + \|D_{\boldsymbol{p}}A_{ij}(\boldsymbol{p},\cdot)\|_{L^{\infty}(\Omega)} \leq N_{\text{eq}} \quad \text{for any } \boldsymbol{p} \in \mathbb{R}^{2} \,, \\ \|A_{ij}\|_{C^{1,\alpha}(\mathbb{R}^{2}\times\overline{\Omega\setminus\mathcal{D}_{\epsilon_{\text{eq}}}})} + s^{5}\|D_{\boldsymbol{p}}^{m}A_{ij}\|_{C^{1,\alpha}(\mathbb{R}^{2}\times(\overline{\Omega\setminus\mathcal{N}_{s}(\Gamma_{\text{sonic}})))} \leq C \quad \text{for each } s \in (0,\frac{\epsilon_{0}}{2}) \end{split}$$

- (ii) $A_{ij}(\boldsymbol{p},\boldsymbol{\xi}) = A_{ij}^{(3)}(\boldsymbol{p},\boldsymbol{\xi}) \text{ for any } (\boldsymbol{p},\boldsymbol{\xi}) \in \mathbb{R}^2 \times (\Omega \cap \mathcal{D}_{\epsilon_{eq}/2}^l), l = 5, 6.$ Furthermore, if $\theta_{l-4} \leq \theta^s + \frac{\sigma_3}{4}$, then $A_{ij}(\boldsymbol{p},\boldsymbol{\xi}) = A_{ij}^{(2)}(\boldsymbol{p},\boldsymbol{\xi})$; whereas, if $\theta_{l-4} \in [\theta^s + \delta, \theta_*]$ for some $\delta \in (0, \frac{\sigma_3}{2})$, then $\lambda_0(\operatorname{dist}(\boldsymbol{\xi}, \Gamma_{\operatorname{sonic}}) + \delta)|\boldsymbol{\kappa}|^2 \leq \sum_{i,j=1}^2 A_{ij}(\boldsymbol{p},\boldsymbol{\xi})\kappa_i\kappa_j \leq \lambda_0^{-1}|\boldsymbol{\kappa}|^2 \quad \text{for any } \boldsymbol{\xi} \in \Omega \cap \mathcal{D}_{\epsilon_{eq}/2}^l,$ $\sup_{\boldsymbol{p} \in \mathbb{R}^2} \|\mathcal{D}_{\boldsymbol{p}}^m A_{ij}(\boldsymbol{p},\cdot)\|_{C^{1,\alpha}(\overline{\Omega \cap \mathcal{D}_{\epsilon_{eq}/2}^l})} \leq C \quad \text{for } m = 0, 1, 2.$
- (iii) Suppose that ϵ from Definition 4.10 satisfies $\epsilon \in (0, \frac{\epsilon_{eq}}{2})$. Then the equation: $\mathcal{N}_{(u,\theta)}(\phi) = 0$ coincides with (3.1) in $\Omega \setminus \mathcal{D}_{\epsilon/10}$. In addition, for l = 5, 6, if $x_{P_0^{l-4}} \ge \frac{\epsilon}{10}$ or $\theta_{l-4} \ge \theta^{s} + \frac{\sigma_3}{2}$, then the equation: $\mathcal{N}_{(u,\theta)}(\phi) = 0$ coincides with (3.1) in $\Omega \setminus \mathcal{D}_{\epsilon/10}^{11-l}$.

Construction of $\mathcal{M}_{(u,\theta)}$ in (4.23). We turn our attention to the definition of the nonlinear linear operator $\mathcal{M}_{(u,\theta)}(\mathbf{p}, z, \boldsymbol{\xi})$ in (4.23). The construction is presented in the following four steps that follow closely [2, §4.4.2].

1. For $g^{\rm sh}$ given by (3.27), set

$$\mathcal{M}_0(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq g^{\mathrm{sh}}(\boldsymbol{p} - \boldsymbol{\xi}, z - \frac{1}{2} |\boldsymbol{\xi}|^2, \boldsymbol{\xi}), \qquad \mathcal{M}_1(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq \mathcal{M}_0(\boldsymbol{p}, z, (\boldsymbol{\xi}, \frac{z}{v_2})).$$

For j = 5, 6, write $\phi_j \coloneqq \varphi_j + \frac{1}{2} |\boldsymbol{\xi}|^2$, which is independent of the η -coordinate since $O_j \in \{\eta = 0\}$. Then $(\xi, \frac{1}{v_2}\phi_j(\boldsymbol{\xi})) \in \{\varphi_j = \varphi_2\}$ for any $\boldsymbol{\xi} \in \mathbb{R}^2$, and \mathcal{M}_1 is homogeneous in the sense that

$$\mathcal{M}_1(D\phi_j(\boldsymbol{\xi}),\phi_j(\boldsymbol{\xi}),\xi) = 0$$
 for all $\boldsymbol{\xi} = (\xi,\eta) \in \mathbb{R}^2$ and $j = 5,6$.

Furthermore, observe that $\boldsymbol{\xi} = (\xi, \eta) \in \Gamma_{\text{shock}}$ if and only if $\boldsymbol{\xi} \in \overline{\Omega}$ with $\eta = \frac{1}{v_2} \phi(\boldsymbol{\xi})$, and hence

(4.30)
$$\begin{cases} \mathcal{M}_1(D\phi(\boldsymbol{\xi}), \phi(\boldsymbol{\xi}), \xi) = \mathcal{M}_0(D\phi(\boldsymbol{\xi}), \phi(\boldsymbol{\xi}), \boldsymbol{\xi}), \\ D_{\boldsymbol{p}}\mathcal{M}_1(D\phi(\boldsymbol{\xi}), \phi(\boldsymbol{\xi}), \xi) = D_{\boldsymbol{p}}\mathcal{M}_0(D\phi(\boldsymbol{\xi}), \phi(\boldsymbol{\xi}), \boldsymbol{\xi}), \end{cases} \text{ for } \boldsymbol{\xi} \in \Gamma_{\text{shock}} \end{cases}$$

2. For the cut-off function ς_{σ} given by (4.28) and a constant $\sigma_{bc} > 0$ to be determined later, we set

(4.31)
$$\mathcal{M}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq \left(\varsigma_{\sigma_{\mathrm{bc}}}^{(5)} \varsigma_{\sigma_{\mathrm{bc}}}^{(6)} + (1 - \varsigma_{\sigma_{\mathrm{bc}}}^{(5)}) \varsigma_{\sigma_{\mathrm{bc}}}^{(6)} + \varsigma_{\sigma_{\mathrm{bc}}}^{(5)} (1 - \varsigma_{\sigma_{\mathrm{bc}}}^{(6)}) \right) \mathcal{M}_{1}(\boldsymbol{p}, z, \boldsymbol{\xi}) \\ + (1 - \varsigma_{\sigma_{\mathrm{bc}}}^{(5)}) (1 - \varsigma_{\sigma_{\mathrm{bc}}}^{(6)}) \mathcal{M}_{0}(\boldsymbol{p}, z, \boldsymbol{\xi}) ,$$

where $\varsigma_{\sigma_{\rm bc}}^{(j)} \coloneqq \varsigma_{\sigma_{\rm bc}}(|(\boldsymbol{p}, z) - (D\phi_j, \phi_j(\boldsymbol{\xi}))|)$ for j = 5, 6. There exist constants $\sigma_{\rm bc}, \delta_{\rm bc}, \bar{\epsilon}_{\rm bc} > 0$ depending only on (γ, v_2) such that, if ϵ from Definition 4.10 satisfies $\epsilon \in (0, \bar{\epsilon}_{\rm bc})$, then $\mathcal{M}(\boldsymbol{p}, z, \boldsymbol{\xi})$ satisfies that, for all $\boldsymbol{\xi} \in \Gamma_{\rm shock}$,

$$\delta_{\rm bc} \leq D_{\boldsymbol{p}} \mathcal{M}(D\phi(\boldsymbol{\xi}), \phi(\boldsymbol{\xi}), \boldsymbol{\xi}) \cdot \boldsymbol{\nu}_{\rm sh}(\boldsymbol{\xi}) \leq \delta_{\rm bc}^{-1}, \qquad D_z \mathcal{M}(D\phi(\boldsymbol{\xi}), \phi(\boldsymbol{\xi}), \boldsymbol{\xi}) \leq -\delta_{\rm bc},$$

where $\nu_{\rm sh}$ is the unit normal vector to $\Gamma_{\rm shock}$ towards the interior of Ω . These results follow from the calculations in [2, Lemma 4.32]. Indeed, the first follows from a direct calculation by using (4.30), [2, Eq. A.18], the fact that \mathcal{M}_0 is C^1 , and the properties in Definition 4.10(vi)–(v). For the second, direct computation gives

$$D_z \mathcal{M}_0(D\phi(\boldsymbol{\xi}), \phi(\boldsymbol{\xi}), \boldsymbol{\xi}) = -
ho^{2-\gamma} D arphi \cdot \boldsymbol{
u}_{
m sh}(\boldsymbol{\xi}) < 0 \qquad ext{for } \boldsymbol{\xi} \in \Gamma_{
m shock} \,,$$

whilst, for j = 5, 6, and any $\boldsymbol{\xi} = (\xi, \eta) \in Q^{\boldsymbol{\theta}}$, we have

$$D_z \mathcal{M}_1(D\phi_j(\boldsymbol{\xi}), \phi_j(\boldsymbol{\xi}), \xi) = -\rho_j^{2-\gamma} \operatorname{dist}(O_j, S_{2j}) + (-1)^j \frac{\rho_j - 1}{v_2} \cos \theta_{2j} < 0,$$

after which an argument similar to [2, Lemma 4.32] applies. The constant, $\sigma_{\rm bc} > 0$, is now fixed.

3. For j = 5, 6, set

(4.32)
$$\mathcal{M}^{(j)}(\boldsymbol{q}, \boldsymbol{z}, \boldsymbol{\xi}) \coloneqq \mathcal{M}(\boldsymbol{q} + \boldsymbol{D}\phi_j, \boldsymbol{z} + \phi_j(\boldsymbol{\xi}), \boldsymbol{\xi}) \,.$$

Then, for the coordinates $(x, y) = \mathbf{x} = \mathcal{R}(\boldsymbol{\xi})$ given by (4.1), define

(4.33)
$$\hat{\mathcal{M}}^{(j)}(q_x, q_y, z, \boldsymbol{x}) \coloneqq \mathcal{M}^{(j)}\big((-1)^j (q_x \cos y + \frac{q_y \sin y}{c_j - x}), -q_x \sin y + \frac{q_y \cos y}{c_j - x}, z, \mathcal{R}^{-1}(\boldsymbol{x})\big).$$

4. Finally, we extend the definition of \mathcal{M} in (4.31) to all $(\boldsymbol{p}, z, \boldsymbol{\xi}) \in \mathbb{R}^2 \times \mathbb{R} \times \overline{\Omega}$. For each $(u, \boldsymbol{\theta}) \in \overline{\mathcal{K}^{\text{ext}}}$ and a constant $\sigma > 0$, let $v_{\sigma}^{(u, \boldsymbol{\theta})} \in C^4(\overline{\Omega})$ be given as in Step **3** in the construction of $\mathcal{N}_{(u, \boldsymbol{\theta})}$ above. We define a linear operator $\mathcal{L}_{\sigma}^{(u, \boldsymbol{\theta})}(\boldsymbol{p}, z, \boldsymbol{\xi})$ by

$$\mathcal{L}^{(u,\theta)}_{\sigma}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq \mathcal{M}(Dv^{(u,\theta)}_{\sigma}(\boldsymbol{\xi}), v^{(u,\theta)}_{\sigma}(\boldsymbol{\xi}), \boldsymbol{\xi}) + D_{\boldsymbol{p}}\mathcal{M}(Dv^{(u,\theta)}_{\sigma}(\boldsymbol{\xi}), v^{(u,\theta)}_{\sigma}(\boldsymbol{\xi}), \boldsymbol{\xi}) \cdot \boldsymbol{p} + D_{z}\mathcal{M}(Dv^{(u,\theta)}_{\sigma}(\boldsymbol{\xi}), v^{(u,\theta)}_{\sigma}(\boldsymbol{\xi}), \boldsymbol{\xi})z,$$

The operator, $\mathcal{M}_{(u,\theta)}(\boldsymbol{p}, z, \boldsymbol{\xi})$, in (4.23) is then defined by

(4.34)
$$\mathcal{M}_{(u,\theta)}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq \varsigma_{\sigma} \mathcal{M}(\boldsymbol{p}, z, \boldsymbol{\xi}) + (1 - \varsigma_{\sigma}) \mathcal{L}_{\sigma}^{(u,\theta)}(\boldsymbol{p} - Dv_{\sigma}^{(u,\theta)}(\boldsymbol{\xi}), z - v_{\sigma}^{(u,\theta)}(\boldsymbol{\xi}), \boldsymbol{\xi})$$

for $\varsigma_{\sigma} \coloneqq \varsigma_{\sigma}(|(\boldsymbol{p}, z) - (Dv_{\sigma}^{(u,\theta)}(\boldsymbol{\xi}), v_{\sigma}^{(u,\theta)}(\boldsymbol{\xi}))|)$ given by (4.28). We have the following lemma concerning the properties of the nonlinear operator $\mathcal{M}_{(u,\theta)}(\boldsymbol{p}, z, \boldsymbol{\xi})$; see [2, Lemma 4.34].

Lemma 4.13. Let $\bar{\epsilon}_{bc}$ be the constant from Step 2 above. Then there exist positive constants $\epsilon_{bc} \in (0, \bar{\epsilon}_{bc}), \delta_1^{(2)}, N_1^{(1)}, \delta_{bc}, C, C_{\theta_*}, and \epsilon_{\mathcal{M}} \in (0, \epsilon_{bc}]$ with $(\epsilon_{bc}, \delta_1^{(2)}, N_1^{(1)}, \delta_{bc}, C)$ depending only on $(\gamma, v_2), \epsilon_{\mathcal{M}}$ depending on (γ, v_2, θ_*) , and C_{θ_*} depending on $(\gamma, v_2, \theta_*, \alpha)$ such that, whenever parameters $(\epsilon, \delta_1, N_1)$ from Definition 4.10 are chosen from $(0, \bar{\epsilon}_{bc}] \times (0, \delta_1^{(2)}] \times [N_1^{(1)}, \infty)$, the following statements hold: For each $(u, \theta) \in \overline{\mathcal{K}^{ext}}$, operator $\mathcal{M}_{(u,\theta)} : \mathbb{R}^2 \times \mathbb{R} \times \overline{\Omega} \to \mathbb{R}$ given by (4.34) with $\sigma = \sigma_{cf} \equiv \sqrt{\delta_1}$ satisfies

(i)
$$\mathcal{M}_{(u,\theta)} \in C^3(\mathbb{R}^2 \times \mathbb{R} \times \overline{\Omega})$$
 and, for all $(\mathbf{p}, z, \boldsymbol{\xi}) \in \mathbb{R}^2 \times \mathbb{R} \times \overline{\Omega}$ and $\boldsymbol{\xi}_{\mathrm{sh}} \in \overline{\Gamma_{\mathrm{shock}}}$,
 $\|(\mathcal{M}_{(u,\theta)}(\mathbf{0}, 0, \cdot), D^m_{(\mathbf{p}, z)} \mathcal{M}_{(u,\theta)}(\mathbf{p}, z, \cdot))\|_{C^3(\overline{\Omega})} \leq C_{\theta_*} \quad \text{for } m = 1, 2, 3,$
 $\mathcal{M}_{(u,\theta)}(\mathbf{p}, z, \boldsymbol{\xi}) = \mathcal{M}(\mathbf{p}, z, \boldsymbol{\xi}) \quad \text{for } |(\mathbf{p}, z) - (D\phi(\boldsymbol{\xi}), \phi(\boldsymbol{\xi}))| < \frac{1}{2}\sqrt{\delta_1},$
 $|D_{(\mathbf{p}, z)} \mathcal{M}_{(u,\theta)}(\mathbf{p}, z, \boldsymbol{\xi}) - D_{(\mathbf{p}, z)} \mathcal{M}(D\phi(\boldsymbol{\xi}), \phi(\boldsymbol{\xi}), \boldsymbol{\xi})| \leq C\sqrt{\delta_1},$
 $\delta_{\mathrm{bc}} \leq D_{\mathbf{p}} \mathcal{M}_{(u,\theta)}(\mathbf{p}, z, \boldsymbol{\xi}_{\mathrm{sh}}) \cdot \boldsymbol{\nu}_{\mathrm{sh}} \leq \frac{1}{\delta_{\mathrm{bc}}}, \quad D_z \mathcal{M}_{(u,\theta)}(\mathbf{p}, z, \boldsymbol{\xi}_{\mathrm{sh}}) \leq -\delta_{\mathrm{bc}},$

where $\mathcal{M}(\mathbf{p}, z, \boldsymbol{\xi})$ is defined by (4.31), and $\boldsymbol{\nu}_{sh}$ is the unit normal vector to Γ_{shock} pointing into Ω .

(ii) Denote
$$\mathcal{B}_{\sigma,\Gamma_{\mathrm{shock}}}^{(u,\theta)}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq \mathcal{L}_{\sigma}^{(u,\theta)}(\boldsymbol{p} - Dv_{\sigma}^{(u,\theta)}(\boldsymbol{\xi}), z - v_{\sigma}^{(u,\theta)}(\boldsymbol{\xi}), \boldsymbol{\xi})$$
 and set
 $\mathcal{B}_{\sigma,\Gamma_{\mathrm{shock}}}^{(u,\theta)}(\boldsymbol{p}, z, \boldsymbol{\xi}) \coloneqq (b_{1}^{\mathrm{sh}}(\boldsymbol{\xi}), b_{2}^{\mathrm{sh}}(\boldsymbol{\xi}), b_{0}^{\mathrm{sh}}(\boldsymbol{\xi}), h^{\mathrm{sh}}(\boldsymbol{\xi})) \cdot (p_{1}, p_{2}, z, 1)$

 $Then \ \|(b_i^{\mathrm{sh}},h^{\mathrm{sh}})\|_{C^3(\overline{\Gamma_{\mathrm{shock}}})} \leq C_{\theta_*} \ for \ i=0,1,2, \ and, \ for \ all \ (\boldsymbol{p},z,\boldsymbol{\xi}) \in \mathbb{R}^2 \times \mathbb{R} \times \overline{\Omega},$

$$\begin{split} \| \big(\mathcal{M}_{(u,\boldsymbol{\theta})} - \mathcal{B}_{\sigma_{\mathrm{cf}},\Gamma_{\mathrm{shock}}} \big) (\boldsymbol{p}, z, \boldsymbol{\xi}) \| &\leq C \sqrt{\delta_1} \left| (\boldsymbol{p}, z) - \big(Dv_{\sigma_{\mathrm{cf}}}^{(u,\boldsymbol{\theta})}(\boldsymbol{\xi}), v_{\sigma_{\mathrm{cf}}}^{(u,\boldsymbol{\theta})}(\boldsymbol{\xi}) \big) \right|, \\ \left| D_{(\boldsymbol{p}, z)} \big(\mathcal{M}_{(u,\boldsymbol{\theta})} - \mathcal{B}_{\sigma_{\mathrm{cf}},\Gamma_{\mathrm{shock}}} \big) (\boldsymbol{p}, z, \boldsymbol{\xi}) \right| &\leq C \sqrt{\delta_1} \,. \end{split}$$

(iii) $\mathcal{M}_{(u,\theta)}$ is homogeneous in the sense that, for j = 5, 6,

$$\mathcal{M}_{(u,\boldsymbol{\theta})}(D\phi_j(\boldsymbol{\xi}),\phi_j(\boldsymbol{\xi}),\boldsymbol{\xi}) = 0 \qquad for \ all \begin{cases} \boldsymbol{\xi} \in \Gamma_{\text{shock}} \cap \mathcal{D}_{\epsilon_{\mathcal{M}}} & for \ any \ \boldsymbol{\theta} \in [0,\theta_*]^2 ,\\ \boldsymbol{\xi} \in \Gamma_{\text{shock}} & when \ \max\{\theta_1,\theta_2\} \in [0,\frac{\delta_1}{N_1}] . \end{cases}$$

(iv) For coordinates $(x, y) = \mathcal{R}(\boldsymbol{\xi})$ given by (4.1), j = 5, 6, and $\boldsymbol{\xi} \in \Gamma_{\text{shock}} \cap \mathcal{D}^{j}_{\epsilon_{\text{bc}}}$, define $\hat{\mathcal{M}}^{(j)}_{(u,\theta)}(q_{x}, q_{y}, z, x, y)$ by (4.33) with \mathcal{M} replaced by $\mathcal{M}_{(u,\theta)}$ in (4.32). Then, for j = 5, 6, $\hat{\mathcal{M}}^{(j)}_{(u,\theta)}$ satisfies the following properties when $\Gamma_{\text{shock}} \cap \mathcal{D}^{j}_{\epsilon_{\text{bc}}}$ is non-empty: $\|\hat{\mathcal{M}}^{(j)}_{(u,\theta)}\|_{C^{3}(\mathbb{R}^{2} \times \mathbb{R} \times \mathcal{R}(\overline{\Gamma_{\text{shock}} \cap \mathcal{D}^{j}_{\epsilon_{\text{bc}}}))} \leq C_{\theta_{*}},$ $\hat{\mathcal{M}}^{(j)}_{(u,\theta)}(\boldsymbol{q}, z, x, y) = \hat{\mathcal{M}}^{(j)}(\boldsymbol{q}, z, x, y) \qquad in \, \mathcal{R}(\Gamma_{\text{shock}} \cap \mathcal{D}^{j}_{\epsilon_{\text{bc}}}) \text{ for any } |(\boldsymbol{q}, z)| \leq \frac{\delta_{\text{bc}}}{C},$ $\partial_{a} \hat{\mathcal{M}}^{(j)}_{(u,\theta)}(\boldsymbol{q}, z, x, y) \leq -\delta_{\text{bc}} \qquad in \, \mathcal{R}(\Gamma_{\text{shock}} \cap \mathcal{D}^{j}_{\epsilon_{\mathcal{M}}}) \text{ for any } (\boldsymbol{q}, z) \in \mathbb{R}^{2} \times \mathbb{R},$

with $\partial_a = \partial_{q_x}$, ∂_{q_y} or ∂_z , provided that $\Gamma_{\text{shock}} \cap \mathcal{D}^j_{\epsilon_{\mathcal{M}}}$ is non-empty.

(v) $\mathcal{M}_{(u,\theta)}(D\phi(\boldsymbol{\xi}),\phi(\boldsymbol{\xi}),\boldsymbol{\xi}) = 0$ on Γ_{shock} if and only if $\varphi = \phi - \frac{1}{2}|\boldsymbol{\xi}|^2$ satisfies the Rankine– Hugoniot jump condition (3.27) on $\Gamma_{\text{shock}} = \{\varphi = \varphi_2\}.$

4.2.2. Well-posedness of the boundary value problem (4.23). Since the complete definition of the iteration boundary value problem (4.23) has been given, we now consider its well-posedness.

Lemma 4.14. Fix $\gamma \geq 1$, $v_2 \in (v_{\min}, 0)$, and $\theta_* \in (0, \theta^d)$. Let constant σ_2 be from Step 1 of the proof of Proposition 3.14. Let $\bar{\alpha} \in (0, \frac{1}{3}]$ and $\epsilon_0 > 0$ be from Proposition 4.6 and Lemma 4.8(ii) respectively, and let $\alpha \in (0, \frac{\bar{\alpha}}{2}]$ be from Definition 4.10. Then there exist constants $\epsilon^{(w)} \in (0, \epsilon_0]$, $\delta_1^{(w)} \in (0, 1)$, $N_1^{(w)} \geq 1$, and $\alpha_1^* \in (0, \bar{\alpha}]$ depending only on (γ, v_2, θ_*) such that, whenever parameters $(\epsilon, \delta_1, N_1)$ in Definition 4.10 are chosen with $\epsilon \in (0, \epsilon^{(w)}]$, $\delta_1 \in (0, \delta_1^{(w)}]$, and $N_1 \geq N_1^{(w)}$, the following statements hold:

Case (i). If $\max\{\theta_1, \theta_2\} \leq \theta^s + \sigma_2$, then the boundary value problem (4.23) associated with $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}} \cap \{\max\{\theta_1, \theta_2\} \leq \theta^s + \sigma_2\}$ has a unique solution $\hat{\phi} \in C^2(\Omega) \cap C^1(\overline{\Omega} \setminus (\Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6)) \cap C^0(\overline{\Omega})$. Moreover, there exists a constant C > 0 depending only on $(\gamma, v_2, \theta_*, \alpha)$ such that solution $\hat{\phi}$ satisfies

(4.35)
$$\|\hat{\phi}\|_{L^{\infty}(\Omega)} \leq C, \qquad |\hat{\phi}(\boldsymbol{\xi}) - \phi_{\boldsymbol{\theta}}(\boldsymbol{\xi})| \leq C \operatorname{dist}(\boldsymbol{\xi}, \Gamma_{\operatorname{sonic}}^{5} \cup \Gamma_{\operatorname{sonic}}^{6}) \quad in \ \Omega$$

for $\phi_{\boldsymbol{\theta}} := \max\{\varphi_5, \varphi_6\} + \frac{1}{2} |\boldsymbol{\xi}|^2$. Furthermore, for each $d \in (0, \epsilon_0)$, there exists a constant $C_d > 0$ depending only on $(\gamma, v_2, \theta_*, \alpha, d)$ such that

(4.36)
$$\|\tilde{\phi}\|_{2,\alpha_1^*,\Omega\setminus\mathcal{D}_d} \le C_d.$$

Case (ii). For each $\delta \in (0, \frac{\sigma_2}{2})$, whenever $\max\{\theta_1, \theta_2\} \in [\theta^s + \delta, \theta_*]$, the boundary value problem (4.23) associated with $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}} \cap \{\max\{\theta_1, \theta_2\} \geq \theta^s + \delta\}$ has a unique solution $\hat{\phi} \in C^2(\Omega) \cap C^1(\overline{\Omega} \setminus (\Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6)) \cap C^0(\overline{\Omega})$. Moreover, there exists a constant C' > 0 depending only on $(\gamma, v_2, \theta_*, \alpha, \delta)$ such that solution $\hat{\phi}$ satisfies (4.35), while there exists a constant $C'_d > 0$ depending only on $(\gamma, v_2, \theta_*, \alpha, \delta, d)$ such that (4.36) holds.

Proof. For Case (i), we fix $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}} \cap \{\max\{\theta_1, \theta_2\} \leq \theta^s + \sigma_2\}$ and fix a constant h > 0. Using \mathcal{G}_1^{θ} defined by (4.7), we rewrite the boundary value problem (4.23) associated with (u, θ) as a new boundary value problem for the unknown function:

$$v(\boldsymbol{x}) \coloneqq \hat{\phi} \circ (L_h \circ \mathcal{G}_1^{\boldsymbol{\theta}})^{-1}(\boldsymbol{x}) \qquad \text{for any } \boldsymbol{x} \in L_h \circ \mathcal{G}_1^{\boldsymbol{\theta}}(\Omega(u, \boldsymbol{\theta})),$$

where $\boldsymbol{x} \coloneqq L_h(s, t') = (\frac{s+1}{2}h, t')$ and $(s, t') = \mathcal{G}_1^{\boldsymbol{\theta}}(\boldsymbol{\xi})$ for any $\boldsymbol{\xi} \in \Omega(u, \boldsymbol{\theta})$. We write

 $\Gamma_0 \coloneqq L_h \circ \mathcal{G}_1^{\boldsymbol{\theta}}(\Gamma_{\text{sonic}}^6), \ \Gamma_1 \coloneqq L_h \circ \mathcal{G}_1^{\boldsymbol{\theta}}(\Gamma_{\text{shock}}(u, \boldsymbol{\theta})), \ \Gamma_2 \coloneqq L_h \circ \mathcal{G}_1^{\boldsymbol{\theta}}(\Gamma_{\text{sonic}}^5), \ \Gamma_3 \coloneqq L_h \circ \mathcal{G}_1^{\boldsymbol{\theta}}(\Gamma_{\text{sym}}).$ Moreover, we take

$$g_{\rm so}(\boldsymbol{x}) \coloneqq (1-\varsigma_h)\phi_5 \circ (L_h \circ \mathcal{G}_1^{\boldsymbol{\theta}})^{-1}(\boldsymbol{x}) + \varsigma_h \phi_6 \circ (L_h \circ \mathcal{G}_1^{\boldsymbol{\theta}})^{-1}(\boldsymbol{x}) \,,$$

where $\varsigma_h \coloneqq \varsigma_1(\frac{1}{2} + \frac{2}{h}x_1)$, and the cut-off function $\varsigma_1(\cdot) \in C^{\infty}(\mathbb{R})$ is given by (4.28). Using Lemmas 4.2 and 4.12–4.13, we can choose suitable constants $\epsilon^{(w)} \in (0, \epsilon], \delta_1^{(w)} > 0$, and $N_1^{(w)} \ge 1$ such that the new boundary value problem for v satisfies the conditions in Proposition B.1 when $\epsilon \in (0, \epsilon^{(w)}], \delta_1 \in (0, \delta_1^{(w)}]$, and $N_1 \ge N_1^{(w)}$. Therefore, Proposition B.1 gives the existence and uniqueness of a solution $\hat{\phi} \in C(\overline{\Omega}) \cap C^{2,\alpha_1}(\overline{\Omega} \setminus (\Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6))$ of the boundary value problem (4.23) satisfying (4.35)–(4.36).

For Case (ii), fix any $\delta \in (0, \frac{\sigma_2}{2})$. In the subcase: $\min\{\theta_1, \theta_2\} < \theta^s + \delta \leq \max\{\theta_1, \theta_2\}$ (without loss of generality, we consider the case: $\theta_1 < \theta^s + \delta \leq \theta_2$), one can follow the same analysis as above to check all the conditions for Proposition B.2, from which we obtain the existence of a unique $\hat{\phi}$ satisfying (4.35)–(4.36) for any $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}} \cap \{\theta_1 < \theta^s + \delta \leq \theta_2\}$. In the other subcase: $\min\{\theta_1, \theta_2\} \in [\theta^s + \delta, \theta_*]$, we can apply Proposition B.3 to prove the same results.

For each $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}}$, the corresponding pseudo-subsonic region $\Omega = \Omega(u, \theta)$ depends continuously on (u, θ) . We rewrite (4.23) as a boundary value problem for

(4.37)
$$\hat{u}(s,t) \coloneqq (\hat{\phi} - \frac{1}{2} |\boldsymbol{\xi}|^2 - \varphi_{\boldsymbol{\theta}}^*) \circ \mathfrak{F}_{(u,\boldsymbol{\theta})}(s,t) \quad \text{in } \mathcal{Q}^{\text{iter}},$$

where φ_{θ}^* is given by Definition 4.4, and $\mathfrak{F} = \mathfrak{F}_{(u,\theta)}$ is given by Definition 4.7(ii). We obtain

$$(4.38) \begin{cases} \sum_{i,j=1}^{2} \mathcal{A}_{ij}^{(u,\theta)}(D_{(s,t)}\hat{u},s,t)\partial_{i}\partial_{j}\hat{u} + \sum_{i=1}^{2} \mathcal{A}_{i}^{(u,\theta)}(D_{(s,t)}\hat{u},s,t)\partial_{i}\hat{u} = f^{(u,\theta)} & \text{in } \mathcal{Q}^{\text{iter}}, \\ \mathcal{M}_{(u,\theta)}(D_{(s,t)}\hat{u},\hat{u},s) = 0 & \text{on } \partial_{\text{sh}}\mathcal{Q}^{\text{iter}} \\ \hat{u} = 0 & \text{on } \partial_{\text{so}}\mathcal{Q}^{\text{iter}}, \\ \mathcal{B}_{(u,\theta)}^{(w)}(D_{(s,t)}\hat{u},s) \coloneqq \sum_{i=1}^{2} b_{i}^{(w)}(s)\partial_{i}\hat{u} = 0 & \text{on } \partial_{\text{w}}\mathcal{Q}^{\text{iter}}, \end{cases}$$

where $(\partial_1, \partial_2) \coloneqq (\partial_s, \partial_t)$ and

$$\partial_{\mathrm{sh}}\mathcal{Q}^{\mathrm{iter}} \coloneqq (-1,1) \times \{1\}, \quad \partial_{\mathrm{so}}\mathcal{Q}^{\mathrm{iter}} \coloneqq \{-1,1\} \times (0,1), \quad \partial_{\mathrm{w}}\mathcal{Q}^{\mathrm{iter}} \coloneqq (-1,1) \times \{0\}.$$

From Lemmas 4.8 and 4.12–4.14, we can obtain the following results; see [2, Lemma 4.36].

Lemma 4.15. For every $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}}$, let $\mathcal{A}_{ij}^{(u,\theta)}, \mathcal{A}_i^{(u,\theta)}, \mathcal{M}_{(u,\theta)}, \mathcal{M}_{(u,\theta)}, \mathcal{B}_{(u,\theta)}^{(w)}$, and $b_i^{(w)}$ be given as in (4.38). Then the following properties hold:

- (i) $\mathcal{A}_{ij}^{(u,\theta)}, \mathcal{A}_{i}^{(u,\theta)} \in C(\mathbb{R}^2 \times \mathcal{Q}^{\text{iter}}), f^{(u,\theta)} \in C(\mathcal{Q}^{\text{iter}}), \mathcal{M}_{(u,\theta)} \in C(\mathbb{R}^2 \times \mathbb{R} \times \partial_{\text{sh}} \mathcal{Q}^{\text{iter}}), and \mathscr{B}_{(u,\theta)}^{(w)} \in C(\mathcal{Q}^{\text{iter}})$ $C(\mathbb{R}^2 \times \mathbb{R} \times \partial_w \mathcal{Q}^{\text{iter}});$
- (ii) If a sequence $\{(u_k, \theta_k)\}_{k \in \mathbb{N}} \subseteq \overline{\mathcal{K}^{\text{ext}}}$ converges to $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}}$ in $C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2$, then the following sequences converge uniformly: • $(\mathcal{A}_{ij}^{(u_k,\theta_k)}, \mathcal{A}_i^{(u_k,\theta_k)}) \rightarrow (\mathcal{A}_{ij}^{(u,\theta)}, \mathcal{A}_i^{(u,\theta)})$ on compact subsets of $\mathbb{R}^2 \times \mathcal{Q}^{\text{iter}}$,

 - $f^{(u_k,\theta_k)} \to f^{(u,\theta)}$ on compact subsets of $\mathcal{Q}^{\text{iter}}$, $\mathcal{M}_{(u_k,\theta_k)} \to \mathcal{M}_{(u,\theta)}$ on compact subsets of $\mathbb{R}^2 \times \mathbb{R} \times \partial_{\text{sh}} \mathcal{Q}^{\text{iter}}$, $\mathscr{B}^{(w)}_{(u_k,\theta_k)} \to \mathscr{B}^{(w)}_{(u,\theta)}$ on compact subsets of $\mathbb{R}^2 \times \mathbb{R} \times \partial_{\text{w}} \mathcal{Q}^{\text{iter}}$.

Corollary 4.16. Let $\bar{\alpha} \in (0,1)$ be from Proposition 4.6, and let $\alpha_1^* \in (0,\bar{\alpha}], \epsilon^{(w)}, \delta_1^{(w)}$, and $N_1^{(w)}$ be from Lemma 4.14. Let ϵ, δ_1 , and N_1 from Definition 4.10 satisfy $\epsilon \in (0, \epsilon^{(w)}], \delta_1 \in (0, \delta_1^{(w)}]$ and $N_1 \ge N_1^{(w)}$.

(i) For each $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}}$, $\hat{\phi}$ solves the boundary value problem (4.23) if and only if \hat{u} given by (4.37) solves the boundary value problem (4.38). Thus, (4.38) has a unique solution $\hat{u} \in C^2(\mathcal{Q}^{\text{iter}}) \cap C^1(\overline{\mathcal{Q}^{\text{iter}}} \setminus \overline{\partial_{so}\mathcal{Q}^{\text{iter}}}) \cap C(\overline{\mathcal{Q}^{\text{iter}}}).$ Furthermore, there exists a constant $C \ge 1$ depending on $(\gamma, v_2, \theta_*, \alpha)$ such that

$$|\hat{u}(s,t)| \leq C(1-|s|)$$
 in $\mathcal{Q}^{\text{iter}}$.

For each $\hat{d} \in (0, \frac{1}{2})$, there exists $C_{\hat{d}} > 0$ depending on $(\gamma, v_2, \theta_*, \hat{d}, \alpha)$ such that

$$\|\hat{u}\|_{2,\alpha_1^*,\mathcal{Q}^{\text{iter}}\cap\{1-|s|>\hat{d}\}} \le C_{\hat{d}}.$$

- (ii) Let $\{(u_k, \boldsymbol{\theta}_k)\}_{k \in \mathbb{N}} \subseteq \overline{\mathcal{K}^{\text{ext}}}$ converge to $(u, \boldsymbol{\theta}) \in \overline{\mathcal{K}^{\text{ext}}}$ in $C^1(\overline{\mathcal{Q}^{\text{iter}}}) \times [0, \theta_*]^2$, and let \hat{u}_k be the solution of the boundary value problem (4.38) associated with (u_k, θ_k) . Then there exists a unique solution $\hat{u} \in C^2(\mathcal{Q}^{\text{iter}}) \cap C^1(\overline{\mathcal{Q}^{\text{iter}}} \setminus \overline{\partial_{s_0}\mathcal{Q}^{\text{iter}}}) \cap C(\overline{\mathcal{Q}^{\text{iter}}})$ of the boundary value problem (4.38) associated with (u, θ) . Moreover, \hat{u}_k converges to \hat{u} uniformly in $\overline{\mathcal{Q}^{\text{iter}}}$ and, for any $\alpha' \in [0, \alpha_1^*)$,
 - $\hat{u}_k \to \hat{u}$ in $C^{1,\alpha'}(K)$ for any compact subset $K \subseteq \overline{\mathcal{Q}^{\text{iter}}} \setminus \overline{\partial_{\text{so}}\mathcal{Q}^{\text{iter}}}$, $\hat{u}_k \to \hat{u}$ in $C^{2,\alpha'}(K)$ for any compact subset $K \subseteq \overline{\mathcal{Q}^{\text{iter}}}$.
- (iii) If $(u, \theta) \in \overline{\mathcal{K}}$, then (u, θ) satisfies Definition 4.10(vii) with nonstrict inequality in (4.25).

Remark 4.2. For a constant M > 0, define a set \mathcal{K}_M^E by

$$\mathcal{K}_{M}^{E} \coloneqq \left\{ (u, \boldsymbol{\theta}) \in C_{(*)}^{2, \alpha}(\mathcal{Q}^{\mathrm{iter}}) \times [0, \boldsymbol{\theta}_{*}]^{2} : \begin{array}{c} (u, \boldsymbol{\theta}) \text{ satisfies Definition 4.10(ii)-(v)} \\ \mathrm{and} \|u\|_{2, \alpha, \mathcal{Q}^{\mathrm{iter}}}^{(*)} \leq M \end{array} \right\}.$$

Let $\overline{\mathcal{K}_M^E}$ be the closure of \mathcal{K}_M^E in $C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2$. Then Lemma 4.15 and Corollary 4.16 still hold when $\overline{\mathcal{K}^{\text{ext}}}$ is replaced by $\overline{\mathcal{K}^E_M}$ for some M > 0.

4.2.3. Properties of the iteration set. From the well-posedness of the iteration boundary value problem (4.23), we can obtain the following a priori estimates on the iteration set. Let $\bar{\alpha} \in (0, \frac{1}{3}]$ be the constant from Proposition 4.6, and let $(\epsilon^{(w)}, \delta_1^{(w)}, N_1^{(w)})$ be the constants from Lemma 4.14.

Lemma 4.17 (A priori estimates). There exist positive constants $\alpha^{(ap)} \in (0, \frac{\bar{\alpha}}{2}), \epsilon^{(ap)} \in (0, \epsilon^{(w)}],$ $\begin{array}{l} \delta_{1}^{(\mathrm{ap})} \in (0, \delta_{1}^{(\mathrm{w})}], \ N_{1}^{(\mathrm{adm})} \geq N_{1}^{(\mathrm{w})}, \ and \ \delta_{3}^{(\mathrm{ap})} \ with \ (\alpha^{(\mathrm{ap})}, \epsilon^{(\mathrm{ap})}, \delta_{1}^{(\mathrm{ap})}) \ depending \ only \ on \ (\gamma, v_{2}, \theta_{*}, \delta_{1}), \\ N_{1}^{(\mathrm{adm})} \ depending \ only \ on \ (\gamma, v_{2}, \theta_{*}, \delta_{1}), \ and \ \delta_{3}^{(\mathrm{ap})} \ depending \ only \ on \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2}, N_{1}) \ such \ (\gamma, v_{2}, \theta_{*}, \delta_{1}, \delta_{2},$ that, whenever parameters $(\alpha, \epsilon, \delta_1, \delta_3, N_1)$ in Definition 4.10 are chosen according to

- $(\alpha, \epsilon, \delta_1) \in (0, \alpha^{(\mathrm{ap})}] \times (0, \epsilon^{(\mathrm{ap})}] \times (0, \delta_1^{(\mathrm{ap})}]$,
- $N_1 \in [N_1^{(\text{adm})}(\delta_1), \infty)$,
- $\delta_3 \in (0, \delta_3^{(ap)}(\delta_1, \delta_2, N_1)],$

with parameter $\delta_2 > 0$ to be determined later in §4.3, the following statements hold:

- (i) For any admissible solution φ corresponding to parameter $\boldsymbol{\theta} \in \Theta \cap \{\theta_1, \theta_2 \leq \theta_*\}$, function $u = u^{(\varphi, \boldsymbol{\theta})}$ given by (4.14) satisfies that $(u, \boldsymbol{\theta}) \in \mathcal{K}$.
- (ii) There exists a constant C > 0 depending only on (γ, v_2, θ_*) such that, for each $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}}$, the unique solution $\hat{\phi} \in C^2(\Omega) \cap C^1(\overline{\Omega} \setminus \Gamma_{\text{sonic}}) \cap C^0(\overline{\Omega})$ of the boundary value problem (4.23) associated with (u, θ) satisfies

(4.39)
$$\|\hat{u}\|_{2,2\alpha,\mathcal{Q}^{\text{iter}}}^{(*)} \leq C,$$

for $\hat{u}: \overline{\mathcal{Q}^{\text{iter}}} \to \mathbb{R}$ given by (4.24), whenever $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}}$ satisfies

(4.40)
$$\|u^{\sharp} - u\|_{C^{1}(\overline{\mathcal{Q}^{\text{iter}}})} + |\boldsymbol{\theta}^{\sharp} - \boldsymbol{\theta}| \le \delta^{\sharp}$$

for some $(u^{\sharp}, \theta^{\sharp}) \in \overline{\mathcal{K}}$ and some sufficiently small positive constant δ^{\sharp} depending only on $(\gamma, v_2, \theta_*, \delta_2, \delta_3, u^{\sharp}, \theta^{\sharp})$.

Proof. We describe the proof in four steps, which follow [2, Corollary 4.40, Lemma 4.42, Lemma 4.44, Corollary 4.45] closely.

1. The proof of statement (i) above follows similarly to [2, Corollary 4.40]. Indeed, let $u = u^{(\varphi, \theta)}$ be given by (4.14) for any admissible solution φ corresponding to $\theta \in \Theta \cap [0, \theta_*]^2$. Then (u, θ) automatically satisfies properties (ii)–(v) of Definition 4.10 through the choices of constants $\{N_i\}_{i=2}^5, \sigma_2, \mu_0, \mu_1, \tilde{\mu}, \rho^*(\gamma), \text{ and } C_{\text{ub}} \text{ in Definition 4.10(iii)–(v).}$

Property (i) of Definition 4.10 follows from the choice of N_0 in Definition 4.10(i) and the choice of N_1 below, in which we use an argument similar to [2, Lemma 4.39]. Indeed, by Lemma 4.15, Corollary 4.16, and Remark 4.2, for any $\alpha \in (0, \frac{\bar{\alpha}}{2}]$, we have the following continuity property of admissible solutions: For any sequence $\{\boldsymbol{\theta}_k\}_{k\in\mathbb{N}} \subseteq \Theta$ with $\boldsymbol{\theta}_k \to \mathbf{0}$ as $k \to \infty$, let $\varphi^{(\boldsymbol{\theta}_k)}$ be any admissible solution corresponding to $\boldsymbol{\theta}_k$, and let $u^{(\boldsymbol{\theta}_k)}$ be given by (4.14) with $\varphi = \varphi^{(\boldsymbol{\theta}_k)}$ and $\boldsymbol{\theta} = \boldsymbol{\theta}_k$. Then there exists a subsequence of $\{u^{(\boldsymbol{\theta}_k)}\}_{k\in\mathbb{N}}$ converging in $C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}})$ to $u^{(\text{norm})} = 0$. Using this continuity property, for any $\delta_1 \in (0, \delta_1^{(w)}]$, there exists $N_1^{(\text{adm})} \ge N_1^{(w)}$ depending on $(\gamma, v_2, \theta_*, \delta_1)$ such that $\|u - u^{(\text{norm})}\|_{2,\alpha,\mathcal{Q}^{\text{iter}}}^{(*)} < \frac{\delta_1}{2}$ whenever $\max\{\theta_1, \theta_2\} \in (0, \frac{2\delta_1}{N_1^{(\text{adm})}}]$.

Therefore, we have shown that $(u, \theta) \in \mathcal{K}^{\text{ext}}$. Finally, property (vii) of Definition 4.10 now follows from Corollary 4.16 because $(u, \theta) \in \mathcal{K}^{\text{ext}}$. We conclude that $(u, \theta) \in \mathcal{K}$.

2. It remains to prove statement (ii). Fix any $(u^{\sharp}, \theta^{\sharp}) \in \overline{\mathcal{K}}$. We claim that there exist constants $\epsilon^{(\text{lb})} \in (0, \epsilon^{(\text{w})})$ depending only on $(\gamma, v_2, \theta_*), \delta_3^{(\text{ap})} > 0$ depending only on $(\gamma, v_2, \theta_*, \delta_1, \delta_2, N_1)$, and δ^{\sharp} depending only on $(\gamma, v_2, \theta_*, \delta_2, \delta_3, u^{\sharp}, \theta^{\sharp})$ such that $\hat{\phi} - (\varphi_{\theta}^* + \frac{1}{2} |\boldsymbol{\xi}|^2) > 0$ in Ω , whenever $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}}$ satisfies (4.40). The proof of this claim follows from that for [2, Lemma 4.42], which is split into two cases: (i) $\max\{\theta_1, \theta_2\} \in [\frac{2\delta_1}{N_1^2}, \theta_*]$, and (ii) $\max\{\theta_1, \theta_2\} \in [0, \frac{2\delta_1}{N_1^2}]$. In particular, we can choose $\delta_3^{(\text{ap})} = \frac{\delta_1 \delta_2}{2N_1^2}$ and $\epsilon^{(\text{lb})} = \overline{k}^{-1} \inf_{\theta \in \overline{\Theta}} \{\hat{c}_5, \hat{c}_6\}$ for constant $\overline{k} > 1$ that has been fixed after Definition 4.4 such that (4.10) holds.

3. We obtain the *a priori* estimates for $\hat{\phi}$ near Γ_{sonic} . Let coordinates $(x, y) = \mathcal{R}(\boldsymbol{\xi})$ be given by (4.1). For j = 5, 6, we have the following estimates (with estimates (b)–(c) below only valid when $\theta_* \in (\theta^s, \theta^d)$):

(a) Whenever $\theta_{j-4} \in [0, \theta^{s})$, for each $\alpha' \in (0, 1)$, there exist positive constants $\epsilon_{p} \in (0, \epsilon_{0}]$ and $C_{\alpha'}$ depending only on $(\gamma, v_{2}, \theta_{*}, \alpha')$ with

$$\|(\hat{\phi} - \phi_j) \circ \mathcal{R}^{-1}\|_{2,\alpha',\mathcal{R}(\Omega \cap \mathcal{D}^j_{\epsilon_{\mathrm{D}}})}^{(2),(\mathrm{par})} \leq C_{\alpha'};$$

(b) There exists a constant $\delta_{\mathbf{p}} \in (0, \theta_* - \theta^{\mathbf{s}})$ depending only on (γ, v_2, θ_*) such that, whenever $\theta_{j-4} \in [\theta^{\mathbf{s}}, \theta^{\mathbf{s}} + \delta_{\mathbf{p}}]$, for each $\alpha' \in (0, 1)$, there exist positive constants $\epsilon_{\mathbf{p}} \in (0, \epsilon_0]$ depending only on (γ, v_2, θ_*) and $C_{\alpha'}$ depending only on $(\gamma, v_2, \theta_*, \alpha')$ so that

$$\|\hat{\phi} - \phi_j\|_{C^{2,\alpha'}(\Omega \cap \mathcal{D}^j_{\epsilon_{\mathrm{p}}})} \le C_{\alpha'}, \qquad D^m(\hat{\phi} - \phi_j)(P_0^{j-4}) = 0 \quad \text{for } m = 0, 1, 2;$$

(c) For $\delta_{p} > 0$ as above, there exist constants $\hat{\alpha} \in (0, \frac{1}{3}]$ and C > 0 depending only on (γ, v_2, θ_*) such that, whenever $\theta_{j-4} \in [\theta^{s} + \frac{1}{2}\delta_{p}, \theta_*]$,

$$\|\hat{\phi} - \phi_j\|_{2,\hat{\alpha},\Omega \cap \mathcal{D}^j_{\epsilon_0}}^{(-1-\hat{\alpha}),\{P_0^{j-4}\}} \le C, \qquad D^m(\hat{\phi} - \phi_j)(P_0^{j-4}) = 0 \quad \text{for } m = 0, 1$$

Using Step 2, the proof of these *a priori* estimates follows from that for [2, Lemma 4.44], whereby it is necessary to further reduce constants ϵ and δ_1 depending only on (γ, v_2, θ_*) , and we define $\alpha^{(ap)} := \frac{1}{2} \min\{\alpha_1^*, \hat{\alpha}\}$ with α_1^* given in Lemma 4.14. In particular, the estimate in (i) is similar to that for Propositions 3.12–3.13, with the free boundary replaced by a fixed boundary. The estimate in (ii) is similar to that for Proposition 3.14, with the free boundary replaced again by a fixed boundary. Finally, the estimate in (iii) is similar to that for Proposition 3.16.

4. Following the method from [2, Proposition 4.12], we are able to combine the estimates from Step **3** with the interior estimates from Lemma 4.14 to obtain (4.39). \Box

Let parameters $(\alpha, \epsilon, \delta_1, \delta_3, N_1)$ in Definition 4.10 be chosen as in Lemma 4.17. It is simple to verify that $\mathcal{K}^{\text{ext}} \subseteq C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2$ is relatively open. Indeed, the argument is similar to that for [10, Lemmas 12.8.1 and 17.5.1] and [2, Lemma 4.41]. Furthermore, following the proofs of [10, Propositions 12.8.2 and 17.5.3] and [2, Lemma 4.46], and applying the *a priori* estimates from Lemma 4.17, we conclude that the iteration set $\mathcal{K} \subseteq C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2$ is relatively open.

Proposition 4.18 (Openness of the iteration set). For the choice of parameters $(\alpha, \epsilon, \delta_1, \delta_3, N_1)$ given in Lemma 4.17, the iteration set \mathcal{K} defined by Definition 4.10 is a relatively open subset of $C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2$.

4.3. Definition of the iteration map. Fix $\theta_* \in (0, \theta^d)$. For the iteration set \mathcal{K} given by Definition 4.10, let parameters $(\alpha, \epsilon, \delta_1, \delta_3, N_1)$ be chosen as in Lemma 4.17. Define

 $\mathcal{K}(\boldsymbol{\theta}) \coloneqq \left\{ u \in C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) : (u,\boldsymbol{\theta}) \in \mathcal{K} \right\} \qquad \text{for each } \boldsymbol{\theta} \in [0,\theta_*]^2 \,,$

and similarly for $\overline{\mathcal{K}}(\boldsymbol{\theta})$. We now define an iteration map $\mathcal{I} : \overline{\mathcal{K}} \to C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}})$ satisfying the following properties:

- (a) For each $\boldsymbol{\theta} \in [0, \theta_*]^2$, there exists $u \in \mathcal{K}(\boldsymbol{\theta})$ satisfying $\mathcal{I}(u, \boldsymbol{\theta}) = u$;
- (b) If u satisfies $\mathcal{I}(u, \theta) = u$, then $\varphi^{(u, \theta)}$ defined in Definition 4.7(iii) is an admissible solution corresponding to θ .

Let $\hat{\phi} = \hat{\varphi}^{(u,\theta)} + \frac{1}{2} |\boldsymbol{\xi}|^2 \in C^2(\Omega) \cap C^1(\overline{\Omega})$ solve the iteration boundary value problem (4.23) associated with (u, θ) in Definition 4.10. Accordingly, from the definition of $\hat{u} : \overline{\mathcal{Q}^{\text{iter}}} \to \mathbb{R}$ given by (4.24), we have

$$\hat{arphi}^{(u,oldsymbol{ heta})}(oldsymbol{\xi}) = \hat{u} \circ (\mathfrak{F}_{(u,oldsymbol{ heta})})^{-1}(oldsymbol{\xi}) + arphi_{oldsymbol{ heta}}^*(oldsymbol{\xi})$$
 ,

where $\varphi_{\theta}^{*}(\boldsymbol{\xi})$ is given by Definition 4.4. We also define functions (w, w_2, \hat{w}) by

(4.41)
$$(w, w_2, \hat{w})(s, t') \coloneqq (\varphi - \varphi_{\boldsymbol{\theta}}^*, \varphi_2 - \varphi_{\boldsymbol{\theta}}^*, \hat{\varphi}^{(u, \boldsymbol{\theta})} - \varphi_{\boldsymbol{\theta}}^*) \circ (\mathcal{G}_1^{\boldsymbol{\theta}})^{-1}(s, t').$$

From (4.12) and the implicit function theorem, for any $\boldsymbol{\theta} \in \overline{\Theta}$, there exists a unique function $\mathfrak{g}_2: [-1,1] \to \overline{\mathbb{R}_+}$ satisfying $w_2(s,\mathfrak{g}_2(s)) = 0$ on [-1,1]. Then, for $Q^{\boldsymbol{\theta}}$ defined by Definition 4.1,

$$\{(s,t') : -1 < s < 1, t' = \mathfrak{g}_2(s)\} \subseteq \mathcal{G}_1^{\boldsymbol{\theta}}(Q^{\boldsymbol{\theta}}).$$

It follows from (4.11), Lemma 4.2, and Definition 4.4 that $\|\mathfrak{g}_2\|_{C^3([-1,1])} \leq C$ for some constant C > 0 depending only on (γ, v_2) .

For any $g \in C^{0,1}([-1,1])$ satisfying g(s) > 0 for all $s \in (-1,1)$, introduce the following sets: $R_{\infty} := (-1,1) \times \mathbb{R}_+$ and

(4.42)
$$R_g \coloneqq \{(s,t') \in \mathbb{R}^2_+ : -1 < s < 1, \ 0 < t' < g(s)\} \\ \Sigma_g \coloneqq \{(s,t') \in \mathbb{R}^2_+ : -1 < s < 1, \ t' = g(s)\}.$$

For each $(u, \theta) \in \overline{\mathcal{K}}$, $\mathfrak{g}_{\mathrm{sh}} = \mathfrak{g}_{\mathrm{sh}}^{(u,\theta)} : [-1,1] \to \overline{\mathbb{R}_+}$ given by Definition 4.7 is a Lipschitz function on [-1,1] and is positive on (-1,1). Then we can define sets $R_{\mathfrak{g}_{\mathrm{sh}}}$ and $\Sigma_{\mathfrak{g}_{\mathrm{sh}}}$ with $g = \mathfrak{g}_{\mathrm{sh}}$ in (4.42). Note that w and \hat{w} are defined on $R_{\mathfrak{g}_{\mathrm{sh}}}$, while w_2 is defined on R_{∞} .

The regularized distance function $\delta_{\mathfrak{g}_{sh}} \in C^{\infty}(\overline{R_{\infty}} \setminus \overline{R_{\mathfrak{g}_{sh}}})$ is given by Lemma B.4. Let $C_{rd} > 0$ be the constant from Lemma B.4, which depends only on $\text{Lip}[\mathfrak{g}_{sh}]$. We define

(4.43)
$$\delta^*_{\mathfrak{g}_{\mathrm{sh}}}(s,t') \coloneqq C_{\mathrm{rd}}^{-1}\delta_{\mathfrak{g}_{\mathrm{sh}}}(s,t') \quad \text{for any } (s,t') \in \overline{R_{\infty}} \setminus \overline{R_{\mathfrak{g}_{\mathrm{sh}}}} \,.$$

Then, for all $(s, t') \in R_{(1+\kappa)\mathfrak{g}_{sh}} \setminus \overline{R_{\mathfrak{g}_{sh}}}$ and all $\lambda \in [1, 2]$,

(4.44)
$$(s,t'-\lambda\delta^*_{\mathfrak{g}_{\mathrm{sh}}}(s,t')) \in \{s\} \times \left[\frac{\mathfrak{g}_{\mathrm{sh}}(s)}{3}, \,\mathfrak{g}_{\mathrm{sh}}(s) - (t'-\mathfrak{g}_{\mathrm{sh}}(s))\right] \Subset R_{\mathfrak{g}_{\mathrm{sh}}},$$

where constant $\kappa \in (0, \frac{1}{3}]$ depends only on Lip[\mathfrak{g}_{sh}].

Let $\{(u_k, \boldsymbol{\theta}_k)\}_{k \in \mathbb{N}} \subseteq \check{\overline{\mathcal{K}^{\text{ext}}}}$ converge to $(u, \boldsymbol{\theta})$ in $C^{2, \alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2$. Note that

$$\|\mathfrak{g}_{\mathrm{sh}}^{(u,\theta)}\|_{2,\alpha,(-1,1)}^{(-1-\alpha),\{-1,1\}} \le CN_0$$

where $N_0 > 0$ is from Definition 4.10(i), and C > 0 depends only on (γ, v_2, α) . Moreover, for \mathfrak{g}_5 and \mathfrak{g}_6 defined in Proposition 4.3(ii),

$$\frac{\mathrm{d}^m}{\mathrm{d}s^m}(\mathfrak{g}_{\mathrm{sh}}^{(u,\theta)} - \mathfrak{g}_5)(1) = \frac{\mathrm{d}^m}{\mathrm{d}s^m}(\mathfrak{g}_{\mathrm{sh}}^{(u,\theta)} - \mathfrak{g}_6)(-1) = 0 \qquad \text{for } m = 0, 1.$$

Using Lemma 4.8(iii) and (v)–(vi), we see that $\mathfrak{g}_{sh}^{(u_k, \theta_k)}$ converges to $\mathfrak{g}_{sh}^{(u, \theta)}$ in $C^{0,1}([-1, 1])$ and

$$\lim_{k \to \infty} \|\delta_{\mathfrak{g}_{\mathrm{sh}}^{(u_k, \theta_k)}} - \delta_{\mathfrak{g}_{\mathrm{sh}}^{(u, \theta)}}\|_{C^m(K)} = 0$$

for any compact set $K \subseteq \overline{R_{\infty}} \setminus \overline{R_{g_{sh}}}$ and $m \in \mathbb{N} \cup \{0\}$, which follows by Lemma B.4.

From [10, Lemma 13.9.2], there exists a function $\Psi \in C_{c}^{\infty}(\mathbb{R})$ satisfying

supp
$$\Psi \subseteq [1,2]$$
, $\int_{-\infty}^{\infty} \lambda^m \Psi(\lambda) \, d\lambda = 1 - \operatorname{sgn}(m)$ for $m = 0, 1, 2$.

Definition 4.19 (Extension operator). For each $(u, \theta) \in \overline{\mathcal{K}^{\text{ext}}}$, write $\mathfrak{g}_{\text{sh}} = \mathfrak{g}_{\text{sh}}^{(u,\theta)}$, and let $\delta_{\mathfrak{g}_{\text{sh}}}^*$ be defined by (4.43) satisfying (4.44) with constants $C_{\rm rd} > 0$ and $\kappa \in (0, \frac{1}{3}]$ depending only on (γ, v_2, θ_*) . For any $v \in C^0(\overline{R_{g_{sh}}}) \cap C^2(R_{g_{sh}} \cup \Sigma_{g_{sh}})$, define the extension of v(s, t') by

$$\mathcal{E}_{\mathfrak{g}_{\mathrm{sh}}}(v)(s,t') \coloneqq \begin{cases} v(s,t') & \text{for } (s,t') \in \overline{R}_{\mathfrak{g}_{\mathrm{sh}}}, \\ \int_{-\infty}^{\infty} v(s,t'-\lambda \delta_{\mathfrak{g}_{\mathrm{sh}}}^*(s,t')) \Psi(\lambda) \, \mathrm{d}\lambda & \text{for } (s,t') \in R_{(1+\kappa)\mathfrak{g}_{\mathrm{sh}}} \setminus \overline{R}_{\mathfrak{g}_{\mathrm{sh}}}. \end{cases}$$

For simplicity, we use the following notation: For any $a, b \in [-1, 1]$, denote

$$R_{\mathfrak{g}_{\mathrm{sh}}}^{(u,\boldsymbol{\theta})}[a,b] \coloneqq \left\{ (s,t') \in \mathbb{R}^2_+ : a < s < b, 0 < t' < \mathfrak{g}_{\mathrm{sh}}^{(u,\boldsymbol{\theta})}(s) \right\}$$

We often use the notation $R_{\mathfrak{g}_{sh}}$ to represent $R_{\mathfrak{g}_{sh}}^{(u,\theta)}$. Combining the above with Lemma 4.8(iii) and the property that the uniform bound of Lip $[\mathfrak{g}_{sh}]$ depends only on (γ, v_2, θ_*) , we obtain the following proposition, for which the details of the proof can be found in [10, Lemma 13.9.6, and Theorems 13.9.5 and 13.9.8] for each case, respectively.

Proposition 4.20 (Properties of the extension operator \mathcal{E}). Fix $\alpha \in (0,1)$. For each $(u, \theta) \in$ $\overline{\mathcal{K}^{\text{ext}}}$, let the extension operator $\mathcal{E}_{\mathfrak{g}_{\text{sh}}}: C^2(R_{\mathfrak{g}_{\text{sh}}} \cup \Sigma_{\mathfrak{g}_{\text{sh}}}) \to C^2(R_{(1+\kappa)\mathfrak{g}_{\text{sh}}})$ be given by Definition 4.19. We introduce the notation for the following three different cases:

(a) Fix any (b_1, b_2) with $-1 < b_1 < b_2 < 1$, and $C = C_{int} > 0$ depending only on $(\gamma, v_2, \theta_*, \alpha)$. For any $\alpha' \in (0, \alpha)$, denote the function spaces:

$$X \coloneqq C^{2,\alpha}(R^{(u,\boldsymbol{\theta})}_{\mathfrak{g}_{\mathrm{sh}}}[b_1, b_2]), \quad Y \coloneqq C^{2,\alpha}(R^{(u,\boldsymbol{\theta})}_{(1+\kappa)\mathfrak{g}_{\mathrm{sh}}}[b_1, b_2]), \quad Y^- \coloneqq C^{2,\alpha'}(R^{(u,\boldsymbol{\theta})}_{(1+\frac{\kappa}{2})\mathfrak{g}_{\mathrm{sh}}}[b_1, b_2]);$$

(b) Fix $\sigma > 0$ and $\epsilon \in (0, \frac{1}{4}]$. Fix $(b_1, b_2) = (-1, -1 + \epsilon)$ or $(1 - \epsilon, 1)$, and $C = C_{\text{par}} > 0$ depending only on $(\gamma, v_2, \theta_*, \alpha, \sigma)$. For any $\alpha' \in (0, \alpha)$ and $\sigma' \in (0, \sigma)$, denote the function spaces:

$$\begin{split} X &\coloneqq C^{2,\alpha}_{(\sigma),(\mathrm{par})}(R^{(u,\theta)}_{\mathfrak{g}_{\mathrm{sh}}}[b_1,b_2])\,, \qquad Y \coloneqq C^{2,\alpha}_{(\sigma),(\mathrm{par})}(R^{(u,\theta)}_{(1+\kappa)\mathfrak{g}_{\mathrm{sh}}}[b_1,b_2])\\ Y^- &\coloneqq C^{2,\alpha'}_{(\sigma'),(\mathrm{par})}(R^{(u,\theta)}_{(1+\frac{\kappa}{2})\mathfrak{g}_{\mathrm{sh}}}[b_1,b_2])\,; \end{split}$$

(c) Fix (b_1, b_2) with either $(b_1, b_2) = (b_1^{(1)}, b_2^{(1)}) = (\frac{1}{2}, 1)$ or $(b_1, b_2) = (b_1^{(2)}, b_2^{(2)}) = (-1, -\frac{1}{2})$, and $C = C_{\text{sub}} > 0$ depending only on $(\gamma, v_2, \theta_*, \alpha)$. For any $\alpha' \in (0, \alpha)$, denote the function spaces:

$$\begin{split} X &\coloneqq C^{2,\alpha}_{(-1-\alpha),\{s=(-1)^{i-1}\}} (R^{(u,\theta)}_{\mathfrak{g}_{\mathrm{sh}}}[b^{(i)}_{1}, b^{(i)}_{2}]) \,, \quad Y \coloneqq C^{2,\alpha}_{(-1-\alpha),\{s=(-1)^{i-1}\}} (R^{(u,\theta)}_{(1+\kappa)\mathfrak{g}_{\mathrm{sh}}}[b^{(i)}_{1}, b^{(i)}_{2}]) \\ Y^{-} &\coloneqq C^{2,\alpha'}_{(-1-\alpha'),\{s=(-1)^{i-1}\}} (R^{(u,\theta)}_{(1+\frac{\kappa}{2})\mathfrak{g}_{\mathrm{sh}}}[b^{(i)}_{1}, b^{(i)}_{2}]) \,. \end{split}$$

Then the extension operator $\mathcal{E}_{\mathfrak{g}_{sh}}$ satisfies the following:

(i) There exists C > 0 for each case above such that $\|\mathcal{E}_{g_{sh}}(v)\|_{Y} \leq C\|v\|_{X}$. Furthermore, for Case (c), if $(v, Dv) = (0, \mathbf{0})$ on $\overline{R_{g_{sh}}^{(u, \theta)}} \cap \{x_1 = (-1)^{i-1}\}$ for i = 1, 2, then

$$\left(\mathcal{E}_{\mathfrak{g}_{\mathrm{sh}}}(v), D\mathcal{E}_{\mathfrak{g}_{\mathrm{sh}}}(v)\right) = (0, \mathbf{0}) \qquad on \ \overline{R_{(1+\kappa)\mathfrak{g}_{\mathrm{sh}}}^{(u, \theta)}} \cap \left\{x_1 = (-1)^{i-1}\right\}.$$

- (ii) For each case, $\mathcal{E}_{g_{sh}}: X \to Y$ is linear and continuous.
- (iii) Suppose that $\{(u_k, \boldsymbol{\theta}_k)\}_{k \in \mathbb{N}} \subseteq \overline{\mathcal{K}^{\text{ext}}}$ converges to $(u, \boldsymbol{\theta})$ in $C_{(*)}^{2,\tilde{\alpha}}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2$ for some $\tilde{\alpha} \in (0, 1)$. Write X_k for the function space X with $(u, \boldsymbol{\theta})$ replaced by $(u_k, \boldsymbol{\theta}_k)$. If sequence $\{v_k\}_{k \in \mathbb{N}} \subseteq X_k$ satisfies $\|v_k\|_{X_k} \leq M$ for all $k \in \mathbb{N}$ for some constant M > 0 and $\{v_k\}_{k \in \mathbb{N}}$ converges uniformly to v on any compact set $K \in R_{\mathfrak{g}_{sh}}^{(u, \boldsymbol{\theta})}$ for some $v \in X$, then

$$\mathcal{E}_{\mathfrak{g}^{(u_k,\theta_k)}_{\mathrm{sh}}}(v_k) \to \mathcal{E}_{\mathfrak{g}^{(u,\theta)}_{\mathrm{sh}}}(v) \qquad in \; Y^- \,,$$

where
$$\mathcal{E}_{\mathfrak{g}_{sh}^{(u_k,\theta_k)}}(v_k)$$
 is well defined on $R_{(1+\frac{\kappa}{2})\mathfrak{g}_{sh}}^{(u,\theta)}[b_1,b_2]$ for large enough k.

The proof of the following result is similar to [2, Lemma 5.5].

Lemma 4.21. Let parameters $(\alpha, \epsilon, \delta_1, \delta_3, N_1)$ in Definition 4.10 be chosen as in Lemma 4.17, with $\delta_2 > 0$ to be specified later. Then there exists a constant $\delta_3^{(imp)} > 0$ depending only on $(\gamma, v_2, \theta_*, \delta_2)$ such that, if $\delta_3 \in (0, \delta_3^{(imp)})$, then, for each $(u, \theta) \in \overline{\mathcal{K}}$, there is a unique function $\hat{\mathfrak{g}}_{sh} : [-1, 1] \to \overline{\mathbb{R}_+}$ such that

(4.45)
$$\left(w_2 - \mathcal{E}_{\mathfrak{g}_{\mathrm{sh}}^{(u,\theta)}}(\hat{w})\right)(s,\hat{\mathfrak{g}}_{\mathrm{sh}}(s)) = 0 \quad \text{for all } s \in [-1,1].$$

Furthermore, there exists a constant C > 0 depending only on (γ, v_2, θ_*) such that $\hat{\mathfrak{g}}_{sh}$ satisfies

(4.46)
$$\begin{cases} \|\hat{\mathfrak{g}}_{\rm sh} - \mathfrak{g}_2\|_{2,2\alpha,(-1,1)}^{(-1-2\alpha),\{-1,1\}} \leq C, & \|\hat{\mathfrak{g}}_{\rm sh} - \mathfrak{g}_{\rm sh}^{(u,\theta)}\|_{2,\frac{\alpha}{2},(-1,1)} \leq C\delta_3, \\ \frac{\mathrm{d}^m}{\mathrm{d}s^m}(\hat{\mathfrak{g}}_{\rm sh} - \mathfrak{g}_2)(\pm 1) = \frac{\mathrm{d}^m}{\mathrm{d}s^m}(\hat{\mathfrak{g}}_{\rm sh} - \mathfrak{g}_{\rm sh}^{(u,\theta)})(\pm 1) = 0 & \text{for } m = 0, 1. \end{cases}$$

Definition 4.22 (Iteration map). Let parameters $(\alpha, \epsilon, \delta_1, \delta_3, N_1)$ in Definition 4.10 be chosen as in Lemma 4.17, with $\delta_2 > 0$ to be specified later. Furthermore, let $\delta_3 \in (0, \delta_3^{(imp)})$ for constant $\delta_3^{(imp)} > 0$ from Lemma 4.21. For each $(u, \theta) \in \overline{\mathcal{K}}$, let $\tilde{u} : \overline{\mathcal{Q}^{iter}} \to \mathbb{R}$ be given by

$$\tilde{u} \coloneqq \mathcal{E}_{\mathfrak{g}_{\mathrm{sh}}^{(u,\theta)}}(\hat{w}) \circ (G_{2,\hat{\mathfrak{g}}_{\mathrm{sh}}})^{-1},$$

where $G_{2,\hat{\mathfrak{g}}_{sh}}$ is given by (4.13) with $\hat{\mathfrak{g}}_{sh}$: $[-1,1] \to \mathbb{R}_+$ from (4.45). Then define the iteration map $\mathcal{I}: \overline{\mathcal{K}} \to C^{2,\alpha}_{(*)}(\mathcal{Q}^{iter})$ by

$$\mathcal{I}(u,\boldsymbol{\theta}) = \tilde{u}$$

Note that map $\mathcal{I}: \overline{\mathcal{K}} \to C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}})$ defined as above is reasonable, since

(4.47)
$$\|\mathcal{I}(u,\boldsymbol{\theta})\|_{2,2\alpha,\mathcal{Q}^{\text{iter}}}^{(*)} \leq C \quad \text{for all } (u,\boldsymbol{\theta}) \in \mathcal{K}$$

for C > 0 depending only on (γ, v_2, θ_*) , which follows from Propositions 4.18 and 4.20, and Lemma 4.21.

4.4. Proof of Theorem 2.1: Existence of admissible solutions. Fix $\theta_* \in (0, \theta^d)$. For the iteration map \mathcal{I} defined in Definition 4.22, we now prove that, for any fixed point $u \in \overline{\mathcal{K}}(\theta)$ of $\mathcal{I}(\cdot, \boldsymbol{\theta})$ for some $\boldsymbol{\theta} \in \Theta \cap [0, \theta_*]^2$, then $\varphi = \varphi^{(u, \boldsymbol{\theta})}$ defined in Definition 4.7(iii) is an admissible solution corresponding to $\boldsymbol{\theta}$ in the sense of Definition 2.11.

Proposition 4.23 (Fixed points of map $\mathcal{I}(\cdot, \boldsymbol{\theta})$). Let parameters $(\alpha, \epsilon, \delta_1, \delta_3, N_1)$ in Definition 4.10 be chosen as in Definition 4.22. Then parameters (ϵ, δ_1) can be chosen even smaller, depending only on (γ, v_2, θ_*) , such that, for each $\boldsymbol{\theta} \in \Theta \cap [0, \theta_*]^2$, $u \in \mathcal{K}(\boldsymbol{\theta})$ is a fixed point of $\mathcal{I}(\cdot,\boldsymbol{\theta}):\overline{\mathcal{K}}(\boldsymbol{\theta})\to C^{2,\alpha}_{(*)}(\mathcal{Q}^{\mathrm{iter}}) \text{ if and only if } \varphi=\varphi^{(u,\boldsymbol{\theta})} \text{ defined in Definition 4.7(iii) is an admissible}$ solution corresponding to $\boldsymbol{\theta}$ in the sense of Definition 2.11 after extending φ to \mathbb{R}^2_+ via (2.41).

Proof. Since the proof is similar to that for [2, Proposition 5.8], we omit the details of the proof here and sketch only the main ideas in the following four steps.

1. For $\theta \in \Theta \cap [0, \theta_*]^2$, let φ be an admissible solution corresponding to θ , and let u := $u^{(\varphi,\theta)}$ be given by (4.14), which satisfies $u \in \mathcal{K}(\theta)$ by Lemma 4.17(i). It follows directly from Definition 4.7(i) and the boundary condition on Γ_{shock} that $\mathfrak{g}_{\text{sh}}^{(u,\theta)} = \mathfrak{g}_{\text{sh}}$. It is clear that $\hat{\phi} :=$ $\phi = \varphi + \frac{1}{2} |\boldsymbol{\xi}|^2$ solves the iteration boundary value problem (4.23), which indicates that $\hat{w} = w$ in (4.41). Then $\hat{\mathfrak{g}}_{sh} = \mathfrak{g}_{sh}$ by solving (4.45). We conclude that

$$\mathcal{I}(u,\boldsymbol{\theta}) = \mathcal{E}_{\mathfrak{g}_{\mathrm{sh}}^{(u,\boldsymbol{\theta})}}(\hat{w}) \circ (G_{2,\hat{\mathfrak{g}}_{\mathrm{sh}}})^{-1} = \mathcal{E}_{\mathfrak{g}_{\mathrm{sh}}}(w) \circ (G_{2,\mathfrak{g}_{\mathrm{sh}}})^{-1} = u\,,$$

so $u \in \mathcal{K}(\boldsymbol{\theta})$ is a fixed point of $\mathcal{I}(\cdot, \boldsymbol{\theta})$.

2. On the other hand, for any fixed point u of map $\mathcal{I}(\cdot, \boldsymbol{\theta}) : \overline{\mathcal{K}}(\boldsymbol{\theta}) \to C^{2,\alpha}_{(*)}(\mathcal{Q}^{\mathrm{iter}})$, let $\varphi^{(u,\boldsymbol{\theta})}$ be given by Definition 4.7(iii). We now show that $\varphi^{(u,\theta)}$ is an admissible solution in the sense of Definition 2.11 corresponding to θ . Using Definition 4.7(i), (4.45), and Definition 4.22, we have

$$w_2(s, \mathfrak{g}_{\mathrm{sh}}^{(u,\theta)}) = u(s, 1) = \mathcal{E}_{\mathfrak{g}_{\mathrm{sh}}^{(u,\theta)}}(\hat{w})(s, \hat{\mathfrak{g}}_{\mathrm{sh}}) = w_2(s, \hat{\mathfrak{g}}_{\mathrm{sh}}),$$

from which we obtain that $\hat{\mathfrak{g}}_{\rm sh} = \mathfrak{g}_{\rm sh}^{(u,\theta)}, \, \hat{u} = u$, and $\hat{\varphi}^{(u,\theta)} = \varphi^{(u,\theta)}$.

Define $v \coloneqq \hat{\varphi}^{(u,\theta)} - \varphi_2$. Then v solves the following strictly elliptic equation:

$$\mathcal{L}_{(u,\boldsymbol{\theta})}(v) = \sum_{i,j=1}^{2} A_{ij}(D\hat{\phi}(\boldsymbol{\xi}), \boldsymbol{\xi})\partial_{i}\partial_{j}v = 0.$$

From the boundary conditions of the iteration boundary value problem (4.23) and $\hat{u} = u$ above, we see that $v \leq 0$ on $\Gamma_{\text{shock}} \cup \Gamma_{\text{sonic}}$ and $\partial_{\eta} v = -v_2 > 0$ on Γ_{sym} . From the maximum principle and Hopf's lemma, we obtain that $\hat{\varphi}^{(u,\theta)} \leq \varphi_2$ on $\overline{\Omega}$. When $\max\{\theta_1, \theta_2\} \in [\frac{2\delta_1}{N_1^2}, \theta_*]$, we see that $\hat{\varphi}^{(u,\theta)} \geq \max\{\varphi_5, \varphi_6\}$ in Ω , which follows directly

from (4.21) in Definition 4.10 and Step 1 of Lemma 4.17.

When $\max\{\theta_1, \theta_2\} \in (0, \frac{2\delta_1}{N_1^2}]$, we can show the same result by applying the maximum principle as in the proof of [2, Lemma 4.42]. Then inequality (2.42) in Definition 2.11(iv) is proved.

Similarly, for inequality (2.43) in Definition 2.11(iv), we also split into two cases. Denote $w \coloneqq \partial_{\boldsymbol{e}_{S_{25}}}(\varphi_2 - \varphi^{(u,\boldsymbol{\theta})})$. We introduce (X_1, X_2) -coordinates by $\boldsymbol{\xi} \coloneqq X_1 \boldsymbol{e}_{S_{25}} + X_2 \boldsymbol{e}_{S_{25}}^{\perp}$. From the equation for v, one can derive the following strictly elliptic equation for $w = -\partial_{X_1} v$:

$$\hat{\mathcal{L}}_{(u,\boldsymbol{\theta})}(w) \coloneqq \sum_{i,j=1}^{2} \hat{A}_{ij} \partial_{X_i X_j} w + \sum_{i=1}^{2} \hat{A}_i \partial_{X_i} w = 0 \qquad \text{in } \Omega$$

When $\max\{\theta_1, \theta_2\} \in [\frac{2\delta_1}{N_1^2}, \theta_*]$, it follows from (4.21) in Definition 4.10 that w < 0 in $\overline{\Omega} \setminus \mathcal{D}_{\epsilon/10}^5$. In domain $\Omega \cap \mathcal{D}^5_{\epsilon}$, one can check

(4.48)
$$\begin{cases} w = 0 & \text{on } \Gamma_{\text{sonic}}^5, \\ \boldsymbol{b}_{w} \cdot Dw = 0 & \text{with } \boldsymbol{b}_{w} \cdot \boldsymbol{\nu} > 0 \text{ on } \Gamma_{\text{sym}}, \\ \boldsymbol{b}_{\text{sh}} \cdot Dw = 0 & \text{with } \boldsymbol{b}_{\text{sh}} \cdot \boldsymbol{\nu} > 0 \text{ on } \Gamma_{\text{shock}}, \end{cases}$$

where $\boldsymbol{\nu}$ is the unit normal vector on $\Gamma_{\text{sym}} \cup \Gamma_{\text{shock}}$ pointing to Ω . Note that we have used [10, Lemma 13.4.5] to derive the boundary condition on Γ_{shock} . By the maximum principle and Hopf's lemma, we obtain that $w \leq 0$ in $\Omega \cap \mathcal{D}_{\epsilon}^{5}$ when $\epsilon \in (0, \epsilon_{\text{sh}}]$, for some sufficiently small $\epsilon_{\text{sh}} > 0$ depending only on (γ, v_2, θ_*) .

When $\max\{\theta_1, \theta_2\} \in (0, \frac{2\delta_1}{N_1^2}]$, using property (i) of Definition 4.10, one can follow the same procedure as the first case to derive that $w \leq 0$ in Ω when $\delta_1 \in (0, \delta_{\rm sh}]$, for some $\delta_{\rm sh} > 0$ depending only on (γ, v_2, θ_*) . Therefore, $\hat{\varphi}^{(u,\theta)}$ satisfies (2.43) when $\epsilon \in (0, \epsilon_{\rm sh}]$ and $\delta_1 \in (0, \delta_{\rm sh}]$, since the argument above also works for $\partial_{\boldsymbol{e}_{S_{26}}}(\varphi_2 - \varphi^{(u,\theta)})$.

3. Now we check that $\mathcal{N}_{(u,\theta)}(\phi) = 0$ in Definition 4.10(vii) coincides with (3.1) in Ω . From Lemma 4.12(iii), this property holds in $\Omega \setminus \mathcal{D}_{\epsilon/10}$, so it remains to show that this property holds in $\Omega \cap \mathcal{D}_{\epsilon/2}$. We only consider domain $\Omega \cap \mathcal{D}_{\epsilon/2}^5$, since the proof is the same for domain $\Omega \cap \mathcal{D}_{\epsilon/2}^6$ due to the symmetry between θ_1 and θ_2 in the problem.

For any $\boldsymbol{\xi} \in \Omega \cap \mathcal{D}_{\epsilon}$, let coordinates $(x, y) = \mathcal{R}(\boldsymbol{\xi})$ be given by (4.1). Using the monotone property (4.20), we choose a constant $\epsilon'_{\rm sh} \in (0, \epsilon_{\rm sh}]$ sufficiently small, depending only on (γ, v_2, θ_*) , such that $\theta_1 \leq \theta^{\rm s} + \min\{\frac{\sigma_3}{2}, \delta_{\rm p}\}$ whenever $x_{P_0^1} < \frac{\epsilon'_{\rm sh}}{10}$, for $\delta_{\rm p} > 0$ the constant from Step **3**(b) in the proof of Lemma 4.17.

Choose $\epsilon \in (0, \epsilon'_{\text{sh}}]$, and suppose $x_{P_0^1} < \frac{\epsilon}{10}$. Denote $\tilde{w} \coloneqq Ax - \partial_x \psi(x, y)$ for $A \coloneqq \frac{1}{1+\gamma}(2 - \frac{\mu_0}{5})$ and $\psi = (\varphi^{(u,\theta)} - \varphi_5) \circ \mathcal{R}^{-1}$. It is clear that $\tilde{w} = 0$ on $\mathcal{R}(\Gamma^5_{\text{sonic}})$, and $\tilde{w}_y = 0$ on $\mathcal{R}(\Gamma_{\text{sym}} \cap \partial \mathcal{D}^5_{\epsilon/2})$. From Lemma 4.13(i) and (iii)–(iv), the boundary condition: $\mathcal{M}_{(u,\theta)}(D\hat{\phi}, \hat{\phi}, \boldsymbol{\xi}) = 0$ in (4.23) can be written as

$$(b_1, b_2, b_0) \cdot (D\hat{\phi}, \hat{\phi}) = 0$$
 on $\Gamma_{\text{shock}} \cap \mathcal{D}^5_{\epsilon}$,

for some $b_0, b_1, b_2 \in [-\frac{1}{\delta_{\rm bc}}, -\delta_{\rm bc}]$. Using the *a priori* estimates (4.39) in the case: $\theta_1 \leq \theta^{\rm s} + \delta_{\rm p}$, we see that $|\psi_x| \leq \frac{1}{\delta_{\rm bc}^2}(|\psi_y| + |\psi|) \leq Cx^{\frac{3}{2}}$ for any $(x, y) \in \mathcal{R}(\Gamma_{\rm shock} \cap \mathcal{D}_{\epsilon}^5)$. Reducing $\epsilon'_{\rm sh} > 0$ further, depending only on (γ, v_2) , we obtain that $\tilde{w} > 0$ on $\mathcal{R}(\Gamma_{\rm shock} \cap \mathcal{D}_{\epsilon}^5)$ for any $\epsilon \in (0, \epsilon'_{\rm sh}]$.

By Lemma 4.12(ii) and the fact that $\hat{\phi} = \phi$, we have

$$\sum_{i,j=1}^{2} A_{ij}(D\hat{\phi},\boldsymbol{\xi})\partial_{i}\partial_{j}\hat{\phi} = \sum_{i,j=1}^{2} A_{ij}^{(3)}(D\hat{\phi},\boldsymbol{\xi})\partial_{i}\partial_{j}\hat{\phi} = c_{\boldsymbol{\theta}}\mathcal{N}_{(u,\boldsymbol{\theta})}^{\text{polar}}(\hat{\phi}),$$

from where we can derive the following equation for \tilde{w} :

$$a_{11}\tilde{w}_{xx} + 2a_{12}\tilde{w}_{xy} + a_{22}\tilde{w}_{yy} + a_1\tilde{w}_x + a_2\tilde{w}_y = -A((\gamma+1)A - 1) + E(x,y) \quad \text{in } \mathcal{R}(\Omega \cap \mathcal{D}^5_{\epsilon/2}).$$

From expression (3.26) and the *a priori* estimates (4.39) in Lemma 4.17, reducing $\epsilon'_{\rm sh} > 0$ further if necessary, we obtain directly that $E(x, y) < A((\gamma + 1)A - 1)$ in $\mathcal{R}(\Omega \cap \mathcal{D}^5_{\epsilon/2})$ for any $\epsilon \in (0, \epsilon'_{\rm sh}]$. Therefore, the property: $\tilde{w} = Ax - \partial_x \psi(x, y) \ge 0$ holds in $\mathcal{R}(\Omega \cap \mathcal{D}^5_{\epsilon/2})$, which follows from Lemma 4.12(i), the maximum principle, and Hopf's lemma.

Now we show the inequality: $\partial_x \psi(x, y) \ge -Ax$. From Step 2, we have

$$\left(\partial_{e_{S_{25}}}(\varphi_2-\varphi)\right)\circ\mathcal{R}^{-1}=\psi_x\cos(\theta_{25}-y)-\frac{\sin(\theta_{25}-y)}{c_5-x}\psi_y\leq 0\qquad\text{in }\mathcal{R}(\Omega\cap\mathcal{D}^5_{\epsilon/2})\,.$$

This implies that

$$\psi_x \ge -\frac{\tan(\frac{\pi}{2} - \tilde{\omega}_0)}{c_5 - x} \psi_y \ge -Cx^{\frac{3}{2}} \quad \text{in } \mathcal{R}(\Omega \cap \mathcal{D}_{\epsilon/2}^5) \,,$$

by using property (4.3) and the *a priori* estimates (4.39) in Lemma 4.17. Then $\partial_x \psi(x, y) \ge -Ax$ after further reducing $\epsilon'_{\rm sh} > 0$. Thus, $|\psi_x| \le Ax$ is proved, which indicates that $\mathcal{N}_{(u,\theta)}(\phi) = 0$ in Definition 4.10(vii) coincides with (3.1) in $\Omega \cap \mathcal{D}^5_{\epsilon/2}$. Moreover, if $x_{P_0^1} \ge \frac{\epsilon}{10}$, the same conclusion holds in $\Omega \cap \mathcal{D}^5_{\epsilon/2} \subseteq \Omega \setminus \mathcal{D}^6_{\epsilon/10}$ by Lemma 4.12(iii).

4. Definition 2.11(iii) follows directly from Lemma 4.12(i) and the fact proved in Step **3**. Let the sound speed $c(|D\varphi|, \varphi)$ be given by (3.2). The strict ellipticity implies that

$$M \coloneqq \frac{|\partial_{\boldsymbol{\nu}} \varphi(\boldsymbol{\xi})|}{c(|D\varphi(\boldsymbol{\xi})|, \varphi(\boldsymbol{\xi}))} < 1 \quad \text{on } \Gamma_{\text{shock}} \,.$$

Define $M_2 := |\partial_{\nu} \varphi_2(\boldsymbol{\xi})|$. Similar to [2, Eq. (2.4.9)], we have the following equality:

$$(1 + \frac{\gamma - 1}{2}M^2)M^{-\frac{2(\gamma - 1)}{\gamma + 1}} = (1 + \frac{\gamma - 1}{2}M_2^2)M_2^{-\frac{2(\gamma - 1)}{\gamma + 1}}$$

Then $M_2 > 1$, and so $|D\varphi_2| > 1 = c(|D\varphi_2|, \varphi_2)$ holds on Γ_{shock} , *i.e.*, $\Gamma_{\text{shock}} \subseteq \mathbb{R}^2_+ \setminus \overline{B_{c_2}(O_2)}$. Finally, for the property: $\Gamma_{\text{shock}} \subseteq \{\xi^{P_3} < \xi < \xi^{P_2}\}$, we follow the same proof as in Lemma 3.2 by using all the properties above. Therefore, all parts of Definition 2.11 have been verified, which implies that $\varphi^{(u,\theta)}$ is an admissible solution in the sense of Definition 2.11.

Proof of Theorem 2.1. We now complete the proof of Theorem 2.1 in five steps.

1. We prove that the iteration map $\mathcal{I}: \overline{\mathcal{K}} \to C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}})$ is a compact map in the sense of Definition B.5. Let $\{(u_k, \theta_k)\}_{k \in \mathbb{N}} \subseteq \mathcal{K}$ converge to (u, θ) in $C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}) \times [0, \theta_*]^2$. For each $k \in \mathbb{N}$, write $\Omega^{(k)} \coloneqq \Omega(u_k, \theta_k)$ and $\mathfrak{g}^{(k)}_{\text{sh}} \coloneqq \mathfrak{g}^{(u_k, \theta_k)}_{\text{sh}}$, as given by Definition 4.7, and $\Gamma^{(k)}_{\text{sonic}} \coloneqq \Gamma^5_{\text{sonic}} \cup \Gamma^6_{\text{sonic}}$ for the sonic arcs related to parameters θ_k . By Lemma 4.14, there exists a unique solution $\hat{\phi}^{(k)} \in C^2(\Omega^{(k)}) \cap C^1(\overline{\Omega^{(k)}} \setminus \Gamma^{(k)}_{\text{sonic}}) \cap C^0(\overline{\Omega^{(k)}})$ of the iteration boundary value problem (4.23) related to (u_k, θ_k) . For any $(s, t') \in R_{\mathfrak{g}^{(k)}_{\text{sh}}}$, define

$$\hat{w}^{(k)}(s,t') \coloneqq \left(\hat{\phi}^{(k)} + \frac{1}{2}|\boldsymbol{\xi}|^2 - \varphi^*_{\boldsymbol{\theta}_k}\right) \circ (\mathcal{G}_1^{\boldsymbol{\theta}_k})^{-1}(s,t'),$$

where $R_{\mathfrak{g}_{sh}^{(k)}}$, $\varphi_{\theta_k}^*$, and $\mathcal{G}_1^{\theta_k}$ are given by (4.42), Definition 4.4, and (4.7), respectively. Then

$$\hat{w}^{(k)} = \hat{u}^{(k)} \circ G_{2,\mathfrak{g}_{\mathrm{sb}}^{(k)}}$$

for $\hat{u}^{(k)}: \overline{\mathcal{Q}^{\text{iter}}} \to \mathbb{R}$ given by (4.24) with $\hat{\phi} = \hat{\phi}^{(k)}$, and $G_{2,\mathfrak{g}_{\text{sh}}^{(k)}}$ given by (4.13). Subsequently, we obtain functions $\hat{\mathfrak{g}}_{\text{sh}}^{(k)}: [-1,1] \to \overline{\mathbb{R}_+}$ via Lemma 4.21.

Since $(u_k, \boldsymbol{\theta}_k)$ converges to $(u, \boldsymbol{\theta})$ in $C^{1,\alpha}(\mathcal{Q}^{\text{iter}}) \times [0, \boldsymbol{\theta}_*]^2$, we find that $\mathfrak{g}_{\text{sh}}^{(k)} \to \mathfrak{g}_{\text{sh}}$ in $C^{1,\alpha}([-1, 1])$ by Lemma 4.8(iii). Fix any compact subset $K \subseteq R_{\mathfrak{g}_{\text{sh}}}$. From (4.13), Definition 4.10(iii), and the convergence of $\{\mathfrak{g}_{\text{sh}}^{(k)}\}_{k\in\mathbb{N}}$, we see that $G_{2,\mathfrak{g}_{\text{sh}}^{(k)}} \to G_{2,\mathfrak{g}_{\text{sh}}}$ in $C^{1,\alpha}(K)$. Then, by Corollary 4.16 and the convergence of $\{\hat{u}^{(k)}\}_{k\in\mathbb{N}}$, we obtain that $\hat{w}^{(k)} \to \hat{w}$ in $C^{1,\alpha}(K)$.

By Lemma 4.17, Proposition 4.20(iii), and the local convergence of $\{\hat{w}^{(k)}\}_{k\in\mathbb{N}}$, for any $b_1, b_2 \in [-1, 1]$ chosen as in each case of Proposition 4.20(a)–(c), $\{\mathcal{E}_{\mathfrak{g}_{sh}^{(k)}}(\hat{w}^{(k)})\}_{k\in\mathbb{N}}$ converges to $\mathcal{E}_{\mathfrak{g}_{sh}}(\hat{w})$ in the corresponding function space. Combining the statement above with the convergence of $\{\hat{\mathfrak{g}}_{sh}^{(k)}\}_{k\in\mathbb{N}}$ from Lemma 4.21, we conclude that

$$\mathcal{E}_{\mathfrak{g}_{\mathrm{sh}}^{(k)}}(\hat{w}^{(k)}) \circ (G_{2,\hat{\mathfrak{g}}_{\mathrm{sh}}^{(k)}})^{-1} \to \mathcal{E}_{\mathfrak{g}_{\mathrm{sh}}}(\hat{w}) \circ (G_{2,\hat{\mathfrak{g}}_{\mathrm{sh}}})^{-1} \qquad \text{in } C^{2,\alpha}_{(*)}(\mathcal{Q}^{\mathrm{iter}}) \,.$$

Since $C_{(*)}^{2,2\alpha}(\mathcal{Q}^{\text{iter}})$ is compactly embedded into $C_{(*)}^{2,\alpha}(\mathcal{Q}^{\text{iter}})$, we obtain from (4.47) that $\mathcal{I}(U)$ is precompact in $C_{(*)}^{2,\alpha}(\mathcal{Q}^{\text{iter}})$ for any bounded subset $U \subseteq \overline{\mathcal{K}}$ so that the iteration map \mathcal{I} is compact.

2. Next, we show that constants $N_1, \delta_2 > 0$ in Definition 4.10 can be chosen, depending only on (γ, v_2, θ_*) , such that, for any $\boldsymbol{\theta} \in \Theta \cap [0, \theta_*]^2$, no fixed point of $\mathcal{I}(\cdot, \boldsymbol{\theta})$ lies on boundary $\partial \mathcal{K}(\boldsymbol{\theta})$, where $\partial \mathcal{K}(\boldsymbol{\theta})$ is considered relative to $C_{(*)}^{2,\alpha}(\mathcal{Q}^{\text{iter}})$.

Suppose that $(u, \theta) \in \overline{\mathcal{K}}$ satisfies $\mathcal{I}(u, \theta) = u$. For $\varphi = \varphi^{(u, \theta)}$ defined in $\Omega = \Omega(u, \theta)$ by Definition 4.7(iii), extend φ to \mathbb{R}^2_+ via (2.41). Then φ is an admissible solution corresponding to parameters $\theta \in \Theta \cap [0, \theta_*]^2$ by Proposition 4.23.

From (3.17), Propositions 3.4, 3.8, and 4.3(iv), and Lemma 3.3, constants $(N_2, \tilde{\mu}, \rho^*(\gamma), C_{ub})$ are fixed so that any admissible solution satisfies the strict inequalities in Definition 4.10(ii) and (vi)–(v). Moreover, from Lemma 4.17(i), u satisfies the strict inequality given in Definition 4.10(i) when $N_1 \geq N_1^{(adm)}$.

For condition (iv) of Definition 4.10, constants $(\mu_0, \mu_1, N_4, N_5, \sigma_2)$ are fixed such that any admissible solution corresponding to $\boldsymbol{\theta} \in \Theta \cap [0, \theta_*]^2$ satisfies the strict inequalities (4.22). Next, we focus on the strict inequalities (4.21). It is direct to see that the strict inequalities (4.21) hold

when $\max\{\theta_1, \theta_2\} < \frac{\delta_1}{N_1^2}$, since $\mathscr{K}_2(\max\{\theta_1, \theta_2\}) < 0$ and inequality (2.42) in Definition 2.11(iv) and Lemma 3.1(ii) hold for any admissible solution corresponding to $\boldsymbol{\theta} \in \Theta \cap [0, \theta_*]^2$. When $\max\{\theta_1, \theta_2\} \geq \frac{\delta_1}{N_1^2}$, we similarly have

$$\begin{aligned} \varphi - \max\{\varphi_5, \varphi_6\} > 0 & \text{ in } \overline{\Omega} \setminus (\mathcal{D}^5_{\epsilon/10} \cup \mathcal{D}^6_{\epsilon/10}) \,, \\ \partial_{e_{S_{2i}}}(\varphi_2 - \varphi) < 0 & \text{ in } \overline{\Omega} \setminus \mathcal{D}^j_{\epsilon/10} \text{ for } j = 5, 6 \,. \end{aligned}$$

By Corollary 3.10, and Propositions 3.12–3.14 and 3.16, we obtain that $\varphi \in C^{1,\alpha}(\Omega)$. Combining this with Lemma 3.17, there exist both $\tau_0 > 0$ depending only on (γ, v_2, θ_*) and $\tau_1 > 0$ depending only on $(\gamma, v_2, \theta_*, \delta_1, N_1)$ such that, for all $\boldsymbol{\theta} \in \Theta \cap \{\frac{\delta_1}{N_1^2} \leq \max\{\theta_1, \theta_2\} \leq \theta_*\}$,

$$\psi = \varphi - \max\{\varphi_5, \varphi_6\} \ge \tau_0 \qquad \text{in } \overline{\Omega} \setminus (\mathcal{D}^5_{\epsilon/10} \cup \mathcal{D}^6_{\epsilon/10}),$$
$$\partial_{e_{S_{2,i}}}(\varphi_2 - \varphi) \le -\tau_1 \qquad \text{in } \overline{\Omega} \setminus \mathcal{D}^j_{\epsilon/10} \text{ for } j = 5, 6.$$

For any fixed $\delta_1 > 0$, we can choose $\delta_2 > 0$ small enough such that

$$\mathscr{K}_{2}(\max\{\theta_{1},\theta_{2}\}) \leq \frac{\delta_{1}\delta_{2}}{N_{1}^{2}} < \min\{\tau_{0},\tau_{1}\} \qquad \text{for all } \boldsymbol{\theta} \in \Theta \cap \{\frac{\delta_{1}}{N_{1}^{2}} \leq \max\{\theta_{1},\theta_{2}\} \leq \theta_{*}\},$$

which implies that the strict inequalities (4.21) hold when $\max\{\theta_1, \theta_2\} \geq \frac{\delta_1}{N^2}$.

Finally, let $(\alpha, \epsilon, \delta_1, \delta_3, N_1)$ be chosen as in Lemma 4.17 with δ_2 chosen as above. Then the strict inequality (4.25) of Definition 4.10(vii) holds by Proposition 4.18. Therefore, $u \in \mathcal{K}(\boldsymbol{\theta})$ for any $\boldsymbol{\theta} \in \Theta \cap [0, \theta_*]^2$, which completes this step.

3. Now, we can fix constants $(\alpha, \epsilon, \delta_1, \delta_2, \delta_3, N_1)$ in Definition 4.10 as follows:

- Choose constants $(\alpha, \epsilon, \delta_1, \delta_3, N_1)$ as in Lemma 4.17 and reduce (ϵ, δ_1) as in Proposition 4.23, depending only on (γ, v_2, θ_*) ;
- Fix δ_2 as in Step 2, depending only on $(\gamma, v_2, \theta_*, \delta_1, N_1)$;
- Adjust $\delta_3 > 0$ as in Definition 4.22, depending only on $(\gamma, v_2, \theta_*, \delta_2)$.

4. Leray-Schauder degree theory. When $\boldsymbol{\theta} = \mathbf{0}$, $\varphi_5 = \varphi_6 = \varphi_0$ with φ_0 defined in §2.1.3, and the iteration boundary value problem in the (s, t')-coordinates is a homogeneous problem. From Lemma 4.14, the iteration boundary value problem (4.23) corresponding to any $(u, \mathbf{0}) \in \overline{\mathcal{K}}$ has a unique solution $\hat{\phi} = \frac{1}{2} |\boldsymbol{\xi}|^2 + \varphi_{\mathbf{0}}^* = v_2 \eta_0$, where $\eta_0 > 0$ is defined in §2.1.3. It follows from Definition 4.10 that the unique fixed point $u \equiv 0$ of $\mathcal{I}(\cdot, \mathbf{0})$ lies in $\mathcal{K}(\mathbf{0})$, and $\operatorname{Ind}(\mathcal{I}(\cdot, \mathbf{0}), \mathcal{K}(\mathbf{0})) = 1$. From the arguments above, map $\mathcal{I}(\cdot, \boldsymbol{\theta}) : \overline{\mathcal{K}(\boldsymbol{\theta})} \to C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}})$ satisfies $\mathcal{I}(\cdot, \boldsymbol{\theta}) \in \mathcal{V}(\mathcal{K}(\boldsymbol{\theta}), C^{2,\alpha}_{(*)}(\mathcal{Q}^{\text{iter}}))$ from Definition B.6 in Appendix B. From Theorem B.2 and for any $t \in [0, 1]$, we have

$$\mathbf{Ind}(\mathcal{I}(\cdot, t\boldsymbol{\theta}), \mathcal{K}(t\boldsymbol{\theta})) = \mathbf{Ind}(\mathcal{I}(\cdot, \mathbf{0}), \mathcal{K}(\mathbf{0})) \quad \text{for all } \boldsymbol{\theta} \in [0, \theta_*]^2.$$

We obtain the existence of a fixed point of map $\mathcal{I}(\cdot, \boldsymbol{\theta})$, which is equivalent to the existence of an admissible solution corresponding to $\boldsymbol{\theta} \in [0, \theta_*]^2$ in the sense of Definition 2.11 by Proposition 4.23. The arbitrary choice of $\theta_* \in (0, \theta^d)$ completes the proof of the existence of admissible solutions for any $\boldsymbol{\theta} \in [0, \theta^d)^2$.

5. Proof of the continuity at $\boldsymbol{\theta} = \mathbf{0}$. Let φ_{norm} be the admissible solution for $\boldsymbol{\theta} = \mathbf{0}$, which is given by (2.24), and let $\Omega^{(0)}$ be the corresponding pseudo-subsonic domain. For any sequence $\{\boldsymbol{\theta}_k\}_{k\in\mathbb{N}} \subseteq \Theta$ with $\boldsymbol{\theta}_k \to \mathbf{0}$ as $k \to \infty$, let $\varphi^{(\boldsymbol{\theta}_k)}$ be an admissible solution corresponding to $\boldsymbol{\theta}_k$, and let $\Omega^{(k)}$ be the corresponding pseudo-subsonic domain.

Let $G \subseteq \mathbb{R}^2_+$ be any open set satisfying $\overline{G} \in \mathbb{R}^2_+$. Choose any compact subset $K \in G \setminus \partial \Omega^{(0)}$. Then, by Lemma 3.17(ii), passing to a subsequence (still denoted) $k, K \cap \Omega^{(0)} \in \Omega^{(k)}$ and $K \setminus \Omega^{(0)} \in G \setminus \Omega^{(k)}$. Using Proposition 4.3(iv), Proposition 4.6, Lemma 4.8(v), and the continuity property of admissible solutions given in Step 1 of the proof of Lemma 4.17, we have

$$\|\varphi^{(\boldsymbol{\theta}_k)} - \varphi_{\operatorname{norm}}\|_{1,\alpha,K \cap \Omega^{(0)}} \to 0$$
 as $k \to \infty$.

Similarly, using (2.24), Proposition 2.6(i), and (2.41), we have

$$\|\varphi^{(\boldsymbol{\theta}_k)} - \varphi_{\operatorname{norm}}\|_{W^{1,1}(K \setminus \Omega^{(\mathbf{0})})} \to 0 \quad \text{as } k \to \infty.$$

Finally, by (2.41) and the uniform bound in Lemma 3.3, there exists a constant C > 0 depending only on (γ, v_2, G) such that $\|\varphi^{(\theta_k)} - \varphi_{\text{norm}}\|_{0,1,G} \leq C$ for all $k \in \mathbb{N}$, which leads to

$$\|\varphi^{(\boldsymbol{\theta}_k)} - \varphi_{\operatorname{norm}}\|_{W^{1,1}(G)} \le \|\varphi^{(\boldsymbol{\theta}_k)} - \varphi_{\operatorname{norm}}\|_{W^{1,1}(K)} + 2C\operatorname{meas}(G \setminus K).$$

This completes the proof of Theorem 2.1.

5. Optimal Regularity of Solutions and Convexity of Free Boundaries

With the *a priori* estimates obtained in §3–§4, we can now give the complete proofs of Theorems 2.2–2.3, respectively.

5.1. Proof of Theorem 2.2: Optimal regularity of solutions. This section is devoted to the complete proof of Theorem 2.2. Let φ be an admissible solution in the sense of Definition 2.11.

Proof. Fix $\theta \in \Theta$. By the symmetry of the problem, it suffices to prove the statement near Γ_{sonic}^5 .

1. Proof of Theorem 2.2(i). First, it follows from Lemmas 3.6 and 3.9 that Γ_{shock} is C^{∞} in its relative interior and $\varphi \in C^{\infty}(\overline{\Omega} \setminus \Gamma_{\text{sonic}}^5 \cup \Gamma_{\text{sonic}}^6)$. By Definition 2.9, Γ_{sonic}^5 is a closed arc of a circle when $\theta_1 \in [0, \theta^{\text{s}})$, and becomes a point P_1 when $\theta_1 \in [\theta^{\text{s}}, \theta^{\text{d}})$. If $\theta_1 \in [0, \theta^{\text{s}})$, it follows from Propositions 3.12–3.13 that φ is $C^{1,1}$ up to Γ_{sonic}^5 , whilst, if $\theta_1 \in [\theta^{\text{s}}, \theta^{\text{d}})$, then Propositions 3.14 and 3.16 imply that φ is $C^{1,\alpha}$ up to $\Gamma_{\text{sonic}}^5 = \{P_1\}$ for some $\alpha \in (0, 1)$. Furthermore, from Propositions 3.12–3.14 and 3.16, we see that $\overline{\Gamma_{\text{shock}} \cup S_{25}^{\text{seg}}}$ is a $C^{1,\alpha}$ -curve, including at P_2 , for any $\theta_1 \in [0, \theta^{\text{d}})$. It remains to show that $\overline{\Gamma_{\text{shock}} \cup S_{25}^{\text{seg}}}$ is a $C^{2,\alpha}$ -curve when $\theta_1 \in [0, \theta^{\text{s}})$.

For $\theta_1 \in [0, \theta^s)$, let coordinates $(x, y) = \mathcal{R}(\boldsymbol{\xi})$ be given by (3.24). Let $\bar{\varepsilon} > 0$, $f_{5,0}$, and $f_{5,sh}$ be from Step 1 of the proof of Proposition 3.12. We extend $f_{5,sh}$ to $(-\bar{\varepsilon}, \bar{\varepsilon})$ by

(5.1)
$$f_{5,\mathrm{sh}}(x) \coloneqq f_{5,0}(x) \qquad \text{for } x \in (-\bar{\varepsilon}, 0].$$

so that

(5.2)
$$(f_{5,\rm sh} - f_{5,0})(0) = (f_{5,\rm sh} - f_{5,0})'(0) = 0.$$

Define $\psi \coloneqq (\varphi - \varphi_5) \circ \mathcal{R}^{-1}$ and $\bar{\psi}_{25} \coloneqq (\varphi_2 - \varphi_5) \circ \mathcal{R}^{-1}$. Then $\bar{\psi}_{25}(x, f_{5,0}(x)) = 0$, which implies $\bar{\psi}_{25}(x, f_{5,\text{sh}}(x)) - \bar{\psi}_{25}(x, f_{5,0}(x)) = \psi(x, f_{5,\text{sh}}(x)) \quad \text{for } x \in (0, \bar{\varepsilon}).$

Differentiating the above equality with respect to x twice, we obtain the following expression:

(5.3)
$$(f_{5,\rm sh} - f_{5,0})''(x) = \frac{A_1(x) + A_2(x) + A_3(x)}{\partial_y \bar{\psi}_{25}(x, f_{5,\rm sh}(x))}$$

where, for $(a_0, a_1, a_2) := (1, 2, 1)$,

$$A_{1}(x) \coloneqq \sum_{k=0}^{2} a_{k} \left((f_{5,0}(x))^{k} \partial_{x}^{2-k} \partial_{y}^{2} \bar{\psi}_{25}(x, f_{5,0}(x)) - (f_{5,\mathrm{sh}}(x))^{k} \partial_{x}^{2-k} \partial_{y}^{2} \bar{\psi}_{25}(x, f_{5,\mathrm{sh}}(x)) \right),$$

$$A_{2}(x) \coloneqq \left(\partial_{y} \bar{\psi}_{25}(x, f_{5,0}(x)) - \partial_{y} \bar{\psi}_{25}(x, f_{5,\mathrm{sh}}(x)) \right) f_{5,0}''(x),$$

$$A_{3}(x) \coloneqq -f_{5,\mathrm{sh}}''(x) \partial_{y} \psi(x, f_{5,\mathrm{sh}}(x)) - \sum_{k=0}^{2} a_{k} (f_{5,\mathrm{sh}}'(x))^{k} \partial_{x}^{2-k} \partial_{y}^{k} \psi(x, f_{5,\mathrm{sh}}(x)) .$$

By (5.2), $A_1(0) = A_2(0) = 0$. We differentiate the boundary condition in the tangential direction along Γ_{shock} and apply the arguments in Step 1 of the proof of Proposition 3.12 to obtain that there exists a constant C > 0 such that

$$|\psi_{xx}| \le C |(\psi, \psi_x, \psi_y, \psi_{xy}, \psi_{yy})| \quad \text{on } \mathcal{R}(\Gamma_{\text{shock}} \cap \partial \Omega_{\bar{\varepsilon}}^5),$$

where $\Omega_{\overline{\varepsilon}}^{5}$ is from Step 1 of the proof of Proposition 3.12. From the above estimate and Proposition 3.12, we obtain that $\psi_{xx}(x, f_{5,\text{sh}}(x)) \to 0$ as $x \to 0^{+}$, which implies that $A_{3}(x) \to 0$ as $x \to 0^{+}$. Hence, $\partial_{y}\bar{\psi}_{25}(x, f_{5,\text{sh}}(x)) \neq 0$ on $\mathcal{R}(\Gamma_{\text{shock}} \cap \partial \Omega_{\overline{\varepsilon}}^{5})$. Then we conclude from (5.3) that $(f_{5,\text{sh}} - f_{5,0})''(0) = 0$. This implies that the extension of $f_{5,\text{sh}}$ given by (5.1) is in $C^{2}([-\overline{\varepsilon}, \overline{\varepsilon}])$. Furthermore, we conclude from Proposition 3.12 that the extension of $f_{5,\text{sh}}$ given by (5.1) is in $C^{2,\alpha}((-\overline{\varepsilon}, \overline{\varepsilon}))$ for any $\alpha \in (0, 1)$. This implies that $\overline{\Gamma_{\text{shock}} \cup S_{25}^{\text{seg}}}$ is $C^{2,\alpha}$ for any $\alpha \in (0, 1)$, including at point $P_{2} = \mathcal{R}^{-1}(0, f_{5,\text{sh}}(0))$. This completes the proof of statement (i).

2. Proof of Theorem 2.2(ii)–(iii). It suffices to consider the case that $\theta_1 \in [0, \theta^s)$. We can apply Theorem B.3 to obtain the regularity of ψ in the neighbourhood of $\mathcal{R}(\Gamma_{\text{sonic}}^5)$. Then the admissible solution φ satisfies Theorem 2.2(ii)–(iii).

3. Proof of Theorem 2.2(iv). By Propositions 3.12–3.13, $\Gamma_{\text{shock}} \cap \partial \Omega_{\overline{\varepsilon}}^5$ can be represented as the graph of $y = f_{5,\text{sh}}(x)$ for $0 \le x \le \overline{\varepsilon}$.

Let $\{y_m^{(1)}\}_{m\in\mathbb{N}}$ be a sequence satisfying that $0 < y_m^{(1)} < f_{5,\mathrm{sh}}(0)$ for each $m \in \mathbb{N}$, and $y_m^{(1)} \rightarrow f_{5,\mathrm{sh}}(0)$ as $m \to \infty$. By Theorem 2.2(iii), we can choose a sequence $\{x_m^{(1)}\}_{m\in\mathbb{N}}$ such that $(x_m^{(1)}, y_m^{(1)}) \in \mathcal{R}(\Omega_{\bar{\varepsilon}}^5), 0 < x_m^{(1)} < \frac{1}{m}$, and

$$\left|\psi_{xx}(x_m^{(1)}, y_m^{(1)}) - \frac{1}{\gamma+1}\right| \le \frac{1}{m} \quad \text{for each } m \in \mathbb{N}.$$

It follows from Step 1 of the proof of Proposition 3.12 that $0 < y_m^{(1)} < f_{5,sh}(0) < f_{5,sh}(x_m^{(1)})$ for each $m \in \mathbb{N}$ so that

(5.4)
$$\lim_{m \to \infty} (x_m^{(1)}, y_m^{(1)}) = (0, f_{5, \text{sh}}(0)), \qquad \lim_{m \to \infty} |\psi_{xx}(x_m^{(1)}, y_m^{(1)})| = \frac{1}{\gamma + 1}$$

On the other hand, there exists a constant $\varepsilon \in (0, \overline{\varepsilon}]$ such that the boundary condition on $\Gamma_{\text{shock}} \cap \partial \Omega_{\overline{\varepsilon}}^5$ can be written as $\psi_x + b_1 \psi_y + b_0 \psi = 0$ with $(b_0, b_1) = (b_0, b_1)(\psi_x, \psi_y, \psi, x, f_{5,\text{sh}}(x))$ for any $x \in (0, \varepsilon)$. Let $\omega > 0$ from Step 1 of the proof of Proposition 3.12. Then

$$\{(x, f_{5, \mathrm{sh}}(x) - \frac{\omega}{10}x) : 0 < x < \varepsilon\} \subseteq \mathcal{R}(\Omega^{5}_{\overline{\varepsilon}})$$

Denote $F(x) \coloneqq \psi_x(x, f_{5,sh}(x) - \frac{\omega}{10}x)$. By the geometric relation given above, we have

$$F(x) = \psi_x(x, f_{5,\text{sh}}(x)) - \frac{\omega}{10}x \int_0^1 \psi_{xy}(x, f_{5,\text{sh}}(x) - \frac{\omega}{10}tx) dt$$

= $-(b_1\psi_y + b_0\psi)(x, f_{5,\text{sh}}(x)) - \frac{\omega}{10}x \int_0^1 \psi_{xy}(x, f_{5,\text{sh}}(x) - \frac{\omega}{10}tx) dt$ for $x \in (0, \varepsilon)$.

Then Proposition 3.12 implies that F(0) = 0, $F \in C([0,\varepsilon]) \cap C^1((0,\varepsilon))$, $\lim_{x \to 0+} \frac{F(x)}{x} = 0$ so that, by the mean value theorem, there exists a sequence $\{x_m^{(2)}\}_{m \in \mathbb{N}} \subseteq (0,\varepsilon)$ such that

(5.5)
$$\lim_{m \to \infty} x_m^{(2)} = 0, \qquad F'(x_m^{(2)}) = 0$$

For each $m \in \mathbb{N}$, define $y_m^{(2)} \coloneqq f_{5,\text{sh}}(x_m^{(2)}) - \frac{\omega}{10}x_m^{(2)}$, so that $\{(x_m^{(2)}, y_m^{(2)})\}_{m \in \mathbb{N}} \subseteq \mathcal{R}(\Omega_{\bar{\varepsilon}}^5)$. By the definition of $F(\cdot)$ and (5.5), we have

(5.6)
$$\lim_{m \to \infty} \psi_{xx}(x_m^{(2)}, y_m^{(2)}) = \lim_{m \to \infty} F'(x_m^{(2)}) - \lim_{m \to \infty} (f_{5, \text{sh}}(x_m^{(2)}) - \frac{\omega}{10}) \psi_{xy}(x_m^{(2)}, y_m^{(2)}) = -\lim_{m \to \infty} (f_{5, \text{sh}}(x_m^{(2)}) - \frac{\omega}{10}) \psi_{xy}(x_m^{(2)}, y_m^{(2)}).$$

Since $\lim_{m \to \infty} (x_m^{(2)}, y_m^{(2)}) = (0, f_{5,\text{sh}}(0))$, we combine (5.6) with Proposition 3.12 to obtain

$$\lim_{m \to \infty} \psi_{xx}(x_m^{(2)}, y_m^{(2)}) = 0.$$

We have demonstrated that there are two sequences, $\{(x_m^{(1)}, y_m^{(1)})\}_{m \in \mathbb{N}}$ and $\{(x_m^{(2)}, y_m^{(2)})\}_{m \in \mathbb{N}}$, taking values in $\mathcal{R}(\Omega_{\varepsilon}^5)$, such that the limits of both sequences are $(0, f_{5,\text{sh}}(0))$, but

$$\lim_{m \to \infty} \psi_{xx}(x_m^{(1)}, y_m^{(1)}) \neq \lim_{m \to \infty} \psi_{xx}(x_m^{(2)}, y_m^{(2)})$$

This completes the proof of Theorem 2.2(iv).

5.2. **Proof of Theorem 2.3: Convexity of free boundaries and transonic shocks.** We now discuss the geometric properties of the transonic shock as a free boundary. In Chen-Feldman-Xiang [12], a general framework was developed under which the self-similar transonic shock, as a free boundary, is proved to be uniformly convex for the potential flow equation; see Framework (A) in Appendix B.5. In this subsection, we apply this framework to prove the uniform convexity of the transonic shock in the four-shock interaction problem, so it suffices to prove that the admissible solutions satisfy the conditions in Theorems B.4–B.5.

Lemma 5.1. The following statements hold:

- (i) Any admissible solution in the sense of Definition 2.11 satisfies the conditions of Theorems B.4–B.5.
- (ii) Any global weak solution of the four-shock interaction problem in the sense of Definition 2.1 —satisfying also all properties of Definition 2.11 except (2.43) and with transonic shock Γ_{shock} being a strictly convex graph—satisfies condition (2.43) of Definition 2.11.

Proof. We divide the proof into six steps. We start with the proof of assertion (i).

1. Claim: Region Ω satisfies the conditions in Framework (A). Indeed, for $\boldsymbol{\theta} \in [0, \theta^s)^2$, the required piecewise regularity holds, since Γ_{sym} is a straight segment, while Γ_{sonic}^5 and Γ_{sonic}^6 are the arcs of the sonic circles, and Γ_{shock} has the regularity stated in Theorem 2.2(i). For any incident angles $\boldsymbol{\theta} \in \Theta$, the fact that all angles of the corners of Ω are less than π can be verified as follows:

If $\theta_1 \in [0, \theta^s)$, since $\overline{S_{25}^{\text{seg}} \cup \Gamma_{\text{shock}}}$ is a $C^{2,\alpha}$ -curve for any $\alpha \in (0, 1)$, including at points P_2 , it follows that Γ_{shock} is tangential to S_{25}^{seg} at P_2 . Meanwhile, for any $\boldsymbol{\theta} \in \Theta$, $\partial B_{c_5}(O_5)$ and S_{25} always have two intersection points including P_2 . Therefore, the meeting angles at the two corners $\{P_1, P_2\}$ belong to $[0, \pi)$ when $\theta_1 \in (0, \theta^s)$. If $\theta_1 \in [\theta^s, \theta^d)$, we see that $P_2 = P_1 = P_0^1$. From (2.38), the meeting angle at P_0^1 is $\pi - \theta_{25} \in (0, \frac{\pi}{2})$, which satisfies

$$\pi - \theta_{25} \ge \delta_{5,6}^{(\theta^{\mathrm{s}})} > 0$$
.

Similarly, the meeting angles at the two corners $\{P_3, P_4\}$ also belong to $(0, \pi)$ for any $\theta_2 \in [0, \theta^d)$.

2. The entropy condition (C-1) in Theorem B.4 follows directly from the properties of Definition 2.11, where state (0) in Theorem B.4 is state (2) in our problem. From the regularity of φ and Γ_{shock} in Theorem 2.2(i), we see that conditions (C-2) and (C-4) of Theorem B.4 hold. Property (iii) of Definition 2.11 implies that condition (C-3) of Theorem B.4 holds.

3. We use the notation for the endpoints of Γ_{shock} from Framework (A), that is, $A := P_2$ and $B := P_3$, where $P_{i+1} \equiv P_0^i$ when $\theta_i \in [\theta^s, \theta^d)$ for i = 1, 2. Then, by the properties of Definition 2.11, we see that $\tau_A = e_{S_{25}}$ and $\tau_B = e_{S_{26}}$, and it is clear that $\tau_A \neq \pm \tau_B$ for all $\theta \neq 0$. Accordingly, let Con be given by (B.14). Combining property (2.43) with the fact that Γ_{shock} is the level set: $\{\varphi - \varphi_2 = 0\}$, we obtain that $\{P + \text{Con}\} \cap \Omega = \emptyset$ for all $P \in \Gamma_{\text{shock}}$. Thus, condition (C-5) of Theorem B.4 is satisfied.

4. For condition (C-6) of Theorem B.4, we show that Case (C-6c) holds with $\phi = \varphi - \varphi_2$, $e = (0,1) \in \text{Con}, \Gamma_1 \coloneqq \Gamma_{\text{sonic}}^6 \cup \Gamma_{\text{sym}} \cup \Gamma_{\text{sonic}}^5$, and $\Gamma_2 = \emptyset$. For j = 5, 6, write $w \coloneqq \partial_e(\varphi - \varphi_j)$, which satisfies a strictly elliptic equation in Ω by taking the derivative of (3.1) in direction e, and using Definition 2.11(iii). By conditions (iii)–(iv) of Problem 2.10, w = 0 on $\Gamma_{\text{sym}} \cup \Gamma_{\text{sonic}}^j$, which is a global maximum by Lemma 3.1(iii). Therefore, if $P \in \Gamma_{\text{sym}} \cup \Gamma_{\text{sonic}}^j$ is a point of local minimum for w, then w = 0 on $B_r(P) \cap \Omega$ for some r > 0, and so w = 0 in Ω by the strong maximum principle, which is a contradiction to Lemma 3.1(i). It follows that $\phi_e \equiv w - v_2$ cannot attain a local minimum on Γ_1 .

5. We now show that conditions (C-7)–(C-10) are satisfied with $\hat{\Gamma}_0 = \Gamma_{\text{sonic}}^5 \setminus \{P_2\}$ for $\theta_1 \in [0, \theta^s)$ (resp. $\hat{\Gamma}_0 = \emptyset$ for $\theta_1 \in [\theta^s, \theta^d)$), $\hat{\Gamma}_1 = \Gamma_{\text{sym}}^0$, $\hat{\Gamma}_2 = \emptyset$, and $\hat{\Gamma}_3 = \Gamma_{\text{sonic}}^6 \setminus \{P_3\}$ for $\theta_2 \in [0, \theta^s)$ (resp. $\hat{\Gamma}_3 = \emptyset$ for $\theta_2 \in [\theta^s, \theta^d)$).

Indeed, (C-7) clearly holds. Also, (C-8) holds since $D\varphi = D\varphi_5$ on Γ_{sonic}^5 for $\theta_1 \in [0, \theta^s)$ so that $\phi_e = \partial_e(\varphi_5 - \varphi_2)$ is a constant. Similarly, $D\varphi = D\varphi_6$ on Γ_{sonic}^6 for $\theta_2 \in [0, \theta^s)$. Condition (C-9) on $\hat{\Gamma}_1 = \Gamma_{\text{sym}}^0$ can be checked as follows: If $e \neq (0, \pm 1)$, then, as in Step **6** of the proof of [12,

Lemma 7.8], $\phi_{\boldsymbol{e}}$ cannot attain its local minima or maxima on Γ_{sym}^{0} . In the other case, when $\boldsymbol{e} = (0, \pm 1)$, the slip boundary condition $\partial_{\eta}\phi = 0$ on Γ_{sym}^{0} implies that $\phi_{\boldsymbol{e}}$ is constant on Γ_{sym}^{0} , which verifies (C-9). Case (C-10a) of (C-10) clearly holds here since $\hat{\Gamma}_{2} = \emptyset$. This concludes the proof of assertion (i).

6. It remains to prove assertion (ii): Any solution satisfying all the properties of Definition 2.11 except (2.43), and with transonic shock Γ_{shock} being a strictly convex graph in the sense of (B.15)–(B.16), satisfies condition (2.43) of Definition 2.11.

We apply (B.15) in the present case with $(A, B) = (P_2, P_3)$ and e = (0, 1) such that $\xi(S, T) = (-T, S)$. Then, using the properties of Definition 2.11(i),

$$e_{S_{25}} = rac{(-1, f'(T_A))}{|(-1, f'(T_A))|}, \qquad e_{S_{26}} = -rac{(-1, f'(T_B))}{|(-1, f'(T_B))|}.$$

Also, from the strict concavity of f in the sense of (B.16), we obtain that $f'(T_A) > f'(T) > f'(T_B)$ and $f(T) < f(T_1) + f'(T_1)(T - T_1)$ for all $T, T_1 \in (T_A, T_B)$. From this, we see that $\{P + \operatorname{Con}\} \cap \Omega = \emptyset$ for any $P \in \Gamma_{\operatorname{shock}}$. Since $\varphi \leq \varphi_2$ in Ω from Definition 2.11 and $\varphi = \varphi_2$ on $\Gamma_{\operatorname{shock}}$, we obtain that $\partial_e \varphi \geq \partial_e \varphi_2$ for any $e \in \operatorname{Con}$, which implies (2.43).

A. PROOF OF LEMMA 2.5 AND RELATED PROPERTIES OF SOLUTIONS

A.1. Proof of Lemma 2.5: Monotonicity of critical angles for 2-D steady potential flow. The proof of Lemma 2.5 is given in four steps.

1. The shock polar for 2-D steady potential flow connecting a constant supersonic upstream state $U_{\infty} = (\rho_{\infty}, u_{\infty}, 0)$ to a constant downstream state $U = (\rho, u, v)$ is given by

(A.1)
$$u = u_{\infty} - \frac{2(h(\rho) - h(\rho_{\infty}))}{\rho + \rho_{\infty}} \frac{\rho}{u_{\infty}}, \quad v^2 = u_{\infty}^2 - u^2 - 2(h(\rho) - h(\rho_{\infty})),$$

where $h(\rho) \coloneqq \frac{1}{\gamma-1}(\rho^{\gamma-1}-1)$ for $\gamma > 1$, and $h(\rho) = \ln \rho$ for $\gamma = 1$; see *e.g.* [19, §17.2] for the derivation. The downstream density $\rho \in (\rho_{\infty}, \overline{\rho})$ is a parameter along the shock polar, where the maximal density $\overline{\rho} > \rho_{\infty}$ is determined uniquely by $v(\overline{\rho}) = 0$ in (A.1), which satisfies

$$M_{\infty}^{2}\left(1-\left(\frac{\rho_{\infty}}{\overline{\rho}}\right)^{2}\right)=\frac{2\left(h(\overline{\rho})-h(\rho_{\infty})\right)}{c_{\infty}^{2}}$$

for $M_{\infty} \coloneqq \frac{u_{\infty}}{c_{\infty}} > 1$, the Mach number of the upstream state that is assumed to be supersonic, and c_{∞} the sonic speed. In the following, we denote by $w \coloneqq \tan \theta_{\text{stdy}}$, where $\theta_{\text{stdy}} \in (0, \frac{\pi}{2})$ is the angle between the velocities of the upstream and the downstream flows, and by $\tau \coloneqq \frac{\rho}{\rho_{\infty}} \in (1, \overline{\tau})$ the ratio between the downstream and upstream densities, with $\overline{\tau} \coloneqq \frac{\overline{\rho}}{\rho_{\infty}}$. Using the polytropic pressure law with $\gamma \ge 1$, we have

(A.2)
$$w = \frac{v}{u} = \frac{(2h(\tau))^{1/2} (M_{\infty}^2 (1 - \tau^{-2}) - 2h(\tau))^{1/2}}{M_{\infty}^2 (1 + \tau^{-1}) - 2h(\tau)}$$

From [10, Lemma 7.3.2], there exist both a unique detachment angle $\theta_{\text{stdy}}^{d} \in (0, \frac{\pi}{2})$ and a unique sonic angle $\theta_{\text{stdy}}^{s} \in (0, \theta_{\text{stdy}}^{d})$, which are characterized by

$$w^{\mathrm{d}} \coloneqq \tan \theta^{\mathrm{d}}_{\mathrm{stdy}} = \sup \left\{ w(\tau) \, : \, \tau \in (1, \overline{\tau}) \right\}, \qquad w^{\mathrm{s}} \coloneqq \tan \theta^{\mathrm{s}}_{\mathrm{stdy}} = \frac{v}{u} \quad \mathrm{when} \, \, u^2 + v^2 = \rho^{\gamma - 1} \, .$$

By differentiation of (A.2) with respect to $\tau \in (1, \overline{\tau})$, it follows that $w^{d} = w(\tau_{d})$, where $\tau_{d} \in (1, \overline{\tau})$ satisfies

(A.3)
$$M_{\infty}^{2} \left(2h(\tau_{\rm d}) + (\tau_{\rm d}^{2} - 1)h'(\tau_{\rm d}) \right) - 2h(\tau_{\rm d}) \left(2h(\tau_{\rm d}) + \tau_{\rm d}(\tau_{\rm d} + 1)h'(\tau_{\rm d}) \right) = 0.$$

Similarly, by routine calculation, it follows that $w^{s} = w(\tau_{s})$, where $\tau_{s} \in (1, \overline{\tau})$ satisfies

(A.4)
$$(\gamma + 1)h(\tau_{\rm s}) = M_{\infty}^2 - 1.$$

For the polytropic gas, it is clear that the detachment angle and sonic angle $\theta_{\text{stdy}}^{\text{d}}$ and $\theta_{\text{stdy}}^{\text{s}}$ depend only on γ and M_{∞} . We also observe the following useful identities:

2. For the detachment angle, we combine (A.2) with (A.3) to obtain a parametric description of the detachment angle and the upstream Mach number in terms of τ_d :

(A.6)
$$w^{d} = \frac{1}{2} \frac{\left(\tau_{d}(\tau_{d}^{2}-1)h'-2h\right)^{1/2} \left((\tau_{d}^{2}-1)h'+2h\right)^{1/2}}{\tau_{d}(\tau_{d}+1)h'+h}, \qquad M_{\infty}^{2} = \frac{2h\left(2h+\tau_{d}(\tau_{d}+1)h'\right)}{2h+(\tau_{d}^{2}-1)h'},$$

where h, h', and h'' are evaluated at $\tau = \tau_d$. A direct calculation shows that the signs of $(w^d)_{\tau_d}$ and $(M^2_{\infty})_{\tau_d}$ are equal to the sign of the quantity:

$$f_{\rm d}(\tau_{\rm d}) \coloneqq 2h^2 \left(3 + (\tau_{\rm d}+1)\frac{h''}{h'}\right) + (3\tau_{\rm d}-5)(\tau_{\rm d}+1)hh' + \tau_{\rm d}(\tau_{\rm d}+1)(\tau_{\rm d}^2-1)(h')^2.$$

Applying (A.5) evaluated at $\tau = \tau_d$, along with $\gamma \ge 1$, $\tau_d > 1$, and h > 0, we obtain

$$\begin{split} f_{\rm d}(\tau_{\rm d}) &= \big(3(\tau_{\rm d} - \tau_{\rm d}^{-1}) + \big(3(\gamma - 1)(\tau_{\rm d} - \tau_{\rm d}^{-1}) + 2(2 - \tau_{\rm d}^{-1})\big)h\big)h + (1 + \tau_{\rm d}^{-1})\big(\tau_{\rm d}^2(\tau_{\rm d}^2 - 1)(h')^2 - 2h\big) \\ &> (1 + \tau_{\rm d}^{-1})\tilde{f}_{\rm d}(\tau_{\rm d})\,, \end{split}$$

where $\tilde{f}_{\rm d}(\tau_{\rm d}) \coloneqq \tau_{\rm d}^2(\tau_{\rm d}^2-1)(h')^2 - 2h$. We note that $\tilde{f}_{\rm d}(1) = 0$ and

$$\tilde{f}'_{\rm d}(\tau) = 2h'(\tau) ((\gamma - 1)\tau(\tau^2 - 1)h'(\tau) + \tau^{\gamma + 1} - 1) > 0$$
 for any $\tau > 1$.

It follows that $\tilde{f}_{\rm d}(\tau_{\rm d}) > 0$ and $f_{\rm d}(\tau_{\rm d}) > 0$ so that $(w^{\rm d})_{\tau_{\rm d}} > 0$ and $(M^2_{\infty})_{\tau_{\rm d}} > 0$ for all $\tau_{\rm d} > 1$. Using the chain rule, we conclude that $w^{\rm d}$ and $\theta^{\rm d}_{\rm stdy}$ are strictly increasing with respect to $M_{\infty} > 1$.

3. For the sonic angle, we combine (A.2) with (A.4) to give a parametric description of the sonic angle and upstream Mach number in terms of τ_s :

(A.7)
$$w^{s} = \frac{\left(2h\right)^{1/2} \left(\left(\tau_{s}^{2}-1\right)\tau_{s}^{\gamma-1}-2h\right)^{1/2}}{\left(\tau_{s}+1\right)\tau_{s}^{\gamma-1}+2h}, \qquad M_{\infty}^{2} = 1+(\gamma+1)h$$

where h and h' are evaluated at $\tau = \tau_s$. It is clear that $(M_{\infty}^2)_{\tau_s} > 0$ since h' > 0, whilst a direct calculation shows that the sign of $(w^s)_{\tau_s}$ is equal to the sign of the quantity

$$f_{\rm s}(\tau_{\rm s}) \coloneqq 2h \big(1 + (\gamma - 1)h \big) \big(1 + (\gamma + 1)h \big) + (\tau_{\rm s} + 1) \big((\tau_{\rm s} - 1)\tau_{\rm s}^{\gamma - 1} - 2h \big) h' \,.$$

Applying (A.5) evaluated at $\tau = \tau_{\rm s}$, along with $\gamma \ge 1$, $\tau_{\rm s} > 1$, and $h'(\tau_{\rm s}) = \tau_{\rm s}^{\gamma-2} > 0$, we obtain

$$f_{\rm s}(\tau_{\rm s}) = \left(2(\gamma+1)\tau_{\rm s}h^2 + \tau_{\rm s}(\tau_{\rm s}^2-1)h' - 2h\right)h' > \tilde{f}_{\rm s}(\tau_{\rm s})h'$$

where $\tilde{f}_{\rm s}(\tau_{\rm s}) \coloneqq \tau_{\rm s}(\tau_{\rm s}^2 - 1)h' - 2h$. Notice that $\tilde{f}_{\rm s}(1) = 0$ and

1 10

$$\tilde{f}_{\rm s}'(\tau)=(\gamma+1)(\tau^2-1)h'(\tau)>0\qquad {\rm for \ any}\ \tau>1.$$

Then it follows that $f_s(\tau_s) > 0$ and $f_s(\tau_s) > 0$ so that $(w^s)_{\tau_s} > 0$. We conclude via the chain rule that w^s and hence θ^s_{stdy} are strictly increasing with respect to $M_{\infty} > 1$.

4. The limiting values stated in the lemma can be checked directly from (A.6)–(A.7), now that the monotonicity of (w^d, M_{∞}^2) with respect to τ_d in (A.6) and the monotonicity of (w^s, M_{∞}^2) with respect to τ_s in (A.7) have been verified.

A.2. Monotonicity properties with respect to the incident angles. In Lemma 2.3, we have shown that the pseudo-Mach number of state (2) at the reflection point P_0^1 is a strictly decreasing function of $\theta_1 \in (0, \theta^{cr})$. Subsequently, in Proposition 2.6, we have used Lemmas 2.3 and 2.5 to prove the existence of the unique detachment angle θ^d and sonic angle θ^s . In Remark 2.1(i), we have stated the relation between θ^d , θ^s , and θ^{cr} , which depends on the choice of (γ, v_2) . The following lemma provides the proof of this statement. **Lemma A.1.** For any $\gamma > 1$, there exist constants v_2^d and v_2^s , with $v_{\min} < v_2^s < v_2^d < 0$, uniquely determined by γ such that

(A.8)
$$\operatorname{sgn}\left(\hat{\theta}_{25}(\theta^{\mathrm{cr}}; v_2) - \theta^{\mathrm{d}}_{\mathrm{stdy}}(1, |D\varphi_2(P^1_{0,\mathrm{cr}})|)\right) = \operatorname{sgn}(v_2 - v_2^{\mathrm{d}}),$$

(A.9)
$$\operatorname{sgn}\left(\hat{\theta}_{25}(\theta^{\mathrm{cr}};v_2) - \theta^{\mathrm{s}}_{\mathrm{stdy}}(1, |D\varphi_2(P^1_{0,\mathrm{cr}})|)\right) = \operatorname{sgn}(v_2 - v^{\mathrm{s}}_2),$$

where $P_{0,\mathrm{cr}}^1 \coloneqq P_0^1|_{\theta_1=\theta^{\mathrm{cr}}}$, and we define $\theta_{\mathrm{stdy}}^d(1, M_\infty) \equiv \theta_{\mathrm{stdy}}^s(1, M_\infty) \coloneqq 0$ for any $0 \le M_\infty \le 1$. Furthermore, when $\gamma = 1$, we define $v_{\min} = v_2^s = v_2^d = -\infty$.

Proof. We divide the proof into two steps.

1. For $\gamma > 1$ and $v_2 \in (v_{\min}, 0)$, a direct calculation by using (2.15), (2.18), and Lemma 2.3 gives

(A.10)
$$M_{2,\min} \coloneqq \inf_{\theta_1 \in (0,\theta^{cr})} |D\varphi_2(P_0^1)| = |D\varphi_2(P_{0,cr}^1)| = \frac{v_2 v_{\min}}{\sqrt{v_{\min}^2 - v_2^2}}$$
$$\hat{\theta}_{25}(\theta^{cr}; v_2) = \arcsin\left(\frac{-v_2}{M_{2,\min}}\right) = \arccos\left(|\frac{v_2}{v_{\min}}|\right) > 0.$$

Then $\hat{\theta}_{25}(\theta^{\rm cr}; v_2)$ is strictly increasing with respect to $v_2 \in (v_{\min}, 0)$. We also calculate

(A.11)
$$\lim_{v_2 \to v_{\min}^+} M_{2,\min} = \infty, \quad \lim_{v_2 \to 0^-} M_{2,\min} = 0, \quad \frac{\mathrm{d}M_{2,\min}^2}{\mathrm{d}v_2} = \frac{2v_{\min}^4 v_2}{(v_{\min}^2 - v_2^2)^2} < 0,$$

so that $M_{2,\min}$ is strictly decreasing with respect to $v_2 \in (v_{\min}, 0)$. Also observe that $M_{2,\min} > 1$ if and only if $v_2 \in (v_{\min}, v_{\min})$, for $v_{\min} \coloneqq \frac{v_{\min}}{\sqrt{v_{\min}^2 + 1}}$. Then, by Lemma 2.5, $\theta_{stdy}^d(1, M_{2,\min})$ is strictly decreasing with respect to $v_2 \in (v_{\min}, v_{\min})$. In particular, by (A.10)–(A.11) and [10, Lemma 7.3.3], we have

$$(\hat{\theta}_{25}(\theta^{\mathrm{cr}}; v_2), \theta^{\mathrm{d}}_{\mathrm{stdy}}(1, M_{2,\min})) \to (\arctan(|v_{\min}|), 0) \qquad \text{as} \quad v_2 \to v_{\mathrm{mid}}^-, \\ (\hat{\theta}_{25}(\theta^{\mathrm{cr}}; v_2), \theta^{\mathrm{d}}_{\mathrm{stdy}}(1, M_{2,\min})) \to (0, \frac{\pi}{2}) \qquad \text{as} \quad v_2 \to v_{\mathrm{min}}^+.$$

Then we conclude from the monotonicity properties of $(\hat{\theta}_{25}(\theta^{cr}; v_2), \theta^d_{stdy}(1, M_{2,\min}))$ with respect to $v_2 \in (v_{\min}, v_{mid})$ that there exists a unique value $v_2^d \in (v_{\min}, v_{mid})$, depending only on γ , such that (A.8) holds.

The proof of (A.9) follows via a similar argument by using the fact that $\theta_{stdv}^{s} \in (0, \theta_{stdv}^{d})$ and

$$\begin{split} \theta^{\rm s}_{\rm stdy}(1,M_{2,\min}) &\to 0 & \text{as } v_2 \to v_{\rm mid}^- \,, \\ \theta^{\rm s}_{\rm stdy}(1,M_{2,\min}) &\to \arctan\left(\frac{2}{\gamma-1}\right)^{1/2} & \text{as } v_2 \to v_{\min}^+ \,. \end{split}$$

2. For $\gamma = 1$, from (2.15), we see that $v_{\min} = -\infty$ and $\theta^{cr} = \frac{\pi}{2}$. Taking the limit: $v_{\min} \to -\infty$ in Step **1** above, we obtain that $M_{2,\min} = |v_2|$ and $\hat{\theta}_{25}(\frac{\pi}{2}; v_2) = \frac{\pi}{2}$. From Lemma 2.5, we know that the sonic and detachment angles for 2-D steady potential flow satisfy $0 < \theta^{s}(1, M_{\infty}) < \theta^{d}(1, M_{\infty}) < \frac{\pi}{2}$ for any $M_{\infty} > 1$, which implies that (A.8)–(A.9) hold with $v_2^{d} = v_2^{s} = -\infty$.

Lemma 2.3 is also useful to determine the monotonicity of other quantities related to the four-shock interaction problem. In the following lemma, we show that the reflected shock angles θ_{25} and θ_{26} are monotonic with respect to the incident shock angles θ_1 and θ_2 respectively and, subsequently, that certain intersection points of the reflected shocks are also monotonic functions of the incident angles.

Lemma A.2. Fix $\gamma \geq 1$ and $v_2 \in (v_{\min}, 0)$.

- (i) For any incident angle $\theta_1 \in [0, \theta^d)$, let the reflected shock angle $\theta_{25} \in (\frac{\pi}{2}, \pi]$ be given by (2.34). Then θ_{25} is a continuous, strictly decreasing function of θ_1 , and the map: $\theta_1 \mapsto \theta_{25}$ is C^{∞} -smooth on $\theta_1 \in [0, \theta^d)$.
- (ii) For the right-most intersection $P_2 = (\xi^{P_2}, \eta^{P_2})$ of $\partial B_{c_5}(O_5)$ and S_{25} as given in Definition 2.9, and for the η -intercept of S_{25} which is denoted here by a_{25} : η^{P_2} is strictly decreasing with respect to $\theta_1 \in (0, \theta^{\rm s})$, and a_{25} is strictly increasing with $\theta_1 \in (0, \theta^{\rm d})$.

Proof. For (i), the continuity and smoothness of the map: $\theta_1 \mapsto \theta_{25}$ follows due to Proposition 2.6 and (2.34), whilst the strict monotonicity follows directly from Lemma 2.3 and [2, Lemma 2.17]. Indeed, suppose that $\theta_1, \tilde{\theta}_1 \in [0, \theta^d)$ are such that $\theta_{25}(\theta_1) = \theta_{25}(\tilde{\theta}_1)$. Then, by [2, Lemma 2.17], the uniform state (5) is uniquely determined by $(v_2, \gamma, \theta_{25})$ so that the location of the reflected shock S_{25} must coincide for both θ_1 and $\tilde{\theta}_1$. In particular, for the pseudo-Mach number of state (2) at the reflection point P_0^1 , as defined by (2.18), we see that $M_2^{(\theta_1)} = M_2^{(\tilde{\theta}_1)}$ so that $\theta_1 = \tilde{\theta}_1$ by Lemma 2.3.

We have shown that the map: $\theta_1 \mapsto \theta_{25}$ is continuous and injective, and therefore strictly monotonic. In particular, by property (i) of Proposition 2.6 and (2.34), $\theta_{25} \to \pi^-$ as $\theta_1 \to 0^+$; whilst, by property (ii) of Proposition 2.6 and (2.34), $\theta_{25} < \pi$ for any $\theta_1 \in (0, \theta^d]$. It follows that the map: $\theta_1 \mapsto \theta_{25}$ is monotonic decreasing.

For (ii), the first result follows directly from part (i) combined with [2, Lemma 2.22], whilst the second result follows directly from part (i) combined with [2, Eqs. (2.4.14) and (2.4.42)]. \Box

Finally, we give the proof that the intersection point P_I of the reflected shocks S_{25} and S_{26} can be extended to a $C(\overline{\Omega})$ -function.

Lemma A.3. For $\theta \in \overline{\Theta} \setminus \{0\}$, denote by $P_I = (\xi^{P_I}, \eta^{P_I})$ the unique intersection point of the reflected shocks S_{25} and S_{26} , as given by (4.9). Then

(A.12)
$$\lim_{\boldsymbol{\theta} \to \mathbf{0}, \boldsymbol{\theta} \in \overline{\Theta} \setminus \mathbf{0}} (\xi^{P_I}, \eta^{P_I}) = (0, \eta_0),$$

where η_0 is given by (2.23).

Proof. By Proposition 2.6 and Lemma A.2(i), limit (A.12) is equivalent to the limit:

$$\lim_{\substack{(\theta_{25},\theta_{26})\to(\pi,0)\\(\theta_{25},\theta_{26})\in\overline{\Theta_{\text{refl}}}\setminus\{(\pi,0)\}}} (\xi^{P_I},\eta^{P_I}) = (0,\eta_0)$$

where $\Theta_{\text{refl}} \coloneqq \{(\theta_{25}, \theta_{26}) : \boldsymbol{\theta} \in \Theta\}$. The proof of the above limit consists of three steps.

1. We first consider the unilateral normal reflection case: $\boldsymbol{\theta} \in (\{0\} \times (0, \theta^d)) \cup ((0, \theta^d) \times \{0\})$. Without loss of generality, we restrict our attention to the case: $\boldsymbol{\theta} \in (0, \theta^d) \times \{0\}$, and denote $(v_{\infty}, \beta) \coloneqq (-v_2, \pi - \theta_{25})$, where θ_{25} depends only on (v_2, γ, θ_1) . We consider the Prandtl-Meyer reflection configuration associated with parameters (v_{∞}, β) , as defined in [2], and reflect this configuration in the vertical axis. By [2, Lemma 2.17], the location of the reflected shock S_{25} is uniquely determined by $(v_2, \gamma, \theta_{25})$, and thus coincides with the Prandtl-Meyer oblique shock $S_{\mathcal{O}}$. Similarly, the (normal) reflected shock $S_{26} = S_0$ coincides with the Prandtl-Meyer normal shock $S_{\mathcal{N}}$, so that the intersection point P_I of S_{25} and S_{26} , as given by (4.9), coincides with the intersection point of $S_{\mathcal{O}}$ and $S_{\mathcal{N}}$, as given by [2, Eq. (4.1.33)]. That is,

$$\left(\xi^{P_{I}},\eta^{P_{I}}\right)=\left(\frac{a_{25}-\eta_{0}}{\tan\theta_{25}},\eta_{0}\right),$$

where $a_{25} := -\xi^{P_0^1} \tan \theta_{25}$ and η_0 are the η -intercepts of S_{25} and the normal reflection S_0 , respectively, as given in §2. By Proposition 2.6(i) and Lemma A.2(ii), $a_{25} \to \eta_0$ as $\theta_{25} \to \pi^-$. Then we may apply L'Hôpital's rule to give

$$\lim_{\theta_{25}\to\pi^{-}}\xi^{P_{I}} = -\lim_{\theta_{25}\to\pi^{-}}\frac{\mathrm{d}a_{25}}{\mathrm{d}\theta_{25}}$$

From [2, Eq. (2.4.14)], $a_{25} = v_2 - q_2 \sec \theta_{25}$, for $q_2 := \operatorname{dist}(S_{25}, O_2)$. By taking the derivative with respect to θ_{25} , we have

$$\frac{\mathrm{d}a_{25}}{\mathrm{d}\theta_{25}} = -\sec\theta_{25}\frac{\mathrm{d}q_2}{\mathrm{d}\theta_{25}} - q_2\sec\theta_{25}\tan\theta_{25} \,.$$

Using [10, Lemma 6.1.2] and (2.23), $1 < \lim_{\theta_{25} \to \pi^-} q_2 = \eta_0 - v_2 < \infty$ so that

$$\lim_{\theta_{25} \to \pi^-} \frac{\mathrm{d}a_{25}}{\mathrm{d}\theta_{25}} = \lim_{\theta_{25} \to \pi^-} \frac{\mathrm{d}q_2}{\mathrm{d}\theta_{25}}.$$

By [2, Eq. (2.4.12)], we have a relation between the pseudo-Mach numbers $(M_2, M_5) \coloneqq (\frac{q_2}{c_2}, \frac{q_5}{c_5})$ for $q_5 \coloneqq \text{dist}(S_{25}, O_5)$ and the given parameters $(\gamma, v_2, \theta_{25})$, expressed as

$$M_2^{\frac{\gamma-1}{\gamma+1}} (M_2^{\frac{2}{\gamma+1}} - M_5^{\frac{2}{\gamma+1}}) = v_2 \sec \theta_{25}.$$

Taking the derivative:

$$D = \lim_{\theta_{25} \to \pi^{-}} \frac{\mathrm{d}}{\mathrm{d}\theta_{25}} \Big(M_2^{\frac{\gamma-1}{\gamma+1}} (M_2^{\frac{2}{\gamma+1}} - M_5^{\frac{2}{\gamma+1}}) \Big) \\ = \lim_{\theta_{25} \to \pi^{-}} \frac{1}{\gamma+1} \Big((\gamma-1) M_2^{-1} v_2 \sec\theta_{25} + 2M_2^{\frac{\gamma-1}{\gamma+1}} (M_2^{\frac{1-\gamma}{1+\gamma}} - M_5^{\frac{1-\gamma}{1+\gamma}} \frac{\mathrm{d}M_5}{\mathrm{d}M_2}) \Big) \frac{\mathrm{d}M_2}{\mathrm{d}\theta_{25}}$$

By [2, Eqs. (2.4.9)–(2.4.10)], we see that $\frac{dM_5}{dM_2} < 0$ for all $M_2 \in (0,\infty) \setminus \{1\}$. Therefore, the bracketed term above is uniformly positive for $\theta_{25} \in (\frac{\pi}{2}, \pi)$ so that

$$\lim_{\theta_{25} \to \pi^-} \frac{\mathrm{d}q_2}{\mathrm{d}\theta_{25}} = \lim_{\theta_{25} \to \pi^-} \frac{\mathrm{d}M_2}{\mathrm{d}\theta_{25}} = 0.$$

2. We now consider the general case $\theta \in \overline{\Theta} \setminus \{0\}$. By direct calculation, we have

$$\xi^{P_I} = \frac{a_{25} - a_{26}}{\tan \theta_{26} - \tan \theta_{25}} \,.$$

where $a_{2j} > 0$ represents the η -intercept of S_{2j} for j = 5, 6. Note that a_{25} depends only on $(v_2, \gamma, \theta_{25})$, whilst a_{26} depends only on $(v_2, \gamma, \theta_{26})$. In particular, from Step 1,

(A.13)
$$0 = \lim_{\theta_{25} \to \pi^{-}} \frac{a_{25} - \eta_0}{\tan \theta_{25}} = \lim_{\theta_{26} \to 0^+} \frac{a_{26} - \eta_0}{\tan \theta_{26}}$$

Let $\{\boldsymbol{\theta}^{(k)}\}_{k\in\mathbb{N}} \subseteq \overline{\Theta} \setminus \{\mathbf{0}\}$ be any sequence of parameters with $\boldsymbol{\theta}^{(k)} \to \mathbf{0}$ as $k \to \infty$. By moving to a subsequence, we may assume any of the following cases: (i) $\theta_1^{(k)} = 0$ for all $k \in \mathbb{N}$; (ii) $\theta_2^{(k)} = 0$ for all $k \in \mathbb{N}$; (iii) both $\theta_1^{(k)} > 0$ and $\theta_2^{(k)} > 0$ for all $k \in \mathbb{N}$. Denote by $\xi_{(k)}^{P_I}$ the ξ -coordinates of the intersection points P_I associated with parameters $\boldsymbol{\theta}^{(k)}$. In cases (i) and (ii), it is clear from the previous step that $\xi_{(k)}^{P_I} \to 0$ as $k \to \infty$, so we focus on case (iii). Using the triangle inequality, the fact that $\tan \theta_{25}$ and $\tan \theta_{26}$ have opposite signs, and (A.13), we obtain

$$|\xi_{(k)}^{P_{I}}| = \left|\frac{a_{25}^{(k)} - a_{26}^{(k)}}{\tan\theta_{26}^{(k)} - \tan\theta_{25}^{(k)}}\right| \le \left|\frac{a_{25}^{(k)} - \eta_{0}}{\tan\theta_{25}^{(k)}}\right| + \left|\frac{a_{26}^{(k)} - \eta_{0}}{\tan\theta_{26}^{(k)}}\right| \to 0 \quad \text{as } k \to \infty$$

3. Finally, we show that $\eta^{P_I} \to \eta_0$ as $\theta \to \mathbf{0}$ with $\theta \in \overline{\Theta} \setminus \{\mathbf{0}\}$. Indeed, it is clear from (A.13) that $a_{25} \to \eta_0$ as $\theta_1 \to 0^+$, and $a_{26} \to \eta_0$ as $\theta_2 \to 0^+$. Furthermore, it follows from Step 1 of the proof of Lemma 3.3 that $\min\{a_{25}, a_{26}\} \leq \eta^{P_I} \leq \max\{a_{25}, a_{26}\}$.

B. Some Known Results Needed for the Proofs

In this appendix, we present some known results that are used in Sections 3–5 above.

B.1. Well-posedness of the iteration boundary value problem. For a constant h > 0 and a function $f_{bd} : [0, h] \to \mathbb{R}_+$, fix a bounded domain $\Omega \subseteq \mathbb{R}^2$ as

(B.1)
$$\Omega \coloneqq \left\{ \boldsymbol{x} \in \mathbb{R}^2 : 0 < x_1 < h, \, 0 < x_2 < f_{\rm bd}(x_1) \right\}$$

where f_{bd} satisfies that, for constants $t_0 \ge 0$, t_h , t_1 , t_2 , t_3 , $M \in (0, \infty)$, and $\alpha \in (0, 1)$,

(B.2)
$$\begin{aligned} f_{\rm bd} &\in C^1([0,h]), \quad f_{\rm bd}(0) = t_0, \quad f_{\rm bd}(h) = t_h, \\ f_{\rm bd}(x_1) &\geq \min\{t_0 + t_1x_1, t_2, t_h - t_3(x_1 - h)\} \quad \text{for all } x_1 \in (0,h), \\ \|f_{\rm bd}\|_{2,\alpha,(0,h)}^{(-1-\alpha),\{0,h\}} &\leq M. \end{aligned}$$

The boundary, $\partial \Omega \coloneqq \bigcup_{k=0}^{3} (\Gamma_k \cup \{P_{k+1}\})$, with vertices and segments:

$$P_1 = (h, 0), \quad P_2 = (h, f_{bd}(h)), \quad P_3 = (0, f_{bd}(0)), \quad P_4 = (0, 0),$$

(B.3)
$$\Gamma_0 = \partial \Omega \cap \{x_1 = 0\}, \quad \Gamma_1 = \partial \Omega \cap \{x_2 = f_{bd}(x_1)\}, \\ \overline{\Gamma_2} = \partial \Omega \cap \{x_1 = h\}, \quad \overline{\Gamma_3} = \partial \Omega \cap \{x_2 = 0\},$$

and Γ_k , k = 0, 1, 2, 3, are the relative interiors of the segments defined above.

Let $g_{so} \in C^{\infty}(\mathbb{R}^2 \setminus \{\frac{h}{3} < x_1 < \frac{2h}{3}\})$ be a piecewise smooth function defined in \mathbb{R}^2 such that

(B.4)

$$\begin{aligned} \|g_{so}\|_{C^{3}(\overline{\Omega\setminus\{\frac{h}{3} < x_{1} < \frac{2h}{3}\})} &\leq C_{so}, \\ g_{so}(\cdot, x_{2}) \quad \text{is linear on } x_{1} \text{ in } \{x_{1} \leq \frac{h}{4}\} \cup \{x_{1} \geq \frac{3h}{4}\} \\ \partial_{x_{2}}g_{so} &= 0 \quad \text{on } \Gamma_{3}. \end{aligned}$$

We consider the following nonlinear problem:

(B.5)
$$\begin{cases} \sum_{i,j=1}^{2} \tilde{A}_{ij}(Du, \boldsymbol{x}) D_{ij}u + \sum_{i=1}^{2} \tilde{A}_{i}(Du, \boldsymbol{x}) D_{i}u = 0 & \text{ in } \Omega, \\ \tilde{B}(Du, u, \boldsymbol{x}) = 0 & \text{ on } \Gamma_{1}, \\ u = g_{\text{so}}(\boldsymbol{x}) & \text{ on } \Gamma_{2} \cup \Gamma_{0}, \\ \boldsymbol{b}^{(\text{w})}(\boldsymbol{x}) \cdot Du = 0 & \text{ on } \Gamma_{3}. \end{cases}$$

For constants $\lambda \in (0,1)$, $M < \infty$, $\alpha \in (0,1)$, $\beta \in [\frac{1}{2},1)$, $\delta \in (0,1)$, $\sigma \in (0,1)$, and $\varepsilon \in (0,\frac{h}{10})$, assume that the following conditions are satisfied:

(i) For any $\boldsymbol{x} \in \Omega$ and $\boldsymbol{p}, \boldsymbol{\mu} \in \mathbb{R}^2$, $\lambda \operatorname{dist}(\boldsymbol{x}, \Gamma_2 \cup \Gamma_0) |\boldsymbol{\mu}|^2 \leq \sum_{i,j=1}^2 \tilde{A}_{ij}(\boldsymbol{p}, \boldsymbol{x}) \mu_i \mu_j \leq \lambda^{-1} |\boldsymbol{\mu}|^2$. Moreover, for any $\boldsymbol{x} \in \Omega \setminus \{\frac{\varepsilon}{2} \leq x_1 \leq h - \frac{\varepsilon}{2}\}$,

$$\lambda |\boldsymbol{\mu}|^2 \le \sum_{i,j=1}^2 \frac{\tilde{A}_{ij}(\boldsymbol{p}, \boldsymbol{x}) \mu_i \mu_j}{\left(\min\{x_1, h - x_1, \delta\}\right)^{2 - \frac{i+j}{2}}} \le \lambda^{-1} |\boldsymbol{\mu}|^2.$$

(ii) $(\tilde{A}_{ij}, \tilde{A})(\boldsymbol{p}, \boldsymbol{x})$ are independent of $\boldsymbol{p} \in \mathbb{R}^2$ on $\Omega \cap \{\varepsilon \leq x_1 \leq h - \varepsilon\}$ with

$$\|\tilde{A}_{ij}\|_{L^{\infty}(\Omega \cap \{\varepsilon \le x_1 \le h - \varepsilon\})} \le \lambda^{-1}, \qquad \|(\tilde{A}_{ij}, \tilde{A})\|_{1,\alpha,\Omega \cap \{\varepsilon \le x_1 \le h - \varepsilon\}} \le M.$$

(iii) For any $\boldsymbol{p} \in \mathbb{R}^2$,

$$\|(\tilde{A}_{ij}, \tilde{A}_i)(\boldsymbol{p}, \cdot)\|_{C^{\beta}(\overline{\Omega} \setminus \{2\varepsilon \le x_1 \le h - 2\varepsilon\})} + \|D_{\boldsymbol{p}}(\tilde{A}_{ij}, \tilde{A}_i)(\boldsymbol{p}, \cdot)\|_{L^{\infty}(\Omega \setminus \{2\varepsilon \le x_1 \le h - 2\varepsilon\})} \le M.$$

(iv) $(\tilde{A}_{ij}, \tilde{A}_i) \in C^{1,\alpha}(\mathbb{R}^2 \times (\overline{\Omega} \setminus (\overline{\Gamma}_2 \cup \overline{\Gamma}_0)))$ and, for any $s \in (0, \frac{h}{4})$,

$$\left\| \left(\tilde{A}_{ij}, \tilde{A}_i \right) \right\|_{1,\alpha, \mathbb{R}^2 \times (\overline{\Omega} \cap \{ s \le x_1 \le h - s \})} \le M \left(\frac{h}{s} \right)^M$$

(v) For each $(\boldsymbol{p}, \boldsymbol{x}) \in \mathbb{R}^2 \times (\overline{\Omega} \setminus \{\varepsilon < x_1 < h - \varepsilon\})$ and i, j = 1, 2, define

$$\hat{\boldsymbol{p}} \coloneqq \boldsymbol{p} - Dg_{\mathrm{so}}(\boldsymbol{x}), \qquad (\tilde{a}_{ij}, \tilde{a}_i)(\hat{\boldsymbol{p}}, \boldsymbol{x}) \coloneqq (\tilde{A}_{ij}, \tilde{A}_i)(\hat{\boldsymbol{p}} + Dg_{\mathrm{so}}(\boldsymbol{x}), \boldsymbol{x}).$$

For any $(\boldsymbol{p}, (x_1, 0)) \in \mathbb{R}^2 \times (\Gamma_3 \setminus \{\varepsilon \le x_1 \le h - \varepsilon\}),\$

$$(\tilde{a}_{11}, \tilde{a}_{22}, \tilde{a}_1)((\hat{p}_1, -\hat{p}_2), (x_1, 0)) = (\tilde{a}_{11}, \tilde{a}_{22}, \tilde{a}_1)((\hat{p}_1, \hat{p}_2), (x_1, 0))$$

For any $(\boldsymbol{p}, \boldsymbol{x}) \in \mathbb{R}^2 \times (\Omega \setminus \{\varepsilon \leq x_1 \leq h - \varepsilon\})$ and i = 1, 2,

$$|\tilde{a}_{ii}(\hat{\boldsymbol{p}},\boldsymbol{x}) - \tilde{a}_{ii}(Dg_{\rm so}(\boldsymbol{x}_{\rm e}),\boldsymbol{x}_{\rm e})| \leq M |\boldsymbol{x} - \boldsymbol{x}_{\rm e}|^{\beta}, \qquad (\tilde{A}_{12},\tilde{A}_{21})(\boldsymbol{p},\boldsymbol{x}_{\rm e}) = 0,$$

either for point $\mathbf{x}_{e} \coloneqq (0, x_{2})$ or point $\mathbf{x}_{e} \coloneqq (h, x_{2})$.

- (vi) For any $\boldsymbol{p} \in \mathbb{R}^2$ and $\boldsymbol{x} \in \Omega \setminus \{\frac{\varepsilon}{2} \leq x_1 \leq h \frac{\varepsilon}{2}\}, \ \tilde{A}_1(\boldsymbol{p}, \boldsymbol{x}) \leq -\lambda$. (vii) The nonlinear boundary condition (B.5)₂ satisfies the following conditions: (vii–a) For any $(\boldsymbol{p}, z, \boldsymbol{x}) \in \mathbb{R}^2 \times \mathbb{R} \times \Gamma_1$, $D_{\boldsymbol{p}} \tilde{B}(\boldsymbol{p}, z, \boldsymbol{x}) \cdot \boldsymbol{\nu}^{(1)}(\boldsymbol{x}) \geq \lambda$, where $\boldsymbol{\nu}^{(1)}$ is the unit normal vector on Γ_1 pointing to Ω .

(vii–b) For any $(\boldsymbol{p}, z) \in \mathbb{R}^2 \times \mathbb{R}$ and k = 1, 2, 3,

$$\begin{split} \|\tilde{B}(Dg_{\mathrm{so}}, g_{\mathrm{so}}, \cdot)\|_{C^{3}(\overline{\Omega}\setminus\{\frac{h}{3}\leq x_{1}\leq \frac{2h}{3}\})} + \|D_{(\boldsymbol{p}, z)}^{k}\tilde{B}(\boldsymbol{p}, z, \cdot)\|_{C^{3}(\overline{\Omega})} \leq M, \\ \|D_{\boldsymbol{p}}\tilde{B}(\boldsymbol{p}, z, \cdot)\|_{C^{0}(\overline{\Omega})} \leq \lambda^{-1}, \\ D_{z}\tilde{B}(\boldsymbol{p}, z, \boldsymbol{x}) \leq -\lambda \qquad \text{for all } \boldsymbol{x} \in \Gamma_{1}, \\ D_{p_{1}}\tilde{B}(\boldsymbol{p}, z, \boldsymbol{x}) \leq -\lambda \qquad \text{for all } \boldsymbol{x} \in \Gamma_{1} \setminus \{\varepsilon \leq x_{1} \leq h - \varepsilon\}. \end{split}$$

(vii–c) There exist $v \in C^3(\overline{\Gamma_1})$ and a nonhomogeneous linear operator:

$$L(\mathbf{p}, z, \mathbf{x}) = \mathbf{b}^{(1)}(\mathbf{x}) \cdot \mathbf{p} + b_0^{(1)}(\mathbf{x})z + g_1(\mathbf{x})$$

defined for $\boldsymbol{x} \in \Gamma_1$ and $(\boldsymbol{p}, z) \in \mathbb{R}^2 \times \mathbb{R}$, satisfying

$$\|v\|_{C^3(\Omega)} + \|(\boldsymbol{b}^{(1)}, b_0^{(1)}, g_1)\|_{C^3(\overline{\Gamma_1})} \le M,$$

such that, for any $(\boldsymbol{p}, z, \boldsymbol{x}) \in \mathbb{R}^2 \times \mathbb{R} \times \Gamma_1$,

$$\begin{split} \left| \tilde{B}(\boldsymbol{p}, z, \boldsymbol{x}) - L(\boldsymbol{p}, z, \boldsymbol{x}) \right| &\leq \sigma \left(|\boldsymbol{p} - Dv(\boldsymbol{x})| + |z - v(\boldsymbol{x})| \right), \\ \left| D_{\boldsymbol{p}} \tilde{B}(\boldsymbol{p}, z, \boldsymbol{x}) - \boldsymbol{b}^{(1)}(\boldsymbol{x}) \right| + \left| D_{z} \tilde{B}(\boldsymbol{p}, z, \boldsymbol{x}) - b^{(1)}_{0}(\boldsymbol{x}) \right| &\leq \sigma \,. \end{split}$$

(vii–d) Obliqueness requirements: For the interior unit normal vector $\boldsymbol{\nu}^{(w)}$ on Γ_3 to Ω ,

$$\begin{split} \boldsymbol{b}^{(\mathrm{w})} \cdot \boldsymbol{\nu}^{(\mathrm{w})} &\geq \lambda \,, \quad b_1^{(\mathrm{w})} \leq 0 \qquad \text{on } \Gamma_3 \,, \\ \boldsymbol{b}^{(\mathrm{w})} &= (0,1) \ \text{on } \Gamma_3 \setminus \{ \varepsilon \leq x_1 \leq h - \varepsilon \} \,, \qquad \| \boldsymbol{b}^{(\mathrm{w})} \|_{C^3(\overline{\Gamma_3})} \leq M \end{split}$$

(vii–e) $\tilde{B}(\mathbf{0}, 0, \cdot) \equiv 0$ on $\Gamma_1 \setminus \{\varepsilon \le x_1 \le h - \varepsilon\}$.

Proposition B.1 ([10, Proposition 4.7.2]). Fix constants $h, t_1, t_2, t_3 \in (0, \infty)$ and $t_0, t_h \in [0, \infty)$. For constants $\lambda > 0$, $M < \infty$, $\alpha \in (0, 1)$, and $C_{so} > 0$, domain $\Omega \subseteq \mathbb{R}^2$ satisfies conditions (B.1)–(B.4). For $\beta \in [\frac{1}{2}, 1)$, $\delta \in (0, 1)$, $\sigma \in (0, 1)$, and $\varepsilon \in (0, \frac{h}{10})$, the nonlinear boundary value problem (B.5) satisfies conditions (i)–(vii). Then there exist $\alpha_1 \in (0, \frac{1}{2})$ depending only on λ , and constants $\delta_0, \sigma \in (0, 1)$ depending only on $(\lambda, M, C_{so}, \alpha, \beta, \varepsilon)$ such that, under further requirement $\delta \in [0, \delta_0)$, the boundary value problem (B.5) has a unique solution $u \in C(\overline{\Omega}) \cap C^1(\overline{\Omega} \setminus (\overline{\Gamma_2} \cup \overline{\Gamma_0})) \cap C^2(\Omega)$. Moreover, u satisfies

(B.6)
$$\|u\|_{0,\overline{\Omega}} \leq C, \qquad |u(\boldsymbol{x}) - g_{\rm so}(\boldsymbol{x})| \leq C \min\{x_1, h - x_1\} \quad in \ \Omega,$$

where C > 0 depends only on $(\lambda, M, C_{so}, \varepsilon)$. Furthermore, $u \in C(\overline{\Omega}) \cap C^{2,\alpha_1}(\overline{\Omega} \setminus \overline{\Gamma_2} \cup \overline{\Gamma_0})$ satisfies (B.7) $\|u\|_{2,\alpha_1,\overline{\Omega} \cap \{s < x_1 < h - s\}} \leq C_s$

for each $s \in (0, \frac{h}{10})$, where $C_s > 0$ depends only on $(\lambda, M, C_{so}, \alpha, \beta, \varepsilon, s)$.

Proposition B.2 ([10, Proposition 4.8.7]). Fix constants $h, t_1, t_2, t_3 \in (0, \infty)$, $t_0 = 0$, and $t_h \ge 0$. For constants $\lambda > 0$, $M < \infty$, $\alpha \in (0, 1)$, and $C_{so} > 0$, domain $\Omega \subseteq \mathbb{R}^2$ satisfies conditions (B.1)–(B.4) with changes: $P_3 = P_4 = (0, 0)$ and $\overline{\Gamma_0} = \{(0, 0)\}$. For $\beta \in [\frac{1}{2}, 1)$, $\delta \in (0, 1)$, $\sigma \in (0, 1)$, and $\varepsilon \in (0, \frac{h}{10})$, the nonlinear boundary value problem (B.5) satisfies conditions (ii), (iv), and (vii) above, and (i^{*}), (iii^{*}), (v^{*}), and (vi^{*}) below:

(i*) For any $\boldsymbol{x} \in \Omega$ and $\boldsymbol{p}, \boldsymbol{\kappa} = (\kappa_1 \kappa_2) \in \mathbb{R}^2$,

$$\min\left\{\lambda \operatorname{dist}(\boldsymbol{x}, \overline{\Gamma_0}) + \delta, \lambda \operatorname{dist}(\boldsymbol{x}, \Gamma_2)\right\} |\boldsymbol{\kappa}|^2 \leq \sum_{i,j=1}^2 \tilde{A}_{ij}(\boldsymbol{p}, \boldsymbol{x}) \kappa_i \kappa_j \leq \lambda^{-1} |\boldsymbol{\kappa}|^2,$$
$$\left\| \left((\tilde{A}_{ij}, \tilde{A}_i) (Dg_{\mathrm{so}}, \cdot), D_{\boldsymbol{p}}^m (\tilde{A}_{ij}, \tilde{A}_i) (\boldsymbol{p}, \cdot) \right) \right\|_{1,\alpha,\Omega \cap \{x_1 < 2\varepsilon\}}^{(-\alpha), \{P_4\}} \leq M \quad \text{for } m = 1, 2.$$

Moreover, for any $\boldsymbol{x} \in \Omega \cap \{h - \frac{\varepsilon}{2} < x_1 < h\},\$

$$\lambda |\mu|^2 \le \sum_{i,j=1}^2 \frac{\tilde{A}_{ij}(\boldsymbol{p}, \boldsymbol{x}) \mu_i \mu_j}{(\min\{h - x_1, \delta\})^{2 - \frac{i+j}{2}}} \le \lambda^{-1} |\mu|^2.$$

(iii*) For any $\boldsymbol{p} \in \mathbb{R}^2$,

$$\|(A_{ij}, A_i)(\boldsymbol{p}, \cdot)\|_{C^{\beta}(\overline{\Omega \cap \{h-2\varepsilon < x_1 < h\}})} + \|D_{\boldsymbol{p}}(A_{ij}, A_i)(\boldsymbol{p}, \cdot)\|_{L^{\infty}(\Omega \cap \{h-2\varepsilon < x_1 < h\})} \le M.$$

(v*) For each $(\boldsymbol{p}, \boldsymbol{x}) \in \mathbb{R}^2 \times (\overline{\Omega} \cap \{h - \varepsilon < x_1 < h\})$ and i, j = 1, 2, define

$$\hat{\boldsymbol{p}} \coloneqq \boldsymbol{p} - Dg_{\mathrm{so}}(\boldsymbol{x}), \qquad (\tilde{a}_{ij}, \tilde{a}_i)(\hat{\boldsymbol{p}}, \boldsymbol{x}) \coloneqq (\tilde{A}_{ij}, \tilde{A}_i)(\hat{\boldsymbol{p}} + Dg_{\mathrm{so}}(\boldsymbol{x}), \boldsymbol{x}).$$

For any $(\mathbf{p}, (x_1, 0)) \in \mathbb{R}^2 \times (\Gamma_3 \cap \{h - \varepsilon < x_1 < h\}),$

$$(\tilde{a}_{11}, \tilde{a}_{22}, \tilde{a}_1)((\hat{p}_1, -\hat{p}_2), (x_1, 0)) = (\tilde{a}_{11}, \tilde{a}_{22}, \tilde{a}_1)((\hat{p}_1, \hat{p}_2), (x_1, 0)).$$

For any
$$(\boldsymbol{p}, \boldsymbol{x}) \in \mathbb{R}^2 \times (\Omega \cap \{h - \varepsilon < x_1 < h\})$$
 and $i = 1, 2,$
 $|\tilde{a}_{ii}(\hat{\boldsymbol{p}}, \boldsymbol{x}) - \tilde{a}_{ii}(Dg_{so}(\boldsymbol{x}_e), \boldsymbol{x}_e)| \le M |\boldsymbol{x} - \boldsymbol{x}_e|^{\beta}, \quad (\tilde{A}_{12}, \tilde{A}_{21})(\boldsymbol{p}, \boldsymbol{x}_e) = 0 \quad for \ \boldsymbol{x}_e \coloneqq (h, x_2)$

(vi*) For any $\boldsymbol{p} \in \mathbb{R}^2$ and $\boldsymbol{x} \in \Omega \cap \{h - \frac{\varepsilon}{2} < x_1 < h\}, \ \tilde{A}_1(\boldsymbol{p}, \boldsymbol{x}) \leq -\lambda$.

Then there exist both $\alpha_1 \in (0, \frac{1}{2})$ depending only on (λ, δ) and $\sigma \in (0, 1)$ depending only on $(\lambda, \delta, M, C_{so}, \alpha, \beta, \varepsilon)$ such that, under the further requirement $\delta \in [0, \delta_0)$, the boundary value problem (B.5) has a unique solution $u \in C(\overline{\Omega}) \cap C^1(\overline{\Omega} \setminus (\overline{\Gamma_2} \cup \overline{\Gamma_0})) \cap C^2(\Omega)$. Moreover, u satisfies

(B.8)
$$||u||_{0,\overline{\Omega}} \leq C$$
, $|u(\boldsymbol{x}) - g_{so}(\boldsymbol{x})| \leq C \min\{x_1, h - x_1\}$ in Ω ,

where C > 0 depends only on $(\lambda, \delta, M, C_{so}, \varepsilon)$. Furthermore, $u \in C(\overline{\Omega}) \cap C^{2,\alpha_1}(\overline{\Omega} \setminus \overline{\Gamma_2} \cup \overline{\Gamma_0})$ satisfies

(B.9)
$$\|u\|_{2,\alpha_1,\overline{\Omega\cap\{s< x_1< h-s\}}} \le C_s$$

for each $s \in (0, \frac{h}{10})$, where $C_s > 0$ depends only on $(\lambda, \delta, M, C_{so}, \alpha, \beta, \varepsilon, s)$. In addition, u satisfies

$$\|u\|_{2,\alpha_1,\Omega\cap\{x_1<\frac{h}{4}\}}^{(-1-\alpha_1),\{P_4\}} < \hat{C},$$

for constant $\hat{C} > 0$ depending only on $(\lambda, \delta, M, \alpha, \varepsilon)$.

Proposition B.3. Fix constants $h, t_1, t_2, t_3 \in (0, \infty)$ and $t_0 = t_h = 0$. For constants $\lambda > 0$, $M < \infty$, $\alpha \in (0, 1)$, and $C_{so} > 0$, domain $\Omega \subseteq \mathbb{R}^2$ satisfies conditions (B.1)–(B.4) with changes:

$$P_3 = P_4 = (0,0), \quad \overline{\Gamma_0} = \{(0,0)\}; \qquad P_2 = P_1 = (h,0), \quad \overline{\Gamma_2} = \{(h,0)\}.$$

For $\beta \in [\frac{1}{2}, 1)$, $\delta \in (0, 1)$, $\sigma \in (0, 1)$, and $\varepsilon \in (0, \frac{h}{10})$, the nonlinear boundary value problem (B.5) satisfies conditions (ii), (iv), and (vii) above, and (i^{*}) below:

(i*) For any $\boldsymbol{x} \in \Omega$ and $\boldsymbol{p}, \boldsymbol{\kappa} = (\kappa_1, \kappa_2) \in \mathbb{R}^2$,

$$\min\left\{\lambda \operatorname{dist}(\boldsymbol{x}, \overline{\Gamma_0}) + \delta, \lambda \operatorname{dist}(\boldsymbol{x}, \Gamma_2) + \delta\right\} |\boldsymbol{\kappa}|^2 \leq \sum_{i,j=1}^2 \tilde{A}_{ij}(\boldsymbol{p}, \boldsymbol{x}) \kappa_i \kappa_j \leq \lambda^{-1} |\boldsymbol{\kappa}|^2,$$
$$\left\| \left((\tilde{A}_{ij}, \tilde{A}_i) (Dg_{\mathrm{so}}, \cdot), D_{\boldsymbol{p}}^m (\tilde{A}_{ij}, \tilde{A}_i) (\boldsymbol{p}, \cdot) \right) \right\|_{1,\alpha,\Omega \setminus \{2\varepsilon < x_1 < h - 2\varepsilon\}}^{(-\alpha), \{P_1, P_4\}} \leq M \quad \text{for } m = 1, 2.$$

Then there exist constants $\alpha_1 \in (0, \frac{1}{2})$ depending only on (λ, δ) , and $\sigma \in (0, 1)$ depending only on $(\lambda, \delta, M, C_{so}, \alpha, \beta, \varepsilon)$ such that, under the further requirement $\delta \in [0, \delta_0)$, the boundary value problem (B.5) has a unique solution $u \in C(\overline{\Omega}) \cap C^1(\overline{\Omega} \setminus (\overline{\Gamma_2} \cup \overline{\Gamma_0})) \cap C^2(\Omega)$. Moreover, u satisfies

(B.10)
$$\|u\|_{0,\overline{\Omega}} \leq C, \qquad |u(\boldsymbol{x}) - g_{\rm so}(\boldsymbol{x})| \leq C \min\{x_1, h - x_1\} \quad in \ \Omega,$$

where C > 0 depends only on $(\lambda, \delta, M, C_{so}, \varepsilon)$. Furthermore, $u \in C(\overline{\Omega}) \cap C^{2,\alpha_1}(\overline{\Omega} \setminus \overline{\Gamma_2} \cup \overline{\Gamma_0})$ satisfies (B.11) $\|u\|_{2,\alpha_1,\overline{\Omega} \cap \{s < x_1 < h - s\}} \leq C_s$

for each $s \in (0, \frac{h}{10})$, where $C_s > 0$ depends only on $(\lambda, \delta, M, C_{so}, \alpha, \beta, \varepsilon, s)$. In addition, u satisfies

$$\|u\|_{2,\alpha_1,\Omega\setminus\{\frac{h}{4}\leq x_1\leq\frac{3h}{4}\}}^{(-1-\alpha_1),\{P_1,P_4\}} < \hat{C},$$

for constant $\hat{C} > 0$ depending only on $(\lambda, \delta, M, \alpha, \varepsilon)$.

B.2. Regularized distance function.

Lemma B.4 ([10, Lemmas 13.9.1]). For any $g(\cdot) \in C^{0,1}([-1,1])$ that is positive on (-1,1), there exists a function $\delta_g \in C^{\infty}(\overline{R_{\infty}} \setminus \overline{R_g})$, the regularized distance, such that

- (i) $|\delta_g(s,t') \operatorname{dist}((s,t'),\Sigma_g)| \leq \frac{1}{2}\operatorname{dist}(\mathbf{x},\Sigma_g)$ for all $(s,t') \in \mathbb{R}^2 \setminus \overline{\Sigma_g}$;
- (ii) $|D^m \delta_g(s,t')| \leq C(m) (\operatorname{dist}((s,t'), \Sigma_g))^{1-m}$ for all $(s,t') \in \mathbb{R}^2 \setminus \overline{\Sigma_g}$ and $m = 1, 2, \cdots$, where C(m) > 0 depends only on m;
- (iii) $\delta_g(s,t') \ge C_{\mathrm{rd}}(t'-g(s))$ for all $(s,t') \in \overline{R_g} \setminus \overline{\Sigma_g}$, where $C_{\mathrm{rd}} > 0$ depends only on $\mathrm{Lip}[g]$;
- (iv) Let $g_k, g \in C^{0,1}([-1,1])$ be uniformly bounded Lipschitz functions for all $k \in \mathbb{N}$, with $g_k(s)$ converging to g(s) uniformly on [-1,1]. Then $\lim_{k \to \infty} \|\delta_{g_k}(s,t') - \delta_g(s,t')\|_{C^m(K)} = 0$ for any $m = 0, 1, 2, \cdots$, and any compact set $K \subseteq \overline{R_\infty} \setminus \overline{R_g}$.

B.3. Leray-Schauder degree theorem.

Definition B.5 (Compact map). Let X and Y be Banach spaces. Fix any open subset $G \subseteq X$. A map $\mathbf{f} : \overline{G} \to Y$ is called compact if

- (i) **f** is continuous;
- (ii) $\mathbf{f}(U)$ is precompact in Y for any bounded subset $U \subseteq \overline{G}$.

Definition B.6. Let X be a Banach space, and let G be a bounded open subset of X. Write V(G, X) to denote the set of all maps $\mathbf{f} : \overline{G} \to X$ satisfying

- (i) **f** is a compact map in the sense of Definition B.5;
- (ii) **f** has no fixed point on the boundary of ∂G .

Definition B.7. Let X be a Banach space, and let G be a bounded open subset of X. Two maps $\mathbf{f}, \mathbf{g} \in V(G, X)$ are called compactly homotopic on ∂G if there exists a map \mathbf{H} with the following properties:

- (i) $\mathbf{H}: \overline{G} \times [0,1] \to X$ is a compact map;
- (ii) $\mathbf{H}(\mathbf{x},\tau) \neq \mathbf{x}$ for all $(\mathbf{x},\tau) \in \partial G \times [0,1];$
- (iii) $\mathbf{H}(\mathbf{x}, 0) = \mathbf{f}(\mathbf{x})$ and $\mathbf{H}(\mathbf{x}, 1) = \mathbf{g}(\mathbf{x})$ for all $\mathbf{x} \in \overline{G}$.

We write $\partial G : \mathbf{f} \cong \mathbf{g}$ if \mathbf{f} and \mathbf{g} are compactly homotopic on ∂G in the sense of Definition B.7, and we call \mathbf{H} a compact homotopy.

Theorem B.1 (Leray-Schauder degree theorem). Let X be a Banach space, and let G be a bounded open subset of X. Then, for each map $\mathbf{f} \in V(G, X)$, a unique integer $\mathbf{Ind}(\mathbf{f}, G)$ can be assigned with the following properties:

- (i) If $\mathbf{f}(\cdot) \equiv \mathbf{x}_0$ on \overline{G} for some fixed $\mathbf{x}_0 \in G$, then $\mathbf{Ind}(\mathbf{f}, G) = 1$;
- (ii) If $\mathbf{Ind}(\mathbf{f}, G) \neq 0$, then there exists a fixed point $\mathbf{x} \in G$ of map \mathbf{f} , that is, $\mathbf{f}(\mathbf{x}) = \mathbf{x}$;
- (iii) $\operatorname{Ind}(\mathbf{f}, G) = \sum_{j=1}^{n} \operatorname{Ind}(\mathbf{f}, G_j)$, whenever $\mathbf{f} \in V(G, X) \cap (\bigcap_{j=1}^{n} V(G_j, X))$, where $G_i \cap G_j = \emptyset$ for $i \neq j$ and $\overline{G} = \bigcup_{i=1}^{n} \overline{G_i}$;
- (iv) If $\partial G : \mathbf{f} \cong \mathbf{g}$, then $\mathbf{Ind}(\mathbf{f}, G) = \mathbf{Ind}(\mathbf{g}, G)$.

Such a number $\mathbf{Ind}(\mathbf{g}, G)$ is called the fixed point index of \mathbf{f} over G.

A generalized homotopy invariance of the fixed point index is given by the following theorem.

Theorem B.2 ([48, §13.6, A4^{*}]). Let X be a Banach space, and let $t_2 > t_1$. Let $U \subseteq X \times [t_1, t_2]$, and let $U_t = \{\mathbf{x} : (\mathbf{x}, t) \in U\}$. Then, for any compact map $\mathbf{h} : \overline{U} \to X$ with $\mathbf{h}(\mathbf{x}, t) \neq \mathbf{x}$ on ∂U ,

$$\operatorname{Ind}(\mathbf{h}(\cdot, t), U_t) = \operatorname{const.} \quad for \ all \ t \in [t_1, t_2],$$

provided that U is a bounded open set in $X \times [t_1, t_2]$.

B.4. Regularity theorem.

Theorem B.3 ([1, Theorem 3.1]). For constants r, R > 0, define $Q_{r,R}^+$ by

$$Q_{r,R}^{+} \coloneqq \left\{ (x,y) \in \mathbb{R}^{2} : 0 < x < r, |y| < R \right\}$$

For positive constants a, b, M, N, and $\kappa \in (0, \frac{1}{4})$, suppose that $\psi \in C(\overline{Q_{r,R}^+}) \cap C^2(Q_{r,R}^+)$ satisfies $\psi > 0, -Mx \le \psi_x \le \frac{2-\kappa}{a}x$, and

$$(2x - a\psi_x + O_1)\psi_{xx} + O_2\psi_{xy} + (b + O_3)\psi_{yy} - (1 + O_4)\psi_x + O_5\psi_y = 0 \qquad in \ Q_{r,R}^+,$$

and $\psi = 0$ on $\partial Q_{r,R}^+ \cap \{x = 0\}$, where $O_k(x, y)$, $k = 1, \dots, 5$, are continuously differentiable and

$$\frac{|O_1(x,y)|}{x^2} + \frac{|D_{(x,y)}O_1(x,y)|}{x} + \sum_{k=2}^5 \left(\frac{|O_k(x,y)|}{x} + |D_{(x,y)}O_k(x,y)|\right) \le N \qquad \text{in } Q_{r,R}^+.$$

Then $\psi \in C^{2,\alpha}(\overline{Q^+_{r/2,R/2}})$ for any $\alpha \in (0,1)$ and

$$\psi_{xx}(0,y) = a^{-1}, \quad \psi_{xy}(0,y) = \psi_{yy}(0,y) = 0 \quad \text{for all } y \in \left(-\frac{R}{2}, \frac{R}{2}\right)$$

B.5. General framework for the convexity of transonic shocks. We state a general framework for the transonic shock as a free boundary, developed in Chen-Feldman-Xiang [12].

Let Ω be an open, bounded, and connected set, and $\partial \Omega = \Gamma_{\text{shock}} \cup \Gamma_1 \cup \Gamma_2$, where the closed curve segment Γ_{shock} is a transonic shock that separates a pseudo-supersonic constant state denoted as state (2) outside Ω from a pseudo-subsonic (non-constant) state inside Ω , and $\Gamma_1 \cup \Gamma_2$ is a fixed boundary.

We now present a more structural framework for domain Ω under consideration.

Framework (A). The structural framework for domain Ω :

- (i) Domain Ω is bounded, with boundary $\partial\Omega$ that is a continuous closed curve without selfintersections. Furthermore, $\partial\Omega$ is piecewise $C^{1,\alpha}$ up to the endpoints of each smooth part for some $\alpha \in (0, 1)$, and the number of smooth parts is finite.
- (ii) At each corner point of $\partial\Omega$, angle θ between the arcs meeting at that point from the interior of Ω satisfies $\theta \in (0, \pi)$.
- (iii) $\partial \Omega = \Gamma_{\text{shock}} \cup \Gamma_1 \cup \Gamma_2$, where Γ_{shock} , Γ_1 , and Γ_2 are connected and disjoint, and both Γ^0_{shock} and $\Gamma_1 \cup \Gamma_2$ are non-empty. Moreover, if $\Gamma_i \neq \emptyset$ for some $i \in \{1, 2\}$, then its relative interior is nonempty, *i.e.*, $\Gamma^0_i \neq \emptyset$.
- (iv) Γ_{shock} includes its endpoints A and B with corresponding unit tangent vectors τ_A and τ_B pointing to the interior of Γ_{shock} , respectively. If $\Gamma_1 \neq \emptyset$, then A is a common endpoint of Γ_{shock} and Γ_1 . If $\Gamma_2 \neq \emptyset$, then B is a common endpoint of Γ_{shock} and Γ_2 .

Let $\phi = \varphi - \varphi_2$. Then ϕ satisfies the following equation:

(B.12)
$$(c^2 - \varphi_{\xi_1}^2)\phi_{\xi_1\xi_1} - 2\varphi_{\xi_1}\varphi_{\xi_2}\phi_{\xi_1\xi_2} + (c^2 - \varphi_{\xi_2}^2)\phi_{\xi_2\xi_2} = 0 \quad \text{in } \Omega ,$$

with the boundary conditions:

(B.13)
$$\phi = 0, \quad \rho(|D\phi + D\varphi_2|^2, \phi + \varphi_2)D(\phi + \varphi_2) \cdot \boldsymbol{\nu} = \rho_2 D\varphi_2 \cdot \boldsymbol{\nu} \quad \text{on } \Gamma_{\text{shock}}$$

If $\tau_A \neq \pm \tau_B$, define the cone:

(B.14)
$$\operatorname{Con} \coloneqq \{ r \boldsymbol{\tau}_A + s \boldsymbol{\tau}_B : r, s > 0 \}$$

Theorem B.4 ([12, Theorem 2.1]). Assume that domain Ω satisfies Framework (A). Assume that $\phi \in C^1(\overline{\Omega}) \cap C^2(\Omega \cup \Gamma^0_{\text{shock}}) \cap C^3(\Omega)$ is a solution of (B.12)–(B.13), which is not a constant state in Ω . Moreover, let ϕ satisfy the following conditions:

- (C-1) The entropy condition holds across Γ_{shock} : $\rho(|D\varphi|^2, \varphi) > \rho_2$ and $\phi_{\nu} < 0$ along Γ_{shock} , where ν is the unit normal vector on Γ_{shock} pointing to Ω ;
- (C-2) There exist constants $C_1 > 0$ and $\alpha_1 \in (0,1)$ such that $\|\phi\|_{1+\alpha_1,\overline{\Omega}} \leq C_1$;
- (C-3) Equation (B.12) is strictly elliptic in $\Omega \cup \Gamma^0_{\text{shock}}$: $c^2 |D\varphi|^2 > 0$ in $\Omega \cup \Gamma^0_{\text{shock}}$;
- (C-4) Γ_{shock} is C^2 in its relative interior;
- (C-5) $\tau_A \neq \pm \tau_B$, and $\{P + \operatorname{Con}\} \cap \Omega = \emptyset$ for any point $P \in \overline{\Gamma_{\operatorname{shock}}};$
- (C-6) There exists a vector $e \in \text{Con such that one of the following conditions holds:}$
 - (C-6a) $\Gamma_1 \neq \emptyset$, and the directional derivative $\phi_{\boldsymbol{e}}$ cannot have a local maximum point on $\Gamma_1^0 \cup \{A\}$ and a local minimum point on Γ_2^0 ;
 - (C-6b) $\Gamma_2 \neq \emptyset$, and ϕ_e cannot have a local minimum point on Γ_1^0 and a local maximum point on $\Gamma_2^0 \cup \{B\}$;
(C-6c) $\phi_{\boldsymbol{e}}$ cannot have a local minimum point on $\Gamma_1 \cup \Gamma_2$; where all the local maximum or minimum points are relative to $\overline{\Omega}$.

Then the free boundary Γ_{shock} is a convex graph in each direction $e \in \text{Con.}$ That is, there exists a concave function $f \in C^{1,\alpha}(\mathbb{R})$ in the orthogonal coordinate system $\boldsymbol{\xi} \eqqcolon S\boldsymbol{e} + T\boldsymbol{e}^{\perp}$ such that

(B.15)
$$\Gamma_{\text{shock}} = \left\{ \boldsymbol{\xi}(S,T) \in \mathbb{R}^2 : S = f(T), T_A < T < T_B \right\},$$
$$\Omega \cap \left\{ \boldsymbol{\xi} : T_A < T < T_B \right\} \subseteq \left\{ \boldsymbol{\xi} : S < f(T) \right\}.$$

Moreover, the free boundary Γ_{shock} is strictly convex in its relative interior in the sense that, if $P = \boldsymbol{\xi}(S,T) \in \Gamma^0_{\text{shock}}$ and f''(T) = 0, then there exists an integer m > 1, independent of the choice of $e \in \text{Con such that}$, for any $n = 2, \dots, 2m - 1$,

(B.16)
$$f^{(n)}(T) = 0, \qquad f^{(2m)}(T) < 0.$$

The number of the points at which f''(T) = 0 is at most finite on each compact subset of Γ^0_{shock} . In particular, the free boundary $\Gamma_{\rm shock}$ cannot contain any straight segment.

Furthermore, under some additional assumptions, we can show that the shock curve is uniformly convex in its relative interior in the sense defined in the following theorem:

Theorem B.5 ([12, Theorem 2.3]). Let Ω and ϕ be as in Theorem B.4. Furthermore, assume that, for any unit vector $e \in \mathbb{R}^2$, the boundary part $\Gamma_1 \cup \Gamma_2$ can be further decomposed so that

- (C-7) $\Gamma_1 \cup \Gamma_2 = \hat{\Gamma}_0 \cup \hat{\Gamma}_1 \cup \hat{\Gamma}_2 \cup \hat{\Gamma}_3$, where some $\hat{\Gamma}_i$ may be empty, $\hat{\Gamma}_i$ is connected for each i = 0, 1, 2, 3, and all curves $\hat{\Gamma}_i$ are located along $\partial \Omega$ in the order of their indices, i.e., non-empty sets $\hat{\Gamma}_i$ and $\hat{\Gamma}_k$, k > j, have a common endpoint if and only if either k = j + 1 or $\hat{\Gamma}_i = \emptyset$ for all $i = j + 1, \dots, k - 1$. Also, the non-empty set $\hat{\Gamma}_i$ with the smallest (resp. largest) index has the common endpoint A (resp. B) with Γ_{shock} . Moreover, if $\hat{\Gamma}_i \neq \emptyset$ for some i = 0, 1, 2, 3, then its relative interior is nonempty: $\hat{\Gamma}_i^0 \neq \varnothing$;
- (C-8) $\phi_{\boldsymbol{e}}$ is constant along $\hat{\Gamma}_0$ and $\hat{\Gamma}_3$;
- (C-9) For i = 1, 2, if $\phi_{\boldsymbol{e}}$ attains its local minimum or maximum relative to $\overline{\Omega}$ on $\hat{\Gamma}_{i}^{0}$, then $\phi_{\boldsymbol{e}}$ is constant along $\hat{\Gamma}_i$;
- (C-10) One of the following two conditions holds:

(C-10a) Either $\hat{\Gamma}_1 = \emptyset$ or $\hat{\Gamma}_2 = \emptyset$;

- (C-10b) Both $\hat{\Gamma}_1$ and $\hat{\Gamma}_2$ are non-empty, and $\hat{\Gamma}_3 = \emptyset$, so that $\hat{\Gamma}_2$ has the common endpoint B with Γ_{shock} . At point B, the following conditions hold:
 - (1*) If $\boldsymbol{\nu}_{\rm sh}(B) \cdot \boldsymbol{e} < 0$, then $\phi_{\boldsymbol{e}}$ cannot attain its local maximum relative to $\overline{\Omega}$ at B:
 - (2*) If $\boldsymbol{\nu}_{sh}(B) \cdot \boldsymbol{e} = 0$, then $\phi_{\boldsymbol{e}}(B) = \phi_{\boldsymbol{e}}(Q^*)$ for the common endpoint Q^* of $\hat{\Gamma}_1$

 $\begin{array}{l} (\mathbf{L}^{-}) \stackrel{\text{subsect}}{\to} \text{subsect} \\ \text{and } \widehat{\Gamma}_{2}^{2}; \\ \text{where } \boldsymbol{\nu}_{\mathrm{sh}}(B) \coloneqq \lim_{\Gamma^{0}_{\mathrm{shock}} \ni P \to B} \boldsymbol{\nu}(P), \text{ which exists since } \Gamma_{\mathrm{shock}} \text{ is } C^{1} \text{ up to } B. \end{array}$

Then the shock function f(T) in (B.15) satisfies f''(T) < 0 for all $T \in (T_A, T_B)$, that is, Γ_{shock} is uniformly convex on closed subsets of its relative interior.

Acknowledgements: The research of Gui-Qiang G. Chen is supported in part by the UK Engineering and Physical Sciences Research Council Awards EP/L015811/1, EP/V008854/1, and EP/V051121/1. The research of Alex Cliffe is supported in part by the UK Engineering and Physical Sciences Research Council Awards EP/N509711/1 and EP/R513295/1. The research of Feimin Huang is supported in part by the National Key R&D Program of China No. 2021YFA1000800, and the National Natural Sciences Foundation of China No. 12288201. The research of Song Liu is supported in part by the Hong Kong Institute of Advanced Study and the GRF grant CityU 11300420. The research of Qin Wang is supported in part by the National Natural Sciences Foundation of China No. 12261100.

References

- M. Bae, G.-Q. Chen, and M. Feldman (2009). Regularity of solutions to regular shock reflection for potential flow. *Invent. Math.*, 175, no. 3: 505–543.
- [2] M. Bae, G.-Q. Chen, and M. Feldman (2023). Prandtl-Meyer Reflection Configurations, Transonic Shocks, and Free Boundary Problems. Research Monograph, 228pp, *Memoirs of the American Mathematical Society*, AMS: Providence, 2023 (to appear). [arXiv:1901.05916].
- [3] A. Bressan (2000). Hyperbolic Systems of Conservation Laws: The One-Dimensional Cauchy Problem. Oxford University Press: Oxford.
- [4] A. Bressan and P. LeFloch (1997). Uniqueness of entropy solutions for systems of conservation laws. Arch. Ration. Mech. Anal., 1140: 301–317.
- [5] A. Bressan, T.-P. Liu and T. Yang (1999). L^1 -stability estimates for $n \times n$ conservation laws. Arch. Ration. Mech. Anal., 149: 1–22.
- [6] T. Chang, G.-Q. Chen, and S. Yang (1995). On the 2-D Riemann problem for the compressible Euler equations, I. Interaction of shocks and rarefaction waves. *Discrete Continuous Dynam. Systems*, 1: 555–84.
- [7] T. Chang, G.-Q. Chen, and S. Yang (2000). On the 2-D Riemann problem for the compressible Euler equations, II. Interaction of contact discontinuities. *Discrete Continuous Dynam. Systems*, 6: 419–430.
- [8] T. Chang and L. Hsiao (1989). The Riemann Problem and Interaction of Waves in Gas Dynamics. New York: John Wiley & Sons.
- [9] G.-Q. Chen and M. Feldman (2010). Global solutions to shock reflection by large-angle wedges for potential flow. Ann. of Math., 171: 1067–1182.
- [10] G.-Q. Chen and M. Feldman (2018). Mathematics of Shock Reflection-Diffraction and von Neumann's Conjectures. Research Monograph, Annals of Mathematics Studies, 197, Princeton University Press.
- [11] G.-Q. Chen and M. Feldman (2022). Multidimensional transonic shock waves and free boundary problems. Bull. Math. Sci., 12(01): 2230002.
- [12] G.-Q. Chen, M. Feldman, and W. Xiang (2020). Convexity of self-similar transonic shocks and free boundaries for the Euler equations for potential flow. Arch. Ration. Mech. Anal., 238, No. 1: 47–124.
- [13] G.-Q. Chen, M. Feldman, and W. Xiang (2020). Uniqueness and stability for the shock reflectiondiffraction problem for potential flow. *Hyperbolic Problems: Theory, Numerics & Applications*, 2–24, AIMS Ser. Appl. Math., **10**, [arXiv:1904.00114].
- [14] G.-Q. Chen, Q. Wang, and S.-G. Zhu (2021). Global solutions of a two-dimensional Riemann problem for the pressure gradient system. Comm. Pure Appl. Anal., 20: 2475–2503.
- [15] S.-X. Chen (1992). Existence of local solution to supersonic flow around a three-dimensional wing. Adv. Appl. Math., 13(3): 273–304.
- [16] S.-X. Chen (1998). Global existence of supersonic flow past a curved convex wedge. J. Partial Diff. Eq., 11: 73–82.
- [17] S.-X. Chen and A.-F. Qu (2012). Two-dimensional Riemann problems for Chaplygin gas. SIAM J. Math. Anal., 443: 2146–2178.
- [18] R. Courant and K. O. Friedrichs (1948). Supersonic Flow and Shock Waves. Springer-Verlag: New York.
- [19] C. M. Dafermos (2016). Hyperbolic Conservation Laws in Continuum Physics. Springer-Verlag: Berlin.
- [20] S. Earnshaw (1860). On the mathematical theory of sound. Trans. Royal Soc. London, 150: 133-148.
- [21] V. Elling and T.-P. Liu (2005). The ellipticity principle for self-similar potential flow. J. Hyperbolic Diff. Eqs., 2: 909–917.
- [22] V. Elling and T.-P. Liu (2008). Supersonic flow onto a solid wedge. Comm. Pure Appl. Math., 61: 1347–1448.
- [23] D. Gilbarg and N. Trudinger (1983). Elliptic Partial Differential Equations of Second Order. 2nd Edition, Springer-Verlag: Berlin.
- [24] J. Glimm (1965). Solution in the large for nonlinear systems of conservation laws. Comm. Pure Appl. Math., 18: 695–715.
- [25] J. Glimm, X. Ji, J. Li, X. Li, P. Zhang, T. Zhang and Y. Zheng (2008). Transonic shock formation in a rarefaction Riemann problem for the 2D compressible Euler equations. SIAM J. Appl. Math., 69(3): 720–742.
- [26] J. Glimm and A. Majda (1991). Multidimensional Hyperbolic Problems and Computations. Springer-Verlag: New York.
- [27] H. Hugoniot (1889). Sur la propagation du mouvement dans les corps et spécialement dans les gaz parfaits. Journal de l'École polytechnique, 58: 1–125.
- [28] A. Kurganov and E. Tadmor (2002). Solution of two-dimensional Riemann problems for gas dynamics without Riemann problem solvers. Numer. Methods Partial Diff. Eqs., 18: 584–608.
- [29] P. D. Lax (1957). Hyperbolic systems of conservation laws II. Comm. Pure Appl. Math., 10: 537–566.
- [30] P. D. Lax (1973). Hyperbolic Systems of Conservation Laws and the Mathematical Theory of Shock Waves, CBMS-RCSM, SIAM: Philiadelphia.
- [31] P. D. Lax (2006). Hyperbolic Partial Differential Equations. Courant Lecture Notes in Mathematics, 14, New York University and American Mathematical Society.
- [32] P. D. Lax and X. Liu (1998). Solution of two-dimensional Riemann problems of gas dynamics by positive schemes. SIAM J. Sci. Comput., 19: 319–340.

- [33] J. Li, T. Zhang, and S. Yang (1998). The Two-dimensional Riemann Problem in Gas Dynamics. Pitman Monographs and Surveys in Pure and Applied Mathematics, 98, Addison Wesley Longman limited.
- [34] J. Li, T. Zhang, and Y. Zheng (2006). Simple waves and a characteristics decomposition of the two-dimensional compressible Euler equation. *Commun. Math. Phys.*, 267: 1–12.
- [35] J. Li and Y. Zheng (2009). Interaction of rarefaction waves of the two-dimensional self-similar Euler equations. Arch. Ration. Mech. Anal., 193: 623–657.
- [36] J. Li and Y. Zheng (2010). Interaction of four rarefaction waves in the bi-symmetric class of the two-dimensional Euler equations. *Commun. Math. Phys.*, **296**: 303–321.
- [37] M. Li, Y. Zheng (2011). Semi-hyperbolic patches of solutions to the two dimensional Euler equations. Arch. Ration. Mech. Anal., 201(3): 1069–1096.
- [38] G. M. Lieberman (1986). Mixed boundary value problem for elliptic and parabolic differential equations of second order. J. Math. Anal. Appl., 113: 422–440.
- [39] G. M. Lieberman (2013). Oblique Derivative Problems For Elliptic Equations. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ.
- [40] T.-P. Liu (1981). Admissible Solutions of Hyperbolic Conservation Laws. Mem. Amer. Math. Soc., 30, No. 240, AMS: Providence.
- [41] T.-P. Liu and T. Yang (1999). L^1 stability for 2×2 systems of hyperbolic conservation laws. J. Amer. Math. Soc., **12**(3): 729–774.
- [42] A. Majda (1984). Compressible Fluid Flow and Systems of Conservation Laws in Several Space Variables. Springer-Verlag: New York.
- [43] W. J. M. Rankine (1870). On the thermodynamic theory of waves of finite longitudinal disturbance. Trans. Royal Soc. London, 160: 277–288.
- [44] B. Riemann (1860). Uber die Fortpflanzung ebener Luftvellen von endlicher Schwingungsweite. *Gött. Abh. Math. Cl.*, **8**: 43–65.
- [45] C. W. Schulz-Rinne (1993). Classification of the Riemann problem for two-dimensional gas dynamics. SIAM J. Math. Anal., 24: 76–88.
- [46] D. Serre (2009). Multidimensional shock interaction for a Chaplygin gas. Arch. Ration. Mech. Anal., 191: 539–577.
- [47] G. G. Stokes (1848). On a difficulty in the theory of sound. *Philosophical Magazine* [3], **33**: 349–356.
- [48] E. Zeidler (1986). Nonlinear Functional Analysis and Its Applications. I Fixed Point Theorems. Translated by P. R. Wadsack, Springer-Verlag: New York.
- [49] T. Zhang and Y. Zheng (1990). Conjecture of the structure of solutions of the Riemann problem for twodimensional gas dynamics systems. SIAM J. Math. Anal., 21: 593–630.

(Gui-Qiang G. Chen) MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, OXFORD, OX2 6GG, UK *Email address*: chengq@maths.ox.ac.uk

(Alex Cliffe) MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, OXFORD, OX2 6GG, UK *Email address*: alex.cliffe@maths.ox.ac.uk

(Feimin Huang) ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA

Email address: fhuang@amt.ac.cn

(Song Liu) DEPARTMENT OF MATHEMATICS, CITY UNIVERSITY OF HONG KONG, HONG KONG, CHINA; ACAD-EMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA *Email address*: liusong@amss.ac.cn

(Qin Wang) DEPARTMENT OF MATHEMATICS, YUNNAN UNIVERSITY, KUNMING, 650091, CHINA *Email address*: mathwq@ynu.edu.cn