

# Guiding Center Derivation of the Generalized Hasegawa-Mima Equation for Drift Wave Turbulence in Curved Magnetic Fields

Naoki Sato<sup>1</sup> and Michio Yamada<sup>2</sup>

<sup>1</sup>*Graduate School of Frontier Sciences,  
The University of Tokyo, Kashiwa, Chiba 277-8561, Japan*

*Email: sato-naoki@edu.k.u-tokyo.ac.jp*

<sup>2</sup>*Research Institute for Mathematical Sciences,  
Kyoto University, Kyoto 606-8502, Japan*

*Email: yamada@kurims.kyoto-u.ac.jp*

May 29, 2023

## Abstract

Recently, a generalized Hasegawa-Mima (gHM) equation describing drift wave turbulence in curved magnetic fields has been derived in [N. Sato and M. Yamada, J. Plasma Phys. (2022), vol. 88, 905880319] for an ion-electron plasma modeled as a two-fluid system. In this work, we show that a mathematically equivalent GHM equation can be obtained within the kinetic framework of guiding center motion, and that the relevant drift wave turbulence ordering can be further relaxed, effectively generalizing the applicability of the equation to any magnetic field geometry and electron spatial density, in the sense that no ordering requirements involve spatial derivatives of the magnetic field or the electron spatial density.

## 1 Introduction

The Hasegawa-Mima (HM) equation [1, 2] is a nonlinear equation describing the turbulent behavior of electric potential and spatial density in a quasi-neutral plasma, made of hot thermalized electrons and cold ions, permeated by a strong, straight, and homogeneous magnetic field, and evolving over time scales long compared with the period of cyclotron motion. The nonlinearity of the HM equation is caused by the convection of the  $\mathbf{E} \times \mathbf{B}$  velocity associated with the polarization drift. In the presence of an electron density gradient, solutions of the linearized HM equation are the characteristic drift waves, whose interaction gives rise to drift wave turbulence. The HM equation shares the same mathematical structure with the quasi-geostrophic equation characterizing atmospheric motion over the surface of rotating planets [3, 4] due to the similarity between the Lorentz force and the Coriolis force, and contains 2-dimensional incompressible vorticity dynamics as a special case [5]. Despite its relative simplicity, the physical significance of the HM equation stems from its ability to capture essential features of 2-dimensional plasma and fluid turbulence [6], including onset of inverse turbulent cascades of energy [7, 8, 9, 10] and self-organization of large scale structures and zonal flows [11, 12, 13, 14, 15].

One of the key assumptions behind the HM equation is that both the background magnetic field and the electron spatial density change over a spatial scale  $L$  that is large compared to the typical turbulence wavelength across the magnetic field  $k_{\perp}^{-1}$ , i.e.  $k_{\perp}L \gg 1$ . This hypothesis effectively restricts the applicability of the HM equation to plasmas with a small density gradient and to magnetic fields with small curvature or field inhomogeneities. However, experimental observations pertaining to plasmas confined by dipole magnetic fields [16, 17] suggest the existence of drift wave turbulence and zonal flows in systems where both the electron spatial density and the magnetic field are characterized by strong gradients over spatial scales comparable to that of electric field and density fluctuations (these low frequency fluctuations are often referred to as

entropy modes [18]). In principle, an accurate description of electromagnetic turbulence in such setting could be obtained with the aid of nonlinear gyrokinetic theory [19, 20, 21]. Nevertheless, it is natural to ask whether the HM equation can be generalized to allow strong magnetic field and density inhomogeneities while maintaining a single governing equation for the electric potential  $\varphi$ . In [22] this question has been answered positively, and a generalized Hasegawa-Mima (gHM) equation has been obtained from a two-fluid model [23] of an ion-electron plasma in the form below (see section 2 for the definition of the physical quantities appearing in the equation):

$$\frac{\partial}{\partial t} \left[ \lambda A_e \varphi - \sigma \nabla \cdot \left( A_e \frac{\nabla_{\perp} \varphi}{B^2} \right) \right] = \nabla \cdot \left[ A_e \left( \sigma \frac{\mathbf{B} \cdot \nabla \times \mathbf{v}_E}{B^2} - 1 \right) \mathbf{v}_E \right]. \quad (1)$$

In particular, the ordering used in [22] to derive the gHM equation (1) only involves one ordering condition on the spatial derivatives of the magnetic field and the electron spatial density, effectively extending the range of the HM equation to general magnetic field geometries (see section 4 for details).

The aim of this paper is twofold. First, we want to show that a guiding center generalized Hasegawa-Mima (GHM) equation (34), mathematically equivalent to the gHM equation (1), can be derived within the framework of guiding center dynamics [24, 25] (here, the uppercase letter G is used to emphasize that the GHM equation can be derived under weaker assumptions, and thus it is more general than the gHM equation). Secondly, we wish to show that the guiding center drift wave turbulence ordering required to derive the GHM equation does not involve conditions on the spatial derivatives of the magnetic field or the electron spatial density, and it is therefore weaker than the two-fluid drift wave turbulence ordering used to derive the gHM equation from a two-fluid model in [22]. This difference does not originate from a discrepancy between two-fluid theory and guiding center theory. In fact, we will see that the new GHM equation can be obtained from a two-fluid ordering equivalent to the guiding center drift wave turbulence ordering by adding tailored higher-order terms to the gHM equation.

The present paper is organized as follows. In section 2, we derive the GHM equation by considering the evolution of the phase space distribution function of a magnetized plasma according to the guiding center equations of motion. In section 3, we discuss the constants of motion of the GHM equation, and obtain a more general form of the generalized enstrophy presented in [22]. In section 4 we examine the relationship between the ordering used to derive the gHM equation (1) from a two-fluid model in [22] and the ordering used here to obtain the GHM equation (34) from guiding center theory. In particular, we show that the guiding-center ordering does not involve any conditions on the spatial derivatives of the magnetic field or the electron spatial density. Concluding remarks are given in section 5.

## 2 Derivation of the GHM equation within the kinetic framework of guiding center motion

In this section, we derive the GHM equation for an ion-electron plasma obeying the guiding-center equations of motion under an appropriate drift wave turbulence ordering.

### 2.1 Guiding center ordering

We consider a guiding-center plasma made of ions and electrons within a region  $\Omega \subseteq \mathbb{R}^3$  permeated by a static magnetic field  $\mathbf{B}(\mathbf{x}) \neq \mathbf{0}$  with modulus  $B$ , and where  $\mathbf{x} = (x, y, z) \in \mathbb{R}^3$  are Cartesian coordinates. Let  $\mathbf{E} = -\nabla\varphi$  denote the electric field, with  $\varphi(\mathbf{x}, t)$  the electric potential ( $t$  is the time variable),  $\mathbf{E}_{\perp}$  the component of  $\mathbf{E}$  perpendicular to  $\mathbf{B}$ , and  $\mathbf{E}_{\parallel}$  the component of  $\mathbf{E}$  parallel to  $\mathbf{B}$ . The small ordering parameter for the guiding-center expansion will be denoted by  $\epsilon > 0$ , the spatial scale of the system by  $L$ , the time scale of the system by  $\tau$ , the ion gyroradius by  $\rho$ , and the ion cyclotron frequency by  $\omega_c = ZeB/m$  where  $Z \in \mathbb{N}$  and  $Ze$  and  $m$  are the ion electric charge and mass respectively. Let  $\mathbf{X} = \mathbf{x} - \rho$  denote the ion guiding center position,  $\mathbf{b} = \mathbf{B}/B$  the unit vector along  $\mathbf{B}$ ,  $\mu = m(\mathbf{v}_{\perp} - \mathbf{v}_E)^2/2B$  the lowest order magnetic moment,  $\mathbf{v} = \dot{\mathbf{x}}$  the charged particle velocity,  $\mathbf{v}_{\perp} = \mathbf{b} \times (\mathbf{v} \times \mathbf{b})$  the charged particle velocity perpendicular

to  $\mathbf{B}$ , and

$$\mathbf{v}_E = \frac{\mathbf{E} \times \mathbf{B}}{B^2}, \quad (2)$$

the  $\mathbf{E} \times \mathbf{B}$  velocity. Then, the ion guiding center equations of motion obtained from the Northrop phase space guiding center Lagrangian [24, 25] are

$$m\dot{\mathbf{b}} = Ze \left( \mathbf{E}' + \dot{\mathbf{X}} \times \mathbf{B}' \right), \quad (3a)$$

$$\dot{\mu} = 0, \quad (3b)$$

$$\dot{\zeta} = \omega_c, \quad (3c)$$

which can be equivalently written as

$$\dot{\mathbf{X}} = u \frac{\mathbf{B}'}{B'_\parallel} + \mathbf{E}' \times \frac{\mathbf{b}}{B'_\parallel}, \quad (4a)$$

$$\dot{u} = \frac{Ze}{m} \frac{\mathbf{B}' \cdot \mathbf{E}'}{B'_\parallel}, \quad (4b)$$

$$\dot{\mu} = 0, \quad (4c)$$

$$\dot{\zeta} = \omega_c, \quad (4d)$$

where  $\zeta$  is the gyrophase and

$$u = \dot{\mathbf{X}} \cdot \mathbf{b}, \quad (5a)$$

$$Ze\varphi' = Ze\varphi + \mu B + \frac{m}{2} \mathbf{v}_E^2, \quad (5b)$$

$$\mathbf{A}' = \mathbf{A} + \frac{m}{Ze} (u\mathbf{b} + \mathbf{v}_E) \quad (5c)$$

$$\mathbf{E}' = -\nabla\varphi' - \frac{\partial \mathbf{A}'}{\partial t}, \quad (5d)$$

$$\mathbf{B} = \nabla \times \mathbf{A}, \quad \mathbf{B}' = \nabla \times \mathbf{A}', \quad B'_\parallel = \mathbf{B}' \cdot \mathbf{b}. \quad (5e)$$

Here, we observe that  $u$  represents the component of the guiding center velocity parallel to  $\mathbf{B}$ . Furthermore, notice that the following guiding center drift velocities  $\mathbf{v}_E$  ( $\mathbf{E} \times \mathbf{B}$  drift),  $\mathbf{v}_\nabla$  ( $\nabla B$  drift), and  $\mathbf{v}_\kappa$  (curvature drift) are contained in the right-hand side of (4a) according to

$$\mathbf{v}'_E = \frac{\mathbf{b} \times \nabla\varphi}{B'_\parallel}, \quad \mathbf{v}'_\nabla = \frac{\mu}{Ze} \frac{\mathbf{b} \times \nabla B}{B'_\parallel}, \quad \mathbf{v}'_\kappa = \frac{mu^2}{ZeB'_\parallel} \nabla \times \mathbf{b}, \quad (6)$$

where the  $'$  symbol is used to emphasize that the correction  $B'_\parallel$  of the magnetic field  $B$  caused by the term  $m\nabla \times (u\mathbf{b} + \mathbf{v}_E)/Ze$  in  $\mathbf{B}'$  is used in these formulas. Similarly, the polarization drift  $\mathbf{v}_{\text{pol}}$  is included in (4a) according to

$$\mathbf{v}'_{\text{pol}} = \frac{m}{Ze} \frac{\mathbf{b} \times \frac{\partial \mathbf{v}_E}{\partial t}}{B'_\parallel} + \mathbf{v}'_E - \mathbf{v}_E + \frac{m}{2Ze} \frac{\mathbf{b} \times \nabla \mathbf{v}_E^2}{B'_\parallel}. \quad (7)$$

The physical meaning carried by this expression will become clear later. Equation (4a) also includes a further drift term

$$\mathbf{v}'_* = \frac{mu}{ZeB'_\parallel} \nabla \times \mathbf{v}_E, \quad (8)$$

which originates from the effective magnetic field  $\frac{m}{Ze} \nabla \times \mathbf{v}_E$  associated with  $\mathbf{E} \times \mathbf{B}$  motion. The total guiding center velocity can thus be written as

$$\dot{\mathbf{X}} = u \frac{\mathbf{B}}{B_\parallel} + \mathbf{v}_E + \mathbf{v}'_\nabla + \mathbf{v}'_\kappa + \mathbf{v}'_{\text{pol}} + \mathbf{v}'_*. \quad (9)$$

Order	Dimensionless	Fields	Distances	Rates	Velocities
$\epsilon^{-1}$		$\mathbf{B}, \mathbf{E}_\perp$		$\omega_c$	
1		$\mathbf{E}_\parallel$	$L$	$\mathbf{v}/L, \mathbf{v}_E/L, \tau^{-1}$	$\mathbf{v}, \mathbf{v}_E$
$\epsilon$	$\rho/L, (\omega_c\tau)^{-1}$		$\rho$	$\mathbf{v}_\nabla/L, \mathbf{v}_\kappa/L, \mathbf{v}_{\text{pol}}/L$	$\mathbf{v}_\nabla, \mathbf{v}_\kappa, \mathbf{v}_{\text{pol}}$

Table 1: Guiding center ordering required for the existence of the first adiabatic invariant  $\mu$  (see [24]).

For completeness we recall that the gyroradius, which defines the coordinate transformation  $\mathbf{X} = \mathbf{x} - \boldsymbol{\rho}$ , is an oscillatory (gyrophase dependent) term given by

$$\boldsymbol{\rho} = \frac{m}{Ze} \frac{\mathbf{b} \times (\mathbf{v} - \mathbf{v}_E)}{B^2}. \quad (10)$$

This term is removed from the Northrop guiding center phase space Lagrangian density  $\mathcal{L}_{Ngc}$  by appropriate subtraction of total time derivatives (on this point, see section III.C and appendix A of [24]). Here, the Northrop guiding center phase space Lagrangian density  $\mathcal{L}_{Ngc}$  is obtained by expansion of the charged particle phase space Lagrangian density  $\mathcal{L} = \mathcal{L}_{Ngc} + o(\epsilon)$  according to the guiding center ordering. Since the Northrop guiding center phase space Lagrangian density  $\mathcal{L}_{Ngc}$  is independent of  $\boldsymbol{\rho}$ , it is also independent of the gyrophase  $\zeta$ , leading to conservation of the conjugate momentum  $\mu$  by the Noether theorem. The guiding-center ordering required for the conservation of the magnetic moment  $\mu$  is given in table I of [24], which we report in table 1. This ordering represents the starting point that we will use to construct a more restrictive ordering leading to the GHM equation. We conclude by observing that the ordering parameter  $\epsilon$  arises from the physical constant

$$\sigma = \frac{m}{Ze} \ll 1, \quad (11)$$

which is small for elementary particles such as ions and electrons. The constant  $\sigma^{-1}$  always multiplies the electromagnetic fields within the charged particle phase space Lagrangian density  $\mathcal{L}(\mathbf{x}, \mathbf{v}, t) = (\sigma^{-1}\mathbf{A} + \mathbf{v}) \cdot \dot{\mathbf{x}} - \frac{1}{2}\mathbf{v}^2 - \sigma^{-1}\varphi$ , which is the reason why  $\mathbf{E}$  and  $\mathbf{B}$  are treated as large fields in table 1.

## 2.2 Derivation of the GHM equation from a drift wave turbulence ordering within guiding center theory

We start by assuming that the guiding center ordering presented in table 1 holds, and gradually impose additional conditions to obtain the relevant drift wave turbulence ordering. From now on we set  $\tau = \tau_d$ , with  $\tau_d$  the drift turbulence time scale. Recall that  $\tau_d\omega_c \sim \epsilon^{-1}$ . Let  $f(\mathbf{p}, \mathbf{x}, t)$  denote the ion distribution function in the canonical phase space  $(\mathbf{p}, \mathbf{x})$  of charged particle dynamics. The distribution function  $f$  satisfies the Boltzmann equation

$$\frac{\partial f}{\partial t} = -\frac{\partial}{\partial \mathbf{p}} \cdot (\dot{\mathbf{p}}f) - \frac{\partial}{\partial \mathbf{x}} \cdot (\dot{\mathbf{x}}f) + \left(\frac{df}{dt}\right)_c, \quad (12)$$

where the last term on the right-hand side describes particle collisions. Introducing the ion spatial density  $n(\mathbf{x}, t) = \int_{\mathbb{R}^3} f d\mathbf{p}$ , integrating equation (12) with respect to the momentum variables, and assuming  $\lim_{|\mathbf{p}| \rightarrow \infty} f = 0$ , we obtain the ion continuity equation

$$\frac{\partial n}{\partial t} = -\nabla \cdot (\langle \dot{\mathbf{x}} \rangle n), \quad (13)$$

where

$$\langle \dot{\mathbf{x}} \rangle = \frac{1}{n} \int_{\mathbb{R}^3} f \dot{\mathbf{x}} d\mathbf{p} = \frac{1}{n} \int_{\mathbb{R}^3} f (\dot{\mathbf{X}} + \dot{\boldsymbol{\rho}}) d\mathbf{p} = \langle \dot{\mathbf{X}} \rangle, \quad (14)$$

is the ensemble averaged ion velocity at a given position  $\mathbf{x} = \mathbf{X} + \boldsymbol{\rho}$ . Notice that in the last passage we used the fact that by hypothesis  $\boldsymbol{\rho}$  is an oscillatory term such that the ensemble average  $\langle \dot{\boldsymbol{\rho}} \rangle$  identically vanishes.

We also remark that the collision term in (12) vanishes upon integration in momentum space because we assume that collisions result in deflections in velocity space that do not change the local particle number. Now suppose that the parallel velocity  $u$  is small (the time scale  $\tau_b$  of dynamics along  $\mathbf{B}$  is long):

$$u \frac{\tau_d}{L} \sim \frac{\tau_d}{\tau_b} \sim \epsilon^2. \quad (15)$$

Note that consistency with (4b) requires that the component  $E'_{\parallel}$  of  $\mathbf{E}'$  along  $\mathbf{B}'$  is small, i.e.  $\mathbf{B}' \cdot \mathbf{E}' / B'_{\parallel} E_{\perp} \sim \epsilon^3$  (physically, this means that the electric field experienced by a charged particle along the magnetic field is negligible). Let us now consider all contributions to the continuity equation (13) that are greater than  $\epsilon^2$ . From the ordering condition (15) it readily follows that the only surviving terms in (4a) are those involving the  $\mathbf{E} \times \mathbf{B}$  drift velocity and the  $\nabla B$  drift. In particular, observing that  $B'_{\parallel} = B(1 + o(\epsilon))$ , we have

$$\dot{\mathbf{X}} = \mathbf{v}_{\mathbf{E}} + \mathbf{v}'_{\nabla} + \mathbf{v}_{\text{pol}} + o(\epsilon^2), \quad (16)$$

where the polarization drift  $\mathbf{v}_{\text{pol}}$  now has expression

$$\mathbf{v}_{\text{pol}} = \sigma \frac{\mathbf{b} \times \frac{d\mathbf{v}_{\mathbf{E}}}{dt}}{B}, \quad \frac{d\mathbf{v}_{\mathbf{E}}}{dt} = \frac{\partial \mathbf{v}_{\mathbf{E}}}{\partial t} + \mathbf{v}_{\mathbf{E}} \cdot \nabla \mathbf{v}_{\mathbf{E}}, \quad (17)$$

and where we used the fact that

$$\mathbf{v}_{\mathbf{E}} + \mathbf{v}'_{\text{pol}} = \mathbf{v}_{\mathbf{E}} - \sigma \frac{\mathbf{b} \cdot \nabla \times \mathbf{v}_{\mathbf{E}}}{B} \mathbf{v}_{\mathbf{E}} + \sigma \frac{\mathbf{b} \times \left( \frac{\partial \mathbf{v}_{\mathbf{E}}}{\partial t} + \frac{1}{2} \nabla \mathbf{v}_{\mathbf{E}}^2 \right)}{B} + o(\epsilon^2) = \mathbf{v}_{\mathbf{E}} + \mathbf{v}_{\text{pol}} + o(\epsilon^2). \quad (18)$$

We see that the polarization drift  $\mathbf{v}_{\text{pol}}$  is that average particle velocity resulting from a non-vanishing average acceleration  $d\mathbf{v}_{\mathbf{E}}/dt$  across the magnetic field.

In the following, we shall also demand the energy  $\mu B$  of cyclotron dynamics to be small, so that  $\mathbf{v}'_{\nabla}$  becomes a higher order correction. More precisely, denoting with  $\langle \mu B \rangle = n^{-1} \int_{\mathbb{R}^3} f \mu B d\mathbf{p}$  the ensemble averaged kinetic energy of cyclotron dynamics, and defining an associated temperature  $T_c$  according to  $\langle \mu B \rangle = k_B T_c$  with  $k_B$  the Boltzmann constant, we demand  $T_c$  to satisfy

$$\frac{k_B T_c}{\frac{m}{2} \mathbf{v}_{\mathbf{E}}^2} \sim \epsilon. \quad (19)$$

The ordering conditions (15) and (19) can be regarded as the usual drift wave turbulence ordering requirement of cold ions. The guiding-center velocity thus becomes

$$\dot{\mathbf{X}} = \mathbf{v}_{\mathbf{E}} + \mathbf{v}_{\text{pol}} + o(\epsilon^2) = \frac{\mathbf{b} \times \nabla \varphi}{B} + \sigma \frac{\mathbf{b} \times \frac{d\mathbf{v}_{\mathbf{E}}}{dt}}{B} + o(\epsilon^2). \quad (20)$$

To proceed further, it is convenient to introduce the orthogonal gradient operator

$$\nabla_{\perp} = -\mathbf{b} \times (\mathbf{b} \times \nabla). \quad (21)$$

Although the expression (20) is convenient to highlight the usual guiding center drift contributions separately,  $o(\epsilon^2)$  order terms must be added to  $\mathbf{v}_{\mathbf{E}} + \mathbf{v}_{\text{pol}}$  in order for the reduced (drift wave) Hamiltonian  $\chi = \varphi + \frac{\sigma}{2} \mathbf{v}_{\mathbf{E}}^2$  arising from the expansion of the Northrop guiding center Hamiltonian  $H_{Ngc} = \frac{\sigma}{2} u^2 + \varphi + \mu B + \frac{\sigma}{2} \mathbf{v}_{\mathbf{E}}^2 = \chi + o(\epsilon)$  to be an exact constant of motion in the case of time-independent electromagnetic fields. To this end, one can verify that equation (20) can be equivalently written as

$$\dot{\mathbf{X}} = \dot{\mathbf{X}}_{dw} + o(\epsilon^2), \quad \dot{\mathbf{X}}_{dw} = \frac{\mathbf{b} \times \nabla \left( \varphi + \frac{\sigma}{2} \mathbf{v}_{\mathbf{E}}^2 \right)}{B''_{\parallel}} - \sigma \frac{\nabla_{\perp} \varphi_t}{B^2}, \quad B''_{\parallel} = B \left( 1 + \sigma \frac{\mathbf{b} \cdot \nabla \times \mathbf{v}_{\mathbf{E}}}{B} \right), \quad (22)$$

with  $\chi$  an exact integral of the first order term  $\dot{\mathbf{X}}_{dw}$  when  $\varphi_t = \partial \varphi / \partial t = 0$ .

Order	Dimensionless	Fields	Distances	Rates	Velocities
$\epsilon^{-1}$		$\mathbf{B}, \mathbf{E}_\perp$		$\omega_c$	
1		$A_e$	$L$	$\tau_d^{-1}, \mathbf{v}_E/L$	$\mathbf{v}_E$
$\epsilon$	$\lambda\varphi, \rho/L, (\omega_c\tau_d)^{-1}, k_B T_c / \frac{m}{2} \mathbf{v}_E^2$		$\rho$	$\mathbf{v}_{\text{pol}}/L$	$\mathbf{v}_{\text{pol}}$
$\epsilon^2$	$\frac{m}{2} \mathbf{v}_E^2 / k_B T_e, \tau_d / \tau_b$	$E'_\parallel$		$\mathbf{v}_\nabla / L, u/L, \tau_b^{-1}$	$\mathbf{v}_\nabla, u$
$\epsilon^5$				$\mathbf{v}_\kappa / L$	$\mathbf{v}_\kappa$

Table 2: Drift wave turbulence ordering used for the derivation of the GHM equation within the guiding-center framework.

Noting that  $\dot{\mathbf{X}}_{dw}$  is a pure spatial function, the ensemble averaged ion velocity (14) at a given position  $\mathbf{x} = \mathbf{X} + \boldsymbol{\rho}$  is

$$\langle \dot{\mathbf{x}} \rangle = \frac{1}{n} \int_{\mathbb{R}^3} f(\dot{\mathbf{X}} + \dot{\boldsymbol{\rho}}) d\mathbf{p} = \dot{\mathbf{X}}_{dw} + o(\epsilon^2). \quad (23)$$

Next, consider the density  $n_e(\mathbf{x}, t)$  of the electron component. We assume that  $n_e$  follows a Boltzmann distribution with temperature  $T_e$ , i.e.

$$n_e = A_e(\mathbf{x}) \exp\{\lambda\varphi(\mathbf{x}, t)\}, \quad \lambda = \frac{e}{k_B T_e} \quad (24)$$

where  $A_e(\mathbf{x})$  is a spatial function. If we further demand the ion-electron plasma to be quasi-neutral, we have the following condition:

$$Zn(\mathbf{x}, t) = n_e(\mathbf{x}, t). \quad (25)$$

Then, the continuity equation for the ion density reads as

$$Z \left[ \frac{\partial n}{\partial t} + \nabla \cdot (\langle \dot{\mathbf{x}} \rangle n) \right] = \frac{\partial n_e}{\partial t} + \nabla \cdot (\dot{\mathbf{X}}_{dw} n_e) + o(\epsilon^2) = 0, \quad (26)$$

where we used equation (23). Substituting equation (24), equation (26) can be rearranged as

$$\begin{aligned} \lambda A_e \frac{\partial \varphi}{\partial t} &= -\lambda \nabla \varphi \cdot A_e \dot{\mathbf{X}}_{dw} - \nabla \cdot (A_e \dot{\mathbf{X}}_{dw}) + o(\epsilon^2) \\ &= \lambda \sigma A_e \nabla \varphi \cdot \left( \frac{\nabla_\perp \varphi_t}{B^2} + \frac{\mathbf{b} \times \nabla \mathbf{v}_E^2}{2B'_\parallel} \right) - \nabla \cdot \left[ A_e \frac{\mathbf{b} \times \nabla (\varphi + \frac{\sigma}{2} \mathbf{v}_E^2)}{B'_\parallel} \right] + \sigma \nabla \cdot \left( A_e \frac{\nabla_\perp \varphi_t}{B^2} \right) + o(\epsilon^2). \end{aligned} \quad (27)$$

Next, we demand the electron component to be hot compared to the ion component, i.e.

$$\frac{m}{2} \frac{\mathbf{v}_E^2}{k_B T_e} \sim \epsilon^2. \quad (28)$$

Since  $\lambda = e/k_B T_e$ , it follows that  $\lambda\varphi \sim \epsilon$ , while the first term on the right-hand side of equation (27) scales as  $\epsilon^2$ . Equation (27) thus reduces to

$$\frac{\partial}{\partial t} \left[ \lambda A_e \varphi - \sigma \nabla \cdot \left( A_e \frac{\nabla_\perp \varphi}{B^2} \right) \right] = -\nabla \cdot \left[ A_e \frac{\mathbf{b} \times \nabla (\varphi + \frac{\sigma}{2} \mathbf{v}_E^2)}{B'_\parallel} \right] + o(\epsilon^2). \quad (29)$$

Since this equation is correct up to first order in  $\epsilon$ , we can add terms scaling as  $\epsilon^2$  that will be useful later to obtain exact conservation laws:

$$\frac{\partial}{\partial t} \left\{ \lambda A_e \left( \varphi + \frac{\sigma}{2} \mathbf{v}_E^2 \right) - \sigma \nabla \cdot \left[ A_e \frac{\nabla_\perp (\varphi + \frac{\sigma}{2} \mathbf{v}_E^2)}{B^2} \right] \right\} = -\nabla \cdot \left[ A_e \frac{\mathbf{b} \times \nabla (\varphi + \frac{\sigma}{2} \mathbf{v}_E^2)}{B'_\parallel} \right] + o(\epsilon^2). \quad (30)$$

Recalling that the Northrop guiding center Hamiltonian reduces to

$$\chi = \varphi + \frac{\sigma}{2} \mathbf{v}_{\mathbf{E}}^2, \quad (31)$$

in the drift wave turbulence ordering, and that  $B_{\parallel}'' = B(1 + \sigma \frac{\mathbf{b} \cdot \nabla \times \mathbf{v}_{\mathbf{E}}}{B})$  we thus arrive at the equation

$$\begin{aligned} \frac{\partial}{\partial t} \left[ \lambda A_e \chi - \sigma \nabla \cdot \left( A_e \frac{\nabla_{\perp} \chi}{B^2} \right) \right] &= -\nabla \cdot \left( A_e \frac{\mathbf{b} \times \nabla \chi}{B_{\parallel}''} \right) + o(\epsilon^2) \\ &= \nabla \cdot \left\{ A_e \left[ \sigma \frac{\mathbf{b} \cdot \nabla \times \left( \frac{\mathbf{b} \times \nabla \chi}{B} \right)}{B} - 1 \right] \frac{\mathbf{b} \times \nabla \chi}{B} \right\} + o(\epsilon^2). \end{aligned} \quad (32)$$

Notice that  $\chi = \varphi + o(1)$ . If we further define the vector field

$$\mathbf{v}_{\mathbf{E}}^{\chi} = \frac{\mathbf{b} \times \nabla \chi}{B} = \mathbf{v}_{\mathbf{E}} + \sigma \frac{\mathbf{b} \times \nabla \mathbf{v}_{\mathbf{E}}^2}{2B}, \quad (33)$$

equation (32) gives the following closed equation for the variable  $\chi$ ,

$$\frac{\partial}{\partial t} \left[ \lambda A_e \chi - \sigma \nabla \cdot \left( A_e \frac{\nabla_{\perp} \chi}{B^2} \right) \right] = \nabla \cdot \left[ A_e \left( \sigma \frac{\mathbf{B} \cdot \nabla \times \mathbf{v}_{\mathbf{E}}^{\chi}}{B^2} - 1 \right) \mathbf{v}_{\mathbf{E}}^{\chi} \right]. \quad (34)$$

In the following, we shall refer to equation (34) as the guiding center generalized Hasegawa-Mima equation (GHM) to distinguish it from the gHM equation (1) derived in [22] from a two-fluid plasma model. Nonetheless, notice that these two equations share the same mathematical form, with the variable  $\chi$  in (34) replacing  $\varphi$  in (1). The relationship between the ordering conditions used to construct these two models will be discussed in section 4. Here, we observe that in (34) the term  $\nabla \cdot (A_e \mathbf{v}_{\mathbf{E}}^{\chi}) = \nabla \chi \cdot \nabla \times (A_e \mathbf{B}/B^2)$  must scale as  $\sim \epsilon$  to be consistent with the other terms in the equation. Hence, either  $\nabla \times (A_e \mathbf{B}/B^2)$  scales as  $\sim \epsilon^2$ , or the effective electric field  $-\nabla \chi \sim \epsilon^{-1}$  is mostly orthogonal to the vector field  $\nabla \times (A_e \mathbf{B}/B^2) \sim \epsilon$ . We stress however that the behavior of the term  $\nabla \cdot (A_e \mathbf{v}_{\mathbf{E}}^{\chi})$  is a consequence of the ordering used to obtain equation (34), and not an ordering condition required to arrive at (34).

The ordering used to derive the GHM equation (34) is summarized in table 2. Finally, one can verify that equation (34) reduces to the standard HM equation

$$\frac{\partial}{\partial t} \left( \lambda \varphi - \frac{\sigma}{B_0^2} \Delta_{(x,y)} \varphi \right) = \frac{\sigma}{B_0^3} [\varphi, \Delta_{(x,y)} \varphi]_{(x,y)} + \frac{\beta}{B_0} \varphi_y, \quad (35)$$

when  $\mathbf{B} = B_0 \nabla z$ ,  $\log A_e = \log A_{e0} + \beta x$ ,  $B_0, A_{e0}, \beta \in \mathbb{R}$ ,  $\beta L \sim \epsilon$ . Here,  $[f, g]_{(x,y)} = f_x g_y - f_y g_x$ ,  $\Delta_{(x,y)} = \partial_x^2 + \partial_y^2$ , and lower indexes denote partial derivatives, for example  $f_x = \partial f / \partial x$ .

### 3 Conservation laws

In this section, we show that the derived GHM equation (34) preserves both mass and energy. Furthermore, we identify a third invariant associated with the vorticity of the flow in a more general form than the one obtained in [22], and discuss its relationship with the generalized enstrophy encountered in the standard HM equation.

Since the GHM equation (34) and the gHM equation (1) share the same mathematical structure, we already know that the invariants of the GHM equation (34) can be obtained by replacing  $\varphi$  with  $\chi$  in the expressions of the invariants of the gHM equation (1). It is however useful to recall the physical origin of these quantities. First observe that the the total ion mass can be written as

$$\mathcal{M}_{\Omega} = \frac{m}{Z} \int_{\Omega} A_e e^{\lambda \varphi} d\mathbf{x}. \quad (36)$$

Since  $\lambda\varphi \sim \epsilon$ , we may expand the exponential in powers of  $\lambda\varphi$  according to  $e^{\lambda\varphi} = 1 + \lambda\chi + o(\epsilon^2)$  and consider the conservation of the first order term,

$$M_\Omega = \frac{m}{Z} \int_\Omega A_e (1 + \lambda\chi) d\mathbf{x}. \quad (37)$$

Using (34) we have

$$\frac{dM_\Omega}{dt} = \frac{m}{Z} \int_{\partial\Omega} A_e \left[ \sigma \frac{\nabla_\perp \chi_t}{B^2} - \left( 1 - \sigma \frac{\mathbf{B} \cdot \nabla \times \mathbf{v}_E^\chi}{B^2} \right) \mathbf{v}_E^\chi \right] \cdot \mathbf{n} dS = -\frac{m}{Z} \int_{\partial\Omega} A_e \dot{\mathbf{X}}'_{dw} \cdot \mathbf{n} dS, \quad (38)$$

where  $\mathbf{n}$  denotes the unit outward normal to the bounding surface  $\partial\Omega$ ,  $dS$  the surface element on  $\partial\Omega$ , and we defined

$$\dot{\mathbf{X}}'_{dw} = \left( 1 - \sigma \frac{\mathbf{B} \cdot \nabla \times \mathbf{v}_E^\chi}{B^2} \right) \mathbf{v}_E^\chi - \sigma \frac{\nabla_\perp \chi_t}{B^2} = \dot{\mathbf{X}}_{dw} + o(\epsilon^2). \quad (39)$$

The boundary integral (38) vanishes under suitable boundary conditions, such as  $A_e = 0$  on  $\partial\Omega$  or  $\dot{\mathbf{X}}'_{dw} \cdot \mathbf{n} = 0$  on  $\partial\Omega$ .

Next, observe that the leading order ion Hamiltonian is given by

$$H = \frac{m}{2} \mathbf{v}^2 + Ze\varphi = Ze\varphi + \frac{m}{2} \mathbf{v}_E^2 + o(\epsilon). \quad (40)$$

Therefore, at leading order the total ion energy satisfies

$$\mathcal{H}_\Omega = Ze \int_{\Omega \times \mathbb{R}^3} f \left( \varphi + \frac{\sigma}{2} \mathbf{v}_E^2 \right) d\mathbf{p} d\mathbf{x} + o(\epsilon) = e \int_\Omega A_e e^{\lambda\varphi} \left( \varphi + \frac{\sigma}{2} \mathbf{v}_E^2 \right) d\mathbf{x} + o(\epsilon). \quad (41)$$

Dividing this expression by  $k_B T_e$  we obtain

$$\frac{\mathcal{H}_\Omega}{k_B T_e} = \int_\Omega A_e \left( 1 + \lambda\varphi + \frac{1}{2} \lambda^2 \varphi^2 \right) \left( \lambda\varphi + \frac{\sigma}{2} \lambda \mathbf{v}_E^2 \right) d\mathbf{x} + o(\epsilon^4) = \int_\Omega A_e \left( \lambda\varphi + \lambda^2 \varphi^2 + \frac{\sigma}{2} \lambda \mathbf{v}_E^2 \right) d\mathbf{x} + o(\epsilon^4). \quad (42)$$

It follows that

$$\frac{\mathcal{H}_\Omega}{k_B T_e} - \frac{Z}{m} \mathcal{M}_\Omega = \frac{1}{2} \int_\Omega A_e \left( \lambda^2 \varphi^2 + \lambda \sigma \mathbf{v}_E^2 - 2 \right) d\mathbf{x} + o(\epsilon^3) = \frac{1}{2} \int_\Omega A_e \left( \lambda^2 \chi^2 + \lambda \sigma \mathbf{v}_E^2 - 2 \right) d\mathbf{x} + o(\epsilon^3). \quad (43)$$

Since  $A_e$  is a spatial function, we thus expect the GHM energy

$$H_\Omega = \frac{1}{2} \int_\Omega A_e \left( \lambda \chi^2 + \sigma \frac{|\nabla_\perp \chi|^2}{B^2} \right) d\mathbf{x}, \quad (44)$$

to be a constant of motion. From equation (34), one can verify that

$$\begin{aligned} \frac{dH_\Omega}{dt} &= \int_\Omega \chi \frac{\partial}{\partial t} \left[ \lambda A_e \chi - \sigma \nabla \cdot \left( A_e \frac{\nabla_\perp \chi}{B^2} \right) \right] d\mathbf{x} + \sigma \int_{\partial\Omega} A_e \chi \frac{\nabla_\perp \chi_t}{B^2} \cdot \mathbf{n} dS \\ &= \int_{\partial\Omega} A_e \chi \left[ \sigma \frac{\nabla_\perp \chi_t}{B^2} - \left( 1 - \sigma \frac{\mathbf{B} \cdot \nabla \times \mathbf{v}_E^\chi}{B^2} \right) \mathbf{v}_E^\chi \right] \cdot \mathbf{n} dS \\ &= - \int_{\partial\Omega} A_e \chi \dot{\mathbf{X}}'_{dw} \cdot \mathbf{n} dS. \end{aligned} \quad (45)$$

Again, this boundary integral vanishes under suitable boundary conditions, such as  $A_e = 0$  on  $\partial\Omega$ ,  $\chi = 0$  on  $\partial\Omega$ , or  $\dot{\mathbf{X}}'_{dw} \cdot \mathbf{n} = 0$  on  $\partial\Omega$ .

Invariant	Expression	Field Conditions	Boundary Conditions
Mass $M_\Omega$	$\int_\Omega A_e (1 + \lambda\chi) d\mathbf{x}$	none	$A_e \dot{\mathbf{X}}'_{dw} \cdot \mathbf{n} = 0$
Energy $H_\Omega$	$\frac{1}{2} \int_\Omega A_e \left( \lambda\chi^2 + \sigma \frac{ \nabla_\perp \chi ^2}{B^2} \right) d\mathbf{x}$	none	$A_e \chi \dot{\mathbf{X}}'_{dw} \cdot \mathbf{n} = 0$
Enstrophy $W_\Omega$	$\int_\Omega A_e w \left( \lambda\chi - \frac{\sigma}{A_e} \nabla \cdot \left( A_e \frac{\nabla_\perp \chi}{B^2} \right) \right) d\mathbf{x}$	$\nabla \times \left( A_e \frac{\mathbf{B}}{B^2} \right) = \mathbf{0}$	$w A_e \mathbf{v}_E^\chi \cdot \mathbf{n} = 0$

Table 3: Invariants of the GHM equation (34).

An additional invariant, associated with the vorticity  $\nabla \times \mathbf{v}_E^\chi$ , exists when the magnetic field and the electron spatial density satisfy the condition  $\nabla \times (A_e \mathbf{B}/B^2) = \mathbf{0}$ . To see this, define the quantity

$$\omega = \nabla \cdot \left( A_e \frac{\nabla_\perp \chi}{B^2} \right) = A_e \frac{\mathbf{B} \cdot \nabla \times \mathbf{v}_E^\chi}{B^2} + \nabla \chi \cdot \frac{\mathbf{B}}{B^2} \times \left[ \nabla \times \left( A_e \frac{\mathbf{B}}{B^2} \right) \right]. \quad (46)$$

Next, observe that whenever

$$\frac{\mathbf{B}}{B^2} \times \left[ \nabla \times \left( A_e \frac{\mathbf{B}}{B^2} \right) \right] = \mathbf{0}, \quad (47)$$

which implies that  $A_e \mathbf{B}/B^2$  is a Beltrami field, the following identity holds

$$\omega = \nabla \cdot \left( A_e \frac{\nabla_\perp \chi}{B^2} \right) = A_e \frac{\mathbf{B} \cdot \nabla \times \mathbf{v}_E^\chi}{B^2}. \quad (48)$$

The derived GHM equation (34) can thus be written in the form

$$\frac{\partial}{\partial t} [\lambda A_e \chi - \sigma \omega] = -\nabla \cdot \left[ \left( 1 - \sigma \frac{\omega}{A_e} \right) A_e \mathbf{v}_E^\chi \right]. \quad (49)$$

On the other hand, functionals of the form

$$W_\Omega = \int_\Omega A_e w \left( \lambda\chi - \sigma \frac{\omega}{A_e} \right) d\mathbf{x}, \quad (50)$$

where  $w(\lambda\chi - \sigma\omega/A_e)$  is any function of  $\lambda\chi - \sigma\omega/A_e$ , satisfy

$$\frac{dW_\Omega}{dt} = \int_\Omega w' \frac{\partial}{\partial t} (\lambda A_e \chi - \sigma \omega) d\mathbf{x} = - \int_\Omega w' \left[ \nabla \cdot \left( \lambda\chi - \sigma \frac{\omega}{A_e} \right) \cdot A_e \mathbf{v}_E^\chi + \left( 1 - \sigma \frac{\omega}{A_e} \right) \nabla \cdot (A_e \mathbf{v}_E^\chi) \right] d\mathbf{x}, \quad (51)$$

with  $w' = dw/d(\lambda\chi - \sigma\omega/A_e)$ . Now observe that

$$\nabla \cdot (A_e \mathbf{v}_E^\chi) = \nabla \chi \cdot \nabla \times \left( A_e \frac{\mathbf{B}}{B^2} \right). \quad (52)$$

Hence, if we further demand that

$$\nabla \times \left( A_e \frac{\mathbf{B}}{B^2} \right) = \mathbf{0}, \quad (53)$$

which is a special case of (47), we find that

$$\frac{dW_\Omega}{dt} = - \int_{\partial\Omega} w A_e \mathbf{v}_E^\chi \cdot \mathbf{n} dS. \quad (54)$$

This boundary integral vanishes whenever  $w A_e \mathbf{v}_E^\chi \cdot \mathbf{n} = 0$  on  $\partial\Omega$ . The quantity  $W_\Omega$  can be identified with the generalized enstrophy encountered in the standard Hasegawa-Mima equation if the boundary condition

Order	Dimensionless	Fields	Distances	Rates	Velocities
1		$\mathbf{B}, A_e$	$L$	$\omega_c$	
$\epsilon$	$\lambda\varphi, \omega_c^{-1}\partial_t$	$\mathbf{E}_\perp$		$\tau_d^{-1}, \mathbf{v}_E/L$	$\mathbf{v}_E$
$\epsilon^2$	$\tau_d/\tau_b$			$\mathbf{v}_{\text{pol}}/L$	$\mathbf{v}_{\text{pol}}$
$\epsilon^3$	$\sigma  \nabla \mathbf{v}_E^2 \cdot \nabla \times (A_e \frac{\mathbf{B}}{B^2})  / A_e \omega_c$	$E_\parallel, P$		$\tau_b^{-1}, v_\parallel/L$	$v_\parallel$

Table 4: Drift wave turbulence ordering used for the derivation of the gHM equation from a two-fluid model in [22]. Here,  $P$  denotes the ion fluid pressure and  $v_\parallel$  the ion fluid velocity along  $\mathbf{B}$ .

Order	Dimensionless	Fields	Distances	Rates	Velocities
1		$\mathbf{B}, A_e$	$L$	$\omega_c$	
$\epsilon$	$\lambda\varphi, \omega_c^{-1}\partial_t, L\nabla \log B, L\nabla \log A_e$	$\mathbf{E}_\perp$		$\tau_d^{-1}, \mathbf{v}_E/L$	$\mathbf{v}_E$
$\epsilon^2$	$\tau_d/\tau_b$			$\mathbf{v}_{\text{pol}}/L$	$\mathbf{v}_{\text{pol}}$
$\epsilon^3$		$E_\parallel, P$		$\tau_b^{-1}, v_\parallel/L$	$v_\parallel$

Table 5: Drift wave turbulence ordering used for the derivation of the standard HM equation in a straight homogeneous magnetic field from a two-fluid model. Here,  $P$  denotes the ion fluid pressure and  $v_\parallel$  the ion fluid velocity along  $\mathbf{B}$ .

above is satisfied through  $A_e$ , i.e.  $A_e = 0$  on  $\partial\Omega$ . Indeed, choosing  $w = (\lambda\chi - \sigma\omega/A_e)^2$  and integrating by parts gives

$$W_\Omega = 2\lambda H_\Omega + \sigma \int_\Omega \left\{ \lambda A_e \frac{|\nabla_\perp \chi|^2}{B^2} + \sigma A_e^{-1} \left[ \nabla \cdot \left( A_e \frac{\nabla_\perp \chi}{B^2} \right) \right]^2 \right\} d\mathbf{x}. \quad (55)$$

In the following, we shall refer to  $W_\Omega$  as the generalized enstrophy. It is worth observing that the condition (53) implies (Poincaré lemma) that the magnetic field locally defines the normal direction of a surface  $C = \text{constant}$ , i.e.  $\mathbf{B} \propto \nabla C$  for some appropriate function  $C$  and sufficiently small neighborhood  $U \subseteq \Omega$ .

The invariants of the GHM equation are summarized in table 3.

## 4 Relationship with the two-fluid model gHM

In this section, we discuss the relationship between the ordering conditions used to derive the gHM equation (1) from a two-fluid model in [22], and those used above to derive the GHM equation (34) within the kinetic framework of guiding-center dynamics. Recall that the two equations possess the same mathematical form, but the respective orderings are slightly different.

Table 4 summarizes the drift wave turbulence ordering used to derive the gHM equation from a two-fluid model in [22], while the standard drift wave turbulence ordering for the HM equation (35) is given in table 5. Here, we observe that the standard HM ordering of table 5 is stricter than the two-fluid gHM ordering of table 4. In particular, the HM conditions on the spatial changes in  $B$  and  $A_e$ ,  $L\nabla \log B \sim L\nabla \log A_e \sim \epsilon$  are relaxed through the gHM condition

$$\frac{\sigma}{A_e \omega_c} \left| \nabla \mathbf{v}_E^2 \cdot \nabla \times \left( A_e \frac{\mathbf{B}}{B^2} \right) \right| \sim \epsilon^3. \quad (56)$$

If we now compare the GHM ordering of table 2 with the gHM ordering of table 4, one first notices that  $\mathbf{B}$  scales as  $\epsilon^{-1}$  in the GHM case, while it is treated as a  $\epsilon^0 \sim 1$  term in the gHM ordering. This difference does not change the order of the ratio  $v_E/v_{\text{pol}}$ , and it is therefore not essential (in fact, all dimensionless ratios have the same order in both orderings; compare the column ‘dimensionless’ in 2 and 4 with the fluid pressure  $P$  playing the role of the cyclotron temperature  $T_c$ ). The key difference is the absence of any

ordering requirement on derivatives of  $\mathbf{B}$  and  $A_e$  in the GHM ordering, and, in particular, the absence of the condition (56) found in the gHM case. This implies that the GHM model is free from any requirements on the geometry of the magnetic field  $\mathbf{B}$  or the spatial density  $A_e$ , and turbulence may develop over spatial scales  $k_{\perp}^{-1}$  comparable to  $1/|\nabla \log B|$  and  $1/|\nabla \log A_e|$ . This difference does not originate from a discrepancy between two-fluid theory and guiding center theory. Indeed, the ordering condition (56) used in [22] to enforce conservation of energy for the gHM equation (1) can be removed by adding higher order terms to (1) through the same procedure used to arrive at the GHM equation (34). Hence, one can obtain the same equation (34) from two fluid theory as well.

## 5 Concluding remarks

In this study, we derived the GHM equation (34) describing drift wave turbulence in curved magnetic fields within the framework of guiding center dynamics. This equation exhibits the same mathematical structure of the gHM equation (1) derived in [22] from a two-fluid model of an ion-electron plasma: the GHM equation can be obtained from the gHM equation by replacing the electrostatic potential  $\varphi$  with the reduced Northrop guiding center Hamiltonian  $\chi = \varphi + \frac{\sigma}{2} \mathbf{v}_{\mathbf{E}}^2$ . However, the ordering (table 2) used to obtain the GHM equation is weaker than the ordering (table 4) used in [22] to derive the gHM equation. In particular, while the gHM equation relies on the ordering condition (56) to enforce conservation of energy, the GHM ordering does not involve ordering conditions on spatial derivatives of the magnetic field  $\mathbf{B}$  or the electron spatial density  $A_e$ . As explained in section 4, the GHM equation can also be obtained from two-fluid theory as well by adding higher-order terms to the gHM equation. These results imply that the GHM equation is appropriate to describe drift wave turbulence in general magnetic fields and in the presence of strong inhomogeneities in the electron spatial density, including the case in which the scale of spatial change in the magnetic field and the electron spatial density is comparable to that of the turbulent electric field. This setting is expected to be physically relevant in strongly curved and inhomogeneous magnetic fields, such as dipole magnetic fields or the confining magnetic field of a stellarator. Finally, we remark that for practical purposes (e.g. numerical implementation), the solution  $\chi$  of the GHM equation can be used to approximate  $\varphi \approx \chi$  since both  $\chi$  and  $\varphi = \chi + o(1)$  scale as  $\epsilon^{-1}$ .

## Statements and declarations

### Data availability

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

### Funding

The research of NS was partially supported by JSPS KAKENHI Grant No. 21K13851. and 22H04936.

### Competing interests

The authors have no competing interests to declare that are relevant to the content of this article.

## References

- [1] A. Hasegawa and K. Mima, *Pseudo-three-dimensional turbulence in magnetized nonuniform plasma*, Phys. Fluids **21**, 1 (1977).
- [2] A. Hasegawa and K. Mima, *Stationary spectrum of strong turbulence in magnetized nonuniform plasma*, Phys. Rev. Lett. **39**, 4 (1977).

- [3] J. G. Charney, *On the scale of atmospheric motions*, Geof. Publ. **17**, 2 (1948).
- [4] J. G. Charney and P. G. Drazin, *Propagation of planetary-scale disturbances from the lower into the upper atmosphere*, J. Geophys. Res. **66**, 1 (1961).
- [5] W. Horton and A. Hasegawa, *Quasi-two-dimensional dynamics of plasmas and fluids*, Chaos **4**, 227 (1994).
- [6] G. K. Batchelor, *Computation of the energy spectrum in homogeneous two-dimensional turbulence*, The Physics of Fluids **12**, II-233 (1969).
- [7] R. H. Kraichnan and D. Montgomery, *Two-dimensional turbulence*, Rep. Prog. Phys. **43**, 547 (1980).
- [8] R. H. Kraichnan, *Inertial ranges in two-dimensional turbulence*, The Physics of Fluids **10**, 7 (1967).
- [9] M. K. Rivera, W. B. Daniel, S. Y. Chen, and R. E. Ecke, *Energy and enstrophy transfer in decaying two-dimensional turbulence*, Phys. Rev. Lett. **90**, 10 (2003).
- [10] Z. Xiao, M. Wan, S. Chen, and G. L. Eyink, *Physical mechanism of the inverse energy cascade of two-dimensional turbulence: a numerical investigation*, J. Fluid Mech. **619**, pp. 1-44 (2009).
- [11] A. Hasegawa and K. Mima, *Strong turbulence, self-organization and plasma confinement*, Eur. Phys. J. H **43**, pp. 499-521 (2018).
- [12] A. Hasegawa, *Self-organization processes in continuous media*, Adv. Physics **34**, 1, pp. 1-42 (1985).
- [13] W. Horton, *Drift waves and transport*, Rev. Mod. Phys. **71**, 3 (1999).
- [14] A. Fujisawa, K. Itoh, H. Iguchi, K. Matsuoka, S. Okamura, A. Shimizu et al., *Identification of Zonal Flows in a Toroidal Plasma* Phys. Rev. Lett. **93**, 16 (2004).
- [15] P. H. Diamond, A. Hasegawa, and K. Mima, *Vorticity dynamics, drift wave turbulence, and zonal flows: a look back and a look ahead*, Plasma Phys. Control. Fusion **53**, 124001 (2011).
- [16] A. C. Boxer, R. Bergmann, J. L. Ellsworth, D. T. Garnier, J. Kesner, M. E. Mauel, and P. Woskov, *Turbulent inward pinch of plasma confined by a levitated dipole magnet*, Nat. Phys. **6**, pp. 207-212 (2010).
- [17] N. Kenmochi, Y. Yokota, M. Nishiura, H. Saitoh, N. Sato, K. Nakamura, T. Mori, K. Ueda, and Z. Yoshida, *Inward diffusion driven by low frequency fluctuations in self-organizing magnetospheric plasma*, Nucl. Fusion **62**, 026041 (2022).
- [18] D. T. Garnier, M. E. Mauel, T. M. Roberts, J. Kesner, and P. P. Woskov, *Turbulent fluctuations during pellet injection into a dipole confined plasma torus*, Phys. Plasmas **24**, 012506 (2017).
- [19] T. S. Hahm *Nonlinear gyrokinetic equations for turbulence in core transport barriers*, Phys. Plasmas **3**, pp. 4658-4664 (1996).
- [20] T. S. Hahm and W. M. Tang, *Nonlinear theory of collisionless trapped ion modes*, Phys. Plasmas **3**, pp. 242-247 (1996).
- [21] T. S. Hahm, L. Wang, and J. Madsen, *Fully electromagnetic nonlinear gyrokinetic equations for tokamak edge turbulence*, Phys. Plasmas **16**, 022305 (2009).
- [22] N. Sato and M. Yamada, *A generalized Hasegawa Mima equation in curved magnetic fields*, J. Plasma Phys. **88**, 905880319 (2022).
- [23] R. D. Hazeltine and F. L. Waelbroeck, *Fluid description of a plasma*, in The Framework of Plasma Physics, Perseus Books, pp. 49-86 (1998).

- [24] J. R. Cary and A. J. Brizard, *Hamiltonian theory of guiding-center motion*, Rev. Mod. Phys. **81**, pp. 730-732 (2009).
- [25] T. G. Northrop, *Adiabatic Charged-Particle Motion*, Rev. Geophys. **1**, 3 (1963).