DETERMINATION OF THE SYMMETRY GROUP FOR SOME QRT ROOTS

GIORGIO GUBBIOTTI AND YANG SHI

ABSTRACT. We determine the affine Weyl symmetries of some two-dimensional birational maps known as QRT roots arising from Kahan–Hirota–Kimura discretisation of two different reduced Nahm systems. The main finding is that the symmetry types of these discrete systems are subgroups of the Weyl groups for Sakai's discrete Painlevé equations to which the QRT maps are the autonomous limits.

Contents

1.	Introducti	ion	1
2.	Backgrou	nd material	3
3.	The space	e of initial conditions of the maps φ_1 , φ_2 , and φ_s	11
4.	Singular f	ibre of type $A_2^{(1)}$ associated with the map φ_1	18
5.	Singular f	ibre $\zeta_2(s)$ of type $A_1^{(1)}$ associated with the map φ_s	24
6.	Singular f	ibre of type $A_1^{(1)}$ associated with the map φ_2	30
7.	Conclusio	n	33
Ap	pendix A.	Reduced Nahm systems of tetrahedral and octahedral type	34
Ap	pendix B.	Basic properties of the group $\widetilde{W}(E_6^{(1)})$	35
Ap	pendix C.	Basic properties of the group $\widetilde{W}(E_7^{(1)})$	37
Ret	ferences		38

1. INTRODUCTION

In this paper, we consider the geometric and algebraic structure of two QRT roots arising from Kahan–Hirota–Kimura (KHK) discretisation [19, 26, 27] of reduced Nahm systems [20]. We show that these systems admit, as symmetry group, certain subgroups of the affine Weyl groups appearing in Sakai's classification of second-order discrete Painlevé equations [40]. Our results are summarised in the following statements:

Theorem 1.1. Consider the map:

$$\varphi_1: (x, y) \mapsto \left(y, \frac{xy - 1}{2x - y}\right). \tag{1.1}$$

Date: May 29, 2023.

²⁰²⁰ Mathematics Subject Classification. 14H70, 17B22, 39A36.

Key words and phrases. Discrete integrable systems; Weyl groups; QRT roots.

Map φ_1 admits a covariant pencil p_1 of biquadratic curves whose singular fibres configuration is of type $(A_3 \times A_2 \times 2A_1)^{(1)}$. Moreover, to the singular fibre of type $A_2^{(1)}$ is associated a discrete integrable system with symmetry group of type $A_2^{(1)} \subset E_6^{(1)}$. In particular, we show that φ_1 is an element of quasi-translation in the extended affine Weyl group of type $E_6^{(1)}$.

Theorem 1.2. Consider the map:

$$\varphi_2: (x,y) \mapsto \left(y, \frac{2xy^2 - 3xy + 2x - 1}{xy^2 - 2y^2 + 3y - 2}\right).$$
 (1.2)

Map φ_2 admits a covariant pencil p_2 of biquadratic curves whose singular fibres configuration is of type $(3A_3 \times A_1)^{(1)}$. Moreover, the singular fibre of type $A_1^{(1)}$ is associated with a discrete integrable system with a symmetry group of type $B_3^{(1)} \subset E_7^{(1)}$. In particular, we show that φ_2 is an element of quasi-translation in the extended affine Weyl group of type $E_7^{(1)}$.

Theorem 1.2 will be proved using the following statement on a map which is a generalisation of the map φ_2 .

Theorem 1.3. Consider the map:

$$\varphi_s: (x,y) \mapsto \left(y, \frac{3sxy^2 - 3sxy + 2sy^2 - 4xy^2 + sx + 6yx - 4x + 2}{2sxy^2 - 3sy^2 - 2xy^2 + 2sx + 3sy + 4y^2 - s - 6y + 4}\right).$$
(1.3)

Map φ_s admits a covariant pencil p_s of biquadratic curves whose singular fibres configuration is of type $(2A_3 \times 2A_1 \times A_0)^{(1)}$. Moreover to the singular fibre of type $A_1^{(1)}$ is associated with a discrete integrable system with a symmetry group of type $B_3^{(1)} \subset E_7^{(1)}$. In particular, we show that φ_s is an element of quasi-translation in the extended affine Weyl group of type $E_7^{(1)}$.

The two maps φ_1 and φ_2 are two QRT roots related to the Kahan–Hirota–Kimura (KHK) discretisations [19, 26, 31] of so-called reduced Nahm systems of tetrahedral and octahedral type [20]. In particular, Equations (1.1) and (1.2) are obtained from the KHK discretisation of the tetrahedral and octahedral reduced Nahm systems with the method of [30]. The interested reader can find more details on this construction, with the explicit formulas are given in Appendix A. Finally, the map φ_s is a QRT root [11, Introduction] preserving a more general pencil than φ_2 . The map φ_s will be discussed more thoroughly in Section 3.

In [11], the geometry of QRT maps is discussed in the context of Oguiso and Shioda's classification of 74 types of singular fibre configurations of rational elliptic surfaces [32]. On the other hand, it was shown in [6] how different choices of singular fibres in a configuration lead to different autonomous Painlevé equations in Sakai's classification of second-order discrete Painlevé equations [40]. The approach given in [6] allows one to discuss a given QRT system within Sakai's framework.

In this paper, we analyse the two discrete systems given by the maps φ_1 and φ_2 in Equations (1.1) and (1.2) with tools from algebraic geometry and theory of the Weyl group. That is, we build the space of initial values and obtain the corresponding singular fibre configurations of the two maps. We find that the map φ_2 has a peculiar geometric structure, which is different from the canonical one presented in the foundational paper [40] and in the review [28]. For some choices of singular fibres from the two configurations, we study in detail the symmetries of

the associated discrete systems. In particular, we show that these discrete systems admit symmetries that do not appear explicitly in Sakai's classification. Such two-dimensional discrete integrable systems have been found in different contexts [1, 6, 7, 24, 29, 44, 46]. In [42], it was found that these symmetries arise as normalizers of certain parabolic subgroups of Sakai's Weyl groups using the normalizer theory of of Coxeter groups [5, 21]. Sakai's equations are realised as birational representations of affine Weyl groups of ADE type in the plane, see [41, Proposition 5.15]. In particular, the map that gives the iteration/dynamics of Sakai's equation corresponds to translation along a shortest weight on the weight lattice of the corresponding Weyl group [28, 40]. Here we show that the QRT roots considered here are quasitranslations. That is, an element of an affine Weyl group of infinite order, which after m number of iterations become a translation in the original Weyl group is called a quasi-translation of order m.

The plan of the paper is the following: in Section 2 we recall some basic facts from the theory of the elimination of the indeterminacies of maps of surfaces and the theory of Weyl groups. In Section 3 we construct the space of initial conditions of the maps φ_1 and φ_2 and identify the associated singular fibres configurations, that is the first statements in Theorems 1.1 and 1.2. Then, we proceed to introduce the map φ_s by constructing the QRT root preserving a pencil generalising the one preserved by φ_2 . We conclude the section by computing the configuration of the singular fibres associated to φ_s , thus proving the first statement of Theorem 1.3. Next, in Section 4 we prove the second Statement of Theorem 1.1. Then, in Section 5 we prove the last statement of Theorem 1.3 where we show that to singular fibre of type $A_1^{(1)}$ associated to the map φ_s (1.3) corresponds to a discrete system of type $B_3^{(1)} \subset E_7^{(1)}$. Using the results of Section 5 in Section 6 we are able to prove the last statement of Theorem 1.2, i.e. that to singular fibre of type $A_1^{(1)}$ associated to the map φ_2 (1.2) corresponds to a discrete system of type $B_3^{(1)} \subset E_7^{(1)}$. This result is obtained by showing that the action of the map φ_2 on the root lattice $E_7^{(1)}$ is the same as the one of φ_s , but on a translated sub-root lattice. Finally, in Section 7 we give some conclusions and outlook for further work.

2. Background material

In this section, we give the background we need for our study of the symmetries of the two maps in Equations (1.1) and (1.2). In particular, we will introduce the concept we need from the theory of elimination of the indeterminacy of surfaces following [2]. Following Sakai [40] this allows us to construct the *space of initial values* of a given map, which is the discrete analogue of Okamoto's description [33, 34] of the continuous Painlevé equations [23]. Moreover, we will need some basic facts from the theory of Weyl groups or more generally, the theory of Coxeter groups [4].

2.1. Birational maps and biquadratic pencils. In this section, we briefly recall some tools from the algebraic geometry of birational maps we need to carry out our constructions, namely the elimination of indeterminacies for maps of complex surfaces, and some generalities on singular fibres of pencils.

2.1.1. Eliminaton of indeterminacies for maps of surfaces. As stated in the introduction, we will consider rational maps of \mathbb{C}^2 to itself in the compactification $\mathbb{P}^1 \times \mathbb{P}^1$. In particular, we will consider birational maps, i.e. rational maps whose inverse is rational, and denote their space by $\operatorname{Bir}(\mathbb{P}^1 \times \mathbb{P}^1)$. An *indeterminacy point* is a point where a map $\varphi \in \operatorname{Bir}(\mathbb{P}^1 \times \mathbb{P}^1)$ is not well defined. Representing locally a map as pairs of rational functions, being defined corresponds to say that one of the entries is in the form 0/0. Indeterminacies are eliminated through the repeated application of a procedure called the blow-up at a given point, which can be described heuristically as the variety obtained by replacing a point with the set of lines passing through it, see [2, Section II.1]. A blow-up is a birational projective morphism, i.e. it is an isomorphism outside the given point, and the preimage of that point is called the *exceptional curve* (usually denoted by the letter E).

Since $\mathbb{P}^1 \times \mathbb{P}^1$ is a ruled surface (in particular, the simplest of the Hirzebruch surfaces), to eliminate the indeterminacies of $\varphi \in \text{Bir}(\mathbb{P}^1 \times \mathbb{P}^1)$ we use the following result from [2, Corollay II.12] adapted to our case:

Theorem 2.1. Let $\varphi \in Bir(S)$ with S surface. Then there exists a surface \widetilde{S} and a map $\widetilde{\varphi} \in Bir(\widetilde{S})$ such that the following diagram commute:

and the morphisms ε is the composition of blow-ups.

The map $\tilde{\varphi} \in \operatorname{Bir}(\tilde{S})$ in Theorem 2.1 is called the *lift* of φ and if no confusion arise we will denote it without the tilde. In the case of $\varphi \in \operatorname{Bir}(\mathbb{P}^1 \times \mathbb{P}^1)$, following Theorem 2.1, we eliminate the indeterminacies blowing up all the points of the following classes:

- indeterminacy points of the map φ and its inverse φ^{-1} ;
- indeterminacy points lying on an exceptional line E after blowing up, called *infinitely near points*;
- images of lines contracted by φ or φ^{-1} .

Remark 2.2. We remark that while in this paper we are following the the classical strategy of elimination of indeterminacies explained in Theorem 2.1 this is not the only possible approach. Indeed, the blow-up of infinitely near points can be achieved with a single blow-up using an approach based on the ideals of these points, see [13].

So, to eliminate the indeterminacies of a map $\varphi \in Bir(\mathbb{P}^1 \times \mathbb{P}^1)$ we have to blow-up a set of the form¹:

$$\operatorname{Sing}(\varphi) = \{b_{1,\dots,i_{1}}, b_{i_{1}+1,\dots,i_{2}},\dots, b_{i_{L-1}+1,\dots,i_{L}}\},$$
(2.2)

where $i_1 < i_2 < \ldots < i_{L-1} < i_L = M$, L, M are positive integers, and we denoted some infinitely near points by $b_{i_k,\ldots,i_{k+1}-1}$. Then on the space obtained by blowing up the set $\operatorname{Sing}(\varphi)$, written as B_{φ} , the map $\widetilde{\varphi}$ is an automorphism. The smooth variety B_{φ} is called the *space of initial values of the map* φ [40, 45]. This definition is stated in analogy with the same definition for continuous systems given by Okamoto [33, 34].

¹This is a slight abuse of notation because after blowing up points are replaced by lines, but we use them to keep track of which points originated a "chain" of successive blow-ups.

2.1.2. The Picard group. Given a smooth (quasi-)projective variety X, recall that [2, Fact I.1] the Picard group of X, Pic(X), is the group of isomorphism classes of line bundles on X. Line bundles can be identified with divisors up to linear equivalence, and throughout the paper we will make use of this identification. In the case of a surface B obtained blowing up $\mathbb{P}^1 \times \mathbb{P}^1$ M times the Picard group is the free \mathbb{Z} -module generated by the proper transform of classes of equivalence of horizontal and vertical lines:

$$H_x = \varepsilon^* \{ x = \text{const} \}, \quad H_y = \varepsilon^* \{ y = \text{const} \}, \quad (2.3)$$

and the exceptional lines E_1, E_2, \ldots, E_M :

$$\operatorname{Pic}(B) = \operatorname{Span}_{\mathbb{Z}}(H_x, H_y, E_1, \dots, E_M).$$

$$(2.4)$$

Notation 2.3. We will denote elements of the Picard group by capital letters while their (possible) representatives (complete linear systems of) curves will be denoted by small letters, e.g. $D \in \text{Pic}(B)$ and $d \in \mathbb{P}^1 \times \mathbb{P}^1$ is a curve.

The Picard group is equipped with a symmetric bilinear form (,) with integer values called the *intersection form*, see [2, Definition I.3 and Theorem I.4]. Let B be a surface as above, and then the intersection form satisfies the following rules:

$$(H_x, H_x) = (H_y, H_y) = (H_x, E_i) = (H_y, E_i) = 0, (H_x, H_y) = 1, \quad (E_i, E_j) = -\delta_{i,j},$$
(2.5)

where $i, j = 1, \ldots, M$, see Equation (2.4).

We will denote the action induced by a morphism $\varphi: B_{\varphi} \to B_{\varphi}$ on the Picard group by $\varphi^*: \operatorname{Pic}(B_{\varphi}) \to \operatorname{Pic}(B_{\varphi})$. This action is *linear* and can be represented by a matrix with integer entries.

2.1.3. Rational elliptic fibrations and minimality. It is known that integrable maps in the plane are related to rational elliptic surfaces or their generalisation [3, 7, 47]. We recall that a surface is elliptic if it admits an elliptic fibration. Such a surface is a rational elliptic surface if the fibres are connected, see [11, Theorem 9.1.3 and Definition 9.1.4]. This implies that there these maps are birationally equivalent to a map φ_{\min} such that $|\text{Sing}(\varphi_{\min})| = 8$, that is $i_L = 8$ in equation (2.2), see also [8]. For a "generic" map whose space of initial values is a rational elliptic surface we have $|\text{Sing}(\varphi)| \ge 8$. From now on, we will assume that we are considering the "minimal" map such that this lower bound is obtained, i.e. $\varphi = \varphi_{\min}$. In the case of a rational elliptic surface B_{φ} the anti-canonical divisor takes the following form:

$$-K_{B_{\varphi}} = 2H_x + 2H_y - \sum_{k=1}^{8} E_i.$$
(2.6)

Moreover, in this case the anti-canonical divisor, or a multiple of it, is represented by a *pencil of biquadratic curves* whose base points are the points b_k with k = 1, ..., 8. We recall that a point $b \in \mathbb{P}^1 \times \mathbb{P}^1$ is a *base point* for a pencil p if $p(b; \mu : \nu) = 0$ regardless of the value of $[\mu : \nu] \in \mathbb{P}^1$. The map on the Picard group φ^* preserves the anti-canonical divisor, that is:

$$\varphi^*(-K_{B_{\varphi}}) = -K_{B_{\varphi}}.$$
(2.7)

This, in turn implies that taking the ratio of the biquadratic polynomials defining the pencil p, i.e. the rational function $\mathcal{H} = q/r$, we obtain an invariant for the map $\varphi \in \operatorname{Bir}(\mathbb{P}^1 \times \mathbb{P}^1)$, or a k-invariant in the case the map does not preserve the fibres of the pencil [7]. In the rest of the paper we will only consider fibre-preserving actions, we will not need the notion of k-invariant we will not enter into the details of this approach, but we refer the interested reader to the papers [12, 16, 17, 39]. So, from now on when we will say *covariant pencil* p we will mean that $\varphi^*(p) = \kappa p$ for some κ bihomogeneous polynomial and additionally the fibres of the pencil are preserved.

Remark 2.4. An alternative construction of integrable maps starts from a given pencil of biquadratic maps and then considers the blow-up of its base points. This yields the rational elliptic surface B_{φ} again. The map $\varphi \in Bir(\mathbb{P}^1 \times \mathbb{P}^1)$ is obtained from the pencil through the application of some properly defined involutions, see [11, 37, 38].

2.1.4. *Singular fibres and root systems.* Before going on we recall the following definition of singular fibres of a pencil:

Definition 2.5. Let

$$p(x, y; \mu:\nu) = \mu p_0(x, y) + \nu p_1(x, y), \qquad (2.8)$$

be a *pencil* of plane curves. A point $(x_0, y_0; \mu' : \nu') \in \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ such that

$$p(x_0, y_0; \mu': \nu') = \frac{\partial p}{\partial x}(x_0, y_0; \mu': \nu') = \frac{\partial p}{\partial y}(x_0, y_0; \mu': \nu') = 0, \qquad (2.9)$$

where the derivatives are taken in a properly chosen chart, is called a *singular point* for the pencil (2.8). Given a pencil of plane curves p if the curve $p(x, y; \mu' : \nu')$, with fixed $[\mu' : \nu'] \in \mathbb{P}^1$ contains a singular point then it is called a *singular fibre* of the pencil (2.8).

The singular fibres of rational elliptic surfaces have been extensively studied in the literature, see [41] for a general overview and [11, Chap. 7] for some applications to integrable systems. Here, following [28], we limit ourselves to note that to a singular fibres of a pencil p on the space B_{φ} the anti-canonical divisor splits into several *rational* components. That is, given a singular fibre $[\mu' : \nu'] \in \mathbb{P}^1$ there exists some divisors D_j , $j = 0, \ldots, \kappa$ such that:

$$-K_{B_{\varphi}} = \sum_{j=0}^{\kappa} D_j, \qquad (2.10)$$

which realise this singular fibre. Moreover, the intersection matrix:

$$C = (-(D_i, D_j))_{i,j=0}^{\kappa}, \qquad (2.11)$$

is a generalised Cartan matrix of a Coxeter group of rank κ , see [41, Proposition 5.15]. We call the type of root system defined by this generalised Cartan matrix *the* surface type, denoted by Λ , for a singular fibre for the map φ . That is, the divisors D_k correspond to the simple roots of Λ . The orthogonal complement of Λ , Λ^{\perp} , is another root system, called the symmetry type of the map φ . Note that the surface type associated with the map φ depends on the choice of the singular fibre as noted in [8]. Indeed, the possible surface type associated with the singular fibres are those contained in [41, Theorems 8.8 and 8.9].

To summarise, for a given birational map $\varphi \in Bir(\mathbb{P}^1 \times \mathbb{P}^1)$, one can eliminate its indeterminacies and those of its inverse φ^{-1} and build the space of initial conditions B_{φ} . If the map is integrable, then B_{φ} is a rational elliptic surface, and to each of its singular fibres there are associated with two orthogonal root systems Λ and Λ^{\perp} . The simple roots of these systems give the surface and the symmetry type, respectively, and can both be represented by Dynkin diagrams. The pullback φ^* acts on the simple roots of Λ as a permutation, while in general, it acts on the simple roots of Λ^{\perp} as a quasi-translation. In the next subsection, we will recall some facts from the theory of Weyl groups needed to discuss the natures of the maps φ_1 and φ_2 and the symmetries of the associated discrete systems.

2.2. Weyl group. The symmetry groups of affine Weyl type for the two discrete systems given by maps (1.1) and (1.2) can be constructed using the geometric characterisation given in Section 2.1. Here we recall necessary facts and properties of the Weyl group in order for us to construct the groups of symmetries and the corresponding group elements that are associated to the two maps (1.1) and (1.2).

2.2.1. Dynkin diagram and root system. Let Γ be a Dynkin diagram with n nodes, its associated Coxeter group is a generating set,

$$W = W(\Gamma) = \langle s_i \mid 1 \le i \le n \rangle \tag{2.12}$$

with *defining relations*:

$$s_i^2 = 1, \quad (s_i s_j)^{m_{ij}} = 1, \ 1 \le i < j \le n.$$
 (2.13)

When $m_{ij} \in \{2, 3, 4, 6\}$, known as the *crystallographic condition*, W is called a Weyl group. Each node of Γ , labelled *i* represents the generator s_i of order 2. The parameter m_{ij} takes value of: 2, 3, 4 or 6 when two nodes labeled *i* and *j* are: disconnected, joined by a single, a double, or a triple bond, respectively. Diagrams which have only single bonds are called *simply-laced*, they are of types A_n , D_n , E_6 , E_7 and E_8 . The non-simply laced types are B_n , C_n , F_4 and G_2 (see Humphries [22, Sec. 2.5] for example). Let $\Gamma^{(1)}$ be the affine extension of Γ (with an extra node labelled 0), then $W^{(1)} = \langle s_i | 0 \leq i \leq n \rangle$ is the associated affine Weyl group.

Let $V^{(1)}$ be an n + 1-dimensional real vector space with a basis,

$$\Delta^{(1)} = \Delta(\Gamma^{(1)}) = \{ \alpha_i \in V^{(1)} \mid 0 \le i \le n \},$$
(2.14)

known as the simple system, and α_i $(0 \le i \le n)$ are the simple roots of $W^{(1)}$. Note that $\Delta = \Delta(\Gamma) = \{\alpha_i \in V^{(1)} \mid 1 \le i \le n\}$ is the simple system of W, and $V = \text{Span}(\Delta)$ is a subspace of $V^{(1)}$.

A semidefinite symmetric bilinear form on $V^{(1)}$ can be defined using the *Cartan* matrix of type $\Gamma^{(1)}$, $C(\Gamma^{(1)}) = (a_{ij})_{1 \le i,j \le n,0}$ by,

$$\alpha_i \cdot \alpha_j = \frac{|\alpha_j|^2 a_{ij}}{2}, \quad \text{for} \quad i, j \in \{0, 1, ..., n\},$$
(2.15)

where the quantity $\alpha_j \cdot \alpha_j = |\alpha_j|^2$ gives the usual interpretation of squared length of α_j . The generator $s_j \in W^{(1)}$ known as a *simple reflection*, can be realised as the reflection along $\alpha_j \in V^{(1)}$, its action on $\Delta^{(1)}$ given by

$$s_j(\alpha_i) = s_{\alpha_j}(\alpha_i) = \alpha_i - \frac{2(\alpha_i \cdot \alpha_j)}{\alpha_j \cdot \alpha_j} \alpha_j = \alpha_i - a_{ij}\alpha_j, \quad \text{for all} \quad i, j \in \{0, 1, ..., n\}, (2.16)$$

where a_{ij} is the (i, j)-entry of $C(\Gamma^{(1)})$.

Note that nodes of a Dynkin diagram can be equivalently associated with either simple reflections or simple roots of the Weyl group.

There is a null root δ in $V^{(1)}$ such that

$$\alpha_i \cdot \delta = 0 \quad \text{for all} \quad \alpha_i \in V^{(1)}, \quad 0 \le i \le n.$$
(2.17)

In particular, we have

$$\delta = \alpha_0 + \tilde{\alpha} = \alpha_0 + \sum_{i=1}^n c_i \alpha_i = \sum_{i=0}^n c_i \alpha_i, \qquad (2.18)$$

where $\tilde{\alpha}$ is the *highest long root* of W, and values of the c_i 's can be found in any classic textbooks on Coxeter groups such as Bourbaki [4, Planche I–X]. Finally, the root system of $W^{(1)}$ is given by

$$\Phi^{(1)} = \Phi(\Gamma^{(1)}) = W^{(1)} \cdot \Delta^{(1)} = \{ \alpha + m\delta \mid \alpha \in \Phi, \ m \in \mathbb{Z} \},$$
(2.19)

where $\Phi = W.\Delta$ is the finite root system of W in V. Moreover, we have $\Phi^{(1)} = \Phi^{(1)}_+ \cup \Phi^{(1)}_-$ where,

$$\Phi_{+}^{(1)} = \{ \alpha = \sum_{i=0}^{n} \lambda_{i} \alpha_{i} \mid \mathbb{Z} \ni \lambda_{i} \ge 0, \alpha_{i} \in \Delta^{(1)} \} \text{ and } \Phi_{-}^{(1)} = \{ -\alpha \mid \alpha \in \Phi_{+}^{(1)} \}, \quad (2.20)$$

and Φ_+ is the positive root system.

In general, the element of reflection along any root $\beta \in \Phi^{(1)}$, denoted by $s_{\beta} \in W^{(1)}$, acts on $V^{(1)}$ by,

$$s_{\beta}(v) = v - \frac{2v \cdot \beta}{\beta \cdot \beta} \beta, \qquad (2.21)$$

for all $v \in V^{(1)}$. Furthermore, s_{β} is related to a simple reflection s_i by the formula

$$s_{\beta} = s_{w(\alpha_i)} = w s_i w^{-1}, \quad \text{for some} \quad w(\alpha_i) = \beta, \quad w \in W^{(1)}.$$
(2.22)

2.2.2. Dual space, coroots and weights. By Equation (2.18), we see that $\{\alpha_1, \alpha_2, \ldots, \alpha_n, \delta\}$ forms another basis of $V^{(1)}$. Let $V^{(1)*}$ be an (n+1)-dimensional real vector space, and $\langle , \rangle : V^{(1)} \times V^{(1)*} \to \mathbb{R}$ be a bilinear pairing between $V^{(1)}$ and $V^{(1)*}$. Let $\{h_1, \ldots, h_n, h_\delta\}$ be a basis of $V^{(1)*}$ dual to $\{\alpha_1, \alpha_2, \ldots, \alpha_n, \delta\} \in V^{(1)}$. This is,

$$\langle \alpha_i, h_j \rangle = \delta_{ij}, \tag{2.23a}$$

$$\langle \alpha_i, h_\delta \rangle = \langle \delta, h_j \rangle = 0, \quad \text{for} \quad 1 \le i, j \le n,$$
 (2.23b)

$$\langle \delta, h_\delta \rangle = 1. \tag{2.23c}$$

The group $W^{(1)}$ acts on $V^{(1)*}$ via the contragredient action:

$$\langle w^{-1}v,h\rangle = \langle v,wh\rangle, \text{ for } v \in V^{(1)}, h \in V^{(1)*}, w \in W^{(1)}.$$
 (2.24)

The vectors $h_j \in V^{(1)*}$ $(1 \le j \le n)$ are called the fundamental weights of $W^{(1)}$, and the weight lattice is given by,

$$P = \operatorname{Span}_{\mathbb{Z}}(h_1, \dots, h_n).$$
(2.25)

 $W^{(1)}$ acts on h_j $(1 \le j \le n)$ by,

$$s_i(h_j) = \begin{cases} h_j, & \text{for } i \neq j, \\ h_j - \sum_{k=1}^n a_{ki}h_k, & \text{for } i = j, & \text{for } 1 \le i, j \le n \end{cases}$$
(2.26a)

and

$$s_0(h_j) = h_j + c_j \sum_{k=1}^n a_{k0} h_k,$$
 (2.26b)

where c_i 's are the coefficients of α_i in δ , and a_{ij} are the entries of $C(\Gamma^{(1)})$.

In [43], we discussed some useful properties and formulas of the Weyl groups that enable us to describe quantitatively different types of translational elements of the Weyl group. Here, we simply state the results needed from [43]. First we define an *n*-dimensional hyperplane in $V^{(1)*}$,

$$X_k = \{ h \in V^{(1)*} | \langle \delta, h \rangle = k, k \in \mathbb{R} \}.$$

$$(2.27)$$

Note that, X_0 is an *n*-dimensional vector space, and by Equation (2.26a) we see that $\{h_j \mid 1 \leq j \leq n\}$ forms a basis of X_0 . Moreover, we have $X_1 = h_{\delta} + X_0$ is the affine plane on which translations of the affine Weyl group can be realised.

For each $\beta \in \Phi^{(1)}$, let

$$\beta^{\vee} = \frac{2\beta}{\beta \cdot \beta} = \frac{2\beta}{|\beta|^2}.$$
(2.28)

The set of simple coroots $\{\pi(\alpha_i^{\vee}) \in V^{(1)*} \mid 1 \leq j \leq n\}$ is a basis of X_0 given by,

$$\pi(\alpha_j^{\vee}) = \sum_{k=1}^n a_{kj} h_k = \sum_{k=1}^n \left(C(\Gamma)^T \right)_{jk} h_k, \quad \text{for} \quad 1 \le j \le n,$$
(2.29)

or we have

$$h_{i} = \sum_{k=1}^{n} \left(C(\Gamma)^{T} \right)_{ik}^{-1} \pi(\alpha_{k}^{\vee}), \quad \text{for} \quad 1 \le i \le n.$$
(2.30)

The following proposition enables us to discuss the lengths of vectors in the subspace $X_0 \subset V^{(1)*}$.

Proposition 2.6. The bilinear pairing \langle , \rangle given in Equation (2.23) can be restricted to a symmetric and positive definite bilinear form $(,): X_0 \times X_0 \to \mathbb{R}, X_0 \in V^{(1)*},$ satisfying:

$$\left(\pi(\alpha_i^{\vee}), \pi(\alpha_j^{\vee})\right) = \frac{2}{|\alpha_i|^2} a_{ij}, \quad 1 \le i, j \le n.$$

$$(2.31)$$

In the $\{h_j \mid 1 \le j \le n\}$ basis we have,

$$(h_i, h_j) = \sum_{k=1}^n \left(C(\Gamma)^T \right)_{ik}^{-1} \frac{2}{|\alpha_k|^2} \delta_{kj}, \quad 1 \le i, j \le n.$$
(2.32)

Remark 2.7. One can identify the simple coroots with simple roots by $\pi(\alpha_i^{\vee}) = 2\alpha_i/|\alpha_i|^2$. Then we see that for simply-laced type systems, we have $\pi(\alpha_j^{\vee}) = \alpha_j$ $(1 \le j \le n)$, and $\pi(\alpha_0^{\vee}) = -\tilde{\alpha}$.

Finally, let l(w) be the *length* of $w \in W^{(1)}$, we have,

$$l(ws_{\alpha}) = \begin{cases} l(w) + 1, & \text{if } w(\alpha) \in \Phi_{+}^{(1)} \\ l(w) - 1, & \text{if } w(\alpha) \in \Phi_{-}^{(1)}, \end{cases}$$
(2.33)

where $\alpha \in \Delta^{(1)}$. Equation (2.33) can be applied repeatedly until we can write w as a product of k simple reflections,

$$ws_{l_1}...s_{l_k} = 1, \quad l_1,...,l_k \in \{0, 1, ..., n\},$$
that is, $w = s_{l_k}...s_{l_1}.$
(2.34)

2.2.3. Translations. It is well-known that $W^{(1)}$ suitably extended contains an abelian group of translations on the weight lattice (which is isomorphic to the weight lattice P),

$$\widetilde{W}^{(1)} = A \ltimes W^{(1)} = W \ltimes P = W \ltimes \langle u_j \mid 1 \le j \le n \rangle, \tag{2.35}$$

where A is a group of diagram automorphisms of the affine Dynkin diagram $\Gamma^{(1)}$, and $u_j \in \widetilde{W}^{(1)}$ is defined to be the element that acts on X_1 by a translation of the fundamental weight h_j ,

$$u_j(X_1) = X_1 + h_j, \quad h_j \in X_0, \quad 1 \le j \le n.$$
 (2.36)

In general, actions of u_i on $V^{(1)*}$ and $V^{(1)}$ are given by the following proposition.

Proposition 2.8. The element $u_i \in \widetilde{W}^{(1)}$ acts on $V^{(1)*}$ by

$$u_j(h_i) = h_i, \quad and \quad u_j(h_\delta) = h_\delta + h_j, \qquad 1 \le i, j \le n,$$
(2.37a)

while on $V^{(1)}$ we have for $1 \le i, j \le n$,

$$u_j(\alpha_i) = \begin{cases} \alpha_i, & \text{for } i \neq j, \\ \alpha_j - \delta, & \text{for } i = j, \end{cases}$$
(2.38a)

and

$$u_j(\alpha_0) = \alpha_0 + c_j \delta, \tag{2.38b}$$

where c_j is the coefficient of α_j in δ .

In particular, if we have $s(h_j) = \sum_{i=1}^n n_{ij}h_i$ for $s \in W$ and some $n_{ij} \in \mathbb{Z}$ then,

$$su_j s^{-1} = \prod_{i=1}^n u_i^{n_{ij}}.$$
 (2.39)

Finally, any element u of $\widetilde{W}^{(1)}$ can be written in the form aw, with $a \in A$, and $w \in W^{(1)}$ with l(w) = k. That is, we have

$$u = aw = as_{l_k}...s_{l_1}, \quad l_1, ..., l_k \in \{0, 1, ..., n\},$$
(2.40)

where s_i are the simple reflections of $W^{(1)}$.

Remark 2.9. We will use the notation $\alpha_i + ... + \alpha_j = \alpha_{i...j}$ for a sum of simple roots, and $s_{i...s_j} = s_{i...j}$ for a product of simple reflections, from time to time for simplicity. For example, the null root of $E_6^{(1)}$ (see Equation (B.2)) can be written as $\delta = \alpha_{015224466333}$, while a product of simple reflections such as $s_{4s_5s_3s_4}$ is now s_{4534} .

2.2.4. Normalizers and quasi-translations. For $\widetilde{W}^{(1)}$ with a simple system $\Delta^{(1)}$, let $J \subset \Delta^{(1)}$, the corresponding reflection group $W_J = \langle s_i \mid \alpha_i \in J \rangle$ is called the *standard* parabolic subgroup of $\widetilde{W}^{(1)}$. The normaliser of W_J in $\widetilde{W}^{(1)}$ is defined by

$$N(W_J) = \{ g \in \widetilde{W}^{(1)} \mid g^{-1} W_J g = W_J \}.$$
(2.41)

In [5, 21] it is shown that

$$N(W_J) = N_J \ltimes W_J, \quad \text{where} \quad N_J = \{ w \in \widetilde{W}^{(1)} \mid wJ = J \}.$$

$$(2.42)$$

In particular, the group N_J is generated by *R*-elements and *M*-elements. The R-elements, also known as *quasi-reflections*, are in general involutions that act permutatively on the subset J, whereas the M-elements permute the R-elements. Together, they generate a group of extended affine Weyl type for which translations

(or quasi-translations) can be defined. In Sections 4.2 and 5.2 we construct the corresponding $N(W_J)$'s of which the QRT roots are elements.

Let $V_J = \text{Span}(J)$, and V_J^{\perp} be the orthogonal complement of V_J in $V^{(1)}$, that is, $V^{(1)} = V_J \bigoplus V_J^{\perp}$. Then on V_J^{\perp} , the R-elements and quasi-translations behave as reflections and translations, respectively. In Sections 4.2.1 and 5.2.1, we find a basis for V_J^{\perp} , which forms a root system for N_J and a subroot system of Λ^{\perp} . Using properties of this subroot system we show that QRT roots are elements of quasi-translations in $N(W_J)$.

3. The space of initial conditions of the maps φ_1 , φ_2 , and φ_s

In this section, we build the space of initial conditions of the maps φ_1 (1.1) and φ_2 (1.2), and introduce the map φ_s (1.3) as a generalisation of φ_2 . Additionally, we identify the configuration of the singular fibres of the pencils associated to these three maps within Oguiso and Shioda's classification of 74 types of singular fibre configurations of rational elliptic surfaces [32]. We will mainly follow the exposition of this theory from the book [41].

3.1. The space of initial values of φ_1 . In this section, we prove the first statement of Theorem 1.1 on the existence of a covariant pencil with singular fibres of type $(A_3 \times A_2 \times 2A_1)^{(1)}$ for the map φ_1 (1.1). We do so by constructing a space of initial values for such a map. Following Section 2, a space of initial values of the map φ_1 (1.1) is constructed blowing up its singular locus. By direct computations, we have:

$$S_1 \coloneqq \operatorname{Sing} \varphi_1 = (b_1, \dots, b_8) \tag{3.1}$$

where:

$$b_{1}:(x,y) = \left(\sqrt{2}, \frac{1}{\sqrt{2}}\right), \quad b_{2}:(x,y) = \left(-\sqrt{2}, -\frac{1}{\sqrt{2}}\right),$$

$$b_{3}:(x,y) = \left(-\frac{1}{\sqrt{2}}, -\sqrt{2}\right), \quad b_{4}:(x,y) = \left(\frac{1}{\sqrt{2}}, \sqrt{2}\right),$$

$$b_{5}:(x,y) = \left(\frac{1}{\sqrt{2}}, 0\right), \quad b_{6}:(x,y) = \left(-\frac{1}{\sqrt{2}}, 0\right),$$

$$b_{7}:(x,y) = \left(0, -\frac{1}{\sqrt{2}}\right), \quad b_{8}:(x,y) = \left(0, \frac{1}{\sqrt{2}}\right).$$

(3.2)

We remark that not all the base points are immediately visible as indeterminacy points of φ_1 and its inverse, but some appear as a contraction of the exceptional lines associated with a point after blowing up. We call E_i the exceptional line associated to the blowing up of the indeterminacy point b_i . Then we have that b_i , i = 5, 6, 7, 8, appear as image of the exceptional lines E_1 , E_2 and E_3 , E_4 under φ_1 and φ_1^{-1} respectively.

Then, a space of initial values is $X_1 = B_{\varphi_1}$, a rational elliptic surface. A biquadratic pencil is defined by the condition of passing through the points b_i :

$$p_1(x,y;\mu:\nu) = \mu[\sqrt{2}(x-y)+1][\sqrt{2}(x-y)-1] + \nu xy(xy-1).$$
(3.3)

By construction this pencil is covariant with respect to the map φ_1 (1.1).

Then, from Definition 2.5 we have that there are four distinct singular fibres corresponding to the values of $[\mu : \nu]$ shown in the first column of Table 1. The irreducible components of the associated singular fibres are shown in the second column of the same table.

Point $[\mu:\nu]$	Irred. comp. d_i	Repr. in $\operatorname{Pic}(X_1)$ D_i	Surf./symm. type Λ/Λ^{\perp}
[0:1]	$ \{ x = 0 \} \\ \{ y = 0 \} \\ \{ xy = 1 \} $	$H_{y} - E_{5} - E_{6}$ $H_{x} - E_{7} - E_{8},$ $H_{x} + H_{y} - E_{1} - E_{2} - E_{3} - E_{4}$	$A_2^{(1)}/E_6^{(1)}$
[1:0]	$ \begin{cases} y - x = -1/\sqrt{2} \\ y - x = 1/\sqrt{2} \end{cases} $	$ \begin{array}{l} H_x + H_y - E_1 - E_3 - E_5 - E_7 \\ H_x + H_y - E_2 - E_4 - E_6 - E_8 \end{array} $	$A_1^{(1)}/E_7^{(1)}$
[1:4]	$ \begin{cases} x = -1/\sqrt{2} \\ \{y = 1/\sqrt{2} \\ \{x = 1/\sqrt{2} \\ \{y = -1/\sqrt{2} \} \end{cases} $	$H_{x} - E_{3} - E_{6}$ $H_{y} - E_{1} - E_{8}$ $H_{x} - E_{4} - E_{5}$ $H_{y} - E_{2} - E_{7}$	$A_3^{(1)}/D_5^{(1)}$
[1:16]	$ \{4xy + \sqrt{2}(x+y) = 1\} \{4xy - \sqrt{2}(x+y) = 1\} $	$\begin{array}{l} H_x + H_y - E_2 - E_3 - E_5 - E_8 \\ H_x + H_y - E_1 - E_4 - E_6 - E_7 \end{array}$	$A_1^{(1)}/E_7^{(1)}$

TABLE 1. Irreducible components of the singular fibres in the pencil associated to (3.4), their representative in the Picard groups, and the associated surface and symmetry type.

Remark 3.1. From the covariant pencil (3.3) we can build the rational function:

$$\mathcal{H}_1 = \frac{\left[\sqrt{2}(x-y)+1\right]\left[\sqrt{2}(x-y)-1\right]}{xy(xy-1)}$$
(3.4)

which is an invariant for the map φ_1 (1.1). This invariant was shown first in [15], where it was derived from the known invariant for the map (A.8) using the birational conjugation with the map (A.11a).

We have associated to X_1 the Picard group:

$$\operatorname{Pic}(X_1) = \operatorname{Span}_{\mathbb{Z}}(H_x, H_y, E_1, \dots, E_8).$$
(3.5)

The pullback of φ_1 act linearly on the Picard group of X_1 as follows:

$$\varphi_1^*: \begin{pmatrix} H_x, H_y, E_1, E_2, E_3, \\ E_4, E_5, E_6, E_7, E_8 \end{pmatrix} \mapsto \begin{pmatrix} H_x + H_y - E_1 - E_2, H_x, E_5, E_6, H_x - E_2, \\ H_x - E_1, E_7, E_8, E_3, E_4 \end{pmatrix}.$$
(3.6)

Furthermore, the representatives on the Picard group of the irreducible components of the singular fibres are given in the third column of Table 1. Using the intersection form (2.5), we can find the associated surface type. This is given in the fourth and last column of Table 1. This implies that the rational elliptic fibration has a singular fibre configuration of type $(A_3 \times A_2 \times 2A_1)^{(1)}$, which is number 59 in Oguiso and Shioda's list [41, Table 8.2]. This ends the proof of the first statement of Theorem 1.1. The structure of the singular fibres is summarised in Figure 1.

The symmetry type Λ^{\perp} of a map is the orthogonal complement of the surface type in the Picard group. In particular, in the case of $\Lambda = A_2^{(1)}$ we have $\Lambda^{\perp} = E_6^{(1)}$. In Section 4 we will show that map φ_1 (1.1), when considered on a $A_2^{(1)}$ type surface admits $A_2^{(1)} \subset E_6^{(1)}$ type symmetry.



FIGURE 1. A graphical representation of the singular fibres of the pencil p_1 .

3.2. The space of initial values of φ_2 . In this section, we prove the first statement of Theorem 1.2 on the existence of a covariant pencil with singular fibres of type $(3A_3 \times A_1)^{(1)}$ for the map φ_2 (1.2). Again, we construct a space of initial values for such a map. Following Section 2 a space of initial values of the map φ_2 (1.2) is constructed by blowing up its singular locus. By direct computations, we have:

$$S_2 \coloneqq \operatorname{Sing} \varphi_2 = (c_1, \dots, c_{7,8})$$
 (3.7)

where:

$$c_{1}:(x,y) = \left(-\frac{1+i\sqrt{3}}{2}, \frac{1-i\sqrt{3}}{2}\right), \quad c_{2}:(x,y) = \left(\frac{1-i\sqrt{3}}{2}, -\frac{1+i\sqrt{3}}{2}\right),$$

$$c_{3}:(x,y) = \left(-\frac{1-i\sqrt{3}}{2}, \frac{1+i\sqrt{3}}{2}\right), \quad c_{4}:(x,y) = \left(\frac{1+i\sqrt{3}}{2}, -\frac{1-i\sqrt{3}}{2}\right),$$

$$c_{5}:(x,y) = \left(-\frac{1+i\sqrt{3}}{2}, -\frac{1-i\sqrt{3}}{2}\right), \quad c_{6}:(x,y) = \left(-\frac{1-i\sqrt{3}}{2}, -\frac{1+i\sqrt{3}}{2}\right),$$

$$c_{7}:(x,y) = (1,1), \quad c_{8}:\left(x-1, \frac{y-1}{x-1}\right) = (0,-1).$$
(3.8)

We remark that not all the base points are immediately visible as indeterminacy points of φ_2 (1.2) and its inverse, but some appear as a contraction of the exceptional lines associated to a point after blowing up. We call F_i the exceptional line associated to the blowing up of the indeterminacy point c_i . Then, c_i , i = 5, 6, appear as image of the exceptional lines F_1 and F_4 under φ_2 and φ_2^{-1} respectively. Finally, c_8 arises after blowing-up c_7 in the coordinate system specified in its defining equation, i.e. c_7 and c_8 are infinitely near points.

Then, a space of initial values is $X_2 = B_{\varphi_2}$, a rational elliptic surface. A biquadratic pencil p_2 is defined by the condition of passing through the points c_i :

$$p_2(x, y; \mu : \nu) = \mu(x^2y^2 - 2x^2y - 2xy^2 + 4x^2 - 2xy + 4y^2 + 2x - 2y + 1) + \nu(xy - 1)(xy + 1).$$
(3.9)

Point	Irred. comp.	Repr. in $\operatorname{Pic}(X_2)$	Surf./symm. type
$\lceil n \cdot n \rceil$	u_i	$ u_i $	11/11
[0:1]	$\{xy + 1 = 0\},\ \{xy - 1 = 0\},\$	$\begin{array}{l} H_{x}+H_{y}-F_{1}-F_{2}-F_{3}-F_{4}\\ H_{x}+H_{y}-F_{5}-F_{6}-F_{7}-F_{8} \end{array}$	$A_1^{(1)}/E_7^{(1)}$
[1:0]	$\begin{cases} xy - (1 + i\sqrt{3})x - (1 - i\sqrt{3})y + 1 = 0 \\ xy - (1 - i\sqrt{3})x - (1 + i\sqrt{3})y + 1 = 0 \\ \frac{y-1}{x-1} = 0 \\ 0 \\ \end{bmatrix} \cup \begin{cases} \frac{x-1}{y-1} = 0 \\ \frac{y-1}{y-1} = 0 \end{cases}$	$H_x + H_y - F_2 - F_3 - F_5 - F_7$ $H_x + H_y - F_1 - F_4 - F_6 - F_7$ $F_7 - D_8$	$A_2^{(1)}/E_6^{(1)}$
$\left[-\mathrm{i}\sqrt{3},9 ight]$	$\begin{cases} y = -\frac{1+i\sqrt{3}}{2} \\ x = -$	$H_{y} - F_{2} - F_{6},$ $H_{x} - F_{1} - E_{5}$ $H_{x} + H_{y} - F_{3} - F_{4} - F_{7} - F_{8}$	$A_2^{(1)}/E_6^{(1)}$
$[i\sqrt{3},9]$	$\begin{cases} x = -\frac{1-i\sqrt{3}}{2} \\ y = -\frac{1-i\sqrt{3}}{2} \\ 7xy + (2i\sqrt{3} - 4)(x + y) - 4i\sqrt{3} + 1 = 0 \end{cases}$	$\begin{array}{c} H_{x}-F_{3}-F_{6}\\ H_{y}-F_{4}-F_{5}\\ H_{x}+H_{y}-E_{1}-E_{2}-E_{7}-E_{8} \end{array}$	$A_2^{(1)}/E_6^{(1)}$

TABLE 2. Irreducible components of the singular fibres in the pencil associated to (3.9), their representative in the Picard groups, and the associated surface and symmetry type.

By construction, this pencil is covariant with respect to the map φ_2 (1.2).

Then, from Definition 2.5 we have that there are four singular fibres corresponding to the four values of $[\mu : \nu]$ shown in the first column of Table 2. The irreducible components of the associated singular fibres are shown in the second column of the same table.

Remark 3.2. We remark that in the singular fibres presented in Table 2 the arrangement of the base points is different from the standard one presented e.g. in [28]. In particular, the presence of the infinitely near points b_7 and b_8 implies that X_2 is not isomorphic to the elliptic fibration given in [28].

Remark 3.3. From the covariant pencil (3.9) we can build the rational function:

$$\mathcal{H}_2 = \frac{x^2y^2 - 2x^2y - 2xy^2 + 4x^2 - 2xy + 4y^2 + 2x - 2y + 1}{(xy - 1)(xy + 1)},$$
(3.10)

which is an invariant for the map φ_2 (1.2). By direct computation it is possible to prove that this invariant arises from the one obtained from the construction in [9] for the map (A.9) using the birational conjugation with the map (A.11b).

Associated to X_2 we have the Picard group:

$$\operatorname{Pic}(X_2) = \operatorname{Span}_{\mathbb{Z}}(H_x, H_y, F_1, \dots, F_8), \qquad (3.11)$$

On the Picard group of X_2 , the pullback of φ_2 act linearly as follows:

$$\varphi_{2}^{*}:\begin{pmatrix} H_{x}, H_{y}, \\ F_{1}, F_{2}, F_{3}, F_{4}, \\ F_{5}, F_{6}, F_{7}, F_{8} \end{pmatrix} \mapsto \begin{pmatrix} 2H_{x} + H_{y} - F_{2} - F_{4} - F_{7} - F_{8}, H_{x}, \\ H_{x} - F_{2}, F_{5}, H_{x} - F_{4}, F_{6}, \\ F_{3}, F_{1}, H_{x} - F_{8}, H_{x} - F_{7} \end{pmatrix}.$$
(3.12)

Furthermore, the representatives on the Picard group of the irreducible components of the singular fibres are shown in the third column of Table 2. Using the intersection form (2.5), we can find the associated surface type. This is given in the fourth and last column of Table 2. This implies that the singular fibres configuration is of type $(3A_2 \times A_1)^{(1)}$, type 61 in Oguiso and Shioda's list [41, Table 8.2]. This ends of the proof of the first statement of Theorem 1.2. The structure of the singular fibres is summarised in Figure 2.



FIGURE 2. A graphical representation of the singular fibres of p_2 .

From Table 2 we see that the full symmetry of a system on a surface of type $\Lambda = A_1^{(1)}$ is $\Lambda^{\perp} = E_7^{(1)}$. Then, in Section 5 we show that map φ_2 (1.2), when considered on a $A_1^{(1)}$ type surface admits $B_3^{(1)} \subset E_7^{(1)}$ type symmetry.

3.3. The map φ_s and its initial value space. In Remark 3.2 we noticed that the point configuration of that $A_1^{(1)}$ surface is different with respect to the standard one considered for instance in [28]. So, a natural question to ask is if we can recover a more general pencil with a similar structure, but no infinitely near points.

Consider a generic biquadratic polynomial and impose it to pass through the points c_i with i = 1, ..., 6. By direct computation, is shown that the obtained curve can be parametrised as:

$$p_{s} = \mu \left(x^{2}y^{2} - 2x^{2}y - 2xy^{2} + 4x^{2} - 2xy + 4y^{2} - 2x - 2y + 1 \right) + \nu \left[s \left(x^{2} + y^{2} + 1 \right) - (1 - s) \left(x^{2}y^{2} - 1 \right) \right].$$
(3.13)

In Eq. (3.13) where $[\mu : \nu] \in \mathbb{P}^1$ are the parameters of the pencil and s is an additional parameter. For $s \neq 0$ we have a pencil with eight distinct base points. For sake of simplicity we denote the base points by l_i and we list them as follows:

$$l_{1}(s):(x,y) = \left(\frac{(5-3\sqrt{3}i)s^{2}-32s+32}{26s^{2}-56s+32}, \frac{(5s^{2}+3i\sqrt{3})s^{2}-32s+32}{26s^{2}-56s+32}\right),$$

$$l_{2}(s):(x,y) = \left(\frac{(5s^{2}+3i\sqrt{3})s^{2}-32s+32}{26s^{2}-56s+32}, \frac{(5-3\sqrt{3}i)s^{2}-32s+32}{26s^{2}-56s+32}\right)$$

$$l_{3} = c_{5}, \ l_{4} = c_{6}, \ l_{5} = c_{2}, \ l_{6} = c_{4}, \ l_{7} = c_{3}, \ l_{8} = c_{1}.$$

$$(3.14)$$

When s = 0 the two base points $l_1(s)$ and $l_2(s)$ coalesce and we get the base points $c_{7,8}$. Note that, however the limit is singular.

We now define a map which is covariant on the pencil (3.13). Since the pencil p_s (3.13) is symmetric under the involution $\iota:(x,y) \mapsto (y,x)$, we can adopt the construction of the QRT root as explained in [18, §6.3]. Consider the rational function obtained by taking the coefficients of the biquadratic polynomial p_s (3.13) with respect to μ and ν :

$$\mathcal{H}_{s} = \frac{x^{2}y^{2} - 2x^{2}y - 2xy^{2} + 4x^{2} - 2xy + 4y^{2} - 2x - 2y + 1}{s(x^{2} + y^{2} + 1) - (1 - s)(x^{2}y^{2} - 1)}.$$
(3.15)

Considering the map:

$$v_s:(x,y)\mapsto(x,y'),\tag{3.16}$$

with y' non-trivial solution of:

$$\mathcal{H}_s(x,y') = \mathcal{H}_s(x,y). \tag{3.17}$$

By direct computation from (3.17) we obtain a trivial solution y' = y, while the non-trivial one is:

$$y' = \frac{3sx^2y + 2sx^2 - 3sxy - 4x^2y + sy + 6xy - 4y + 2}{2sx^2y - 3sx^2 - 2x^2y + 3sx + 2sy + 4x^2 - s - 6x + 4},$$
(3.18)

and the map v_s is an involution. The composition $\iota \circ v_s$ is then explicitly given by the maps φ_s of Equation (1.3). This is a map of infinite order, called a QRT root, is by construction covariant on p_s (3.13) and invariant on \mathcal{H}_s (3.15).

Now, by construction, the space of the initial conditions of the map φ_s (1.3), namely $X_s = B_{\varphi_s}$, is constructed by blowing up $\mathbb{P}^1 \times \mathbb{P}^1$ on the points l_i . Indeed, by construction

$$S_s = \operatorname{Sing} \varphi_s = (l_1, \dots, l_8). \tag{3.19}$$

Again, by construction X_s is a rational elliptic surface. We denote the exceptional lines corresponding to the blow up of the points l_i by L_i .

Associated to X_s we have the Picard group:

$$\operatorname{Pic}(B_{\varphi_s}) = \operatorname{Span}_{\mathbb{Z}}(H_x, H_y, L_1, \dots, L_8).$$
(3.20)

Then the action on $\operatorname{Pic}(B_{\varphi_s})$ is given by:

$$\varphi_s^* : \begin{pmatrix} H_x, H_y, \\ L_1, L_2, L_3, L_4, \\ L_5, L_6, L_7, L_8 \end{pmatrix} \mapsto \begin{pmatrix} 2H_x + H_y - L_1 - L_2 - L_5 - L_6, H_x, \\ H_x - L_2, H_x - L_1, L_7, L_8, \\ L_3, L_4, H_x - L_6, H_x - L_5 \end{pmatrix}.$$
(3.21)

Then, from Definition 2.5 we find that the pencil p_s (3.13) has five singular fibres corresponding to the following values of $[\mu : \nu] \in \mathbb{P}^1$:

$$\zeta_1 = [1:0], \tag{3.22a}$$

$$\zeta_2(s) = [-s/4, 1], \tag{3.22b}$$

$$\zeta_3(s) = [i(s-2)\sqrt{3} - 3s, 18], \qquad (3.22c)$$

$$\zeta_4(s) = [-i(s-2)\sqrt{3} - 3s, 18], \qquad (3.22d)$$

$$\zeta_5(s) = [(1-s)(s^2 - s + 1), s^3]. \tag{3.22e}$$

This information, along with the irreducible components of the associated singular fibres are shown in the first and second column of Table 3.

Remark 3.4. In the form of the singular values of the pencil p_s (3.13) we choose to underline that the point ζ_1 is independent from s, while $\zeta_i(s)$, for i = 2, 3, 4, 5 the dependence is explicit. The reason will more evident from the discussion we will make in Remark 3.5.

Furthermore, the representatives on the Picard group of the irreducible components of the singular fibres are shown in the third column of Table 3. Using the intersection form (2.5) we can find the associated surface type. This is given in the fourth and last column of Table 3. This implies that the singular fibre structure is $(2A_2 \times 2A_1 \times A_0^*)^{(1)}$, that is case 40 from [41, Table 8.2]². The fibre of type $A_0^{(1)*}$ has as representative the following irreducible curve in $\mathbb{P}^1 \times \mathbb{P}^1$ the nodal biquadric:

$$p_{5}^{*} = (s^{2} + 1) (s - 1)^{2} x^{2} y^{2} + 2 (s - 1) (s^{2} - s + 1) x^{2} y + (s^{2} - 2s + 2)^{2} x^{2} + 2 (s - 1) (s^{2} - s + 1) x y^{2} + 2 (s - 1) (s^{2} - s + 1) x y + 2 (s - 1) (s^{2} - s + 1) x + (s^{2} - 2s + 2)^{2} y^{2} + 2 (s - 1) (s^{2} - s + 1) y + 2s^{2} - 2s + 1,$$
(3.23)

having a node in the point:

$$n_s:(x,y) = \left(\frac{1}{1-s}, \frac{1}{1-s}\right).$$
(3.24)

The fibre structure is graphically represented in Figure 3. The ends the proof of the first statement of Theorem 1.3.

Remark 3.5. We remark that as $s \to 0$ we have that the singular fibre $\zeta_5(s)$ collide with the fibre $\zeta_1 = [0:1]$. That is, they originate the second fibre of the map φ_2 , listed in Table 2: the collision of a fibre of surface type $A_0^{(1)*}$ and $A_1^{(1)}$ generates a

²Note that in that classification the contribution from $A_0^{(1)*}$ fibre is not present since it does not contribute to the Mordell–Weil lattice.

fibre of surface type $A_3^{(1)}$. This is because both the irreducible components of the fibre in ζ_1 , see Table 3 pass through the point $(x, y) = (1, 1) = b_7$ which becomes a base point for the pencil as $s \to 0$. Note also that the nodal point n_s (3.24) of the curve p_5^* (3.23) tend to new base point c_7 . The fibre structure is then changed to $A_3^{(1)}$ because the proper transform of the irreducible components of the fibre in ζ_1 and the exceptional line on which $c_{7,8}$ lie all intersect on a single point, see Figure 2.



FIGURE 3. A graphical representation of the singular fibres of p_s .

4. Singular fibre of type $A_2^{(1)}$ associated with the map φ_1

In this section we compute the action of the map φ_1 (1.1) on the orthogonal complement of the singular fibre [0:1] of type $A_2^{(1)}$ of pencil p_1 (3.3). Then, we compute the normalizer of the Weyl group $W(4A_1)$ in the extended Weyl group $\widetilde{W}(E_6^{(1)})$. This will yield that the map φ_1 (1.1) on such a singular fibre has symmetry type $A_2^{(1)} \subset E_6^{(1)}$, thus ending the proof of Theorem 1.1.

4.1. Action on the simple roots of $E_6^{(1)}$ and decomposition in $\widetilde{W}(E_6^{(1)})$. Following Table 1, we have that the singular fiber of the pencil (3.3) at the point $[\mu:\nu] = [0:1]$ is of type $A_2^{(1)}$ and is generated by the following divisors in $\operatorname{Pic}(X_1)$: $D_0 = H_y - E_5 - E_6$, $D_1 = H_x - E_7 - E_8$, $D_2 = H_x + H_y - E_1 - E_2 - E_3 - E_4$. (4.1) The three divisors are permuted by the pullback map φ_1^* , that is $\varphi_1^*(D_i) = D_{i+1}$, $i \in \mathbb{Z}/3\mathbb{Z}$. The rational elliptic surface X_1 with this choice of the singular fibre is shown in Figure 4.

The orthogonal complement of $A_2^{(1)}$ type singular fiber in Pic (X_1) is generated by:

$$\alpha_0 = E_5 - E_6, \quad \alpha_1 = E_8 - E_7, \quad \alpha_2 = H_y - E_1 - E_8, \quad \alpha_3 = E_1 - E_2, \\ \alpha_4 = E_2 - E_3, \quad \alpha_5 = E_3 - E_4, \quad \alpha_6 = H_x - E_1 - E_5.$$
(4.2)

The set $\Delta^{(1)} = \Delta(E_6^{(1)}) = \{\alpha_j \mid 0 \le j \le 6\}$ is a simple system of type $E_6^{(1)}$, its Dynkin diagram is given in Figure 5. The anti-canonical divisor corresponds to the null root of the $E_6^{(1)}$ system:

$$-K_{X_1} = \delta = \alpha_{015224466333}. \tag{4.3}$$

Let $V^{(1)}$ be a vector space with basis $\Delta^{(1)}$, equipped with a symmetric bilinear form given by Equations (2.15) and (B.1). The corresponding extended affine Weyl group is,

$$\widetilde{W}(E_6^{(1)}) = \langle s_i \mid 0 \le i \le 6 \rangle \rtimes \langle \sigma \rangle = W(E_6) \ltimes P = W(E_6) \ltimes \langle u_j \mid 1 \le j \le 6 \rangle, \quad (4.4)$$

where u_j is the translation by the fundamental weight h_j $(1 \le j \le 6)$ of the $E_6^{(1)}$ system. The action of $s_j \in W(E_6^{(1)})$ on $\Delta^{(1)}$ is given by Equations (2.16) and (B.1). See Appendix B for basic data and properties of $\widetilde{W}(E_6^{(1)})$.



FIGURE 4. The space of initial values of (1.1) interpreted as elliptic surface of type $A_2^{(1)}$.

Using Equations (3.6) and (4.2) we have φ_1^* acts on $\Delta^{(1)}$ by:

$$\varphi_1^*: (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6) \mapsto (-\alpha_1, -\alpha_5, \alpha_{3456}, \alpha_0, -\alpha_{036}, \alpha_3, \alpha_{1236}). \tag{4.5}$$

Computing four iterations of the map φ_1^* we obtain:

$$\left(\varphi_1^*\right)^4 : \left(\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6\right) \mapsto \left(\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4 - \delta, \alpha_5, \alpha_6 + \delta\right).$$
(4.6)

By Equation (2.38) we recognise that $(\varphi_1^*)^4 \in P$ is the translation by $h_4 - h_6$, to which φ_1^* is a quasi-translation of order four in $\widetilde{W}(E_6^{(1)})$. It can be checked that



FIGURE 5. $\Gamma(E_6^{(1)})$ with diagram automorphism $\sigma = (150)(246)$.

 $|h_4 - h_6|^2 = \frac{4}{3}$ using Equation (B.4). Finally, φ_1^* has the following decomposition in terms of the generators of $\widetilde{W}(E_6^{(1)})$ using Equations (2.33) and (2.40):

$$\varphi_1^* = \sigma s_{453401}. \tag{4.7}$$

4.2. The normalizer of $W(4A_1)$ in $\widetilde{W}(E_6^{(1)})$. The initial observation on φ_1^* comes from its actions on $\Delta^{(1)}$ given in Equation (4.5), that is φ_1^* acts on the subset $\{\alpha_0, \alpha_1, \alpha_3, \alpha_5\} \subset \Delta^{(1)}$ as a permutation of order four modulo the minus signs. This prompts us to take

$$J = \{\alpha_0, \alpha_1, \alpha_3, \alpha_5\} \cong 4A_1, \tag{4.8}$$

and consider the normalizer of $W_J = \langle s_0, s_1, s_3, s_5 \rangle$ in $\widetilde{W}(E_6^{(1)})$.

Proposition 4.1. The normalizer of W_J with $J = \{\alpha_0, \alpha_1, \alpha_3, \alpha_5\}$ in $\widetilde{W}(E_6^{(1)})$ is given by

$$N(W_J) = N_J \ltimes W_J = (\langle b_1, b_2, b_0 \rangle \rtimes \langle \sigma \rangle) \ltimes \langle s_0, s_1, s_3, s_5 \rangle \cong \widetilde{W}(A_2^{(1)}) \ltimes W(4A_1),$$
(4.9)

with

$$b_1 = s_{4534}, \quad b_2 = s_{6036}, \quad b_0 = s_{2312}.$$
 (4.10)

Proof. Recall that $N(W_J) = N_J \ltimes W_J$, where N_J generated by the R- and Melements, is the group of all elements of $\widetilde{W}(E_6^{(1)})$ that act permutatively on the set J. Here we have $J = \{\alpha_0, \alpha_1, \alpha_3, \alpha_5\} \cong 4A_1$, and $W_J = \langle s_0, s_1, s_3, s_5 \rangle \cong W(4A_1)$. Following the procedure of computing normalizers given in [5, 21], we find that the R-elements (that is, quasi-reflections) are given by,

$$b_1 = s_{4534}, \quad b_2 = s_{6036}, \quad b_0 = s_{2312},$$
 (4.11)

where b_1, b_2, b_0 act on the subset J as permutations of order two, written as a permutation on the index set of J we have,

$$b_1 = (35), \quad b_2 = (03), \quad b_0 = (13).$$
 (4.12)

The M-element is given by σ (action on $\Delta^{(1)}$ in Equation (B.6)), that is its action on J is,

$$\sigma = (150). \tag{4.13}$$

It can be verified using the defining relations of $\widetilde{W}(E_6^{(1)})$ that b_i 's are involutions that satisfy the defining relations (4.14a) for an affine Weyl group of type $A_2^{(1)}$, while σ acts on the b_i 's by Equation (4.14b).

$$b_j^2 = 1, \ (b_j b_{j+1})^3 = 1, \ (j \in \mathbb{Z}/3\mathbb{Z}),$$
 (4.14a)

$$\sigma^3 = 1, \ \sigma b_j = b_{j+1}\sigma. \tag{4.14b}$$

That is, together the R- and M-elements generate N_J , an extended affine Weyl group of type $A_2^{(1)}$,

$$N_J = \langle b_1, b_2, b_0 \rangle \rtimes \langle \sigma \rangle \cong \widetilde{W}(A_2^{(1)}). \tag{4.15}$$

Consider the expression for φ_1^* in Equation (4.7), we see that $\varphi_1^* = \sigma s_{4534} s_0 s_1 = \sigma b_1 s_0 s_1$ is certainly an element of the subgroup $N(W_J)$ of $\widetilde{W}\left(E_6^{(1)}\right)$ given by Equation (4.9).

In what follows we further clarify the nature of φ_1^* , showing that there is a conjugation that takes φ_1^* to an element of quasi-translation in $N(W_J)$.

4.2.1. Sub-root system and translations in $N_J \cong \widetilde{W}(A_2^{(1)})$. First we find a subspace $V_J^{\perp} \subset V^{(1)}$ on which $b_i \in N_J$ $(0 \le i \le 2)$ can be realised as simple reflections of $A_2^{(1)}$ type. We then construct an element of translation in $\widetilde{W}(A_2^{(1)})$ and find its relation to the φ_1^* given in Equation (4.7).

Proposition 4.2. The root system for N_J given in Equation (4.15) is generated by $\beta = \{\beta_i \mid 0 \le i \le 2\}$ with

$$\beta_0 = \alpha_2 + \frac{\alpha_{13}}{2}, \quad \beta_1 = \alpha_4 + \frac{\alpha_{35}}{2}, \quad \beta_2 = \alpha_6 + \frac{\alpha_{03}}{2}, \quad (4.16)$$

is of $A_2^{(1)}$ type, and $V_J^{\perp} = \text{Span}(\beta)$. The group $N_J = \langle b_1, b_2, b_0 \rangle \rtimes \langle \sigma \rangle$ can be realised as $\widetilde{W}(A_2^{(1)})$ on V_J^{\perp} with b_i acting as the reflection along the root β_i , and σ is the diagram automorphism of $\Gamma(A_2^{(1)})$ for the β -system, see Figure 6.

Proof. Let $\beta = \{\beta_i \mid 0 \le i \le 2\}$ with the β_i 's be as given in Equation (4.16). The bilinear form on $V^{(1)}$ in the $\Delta^{(1)}$ basis given by Equations (2.15) and (B.1) is used to check that the β -system is or dimension three and orthogonal to V_J . That is, $V_J^{\perp} = \text{Span}(\beta)$. Moreover, we found that $|\beta_i|^2 = 2$, so $\beta_j^{\vee} = \beta_j$ for $(0 \le i \le 2)$.

Computing the values of $(\beta_i \cdot \beta_j^{\vee})$ for the β -system (4.16) again using the bilinear form on $V^{(1)}$ we have,

$$C(A_2^{(1)}) = (\beta_i \cdot \beta_j^{\vee})_{1 \le i, j \le 2, 0} = (\beta_i \cdot \beta_j)_{1 \le i, j \le 2, 0} = \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}.$$
 (4.17)

The matrix in Equation (4.17) is the Cartan matrix of $A_2^{(1)}$ type. That is, β -system is of $A_2^{(1)}$ type, its Dynkin diagram given in Figure 6. The null root of this $A_2^{(1)}$ system is given by

$$\delta_{A_2} = \beta_0 + \beta_1 + \beta_2 = \frac{1}{2} \left(\alpha_0 + \alpha_1 + \alpha_5 + 2 \left(\alpha_2 + \alpha_4 + \alpha_6 \right) + 3\alpha_3 \right) = \frac{\delta}{2}, \quad (4.18)$$

where δ (Equation (4.3)) is the null root of the $E_6^{(1)}$ root system. Next, we look at the actions of b_i ($0 \le i \le 2$) on the $V_J \bigoplus V_J^{\perp}$ basis of $V^{(1)}$, that is $J \cup \beta =$

 $\{\alpha_0, \alpha_1, \alpha_3, \alpha_5, \beta_1, \beta_2, \beta_0, \}$. These are computed using Equation (4.10) by composing the actions of $s_j \in W(E_6^{(1)})$ on $V^{(1)}$ given by Equations (2.16) and (B.1). We have,

$$b_1:(\alpha_0,\alpha_1,\alpha_3,\alpha_5,\beta_1,\beta_2,\beta_0) \mapsto (\alpha_0,\alpha_1,\alpha_5,\alpha_3,-\beta_1,\beta_2+\beta_1,\beta_0+\beta_1), \tag{4.19a}$$

$$b_2:(\alpha_0,\alpha_1,\alpha_3,\alpha_5,\beta_1,\beta_2,\beta_0) \mapsto (\alpha_0,\alpha_1,\alpha_3,\alpha_5,\beta_1+\beta_2,-\beta_2,\beta_0+\beta_2), \tag{4.19b}$$

$$b_0:(\alpha_0,\alpha_1,\alpha_3,\alpha_5,\beta_1+\beta_2,\beta_2,\beta_0)\mapsto(\alpha_0,\alpha_1,\alpha_3,\alpha_5,\beta_1+\beta_0,\beta_2+\beta_0,-\beta_0). \quad (4.19c)$$

From Equation (4.19), we see that b_i $(0 \le i \le 2)$ acts on the β -system exactly as the reflection along the root β_i of $A_2^{(1)}$ type, while acting permutatively on J. The action of σ on the β -system is computed using Equations (4.16) and (B.6), and we have

$$\sigma:(\beta_0,\beta_1,\beta_2)\mapsto(\beta_1,\beta_2,\beta_0),\tag{4.20}$$

that is, σ is an order three diagram automorphism of the type $A_2^{(1)} \beta$ -system. Hence we see that $N_J = \langle b_1, b_2, b_0 \rangle \rtimes \langle \sigma \rangle$ can be realised as $\widetilde{W}(A_2^{(1)})$ on the subspace $V_J^{\perp} = \text{Span}(\beta)$.



FIGURE 6. $\Gamma(A_2^{(1)})$ with diagram automorphism σ .

Recall that by Equation (2.35), for an extended affine Weyl group we have the following decomposition $\widetilde{W}^{(1)} = W \ltimes P$. That is,

$$N_J = \langle b_1, b_2, b_0 \rangle \rtimes \langle \sigma \rangle \cong \widetilde{W}\left(A_2^{(1)}\right) = W\left(A_2\right) \ltimes P_{A_2} = \langle b_1, b_2 \rangle \ltimes \langle U_1, U_2 \rangle, \quad (4.21)$$

where U_i is the translation by the fundamental weight H_i (i = 1, 2) of this $A_2^{(1)}$ type β -system. In particular we have [43]:

$$U_1 = \sigma b_2 b_1, \quad U_2 = \sigma^{-1} b_1 b_2. \tag{4.22}$$

 H_i (i = 1, 2) are related to the finite simple roots of the β -system by Equations (2.29) and (2.30),

$$\begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} = C(A_2)^T \begin{pmatrix} H_1 \\ H_2 \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \begin{pmatrix} H_1 \\ H_2 \end{pmatrix},$$
(4.23)

and

$$\begin{pmatrix} H_1 \\ H_2 \end{pmatrix} = \left(C(A_2)^T \right)^{-1} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}.$$
 (4.24)

In particular we have,

$$H_2 = \frac{1}{3}(\beta_1 + 2\beta_2) = \frac{1}{3}\left(\alpha_4 + \frac{\alpha_{35}}{2} + 2\left(\alpha_6 + \frac{\alpha_{03}}{2}\right)\right) = \frac{1}{2}(h_6 - h_2), \quad (4.25)$$

where we have used Equation (B.3) for the last expression in terms of fundamental weights h_i ($1 \le i \le 6$) of $E_6^{(1)}$. So we have,

$$2H_2 = h_6 - h_2. \tag{4.26}$$

Moreover, $|2H_2|^2 = |h_6 - h_2|^2 = \frac{4}{3}$. Elements U_2 and $(U_2)^2$ act on the simple system of $E_6^{(1)}$ by

$$U_{2}:(\alpha_{0},\alpha_{1},\alpha_{2},\alpha_{3},\alpha_{4},\alpha_{5},\alpha_{6}) \mapsto (\alpha_{1},\alpha_{0},\alpha_{1223346},\alpha_{5},\alpha_{4},\alpha_{3},-\alpha_{12345}), \quad (4.27)$$

and

$$(U_2)^2: (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6) \mapsto (\alpha_0, \alpha_1, \alpha_2 + \delta, \alpha_3, \alpha_4, \alpha_5, \alpha_6 - \delta).$$
(4.28)

By Equation (2.38) we recognise that $U_2^2 \in P$ is a translation by $h_6 - h_2$ in $\widetilde{W}(E_6^{(1)})$. The nature of element U_2 becomes more evident in the $J \cup \beta$ basis of $V^{(1)}$,

$$U_{2}: \begin{pmatrix} \alpha_{0}, \alpha_{1}, \alpha_{3}, \alpha_{5}, \\ \beta_{1}, \beta_{2}, \beta_{0} \end{pmatrix} \mapsto \begin{pmatrix} \alpha_{1}, \alpha_{0}, \alpha_{5}, \alpha_{3}, \\ \beta_{1}, \beta_{2} - \delta_{A_{2}}, \beta_{0} + \delta_{A_{2}} \end{pmatrix}.$$
(4.29)

Again by Equation (2.38) we see that U_2 acts on the β -system as a translation by H_2 . However, it also acts on the set J as a permutation of order two (01)(35), hence can not be an element of translation in $\widetilde{W}(E_6^{(1)})$. On the other hand, we have

$$(U_2)^2 : \begin{pmatrix} \alpha_0, \alpha_1, \alpha_3, \alpha_5, \\ \beta_1, \beta_2, \beta_0 \end{pmatrix} \mapsto \begin{pmatrix} \alpha_0, \alpha_1, \alpha_3, \alpha_5, \\ \beta_1, \beta_2 - 2\delta_{A_2}, \beta_0 + 2\delta_{A_2} \end{pmatrix}.$$
 (4.30)

That is, U_2^2 acts on the β -system as a translation by $2H_2 = h_6 - h_2$ but also fixing J. Hence, U_2^2 is a translation by $2H_2 = h_6 - h_2$ in $\widetilde{W}\left(E_6^{(1)}\right)$ as we knew from Equation (4.28).

Let us denote by u_{ij} the element of translation by $h_i - h_j$ in $\widetilde{W}\left(E_6^{(1)}\right)$. Then we have $u_{ij} = u_i u_j^{-1}$ and $(u_{ij})^{-1} = u_{ji}$. By Equation (4.6) we have $(\varphi_1^*)^4 = u_{46}$, that is φ_1^* is a quasi-translation of order four for u_{46} , and by Equation (4.30) U_2 is a quasi-translation of order two for u_{62} .

A quasi-translation of order four for u_{62} can be constructed simply by a rewriting of U_2 using the defining relations (4.14b) of $\widetilde{W}(A_2^{(1)})$,

$$U_2 = \sigma^{-1}b_1b_2 = \sigma^2 b_1b_2 = \sigma b_2\sigma b_2 = (\sigma b_2)^2.$$
(4.31)

Let

$$\rho = \sigma b_2 = \sigma s_{6036}, \quad \text{and we have} \quad \rho^2 = U_2, \quad \rho^4 = u_{62}.$$
(4.32)

A conjugation that relates the two order four quasi-translations $\varphi_1^* = \sigma b_1 s_0 s_1$ and $\rho = \sigma b_2$ is found by the following observations.

Proposition 4.3. $u_{62} = \rho^4$ is related to $u_{46} = (\varphi_1^*)^4$ by a conjugation of $b_2 \in W(A_2)$,

$$b_2 u_{62} b_2^{-1} = u_{46}. ag{4.33}$$

Proof. First, we knew that vectors $h_6 - h_2$ and $h_4 - h_6$ are of the same length (that is, squared length of $\frac{4}{3}$). In fact, they belong to the same orbit under the actions of $W(A_2)$,

$$b_2(H_2) = b_2\left(\frac{h_6 - h_2}{2}\right) = H_2 - (-H_1 + 2H_2) = H_1 - H_2$$

= $\frac{1}{3}(\beta_1 - \beta_2) = \frac{1}{2}(h_4 - h_6),$ (4.34)

where we have used Equation (2.26a) and the Cartan matrix (4.17) to compute the action of b_2 on H_2 . Then by Equation (2.39) we have Equation (4.33).

Proposition 4.4. Any conjugation of $b_2\rho b_2^{-1} = \sigma b_1$ by an element of $W_J = \langle s_0, s_1, s_3, s_5 \rangle$ is a quasi-translation of order four for u_{46} . In particular we found,

$$s_1 b_2 \rho(s_1 b_2)^{-1} = \varphi_1^*. \tag{4.35}$$

That is, φ_1^* (given in Equation (4.7)) is related to the order four quasi-translation ρ via a conjugation of s_1b_2 .

Proof. By Proposition 4.3 and Equation (4.32),

$$u_{46} = b_2 u_{62} b_2^{-1} = b_2 \rho^4 b_2^{-1} = \left(b_2 \rho b_2^{-1}\right)^4, \tag{4.36}$$

where we have,

$$b_2 \rho b_2^{-1} = b_2 \sigma b_2 b_2^{-1} = b_2 \sigma = \sigma b_1.$$
(4.37)

That is,

$$\left(b_2\rho b_2^{-1}\right)^4 = (\sigma b_1)^4 = u_{46}.$$
(4.38)

Recall from Proposition 4.1 that

$$N(W_J) = N_J \ltimes W_J = (\langle b_1, b_2, b_0 \rangle \rtimes \langle \sigma \rangle) \ltimes \langle s_0, s_1, s_3, s_5 \rangle \cong \widetilde{W}(A_2^{(1)}) \ltimes W(4A_1).$$
(4.39)

The root system of $N_J \beta$, is orthogonal to the root system of $W_J J$. This means that any conjugation of $r \in N_J$ by $s \in W_J$, srs^{-1} does not change r's action on the β -system. It only results an even number of sign changes (an action of order at most four) in the simple roots of J observed earlier in Section 4.2, this difference in action is not seen after four iterations. Equation (4.35) is a consequence of this observation and can be verified by direct computation:

$$s_1 b_2 \rho(s_1 b_2)^{-1} = s_1 b_2 \rho b_2^{-1} s_1^{-1} = s_1 \sigma b_1 s_1^{-1} = s_1 \sigma b_1 s_1 = \sigma b_1 s_0 s_1 = \varphi_1^*, \qquad (4.40)$$

where we have used the defining relations given in Equation (4.14a).

Hence we have shown that the map φ_1^* , as given by Equation (4.7), is an quasitranslation in a normalizer of $W(4A_1)$ in $\widetilde{W}(E_6^{(1)})$ (given by Equation (4.9)), and we completed the proof of Theorem 1.1.

5. Singular fibre $\zeta_2(s)$ of type $A_1^{(1)}$ associated with the map φ_s

In this section we compute the action of the map φ_s (1.3) on the orthogonal complement of the singular fibre [0:1] of type $A_1^{(1)}$ of pencil p_s (3.13). Then, we compute the normalizer of the Weyl group $W(4A_1)$ in the extended Weyl group $\widetilde{W}(E_7^{(1)})$. This will yield that the map φ_s (1.3) on such a singular fibre has symmetry type $B_3^{(1)} \subset E_7^{(1)}$, a non-simply laced Coxeter group, thus ending the proof of Theorem 1.3.

5.1. Action on the simple roots of $E_7^{(1)}$ and decomposition in $\widetilde{W}(E_7^{(1)})$. Following Table 2 we have that the singular fibre of the pencil (3.9) at the point $[\mu:\nu] = [0:1]$ is of type $A_1^{(1)}$ and it is generated by the following divisors in $\operatorname{Pic}(X_s)$:

$$D_0 = H_x + H_y - L_1 - L_2 - L_3 - L_4, \quad D_1 = H_x + H_y - L_5 - L_6 - L_7 - L_8.$$
(5.1)

These two divisors are permuted by the pullback map φ_s^* , that is $\varphi_s^*(D_i) = D_{(i+1)}$, $i \in \mathbb{Z}/2\mathbb{Z}$. The rational elliptic surface X_s with this choice of the singular fibre is shown in Figure 7. The orthogonal complement of $A_1^{(1)}$ in Pic (X_s) is generated by:

$$\alpha_0 = L_7 - L_8, \quad \alpha_1 = L_6 - L_7, \quad \alpha_2 = H_x - H_y, \quad \alpha_3 = L_5 - L_6, \\ \alpha_4 = H_y - L_1 - L_5, \quad \alpha_5 = L_1 - L_2, \quad \alpha_6 = L_2 - L_3, \quad \alpha_7 = L_3 - L_4.$$
(5.2)

The set $\Delta^{(1)} = \Delta(E_7^{(1)}) = \{\alpha_j \mid 0 \le j \le 7\}$ is a simple system of type $E_7^{(1)}$, its Dynkin diagram given in Figure 8. The anti-canonical divisor corresponds to the following null-root in the $E_7^{(1)}$ root system:

$$-K_{X_2} = \delta = \alpha_{071122663335554444}.$$
(5.3)

Let $V^{(1)}$ be a vector space with basis $\Delta^{(1)}$, equipped with a symmetric bilinear form given by Equations (2.15) and (C.1). The corresponding extended affine Weyl group is,

$$\widetilde{W}(E_7^{(1)}) = \langle s_i \mid 0 \le i \le 7 \rangle \rtimes \langle \sigma \rangle = W(E_7) \ltimes P = W(E_7) \ltimes \langle u_j \mid 1 \le j \le 7 \rangle, \quad (5.4)$$

where u_j is the translation by the fundamental weight of the $E_7^{(1)}$ system, h_j $(1 \le j \le 7)$. In particular, the action of $s_j \in W(E_7^{(1)})$ on $\Delta^{(1)}$ is given by Equations (2.16) and (C.1). See Appendix C for basic data and properties of $\widetilde{W}(E_7^{(1)})$.



FIGURE 7. The space of initial values of (1.3) interpreted as elliptic surface of type $A_1^{(1)}$.

Using Equations (3.12) and (5.2) we have φ_s^* acts on $\Delta^{(1)}$ by:

$$\varphi_s^*: \begin{pmatrix} \alpha_0, \alpha_1, \alpha_2, \alpha_3, \\ \alpha_4, \alpha_5, \alpha_6, \alpha_7 \end{pmatrix} \mapsto \begin{pmatrix} \alpha_3, -\alpha_{234567}, \alpha_{23445}, \alpha_7, \\ \alpha_6, \alpha_5, \alpha_{1234}, \alpha_0 \end{pmatrix}.$$
(5.5)

Compute three iterations of the map φ_s^* we obtain:

$$(\varphi_s^*)^3 : \begin{pmatrix} \alpha_0, \alpha_1, \alpha_2, \alpha_3, \\ \alpha_4, \alpha_5, \alpha_6, \alpha_7 \end{pmatrix} \mapsto \begin{pmatrix} \alpha_0, \alpha_1 - \delta, \alpha_2 + \delta, \alpha_3, \\ \alpha_4, \alpha_5, \alpha_6, \alpha_7 \end{pmatrix}.$$
 (5.6)

By Equation (2.38) we recognise that $(\varphi_s^*)^3 \in P$ is the translation by $h_1 - h_2$, and φ_s^* , a quasi-translation of order three in $\widetilde{W}(E_7^{(1)})$. Moreover, using Equation (C.4)



FIGURE 8. The Dynkin diagram of the Weyl group $\widetilde{W}(E_7^{(1)})$ in Bourbaki's numbering.

one can check that $|h_1 - h_2|^2 = \frac{3}{2}$, hence it is in the same $W(E_7)$ -orbit of h_7 . Finally, φ_s^* has the following decomposition in terms of the generators of $\widetilde{W}(E_7^{(1)})$ using Equations (2.33) and (2.40):

$$\varphi_s^* = \sigma s_2 s_4 s_5 s_3 s_4 s_1 s_3 s_0 s_1. \tag{5.7}$$

5.2. Normalizer of $W(4A_1)$ in $\widetilde{W}(E_7^{(1)})$. The initial observation on φ_s^* comes from its actions on $\Delta^{(1)}$ given in Equation (5.5), that is, φ_s^* permutes the subset $\{\alpha_0, \alpha_3, \alpha_5, \alpha_7\} \subset \Delta^{(1)}$. This prompts us to take

$$J = \{\alpha_0, \alpha_3, \alpha_5, \alpha_7\} \cong 4A_1, \tag{5.8}$$

and look at the normalizer of $W_J = \langle s_0, s_3, s_5, s_7 \rangle$ in $\widetilde{W}(E_7^{(1)})$.

Proposition 5.1. The normalizer of W_J with $J = \{\alpha_0, \alpha_3, \alpha_5, \alpha_7\}$ in $\widetilde{W}(E_7^{(1)})$ is given by $N(W_J) = N_J \ltimes W_J = (\langle b_1, b_2, b_3, b_0 \rangle \rtimes \langle \sigma \rangle) \ltimes \langle s_0, s_3, s_5, s_7 \rangle$

$$V(W_J) = N_J \ltimes W_J = (\langle b_1, b_2, b_3, b_0 \rangle \rtimes \langle \sigma \rangle) \ltimes \langle s_0, s_3, s_5, s_7 \rangle$$

$$\cong \widetilde{W} \left(B_3^{(1)} \right) \ltimes W(4A_1),$$
(5.9)

with

$$b_1 = s_{6576}, \quad b_2 = s_{4534}, \quad b_3 = s_2, \quad b_0 = s_{1301}.$$
 (5.10)

Proof. Recall that $N(W_J) = N_J \ltimes W_J$, where N_J generated by the R- and Melements, is the group of all elements of $\widetilde{W}(E_7^{(1)})$ that act permutatively on the subset $J \subset \Delta^{(1)}$. Here we have $J = \{\alpha_0, \alpha_3, \alpha_5, \alpha_7\} \cong 4A_1$, and $W_J = \langle s_0, s_3, s_5, s_7 \rangle \cong$ $W(4A_1)$. Following the procedure of computing normalizers given in [5, 21], we find that the R-elements are given by,

$$b_1 = s_{6576}, \quad b_2 = s_{4534}, \quad b_3 = s_2, \quad b_0 = s_{1301}.$$
 (5.11)

where b_1, b_2, b_0 act on the subset J as permutations of order two, written as a permutation on the index set of J we have,

$$b_1 = (57), \quad b_2 = (35), \quad b_0 = (03).$$
 (5.12)

The M-element is given by σ (its actions on $\Delta^{(1)}$ are given in Equation (C.6)), its action on J is,

$$\sigma = (35)(07). \tag{5.13}$$

It can be verified using the defining relations of $\widetilde{W}(E_7^{(1)})$ that b_i 's are involutions that satisfy the defining relations (5.14a) for an affine Weyl group of type $B_3^{(1)}$, while σ is the diagram automorphism of $\Gamma(B_3^{(1)})$ (see Figure 9), and it acts on the b_i 's by Equation (5.14b),

$$b_j^2 = 1, \ (j \in \{0, 1, 2, 3\}), \ (b_1 b_2)^3 = (b_1 b_3)^2 = (b_2 b_3)^4 = 1,$$
 (5.14a)
 $(b_0 b_2)^3 = (b_0 b_3)^2 = (b_0 b_1)^2 = 1$

$$\sigma^2 = 1, \ \sigma b_0 = b_1 \sigma. \tag{5.14b}$$

That is, together the R- and M-elements generate N_J , an extended affine Weyl group of type $B_3^{(1)}$,

$$N_J = \langle b_1, b_2, b_3, b_0 \rangle \rtimes \langle \sigma \rangle \cong \widetilde{W}(B_3^{(1)}).$$
(5.15)

This ends the proof.



FIGURE 9. Dynkin diagram of affine B_3 type with the diagram automorphism, $\widetilde{\Gamma}(B_3^{(1)})$.

5.2.1. Sub-root system and translations in $N_J \cong \widetilde{W}(B_3^{(1)})$. First we find a subspace $V_J^{\perp} \subset V^{(1)}$ on which $b_i \in N_J$ $(0 \le i \le 3)$ can be realised as simple reflections of $B_3^{(1)}$ type. We construct an element of translation in $\widetilde{W}(B_3^{(1)})$ and find its relation to the element φ_s^* given in Equation (5.7).

Proposition 5.2. The root system for N_J given in Equation (5.15) generated by $\beta = \{\beta_i \mid 0 \le i \le 3\}$ with

$$\beta_0 = \alpha_{0113}, \quad \beta_1 = \alpha_{5667}, \quad \beta_2 = \alpha_{3445}, \quad \beta_3 = \alpha_2, \tag{5.16}$$

is of type $B_3^{(1)}$, and we have $V_J^{\perp} = \text{Span}(\beta)$. The group $N_J = \langle b_1, b_2, b_3, b_0 \rangle \rtimes \langle \sigma \rangle$ can be realised as $\widetilde{W}(B_3^{(1)})$ on V_J^{\perp} with b_i acting as the reflection along the root β_i , while σ exchanges β_0 and β_1 , which is the diagram automorphism of $\Gamma(B_3^{(1)})$ for the β -system (see Figure 9).

Proof. Let $\beta = \{\beta_i \mid 0 \le i \le 3\}$ be as given in Equation (5.16). The bilinear form on $V^{(1)}$ in $\Delta^{(1)}$ basis given by Equations (2.15) and (C.1) is used to check that the β -system is of dimension four and orthogonal to V_J . That is, $V_J^{\perp} = \text{Span}(\beta)$. Moreover, we found that $|\beta_i|^2 = 4$, so $\beta_j^{\vee} = \frac{\beta_j}{2}$ (for $0 \le i \le 2$), and $|\beta_3|^2 = 2$ so $\beta_3^{\vee} = \beta_3$.

Computing the values of $\beta_i \cdot \beta_j^{\vee}$ for the β -system (5.16) using the bilinear form on $V^{(1)}$ we have,

$$C(B_3^{(1)}) = (\beta_i \cdot \beta_j^{\vee})_{1 \le i, j \le 3, 0} = \begin{pmatrix} 2 & -1 & 0 & -1 \\ -1 & 2 & -2 & 0 \\ 0 & -1 & 2 & 0 \\ -1 & 0 & 0 & 2 \end{pmatrix}.$$
 (5.17)

The matrix in Equation (5.17) is the Cartan matrix of type $B_3^{(1)}$. That is, β -system is of $B_3^{(1)}$ type, its Dynkin diagram given in Figure 9. The null root of this $B_3^{(1)}$ system given by

$$\delta_{B_3} = \beta_0 + \beta_1 + 2\beta_2 + 2\beta_3 = \alpha_{071122663335554444} = \delta, \tag{5.18}$$

where δ (Equation (5.3)) is the null root of the $E_7^{(1)}$ root system. The action of σ on the β -system is computed using Equations (5.16) and (C.6), and we have

$$\sigma: (\beta_0, \beta_1, \beta_2, \beta_3) \mapsto (\beta_1, \beta_0, \beta_2, \beta_3), \tag{5.19}$$

that is, σ is an order two diagram automorphism of this type $B_3^{(1)} \beta$ -system. Hence we see that $N_J = \langle b_1, b_2, b_3, b_0 \rangle \rtimes \langle \sigma \rangle$ can be realised as $\widetilde{W}(B_3^{(1)})$ on the subspace $V_J^{\perp} = \text{Span}(\beta)$.

Recall that by Equation (2.35), for an extended affine Weyl group we have the following decomposition $\widetilde{W}^{(1)} = W \ltimes P$. That is,

$$N_J = \langle b_1, b_2, b_3, b_0 \rangle \rtimes \langle \sigma \rangle \cong \widetilde{W} \left(B_3^{(1)} \right) = W \left(B_3 \right) \bowtie P_{B_3} = \langle b_1, b_2, b_3 \rangle \bowtie \langle U_1, U_2, U_3 \rangle,$$

$$(5.20)$$

where U_i is the translation by the fundamental weight H_i (i = 1, 2, 3) of the type $B_3^{(1)} \beta$ -system. In particular we have [43]:

$$U_1 = \sigma b_{12321}, \quad U_2 = b_{02321232}, \quad U_3 = \sigma b_{123023123}. \tag{5.21}$$

 H_i (i = 1, 2, 3) are related to the finite simple roots of the β -system by Equations (2.29) and (2.30),

$$\begin{pmatrix} \pi(\beta_1^{\vee}) \\ \pi(\beta_2^{\vee}) \\ \pi(\beta_3^{\vee}) \end{pmatrix} = C(B_3)^T \begin{pmatrix} H_1 \\ H_2 \\ H_3 \end{pmatrix} = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -2 & 2 \end{pmatrix} \begin{pmatrix} H_1 \\ H_2 \\ H_3 \end{pmatrix},$$
(5.22)

and

$$\begin{pmatrix} H_1 \\ H_2 \\ H_3 \end{pmatrix} = \left(C(B_3)^T \right)^{-1} \begin{pmatrix} \pi(\beta_1^{\vee}) \\ \pi(\beta_2^{\vee}) \\ \pi(\beta_3^{\vee}) \end{pmatrix} = \begin{pmatrix} 1 & 1 & \frac{1}{2} \\ 1 & 2 & 1 \\ 1 & 2 & \frac{3}{2} \end{pmatrix} \begin{pmatrix} \pi(\beta_1^{\vee}) \\ \pi(\beta_2^{\vee}) \\ \pi(\beta_3^{\vee}) \end{pmatrix}.$$
(5.23)

Recall that for non-simply-laced systems one can identify $\pi(\beta_i^{\vee})$ with $\frac{2\beta_i}{|\beta_i|^2}$, so we can express coroots of the $B_3^{(1)}$ system in terms of the $E_7^{(1)}$ simple roots,

$$\pi(\beta_0^{\vee}) = \frac{\alpha_{03}}{2} + \alpha_1, \quad \pi(\beta_1^{\vee}) = \frac{\alpha_{57}}{2} + \alpha_6, \quad \pi(\beta_2^{\vee}) = \frac{\alpha_{35}}{2} + \alpha_4, \quad \pi(\beta_3^{\vee}) = \alpha_2.$$
(5.24)

In particular, we have

$$H_{3} = \pi(\beta_{1}^{\vee}) + 2\pi(\beta_{2}^{\vee}) + \frac{3}{2}\pi(\beta_{3}^{\vee}),$$

$$= \frac{3}{2}\alpha_{2} + \alpha_{3} + 2\alpha_{4} + \frac{3}{2}\alpha_{5} + \alpha_{6} + \frac{1}{2}\alpha_{7},$$

$$= -h_{1} + h_{2}.$$
 (5.25)

where we have used Equation (C.3) for the last expression in terms of fundamental weights $h_i (1 \le i \le 7)$ of $E_7^{(1)}$. That is, $U_3 \in P$ defined in Equation (5.21) is a translation by $-h_1 + h_2$ in $\widetilde{W}(E_7^{(1)})$.

Proposition 5.3. The map φ_s^* given in Equation (5.7) is an element of quasitranslation of order three in the $N_J \cong \widetilde{W}(B_3^{(1)})$ subgroup of $\widetilde{W}(E_7^{(1)})$ given in Proposition 5.1. We have,

$$\varphi_s^* = \sigma s_2 s_{4534} s_{1301} = \sigma b_{320}, \tag{5.26}$$

where $(\varphi_s^*)^3 = U_3^{-1}$ with U_3 given by Equation (5.21).

Proof. The expression given in Equation (5.26) in terms of the generators of N_J can be written down directly by inspection on Equations (5.7) and (5.10). To show that φ_s^* is an element of quasi-translation of N_J , recall that by Equation (5.6) $(\varphi_s^*)^3 \in P$ is the translation by $h_1 - h_2$. That is, φ_s^* is a quasi-translation of order three to the element U_3^{-1} . A quasi-translation of order three for U_3^{-1} can be constructed simply by a rewriting of U_3^{-1} using the defining relations (5.14b) of $\widetilde{W}(B_3^{(1)})$,

$$U_{3}^{-1} = b_{321320321}\sigma,$$

= $\sigma b_{320}b_{321}b_{320},$
= $\sigma b_{320}b_{321}\sigma^{2}b_{320},$
= $\sigma b_{320}\sigma b_{320}\sigma b_{320}.$ (5.27)

In fact, we have

$$\varphi_s^* = \sigma s_2 s_{4534} s_{1301} = \sigma b_{320}, \quad (\varphi_s^*)^3 = U_3^{-1}.$$
(5.28)

The action of φ_s^* on the $J \cup \beta = \{\alpha_0, \alpha_3, \alpha_5, \alpha_7, \beta_1, \beta_2, \beta_3, \beta_0\}$ basis is given by,

$$\varphi_s^* : \begin{pmatrix} \alpha_0, \alpha_3, \alpha_5, \alpha_7, \\ \beta_1, \beta_2, \beta_3, \beta_0 \end{pmatrix} \mapsto \begin{pmatrix} \alpha_3, \alpha_7, \alpha_5, \alpha_0, \\ \beta_{2330}, \beta_1, \beta_{23}, -\beta_{1233} \end{pmatrix},$$
(5.29)

moreover we have,

$$\left(\varphi_s^*\right)^3: \begin{pmatrix} \alpha_0, \alpha_3, \alpha_5, \alpha_7, \\ \beta_1, \beta_2, \beta_3, \beta_0 \end{pmatrix} \mapsto \begin{pmatrix} \alpha_0, \alpha_3, \alpha_5, \alpha_7, \\ \beta_1, \beta_2, \beta_3 + \delta, \beta_0 - 2\delta \end{pmatrix},$$
(5.30)

thus ending the proof.

Hence, the proof of last statement of Theorem 1.3 follows from Proposition 5.3.

6. Singular fibre of type $A_1^{(1)}$ associated with the map φ_2

In this section we compute the action of the map φ_2 (1.2) on the orthogonal complement of the singular fibre [0:1] of type $A_2^{(1)}$ of pencil p_2 (3.3). Then, we compute the normalizer of the Weyl group $W(4A_1)$ in the extended Weyl group $\widetilde{W}(E_6^{(1)})$. In this case, this will not be directly as in the Section 5, but we will use the results of that section to show that the mentioned action is on a translated root lattice of type $B_3^{(1)}$. So, the final outcome of this section is that the map φ_2 (1.2) on the singular fibre [0:1] has symmetry type $B_3^{(1)} \subset E_7^{(1)}$, thus ending the proof of Theorem 1.2.

6.1. Action on the simple roots of $E_7^{(1)}$ and decomposition in $\widetilde{W}(E_7^{(1)})$. Following Table 2 we have that the singular fiber of the pencil (3.9) at the point $[\mu:\nu] = [0:1]$ is of type $A_1^{(1)}$ and it is generated by the following divisors in Pic(X_2):

$$D_0 = H_x + H_y - F_1 - F_2 - F_3 - F_4, \quad D_1 = H_x + H_y - F_5 - F_6 - F_7 - F_8.$$
(6.1)

These two divisors are permuted by the pullback map φ_2^* , that is $\varphi_2^*(D_i) = D_{(i+1)}$, $i \in \mathbb{Z}/2\mathbb{Z}$. The rational elliptic surface X_2 with this choice of the singular fibre is shown in Figure 10. The orthogonal complement of $A_1^{(1)}$ in Pic (X_2) is generated by the same formula as Equation (5.2) with $F_i \leftrightarrow L_i$.

The set $\Delta^{(1)} = \Delta(E_7^{(1)}) = \{\alpha_j \mid 0 \le j \le 7\}$ is a simple system of type $E_7^{(1)}$, its Dynkin diagram given in Figure 8. The anti-canonical divisor corresponds to the following null-root in the $E_7^{(1)}$ root system:

$$-K_{X_2} = \delta = \alpha_{071122663335554444}. \tag{6.2}$$

Let $V^{(1)}$ be a vector space with basis $\Delta^{(1)}$, equipped with a symmetric bilinear form given by Equations (2.15) and (C.1). The corresponding extended affine Weyl group is,

$$\widetilde{W}(E_7^{(1)}) = \langle s_i \mid 0 \le i \le 7 \rangle \rtimes \langle \sigma \rangle = W(E_7) \ltimes P = W(E_7) \ltimes \langle u_j \mid 1 \le j \le 7 \rangle, \quad (6.3)$$

where u_j is the translation by the fundamental weight of the $E_7^{(1)}$ system, h_j $(1 \le j \le 7)$. In particular, the action of $s_j \in W(E_7^{(1)})$ on $\Delta^{(1)}$ is given by Equations (2.16) and (C.1). See Appendix C for basic data and properties of $\widetilde{W}(E_7^{(1)})$.

Using Equations (3.12) and (5.2) we have φ_s^* acts on $\overline{\Delta}^{(1)}$ by:

$$\varphi_{2}^{*}: \begin{pmatrix} \alpha_{0}, \alpha_{1}, \alpha_{2}, \alpha_{3}, \\ \alpha_{4}, \alpha_{5}, \alpha_{6}, \alpha_{7} \end{pmatrix} \mapsto \begin{pmatrix} \alpha_{0}, -\alpha_{123450}, \alpha_{11233445670}, -\alpha_{6}, \\ \alpha_{2456}, -\alpha_{24}, -\alpha_{567}, \alpha_{234567} \end{pmatrix}.$$
(6.4)

Computing three iterations of the map φ_2^* we obtain:

$$(\varphi_2^*)^3 : \begin{pmatrix} \alpha_0, \alpha_1, \alpha_2, \alpha_3, \\ \alpha_4, \alpha_5, \alpha_6, \alpha_7 \end{pmatrix} \mapsto \begin{pmatrix} \alpha_0, \alpha_1, \alpha_2 + \delta, \alpha_3, \\ \alpha_4, \alpha_5 - \delta, \alpha_6, \alpha_7 + \delta \end{pmatrix}.$$
(6.5)

By Equation (2.38), we recognise that $(\varphi_2^*)^3 \in P$ is the translation by $-h_2 + h_5 - h_7$, and φ_2^* , a quasi-translation of order three in $\widetilde{W}(E_7^{(1)})$. Using Equation (C.4), we find that $|-h_2 + h_5 - h_7|^2 = \frac{3}{2}$. Finally, φ_2^* has the following decomposition in terms of the generators of $\widetilde{W}(E_7^{(1)})$ using Equations (2.33) and (2.40):

$$\varphi_2^* = \sigma s_{6524310765431463456}. \tag{6.6}$$



FIGURE 10. The space of initial values of (1.2) interpreted as elliptic surface of type $A_1^{(1)}$.

6.2. Normalizer of $W(4A_1)$ in $\widetilde{W}(E_7^{(1)})$. The initial observation on φ_2^* given in Equation (6.6), is that like the φ_s^* discussed in Section 5 (where φ_s^* was shown to be an element of the normalizer for a standard parabolic subgroup W_J with $J = \{\alpha_0, \alpha_3, \alpha_5, \alpha_7\} \cong 4A_1\}$, φ_2^* is an order three quasi-translation by a vector of squared length $\frac{3}{2}$. However, unlike φ_s^* the actions of φ_2^* on $\Delta^{(1)}$ given in Equation (6.4) shows no clear permutations on any subset of $\Delta^{(1)}$. Here we show that φ_2^* in fact is an element of the normalizer for another type $4A_1$ parabolic subgroup W_{uJ} for some $u \in \widetilde{W}(E_7^{(1)})$. To find the appropriate u in $W(E_7^{(1)})$ we use the fact that the two squared length $\frac{3}{2}$ vectors associate to φ_2^* and φ_s^* are related by,

$$u^{-1}(h_1 - h_2) = -h_2 + h_5 - h_7$$
, where $u = s_{5436542} \in \widetilde{W}\left(E_7^{(1)}\right)$. (6.7)

By the contragredient action of $\widetilde{W}\left(E_7^{(1)}\right)$ we have,

$$u\Delta^{(1)} \colon \begin{pmatrix} \alpha_0, \alpha_1, \alpha_2, \alpha_3, \\ \alpha_4, \alpha_5, \alpha_6, \alpha_7 \end{pmatrix} \mapsto \begin{pmatrix} \alpha_0, \alpha_{1345}, -\alpha_{2344556}, \alpha_6, \\ \alpha_{245}, \alpha_3, \alpha_4, \alpha_{567}, \end{pmatrix}.$$
 (6.8)

That is, we have

$$uJ = u\{\alpha_0, \alpha_3, \alpha_5, \alpha_7\} = \{uJ_1, uJ_2, uJ_3, uJ_4\} = \{\alpha_0, \alpha_6, \alpha_3, \alpha_{567}\},$$
(6.9)

which is another type $4A_1$ sub-root system of the $E_7^{(1)}$ root system $\Phi^{(1)}$.

Proposition 6.1. The normalizer of W_{uJ} with $uJ = \{\alpha_0, \alpha_6, \alpha_3, \alpha_{567}\} \subset \Phi^{(1)}$ in $\widetilde{W}(E_7^{(1)})$ is given by

$$N(W_{uJ}) = W_{uJ} \rtimes N_{uJ} = \langle s_0, s_6, s_3, s_{76567} \rangle \rtimes \left(\langle ub_1, ub_2, ub_3, ub_0 \rangle \rtimes \langle u\sigma u^{-1} \rangle \right)$$
(6.10)
$$\cong W(4A_1) \rtimes \widetilde{W} \left(B_3^{(1)} \right),$$

with,

$$ub_1 = us_{6576}u^{-1}, \quad ub_2 = us_{4534}u^{-1}, \quad ub_3 = us_2u^{-1}, \quad ub_0 = us_{1301}u^{-1},$$
 (6.11)
where $u = s_{5436542}.$

Proof. By Equations (2.22) and (6.9), generators of the parabolic subgroup W_{uJ} are obtain from generators of W_J by a conjugation of $u = s_{5436542}$, that is,

$$W_{uJ} = uW_J u^{-1} = \langle us_0 u^{-1}, us_3 u^{-1}, us_5 u^{-1}, us_7 u^{-1} \rangle = \langle s_0, s_6, s_3, s_{76567} \rangle.$$
(6.12)
By [5], the normalizer of W_{uJ} is obtained from the normalizer of W_J by a conjugation of u .

Proposition 6.2. The root system for $N(W_{uJ})$ of Proposition 6.1 is given by

$$uJ \cup u\beta = \{uJ_1, uJ_2, uJ_3, uJ_4\} \cup \{u\beta_1, u\beta_2, u\beta_3, u\beta_0\} \cong 4A_1 \times B_3^{(1)}, \tag{6.13}$$

with

$$uJ_{1} = \alpha_{0}, \quad uJ_{2} = \alpha_{6},$$

$$uJ_{3} = \alpha_{3}, \quad uJ_{4} = \alpha_{567},$$

$$u\beta_{1} = \alpha_{344567}, \quad u\beta_{2} = \alpha_{22344556},$$

$$\beta_{2} = -\alpha_{22344556}, \quad u\beta_{0} = \alpha_{1122445560}.$$

(6.14)

$$\begin{split} & u\beta_3 = -\alpha_{2344556}, \quad u\beta_0 = \alpha_{1133445560}, \\ & where \ u\beta_0 + u\beta_1 + 2u\beta_2 + 2u\beta_3 = \alpha_{071122663335554444} = \delta. \\ & The \ group \ N_{uJ} \ can \ be \\ & realised \ as \ \widetilde{W}(B_3^{(1)}) \ with \ ub_i \ acting \ as \ reflection \ along \ the \ root \ u\beta_i, \ while \ u\sigma u^{-1} \\ & exchanges \ u\beta_0 \ and \ u\beta_1, \ see \ Figure \ 9. \end{split}$$

Proposition 6.3. The map φ_2^* is an element of quasi-translation of order three in the

$$N(W_{uJ}) \cong W(4A_1) \rtimes \widetilde{W}(B_3^{(1)}) \tag{6.15}$$

subgroup of $\widetilde{W}(E_7^{(1)})$ give in Proposition 6.1. We have,

$$\varphi_2^* = u\sigma u^{-1} s_0 s_3 u b_{3231203}. \tag{6.16}$$

Proof. The actions of φ_2^* on the $uJ \cup u\beta$ basis is obtained from that of Equation (6.4) by a change of basis,

$$\varphi_{2}^{*}: \begin{pmatrix} uJ_{1}, uJ_{2}, uJ_{3}, uJ_{4}, \\ u\beta_{1}, u\beta_{2}, u\beta_{3}, u\beta_{0} \end{pmatrix} \mapsto \begin{pmatrix} uJ_{1}, -uJ_{4}, -uJ_{2}, uJ_{3}, \\ u\beta_{2}, u\beta_{1222333300}, -u\beta_{230}, -u\beta_{1222333300} \end{pmatrix}.$$
(6.17)

Moreover, we have

$$(\varphi_{2}^{*})^{3} : \begin{pmatrix} uJ_{1}, uJ_{2}, uJ_{3}, uJ_{4}, \\ u\beta_{1}, u\beta_{2}, u\beta_{3}, u\beta_{0} \end{pmatrix} \mapsto \begin{pmatrix} uJ_{1}, uJ_{2}, uJ_{3}, uJ_{4}, \\ u\beta_{1}, u\beta_{2}, u\beta_{3} + \delta, u\beta_{0} - 2\delta \end{pmatrix}.$$
 (6.18)

We use Equation (2.33) on φ_2^* 's action on the $uJ \cup u\beta$ -system given in Equation (6.17) to write φ_2^* as a product of generators of $N(W_{uJ})$. We find,

$$\varphi_2^* = u\sigma u^{-1} s_0 s_3 u b_{3231203}, \tag{6.19}$$

thus ending the proof.

Hence, the proof of last statement of Theorem 1.2 follows from Proposition 6.3.

7. Conclusion

In this paper, we investigated the symmetries of three plane integrable systems given by Equations (1.1) to (1.3). The first two systems, φ_1 (1.1) and φ_2 (1.2) arise from the KHK discretisation of the tetrahedral and octahedral reduced Nahm systems [20]. Such systems and some generalisations have been discussed extensively in the literature [8, 9, 14, 15, 35, 48] (see Appendix A), but a close inspection of their action on singular fibres was never discussed. The last map, φ_s (1.3), is a generalisation of the map φ_2 we introduced in Section 3 through the QRT root construction (see [18, §6.3] and [11, Introduction]).

In this paper, in light of the theory of Oguiso and Shioda [32] and its application to plane discrete integrable systems, see [6] and [11, Chap. 7], we found out that these three systems possess singular fibre structure of type $(A_3 \times A_2 \times 2A_1)^{(1)}$, $(3A_3 \times A_1)^{(1)}$, and $(2A_1 \times 2A_2 \times A_0^*)^{(1)}$ respectively. After this construction, we concentrated our study on three specific singular

After this construction, we concentrated our study on three specific singular fibres and their symmetry group, i.e. their orthogonal component in the Picard group, namely the fibre [0:1] of type $A_2^{(1)}$ for φ_1 , the fibre [0:1] of type $A_1^{(1)}$ for the map φ_2 , and the fibre [0:1] of type $A_1^{(1)}$ for the map φ_s . The result is that on these singular fibres the symmetry group is a proper subgroup of symmetry group, forming a sublattice in the Picard group. This situation is not completely unusual, for instance see the papers [1, 6, 7, 24, 29, 44, 46]. However, our result is relevant because for the first time we find the non-simply laced Coxeter group $B_3^{(1)} \subset E_7^{(1)} = (A_1^{(1)})^{\perp}$ as symmetry group for both the maps φ_2 and φ_s . Moreover, our result is obtained within the algorithmic framework of the normaliser of parabolic subgroups [5, 21], which already proved fruitful in interpreting known results [42, 43]. In this sense, we can say that non-simply laced symmetry groups were actually hidden in plain sight in known integrable plane systems.

Further research is ongoing about the symmetry types associated to the other singular fibres. Furthermore, we note that it is also interesting to consider the (possible) subgroups of elements of the generic fibres, i.e. the elliptic fibre of type $A_0^{(1)}$. Considering the deautonomisation of these systems we could then possibly build new "degenerate" elliptic discrete Painlevè equations as it was done for instance in [1, 6, 25]. In general, for each singular fibre the deautonomisation procedure of [6] might be applied.

To conclude, what we laid down in this paper can also evolve into an ambitious program of building the symmetry group of all the QRT maps and QRT root associated with given elements of the classification given by Oguiso and Shioda's classification of singular fibre configurations of rational elliptic surfaces [32], following what was done in [11, Chap. 7].

Acknowledgments. GG acknowledges support of the GNFM through Progetto Giovani GNFM 2023: "Strutture variazionali e applicazioni delle equazioni alle differenze ordinarie", CUP_E53C22001930001.

We thank Prof. Bert Van Geemen and Dr. Michele Graffeo for their assistance and stimulating discussion during the preparation of this paper.

Appendix A. Reduced Nahm systems of tetrahedral and octahedral type

In [20], some special cases of Nahm's equations with particular symmetries studied in connection with the theory of monopoles. Due to the symmetry of the associated Nahm matrices the systems are called the *tetrahedral Nahm system*, *octahedral Nahm system*, and *icosahedral Nahm system*. In particular, we adopt the form of the systems of *tetrahedral Nahm system* and *octahedral Nahm system* from [35]:

$$\dot{x} = x^2 - y^2, \quad \dot{y} = -2xy,$$
 (A.1a)

$$\dot{x} = 2x^2 - 48y^2, \quad \dot{y} = -6xy - 8y^2.$$
 (A.1b)

Consider now the family of Hamiltonian systems:

$$H_{a,b,c} = x^a y^b \left(kx + ly\right)^c,\tag{A.2}$$

see for instance [9]. The corresponding Hamilton equations are given by:

$$\begin{pmatrix} \dot{x} \\ \dot{y} \end{pmatrix} = x^{1-a} y^{1-b} (kx+ly)^{1-c} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \nabla H_{a,b,c} = \begin{pmatrix} x(bkx+bly+cly), \\ -y(akx+aly+ckx) \end{pmatrix}.$$
 (A.3)

This system is *quadratic* and *integrable* for all triples (a, b, c) (it preserves the invariant (A.2)). By direct computation we have the following result:

Lemma A.1. The Hamiltonian systems (A.3) with (a, b, c) = (1, 1, 1) is mapped to the tetrahedral Nahm system (A.1a) through the following transformation:

$$(x,y) \to \left(\frac{\sqrt{3}(4kl)^{1/3}}{2k}x - \frac{(4kl)^{1/3}}{2k}y, \frac{(4kl)^{1/3}}{l}y\right), \quad t \to \frac{\sqrt{(3)(4kl)^{1/3}}}{2}t, \qquad (A.4)$$

while the Hamiltonian systems (A.3) with (a, b, c) = (1, 1, 2) is mapped to the octahedral Nahm systems (A.1b) through the following transformation:

$$(x,y) \to \left(-\frac{1+\mathrm{i}}{5}\left(\frac{5}{k}\right)^{3/4}l^{1/4}x - \frac{3}{10}(1+\mathrm{i})l^{1/4}\left(\frac{5}{k}\right)^{3/4}y, \frac{1+\mathrm{i}}{2}\left(\frac{5}{l}\right)^{3/4}k^{1/4}y\right) \quad (A.5a)$$

$$t \to -\frac{1+i}{10} 5^{3/4} (lk)^{1/4} t$$
 (A.5b)

Let us now turn to the genesis of the maps φ_1 and φ_2 . As stated in the introduction the KHK discretisation [19, 26, 27, 31] is a discretisation method for systems of first-order differential equations.

Definition A.2. Consider a system of first-order ordinary differential equations:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}), \quad \mathbf{x}: \mathbb{R}^N \to \mathbb{R}, \quad \mathbf{f}: \mathbb{R}^N \to \mathbb{R}^N, \quad N \in \mathbb{N}, t \in \mathbb{R}.$$
 (A.6)

Then the KHK discretisation of the system (A.6) is given by:

$$\frac{\mathbf{x}' - \mathbf{x}}{h} = 2\mathbf{f}\left(\frac{\mathbf{x}' + \mathbf{x}}{2}\right) - \frac{\mathbf{f}(\mathbf{x}') + \mathbf{f}(\mathbf{x})}{2},\tag{A.7}$$

where $\mathbf{x} = \mathbf{x}(nh)$, $\mathbf{x}' = \mathbf{x}((n+1)h)$, and h is a infinitesimal parameter, i.e. $h \to 0^+$.

If the function **f** is *quadratic*, then the discrete system defined by Equation (A.7) give rise to a birational map of \mathbb{R}^N , see [10, 35]. In such a case, this birational map can be considered to be a restriction of a map over \mathbb{C}^N , but dynamical properties are better understood over compact spaces to take into account points escaping at

infinity. So, it is costumary to consider the maps over some compactification of \mathbb{C}^N . The usual choices are \mathbb{P}^M or $(\mathbb{P}^1)^{\times M}$. In this paper, we will use the latter, but each choice will yield the same final results.

Now following [30] we show that the KHK maps corresponding to the discretisation cases (a, b, c) = (1, 1, 2) and (a, b, c) = (1, 1, 1) of the systems (A.3) can be brought to the form (1.1) and (1.2). That is, denoting these two maps by $\Phi_{(1,1,1),h} \in \operatorname{Bir}(\mathbb{P}^1 \times \mathbb{P}^1)$ and $\Phi_{(1,1,2),h} \in \operatorname{Bir}(\mathbb{P}^1 \times \mathbb{P}^1)$ respectively, we have by a direct computation:

Proposition A.3. The maps $\Phi_{(1,1,2),h} \in Bir(\mathbb{P}^1 \times \mathbb{P}^1)$:

$$\Phi_{(1,1,2),h}:(x,y) \mapsto \left(\frac{x\left[2 + (3kx + 5ly)h\right]}{2 + (kx + ly) - (3k^2x^2 + 2klxy + 3l^2y^2)h^2} \\ \frac{y\left[2 - (5kx + 3ly)\right]}{2 + (kx + ly) - (3k^2x^2 + 2klxy + 3l^2y^2)h^2}\right)^T$$
(A.8)

and $\Phi_{(1,1,1),h} \in \operatorname{Bir}(\mathbb{P}^1 \times \mathbb{P}^1)$:

$$\boldsymbol{\Phi}_{(1,1,1),h}:(x,y) \mapsto \left(\frac{x[1+(kx+2ly)h]}{1-h^2k^2x^2-h^2klxy-h^2l^2y^2}, \frac{y[1-(2kx-ly)h]}{1-h^2k^2x^2-h^2klxy-h^2l^2y^2}\right)^T$$
(A.9)

are conjugated to the maps φ_1 and φ_2 respectively, through a projective coallineation. That is:

$$\varphi_1 = \chi_1^{-1} \circ \mathbf{\Phi}_{(1,1,2),h} \circ \chi_1, \quad and \quad \varphi_2 = \chi_2^{-1} \circ \mathbf{\Phi}_{(1,1,1),h} \circ \chi_2.$$
 (A.10)

where:

$$\chi_1: (x,y) \mapsto \left(\frac{1}{2kh} \frac{2(x-y) + \sqrt{2}}{x+y}, -\frac{1}{2lh} \frac{2(x-y) - \sqrt{2}}{x+y}\right),$$
(A.11a)

$$\chi_2: (x,y) \mapsto \left(\frac{2}{x+y}, -\frac{hk-x+y}{hl(x+y)}\right).$$
(A.11b)

Remark A.4. We remark that the literature on the KHK discretisation of the Nahm systems (A.1) is extensive. In this remark we collect some sparse facts that are worth mentioning:

- The integrability of the KHK discretisation of the three kinds of reduced Nahm systems, tetrahedral, octahedral and icosahedral, was proved by direct computation in [35]. A Lax pair of these systems was introduced in [14].
- In [8] it was noted that these systems are not minimal, in the sense of Section 2, and can be reduced to QRT roots.
- Generalisations of these systems in N-dimensions were presented in [36].
- From [48] it is known that the only integrable cases of KHK discretisation of the system (A.3) are obtained if (a, b, c) = (1, 1, 1), (a, b, c) = (1, 1, 2), (a, b, c) = (1, 2, 3) and their permutations.
- The map χ_1 given in (A.11a) is a modification of a map already presented in [15].

Appendix B. Basic properties of the group $\widetilde{W}(E_6^{(1)})$

The Dynkin diagram of type $E_6^{(1)}$, $\Gamma(E_6^{(1)})$ is given in Figure 5. Its simple system $\Delta^{(1)} = \{\alpha_j \mid 0 \le j \le 6\}$ forms a basis for an 7-dimensional real vector space $V^{(1)}$,

equipped with a bilinear form given by Equation (2.15) and the generalized Cartan matrix of type $E_6^{(1)}$,

$$C(E_6^{(1)}) = (a_{ij})_{1 \le i,j \le 6,0} = (\alpha_i \cdot \alpha_j^{\vee})_{1 \le i,j \le 6,0},$$

= $(\alpha_i \cdot \alpha_j)_{1 \le i,j \le 6,0},$
= $\begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & -1 \\ 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & -1 & 2 & 0 \\ 0 & 0 & -1 & 0 & 0 & 2 \end{pmatrix}$. (B.1)

Defining relations of $W(E_6^{(1)}) = \langle s_i | 0 \le i \le 6 \rangle$ can be read off $\Gamma(E_6^{(1)})$. Generators $s_j \in W(E_6^{(1)})$ act on $V^{(1)}$ by Equation (2.16), where a_{ij} is the (i, j)-entry of $C(E_6^{(1)})$ in Equation (B.1). We have the null root and the highest root of $E_6^{(1)}$ given by

$$\delta = \alpha_0 + \tilde{\alpha} = \alpha_0 + \alpha_1 + \alpha_5 + 2(\alpha_2 + \alpha_4 + \alpha_6) + 3\alpha_3. \tag{B.2}$$

A dual space $V^{(1)*}$ with basis $\{h_1, \ldots, h_6, h_\delta\}$ is given by Equation (2.23). The subspace $X_0 \subset V^{(1)*}$ has two bases: the set of simple coroots $\{\pi(\alpha_j^{\vee}) \mid 1 \leq j \leq 6\}$, and the set of fundamental weights $\{h_j \mid 1 \leq j \leq 6\}$. They are related by Equation (2.29),

$$\alpha_j = \sum_{k=1}^6 \left(C(E_6) \right)_{kj} h_k = \sum_{k=1}^6 a_{jk} h_k, \quad 1 \le j \le 6,$$
(B.3)

where we have used the fact that $\Gamma(E_6^{(1)})$ is simply-laced, so $C(E_6)$ is symmetric and we have identified the simple coroots $\pi(\alpha_j^{\vee})$ with the simple roots α_j for $1 \le j \le 6$. The squared lengths of $h_j(1 \le j \le 6)$ is given by the diagonal entries of $C(E_6)^{-1}$ in Equation (B.4),

$$((h_i, h_j))_{1 \le i, j \le 6} = C(E_6)^{-1} = \begin{pmatrix} \frac{4}{3} & \frac{5}{3} & 2 & \frac{4}{3} & \frac{2}{3} & 1\\ \frac{5}{3} & \frac{10}{3} & 4 & \frac{8}{3} & \frac{4}{3} & 2\\ 2 & 4 & 6 & 4 & 2 & 3\\ \frac{4}{3} & \frac{8}{3} & 4 & \frac{10}{3} & \frac{5}{3} & 2\\ \frac{2}{3} & \frac{4}{3} & 2 & \frac{5}{3} & \frac{4}{3} & 1\\ 1 & 2 & 3 & 2 & 1 & 2 \end{pmatrix}.$$
 (B.4)

That is they are $\frac{4}{3}$, $\frac{10}{3}$, 6, $\frac{10}{3}$, $\frac{4}{3}$ and 2, respectively. Moreover, by Equation (2.30) we have

$$h_i = \sum_{k=1}^{6} C(E_6)_{ik}^{-1} \alpha_k \quad \text{for} \quad 1 \le i \le n.$$
 (B.5)

with $C(E_6)^{-1}$ in Equation (B.4). The group A of diagram automorphisms of $\Gamma(E_6^{(1)})$ is generated by σ , written as a permutation on the index set of $\Delta^{(1)}$ we have,

$$\sigma = (150)(246), \tag{B.6}$$

so that $(\sigma)^3 = 1$. Moreover, we have

$$\sigma s_{\{1,2,3,4,5,6,0\}} = s_{\{5,4,3,6,0,2,1\}} \sigma. \tag{B.7}$$

That is, σ permutes the simple reflections by conjugation, for example, we have $\sigma s_1 \sigma^{-1} = s_5$ from Equation (B.7). The extended affine Weyl group of type E_6 is given by

$$\widetilde{W}(E_6^{(1)}) = \langle s_i \mid 0 \le i \le 6 \rangle \rtimes \langle \sigma \rangle = W(E_6) \ltimes \langle u_j \mid 1 \le j \le 6 \rangle, \tag{B.8}$$

where u_j is the element of translation by h_j .

Appendix C. Basic properties of the group $\widetilde{W}(E_7^{(1)})$

The Dynkin diagram of type $E_7^{(1)}$, $\Gamma(E_7^{(1)})$ is given in Figure 8. Its simple system $\Delta^{(1)} = \{\alpha_j \mid 0 \le j \le 7\}$ forms a basis for an 8-dimensional real vector space $V^{(1)}$ equipped with a bilinear form given by Equation (2.15) and the generalized Cartan matrix of type $E_7^{(1)}$:

$$C(E_7^{(1)}) = (a_{ij})_{1 \le i,j \le 7,0} = (\alpha_i \cdot \alpha_j^{\vee})_{1 \le i,j \le 7,0},$$

$$= (\alpha_i \cdot \alpha_j)_{1 \le i,j \le 7,0},$$

$$= \begin{pmatrix} 2 & 0 & -1 & 0 & 0 & 0 & 0 & -1 \\ 0 & 2 & 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 2 & -1 & 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 2 \end{pmatrix}.$$
(C.1)

Defining relations for $W(E_7^{(1)}) = \langle s_i | 0 \le i \le 7 \rangle$ can be read off $\Gamma(E_7^{(1)})$. Generators $s_j \in W(E_7^{(1)})$ act on $V^{(1)}$ by Equation (2.16), where a_{ij} is the (i, j)-entry of $C(E_7^{(1)})$ from Equation (C.1). We have the null root and the highest root of $E_7^{(1)}$ given by

$$\delta = \alpha_0 + \tilde{\alpha} = \alpha_0 + \alpha_7 + 2(\alpha_2 + \alpha_1 + \alpha_6) + 3(\alpha_3 + \alpha_5) + 4\alpha_4.$$
(C.2)

A dual space $V^{(1)*}$ with basis $\{h_1, \ldots, h_7, h_\delta\}$ is given by Equation (2.23). The subspace $X_0 \subset V^{(1)*}$ has two bases: the set of simple coroots $\{\pi(\alpha_j^{\vee}) \mid 1 \leq j \leq 7\}$, and the set of fundamental weights $\{h_j \mid 1 \leq j \leq 7\}$, and they are related by Equation (2.29),

$$\alpha_j = \sum_{k=1}^7 \left(C(E_7) \right)_{kj} h_k = \sum_{k=1}^7 a_{kj} h_k, \quad 1 \le j \le 7,$$
(C.3)

where we have used the fact that $\Gamma(E_7^{(1)})$ is simply-laced, so $C(E_7)$ is symmetric and we have identified the simple coroots $\pi(\alpha_j^{\vee})$ with the simple roots α_j for $1 \le j \le 7$. The squared lengths of $\{h_j \mid 1 \le j \le 7\}$ is given by the diagonal entries of $C(E_7)^{-1}$ in Equation (C.4),

$$((h_i, h_j))_{1 \le i, j \le 7} = C(E_7)^{-1} = \begin{pmatrix} 2 & 2 & 3 & 4 & 3 & 2 & 1 \\ 2 & \frac{7}{2} & 4 & 6 & \frac{9}{2} & 3 & \frac{3}{2} \\ 3 & 4 & 6 & 8 & 6 & 4 & 2 \\ 4 & 6 & 8 & 12 & 9 & 6 & 3 \\ 3 & \frac{9}{2} & 6 & 9 & \frac{15}{2} & 5 & \frac{5}{2} \\ 2 & 3 & 4 & 6 & 5 & 4 & 2 \\ 1 & \frac{3}{2} & 2 & 3 & \frac{5}{2} & 2 & \frac{3}{2} \end{pmatrix}.$$
 (C.4)

That is, they are 2, $\frac{7}{2}$, 6, 12, $\frac{15}{2}$, 4, $\frac{3}{2}$. Moreover, by Equation (2.30) we have

$$h_i = \sum_{k=1}^{7} C(E_7)_{ik}^{-1} \alpha_k \quad \text{for} \quad 1 \le i \le n,$$
 (C.5)

with $C(E_7)^{-1}$ given in Equation (C.4). The group A of diagram automorphisms of $\Gamma(E_7^{(1)})$ is generated by σ , written as a permutation on the index set of $\Delta^{(1)}$ we have:

$$\sigma = (07)(16)(35) \tag{C.6}$$

so that $(\sigma)^2 = 1$. Moreover, we have

$$\sigma s_{\{1,2,3,4,5,7,0\}} = s_{\{6,2,5,4,3,1,0\}} \sigma. \tag{C.7}$$

The extended affine Weyl group of type E_7 is then

$$\widetilde{W}(E_7^{(1)}) = \langle s_i \mid 0 \le i \le 7 \rangle \rtimes \langle \sigma \rangle = W(E_7) \ltimes \langle u_j \mid 1 \le j \le 7 \rangle, \tag{C.8}$$

where u_j is the element of translation by h_j .

References

- J. Atkinson, P. Howes, N. Joshi, and N. Nakazono. "Geometry of an elliptic difference equation related to Q4". In: J. Lond. Math. Soc. (2) 93.3 (2016), pp. 763–784.
- [2] A. Beauville. Complex algebraic surfaces. Second. Vol. 34. London Mathematical Society Student Texts. Translated from the 1978 French original by R. Barlow, with assistance from N. I. Shepherd-Barron and M. Reid. Cambridge University Press, Cambridge, 1996, pp. x+132.
- M. P. Bellon. "Algebraic Entropy of Birational Maps with Invariant Curves". In: Lett. Math. Phys. 50.1 (1999), pp. 79–90.
- [4] N. Bourbaki. Groupes et algèbres de Lie, Chapitre IV-VI. Hermann, Paris, 1968, 288 pp. (loose errata).
- [5] B. Brink and R. B. Howlett. "Normalizers of parabolic subgroups in Coxeter groups". In: *Invent. Math.* 136.2 (1999), pp. 323–351.
- [6] A. S. Carstea, A. Dzhamay, and T. Takenawa. "Fiber-dependent deautonomization of integrable 2D mappings and discrete Painlevé equations". In: J. Phys. A: Math. Theor. 50 (2017), 405202, (41pp).
- [7] A. S. Carstea and T. Takenawa. "A classification of two-dimensional integrable mappings and rational elliptic surfaces". In: J. Phys. A 45 (2012), 155206 (15pp).
- [8] A. S. Carstea and T. Takenawa. "A note on minimization of rational surfaces obtained from birational dynamical systems". In: J. Nonlinear Math. Phys. 20.suppl. 1 (2013), pp. 17–33. ISSN: 1402-9251.
- [9] E. Celledoni, R. I. McLachlan, D. I. McLaren, B. Owren, and G. R. W. Quispel. "Two classes of quadratic vector fields for which the Kahan discretization is integrable". In: *MI Lecture Notes* 74 (2017), pp. 60–62.
- [10] E. Celledoni, R. I. McLachlan, B. Owren, and G. R. W. Quispel. "Geometric properties of Kahan's method". In: J. Phys. A: Math. Theor. 46.2 (2013), p. 025201.
- J. Duistermaat. Discrete Integrable Systems: QRT Maps and Elliptic Surfaces. Springer Monographs in Mathematics. Springer New York, 2011. ISBN: 9781441991263.

- [12] M. Graffeo and G. Gubbiotti. Growth and complexity in the Cremona cube group. 2023. arXiv: 2301.09129 [math.GA].
- [13] M. Graffeo and A. T. Ricolfi. "On the Behrend function and the blowup of some fat points". In: Adv. Math. (2023).
- [14] G. Gubbiotti. "Lax pairs for the discrete reduced Nahm systems". In: Math. Phys. Anal. Geom. 24 (2021), 9 (13pp).
- [15] G. Gubbiotti and N. Joshi. "Space of initial values of a map with a quartic invariant". In: Bull. Aus. Mat. Soc. (2020), pp. 1–12.
- [16] F. Haggar, G. Byrnes, G. Quispel, and H. Capel. "k-integrals and k-Lie symmetries in discrete dynamical systems". In: *Physica A* 233.1 (1996), pp. 379– 394.
- [17] J. Hietarinta. "Definitions and Predictions of Integrability for Difference Equations". In: Symmetries and Integrability of Difference Equations. Ed. by D. Levi, P. Olver, Z. Thomova, and P. Winternitz. London Mathematical Society Lecture Notes series. Cambridge: Cambridge University Press, 2011, pp. 83–114.
- [18] J. Hietarinta, N. Joshi, and F. Nijhoff. Discrete Systems and Integrability. Cambridge Texts in Applied Mathematics. Cambridge University Press, 2016.
- [19] R. Hirota and K. Kimura. "Discretization of the Euler Top". In: J. Phys. Soc. Japan 69.3 (2000), pp. 627–630.
- [20] N. J. Hitchin, N. S. Manton, and M. K. Murray. "Symmetric monopoles". In: Nonlinearity 8 (1995), pp. 661–692.
- [21] R. B. Howlett. "Normalizers of parabolic subgroups of reflection groups". In: J. London Math. Soc. (2) 21.1 (1980), pp. 62–80.
- [22] J. E. Humphreys. Reflection groups and Coxeter groups. Vol. 29. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1990, pp. xii+204. ISBN: 0-521-37510-X.
- [23] E. L. Ince. Ordinary Differential Equations. Dover Books on Mathematics. New York: Dover Publications, 1957.
- [24] N. Joshi, K. Kajiwara, T. Masuda, N. Nakazono, and Y. Shi. "Geometric description of a discrete power function associated with the sixth Painlevé equation". In: Proc. A. 473 (2017).
- [25] N. Joshi and N. Nakazono. "Elliptic Painlevé equations from next-nearest-neighbor translations on the $E_8^{(1)}$ lattice". In: J. Phys. A 50.30 (2017), 305205 (17pp.) ISSN: 1751-8113.
- [26] W. Kahan. Unconventional numerical methods for trajectory calculations. Unpublished lecture notes. 1993.
- [27] W. Kahan and R.-C. Li. "Unconventional schemes for a class of ordinary differential equations - with applications to the Korteweg-de Vries equation". In: J. Comp. Phys. 134 (1997), pp. 316–331.
- [28] K. Kajiwara, M. Noumi, and Y. Yamada. "Geometric aspects of Painlevé equations". In: J. Phys. A 50 (2017), 073001 (164pp).
- [29] K. Kajiwara, N. Nakazono, and T. Tsuda. "Projective Reduction of the Discrete Painlevè System of Type $(A_2 + A_1)^{(1)}$ ". In: Int. Math. Res. Not. 2011.4 (2011), pp. 930–966.
- [30] P. H. van der Kamp, E. Celledoni, R. I. McLachlan, D. I. McLaren, B. Owren, and G. R. W. Quispel. "Three classes of quadratic vector fields for which the

Kahan discretisation is the root of a generalised Manin transformation". In: J. Phys. A: Math. Theor. 52 (2019), 045204 (10pp).

- [31] K. Kimura and R. Hirota. "Discretization of the Lagrange top". In: J. Phys. Soc. Japan 69 (2000), pp. 3193–3199.
- [32] K. Oguiso and T. Shioda. "The Mordell-Weil lattice of a rational elliptic surface". In: Comment. Math. Univ. St. Pauli 40 (Jan. 1991).
- [33] K. Okamoto. "Sur les feuilletages associés aux équations du second ordre à points critiques fixes de P. Painlevé". In: C. R. Acad. Sci. Paris Sér. A-B 285.12 (1977), A765–A767.
- [34] K. Okamoto. "Sur les feuilletages associés aux équation du second ordre à points critiques fixes de P. Painlevé. espaces des conditions initiales". In: Japan. J. Math New ser. 5 (1979), pp. 1–79.
- [35] M. Petrera, A. Pfadler, and Y. B. Suris. "On Integrability of Hirota-Kimura Type Discretizations". In: *Regul. Chaot. Dyn.* 16 (2011), pp. 245–289.
- [36] M. Petrera and R. Zander. "New classes of quadratic vector fields admitting integral-preserving Kahan-Hirota-Kimura discretizations". In: J. Phys. A: Math. Theor. 50 (2017), 205203, (13pp).
- [37] G. R. W. Quispel, J. A. G. Roberts, and C. J. Thompson. "Integrable mappings and soliton equations". In: *Phys. Lett. A* 126 (1988), p. 419.
- [38] G. R. W. Quispel, J. A. G. Roberts, and C. J. Thompson. "Integrable mappings and soliton equations II". In: *Physica D* 34.1 (1989), pp. 183–192.
- [39] J. A. G. Roberts and D. Jogia. "Birational maps that send biquadratic curves to biquadratic curves". In: J. Phys. A: Math. Theor. 48 (2015), 08FT02.
- [40] H. Sakai. "Rational surfaces associated with affine root systems and geometry of the Painlevé Equations". In: Comm. Math. Phys. 220.1 (2001), pp. 165–229.
- [41] M. Schütt and T. Shioda. Mordell-Weil Lattices. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics. Springer Nature Singapore, 2019.
- [42] Y. Shi. "Two variations on $(A_3 \times A_1 \times A_1)^{(1)}$ type discrete Painlevé equations". In: *Proc. R. Soc. A* 475.2229 (2019), 20190299 (18pp).
- [43] Y. Shi. Translations in affine Weyl groups and their applications in discrete integrable systems. 2022. arXiv: 2210.13736 [math-ph].
- [44] A. Stokes. "Full-parameter discrete Painlevé systems from non-translational Cremona isometries". In: J. Phys. A: Math. Theor. 51.49 (2018), p. 495206.
- [45] T. Takenawa. "Algebraic entropy and the space of initial values for discrete dynamical systems". In: J. Phys. A: Math. Gen. 34 (2001), p. 10533.
- [46] T. Takenawa. "Weyl group symmetry of type D₅⁽¹⁾ in the q-Painlevé V equation". In: Funkcial. Ekvac. 46.1 (2003), pp. 173–186.
- [47] T. Tsuda. "Integrable mappings via rational elliptic surfaces". In: J. Phys. A: Math. Gen. 37 (2004), p. 2721.
- [48] R. Zander. On the singularity structure of Kahan discretizations of a class of quadratic vector fields. 2021.

(G. Gubbiotti) Dipartimento di Matematica "Federigo Enriques", Università degli Studi di Milano, Via C. Saldini 50, 20133 Milano, Italy & INFN Sezione di Milano, Via G. Celoria 16, 20133 Milano, Italy

 $Email \ address: \verb"giorgio.gubbiotti@unimi.it"$

(Y. Shi) College of science and Engineering, Flinders at Tonsley, Flinders University, SA 5042, Australia

 $Email \ address: \verb"yang.shi@flinders.edu.au"$

Point	Irred. comp.	Repr. in $\operatorname{Pic}(X_2)$	Surf./symm. type
$[\mu:\nu]$	d_i	D_i	Λ/Λ^{\perp}
ζı	$\begin{cases} xy - (1 + i\sqrt{3})x - (1 - i\sqrt{3})y + 1 = 0 \\ xy - (1 - i\sqrt{3})x - (1 + i\sqrt{3})y + 1 = 0 \end{cases},$	$\begin{array}{l} H_x + H_y - L_1 - L_3 - L_5 - L_7 \\ H_x + H_y - L_2 - L_4 - L_6 - L_8 \end{array}$	$A_1^{(1)}/E_7^{(1)}$
$\zeta_2(s)$	$\left\{xy + 2\frac{sx}{3s - 4} + 2\frac{sy}{3s - 4} = \frac{s - 4}{3s - 4}\right\}$ $\left\{xy + 1 = 0\right\}$	$H_x + H_y - L_1 - L_2 - L_3 - L_4$ $H_x + H_y - L_5 - L_6 - L_7 - L_8$	$A_2^{(1)}/E_6^{(1)}$
$\zeta_3(s)$	$\begin{cases} x = -\frac{1 - i\sqrt{3}}{2} \\ y = -\frac{1 - i\sqrt{3}}{2} \end{cases}$	$H_x - L_4 - L_7,$ $H_y - L_3 - L_6$	$A_1^{(1)}/E_7^{(1)}$
	$\left\{\begin{array}{c} xy + \frac{1}{14} \frac{\left(13\sqrt{3} - 5\right)\left(4i\sqrt{3} - 7s + 8\right)x}{i\sqrt{3} - 19s + 23} \\ + \frac{1}{14} \frac{\left(13i\sqrt{3} - 5\right)\left(4i\sqrt{3} - 7s + 8\right)y}{i\sqrt{3} - 19s + 23} - \frac{1}{2} \frac{\left(5i\sqrt{3} + 1\right)\left(i\sqrt{3} - s + 5\right)}{i\sqrt{3} - 19s + 23} \end{array}\right\}$	$H_x + H_y - L_1 - L_2 - L_5 - L_8$	
$\zeta_4(s)$	$\begin{cases} x = -\frac{1+i\sqrt{3}}{2} \\ y = -\frac{1+i\sqrt{3}}{2} \end{cases}$	$H_x - L_3 - L_8$ $H_y - L_4 - L_5$	$A_2^{(1)}/E_6^{(1)}$
	$\begin{cases} xy - \frac{1}{14} \frac{(13i\sqrt{3} + 5)(4i\sqrt{3} + 7s - 8)x}{i\sqrt{3} + 19s - 23} \\ -\frac{1}{14} \frac{(13i\sqrt{3} + 5)(4i\sqrt{3} + 7s - 8)y}{i\sqrt{3} + 18s - 23} + \frac{1}{2} \frac{(5i\sqrt{3} - 1)(s + i\sqrt{3} - 5)}{i\sqrt{3} + 19s - 23} = 0 \end{cases}$	$H_x + H_y - L_1 - L_2 - L_6 - L_7$	
$\zeta_5(s)$	$\{p_5^*=0\}$	$-K_{X_s}$	$A_0^{(1)*}/E_8^{(1)}$

TABLE 3. Irreducible components of the singular fibres in the pencil associated to (3.13), their representative in the Picard groups, and the associated surface and symmetry type. The biquadratic polynomial p_5^* is given in Equation (3.23).