

Emergent quantum probability from full quantum dynamics and the role of energy conservation

Chen Wang

Department of Physics, Zhejiang Normal University, Jinhua, Zhejiang 321004, China

Jincheng Lu

*Jiangsu Key Laboratory of Micro and Nano Heat Fluid Flow Technology and Energy Application,
School of Physical Science and Technology, Suzhou University of Science and Technology, Suzhou, 215009, China*

Jian-Hua Jiang*

Suzhou Institute for Advanced Research, University of Science and Technology of China, Suzhou, 215123, China

(Dated: June 2, 2023)

We propose and study a toy model for the quantum measurements that yield the Born's rule of quantum probability. In this model, the electrons interact with local photon modes and the photon modes are dissipatively coupled with local photon reservoirs. We treat the interactions of the electrons and photons with full quantum mechanical description, while the dissipative dynamics of the photon modes are treated via the Lindblad master equation. By assigning double quantum dot setup for the electrons coupling with local photons and photonic reservoirs, we show that the Born's rule of quantum probability can emerge directly from microscopic quantum dynamics. We further discuss how the microscopic quantities such as the electron-photon couplings, detuning, and photon dissipation rate determine the quantum dynamics. Surprisingly, in the infinite long time measurement limit, the energy conservation already dictates the emergence of the Born's rule of quantum probability. For finite-time measurement, the local photon dissipation rate determines the characteristic time-scale for the completion of the measurement, while other microscopic quantities affect the measurement dynamics. Therefore, in genuine measurements, the measured probability is determined by both the local devices and the quantum mechanical wavefunction.

PACS numbers:

I. INTRODUCTION

The emergence of probability is one of the key features in quantum mechanics that has been puzzling since the birth of quantum mechanics [1, 2]. In 1926 Max Born proposed his theory where quantum states are interpreted as probability waves [3, 4]. This probability wave theory connects the probability in quantum measurements with the well-known quantum probability, i.e., the square of the absolute value of the wavefunction. This interpretation, though gives ease in using quantum theory to compute experimental observables, it never fully satisfied the questioning minds in the past century [5–8]. In particular, how could a wave be probabilistic when the wave equation (i.e., the Schrödinger equation) is deterministic? What if the connection between the quantum wavefunction and the measured probability is an emergent phenomenon, i.e., it emerges from the quantum dynamics of the measurement process instead of a direct connection as interpreted by Born?

In the past decades, it has been accepted that probability emerges in quantum mechanics due to the interaction between the quantum system and the measurement instrument [9, 10]. The instrument generally contains very large degrees of freedom and serves as an environment interacting with the quantum system. Such interaction causes quantum decoherence that leads to the loss of certainty and the emergence of quan-

tum probability [12–15]. Several models have been proposed to explore the underlying physics of quantum measurement [16], e.g., the Coleman-Hepp model [17], the Gaveau-Schulman model [18], and the Curie-Weiss model [19]. In particular, the Curie-Weiss model consisting of a spin 1/2 measured by coupled-spins interacting with one phonon reservoir, was used to elucidate the practical measurement in spin systems [16]. However, a simple microscopic model of quantum measurements which can demonstrate the emergence of Born's quantum probability from a full quantum mechanical description is still helpful in understanding the underlying physics of quantum measurements.

Here, we propose such a model by considering the detection of electron's probability in a double quantum dot setup via the local electron-photon interactions [20–23]. In this model, electrons interact with local photon modes and the photon modes dissipate their energy into local photonic reservoirs. We treat the electron-photon interaction in a fully quantum mechanical framework and describe the photon dissipation via the Lindblad master equation [26, 27]. We propose to measure the quantum system by monitoring the photonic energy dissipation. By solving the quantum dynamic equations, we find that in the infinite long time limit, the photonic energy dissipation into the reservoir is equal to the input electron energy. Since the latter is proportional to the square of the absolute value of the wavefunction, the energy conservation already dictates that the measured quantum probability is equal to the Born's rule of quantum probability. For finite time measurements, the local photon dissipation rate determines the characteristic timescale for the completion of the measurement, while other microscopic quantities affect the

*Electronic address: joejhjiang@hotmail.com




FIG. 1: (Color online) (a) The schematic illustration of the electron-photon model at Eq. (1). It is composed of two individual quantum dots, each owning one electron and interacting with single-mode photonic resonator. (b) The description of quantum probability readout from the quantum dots at Eq. (7) via the electron-photon interaction λ_v and the photon damping process κ_v . (c) The microscopic energy conservation process during the measurement process at Eq. (8), i.e., the energy loss from the electron-photon hybrid system is identical with the gain into the reservoir, which is characterized as the energy current J_v .

measurement dynamics. We further discuss how electron-photon detuning, strong electron-photon interaction, and finite reservoir temperature affect the measurement dynamics. Our work reveals the underlying microscopic quantum physics of quantum measurements and can serve as a prototype model to study the origin of the Born's rule on the quantum probability. Throughout this work, we set Planck constant $\hbar \equiv 1$.

II. MEASUREMENT SETUP

We begin by introducing a generic quantum system that consists of two individual quantum dots interacting with one photonic resonator [28–30], to describe the probability measurement of the electron wavefunction through the electron-photon interactions, where the photonic resonators may be treated as the probability detectors [31, 32]. Then, we analyze the dissipative dynamics of the electron-photon interacting model to practically read out information on the quantum probability of the electron wavefunction in each quantum dot. We note that the measurement description of the electron-photon interaction and photon dissipation is fully quantum.

A. Electron-photon Model

The electron-photon system is modeled as two quantum dots each owning single electron, separately coupled to two

photonic resonators, shown in Fig. 1(a). The Hamiltonian is described as $H_s = \sum_{v=l,r} H_{s,v}$. The v -th system component is specified as

$$H_{s,v} = \sum_{j=e,g} E_{j,v} |j_v\rangle \langle j_v| + \omega_{0,v} a_v^\dagger a_v + \lambda_v (a_v^\dagger |g_v\rangle \langle e_v| + a_v |e_v\rangle \langle g_v|), \quad (1)$$

where $|e_v\rangle$ and $|g_v\rangle$ are the excited and ground states with the corresponding occupation energies $E_{v,e}$ and $E_{v,g}$ in the v -th quantum dot, a_v^\dagger (a_v) is the creating (annihilating) operator of the photonic resonator with the photon energy $\omega_{0,v}$, λ_v denotes the electron-photon coupling strength. In the following, we set $E_{e,l} = E_{e,r} = 0$ without losing any generality.

Considering $H_{s,v}$ in the subspace $\{|n_v - 1, e_v\rangle, |n_v, g_v\rangle\}$ with n_v being the photon excitation number, it is straightforward to find that the n -th eigenvalues of the electron-photon model are given by $E_{n_v, \pm} = \omega_{0,v} n_v + \frac{E_{g,v} - \omega_{0,v}}{2} \pm \sqrt{(\frac{\omega_{0,v} + E_{g,v}}{2})^2 + \lambda_v^2 n_v}$, and the eigenvectors are given by

$$|\psi_{n_v, +}\rangle = \cos \frac{\theta_{n_v}}{2} |n_v, g_v\rangle + \sin \frac{\theta_{n_v}}{2} |n_v - 1, e_v\rangle, \quad (2a)$$

$$|\psi_{n_v, -}\rangle = -\sin \frac{\theta_{n_v}}{2} |n_v, g_v\rangle + \cos \frac{\theta_{n_v}}{2} |n_v - 1, e_v\rangle, \quad (2b)$$

with $\tan \theta_{n_v} = 2\lambda_v \sqrt{n_v} / (\omega_{0,v} + E_{g,v})$ ($n_v \geq 1$). Moreover, we define the effective ground state $|\psi_{0_v}\rangle = |0, g_v\rangle$ and the corresponding energy $E_{0,v} = E_{g,v}$. It need note that the real bare ground state actually is $|0, g_v\rangle$. However, this state will never take participate into the dissipative dynamics of the dot-photon system.

B. Dynamics

We consider master dynamics of the electron-photon model with weak photon dissipation [26, 27, 30]

$$\frac{d\rho_s}{dt} = -i[H_s, \rho_s] - \sum_{v=l,r} \frac{\kappa_v}{2} [a_v^\dagger a_v \rho_s + \rho_s a_v^\dagger a_v - 2a_v \rho_s a_v^\dagger], \quad (3)$$

where κ_v means the dissipation strength of photons from the v -th resonator into the corresponding zero-temperature environment (e.g., the v -th photon counter). Here, the dissipation of quantum dots is assumed negligible compared to the photon damping strength. Then, we select the initial state of the electron-photon model. There exists one electron in each quantum dot, i.e., $|\phi_e(0)\rangle = C_l |e_l, g_r\rangle + C_r |g_l, e_r\rangle$, and resonators are initially both in vacuum before the procedures of the probability readout. Hence, the initial wavefunction is generally described as

$$|\phi(0)\rangle = C_l |0_l, e_l\rangle \otimes |0_r, g_r\rangle + C_r |0_l, g_l\rangle \otimes |0_r, e_r\rangle, \quad (4)$$

where C_v is the wavefunction coefficient.

We focus on the individually measuring the quantum probability in the local dots [33, 34]. Consequently, the initial

reduced density matrix of the v -th system component is obtained as

$$\rho_v(0) = \mathbf{P}_v |0_v, e_v\rangle \langle 0_v, e_v| + \mathbf{P}_{\bar{v}} |0_v, g_v\rangle \langle 0_v, g_v|, \quad (5)$$

where the Born's quantum probabilities are expressed as

$$\mathbf{P}_v = |C_v|^2, \quad (6)$$

with $\bar{v} = r$ for $v = l$ and $\bar{v} = l$ for $v = r$. Hence, the dissipative dynamics of the v -th component is expressed as

$$\frac{d\rho_v}{dt} = -i[H_{s,v}, \rho_v] - \frac{\kappa_v}{2} [a_v^\dagger a_v \rho_v + \rho_v a_v^\dagger a_v - 2a_v \rho_v a_v^\dagger], \quad (7)$$

with $\rho_{l(r)} = \text{Tr}_{r(l)}\{\rho_s\}$, which is also exhibited in Fig. 1(b). Moreover, the dynamics is constraint to the lowest three eigenstates, i.e., $|\psi_{1_v,+}\rangle$, $|\psi_{1_v,-}\rangle$, and $|\psi_{0_v}\rangle$. And it is straightforward to show the inverse relations $|0_v, e_v\rangle = (\sin \frac{\theta_v}{2} |\psi_{1_v,+}\rangle + \cos \frac{\theta_v}{2} |\psi_{1_v,-}\rangle)$ and $|1_v, g_v\rangle = (\cos \frac{\theta_v}{2} |\psi_{1_v,+}\rangle - \sin \frac{\theta_v}{2} |\psi_{1_v,-}\rangle)$, with $\tan \theta_v = 2\lambda_v/(\omega_{0,v} + E_{g,v})$.

III. QUANTUM PROBABILITY MEASUREMENTS

In this section, we investigate the probability measurement at weak and strong electron-photon couplings and finite temperature reservoir, respectively.

A. Measurement with weak electron-photon coupling

We first study the probability measurement based on the quantum dots and photonic resonators in the long time evolution limit, i.e., complete measurement. Then, we further analyze the finite time behavior of the emitted photons, i.e., finite time measurement. Finally, we briefly discuss the influence of finite temperature reservoir on measured probability.

1. Complete measurement

We investigate the probability measurement from the energy perspective in the double quantum dot setup via the interaction with local photon resonators, with the initial state at Eq. (4). Specifically, the energy flow from the v -th system component into the dissipative reservoir (e.g., the v -th photon counter) can be defined as

$$J_v = -d\langle H_{s,v} \rangle(t)/dt, \quad (8)$$

which quantifies the energy loss of the v -th component, shown in Fig. 1(c). Meanwhile, in the eigenspace of $H_{s,v}$, the expression of the energy flow can be reexpressed as $J_v = -[E_{+,v}(dP_{v,+}(t)/dt) + E_{-,v}(dP_{v,-}(t)/dt)]$, with the density matrix elements denoting $P_{v,\pm}(t) = \langle \psi_{1_v,\pm} | \rho_v(t) | \psi_{1_v,\pm} \rangle$ and $P_{v,+}(t) = \langle \psi_{1_v,+} | \rho_v(t) | \psi_{1_v,-} \rangle$, and energies given by

$$E_{+(-),v} = [(\omega_{0,v} - E_{g,v}) \pm \sqrt{(\omega_{0,v} + E_{g,v})^2 + 4\lambda_v^2}]/2, \quad (9)$$

which characterizes the microscopic energy transport processes, e.g., $E_{+,v}[-dP_{v,+}(t)/dt]$ characterizing the relaxation of the system from the channel $|\psi_{1_v,+}\rangle$ to $|\psi_{0_v}\rangle$ by releasing the energy $E_{+,v}$ and one excitation particle. After sufficient long time evolution, it is straightforward to find that the total energy loss of the quantum system is obtained as $\int_0^\infty dt [-d\langle H_v \rangle(t)/dt] = -E_{g,v}|C_v|^2$. Simultaneously, the microscopic transferred energy to the reservoir is given by integrating the current

$$-\int_0^\infty dt \left[E_{+,v} \frac{dP_{v,+}(t)}{dt} + E_{-,v} \frac{dP_{v,-}(t)}{dt} \right] = -E_{g,v}|C_v|^2 \quad (10)$$

This clearly demonstrates the microscopic energy conservation, i.e., the energy of the electron in the v -th dot is fully dissipated into the corresponding reservoir via two eigenchannels.

Furthermore, it is interesting to find that Based on the master equation, the energy flow at Eq. (8) can be reexpressed as

$$J_v = E_{+,v}\kappa_v \left[\cos^2 \frac{\theta_v}{2} P_{v,+}(t) - \cos \frac{\theta_v}{2} \sin \frac{\theta_v}{2} \text{Re}\{P_{v,+}\} \right] + E_{-,v}\kappa_v \left[\sin^2 \frac{\theta_v}{2} P_{v,-}(t) - \cos \frac{\theta_v}{2} \sin \frac{\theta_v}{2} \text{Re}\{P_{v,+}\} \right]. \quad (11)$$

Both the populations and coherence terms contribute to the energy flow. Moreover, the dynamics of the photon number in the v -th resonator is described as

$$\begin{aligned} \langle a_v^\dagger a_v \rangle(t) &= \cos^2 \frac{\theta_v}{2} P_{v,+}(t) + \sin^2 \frac{\theta_v}{2} P_{v,-}(t) \quad (12) \\ &\quad - 2 \cos \frac{\theta_v}{2} \sin \frac{\theta_v}{2} \text{Re}[P_{v,+}(t)]. \end{aligned}$$

Then, we consider the dynamical equation at Eq. (7) and integrate the photonic energy flow $\int_0^\infty J_v dt$ at Eq. (11), which is connected with the countered photon numbers (see Appendix for the details). By defining the measured quantum probability

$$\mathcal{P}_v(t) \equiv \kappa_v \int_0^\infty dt \langle a_v^\dagger a_v \rangle(t), \quad (13)$$

the complete quantum probability measurement inherited from microscopic energy conservation is established as

$$\mathcal{P}_v(\infty) = \mathbf{P}_v, \quad (14)$$

with $\mathbf{P}_v = |C_v|^2$. This relation denotes that by practically counting the emitted photon number from the individual resonator to the external sink (i.e., the measured probability), we are able to correctly detect the quantum probability of the electron wavefunction in each dot, once the dissipation rate κ_v is available. And the measured probability is irrelevant with the electron-photon system parameters, e.g., $\omega_{0,v}$, $E_{g,v}$, and λ_v . Therefore, we conclude that the microscopic energy conservation is able to yield the complete probability measurement.

Alternatively, from the particle conservation perspective, i.e., from the excitation particles in the hybrid system to

detected photon particles via the photon dissipation processes, we find the dynamics of the ground state population ($P_{v,0}(t) = \langle \psi_{0v} | \rho_v(t) | \psi_{0v} \rangle$) is tightly related with the photon flow, i.e., $dP_{v,0}(t)/dt = \kappa_v \langle a_v^\dagger a_v \rangle(t)$. This demonstrates that the probability of electrons transport into the ground state can be read out via the emitted photon flows $\kappa_v \langle a_v^\dagger a_v \rangle(t)$. After the long time evolution the complete probability measurement at Eq. (14) can be fully recovered, considering $P_{v,0}(0) = |C_{\bar{v}}|^2$ and $P_{v,0}(\infty) = 1$.

2. Finite time measurement

Here, we study the probability measurement at finite time dynamics. Since the complete probability measurement holds individually for each electron-photon interacting component, we consider the initial state at Eq. (4). It should note that even for such a concise reduced density matrix Eq. (5), the general solution of dissipative dynamics is hard to obtain, due the coupling between diagonal and off-diagonal elements of the electron-photon system. While if we include the secular approximation, the equations of motion of system density matrix are reduced to $dP_{v,+}/dt \approx -\kappa_v \cos^2 \frac{\theta_v}{2} P_{v,+}$, $dP_{v,-}/dt \approx -\kappa_v \sin^2 \frac{\theta_v}{2} P_{v,-}$, and $dP_{v,+ -}/dt \approx -(i\Lambda_v P_{v,+ -} + \frac{\kappa_v}{2} P_{v,+ -})$, with the energy gap $\Lambda_v = \sqrt{(\omega_{0,v} + E_{g,v})^2 + 4\lambda_v^2}$ between $|\psi_{1v,+}\rangle$ and $|\psi_{1v,-}\rangle$. The dynamics of density matrix elements are given by $P_{v,+}(t) \approx |C_v|^2 \exp(-\kappa_v \cos^2 \frac{\theta_v}{2} t) \sin^2 \frac{\theta_v}{2}$, $P_{v,-}(t) \approx |C_v|^2 \exp(-\kappa_v \sin^2 \frac{\theta_v}{2} t) \cos^2 \frac{\theta_v}{2}$, $P_{v,+ -}(t) \approx |C_v|^2 \exp[-(iE_{v,+ -} + \kappa_v/2)t] \cos \frac{\theta_v}{2} \sin \frac{\theta_v}{2}$. Then, the accumulated photonic energy base on Eq. (11) into the dissipative reservoir is approximated as

$$\begin{aligned} \int_0^t J_v(s) ds &\approx |C_v|^2 \left[E_{+,v} \sin^2 \frac{\theta_v}{2} (1 - e^{-(\cos^2 \frac{\theta_v}{2}) \kappa_v t}) \right. \\ &\quad + E_{-,v} \cos^2 \frac{\theta_v}{2} (1 - e^{-(\sin^2 \frac{\theta_v}{2}) \kappa_v t}) \\ &\quad - \kappa_v \frac{\sin^2 \theta_v}{2} (E_{+,v} + E_{-,v}) \\ &\quad \left. \times \text{Re} \left\{ \frac{1 - e^{-(i\Lambda_v + \frac{\kappa_v}{2})t}}{i\Lambda_v + \frac{\kappa_v}{2}} \right\} \right]. \end{aligned} \quad (15)$$

After long time evolution $t \rightarrow \infty$, the accumulated photonic energy is given by $\int_0^\infty J_v(s) ds \approx |C_v|^2 (E_{+,v} \sin^2 \frac{\theta_v}{2} + E_{-,v} \cos^2 \frac{\theta_v}{2}) \approx -E_{g,v} |C_v|^2$ under the condition $\lambda_v \gg \kappa_v$, which approaches the total energy dissipated to the reservoir.

Similarly, the measured quantum probability, i.e., the detected photon numbers at Eq. (13), is generally given by

$$\begin{aligned} \mathcal{P}_v(t) &\approx |C_v|^2 \frac{\sin^2 \theta_v}{4} \left[\frac{1 - e^{-(\cos^2 \frac{\theta_v}{2}) \kappa_v t}}{\cos^2 \frac{\theta_v}{2}} \right. \\ &\quad \left. + \frac{1 - e^{-(\sin^2 \frac{\theta_v}{2}) \kappa_v t}}{\sin^2 \frac{\theta_v}{2}} - 2\kappa_v \text{Re} \left\{ \frac{1 - e^{-(i\Lambda_v + \frac{\kappa_v}{2})t}}{\frac{\kappa_v}{2} + i\Lambda_v} \right\} \right], \end{aligned}$$

In the limit $t \rightarrow \infty$, we find that $\kappa_v \int_0^\infty \langle a_v^\dagger a_v \rangle(s) ds \approx [1 -$




FIG. 2: (Color online) The finite time dynamics of (a) photon numbers and (b) energy flow with various quantum dot energies. The initial state is selected as $|\phi(0)\rangle = |0_l, e_l\rangle \otimes |0_r, g_r\rangle$. The system parameters are given by $\omega_0 = 1$, $E_e = 0$, $\lambda = 0.01$, and $\kappa = 0.001$.

$\frac{\cos^2 \frac{\theta_v}{2} \sin^2 \frac{\theta_v}{2} \kappa_v^2}{(\kappa_v/2)^2 + E_{v,+ -}^2}$. In the coupling regime comparatively stronger than the dissipation strength $\lambda_v \gg \kappa_v$, we obtain $\kappa_v \int_0^\infty \langle a_v^\dagger a_v \rangle(s) ds \approx |C_v|^2$, which consistently recovers the complete probability measurement. For convenience, we select system parameters as $\omega_{0,v} = \omega_0$, $E_{g,v} = E_g$, $\lambda_v = \lambda$, and the dissipative strength $\kappa_v = \kappa$.

Then, we plot Fig. 2 to study the finite time dynamics of the dissipated energy and the photon numbers with $C_l = 1$ and $C_r = 0$. Thus, only the l -th part component contributes to the dissipative dynamics. In particular at resonance ($E_g = -\omega_0$), it is found that the average photon number and the photonic energy flow are reduced to

$$\langle a_l^\dagger a_l \rangle(t) \approx \exp(-\kappa t/2) (1 - \cos 2\lambda t)/2, \quad (16a)$$

$$J_l(t) \approx \kappa (-E_g) \exp(-\kappa t/2) (1 - \cos 2\lambda t)/2, \quad (16b)$$

which both exhibits periodic damping oscillation behavior with the period $T = \pi/\lambda$. Accordingly, the measured quantum probability and dissipated energy show intermittent increase. In particular, the n -th ‘‘quasi-step’’ [35] with the amplitude

$$\mathcal{P}(nT) \approx [1 - e^{-n\pi\kappa/2\lambda}], \quad (17a)$$

$$\frac{1}{-E_g} \int_0^{nT} J_l(s) ds \approx [1 - e^{-n\pi\kappa/2\lambda}], \quad (17b)$$




FIG. 3: (Color online) The finite time dynamics of photon number and accumulation with various photon dissipation strengths. The initial state is the same as Fig. 2. The system parameters are given by $\omega_0 = 1$, $E_e = 0$, $E_g = -1$, and $\lambda = 0.01$.

are shown around the time $t = nT$, where the photon counter detects the negligible photon flow. From Eqs. (17a) and (17b) it is known that the characteristic measurement number of times can be quantified as $n_c \approx 2\lambda/\kappa$ and the corresponding time is given by $t_c = n_c T \approx 2\pi/\kappa$. Then the measurement outcome of emitted photons number and energy gradually becomes complete, which approaches the complete measurement results. While at a given finite measurement time, the effectiveness and systematic error of quantum probability measurement generally rely on the specific properties of the measurement devices, which are typically characterized as λ and κ .

Moreover, we analyze the influence of the bias condition ($E_g \neq -\omega_0$) on the dissipative dynamics of photons. Both the photon numbers and energy flow are monotonically suppressed with the increase of energy bias $|\omega_0 + E_g|$ (though not shown here, such monotonic suppression also persists in the regime $(-E_g > \omega_0)$). Hence, the fast detection of the complete quantum probability and the underlying conservation favors the resonant condition.

We also study the effect of the photon dissipation on the dynamics of photons numbers at resonance in Fig. 3, considering the analogous behaviors of energy flow as shown in Eqs. (16b) and (17b). For the average photon number, it is shown that the profile is dramatically suppressed with the increase of the dissipation strength κ , whereas the oscillating period is robust against the dissipation process. This is consistent with Eq. (16a) and quantified by the characteristic time $2\pi/\kappa$. Accordingly, the emitted photon numbers exhibit fast accumulation in the photon counter. Therefore, we conclude that enhancing the photon dissipation is helpful for the effi-




FIG. 4: (Color online) The finite time dynamics of photon number with strong electron-photon interaction (a) at $\lambda = 0.5\omega_0$ and (b) by tuning coupling strength λ . The inset in (b) denotes the photon number as a function of electron-photon coupling strength, where the circles and dashed lines are calculated from full numerical calculation at the time $\kappa t = 10$ and $(\lambda/\omega_0)^2$, respectively. The initial state is the same as Fig. 2. The system parameters are given by $\omega_0 = 1$, $E_e = 0$, and $E_g = -1$.

cient measurement of the emitted photons and energy.

B. Measurement with strong electron-photon coupling

When the electron-photon coupling becomes strong and even comparable with the bare frequency of the optical resonator, the counter-rotating terms of the electron-photon interaction should be considered [36–38]. The system then becomes analogous with the seminal quantum Rabi model [39–41], is modified to

$$H_{s,v} = \sum_{j=e,g} E_{j,v} |j_v\rangle \langle j_v| + \omega_{0,v} a_v^\dagger a_v + \lambda_v (a_v^\dagger + a_v) (|g_v\rangle \langle e_v| + |e_v\rangle \langle g_v|), \quad (18)$$

Due to the inclusion of the counter-rotating terms $(a_v^\dagger |e_v\rangle \langle g_v| + a_v |g_v\rangle \langle e_v|)$, the particle conservation breaks down, whereas the energy conservation is still valid. Accordingly, we find that the analytical eigensolution of $H_{s,v}$ is hard to obtain until now, except the solution mapped to the special function via Bargmann algebra [39] and photonic coherent states [42]. Here, we include the master equation Eq. (3) to

study the influence of strong electron-photon interaction on probability measurement.

With the initial state $|\phi(0)\rangle = |0_l, e_l\rangle \otimes |0_r, g_r\rangle$, we first plot Fig. 4(a) to study the short-time dissipative dynamics of average photon number $\langle a_l^\dagger a_l \rangle(t)$ at strong electron-photon coupling, e.g., $\lambda = 0.5\omega_0$, by comparing with the counterpart based on the system Hamiltonian at Eq. (1). It is interesting to find that the photon number shows the aperiodic oscillation due to the effect of counter-rotating terms, which is significantly distinct from the one with the electron-photon interaction ($a_l^\dagger |g_l\rangle \langle e_l| + a_l |e_l\rangle \langle g_l|$). Furthermore, the peak of photon number breaches the unity, which is unavailable in absence of counter-rotating terms.

Next, we study the photon number dynamics at long time evolution in Fig. 4(b). It is shown that by increasing the electron-photon coupling strength, the aperiodicity degree becomes increasingly apparent. Moreover, it is interesting to find that after long time evolution the photon number becomes nonvanishing. This directly demonstrates the breakdown of the complete measurement at Eq. (14) with weak electron-photon interaction. We plot the inset of Fig. 4(b) to quantify the robustness of the photon number. $\langle a_l^\dagger a_l \rangle(t \rightarrow \infty)$ becomes monotonic enhancement with the increase of λ .

As is known that after long time evolution the system may reach the steady state, which generally may not be the eigenstate, e.g., the ground state at zero temperature. In particular for rather strong electron-photon coupling, by considering the degenerate perturbation theory, the eigenvectors of $H_{s,l}$ at Eq. (18) are estimated as

$$|\psi_{n,\pm}\rangle \approx \frac{1}{\sqrt{2}} \left(\exp\left[-\frac{\lambda}{\omega_0}(a_l^\dagger - a_l)\right] \frac{(a_l^\dagger)^n}{\sqrt{n!}} |0\rangle_a \otimes |-\rangle \right. \\ \left. \pm \exp\left[\frac{\lambda}{\omega_0}(a_l^\dagger - a_l)\right] \frac{(a_l^\dagger)^n}{\sqrt{n!}} |0\rangle_a \otimes |+\rangle \right), \quad (19)$$

with $|\pm\rangle = (|e_l\rangle \pm |g_l\rangle)/\sqrt{2}$ and $a_l|0\rangle_a = 0$. And the corresponding eigenvalues become $E_{n,\pm} \approx \omega_0 n \mp |D_{nn}(2\lambda/\omega_0)|^2 E_g/2 - \lambda^2/\omega_0 + E_g/2$, with the coherent state overlap coefficient $D_{nn} = e^{-2(\lambda/\omega_0)^2} \sum_{r=0}^n (-1)^r n! (2\lambda/\omega_0)^{2n-2r} / [(n-r)!(n-r)!r!]$ [43]. Then, the transition rates between two eigenstate populations via the master equation could be reestablished. Specifically, by including the secular approximation the population dynamics is given by $dP_{n,\pm}/dt \approx [\kappa(n+1)P_{n+1,\pm} - \kappa n P_{n,\pm}] + \kappa(\lambda/\omega_0)^2 (P_{n,\mp} - P_{n,\pm})$, with the density matrix elements $P_{n,\pm} = \langle \psi_{n,\pm} | \rho_l | \psi_{n,\pm} \rangle$ and the transition coefficients $\langle \psi_{n-1,\pm} | a_l | \psi_{n,\pm} \rangle = \sqrt{n}$ and $\langle \psi_{n-1,\pm} | a_l | \psi_{n,\mp} \rangle = 0$. The dissipative dynamics generally results in the steady state

$$\rho_l(\infty) \approx \frac{1}{2} (|\psi_{0,+}\rangle \langle \psi_{0,+}| + |\psi_{0,-}\rangle \langle \psi_{0,-}|). \quad (20)$$

Thus, the corresponding steady-state photon number is given by $\langle a_l^\dagger a_l \rangle(t \rightarrow \infty) \approx (\lambda/\omega_0)^2$. This partially explain the non-vanishing steady-state value of the average photon number at strong electron-photon coupling in the inset of Fig. 4(b). Therefore, we conclude that strong electron-photon interaction generally breaks the microscopic particle conservation and complete measurement.

C. Finite temperature reservoir

We study the effect of finite-temperature reservoir on the quantum measurement of probability. At weak electron-photon coupling $\lambda_v/\omega_{0,v} \ll 1$, the master equation for dissipative dynamics of the electron-photon model can be described as

$$\frac{d\rho_s}{dt} = -i \sum_{v=l,r} [H_{s,v}, \rho_s] \quad (21) \\ + \sum_{v=l,r} \kappa_v (1 + \bar{n}_v) \mathcal{L}[a_v] \rho_s + \sum_{v=l,r} \kappa_v \bar{n}_v \mathcal{L}[a_v^\dagger] \rho_s,$$

where the dissipator denotes $\mathcal{L}[a_v] \rho_s = (2a_v \rho_s a_v^\dagger - a_v^\dagger a_v \rho_s - \rho_s a_v^\dagger a_v)/2$. Here, we empirically describe the incoherent coupling between the photon resonator and the bosonic reservoir at the temperature T_v with the Bose-Einstein distribution function $\bar{n}_v = 1/[\exp(\omega_{0,v}/k_B T_v) - 1]$ [44, 45]. We select the initial state of the reduced $H_{s,v}$ as

$$\rho_v(0) = (|C_v|^2 |e_v\rangle \langle e_v| + |C_{\bar{v}}|^2 |g_v\rangle \langle g_v|) \quad (22) \\ \otimes \frac{e^{-\beta_v \omega_{0,v} a_v^\dagger a_v}}{\mathcal{Z}_v},$$

with the partition function $\mathcal{Z}_v = 1/[1 - \exp(-\beta_v \omega_{0,v})]$ and $\beta_v = 1/k_B T_v$.

We investigate the energy flow into the v -th reservoir. At low temperature regime, it is numerically found that the dissipative dynamics of the flow is typically contributed by three lowest eigenstates, i.e., $|\psi_{1_v,\pm}\rangle$ and $|\psi_{0_v}\rangle$. Accordingly, relation between the energy flow and the photon number can be expressed as

$$J_v \approx \frac{(\omega_{0,v} - E_{g,v})}{2} [\kappa_v (1 + \bar{n}_v) \langle a_v^\dagger a_v \rangle - \kappa_v \bar{n}_v P_{0,v}] \\ + \frac{\Lambda_v}{2} \frac{d(P_{-,v} - P_{+,v})}{dt}, \quad (23)$$

with $P_{\pm,v} = \langle \psi_{1_v,\pm} | \rho_v | \psi_{1_v,\pm} \rangle$, $P_{0,v} = \langle \psi_{0_v} | \rho_v | \psi_{0_v} \rangle$, and $\Lambda_v = \sqrt{(\omega_{0,v} + E_{g,v})^2 + 4\lambda_v^2}$. Moreover, the total energy loss of the quantum system is obtained as $\int_0^\infty J_v dt = [\langle H_{s,v} \rangle(0) - \langle H_{s,v} \rangle(\infty)]$, which is approximated as $\int_0^\infty J_v dt \approx E_{g,v} (1/[1 + \exp(-\beta_v E_{g,v})] - |C_v|^2)$. Hence, based on the microscopic energy conservation law, we obtain the effective probability measurement by integrating the current at Eq. (23) as

$$\kappa_v \int_0^\infty dt [\langle a_v^\dagger a_v \rangle(t) - \frac{\bar{n}_v P_{0,v}(t)}{1 + \bar{n}_v}] \approx \frac{|C_v|^2 - 1/\mathcal{Z}'_v}{(1 + \bar{n}_v)^2}, \quad (24)$$

with $\mathcal{Z}'_v = (1 + e^{-\beta_v E_{g,v}})$. If the temperature approaches zero, i.e., $\bar{n}_v = 0$, the effective relation is naturally reduced to the complete probability measurement relation, i.e., $\kappa_v \int_0^\infty dt \langle a_v^\dagger a_v \rangle(t) = |C_v|^2$ at Eq. (14).

Alternatively, the dynamical equation of the ground state probability is described as

$$\frac{d}{dt} P_{0,v} \approx \kappa(1 + \bar{n}_v) \langle a_v^\dagger a_v \rangle - \kappa_v \bar{n}_v P_{0,v}, \quad (25)$$

in which the change of the probability is contributed by the photon flow into the reservoir and the backward flow from the reservoir thermal noise. Straightforwardly, the effective probability measurement at Eq. (24) can be recovered from Eq. (25) by considering $P_{0,v}(0) \approx |C_v|^2 / \mathcal{Z}_v$ and $P_{0,v}(\infty) \approx 1 / \mathcal{Z}_v \mathcal{Z}'_v$.

IV. CONCLUSION

In summary, we investigate a practical measurement of the quantum probability and its underlying microscopic conservation laws through quantum electron-photon interaction and quantum photon dissipation, connecting the quantum probability with the probabilistic Born's rule. We model the measurement setup with two individual quantum dots, each with an electron occupation interacting with a single-mode photonic resonator, which is ultimately dissipated to a reservoir, e.g., a photon counter.

For the quantum measurement with weak electron-photon coupling, we first study the complete probability measurement at Eq. (14), which exactly connects the quantum probability of the electron wavefunction in the dot with the counted photon numbers. The measurement rule is protected by the microscopic energy conservation law with electron-photon energy exchange processes. Such rule can be alternatively recovered under the microscopic particle conservation law. This complete probability measurement relation is impervious to the quantum system parameters, such as $E_{e(g),v}$, $\omega_{0,v}$, and λ_v . Next, we study the finite time probability measurement at finite time dynamics. Under the secular approximation, at resonance ($E_{e,v} - E_{g,v} = \omega_{0,v}$) the photon numbers at Eq. (16a) exhibit the periodic oscillating damping behavior, which are determined by the electron-photon coupling and photon dissipation strengths. The emitted photon number shows intermittent increase feature at Eq. (17a), with quasi-steps appearing around the times $t_v = n\pi / \lambda_v$. The characteristic measurement time is quantified as $t_{c,v} \approx 2\pi / \kappa_v$, where the measurement outcome of emitted photons numbers and energy becomes effectively complete. Furthermore, it is found that the bias condition ($(E_{e,v} - E_{g,v}) \neq \omega_{0,v}$) blocks the efficient measurement of the quantum probability.

We further analyze the applicability of quantum measurement in the strongly-coupled electron-photon model and consider the counter-rotating terms in $H_{s,v}$ Eq. (18), which straightforwardly breaks the particle number conservation. In short-time dynamics, by comparing the photon number dynamics with the counterpart in absence of counter-rotating terms, we observe two distinct features: (1) the photon number exhibits the aperiodic oscillations, and (2) the peak of photon number breaches the unity. While in long time dynamics, it is found that (1) the aperiodicity degree of photon number oscillating behavior becomes apparent with the increase of electron-photon coupling, and (2) the photon number still becomes nonvanishing, due to the formation of the mixture of photonic coherent states, which demonstrates the breakdown of the complete probability measurement relation.

Moreover, we investigate the influence of finite temperature reservoir on the quantum measurement and explore the effective

measurement relation connecting the counted photon flux with quantum probability and thermal statistics.

We hope that the reported measurements of quantum probability and the microscopic conservation laws may provide general avenues for practically detecting quantum probability.

V. ACKNOWLEDGEMENTS

This work was supported by the funding for the National Natural Science Foundation of China under Grants No. 12125504, No. 12074281, and No. 11704093, Jiangsu Key Disciplines of the Fourteenth Five-Year Plan (Grant No. 2021135), and the Opening Project of Shanghai Key Laboratory of Special Artificial Microstructure Materials and Technology.

Appendix: Derivation of complete quantum probability measurement relation

In the eigenbasis of $H_{s,v}$, the energy current $J_v = -d\langle H_{s,v} \rangle / dt$ is expressed as

$$J_v = -E_{+,v} \frac{dP_{v,+}(t)}{dt} - E_{-,v} \frac{dP_{v,-}(t)}{dt}, \quad (\text{A.1})$$

with the eigen-energies $E_{\pm,v}$ at Eq. (9). Including the dynamics of density matrix operator in the eigenbasis of $H_{s,v}$

$$\frac{d}{dt} P_{v,+} = -\kappa \cos^2 \frac{\theta}{2} P_{v,+} + \frac{\kappa}{2} \sin \theta \text{Re}[P_{v,+}], \quad (\text{A.2a})$$

$$\frac{d}{dt} P_{v,-} = -\kappa \sin^2 \frac{\theta}{2} P_{v,-} + \frac{\kappa}{2} \sin \theta \text{Re}[P_{v,+}], \quad (\text{A.2b})$$

$$\frac{d}{dt} P_{v,+} = -(i\Lambda_v + \frac{\kappa}{2}) P_{v,+} + \frac{\kappa}{2} \sin \theta \text{Re}[P_{v,+}], \quad (\text{A.2c})$$

with $\Lambda_v = E_{+,v} - E_{-,v}$ and $\text{Re}[P_{v,+}] = (P_{v,+} + P_{v,-})/2$, the current is specified as

$$J_v = \frac{\kappa}{2} (2 \cos^2 \frac{\theta}{2} P_{v,+} - \sin \theta \text{Re}[P_{v,+}]) + \frac{\kappa}{2} (2 \sin^2 \frac{\theta}{2} P_{v,-} - \sin \theta \text{Re}[P_{v,+}]). \quad (\text{A.3})$$

Considering the expression of photon number $\langle a_l^\dagger a_l \rangle$ at Eq. (12), the relation between the current and the photon number can be established as

$$J_v = \frac{\kappa}{2} (\omega_{0,v} - E_{g,v}) \langle a_l^\dagger a_l \rangle - \frac{\Lambda_v}{2} \left(\frac{d}{dt} P_{v,+} - \frac{d}{dt} P_{v,-} \right). \quad (\text{A.4})$$

By integrating Eq. (A.4) at both sides, we find $\int_0^\infty J_v dt = -E_{g,v} |C_v|^2$, $P_{v,-}(0) - P_{v,+}(0) = |C_v|^2 \cos \theta$, and $P_{v,-}(\infty) - P_{v,+}(\infty) = 0$. Finally, we obtain the complete quantum measurement relation at Eq. (14).

-
- [1] S. Weinberg, *Ann. Phys. (NY)* **194**, 336 (1989).
- [2] S. Weinberg, *Phys. Rev. Lett.* **62**, 485 (1989).
- [3] M. Born, *Z. Phys.* **38**, 803 (1926).
- [4] M. Born, *The statistical interpretation of quantum mechanics*, Nobel Lecture, (1954).
- [5] U. Sinha, C. Couteau, T. Jennewein, R. Laflamme, and G. Weihs, *Science* **329**, 418 (2010).
- [6] A. Bassi, K. Lochan, S. Satin, T. P. Singh, and H. Ulbricht, *Rev. Mod. Phys.* **85**, 471 (2013).
- [7] P. Shadbolt, J. C. F. Mathews, A. Laing, and Jeremy L. O'Brien, *Nature Physics* **10**, 278 (2014).
- [8] M. O. Pleinert, A. Rueda, E. Lutz, and J. von Zanthier, *Phys. Rev. Lett.* **126**, 190401 (2021).
- [9] J. A. Wheeler and W. H. Zurek, *Quantum Theory and Measurement* (Princeton University Press, 2014).
- [10] H. M. Wiseman and G. J. Milburn, *Quantum Measurement and Control* (Cambridge University Press, Cambridge, 2011).
- [11] A. A. Clerk, M. H. Devoret, S. M. Girvin, Florian Marquardt, and R. J. Schoelkopf, *Rev. Mod. Phys.* **82**, 1155 (2010).
- [12] W. H. Zurek, *Phys. Rev. Lett.* **90**, 120404 (2003).
- [13] W. H. Zurek, *Phys. Rev. A* **71**, 052105 (2005).
- [14] M. Schlosshauer and A. Fine, *Foundations of Physics* **35**, 197 (2005).
- [15] U. Mohrhoff, *Int. J. Quantum Inform.* **2**, 221 (2004).
- [16] A. E. Allahverdyan, R. Balianb, and T. M. Nieuwenhuizen, *Phys. Rep.* **525**, 1 (2013).
- [17] J. S. Bell, *On wave packet reduction in the Coleman-Hepp model. In John S Bell On The Foundations Of Quantum Mechanics* (World Scientific, Singapore, 2001).
- [18] B. Gaveau and L. Schulman, *J. Stat. Phys.* **58**, 1209 (1990).
- [19] A. E. Allahverdyan, R. Balian, and Th. M. Nieuwenhuizen, *Europhys. Lett.* **61**, 452 (2003).
- [20] M. R. Delbecq, V. Schmitt, F. D. Parmentier, N. Roch, J. J. Viennot, G. Feve, B. Huard, C. Mora, A. Cottet, and T. Kontos, *Phys. Rev. Lett.* **107**, 256804 (2011).
- [21] Y. Y. Liu, K. D. Petersson, J. Stehlik, J. M. Taylor, and J. R. Petta, *Phys. Rev. Lett.* **113**, 036801 (2014).
- [22] X. Mi, J. V. Cady, D. M. Zajac, P. W. Deelman, and J. R. Petta, *Science* **355**, 156 (2017).
- [23] T. R. Hartke, Y.-Y. Liu, M. J. Gullans, and J. R. Petta, *Phys. Rev. Lett.* **120**, 097701 (2018).
- [24] J. H. Jiang and S. John, *Phys. Rev. X* **4**, 031025 (2014).
- [25] R. Q. Wang, C. Wang, J. C. Lu, and J. H. Jiang, *Advances in Physics:X* **7**, 2082317 (2022).
- [26] H. J. Carmichael, *Statistical Methods in Quantum Optics 1: Master Equations and Fokker-Planck Equations* (Springer-Verlag, New York, 1999).
- [27] U. Weiss, *Quantum Dissipative Systems* (World Scientific, Singapore, 2012).
- [28] S. Y. Zhu, Z. D. Liu, and X. S. Li, *Phys. Lett. A* **128**, 89 (1988).
- [29] V. V. Albert, *Phys. Rev. Lett.* **108**, 180401 (2012).
- [30] P. Stegman, S. N. Gupta, G. Haran, and J. S. Cao, *ACS Photonics* **9**, 2119 (2022).
- [31] R. Lescanne, S. Deleglise, E. Albertinale, U. Reglade, T. Capelle, E. Ivanov, T. Jacqmin, Z. Leghtas, and E. Flurin, *Phys. Rev. X* **10**, 021038 (2020).
- [32] E. Albertinale, L. Balembois, E. Billaud, V. Ranjan, D. Flanagan, T. Schenkel, D. Esteve, D. Vion, P. Bertet, and E. Flurin, *Nature* **600**, 434 (2021).
- [33] M. A. Nielsen and I. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, 2010).
- [34] J. Cotler and F. Wilczek, *Phys. Rev. Lett.* **124**, 100401 (2020).
- [35] Actually, there exists slight increase around each quasi-step (not exact step) of countered photon numbers, which is contributed by negligible photon flow around nT .
- [36] M. Kulkarni, O. Cotlet, and H. E. Tureci, *Phys. Rev. B* **90**, 125402 (2014).
- [37] B. K. Agarwalla, M. Kulkarni, S. Mukamel, and D. Segal, *Phys. Rev. B* **94**, 121305(R) (2016).
- [38] J. C. Lu, R. Q. Wang, J. Ren, M. Kulkarni, and J. H. Jiang, *Phys. Rev. B* **99**, 035129 (2019).
- [39] D. Braak, *Phys. Rev. Lett.* **107**, 100401 (2011).
- [40] A. F. Kockum, A. Miranowicz, S. De Liberato, S. Savasta, and F. Nori, *Nature Reviews Physics* **1**, 19 (2019).
- [41] P. Forn-Diaz, L. Lamata, E. Rico, J. Kono, and E. Solano, *Rev. Mod. Phys.* **91**, 025005 (2019).
- [42] Q. H. Chen, C. Wang, S. He, T. Liu, and K. L. Wang, *Phys. Rev. A* **86**, 023822 (2012).
- [43] Q. H. Chen, Y. Y. Zhang, T. Liu, and K. L. Wang, *Phys. Rev. A* **78**, 051801(R) (2008).
- [44] J. M. Fink, L. Steffen, P. Studer, Lev S. Bishop, M. Baur, R. Bianchetti, D. Bozyigit, C. Lang, S. Philipp, P. J. Leek, and A. Wallraff, *Phys. Rev. Lett.* **105**, 163601 (2010).
- [45] T. Bonsen, P. Harvey-Collard, M. Russ, J. Dijkema, A. Sammak, G. Scappucci, and L. M. K. Vandersypen, *Phys. Rev. Lett.* **130**, 137001 (2023).