### Riemannian curvature identities on almost Calabi-Yau with torsion 6-manifold and generalized Ricci solitons

S. Ivanov<sup>1</sup> and N. Stanchev<sup>2</sup>

<sup>1</sup> University of Sofia, Faculty of Mathematics and Informatics, blvd. James Bourchier 5, 1164, Sofia, Bulgaria and Institute of Mathematics and Informatics, Bulgarian Academy of Sciences email: ivanovsp@fmi.uni-sofia.bg

> <sup>2</sup> University of Sofia, Faculty of Mathematics and Informatics, blvd. James Bourchier 5, 1164, Sofia, Bulgaria

#### Abstract

It is observed that on a compact almost complex Calabi-Yau with torsion 6-manifold the Nijenhuis tensor is parallel with respect to the torsion connection. If the torsion is closed then the space is a compact generalized gradient Ricci soliton. In this case, the torsion connection is Ricci-flat if and only if either the norm of the torsion or the Riemannian scalar curvature is constant. On a compact almost complex Calabi-Yau with torsion 6-manifold it is shown that the curvature of the torsion connection is symmetric on exchange of the first and the second pairs and has vanishing Ricci tensor if and only if it satisfies the Riemannian first Bianchi identity.

Keywords: torsion connection, SU(3) holonomy, almost Calabi-Yau with torsion, Riemannian Bianchi identities, generalized gradient Ricci solitons.

AMS MSC2020: 53C55, 53C21, 53C29, 53Z05

#### Acknowledgements:

We would like to thank Jeffrey Streets and Ilka Agricola for the extremely useful remarks, comments and suggestions.

The research of S.I. is partially supported by Contract KP-06-H72-1/05.12.2023 with the National Science Fund of Bulgaria, by Contract 80-10-192 / 17.5.2023 with the Sofia University "St.Kl.Ohridski" and the National Science Fund of Bulgaria, National Scientific Program "VIHREN", Project KP-06-DV-7. The research of N.S. is partially financed by the European Union-Next Generation EU, through the National Recovery and Resilience Plan of the Republic of Bulgaria, project SUMMIT BG-RRP-2.004- 0008-C01.

Statements and Declarations: not applicable

## Contents



<span id="page-1-1"></span>

# <span id="page-1-0"></span>1 Introduction

Riemannian manifolds with metric connections having totally skew-symmetric torsion and special holonomy received a lot of interest in mathematics and theoretical physics mainly from supersymmetric string theories and supergravity. The main reason comes from the Hull-Strominger system which describes the supersymmetric background in heterotic string theories [\[86,](#page-26-0) [53\]](#page-24-0). The number of preserved supersymmetries depends on the number of parallel spinors with respect to a metric connection  $\nabla$  with totally skew-symmetric torsion T. The torsion 3-form T is identified with the 3-form field strength in these theories. The presence of a  $\nabla$ −parallel spinor leads to restriction of the holonomy group  $Hol(\nabla)$  of the torsion connection  $\nabla$ . Namely,  $Hol(\nabla)$  has to be contained in  $SU(n)$ ,  $dim = 2n$  [\[86,](#page-26-0) [44,](#page-24-1) [57,](#page-24-2) [58,](#page-24-3) [13,](#page-22-0) [7,](#page-22-1) [8,](#page-22-2) [49\]](#page-24-4), the exceptional group  $G_2$ ,  $dim = 7$  [\[28,](#page-23-0) [45,](#page-24-5) [29\]](#page-23-1), the Lie group  $Spin(7)$ ,  $dim = 8$  [\[45,](#page-24-5) [56\]](#page-24-6). A detailed analysis of the possible geometries is carried out in [\[44\]](#page-24-1).

Complex non-Kähler geometries appear in string compactifications and are studied intensively for a long time [\[86,](#page-26-0) [39,](#page-23-2) [44,](#page-24-1) [45,](#page-24-5) [46,](#page-24-7) [39,](#page-23-2) [38,](#page-23-3) [7,](#page-22-1) [8\]](#page-22-2). Hermitian manifolds have widespread applications in both physics and differential geometry in connection with solutions to the Hull-Strominger system see [\[68,](#page-25-0) [30,](#page-23-4) [31,](#page-23-5) [24,](#page-23-6) [71,](#page-25-1) [72,](#page-25-2) [76,](#page-25-3) [74,](#page-25-4) [75,](#page-25-5) [77,](#page-25-6) [78,](#page-25-7) [22,](#page-23-7) [23,](#page-23-8) [20,](#page-22-3) [19,](#page-22-4) [34,](#page-23-9) [79\]](#page-25-8) and references therein. On a Hermitian manifold, there exists a unique connection which preserves the Hermitian structure and has totally skewsymmetric torsion tensor. Its existence and explicit expression first appeared in Strominger's seminal paper [\[86\]](#page-26-0) in 1986 in connection with the heterotic supersymmetric string background, where he called it the H-connection. Three years later, Bismut formally discussed and used this connection in his local index theorem paper [\[11\]](#page-22-5), which leads to the name Bismut connection in literature. We call this connection the Strominger-Bismut connection. Note that the connection also appeared implicitly earlier([\[88\]](#page-26-1)) and in some literature it was also called the KT connection (Kähler with torsion) or characteristic connection. When the holonomy of the Strominger-Bismut connection is contained in  $SU(n)$  one has the notion of Calabi-Yau manifold with torsion (CYT) spaces and these manifolds are of great interest in string theories and in mathematics since CYT appear in the Hull-Strominger system. Vanishing theorems for the Dolbeault cohomology on compact Hermitian non-Kähler manifold were found in terms of the Strominger-Bismut connection in [\[5,](#page-22-6) [58,](#page-24-3) [57,](#page-24-2) [90,](#page-26-2) [91\]](#page-26-3).

If the torsion 3-form T of  $\nabla$  is closed,  $dT = 0$  which, in Hermitian manifold, is equivalent to the condition  $\partial \bar{\partial} F = 0$ , where F is the fundamental 2-form, the Hermitian metric g is called SKT (strong Kähler with torsion) [\[51\]](#page-24-8) or pluriclosed. The SKT (pluriclosed) metrics have found many applications in both physics, see eg [\[36,](#page-23-10) [52,](#page-24-9) [86,](#page-26-0) [47,](#page-24-10) [48,](#page-24-11) [21\]](#page-22-7) and geometry, see eg [\[61,](#page-24-12) [57,](#page-24-2) [58,](#page-24-3) [25,](#page-23-11) [30,](#page-23-4) [31,](#page-23-5) [24,](#page-23-6) [26,](#page-23-12) [83,](#page-25-9) [82\]](#page-25-10) and references therein. For example in type II string theories, the three form  $T$  is identified with the 3-form field strength and it is required by construction to satisfy  $dT = 0$  (see e.g. [\[45,](#page-24-5) [44\]](#page-24-1)). In a series of papers Streets and Tian [\[83,](#page-25-9) [84,](#page-25-11) [85\]](#page-26-4) introduced a Hermitian Ricci flow under which the pluriclosed or equivalently strong KT structure is preserved, show the relationship of pluriclosed flow to type II string backgrounds and find that pluriclosed flow preserves  $N=(2,2)$  supersymmetry (generalized Kähler geometry). More generally, the geometry of a torsion connection with closed torsion form appears in the framework of the generalized Ricci flow and the generalized (gradient) Ricci solitons developed by Garcia-Fernandez and Streets [\[35\]](#page-23-13) (see the references therein). Compact pluriclosed (strong) CYT spaces are connected with the pluriclosed flow and the equations of motion in compactifications of type II supergravity due to the recent work of Garcia-Fernandez, Jordan and Streets [\[33\]](#page-23-14). It is shown in [\[35,](#page-23-13) Proposition 8.14], (see also [\[33,](#page-23-14) Proposition 2.6] that any compact pluriclosed CYT space is automatically a steady generalized gradient Ricci soliton [\[35,](#page-23-13) Proposition 8.14], (see also [\[33,](#page-23-14) Proposition 2.6]). Moreover, global existence and convergence of pluriclosed flow to a Stromonger-Bismut-flat metric on complex manifolds admitting Strominger-Bismut-flat metric is also shown in [\[33\]](#page-23-14). Generalizations of the pluriclosed condition  $\partial \bar{\partial}F = 0$ on 2n dimensional Hermitian manifolds in the form  $F^{\ell} \wedge \partial \bar{\partial} F^{k} = 0$ ,  $1 \leq k + \ell \leq n - 1$  has been investigated in [\[27,](#page-23-15) [80,](#page-25-12) [32,](#page-23-16) [59\]](#page-24-13) etc.

Hermitian metrics with the Strominger-Bismut connection being Kähler-like, namely, its curvature satisfies the Riemannian first Bianchi identity  $(2.10)$  below) have been studied in [\[6\]](#page-22-8), investigating <span id="page-2-1"></span>this property on 6-dimensional solvmanifolds with holomorphically trivial canonical bundle and it is conjectured by Angela-Otal-Ugarte-Villakampa in [\[6\]](#page-22-8) that such metrics should be SKT (pluriclosed). Support to that can be derived from [Theorem 5][\[87\]](#page-26-5) where the simply connected Hermitian manifolds with flat Strominger-Bismut connection are classified and it easily follows that it is SKT (pluriclosed) with parallel 3-form torsion [\[6\]](#page-22-8). The latter conclusion also follows from the Cartan-Schouten theorem [\[12\]](#page-22-9), (see also [Theorem 2.2][\[3\]](#page-22-10)). The conjecture by Angella-Otal-Ugarte-Villakampa was proved in [\[92\]](#page-26-6). It is also shown in [\[92\]](#page-26-6) that in this case the torsion is closed and  $\nabla$ −parallel,  $dT = \nabla T = 0$ . Hermitian nilmanifolds whose Strominger-Bismut connection is Kähler-like are classified in  $[93]$ . A classification of compact non-Kähler Hermitian manifolds with Kähler-like Strominger-Bismut connection in complex dimension 3 and those with degenerate torsion in higher dimensions is given in  $[90]$ . For complex dimension three  $[90]$ , Theorem 5] shows that if the space is CYT and the Strominger-Bismut connection is Kähler-like then the Strominger-Bismut connection is flat and it is conjectured that this is true in higher dimensions provided the space is compact, [\[90,](#page-26-2) Conjecture 2]. This conjecture was confirmed very recently in [\[95\]](#page-26-8).

Some types of non-complex 6-manifolds have also been invented in the string theory due to the mirror symmetry and T-duality [\[62,](#page-24-14) [42,](#page-23-17) [43,](#page-24-15) [13,](#page-22-0) [14,](#page-22-11) [65,](#page-25-13) [64,](#page-25-14) [50\]](#page-24-16). Almost Hermitian manifolds with totally skewsymmetric Nijenhuis tensor arise as target spaces of a class of  $(2,0)$ -supersymmetric two-dimensional sigma models [\[73\]](#page-25-15). For the consistency of the theory, the Nijenhuis tensor has to be parallel with respect to the torsion connection with holonomy contained in  $SU(n)$ . The known models are group manifolds as well as the nearly Kähler spaces. In general, an almost Hermitian manifold does not admit a metric connection preserving the almost Hermitian structure and having totally skew-symmetric torsion. It is shown in [\[28,](#page-23-0) Theorem 10.1] that this is equivalent to the condition that the Nijenhuis tensor of type  $(0,3)$  is totally skew-symmetric, i.e. this is the class  $G_1$  in Gray-Hervella classification [\[41\]](#page-23-18). In this case, the connection is unique. This class contains the Hermitian manifolds with Strominger-Bismut connection as well as the nearly Kähler spaces. In the nearly Kähler case the torsion connection  $\nabla$  is the characteristic connection considered by Gray,  $[40]$  and the torsion T and the Nijenhuis tensor are  $\nabla$ −parallel,  $\nabla T = \nabla N = 0$  (see e.g. [\[63,](#page-24-17) [9\]](#page-22-12)). If the holonomy of the torsion connection is contained in  $SU(n)$ , that is, the Ricci 2-form  $\rho$  of  $\nabla$  representing the first Chern class is identically zero, then we have the notion of Almost Calabi-Yau manifold with torsion (briefly ACYT).

As it was pointed out in  $[86]$ , the existence of a parallel spinor with respect to a metric connection  $\nabla$  with torsion 3-form, in dimension six, leads to the restriction that its holonomy group has to be contained in  $SU(3)$ . This means one has to consider an  $SU(3)$ -structure, i.e. an almost Hermitian manifold  $(M^6, g, J)$  with topologically trivial canonical bundle trivialized by a  $(3,0)$  with respect to J form  $\Psi = \Psi^+ + \sqrt{-1}\Psi^-$  endowed with a metric connection preserving the  $SU(3)$  structure with totally skew-symmetric torsion. Our main purposes are to investigate the curvature properties of the torsion connection on an ACYT 6-manifold.

We observe in Theorem [4.4](#page-12-0) that on any compact 6-dimensional ACYT the Nijenhuis tensor is  $\nabla$ parallel and deduce in Corollary [6.2](#page-17-1) that a compact ACYT 6-manifold has closed torsion exactly when  $Ric = -\nabla \theta$ . This helps to prove our first main result,

- <span id="page-2-0"></span>**Theorem 1.1.** Let  $(M, q, J, \Psi)$  be a 6-dimensional compact ACYT space with closed torsion,  $dT = 0$ . The following conditions are equivalent:
	- a). The torsion connection is Ricci flat,  $Ric = 0$ ;
	- b). The norm of the torsion is constant,  $d||T||^2 = 0$ ;
	- c). The Lee form is  $\nabla$ -parallel,  $\nabla \theta = 0$ ;
	- d). The torsion connection has vanishing scalar curvature,  $Scal = 0$ ;
	- e). The Riemannian scalar curvature is constant,  $Scal^g = const.$ ;
	- f). The Lee form is co-closed,  $\delta\theta = 0$ .

In each of the six cases above the torsion is co-closed and therefore it is a harmonic 3-form.

In the complex case,  $N = 0$ , we have

<span id="page-3-5"></span>Corollary 1.2. A compact pluriclosed CYT 6-manifold is Strominger-Bismut Ricci flat if and only if either the torsion has a constant norm or the Riemannian scalar curvature is constant.

In particular, the torsion is harmonic.

We show in Theorem [6.5](#page-20-0) that any compact ACYT 6-manifold with closed torsion 3-form is a steady generalized gradient Ricci soliton and this condition is equivalent to a certain vector field to be parallel with respect to the torsion connection. We also find out that this vector field preserves the  $SU(3)$  structure, thus extending the above-mentioned result in [\[33\]](#page-23-14) for compact pluriclosed CYT space to compact ACYT space with closed torsion in dimension six.

Studying the curvature properties of the torsion connection on an ACYT 6-manifold we find sufficient conditions for the torsion 3-form to be closed and harmonic.

<span id="page-3-0"></span>**Theorem 1.3.** Let  $(M, g, J, \Psi)$  be a 6-dimensional ACYT space. The torsion connection  $\nabla$  preserving the SU(3) structure is Ricci flat, has curvature  $R \in S^2 \Lambda^2$  and the Nijenhuis tensor is of constant norm,

<span id="page-3-4"></span>
$$
R(X, Y, Z, V) = R(Z, V, X, Y), \qquad Ric(X, Y) = 0, \qquad ||N||^2 = const.
$$
\n(1.1)

if and only if the torsion T is  $\nabla^g$  and  $\nabla$ -parallel,  $\nabla^g T = \nabla T = 0$ .

In particular, the torsion 3-form is harmonic, the Nijenhuis tensor and the exterior derivative dF of the Kähler 2-form are  $\nabla$ -parallel,

<span id="page-3-2"></span>
$$
\nabla N = \nabla dF = 0. \tag{1.2}
$$

The curvature of the torsion connection obeys the Riemannian first Bianchi identity [\(2.10\)](#page-5-0) and vice-versa.

Taking into account Theorem [4.4](#page-12-0) below, in the compact case, Theorem [1.3](#page-3-0) can be stated in the form

<span id="page-3-1"></span>**Theorem 1.4.** Let  $(M, g, J, \Psi)$  be a compact 6-dimensional ACYT space. The torsion connection  $\nabla$ preserving the  $SU(3)$  structure has curvature  $R \in S^2 \Lambda^2$  with vanishing Ricci tensor

<span id="page-3-3"></span>
$$
R(X, Y, Z, V) = R(Z, V, X, Y), \qquad Ric(X, Y) = 0.
$$
\n(1.3)

if and only if the torsion 3-form T is  $\nabla^g$  and  $\nabla$ -parallel,  $\nabla^g T = \nabla T = 0$ .

In particular the torsion 3-form is harmonic,  $dT = \delta T = 0$  and  $(1.2)$  holds.

The curvature of the torsion connection obeys the Riemannian first Bianchi identity [\(2.10\)](#page-5-0) and vice-versa.

We can reformulate the above result as follows

Corollary 1.5. On a compact ACYT manifold in dimension 6, the next conditions are equivalent:

- The torsion connection has curvature  $R \in S^2 \Lambda^2$  with vanishing Ricci tensor;
- The curvature of the torsion connection is Kähler like;
- The torsion is closed and parallel,  $dT = \nabla^g T = \nabla T = 0$ .

Theorem [1.3](#page-3-0) helps to generalize [\[90,](#page-26-2) Theorem 5] as follows

**Theorem 1.6.** Let  $M^6$  be a CYT whose Strominger-Bismut connection  $\nabla$  satisfies [\(1.3\)](#page-3-3). Then it is Strominger-Bismut Kähler-like and therefore is Strominger-Bismut flat.

We recall,  $[35]$ , that the fixed points of the generalized Ricci flow are Ricci flat metric connections with harmonic torsion 3-form,  $Ric = dT = \delta T = 0$ . In this direction, our results show that a 6dimensional ACYT manifold with a constant norm of the Nijenhuis tensor (in particular any compact ACYT manifold) with Ricci flat torsion connection having curvature  $R \in S^2 \Lambda^2$  is a fixed point of the corresponding generalized Ricci flow. In particular, if the curvature of the torsion connection satisfies the Riemannian first Bianchi identity then it is a fixed point of the generalized Ricci flow. In the compact case, any ACYT 6-manifold with closed torsion and either constant Riemannian scalar curvature or constant norm of the torsion is a fixed point of the generalized Ricci flow.

Note that Hermitian and almost Hermitian manifolds of type  $G_1$  with  $\nabla$ −parallel torsion 3-form are investigated in  $[4, 81]$  $[4, 81]$ , recently in  $[18, 94, 70]$  $[18, 94, 70]$  $[18, 94, 70]$  $[18, 94, 70]$  and a large number of examples are given there. On the

<span id="page-4-4"></span>Hermitian manifold,  $N = 0$ , the condition  $\nabla T = 0$  is precisely expressed in terms of the curvature of the Strominger-Bismut connection  $\nabla$  and its traces ([94, Theorem 1.1]). More generally, Riemannian manifolds equipped with metric connection  $\nabla$  with  $\nabla$ −parallel torsion 3-form are investigated in [\[2,](#page-22-15) [1,](#page-22-16) [16,](#page-22-17) [17\]](#page-22-18). It is conjectured very recently that if a complete Hermitian manifold has Strominger-Bismut parallel curvature and torsion and zero all Ricci tensors then it has to be Strominger-Bismut flat, [\[70,](#page-25-17) Conjecture 1.8. A special case with the additional conditions that the space is Kähler-like and CYT is proved in [\[70,](#page-25-17) Proposition 6.8]. It is worth mentioning  $[2,$  Theorem 4.1] which states that an irreducible complete and simply connected Riemannian manifold of dimension bigger or equal to 5 with ∇−parallel and closed torsion 3-form,  $\nabla T = dT = 0$  (which is equivalent to  $\nabla T = \sigma^T = 0$  due to [\(2.5\)](#page-4-2) below), is a simple compact Lie group or its dual non-compact symmetric space with biinvariant metric, and, in particular, the torsion connection is the flat Cartan connection.

**Remark 1.7.** A combination of the above-mentioned  $(92,$  Theorem 1] with [\[2,](#page-22-15) Theorem 4.1] shows that an irreducible complete and simply connected Hermitian manifold of dimension bigger or equal to 6 which is Kähler-like should be a simple compact Lie group with flat Strominger-Bismut connection. Moreover, due to Theorem [1.4](#page-3-1) and  $[2,$  Theorem 4.1, irreducible simply connected compact 6-dimensional ACYT space satisfying conditions  $(1.3)$  does not exist since there are no simple compact Lie groups of dimension six.

We also note that the main result  $[92,$  Theorem 1 combined with  $[58,$  Corollary 4.7 yields in a straightforward way the next

Corollary 1.8. Let M be a 2n-dimensional compact Hermitian manifold and its Strominger-Bismut connection is Kähler-like. If M is conformally balanced (the Lee form is an exact form) then the space is a Kähler manifold.

Convention 1.9. Everywhere in the paper, we will make no difference between tensors and the corresponding forms via the metric as well as we will use Einstein summation conventions, i.e. repeated Latin indices are summed over.

## <span id="page-4-0"></span>2 Preliminaries

In this section, we recall some known curvature properties of a metric connection with totally skewsymmetric torsion on Riemannian manifold as well as the notions and existence of a metric linear connection preserving a given Almost Hermitian structure and having skew-symmetric torsion from [\[55,](#page-24-18) [28,](#page-23-0) [60\]](#page-24-19).

### <span id="page-4-1"></span>2.1 Metric connection with totally skew-symmetric torsion and its curvature

Let  $(M, g, \nabla)$  be a Riemannian manifold with a metric connection  $\nabla$  and  $T(X, Y) = \nabla_X Y - \nabla_Y X - [X, Y]$ be its torsion. The torsion of type  $(0,3)$  is denoted by the same letter and it is defined by  $T(X, Y, Z)$  $g(T(X, Y), Z)$ . The torsion is totally skew-symmetric if  $T(X, Y, Z) = -T(X, Z, Y)$ .

On a Riemannian manifold  $(M, g)$  of dimension n any metric connection  $\nabla$  with totally skewsymmetric torsion T is connected with the Levi-Civita connection  $\nabla^g$  of the metric g by

<span id="page-4-3"></span>
$$
\nabla^g = \nabla - \frac{1}{2}T. \tag{2.4}
$$

The exterior derivative  $dT$  has the following expression (see e.g. [\[55,](#page-24-18) [58,](#page-24-3) [28\]](#page-23-0))

$$
dT(X, Y, Z, V) = (\nabla_X T)(Y, Z, V) + (\nabla_Y T)(Z, X, V) + (\nabla_Z T)(X, Y, V) + 2\sigma^T(X, Y, Z, V) - (\nabla_V T)(X, Y, Z),
$$
\n(2.5)

<span id="page-4-2"></span>where the 4-form  $\sigma^T$ , introduced in [\[28\]](#page-23-0), is defined by  $\sigma^T(X, Y, Z, V) = \frac{1}{2} \sum_{j=1}^n (e_j \Box T) \wedge (e_j \Box T) (X, Y, Z, V)$ ,  $(e_a\lrcorner T)(X,Y) = T(e_a,X,Y)$  is the interior multiplication and  $\{e_1,\ldots,e_n\}$  is a basis. The properties of the 4-form  $\sigma^T$  are studied in detail in [\[2\]](#page-22-15) where it is shown that  $\sigma^T$  measures the 'degeneracy' of the  $3$ -form  $T$ .

<span id="page-5-8"></span>The identity [\(2.4\)](#page-4-3) yields

<span id="page-5-7"></span>
$$
\nabla^g T = \nabla T + \frac{1}{2}\sigma^T.
$$
\n(2.6)

For the curvature of  $\nabla$  we use the convention  $R(X,Y)Z = [\nabla_X, \nabla_Y]Z - \nabla_{[X,Y]}Z$  and  $R(X,Y,Z,V) =$  $g(R(X, Y)Z, V)$ . It has the well-known properties

<span id="page-5-2"></span>
$$
R(X, Y, Z, V) = -R(Y, X, Z, V) = -R(X, Y, V, Z). \tag{2.7}
$$

<span id="page-5-1"></span>The first Bianchi identity for  $\nabla$  can be written in the form (see e.g. [\[55,](#page-24-18) [58,](#page-24-3) [28\]](#page-23-0))

$$
R(X, Y, Z, V) + R(Y, Z, X, V) + R(Z, X, Y, V)
$$
  
=  $dT(X, Y, Z, V) - \sigma^T(X, Y, Z, V) + (\nabla_V T)(X, Y, Z).$  (2.8)

It is proved in [\[28,](#page-23-0) p.307] that the curvature of a metric connection  $\nabla$  with totally skew-symmetric torsion T satisfies also the identity

$$
R(X, Y, Z, V) + R(Y, Z, X, V) + R(Z, X, Y, V) - R(V, X, Y, Z) - R(V, Y, Z, X) - R(V, Z, X, Y)
$$
  
= 
$$
\frac{3}{2}dT(X, Y, Z, V) - \sigma^{T}(X, Y, Z, V),
$$

which combined with [\(2.8\)](#page-5-1) yields

$$
R(V, X, Y, Z) + R(V, Y, Z, X) + R(V, Z, X, Y) = -\frac{1}{2}dT(X, Y, Z, V) + (\nabla_V T)(X, Y, Z)
$$
(2.9)

<span id="page-5-5"></span>**Definition 2.1.** We say that the curvature R satisfies the Riemannian first Bianchi identity if

<span id="page-5-0"></span>
$$
R(X, Y, Z, V) + R(Y, Z, X, V) + R(Z, X, Y, V) = 0.
$$
\n(2.10)

It is well known algebraic fact that  $(2.7)$  and  $(2.10)$  imply  $R \in S^2 \Lambda^2$ , i.e.

<span id="page-5-3"></span>
$$
R(X, Y, Z, V) = R(Z, V, X, Y)
$$
\n
$$
(2.11)
$$

holds.

**Remark 2.2.** Note that, in general,  $(2.7)$  and  $(2.11)$  do not imply  $(2.10)$ .

It is known that a metric connection  $\nabla$  with totally skew-symmetric torsion T has curvature  $R \in S^2 \Lambda^2$ , i.e. it satisfies [\(2.11\)](#page-5-3) if and only if the covariant derivative of the torsion with respect to the torsion connection is a 4-form.

**Lemma 2.3.** [\[55,](#page-24-18) Lemma 3.4] The next equivalences hold for a metric connection with torsion 3-form

$$
(\nabla_X T)(Y, Z, V) = -(\nabla_Y T)(X, Z, V) \iff R(X, Y, Z, V) = R(Z, V, X, Y)) \iff dT = 4\nabla^g T. \tag{2.12}
$$

We need the very recent result from [\[60\]](#page-24-19)

<span id="page-5-6"></span>**Theorem 2.4.** [\[60,](#page-24-19) Theorem 1.2] A metric connection  $\nabla$  with torsion 3-form T satisfies the Riemannian first Bianchi identity exactly when the next identities hold

$$
dT = -2\nabla T = \frac{2}{3}\sigma^T.
$$

In this case, the torsion 3-form T is parallel with respect to a metric connection with torsion 3-form  $\frac{1}{3}T$ [\[3\]](#page-22-10) and therefore has a constant norm,  $||T||^2 = const.$ 

The Ricci tensors and scalar curvatures of the connections  $\nabla^g$  and  $\nabla$  are related by [\[28,](#page-23-0) Section 2], (see also [\[35,](#page-23-13) Prop. 3.18])

$$
Ric^{g}(X,Y) = Ric(X,Y) + \frac{1}{2}(\delta T)(X,Y) + \frac{1}{4} \sum_{i=1}^{n} (g(T(X,e_i), T(Y,e_i);
$$
  
\n
$$
Scal^{g} = Scal + \frac{1}{4}||T||^{2}, \qquad Ric(X,Y) - Ric(Y,X) = -(\delta T)(X,Y),
$$
\n(2.13)

<span id="page-5-4"></span>where  $\delta = (-1)^{np+n+1} * d*$  is the co-differential acting on p-forms and  $*$  is the Hodge star operator satisfying  $*^2 = (-1)^{p(n-p)}$ .

### <span id="page-6-6"></span><span id="page-6-0"></span>2.2 Torsion connection preserving an almost Hermitian structure

In this section we recall the notions and existence of a metric linear connection preserving a given almost Hermitian structure and having totally skew-symmetric torsion from [\[28\]](#page-23-0).

Let  $(M, g, J)$  be an almost Hermitian 2n-dimensional manifold with Riemannian metric g and an almost complex structure  $J$  satisfying

$$
g(JX, JY) = g(X, Y).
$$

The Nijenhuis tensor N, the Kähler form F and the Lee form  $\theta$  are defined by

<span id="page-6-4"></span>
$$
N = [J, J, -[, .] - J[J, .] -[, J, J], \quad F = g(., J, ), \quad \theta(.) = \delta F(J, ), \tag{2.14}
$$

respectively, where  $\delta = - * d *$  is the co-differential on an even dimensional manifold.

Consequently, the 1-form  $J\theta$  defined by  $J\theta(X) = -\theta(JX)$  is co-closed due to  $\delta(J\theta) = -\delta^2 F = 0$ .

The Nijenhuis tensor of type (0,3), denoted by the same letter, is defined by  $N(X, Y, Z) = q(N(X, Y), Z)$ .

In general, on an almost Hermitian manifold, there not always exists a metric connection with totally skew-symmetric torsion preserving the almost Hermitian structure. It is shown in [\[28,](#page-23-0) Theorem 10.1] that there exists a unique linear connection  $\nabla$  preserving an almost hermitian structure  $(g, J)$ ,

$$
\nabla g = \nabla J = 0
$$

and having totally skew-symmetric torsion  $T$  if and only if the Nijenhuis tensor  $N$  is a 3-form. This is precisely the class  $G_1$  in the Gray-Hervella classification of almost Hermitian manifolds [\[41\]](#page-23-18) and includes Hermitian manifolds,  $N = 0$  as well as nearly Kähler spaces,  $(\nabla^g \chi J)X = 0$ . We call this connection the torsion connection.

The torsion T of the torsion connection is determined by [\[28,](#page-23-0) Theorem 10.1]

<span id="page-6-2"></span>
$$
T = JdF + N = -dF(J, J, J, J) + N = -dF^{+}(J, J, J, J) + \frac{1}{4}N,
$$
\n(2.15)

where  $dF^+$  denotes the  $(1,2)+(2,1)$ -part of  $dF$  with respect to J and satisfies the equality

$$
dF^+(X,Y,Z) = dF^+(JXJ,Y,Z) + dF^+(JX,Y,JZ) + dF^+(X,JY,JZ). \tag{2.16}
$$

<span id="page-6-1"></span>It follows from [\(2.16\)](#page-6-1) that

<span id="page-6-3"></span>
$$
dF^{+}(Je_k, Je_i, e_j)dF^{+}(e_k, e_i, e_j) = \frac{1}{3}||dF^{+}||^{2}.
$$
\n(2.17)

The  $(3,0)+(0,3)$ -part  $dF^-$  obeys the identity

$$
dF^{-}(JX, Y, Z) = dF^{-}(X, JY, Z) = dF^{-}(X, Y, JZ)
$$

and is determined completely by the Nijenhuis tensor  $[37]$ . If N is a three form then (see e.g. [\[37,](#page-23-20) [28\]](#page-23-0))

$$
dF^{-}(X, Y, Z) = \frac{3}{4}N(JX, Y, Z).
$$

The Lee form  $\theta := \delta F \circ J$  of the  $G_1$ - manifold is given in terms of the torsion T as

<span id="page-6-5"></span>
$$
\theta(X) = -\frac{1}{2}T(JX, e_i, Je_i) = \frac{1}{2}g(T(JX), F) = \frac{1}{2}(F\lrcorner T(JX)), \quad \theta_i = \frac{1}{2}J^k{}_iT_{kj\ell}F_{j\ell} \,, \tag{2.18}
$$

where  $\{e_1, \ldots, e_n, e_{n+1} = Je_1, \ldots, e_{2n} = Je_n\}$  is an orthonormal basis.

When the almost complex structure is integrable,  $N = 0$ , the torsion connection was defined and studied by Strominger [\[86\]](#page-26-0) in connection with the heterotic string background and was used by Bismut to prove a local index theorem for the Dolbeault operator on Hermitian non-Kähler manifold  $[11]$ . This formula was applied also in string theory (see e.g. [\[8\]](#page-22-2)). In the Hermitian case, the torsion connection

<span id="page-7-5"></span>is also known as the Strominger-Bismut connection and has attained a lot of consideration both in mathematics and physics (see the Introduction).

In the nearly Kähler case, the torsion connection  $\nabla$  is the characteristic connection considered by Gray, [\[40\]](#page-23-19) and the torsion T and the Nijenhuis tensor are  $\nabla$ -parallel,  $\nabla T = \nabla N = 0$  (see e.g. [\[63,](#page-24-17) [9\]](#page-22-12)).

In general  $G_1$  manifold, the holonomy group of  $\nabla$  is contained in  $U(n)$ . The reduction of the holonomy group of the torsion connection to a subgroup of  $SU(n)$  can be expressed in terms of its Ricci 2-form which represents the first Chern class, namely

$$
\rho(X, Y) = \frac{1}{2}R(X, Y, e_i, Je_i) = 0.
$$

We remark that the formula [\[58,](#page-24-3)  $(3.16)$ ] holds in the general case of a  $G_1$ -manifold (see also [\[28\]](#page-23-0)), namely, the next formula holds true

<span id="page-7-2"></span>
$$
\rho(X, Y) = Ric(X, JY) + (\nabla_X \theta)(JY) + \frac{1}{4} dT(X, Y, e_i, Je_i).
$$
\n(2.19)

Thus, on a  $G_1$  manifold the restricted holonomy group of the torsion connection is contained in  $SU(n)$  if and only if the next condition holds [\[28,](#page-23-0) Theorem 10.5]

<span id="page-7-1"></span>
$$
Ric(X,Y) + (\nabla_X \theta)(Y) - \frac{1}{4}dT(X,JY,e_i,Je_e) = 0.
$$
\n(2.20)

We calculate from  $(2.5)$  using  $(2.15)$  and  $(2.17)$ 

$$
dT(e_j, Je_j, e_i, Je_i) = -8(\nabla_{e_i} \theta)(e_i) + 8|\theta|^2 - 4T(e_i, e_j, e_k)T(Je_i, Je_j, e_k)
$$
  
= 
$$
8\delta\theta + 8||\theta||^2 - \frac{4}{3}||dF^+||^2 + \frac{1}{4}||N||^2 = 8\delta\theta + 8|\theta|^2 - \frac{4}{3}||T||^2 + \frac{1}{3}||N||^2.
$$
 (2.21)

**Remark 2.5.** It follows from [\(2.21\)](#page-7-0) that if the structure is balanced,  $\theta = 0$ , and the torsion is closed,  $dT = 0$  then  $16|dF|^2 = 3|N|^2$ . In the complex case,  $N = 0$ , this confirms the old result first established in  $[5]$  that a balanced Hermitian manifold with closed torsion must be Kähler.

The trace in [\(2.20\)](#page-7-1) together with [\(2.21\)](#page-7-0) and [\(2.13\)](#page-5-4) yield the following formulas for the scalar curvature of the torsion connection and the Riemannian scalar curvature on an ACYT space

<span id="page-7-3"></span>
$$
Scal = 3\delta\theta + 2||\theta||^2 - \frac{1}{3}||dF^+||^2 + \frac{1}{16}||N||^2; \qquad Scal^g = 3\delta\theta + 2||\theta||^2 - \frac{1}{3}||dF^+||^2 + \frac{5}{64}||N||^2. \tag{2.22}
$$

We obtain from  $(2.19)$ ,  $(2.22)$  and  $(2.21)$  the following

**Theorem 2.6.** Let  $(M, q, J)$  be a 2n-dimensional balanced ACYT space.

a) If either the Riemannian scalar curvature or the scalar curvature of the torsion connection vanishes,  $s^g = 0$  or  $s = 0$  then

<span id="page-7-4"></span><span id="page-7-0"></span>
$$
16||dF^+||^2 = 3||N||^2. \tag{2.23}
$$

b) If the Ricci tensor of the torsion connection vanishes,  $Ric = 0$  then  $dT(X, Y, e_i, Je_i) = 0$ . In particular, [\(2.23\)](#page-7-4) holds.

Consequently, we have

Corollary 2.7. On a balanced CYT manifold if either the Riemannian scalar curvature or the scalar curvature of the Strominger-Bismut connection vanishes,  $Scal^g = 0$  or  $Scal = 0$  then it must be a Kähler Ricci flat, i.e. a Calabi-Yau space.

In particular, any balanced CYT space with vanishing Ricci tensor of the Strominger-Bismut connection is Kähler Ricci flat, i.e. it is a Calabi-Yau space.

Following [\[6\]](#page-22-8) we have

<span id="page-8-4"></span>**Definition 2.8** ([\[6\]](#page-22-8)). We say that a curvature tensor R is Kähler like if it resembles the Kähler curvature identities, i.e. in addition to  $(2.7)$  and  $(2.10)$  it satisfies

$$
R(X, Y, JZ, JV) = R(X, Y, Z, V); \tag{2.24}
$$

<span id="page-8-1"></span>Since the torsion connection on an almost Hermitian manifold with skew-symmetric Nijenhuis tensor preserves the Hermitian metric and the complex structure its curvature satisfies [\(2.7\)](#page-5-2) and [\(2.24\)](#page-8-1). Clearly, the curvature of the torsion connection is of Kähler type exactly when it satisfies the Riemannian first Bianchi identity [\(2.10\)](#page-5-0). In this case, the Ricci tensor is symmetric, J-invariant and  $Ric(X, JY) = \rho(X, Y)$ .

## <span id="page-8-0"></span>3 SU(3)-structures

Let  $(M, g, J)$  be an almost Hermitian 6-manifold with a Riemannian metric g and an almost complex structure J, i.e.  $(q, J)$  define an  $U(3)$ -structure.

An  $SU(3)$ -structure is determined by an additional non-degenerate (3,0)-form  $\Psi = \Psi^+ + \sqrt{-1}\Psi^$ of constant norm, or equivalently by a non-trivial spinor. To be more explicit, we may choose a local orthonormal frame  $e_1, \ldots, e_6$ , identifying it with the dual basis via the metric. Write  $e_{i_1 i_2 \ldots i_p}$  for the monomial  $e_{i_1} \wedge e_{i_2} \wedge \cdots \wedge e_{i_p}$ . An  $SU(3)$ -structure is described locally by

$$
\Psi = -(e_1 + \sqrt{-1}e_2) \wedge (e_3 + \sqrt{-1}e_4) \wedge (e_5 + \sqrt{-1}e_6), \quad F = -e_{12} - e_{34} - e_{56},
$$
  
\n
$$
\Psi^+ = -e_{135} + e_{236} + e_{146} + e_{245}, \quad \Psi^- = -e_{136} - e_{145} - e_{235} + e_{246}.
$$
\n(3.25)

<span id="page-8-3"></span>These forms are subject to the compatibility relations

$$
F \wedge \Psi^{\pm} = 0
$$
,  $\Psi^+ \wedge \Psi^- = -\frac{2}{3}F^3$ ;

The subgroup of  $SO(6)$  fixing the forms F and  $\Psi$  simultaneously is  $SU(3)$ . The two forms F and  $\Psi$ determine the metric completely. The Lie algebra of  $SU(3)$  is denoted  $su(3)$ .

The failure of the holonomy group of the Levi-Civita connection  $\nabla^g$  of the metric g to be reduced to  $SU(3)$  can be measured by the intrinsic torsion  $\tau$ , which is identified with  $dF$ ,  $d\Psi^+$  and  $d\Psi^-$  and can be decomposed into five classes [\[15\]](#page-22-19),  $\tau \in W_1 \oplus \cdots \oplus W_5$ . The intrinsic torsion of an  $U(n)$ - structure belongs to the first four components described by Gray-Hervella  $[41]$ . The five components of a  $SU(3)$ -structure are first described by Chiossi-Salamon [\[15\]](#page-22-19) (for interpretation in physics see [\[13,](#page-22-0) [42,](#page-23-17) [43\]](#page-24-15)) and are determined by  $dF, d\Psi^+, d\Psi^-$  as well as  $dF$  and N. We list below those of these classes which we will use later.

 $\tau \in W_1$ : The class of Nearly Kähler (weak holonomy) manifold defined by  $dF$  to be  $(3,0)+(0,3)$ -form.

- $\tau \in W_3$ : The class of balanced Hermitian manifold determined by the conditions  $N = \theta = 0$ .
- $\tau \in W_1 \oplus W_3 \oplus W_4$ : The class called by Gray-Hervella  $G_1$ -manifolds determined by the condition that the Nijenhuis tensor is totally skew-symmetric. This is the precise class which we are interested in.
- $\tau \in W_1 \oplus W_3$ : The class of balanced  $G_1$ -manifolds determined by the condition that the Nijenhuis tensor is totally skew-symmetric and the Lee form vanishes,  $N(X, Y, Z) = -N(X, Z, Y)$ ,  $\theta = 0$ .
- $\tau \in W_3 \oplus W_4$ : The class of Hermitian manifolds,  $N = 0$ .

If all five components are zero then we have a Ricci-flat Kähler (Calabi-Yau) 3-fold.

Let  $(M, g, J, \Psi)$  be an  $SU(3)$  manifold. Letting the four form

$$
\Phi = \frac{1}{2}F^2 = \frac{1}{2}F \wedge F
$$

<span id="page-8-2"></span>we have the relations

$$
*\Phi = -F, \quad *F = -\Phi; \quad *\Psi^+ = \Psi^-, \quad *\Psi^- = -\Psi^+ \n*(\alpha \wedge F) = -\alpha \Box \Phi, \quad \alpha \in \Lambda^1, \quad *(\alpha \wedge \Phi) = J\alpha = -\alpha \Box F, \quad \alpha \in \Lambda^1, \n*(\alpha \wedge \Psi^+) = -\alpha \Box \Psi^- \quad \alpha \in \Lambda^1, \quad *(\alpha \wedge \Psi^-) = \alpha \Box \Psi^+ \quad \alpha \in \Lambda^1, \n*(\beta \wedge \Psi^+) = \beta \Box \Psi^-, \quad \beta \in \Lambda^2, \quad *(\beta \wedge \Psi^-) = -\beta \Box \Psi^+, \quad \beta \in \Lambda^2
$$
\n(3.26)

<span id="page-9-5"></span>Extending the action of the almost complex structure on the exterior algebra,  $J\beta = (-1)^p \beta(J,\ldots,J)$ . for a p-form  $\beta \in \Lambda^p$ , we recall the decomposition of the exterior algebra under the action of  $SU(3)$  from [\[10\]](#page-22-20) (see also [\[67\]](#page-25-18)). Let  $V = T_pM$  be the vector space at a point  $p \in M$  and  $V^*$  be the dual vector space. Obviously  $SU(3)$  acts irreducibly on  $V^*$  and  $\Lambda^5 V^*$ , while  $\Lambda^2 V^*$  and  $\Lambda^3 V^*$  decompose as follows:

$$
\Lambda^2 V^* = \Lambda_1^2 V^* \oplus \Lambda_6^2 V^* \oplus \Lambda_8^2 V^*;
$$
  

$$
\Lambda^3 V^* = \Lambda_{Re}^3 V^* \oplus \Lambda_{Im}^3 \oplus \Lambda_6^3 V^* \oplus \Lambda_{12}^3 V^*,
$$

where

$$
\Lambda_1^2 V^* = \mathbb{R} \cdot F \quad \Lambda_6^2 V^* = \{ \phi \in \Lambda^2 V^* | J \phi = -\phi \}
$$

$$
\Lambda_8^2 V^* = \{ \phi \in \Lambda^2 V^* | J \phi = \phi, \phi \wedge F^2 = 0 \}
$$

$$
\Lambda_{Re}^3 V^* = \mathbb{R} \Psi^+; \qquad \Lambda_{Im}^3 = \mathbb{R} \Psi^-;
$$

$$
\Lambda_6^3 V^* = \{ \alpha \wedge F | \alpha \in \Lambda^1 V^* \};
$$

$$
\Lambda_{12}^3 V^* = \{ \gamma \in \Lambda^3 V^* | \gamma \wedge F = \gamma \wedge \Psi^+ = \gamma \wedge \Psi^- = 0 \}.
$$

Denote by  $S^2$  the space of symmetric 2-tensors which are of type  $(2,0)+(0,2)$  with respect to the almost complex structure  $J$ , i.e.

<span id="page-9-1"></span>
$$
h(X,Y) = h(Y,X), \qquad h(JX,JY) = -h(X,Y). \tag{3.27}
$$

It is known (see [\[10\]](#page-22-20)) that the map  $\gamma : S^2 \to \Lambda^3_{12}$  defined by

<span id="page-9-0"></span>
$$
\gamma(h_{ij}) = h_{ip}\Psi^{+}_{\ \ pjk} + h_{jp}\Psi^{+}_{\ \ pki} + h_{kp}\Psi^{+}_{\ \ pij}, \qquad h_{im} = \gamma^{-1}(B_{ijk}) = \frac{1}{4}B_{ijk}\Psi^{+}_{\ \ mjk} \tag{3.28}
$$

is an isomorphism of  $su(3)$  representations.

Writing

$$
F = \frac{1}{2} F_{ij} e_{ij}, \quad \Psi^+ = \frac{1}{6} \Psi^+_{ijk} e_{ijk}, \quad \Psi^- = \frac{1}{6} \Psi^-_{ijk} e_{ijk}, \quad \Phi = \frac{1}{24} \Phi_{ijkl} e_{ijkl}
$$

we have the obvious identites  $(c.f. [10])$  $(c.f. [10])$  $(c.f. [10])$ 

$$
\Phi_{jslm} = F_{js}F_{lm} + F_{sl}F_{jm} + F_{lj}F_{sm},
$$
\n
$$
\Psi_{ipq}^{+}F_{pq} = 0 = \Psi_{ipq}^{-}F_{pq}, \quad \Phi_{ijkl}\Psi_{jkl}^{+} = 0 = \Phi_{ijkl}\Psi_{jkl}^{-},
$$
\n
$$
F_{ip}F_{pj} = -\delta_{ij}, \quad \Psi_{ijs}^{+}F_{sk} = -\Psi_{ijk}^{-}, \quad \Psi_{ijs}^{-}F_{sk} = \Psi_{ijk}^{+},
$$
\n
$$
\Psi_{ipq}^{+}\Psi_{jpq}^{-} = -4F_{ij}, \quad \Psi_{ipq}^{+}\Psi_{jpq}^{+} = 4\delta_{ij} = \Psi_{ipq}^{-}\Psi_{jpq}^{-},
$$
\n
$$
\Psi_{kls}^{+}\Psi_{ijs}^{-} = \delta_{kj}F_{li} + \delta_{li}F_{kj} - \delta_{ki}F_{lj} - \delta_{lj}F_{ki},
$$
\n
$$
\Psi_{kls}^{+}\Psi_{ijs}^{+} = F_{kj}F_{li} - \delta_{li}\delta_{kj} - F_{ki}F_{lj} + \delta_{lj}\delta_{ki}
$$
\n
$$
\Phi_{ijkl}F_{kl} = 4F_{ij}, \quad \Phi_{ijkl}\Psi_{klp}^{+} = 2\Psi_{ijp}^{+}, \quad \Phi_{ijkl}\Psi_{klp}^{-} = 2\Psi_{ijp}^{-},
$$
\n
$$
\Phi_{ijkl}\Psi_{lqp}^{+} = -F_{ij}\Psi_{pqk}^{-} - F_{jk}\Psi_{pqi}^{-} - F_{ki}\Psi_{pqj}^{-},
$$
\n
$$
\Phi_{ijkl}\Psi_{lqp}^{-} = F_{ij}\Psi_{pqk}^{+} + F_{jk}\Psi_{pqi}^{+} + F_{ki}\Psi_{pqj}^{+},
$$
\n
$$
\Phi_{ijkl}\Phi_{rjkl} = 12\delta_{ir}, \quad \Phi_{ijkl}\Phi_{klqr} = 2F_{ij}F_{qr} - 2\delta_{jq}\delta_{ir} + 2\delta_{iq}\delta_{jr},
$$

<span id="page-9-4"></span><span id="page-9-3"></span>Now, we prove the following algebraic fact which is crucial for our investigations.

**Proposition 3.1.** Let A be a 4-form and define the 3-form  $B = (X \rvert A)$  for any  $X \in T_pM$ . If the 3-form  $B \in \Lambda^3_{12}$  then the four form A vanishes identically,  $A = 0$ 

Proof. According to [\(3.28\)](#page-9-0) and [\(3.27\)](#page-9-1) the tensor

<span id="page-9-2"></span>
$$
C(X, Y, Z) = 4\gamma^{-1}(X \lrcorner A)(Y, Z) = A(X, Y, e_i, e_j)\Psi^+(Z, e_i, e_j)
$$
  
= 
$$
C(X, Z, Y) = -C(Z, X, Y) = -C(Z, Y, X) = C(Y, Z, X) = C(Y, X, Z) = -C(X, Y, Z) \quad (3.30)
$$

<span id="page-10-7"></span>and therefore vanishes. We get from [\(3.30\)](#page-9-2) using [\(3.29\)](#page-9-3)

$$
0 = A_{pqkl}\Psi^{+}{}_{kls}\Psi^{+}{}_{ijs} = A_{pqkl}\Big[F_{kj}F_{li} - \delta_{li}\delta_{kj} - F_{ki}F_{lj} + \delta_{lj}\delta_{ki}\Big] = 2A_{pqkl}F_{kj}F_{li} - 2A_{pqji} \quad (3.31)
$$

The equalities  $(3.30)$  and  $(3.31)$  yield

$$
A(X, Y, JZ, JV) = A(X, Y, Z, V) = A(X, JY, JZ, V) = A(X, JY, Z, JV) = -A(X, Y, JZ, JV)
$$

<span id="page-10-1"></span> $\Box$ 

<span id="page-10-0"></span>Hence  $A = 0$  which completes the proof of Proposition [3.1.](#page-9-4)

## 4 Almost Calabi-Yau with torsion space in dimension 6

The presence of a parallel spinor with respect to a metric connection with torsion 3-form in dimension 6 leads to the reduction to  $SU(3)$ , i.e. the existence of an almost Hermitian structure and a linear connection preserving the almost Hermitian structure with torsion 3-form with holonomy group inside  $SU(3)$ .

The reduction of the holonomy group of the torsion connection to  $SU(3)$ , was investigated intensively in the complex case, i.e. when the induced almost complex structure is integrable  $(N = 0)$  and it is known as the Strominger condition. The Strominger condition (see [\[86\]](#page-26-0)) has different very useful expressions in terms of the five torsion classes (see e.g. [\[13,](#page-22-0) [68,](#page-25-0) [30,](#page-23-4) [31,](#page-23-5) [54\]](#page-24-20)). We present here the result from [\[54\]](#page-24-20) which gives necessary and sufficient condition to have a  $\nabla$ -parallel spinor in dimension 6,

<span id="page-10-6"></span>**Theorem 4.1.** [\[54,](#page-24-20) Theorem 4.1] Let  $(M, g, J, \Psi)$  be a 6-dimensional smooth manifold with an SU(3)structure  $(q, J, \Psi)$ . The next two conditions are equivalent:

- a)  $(M, g, J, \Psi)$  is an ACYT space, i.e. there exists a unique SU(3)-connection  $\nabla$  with torsion 3-form whose holonomy is contained in  $SU(3)$ ,  $\nabla g = \nabla J = \nabla \Psi^+ = \nabla \Psi^- = 0$ ;
- b) The Nijenhuis tensor N is totally skew-symmetric and the following two conditions hold

$$
d\Psi^{+} = \theta \wedge \Psi^{+} - \frac{1}{4}(N, \Psi^{+}) * F; \qquad d\Psi^{-} = \theta \wedge \Psi^{-} - \frac{1}{4}(N, \Psi^{-}) * F. \tag{4.32}
$$

<span id="page-10-4"></span><span id="page-10-2"></span>The torsion is given by

$$
T = - * dF + * (\theta \wedge F) + \frac{1}{4} (N, \Psi^{+}) \Psi^{+} + \frac{1}{4} (N, \Psi^{-}) \Psi^{-}.
$$
 (4.33)

Our first observation is

<span id="page-10-5"></span>**Proposition 4.2.** Let  $(M, g, J, \Psi)$  be an ACYT 6-manifold. The following conditions are equivalent:

- a) The exterior derivative of the Lee form is J-invariant,  $d\theta(X, Y) = d\theta(JX, JY)$ ;
- b) The Nijenhuis tensor is of a constant norm,  $||N||^2 = const.$ ;
- c) The Nijenhuis tensor is parallel with respect to the torsion connection,  $\nabla N = 0$ .

In particular, on a 6-dimensional CYT manifold, the 2-form  $d\theta$  is of type (1,1).

<span id="page-10-3"></span>*Proof.* For the torsion T, we obtain from  $(4.33)$  using  $(3.26)$  that

$$
T = - * dF + *( \theta \wedge F) + \frac{1}{4} (N, \Psi^{+}) \Psi^{+} + \frac{1}{4} (N, \Psi^{-}) \Psi^{-}.
$$
  

$$
= *d * \Phi - \theta \Psi^{+} + \frac{1}{4} (N, \Psi^{+}) \Psi^{+} + \frac{1}{4} (N, \Psi^{-}) \Psi^{-}.
$$
  

$$
= -\delta \Phi - \theta \Psi^{+} + \frac{1}{4} (N, \Psi^{+}) \Psi^{+} + \frac{1}{4} (N, \Psi^{-}) \Psi^{-}.
$$
 (4.34)

The formula for the codifferential of a 4-form in terms of the Levi-Civita connection  $\nabla^g$  and [\(2.4\)](#page-4-3) imply

$$
- \delta \Phi_{klm} = \nabla_j^g \Phi_{jklm} = \nabla_j \Phi_{jklm} - \frac{1}{2} T_{jsk} \Phi_{jslm} + \frac{1}{2} T_{jsl} \Phi_{jskm} - \frac{1}{2} T_{jsm} \Phi_{jskl} = -\frac{1}{2} T_{jsk} \Phi_{jslm} + \frac{1}{2} T_{jsl} \Phi_{jskm} - \frac{1}{2} T_{jsm} \Phi_{jskl}, \quad (4.35)
$$

since  $\nabla \Phi = 0$ .

Substitute  $(4.35)$  into  $(4.34)$  to get the expression for the 3-form torsion T,

<span id="page-11-1"></span>
$$
T_{klm} = -\frac{1}{2}T_{jsk}\Phi_{jslm} + \frac{1}{2}T_{jsl}\Phi_{jskm} - \frac{1}{2}T_{jsm}\Phi_{jskl} - \theta_s\Phi_{sklm} + \lambda\Psi^+_{klm} + \mu\Psi^-_{klm},
$$
(4.36)

where the functions  $\lambda, \mu$  are defined by the scalar products

$$
\lambda = \frac{1}{4}(N, \Psi^+) = \frac{1}{24} N_{ijk} \Psi^+{}_{ijk}, \qquad \mu = \frac{1}{4}(N, \Psi^-) = \frac{1}{24} N_{ijk} \Psi^-{}_{ijk}.
$$
 (4.37)

It is straightforward to check that the Lee form  $\theta$  can be expressed in terms of the torsion T, the 2-form  $F$  and the 4-form  $\Phi$  as follows

<span id="page-11-2"></span><span id="page-11-0"></span>
$$
\theta_i = \frac{1}{2} T_{sjk} F_{jk} F_{si} = \frac{1}{6} T_{jkl} \Phi_{jkli}; \quad \theta_i F_{iq} = -\frac{1}{2} T_{qjk} F_{jk}.
$$
\n(4.38)

Further, we calculate from  $(4.36)$  the functions  $\lambda$  and  $\mu$  in terms of the torsion T using  $(3.29)$  as follows

<span id="page-11-10"></span>
$$
T_{klm}\Psi^{+}{}_{klm} = -\frac{3}{2}T_{jsk}\Phi_{jslm}\Psi^{+}{}_{klm} + 24\lambda = -3T_{jsk}\Psi^{+}{}_{jsk} + 24\lambda \Longrightarrow \lambda = \frac{1}{6}T_{klm}\Psi^{+}{}_{klm};
$$
  
\n
$$
T_{klm}\Psi^{-}{}_{klm} = -\frac{3}{2}T_{jsk}\Phi_{jslm}\Psi^{-}{}_{klm} + 24\mu = -3T_{jsk}\Psi^{-}{}_{jsk} + 24\mu \Longrightarrow \mu = \frac{1}{6}T_{klm}\Psi^{-}{}_{klm}.
$$
\n(4.39)

We have from  $(4.36)$  using the first formula in  $(3.29)$  and  $(4.38)$  that

<span id="page-11-3"></span>
$$
T_{klm} = -\frac{1}{2}T_{jsk}\Phi_{jslm} + \frac{1}{2}T_{jsl}\Phi_{jskm} - \frac{1}{2}T_{jsm}\Phi_{jskl} - \theta_s\Phi_{sklm} + \lambda\Psi^+{}_{klm} + \mu\Psi^-{}_{klm}
$$
  
=  $\theta_sF_{sk}F_{lm} + T_{jsk}F_{jm}F_{ls} - \theta_sF_{sl}F_{km} - T_{jsl}F_{jm}F_{ks} - \theta_sF_{sm}F_{lk} - T_{jsm}F_{jk}F_{ls}$   

$$
- \theta_s[F_{sk}F_{lm} + F_{kl}F_{sm} + F_{ls}F_{km}] + \lambda\Psi^+{}_{klm} + \mu\Psi^-{}_{klm}
$$
  
=  $T_{jsk}F_{jm}F_{ls} - T_{jsl}F_{jm}F_{ks} - T_{jsm}F_{jk}F_{ls} + \lambda\Psi^+{}_{klm} + \mu\Psi^-{}_{klm}. \tag{4.40}$ 

We can write the formula [\(4.40\)](#page-11-3) in the form

<span id="page-11-4"></span>
$$
T(X, Y, Z) = T(JX, JY, Z) + T(JX, Y, JZ) + T(X, JY, JZ) + \lambda \Psi^{+}(X, Y, Z) + \mu \Psi^{-}(X, Y, Z). \tag{4.41}
$$

Hence, we get from  $(2.15)$  and  $(4.41)$ 

$$
N = \lambda \Psi^{+} + \mu \Psi^{-}, \quad ||N||^{2} = 24(\lambda^{2} + \mu^{2}). \tag{4.42}
$$

<span id="page-11-9"></span><span id="page-11-8"></span>**Lemma 4.3.** On a 6-dimensional ACYT manifold, the next identities hold

<span id="page-11-7"></span><span id="page-11-6"></span>
$$
d\mu = Jd\lambda, \quad d\lambda = -Jd\mu,\tag{4.43}
$$

$$
d\theta(X,Y) - d\theta(JX,JY) = -(d\mu \Box \Psi^{-})(X,Y) = -(d\lambda \Box \Psi^{+})(X,Y). \tag{4.44}
$$

*Proof.* Using  $(3.29)$ , we obtain from  $(4.32)$ 

<span id="page-11-5"></span>
$$
0 = d^2\Psi^+ = d\theta \wedge \Psi^+ + \lambda\theta \wedge *F - d\lambda \wedge *F - \lambda d * F = d\theta \wedge \Psi^+ - \lambda\theta \wedge \Phi - \lambda * J\theta - d\lambda \wedge *F
$$
  

$$
= d\theta \wedge \Psi^+ - d\lambda \wedge *F,
$$
  

$$
0 = d^2\Psi^- = d\theta \wedge \Psi^- + \mu\theta \wedge *F - \mu d * F - d\mu \wedge *F = d\theta \wedge \Psi^- - \mu\theta \wedge \Phi - \mu * J\theta - d\mu \wedge *F
$$
  

$$
= d\theta \wedge \Psi^- - d\mu \wedge *F,
$$
  

$$
= d\theta \wedge \Psi^- - d\mu \wedge *F,
$$
  

$$
(4.45)
$$

<span id="page-12-3"></span>where we used the identities

$$
J\theta = \delta F = - *d * F = *(dF \wedge F), *J\theta = -dF \wedge F = d * F.
$$

The equalities  $(4.45)$  and  $(3.26)$  imply

<span id="page-12-1"></span>
$$
d\theta_{ij}\Psi^{-}{}_{ijk} = 2d\lambda_s F_{sk}; \qquad d\theta_{ij}\Psi^{+}{}_{ijk} = -2d\mu_s F_{sk}.
$$
\n(4.46)

Multiply  $(4.46)$  with  $F_{kl}$ , use  $(3.29)$  and  $(4.46)$  to obtain

$$
-2d\mu_s F_{sl} = d\theta_{ij}\Psi^+{}_{ijl} = d\theta_{ij}\Psi^-{}_{ijk}F_{kl} = -2d\lambda_l,
$$

which shows the identities  $(4.43)$  hold.

The equalities  $(4.46)$  yield applying  $(3.26)$  and  $(3.29)$ 

$$
0 = \frac{1}{2}d\theta_{ij}\Psi^{+}{}_{ijk}\Psi^{+}{}_{kab} + d\mu_{s}F_{sk}\Psi^{+}{}_{kab} = \frac{1}{2}d\theta_{ij}\Big[F_{ib}F_{ja} - F_{ia}F_{jb} - \delta_{ib}\delta_{ja} + \delta_{ia}\delta_{jb}\Big] + d\mu_{s}\Psi^{-}{}_{sab}
$$

$$
= -d\theta_{ij}F_{ia}F_{jb} + d\theta_{ab} + d\mu_{s}\Psi^{-}{}_{sab}.
$$

This proves the first equality in [\(4.44\)](#page-11-7). The second equality in [\(4.44\)](#page-11-7) follows from the already established [\(4.43\)](#page-11-6) and [\(3.26\)](#page-8-2). The lemma is proved.  $\Box$ 

The equivalence between a) and c) is a consequence from [\(4.44\)](#page-11-7) and [\(4.42\)](#page-11-8) since  $\nabla \Psi^+ = \nabla \Psi^- = 0$ . Clearly, c) implies b). For the converse, the condition  $||N||^2 = const.$  and [\(4.42\)](#page-11-8) imply

$$
\lambda^2 + \mu^2 = const., \lambda d\lambda + \mu d\mu = 0
$$

which yields applying [\(4.43\)](#page-11-6)

$$
-\lambda J d\mu + \mu J d\lambda = 0, \quad \lambda d\mu - \mu d\lambda = 0.
$$

Thus we obtain a homogeneous linear system with respect to  $d\lambda$  and  $d\mu$ . The determinant of this system is the constant  $\lambda^2 + \mu^2$ . If this constant is zero then  $N = 0$  and  $(4.42)$  gives  $\lambda = \mu = 0$ . If the constant is not zero one concludes  $d\lambda = d\mu = 0$ . Therefore  $\nabla N = 0$  due to [\(4.42\)](#page-11-8). □

Applying the Hopf's maximum principle we obtain

<span id="page-12-0"></span>**Theorem 4.4.** Let  $(M, g, J, \Psi = \Psi^+ + \sqrt{-1}\Psi^-)$  be a 6-dimensional compact ACYT manifold.

Then the Nijenhuis tensor is parallel with respect to the torsion connection and the exterior derivative of the Lee form is J-invariant,

$$
\nabla N = 0, \qquad d\theta(X, Y) = d\theta(JX, JY).
$$

In particular, the Nijenhuis tensor is of a constant norm,  $||N||^2 = const.$ 

*Proof.* From Lemma [4.3,](#page-11-9) we have  $0 = d^2\lambda = -dJd\mu$ . The trace yields  $(dJd\mu)_{ij}F_{ij} = 0$  which is an elliptic second-order operator annihilating constants and we apply the Hopf's maximum principle (see e.g. [\[89,](#page-26-10) [35\]](#page-23-13)) to conclude  $\mu = const.$  Hence  $d\lambda = d\mu = 0$  due to [\(4.43\)](#page-11-6). Now, [\(4.44\)](#page-11-7) and Proposition [4.2](#page-10-5) complete the proof. □

<span id="page-12-2"></span>Corollary 4.5. Let  $(M, g, J, \Psi)$  be a 6-dimensional ACYT manifold.

If the Lee form is closed,  $d\theta = 0$ , then the Nijenhuis tensor is  $\nabla$ -parallel,  $\nabla N = 0$ .

If the Lee form vanishes,  $\theta = 0$  then the Nijenhuis tensor N is parallel with respect to the torsion connection, the torsion 3-form is co-closed and the Ricci tensor of  $\nabla$  is symmetric,

$$
\nabla N = \delta T = 0, \qquad Ric(X, Y) = Ric(Y, X).
$$

*Proof.* If  $d\theta = 0$ , one has  $\nabla N = 0$  due to Proposition [4.2.](#page-10-5)

Suppose  $\theta = 0$ . Then  $d\lambda = d\mu = 0$  due to [\(4.44\)](#page-11-7) and [\(4.33\)](#page-10-2) together with [\(4.32\)](#page-10-4) and [\(3.26\)](#page-8-2) imply

$$
d*T=\lambda d\Psi^--\mu d\Psi^+=(-\lambda\mu+\mu\lambda)\Phi=0.
$$

This completes the proof in view of  $(2.13)$ .

 $\Box$ 

As a consequence of the proof of Proposition [4.2](#page-10-5) and Corollary [4.5](#page-12-2) we characterize Ricci-flat balanced ACYT.

**Theorem 4.6.** Let  $(M, q, J, \Psi)$  be a 6-dimensional balanced ACYT manifold,  $\theta = 0$ . The following conditions are equivalent:

- a) The Ricci tensor of the torsion connection vanishes,  $Ric = 0$ ;
- b) The torsion 3-form is harmonic, i.e. it is closed and co-closed,  $dT = \delta T = 0$ .

*Proof.* Let the assumption a) holds. The condition  $Ric = 0$  implies  $\delta T = 0$ .

On an SU(3) manifold in dimension six, we can express the Ricci tensor in a more appropriate form as follows. Since the torsion connection  $\nabla$  preserves the SU(3)-structure its curvature lies in the Lie algebra  $su(3)$ , i.e. it satisfies

<span id="page-13-1"></span>
$$
R_{ijab}\Psi^{+}{}_{abc} = R_{ijab}\Psi^{-}{}_{abc} = R_{ijab}F_{ab} = 0 \iff R_{ijab}\Phi_{abkl} = -2R_{ijkl}.\tag{4.47}
$$

<span id="page-13-0"></span>We have from  $(4.47)$  using  $(2.9)$ ,  $(4.38)$  and  $(2.5)$  that the Ricci tensor Ric of  $\nabla$  is given by

$$
Ric_{ij} = \frac{1}{2}R_{iabc}\Phi_{jabc} = \frac{1}{6}\Big[R_{iabc} + R_{ibca} + R_{icab}\Big]\Phi_{jabc} = \frac{1}{12}dT_{iabc}\Phi_{jabc} + \frac{1}{6}\nabla_i T_{abc}\Phi_{jabc} = \frac{1}{12}dT_{iabc}\Phi_{jabc} - \nabla_i \theta_j.
$$
 (4.48)

Similarly, we have

<span id="page-13-2"></span>
$$
0 = R_{iabc}\Psi^{+}_{abc} = \frac{1}{3} \Big[ R_{iabc} + R_{ibca} + R_{icab} \Big] \Psi^{+}_{abc} = \frac{1}{6} dT_{iabc}\Psi^{+}_{abc} + \frac{1}{3} \nabla_i T_{abc} \Psi^{+}_{abc},
$$
  
\n
$$
0 = R_{iabc}\Psi^{-}_{abc} = \frac{1}{3} \Big[ R_{iabc} + R_{ibca} + R_{icab} \Big] \Psi^{-}_{abc} = \frac{1}{6} dT_{iabc}\Psi^{-}_{abc} + \frac{1}{3} \nabla_i T_{abc} \Psi^{-}_{abc}.
$$
\n(4.49)

From Proposition [4.5](#page-12-2) we know that the condition  $\theta = 0$  implies  $d\lambda = d\mu = 0$ . Then  $Ric = 0$ , [\(4.48\)](#page-13-1) and [\(4.49\)](#page-13-2) yield

<span id="page-13-3"></span>
$$
dT_{pjkl}\Phi_{jkli} = 12\nabla_p\theta_i = 0, \qquad dT_{pjkl}\Psi^+{}_{jkl} = -12\nabla_p\lambda = 0, \qquad dT_{pjkl}\Psi^-{}_{jkl} = -12\nabla_p\mu = 0. \tag{4.50}
$$

The identities [\(4.50\)](#page-13-3) show that the 3-form  $(X \lrcorner dT) \in \Lambda^3_{12}$ . Then the 4-form  $dT = 0$  due to Proposition [3.1.](#page-9-4) For the converse, if we assume b) then [\(4.48\)](#page-13-1) and  $\theta = 0$  show  $Ric = 0$  which completes the proof.  $\Box$ 

One gets from [\(4.48\)](#page-13-1) and Theorem [2.4](#page-5-6)

<span id="page-13-4"></span>**Theorem 4.7.** Let  $(M, g, J, \Psi)$  be a 6-dimensional ACYT manifold and the torsion connection  $\nabla$  has curvature  $R \in S^2 \Lambda^2$  i.e. [\(2.11\)](#page-5-3) holds.

Then the covariant derivative of the 1-form  $J\theta$  with respect to  $\nabla$  is skew-symmetric,

$$
(\nabla_X J\theta)Y + (\nabla_Y J\theta)X = 0 \Longleftrightarrow (\nabla_X \theta)JY = -(\nabla_Y \theta)JX.
$$

In particular, the vector field dual to the 1-form Jθ is a Killing vector field.

a) If in addition, the Nijenhuis tensor N is of constant norm,  $||N||^2 = const.$  then N is  $\nabla$ -parallel, the covariant derivative of the Lee form  $\theta$  with respect to  $\nabla$  is symmetric and J-invariant,

<span id="page-13-5"></span>
$$
\nabla N = 0, \qquad (\nabla_X \theta) Y = (\nabla_Y \theta) X = (\nabla_{JX} \theta) JY. \tag{4.51}
$$

b) If the Lee form vanishes,  $\theta = 0$ , i.e. the manifold is balanced  $G_1$  space,  $M \in W_1 \oplus W_3$ , then

$$
\nabla T = \nabla dF = \nabla N = 0, \qquad Scal^g = -\frac{1}{3} ||dF^+||^2 + \frac{5}{64} ||N||^2 = const.
$$

<span id="page-14-7"></span>Proof. We get from [\(4.38\)](#page-11-2) using the fact proved in [\[55,](#page-24-18) Lemma 3.4] that the curvature of a metric connection  $\nabla$  with totally skew-symmetric torsion T satisfies [\(2.11\)](#page-5-3) exactly when  $\nabla T$  is a 4-form,

<span id="page-14-2"></span>
$$
\nabla_p \theta_i F_{iq} = -\frac{1}{2} \nabla_p T_{qjk} F_{jk} = \frac{1}{2} \nabla_q T_{pjk} F_{jk} = -\nabla_q \theta_i F_{ip};
$$
  
\n
$$
(\nabla_X \theta) JY = -(\nabla_Y \theta) JX;
$$
  
\n
$$
(\nabla_X J\theta) Y = -(\nabla_X \theta) JY = (\nabla_Y \theta) JX = -(\nabla_Y J\theta) X.
$$
\n(4.52)

<span id="page-14-1"></span>It follows from [\(4.52\)](#page-14-1) that the vector field  $J\theta$  dual to the 1-form  $J\theta$  is Killing since [\(2.4\)](#page-4-3) yields  $(\nabla^g_X J\theta)Y + (\nabla^g_Y J\theta)X = 0$  which proves the first part of Theorem [4.7.](#page-13-4)

Take the covariant derivative of the torsion expressed in  $(4.36)$  with respect to  $\nabla$  to get

$$
\nabla_p T_{klm} = -\frac{1}{2} \nabla_p T_{jsk} \Phi_{jslm} + \frac{1}{2} \nabla_p T_{jsl} \Phi_{jskm} - \frac{1}{2} \nabla_p T_{jsm} \Phi_{jskl} - \nabla_p \theta_s \Phi_{sklm} + \nabla_p \lambda \Psi^+{}_{klm} + \nabla_p \mu \Psi^-{}_{klm}, \quad (4.53)
$$

where we used that all the structure forms  $g, F, \Psi^+, \Psi^-, \Phi$  are  $\nabla$ -parallel.

The fact that  $\nabla T$  is a 4-form, taking the trace in  $(4.53)$  and using  $(4.38)$ , yields

$$
0 = \nabla_p T_{plm} = \frac{1}{2} \nabla_p T_{jsl} \Phi_{jspm} - \frac{1}{2} \nabla_p T_{jsm} \Phi_{jspl} - \nabla_p \theta_s \Phi_{splm} + \nabla_p \lambda \Psi^+_{plm} + \nabla_p \mu \Psi^-_{plm}
$$
  
\n
$$
= -3 \nabla_l \theta_m + 3 \nabla_m \theta_l + \nabla_p \lambda \Psi^+_{plm} + \nabla_p \mu \Psi^-_{plm} + \frac{1}{2} (\nabla_p \theta_s - \nabla_s \theta_p) \Big( F_{ps} F_{lm} + F_{sl} F_{pm} + F_{lp} F_{sm} \Big)
$$
  
\n
$$
= -3 \nabla_l \theta_m + 3 \nabla_m \theta_l + \nabla_p \lambda \Psi^+_{plm} + \nabla_p \mu \Psi^-_{plm} + \frac{1}{2} (\nabla_p \theta_s - \nabla_s \theta_p) \Big( F_{sl} F_{pm} + F_{lp} F_{sm} \Big)
$$
  
\n
$$
= -3 \nabla_l \theta_m + 3 \nabla_m \theta_l + \nabla_p \lambda \Psi^+_{plm} + \nabla_p \mu \Psi^-_{plm} + (\nabla_p \theta_s - \nabla_s \theta_p) F_{sl} F_{pm}, \quad (4.54)
$$

where we applied the general identity  $\nabla_i \theta_s F_{si} = 0$  which is a consequence of the fact that the 1-form  $J\theta$ is co-closed. Indeed, we have  $0 = \delta(J\theta) = -\nabla^g_i (J\theta)_i = -\nabla_i (J\theta)_i = -\nabla_i \theta_s F_{si}$  applying [\(2.4\)](#page-4-3).

Substitute  $(4.52)$  into  $(4.54)$  to get

<span id="page-14-4"></span><span id="page-14-3"></span>
$$
d^{\nabla}\theta_{lm} = \nabla_l \theta_m - \nabla_m \theta_l = \frac{1}{2} [\nabla_p \lambda \Psi^+_{plm} + \nabla_p \mu \Psi^-_{plm}] = \nabla_p \lambda \Psi^+_{plm} = \nabla_p \mu \Psi^-_{plm},\tag{4.55}
$$

where we used  $(4.43)$  to conclude the last equality.

The condition  $||N||^2 = const.$  and Proposition [4.2](#page-10-5) imply  $d\lambda = d\mu = 0$  and one gets  $d^{\nabla} \theta = 0$  from  $(4.55)$ . Then  $(4.52)$  completes the proof of a) in Theorem [4.7.](#page-13-4)

To show the condition  $\theta = 0$  yields  $\nabla T = 0$  we observe that [\(4.44\)](#page-11-7) implies the functions  $\lambda$  and  $\mu$  are constants. In particular  $\nabla N = 0$  due to [\(4.42\)](#page-11-8). Then [\(4.39\)](#page-11-10) together with [\(4.38\)](#page-11-2) imply

$$
\nabla_p T_{jkl} \Phi_{jkli} = 6 \nabla_p \theta_i = 0, \qquad \nabla_p T_{jkl} \Psi^+{}_{jkl} = 6 \nabla_p \lambda = 0, \qquad \nabla_p T_{jkl} \Psi^-{}_{jkl} = 6 \nabla_p \mu = 0. \tag{4.56}
$$

<span id="page-14-5"></span>The identities [\(4.56\)](#page-14-5) show that the 3-form  $(X \rvert \nabla T) \in \Lambda^3_{12}$ . Hence, the 4-form  $\nabla T = 0$  due to Proposition [3.1.](#page-9-4) Then the formulas [\(2.15\)](#page-6-2) and [\(2.22\)](#page-7-3) complete the proof of the Theorem [4.7.](#page-13-4)

### <span id="page-14-0"></span>4.1 Example

We take the next example from [\[54\]](#page-24-20) which supports Theorem [4.7.](#page-13-4)

Let G be the six-dimensional connected simply connected and nilpotent Lie group, determined by the left-invariant 1-forms  $\{e_1, \ldots, e_6\}$  such that

<span id="page-14-6"></span>
$$
de_2 = de_3 = de_6 = 0,
$$
  
\n
$$
de_1 = e_3 \wedge e_6, \quad de_4 = e_2 \wedge e_6, \quad de_5 = e_2 \wedge e_3.
$$
\n(4.57)

Consider the metric on  $G \cong \mathbb{R}^6$  defined by  $g = \sum_{i=1}^6 e_i^2$ . Let  $(F, \Psi)$  be the  $SU(3)$ -structure on G given by [\(3.25\)](#page-8-3). Then  $(G, F, \Psi)$  is an almost complex manifold with a  $SU(3)$ -structure.

<span id="page-15-7"></span>It is easy to verify using  $(2.14)$  and  $(4.57)$  that

<span id="page-15-1"></span>
$$
dF = -3e_{236}, \quad N = -\Psi^-, \quad d\Psi^- = *F, \quad (N, \Psi^-) = -4,
$$
  

$$
\theta = d\Psi^+ = (N, \Psi^+) = 0.
$$
 (4.58)

Hence,  $(G, \Psi, g, J)$  is neither complex nor Nearly Kähler manifold but it fulfills the conditions [\(4.32\)](#page-10-4) of Theorem [4.1](#page-10-6) and therefore there exists a  $SU(3)$  connection with torsion 3-form on  $(G, \Psi, g, J)$ . The expression  $(4.33)$  and  $(4.58)$  give

<span id="page-15-2"></span>
$$
T = -2e_{145} + e_{136} + e_{235} - e_{246}, \quad dT = -2(e_{1256} + e_{3456} + e_{1234}) = 2 * F. \tag{4.59}
$$

The equalities  $(4.59)$  and  $(2.15)$  imply the nonzero essential terms of the torsion connection are (c.f. [\[54\]](#page-24-20))

$$
\nabla_{e_1} e_6 = -e_3, \qquad \nabla_{e_5} e_2 = e_3, \qquad \nabla_{e_4} e_6 = -e_2, \n\nabla_{e_4} e_5 = -e_1, \qquad \nabla_{e_5} e_1 = -e_4, \qquad \nabla_{e_1} e_4 = -e_5.
$$
\n(4.60)

It is easy to verify cf. [\[54\]](#page-24-20)) that the curvature of the torsion connection satisfies [\(2.11\)](#page-5-3) and the torsion tensor T as well as the Nijenhuis tensor N and  $dF$  are parallel with respect to the connection  $\nabla$ .

The coefficients of the structure equations of the Lie algebra given by [\(4.57\)](#page-14-6) are integers. Therefore, the well-known theorem of Malcev [\[69\]](#page-25-19) states that the group G has a uniform discrete subgroup  $\Gamma$  such that  $Nil^6 = G/\Gamma$  is a compact 6-dimensional nil-manifold. The  $SU(3)$ -structure, described above, descends to  $Nil^6$  and therefore we obtain a compact example.

## <span id="page-15-0"></span>5 Proof of Theorem [1.3](#page-3-0) and Theorem [1.4](#page-3-1)

<span id="page-15-6"></span>We begin with the next

- **Lemma 5.1.** Let  $(M, g, J, \Psi)$  be an ACYT 6-manifold satisfying the condition [\(1.1\)](#page-3-4). Then the Lee form  $\theta$  is  $\nabla$ -parallel,  $\nabla \theta = 0$ . In particular, the Lee form is co-closed,  $\delta\theta = 0$ .
- *Proof.* We already know from Theorem [4.7](#page-13-4) a) that  $d\lambda = d\mu = 0$  and [\(4.51\)](#page-13-5) holds. Since  $\nabla T$  is a 4-form, substitute  $(2.5)$  into  $(2.20)$  to get

$$
0 = Ric(X, Y) + (\nabla_X \theta)(Y) - (\nabla_X T)(JY, e_i, Je_e) - \frac{1}{2}\sigma^T(X, JY, e_i, Je_i)
$$
  
= Ric(X, Y) + (\nabla\_X \theta)(Y) + 2(\nabla\_X \theta)Y - \frac{1}{2}\sigma^T(X, JY, e\_i, Je\_i)  
= Ric(X, Y) + 3(\nabla\_X \theta)(Y)) - \frac{1}{2}\sigma^T(X, JY, e\_i, Je\_i). (5.61)

Let  $Ric = 0$ . Then we have from  $(2.20)$ ,  $(5.61)$  and  $(2.18)$ 

$$
-\frac{1}{2}(\nabla_X T)(JY, e_i, Je_i) = (\nabla_X \theta)Y = \frac{1}{4}dT(X, JY, e_i, Je_e) = \frac{1}{6}\sigma^T(X, JY, e_i, Je_i)
$$
  
= 
$$
-\frac{1}{3}T(X, JY, J\theta) - \frac{1}{3}T(X, e_i, e_j)T(JY, Je_i, e_j).
$$
 (5.62)

<span id="page-15-4"></span>Using that  $d^{\nabla} \theta(X,Y) = (\nabla_X \theta)Y - (\nabla_Y \theta)X = -(d\mu \Psi^{-1})(X,Y) = 0$  due to [\(4.44\)](#page-11-7) and [\(4.51\)](#page-13-5), we get

<span id="page-15-3"></span>
$$
d\theta(X,Y) = (\nabla_X^g \theta)Y - (\nabla_Y^g \theta)X = -(d\mu \Box \Psi^-)(X,Y) + T(X,Y,\theta),
$$

which combined with  $(4.44)$  and  $(4.51)$  yields

<span id="page-15-5"></span>
$$
T(X,Y,\theta) - T(JX,JY,\theta) = -3d^{\nabla}\theta(X,Y) = -3(d\mu \Box \Psi^{-})(X,Y) = -3(d\lambda \Box \Psi^{+})(X,Y) = 0.
$$
 (5.63)

We have from [\(5.62\)](#page-15-4), the symmetricity of  $\nabla \theta$  and [\(5.63\)](#page-15-5) that

$$
(\nabla_X \theta)\theta = (\nabla_\theta \theta)X = -\frac{1}{3}T(\theta, JX, J\theta) - \frac{1}{3}T(\theta, e_i, e_j)T(JX, Je_i, e_j) = -\frac{1}{3}T(\theta, e_i, e_j)T(JX, Je_i, e_j)
$$
  
= 
$$
\frac{1}{3}T(\theta, Je_i, e_j)T(JX, e_i, e_j) = -\frac{1}{3}T(\theta, e_i, Je_j)T(JX, e_i, e_j) = -(\nabla_\theta \theta)X = -(\nabla_X \theta)\theta,
$$

which shows  $\nabla ||\theta||^2 = 2(\nabla_X \theta)\theta = 0$  and  $|\theta|^2 = |J\theta|^2 = const.$ Since  $|J\theta|^2 = const.$  we have

$$
0 = \nabla_{e_i} \nabla_{e_i} ||J\theta||^2 = (\nabla_{e_i} \nabla_{e_i} J\theta)(e_j) J\theta(e_j) + ||\nabla J\theta||^2.
$$
 (5.64)

<span id="page-16-0"></span>We use the Killing condition of  $J\theta$  and the Ricci identities for  $\nabla$  to evaluate the first term in [\(5.64\)](#page-16-0)

$$
\begin{split} (\nabla_{e_i} \nabla_{e_i} J \theta)(e_j) \cdot J \theta(e_j) &= -(\nabla_{e_i} \nabla_{e_j} J \theta)(e_i) \cdot J \theta(e_j) \\ &= -(\nabla_{e_j} \nabla_{e_i} J \theta)(e_i) \cdot J \theta(e_j) + R(e_i, e_j, e_k) J \theta(e_k) J \theta(e_j) + T(e_i, e_j, e_k) (\nabla_{e_k} J \theta)e_i \cdot J \theta(e_j) \\ &= -Ric(J\theta, J\theta) + T(e_i, J\theta, e_k) (\nabla_{e_k} J \theta)e_i = T(e_i, J\theta, e_k) (\nabla_{e_k} J \theta)e_i = -T(e_i, J\theta, e_k) (\nabla_{e_k} \theta) J e_i \\ &= T(Je_i, J\theta, e_k) (\nabla_{e_k} \theta)e_i. \end{split} \tag{5.65}
$$

We calculate from [\(5.63\)](#page-15-5), [\(4.51\)](#page-13-5) and [\(5.62\)](#page-15-4)

$$
0 = (\nabla_{e_i} T)(X, e_i, \theta) + T(X, e_i, e_j)(\nabla_{e_i} \theta)(e_j) = (\nabla_{e_i} T)(JX, Je_i, \theta) + T(JX, Je_i, e_j)(\nabla_{e_i} \theta)(e_j)
$$
  
\n
$$
= \frac{1}{3} \sigma^T(JX, Je_i, \theta, e_i) + T(JX, Je_i, e_j)(\nabla_{e_i} \theta)(e_j)
$$
  
\n
$$
= \frac{1}{3} \Big[ T(JX, Je_i, e_k) T(e_k, \theta, e_i) + T(Je_i, \theta, e_k) T(e_k, JX, e_i) + T(\theta, JX, e_k) T(e_k, Je_i, e_i) \Big]
$$
  
\n
$$
- T(Je_i, JX, e_j)(\nabla_{e_j} \theta)(e_i) = \frac{2}{3} \Big[ T(JX, Je_i, e_k) T(e_k, \theta, e_i) - T(\theta, JX, J\theta) \Big] - T(Je_i, JX, e_j)(\nabla_{e_j} \theta)(e_i)
$$
  
\n
$$
= \frac{2}{3} T(JX, Je_i, e_k) T(e_k, \theta, e_i) - T(Je_i, JX, e_j)(\nabla_{e_j} \theta)(e_i).
$$
 (5.66)

Set  $X = \theta$  into [\(5.66\)](#page-16-1) to get applying [\(5.63\)](#page-15-5)

$$
3T(Je_i, J\theta, e_j)(\nabla_{e_j}\theta)(e_i) = 2T(J\theta, Je_i, e_k)T(\theta, e_i, e_k) = -2T(J\theta, e_i, e_k)T(\theta, Je_i, e_k)
$$
  
= 
$$
2T(J\theta, e_i, e_k)T(\theta, e_i, Je_k) = -2T(J\theta, e_i, Je_k)T(\theta, e_i, e_k)
$$
  
= 
$$
-3T(Je_i, J\theta, e_j)(\nabla_{e_j}\theta)(e_i).
$$

Hence,

<span id="page-16-2"></span>
$$
T(Je_i, J\theta, e_j)(\nabla_{e_j}\theta)(e_i) = 0.
$$
\n(5.67)

<span id="page-16-1"></span> $\Box$ 

Substitute [\(5.67\)](#page-16-2) into [\(5.64\)](#page-16-0) to get  $\nabla J\theta = 0 = \nabla \theta$  which completes the proof of Lemma [5.1](#page-15-6)

To finish the proof of Theorem [1.3](#page-3-0) we observe from [\(5.62\)](#page-15-4), [\(4.38\)](#page-11-2), [\(4.39\)](#page-11-10), [\(4.48\)](#page-13-1), [\(4.49\)](#page-13-2) and Lemma [5.1](#page-15-6) the validity of the following identities

$$
2\nabla_p T_{jkl} \Phi_{jkli} = 12\nabla_p \theta_i = 0 = dT_{pjkl} \Phi_{jkli};
$$
  
\n
$$
2\nabla_p T_{jkl} \Psi^+_{jkl} = 12\nabla_p \lambda = 0 = -dT_{pjkl} \Psi^+_{jkl};
$$
  
\n
$$
2\nabla_p T_{jkl} \Psi^-_{jkl} = 12\nabla_p \mu = 0 = -dT_{pjkl} \Psi^-_{jkl}.
$$
\n(5.68)

<span id="page-16-3"></span>The identities [\(5.68\)](#page-16-3) show that the 3-forms  $(X\cup \nabla T) \in \Lambda^3_{12}$  and  $(X\cup dT) \in \Lambda^3_{12}$ . Hence, the four forms  $\nabla T = dT = 0$  due to Proposition [3.1](#page-9-4) and [\(2.5\)](#page-4-2) implies  $\sigma^T = 0$ . Therefore,  $\nabla^g T = 0$  because of [\(2.6\)](#page-5-7). Now Theorem [2.4](#page-5-6) shows that the Riemannian first Bianchi identity [\(2.10\)](#page-5-0) holds.

For the converse, it is a well-known algebraic fact that the Riemannian first Bianchi identity [\(2.10\)](#page-5-0) on an ACYT manifold implies [\(2.11\)](#page-5-3) and the vanishing of the Ricci tensor. This completes the proof of Theorem [1.3.](#page-3-0) Theorem [1.4](#page-3-1) follows from Theorem [1.3](#page-3-0) and Theorem [4.4.](#page-12-0)

## <span id="page-17-0"></span>6 ACYT with closed torsion and generalized Ricci solitons

<span id="page-17-4"></span>We begin with the next

**Theorem 6.1.** Let  $(M, q, J, \Psi)$  be an ACYT 6-manifold. The following conditions are equivalent:

- a). The torsion is closed,  $dT = 0$ ;
- b). The exterior derivative of the Lee form belongs to  $su(3) \cong \Lambda_8^2$  (i.e. it is J-invariant trace-free 2-orm) and the Ricci tensor of the torsion connection is equal to  $-\nabla\theta$ ,

<span id="page-17-2"></span>
$$
d\theta(X,Y) = d\theta(JX,JY), \quad d\theta \lrcorner F = d\theta \wedge \Phi = 0, \qquad Ric = -\nabla \theta; \tag{6.69}
$$

c). The Nijenhuis tensor is parallel with respect to the torsion connection and the Ricci tensor of the torsion connection is equal to  $-\nabla\theta$ ,

$$
\nabla N = 0, \qquad Ric = -\nabla \theta. \tag{6.70}
$$

*Proof.* The condition  $dT = 0$  and the equality [\(4.48\)](#page-13-1) imply  $Ric = -\nabla \theta$ . We get from [\(4.49\)](#page-13-2) and [\(4.39\)](#page-11-10)  $d\lambda = d\mu = 0$  since the torsion is closed. Hence, c) follows from [\(4.42\)](#page-11-8).

Now Proposition [4.2](#page-10-5) implies the first and the third identities in [\(6.69\)](#page-17-2).

To show that the second equality in [\(6.69\)](#page-17-2) follows from c) we observe  $-\delta T_{ij} = Ric_{ij} - Ric_{ji} = d^{\nabla} \theta_{ij}$ . Consequently, we have  $d^{\nabla}\theta_{ij}F_{ij} = \nabla_i\theta_jF_{ij} = 0$  since  $\delta J\theta = \nabla_i\theta_jF_{ij} = \delta^2 F = 0$ . Hence,

<span id="page-17-3"></span>
$$
0 = (d^{\nabla}\theta \lrcorner F).vol = -(\delta T \lrcorner F).vol = \delta T \wedge \Phi = \frac{1}{2}\delta T \wedge F \wedge F.
$$
\n(6.71)

On the other hand, we have from [\(4.33\)](#page-10-2) using  $d\lambda = d\mu = 0$  that

$$
d * T = d\theta \wedge F - \theta \wedge dF + \lambda d\Psi^{-} - \mu d\Psi^{+}
$$
  
=  $d\theta \wedge F - \theta \wedge dF + \lambda [\theta \wedge \Psi^{-} - \mu * F] - \mu [\theta \wedge \Psi^{+} - \lambda * F]$   
=  $d\theta \wedge F - \theta \wedge dF + \lambda \theta \wedge \Psi^{-} - \mu \theta \wedge \Psi^{+},$ 

yielding the identity

$$
d * T \wedge F = d\theta \wedge F \wedge F - \theta \wedge dF \wedge F = d\theta \wedge F \wedge F + \theta \wedge (*J\theta) = d\theta \wedge F \wedge F;
$$
  

$$
d\theta \wedge F \wedge F = d * T \wedge F = -d * T \wedge *\Phi = \Phi \wedge \delta T = 0,
$$

where we apply  $(6.71)$  to obtain the last equality. Hence, b) follows from c).

For the converse, assume b) holds. The third equality in [\(6.69\)](#page-17-2) combined with [\(4.48\)](#page-13-1) yields

$$
dT_{jabc}\Phi_{iabc} = 0.
$$

The first equality in [\(6.69\)](#page-17-2) implies  $d\lambda = d\mu = 0$  due to Proposition [4.2.](#page-10-5) Apply [\(4.49\)](#page-13-2) and [\(4.39\)](#page-11-10) to get

$$
dT_{jabc}\Psi^{+}{}_{abc} = dT_{jabc}\Psi^{-}{}_{abc} = 0.
$$

The last three equalities show that the 3-form  $(X \cup dT) \in \Lambda^3_{12}$ . Hence, the four form  $dT = 0$  due to Proposition [3.1](#page-9-4) which completes the equivalences between a) and b).

In view of Theorem [4.4](#page-12-0) we get from Theorem [6.1](#page-17-4)

<span id="page-17-1"></span>**Corollary 6.2.** Let  $(M, g, J, \Psi)$  be a compact ACYT 6-manifold. The following conditions are equivalent:

- a). The torsion is closed,  $dT = 0$ ;
- b). The Ricci tensor of the torsion connection satisfies  $Ric = -\nabla \theta$ . In particular  $d\theta \in su(3) \cong \Lambda_8^2$ .

As a consequence of Theorem [6.1,](#page-17-4) we obtain

**Corollary 6.3.** Let  $(M, q, J, \Psi)$  be a compact CYT 6-manifold. The following conditions are equivalent:

- a). The CYT space is pluriclosed,  $dT = \partial \bar{\partial} F = 0$ ;
- b). The Ricci tensor of the torsion connection satisfies  $Ric = -\nabla \theta$ .

In particular  $d\theta \in su(3) \cong \Lambda_8^2$ .

### <span id="page-18-11"></span><span id="page-18-0"></span>6.1 Proof of Theorem [1.1](#page-2-0)

*Proof.* The equivalence of a) and c) as well as d) and f) follow from Theorem [6.1.](#page-17-4)

The second Bianchi identity for the torsion connection reads [\[60,](#page-24-19) Proposition 3.5]

<span id="page-18-2"></span>
$$
d(Scal)_j - 2\nabla_i Ric_{ji} + \frac{1}{6}d||T||_i^2 + \delta T_{ab}T_{abj} + \frac{1}{6}T_{abs}dT_{jabc} = 0.
$$
 (6.72)

From Theorem [6.1](#page-17-4) we have taking into account [\(2.13\)](#page-5-4)

<span id="page-18-1"></span>
$$
Ric = -\nabla\theta, \quad Scal = \delta\theta, \quad \delta T_{ij} = \nabla_i \theta_j - \nabla_j \theta_i.
$$
\n(6.73)

Substitute [\(6.73\)](#page-18-1) into [\(6.72\)](#page-18-2) and use  $dT = 0$  to get

<span id="page-18-3"></span>
$$
\nabla_j \delta \theta + 2 \nabla_i \nabla_j \theta_i + \delta T_{ab} T_{abj} + \frac{1}{6} d||T||_j^2 = 0.
$$
\n(6.74)

We evaluate the second term in [\(6.74\)](#page-18-3) using the Ricci identity for  $\nabla$  and [\(6.73\)](#page-18-1) as follows

<span id="page-18-4"></span>
$$
\nabla_i \nabla_j \theta_i = \nabla_j \nabla_i \theta_i - R_{ijis} \theta_s - T_{ijs} \nabla_s \theta_i = -\nabla_j \delta \theta - \nabla_j \theta_s \theta_s - \frac{1}{2} \delta T_{si} T_{sij}.
$$
\n(6.75)

We obtain from  $(6.75)$  and $(6.74)$ 

<span id="page-18-5"></span>
$$
-\nabla_j \delta\theta - 2\nabla_j \theta_s \theta_s + \frac{1}{6} \nabla_j ||T||^2 = 0.
$$
\n(6.76)

Another covariant derivative of [\(6.76\)](#page-18-5) together with [\(6.73\)](#page-18-1) yield

<span id="page-18-9"></span>
$$
\Delta \delta \theta - 2 \nabla_j \nabla_j \theta_s \theta_s - 2||Ric||^2 - \frac{1}{6}\Delta ||T||^2 = 0,
$$
\n(6.77)

where  $\Delta$  is the Laplace operator  $\Delta = -\nabla^g_i \nabla^g_i f = -\nabla_i \nabla_i f$  acting on smooth function f since the torsion of  $\nabla$  is a 3-form.

On the other hand, [\(6.72\)](#page-18-2) yields

<span id="page-18-7"></span>
$$
\nabla_i \nabla_j \theta_i = \nabla_i (\nabla_i \theta_j - \delta T_{ij}) = \nabla_i \nabla_i \theta_j - \nabla_i \delta T_{ij} = \nabla_i \nabla_i \theta_j - \frac{1}{2} \delta T_{ia} T_{iaj},\tag{6.78}
$$

where we used the identity

<span id="page-18-6"></span>
$$
\nabla_i \delta T_{ij} = \frac{1}{2} \delta T_{ia} T_{iaj}.
$$
\n(6.79)

shown in [\[60,](#page-24-19) Proposition 3.2] for any metric connection with skew torsion.

We give a proof of  $(6.79)$  for completeness. The identity  $\delta^2 = 0$  together with  $(2.4)$  and  $(2.6)$  imply

$$
0 = \delta^2 T_k = \nabla^g_i \nabla^g_j T_{ijk} = \nabla^g_i \nabla_j T_{ijk} = \nabla_i \delta T_{ik} + \frac{1}{2} T_{iks} \delta T_{is} = \nabla_i \delta T_{ik} - \frac{1}{2} \delta T_{is} T_{isk}.
$$

The equalities [\(6.78\)](#page-18-7) and [\(6.74\)](#page-18-3) yield

<span id="page-18-8"></span>
$$
\nabla_j \delta \theta + 2 \nabla_i \nabla_i \theta_j + \frac{1}{6} \nabla_j ||T||^2 = 0.
$$
\n(6.80)

The identities  $(6.80)$  and  $(6.77)$  imply

<span id="page-18-10"></span>
$$
\Delta \left( \delta \theta - \frac{1}{6} ||T||^2 \right) + \theta_j \nabla_j \left( \delta \theta + \frac{1}{6} ||T||^2 \right) = ||Ric||^2 \ge 0. \tag{6.81}
$$

Assume b) holds. Since M is compact and  $d||T||^2 = 0$  we may apply the strong maximum principle (see e.g. [\[89,](#page-26-10) [35\]](#page-23-13)) to achieve  $\delta\theta = const. = 0$ . Conversely, the condition  $\delta\theta = 0$  implies  $d||T||^2 = 0$  by the strong maximum principle applied to [\(6.81\)](#page-18-10). Hence b) is equivalent to f).

<span id="page-19-7"></span>Further, assume f) or b). Then  $(2.21)$  implies  $d||\theta||^2 = 0$  since  $dT = \delta\theta = 0 = d||T||^2$  and the norm of the Nijenhuis tensor is constant. We calculate using the Ricci identity for  $\nabla$ , [\(6.73\)](#page-18-1) and [\(6.79\)](#page-18-6)

$$
0 = \frac{1}{2} \nabla_i \nabla_i ||\theta||^2 = \theta_j \nabla_i \nabla_i \theta_j + ||\nabla \theta||^2 = \theta_j \nabla_i \left( \nabla_j \theta_i + \delta T_{ij} \right) + ||\nabla \theta||^2
$$
  
\n
$$
= -\theta_j \nabla_j \delta \theta - R_{ijis} \theta_s \theta_j - \theta_j T_{ijs} \nabla_s \theta_i + \theta_j \nabla_i \delta T_{ij} + ||\nabla \theta||^2
$$
  
\n
$$
= -\nabla_j \theta_s \theta_s \theta_j - \frac{1}{2} \theta_j \delta T_{si} T_{sij} + \frac{1}{2} \theta_j \delta T_{ia} T_{iaj} + ||\nabla \theta||^2
$$
  
\n
$$
= -\frac{1}{2} \theta_j \nabla_j ||\theta||^2 + ||\nabla \theta||^2 = ||\nabla \theta||^2.
$$

Hence, we show that c) follows from b).

For the converse, suppose c) holds,  $\nabla \theta = 0$ . Then b) is a consequence from [\(6.76\)](#page-18-5) and [\(6.73\)](#page-18-1).

Finally, to show the equivalences of b) and e) we use  $(2.13)$  and  $(6.73)$  to write  $(6.81)$  in the form

<span id="page-19-1"></span>
$$
\Delta \left( Scal^g - \frac{5}{12} ||T||^2 \right) + \theta_j \nabla_j \left( Scal^g - \frac{1}{12} ||T||^2 \right) = ||Ric||^2 \ge 0. \tag{6.82}
$$

The strong maximum principle applied to [\(6.82\)](#page-19-1) implies b) is equivalent to e) in the same way as above. □ The proof of Theorem [1.1](#page-2-0) is completed.

### <span id="page-19-0"></span>6.2 Steady generalized Ricci solitons

It is shown in [\[35,](#page-23-13) Proposition 4.28] that a Riemannian manifold  $(M, g, T)$  with a closed torsion  $dT = 0$ is a steady generalized Ricci soliton if there exists a vector field  $X$  and a two form  $B$  such that it is a solution to the equations

<span id="page-19-3"></span>
$$
Ric^g = \frac{1}{4}T^2 - \frac{1}{2}\mathbb{L}_X g, \qquad \delta T = B \tag{6.83}
$$

for B satisfying  $d(B + X\mathcal{F}) = 0$ , where  $\mathbb{L}_X$  is the Lie derivative in the direction of X. In particular  $\Delta_d T = \mathbb{L}_X T.$ 

We also recall the equivalent formulation [\[35,](#page-23-13) Definition 4.31] that a compact Riemannian manifold  $(M, q, T)$  with a closed 3-form T is a steady generalized Ricci soliton with  $k = 0$  if there exists a vector field  $X$  such that

<span id="page-19-5"></span>
$$
Ric^{g} = \frac{1}{4}T^{2} - \frac{1}{2}\mathbb{L}_{X}g, \qquad \delta T = -X \Box T, \qquad dT = 0.
$$
 (6.84)

If the vector field  $X$  is a gradient of a smooth function  $f$  then one has the notion of a steady generalized gradient Ricci soliton.

It is shown in [\[35,](#page-23-13) Proposition 8.14] that any compact pluriclosed (strong) CYT space is automatically a steady generalized Ricci soliton.

We extend this result to a 6-dimensional ACYT space with a closed torsion form.

<span id="page-19-6"></span>**Proposition 6.4.** Let  $(M, g, J, \Psi)$  be a 6-dimensional ACYT manifold with closed torsion form,  $dT = 0$ . Then it is a steady generalized Ricci soliton with  $X = \theta$  and  $B = d\theta - \theta \Box T$ .

*Proof.* As we identify the vector field X with its corresponding 1-form via the metric, we have using  $(2.4)$ 

<span id="page-19-2"></span>
$$
dX_{ij} = \nabla^g i X_j - \nabla^g j X_i = \nabla_i X_j - \nabla_j X_i + X_s T_{sij}.
$$
\n(6.85)

In view of  $(2.13)$ ,  $(2.4)$  and  $(6.85)$  we write the first equation in  $(6.83)$  in the form

<span id="page-19-4"></span>
$$
Ric_{ij} = -\frac{1}{2}\delta T_{ij} - \frac{1}{2}(\nabla_i X_j + \nabla_j X_i) = -\frac{1}{2}\delta T_{ij} - \nabla_i X_j + \frac{1}{2}dX_{ij} - \frac{1}{2}X_s T_{sij}.
$$
 (6.86)

Set  $X = \theta$ ,  $B = d\theta - \theta \mathcal{F}$  and use [\(6.69\)](#page-17-2) to get  $\delta T = B$  and [\(6.86\)](#page-19-4) is trivially satisfied. Hence, [\(6.83\)](#page-19-3) holds since  $d(B + \theta \mathcal{F}) = d^2\theta = 0$ . holds since  $d(B + \theta \Box T) = d^2 \theta = 0$ .

<span id="page-20-5"></span>One fundamental consequence of Perelman's energy formula for Ricci flow is that compact steady solitons for Ricci flow are automatically gradient. Adapting these energy functionals to generalized Ricci flow it is proved in [\[35,](#page-23-13) Chapter 6] that steady generalized Ricci solitons on compact manifolds are automatically gradient with  $k = 0$ , i.e there exists a smooth function f such that the vector field X is equal to the gradient of the function  $f$  and  $(6.84)$  takes the form

<span id="page-20-1"></span>
$$
Ric_{ij}^g = \frac{1}{4}T_{ij}^2 - \nabla^g i \nabla^g j f, \qquad \delta T_{ij} = -df_s T_{sij}, \qquad dT = 0.
$$
\n
$$
(6.87)
$$

The smooth function f is determined with  $u = \exp(-\frac{1}{2}f)$  where u is the first eigenfunction of the Schrödinger operator

<span id="page-20-2"></span>
$$
-4\Delta + Scal^g - \frac{1}{12} ||T||^2 = -4\Delta + Scal + \frac{1}{6} ||T||^2,
$$
\n(6.88)

where the Laplace operator is defined by  $\Delta = -\delta d$  (see [\[35,](#page-23-13) Lemma 6.3, Corollary 6.10, Corollary 6.11]). In terms of the torsion connection  $(6.87)$  can be written in the form (see  $[60]$ )

<span id="page-20-3"></span>
$$
Ric_{ij} = -\nabla_i \nabla_j f, \qquad \delta T_{ij} = -df_s T_{sij}, \qquad dT = 0.
$$
\n(6.89)

Thus, a pluriclosed CYT space is a steady generalized gradient Ricci soliton [\[33,](#page-23-14) Proposition 2.6], (see also [\[35,](#page-23-13) Proposition 8.14]).

We show that similar conclusions hold for a compact closed ACYT manifold of dimension six.

<span id="page-20-0"></span>**Theorem 6.5.** Let  $(M, g, J, \Psi)$  be a 6-dimensional compact ACYT manifold with closed torsion,  $dT = 0$ . The next two conditions are equivalent:

- a).  $(M, g, J, \Psi)$  is a steady generalized gradient Ricci soliton. i.e. there exists a smooth function f determined with  $u = \exp(-\frac{1}{2}f)$  where u is the first eigenfunction of the Schrödinger operator [\(6.88\)](#page-20-2) such that  $(6.87)$ , equivalently  $(6.89)$ , hold.
- b). For f determined by the first eigenfunction u of the Schrödinger operator [\(6.88\)](#page-20-2) with  $u = \exp(-\frac{1}{2}f)$ , the vector field

$$
V = \theta - df \quad is \quad \nabla - parallel, \quad \nabla V = 0.
$$

The  $\nabla$ −parallel vector field V determines d $\theta$  and preserves the SU(3) structure  $(g, J, F, \Psi^+, \Psi^-)$  and the Nijenhuis tensor, i.e

<span id="page-20-4"></span>
$$
d\theta = V \mathbf{I}T, \quad \mathbb{L}_V g = \mathbb{L}_V F = \mathbb{L}_V J = \mathbb{L}_V \Psi^+ = \mathbb{L}_V \Psi^- = \mathbb{L}_V N = 0. \tag{6.90}
$$

The vector field JV is also  $\nabla$ -parallel and therefore Killing.

*Proof.* The first statement a) follows from Proposition  $6.4$  and the general considerations in [\[35,](#page-23-13) Chapter 6], [\[35,](#page-23-13) Lemma 6.3, Corollary 6.10, Corollary 6.11]).

To prove b) follows from a) observe that the condition  $dT = 0$  and  $(2.13)$  imply

$$
Ric = -\nabla \theta, \quad -\delta T = -d^{\nabla} \theta = -d\theta + \theta \Box T
$$

by Theorem  $6.1$  and  $(2.13)$ . The latter combined with  $(6.89)$  yield

$$
\nabla(\theta - df) = 0, \qquad d\theta = V \lrcorner T.
$$

For the converse, b) yields  $\nabla \theta = \nabla df$ , which, combined with Theorem [6.1](#page-17-4) and [\(2.13\)](#page-5-4) gives

$$
Ric_{ij} = -\nabla_i \theta_j = -\nabla_i \nabla_j f, \quad -\delta T_{ij} = Ric_{ij} - Ric_{ji} = df_s T_{sij}
$$

since  $0 = d^2 f = \nabla^g i \nabla^g j f - \nabla^g j \nabla^g i f = \nabla_i \nabla_j f - \nabla_j \nabla_i f + df_s T_{sij}$ . Hence, [\(6.89\)](#page-20-3) holds which proves the equivalences between a) and b).

Further,  $\nabla JV = (\nabla V)J = 0$ . We also have  $(\mathbb{L}_V g)_{ij} = \nabla^g_i V_j + \nabla^g_j V_i = \nabla_i V_j + \nabla_j V_i = 0$  since  $\nabla V = 0$ . Hence V and JV are Killing vector fields.

<span id="page-21-4"></span>We calculate from the definitions of the Lie derivative and the torsion tensor (see e.g.  $[66]$ ) using  $\nabla V = 0$  that

$$
\begin{aligned} (\mathbb{L}_V F)(X, Y) &= V F(X, Y) - F([V, X], Y) - F(X, [V, Y]) \\ &= V F(X, Y) - F(\nabla_V X, Y) - F(X, \nabla_V Y) + T(V, X, e_a) F(e_a, Y) + T(V, Y, e_a) F(X, e_a) \\ &= (\nabla_V F)(X, Y) + d\theta(X, JY) - d\theta(Y, JX) = 0, \quad (6.91) \end{aligned}
$$

where we apply the already proved identity  $d\theta = V \mathcal{F}$  to get the third equality and use  $\nabla F = 0$  and that  $d\theta$  is (1,1) form due to Theorem [6.1](#page-17-4) to achieve the last identity.

Using the general identity

<span id="page-21-0"></span>
$$
g(\mathbb{L}_V J)X,Y) = -(\mathbb{L}_V g)(JX,Y) - (\mathbb{L}_V F)(X,Y),
$$

we conclude  $\mathbb{L}_V J = 0$ .

Similarly to the proof of [\(6.91\)](#page-21-0) we obtain

$$
(\mathbb{L}_V \Psi^+)_{ijk} = d\theta_{is} \Psi^+{}_{sjk} + d\theta_{js} \Psi^+{}_{ski} + d\theta_{ks} \Psi^+{}_{sij};
$$
  

$$
(\mathbb{L}_V \Psi^-)_{ijk} = d\theta_{is} \Psi^-{}_{sjk} + d\theta_{js} \Psi^-{}_{ski} + d\theta_{ks} \Psi^-{}_{sij}.
$$
\n
$$
(6.92)
$$

<span id="page-21-2"></span><span id="page-21-1"></span>We calculate from  $(6.92)$  using the identities  $(3.29)$  that

$$
(\mathbb{L}_V \Psi^+)_{ijk} \Psi^+_{ijk} = 3d\theta_{is} \Psi^+_{ijk} \Psi^+_{ijk} = 12d\theta_{is}\delta_{si} = 0 = (\mathbb{L}_V \Psi^-)_{ijk} \Psi^-_{ijk};
$$
  

$$
(\mathbb{L}_V \Psi^+)_{ijk} \Psi^-_{ijk} = 3d\theta_{is} \Psi^+_{ijk} \Psi^-_{ijk} = -12d\theta_{is} F_{si} = 0 = -(\mathbb{L}_V \Psi^-)_{ijk} \Psi^+_{ijk}.
$$
 (6.93)

since  $d\theta \in su(3)$  due to Theorem [6.1](#page-17-4) which is used to conclude the last equality in [\(6.93\)](#page-21-2).

We know that  $d\theta$  is a (1,1) two form due to Theorem [6.1](#page-17-4) and  $\Psi^+, \Psi^-$  are (3,0)+(0,3) three forms. Apply this to [\(6.92\)](#page-21-1) to conclude that the three forms  $(\mathbb{L}_V \Psi^+), (\mathbb{L}_V \Psi^-)$  are  $(3,0)+(0,3)$  three forms and therefore belong to  $span{\Psi^+, \Psi^-}$  since M is six dimensional. Therefore, [\(6.93\)](#page-21-2) imply  $\mathbb{L}_V \Psi^+ = 0 = \mathbb{L}_V \Psi^-$ . Since  $d\lambda = d\mu = 0$  we get from (4.42)  $\mathbb{L}_V N = 0$  which completes the proof of (6.90).  $\mathbb{L}_V \Psi$ <sup>-</sup>. Since  $d\lambda = d\mu = 0$  we get from [\(4.42\)](#page-11-8)  $\mathbb{L}_V N = 0$  which completes the proof of [\(6.90\)](#page-20-4).

In the case  $N = 0$  one has general identity  $\mathbb{L}_{JV} J = -J\mathbb{L}_V J$  and we have

<span id="page-21-3"></span>**Corollary 6.6.** Let  $(M, g, J, \Psi)$  be a 6-dimensional compact pluriclosed CYT space. The next two conditions are equivalent:

- a).  $(M, q, J, \Psi)$  is a steady generalized gradient Ricci soliton;
- b). The vector field  $V = \theta df$  is  $\nabla$ -parallel,  $\nabla V = 0$ .

The  $\nabla$ −parallel vector field V determines d $\theta$  and preserves the SU(3) structure  $(g, J, F, \Psi^+, \Psi^-)$ , i.e.

$$
d\theta = V \mathbf{I}T, \quad \mathbb{L}_V g = \mathbb{L}_V F = \mathbb{L}_V J = \mathbb{L}_V \Psi^+ = \mathbb{L}_V \Psi^- = 0.
$$

The vector field JV is  $\nabla$ -parallel, Killing and holomorphic.

We remark that Corollary [6.6](#page-21-3) is known in all dimensions due to [\[33,](#page-23-14) Proposition 2.6]. Corollary  $6.6$ on a complex 6-manifold is a part of the 6-dimensional case of [\[33,](#page-23-14) Proposition 2.6] plus additional conclusions that the vector field V is  $\nabla$ -parallel and preserves the  $SU(3)$  structure. Basic examples of compact Hermitian non-Kähler steady generalized gradient Ricci solitons are compact Lie groups endowed with biinvariant metric and the left-invariant Samelson's complex structure. The Strominger-Bismut connection is the left-invariant flat Cartan connection with harmonic torsion 3-form of constant norm and the function  $f$  has to be a constant.

## <span id="page-22-16"></span>References

- [1] I. Agricola, A. Ferreira, Einstein manifolds with skew torsion, The Quarterly Journal of Mathematics, 65, Issue 3, (2014), 717-741, https://doi.org/10.1093/qmath/hat050 [5](#page-4-4)
- <span id="page-22-15"></span><span id="page-22-10"></span>[2] I. Agricola, A. Ferreira, Th. Friedrich, The classification of naturally reductive homogeneous spaces in dimensions  $n \leq 6$ , Diff. Geom. Appl., **39** (201[5](#page-4-4)), 59-92. 5
- <span id="page-22-13"></span>[3] I. Agricola and T. Friedrich, A note on flat metric connections with antisymmetric torsion, Diff. Geom. Appl. 28 (2010), 480-487. [3,](#page-2-1) [6](#page-5-8)
- [4] B. Alexandrov, T. Friedrich and N. Schoemann, Almost Hermitian 6-manifolds revisited, J. Geom. Phys. 53 (2005) 1. [4](#page-3-5)
- <span id="page-22-8"></span><span id="page-22-6"></span>[5] B.Alexandrov, S.Ivanov, Vanishing theorems on Hermitian manifolds, Diff. Geom. Appl. 14 (2001), 251-265. [2,](#page-1-1) [8](#page-7-5)
- <span id="page-22-1"></span>[6] D. Angella, A. Otal, L. Ugarte, R. Villacampa, On Gauduchon connections with K¨ahler-like curvature, Comm. Anal. Geom. 30 (2022), no. 5, 961-1006.[2,](#page-1-1) [3,](#page-2-1) [8,](#page-7-5) [9](#page-8-4)
- [7] K. Becker, M. Becker, K. Dasgupta, P.S. Green, Compactifications of Heterotic Theory on Non-Kahler Complex Manifolds: I, JHEP 0304 (2003) 007. [2](#page-1-1)
- <span id="page-22-2"></span>[8] K. Becker, M. Becker, K. Dasgupta, P.S. Green, E. Sharpe, Compactifications of Heterotic Strings on Non-Kahler Complex Manifolds: II, Nucl. Phys. B678 (2004), 19-100. [2,](#page-1-1) [7](#page-6-6)
- <span id="page-22-12"></span>[9] F. Belgun, A. Moroianu, Nearly Kdhler manifolds with reduced holonomy, Ann. Global Anal. Geom., 19 (2001), pp. 307-319. [3,](#page-2-1) [8](#page-7-5)
- <span id="page-22-20"></span><span id="page-22-5"></span>[10] Lucio Bedulli, Luigi Vezzoni, The Ricci tensor of SU(3)-manifolds, J. Geom. Phys. 57 (2007), n. 4, 1125-1146. [10](#page-9-5)
- <span id="page-22-9"></span>[11] J. M. Bismut, A local index theorem of non-Kähler manifolds, Math. Ann.  $284$  (1989), 681-699. [2,](#page-1-1) [7](#page-6-6)
- [12] E. Cartan, J.A. Schouten, On Riemannian manifolds admitting an absolute parallelism, Proc. Amsterdam 29 (1926), 933-946. [3](#page-2-1)
- <span id="page-22-0"></span>[13] G. L. Cardoso, G. Curio, G. Dall'Agata, D. Lust, P. Manousselis, G. Zoupanos, Non-Kähler String Backgrounds and their Five Torsion Classes, Nucl.Phys. B652 (2003) 5-34. [2,](#page-1-1) [3,](#page-2-1) [9,](#page-8-4) [11](#page-10-7)
- <span id="page-22-11"></span>[14] G. L. Cardoso, G. Curio, G. Dall'Agata, D. Lust, BPS Action and Superpotential for Heterotic String Compactifications with Fluxes, JHEP 0310 (2003) 004. [3](#page-2-1)
- <span id="page-22-19"></span>[15] S. Chiossi, S. Salamon, The intrinsic torsion of  $SU(3)$  and  $G_2$ -structures, Differential Geometry, Valencia 2001, World Sci. Publishing, 2002, pp. 115-133. [9](#page-8-4)
- <span id="page-22-17"></span>[16] I. Chrysikos, Invariant Connections with Skew-Torsion and ∇-Einstein Manifolds, Journal of Lie Theory 26 (2016) 011-048. [5](#page-4-4)
- <span id="page-22-18"></span>[17] I. Chrysikos, Ch. O'Cadiz Gustad, H. Winther, Invariant connections and ∇-Einstein structures on isotropy irreducible spaces, J. Geom. Phys.  $138$  (2019) 2[5](#page-4-4)7-284 5
- <span id="page-22-14"></span>[18] R. Cleyton, A. Moroianu, U. Semmelmann, Metric connections with parallel skew-symmetric torsion, Adv. Math. 378 (2021), Paper No. 107519, 50 pp., https://doi.org/10.1016/j.aim.2020.107519 [4](#page-3-5)
- <span id="page-22-4"></span><span id="page-22-3"></span>[19] T. C. Collins, S. Picard, S.-T. Yau, The Strominger system in the square of a Kähler class, arXiv:211.03784. [2](#page-1-1)
- [20] T. C. Collins, S. Picard, S.-T. Yau, Stability of the tangent bundle through conifold transitions, Comm. Pure Appl. Math. 77 (2024), no. 1, 284-371.[2](#page-1-1)
- <span id="page-22-7"></span>[[2](#page-1-1)1] K. Dasgupta, G. Rajesh and S. Sethi, M theory, orientifolds and G-flux, JHEP 081999023; 2
- <span id="page-23-8"></span><span id="page-23-7"></span>[22] T. Fei, Z. Huang, S. Picard, The Anomaly flow over Riemann surfaces, Int. Math. Res. Not. IMRN(2021), no. 3, 2134-2165. https://doi.org/10.1093/imrn/rnz076 [2](#page-1-1)
- <span id="page-23-6"></span>[23] T. Fei, Z. Huang, S. Picard, A Construction of Infinitely Many Solutions to the Strominger System, J. Differential Geom. Volume 117, Number 1 (2021), 23-39. [2](#page-1-1)
- <span id="page-23-11"></span>[24] M. Fernández, S. Ivanov, L. Ugarte, R. Villacampa, Non-Kähler Heterotic String Compactifications with non-zero fluxes and constant dilaton, Comm. Math. Phys. [2](#page-1-1)88 (2009), 677-697. 2
- <span id="page-23-12"></span>[25] A Fino, M Parton, S Salamon, Families of strong KT structures in six dimensions, Comment. Math. Helv. 79 (2004), 317-340; [2](#page-1-1)
- [26] A. Fino, A. Tomassini, Blow-ups and resolutions of strong K¨ahler with torsion metrics, Adv. Math. 221 (2009), 914-935. [2](#page-1-1)
- <span id="page-23-15"></span><span id="page-23-0"></span>[27] A.Fino, L. Ugarte, On generalized gauduchon metrics, Proc. Edinb. Math. Soc. 56(3) 2013, 733-753. [2](#page-1-1)
- [28] Th. Friedrich, S. Ivanov, Parallel spinors and connections with skew-symmetric torsion in string theory, Asian Journ. Math. 6 (2002), 303 - 336. [2,](#page-1-1) [3,](#page-2-1) [5,](#page-4-4) [6,](#page-5-8) [7,](#page-6-6) [8](#page-7-5)
- <span id="page-23-1"></span>[29] Th. Friedrich, S.Ivanov, Killing spinor equations in dimension 7 and geometry of integrable  $G_2$ manifolds, J. Geom. Phys **48** ([2](#page-1-1)003), 1-11. 2
- <span id="page-23-4"></span>[30] J. X. Fu and S. T. Yau, The theory of superstring with flux on non-Kähler manifolds and the complex Monge-Ampére equation, J. Differential Geom. **78** (2008), no. 3, 369-428. [2,](#page-1-1) [11](#page-10-7)
- <span id="page-23-5"></span>[31] J. X. Fu and S. T. Yau, The theory of superstring with flux on non-Kähler manifolds and the complex Monge-Ampere equation, J. Diff. Geom. 78 (2009), 369–428. [2,](#page-1-1) [11](#page-10-7)
- <span id="page-23-16"></span>[32] J. Fu, Z. Wang, and D. Wu, Semilinear equations, the  $\gamma_k$  function, and generalized Gauduchon metrics, J. Eur. Math. Soc., **15** ([2](#page-1-1)013), 659-680. 2
- <span id="page-23-14"></span>[33] M. Garcia-Fernandez, J. Jordan, J. Streets, Non-Kähler Calabi-Yau geometry and pluriclosed flow, J. Math. Pures Appl. (9) 177 (2023), 329-367, DOI: 10.1016/j.matpur.2023.07.002[.2,](#page-1-1) [4,](#page-3-5) [21,](#page-20-5) [22](#page-21-4)
- <span id="page-23-9"></span>[34] M. Garcia-Fernandez, R. Rubio, C. S. Shahbazi, C. Tipler. Canonical metrics on holomorphic Courant algebroids, Proc. London Math. Soc. 125 (2022), 700-758. [2](#page-1-1)
- <span id="page-23-13"></span><span id="page-23-10"></span>[35] M. Garcia-Fernandez and J. Streets, Generalized Ricci Flow. AMS University Lecture Series 76, 2021. [2,](#page-1-1) [4,](#page-3-5) [6,](#page-5-8) [13,](#page-12-3) [19,](#page-18-11) [20,](#page-19-7) [21](#page-20-5)
- [36] S. J. Gates, Jr., C. M. Hull and M. Rocek, Twisted Multiplets and New Supersymmetric Nonlinear Sigma Models, Nucl. Phys. B 248 (1984) 157. [2](#page-1-1)
- <span id="page-23-20"></span>[37] P. Gauduchon, Hermitian connections and Dirac operators, Boll. Un. Mat. It. 11-B (1997) Suppl. Fasc., 257-288. [7](#page-6-6)
- <span id="page-23-3"></span>[38] J. Gillard, G. Papadopoulos, D. Tsimpis, Anomaly, Fluxes and (2,0) Heterotic-String Compactifications, JHEP 0306 (2003) 035. [2](#page-1-1)
- <span id="page-23-2"></span>[39] E. Goldstein, S. Prokushkin, Geometric Model for Complex Non-Kähler Manifolds with  $SU(3)$  Structure, Commun. Math. Phys. 251 (2004), 65-78. [2](#page-1-1)
- <span id="page-23-19"></span><span id="page-23-18"></span>[40] A. Gray, *The structure of nearly Kähler manifolds*, Math. Ann., 223 (1976), pp. 233-248. [3,](#page-2-1) [8](#page-7-5)
- [41] A. Gray, L. Hervella, The sixteen classes of almost Hermitian manifolds and their linear invariants, Ann. Mat. Pura Appl. (4) 123 (1980), 35–58. [3,](#page-2-1) [7,](#page-6-6) [9](#page-8-4)
- <span id="page-23-17"></span>[42] S. Gurrieri, J. Louis, A. Micu, D. Waldram, Mirror Symmetry in Generalized Calabi-Yau Compactifications, Nucl.Phys. B654 (2003) 61-113 [3,](#page-2-1) [9](#page-8-4)
- <span id="page-24-15"></span><span id="page-24-1"></span>[43] S. Gurrieri, A. Micu, Type IIB Theory on Half-flat Manifolds, Class.Quant.Grav. 20 (2003), 2181- 2192. [3,](#page-2-1) [9](#page-8-4)
- <span id="page-24-5"></span>[44] J.Gauntlett, D.Martelli, D.Waldram, Superstrings with Intrinsic torsion, Phys. Rev. D69 (2004) 086002. [2](#page-1-1)
- <span id="page-24-7"></span>[45] J.Gauntlett, N. Kim, D.Martelli, D.Waldram, Fivebranes wrapped on SLAG three-cycles and related geometry, JHEP 0111 (2001) 018. [2](#page-1-1)
- <span id="page-24-10"></span>[46] J.P. Gauntlett, D. Martelli, S. Pakis, D. Waldram, G-Structures and Wrapped NS5-Branes, Commun. Math. Phys. 247 (2004), 421-445. [2](#page-1-1)
- <span id="page-24-11"></span>[47] J. Gutowski and G. Papadopoulos, Heterotic Black Horizons, JHEP 1007 (2010) 011 [2](#page-1-1)
- [48] J. Gutowski and G. Papadopoulos, Heterotic horizons, Monge-Ampere equation and del Pezzo surfaces, JHEP 1010 (2010) 084 [2](#page-1-1)
- <span id="page-24-4"></span>[49] J. Gutowski, S. Ivanov, G. Papadopoulos, Deformations of generalized calibrations and compact non-Kahler manifolds with vanishing first Chern class, Asian J. Math. 7 (2003), 39-80. [2](#page-1-1)
- <span id="page-24-16"></span>[50] A.S. Haupt, O. Lechtenfeld, E.T. Musaev, Order alpha′ heterotic domain walls with warped nearly Kähler geometry, Journal of High Energy Physics, (JHEP), Volume 2014, article id.  $\#152$ , 28 pp. [3](#page-2-1)
- <span id="page-24-9"></span><span id="page-24-8"></span>[51] P. S. Howe and G. Papadopoulos, *Twistor spaces for HKT manifolds*, Phys. Lett. B **379** (1996) 80 [2](#page-1-1)
- [52] P. S. Howe and G. Sierra, Two-dimensional Supersymmetric Nonlinear Sigma Models With Torsion, Phys. Lett. B 148 (1984) 451. [2](#page-1-1)
- <span id="page-24-20"></span><span id="page-24-0"></span>[53] C.M. Hull, *Compactification of the heterotic superstrings*, Phys. Lett. B 178 (1986) 357-364 [2](#page-1-1)
- [54] P. Ivanov and S. Ivanov, SU(3) instantons and G2, Spin(7) heterotic string solitons, Commun. Math. Phys. 259 (2005) 79-102. [11,](#page-10-7) [15,](#page-14-7) [16](#page-15-7)
- <span id="page-24-18"></span><span id="page-24-6"></span>[55] S. Ivanov, *Geometry of Quaternionic Kähler connections with torsion*, J. Geom. Phys. 41 (2002), 235-257. [5,](#page-4-4) [6,](#page-5-8) [15](#page-14-7)
- [56] S. Ivanov, Connection with torsion, parallel spinors and geometry of Spin(7) manifolds, Math. Res. Lett. 11 (2004), no. 2-3, 171–186. [2](#page-1-1)
- <span id="page-24-2"></span>[57] S. Ivanov and G. Papadopoulos, A no go theorem for string warped compactifications, Phys. Lett. B497 (2001) 309; [2](#page-1-1)
- <span id="page-24-3"></span>[58] S. Ivanov and G. Papadopoulos, Vanishing theorems and string backgrounds, Class. Quant. Grav. 18 (2001) 1089; [2,](#page-1-1) [5,](#page-4-4) [6,](#page-5-8) [8](#page-7-5)
- <span id="page-24-13"></span>[59] S. Ivanov and G. Papadopoulos, Vanishing theorems on  $(l|k)$ -strong Kähler manifolds with torsion, Adv. Math., 237 (2013), 147-164. [2](#page-1-1)
- <span id="page-24-19"></span>[60] S. Ivanov, N. Stanchev, The Riemannian Bianchi identities of metric connections with skew torsion and generalized Ricci solitons, arXiv:2307.03986. [5,](#page-4-4) [6,](#page-5-8) [19,](#page-18-11) [21](#page-20-5)
- <span id="page-24-12"></span>[61] J. Jost and S.T. Yau, A non-linear elliptic system for maps from Hermitian to Riemannian manifolds and rigidity theorems in Hermitian geometry, Acta Math 170 (1993) 221; Corrigendum Acta Math 177 (1994) 307. [2](#page-1-1)
- <span id="page-24-14"></span>[62] S. Kachru, M.B. Schulz, P.K. Tripathy, S.P. Trivedi, New Supersymmetric String Compactifications, JHEP 0303 (2003) 061 [3](#page-2-1)
- <span id="page-24-17"></span>[63] V. Kirichenko, K-spaces of maximal rank, (russian), Mat. Zam., 22 (1977), pp. 465-476. [3,](#page-2-1) [8](#page-7-5)
- <span id="page-25-14"></span><span id="page-25-13"></span>[64] M. Klaput, A. Lukas, C,l Matti, Bundles over Nearly-Kahler Homogeneous Spaces in Heterotic String Theory, JHEP 1109 (2011) 100 . [3](#page-2-1)
- <span id="page-25-20"></span>[65] M. Klaput, A. Lukas, C. Matti and E.E. Svanes, Moduli Stabilising in Heterotic Nearly Kähler Compactifications, JHEP 01 (2013) 015 [3](#page-2-1)
- <span id="page-25-18"></span>[66] S. Kobayashi, K. Nomizu, Foundations of differential geometry. Vol. I, Wiley Classics Lib. Wiley-Intersci. Publ. John Wiley & Sons, Inc., New York, 1996, xii+329 pp. [22](#page-21-4)
- <span id="page-25-0"></span>[67] M. Larfors, A. Lukas, F. Ruehle, Calabi-Yau Manifolds and SU(3) Structure, Journal of High Energy Physics 2019(1), DOI: 10.1007/JHEP01(2019)171 [10](#page-9-5)
- <span id="page-25-19"></span>[68] J. Li, S.-T. Yau, The existence of supersymmetric string theory with torsion, J. Diff. Geom. 70 (2005) 143. [2,](#page-1-1) [11](#page-10-7)
- [69] A. I. Malcev, On a class of homogeneous spaces, reprinted in Amer. Math. Soc. Translations, Series 1, 9 (1962), 276-307. [16](#page-15-7)
- <span id="page-25-17"></span><span id="page-25-1"></span>[70] L. Ni, F. Zheng, A classification of locally Chern homogeneous Hermitian manifolds, arXiv:2301.00579. [4,](#page-3-5) [5](#page-4-4)
- <span id="page-25-2"></span>[71] X. de la Ossa, E. E. Svanes, *Holomorphic Bundles and the Moduli Space of N=1 Heterotic Compactifications*, J. High Energy Phys. [2](#page-1-1)014, no. 10, 123, front matter $+54$  pp. 2
- [72] X. de la Ossa, M. Larfors, E. E. Svanes, Exploring SU(3) Structure Moduli Spaces with Integrable G2 Structures, Adv. Theor. Math. Physics, Volume 19 (2015) Number 4, 837-903. [2](#page-1-1)
- <span id="page-25-15"></span>[73] G. Papadopoulos (2,0)-supersymmetric sigma models and almost complex structures, Nucl.Phys. B448 (1995), 199-219. [3](#page-2-1)
- <span id="page-25-5"></span><span id="page-25-4"></span>[74] D. H. Phong, S. Picard, X. Zhang, Anomaly flows, Comm. Anal. Geom. 26 (2018), no. 4, 955-1008[.2](#page-1-1)
- [75] D. H. Phong, S. Picard, X. Zhang, The anomaly flow and the Fu-Yau equation, Ann. PDE 4 (2018), no. 2, Paper No. 13, 60 pp.[2](#page-1-1)
- <span id="page-25-3"></span>[76] D. H. Phong, S. Picard, X. Zhang, Geometric flows and Strominger systems, Math. Z. 288 (2018), no. 1-2, 101-113.[2](#page-1-1)
- <span id="page-25-6"></span>[77] D. H. Phong, S. Picard, X. Zhang, The Fu-Yau equation with negative slope parameter, Invent. Math. 209 2 (2017), doi:10.1007/s00222-016-0715-z [2](#page-1-1)
- <span id="page-25-7"></span>[78] D. H. Phong, S. Picard, X. Zhang, New curvature flows in complex geometry, Surveys in Differential Geometry 2017. Celebrating the 50th anniversary of the Journal of Differential Geometry, 331-364, Surv. Differ. Geom., 22, Int. Press, Somerville, MA, 2018. [2](#page-1-1)
- <span id="page-25-8"></span>[79] D. H. Phong, Geometric flows from unified string theories, Contribution to Surveys in Differential Geometry, "Forty Years of Ricci flow", edited by H.D. Cao, R. Hamilton, and S.T. Yau, arXiv:2304.02533. [2](#page-1-1)
- <span id="page-25-12"></span>[80] D. Popovici, Limits of projective manifolds under holomorphic deformations: Hodge numbers and strongly Gauduchon metrics, Invent. Math.  $194$  ([2](#page-1-1)013), no. 3, 515-534. 2
- <span id="page-25-16"></span><span id="page-25-10"></span>[81] N. Schoemann, Almost Hermitian structures with parallel torsion, J. Geom. Phys. 57 (2007) 2187. [4](#page-3-5)
- [82] J. Streets, Classification of solitons for pluriclosed flow on complex surfaces, Math. Ann. 375 (2019), no. 3-4, 1555-1595. [2](#page-1-1)
- <span id="page-25-11"></span><span id="page-25-9"></span>[83] J. Streets, G. Tian, A parabolic flow of pluriclosed metrics, Int. Math. Res. Not. IMRN 16 (2010), 3101-3133. [2](#page-1-1)
- [84] J. Streets, G. Tian, Regularity results for pluriclosed flow, Geom. Topol. 17 (2013), no. 4, 2389-2429. [2](#page-1-1)
- <span id="page-26-4"></span>[85] J. Streets, G. Tian,, *Generalized Kähler geometry and the pluriclosed flow*, Nuclear Phys. B 858 (2012), no. 2, 366-376. [2](#page-1-1)
- <span id="page-26-5"></span><span id="page-26-0"></span>[86] A. Strominger, Superstrings With Torsion, Nucl. Phys. B 274, 253 (1986). [2,](#page-1-1) [3,](#page-2-1) [7,](#page-6-6) [11](#page-10-7)
- <span id="page-26-1"></span>[87] Q. Wang, B. Yang, F. Zheng, On Bismut flat manifolds, Trans. Amer. Math. Soc. 373 (2020), no. 8, 5747-5772. [3](#page-2-1)
- <span id="page-26-10"></span>[88] K. Yano. Differential geometry on complex and almost complex spaces. International Series of Monographs in Pure and Applied Mathematics, Vol 49, A Pergamon Press Book. 1965. [2](#page-1-1)
- <span id="page-26-2"></span>[89] K. Yano and S. Bochner, Curvature and Betti Numbers, Ann. of Math. Studies 32, Princeton University Press, 1953. [13,](#page-12-3) [19](#page-18-11)
- [90] S. T. Yau, Q. Zhao, F. Zheng, On Strominger Kähler-like manifolds with degenerate torsion, Trans. Amer. Math. Soc. 376 (2023), no. 5, 3063-3085. [2,](#page-1-1) [3,](#page-2-1) [4](#page-3-5)
- <span id="page-26-6"></span><span id="page-26-3"></span>[91] Y. Ye, Bismut-Einstein metrics on compact complex surfaces, arXiv:2212.04060. [2](#page-1-1)
- [92] Q. Zhao, F. Zheng, Strominger connection and pluriclosed metrics, J. Reine Angew. Math. 796 (2023), 245-267. [3,](#page-2-1) [5](#page-4-4)
- <span id="page-26-9"></span><span id="page-26-7"></span>[93] Q. Zhao, F. Zheng, *Complex nilmanifolds and Kähler-like connections*, J. Geom. Phys. 146 (2019), 103512, 9 pp. [3](#page-2-1)
- <span id="page-26-8"></span>[94] Q. Zhao, F. Zheng, On Hermitian manifolds with Bismut-Srominger parallel torsion, arXiv:2208.03071. [4,](#page-3-5) [5](#page-4-4)
- [95] Q. Zhao, F. Zheng, *Bismut Kähler-like manifolds of dimension 4 and 5*, arXiv:2[3](#page-2-1)03.09267. 3