Network Model with Application to Allergy Diseases

Konrad Furmańczyk^{1,2}, Wojciech Niemiro^{3,4}, Mariola Chrzanowska^{5,2}, Marta Zalewska²

July 14, 2023

Department of Applied Mathematics, Institute of Information Technology,

Warsaw University of Life Sciences, Poland, e-mail correspondence: konrad_furmanczyk@sggw.edu.pl 1

Department of Prevention of Environmental Hazards, Allergology and Immunology,

Medical University of Warsaw, Poland²

Faculty of Mathematics, Informatics and Mechanics University of Warsaw, Poland ³ Faculty of Mathematics and Computer Science Nicolaus Copernicus University, Torun, Poland ⁴ Department of Statistics and Econometrics, Institute of Economics and Finance,

Warsaw University of Life Sciences, Poland ⁵

Abstract

We propose a new graphical model to describe the comorbidity of allergic diseases. We present our model in two versions. First, we introduce a generative model that correctly reflects the variables' causal relationship. Then we propose an approximation of the generative model by another misspecified model that is computationally more efficient and easily interpretable. We will focus on the misspecified version, which we consider more practical. We include in the model two directed graphs, one graph of known dependency between the main binary variables (diseases), and a second graph of the dependence between the occurrence of the diseases and their symptoms. In the model, we also consider additional auxiliary variables. The proposed model is evaluated on a cross-sectional multicentre study in Poland on the ECAP database (www.ecap.pl). An assessment of the stability of the proposed model was obtained using bootstrap and jackknife techniques.

Keywords: Network Model, Bayesian Network, Logistic Regression, Allergy Diseasses

1 Introduction

Modeling dependency between different binary variables is an essential statistical task with many applications in medicine, life sciences, economics, and sociology. The basic statistical tools used in such modeling are the autologistic model ([Besag (1972)]) and network modeling based on the Ising model model ([German(1984)], [Ravikumar et al. (2010)], [Zavlis et al. (2021)], [Abeyasinghe et al. (2020)], [Dimitrakopoulos et al. (2020)], [Briganti and Linkowski (2000)]), which is a special case of autologistic model. More general information of graphical models for discrete data can be found in [Madigan at al. (1995)] and [Maathuis et al. (2019)]. The classical autologistic model ([Besag (1972)]) has been applied many times, e.g., in epidemiology, marketing, agriculture, ecology, forestry, geography, and image analysis ([Gégout-Petit et al.(2019)], [Caragea and Kaiser (2009)], [Shin et al. (2019)], [He et al. (2003)], [Koutsias N (2003)]). [Caragea and Kaiser (2009)] considered a centered autologistic model with more interpretable parameters which describe spatial dependence. [Agaskar and Lu (2013)] propose a binary vec-

tor autologistic regressive model in time and use regularization methods to estimate a sparse

network. Most common approach for estimation of the model parameters is pseudo-likelihood ([Besag (1975)]) estimation. [Zalewska et al. (2010)] described maximum likelihood via MCMC and recommended a heuristic method of estimation (averaging estimators).

Recently in the medical area, [Shin et al. (2019)] invented and applied autologistic network model for a disease progression study. Their model work with complex spatial and temporal dependencies in muscle strength among different muscles. [Shin et al. (2019)] use pseudo-likelihood ([Besag (1975)]) to estimate the model parameters. To overcome a large number of pairwise spatial associations, they apply the least absolute shrinkage and selection operator (LASSO) ([Tibshirani (1996)]).

This paper proposes a new graphical model related to but different from the autologistic model. Our model aims to describe the interdependence of allergic diseases in contrast to most studies that do not consider dependencies between allergies ([Kim et al. (2013)], [Westman et al. (2012)], [Jung et al. (2020)]).

We present our model in two versions. First, we introduce a generative model that correctly reflects the variables' causal relationship. Then we propose an approximation of the generative model by another misspecified model that is computationally more efficient and easily interpretable. We will focus on the misspecified version, which we consider more practical. In both versions of our model, we will consider typical allergic disease symptoms, family history of allergic disease, and control variables as covariate variables. We describe information about the coexistence of certain allergic diseases (binary variables) by a directed acyclic graph (DAG), which will allow us to estimate only those model parameters responsible for the strength of the relationship between individual allergic diseases. The second graph will describe the relationship between particular symptoms and the occurrence of these diseases. In the generative model, the edges lead from diseases to symptoms, corresponding to causal relations. In the misspecified model, we reverse the direction of edges: they lead from symptoms to diseases. Additionally, we consider the potential impact of a family history of allergy on the occurrence of the disease. The applied approach based on the graphs of known interdependencies of binary variables is very flexible. It will allow us to estimate only those parameters for which experts have knowledge about the presence of interdependencies. This approach will avoid the inevitable collinearity between the variables under consideration and significantly reduce computational costs. Our model was naturally divided into separate logistic models for individual allergy diseases. Each individual logistic regression is estimated by standard glm procedure and also by weighted logistic estimation ([King and Zeng (2001)]). This is because we study the so-called rare diseases. The dataset is divided into a learning and testing sample to recommend which estimation method is appropriate, and the ROC curve and average AUC on the testing sample are determined from 20 repetitions via the bootstrap and jackknife method. This approach is general and can be applied to any binary variable dependency model. It can also be extended to the high-dimensional case by adding the Lasso penalty to the conditional likelihood estimation.

The network considered in our study has a relatively small size. Therefore we can compare two versions of our model. For five (different) scenarios of covariates, we compute the 'diagnostic' probability of diseases on given symptoms for both the generative and the misspecified models. The obtained differences are negligible. We can treat the misspecified model as a good approximation for the more logically consistent but computationally expensive generative model. The rest of the paper is organized as follows. In Section 2, we introduce the proposed model with estimation. Section 3 describes the construction of the proposed model to a real big epidemiological data set with estimation results for two methods (standard logistic estimation and weighted logistic estimation), and we compare the generative model with the misspecified model. At the end of this section, we present the evaluation of the proposed model. Section 4 contains a discussion of our approach, and in Section 5, we conclude our work.

2 Network Model

2.1 Genarative model

Our proposed model contains four groups of variables. In the first group, we consider a random vector $\mathbf{Y} = (Y_1, \ldots, Y_p)^T$ with binary components. These variables will determine a patient's presence or absence of a given allergic disease. In our application we describe the interdependencies of p most common allergic diseases. Taking into account the known co-occurrence of diseases, these relationships can be described by a directed graph with the adjacency matrix $\mathbf{A} = (a_{ki})$ as follows: $a_{ki} = 1$ if Y_i is affected by Y_k and otherwise $a_{ki} = 0$. The random variables Y_i for $i = 1, \ldots, p$ are considered as vertices of a graph, where edge (k, i) occurs when Y_k affects Y_i .

In the second group, we have a random vector of symptoms of our diseases $\mathbf{S} = (S_1, \ldots, S_m)^T$. The remaining two groups consist of common factors $\mathbf{F} = (F_1, \ldots, F_l)^T$, which can affect all considered diseases, and a vector of additional covariates $\mathbf{X} = (X_1, \ldots, X_r)^T$ such as sex, age, region of a patient, etc. For example, genetic features can be considered as common factors for allergic diseases. Symptoms S_i can be continuous or discrete random variables. It is usually known which symptoms of diseases are characteristic for each disease. This knowledge can be represented in a similar way as in the case of correlations among diseases by an directed graph with adjacency matrix $\mathbf{B} = (b_{kj})$ such that: $b_{kj} = 1$ if Y_k causes S_j and otherwise $b_{kj} = 0$.

Our generative model is therefore a graphical model that includes diseases \mathbf{Y} , symptoms \mathbf{S} , common factors \mathbf{F} and additional covariates \mathbf{X} . The structure of this graph is described by edges among \mathbf{Y}, \mathbf{S} variables given by matrices \mathbf{A}, \mathbf{B} , and all edges leading from \mathbf{F}, \mathbf{X} variables to all components of \mathbf{Y}, \mathbf{S} . We assume that the graph corresponding to the adjacency matrix \mathbf{A} is acyclic. Consequently, the whole consider graph is a directed acyclic graph (DAG). The conditional probability distribution of \mathbf{Y}, \mathbf{S} is given by

$$P(\mathbf{Y} = \mathbf{y}, \mathbf{S} = \mathbf{s} | \mathbf{F} = \mathbf{f}, \mathbf{X} = \mathbf{x}) = \prod_{i=1}^{p} P(Y_i = y_i | \mathbf{Y}_{pa}(Y_i), \mathbf{F} = \mathbf{f}, \mathbf{X} = \mathbf{x})$$

$$\times \prod_{j=1}^{m} P(S_j = s_j | \mathbf{Y}_{pa}(S_j), \mathbf{F} = \mathbf{f}, \mathbf{X} = \mathbf{x}),$$
(1)

where $\mathbf{Y}_{pa}(Y_i) = \{Y_k : Y_k \to Y_i\}$ is a set of diseases which affect the occurrence of disease Y_i , $\mathbf{Y}_{pa}(S_j) = \{Y_k : Y_k \to S_j\}$ is a set of diseases which cause symptom S_j . We assume the following parametric form of conditional distribution:

$$\log\left(\frac{P(Y_i=1|\mathbf{Y}_{pa}(Y_i), \mathbf{F}=\mathbf{f}, \mathbf{X}=\mathbf{x})}{P(Y_i=0|\mathbf{Y}_{pa}(Y_i), \mathbf{F}=\mathbf{f}, \mathbf{X}=\mathbf{x})}\right) = \omega_{0i} + \sum_{k=1}^p a_{ki}\omega_{ki}Y_k + \mathbf{x}^T\boldsymbol{\alpha}_i + \mathbf{f}^T\boldsymbol{\beta}_i,$$
(2)

$$\log\left(\frac{P(S_j=1|\mathbf{Y}_{pa}(S_j), \mathbf{F}=\mathbf{f}, \mathbf{X}=\mathbf{x})}{P(S_j=0|\mathbf{Y}_{pa}(S_j), \mathbf{F}=\mathbf{f}, \mathbf{X}=\mathbf{x})}\right) = \gamma_{0j} + \sum_{k=1}^p b_{kj} \gamma_{kj} Y_k + \mathbf{x}^T \boldsymbol{\delta}_j + \mathbf{f}^T \boldsymbol{\epsilon}_j.$$
(3)

where model parameters: $\omega_{0i} \in R, \omega_{ki} \in R, \alpha_i \in R^r, \beta_i \in R^l, \gamma_{0j} \in R, \gamma_{kj} \in R, \delta_j \in R^r, \epsilon_j \in R^l$. Since the conditional probability (1) consists of the product of p + m probabilities (factors), the parameters of each factor can be estimated separately by fitting a standard logistic regression procedure.

2.2 Misspecified model

Unfortunately, the model presented in the previous subsection is computationally demanding, and moreover, its parameters are not easy to interpret. We propose using another model that does not reflect causal relations between variables correctly but is computationally easier in a big network and has parameters with simple, intuitive meanings. We say that this model is misspecified. We change the direction of edges joining symptoms and diseases. Entries of adjacency matrix **B** will now be interpreted as follows: $b_{ij} = 1$ indicates the presence of arrow $Y_i \leftarrow S_j$. We assume that the remaining edges of the graph are the same as in generative model. In the misspecified model, equation (1) is replaced by equation (4), and equations (2)-(3) are replaced by equation (5) as follows:

$$P(\mathbf{Y} = \mathbf{y} | \mathbf{S} = \mathbf{s}, \mathbf{F} = \mathbf{f}, \mathbf{X} = \mathbf{x}) = \prod_{i=1}^{p} P(Y_i = y_i | \mathbf{Y}_{pa}(Y_i), \mathbf{S}_{pa}(Y_i), \mathbf{F} = \mathbf{f}, \mathbf{X} = \mathbf{x}), \quad (4)$$

where $\mathbf{S}_{pa}(Y_i) = \{S_j : Y_i \leftarrow S_j\}$ is a set of symptoms related to occurrence of disease Y_i . Similarly, as in generative model, we assume a log-linear form of conditional distributions. To simplify notation, we use the same symbols for the parameters for both models.

$$\log\left(\frac{P(Y_i = 1 | \mathbf{Y}_{pa}(Y_i), \mathbf{S}_{pa}(Y_i), \mathbf{F} = \mathbf{f}, \mathbf{X} = \mathbf{x})}{P(Y_i = 0 | \mathbf{Y}_{pa}(Y_i), \mathbf{S}_{pa}(Y_i), \mathbf{F} = \mathbf{f}, \mathbf{X} = \mathbf{x})}\right) = \omega_{0i} + \sum_{k=1}^{p} a_{ki} \omega_{ki} Y_k + \sum_{j=1}^{m} b_{ij} \gamma_{ij} S_j + \mathbf{x}^T \boldsymbol{\alpha}_i + \mathbf{f}^T \boldsymbol{\beta}_i.$$
(5)

We do not always have appropriate sample sizes for rare diseases and work with imbalanced datasets. In such cases, we may improve prediction accuracy for logistic regression using weighted logistic regression ([King and Zeng (2001)], [Zhang et al. (2021)]) or apply a machine learning algorithm such as use SMOTE Simple Genetic Algorithm ([Tallo (2018)]) to determine the sampling rate of each example in order to get unequal synthetic samples or using undersampling or oversampling ([Zhang et al. (2019)]). However, resampling techniques do not easily transfer to dependent logistic regression equations. For this reason, in the paper, we use weighted regression as in [King and Zeng (2001)]. Following the approach of [King and Zeng (2001)] we penalized misclassification costs of events and non-events differently by penalty weights w_1 and w_0 in the log-likelihood function for each *i* equation

$$min_{\boldsymbol{\theta}_i} \left\{ -w_1 \sum_{j=1}^n y_{ij} log(\sigma(\mathbf{z}_j^T \boldsymbol{\theta}_i)) - w_0 \sum_{j=1}^n (1 - y_{ij}) log(1 - \sigma(\mathbf{z}_j^T \boldsymbol{\theta}_i)) \right\},\$$

where *n* is a sample size, $w_1 = \frac{\tau_i}{\bar{y}_i}$ and $w_0 = \frac{1-\tau_i}{1-\bar{y}_i}$, and τ_i denoting the population fraction of events induced by choice-based sampling and \bar{y}_i denoting the sample proportion of events, $\boldsymbol{\theta}_i$ is a vector of all parameters, \mathbf{z}_j is a vector of all predictors, and $\sigma(x) = \frac{exp(x)}{1+exp(x)}$.

3 A Model of Allergic Diseases

The proposed model of disease interdependence will be used to investigate the prevalence of allergic diseases in Poland based on a big epidemiological study ECAP ([Samoliński et al. (2014)]). The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Ethics Committee of the Medical University of Warsaw, Poland (KB/206/2005).



Figure 1: Graph with adjacency matrix **A**.

The study method involved the use of questionnaires adapted for Central and Eastern Europe based on the European Community Respiratory Health Survey II (ECRHS II) ([Tallo (2018)]) and International Study of Asthma and Allergies in Childhood (ISAAC) ([Strachan (1989)]), which had been used as part of a larger project, titled the Implementation of a System for the Prevention and Early Detection of Allergic Diseases in Poland ([Tesse (2011)]). The project was conducted in eight urban areas and one rural area. The study had two stages; the first stage involved grouping the 22,500 respondents based on their questionnaire responses using of a Personal Digital Assistant (PDA); the second stage involved complementary examination (4,783 patients) of a subgroup of stage I respondents who underwent a medical examination. The final data set contains 18,617 units (cases, response) and 1,225 variables (mostly binary).

3.1 The structure of the model

In the first group for the interdependence study, we selected the following diseases, allergic: Y_1 -atopic asthma, Y_2 -intermittent allergic rhinitis, Y_3 -chronic allergic rhinitis, Y_4 -allergic dermatitis, Y_5 -food allergy. Figure 1 illustrates the assumed dependencies between these allergic diseases. The structure of this graph is based on the expert knowledge taken from the medical literature and on discussion with medical doctors ([Raciborski et al. (2019)], [Krzych-Fałta et al. (2016)]).

In the second group, we consider typical symptoms of those allergic diseases: S_1 -Have you had wheezing or whistling in your chest at any time in the last 12 months?; S_2 - Have you ever had a problem with sneezing or a runny or blocked nose when you did not have fever, a cold, or the flu?; S_3 -Have you ever had eczema or any other form of skin allergy? Additionally we consider in this group history of allergy diseases in the family as a common factors: F_1 -Does anyone in your immediate family suffer from allergies? - mother; F_2 -Does anyone in your immediate family



Figure 2: Graph with adjacency matrix **B** (direction of arrows is as the misspecified model).

suffer from allergies? - father; F_3 -Does anyone in your immediate family suffer from allergies? - siblings of the child being tested; F_4 -Does anyone in your immediate family suffer from allergies? -grandparents on mother's side; F_5 -Does anyone in your immediate family suffer from allergies? - grandparents on father's side. Figure 2 shows a relationship graph of the considered allergic diseases and their typical symptoms. This presence of the edges in this graph is also based on expert knowledge (see remarks concerning Figure 1). Note that the direction of arrows lead from symptoms to diseases which corresponds to the misspecified model.

In the last group, we consider control covariates for respondents such as: X_1 age of patients with three age group: children 6-7 y.o., children 13-14 y.o., and adults (20-44 y.o.). This variable we replaced by new two binary variables: X_1 for children 13-14 y.o. and X_2 for adults., X_3 binary variables with 1 for urban area, X_4 -sex (binary variable, 1 for male).

3.2 Generative and misspecified models of allergy diseases

We recall that the generative model is a Bayesian Network in which diseases cause symptoms. Conditionally on covariates \mathbf{F} and \mathbf{X} the joint probability distribution of (\mathbf{Y}, \mathbf{S}) is determined by the set of conditional probabilities:

 $P(\mathbf{Y}, \mathbf{S}) = P(Y_1|Y_2, Y_3, Y_4) P(Y_2|Y_4) P(Y_3|Y_4) P(Y_4|Y_5) P(Y_5) P(S_1|Y_1) P(S_2|Y_2, Y_3) P(S_3|Y_4).$

(We omitted \mathbf{F} and \mathbf{X} in this formula). For all conditional probabilities, we assume the logistic form of those probabilities. We may estimate all the parameters of the model by estimating each conditional probability separately by standard glm procedure.

Now we turn to the misspecified model. Using the DAG structure of graph with adjacency matrices \mathbf{A}, \mathbf{B} , conditionally on covariates \mathbf{F} and \mathbf{X} the conditional distribution of \mathbf{Y} given

symptoms \mathbf{S} has the form

$$P(\mathbf{Y}|\mathbf{S}) = P(Y_1|Y_2, Y_3, Y_4, S_1)P(Y_2|Y_4, S_2)P(Y_3|Y_4, S_2)P(Y_4|Y_5, S_3)P(Y_5).$$

Now we are going to give specific equations restricting attention to the misspecified model only. The first equation of our model concerns the logit for asthma Y_1 conditionally on symptom of asthma S_1 , family history of allergy diseases F_1, F_2, F_3, F_4, F_5 , and diseases Y_2, Y_3, Y_4 , and control covariates X_1, X_2, X_3, X_4

$$logit_1 = \omega_{01} + \sum_{j=1}^4 \alpha_{j1}X_j + \sum_{j=1}^5 \beta_{j1}F_j + \gamma_{11}S_1 + \sum_{j=2}^4 \omega_{j1}Y_j.$$

The second equation presents the logit for intermittent allergic rhinitis Y_2 conditionally on symptom of allergic rhinitis S_2 , family history of allergy diseases F_1, F_2, F_3, F_4, F_5 and allergic dermatitis Y_4 , and control covariates X_1, X_2, X_3, X_4

$$logit_2 = \omega_{02} + \sum_{j=1}^4 \alpha_{j2} X_j + \sum_{j=1}^5 \beta_{j2} F_j + \gamma_{22} S_2 + \omega_{42} Y_4.$$

Next, we present logit form for chronic allergic rhinitis Y_3 conditionally on symptom of allergic rhinitis S_2 , allergic dermatitis Y_4 , family history of allergy diseases F_1, F_2, F_3, F_4, F_5 , and Y_4 , and control covariates X_1, X_2, X_3, X_4

$$logit_3 = \omega_{03} + \sum_{j=1}^4 \alpha_{j3} X_j + \sum_{j=1}^5 \beta_{j3} F_j + \gamma_{32} S_2 + \omega_{43} Y_4.$$

Logit for allergic dermatitis Y_4 conditionally on symptom of allergic dermatitis S_3 , food allergy Y_5 , family history of allergy diseases F_1, F_2, F_3, F_4, F_5 and control covariates X_1, X_2, X_3, X_4 has the form

$$logit_4 = \omega_{04} + \sum_{j=1}^4 \alpha_{j4} X_j + \sum_{j=1}^5 \beta_{j4} F_j + \gamma_{43} S_3 + \omega_{54} Y_5.$$

In both generative and misspecified models, estimation is carried out for each equation using the standard procedure for estimating logistic regression coefficients.

Next, we compute the 'diagnostic' probabilities of diseases given symptoms for both models with estimated parameters: $P(Y_1 = 1|Y_2, Y_3, Y_4, S_1), P(Y_2 = 1|Y_4, S_2), P(Y_3 = 1|Y_4, S_2), P(Y_4 = 1|Y_5, S_3), P(Y_5)$. It is worth noting that in the case of a large network, it would not be possible to calculate $P(\mathbf{Y}|\mathbf{S}, \mathbf{F}, \mathbf{X})$ or $P(Y_i|\mathbf{S}, \mathbf{F}, \mathbf{X})$ exactly in the generative model. In this situation, the misspecified model has an advantage over the generative model. The two methods can be compared in the case of a small network as that considered here. We consider five scenarios of covariates \mathbf{X}, \mathbf{F} :

- Case 1: rural area, children 13-14 y.o., male, without allergy history in family $F_1 = \ldots = F_5 = 0$ and without symptoms $S_1 = S_2 = S_3 = 0$;
- Case 2: rural area, children 13-14 y.o., male, without allergy history in family $F_1 = \ldots = F_5 = 0$ and with symptoms $S_1 = S_2 = S_3 = 1$;
- Case 3: urban area, children 13-14 y.o., male, without allergy history in family $F_1 = \ldots = F_5 = 0$ and with symptoms $S_1 = S_2 = S_3 = 1$;

case	method	$P(Y_1 Y_2 = 0, Y_3 = 0, Y_4 = 0, S_1)$	$P(Y_2 Y_4 = 0, S_2)$	$P(Y_3 Y_4 = 0, S_2)$	$P(Y_4 Y_5 = 0, S_3)$
Case 1	exact	0.021	0.081	0.077	0.024
	misspec.	0.023	0.085	0.080	0.015
Case 2	exact	0.103	0.282	0.322	0.208
	misspec.	0.088	0.270	0.307	0.081
Case 3	exact	0.074	0.212	0.321	0.248
	misspec.	0.064	0.202	0.306	0.116
Case 4	exact	0.015	0.056	0.074	0.007
	misspec.	0.016	0.060	0.080	0.022
Case 5	exact	0.120	0.332	0.359	0.233
	misspec.	0.130	0.295	0.314	0.210

Table 1: Results for comparison the generative model (exact computation) and misspecified model

Table 2: Results for comparison the generative model (exact) and misspecified model

case	method	$P(Y_1 Y_2 = 1, Y_3 = 1, Y_4 = 1, S_1)$	$P(Y_2 Y_4 = 1, S_2)$	$P(Y_3 Y_4 = 1, S_2)$	$P(Y_4 Y_5 = 1, S_3)$
Case 1	exact	0.597	0.104	0.134	0.024
	misspec.	0.566	0.097	0.125	0,044
Case 2	exact	0.886	0.326	0.461	0.524
	misspec.	0.842	0.299	0.421	0.216
Case 3	exact	0.845	0.250	0.463	0.581
	misspec.	0.793	0.226	0.420	0.290
Case 4	exact	0.509	0.073	0.130	0.029
	misspec.	0.482	0.068	0.124	0.064
Case 5	exact	0.902	0.398	0.511	0.561
	misspec.	0.893	0.326	0.429	0.452

- Case 4: urban area, children 13-14 y.o., male, without allergy history in family $F_1 = \ldots = F_5 = 0$ and without symptoms $S_1 = S_2 = S_3 = 0$;
- Case 5: urban area, children 13-14 y.o., male, with allergy history in family $F_1 = \ldots = F_5 = 1$ and with symptoms $S_1 = S_2 = S_3 = 1$.

The results are presented in Tables 1-2. The obtained difference between the two models is negligible.

3.3 Results of estimation for the misspecified model

Model estimation is performed separately for each equation (see formula (4)-(5)) using the usual glm procedure for logistic regression in the first scenario, and in the second scenario, we use weighted logistic regression (with weights as in Section 2).

$logit_i$	ω_{0i}	α_{1i}	α_{2i}	α_{3i}	α_{4i}	β_{1i}	β_{2i}	β_{3i}	β_{4i}	β_{5i}
i=1	-5.933	0.355	0.216	-0.336	0.412	-0.075	-0.108	0.194	0.029	0.736
i=2	-4.000	0.287	0.364	-0.384	-0.038	-0.045	0.284	0.313	0.090	-0.135
i=3	-4.741	0.335	0.271	-0.004	0.334	0.055	0.186	-0.039	-0.144	-0.022
i=4	-6.073	0.195	-0.623	0.395	-0.103	0.331	0.220	0.061	-0.380	0.470

Table 3: Estimation results for standard logistic regression - Part 1

Table 4: Estimation results for standard logistic regression - Part 2

$logit_i$	γ_{i1}	γ_{i2}	γ_{i3}	ω_{2i}	ω_{3i}	ω_{4i}	ω_{5i}
i=1	1.412	-	-	1.265	2.040	0.713	-
i=2	-	1.379	-	-	-	0.143	-
i=3	-	1.628	-	-	-	0.496	-
i=4	-	-	1.780	-	-	-	1.132

According to work of [Samoliński et al. (2014)] we assume that the population fraction in Poland for considered allergy diseases are as follows $\tau_1 = 11\%$, $\tau_2 = 20\%$, $\tau_3 = 4\%$, $\tau_4 = 7\%$, $\tau_5 = 10\%$.

The results for estimation for two scenarios are given in Tables 3-6. The standard errors for standard logistic regression coefficients estimation are given in Tables 7-8. Next, we present the odds ratio with the asymptotic 0.95 confidence interval (CI) for the standard logistic regression and the odds ratio for weighted logistic regression (see Tables 9-14).

3.4 Evaluation of the misspecified model

The parameter estimator's accuracy and precision and the model's robustness for both scenarios were assessed using bootstrap and jackknife techniques. We draw 20 ordinary non-parametric bootstrap samples, calculate regression coefficients on each bootstrap sample, treat the whole real sample as a test sample, draw the ROC for it, and calculate the AUC. We also used the jackknife (10-fold cross-validation) method: we drew 10% of the sample for testing and treated the other 90% as a training sample. We repeated the experiment 20 times. Table 15 shows the AUC values for the weighted logistic regression estimate (weight), and the averaged AUC values

Table 5: Estimation results for weighted logistic regression - Part 1

$logit_i$	ω_{0i}	α_{1i}	α_{2i}	α_{3i}	α_{4i}	β_{1i}	β_{2i}	β_{3i}	β_{4i}	β_{5i}
i=1	-5.053	0.356	0.206	-0.349	0.447	-0.039	-0.092	0.181	0.024	0.737
i=2	-3.544	0.289	0.355	-0.377	-0.042	-0.041	0.285	0.310	0.088	-0.154
i=3	-6.212	0.319	0.263	0.024	0.341	0.061	0.186	-0.042	-0.163	0.001
i=4	-3.301	0.139	-0.646	0.278	-0.104	0.414	0.322	-0.009	-0.480	0.409

Table 6: Estimation results for weighted logistic regression - Part 2

$logit_i$	γ_{i1}	γ_{i2}	γ_{i3}	ω_{2i}	ω_{3i}	ω_{4i}	ω_{5i}
i=1	1.392	-	-	1.392	2.085	0.753	-
i=2	-	1.377	-	-	-	0.140	-
i=3	-	1.628	-	-	-	0.450	-
i=4	-	-	1.800	-	-	-	1.302

Table 7: The standard errors of estimation for standard logistic regression - Part 1 $\,$

$logit_i$	ω_{0i}	α_{1i}	α_{2i}	α_{3i}	α_{4i}	β_{1i}	β_{2i}	β_{3i}	β_{4i}	β_{5i}
i=1	0.350	0.197	0.190	0.219	0.149	0.206	0.228	0.180	0.304	0.304
i=2	0.212	0.121	0.113	0.133	0.088	0.123	0.134	0.107	0.184	0.239
i=3	0.223	0.116	0.110	0.142	0.086	0.117	0.130	10.110	0.185	0.224
i=4	0.393	0.145	0.163	0.244	0.127	0.150	0.169	0.151	0.243	0.260

Table 8: The standard errors of estimation for standard logistic regression - Part 2

$logit_i$	γ_{i1}	γ_{i2}	γ_{i3}	ω_{2i}	ω_{3i}	ω_{4i}	ω_{5i}
i=1	0.151	-	-	0.184	0.158	0.222	-
i=2	-	0.096	-	-	-	0.164	-
i=3	-	0.096	-	-	-	0.148	-
i=4	-	-	0.167	-	-	-	0.147

Table 9: The OR with 0.95 CI for estimation results for standard logistic regression - Part 1

$logit_i$	$\exp(\alpha_{1i})$	$\exp(\alpha_{2i})$	$\exp(\alpha_{3i})$	$\exp(\alpha_{4i})$
i=1	1.426(0.969; 2.098)	1.121(0.855;1.801)	0.715(0.465;1.098)	1.510(1.127;2.022)
i=2	1.332(1.051; 1.689)	1.439(1.153;1.796)	0.681(0.525; 0.884)	0.963(0.810; 1.144)
i=3	1.398(1.114;1.755)	1.311(1.057; 1.627)	0.996(0.754; 1.316)	1.397(1.180; 1.653)
i=4	1.215(0.915; 1.615)	0.536(0.390; 0.738)	1.484(0.920; 2.395)	0.902(0.703;1.157)

Table 10: The OR with 0.95 CI for estimation results for standard logistic regression - Part 2

$logit_i$	$\exp(\beta_{1i})$	$\exp(\beta_{2i})$	$\exp(\beta_{3i})$	$\exp(\beta_{i4})$	$\exp(\beta_{i5})$
i=1	0.928(0.620;1.389)	0.898(0.574;1.403)	1.214(0.853;1.728)	1.029(0.567; 1.868)	2.088(1.150;3.788)
i=2	0.956(0.751;1.217)	1.328(1.022; 1.727)	1.368(1.109; 1.687)	1.094(0.763; 1.569)	0.874(0.547; 1.396)
i=3	1.057(0.840; 1.329)	1.204(0.934;1.554)	0.962(0.775;1.193)	0.866(0.603; 1.244)	0.978(0.631; 1.517)
i=4	1.392(1.038;1.868)	1.246(0.895; 1.735)	1.063(0.791; 1.429)	0.684(0.425;1.101)	1.600(0.961; 2.663)

Table 11: The OR with 0.95 CI for estimation results for standard logistic regression - Part 3 $\,$

$logit_i$	$\exp(\gamma_{i1})$	$\exp(\gamma_{i2})$	$\exp(\gamma_{i3})$
i=1	4.104(3.053;5.518)	-	-
i=2	-	4.015(3.326; 4.846)	-
i=3	-	5.094(4.220; 6.148)	-
i=4	-	-	5.930(4.275; 8.226)

Table 12: The OR with 0.95 CI for estimation results for standard logistic regression - Part 4

$logit_i$	$\exp(\omega_{2i})$	$\exp(\omega_{3i})$	$\exp(\omega_{4i})$	$\exp(\omega_{5i})$
i=1	3.543(2.470; 5.082)	7.691(5.642;10.482)	2.040(1.320; 3.152)	-
i=2	-	-	1.154(0.837; 1.591)	-
i=3	-	-	1.642(1.229; 2.195)	-
i=4	-	-	-	3.102(2.325; 4.138)

Table 13: The OR for estimation results for weighted logistic regression - Part 1

$logit_i$	$\exp(\alpha_{1i})$	$\exp(\alpha_{2i})$	$\exp(\alpha_{3i})$	$\exp(\alpha_{4i})$	$\exp(\beta_{1i})$	$\exp(\beta_{2i})$	$\exp(\beta_{3i})$	$\exp(\beta_{4i})$	$\exp(\beta_{5i})$
i=1	1.428	1.229	0.705	1.564	0.962	0.912	1.198	1.024	2.090
i=2	1.335	1.426	0.686	0.959	0.960	1.330	1.363	1.092	0.857
i=3	1.376	1.301	1.024	1.406	1.063	1.204	0.959	0.850	1.001
i=4	1.149	0.524	1.320	0.901	1.513	1.380	0.991	0.619	1.505

Table 14: The OR for estimation results for weighted logistic regression - Part 2 $\,$

$logit_i$	$\exp(\gamma_{i1})$	$\exp(\gamma_{i2})$	$\exp(\gamma_{i3})$	$\exp(\omega_{2i})$	$\exp(\omega_{3i})$	$\exp(\omega_{4i})$	$\exp(\omega_{5i})$
i=1	4.023	-	-	4.023	8.045	2.123	-
i=2	-	3.963	-	-	-	1.150	-
i=3	-	5.094	-	-	-	1.568	-
i=4	-	-	6.050	-	-	-	3.677

$logit_i$	w eight	boot+weight	boot	jackkn+weight	jackkn
i=1	0.8406	0.8258	0.8470	0.8231	0.8165
i=2	0.6704	0.6907	0.6986	0.6700	0.6857
i=3	0.7234	0.7196	0.7201	0.7259	0.7215
i=4	0.7971	0.7936	0.7931	0.7861	0.7921

Table 15: AUC for each logit



Figure 3: ROC for *logit*₁ for unweighted (black curve) and weighted (blue curve) estimation.

for bootstrap (with and without weights) and jackknife (with and without weights).

In Figure 3-7, we may assess the stability and good accuracy and precision of the first logit equation. The rest of the results (Fig1S-Fig15S) are collected in supplementary materials on GitHub ([Supplement (2021)]). The ROC curves for the standard logistic regression and weighted estimations are almost identical except for the case of $logit_2$ (Fig 1S), where better results are obtained from the standard glm estimation method. Using bootstrap and jackknife, even for a few repetitions, showed quite good stability of the obtained results. The average value of AUC (Tab 15) confirmed this tendency. In general, the method without weights gave better AUC results except in the case of $logit_4$. However, the differences were quite negligible. When the jackknife method was used, also clear difference was not observed.



Figure 4: ROC for $logit_1$ for unweighted estimation-bootstrap.



Figure 5: ROC for $logit_1$ for weighted estimation-bootstrap.



Figure 6: ROC for $logit_1$ for unweighted estimation-jackknife.



Figure 7: ROC for $logit_1$ for weighted estimation-jackknife.

3.5 Results of statistical inference

Since the differences in estimation by the two considered methods turned out to be negligible, we will make model inference based on the parameter estimation without weighting. Next, we present the most influential factors affecting each of considered disease, such as other diseases, symptoms, and other additional variables. Based on Table 9-12, we may conclude that atopic asthma Y_1 is most associated with its symptom: wheezing or whistling in chest S_1 (OR =4.104), intermittent allergic rhinitis Y_2 (OR = 3.543), chronic allergic rhinitis Y_3 (OR = 7.691), grandparents on father's side has allergy F_5 (OR = 2.088) and allergic dermatitis Y_4 (OR =2.040). The intermittent allergic rhinitis Y_2 is most associated with its symptom: problem with sneezing or a runny or blocked nose S_2 (OR = 4.015) and allergic dermatitis Y_4 (OR = 1.154). The chronic allergic rhinitis Y_3 is most associated with its symptom: problem with sneezing or a runny or blocked nose S_2 (OR = 5.094) and allergic dermatitis Y_4 (OR = 1.642). The allergic dermatitis Y_4 is most associated with its symptom: problem with sneezing or a runny or blocked nose S_2 (OR = 5.094) and allergic dermatitis Y_4 (OR = 1.642). The allergic dermatitis Y_4 is most associated with its symptom: problem with sneezing or a runny or blocked nose S_2 (OR = 5.094) and allergic dermatitis Y_4 (OR = 1.642). The allergic dermatitis Y_4 is most associated with its symptom: having eczema or skin allergy S_3 (OR = 5.930) and food allergy Y_5 (OR = 3.102).

4 Discussion

Multimorbidity in allergy has been studied, among others, by [Kim et al. (2013)],

[Westman et al. (2012)], [Jung et al. (2020)].

This problem has been also considered in [Raciborski et al. (2019)], [Krzych-Fałta et al. (2016)] in a cross-sectional multicentre study in Poland on the ECAP database ([Samoliński et al. (2014)]). These studies were based on the fitting of single logistic models that did not take into account the correlations between the studied diseases. In the present approach, which is based on the conditional graphical models, we took into account such dependency through a directed acyclic graph of the relationships between the diseases under study and a directed graph of the dependency between the diseases and their symptoms. The presented model of the relationship between allergic diseases is based on discussions with medical doctors dealing with allergies and is a certain generalization of papers [Raciborski et al. (2019)], [Krzych-Fałta et al. (2016)]. The proposed model can be used in studies of associations of other diseases and, in general, in the study of correlations of complex systems. In this study, we used estimation based on a separate logistic regression estimation for the individual equations of the model as well as a weighted version of this estimation. In the case of biased data and rare diseases, we recommend using weighted logistic regression. In our case, the obtained results were very close using standard logistic and weighted logistic regressions. Due to the nature of our task, we considered the low-dimensional case where the number of observations n is greater than the number of features p. Naturally, the proposed approach can be generalized to the high-dimensional case p > n by adding the Lasso or Ridge penalty for log-likelihood for each logit model separately.

5 Conclusions

In our study, we presented some conditional graphical model in two versions, taking into account the known associations between allergic diseases and the symptoms of these diseases together with additional factors such as the family history of allergic diseases, and introduced additional control variables into the model. We compare two versions of our model, one in which diseases cause their symptoms (the generative model) and the misspecified model where the direction of edges is opposite. For five different scenarios of covariates, we compute the 'diagnostic' probability of diseases on given symptoms for both model versions. The obtained differences are negligible. The misspecified model approximates these diagnostic probabilities of allergic diseases very well and is less computationally expensive than calculating the exact inverse probabilities in the generative model. We will focus on the misspecified version, which we consider more practical. Parameter estimation for both versions of model were performed using the standard glm procedure for each logistic regression separately. Due to the rarity of the diseases considered, logistic regression weighting was proposed ([King and Zeng (2001)]). Evaluation of the model using bootstrap and jackknife techniques yielded average AUCs ranging from 0.67 to 0.84 (Table 15), indicating fairly high stability of the results. In general, weighting did not really help in estimating model parameters, except for the equation for $logit_4$.

The proposed model can be easily extended by adding other potential factors influencing the occurrence of the diseases or for general dependency complex model of binary variables.

Conflict of interest: The authors declare no conflict of interest.

References

- [Abeyasinghe et al. (2020)] Abeyasinghe, P.M. and Aiello, M. and Nichols, E.S. and Cavaliere, C. and Fiorenza, S. and Masotta, O. and Borrelli, P. and Owen, A.M. and Estraneo, A. and Soddu, A. (2020), Consciousness and the dimensionality of DOC patients via the generalized ising model, *Journal of Clinical Medicine* 9(5) doi=10.3390/jcm9051342
- [Agaskar and Lu (2013)] Agaskar, A. and Lu, Y.M. (2013), Alarm: a logistic auto-regressive model for binary processes on networks, *IEEE Global Conference on Signal and Information Processing, Austin, TX, 305–308* https://doi.org/10.1109/GlobalSIP.2013.6736876
- [Besag (1972)] Besag J. E. (1972), Nearest-Neighbour Systems and the Auto-Logistic Model for Binary Data Journal of the Royal Statistical Society. Series B (Methodological), 34(1), 75-83
- [Besag (1975)] Besag J. E. (1975), Statistical analysis of non-lattice data, The Statistician, 24(3), 179–195
- [Briganti and Linkowski (2000)] Briganti, G. and Linkowski, P. (2000), Exploring network structure and central items of the Narcissistic Personality Inventory, International Journal of Methods in Psychiatric Research, 29(1), doi: 10.1002/mpr.1810
- [Caragea and Kaiser (2009)] Caragea, P.C. and Kaiser, M.S. (2009), Autologistic models with interpretable parameters, Journal of Agricultural, Biological, and Environmental Statistics, 14, 281–300
- [Dimitrakopoulos et al. (2020)] Dimitrakopoulos, P., Sfikas, G., Nikou, C., (2020), ISING-GAN: Annotated Data Augmentation with a Spatially Constrained Generative Adversarial Network, Proceedings - International Symposium on Biomedical Imaging, 1600-1603, doi: 10.1109/ISBI45749.2020.9098618
- [German(1984)] Geman, S. and Geman, D. (1984), Stochastic relaxation, Gibbs distributions, and the Bayesian restoration of images, *IEEE Trans. PAMI 6 721–741*
- [Gégout-Petit et al.(2019)] Gégout-Petit A, Guérin-Dubrana L, Li S (2019), A new centered spatio-temporal autologistic regression model with an application to local spread of plant diseases, Spat. Stat. 31 100361

- [Haspeslagh et al. (2018)] Haspeslagh E, Heyndickx I, Hammad H et al. (2018), The hygiene Hypothesis: Immunological mechanisms of airway tolerance, Curr. Opin. Immunol. 54, 102-108
- [He et al. (2003)] He, F., Zhou, J. & Zhu, H. (2003), The hygiene Hypothesis: Autologistic regression model for the distribution of vegetation, *JABES 8, 205*
- [Jung et al. (2020)] Jung, S., Lee, S. Y., Yoon, J., Cho, H. J., Kim, Y. H., Suh, D. I., Yang, S. I., Kwon, J. W., Jang, G. C., Sun, Y. H., Woo, S. I., Youn, Y. S., Park, K. S., Lee, E., Cho, H. J., Kook, M. H., Yi, H. R., Chung, H. L., Kim, J. H., Kim, H. Y., Hong, S. J. (2020), Risk Factors and Comorbidities Associated With the Allergic Rhinitis Phenotype in Children According to the ARIA Classification, *Allergy, asthma & immunology research vol. 12,1* (2020): 72-85 doi:10.4168/aair.2020.12.1.72
- [Kim et al. (2013)] Kim, H. Y., Kwon, E. B., Baek, J. H., Shin, Y. H., Yum, H. Y., Jee, H. M., Yoon, J. W., Han, M. Y. (2013), Prevalence and comorbidity of allergic diseases in preschool children, Korean Journal of Pediatrics, 56(8), 338–342 doi.org/10.3345/kjp.2013.56.8.338
- [King and Zeng (2001)] King G and Zeng L, (2001), Logistic regression in rare events data, Polit. Anal. 9 (2001), 137–163
- [Krzych-Fałta et al. (2016)] Krzych-Fałta E, Furmańczyk K, Piekarska B, Tomaszewska A, Sybilski A, Samoliński B. (2016), Allergies in urban versus countryside settings in Poland as part of the Epidemiology of the Allergic Diseases in Poland (ECAP) study—challenge the early differential diagnosis, Adv Dermatol Allergol., 33(5), 359–368 doi: 10.5114/pdia.2016.61338 - DOI - PMC – PubMed
- [Koutsias N (2003)] Koutsias N (2003), An autologistic regression model for increasing the accuracy of burned surface mapping using Landsat Thematic Mapper data, International Journal of Remote Sensing 24(10):2199-2204
- [Maathuis et al. (2019)] Maathuis, M. Drton, M. Lauritzen, S., Wainwright, M., eds. (2019), Handbook of Graphical Models, *Chapman & Hall/CRC Press*, 2019
- [Madigan at al. (1995)] Madigan D, York J, Allard D (1995), Bayesian graphical models for discrete data, Int Stat Rev 63(2):215
- [Raciborski et al. (2019)] Raciborski F, Bousuet J, Namyslowski A, Krzych-Falta E, Tomaszewska A, Piekarska B, Samel-Kowalik P, Bialoszewski A, Walkiewicz A, Lipiec A, Wojas O, Samolinski K, Szylling A, Zielinski W, Sybilski A, Grabczewska A, Samolinski B (2019), Dissociating polysensitization and multimorbidity in children and adults from a Polish general population cohort, *Clinical and Translational Allergy Vol 9, No 1*, doi: 10.1186/s13601-019-0246-y
- [Ravikumar et al. (2010)] Ravikumar, P., Wainwright, M. J. and Lafferty, J. (2010). Highdimensional Ising model selection using l1-regularized logistic regression, Ann. Statist. 38 1287–1319
- [Samoliński et al. (2014)] Samoliński B, Raciborski F, Lipiec A i wsp.. (2014), Epidemiologia Chorób Alergicznych w Polsce (ECAP), Alergol Pol 2014; 1: 10-8
- [Shin et al. (2019)] Shin YE, Sang H, Liu D, Ferguson TA, Song PXK. (2019), Autologistic network model on binary data for disease progression study, *Biometrics*. 75(4), 1310-1320 doi: 10.1111/biom.13111

- [Strachan (1989)] Strachan D. P. (1989), Hay fever, hygiene, and household size, BMJ (Clinical research ed.), 299(6710), 1259–1260 doi.org/10.1136/bmj.299.6710.1259
- [Supplement (2021)] Supplement (2021) Furmańczyk K, Niemiro W, Chrzanowska M, Zalewska M, Supplementary Material to the paper 'Network Model with Application to Allergy Diseases' https://github.com/kfurmanczyk/Network-_Allergy/blob/main/Supplement.pdf
- [Tallo (2018)] Tallo T. E. and A. Musdholifah A. (2018), The Implementation of Genetic Algorithm in Smote (Synthetic Minority Oversampling Technique) for Handling Imbalanced Dataset Problem, 2018 4th International Conference on Science and Technology (ICST), 2018, pp. 1-4, doi: 10.1109/ICSTC.2018.8528591
- [Tesse (2011)] Tesse R, Pandey RC, Kabesch M. (2011), Genetic variations in toll-like receptor pathway genes influence asthma and atopy, Allergy 2011,66, 307-318
- [Tibshirani (1996)] Tibshirani, R. (1996), Regression shrinkage and selection via the lasso, Journal of the Royal Statistical Society, 58(1), 267–288
- [Westman et al. (2012)] Westman M, Stjärne P., Asarnoj A., Kull I., van Hage M., Wickman M., Toskala E. (2012), Natural course and comorbidities of allergic and nonallergic rhinitis in children, Journal of Allergy and Clinical Immunology 129(2), 403-408 doi.org/10.1016/j.jaci.2011.09.036
- [Zalewska et al. (2010)] Zalewska M., Niemiro W., Samoliński B, (2010), MCMC imputation in autologistic model, Monte Carlo Methods and Applications, De Gruyter, vol. 16(3-4), 421-438
- [Zavlis et al. (2021)] Zavlis, O., Butter, S., Bennett, K., Hartman, T.K., Hyland, P., Mason, L., McBride, O., Murphy, J., Gibson-Miller, J., Levita, L., Martinez, A.P., Shevlin, M., Stocks, T.V.A., Vallieres, F., Bentall, R.P. (2021), How Does the COVID-19 Pandemic Impact on Population Mental Health? A Network Analysis of COVID Influences on Depression, Anxiety and Traumatic Stress in the UK Population, *Psychological Medicine*, 16 Mar 2021, 1-9, doi=10.1017/S0033291721000635
- [Zhang et al. (2019)] Zhang H., Li Z., Shahriar H., Tao L., Bhattacharya P., Qian Y, (2019), Improving Prediction Accuracy for Logistic Regression on Imbalanced Datasets, *IEEE 43rd* Annual Computer Software and Applications Conference (COMPSAC), 2019, 918-919 doi: 10.1109/COMPSAC.2019.00140
- [Zhang et al. (2021)] Zhang L., Geisler T., Ray H., Xie Y (2021), Improving logistic regression on the imbalanced data by a novel penalized log-likelihood function, *Journal of Applied Statistics*, 1-21