

July 18, 2023

Transverse single spin asymmetry for very forward neutron production in polarized $p + p$ collisions at $\sqrt{s} = 510$ GeV

MINHO KIM

RIKEN

In the high-energy $p + p$ collisions, the transverse single spin asymmetry for very forward neutron production has been interpreted by an interference between π (spin flip) and a_1 (spin non-flip) exchange with a non-zero phase shift. The π and a_1 exchange model predicted the neutron asymmetry would increase in magnitude with transverse momentum (p_T) in $p_T < 0.4$ GeV/ c . In June 2017, the RHICf experiment installed an electromagnetic calorimeter at the zero-degree area of the STAR experiment at the Relativistic Heavy Ion Collider and measured the neutron asymmetry in a wide p_T range of $0 < p_T < 1$ GeV/ c from polarized $p + p$ collisions at $\sqrt{s} = 510$ GeV. The RHICf data allows us to investigate the kinematic dependence of the neutron asymmetry in detail, which not only can test the π and a_1 exchange model in the higher p_T range but also can study the \sqrt{s} dependence by comparing with the previous measurements. We present the preliminary result and analysis status of the neutron asymmetry measured by the RHICf experiment. In order to understand the RHICf result, a theoretical calculation other than Reggeon exchange will also be discussed.

PRESENTED AT

DIS2023: XXX International Workshop on Deep-Inelastic
Scattering and Related Subjects,
Michigan State University, USA, 27-31 March 2023



1 Introduction

In polarized $p + p$ collisions, transverse single spin asymmetry (A_N) is defined by a left-right cross section asymmetry,

$$A_N = \frac{\sigma_L - \sigma_R}{\sigma_L + \sigma_R},$$

where the $\sigma_{L(R)}$ is the cross section of a specific particle which is produced in the left (right) side with respect to the beam polarization. Since a large A_N for very forward ($\eta > 6$) neutron production has been discovered by a polarimeter development experiment [1], it has been measured by the PHENIX experiment at three different collision energies, 62, 200, and 500 GeV [2]. One pion exchange (OPE) model which had successfully described the cross section of the neutron production introduced an interference between the spin flip π exchange and spin non-flip a_1 exchange with a relative phase shift to reproduce the experimental data [3]. The π and a_1 exchange model reproduced the data well predicting that the A_N increases in magnitude with transverse momentum (p_T) with little longitudinal momentum fraction (x_F) dependence. Recently, the neutron A_N measured by the PHENIX experiment at $\sqrt{s} = 200$ GeV was unfolded to precisely study the kinematic dependence of the neutron A_N and showed a consistency with the model prediction [4]. However, kinematic range of the experimental measurement and model calculation was narrow compared to that of very forward neutron production.

The RHICf experiment [5] measured the A_N for very forward neutron production in a wide p_T coverage of $0.0 < p_T < 1.0$ GeV/ c from polarized $p + p$ collisions at $\sqrt{s} = 510$ GeV. The RHICf data allows us to test the validity of the π and a_1 exchange model in the higher p_T region. Since \sqrt{s} of the RHICf experiment is different from those of previous measurements, one can also study the \sqrt{s} dependence of the neutron A_N .

2 RHICf experiment

In June 2017, The RHICf experiment measured the A_N for very forward neutron production in polarized $p + p$ collisions at $\sqrt{s} = 510$ GeV at the Relativistic Heavy Ion Collider (RHIC). An electromagnetic calorimeter (RHICf detector) [6] was installed in front of the STAR Zero-Degree Calorimeter (ZDC) [7], which was located 18 m away from the beam interaction point. We also installed a thin scintillator counter (FC) in front of the RHICf detector to analyze the charged hadron background. The RHICf detector consists of small and large sampling towers with 20 and 40 mm dimensions, respectively. Both towers are composed of 17 tungsten absorbers with 1.6 interaction length in total, 16 GSO plates for energy measurement, and 4 layers of GSO bar hodoscope for position measurement. Position and energy resolutions for 250 GeV

neutron are about 1 mm and 39% when the incident position of the neutron is the center of a tower. A shower trigger which was operated when the energy deposits of three successive GSO plate layers are larger than 45 MeV was used for the neutron measurement. For more detailed detector configuration and performance, see Ref. [8], [9] and [10].

We requested large β^* value of 8 m at operation to make the angular beam divergence small. Corresponding luminosity at $\sqrt{s} = 510$ GeV was $\sim 10^{31}$ cm $^{-2}$ s $^{-1}$ which was smaller than that in usual RHIC operation. We also requested a radial polarization which was 90°-rotated from that of usual RHIC operation to reach the maximal p_T range by moving the detector vertically. We took the data during about 28 hours with three detector positions where the beam headed the center of large tower, center of small tower, and 24 mm below the center of small tower.

3 Data analysis

Basically, only the shower triggered events were analyzed to avoid any bias from the different trigger efficiencies. The shower trigger is sensitive not only to the neutron events but also to the photon events. In order to separate the neutron events from the photon background, a cut condition of $L_{90\%} - 0.15L_{20\%} > 21$ was applied. $L_{x\%}$ is defined by longitudinal depth of detector where the accumulated energy deposit reaches $x\%$ of the total. The above condition was optimized taking into account the neutron purity and efficiency. Events where the hadronic shower developed in the deeper GSO plates were rejected by applying for $L_{90\%} < 37$ to improve the energy resolution of neutron. Energy resolution of 250 GeV neutron was improved from 39% to 30% accordingly.

Since the RHICf detector has insufficient interaction length for the neutron energy measurement, kinematic values of neutron were unfolded by using Bayesian unfolding [13]. For prior, a Monte Carlo (MC) sample where neutrons from 0 to 255 GeV were uniformly generated on the detector was used to avoid any bias from the particular particle productions. We repeated the iteration until the χ^2 change between two outputs of consecutive iterations became smaller than 1. We generated finite asymmetry by assigning up and down spin patterns to each event and also confirmed that the unfolded distributions reproduced the input A_N well within the statistical uncertainty.

Background A_{NS} from the photon and charged hadron were subtracted using the following equation after unfolding,

$$A_N^{\text{neu}} = \left(\frac{N_{\text{trig}}}{N_{\text{neu}}}\right)A_N^{\text{trig}} - \left(\frac{N_{\text{pho}}}{N_{\text{neu}}}\right)A_N^{\text{pho}} - \left(\frac{N_{\text{had}}}{N_{\text{neu}}}\right)A_N^{\text{had}},$$

where the notations “neu”, “pho”, “had” and “trig” mean the neutron, photon, charged hadron and triggered events, respectively, and N is the number of mea-

sured events. Ratios between different types of events were estimated referring to the QGSJET II-04 sample. A_N^{pho} was calculated using the photon-enhanced sample of data. In the preliminary result, the FC and A_N^{had} were not analyzed, thereby A_N variation between when the value of A_N^{had} was -1 and $+1$ was assigned as one of the systematic uncertainties. For additional sources of the systematic uncertainty, uncertainties of beam center calculation, polarization and unfolding were considered.

4 Results

Fig. 1 shows the preliminary result of the A_N for very forward neutron production. Fig. 1 (a) shows the neutron A_N as a function of p_T in different x_F ranges. In the two higher x_F ranges, the A_N increases in magnitude with p_T without x_F dependence as the model predicted. In $p_T < 0.25$ GeV/ c , they are consistent with those of PHENIX [4]. No \sqrt{s} dependence was observed in the neutron A_N . A_N s in the lowest x_F range show different behavior compared to those in the higher x_F ranges, which indicates a x_F dependence that hasn't been expected by the π and a_1 exchange model. Fig. 1 (b) shows the neutron A_N as a function of x_F in different p_T ranges. In $p_T < 0.25$ GeV/ c , A_N s are flat without x_F dependence. However, in $p_T > 0.25$ GeV/ c , clear x_F dependence is observed.

In order to understand the x_F dependence of the neutron A_N , spin effect by an absorptive correction in the single π exchange was studied. The absorptive correction is an elastic interaction from the initial state proton or final state multi particles. Since phase shift by the absorptive correction starts to deviate from $p_T \sim 0.3$ GeV/ c , it has large A_N in the higher p_T region. Fig. 2 shows a comparison between the RHICf data and prediction by the absorptive correction in $0.46 < x_F < 0.64$. The absorptive correction doesn't reproduce the RHICf data well. However, it shows a possible origin of the x_F dependence. Since the finite neutron A_N also comes from the π and a_1 exchange, we expect that more comprehensive theoretical calculation could reproduce the RHICf data.

To finalize the result, we analyzed the FC and the charged hadron background was suppressed by by rejecting the events where ADC of the FC has finite value. ADC distribution of the FC was reproduced in the simulation and effect of the charged hadron contamination after applying for the above condition was estimated. Since the background study has been recently completed, the final result of the neutron A_N will be released soon.

5 Summary

In June 2017, the RHICf experiment measured the A_N for very forward neutron production in polarized $p + p$ collisions at $\sqrt{s} = 510$ GeV and presented the preliminary




Figure 1: A_N for very forward neutron production (a) as a function of p_T in different x_F ranges and (b) as a function of x_F in different p_T ranges.

result. In $0.46 < x_F$, the A_N increases in magnitude with p_T as the model predicted. No \sqrt{s} was observed when the RHICf data points were compared with those of PHENIX result. In $p_T < 0.25$ GeV/c, A_N s are flat without x_F dependence. However, in $p_T > 0.25$ GeV/c, a clear x_F dependence was observed. In the preliminary result, effect of the charged hadron events was not analyzed in detail. However, since it has been complemented, we will finalize the result soon.




Figure 2: Comparison between the RHICf data and prediction by the absorptive correction in the single π exchange in $0.46 < x_F < 0.64$.

References

- [1] Y. Fukao *et al.*, Phys. Lett. B **650**, 325 (2007).
- [2] K. Tanida (PHENIX Collaboration), J. Phys. Conf. Ser. **295**, 012097 (2011).
- [3] B. Z. Kopeliovich, I. K. Potashnikova, I. Schmidt, and J. Soffer, Phys. Rev. D **84**, 114012 (2011).
- [4] U. A. Acharya *et al.* (PHENIX Collaboration), Phys. Rev. D **103**, 052009 (2021).
- [5] RHICf Collaboration: LOI, arXiv: 1409.4860v1.
- [6] RHICf Collaboration, JINST **16** P10027 (2021).
- [7] C. Adler *et al.*, Nucl. Instrum. Meth. A **470**, 488 (2001).
- [8] T. Suzuki *et al.*, JINST **8**, T01007 (2013).
- [9] Y. Makino *et al.*, JINST **12**, P03023 (2017).
- [10] K. Kawade *et al.*, JINST **9**, P03016 (2014).
- [11] M. H. Kim *et al.* (RHICf Collaboration), Phys. Rev. Lett. **124**, 252501 (2020).
- [12] S. Ostapcheno, Nucl. Phys. B, Proc. Suupl. **151**, 143 (2006).
- [13] G. D’Agostini, Nucl. Instrum. Meth. A **362**, 487 (1995).