Combining Finite Combination Properties: Finite Models and Busy Beavers

Guilherme V. Toledo¹, Yoni Zohar¹, and Clark Barrett²

¹Bar-Ilan University ²Stanford University

Abstract. This work is a part of an ongoing effort to understand the relationships between properties used in theory combination. We here focus on including two properties that are related to shiny theories: the finite model property and stable finiteness. For any combination of properties, we consider the question of whether there exists a theory that exhibits it. When there is, we provide an example with the simplest possible signature. One particular class of interest includes theories with the finite model property that are not finitely witnessable. To construct such theories, we utilize the Busy Beaver function.¹

1 Introduction

The story of this paper begins with [9], where it was shown that the theory of algebraic datatypes, useful for modeling data structures like lists and trees, can be combined with any other theory, using the polite combination method [8]. This combination method offers a way to combine decisions procedures of two theories into a decision procedure for the combined theory, with different assumptions than those of the earlier Nelson-Oppen approach [6]. In particular, it was proven that the theory admits a technical property concerning cardinalities of models, called strong politeness [3]. It was noted in [9] that proving strong politeness for this theory seemed much harder than proving *politeness*, a similar but simpler property. Therefore, the proof was split into three steps: (i) a class of theories was identified in which politeness and strong politeness coincide; (ii) the theory of algebraic datatypes was shown to be in this class; and (iii) this theory was proven to be polite. This proof technique raised the following question: does politeness imply strong politeness? An affirmative answer to this question would simplify strong politeness proofs that follow such steps, as only the last step would be needed. Unfortunately, the answer to this question was shown in [10] to be negative, in its most general form. However, an affirmative answer was given for theories over one-sorted empty signatures, where politeness and strong politeness do coincide.

Seeing that relationships between model-theoretic properties of theories (like politeness and strong politeness) are non-trivial, and can have a big impact on proofs in the field of theory combination, we have recently initiated a more general research plan: to systematically determine the relationships between model-theoretic properties that relate to theory combination. An analysis of such properties can, for example, simplify proofs, in cases where a property follows from a combination of other properties.

¹ Funded in part by NSF-BSF grant numbers 2110397 (NSF) and 2020704 (BSF) and ISF grant number 619/21.

In the first stage of this plan [12], we studied the relationships between all properties that relate to either polite or Nelson-Oppen combination, namely: stable infiniteness, smoothness, finite witnessability, strong finite witnessability, and convexity. The first two properties relate to the ability to enlarge cardinalities of models, while the next two require a computable *witness* function that restricts the models of a formula based on its variables. The last property relies on the ability to deduce an equality from a disjunction of equalities. The result of [12] was a comprehensive table: nearly every combination of these properties (e.g., theories that are smooth and stably infinite but do not admit the other properties) was either proved to be infeasible, or an example for it was given.

In this paper we continue with this plan by adding two properties: the finite model property and stable finiteness, both related to shiny theories [11]. The former requires finite models for satisfiable formulas, and the latter enforces bounds on them.

Of course, the theories from [12] can be reused. For these, one only needs to determine if they admit the finite model property and/or stable finiteness. The results and examples from [12] are, however, not enough. Given that the number of considered combinations is doubled with the addition of each property, new theories need to be introduced in order to exemplify the new possibilities, and new impossible combinations can be found. Hence, in this paper we provide several impossibility results for the aforementioned properties, as well as examples of theories for possible combinations. The overall result is a new table which extends that of [12] with two new columns corresponding to the finite model property and stable finiteness.²

The most interesting combinations that we study are theories that admit the finite model property but not finite witnessability. While both properties deal with finite models, the latter has a computable element to it, namely the witness function. In separating these properties, we found it useful to define theories that are based on the *Busy Beaver* function, a well known function from computability theory, that is not only non-computable, but also grows eventually faster than any computable function.

Outline: Section 2 reviews many-sorted logics and theory combination properties. Section 3 identifies combinations that are contradictory; Section 4 constructs the extended table of combinations, and describes the newly introduced theories. Section 5 gives final remarks and future directions this work can take.

2 Preliminary Notions

2.1 Many-sorted Logic

A many-sorted signature Σ is a triple $(S_{\Sigma}, \mathcal{F}_{\Sigma}, \mathcal{P}_{\Sigma})$ where: S_{Σ} is a countable set of sorts; \mathcal{F}_{Σ} is a countable set of function symbols; and \mathcal{P}_{Σ} is a countable set of predicate symbols containing, for each $\sigma \in S_{\Sigma}$, an equality $=_{\sigma}$. When σ is clear from the context, we write =. Every function symbol has an *arity* of the form $\sigma_1 \times \cdots \times \sigma_n \rightarrow \sigma$, and every predicate symbol one of the form $\sigma_1 \times \cdots \times \sigma_n$, where $\sigma_1, \ldots, \sigma_n, \sigma \in S_{\Sigma}$; equalities $=_{\sigma}$ have arity $\sigma \times \sigma$.

² While we use several results from [12], we do not assume here any familiarity with that paper. All required results are mentioned here explicitly.

A signature that has no functions and only the equalities as predicates is called *empty*. Many-sorted signatures Σ where S_{Σ} has only one element are called *one-sorted*.

For any sort in S_{Σ} we assume a countably infinite set of variables, and distinct sorts have disjoint sets of variables; we then define first-order terms, formulas, and literals in the usual way. The set of free variables of sort σ in a formula φ is denoted by $vars_{\sigma}(\varphi)$, while $vars(\varphi)$ will denote $\bigcup_{\sigma \in S_{\Sigma}} vars_{\sigma}(\varphi)$.

 Σ -Structures \mathbb{A} are defined as usual, by interpreting sorts (denoted by $\sigma^{\mathbb{A}}$), functions $(f^{\mathbb{A}})$ and predicate symbols $(P^{\mathbb{A}})$, with the restrictions that equality symbols are interpreted as identities. A Σ -interpretation \mathcal{A} is an extension of a Σ -structure \mathbb{A} with interpretations to variables. If \mathbb{A} is the underlying Σ -structure of a Σ -interpretation \mathcal{A} , we say that \mathcal{A} is an interpretation on \mathbb{A} . For simplicity, and because the use of structures is sparse in this paper, we will usually denote both structures and interpretations by using the same font, \mathcal{A} , \mathcal{B} and so on. $\alpha^{\mathcal{A}}$ is the value taken by a Σ -term α in a Σ -interpretation \mathcal{A} , and if Γ is a set of terms, we simply write $\Gamma^{\mathcal{A}}$ for $\{\alpha^{\mathcal{A}} : \alpha \in \Gamma\}$.

We write $\mathcal{A} \models \varphi$ if the Σ -interpretation \mathcal{A} satisfies the Σ -formula φ ; φ is then said to be *satisfiable* if it is satisfied by some interpretation \mathcal{A} . The formulas found in Figure 1 will be useful in the sequel. A Σ -interpretation \mathcal{A} : satisfies $\psi_{\geq n}^{\sigma}$ iff $|\sigma^{\mathcal{A}}| \geq n$; satisfies $\psi_{\leq n}^{\sigma}$ iff $|\sigma^{\mathcal{A}}| \leq n$; and satisfies $\psi_{=n}^{\sigma}$ iff $|\sigma^{\mathcal{A}}| = n$. For simplicity, when dealing with one-sorted signatures, we may drop the sort σ from the cardinality formulas.

$$\psi_{\geq n}^{\sigma} = \exists \ \overrightarrow{x}. \ \bigwedge_{1 \leq i < j \leq n} \neg (x_i = x_j) \quad \psi_{\leq n}^{\sigma} = \exists \ \overrightarrow{x}. \ \forall \ y. \ \bigvee_{i=1}^n y = x_i \quad \psi_{=n}^{\sigma} = \psi_{\geq n}^{\sigma} \land \psi_{\leq n}^{\sigma}$$

Fig. 1: Cardinality Formulas. \overrightarrow{x} stands for x_1, \ldots, x_n , all variables of sort σ .

A Σ -theory \mathcal{T} is a class of all Σ -interpretations (called \mathcal{T} -interpretations) that satisfy some set $Ax(\mathcal{T})$ of closed formulas called the *axiomatization* of \mathcal{T} ; the structures underlying these interpretations will be called the *models* of \mathcal{T} .

A formula is \mathcal{T} -satisfiable if it is satisfied by some \mathcal{T} -interpretation and, analogously, a set of formulas is \mathcal{T} -satisfiable if there is a \mathcal{T} -interpretation that satisfies all of them simultaneously. Two formulas are \mathcal{T} -equivalent when a \mathcal{T} -interpretation satisfies the first iff it satisfies the second. We write $\models_{\mathcal{T}} \varphi$, and say that φ is \mathcal{T} -valid if $\mathcal{A} \models \varphi$ for all \mathcal{T} -interpretations \mathcal{A} .

2.2 Theory Combination Properties

Let Σ be a signature, \mathcal{T} a Σ -theory and $\mathcal{S} \subseteq \mathcal{S}_{\Sigma}$. We define several properties \mathcal{T} may have with respect to S.

Convexity, Stable Infiniteness, and Smoothness \mathcal{T} is *convex* with respect to S if for any conjunction of Σ -literals ϕ and any finite set of variables $\{u_1, v_1, \ldots, u_n, v_n\}$ of sorts in S with $\models_{\mathcal{T}} \phi \to \bigvee_{i=1}^n u_i = v_i$, one has $\models_{\mathcal{T}} \phi \to u_i = v_i$ for some i. \mathcal{T} is stably infinite with respect to S if for every \mathcal{T} -satisfiable quantifier-free Σ -formula there is a \mathcal{T} -interpretation \mathcal{A} satisfying it such that $|\sigma^{\mathcal{A}}|$ is infinite for each $\sigma \in S$. \mathcal{T} is smooth with respect to S if for every quantifier-free formula, \mathcal{T} -interpretation \mathcal{A} that satisfies it, and function κ from S to the class of cardinals such that $\kappa(\sigma) \geq |\sigma^{\mathcal{A}}|$ for each $\sigma \in S$, there is a \mathcal{T} -interpretation \mathcal{B} that satisfies it with $|\sigma^{\mathcal{B}}| = \kappa(\sigma)$ for each $\sigma \in S$.

(Strong) Finite witnessability For finite sets of variables V_{σ} of sort σ for each $\sigma \in S$, and equivalence relations E_{σ} on V_{σ} , the arrangement on $V = \bigcup_{\sigma \in S} V_{\sigma}$ induced by $E = \bigcup_{\sigma \in S} E_{\sigma}$, denoted by δ_V or δ_V^E , is the formula $\delta_V = \bigwedge_{\sigma \in S} [\bigwedge_{x \in \sigma y} (x = y) \land$ $\bigwedge_{x \in \overline{E_{\sigma}y}} \neg (x = y)]$, where $\overline{E_{\sigma}}$ denotes the complement of the equivalence relation E_{σ} .

 \mathcal{T} is *finitely witnessable* with respect to S when there exists a computable function wit, called a witness, from the quantifier-free Σ -formulas to themselves that satisfies, for every ϕ : (i) ϕ and $\exists \vec{w} . wit(\phi)$ are \mathcal{T} -equivalent, for $\vec{w} = vars(wit(\phi)) \setminus vars(\phi)$; and (ii) if $wit(\phi)$ is \mathcal{T} -satisfiable, there exists a \mathcal{T} -interpretation \mathcal{A} satisfying $wit(\phi)$ such that $\sigma^{\mathcal{A}} = vars_{\sigma}(wit(\phi))^{\mathcal{A}}$ for each $\sigma \in S$.

Strong finite witnessability is defined similarly to finite witnessability, replacing (ii) by: (ii)' given a finite set of variables V and an arrangement δ_V on V, if $wit(\phi) \wedge \delta_V$ is \mathcal{T} -satisfiable, there exists a \mathcal{T} -interpretation \mathcal{A} that satisfies $wit(\phi) \wedge \delta_V$ with $\sigma^{\mathcal{A}} = vars_{\sigma}(wit(\phi) \wedge \delta_V)^{\mathcal{A}}$ for all $\sigma \in S$. If \mathcal{T} is smooth and (strongly) finitely witnessable with respect to \mathcal{S} , then it is (strongly) polite with respect to \mathcal{S} .

Finite Model Property and Stable Finiteness \mathcal{T} has the *finite model property* with respect to S if for every quantifier-free \mathcal{T} -satisfiable Σ -formula, there exists a \mathcal{T} interpretation \mathcal{A} that satisfies it with $|\sigma^{\mathcal{A}}|$ finite for each $\sigma \in S$. \mathcal{T} is *stably finite* with respect to S if, for every quantifier-free Σ -formula and \mathcal{T} -interpretation \mathcal{A} that satisfies it, there exists a \mathcal{T} -interpretation \mathcal{B} that satisfies it with: $|\sigma^{\mathcal{B}}|$ finite for each $\sigma \in S$; and $|\sigma^{\mathcal{B}}| \leq |\sigma^{\mathcal{A}}|$ for each $\sigma \in S$. Clearly, stable finiteness implies the finite model property:

Theorem 1. If \mathcal{T} is stably finite w.r.t. S, then it has the finite model property w.r.t. S.

We shall write SI for stably infinite; SM for smooth; FW (SW) for (strong) finitely witnessable; CV for convex; FM for the finite model property; and SF for stably finite.

3 Relationships between model-theoretic properties

In this section we study the connections between finiteness properties related to theory combination: the finite model property, stable finiteness, finite witnessability, and strong finite witnessability. We show how these properties are related to one another. In Section 3.1, we provide general results that hold for all signatures. Then, in Section 3.2, we focus on empty signatures, in which we are able to find more connections.

3.1 General Signatures

Finite witnessability, as well as its strong variant, were introduced in the context of polite theory combination. In contrast, the study of shiny theories utilizes the notions of the finite model property, as well as stable finiteness. It was shown in [2] that for theories with a decidable quantifier-free satisfiability problem, shiny theories and strongly polite theories are one and the same. This already showed some connections between the aforementioned finiteness properties. However, that analysis also relied on smoothness, the decidability of the quantifier-free satisfiability problem of the studied theories, as well as the computability of the *mincard* function, the function that computes the minimal sizes of domains in models of a given formula in these theories.

Here we focus purely on the finiteness properties, and show that even without any other assumptions, they are closely related. Considering finite witnessability and the finite model property, notice that any witness ensures that some formulas always have finite models. Using the equivalence of the existential closure of such formulas to the formulas that are given to the witness, one gets the following result, according to which finite witnessability implies the finite model property.

Theorem 2. Any Σ -theory \mathcal{T} finitely witnessable with respect to $S \subseteq S_{\Sigma}$ also has the finite model property with respect to S.

Strong finite witnessability is a stronger property than finite witnessability, obtained by requiring finite models in the presence of arrangements. This requirement allows one to conclude stable finiteness for it, as the finer control on cardinalities that is required for stable finiteness can be achieved with the aid of arrangements. The following result is proved in Lemma 3.6 of [2], although under the assumption that the theory is smooth, something that is not actually used in their proof.

Theorem 3. Any Σ -theory \mathcal{T} strongly finitely witnessable with respect to $S \subseteq S_{\Sigma}$ is also stably finite with respect to S.

Clearly, stable finiteness implies the finite model property (Theorem 1). The converse does not generally hold, as we will see in Section 4. However, when these properties are considered with respect to a single sort, they actually coincide:

Theorem 4. If a Σ -theory \mathcal{T} has the finite model property with respect to a set of sorts S with |S| = 1, then \mathcal{T} is also stably finite with respect to S.

Theorems 2 and 3 are visualized in the Venn diagram of Figure 2, where, for example, theories that are strongly finitely witnessable are clearly inside the intersection of finitely witnessable theories and stably finite theories.

When only one sort is considered, the picture is much simpler, and is described in Figure 3. There, the finite model property and stable finiteness populate the same region, as ensured by Theorem 4. Notice that the results depicted in Figure 3 hold for one-sorted and many-sorted signatures. The key thing is that the properties are all w.r.t. one of the sorts.

3.2 Empty Signatures

Figures 2 and 3 show a complete picture of the relationships between the properties studied in this section, for arbitrary signatures. However, when this generality is relaxed,




Fig. 2: Finiteness properties: general case. Fig. 3: Finiteness properties w.r.t. one sort.

several other connections appear. For this section, we require that the signatures are empty, and that they have a finite set of sorts. We further require that the properties in question hold for the entire set of sorts, not for any subset of it.

Table 1 defines the 5 signatures that will be used in the examples found in Section 4, and that will also appear in some of the results shown below: the empty signatures Σ_1 , Σ_2 and Σ_3 , with sets of sorts $\{\sigma\}$, $\{\sigma, \sigma_2\}$ and $\{\sigma, \sigma_2, \sigma_3\}$, respectively; and the signatures Σ_s and Σ_s^2 with one function s of arity $\sigma \to \sigma$, and sets of sorts $\{\sigma\}$ and $\{\sigma, \sigma_2\}$, respectively. Notice these are the simplest possible signatures when we order those by establishing: first, that the signature with fewer sorts is simpler; and second, that if two signatures have the same number of sorts, the one with fewer function symbols is simpler. We are free not to consider predicates, as they are at least as expressive as functions themselves; furthermore, we do not consider the problem of defining which of two signatures with the same numbers of sorts and function symbols is simpler, choosing rather to add only functions from a sort to itself.

Signature	Sorts	Function Symbols
Σ_1	$\{\sigma\}$	Ø
Σ_2	$\{\sigma,\sigma_2\}$	Ø
Σ_3	$\{\sigma, \sigma_2, \sigma_3\}$	Ø
Σ_s	$\{\sigma\}$	$\{s:\sigma\to\sigma\}$
Σ_s^2	$\{\sigma,\sigma_2\}$	$\{s:\sigma\to\sigma\}$

Table 1: Signatures that will be used throughout the paper.

First, in such a setting, we have that the finite model property implies finite witnessability, in the presence of smoothness.

Theorem 5. If Σ is an empty signature with a finite set of sorts S_{Σ} , and the Σ -theory \mathcal{T} has the finite model property and is smooth with respect to S_{Σ} , then \mathcal{T} is also finitely witnessable with respect to S_{Σ} .




Fig. 4: Interplay between SM, FW (SW) and FM (SF) w.r.t. S_{Σ} in an empty signature.

Next, we show that stable finiteness and smoothness together, imply strong finite witnessability.

Theorem 6. If Σ is an empty signature with a finite set of sorts S_{Σ} , and the Σ -theory T is stably finite and smooth with respect to S_{Σ} , then T is also strongly finitely witnessable with respect to S_{Σ} .

While Theorem 2 and Theorem 3 establish certain unconditional relations between finite witnessability and the finite model property, and strong finite witnessability and stable finiteness, the converses shown to hold in Theorem 5 and Theorem 6 demand smoothness and that the properties hold with respect to the entire set of sorts. In that case, the situation can be represented by the diagram found in Figure 4, showing clearly that a smooth theory that also has the finite model property (respectively, is stably finite), cannot not be finitely witnessable (strongly finitely witnessable).

Lastly, regarding the empty signatures Σ_1 , Σ_2 and Σ_3 , the following theorem shows that Σ_3 is sometimes necessary.

Theorem 7. There are no Σ_1 or Σ_2 -theories \mathcal{T} that are, simultaneously, neither stably infinite nor stably finite, but are convex and have the finite model property, with respect to the entire set of their sorts.

Hence, to exhibit such theories, one has to consider three-sorted theories.

4 A taxonomy of examples

In [12], we have created a table, in which for every possible combinations of properties from $\{$ **SI**, **SM**, **FW**, **SW**, **CV** $\}$ we either gave an example of a theory in this combination, or proved a theorem that shows there is no such example, with the exception of theories that are stably infinite and strongly finitely witnessable but not smooth. Such theories, referred to in [12] as *Unicorn Theories* (due to our conjecture that they do not exist) were left for future work, and are still left for future work, as the focus of the current paper is the integration of finiteness properties, namely **FM** and **SF** to the table.

And indeed, the goal of this section is to add two columns to the table from [12]: one for the finite model property and one for stable finiteness. The extended table is Table 2. We do not assume familiarity with [12], and describe the entire resulting table (though focusing on the new results).

			Em	pty	Non-empty						
SI	SM	FW	SW	CV	FM	SF	One-sorted	Many-sorted	One-sorted	Many-sorted	$N^{\underline{o}}$
			Ŧ	T	Т	T	$\mathcal{T}_{\geq n}$	$(\mathcal{T}_{\geq n})^2$	$(\mathcal{T}_{\geq n})_s$	$((\mathcal{T}_{\geq n})^2)_s$	1
			Т	F	Т	T		2]	$(\mathcal{T}_{\geq n})_{\vee}$	$((\mathcal{T}_{\geq n})^2)_{\vee}$	2
		Т		т	Т	Т	Theorem 6		\mathcal{T}_f	$(\mathcal{T}_f)_s$	3
		1	T	Т		F	Theorem 4	$\mathcal{T}_{2,3}$	Theorem 4	$(\mathcal{T}_{2,3})_s$	4
			F	L.	Т	T	F 1	01	\mathcal{T}_{f}^{s}	$(\mathcal{T}_{f}^{s})^{2}$	5
	T			F		F	[]	2]	Theorem 4	$(\mathcal{T}_{2,3})_{ee}$	6
	1				Т	T	Theo	nom 5	$\mathcal{T}^s_{\varsigma}$	$\mathcal{T}_{\varsigma}^{=}$	7
				T		F	Theorem 5		Theorem 4	$\mathcal{T}_{\varsigma}^2$	8
		F	F		F	F	\mathcal{T}_{∞}	$(\mathcal{T}_{\infty})^2$	$(\mathcal{T}_\infty)_s$	$((\mathcal{T}_{\infty})^2)_s$	9
		1	1		Т	T	[12]		$\mathcal{T}_{\varsigma}^{\vee}$	$(\mathcal{T}^{\vee}_{\varsigma})^2$	10
				F	1	F			Theorem 4	$\mathcal{T}_{\varsigma\vee}^{=}$	11
T					F	F			$(\mathcal{T}_\infty)_ee$	$((\mathcal{T}_{\infty})^2)_{\vee}$	12
			Т	Т	Т	T		Unicorn			13
				F	Т	T			COIII		14
		Т		Т	Т	T	$\mathcal{T}^\infty_{even}$	$(\mathcal{T}^\infty_{even})^2$	$(\mathcal{T}^\infty_{even})_s$	$((\mathcal{T}^{\infty}_{even})^2)_s$	15
		1	F	1		F	Theorem 4	\mathcal{T}^∞	Theorem 4	$(\mathcal{T}^{\infty})_s$	16
				F	Т	T	[12]		$(\mathcal{T}^\infty_{even})_ee$	$((\mathcal{T}^{\infty}_{even})^2)_{\vee}$	17
	F					F			Theorem 4	$(\mathcal{T}^\infty)_{ee}$	18
			F	Т	Т	T	\mathcal{T}_{ς}	$(\mathcal{T}_{\varsigma})^2$	$(\mathcal{T}_{\varsigma})_s$	$((\mathcal{T}_{\varsigma})^2)_s$	19
						F	Theorem 4	$\mathcal{T}^\infty_\varsigma$	Theorem 4	$(\mathcal{T}^\infty_\varsigma)_s$	20
		F		F	F	F	$\mathcal{T}_{n,\infty}$	$(\mathcal{T}_{n,\infty})^2$	$(\mathcal{T}_{n,\infty})_s$	$((\mathcal{T}_{n,\infty})^2)_s$	21
					Т	T	[12]		$(\mathcal{T}_{\varsigma})_{ee}$	$((\mathcal{T}_{\varsigma})^2)_{ee}$	22
						F			Theorem 4	$(\mathcal{T}^{\infty}_{\varsigma})_{\lor}$	23
					F	F			$(\mathcal{T}_{n,\infty})_ee$	$((\mathcal{T}_{n,\infty})^2)_{\vee}$	24
		Т	Т	T	Т	Т	$\mathcal{T}_{\leq 1}$	$\frac{(\mathcal{T}_{\leq 1})^2}{(\mathcal{T}_{\leq n})^2}$ $\frac{\mathcal{T}_1^{odd}}{\mathcal{T}_1^{odd}}$	$(\mathcal{T}_{\leq 1})_s$	$((\mathcal{T}_{\leq 1})^2)_s$	25
				F	Т	Т	$\mathcal{T}_{\leq n}$	$(\mathcal{T}_{\leq n})^2$	$(\mathcal{T}_{\leq n})_s$	$((\mathcal{T}_{\leq n})^2)_s$	26
			F	T	T	Т	[12]	\mathcal{T}_1^{odd}	$\mathcal{T}_{odd}^{ eq}$	$(\mathcal{T}_1^{odd})_s$	27
						F	Theorem 4	$\mathcal{T}^3_{2,3}$	Theorem 4	$\mathcal{T}^{\infty}_{\neq}$	28
				F	Т	Т	$\mathcal{T}_{m,n}$	$(\mathcal{T}_{m,n})^2$	$(\mathcal{T}_{m,n})_s$	$((\mathcal{T}_{m,n})^2)_s$	29
F	F					F	Theorem 4	$\mathcal{T}_{m,n}^\infty$	Theorem 4	$(\mathcal{T}_{m,n}^{\infty})_s$	30
		F	F	Т	T	T		$\mathcal{T}_1^{\varsigma}$	$\mathcal{T}_{\varsigma,1}^{ eq}$	$(\mathcal{T}_1^{\varsigma})^2$	31
						F		$\mathcal{T}^{\infty,3}_{\varsigma}$	Theorem 4	$\mathcal{T}^{\infty}_{\varsigma \neq}$	32
					F	F	[12]	\mathcal{T}_1^{∞}	$\mathcal{T}_{1,\infty}^{ eq}$	$(\mathcal{T}_1^\infty)_s$	33
				г	T	T		$\mathcal{T}_n^{\varsigma}$	$\mathcal{T}_{\varsigma}^{\neq}$	$(\mathcal{T}_n^{\varsigma})_s$	34
				F		F		$\mathcal{T}_{m,n}^{\varsigma}$	Theorem 4 π^{\pm}	$(\mathcal{T}_{m,n}^{\varsigma})_s$	35
					F	F		\mathcal{T}_2^∞	$\mathcal{T}_{2,\infty}^{ eq}$	$(\mathcal{T}_2^\infty)_s$	36

Table 2: Summary of all possible combinations of theory properties. Red cells represent impossible combinations. In lines 26 and 34, n > 1; in lines 29, 30 and 35, m > 1, n > 1 and |m - n| > 1.

This section is structured as follows: In Section 4.1 we describe the structure of Table 2. In Sections 4.2 to 4.4 we provide details about the axiomatizations of theories that populate it. Finally, in Section 4.5, we reuse operators from [12], prove that they preserve the finite model property and stable finiteness, and show how they are used in order to generate more theories for Table 2.

4.1 The Table

The columns left to the vertical double-line of Table 2 correspond to possible combinations of properties. In them, T means that the property holds, while F means that it does not. The first 5 columns correspond to properties already studied in [12], and the next two columns correspond to **FM** and **SF**. The columns right to the vertical double-line correspond to possible signatures: empty or non-empty, and one-sorted or many-sorted. White cells correspond to cases where a theory with the combination of properties induced by the row exists in a signature that is induced by the column. In such a case, the name of the theory is written. The theories themselves are defined in Figures 5, 7 and 8, axiomatically. Red cells correspond to the cases where there is no such theory. In such a case, the theorem that excludes this possibility is written. If that theorem is from [12], we simply write [12].

Example 1. Line 1 of Table 2 corresponds to theories that admit all studied properties. We see that there is such a theory in each of the studied types of signatures (e.g., for the empty one-sorted signature, the theory $\mathcal{T}_{\geq n}$ exhibits all properties). In contrast, line 3 corresponds to theories that admit all properties but strong finite witnessability. We see that such theories exist in non-empty signatures, but not in empty signatures. This is thanks to Theorem 6.

Section 3, as well as results from [12], make some potential rows of Table 2 completely red. To allow this table to fit a single page, we chose to erase such rows. For example, by Theorem 1, there are no theories that are stably finite but do not have the finite model property, in any signature. Thus, no rows that represent such theories appear in the table.

In the remainder of this section, we describe the various theories that populate the cells of the table. Fortunately, all theories from [12] can be reused to exhibit also the new properties **SF** and **FM**, or their negations. These are described in Section 4.2. However, the theories from [12] alone are not enough. Hence we introduce several new theories in Sections 4.3 and 4.4. Some of them are relatively simple, and are described in Section 4.3. Most of them, however, are more complex, and rely on the Busy Beaver function from theoretical computer science. We discuss these theories in Section 4.4.

4.2 Theories from [12]

For completeness, we include in Figure 5 the axiomatizations of all theories from [12] that are used in Table 2 (Figure 6 includes the definitions of formulas that are abbreviated in Figure 5, such as $\psi_{\geq n}^{=}$ from the definition of \mathcal{T}_{f}). For lack of space, however, we refrain from elaborating on these theories, and refer the reader to their detailed description in [12]. For the theories of Figure 5, whether they admit the properties from

Name	Sig.	Axiomatization		_	
$\mathcal{T}_{\geq n}$	Σ_1	$\{\psi_{\geq n}\}$	Name	Sig.	Axiomatization
$\mathcal{T}^\infty_{even}$	Σ_1	$\{\neg\psi_{=2k+1}: k \in \mathbb{N}\}$	$\mathcal{T}_{2,3}$	Σ_2	$\{(\psi_{=2}^{\sigma} \land \psi_{\geq k}^{\sigma_2}) \lor (\psi_{\geq 3}^{\sigma} \land \psi_{\geq 3}^{\sigma_2}) : k \in \mathbb{N}\}$
\mathcal{T}_∞	Σ_1	$\{\neg\psi_{=2k+1}: k \in \mathbb{N}\}$ $\{\psi_{\geq k}: k \in \mathbb{N}\}$	\mathcal{T}_2^∞		
$\mathcal{T}_{n,\infty}$	Σ_1	$\{\psi_{=n} \lor \psi_{\geq k} : k \in \mathbb{N}\}$	\mathcal{T}_1^{odd}	Σ_2	$\{\psi_{=1}^{\sigma}\} \cup \{\neg \psi_{=2k}^{\sigma_2} : k \in \mathbb{N}\}$
$\mathcal{T}_{\leq n}$	Σ_1	$\{\psi_{\leq n}\}$	\mathcal{T}_1^∞	Σ_2	$\{\psi_{=1}^{\sigma}\} \cup \{\psi_{\geq k}^{\sigma_2} : k \in \mathbb{N}\}$
$\mathcal{T}_{m,n}$	Σ_1	$\{\psi_{=m} \lor \psi_{=n}\}$			

Name	Sig.	Axiomatization
\mathcal{T}_{f}	Σ_s	$ \{ [\psi_{\geq f_1(k)}^{=} \land \psi_{\geq f_0(k)}^{\neq}] \lor \bigvee_{i=1}^k [\psi_{=f_1(i)}^{=} \land \psi_{=f_0(i)}^{\neq}] : k \in \mathbb{N} \setminus \{0\} \} $ $ Ax(\mathcal{T}_f) \cup \{\psi_{\lor}\} $
-		
$\mathcal{T}_{2,\infty}^{ eq}$	Σ_s	$\{[\psi_{=2} \land \forall x. p(x)] \lor [\psi_{\geq k} \land \forall x. \neg p(x)] : k \in \mathbb{N}\}$
$\mathcal{T}_{odd}^{ eq}$	Σ_s	$\{\psi_{=1} \lor [\neg \psi_{=2k} \land \forall x. \neg p(x)] : k \in \mathbb{N}\}$
$\mathcal{T}_{1,\infty}^{ eq}$	Σ_s	$\{\psi_{=1} \lor [\psi_{\geq k} \land \forall x. \neg p(x)] : k \in \mathbb{N}\}$

Fig. 5: Theories for Table 2 that were studied in [12]; p(x) stands for s(x) = x. In \mathcal{T}_f , f is any non-computable function from the positive integers to $\{0, 1\}$, such that for every $k \ge 0$, f maps half of the numbers between 0 and 2^k to 1, and the other half to 0. In [12], such a function was proven to exist.

{SI, SM, FW, SW, CV} or not was already established in [12]. For each of them, here, we also check and prove whether they admit the new properties FM and SF.

For example, for each n, $\mathcal{T}_{\geq n}$ consists of all Σ_1 -structures that have at least n elements. This theory was shown in [12] to be strongly finitely witnessable, and so by Theorem 3 it is also stably finite. Then, by Theorem 1, it also admits the finite model property.

It is worth mentioning that $\mathcal{T}_{2,3}$ was first introduced in [2], in the context of shiny theories, where it was shown to have the finite model property, while not being stably finite. An alternative proof of this fact goes as follows: it was proven in [10] that $\mathcal{T}_{2,3}$ is: (*i*) finitely witnessable; (*ii*) not strongly finitely witnessable; and (*iii*) smooth. By Theorem 2 and (*i*), it also has the finite model property. But since it is over an empty signature, by (*ii*), (*iii*) and Theorem 6, we have that it cannot be stably finite.

4.3 New Theories: The Simple Cases

While the theories from Figure 5 suffice to populate many cells of Table 2, they are not enough. Hence we describe new theories, not taken from [12]. The simplest theories that we have added can be found in Figure 7, and are described below.

 \mathcal{T}^{∞} is a theory with three distinct groups of models: its first group consists of models \mathcal{A} that have $|\sigma^{\mathcal{A}}| = 1$ and $\sigma_2^{\mathcal{A}}$ infinite; its second group, of models \mathcal{A} where both

$$\begin{split} \psi_{\geq n}^{=} &= \exists \overrightarrow{x}. \bigwedge_{i=1}^{n} p(x_{i}) \wedge \delta_{n} \qquad \psi_{=n}^{=} = \exists \overrightarrow{x}. \left[\bigwedge_{i=1}^{n} p(x_{i}) \wedge \delta_{n} \wedge \forall x. \left[p(x) \rightarrow \bigvee_{i=1}^{n} x = x_{i} \right] \right] \\ \psi_{\geq n}^{\neq} &= \exists \overrightarrow{x}. \bigwedge_{i=1}^{n} \neg p(x_{i}) \wedge \delta_{n} \qquad \psi_{=n}^{\neq} = \exists \overrightarrow{x}. \left[\bigwedge_{i=1}^{n} \neg p(x_{i}) \wedge \delta_{n} \wedge \forall x. \left[\neg p(x) \rightarrow \bigvee_{i=1}^{n} x = x_{i} \right] \right] \\ \psi_{\vee} &= \forall x. \left[(s(s(x)) = x) \lor (s(s(x)) = s(x)) \right] \end{split}$$

Fig. 6: Formulas for Σ_s -theories. \overrightarrow{x} stands for x_1, \ldots, x_n . δ_n stands for $\bigwedge_{1 \le i \le j \le n} \neg (x_i = x_j)$, and p(x) stands for s(x) = x.

Name	Signature	Axiomatization
\mathcal{T}^∞	Σ_2	$\{(\psi_{=1}^{\sigma} \land \psi_{\geq k}^{\sigma_2}) \lor diag^{\sigma, \sigma_2}(k+2) : k \in \mathbb{N}\}$
$\mathcal{T}_{m,n}^\infty$	Σ_2	$\{\psi_{=\max\{m,n\}}^{\sigma} \lor (\psi_{=\min\{m,n\}}^{\sigma} \land \psi_{\geq k}^{\sigma_2}) : k \in \mathbb{N}\}$
$\mathcal{T}^\infty_{\neq}$	Σ_s^2	$\{(\psi_{=1}^{\sigma} \land \psi_{\geq k}^{\sigma_2}) \lor (diag^{\sigma,\sigma_2}(k+2) \land \forall x. \neg p(x) : k \in \mathbb{N}\}$
$\mathcal{T}^3_{2,3}$	Σ_3	$\{\psi_{=1}^{\sigma_3}\} \cup \{(\psi_{=2}^{\sigma} \wedge \psi_{\geq k}^{\sigma_2}) \lor (\psi_{\geq 3}^{\sigma} \wedge \psi_{\geq 3}^{\sigma_2}) : k \in \mathbb{N}\}$

Fig. 7: Simple theories for Table 2. $diag^{\sigma,\sigma_2}(k+2)$, for any $k \in \mathbb{N}$, stands for the formula $(\psi_{\geq k+2}^{\sigma} \wedge \psi_{\geq k+2}^{\sigma_2}) \vee \bigvee_{i=2}^{k+2} (\psi_{=i}^{\sigma} \wedge \psi_{=i}^{\sigma_2})$, and p(x) stands for s(x) = x.

 $\sigma^{\mathcal{A}}$ and $\sigma_{2}^{\mathcal{A}}$ are infinite; and its third group, of models \mathcal{A} where $|\sigma^{\mathcal{A}}| = |\sigma_{2}^{\mathcal{A}}|$ is any value $k \geq 2$. In its axiomatization, one finds the formula $diag^{\sigma,\sigma_{2}}(k+2)$, equal to $(\psi_{\geq k+2}^{\sigma} \wedge \psi_{\geq k+2}^{\sigma_{2}}) \vee \bigvee_{i=2}^{k+2} (\psi_{=i}^{\sigma} \wedge \psi_{=i}^{\sigma_{2}})$ for $k \in \mathbb{N}$: that formula characterizes the models \mathcal{A} of \mathcal{T}^{∞} that lie in the diagonal, that is, where $|\sigma^{\mathcal{A}}| = |\sigma_{2}^{\mathcal{A}}|$ (and this value is greater than 1).

 $\mathcal{T}_{m,n}^{\infty}$ is a theory that depends on two distinct positive integers m and n, and without loss of generality let us suppose m > n, when the theory has two types of models \mathcal{A} : in the first, $|\sigma^{\mathcal{A}}|$ equals m, while $\sigma_2^{\mathcal{A}}$ can be anything; in the second, $|\sigma^{\mathcal{A}}|$ equals n, and then $\sigma_2^{\mathcal{A}}$ must be infinite.

The models \mathcal{A} of the Σ_s^2 -theory $\mathcal{T}_{\neq}^{\infty}$ have either: $|\sigma^{\mathcal{A}}| = 1$, $|\sigma_2^{\mathcal{A}}| \ge \omega$ and $s^{\mathcal{A}}$ the identity function; both $\sigma^{\mathcal{A}}$ and $\sigma_2^{\mathcal{A}}$ infinite, and $s^{\mathcal{A}}$ with no fixed points; or $|\sigma^{\mathcal{A}}| = |\sigma_2^{\mathcal{A}}|$ equal to any number in $\mathbb{N} \setminus \{0, 1\}$, and again $s^{\mathcal{A}}$ with no fixed points.

Finally, $\mathcal{T}_{2,3}^3$ is made up of just the models \mathcal{A} of $\mathcal{T}_{2,3}$ (see Figure 5) with an extra domain associated to the new sort σ_3 such that $|\sigma_3^{\mathcal{A}}| = 1$.

4.4 New Theories: The Busy Beaver

So far we have seen that the theories from [12], together with a small set of simple new theories, can already get us quite far in filling Table 2. However, for several com-

Name	Signature	Axiomatization
\mathcal{T}_{ς}	Σ_1	$\{\psi_{\geq\varsigma(k+2)}\vee\bigvee_{i=2}^{k+2}\psi_{=\varsigma(i)}:k\in\mathbb{N}\}$
$\mathcal{T}^\infty_\varsigma$	Σ_2	$\{(\psi_{=1}^{\sigma} \wedge \psi_{\geq k}^{\sigma_2}) \lor diag_{\varsigma}^{\sigma, \sigma_2}(k+2) : k \in \mathbb{N}\}$
$\mathcal{T}_n^{\varsigma}$	Σ_2	$\{\psi_{=n}^{\sigma}\} \cup \{\psi_{\geq \varsigma(k+2)}^{\sigma_2} \lor \bigvee_{i=2}^{k+2} \psi_{=\varsigma(i)}^{\sigma_2} : k \in \mathbb{N}\}$
$\mathcal{T}_{m,n}^{\varsigma}$	Σ_2	$\{(\psi_n^{\sigma} \wedge \psi_{\geq k}^{\sigma_2}) \lor (\psi_m^{\sigma} \wedge \psi_{\geq \varsigma(k+1)}^{\sigma_2}) \lor \bigvee_{i=1}^{k+1} (\psi_m^{\sigma} \wedge \psi_{=\varsigma(i)}^{\sigma_2}) : k \in \mathbb{N}\}$
$\mathcal{T}^s_{\varsigma}$	Σ_s	$\{(\psi_{\geq k+1} \land \psi_{\geq \varsigma^{-1}(k+1)}^{=}) \lor \bigvee_{i=1}^{k+1} (\psi_{=i} \land \psi_{=\varsigma^{-1}(i)}^{=}) : k \in \mathbb{N}\}$
$\mathcal{T}_{\varsigma}^{\neq}$	Σ_s	$\{(\psi_{=2} \land \forall x. \neg p(x)) \lor ((\psi_{\geq \varsigma(k+2)} \lor \bigvee_{i=2}^{k+2} \psi_{=\varsigma(i)}) \land \forall x. p(x)) : k \in \mathbb{N}\}$
$\mathcal{T}_{\varsigma,1}^{ eq}$	Σ_s	$\{\psi_{=1} \lor ((\psi_{\geq \varsigma(k+2)} \lor \bigvee_{i=2}^{k+2} \psi_{=\varsigma(i)}) \land \forall x. \neg p(x)) : k \in \mathbb{N}\}$
$\mathcal{T}_\varsigma^\vee$	Σ_s	$\{\psi_{\vee}\} \cup \{(\psi_{\geq k+1} \land \psi_{\geq \varsigma^{-1}(k+1)}^{=}) \lor \bigvee_{i=1}^{k+1} (\psi_{=i} \land \psi_{=\varsigma^{-1}(i)}^{=}) : k \in \mathbb{N}\}$
$\mathcal{T}_{\varsigma}^{=}$	Σ_s^2	$\{\psi_{\geq k+2}^{=} \to \psi_{\geq \varsigma(k+2)}^{\sigma_2} : k \in \mathbb{N}\}$
$\mathcal{T}_{\varsigma}^2$	Σ_s^2	$\{(\psi_{=1}^{\sigma} \wedge \psi_{\geq k}^{\sigma_2}) \lor (\psi_{\geq k+2}^{=} \rightarrow \psi_{\geq \varsigma(k+2)}^{\sigma_2}) : k \in \mathbb{N}\}$
$\mathcal{T}_{\varsigma\vee}^{=}$	Σ_s^2	$\{\psi_{\vee}\} \cup \{(\psi_{=1}^{\sigma} \land \psi_{\geq k}^{\sigma_2}) \lor (\psi_{\geq k+2}^{=} \to \psi_{\geq \varsigma(k+2)}^{\sigma_2}) : k \in \mathbb{N}\}$
$\mathcal{T}^\infty_{\varsigma eq}$	Σ_s^2	$\{(\psi_{=1}^{\sigma} \land \psi_{\geq k}^{\sigma_2}) \lor (diag_{\varsigma}^{\sigma,\sigma_2}(k+2) \land \forall x. \neg p(x)) : k \in \mathbb{N}\}$
$\mathcal{T}^{\infty,3}_{\varsigma}$	Σ_3	$\{\psi_{=1}^{\sigma_3}\} \cup \{(\psi_{=1}^{\sigma} \land \psi_{\geq k}^{\sigma_2}) \lor diag_{\varsigma}^{\sigma,\sigma_2}(k+2) : k \in \mathbb{N}\}$

Fig. 8: Busy Beaver Theories for Table 2. $diag_{\varsigma}^{\sigma,\sigma_2}(k+2)$ stands, for each $k \in \mathbb{N}$, for $(\psi_{\geq\varsigma(k+2)}^{\sigma} \land \psi_{\geq\varsigma(k+2)}^{\sigma_2}) \lor \bigvee_{i=2}^{k+2} (\psi_{=\varsigma(i)}^{\sigma} \land \psi_{=\varsigma(i)}^{\sigma_2})$, and p(x) for s(x) = x; in $\mathcal{T}_{m,n}^{\varsigma}$, we assume w.l.g. $m \ge n$.

binations, it seems that more complex theories are needed. For this purpose, we utilize the well-known Busy Beaver function, and define various theories based on it. In this section, we describe these theories. First, in Section 4.4.1, we review the Busy Beaver function, and explain why it is useful in our context. Then, in Sections 4.4.2 to 4.4.6, we describe the theories that make use of it, separated according to their signatures.

4.4.1 On The Busy Beaver Function The Busy Beaver function, here denoted ς , is an old acquaintance of theoretical computer scientists: essentially, given any $n \in \mathbb{N}$, $\varsigma(n)$ is the maximum number of 1's a Turing machine with at most n states can write to it's tape when it halts, if the tape is initialized to be all 0's. Somewhat confusingly, any Turing machine that achieves that number is also called a Busy Beaver.

It is possible to prove that $\varsigma(n) \in \mathbb{N}$ for any $n \in \mathbb{N}$ (see [7]), and so we may write $\varsigma : \mathbb{N} \to \mathbb{N}$; furthermore, ς is increasing. But the very desirable property of ς is that it is not only increasing, but actually very rapidly increasing.

More formally, Radó proved, in the seminal paper [7], that ς grows asymptotically faster than any computable function (being, therefore, non-computable). That is, for every computable function $f : \mathbb{N} \to \mathbb{N}$, there exists $N \in \mathbb{N}$ such that $\varsigma(n) > f(n)$ for all $n \ge N$. Despite that, the Busy Beaver starts somewhat slowly: $\varsigma(0) = 0$, $\varsigma(1) = 1$, $\varsigma(2) = 4$, $\varsigma(3) = 6$ and $\varsigma(4) = 13$; the exact value of $\varsigma(5)$ (and actually $\varsigma(n)$ for any $n \ge 5$) is not known, but is at least 4098 ([4]). The fact that ς grows eventually faster than any computable function is a great property to have when constructing theories that admit the finite model property, while not being finitely witnessable. Roughly speaking, if the cardinalities of models of a theory are related to ς , this guarantees that it has models of sufficiently large finite size, while not being finitely witnessable since its models grow too fast: by carefully choosing formulas ϕ_n that hold only in the "*n*-th model" of the theory (when ordered by cardinality), the number of variables of $wit(\phi_n)$ offers an upper bound to $\varsigma(n)$ and is therefore not computable, leading to a contradiction with the fact that *wit* is supposed to be computable. Notice that, despite the dependency of our theories on the Busy Beaver, the function is not actually part of their signatures.

Now we present the theories that are based on ς . These theories are axiomatized in Figure 8.

4.4.2 A Σ_1 -Theory The most basic Busy Beaver theory is \mathcal{T}_{ς} . This is the Σ_1 -theory whose models have cardinality $\varsigma(k)$, for some $k \ge 2$, or are infinite: that is, \mathcal{T}_{ς} has models with 4 elements, 6, 13 and so on. This theory forms the basis to all other theories of this section, that are designed to admit various properties from Table 2.

By itself, T_c has the finite model property while not being (strongly) finitely witnessable. It was in fact constructed precisely to exhibit this. As it turns out, it is also not smooth, but does satisfy all other properties. To populate other rows in the table that correspond to theories with other combinations of properties, more theories are needed, with richer signatures.

4.4.3 Σ_2 -Theories To fill the rows that correspond to other combinations, we introduce several Σ_2 theories.

The Σ_2 -theory $\mathcal{T}_{\varsigma}^{\infty}$ is more complex. It has, essentially, three classes of models: the first is made up of structures \mathcal{A} where $|\sigma^{\mathcal{A}}| = 1$ and $\sigma_2^{\mathcal{A}}$ is infinite; the second, of structures where both $\sigma^{\mathcal{A}}$ and $\sigma_2^{\mathcal{A}}$ are infinite; and the third, of structures where $|\sigma^{\mathcal{A}}| =$ $|\sigma_2^{\mathcal{A}}|$ is a finite value that equals $\varsigma(k)$, for some $k \geq 2$. The formula $diag_{\varsigma}^{\sigma,\sigma_2}(k+2)$, for $k \geq 2$, in the axiomatization equals $(\psi_{\geq\varsigma(k+2)}^{\sigma} \wedge \psi_{\geq\varsigma(k+2)}^{\sigma_2}) \vee \bigvee_{i=2}^{k+2} (\psi_{=\varsigma(i)}^{\sigma} \wedge \psi_{=\varsigma(i)}^{\sigma_2})$ and is similar to $diag^{\sigma,\sigma_2}(k+2)$ from \mathcal{T}^{∞} , characterizing the models \mathcal{A} where $|\sigma^{\mathcal{A}}| =$ $|\sigma_2^{\mathcal{A}}|$ and that value, if it is not infinite, equals $\varsigma(k+2)$.

For each n > 0, $\mathcal{T}_n^{\varsigma}$ has as interpretations those \mathcal{A} with $|\sigma^{\mathcal{A}}| = n$, and $|\sigma_2^{\mathcal{A}}|$ either infinite or equal to $\varsigma(k)$, for some $k \ge 2$ (so $(|\sigma^{\mathcal{A}}|, |\sigma_2^{\mathcal{A}}|)$ may equal (n, 4), (n, 6), (n, 13) and so on).

 $\mathcal{T}_{m,n}^{\varsigma}$ is a Σ_2 -theory that can be seen as some sort of combination of $\mathcal{T}_{m,n}^{\varsigma}$ and $\mathcal{T}_n^{\varsigma}$, dependent on two distinct positive integers m and n. Consider the case where the former is the greater of the two (the other cases are similar). In this case, we may divide its interpretations \mathcal{A} into three classes: those with $|\sigma^{\mathcal{A}}| = n$ and $\sigma_2^{\mathcal{A}}$ infinite; those with $|\sigma^{\mathcal{A}}| = m$ and $\sigma_2^{\mathcal{A}}$ infinite; does with $|\sigma^{\mathcal{A}}| = m$ and $|\sigma_2^{\mathcal{A}}|$ equal to some $\varsigma(k)$, for $k \geq 2$.

4.4.4 Σ_s -Theories For some lines of Table 2, e.g. line 7, empty signatures are not enough for presenting examples. Hence we also introduce Σ_s -theories.

We start with \mathcal{T}_{c}^{s} , which is, arguably, the most confusing theory we here define: we are forced to appeal not only to the special cardinality formulas found in Figure 6, but also to the function ς^{-1} , which is a left inverse of ς . More formally, $\varsigma^{-1} : \mathbb{N} \to \mathbb{N}$ is the only function such that $\varsigma^{-1}(k) = \min\{l : \varsigma(l+1) > \varsigma(k)\}$: so $\varsigma^{-1}(0) = 0$, $\varsigma^{-1}(1) = \varsigma^{-1}(2) = \varsigma^{-1}(3) = 1$, $\varsigma^{-1}(4) = \varsigma^{-1}(5) = 2$, $\varsigma^{-1}(6) = \cdots = \varsigma^{-1}(12) = 3$, $\varsigma^{-1}(13) = \cdots = \varsigma^{-1}(4097) = 4$, and further values of ς^{-1} are currently unknown. From the definition of ς^{-1} , we have that $\varsigma(\varsigma^{-1}(k)) \leq k$ and $\varsigma^{-1}(\varsigma(k)) = k$. ς^{-1} is not computable given that, since $\varsigma^{-1}(k) = \min\{l : \varsigma(l+1) > \varsigma(k)\}$ by definition, $\varsigma^{-1}(k+1) \neq \varsigma^{-1}(k)$ iff k+1 is a value of ς : so, an algorithm to compute the values of ς could be obtained by simply computing the values of ς^{-1} and checking where there is a change.

 $\mathcal{T}_{\varsigma}^{s}$ is then the Σ_{s} -theory with models \mathcal{A} with any cardinality $k+1 \geq 1$, such that $s^{\mathcal{A}}(a) = a$ holds for precisely $\varsigma^{-1}(k+1)$ elements of \mathcal{A} , and so $s^{\mathcal{A}}(a) \neq a$ holds for $k+1-\varsigma^{-1}(k+1)$ elements, being the function $k \mapsto k+1-\varsigma^{-1}(k+1)$ itself non-decreasing, given that $\varsigma^{-1}(k+1)$ can equal either $\varsigma^{-1}(k)$ or $\varsigma^{-1}(k) + 1$.

Example 2. We mention some $\mathcal{T}_{\varsigma}^{s}$ -structures as examples: a structure \mathcal{A} with $|\sigma^{\mathcal{A}}| = 1$ and $s^{\mathcal{A}}$ the identity; a structure \mathcal{B} with $|\sigma^{\mathcal{B}}| = 2$ and $s^{\mathcal{A}}$ a constant function; a structure \mathcal{C} with $|\sigma^{\mathcal{C}}| = 3$ (say $\sigma^{\mathcal{C}} = \{a, b, c\}$) and $s^{\mathcal{C}}$ the identity for only one of these elements (e.g., s^{C} can be a constant function, but now there are further possibilities such as $s^{\mathcal{C}}(a) = s^{\mathcal{C}}(b) = a$ and $s^{\mathcal{C}}(c) = b$; and a structure \mathcal{D} with $|\sigma^{\mathcal{D}}| = 4$ (say $\sigma^{\mathcal{D}} = \{a, b, c, d\}$ and $s^{\mathcal{D}}$ the identity for only two of these elements (e.g., $s^{\mathcal{D}}(a) =$ $s^{\mathcal{D}}(b) = s^{\mathcal{D}}(c) = a \text{ and } s^{\mathcal{D}}(d) = d);$

Next, we continue to describe other Σ_s theories.

 \mathcal{T}_{s}^{\neq} has essentially two classes of models \mathcal{A} : those with $|\sigma^{\mathcal{A}}| = 2$ and $s^{\mathcal{A}}$ never the identity; and those with $|\sigma^{\mathcal{A}}|$ equal to $\varsigma(k)$ or infinite, for some $k \geq 2$, and $s^{\mathcal{A}}$ the identity.

 $\mathcal{T}_{s,1}^{\neq}$ is very similar to \mathcal{T}_{s}^{\neq} : the difference lies on where s will be the identity: while in $\mathcal{T}_{\varsigma}^{\neq}$ the function s is the identity for all interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}| > 2$, s in $\mathcal{T}_{\varsigma,1}^{\neq}$ is the identity only for the interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}| = 1$. So, in $\mathcal{T}_{\varsigma,1}^{\neq}$, we have a model \mathcal{A} with $|\sigma^{\mathcal{A}}| = 1$ and $s^{\mathcal{A}}$ the identity, and then models \mathcal{A} with $|\sigma^{\mathcal{A}}| = \varsigma(k)$ for some $k \geq 2$ or infinite, and $s^{\mathcal{A}}(a)$ anything but a.

The Σ_s -theory $\mathcal{T}_{\varsigma}^{\vee}$ is then just $\mathcal{T}_{\varsigma}^s$, satisfying in addition the formula ψ_{\vee} (see Figure 6). It has models A of any finite cardinality k + 1, as long as $\varsigma^{-1}(k + 1)$ of these elements a satisfy $s^{\mathcal{A}}(a) = a$, or infinite cardinalities, as long as the number of elements a satisfying $s^{\mathcal{A}}(a) = a$ is infinite; additionally, $s^{\mathcal{A}}(s^{\mathcal{A}}(a))$ must always equal either $s^{\mathcal{A}}(a)$ or a itself.

4.4.5 Σ_s^2 -Theories Now for theories in a many-sorted non-empty signature. The Σ_s^2 -theory $\mathcal{T}_s^=$ appears simple, but is actually quite tricky: starting by the easy case, if $\sigma^{\mathcal{A}}$ has infinitely many elements *a* satisfying $s^{\mathcal{A}}(a) = a$, $\sigma_2^{\mathcal{A}}$ is also infinite. If, however, the number of elements $a \in \sigma^{\mathcal{A}}$ satisfying $s^{\mathcal{A}}(a) = a$ is finite (notice that, even if this is the case, $\sigma^{\mathcal{A}}$ may still be infinite) and equal to some k + 2, then $\sigma_2^{\mathcal{A}}$ has at least $\varsigma(k+2)$ elements. So, to give a better example, suppose $\sigma^{\mathcal{A}}$ has 2

elements satisfying $s^{\mathcal{A}}(a) = a$: then $\sigma_2^{\mathcal{A}}$ has at least $\varsigma(2) = 4$ elements, but may have any cardinality up to, and including, infinite ones; notice that in this example $\sigma^{\mathcal{A}}$ may be infinite as well, as long as only two of the elements satisfy $s^{\mathcal{A}}(a) = a$.

 $\mathcal{T}_{\varsigma}^2$ is the same as $\mathcal{T}_{\varsigma}^=$, but with extra models \mathcal{A} where $|\sigma^{\mathcal{A}}| = 1$ and $|\sigma_2^{\mathcal{A}}| \ge \omega$ (of course, then we have that $s^{\mathcal{A}}$ is the identity).

 $\mathcal{T}_{\varsigma\vee}^{=}$ is then the same as $\mathcal{T}_{\varsigma}^{2}$, with the added validity of the formula ψ_{\vee} ; So the models of $\mathcal{T}_{\varsigma\vee}^{=}$ are just models of $\mathcal{T}_{\varsigma}^{2}$ satisfying that $s^{\mathcal{A}}(s^{\mathcal{A}}(a))$ equals either $s^{\mathcal{A}}(a)$ or a itself. $\mathcal{T}_{\varsigma\neq}^{\infty}$ is just the Σ_{2} -theory $\mathcal{T}_{\varsigma}^{\infty}$ with the added function s such that, if $|\sigma^{\mathcal{A}}| = 1$, $s^{\mathcal{A}}$

is the identity; and if $|\sigma^{\mathcal{A}}| > 1$, $s^{\mathcal{A}}(a)$ is anything but a.

4.4.6 A Σ_3 -Theory Finally, $\mathcal{T}_{\varsigma}^{\infty,3}$ is obtained by adding a sort with a single element to the Σ_2 -theory $\mathcal{T}_{\varsigma}^{\infty}$, similarly to the definition of $\mathcal{T}_{2,3}^3$, that was based on the Σ_2 -theory $\mathcal{T}_{2,3}$ (see Section 4.3).

4.5 **Theory Operators**

There are two types of theories in Table 2: The first consists of base theories, such as $\mathcal{T}_{>n}$, that are axiomatized in Figures 5, 7 and 8. The second is obtained from the first, by applying several operators on theories. For example, the theories $(\mathcal{T}_{>n})^2$, $(\mathcal{T}_{>n})_s$, $((\mathcal{T}_{\geq n})^2)_s$, are all obtained from the base theory $\mathcal{T}_{\geq n}$. So far we have only described the theories of the first type. In this section we explain the theories of the second type.

The operators that are used in Table 2 were defined in [12], in order to be able to systematically generate examples in various signatures. For example, if \mathcal{T} is a Σ_1 theory, then $(\mathcal{T})^2$ is a Σ_2 -theory with the same axiomatization as \mathcal{T} , that is, the second sort is completely free and is not axiomatized in any way. For completeness sake, we include the definitions of these operators here:

Definition 1 (Theory Operators from [12]).

- 1. If \mathcal{T} is a Σ_1 -theory, then $(\mathcal{T})^2$ is the Σ_2 -theory axiomatized by $Ax(\mathcal{T})$.
- 2. Let Σ_n be an empty signature with sorts $S = \{\sigma_1, \ldots, \sigma_n\}$, and let \mathcal{T} be a Σ_n theory. The signature Σ_s^n has sorts S and a single unary function symbol s of arity $\sigma_1 \to \sigma_1$, and $(\mathcal{T})_s$ is the Σ_s^n -theory axiomatized by $Ax(\mathcal{T}) \cup \{\forall x. [s(x) = x]\},\$ where x is a variable of sort σ_1 .
- 3. Let \mathcal{T} be a theory over an empty signature with sorts $S = \{\sigma_1, \ldots, \sigma_n\}$. Then $(\mathcal{T})_{\vee}$ is the Σ_s^n -theory axiomatized by $Ax(\mathcal{T}) \cup \{\psi_{\vee}\}$ (see Figure 6.

It was proven in [12] that these operators preserve the properties SI, SM, FW, SW, CV, and the lack of them. Here we prove that the same holds for FM and SF as well.

Theorem 8. Let \mathcal{T} be a Σ_1 -theory. Then: \mathcal{T} is **FM**, or **SF**, w.r.t. $\{\sigma\}$ if and only if $(\mathcal{T})^2$ is, respectively, **FM**, or **SF** w.r.t. $\{\sigma, \sigma_2\}$.

Theorem 9. If \mathcal{T} is a theory over an empty signature Σ_n with sorts $S = \{\sigma_1, \ldots, \sigma_n\}$, then: \mathcal{T} is **FM**, or **SF**, w.r.t. S if and only if $(\mathcal{T})_s$ is, respectively, **FM**, or **SF**, w.r.t. S.

Theorem 10. If \mathcal{T} is a theory over an empty signature Σ_n with sorts $S = \{\sigma_1, \ldots, \sigma_n\}$, then: \mathcal{T} is **FM**, or **SF**, w.r.t. S if and only if $(\mathcal{T})_{\vee}$ is, respectively, **FM**, or **SF**, w.r.t. S.

Thus, in various cases, theories need not be invented from scratch, but can be generated from other theories. For example, the theory $\mathcal{T}_{\geq n}$ exhibits all studied properties, but is defined in a one-sorted signature. Using the operators, we obtain variants of this theory in all signature types, namely $(\mathcal{T}_{\geq n})^2$ for empty many-sorted signatures, $(\mathcal{T}_{\geq n})_s$ for non-empty one-sorted signatures, and $((\mathcal{T}_{\geq n})^2)_s$ for non-empty many-sorted signatures. The properties of the theories generated using these operators are guaranteed by Theorems 8 and 9, as well as the corresponding results from [12].

In two cases of theories defined using the Busy Beaver function, $\mathcal{T}_{\varsigma}^{\vee}$ and $\mathcal{T}_{\varsigma}^{=}$, we cannot obtain them by relying on Theorem 10 from, respectively, $\mathcal{T}_{\varsigma}^{s}$ and $\mathcal{T}_{\varsigma}^{=}$, since the signatures of the latter theories are not empty. Curiously, adding ψ_{\vee} to their axiomatizations still has the desirable outcome, but we prove this separately, without relying on Theorem 10. Extending Theorem 10 to non-empty signatures is left for future work.

The number of combinations of properties that we consider, together with the possible types of the signatures, adds up to $2^9 = 512$. Our negative results from Section 3 guarantee that only ~15% of the actual table can be filled with examples. The remaining ~85% are either colored in red or are excluded from the table for space considerations. As for the examples that can be given, notice that there are in total an astonishing number of 78 theories in our table. But, thanks to the theory operators of Definition 1, only 33 of them (~42%) had to be concretely axiomatized in Figures 5, 7 and 8. The remaining 45 theories were defined using the operators.

5 Conclusion

We examined, in addition to all properties considered in [12], the finite model property, and stable finiteness. Interesting restrictions for the combinations involving these properties were established. We also found interesting theories to fill in our table of combinations, most prominently those involving the Busy Beaver function as well as its inverse.

One possible direction this research could take is reasonably clear: considering the computability of the *mincard* function, what will, most probably, double the number of theories to be taken into consideration. Further interesting properties that could be considered include the decidability of the theory's axiomatization, or even its finiteness, and the satisfiability problem of the theory with respect to quantifier-free formulas.

Second, some of the negative results in [12] and in the present paper only hold with respect to the entire set of sorts S_{Σ} . We plan to study if they hold also with respect to proper subsets of sorts, and if they do not, to provide counterexamples to those generalizations.

References

[1] Aaron R. Bradley and Zohar Manna. *The Calculus of Computation*. Springer Berlin Heidelberg, 2007. DOI: 10.1007/978-3-540-74113-8.

- Filipe Casal and João Rasga. "Many-Sorted Equivalence of Shiny and Strongly Polite Theories". In: *Journal of Automated Reasoning* 60.2 (2018), pp. 221–236. ISSN: 1573-0670. DOI: 10.1007/s10817-017-9411-y.
- [3] Dejan Jovanović and Clark Barrett. *Polite Theories Revisited*. Tech. rep. TR2010-922. Depatrment of Computer Science, New York University, Jan. 2010.
- [4] Heiner Marxen and Jürgen Buntrock. "Attacking the Busy Beaver 5". In: *Bull. EATCS* 40 (1990), pp. 247–251.
- [5] María Monzano. "Introduction to Many-sorted Logic". In: *Many-sorted Logic and its Applications*. Ed. by K. Meinke and J. V. Tucker. Wiley professional computing. Wiley, 1993.
- [6] Greg Nelson and Derek C. Oppen. "Simplification by Cooperating Decision Procedures". In: ACM Trans. Program. Lang. Syst. 1.2 (1979), pp. 245–257. ISSN: 0164-0925. DOI: 10.1145/357073.357079.
- T. Radó. "On non-computable functions". In: *The Bell System Technical Journal* 41.3 (1962), pp. 877–884. DOI: 10.1002/j.1538–7305.1962.tb00480.x.
- [8] Silvio Ranise, Christophe Ringeissen, and Calogero G. Zarba. "Combining data structures with nonstably infinite theories using many-sorted logic". In: 5th International Workshop on Frontiers of Combining Systems - FroCoS'05. Ed. by Bernard Gramlich. Vol. 3717. Lecture Notes in Artificial Intelligence. Vienna/Austria: Springer, Sept. 2005, pp. 48–64. DOI: 10.1007/11559306.
- [9] Ying Sheng et al. "Polite Combination of Algebraic Datatypes". In: J. Autom. *Reason.* 66.3 (2022), pp. 331–355.
- [10] Ying Sheng et al. "Politeness and Stable Infiniteness: Stronger Together". In: *Automated Deduction – CADE 28*. Ed. by André Platzer and Geoff Sutcliffe. Cham: Springer International Publishing, 2021, pp. 148–165. ISBN: 978-3-030-79876-5.
- [11] Cesare Tinelli and Calogero Zarba. Combining decision procedures for theories in sorted logics. Tech. rep. 04-01. Department of Computer Science, The University of Iowa, Feb. 2004.
- [12] Guilherme Vicentin de Toledo, Yoni Zohar, and Clark Barrett. Combining Combination Properties: An Analysis of Stable Infiniteness, Convexity, and Politeness. Accepted to CADE 2023. 2023. arXiv: 2305.02384 [cs.L0].

A Preliminaries of the Appendix

The following are basic results in model theory of many-sorted logic that may be found in [5].

Lemma 1 ([5]). Let Σ be a signature; a set of Σ -formulas Γ is satisfiable if and only if each of its finite subsets Γ_0 is satisfiable.

Lemma 2 ([5]). Let Σ be a signature and Γ a set of Σ -formulas: if Γ is satisfiable, there exists an interpretation \mathcal{A} which satisfies Γ and where $\sigma^{\mathcal{A}}$ is countable whenever it is infinite.

The two following theorems are results from [12].

Theorem 11. Any theory over an empty signature that is stably infinite w.r.t. the set of all of its sorts is convex w.r.t. any set of sorts.

Theorem 12. Every one-sorted, non-stably-infinite theory \mathcal{T} with an empty signature is finitely witnessable w.r.t. its only sort.

We shall also make use of the following fact, found as a corollary of exercise 10.5 in Section 10 of [1].

Theorem 13. If \mathcal{T} is a theory over a one-sorted signature with only one unary function, and if \mathcal{T} is axiomatized by the empty set (that is, its functions are uninterpreted), then \mathcal{T} is convex.

Given a function symbol s of arity $\sigma \to \sigma$ and a variable x of sort σ , it is useful to define, recursively, the term $s^k(x)$ (for $k \in \mathbb{N}$) as follows: $s^0(x) = x$, and $s^{k+1}(x) = s(s^k(x))$.

B Proof of Theorem 2

Theorem 2. Any Σ -theory \mathcal{T} finitely witnessable with respect to $S \subseteq S_{\Sigma}$ also has the finite model property with respect to S.

Proof. Let *wit* be the witness for \mathcal{T} with respect to S, ϕ be a quantifier-free Σ -formula, and \mathcal{A} a \mathcal{T} -interpretation that satisfies ϕ : *wit* is a witness, and therefore ϕ and $\exists \vec{x} . wit(\phi)$ (for $\vec{x} = vars(wit(\phi)) \setminus vars(\phi)$) must be \mathcal{T} -equivalent, and so we must have that \mathcal{A} satisfies $\exists \vec{x} . wit(\phi)$; and then there must exist a \mathcal{T} -interpretation \mathcal{A}' , different from \mathcal{A} at most on \vec{x} , that satisfies $wit(\phi)$. Using again that *wit* is a witness, there is a \mathcal{T} -interpretation \mathcal{B} , with $\sigma^{\mathcal{B}} = vars_{\sigma}(wit(\phi))^{\mathcal{B}}$ for each $\sigma \in S$, that satisfies ϕ . But, because $vars_{\sigma}(wit(\phi))$, and thus $vars_{\sigma}(wit(\phi))^{\mathcal{B}}$, must be finite, we get $|\sigma^{\mathcal{B}}|$ is finite for each $\sigma \in S$, proving \mathcal{T} has the finite model property with respect to S. \Box

C Proof of Theorem 3

Theorem 3. Any Σ -theory \mathcal{T} strongly finitely witnessable with respect to $S \subseteq S_{\Sigma}$ is also stably finite with respect to S.

Proof. Let *wit* be the strong witness of \mathcal{T} with respect to S, ϕ be a quantifier-free Σ -formula, and \mathcal{A} a \mathcal{T} -interpretation that satisfies ϕ : we have that \mathcal{A} satisfies $\exists \vec{x} . wit(\phi)$, for $\vec{x} = vars(wit(\phi)) \setminus vars(\phi)$; so there must exist a \mathcal{T} -interpretation \mathcal{A}' , differing from \mathcal{A} at most on \vec{x} , that satisfies $wit(\phi)$. Let V be the set of variables of sort in S on $wit(\phi)$ or ϕ , and δ_V be the arrangement on V such that x is related to y iff $x^{\mathcal{A}'} = y^{\mathcal{A}'}$, which is clearly satisfied by \mathcal{A}' ; by strong finite-witnessability, there must exist a \mathcal{T} -interpretation \mathcal{B} that satisfies $wit(\phi) \wedge \delta_V$ with $\sigma^{\mathcal{B}} = vars_{\sigma}(wit(\phi) \wedge \delta_V)^{\mathcal{B}}$ for each $\sigma \in S$. Because \mathcal{B} satisfies $wit(\phi)$, it also satisfies $\exists \vec{x} . wit(\phi)$ and ϕ ; and $|\sigma^{\mathcal{B}}|$ is finite for each $\sigma \in S$, given $vars_{\sigma}(wit(\phi) \wedge \delta_V)$, and thus $vars_{\sigma}(wit(\phi) \wedge \delta_V)^{\mathcal{B}}$, are finite; finally, because \mathcal{B} satisfies x = y (for x and y variables of $wit(\phi)$ or ϕ) iff \mathcal{A}' satisfies x = y, we have that $|vars_{\sigma}(wit(\phi) \wedge \delta_V)^{\mathcal{B}}| = |vars_{\sigma}(wit(\phi) \wedge \delta_V)^{\mathcal{A}'}|$, and so

$$|\sigma^{\mathcal{B}}| = |vars_{\sigma}(wit(\phi) \land \delta_{V})^{\mathcal{B}}| = |vars_{\sigma}(wit(\phi) \land \delta_{V})^{\mathcal{A}'}| \le |\sigma^{\mathcal{A}'}| = |\sigma^{\mathcal{A}}|,$$

proving \mathcal{T} is stably finite with respect to S.

D Proof of Theorem 4

Theorem 4. If a Σ -theory \mathcal{T} has the finite model property with respect to a set of sorts S with |S| = 1, then \mathcal{T} is also stably finite with respect to S.

Proof. Let ϕ be a quantifier-free Σ -formula, and \mathcal{A} a \mathcal{T} -interpretation that satisfies ϕ : because \mathcal{T} has the finite model property, there is a \mathcal{T} -interpretation \mathcal{B} that satisfies ϕ with $|\sigma^{\mathcal{B}}|$ finite. There are then two cases to consider: if $|\sigma^{\mathcal{B}}| \leq |\sigma^{\mathcal{A}}|$, we already have that \mathcal{T} is stably finite; if $|\sigma^{\mathcal{B}}| > |\sigma^{\mathcal{A}}|$, then \mathcal{A} is a \mathcal{T} -interpretation that satisfies ϕ with $|\sigma^{\mathcal{A}}|$ finite (and this value is equal to or less than $|\sigma^{\mathcal{A}}|$), so \mathcal{T} is, again, stably finite.

E Proof of Theorem 5

Lemma 3. Let n be a natural number, and consider any subset A of \mathbb{N}^n equipped with the order such that $(p_1, \ldots, p_n) \leq (q_1, \ldots, q_n)$ iff $p_i \leq q_i$ for all $1 \leq i \leq n$: then A possesses at most a finite number of minimal elements under this order.

Proof. We will prove the result by induction on n, the result being obviously true if n = 0. So assume the result is true for n, and we shall prove it also holds for n + 1. We start by fixing a minimal element $(p_1, \ldots, p_n, p_{n+1})$ of A (if there are none, we are done): for any other minimal element $(q_1, \ldots, q_n, q_{n+1})$, we have that there must exist distinct $1 \le i, j \le n + 1$ such that $p_i > q_i$, and $p_j < q_j$; this way, $(q_1, \ldots, q_n, q_{n+1})$ is a minimal element of

$$A_{q_i}^i = \{(r_1, \dots, r_n, r_{n+1}) \in A : r_i = q_i\}$$

with the order induced by the order on A. By induction hypothesis, given this set may be simply considered as a subset of \mathbb{N}^n if we omit the coordinate *i*, the set of minimal elements of $A_{q_i}^i$ is finite. So every minimal element of A is either $(p_1, \ldots, p_n, p_{n+1})$, or an element of some $\min(A_{q_i}^i)$, the set of minimal elements of $A_{q_i}^i$: there are n + 1 possible values for *i* (*i.e.* 1 through n + 1), and $p_i - 1$ values for q_i (namely 1 through $p_i - 1$, given that $q_i < p_i$); so we can bound the number of minimal elements of A by the sum

$$1 + \sum_{i=1}^{m} \sum_{q_i=1}^{p_i-1} |\min(A_{q_i}^i)|,$$

which is finite, since it is a finite sum of finite numbers $|\min(A_{q_i}^i)|$. The result is then true for n + 1.

Theorem 5. If Σ is an empty signature with a finite set of sorts S_{Σ} , and the Σ -theory \mathcal{T} has the finite model property and is smooth with respect to S_{Σ} , then \mathcal{T} is also finitely witnessable with respect to S_{Σ} .

Proof. Suppose $S_{\Sigma} = \{\sigma_1, \ldots, \sigma_n\}$. Consider the set

$$A = \{ (|\sigma_1^{\mathcal{A}}|, \dots, |\sigma_n^{\mathcal{A}}|) : \mathcal{A} \text{ is a } \mathcal{T} \text{-interpretation} \} \cap \mathbb{N}^n,$$

which is not empty because \mathcal{T} has the finite model property: by Lemma 3 there exist \mathcal{T} -interpretations \mathcal{A}_1 through \mathcal{A}_m that correspond to the minimal elements of A (notice we are using that two Σ -structures \mathcal{A} and \mathcal{B} are isomorphic iff $|\sigma_i^{\mathcal{A}}| = |\sigma_i^{\mathcal{B}}|$ for each $1 \leq i \leq n$, given the signature is empty; this is necessary as otherwise one would need to consider A as a multiset to account for non-isomorphic models \mathcal{A} and \mathcal{B} that still have minimal cardinalities $(|\sigma_1^{\mathcal{A}} \dots, |\sigma_n^{\mathcal{A}}|) = (|\sigma_1^{\mathcal{B}}|, \dots, |\sigma_n^{\mathcal{B}}|)$, and in this case Lemma 3 does not necessarily hold); so we define $m_{i,j} = |\sigma_i^{\mathcal{A}_j}|$ and $m_i = \max\{m_{i,j} : 1 \leq j \leq m\}$. Once we fix a quantifier-free Σ -formula ϕ , take fresh variables $\{x_i^1, \dots, x_{m_i}^i\}$ of sort σ_i , and consider the witness

$$wit(\phi) = \phi \wedge \delta, \quad \text{where} \quad \delta = \bigwedge_{i=1}^n \bigwedge_{1 \le j < k \le m_i} (x_j^i = x_k^i).$$

To prove ϕ and $\exists \vec{x} . wit(\phi)$ are \mathcal{T} -equivalent is easy: start with a \mathcal{T} -interpretation \mathcal{A} that satisfies ϕ , and since δ is satisfied by all \mathcal{T} -interpretations by design, \mathcal{A} satisfies $\phi \land \delta$, and thus $\exists \vec{x} . wit(\phi)$. Conversely, if the \mathcal{T} -interpretation \mathcal{A} satisfies $\exists \vec{x} . wit(\phi)$, given that none of the variables in \vec{x} occur in ϕ , \mathcal{A} satisfies $\phi \land \exists \vec{x} . \delta$, and thus ϕ .

Now, suppose the \mathcal{T} -interpretation \mathcal{A} satisfies $wit(\phi)$: we then take sets A_i , for each $1 \le i \le n$, disjoint from each other and $\sigma_i^{\mathcal{A}}$, with

$$\max\{0, \quad m_i - |vars_{\sigma_i}(\phi)^{\mathcal{A}}|\}$$

elements. Finally, we define an interpretation \mathcal{B} by making: $\sigma_i^{\mathcal{B}} = vars_{\sigma_i}(\phi)^{\mathcal{A}} \cup A_i$ (which has at least m_i elements for every sort σ_i , what makes of \mathcal{B} a \mathcal{T} -interpretation, given this theory is smooth on an empty signature). And, for the evaluation of the variables, we demand that: $x^{\mathcal{B}} = x^{\mathcal{A}}$ for all variables x in ϕ ; the map taking x_j^i in $\{x_1^{i_1}, \ldots, x_{m_i}^{i_i}\}$ to $(x_j^{i_j})^{\mathcal{B}}$ in A_i is a surjection for each $1 \leq i \leq n$ (what is possible by definition of A_i); and for other variables $x, x^{\mathcal{B}}$ is defined arbitrarily. Then it is true that: \mathcal{B} satisfies ϕ , since the truth-value of ϕ , as a formula on the empty signature, is determined by the truth-value of its atomic subformulas x = y, and $x^{\mathcal{B}} = y^{\mathcal{B}}$ iff $x^{\mathcal{A}} = y^{\mathcal{A}}$ (since $x^{\mathcal{B}} = x^{\mathcal{A}}$ and $y^{\mathcal{B}} = y^{\mathcal{A}}$); \mathcal{B} satisfies $wit(\phi)$, since δ is \mathcal{T} -valid and \mathcal{B} is a \mathcal{T} -interpretation (given \mathcal{T} is smooth); and $\sigma^{\mathcal{B}} = vars_{\sigma}(wit(\phi))^{\mathcal{B}}$ for each $\sigma \in S_{\Sigma}$, by definition of \mathcal{B} .

F Proof of Theorem 6

Theorem 6. If Σ is an empty signature with a finite set of sorts S_{Σ} , and the Σ -theory \mathcal{T} is stably finite and smooth with respect to S_{Σ} , then \mathcal{T} is also strongly finitely witnessable with respect to S_{Σ} .

Proof. Suppose $S_{\Sigma} = \{\sigma_1, \ldots, \sigma_n\}$. Consider the set (also used in Theorem 5)

$$A = \{ (|\sigma_1^{\mathcal{A}}|, \dots, |\sigma_n^{\mathcal{A}}|) : \mathcal{A} \text{ is a } \mathcal{T} \text{-interpretation} \} \cap \mathbb{N}^n :$$

A is not empty because \mathcal{T} is stably finite, and by Lemma 3 there exist \mathcal{T} -interpretations \mathcal{A}_1 through \mathcal{A}_m such that $(|\sigma_1^{\mathcal{A}_1}|, \ldots, |\sigma_n^{\mathcal{A}_n}|)$ through $(|\sigma_1^{\mathcal{A}_m}|, \ldots, |\sigma_n^{\mathcal{A}_m}|)$ are the minimal elements of A (under the order described in Lemma 3). We then take $m_{i,j} = |\sigma_i^{\mathcal{A}_j}|$ and $m_i = \max\{|\sigma_i^{\mathcal{A}_j}| : 1 \le j \le m\}$. Given a quantifier-free Σ -formula ϕ , we take fresh variables x_1^i through $x_{m_i}^i$ of sort σ_i , and consider the strong witness

$$wit(\phi) = \phi \wedge \delta, \quad \text{where} \quad \delta = \bigvee_{j=1}^{m} \bigwedge_{i=1}^{n} \bigwedge_{1 \le k < l \le m_{i,j}} \neg (x_k^i = x_l^i).$$

If a \mathcal{T} -interpretation \mathcal{A} satisfies ϕ , take the set of variables V of ϕ , and the arrangement δ_V on V such that x is related to y iff $x^{\mathcal{A}} = y^{\mathcal{A}}$, and it is clear that \mathcal{A} satisfies δ_V : we can find a \mathcal{T} -interpretation \mathcal{B} that satisfies δ_V with $|\sigma_i^{\mathcal{B}}|$ finite and satisfying $|\sigma_i^{\mathcal{B}}| \leq |\sigma_i^{\mathcal{A}}|$ for each $1 \leq i \leq n$. Then, we can construct a \mathcal{T} -interpretation \mathcal{B}' , that differs from \mathcal{B} at most on $\overrightarrow{x} = vars(wit(\phi)) \setminus vars(\phi) = vars(\delta)$, where δ is satisfied (since we must have $(|\sigma_1^{\mathcal{B}'}|, \ldots, |\sigma_n^{\mathcal{B}'}|) \geq (|\sigma_1^{\mathcal{A}_j}|, \ldots, |\sigma_n^{\mathcal{A}_j}|)$ for some $1 \leq j \leq m$), and thus \mathcal{B}' satisfies δ and δ_V , the last formula given \mathcal{B} satisfies δ_V and \mathcal{B}' agrees with \mathcal{B} on the values assigned to the variables in δ_V . Finally, using the smoothness of \mathcal{T} and the fact that we are dealing with an empty signature, we can find a \mathcal{T} -interpretation \mathcal{C} that satisfies δ_V and δ whose underlying structure is isomorphic to that of \mathcal{A} (that is, $|\sigma_i^{\mathcal{C}}| = |\sigma_i^{\mathcal{A}}|$ for each $1 \leq i \leq n$); defining the \mathcal{T} -interpretation \mathcal{A}' , that differs from \mathcal{A} at most on the values assigned to \overrightarrow{x} , such that $(x_k^i)^{\mathcal{A}'}$ equals $(x_l^i)^{\mathcal{A}'}$ iff $(x_k^i)^{\mathcal{C}}$ equals $(x_l^i)^{\mathcal{C}}$, we get that \mathcal{A}' satisfies $\phi \wedge \delta$, and so \mathcal{A} satisfies $\exists \overrightarrow{x} . wit(\phi)$. Conversely, if \mathcal{A} satisfies $\exists \overrightarrow{x} . wit(\phi)$, again given the fact ϕ has none of the variables in \overrightarrow{x} , \mathcal{A} satisfies $\phi \wedge \exists \overrightarrow{x} . \delta$ and thus ϕ , proving ϕ and $\exists \overrightarrow{x} . wit(\phi)$ are \mathcal{T} -equivalent.

Now, let V be a set of variables with arrangement δ_V , and suppose the \mathcal{T} -interpretation \mathcal{A} satisfies $wit(\phi) \wedge \delta_V$. We define a new \mathcal{T} -interpretation \mathcal{B} by making: $\sigma_i^{\mathcal{B}} = vars_{\sigma_i}(wit(\phi) \wedge \delta_V)^{\mathcal{A}}$ (which has at least $m_{i,j}$ elements, $(x_1^i)^{\mathcal{A}}$ through $(x_{m_{i,j}}^i)^{\mathcal{A}}$, for every sort σ_i and j an index such that \mathcal{A} satisfies $\bigwedge_{i=1}^n \bigwedge_{1 \leq k < l \leq m_{i,j}} \neg (x_k^i = x_l^i)$, what makes of \mathcal{B} indeed a \mathcal{T} -interpretation, given this theory is smooth on an empty signature); $x^{\mathcal{B}} = x^{\mathcal{A}}$ for all variables x in $wit(\phi) \wedge \delta_V$; and arbitrarily for other variables. Then it is true that \mathcal{B} satisfies $wit(\phi) \wedge \delta_V$, and has $\sigma^{\mathcal{B}} = vars_{\sigma}(wit(\phi) \wedge \delta_V)^{\mathcal{B}}$ for each $\sigma \in S_{\Sigma}$.

G Proof of Theorem 7

Theorem 7. There are no Σ_1 or Σ_2 -theories \mathcal{T} that are, simultaneously, neither stably infinite nor stably finite, but are convex and have the finite model property, with respect to the entire set of their sorts.

Proof. If \mathcal{T} is a Σ_1 -theory, we just need to apply Theorem 12.

Assume then \mathcal{T} is a Σ_2 -theory that has the finite model property with respect to $\{\sigma, \sigma_2\}$, but is neither stably infinite nor stably finite with respect to that set: we will prove that then \mathcal{T} is not convex. Because \mathcal{T} is not stably infinite, Lemma 1 guarantees that there exists an $M \in \mathbb{N} \setminus \{0\}$ such that there are no \mathcal{T} -interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}|, |\sigma_2^{\mathcal{A}}| \geq M$.

If \mathcal{T} has no interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}| > M$ (the case with no interpretations where $|\sigma_2^{\mathcal{A}}| > M$ is analogous), because \mathcal{T} is not stably finite there must exist a \mathcal{T} -interpretation \mathcal{A}_0 with $|\sigma_2^{\mathcal{A}_0}| \ge \omega$ (and $|\sigma^{\mathcal{A}_0}| \le M$) such that there are no \mathcal{T} -interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}| \le |\sigma^{\mathcal{A}_0}|$ and $|\sigma_2^{\mathcal{A}}|$ finite. But, because \mathcal{T} has the finite model property, there must exist a \mathcal{T} -interpretation \mathcal{A}_1 with finite domains that satisfies $\neg(u = v)$, for u and v variables of sort σ_2 (since this formula is satisfied by \mathcal{A}_0): of course, then $|\sigma_2^{\mathcal{A}_1}| \ge 2$ and $|\sigma^{\mathcal{A}_1}| > |\sigma^{\mathcal{A}_0}| \ge 1$ (by the definition of \mathcal{A}_0). Therefore, $|\sigma^{\mathcal{A}_1}| \ge 2$, and we can prove \mathcal{T} is not convex: take variables x_1 through x_{M+1} of sort σ , and it is clear that $\vdash_{\mathcal{T}} \bigvee_{i=1}^{M} \bigvee_{j=i+1}^{M+1} x_i = x_j$, but we cannot have $\vdash_{\mathcal{T}} x_i = x_j$ for any pair $1 \le i < j \le M + 1$ since there is a \mathcal{T} - interpretation (\mathcal{A}_1) with at least two elements in its domain of sort σ .

So, for \mathcal{T} to be convex, we must have two \mathcal{T} -interpretations \mathcal{A}_0 and \mathcal{A}_1 with $|\sigma^{\mathcal{A}_0}| > M$ and $|\sigma_2^{\mathcal{A}_1}| > M$, but again we reach a contradiction, since then $|\sigma^{\mathcal{A}_0}| > 1$ and $|\sigma_2^{\mathcal{A}_1}| > 1$. Indeed, take variables x_1 through x_{M+1} of sort σ , and u_1 through u_{M+1} of sort σ_2 , and we have that

$$\vdash_{\mathcal{T}} \bigvee_{i=1}^{M} \bigvee_{j=i+1}^{M+1} x_i = x_j \lor \bigvee_{p=1}^{M} \bigvee_{q=p+1}^{M+1} u_p = u_q,$$

but neither $\vdash_{\mathcal{T}} x_i = x_j$ nor $\vdash_{\mathcal{T}} u_p = u_q$ for pairs $1 \le i < j \le M + 1$ or $1 \le p < q \le M + 1$: in the first case, because there exists a \mathcal{T} -interpretation with at least two elements in its domain of sort $\sigma(\mathcal{A}_0)$; in the second because there exists a \mathcal{T} -interpretation with at least two elements in its domain of sort $\sigma_2(\mathcal{A}_1)$.

H Proof of Theorem 8

Lemma 4 is proven in [12] and only restated here, mostly for context. In what follows, given a Σ_2 -interpretation \mathcal{A} , \mathcal{A}_1 is the Σ_1 -interpretation with $\sigma^{\mathcal{A}_1} = \sigma^{\mathcal{A}}$; and, given a quantifier-free Σ_2 -formula ϕ and a Σ_2 -interpretation \mathcal{A} , $\phi_{\mathcal{A}}$ is the Σ_1 -formula where we replace every atomic subformula u = v of ϕ with variables of sort σ_2 for a Σ_1 -tautology (such as x = x, for x of sort σ) if \mathcal{A} satisfies u = v, and otherwise for a Σ_1 -contradiction (such as $\neg(x = x)$).

Lemma 4. Take a Σ_2 -interpretation A. It is then true that, for any Σ_1 -formula φ , A_1 satisfies φ if, and only if, A satisfies φ .

Theorem 8. Let \mathcal{T} be a Σ_1 -theory. Then: \mathcal{T} is **FM**, or **SF**, w.r.t. $\{\sigma\}$ if and only if $(\mathcal{T})^2$ is, respectively, **FM**, or **SF** w.r.t. $\{\sigma, \sigma_2\}$.

- *Proof.* 1. If \mathcal{T} has the finite model property, given it is one-sorted, we may use Theorem 4 to prove \mathcal{T} is also stably finite. So we may prove that this, in turn, implies $(\mathcal{T})^2$ is stably finite, and therefore has the finite model property.
- 2. Suppose now *T* is stably finite, let φ be a quantifier-free Σ₂-theory, and let *A* be a (*T*)²-interpretation that satisfies φ. The facts *A*₁ satisfies the (quantifier-free) formula φ_A, and *T* is stably finite, imply there exists a *T*-interpretation *C* that satisfies φ_A with |σ^C| finite and satisfying |σ^C| ≤ |σ^{A₁}| = |σ^A|. We then define the (*T*)²-interpretation *B* with *B*₁ = *C* and: σ₂^B = vars_{σ2}(φ)^A (which is a finite set that satisfies |σ₂^B| ≤ |σ₄^A|); and x^B = x^A for every variable x ∈ vars_{σ2}(φ) (and arbitrarily otherwise). We then have that, for every atomic subformula x = y of φ with x and y of sort σ₂, *B* satisfies x = y iff *A* satisfies x = y, meaning φ_B = φ_A and, therefore, that *B* is a (*T*)²-interpretation that satisfies φ with |σ^B| ≤ |σ^A|, |σ₂^B| ≤ |σ₂^A|, and both finite.

- Assume now (*T*)² has the finite model property, let φ be a quantifier-free Σ₁-formula, and C be a *T*-interpretation that satisfies φ. Then A such that A₁ = C and |σ₂^A| = 1 satisfies φ, meaning there exists a *T*-interpretation B that satisfies φ with both |σ^B| and |σ₂^B| finite. Then B₁ is a *T*-interpretation that satisfies φ, with |σ^{B₁}| = |σ^B| finite.
- 2. Assume now $(\mathcal{T})^2$ is stably finite, and thus has the finite model property: as we proved above, this implies \mathcal{T} has the finite model property; but, since \mathcal{T} is one sorted, Theorem 4 tells us that \mathcal{T} must also be stably finite.

I Proof of Theorem 9

Lemma 5 was proven in [12], and we restate it here as it defines some of the symbols we use.

Lemma 5. Let Σ_n be the empty signature with n sorts $S = \{\sigma_1, \ldots, \sigma_n\}$, and Σ_s^n the signature with sorts S and only one function symbol s of arity $\sigma_1 \rightarrow \sigma_1$. Given a Σ_n -interpretation C, we define a Σ_s^n -interpretation s(C) by making:

1. $\sigma^{s(C)} = \sigma^{C}$ for each $\sigma \in S$; 2. $s^{s(C)}(a) = a$ for all $a \in \sigma_{1}^{C}$; 3. and $x^{s(C)} = x^{C}$ for every variable x.

Reciprocally, given any Σ_s^n -interpretation \mathcal{A} , we may consider the Σ_n -interpretation $\mathcal{O}(\mathcal{A})$ with:

1. $\sigma^{\Theta(\mathcal{A})} = \sigma^{\mathcal{A}}$ for each $\sigma \in S$;

2. and $x^{\mathcal{O}(\mathcal{A})} = x^{\mathcal{A}}$, for every variable x.

Finally, given a Σ_s^n -formula φ , we repeatedly replace each occurrence of s(x) in φ by x until we obtain a Σ_n -formula φ_0 . Then, it is true that a Σ_s^n -interpretation \mathcal{A} that satisfies $\forall x. [s(x) = x]$ (where x is of sort σ_1) also satisfies φ iff $\theta(\mathcal{A})$ satisfies φ_0 ; of course, given that for any Σ_n -interpretation \mathcal{C} , $\theta(s(\mathcal{C})) = \mathcal{C}$, \mathcal{C} satisfies a Σ_n -formula φ iff $s(\mathcal{C})$ satisfies φ (since $\varphi_0 = \varphi$).

Theorem 9. If \mathcal{T} is a theory over an empty signature Σ_n with sorts $S = \{\sigma_1, \ldots, \sigma_n\}$, then: \mathcal{T} is **FM**, or **SF**, w.r.t. S if and only if $(\mathcal{T})_s$ is, respectively, **FM**, or **SF**, w.r.t. S.

- *Proof.* 1. If \mathcal{T} has the finite model property, let us take a quantifier-free Σ_n^s -formula ϕ , and a $(\mathcal{T})_s$ -interpretation \mathcal{A} that satisfies ϕ . We then have that $\mathcal{O}(\mathcal{A})$ satisfies ϕ_0 , and so there exists a \mathcal{T} -interpretation \mathcal{C} , with $|\sigma^{\mathcal{A}}|$ finite for each $\sigma \in S$, that satisfies ϕ_0 . Of course, then $s(\mathcal{A})$ is a $(\mathcal{T})_s$ -interpretation, finite in every one of its domains, that satisfies ϕ .
- Suppose *T* is stably finite, let φ be a quantifier-free Σ_n^s-formula, and *A* a (*T*)_s-interpretation that satisfies φ, meaning Ø(*A*) satisfies φ₀. Because φ₀ is quantifier-free, and *T* is stably finite, there exists a *T*-interpretation *C* that satisfies φ₀ with |σ^C| ≤ |σ^{0(A)}| = |σ^A| finite for each σ ∈ S. And then s(*C*) is a (*T*)_s-interpretation that satisfies φ (since Ø(s(*C*)) = *C*), with |σ^{s(C)}| = |σ^C| ≤ |σ^A| finite for each σ ∈ S.
- Suppose (*T*)_s has the finite model property, let φ be a quantifier-free Σ_n-formula, and C a *T*-interpretation that satisfies φ. Then s(C) satisfies φ, and since (*T*)_s has the finite model property there exists a (*T*)_s-interpretation *A* that satisfies φ with |σ^A| finite for each σ ∈ S. Of course, Ø(A) is then a *T*-interpretation that satisfies φ with |σ^{Q(A)}| = |σ^A| finite for each σ ∈ S.
- Assume now (*T*)_s is stably finite, let φ be a quantifier-free Σ_n-formula, and C be a *T*-interpretation that satisfies φ: s(C) is then a (*T*)_s-interpretation that satisfies φ, and there must then exist a (*T*)_s-interpretation *A*, with |σ^A| finite and |σ^A| ≤ |σ^{s(C)}| for each σ ∈ S, that satisfies φ; of course, Ø(A) is then a *T*-interpretation that satisfies φ with |σ^{θ(A)}| finite and |σ^{θ(A)}| ≤ |σ^C| for each σ ∈ S.

J Proof of Theorem 10

We will use some of the symbols found in Lemma 5. Moreover, given a Σ_s^n formula ϕ and a Σ_s^n -interpretation \mathcal{A} that satisfies ϕ , we define the auxiliary formula $\phi_{\mathcal{A}}^{\dagger}$. To do that, let: $vars_{\sigma_1}(\phi) = \{z_1, \ldots, z_m\}$; M_i be the maximum of j such that $s^j(z_i)$ appears in ϕ ; and $V = \{y_{i,j} : 1 \le i \le m, 0 \le j \le M_i + 2\}$ be a set of fresh variables of sort σ_1 .

- 1. ϕ^{\dagger} is defined by replacing each atomic subformula $s^{j}(z_{i}) = s^{q}(z_{p})$ of ϕ by $y_{i,j} = y_{p,q}$;
- 2. \mathcal{A}' is the \mathcal{T} -interpretation that differs from \mathcal{C} at most on V, where $y_{i,j}^{\mathcal{A}'} = (s^{\mathcal{A}})^j (z_i^{\mathcal{A}})$;
- 3. δ_V is the arrangement induced on V by making x related to y iff $x^{\mathcal{A}'} = y^{\mathcal{A}'}$.

We then make $\phi^{\dagger}_{A} = \phi^{\dagger} \wedge \delta_{V}$.

Theorem 10. If \mathcal{T} is a theory over an empty signature Σ_n with sorts $S = \{\sigma_1, \ldots, \sigma_n\}$, then: \mathcal{T} is **FM**, or **SF**, w.r.t. S if and only if $(\mathcal{T})_{\vee}$ is, respectively, **FM**, or **SF**, w.r.t. S.

Proof. 1. Suppose \mathcal{T} has the finite model property, let ϕ be a quantifier-free Σ_s^n -formula, and \mathcal{A} be a $(\mathcal{T})_{\vee}$ -interpretation that satisfies ϕ : there is then an interpretation \mathcal{A}' differing from \mathcal{A} at most on the value assigned to variables (thus making of \mathcal{A}' a $(\mathcal{T})_{\vee}$ -interpretation) that satisfies the Σ_n formula $\phi_{\mathcal{A}}^{\dagger}$. Then $\mathcal{O}(\mathcal{A}')$ is a \mathcal{T} -interpretation that satisfies $\phi_{\mathcal{A}}^{\dagger}$ and, by assumption on \mathcal{T} , there exists a \mathcal{T} -interpretation \mathcal{C} that satisfies $\phi_{\mathcal{A}}^{\dagger}$ with $\sigma_k^{\mathcal{C}}$ finite for each $1 \leq k \leq n$.

Now, we define an interpretation \mathcal{B} . First, we make $\sigma_k^{\mathcal{B}} = \sigma_k^{\mathcal{C}}$ for all $1 \le k \le n$, so \mathcal{B} has all domains finite. Then, $s^{\mathcal{B}}(y_{i,j}^{\mathcal{C}}) = y_{i,j+1}^{\mathcal{C}}$ for each $1 \le i \le m$ and $0 \le j \le M_i + 1$ (notice that that also defines $s^{\mathcal{B}}(y_{i,M_i+2}^{\mathcal{C}})$, since y_{i,M_i+2} must equal either y_{i,M_i} or y_{i,M_i+1} in \mathcal{C}), and $s^{\mathcal{B}}(a) = a$ for each $a \in \sigma_1^{\mathcal{C}} \setminus vars_{\sigma_1}(\phi_A^{\dagger})^{\mathcal{C}}$ (so that \mathcal{B} always satisfies ψ_{\vee} , given \mathcal{A} satisfies that formula and therefore $y_{i,j+2}$ equals either $y_{i,j}$ or $y_{i,j+1}$ in \mathcal{A}' , and thus in \mathcal{C}): this way, \mathcal{B} is then a \mathcal{T} -interpretation that satisfies ψ_{\vee} , or in other words, a $(\mathcal{T})_{\vee}$ -interpretation. Finally, we make $z_i = y_{i,0}^{\mathcal{C}}$, and arbitrary for other variables: this way \mathcal{B} satisfies ϕ , and we have proven $(\mathcal{T})_{\vee}$ has the finite model property.

2. The proof is the same as the one above if \mathcal{T} is stably finite: let ϕ be a quantifier-free Σ_s^n -formula, \mathcal{A} be a $(\mathcal{T})_{\vee}$ -interpretation that satisfies ϕ , and \mathcal{A}' be a $(\mathcal{T})_{\vee}$ -interpretation differing from \mathcal{A} at most on the value assigned to variables that satisfies the Σ_n formula ϕ_A^{\dagger} .

 $\mathcal{O}(\mathcal{A}')$ is then a \mathcal{T} -interpretation that satisfies $\phi_{\mathcal{A}}^{\dagger}$ and, by the fact \mathcal{T} is assumed to be stably finite, there exists a \mathcal{T} -interpretation \mathcal{C} that satisfies $\phi_{\mathcal{A}}^{\dagger}$ with all domains finite, and $|\sigma_{k}^{\mathcal{C}}| \leq |\sigma_{k}^{\mathcal{O}(\mathcal{A}')}| = |\sigma_{k}^{\mathcal{A}}|$ for each $1 \leq k \leq n$. We then carefully define an interpretation \mathcal{B} with the same domains as \mathcal{C} that turns out to be a $(\mathcal{T})_{\vee}$ -interpretation that satisfies ϕ , and since $|\sigma_{k}^{\mathcal{B}}| = |\sigma_{k}^{\mathcal{C}}| \leq |\sigma_{k}^{\mathcal{A}}|$ we are done.

Now we deal with the cases where $(\mathcal{T})_{\vee}$ has the desired properties, which is much easier.

- If (*T*)_∨ has the finite model property, let φ be a Σ_n-formula and C be a *T*-interpretation that satisfies φ: s(C) is then a (*T*)_∨-interpretation that satisfies φ by Lemma 5; so there must exist a (*T*)_∨-interpretation *A*, with all domains finite, that satisfies φ. Again by Lemma 5, we have Ø(*A*) is a *T*-interpretation that satisfies φ, and its domains are finite, so we are done.
- Finally, suppose (*T*)_∨ is stably finite, and let φ be a Σ_n-formula and C be a *T*-interpretation that satisfies φ. By Lemma 5, s(C) is a (*T*)_∨-interpretation that satisfies φ, and given (*T*)_∨ is stably finite, there must exist a (*T*)_∨-interpretation A, with all domains finite, that satisfies φ and |σ_k^A| ≤ |σ_k^{s(C)}| = |σ_k^C| for each 1 ≤ k ≤ n. Using Lemma 5, Ø(A) is a *T*-interpretation that satisfies φ, its domains are finite, and |σ_k^{Ø(A)}| = |σ_k^A| ≤ |σ_k^C| ≤ |σ_k^C|

K Proofs for Theories from Figure 5

For the theories of Figure 5, most properties were proven in [12]. Here we only consider the new properties: stable finiteness and the finite model property.

K.1 $\mathcal{T}_{\geq n}$

Lemma 6. The Σ_1 -theory $\mathcal{T}_{\geq n}$ is stably finite, and thus has the finite model property, with respect to its only sort, for every $n \in \mathbb{N} \setminus \{0\}$.

Proof. From [12], $\mathcal{T}_{\geq n}$ is strongly finitely witnessable, and from Theorem 3 we get that $\mathcal{T}_{\geq n}$ is stably finite, and thus has the finite model property by Theorem 1.

K.2 $\mathcal{T}^{\infty}_{even}$

Lemma 7. The Σ_1 -theory $\mathcal{T}_{even}^{\infty}$ is stably finite, and thus has the finite model property, with respect to its only sort.

Proof. From [12], $\mathcal{T}_{even}^{\infty}$ is finitely witnessable, and from Theorem 2 we get that $\mathcal{T}_{even}^{\infty}$ has the finite model property; by Theorem 4, and the fact $\mathcal{T}_{even}^{\infty}$ is a Σ_1 -theory, we get it is also stably finite.

K.3 T_{∞}

Lemma 8. The Σ_1 -theory \mathcal{T}_{∞} does not have the finite model property, and thus is not stably finite, with respect to $\{\sigma\}$.

Proof. Obvious, since \mathcal{T}_{∞} has models, but none of them are finite.

K.4 $\mathcal{T}_{n,\infty}$

Lemma 9. The Σ_1 -theory $\mathcal{T}_{n,\infty}$ does not have the finite model property, and thus is not stably finite, with respect to $\{\sigma\}$.

Proof. Consider the formula $\bigwedge_{1 \le i < j \le n+1} \neg (x_i = x_j)$. it is satisfied by any infinite model of $\mathcal{T}_{n,\infty}$, but it cannot be satisfied by any finite $\mathcal{T}_{n,\infty}$ -interpretation \mathcal{A} , since necessarily $|\sigma^{\mathcal{A}}| = n$.

K.5 $\mathcal{T}_{< n}$

Lemma 10. The Σ_1 -theory $\mathcal{T}_{< n}$ is stably finite, and thus has the finite model property, with respect to $\{\sigma\}$.

Proof. Let ϕ be a quantifier-free Σ_1 -formula, and \mathcal{A} a $\mathcal{T}_{\leq n}$ -interpretation that satisfies ϕ : then \mathcal{A} is already a finite $\mathcal{T}_{\leq n}$ -interpretation that satisfies ϕ with $|\sigma^{\mathcal{A}}| \leq |\sigma^{\mathcal{A}}|$.

K.6 $\mathcal{T}_{m,n}$

Lemma 11. The Σ_1 -theory $\mathcal{T}_{m,n}$ is stably finite, and thus has the finite model property, with respect to $\{\sigma\}$.

Proof. Let ϕ be a quantifier-free Σ_1 -formula, and \mathcal{A} a $\mathcal{T}_{m,n}$ -interpretation that satisfies ϕ : of course \mathcal{A} is already a finite $\mathcal{T}_{m,n}$ -interpretation that satisfies ϕ .

K.7 $T_{2,3}$

Lemma 12. The Σ_2 -theory $\mathcal{T}_{2,3}$ has the finite model property with respect to $\{\sigma\}$.

Proof. From [10], $\mathcal{T}_{2,3}$ is finitely witnessable; and from Theorem 2, we get that it has also the finite model property.

Lemma 13. The Σ_2 -theory $\mathcal{T}_{2,3}$ is not stably finite with respect to $\{\sigma\}$.

Proof. From [10], $\mathcal{T}_{2,3}$ is smooth but not strongly finitely witnessable with respect to $\{\sigma, \sigma_2\}$: since the signature is empty, by Theorem 6, we conclude it cannot be stably finite.

K.8 \mathcal{T}_2^{∞}

Lemma 14. The Σ_2 -theory \mathcal{T}_2^{∞} does not have the finite model property, and thus is not stably finite, with respect to $\{\sigma, \sigma_2\}$.

Proof. Obvious, given \mathcal{T}_2^{∞} is not contradictory, but also has no interpretations \mathcal{A} with $|\sigma_2^{\mathcal{A}}|$ finite.

K.9 \mathcal{T}_1^{odd}

Lemma 15. The Σ_2 -theory \mathcal{T}_1^{odd} is stably finite, and thus has the finite model property, with respect to $\{\sigma, \sigma_2\}$.

Proof. Take a quantifier-free Σ_2 -formula ϕ and a \mathcal{T}_1^{odd} -interpretation \mathcal{A} that satisfies ϕ : then, let $n = |vars_{\sigma_2}(\phi)^{\mathcal{A}}|$, and N = n if n is odd, and N = n + 1 if n is even. Take a set A with N - n elements disjoint from the domains of \mathcal{A} (notice A may be empty), and define the \mathcal{T}_1^{odd} -interpretation \mathcal{B} such that: $\sigma^{\mathcal{B}} = \sigma^{\mathcal{A}}$ (and so $|\sigma^{\mathcal{B}}| = 1$); $\sigma^{\mathcal{B}} = vars_{\sigma_2}(\phi)^{\mathcal{A}} \cup A$ (which has necessarily N elements, what makes of \mathcal{B} indeed a \mathcal{T}_1^{odd} -interpretation); $x^{\mathcal{B}} = x^{\mathcal{A}}$ for all variables x in ϕ ; and arbitrarily otherwise. Obviously \mathcal{B} satisfies ϕ , is finite in both domains and has the property that $|\sigma^{\mathcal{B}}| \leq |\sigma^{\mathcal{A}}|$ and $|\sigma^{\mathcal{B}}_2| \leq |\sigma^{\mathcal{A}}_2|$ (notice that, even if N = n + 1, since $|\sigma^{\mathcal{A}}_2|$ must be an odd, or infinite, number we have $|\sigma^{\mathcal{A}}_2| \geq N$).

K.10 \mathcal{T}_1^{∞}

Lemma 16. The Σ_2 -theory \mathcal{T}_1^{∞} does not have the finite model property, and thus is not stably finite, with respect to $\{\sigma, \sigma_2\}$.

Proof. Obvious, since \mathcal{T}_1^{∞} has models, and yet has no models \mathcal{A} where $\sigma_2^{\mathcal{A}}$ is finite.

K.11 T_f

Lemma 17. The Σ_s -theory \mathcal{T}_f is stably finite, and thus has the finite model property, with respect to $\{\sigma\}$.

Proof. \mathcal{T}_f is finitely witnessable with respect to $\{\sigma\}$ (proven in [12]), and thus by Theorem 2 \mathcal{T}_f has the finite model property; and, since Σ_s is a one-sorted signature, by Theorem 4 we obtain \mathcal{T}_f is also stably finite.

K.12 \mathcal{T}_f^s

Lemma 18. The Σ_s -theory \mathcal{T}_{odd}^{\neq} is stably finite, and thus has the finite model property, with respect to $\{\sigma\}$.

Proof. Because of Theorem 4, we just need to prove \mathcal{T}_f^s has the finite model property; and because of Theorem 2, we just use the fact that \mathcal{T}_f^s is finitely witnessable, as proved in [12].

K.13 $\mathcal{T}_{2,\infty}^{\neq}$

Lemma 19. The Σ_s -theory $\mathcal{T}_{2,\infty}^{\neq}$ does not have the finite model property, and thus is not stably finite, with respect to $\{\sigma\}$.

Proof. There is a $\mathcal{T}_{2,\infty}^{\neq}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = \omega$ (write, for simplicity, $\sigma^{\mathcal{A}} = \{a_i \mid i \in \mathbb{N}\}$), and $s^{\mathcal{A}}(a_i) = a_{i+1}$ for all $i \in \mathbb{N}$ (and any assignment of the variables) that satisfies $\neg(s(x) = x)$. However, the only finite $\mathcal{T}_{2,\infty}^{\neq}$ -interpretations satisfy $\forall x. (s(x) = x)$, and thus cannot satisfy $\neg(s(x) = x)$.

K.14 \mathcal{T}_{odd}^{\neq}

Lemma 20. The Σ_s -theory \mathcal{T}_{odd}^{\neq} is stably finite, and thus has the finite model property, with respect to $\{\sigma\}$.

Proof. Because the signature is one-sorted, according to Theorem 4 we only need to show the theory has the finite model property. So take a quantifier-free Σ_s -formula ϕ and a \mathcal{T}_{odd}^{\neq} -interpretation \mathcal{A} that satisfies ϕ : we can assume \mathcal{A} has at least 3 elements since, otherwise, ϕ would already be known to be satisfied by a finite \mathcal{T}_{odd}^{\neq} -interpretation. Take as well the variables x_1 through x_n in ϕ , and the maximum M_i , for each i, of the values j such that $s^j(x_i)$ appears in ϕ , and define

$$m = |\{s^{j}(x_{i}) : 1 \le i \le n, 0 \le j \le M_{i} + 1\}^{\mathcal{A}}|,$$

and there are two cases to consider: if m is odd, make M = m + 2, and if m is even, make M = m + 1; take as well a set A with M - m elements disjoint from σ^{A} . We then define an interpretation \mathcal{B} by making:

$$\sigma^{\mathcal{B}} = \{s^j(x_i) : 1 \le i \le n, 0 \le j \le M_i + 1\}^{\mathcal{A}} \cup A;$$

 $s^{\mathcal{B}}(a) = s^{\mathcal{A}}(a)$ for all $a \in \{s^{j}(x_{i}) : 1 \leq i \leq n, 0 \leq j \leq M_{i}\}^{\mathcal{A}}$, and $s^{\mathcal{B}}(a)$ any value in A different from a (of which there is one, since A has at least 2 elements) for each $a \in A$ or $a = (s^{\mathcal{A}})^{M_{i}+1}(x_{i}^{\mathcal{A}})$ not in $\{s^{j}(x_{i}) : 1 \leq i \leq n, 0 \leq j \leq M_{i}\}^{\mathcal{A}}$; $x^{\mathcal{B}} = x^{\mathcal{A}}$ for each variable x of ϕ , and arbitrarily otherwise. \mathcal{B} has then an odd number of elements (thus finite) greater or equal than 3, and satisfies $\neg(s(x) = x)$ for all x, meaning it is a \mathcal{T}_{odd}^{\neq} -interpretation; furthermore, it satisfies ϕ , and so we are done.

K.15 $\mathcal{T}_{1,\infty}^{\neq}$

Lemma 21. The Σ_s -theory $\mathcal{T}_{1,\infty}^{\neq}$ does not have the finite model property, and thus is not stably finite, with respect to $\{\sigma\}$.

Proof. Take a variable x and consider the quantifier-free formula $\neg(s(x) = x)$, satisfied by the infinite $\mathcal{T}_{1,\infty}^{\neq}$ -interpretations: because the only finite $\mathcal{T}_{1,\infty}^{\neq}$ -interpretation has only one element, it must satisfy s(x) = x instead, and so this theory cannot have the finite model property.

L Proofs for Theories from Figure 7

L.1 \mathcal{T}^{∞}

Lemma 22. The Σ_2 -theory \mathcal{T}^{∞} is stably infinite with respect to $\{\sigma, \sigma_2\}$.

Proof. Let ϕ be a quantifier-free Σ_2 -formula, and \mathcal{A} a \mathcal{T}^{∞} -interpretation that satisfies ϕ . Given two infinite sets A and B, disjoint from each other and $\sigma^{\mathcal{A}}$ and $\sigma_2^{\mathcal{A}}$, we define a \mathcal{T}^{∞} -interpretation \mathcal{B} by making: $\sigma^{\mathcal{B}} = \sigma^{\mathcal{A}} \cup A$; $\sigma_2^{\mathcal{B}} = \sigma_2^{\mathcal{A}} \cup B$; $x^{\mathcal{B}} = x^{\mathcal{A}}$ for every variable x of sort σ ; and $u^{\mathcal{B}} = u^{\mathcal{A}}$ for every variable u of sort σ_2 . Then it is true that \mathcal{B} satisfies ϕ , and both $\sigma^{\mathcal{B}}$ and $\sigma_2^{\mathcal{B}}$ are infinite.

Lemma 23. The Σ_2 -theory \mathcal{T}^{∞} is not smooth with respect to $\{\sigma, \sigma_2\}$.

Proof. Let x be a variable of sort σ , and it is clear that x = x is satisfied by a \mathcal{T}^{∞} -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = |\sigma_2^{\mathcal{A}}| = 2$. But there are no \mathcal{T}^{∞} -interpretations \mathcal{B} with $|\sigma^{\mathcal{B}}| = 3$ and $|\sigma_2^{\mathcal{B}}| = 2$, so \mathcal{T}^{∞} cannot be smooth.

Lemma 24. The Σ_2 -theory \mathcal{T}^{∞} is not stably finite with respect to $\{\sigma, \sigma_2\}$.

Proof. Let x be a variable of sort σ , and \mathcal{A} be a \mathcal{T}^{∞} -interpretation with $|\sigma^{\mathcal{A}}| = 1$ and $|\sigma_2^{\mathcal{A}}| = \omega$: it is clear that \mathcal{A} satisfies x = x. But there are no \mathcal{T}^{∞} -interpretations \mathcal{B} with $|\sigma^{\mathcal{B}}| = 1$ and $|\sigma_2^{\mathcal{B}}|$ finite, so \mathcal{T}^{∞} cannot be stably finite.

Lemma 25. The Σ_2 -theory \mathcal{T}^{∞} is finitely witnessable with respect to $\{\sigma, \sigma_2\}$.

Proof. Given a quantifier-free Σ_2 -formula ϕ , consider the witness

wit
$$(\phi) = \phi \land \bigwedge_{i=1}^m x_i = x_i \land \bigwedge_{i=1}^m u_i = u_i,$$

where

$$m = \max\{|vars_{\sigma}(\phi)^{\mathcal{A}}|, |vars_{\sigma_2}(\phi)^{\mathcal{A}}|, 2\},\$$

 x_1 through x_m are fresh variables of sort σ , and u_1 through u_m are fresh variables of sort σ_2 . Given $\bigwedge_{i=1}^m x_i = x_i \land \bigwedge_{i=1}^m u_i = u_i$ is a tautology, and ϕ possesses none of the variables in $\vec{x} = vars(wit(\phi)) \setminus vars(\phi)$, we have that ϕ and $\exists \vec{x} . wit(\phi)$ are \mathcal{T}^{∞} -equivalent.

Now, assume \mathcal{A} is a \mathcal{T}^{∞} -interpretation that satisfies $wit(\phi)$. Take sets A and B, disjoint from each other and $\sigma^{\mathcal{A}}$ and $\sigma^{\mathcal{A}}_2$, with $m - |vars_{\sigma}(\phi)^{\mathcal{A}}|$ and $m - |vars_{\sigma_2}(\phi)^{\mathcal{A}}|$ elements, respectively. We define the \mathcal{T}^{∞} -interpretation \mathcal{B} by making: $\sigma^{\mathcal{B}} = vars_{\sigma}(\phi)^{\mathcal{A}} \cup A$; $\sigma^{\mathcal{B}}_2 = vars_{\sigma_2}(\phi)^{\mathcal{A}} \cup B$; $x^{\mathcal{B}} = x^{\mathcal{A}}$ for $x \in vars_{\sigma}(\phi)$, and $x_i \in \{x_i : 1 \le i \le m\} \mapsto x_i^{\mathcal{B}} \in \sigma^{\mathcal{B}}$ a surjection; and $u^{\mathcal{B}} = u^{\mathcal{A}}$ for $u \in vars_{\sigma_2}(\phi)$, and $u_i \in \{u_i : 1 \le i \le m\} \mapsto u_i^{\mathcal{B}} \in \sigma^{\mathcal{B}}_2$ also a surjection (and arbitrary for other variables). This way, \mathcal{B} satisfies ϕ , and in addition has $\sigma^{\mathcal{B}} = vars_{\sigma}(wit(\phi))^{\mathcal{B}}$.

Lemma 26. The Σ_2 -theory \mathcal{T}^{∞} has the finite model property with respect to $\{\sigma, \sigma_2\}$.

Proof. Follows from Theorem 2 and Lemma 25.

Lemma 27. The Σ_2 -theory \mathcal{T}^{∞} is not strongly finitely witnessable with respect to $\{\sigma, \sigma_2\}$.

Proof. Follows from Theorem 3 and Lemma 24.

Lemma 28. The Σ_2 -theory \mathcal{T}^{∞} is convex with respect to $\{\sigma, \sigma_2\}$.

Proof. Since \mathcal{T}^{∞} is stably infinite, and Σ_2 is an empty signature, we simply apply Theorem 11.

L.2 $\mathcal{T}_{m,n}^{\infty}$

Lemma 29. The Σ_2 -theory $\mathcal{T}_{m,n}^{\infty}$ is not stably infinite, and thus not smooth, with respect to $\{\sigma, \sigma_2\}$ for any $m, n \in \mathbb{N}$.

Proof. Obvious, since $\mathcal{T}_{m,n}^{\infty}$ is not contradictory, but it does not have any interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}|$ infinite. \Box

Lemma 30. The Σ_2 -theory $\mathcal{T}_{m,n}^{\infty}$ is not stably finite with respect to $\{\sigma, \sigma_2\}$ for any $m, n \in \mathbb{N}$ such that $m \neq n$.

Proof. Take a variable x of sort σ , and we know x = x is satisfied by any $\mathcal{T}_{m,n}^{\infty}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = \min\{m, n\}$ (and, necessarily, $|\sigma_2^{\mathcal{A}}| \ge \omega$): but, since there are no $\mathcal{T}_{m,n}^{\infty}$ -interpretations \mathcal{B} with $|\sigma^{\mathcal{B}}| \le \min\{m, n\}$ and $|\sigma_2^{\mathcal{B}}|$ finite, $\mathcal{T}_{m,n}^{\infty}$ cannot be stably finite.

Lemma 31. The Σ_2 -theory $\mathcal{T}_{m,n}^{\infty}$ is finitely witnessable with respect to $\{\sigma, \sigma_2\}$ for any $m, n \in \mathbb{N}$.

Proof. Let ϕ be a quantifier-free Σ_2 -formula, and consider the witness

$$wit(\phi) = \phi \wedge \bigwedge_{i=1}^{\max\{m,n\}} x_i = x_i,$$

where x_1 through $x_{\max\{m,n\}}$ are fresh variables of sort σ ; this set of variables will be denoted by \overrightarrow{x} , and it equals $vars(wit(\phi)) \setminus vars(\phi)$. Suppose first that the $\mathcal{T}_{m,n}^{\infty}$ -interpretation \mathcal{A} satisfies ϕ : because $\bigwedge_{i=1}^{\max\{m,n\}} x_i = x_i$ is a tautology, \mathcal{A} also satisfies that formula, and so \mathcal{A} satisfies $wit(\phi)$ and thus $\exists \overrightarrow{x} . wit(\phi)$. The reciprocal is also true because ϕ has none of the variables in \overrightarrow{x} , and so ϕ and $\exists \overrightarrow{x} . wit(\phi)$ are $\mathcal{T}_{m,n}^{\infty}$ -equivalent.

Now, assume \mathcal{A} is a $\mathcal{T}_{m,n}^{\infty}$ -interpretation that satisfies $wit(\phi)$; take a set A with $\max\{m,n\} - |\sigma^{\mathcal{A}}|$ elements, and disjoint from $\sigma^{\mathcal{A}}$ and $\sigma_2^{\mathcal{A}}$. We define a $\mathcal{T}_{m,n}^{\infty}$ -interpretation \mathcal{B} by making: $\sigma^{\mathcal{B}} = \sigma^{\mathcal{A}} \cup A$; $\sigma_2^{\mathcal{B}} = vars_{\sigma_2}(\phi)^{\mathcal{A}}$; $x^{\mathcal{B}} = x^{\mathcal{A}}$ and $u^{\mathcal{B}} = u^{\mathcal{A}}$ for all variables x and u, of sorts σ and σ_2 respectively, in ϕ ; $x_i \mapsto x_i^{\mathcal{B}}$ any surjective map, for x_i in \overrightarrow{x} (and arbitrarily otherwise). Then it is true that \mathcal{B} satisfies $wit(\phi)$ (since it satisfies ϕ), and $\sigma^{\mathcal{B}} = vars_{\sigma}(wit(\phi))^{\mathcal{B}}$ and $\sigma_2^{\mathcal{B}} = vars_{\sigma_2}(wit(\phi))^{\mathcal{B}}$.

Lemma 32. The Σ_2 -theory $\mathcal{T}_{m,n}^{\infty}$ has the finite model property with respect to $\{\sigma, \sigma_2\}$ for any $m, n \in \mathbb{N}$.

Proof. Follows from Theorem 2 and Lemma 31.

Lemma 33. The Σ_2 -theory $\mathcal{T}_{m,n}^{\infty}$ is not strongly finitely witnessable with respect to $\{\sigma, \sigma_2\}$ for any $m, n \in \mathbb{N}$ such that $m \neq n$.

Proof. Follows from Theorem 3 and Lemma 30.

Lemma 34. The Σ_2 -theory $\mathcal{T}_{m,n}^{\infty}$ is not convex with respect to $\{\sigma, \sigma_2\}$ if $\max\{m, n\} > 1$.

Proof. Let $M = \max\{m, n\} > 1$ and take M + 1 variables x_1 trough x_{M+1} , of sort σ : it is then true that $\vdash_{\mathcal{T}_{m,n}^{\infty}} \bigvee_{i=1}^{M} \bigvee_{j=i+1}^{M} x_i = x_j$, by the pigeonhole principle, but we cannot get $\vdash_{\mathcal{T}_{m,n}^{\infty}} x_i = x_j$ for any $1 \le i < j \le M + 1$ since there is a $\mathcal{T}_{m,n}^{\infty}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = M > 1$.

L.3 $\mathcal{T}^{\infty}_{\neq}$

Lemma 35. The Σ_s^2 -theory $\mathcal{T}_{\neq}^{\infty}$ is not stably infinite, and thus not smooth, with respect to $\{\sigma, \sigma_2\}$.

Proof. s(x) = x is satisfied by the $\mathcal{T}_{\neq}^{\infty}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = 1$ and $|\sigma_{2}^{\mathcal{A}}| = \omega$, but all $\mathcal{T}_{\neq}^{\infty}$ -interpretations \mathcal{B} with $|\sigma^{\mathcal{A}}| \geq 2$ satisfy instead $\forall x. \neg(s(x) = x)$.

Lemma 36. The Σ_s^2 -theory $\mathcal{T}_{\neq}^{\infty}$ is finitely witnessable with respect to $\{\sigma, \sigma_2\}$.

Proof. The proof is similar to the one of Lemma 25, some care being necessary to give witnesses to the function s; see the proof that \mathcal{T}_{odd}^{\neq} is finitely witnessable in [12] for how this can be done.

Lemma 37. The Σ_s^2 -theory $\mathcal{T}_{\neq}^{\infty}$ is not stably finite with respect to $\{\sigma, \sigma_2\}$.

Proof. Obvious, since $\mathcal{T}_{\neq}^{\infty}$ has an interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = 1$ and $|\sigma_2^{\mathcal{A}}| = \omega$, while there are no $\mathcal{T}_{\neq}^{\infty-}$ interpretations \mathcal{B} with $|\sigma^{\mathcal{B}}| = 1$ and $\sigma_2^{\mathcal{B}}$ finite.

Lemma 38. The Σ_s^2 -theory $\mathcal{T}_{\neq}^{\infty}$ has the finite model property with respect to $\{\sigma, \sigma_2\}$.

Proof. Follows from Lemma 36 and Theorem 2.

Lemma 39. The Σ_s^2 -theory $\mathcal{T}_{\neq}^{\infty}$ is not strongly finitely witnessable with respect to $\{\sigma, \sigma_2\}$.

Proof. Follows from Theorem 3 and Lemma 37.

Lemma 40. Let \mathcal{T} be a stably infinite Σ_n -theory (the sorts of Σ_n are $S = \{\sigma_1, \ldots, \sigma_n\}$) with no interpretations \mathcal{A} such that $|\sigma_1^{\mathcal{A}}| = 1$: \mathcal{T} is necessarily convex because of Theorem 11.

Then the Σ_s^n -theory \mathcal{T}_{\neg} , where Σ_s^n is the signature with sorts S and one function symbol $s : \sigma_1 \to \sigma_1$, and \mathcal{T}_{\neg} is axiomatized by $Ax(\mathcal{T}) \cup \{ \forall x. \neg (s(x) = x) \}$, is convex with respect to its only sort.

Proof. Given a Σ_s^n -cube ϕ , we change the name of every variable x of sort σ_1 in ϕ to x_0 , and replace each occurrence of $s^{i+1}(x)$ in ϕ by a new variable x_{i+1} , adding the literal $\neg(x_{i+1} = x_i)$ to ϕ : the result is the Σ_n -cube ϕ' .

It is clear that, if ϕ is \mathcal{T}_{\neg} -satisfiable, ϕ' is \mathcal{T} -satisfiable: indeed, if \mathcal{A} satisfies ϕ , take the Σ_n -interpretation $\mathcal{O}(\mathcal{A})$ with $\sigma_i^{\emptyset(\mathcal{A})} = \sigma_i^{\mathcal{A}}$ for each $1 \leq i \leq n$; by changing the value of $x_i^{\emptyset(\mathcal{A})}$ (for x of sort σ_1) so that it equals $(s^{\mathcal{A}})^i(x^{\mathcal{A}})$, the resulting \mathcal{T} -interpretation \mathcal{C} then satisfies ϕ' .

Conversely, if the \mathcal{T} -interpretation \mathcal{C} satisfies ϕ' , we define a \mathcal{T}_{\neg} -interpretation \mathcal{A} by making: $\sigma_i^{\mathcal{A}} = \sigma_i^{\mathcal{C}}$ for each $1 \leq i \leq n$; $s^{\mathcal{A}}(x_i^{\mathcal{C}}) = x_{i+1}^{\mathcal{C}}$ if x_{i+1} occurs in ϕ' , and otherwise any value different from $x_i^{\mathcal{C}}$ itself; and $x^{\mathcal{A}} = x_0^{\mathcal{C}}$ for any variable x in ϕ , and $x^{\mathcal{A}} = x^{\mathcal{C}}$ otherwise. It is then easy to show \mathcal{A} satisfies ϕ . So assume that $\vdash_{\mathcal{T}_{\neg}} \phi \to \bigvee_{j=1}^{m} x_j = y_j$: then $\vdash_{\mathcal{T}} \phi' \to \bigvee_{j=1}^{m} x_j = y_j$, since otherwise there exists a \mathcal{T} -interpretation that satisfies ϕ' but not $\bigvee_{j=1}^{m} x_j = y_j$, meaning there is a \mathcal{T}_{\neg} -interpretation that satisfies ϕ but not

 $\bigvee_{j=1}^{m} x_j = y_j$ and thus leading to a contradiction. Given \mathcal{T} is convex, $\vdash_{\mathcal{T}} \phi' \to x_j = y_j$ for some $1 \leq j \leq m$, and hence $\phi' \land \neg(x_j = y_j)$ is not \mathcal{T} -satisfiable. It follows that $\phi \land \neg(x_j = y_j)$ is not \mathcal{T}_{\neg} -satisfiable, and so $\vdash_{\mathcal{T}_{\neg}} \phi \rightarrow x_j = y_j$, showing this theory is in fact convex.

Lemma 41. The Σ_s^2 -theory $\mathcal{T}_{\neq}^{\infty}$ is convex with respect to $\{\sigma, \sigma_2\}$.

Proof. Suppose that ϕ is a Σ_s^2 -formula that is also a conjunction of literals, and that $\vdash_{\mathcal{T}_{\neq}^{\infty}} \phi \to \bigvee_{i=1}^n x_i = y_i$. If ϕ is only true in $\mathcal{T}_{\neq}^{\infty}$ -interpretations \mathcal{A} where $|\sigma^{\mathcal{A}}| = 1$ we are done: indeed, in that case we may assume none of the pairs (x_i, y_i) are of sort σ (otherwise we would simply have $\vdash_{\mathcal{T}_{\neq}^{\infty}} \phi \to x_i = y_i$), take the conjunction ϕ' of literals in ϕ with variables of sort σ_2 and it is true that $\vdash_{\mathcal{T}_{\infty}} \phi' \to \bigvee_{i=1}^{n^-} x_i = y_i$, if one sees \mathcal{T}_{∞} as a theory with the sort σ_2 ; since \mathcal{T}_{∞} is convex, it follows that $\vdash_{\mathcal{T}_{\infty}} \phi' \to x_i = y_i$ for some $1 \leq i \leq n$, and thus $\vdash_{\mathcal{T}_{\neq}^{\infty}} \phi \to x_i = y_i$.

So we can assume that ϕ is true in some $\mathcal{T}_{\neq}^{\infty}$ -interpretation \mathcal{A}_0 with $|\sigma^{\mathcal{A}_0}| \geq 2$: by Lemma 2 and the fact the theory \mathcal{T} , whose interpretations are all $\mathcal{T}_{\neq}^{\infty}$ -interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}| \geq 2$, axiomatized by

$$\{\left((\psi_{k+2}^{\sigma} \wedge \psi_{\geq k+2}^{\sigma_2}) \lor \bigvee_{i=2}^{k+2} (\psi_{=i}^{\sigma} \wedge \psi_{=i}^{\sigma_2})\right) \land \forall x. \ \neg (s(x) = x) : k \in \mathbb{N}\},$$

is stably infinite, we may assume that $|\sigma^{\mathcal{A}_0}| = |\sigma_2^{\mathcal{A}_0}| = \omega$. Now, given that $\vdash_{\mathcal{T}^{\infty}_{\neq}} \phi \to \bigvee_{i=1}^n x_i = y_i$, we have that $\vdash_{\mathcal{T}} \phi \to \bigvee_{i=1}^n x_i = y_i$, and since the theory \mathcal{T} is convex by Lemma 40, we have $\vdash_{\mathcal{T}} \phi \to x_i = y_i$ for some $1 \leq i \leq n$, and of course every $\mathcal{T}_{\neq}^{\infty}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| \geq 2$ then satisfies $\phi \to x_i = y_i$.

Suppose, however, there is a $\mathcal{T}_{\neq}^{\infty}$ -interpretation \mathcal{A}_1 with $|\sigma^{\mathcal{A}_1}| = 1$ that does not satisfy $\phi \to x_i = y_i$, and thus satisfies ϕ but not $x_i = y_i$ (of course, for that to happen, x_i and y_i must be variables of sort σ_2): again by Lemma 2, we may assume $|\sigma_2^{\mathcal{A}_1}| = \omega$. We then define an interpretation \mathcal{B} such that: $\sigma^{\mathcal{B}} = \sigma^{\mathcal{A}_0}$; $\sigma_2^{\mathcal{B}} = \sigma_2^{\mathcal{A}_1}$ (so $|\sigma^{\mathcal{B}}| = |\sigma_2^{\mathcal{B}}| = \omega$); $s^{\mathcal{B}} = s^{\mathcal{A}_0}$ (so \mathcal{B} satisfies $\forall x. \neg(s(x) = x)$, and is thus a $\mathcal{T}_{\neq}^{\infty}$ -interpretation); $x^{\mathcal{B}} = x^{\mathcal{A}_0}$ for all variables x of sort σ ; and $u^{\mathcal{B}} = u^{\mathcal{A}_1}$ for all variables u of sort σ_2 . Then \mathcal{B} satisfies ϕ : any literal with variables of sort σ in ϕ is satisfied by A_0 and, by construction, by \mathcal{B} ; and any literals with variables of sort σ_2 in ϕ is satisfied by A_1 and, again by construction, by B; and since ϕ is a conjunction of literals, it is then satisfied by B. But B does not satisfy $x_i = y_i$ while still being a \mathcal{T} -interpretation, what is a contradiction.

L.4 $\mathcal{T}^3_{2,3}$

Lemma 42. The Σ_3 -theory $\mathcal{T}_{2,3}^3$ is not stably infinite, and thus not smooth, with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. Obvious, since it is not contradictory, but has no models \mathcal{A} with $|\sigma_3^{\mathcal{A}}|$ infinite.

Lemma 43. The Σ_3 -theory $\mathcal{T}_{2,3}^3$ is not stably finite with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. Consider any quantifier-free Σ_3 -formula ϕ that is satisfied by the $\mathcal{T}^3_{2,3}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = 2$, $|\sigma_2^{\mathcal{A}}| = \omega$ and $|\sigma_2^{\mathcal{A}}| = 1$: there are no $\mathcal{T}^3_{2,3}$ -interpretations \mathcal{B} such $|\sigma^{\mathcal{B}}| \le |\sigma^{\mathcal{A}}|$ and $|\sigma^{\mathcal{B}}_2| \le |\sigma^{\mathcal{A}}_2|$ and all two of these second sec are finite, so $\mathcal{T}_{2,3}^3$ cannot be stably finite.

Lemma 44. The Σ_3 -theory $\mathcal{T}_{2,3}^3$ is finitely witnessable with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. Take as witness, given a quantifier-free Σ_3 -formula ϕ ,

wit
$$(\phi) = \phi \wedge \delta$$
, where $\delta = \bigwedge_{i=1}^{3} x_i = x_i \wedge \bigwedge_{j=1}^{3} u_j = u_j \wedge (t = t)$,

 x_i are fresh variables of sort σ , u_j of sort σ_2 , and t of sort σ_3 . Of course, if $\overrightarrow{x} = vars(wit(\phi)) \setminus vars(\phi)$, we have that ϕ and $\exists \overrightarrow{x} . wit(\phi) = \phi \land \exists \overrightarrow{x} . \delta$ are $\mathcal{T}^3_{2,3}$ -equivalent since δ is a tautology.

Now, suppose \mathcal{A} is a $\mathcal{T}_{2,3}^3$ -interpretation that satisfies $wit(\phi)$: take $m = |vars_{\sigma}(\phi)^{\mathcal{A}}|$, $n = |vars_{\sigma_2}(\phi)^{\mathcal{A}}|$, $M = \max\{0, 3 - m\}$, $N = \max\{0, 3 - n\}$, and sets A and B with cardinalities, respectively, M and N, disjoint from each other and $\sigma^{\mathcal{A}}$. We define an interpretation \mathcal{B} such that: $\sigma^{\mathcal{B}} = vars_{\sigma}(\phi)^{\mathcal{A}} \cup A$, $\sigma_2^{\mathcal{B}} = vars_{\sigma_2}(\phi)^{\mathcal{A}} \cup B$, and $\sigma_3^{\mathcal{B}} = \sigma_3^{\mathcal{A}}$; $x^{\mathcal{B}} = x^{\mathcal{A}}$ for any variable x in $vars(\phi)$; $x_i \in \{x_1, x_2, x_3\} \mapsto x_i^{\mathcal{B}} \in A$ a surjection, and $y_j \in \{y_1, y_2, y_3\} \mapsto y_j^{\mathcal{B}} \in B$ also a surjection; and arbitrarily for other variables. Then we have that \mathcal{B} is a $\mathcal{T}_{2,3}^3$ -interpretation that satisfies $wit(\phi)$, since it satisfies ϕ , with $\tau^{\mathcal{B}} = vars_{\tau}(wit(\phi))^{\mathcal{B}}$ for each $\tau \in \{\sigma, \sigma_2, \sigma_3\}$.

Lemma 45. The Σ_3 -theory $\mathcal{T}_{2,3}^3$ has the finite model property with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. Follows from Theorem 2 and Lemma 44.

Lemma 46. The Σ_3 -theory $\mathcal{T}_{2,3}^3$ is not strongly finitely witnessable with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. Suppose we have a strong witness *wit*: for any quantifier-free formula ϕ that is satisfied by the $\mathcal{T}_{2,3}^3$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = 2$ and $|\sigma_2^{\mathcal{A}}| = \omega$, there is a $\mathcal{T}_{2,3}^3$ -interpretation \mathcal{A}' differing from \mathcal{A} at most on the value assigned to $\overrightarrow{x} = vars(wit(\phi)) \setminus vars(\phi)$ that satisfies $wit(\phi)$. Of course, if $V = vars(wit(\phi))$, and δ_V is the arrangement such that x is related to y iff $x^{\mathcal{A}'} = y^{\mathcal{A}'}$, then \mathcal{A}' satisfies $wit(\phi) \wedge \delta_V$. Of course, since wit is a strong witness, there is a \mathcal{T} -interpretation \mathcal{B} that satisfies $wit(\phi) \wedge \delta_V$ with $\sigma^{\mathcal{B}} = vars_{\sigma}(wit(\phi) \wedge \delta_V)^{\mathcal{B}}$ and $\sigma_2^{\mathcal{B}} = vars_{\sigma_2}(wit(\phi) \wedge \delta_V)^{\mathcal{B}}$ (and also $\sigma_3^{\mathcal{B}} = vars_{\sigma_3}(wit(\phi) \wedge \delta_V)^{\mathcal{B}}$, but that is irrelevant right now): but if $vars_{\sigma}(wit(\phi) \wedge \delta_V)^{\mathcal{B}}$ has only one element, we have a contradiction since no $\mathcal{T}_{2,3}^3$ -interpretation has only one element in its domain of sort σ ; and if $vars_{\sigma}(wit(\phi) \wedge \delta_V)^{\mathcal{B}}$ has two elements, \mathcal{B} is a $\mathcal{T}_{2,3}^3$ -interpretation with $|\sigma^{\mathcal{B}}| = 2$ and $|\sigma_2^{\mathcal{B}}|$ finite, which is again a contradiction, and we are done.

Lemma 47. The Σ_3 -theory $\mathcal{T}_{2,3}^3$ is convex with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. We can safely ignore variables of sort σ_3 , since all $\mathcal{T}_{2,3}^3$ -interpretations have domains of sort σ_3 with cardinality 1: what is left is essentially $\mathcal{T}_{2,3}$. Using this theory is convex, one can with some effort prove the same holds for $\mathcal{T}_{2,3}^3$.

M Proofs for Theories in Figure 8

First we prove the following lemma:

Lemma 48. The following holds for ς :

- 1. ς is increasing.
- 2. For all $n \in \mathbb{N} \setminus \{0, 1\}$, $\varsigma(n) > n$.

3. For any computable function $f : \mathbb{N} \to \mathbb{N}$, there exists $N \in \mathbb{N}$ such that $\varsigma(n) > f(n)$ for all $n \ge N$.

- *Proof.* 1. Trivial, as one can take a n + 1-states Turing machine whose first n states emulate the Turing machine that prints $\varsigma(n)$ 1's, and make the new state print a single new 1.
- 2. Trivial, from the last item and the fact that $\varsigma(2) > 2$.
- 3. See [7].

M.1 T_{ς}

Lemma 49. The Σ_1 -theory \mathcal{T}_{ς} is stably infinite with respect to its only sort.

Proof. It is a theory over an empty signature, and there is a \mathcal{T}_{ς} -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = \omega$, to which any finite \mathcal{T}_{ς} -interpretation can be extended.

Lemma 50. The Σ_1 -theory \mathcal{T}_{ς} is not smooth with respect to its only sort.

Proof. There is a \mathcal{T}_{ς} -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = 4$, but no $\mathcal{T}_{\varsigma}^{\infty}$ -interpretations \mathcal{B} with $|\sigma^{\mathcal{B}}| = 5$.

Lemma 51. If \mathcal{T} is a Σ_1 -theory such that, for every $n \in \mathbb{N}$, there exists a \mathcal{T} -interpretation \mathcal{A} with $n < |\sigma^{\mathcal{A}}| < \omega$, then \mathcal{T} has the finite model property with respect to σ .

Proof. Let ϕ be a quantifier-free Σ_1 -formula that is satisfied by some \mathcal{T} -interpretation \mathcal{A} , and let n be $|vars(\phi)^{\mathcal{A}}|$: by hypothesis, there exists a \mathcal{T} -interpretation \mathcal{B} with $n < |\sigma^{\mathcal{B}}| < \omega$. Since $n < |\sigma^{\mathcal{B}}|$, there exists an injective function $f : vars(\phi)^{\mathcal{A}} \to \sigma^{\mathcal{B}}$; so we define an interpretation \mathcal{B}' , different from \mathcal{B} only on $vars(\phi)$, where $x^{\mathcal{B}'} = f(x^{\mathcal{A}})$ (for each $x \in vars(\phi)$).

Now, every atomic subformula of ϕ is of the form x = y: and \mathcal{A} satisfies x = y iff $x^{\mathcal{A}} = y^{\mathcal{A}}$, what happens iff $f(x^{\mathcal{A}}) = f(y^{\mathcal{A}})$ (since f is injective), what in turns happens iff $x^{\mathcal{B}'} = y^{\mathcal{B}'}$. So \mathcal{B}' satisfies an atomic subformula of ϕ iff \mathcal{A} satisfies the same subformula, and since ϕ is a quantifier-free formula in an empty signature (and therefore its truth value is entirely determined by the truth values of its atomic subformulas), ϕ is satisfied by \mathcal{B}' . Given $|\sigma^{\mathcal{B}'}| = |\sigma^{\mathcal{B}}| < \omega$, we have finished the proof.

Lemma 52. The Σ_1 -theory \mathcal{T}_S is stably finite, and thus has the finite model property, with respect to its only sort.

Proof. Because of Lemma 48, for any $n \in \mathbb{N}$, there exists an $m \in \mathbb{N} \setminus \{0, 1\}$ such that $\varsigma(m) > n$, and so the \mathcal{T}_{ς} -interpretation \mathcal{A} with $\varsigma(m)$ elements satisfies $n < |\sigma^{\mathcal{A}}| < \omega$. We can then apply Lemma 51 to obtain that \mathcal{T}_{ς} has the finite model property, and then use Theorem 4 to obtain \mathcal{T}_{ς} is also stably finite.

Lemma 53. The Σ_1 -theory \mathcal{T}_{ς} is not finitely witnessable, and thus not strongly finitely witnessable, with respect to *its only sort.*

Proof. Suppose, by contradiction, that \mathcal{T}_{ς} has a witness *wit*, which is a computable function. We then define a function $f : \mathbb{N} \to \mathbb{N}$ by making f(0) = 0 and $f(n+1) = |vars_{\sigma}(wit(\psi_{\geq f(n)+1}))|$. Clearly, f is computable.

But we also show by induction that, for all $n \in \mathbb{N}$, $f(n) \ge \zeta(n)$, what proves f is non computable given Lemma 48, thus leading to a contradiction. It is clear that $f(0) \ge \zeta(0) = 0$; assuming we have successfully proved $f(n) \ge \zeta(n)$, we know that $\psi_{\ge f(n)+1}$ is \mathcal{T}_{ς} -satisfiable, since this theory has an infinite model, and therefore so is $wit(\psi_{\ge f(n)+1})$. Given wit is a witness, there must exist a \mathcal{T}_{ς} -interpretation \mathcal{A} that satisfies $wit(\psi_{\ge f(n)+1})$ with $\sigma^{\mathcal{A}} = vars_{\sigma}(wit(\psi_{\ge f(n)+1}))^{\mathcal{A}}$. But, by the definition of f,

$$f(n+1) = |vars_{\sigma}(wit(\psi_{\geq f(n)+1}))| \geq |vars_{\sigma}(wit(\psi_{\geq f(n)+1}))^{\mathcal{A}}| = |\sigma^{\mathcal{A}}|;$$

and since \mathcal{A} satisfies $wit(\psi_{\geq f(n)+1})$, it must also satisfy $\exists \vec{x} . wit(\psi_{\geq f(n)+1})$, for $\vec{x} = vars(wit(\psi_{\geq f(n)+1})) \setminus vars(\psi_{\geq f(n)+1})$, and thus $\psi_{\geq f(n)+1}$, meaning $|\sigma^{\mathcal{A}}| \geq f(n) + 1$. But \mathcal{A} , being a finite \mathcal{T}_{ς} -interpretation, must satisfy $|\sigma^{\mathcal{A}}| = \varsigma(m)$ for some $m \in \mathbb{N}$; since

$$\varsigma(m) = |\sigma^{\mathcal{A}}| > f(n) \ge \varsigma(n)$$

m > n according to Lemma 48, and therefore $m \ge n+1$, proving $f(n+1) \ge |\sigma^{\mathcal{A}}| \ge \varsigma(n+1)$.

Lemma 54. The Σ_1 -theory \mathcal{T}_{ς} is convex with respect to its only sort.

Proof. T_{s} is stably infinite, and is defined on an empty signature: according to Theorem 11, it is then convex.

M.2 $\mathcal{T}_{\varsigma}^{\infty}$

Lemma 55. The Σ_2 -theory $\mathcal{T}_{\varsigma}^{\infty}$ is stably infinite with respect to $\{\sigma, \sigma_2\}$.

Proof. Obvious, since it is a theory over an empty signature, and there is a $\mathcal{T}_{\varsigma}^{\infty}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = |\sigma_2^{\mathcal{A}}| = \omega$.

Lemma 56. The Σ_2 -theory $\mathcal{T}_{\varsigma}^{\infty}$ is not smooth with respect to $\{\sigma, \sigma_2\}$.

Proof. Obvious, since there is a $\mathcal{T}_{\varsigma}^{\infty}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = |\sigma_{2}^{\mathcal{A}}| = 4$, but no $\mathcal{T}_{\varsigma}^{\infty}$ -interpretations \mathcal{B} with $|\sigma^{\mathcal{B}}| = 5$ and $|\sigma_{2}^{\mathcal{B}}| = 4$.

Lemma 57. The Σ_2 -theory $\mathcal{T}_{\varsigma}^{\infty}$ has the finite model property with respect to $\{\sigma, \sigma_2\}$.

Proof. Let ϕ be a quantifier-free Σ_2 -formula, and \mathcal{A} a $\mathcal{T}_{\varsigma}^{\infty}$ -interpretation that satisfies ϕ . Take then an $n \in \mathbb{N} \setminus \{0,1\}$ such that

$$n \ge \max\{|vars_{\sigma}(\phi)^{\mathcal{A}}|, |vars_{\sigma_2}(\phi)^{\mathcal{A}}|\},\$$

which exists because both $vars_{\sigma}(\phi)$ and $vars_{\sigma_2}(\phi)$ are finite, and we know that $\varsigma(n) > n$ from Lemma 48 (since $n \in \mathbb{N} \setminus \{0, 1\}$). Take then the $\mathcal{T}_{\varsigma}^{\infty}$ -interpretation \mathcal{B} with $|\sigma^{\mathcal{B}}| = |\sigma_2^{\mathcal{B}}| = \varsigma(n)$ and, for injective functions $f : vars_{\sigma}(\phi)^{\mathcal{A}} \to \sigma^{\mathcal{B}}$ and $f_2 : vars_{\sigma_2}(\phi)^{\mathcal{A}} \to \mathcal{B}$, $x^{\mathcal{B}} = f(x^{\mathcal{A}})$ and $u^{\mathcal{B}} = f_2(u^{\mathcal{A}})$ for $x \in vars_{\sigma}(\phi)$ and $u \in vars_{\sigma_2}(\phi)$ (and arbitrary for other variables). Then, for any atomic subformula x = y or u = v of ϕ , where x and y are of sort σ , and u and v are of sort σ_2 , we have that \mathcal{B} satisfies x = y (respectively u = v) iff \mathcal{A} satisfies x = y (u = v), meaning \mathcal{B} satisfies ϕ . Of course, both $|\sigma^{\mathcal{B}}|$ and $|\sigma_2^{\mathcal{B}}|$ equal $\varsigma(n)$, and are thus finite, finishing the proof.

Lemma 58. The Σ_2 -theory $\mathcal{T}_{\varsigma}^{\infty}$ is not stably finite with respect to $\{\sigma, \sigma_2\}$.

Proof. Consider the Σ_2 -formula x = x, where x is a variable of sort σ , that is satisfied by the $\mathcal{T}_{\varsigma}^{\infty}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = 1$ and $|\sigma_2^{\mathcal{A}}| = \omega$. But there is no $\mathcal{T}_{\varsigma}^{\infty}$ -interpretation \mathcal{B} that satisfies x = x with $|\sigma^{\mathcal{B}}| \leq |\sigma^{\mathcal{A}}|$, $|\sigma_2^{\mathcal{B}}| \leq |\sigma_2^{\mathcal{A}}|$ and both finite, since there are no $\mathcal{T}_{\varsigma}^{\infty}$ -interpretations \mathcal{B} with $|\sigma^{\mathcal{B}}| = 1$ and $|\sigma_2^{\mathcal{B}}|$ finite, given that $\varsigma(k+2) \geq 4$ for $k \in \mathbb{N}$.

Lemma 59. The Σ_2 -theory $\mathcal{T}_{\varsigma}^{\infty}$ is not finitely witnessable, and thus not strongly finitely witnessable, with respect to $\{\sigma, \sigma_2\}$.

Proof. We proceed as in the proof of Lemma 53, supposing, by contradiction, that $\mathcal{T}_{\varsigma}^{\infty}$ has a witness *wit*: we then define a function $f: \mathbb{N} \to N$ by making f(0) = 0, f(1) = 1 and $f(n+1) = |vars_{\sigma}(wit(\psi_{\geq f(n)+1}^{\sigma}))|$.

Lemma 60. The Σ_2 -theory $\mathcal{T}_{\varsigma}^{\infty}$ is convex with respect to $\{\sigma, \sigma_2\}$.

Proof. Since $\mathcal{T}_{\varsigma}^{\infty}$ is stably infinite and Σ_2 is empty, Theorem 11 guarantees it is convex.

M.3 T_n^{ς}

Lemma 61. The Σ_2 -theory $\mathcal{T}_n^{\varsigma}$ is not stably infinite, and thus not smooth, with respect to $\{\sigma, \sigma_2\}$ for any $n \ge 1$.

Proof. Obvious, since $\mathcal{T}_n^{\varsigma}$ is not contradictory, but there are no $\mathcal{T}_n^{\varsigma}$ -interpretations \mathcal{A} such that $|\sigma^{\mathcal{A}}|$ and $|\sigma_2^{\mathcal{A}}|$ are both infinite.

Lemma 62. The Σ_2 -theory $\mathcal{T}_n^{\varsigma}$ is stably finite, and thus has the finite model property, with respect to $\{\sigma, \sigma_2\}$ for any $n \ge 1$.

Proof. Let ϕ be a quantifier-free Σ_2 -formula, and \mathcal{A} a $\mathcal{T}_n^{\varsigma}$ -interpretation that satisfies ϕ . Take then $\varsigma(m)$ as the minimum value of $\varsigma(k)$ such that

$$\varsigma(k) \ge \min\{2, |vars_{\sigma_2}(\phi)^{\mathcal{A}}|\}.$$

Take then the $\mathcal{T}_n^{\varsigma}$ -interpretation \mathcal{B} with $\sigma^{\mathcal{B}} = \sigma^{\mathcal{A}}$, $|\sigma_2^{\mathcal{B}}| = \varsigma(m)$ and, for an injective function $f_2 : vars_{\sigma_2}(\phi)^{\mathcal{A}} \to \mathcal{B}$, $x^{\mathcal{B}} = x^{\mathcal{A}}$ and $u^{\mathcal{B}} = f_2(u)$ for $x \in vars_{\sigma}(\phi)$ and $u \in vars_{\sigma_2}(\phi)$ (and arbitrary for other variables). This way, \mathcal{B} satisfies ϕ , and of course, $|\sigma^{\mathcal{B}}| = |\sigma^{\mathcal{A}}|$ and $|\sigma_2^{\mathcal{B}}| \le |\sigma_2^{\mathcal{A}}|$ are both finite, finishing the proof.

Lemma 63. The Σ_2 -theory $\mathcal{T}_n^{\varsigma}$ is not finitely witnessable, and thus not strongly finitely witnessable, with respect to $\{\sigma, \sigma_2\}$ for any $n \ge 1$.

Proof. We proceed as in the proof of Lemma 53, supposing, by contradiction, that $\mathcal{T}_n^{\varsigma}$ has a witness *wit*: we then define a function $f : \mathbb{N} \to N$ by making f(0) = 0, f(1) = 1 and $f(n+1) = |vars_{\sigma_2}(wit(\psi_{\geq f(n)+1}^{\sigma_2}))|$.

Lemma 64. The Σ_2 -theory $\mathcal{T}_n^{\varsigma}$ is not convex with respect to $\{\sigma, \sigma_2\}$ for n > 1.

Proof. Take n + 1 variables x_1 trough x_{n+1} of sort σ , and it is clear that $\vdash_{\mathcal{T}_n^{\varsigma}} \bigvee_{i=1}^n \bigvee_{j=i+1}^{n+1} x_i = x_j$; we cannot, however, derive $\vdash_{\mathcal{T}_n^{\varsigma}} x_i = x_j$ for any $1 \le i < j \le n+1$ since there is a \mathcal{T} -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = n \ge 2$, implying $\mathcal{T}_n^{\varsigma}$ is not convex.

M.3.1 $\mathcal{T}_1^{\varsigma}$ As proved in Appendix M.3, $\mathcal{T}_1^{\varsigma}$ is neither stably infinite, smooth, finitely witnessable nor strongly finitely witnessable, but it is stably finite and has the finite model property with respect to $\{\sigma, \sigma_2\}$.

Lemma 65. The Σ_2 -theory $\mathcal{T}_1^{\varsigma}$ is convex with respect to $\{\sigma, \sigma_2\}$.

Proof. If we disregard the second sort of $\mathcal{T}_1^{\varsigma}$, it becomes simply \mathcal{T}_{ς} : given this theory is stably-infinite in an empty signature, from Theorem 11 we obtain it is convex.

M.4 $\mathcal{T}_{m,n}^{\varsigma}$

Lemma 66. The Σ_2 -theory $\mathcal{T}_{m,n}^{\varsigma}$ is not stably infinite, and thus not smooth, with respect to $\{\sigma, \sigma_2\}$, for all $m, n \in \mathbb{N}$.

Proof. Obvious, given $\mathcal{T}_{m,n}^{\varsigma}$ is not contradictory, but has no interpretations \mathcal{A} where both $|\sigma^{\mathcal{A}}|$ and $|\sigma_{2}^{\mathcal{A}}|$ are infinite.

Lemma 67. The Σ_2 -theory $\mathcal{T}_{m,n}^{\varsigma}$ has the finite model property with respect to $\{\sigma, \sigma_2\}$ for all $m, n \in \mathbb{N}$.

Proof. Let ϕ be a quantifier-free Σ_2 -formula, and \mathcal{A} a $\mathcal{T}^{\varsigma}_{m,n}$ -interpretation that satisfies ϕ . Take sets A and B, disjoint from each other and $\sigma^{\mathcal{A}}$ and $\sigma^{\mathcal{A}}_2$, with $\max\{m,n\} - |vars_{\sigma}(\phi)^{\mathcal{A}}|$ and $\varsigma(k) - |vars_{\sigma_2}(\phi)^{\mathcal{A}}|$ elements, respectively, where k is the minimum natural number (greater than 1) such that $\varsigma(k) \geq |vars_{\sigma_2}(\phi)^{\mathcal{A}}|$. We then define a $\mathcal{T}^{\varsigma}_{m,n}$ -interpretation \mathcal{B} by making: $\sigma^{\mathcal{B}} = vars_{\sigma}(\phi)^{\mathcal{A}} \cup A$ (which has $\max\{m,n\}$ elements); $\sigma^{\mathcal{B}}_2 = vars_{\sigma_2}(\phi)^{\mathcal{A}} \cup B$ (that has, then, $\varsigma(k)$ elements); $x^{\mathcal{B}} = x^{\mathcal{A}}$ and $u^{\mathcal{B}} = u^{\mathcal{A}}$ for all variables x and u in ϕ of sorts, respectively, σ and σ_2 ; and arbitrarily for other variables. It is then clear that \mathcal{B} is a $\mathcal{T}^{\varsigma}_{m,n}$ -interpretation, with both $|\sigma^{\mathcal{B}}|$ and $|\sigma^{\mathcal{B}}_2|$ finite, that satisfies ϕ .

Lemma 68. The Σ_2 -theory $\mathcal{T}_{m,n}^{\varsigma}$ is not stably finite with respect to $\{\sigma, \sigma_2\}$ for all $m, n \in \mathbb{N}$ such that $m \neq n$.

Proof. Obvious, since x = x, for x a variable of sort σ , is satisfied by the $\mathcal{T}_{m,n}^{\varsigma}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = \min\{m, n\}$ and $|\sigma_2^{\mathcal{A}}| = \omega$, but there are no $\mathcal{T}_{m,n}^{\varsigma}$ -interpretations with $|\sigma^{\mathcal{B}}|$ and $|\sigma_2^{\mathcal{B}}|$ finite, and $|\sigma^{\mathcal{B}}| \leq |\sigma^{\mathcal{A}}|$ and $|\sigma_2^{\mathcal{B}}| \leq |\sigma_2^{\mathcal{B}}|$.

Lemma 69. The Σ_2 -theory $\mathcal{T}_{m,n}^{\varsigma}$ is not finitely witnessable, and thus not strongly finitely witnessable, with respect to $\{\sigma, \sigma_2\}$ for all $m, n \in \mathbb{N}$.

Proof. Suppose $\mathcal{T}_{m,n}^{\varsigma}$ has a (necessarily computable) witness *wit*, and consider the function $f : \mathbb{N} \to \mathbb{N}$ such that f(0) = 0, f(1) = 1 and, for k > 1,

$$f(k) = |vars(wit(\delta_k))|, \quad \text{where} \quad \delta_k = \bigwedge_{1 \le i < j \le \max\{m,n\}} \neg (x_i = x_j) \land \bigwedge_{1 \le i < j \le f(k) + 1} \neg (u_i = u_j),$$

where x_1 through $x_{\max\{m,n\}}$ are variables of sort σ , and the u_i are variables of sort σ_2 : we prove by induction that $f(k) \ge \varsigma(k)$, and thus f is not computable, leading to a contradiction with its definition that uses only computable

notions. This is obvious for k = 0 and k = 1, so assume it holds for an arbitrary k. Since δ_{k+1} holds (for all $k \in \mathbb{N}$) in some interpretation on the $\mathcal{T}^{s}_{\varsigma}$ -model with $\max\{m, n\}$ elements in the domain of sort σ and ω elements in the domain of sort σ_2 , so does $wit(\delta_{k+1})$: hence there must exist a $\mathcal{T}^{s}_{\varsigma}$ -interpretation \mathcal{A} that satisfies $wit(\delta_{k+1})$ (and thus δ_{k+1}) with $\sigma^{\mathcal{A}} = vars(wit(\delta_{k+1}))^{\mathcal{A}}$ and $\sigma_2^{\mathcal{A}} = vars_{\sigma_2}(wit(\delta_{k+1}))^{\mathcal{A}}$. So $|\sigma^{\mathcal{A}}| \ge \max\{m, n\}$ (and thus $|\sigma^{\mathcal{A}}| = \max\{m, n\}$), and $|\sigma_2^{\mathcal{A}}| \ge f(k) + 1 \ge \varsigma(k) + 1$, forcing f(k+1) to be equal to or greater than $\varsigma(k+1)$, as we wished to prove.

Lemma 70. The Σ_2 -theory $\mathcal{T}_{m,n}^{\varsigma}$ is not convex with respect to $\{\sigma, \sigma_2\}$ if $\max\{m, n\} > 1$.

Proof. Take $\max\{m, n\} + 1$ variables x_1 trough $x_{\max\{m, n\}+1}$ of sort σ : it is then true that

$$\vdash_{\mathcal{T}_{m,n}^{\varsigma}} \bigvee_{i=1}^{\max\{m,n\}-1} \bigvee_{j=i+1}^{\max\{m,n\}} x_i = x_j,$$

but we cannot get $\vdash_{\mathcal{T}_{m,n}^{\varsigma}} x_i = x_j$ for any $1 \le i < j \le \max\{m,n\} + 1$ since there is a $\mathcal{T}_{m,n}^{\varsigma}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = \max\{m,n\} > 1$ and $|\sigma_2^{\mathcal{A}}| = \omega$.

M.5 $\mathcal{T}_{\varsigma}^{s}$

Lemma 71. The Σ_s -theory \mathcal{T}_s^s is smooth, and thus stably infinite, with respect to $\{\sigma\}$.

Proof. Let ϕ be a quantifier-free Σ_s -formula, \mathcal{A} a \mathcal{T}_s^s -interpretation that satisfies ϕ , and $\kappa(\sigma)$ a cardinal equal to or grater than $|\sigma^{\mathcal{A}}|$. There are then two cases to consider.

- If κ(σ) is infinite, consider a set A with cardinality κ(σ) and disjoint from σ^A: we define a T_ζ^s-interpretation B by making σ^B = σ^A ∪ A (which has cardinality κ(σ)); s^B(a) = s^A(a) for all a ∈ σ^A, and s^B(a) = a for all a ∈ A (meaning B has infinite elements a satisfying s^B(a) = a, and thus is indeed a T_ζ^s-interpretation); and x^B = x^A for all variables x (meaning B satisfies φ).
- 2. If $\kappa(\sigma)$ is finite, so is $|\sigma^{\mathcal{A}}|$: take then sets A and B, disjoint from each other and $\sigma^{\mathcal{A}}$, with $\varsigma^{-1}(\kappa(\sigma)) \varsigma^{-1}(|\sigma^{\mathcal{A}}|)$ and $\kappa(s) (|\sigma^{\mathcal{A}}| + |A|)$ elements, respectively. We then define an interpretation \mathcal{B} by making: (a) $\sigma^{\mathcal{B}} = \sigma^{\mathcal{A}} \cup A \cup B$ (which has $\kappa(\sigma)$ elements);
 - (b) $s^{\mathcal{B}}(a) = s^{\mathcal{A}}(a)$ for all $a \in \sigma^{\mathcal{A}}$, $s^{\mathcal{B}}(a) = a$ for all $a \in A$, and $s^{\mathcal{B}}(a)$ any element of $\sigma^{\mathcal{B}} \setminus \{a\}$ for $a \in B$ (notice that \mathcal{B} possesses $\varsigma^{-1}(|\sigma^{\mathcal{A}}|)$ elements in $\sigma^{\mathcal{A}}$ where $s^{\mathcal{B}}$ is the identity, and $\varsigma^{-1}(\kappa(\sigma)) - \varsigma^{-1}(|\sigma^{\mathcal{A}}|)$ more on A, to a total of $\varsigma^{-1}(\kappa(\sigma))$, hat makes of \mathcal{B} a $\mathcal{T}_{\varsigma}^{s}$ -interpretation);
 - (c) and $x^{\mathcal{B}} = x^{\mathcal{A}}$ for all variables x (meaning \mathcal{B} satisfies ϕ).

Lemma 72. The Σ_s -theory \mathcal{T}_s^s is stably finite, and thus has the finite model property, with respect to $\{\sigma\}$.

Proof. Because of Theorem 4, we only need to prove that $\mathcal{T}_{\varsigma}^{s}$ has the finite model property. So take a quantifierfree Σ_{s} -formula ϕ and an infinite $\mathcal{T}_{\varsigma}^{s}$ -interpretation \mathcal{A} that satisfies ϕ . Let $vars(\phi) = \{z_{1}, \ldots, z_{n}\}$, and for each z_{i} , let M_{i} be the maximum of j such that the term $s^{j}(z_{i})$ occurs in ϕ . Take then

$$\alpha(\phi)^{\mathcal{A}} = \{ (s^{\mathcal{A}})^{j}(z_{i}^{\mathcal{A}}) : 1 \le i \le n, 0 \le j \le M_{i} \}, \quad Id^{\mathcal{A}}(\phi) = \{ a \in \alpha(\phi)^{\mathcal{A}} : s^{\mathcal{A}}(a) = a \},$$
$$\alpha(\phi)^{\mathcal{A}}_{+1} = \{ (s^{\mathcal{A}})^{j}(z_{i}^{\mathcal{A}}) : 1 \le i \le n, 0 \le j \le M_{i} + 1 \},$$

 $p = |\alpha(\phi)^{\mathcal{A}}|, q = |\alpha(\phi)^{\mathcal{A}}_{+1}|, r = |Id^{\mathcal{A}}(\phi)|, \text{ and } m = \max\{\varsigma(-p+q+r), q\}.$ We finally take sets A and B, disjoint from each other and $\sigma^{\mathcal{A}}$, with, respectively, $\varsigma^{-1}(m) + p - q - r$ elements and $m - p + r - \varsigma^{-1}(m)$ elements. We then define an interpretation \mathcal{B} with:

1. $\sigma^{\mathcal{B}} = \alpha(\phi)_{+1}^{\mathcal{A}} \cup A \cup B$ (which has *m* elements);

- 2. $s^{\mathcal{B}}(a) = \sigma^{\mathcal{A}}(a)$ for all $a \in \alpha(\phi)^{\mathcal{A}}$ (which has r elements that satisfy $s^{\mathcal{B}}(a) = a$, namely those in $Id^{\mathcal{A}}(\phi)$), $\sigma^{\mathcal{A}}(a) = a$ for all $a \in A$ or $a \in \alpha(\phi)^{\mathcal{A}}_{+1} \setminus \alpha(\phi)^{\mathcal{A}}$ (there are, respectively, $\varsigma^{-1}(m) + p - q - r$ and q - p, leading to a total of $\varsigma^{-1}(m)$ elements in \mathcal{B} where $\sigma^{\mathcal{B}}$ is the identity, meaning \mathcal{B} is indeed a $\mathcal{T}^{\varsigma}_{\varsigma}$ -interpretation), and $s^{\mathcal{B}}(a)$ any value in $\sigma^{\mathcal{B}} \setminus \{a\}$ for $a \in B$;
- 3. $x^{\mathcal{B}} = x^{\mathcal{A}}$ for all variables x in ϕ , and otherwise arbitrary.

It is obvious that \mathcal{B} is finite, and furthermore one can show that it additionally satisfies ϕ .

Lemma 73. The Σ_s -theory \mathcal{T}_s^s is not finitely witnessable, and thus not strongly finitely witnessable, with respect to $\{\sigma\}$.

Proof. Suppose $\mathcal{T}_{\varsigma}^{s}$ indeed has a witness *wit* and consider the function $f : \mathbb{N} \to \mathbb{N}$ such that f(0) = 0 and, for n > 0,

$$f(n) = |vars(wit(\delta_n))|, \quad \text{where} \quad \delta_n = \bigwedge_{1 \le i < j \le n} \neg (x_i = x_j) \land \bigwedge_{i=1}^n (s(x_i) = x_i) :$$

since δ_n holds in some infinite $\mathcal{T}_{\varsigma}^{s}$ -interpretations, so does $wit(\phi)$, and thus there must exist a $\mathcal{T}_{\varsigma}^{s}$ -interpretation \mathcal{A} that satisfies $wit(\delta_n)$ (and thus δ_n) with $\sigma^{\mathcal{A}} = vars(wit(\delta_n))^{\mathcal{A}}$. Then, there i a finite number $m \ge n$ of elements in \mathcal{A} satisfying $s^{\mathcal{A}}(a) = a$, meaning that \mathcal{A} satisfies $\psi_{=m}^{=}$ and thus $\psi_{\ge \varsigma(m)}$. In other words, $f(n) \ge \varsigma(m) \ge \varsigma(n)$, and so f cannot be computable, contradicting its definition.

Lemma 74. Let \mathcal{T} be the Σ_s -theory with all Σ_s -structures as models: if the quantifier-free Σ_s -formula ϕ is \mathcal{T} -satisfiable, it is \mathcal{T}_s^s -satisfiable.

Proof. Let ϕ be a quantifier-free, \mathcal{T} -satisfiable formula, and let \mathcal{A} be a \mathcal{T} -interpretation that satisfies ϕ ; because \mathcal{T} is stably infinite, we can obtain a \mathcal{T} -interpretation \mathcal{B} that satisfies ϕ with $|\sigma^{\mathcal{B}}| \geq \omega$. Then let m be the number of elements a in $\sigma^{\mathcal{B}}$ such that $s^{\mathcal{B}}(a) = a$, and n be the number of elements a such that $s^{\mathcal{B}}(a) \neq a$.

Because $|\sigma^{\mathcal{B}}| \ge \omega$, either *m* or *n* is infinite, and we define \mathcal{C} by making $\sigma^{\mathcal{C}} = \sigma^{\mathcal{B}} \cup \{a_n : n \in \mathbb{N}\} \cup \{b_n : n \in \mathbb{N}\}$, where the latter sets are disjoint from $\sigma^{\mathcal{B}}$ and each other; $s^{\mathcal{C}}(a) = s^{\mathcal{B}}(a)$ for all $a \in \sigma^{\mathcal{B}}$, $s^{\mathcal{C}}(a_n) = a_n$ and $s^{\mathcal{C}}(b_n) = a_n$; and $x^{\mathcal{C}} = x^{\mathcal{B}}$ for all variables *x*. Then \mathcal{C} not only satisfies ϕ , but is also a $\mathcal{T}_{\varsigma}^s$ -interpretation, meaning ϕ is also $\mathcal{T}_{\varsigma}^s$ -satisfiable.

Lemma 75. The Σ_s -theory \mathcal{T}_s^s is convex with respect to $\{\sigma\}$.

Proof. For a proof by contradiction, take a conjunction of Σ_s -literals ϕ and assume that, although $\vdash_{\mathcal{T}^s_{\varsigma}} \phi \rightarrow \bigvee_{i=1}^n x_i = y_i$, we have $\not\vdash_{\mathcal{T}^s_{\varsigma}} \phi \rightarrow x_i = y_i$ for each $1 \leq i \leq n$. So we can find $\mathcal{T}^s_{\varsigma}$ -interpretations \mathcal{A}_i that satisfy, simultaneously, ϕ and $\neg(x_i = y_i)$.

Now, we state that $\vdash_{\mathcal{T}} \phi \to \bigvee_{i=1}^{n} x_i = y_i$, for \mathcal{T} the Σ_s -theory of the uninterpreted function s: if that were not true, we should be able to find a \mathcal{T} -interpretation \mathcal{A} that satisfies ϕ but not $\bigvee_{i=1}^{n} x_i = y_i$, meaning $\phi \land \neg \bigvee_{i=1}^{n} x_i = y_i$ is \mathcal{T} -satisfiable, and by Lemma 74 that would imply this formula is $\mathcal{T}_{\varsigma}^s$ -satisfiable, contradicting our assumptions. And, because all $\mathcal{T}_{\varsigma}^s$ -interpretations are \mathcal{T} -interpretations, we get that $\phi \land \neg (x_i = y_i)$ is \mathcal{T} -satisfiable for each $1 \leq i \leq n$.

Given Theorem 13 and that $\vdash_{\mathcal{T}} \phi \to \bigvee_{i=1}^{n} x_i = y_i$, there must exist $1 \le i \le n$ such that $\vdash_{\mathcal{T}} \phi \to x_i = y_i$, contradicting that $\phi \land \neg(x_i = y_i)$ is \mathcal{T} -satisfiable. The conclusion must be that $\mathcal{T}_{\varsigma}^s$ is convex.

M.6 $\mathcal{T}_{\varsigma}^{\neq}$

Lemma 76. The Σ_s -theory \mathcal{T}_c^{\neq} is not stably infinite, and thus not smooth, with respect to $\{\sigma\}$.

Proof. Any $\mathcal{T}_{\varsigma}^{\neq}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = 2$ satisfies $\neg(s(x) = x)$, while all infinite $\mathcal{T}_{\varsigma}^{\neq}$ -interpretations satisfy instead s(x) = x.

Lemma 77. The Σ_s -theory \mathcal{T}_s^{\neq} is stably finite, and thus has the finite model property, with respect to $\{\sigma\}$.

Proof. Let ϕ be a quantifier-free Σ_s -formula, and \mathcal{A} a $\mathcal{T}_{\varsigma}^{\neq}$ -interpretation that satisfies ϕ : we may assume \mathcal{A} is infinite, since otherwise there would be nothing to prove, and so \mathcal{A} satisfies $\forall x. (s(x) = x)$. Take then the smallest m such that $\varsigma(m) \geq \max\{|vars(\phi)^{\mathcal{A}}|, 3\}$, and a set \mathcal{A} disjoint from $\sigma^{\mathcal{A}}$ with $\varsigma(m) - |vars(\phi)^{\mathcal{A}}|$ elements. We define the interpretation \mathcal{B} by: $\sigma^{\mathcal{B}} = vars(\phi)^{\mathcal{A}} \cup \mathcal{A}$ (so \mathcal{B} has at least $\varsigma(m) \geq 3$ elements, but $|\sigma^{\mathcal{B}}| \leq |\sigma^{\mathcal{A}}|$); $s^{\mathcal{B}}(a) = a$ for all $a \in \sigma^{\mathcal{B}}$ (so \mathcal{B} satisfies $\forall x. (s(x) = x)$ and is, therefore, a $\mathcal{T}_{\varsigma}^{\neq}$ -interpretation); $x^{\mathcal{B}} = x^{\mathcal{A}}$ for all variables x in ϕ (so \mathcal{B} satisfies ϕ); and arbitrarily for other variables.

Lemma 78. The Σ_s -theory $\mathcal{T}_{\varsigma}^{\neq}$ is not finitely witnessable, and thus not strongly finitely witnessable, with respect to $\{\sigma\}$.

Proof. The proof is very similar to that of Lemma 53: if $\mathcal{T}_{\varsigma}^{\neq}$ indeed has a witness *wit*, we define a function $f : \mathbb{N} \to N$ by making f(0) = 0, f(1) = 1, and for $n \ge 1$ $f(n+1) = |vars_{\sigma}(wit(\psi_{\ge f(n)+1}))|$. Because *wit* is computable, so must be f, but one can prove that f grows at least as fast as ς , so we reach a contradiction.

Lemma 79. The Σ_s -theory \mathcal{T}_s^{\neq} is not convex with respect to $\{\sigma\}$.

Proof. Take variables x, y, w and z, and consider the conjunction of literals ϕ equal to $\neg(s(x) = x) \land \neg(y = w)$: it is then true that $\vdash_{\mathcal{T}_{\varsigma}^{\neq}} \phi \to (y = z) \lor (w = z)$, since $\neg(s(x) = x)$ is only satisfied in a $\mathcal{T}_{\varsigma}^{\neq}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = 2$, and by the pigeonhole principle, if $\neg(y = w)$, then either y = z or w = z. But we cannot get either $\vdash_{\mathcal{T}_{\varsigma}^{\neq}} \phi \to y = z$ nor $\vdash_{\mathcal{T}_{\varsigma}^{\neq}} \phi \to w = z$, since \mathcal{A} has $|\sigma^{\mathcal{A}}| = 2$, and we can change the value assigned to z while keeping ϕ satisfied.

M.7 $\mathcal{T}_{s,1}^{\neq}$

Lemma 80. The Σ_s -theory $\mathcal{T}_{s,1}^{\neq}$ is not stably infinite, and thus is not smooth, with respect to $\{\sigma\}$.

Proof. Obvious, since the $\mathcal{T}_{\varsigma,1}^{\neq}$ -interpretation with one element in its domain satisfies s(x) = x, while all infinite $\mathcal{T}_{\varsigma,1}^{\neq}$ -interpretation satisfy instead $\neg(s(x) = x)$.

Lemma 81. The Σ_s -theory $\mathcal{T}_{s,1}^{\neq}$ is stably finite, and thus has the finite model property, with respect to $\{\sigma\}$.

Proof. Because of Theorem 4, we just need to prove $\mathcal{T}_{\varsigma,1}^{\neq}$ has the finite model property. We then take a quantifierfree Σ_s -formula ϕ , and a $\mathcal{T}_{\varsigma,1}^{\neq}$ -interpretation \mathcal{A} that satisfies ϕ , and without loss of generality we assume that \mathcal{A} has at least 4 elements, and thus $s^{\mathcal{A}}$ is never the identity. If x_1 through x_n are the variables in ϕ , and M_i is the maximum of the values j such that $s^j(x_i)$ appears in ϕ , we define

$$A = \{s^{j}(x_{i}) : 1 \le i \le n, 0 \le j \le M_{i}\}^{\mathcal{A}}, \quad A^{+} = \{s^{j}(x_{i}) : 1 \le i \le n, 0 \le j \le M_{i} + 1\}^{\mathcal{A}}$$

and $m = |A^+|$: then we take the least M on the image of ς such that $M \ge m$; we also take a set B with M - m elements disjoint from σ^A . We then define an interpretation \mathcal{B} by making: $\sigma^{\mathcal{B}} = A^+ \cup B$; $s^{\mathcal{B}}(a) = s^A(a)$ for all $a \in A$, and $s^{\mathcal{B}}(a)$ any value different from a, for each $a \in B$ or $a \in A^+ \setminus A$; $x^{\mathcal{B}} = x^A$ for each variable x of ϕ , and arbitrarily otherwise. \mathcal{B} has then M (finite) number of elements, that is a value of ς , and satisfies $\neg(s(x) = x)$ for all x, meaning it is a $\mathcal{T}_{\varsigma,1}^{\neq}$ -interpretation; furthermore, it satisfies ϕ , and so we are done.

Lemma 82. The Σ_s -theory $\mathcal{T}_{\varsigma,1}^{\neq}$ is not finitely witnessable, and thus is not strongly finitely witnessable, with respect to $\{\sigma\}$.

Proof. This follows Lemma 53: suppose we have a witness *wit*, necessarily computable, and define a function $f : \mathbb{N} \to \mathbb{N}$ such that f(0) = 0, f(1) = 1 and, assuming f(n) defined,

$$f(n+1) = |vars_{\sigma}(wit(\bigwedge_{i=1}^{f(n)} \bigwedge_{j=i+1}^{f(n)+1} \neg (x_i = x_j)))|$$

which is clearly computable. The contradiction is that f grows at least as fast as ς , and by Lemma 48 cannot be computable.

Lemma 83. The Σ_s -theory $\mathcal{T}_{\varsigma,1}^{\neq}$ is convex with respect to $\{\sigma\}$.

Proof. This is a very long and dull proof: however, an essentially equal proof may be found in [12] when we prove that \mathcal{T}_{odd}^{\neq} is convex.

M.8 T_{c}^{\vee}

The proofs that $\mathcal{T}_{\varsigma,1}^{\neq}$ is smooth (and thus stably infinite) and stably finite (and that it therefore has the finite model property), but not finitely witnessable (and thus is not strongly finitely witnessable) are very similar to those of the correspondent properties of $\mathcal{T}_{\varsigma}^{s}$, in Appendix M.5.

Lemma 84. The Σ_s -theory \mathcal{T}_s^{\vee} is not convex with respect to $\{\sigma\}$.

Proof. Because $\mathcal{T}_{\varsigma}^{\lor}$ satisfies ψ_{\lor} , given the cube ϕ equal to $(y = s(x)) \land (z = s(y))$, we have that $\vdash_{\mathcal{T}_{\varsigma}^{\lor}} \phi \rightarrow (x = z) \lor (y = z)$. Take then the $\mathcal{T}_{\varsigma}^{\lor}$ -interpretations \mathcal{A} and \mathcal{B} with: $\sigma^{\mathcal{A}} = \{a, b, c, d\}$; $s^{\mathcal{A}}(a) = s^{\mathcal{B}}(a) = a$, $s^{\mathcal{A}}(b) = s^{\mathcal{B}}(b) = b$, $s^{\mathcal{A}}(c) = a$, $s^{\mathcal{A}}(d) = b$, $s^{\mathcal{B}}(c) = d$ and $s^{\mathcal{B}}(d) = c$; $x^{\mathcal{A}} = c$, $y^{\mathcal{A}} = z^{\mathcal{A}} = a$, $x^{\mathcal{B}} = z^{\mathcal{B}} = c$ and $y^{\mathcal{B}} = d$.

Then both \mathcal{A} and \mathcal{B} satisfy ϕ , but \mathcal{A} does not satisfy x = z, while \mathcal{B} does not satisfy y = z.

M.9 $T_s^{=}$

Lemma 85. The Σ_s^2 -theory $\mathcal{T}_s^=$ is smooth, and thus stably infinite, with respect to $\{\sigma, \sigma_2\}$.

Proof. Take a quantifier-free Σ_s^2 -formula ϕ , a $\mathcal{T}_{\varsigma}^{=}$ -interpretation \mathcal{A} that satisfies ϕ , and cardinals $\kappa(\sigma) \geq |\sigma^{\mathcal{A}}|$ and $\kappa(\sigma_2) \geq |\sigma_2^{\mathcal{A}}|$; take as well sets A and B with, respectively, $\kappa(\sigma) - |\sigma^{\mathcal{A}}|$ and $\kappa(\sigma_2) - |\sigma_2^{\mathcal{A}}|$ elements, disjoint from each other and the domains of \mathcal{A} .

We the define the interpretation \mathcal{B} as follows: $\sigma^{\mathcal{B}} = \sigma^{\mathcal{A}} \cup A$ and $\sigma_2^{\mathcal{B}} = \sigma_2^{\mathcal{A}} \cup B$ (so $|\sigma^{\mathcal{B}}| \ge |\sigma^{\mathcal{A}}|$); $s^{\mathcal{B}}(a) = s^{\mathcal{A}}(a)$ for each $a \in \sigma^{\mathcal{A}}$, and $\sigma^{\mathcal{B}}(a)$ is anything but a for $a \in A$ (this way, the number k + 1 of elements in $\sigma^{\mathcal{B}}$ satisfying $s^{\mathcal{B}}(a) = a$ is the same as the number of elements in $\sigma^{\mathcal{A}}$ satisfying $s^{\mathcal{A}}(a) = a$; since $|\sigma^{\mathcal{A}}| \ge \varsigma(k+1)$, $|\sigma^{\mathcal{B}}| \ge \varsigma(k+1)$, and thus \mathcal{B} is a $\mathcal{T}_{\varsigma}^{=}$ -interpretation); and $x^{\mathcal{B}} = x^{\mathcal{A}}$ for every variable x. This way \mathcal{B} satisfies ϕ , $|\sigma^{\mathcal{B}}| = \kappa(\sigma)$ and $|\sigma_2^{\mathcal{B}}| = \kappa(\sigma_2)$.

Lemma 86. The Σ_s^2 -theory $\mathcal{T}_{\varsigma}^{=}$ is not finitely witnessable, and thus not strongly finitely witnessable, with respect to $\{\sigma, \sigma_2\}$.

Proof. Suppose we have a witness wit. We define a function $f : \mathbb{N} \to \mathbb{N}$ by making f(0) = 0, f(1) = 1, and assuming f(n) defined,

$$f(n+1) = |vars_{\sigma_2}(wit(\bigwedge_{i=1}^{n+1}(s(x_i) = x_i) \land \bigwedge_{i=1}^n \bigwedge_{j=i+1}^{n+1} \neg (x_i = x_j)))|,$$

which is computable: after all, producing the formula inside the witness is easily seen to be computable, as well as is producing its witness (given *wit* is computable), its set of variables and the cardinality of that set. So f is a computable function.

However, since $\delta_{n+1} = \bigwedge_{i=1}^{n+1} (s(x_i) = x_i) \land \bigwedge_{i=1}^n \bigwedge_{j=i+1}^{n+1} \neg (x_i = x_j \text{ is only satisfied in a } \mathcal{T}_{\varsigma}^{=}\text{-interpretation } \mathcal{A}$ that satisfies $\psi_{\geq n+1}^{=}$, $wit(\delta_{n+1})$ is only satisfied in a $\mathcal{T}_{\varsigma}^{=}\text{-interpretation } \mathcal{A}'$ with $|\sigma^{\mathcal{A}'}| \ge \varsigma(n+1)$. Because there must exist a $\mathcal{T}_{\varsigma}^{=}\text{-interpretation } \mathcal{B}$ that satisfies $wit(\delta_{n+1})$ with $\sigma^{\mathcal{B}} = vars_{\sigma}(wit(\delta_{n+1}))^{\mathcal{B}}$, we obtain that $f(n+1) = |vars_{\sigma}(wit(\delta_{n+1}))| \ge |\sigma^{\mathcal{B}}| \ge \varsigma(n+1)$, and by Lemma 48 f cannot be computable, leading to a contradiction.

Lemma 87. Let \mathcal{T} be the Σ_s^2 -theory of uninterpreted functions: then a quantifier-free Σ_s^2 -formula ϕ is $\mathcal{T}_{\varsigma}^{=}$ -satisfiable if it is \mathcal{T} -satisfiable.

Proof. See Lemma 74 for a similar proof.

Lemma 88. The Σ_s^2 -theory $\mathcal{T}_s^=$ is convex with respect to $\{\sigma, \sigma_2\}$.

Proof. Given Theorem 13, it is easy to prove that the Σ_s^2 -theory of uninterpreted functions is also convex; by applying Lemma 87, we get that $\mathcal{T}_s^=$ is convex much like in the proof of Lemma 75.

Lemma 89. The Σ_s^2 -theory $\mathcal{T}_{\varsigma}^{=}$ is stably finite, and thus has the finite model property, with respect to $\{\sigma, \sigma_2\}$.

Proof. Take a quantifier-free Σ_s^2 -formula ϕ , and a $\mathcal{T}_{\varsigma}^{=}$ -interpretation \mathcal{A} that satisfies ϕ . Let $vars_{\sigma}(\phi) = \{x_1, \ldots, x_n\}$ and, for each $i \in [1, n]$, let M_i be the maximum of j such that $s^j(x_i)$ appears in ϕ . We also define

$$A = \{s^{j}(x_{i}) : 1 \le i \le n, 0 \le j \le M_{i}\}^{\mathcal{A}} \text{ and } A^{+} = \{s^{j}(x_{i}) : 1 \le i \le n, 0 \le j \le M_{i} + 1\}^{\mathcal{A}}.$$

Now, let *m* be the maximum between 2 and the number of elements *a* in *A* that satisfy $s^{\mathcal{A}}(a) = a$: notice it is equal to or less than the number of elements *a* in $\sigma^{\mathcal{A}}$ that satisfy $s^{\mathcal{A}}(a) = a$. Take as well a set *B* with $\max\{0, \varsigma(m) - |vars_{\sigma}(\phi)^{\mathcal{A}}|\}$ elements.

We define an interpretation \mathcal{B} such that: $\sigma^{\mathcal{B}} = vars_{\sigma}(\phi)^{\mathcal{A}} \cup B$ and $\sigma_2^{\mathcal{B}} = A^+$ (so $|\sigma^{\mathcal{B}}| = \varsigma(m)$); $s^{\mathcal{B}}(a) = s^{\mathcal{A}}(a)$ for each $a \in A$, and $s^{\mathcal{B}}(a)$ is anything but a for $a \in A^+ \setminus A$ (so the number of elements $a \in \sigma^{\mathcal{B}}$ satisfying $s^{\mathcal{B}}(a) = a$ is m, making of \mathcal{B} a $\mathcal{T}_{\varsigma}^{=}$ -interpretation); and $x^{\mathcal{B}} = x^{\mathcal{A}}$ for any variable x in ϕ , and arbitrary otherwise.

This way \mathcal{B} satisfies ϕ , and has both $\sigma^{\mathcal{B}}$ and $\sigma_2^{\mathcal{B}}$ finite. Furthermore, $\sigma_2^{\mathcal{B}} \subseteq \sigma_2^{\mathcal{A}}$, so $|\sigma_2^{\mathcal{B}}| \leq |\sigma_2^{\mathcal{A}}|$, and since the number of elements $a \in \sigma^{\mathcal{B}}$ satisfying $s^{\mathcal{B}}(a) = a$ is m, equal to or less than the number m' of elements $a \in \sigma^{\mathcal{A}}$ satisfying $s^{\mathcal{A}}(a) = a$, we have that $|\sigma^{\mathcal{B}}| = \varsigma(m) \leq \varsigma(m') \leq |\sigma^{\mathcal{A}}|$, finishing the proof.

M.10 T_c^2

All proofs for $\mathcal{T}_{\varsigma}^2$ are very similar to those of $\mathcal{T}_{\varsigma}^\infty$ in Appendix M.2, with the exception of the proof that $\mathcal{T}_{\varsigma}^2$ is not stably finite.

Lemma 90. The Σ_s^2 -theory \mathcal{T}_s^2 is not stably finite with respect to $\{\sigma, \sigma_2\}$.

Proof. Obvious, since $\mathcal{T}_{\varsigma}^2$ has interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}| = 1$ and $|\sigma_2^{\mathcal{A}}| = \omega$, but no interpretations \mathcal{B} with $|\sigma^{\mathcal{B}}| = 1$ and $\sigma_2^{\mathcal{B}}$ finite.

M.11 $\mathcal{T}_{s\vee}^{=}$

The proofs that $\mathcal{T}_{\varsigma\vee}^{=}$ is smooth (and therefore stably infinite), not finitely witnessable (and thus not strongly finitely witnessable) and has the finite model property are very similar to the corresponding proofs of $\mathcal{T}_{\varsigma}^{=}$ in Appendix M.9. The proof that $\mathcal{T}_{\varsigma\vee}^{=}$ is not convex follows the proof that $\mathcal{T}_{\varsigma\vee}^{\vee}$ is not convex, in Lemma 84; and, finally, the proof that $\mathcal{T}_{\varsigma\vee}^{=}$ is not stably finite, is almost the same as that of Lemma 90.

M.12 $\mathcal{T}_{s\neq}^{\infty}$

Lemma 91. The Σ_s^2 -theory $\mathcal{T}_{s\neq}^{\infty}$ is not stably infinite, and thus not smooth, with respect to $\{\sigma, \sigma_2\}$.

Proof. Obvious: while s(x) = x is satisfied by any $\mathcal{T}_{\varsigma\neq}^{\infty}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = 1$ and $|\sigma_2^{\mathcal{A}}| = \omega$, all $\mathcal{T}_{\varsigma\neq}^{\infty}$ -interpretations \mathcal{B} with $|\sigma^{\mathcal{A}}|$ greater than one satisfy instead $\forall x. \neg(s(x) = x)$.

Lemma 92. The Σ_s^2 -theory $\mathcal{T}_{\varsigma\neq}^{\infty}$ is not finitely witnessable, and thus not strongly finitely witnessable, with respect to $\{\sigma, \sigma_2\}$.

Proof. The proof is similar to that of Lemma 53: if there is a witness *wit*, define a function $f : \mathbb{N} \to \mathbb{N}$ by making f(0) = 0, f(1) = 1, and assuming f(n) defined,

$$f(n+1) = |vars_{\sigma}(wit(\bigwedge_{i=1}^{f(n)}\bigwedge_{j=i+1}^{f(n)+1}\neg(x_i = x_j)))|,$$

where x_1 through $x_{f(n)+1}$ are variables of sort σ . With this, f is supposed to be computable, but grows faster than ς , contradicting Lemma 48.

Lemma 93. The Σ_s^2 -theory $\mathcal{T}_{\varsigma\neq}^{\infty}$ is convex with respect to $\{\sigma, \sigma_2\}$.

Proof. This proof is very similar to the one of Lemma 41. We start by taking a Σ_s^2 -cube ϕ , and assuming that $\vdash_{\mathcal{T}_{\varsigma\neq}^{\infty}} \phi \to \bigvee_{i=1}^n x_i = y_i$. We can also assume that ϕ is true in some $\mathcal{T}_{\varsigma\neq}^{\infty}$ -interpretation \mathcal{A}_0 with $|\sigma^{\mathcal{A}_0}| > 1$: otherwise, given the theory whose interpretations are all $\mathcal{T}_{\varsigma\neq}^{\infty}$ -interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}| = 1$ is essentially \mathcal{T}_{∞} , which is convex, we would be done; we can take $|\sigma^{\mathcal{A}_0}| = \omega$ by Lemma 2.

Now, consider the theory whose interpretations are all $\mathcal{T}_{\varsigma\neq}^{\infty}$ -interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}| > 1$, and since it is stably infinite, and convex by Lemma 40, we have that there exists $1 \leq i \leq n$ such that $\phi \to x_i = y_i$ all $\mathcal{T}_{\varsigma\neq}^{\infty}$ -interpretations \mathcal{A} with $|\sigma^{\mathcal{A}}| > 1$; in particular, by \mathcal{A}_1 . But suppose there is a $\mathcal{T}_{\varsigma\neq}^{\infty}$ -interpretation \mathcal{A}_1 with $|\sigma^{\mathcal{A}_1}| = 1$, and without loss of generality $|\sigma_2^{\mathcal{A}_1}| = \omega$, that satisfies ϕ and not $x_i = y_i$: we then build a $\mathcal{T}_{\varsigma\neq}^{\infty}$ -interpretation \mathcal{B} with $\sigma^{\mathcal{B}} = \sigma^{\mathcal{A}_0}$, $\sigma_2^{\mathcal{B}} = \sigma_2^{\mathcal{A}_1}$ and $s^{\mathcal{B}} = s^{\mathcal{A}_0}$ that satisfies ϕ and not $x_i = y_i$, leading to a contradiction.

Lemma 94. The Σ_s^2 -theory $\mathcal{T}_{\varsigma\neq}^{\infty}$ has the finite model property with respect to $\{\sigma, \sigma_2\}$.

Proof. Follows the steps of Lemma 57.

Lemma 95. The Σ_s^2 -theory $\mathcal{T}_{\varsigma\neq}^{\infty}$ is not stably finite with respect to $\{\sigma, \sigma_2\}$.

Proof. Obvious: while there is a $\mathcal{T}_{\varsigma\neq}^{\infty}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = 1$ and $|\sigma_{2}^{\mathcal{A}}| = \omega$, there are no $\mathcal{T}_{\neq}^{\infty}$ -interpretations \mathcal{B} with $|\sigma^{\mathcal{B}}| = 1$ and $|\sigma_{2}^{\mathcal{B}}| < \omega$.

M.13 $T_c^{\infty,3}$

Lemma 96. The Σ_3 -theory $\mathcal{T}_{\varsigma}^{\infty,3}$ is not stably infinite, and thus not smooth, with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. Obvious, since it is not contradictory, but has no models \mathcal{A} with $|\sigma_3^{\mathcal{A}}|$ infinite.

Lemma 97. The Σ_3 -theory $\mathcal{T}_{\varsigma}^{\infty,3}$ has the finite model property with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. Essentially the same proof as that of Lemma 57.

Lemma 98. The Σ_3 -theory $\mathcal{T}_{\varsigma}^{\infty,3}$ is not stably infinite with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. Consider any quantifier-free Σ_3 -formula ϕ that is satisfied by the $\mathcal{T}_{\varsigma}^{\infty,3}$ -interpretation \mathcal{A} with $|\sigma^{\mathcal{A}}| = |\sigma_3^{\mathcal{A}}| = 1$ and $|\sigma_2^{\mathcal{A}}| = \omega$: since there are no $\mathcal{T}_{\varsigma}^{\infty,3}$ -interpretations \mathcal{B} such that $|\sigma^{\mathcal{B}}| = 1$ and $|\sigma_2^{\mathcal{B}}|$ is finite, $\mathcal{T}_{\varsigma}^{\infty,3}$ cannot be stably finite.

Lemma 99. The Σ_3 -theory $\mathcal{T}_{\varsigma}^{\infty,3}$ is not finitely witnessable, and thus not strongly finitely witnessable, with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. See Lemma 59 for an essentially equal proof.

Lemma 100. The Σ_3 -theory $\mathcal{T}_{\varsigma}^{\infty,3}$ is convex with respect to $\{\sigma, \sigma_2, \sigma_3\}$.

Proof. Take a conjunction of literals ϕ , and suppose $\vdash_{\mathcal{T}_{\varsigma}^{\infty,3}} \phi \to \bigvee_{i=1}^{n} x_i = y_i$: if any pair (x_i, y_i) is of sort σ_3 we are done, since then $\vdash_{\mathcal{T}_{\varsigma}^{\infty,3}} \phi \to x_i = y_i$. So we may assume that some all x_i and y_i are of sorts in $\{\sigma, \sigma_2\}$. We can also assume no literal of ϕ has variables of sort σ_3 , since then ϕ is either a tautology or a contradiction, cases very easy to consider.

So ϕ , under these assumptions, is actually a Σ_2 -formula, as well as $\bigvee_{i=1}^n x_i = y_i$: not only that, but $\vdash_{\mathcal{T}^{\infty}_{\zeta}} \phi \to \bigvee_{i=1}^n x_i = y_i$; indeed, if there is a $\mathcal{T}^{\infty}_{\zeta}$ -interpretation \mathcal{A} that satisfies ϕ but not $\bigvee_{i=1}^n x_i = y_i$, the $\mathcal{T}^{\infty,3}_{\zeta}$ -interpretation \mathcal{B} with $\sigma^{\mathcal{B}} = \sigma^{\mathcal{A}}$ and $\sigma^{\mathcal{B}}_2 = \sigma^{\mathcal{A}}_2$ would satisfy ϕ but not $\bigvee_{i=1}^n x_i = y_i$. But $\mathcal{T}^{\infty}_{\zeta}$ is convex, as proven in Lemma 60, and so $\vdash_{\mathcal{T}^{\infty}_{\zeta}} \phi \to x_i = y_i$ for some $1 \leq i \leq n$. Of course it follows that $\vdash_{\mathcal{T}^{\infty,3}_{\zeta}} \phi \to x_i = y_i$, and so $\mathcal{T}^{\infty,3}_{\zeta}$ is convex.