### DANIEL S. FREED

ABSTRACT. The anomaly of a quantum field theory is an expression of its projective nature. This starting point quickly leads to its manifestation as a special kind of field theory: a once-categorified invertible theory. We arrive at this statement through a general discussion of projectivity and a discussion of projectivity in quantum mechanics. We conclude with a general formula for the anomaly of a free spinor field.

#### CONTENTS

1. Projective spaces, linearization, and symmetry	2
2. Quantum mechanics as a projective system	Ę
3. Quantum field theory as a projective system	8
4. Anomalies as an obstruction to quantization	13
5. Anomaly of a spinor field	15
Appendix A. On the definition of <b>Proj</b>	17
References	18

This note<sup>1</sup> is based on the thesis:

## (A) QUANTUM THEORY IS PROJECTIVE. QUANTIZATION IS LINEAR.

Hence our answer to the titular question:

As such, the anomaly is a feature, not a bug [tH]. It is an obstruction only when quantizing (§4).

Anomalies in quantum field theory have a long history, dating back to work of Steinberger [St], Adler [Ad], and Bell-Jackiw [BJ]; see [AgVm] for a recent survey and many additional references. (We do not attempt an account of that history here.) The idea that an anomaly enjoys the locality properties of a field theory was not explicit in the literature of the 20th century, at least to my knowledge, though the "inflow" mechanism of Callen-Harvey [CH] is a precursor. The first

Date: July 14, 2023.

This material is based on work supported by the National Science Foundation under Grant Number DMS-2005286 and by the Simons Foundation Award 888988 as part of the Simons Collaboration on Global Categorical Symmetries. I thank my longtime collaborators Mike Hopkins, Greg Moore, Constantin Teleman and many others for influential conversations about anomalies, which have taken place over a long period. I also thank Theo Johnson-Freyd for detailed feedback on an earlier draft.

<sup>&</sup>lt;sup>1</sup>This is a written account of online talks given in February, 2023 in Arthur Jaffe's *Picture Language* seminar and in Roger Picken's TQFT Club seminar.

indications for me came from reading [W1]. This and many more influences led to [FT, F1], in which anomalies appear as invertible field theories. (The first brief mention of this formulation is at the end of [FHT].) Nonetheless, this previous work does not tie anomalies in quantum field theory to projectivity, which is the main point of this note. Significantly, the new viewpoint in terms of projective field theories leads us to define an anomaly as a *once-categorified invertible field theory*.

Two myths about anomalies. First, it is often said that anomalies are only associated to symmetries. However, the theory of a free spinor field (and no other fields) has an anomaly. For example, an anomaly is present in the minimal 1-dimensional theory with a single spinor field, and there is no obvious symmetry to which to ascribe the anomaly. Second, one might think that anomalies are only caused by fermionic fields. Not so! For example, the flavor symmetry of QCD is anomalous—indeed, that anomaly involves fermions—but the anomaly persists in the effective theory of pions, which is a bosonic theory in the sense that there are no fluctuating fermionic fields.

# 1. Projective spaces, linearization, and symmetry

We review the interplay between linear and projective geometry, emphasizing the structure of symmetries. The key concept is the *projectivity* of a projective representation. We conclude with a discussion of the question: *What is projective space*?

**Projectivization of a linear space.** Let W be a vector space. For the application to quantum theory we work over the field of complex numbers, but the exposition generalizes to arbitrary fields. For simplicity, assume that W is finite dimensional. Associated to W are two objects of a "projective nature". The projective space  $\mathbb{P}(W)$  is the space of lines  $L \subset W$ . (A line is a 1-dimensional linear subspace.) The algebra  $\operatorname{End}(W)$  is the space of linear maps  $T: W \to W$ . One expression of the projective nature of  $\mathbb{P}(W)$  and  $\operatorname{End}(W)$  is that they do not change when W is replaced by the tensor product  $W \otimes K$  with a line K. Namely, there are *canonical* isomorphisms

(1.1) 
$$\begin{array}{ccc} \mathbb{P}(W) \longrightarrow \mathbb{P}(W \otimes K) & \operatorname{End}(W) \longrightarrow \operatorname{End}(W \otimes K) \\ L & \longmapsto & L \otimes K & T & \longmapsto & T \otimes \operatorname{id}_K \end{array}$$

These isomorphisms suggest that the algebra  $\operatorname{End}(W)$  is canonically associated to the projective space  $\mathbb{P}(W)$  in place of the vector space W, a notion we explain in footnote <sup>4</sup> below.

**Projective symmetries.** A linear symmetry of W induces a projective symmetry of  $\mathbb{P}(W)$ , and conversely a projective symmetry of  $\mathbb{P}(W)$  has a  $\mathbb{C}^{\times}$ -torsor<sup>2</sup> of lifts to a linear symmetry of W. The relationship between linear and projective symmetries is encoded in the short exact sequence of Lie groups

$$(1.2) \qquad \qquad \mathbb{C}^{\times} \longrightarrow \mathrm{GL} \longrightarrow \mathrm{PGL}$$

<sup>&</sup>lt;sup>2</sup>A  $\mathbb{C}^{\times}$ -torsor is a manifold equipped with a simply transitive action of  $\mathbb{C}^{\times}$ . Any projective symmetry lifts to a linear symmetry, unique up to composition with a *homothety*: scaling by a nonzero complex number  $\lambda \in \mathbb{C}^{\times}$ .

which is a central extension. We use notation for the finite dimensional model space  $W = \mathbb{C}^{n+1}$ , though we drop the dimension n; with appropriate topologies these can also be taken to be groups of symmetries of infinite dimensional spaces. A Lie group G of *projective* symmetries is specified by a Lie group homomorphism  $G \to PGL$ . It induces a pull back central extension of G

(1.3) 
$$\begin{array}{c} \mathbb{C}^{\times} \longrightarrow \operatorname{GL} \longrightarrow \operatorname{PGI} \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ \mathbb{C}^{\times} \longrightarrow \widetilde{G} \longrightarrow G \end{array}$$

together with a *linear* action of the centrally extended group  $\widetilde{G}$ .

A linearization of the projective action is a lift of  $G \to PGL$  to a homomorphism  $G \to GL$ . A linearization is equivalent to a *splitting* of the induced central extension:

(1.4) 
$$\begin{array}{c} \mathbb{C}^{\times} \longrightarrow \operatorname{GL} \longrightarrow \operatorname{PGL} \\ \| & & \\ & & \\ \mathbb{C}^{\times} \longrightarrow \widetilde{G} \xrightarrow{} G \end{array}$$

The *obstruction* to a linearization is measured by a homomorphism  $\alpha$  from G to a classifying space for the center  $\mathbb{C}^{\times}$ :

(1.5) 
$$\begin{array}{c} \mathbb{C}^{\times} \longrightarrow \operatorname{GL} \longrightarrow \operatorname{PGL} \longrightarrow B\mathbb{C}^{\times} \\ \| & \stackrel{\wedge}{\underset{I}{\longrightarrow}} & \stackrel{\wedge}{\underset{I}{\longrightarrow}} & \stackrel{\wedge}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow}} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{}{\underset{I}{\longrightarrow} & \stackrel{\circ}{\underset{I}{\longrightarrow} & \stackrel{}{\underset{I}{\longrightarrow} & \stackrel{}{\underset{I}{\longrightarrow} & \stackrel{}{\underset{I}{\longrightarrow}} & \stackrel{}{\underset{I}{\longrightarrow} & \stackrel{}{\underset{I}{\longrightarrow} & \stackrel{}{\underset{I}{\longrightarrow}{\longrightarrow} & \stackrel{}{\underset{I}{\longrightarrow} & \stackrel{}{\underset{I}{\longrightarrow} & \stackrel{}{\underset{I}{\longrightarrow} &$$

One geometric model for  $B\mathbb{C}^{\times}$  is the category of  $\mathbb{C}^{\times}$ -torsors and isomorphisms of  $\mathbb{C}^{\times}$ -torsors. The category  $B\mathbb{C}^{\times}$  is a groupoid—all morphisms are invertible—and furthermore it carries a symmetric tensor product. It is in this sense that we have a homomorphism  $\alpha \colon G \to B\mathbb{C}^{\times}$ : for each  $g \in G$  there is a  $\mathbb{C}^{\times}$ -torsor  $L_g$ ; for each pair  $g_1, g_2 \in G$  there is an isomorphism  $L_{g_1} \otimes L_{g_2} \to L_{g_1g_2}$ ; and for each triple  $g_1, g_2, g_3 \in G$  there is a compatibility (associativity, cocycle condition) among these isomorphisms. In short, a homomorphism  $G \to B\mathbb{C}^{\times}$  is a central extension of G by  $\mathbb{C}^{\times}$ . The torsor  $L_g$  is  $\pi^{-1}(g)$  in the central extension. The homomorphism  $\alpha \colon G \to B\mathbb{C}^{\times}$  is the projectivity of the projective representation; it measures the deviation from linearizability.

One can turn the story around. Begin with the projectivity  $\alpha$ :

and then define a projective representation with projectivity  $\alpha$  to be a homomorphism  $\bar{\rho} \colon G \to \text{PGL}$  that makes the diagram

 $B\mathbb{C}^{\times}$ 



commute. As argued above, the projectivity  $\alpha$  is equivalent to a central extension of G by  $\mathbb{C}^{\times}$ , and then a projective representation  $\bar{\rho}$  of G is equivalent to a linear representation  $\rho$  of  $\tilde{G}$  in which the central  $\mathbb{C}^{\times}$  acts by scalar multiplication:



A linearization of  $\bar{\rho}$  is equivalent to a splitting of the central extension, which in other terms is a *trivialization* of the projectivity  $\alpha$ :

(1.9) 
$$\begin{array}{c} \mathbb{C}^{\times} \longrightarrow \operatorname{GL} \longrightarrow \operatorname{PGL} \longrightarrow B\mathbb{C}^{\times} \\ & & & & & & \\ & & & & & & & \\ \mathbb{C}^{\times} \longrightarrow \widetilde{G} \longrightarrow G \xrightarrow{\overline{\rho}} & & & & \\ & & & & & & & \\ \end{array}$$

Cohomological interpretation. The homotopy class or isomorphism class (the appropriate phrase depends on the model of  $B\mathbb{C}^{\times}$ ) of the projectivity  $\alpha$  lies in a cohomology group  $H^2(G; \mathbb{C}^{\times})$ . The precise type of cohomology depends on the nature of the group G. For example, if G is a finite group, then we can use standard group cohomology of Eilenberg-MacLane, which is equivalent to the singular cohomology of a space that realizes the classifying space BG. If G is a Lie group, then a different type of cohomology is needed. As intimated several times now, the central extension

is a "cocycle" for the cohomology class. Splittings of the central extension—equivalently trivializations of the projectivity  $\alpha$ —form a torsor over characters  $\lambda$  of G:

$$(1.11) \qquad \qquad \mathbb{C}^{\times} \xrightarrow{\widetilde{G}} G \xrightarrow{\alpha} 1$$

Characters are *invertible* linear representations, or in terms of cohomology, elements of  $H^1(G; \mathbb{C}^{\times})$ . Recall that the class of the projectivity lies in the next higher cohomology group  $H^2(G; \mathbb{C}^{\times})$ . This leads to the main takeaway:

(1.12) The projectivity of a projective representation is a "suspended" or "delooped" or "once-categorified" invertible linear representation.

What is a projective space? In other words, how do we define a projective space without expressing it as the projectivization  $\mathbb{P}(W)$  of a linear space W? Our answer is a special case of a definition of Klein-Cartan. A model geometry  $H \subset X$  consists of a smooth manifold X equipped with a left action of a Lie group H. For example, the model for *n*-dimensional complex linear geometry is  $\operatorname{GL}_n \mathbb{C} \subset \mathbb{C}^n$ . The model for *n*-dimensional hyperbolic geometry is  $\operatorname{O}_{n,1} \subset \mathbb{H}^n$ , where  $\mathbb{H}^n$  is hyperbolic space. An instance of  $(H \subset X)$ -geometry is specified by a right H-torsor T; the associated manifold

carries this geometry. Here  $T \times_H X$  is the mixing construction:  $T \times_H X = (T \times X)/H$ , where  $h \in H$  transforms  $(t, x) \in T \times X$  by  $(t, x) \cdot h = (t \cdot h, h^{-1} \cdot x)$ . (Example: for *n*-dimensional complex linear geometry,  $X_T$  is a complex vector space and T is the right  $\operatorname{GL}_n\mathbb{C}$ -torsor of isomorphisms  $\mathbb{C}^n \xrightarrow{\cong} X_T$ : bases of  $X_T$ .) If S is a smooth manifold, then a family of  $(H \subset X)$ -geometries is specified by a principal H-bundle  $P \to S$ . The geometry lives on the associated fiber bundle  $X_P \to S$ . Furthermore, we allow S to be a smooth stack. Example: if S = \*//G for a Lie group G, then an  $(H \subset X)$ -geometry over S is a single  $(H \subset X)$ -geometry equipped with a G-action.

With this in mind, the model for *n*-dimensional complex projective geometry has  $X = \mathbb{CP}^n$ . There are several possibilities for the group H. For complex algebraic geometry  $H = \mathrm{PGL}_{n+1}\mathbb{C}$  is the group of all holomorphic automorphisms of  $\mathbb{CP}^n$ . In Kähler geometry  $H = \mathrm{PU}_{n+1}$  is the group of holomorphic isometries of  $\mathbb{CP}^n$ . There are infinite dimensional analogs. We will see below that neither is the correct choice for the model geometry for quantum mechanics.

## 2. Quantum mechanics as a projective system

We begin with the usual linear description of quantum mechanics, and then point out its projective nature. Wigner's theorem is used to determine the correct model geometry for quantum mechanics.

Quantum mechanics as a linear system. Typically, one gives the state space  $\mathcal{H}$ , assumed to be a complex separable Hilbert space, together with a self-adjoint operator H on  $\mathcal{H}$ , the Hamiltonian. From this data one defines the space  $\mathbb{P}\mathcal{H}$  of pure states, a convex space of all states, and an algebra of observables. The space of pure states comes equipped with a function

(2.1) 
$$p: \mathbb{PH} \times \mathbb{PH} \longrightarrow [0,1]$$
$$L_0, L_1 \longmapsto |\langle \psi_0, \psi_1 \rangle|^2$$

that encodes transition probabilities between pure states. Here  $\psi_i \in L_i$ , i = 0, 1, are unit norm vectors in the lines  $L_0, L_1 \in \mathbb{PH}$ .

Suppose given times (real numbers)  $t_0 < t_1 < \cdots < t_n < t_f$ , initial and final pure states  $L_0, L_f \in \mathbb{PH}$ , and observables<sup>3</sup>  $A_1, \ldots, A_n \in \text{End}(\mathcal{H})$ . Then the basic quantity in the theory is

(2.2) 
$$p\left(L_f, e^{-i(t_f - t_n)H/\hbar} A_n \cdots e^{-i(t_2 - t_1)H/\hbar} A_1 e^{-i(t_1 - t_0)H/\hbar} L_0\right) \in [0, 1]$$

This probability is the norm square of a complex amplitude. If we choose unit vectors  $\psi_0 \in L_0$ ,  $\psi_f \in L_f$ , then the amplitude is the complex number

(2.3) 
$$\langle \psi_f, e^{-i(t_f - t_n)H/\hbar} A_n \cdots e^{-i(t_2 - t_1)H/\hbar} A_1 e^{-i(t_1 - t_0)H/\hbar} \psi_0 \rangle_{\mathcal{H}} \in \mathbb{C}$$

Of course, we should write the amplitude in terms of the states  $L_0, L_f$ , not in terms of the auxiliary choices  $\psi_0, \psi_f$ . This is straightforward: as a function of  $\psi_0, \psi_f$ , (2.3) is an element of the hermitian line  $(\overline{L_f} \otimes L_0)^* \cong L_f \otimes \overline{L_0}$ .

It is striking that ordinary amplitudes in quantum mechanics are elements of complex lines rather than complex numbers.

Quantum mechanics as a projective system. As explained in §1, the space  $\mathbb{PH}$  of pure states and the space  $\mathrm{End}(\mathcal{H})$  of observables only depend on the projective space that underlies the linear space  $\mathcal{H}$ . Therefore, suppose a projective space  $\mathbb{P}$  in the sense of (1.13) is given. (Shortly we determine precisely what model geometry we need.) Then  $\mathbb{P}$  is the space of pure states of a quantum mechanical system, and there is an associated algebra  $\mathscr{A}_{\mathbb{P}}$  of observables.<sup>4</sup> The transition probability function

(2.4) 
$$p: \mathbb{P} \times \mathbb{P} \longrightarrow [0,1]$$
$$\sigma_0, \sigma_1 \longmapsto |\langle \psi_0, \psi_1 \rangle|_{\mathcal{H}}^2$$

is defined by choosing a linearization  $\mathbb{P} \xrightarrow{\cong} \mathbb{P}\mathcal{H}$  and choosing unit norm vectors  $\psi_0, \psi_1$  in the lines of  $\mathcal{H}$  that correspond to  $\sigma_0, \sigma_1$ . One checks<sup>5</sup> that (2.4) is independent of these choices.

With this setup in place, the analog of the probability (2.2) is

(2.5) 
$$p\left(\sigma_f, e^{-i(t_f - t_n)H/\hbar} A_n \cdots e^{-i(t_2 - t_1)H/\hbar} A_1 e^{-i(t_1 - t_0)H/\hbar} \sigma_0\right) \in [0, 1]$$

The amplitude (2.3) is also well-defined in this projective setup; it lives in a hermitian line  $\mathcal{L}_f \otimes \overline{\mathcal{L}_0}$  that is defined using (2.3) and the gerbe of linearizations of  $\mathbb{P}$ . As before, the probability (2.5) is the norm square of this amplitude.

<sup>&</sup>lt;sup>3</sup>For us 'End( $\mathcal{H}$ )' denotes *bounded* operators, and for this formal discussion we restrict to bounded observables.

<sup>&</sup>lt;sup>4</sup>If  $H \subset X$  is the model geometry, with  $X = \mathbb{CP} = \mathbb{P}(\mathcal{H})$  a model complex projective space of finite or infinite dimension, so  $\mathcal{H} = \mathbb{CP}^N$  or  $\mathcal{H}$  is a standard infinite dimensional Hilbert space; and if T is the right H-torsor that specifies the projective space  $\mathbb{P} = T \times_H \mathbb{CP}$ ; then  $\mathscr{A}_{\mathbb{P}} = T \times_H \operatorname{End}(\mathcal{H})$ , where H acts by conjugation on  $\operatorname{End}(\mathcal{H})$ .

<sup>&</sup>lt;sup>5</sup>Use that H = PQ, and that a linearization of the projective space associated to a PQ-torsor T is associated to a lift of T to a Q-torsor. (The Lie groups Q and PQ are defined in the next subsection.) The collection of lifts, and so of linearizations, forms a  $\mathbb{T}$ -gerbe. See footnote <sup>7</sup> below for one model of a gerbe.

The symmetry/structure group of quantum mechanics. This is the group PQ of permutations of the points of a projective space  $\mathbb{P}$  that preserve the function (2.4), which is the basic data of quantum mechanics. Fix a linearization  $\mathbb{P} \xrightarrow{\cong} \mathbb{PH}$ ; then the group  $\mathrm{PQ} = \mathrm{Aut}(\mathbb{P}, p)$  of maps  $\mathbb{P} \longrightarrow \mathbb{P}$ preserving p is the isometry group of the Fubini-Study metric  $d: \mathbb{PH} \times \mathbb{PH} \longrightarrow \mathbb{R}^{\geq 0}$ . This follows from the well-known formula (see [BH] for a discussion and references)

$$(2.6)\qquad\qquad\qquad\cos(d)=2p-1$$

The structure group PQ we seek is the isometry group of a model Fubini-Study projective space.

**Example 2.7.** If  $\mathcal{H} = \mathbb{C}^2$ , then  $\mathbb{P} = \mathbb{CP}^1$  is the complex projective line, which is diffeomorphic to the 2-sphere  $S^2$ , and the Fubini-Study metric transports to the round metric on  $S^2$ . The isometry group in this case is the orthogonal group O<sub>3</sub>. Observe the group extensions

$$\begin{array}{cccc} \mathbb{T} & \longrightarrow & \mathrm{U}_2 \longrightarrow & \mathrm{SO}_3 \\ \\ \mathbb{T} & \longrightarrow & \mathrm{Q}_2 \longrightarrow & \mathrm{O}_3 = \mathrm{PQ}_2 \end{array}$$

with kernel the group  $\mathbb{T} \subset \mathbb{C}^{\times}$  of unit norm complex numbers. The first is a central extension of the identity component SO<sub>3</sub> of the Lie group PQ<sub>2</sub> = O<sub>3</sub>. Elements of SO<sub>3</sub> act projectively on  $\mathbb{C}^2$ , and U<sub>2</sub> is the group of lifts to isometries of  $\mathbb{C}^2$ . The second extension in (2.8) is not central. The extended group Q<sub>2</sub> consists of unitary and antiunitary automorphisms of  $\mathbb{C}^2$ , and antiunitary automorphisms anticommute with elements of the kernel  $\mathbb{T}$ .

The following result, known as Wigner's theorem, is undoubtedly due to Cartan for finite dimensional projective spaces (in the form of determining the isometry group; see [Lo]), and in the infinite dimensional case appears first in [vNW] without proof; see [Bo] for a history and [F2] for two geometric proofs and references to earlier proofs. Let  $\mathbb{P}$  be the projectivization of  $\mathcal{H} = \mathbb{C}^N$  in the finite dimensional case and of a model complex separable Hilbert space  $\mathcal{H}$  in the infinite dimensional case. Let PQ denote the group of automorphisms of  $\mathbb{P}\mathcal{H}$  that preserve the function (2.4).

**Theorem 2.9** (von Neumann–Wigner). The group PQ of projective quantum mechanical symmetries fits into a group extension

where Q is the group of unitary and antiunitary transformations of  $\mathcal{H}$ .

The import of the theorem is the surjectivity of the map  $Q \to PQ$ . Each of Q and PQ is a Lie group with two components; the identity component of Q is the unitary group. Let  $Q_N, PQ_N$  be these Lie groups for  $\mathcal{H} = \mathbb{C}^N$ , and let  $Q_{\infty}, PQ_{\infty}$  be these groups in the infinite dimensional case.

This discussion is summarized by the statement  $(n \in \mathbb{Z}^{>0} \cup \{\infty\})$ :

## 3. Quantum field theory as a projective system

Now we arrive at the central section of the paper. We begin by reviewing the Segal axioms for Wick-rotated field theory, in which a field theory is a linear representation of a bordism category. Then, following the main theme (A), we indicate the variation in which a field theory is a projective representation of a bordism category. The anomaly is the projectivity of that representation, and our main conclusion—analogous to (1.12)—is that the anomaly is a once-categorified invertible field theory. We argue that the physical content of a quantum field theory is contained in a projective theory rather than a linear theory. In most cases we know, the anomaly extends to a one higher dimensional ordinary invertible field theory, as we indicate at the end of this section.

Wick-rotated field theory as a linear representation. Segal [S1] first introduced his axiom system in the context of 2-dimensional conformal field theory. It was taken over to topological field theory by Atiyah [A], and subsequently extended in many directions. Over time it became clear that the axioms should apply to general quantum field theories. This general context is the subject of a recent paper by Kontsevich-Segal [KS]. For our purposes here, we emphasize the formal structure.

There are two discrete parameters that specify the "type" of field theory. The first is a positive integer n, which in Lorentz signature is the dimension of spacetime. The second is the collection  $\mathcal{F}$  of background fields. (In this axiom system there are no fluctuating fields—they have already been integrated out—but see §4.) Let  $\mathbf{Man}_n$  denote the category of smooth *n*-manifolds and local diffeomorphisms, and let **sSet** denote the category of simplicial sets.

Definition 3.1. A Wick-rotated field is a sheaf

$$(3.2) \qquad \qquad \mathcal{F}\colon \mathbf{Man}_n^{\mathrm{op}} \longrightarrow \mathbf{sSet}$$

One needs extra structure on  $\operatorname{Man}_n$  to formulate the sheaf condition, which as usual is in terms of open covers; see [FH1] for one exposition. Some fields take values in the category of sets: Riemannian metrics,  $\mathbb{R}$ -valued functions, N-valued functions for a fixed smooth manifold N, orientations, etc. Others take values in groupoids: G-connections for a fixed Lie group G, spin structures, etc. Still others take values in higher groupoids: B-fields, C-fields, etc. The choice of **sSet** as codomain in (3.2) accommodates all of these. In our usage,  $\mathcal{F}$  is a finite set of fields, such as  $\mathcal{F} = \{ \text{orientations, Riemannian metrics} \}$ . For an n-manifold M, the space of sections  $\mathcal{F}(M)$  of the sheaf is the space (simplicial set) of fields on M.

Given  $n, \mathcal{F}$ , we define a bordism category  $\mathbf{Bord}_{\langle n-1,n\rangle}(\mathcal{F})$ . An object is a closed (n-1)manifold Y, embedded in a germ  $\hat{Y}$  of an n-manifold, equipped with a coorientation in that germ, and equipped with an object of  $\mathcal{F}(\hat{Y})$ . (The coorientation is the Wick-rotated remnant of an "arrow of time" of a spacelike hypersurface.) A morphism  $X: Y \to Y'$  is a compact n-manifold with boundary equipped with a partition into incoming and outgoing components, a germ  $\hat{X}$  of an embedding of the boundary into an n-manifold that extends the germ of a collar neighborhood of the boundary, isomorphisms of the incoming boundary germs with  $\hat{Y}$  and the outgoing boundary germs with  $\hat{Y}'$ , and an object of  $\mathcal{F}(\hat{X})$  together with isomorphisms of its restrictions to the boundary germs with the background fields on  $\hat{Y}$  and  $\hat{Y}'$ . A sample object and morphism are depicted in Figure 1. Composition is by gluing, and disjoint union provides a symmetric monoidal structure.



FIGURE 1. An object and a morphism in the bordism category  $\mathbf{Bord}_n(\mathcal{F})$ . Germs of *n*-manifolds are represented by the shadings, and coorientations by the red arrows. Background fields are not depicted.

Remark 3.3. A field or collection of fields (3.2) is topological if the sheaf that defines it is locally constant. Examples include orientations, spin structures, G-bundles for a finite group G, etc. Non-examples include  $\mathbb{R}$ -valued functions, Riemannian metrics, G-connections for a positive dimensional Lie group G, etc. A locally constant sheaf is equivalent to an n-dimensional tangential structure in the sense introduced into bordism theory by Lashof [La]. For topological fields the bordism category was first introduced by Milnor in his proof of the h-cobordism theorem [M].

Let **Vect** denote a suitable category of topological vector spaces and linear maps; see [KS,  $\S3$ ] for details. The important point is that the linear maps are *nuclear*, which is a form of smallness.<sup>6</sup>

Axiom System 3.4. A (Wick-rotated) field theory

is a linear representation of a bordism category.

Here 'linear representation' means a symmetric monoidal functor of symmetric monoidal categories, which we simply call a 'homomorphism'. So F assigns a vector space (state space) F(Y) to every closed (n-1)-manifold and a linear map  $F(X): F(Y) \to F(Y')$  to every bordism  $X: Y \to Y'$ .

Remark 3.6.

- (1) A field theory is *topological* if it factors through a bordism category  $\mathbf{Bord}_{\langle n-1,n\rangle}(\mathcal{F}')$  in which  $\mathcal{F}'$  is a locally constant sheaf.
- (2) For topological theories there is an extended notion of *locality*.
- (3) Just as representations of Lie groups are not necessarily unitarizable (example:  $SL_2\mathbb{R}$ ), so too the notion of *unitarity* is not part of Axiom System 3.4. Rather, unitarity appears in the form of reflection positivity, which is a structure on F that we do not elaborate here.

Wick-rotated QFT as a projective representation. As in the case of quantum mechanics ( $\S$ 2), quantum field theory is a projective system, and so (3.5) should take values in a suitable category of projective spaces and projective linear maps.

<sup>&</sup>lt;sup>6</sup>In quantum mechanics, for example, in real time the evolution is by unitary operators  $e^{-itH/\hbar}$ ,  $t \in \mathbb{R}$ , whereas Wick-rotated "evolution" is by smoothing maps  $e^{-\tau H/\hbar}$ ,  $\tau \in \mathbb{R}^{>0}$ , which in particular are nuclear (trace class).

Remark 3.7. At the end of §1 we gave several model geometries for complex projective space. Since unitarity is not built into Axiom System 3.4, the appropriate model for the projective variation of the axioms is that of complex algebraic geometry, namely  $\operatorname{PGL}_{n+1}\mathbb{C} \subset \mathbb{CP}^n$ , where  $n \in \mathbb{Z}^{>0} \cup \{\infty\}$ . (For  $n = \infty$  we use ' $\mathbb{CP}^{\infty}$ ' for the projective space of a distinguished object of **Vect**, in the same sense that  $\mathbb{C}^N$  is a distinguished N-dimensional complex vector space.)

Let **Line** be the category of 1-dimensional complex vector spaces and invertible linear maps between them; it is a categorified version of the group  $\mathbb{C}^{\times}$ . We seek a categorification of the short exact sequence (1.2), which we write as:

$$(3.8) \qquad \qquad \text{Line} \longrightarrow \text{Vect} \longrightarrow \text{Proj}$$

Heuristically, **Proj** is a category of complex projective spaces and projective linear maps, and it should be based on the particular topological vector spaces and continuous linear maps used to define the category **Vect**. More precisely, the Picard groupoid **Line**, which is a model for  $B\mathbb{C}^{\times}$ , acts on **Vect** and **Proj** is the quotient of this action. This quotient is naturally a 2-category, not a 1-category. See Appendix A for more indications of the definition of **Proj**. As in (1.5) there is a classifying space for **Line** and an extension of the sequence (3.8):

$$(3.9) \qquad \qquad \mathbf{Line} \longrightarrow \mathbf{Vect} \longrightarrow \mathbf{Proj} \longrightarrow B\mathbf{Line}$$

We have the ingredients in place to mimic the discussion in §1 that begins with (1.3). Namely, a projective field theory  $\overline{F}$  is a homomorphism (symmetric monoidal functor) into **Proj**:

Its projectivity, or anomaly  $\alpha$ , is the composition with the map to BLine:



Since  $\alpha$  is a map out of a bordism category, it is a field theory. Shortly we identify its precise nature. At this stage we simply emphasize the analogy with the map  $\alpha$  in (1.5), which measures the projectivity of a projective representation of a Lie group. Analogous to (1.4), a projective field theory induces a central extension of a bordism category and a linear representation  $\tilde{F}$  of the centrally extended bordism category:



This version of a projective field theory—a linear representation of a centrally extended bordism category—already appears in Segal's original paper, namely [S1, Definition (5.2)] (which in fact is more general). A trivialization of the anomaly is equivalent to a lift of the projective theory  $\tilde{F}$  to a linear theory F:



Trivializations of  $\alpha$ —if they exist—form a torsor over homomorphisms  $\lambda : \mathbf{Bord}_{(n-1,n)}(\mathcal{F}) \to \mathbf{Line}$ :

(3.14) 
$$\begin{array}{c} \text{Line} \longrightarrow \text{Vect} \longrightarrow \text{Proj} \longrightarrow B\text{Line} \\ & & & & \\ & & & & & \\ & & &$$

Anomaly as a once-categorified invertible field theory. We begin by elucidating the classifying space BLine, which we model as a 2-category, just as  $B\mathbb{C}^{\times}$  is modeled as a 1-category after (1.5). A simple model is the 2-groupoid  $B^2\mathbb{C}^{\times}$  which has a single object, a single 1-morphism, and the group  $\mathbb{C}^{\times}$  of 2-morphisms. A more robust model is the 2-category of  $\mathbb{C}^{\times}$ -gerbes,<sup>7</sup> or equivalently of invertible Vect-modules.

### **Definition 3.15.** Fix $n, \mathcal{F}$ .

(1) An *invertible field theory* is a homomorphism

$$(3.16) \qquad \qquad \lambda \colon \mathbf{Bord}_{(n-1,n)}(\mathcal{F}) \longrightarrow \mathbf{Line}$$

(2) A once-categorified invertible field theory is a homomorphism

$$(3.17) \qquad \qquad \alpha \colon \mathbf{Bord}_{\langle n-1,n \rangle}(\mathfrak{F}) \longrightarrow B\mathbf{Line}$$

Let Y be an object of  $\operatorname{Bord}_{\langle n-1,n\rangle}(\mathcal{F})$ , which is essentially a closed (n-1)-manifold. (The additional data of an object—the germ of a cooriented embedding into an n-manifold together with a background field—is spelled out following Definition 3.1.) Then  $\lambda(Y)$  is a complex line, the 1-dimensional state space of the invertible field theory  $\lambda$  on Y. By contrast,  $\alpha(Y)$  is a gerbe. If  $\alpha$  is the anomaly of a projective theory  $\overline{F}$ , then  $\alpha(Y)$  measures the projectivity of the projective space  $\overline{F}(Y)$ . Now suppose X is a closed n-manifold equipped with a background field. Then  $\lambda(X)$  is a nonzero complex number, whereas  $\alpha(X)$  is a complex line. If  $\overline{F}$  is a projective theory with anomaly  $\alpha$ , then  $\overline{F}(X)$  is an element of the complex line  $\alpha(X)$ . (Recall that amplitudes in quantum mechanics are naturally elements of complex lines—see the text that follows (2.5)—so it is natural that partition functions of quantum field theories are also elements of complex lines.)

<sup>&</sup>lt;sup>7</sup>Here is one model of a  $\mathbb{C}^{\times}$ -gerbe. The data is a set  $\mathcal{A}$ ; for each pair  $\alpha_0, \alpha_1 \in \mathcal{A}$  a complex line  $L_{\alpha_0\alpha_1}$ ; and for each triple  $\alpha_0, \alpha_1, \alpha_2 \in \mathcal{A}$  an isomorphism of lines  $\theta_{\alpha_0,\alpha_1,\alpha_2} \colon \mathbb{C} \xrightarrow{\cong} L_{\alpha_1\alpha_2} \otimes L_{\alpha_0\alpha_2}^{-1} \otimes L_{\alpha_0\alpha_1}$ . For each quartet  $\alpha_0, \alpha_1, \alpha_2, \alpha_3 \in \mathcal{A}$  there is a condition on the  $\theta$ 's. A projective space  $\mathbb{P}$  gives rise to a gerbe in which  $\mathcal{A}$  is a set of linearizations of  $\mathbb{P}$ .

**Interlude: Remarks on the physics of projective theories.** We make two observations that may help locate our picture among more familiar ideas in quantum field theory.

The first is an interpretation of conversations with Nati Seiberg. Namely, one might consider two (linear) quantum field theories F, F' to encode the same physics if there is an invertible theory  $\lambda$  on the same background fields such that<sup>8</sup>  $F' = F \otimes \lambda$ . After all, traditional correlation functions are ratios, which in a path integral take the schematic form

(3.18) 
$$\frac{\int D\phi \ e^{-S(\phi)}\phi(x_1)\dots\phi(x_k)}{\int D\phi \ e^{-S(\phi)}},$$

and such ratios are unchanged under tensoring by an invertible theory. In other terms, the categorical group of invertible field theories operates on the collection of quantum field theories, and theories in the same orbit encode the same physics. Now simply observe that the orbits are projective theories.<sup>9</sup> Therefore, it is the projective theory that encodes physical information.

The second remark is about the following piece of common lore: A gapped quantum system is well-approximated at low energy by a topological field theory. In fact, the correct statement is that it is well-approximated by a projective field theory which is topological: a linear field theory approximation may not be topological, but its projectivization is. A showpiece example that illustrates why projectivity is necessary is 3-dimensional Yang-Mills with a nondegenerate Chern-Simons term. Fix a compact Lie group G. A general level is a cocycle that represents a class  $\lambda \in H^4(BG; \mathbb{Z})$ . For  $e \in \mathbb{R}^{>0}$  consider the Wick-rotated theory with lagrangian

(3.19) 
$$L = \frac{1}{4e^2} F_A \wedge *F_A + \Gamma_{\lambda}(A),$$

where  $\Gamma_{\lambda}(A)$  is the Chern-Simons term. For this 3-dimensional theory, the background fields are

$$(3.20) \qquad \qquad \mathcal{F} = \{ \text{orientations, Riemannian metrics} \}.$$

The following claims are implicit in [W2].

### Claim 3.21.

- (1) The lagrangian (3.19) determines a family of Wick-rotated field theories parametrized by  $e \in \mathbb{R}^{>0}$ .
- (2) The singular limit as  $e \to \infty$  exists and defines a field theory

(3) The underlying projective theory  $\overline{F}$  is topological, i.e., it factors through  $\mathbf{Bord}_{(2,3)}(\mathcal{F}')$  for

$$(3.23) \qquad \qquad \mathcal{F}' = \{\text{orientations}\}\$$

(4) The factored theory  $\overline{F}' \colon \mathbf{Bord}_{\langle 2,3\rangle}(\mathfrak{F}') \to \mathbf{Proj}$  has a nontrivial (framing) anomaly.

 $<sup>^8{\</sup>rm The\ tensor\ product\ operation\ on\ theories\ is\ often\ called\ 'stacking'.}$ 

<sup>&</sup>lt;sup>9</sup>As stated here, they are projective theories with trivializable anomaly. We can modify the discussion to incorporate nontrivializable anomalies by passing to centrally extended bordism categories.

Extension of an anomaly theory. Often it happens that the anomaly theory  $\alpha$ , which is a oncecategorified invertible *n*-dimensional theory, can be presented as the truncation of an invertible (n + 1)-dimensional theory  $\tilde{\alpha}$ :

Here  $\tilde{\mathcal{F}}$  is a sheaf of (n + 1)-dimensional fields. In this case, a field theory F with anomaly  $\alpha$  can be realized as a boundary theory of the invertible field theory  $\tilde{\alpha}$ . We give an example of this at the end of the paper: a free spinor field.

### 4. Anomalies as an obstruction to quantization

In this section we explain the second half of Slogan (A), which was introduced at the start:

QUANTUM THEORY IS PROJECTIVE. QUANTIZATION IS LINEAR.

Fix a dimension n, let  $\mathcal{F}$  be a sheaf of n-dimensional fields, and suppose  $\overline{F} : \mathbf{Bord}_{\langle n-1,n \rangle}(\mathcal{F}) \to \mathbf{Proj}$ is an n-dimensional (projective) theory over  $\mathcal{F}$ . We want to integrate out some fields in  $\mathcal{F}$ . Hence assume that  $\mathcal{F}$  is the total space of a fiber bundle

$$(4.1) \qquad \qquad \pi \colon \mathcal{F} \longrightarrow \mathcal{G}$$

of fields. For the quantization of  $\overline{F}$ , we view the fibers of  $\pi$  as *fluctuating* fields and the base of  $\pi$  as *background* fields. A typical example is the fiber bundle

(4.2) {orientations, Riemannian metrics, *G*-connections} 
$$\sqrt[]{\pi}$$
 {orientations, Riemannian metrics}

for G a Lie group. In this case we integrate over G-connections (gauge fields) and treat orientations and Riemannian metrics as background fields. Quantization of  $\overline{F}$  along  $\pi$ —integration over the fluctuating fields—should produce a theory  $\overline{G}$ : **Bord**<sub> $\langle n-1,n \rangle$ </sub>( $\mathcal{G}$ )  $\rightarrow$  **Proj**. Of course, this integration process has all the analytic interest and difficulties of quantum field theory. Our limited goal is to discuss descent of the projectivity (anomaly) along  $\pi$ .

To see the issue, consider a closed *n*-manifold X. Integration over fluctuating fields on X is depicted in Figure 2. Here  $g \in \mathcal{G}(X)$  is a background field on X, and the fiber of  $\pi$  over g consists



FIGURE 2. The Feynman path integral on a closed n-manifold X

of background + fluctuating fields  $f \in \mathcal{F}(X)$  that map to g under  $\pi$ . One must integrate the partition function of  $\overline{F}$  over this fiber. The partition function  $\overline{F}(X)$  is a section of a line bundle  $\alpha(X) \to \mathcal{F}(X)$ . One is instructed to integrate this section over the fibers of  $\pi$ . However, that does not make formal sense: we cannot add elements of different complex lines.<sup>10</sup> To carry out the integration we need an isomorphism of the line bundle on each fiber to a constant line bundle: the fiber over  $g \in \mathcal{G}(X)$  must be identified with a fixed line that depends only on g. In other words, we must descend the line bundle  $\alpha(X) \to \mathcal{F}(X)$  to a line bundle over  $\mathcal{G}(X)$ . (Descent is a parametrized trivialization along the fibers.) This is the argument for the anomaly as an obstruction to quantization that dates back to the 1980's. Also from that period is the variation of this argument for a closed (n-1)-manifold Y, in which case one is descending a gerbe. This is the Hamiltonian anomaly [FS, NAg, Fa, S2].

An extension of this discussion shows that we must descend the entire anomaly theory  $\alpha$  to a once-categorified invertible theory over  $\mathcal{G}$ . Trivialization of the anomaly on each fiber lifts from the projective to the linear, which is the origin of the slogan (A). This descent has an existence and uniqueness theory. The existence question—anomaly as obstruction—is the basis of the English word 'anomaly' for the projectivity of a field theory.

A descent  $\beta$  of the anomaly theory  $\alpha$  fits into the diagram



The data of the descent also includes an isomorphism

(4.4) 
$$\theta \colon \alpha \xrightarrow{\cong} \pi^* \beta$$

 $<sup>^{10}</sup>$ Of course, the integrand must be a density on the fiber, not a function, but the important point is that it take values in a *fixed* line on each fiber, not a variable line.

A change of isomorphism  $\theta$  is multiplication by an invertible field theory  $\lambda \colon \mathbf{Bord}_{\langle n-1,n \rangle}(\mathfrak{G}) \to \mathbf{Line}$ .

Remark 4.5. The descent of the anomaly is a tractable—and often very useful—piece of data for the quantization of  $\overline{F}$  along  $\pi$ . If one carries out that quantization to produce a theory  $\overline{G}$  over  $\mathcal{G}$ , then the anomaly theory of  $\overline{G}$  is *not* necessarily  $\beta$ : the quantization process may introduce new contributions to the projectivity. For example, if  $\overline{F}$  is a classical (invertible) theory of a free spinor field—with a spin structure, Riemannian metric, and possibly a connection as background fields then the anomaly  $\alpha$  is trivial and  $\beta$  can be taken to be trivial as well. But the quantum theory  $\overline{G}$ of a free spinor field has a nontrivial anomaly.

### 5. Anomaly of a spinor field

As an illustration of anomalies, we review the general formula for the anomaly of a free *n*-dimensional spinor field; this formula appears in [FH2, §9.2.5]. This anomaly theory extends to a full (n + 1)-dimensional invertible theory; as explained earlier, the anomaly is the truncation to a once-categorified *n*-dimensional theory. The formula for the extended theory is a map of spectra in stable homotopy theory, and so we begin with a quick review of the representation of invertible field theories as maps of spectra. Then we review the data that defines a free spinor field and conclude with the formula (5.8) for the anomaly theory. We remark that the formula is a conjecture until more foundations (with differential cohomology) are developed for invertible field theories and until the general Wick-rotated free spinor field is completely constructed in the framework of Axiom System 3.4.

The account here is overly laconic, included mainly to illustrate Remark 5.2(1) below. As compensation, we refer to [FH2] for background, details, and references.

Invertible field theories and stable homotopy theory. The basic idea is that an invertible theory

(5.1) 
$$\lambda \colon \mathbf{Bord}_{\langle n-1,n \rangle}(\mathcal{F}) \longrightarrow \mathbf{Line}$$

maps into a groupoid: all morphisms in the category **Line** are invertible. Therefore,  $\lambda$  factors through the quotient of **Bord**<sub> $\langle n-1,n \rangle$ </sub>( $\mathcal{F}$ ) in which all morphisms are inverted. At that point we obtain a map of groupoids, which is equivalently a map of spaces. (For a leisurely exposition of this equivalence, see [Po].) Furthermore, the symmetric monoidal structure on the domain and codomain in (5.1) mean that the spaces are equipped with an infinite loop structure. Infinite loop spaces are equivalent to (connective) spectra. In the end, then, an invertible field theory is represented as a map of spectra.

## Remark 5.2.

- (1) The passage from categories to spectra makes invertible field theories amenable to a suite of mathematical tools.
- (2) The same passage applies to once-categorified invertible theories.

(3) Full locality is well-developed for *topological* field theories, and for invertible theories that are not necessarily topological we implicitly assume full locality in what follows. In the general invertible, possibly non-topological, case we need *differential* versions of spectra; see [ADH] and references therein.

For a free spinor field we have background fields

(5.3) 
$$\mathcal{F} = \{\text{spin structures, Riemannian metrics}\}.$$

If we drop the Riemannian metrics, then the spectrum obtained from the (extended) bordism category is the Thom spectrum MSpin. In some cases the anomaly theory we are after is topological, and then MSpin is the domain of the anomaly theory. In general, the domain is a differential version.

There is a universal codomain for invertible field theories, characterized in the topological case by the universal property that the partition functions determine the theory. We use a differential variant, which for *m*-dimensional theories is a differential version of the Anderson dual to the sphere spectrum  $I\mathbb{Z}$ , shifted according to the dimension. With differential variants implicit, an (n + 1)-dimensional invertible theory with background fields (5.3) is a spectrum map

(5.4) 
$$\alpha \colon M\mathrm{Spin} \longrightarrow \Sigma^{n+2} I\mathbb{Z}$$



FIGURE 3. Minkowski spacetime  $\mathbb{M}^n$  and the vector space  $\mathbb{R}^{1,n-1}$ 

Spinor field data. This data is given in the relativistic setting. Fix a spacetime dimension n. Let  $\mathbb{M}^n$  denote standard Minkowski spacetime. It is an *n*-dimensional affine space acted upon simply transitively by the vector group  $\mathbb{R}^{1,n-1}$  of translations; see Figure 3.  $\mathbb{M}^n$  is equipped with a translation-invariant Lorentz metric and a choice  $C \subset \mathbb{R}^{1,n-1}$  of a component of (forward) timelike vectors—a time-orientation. The spin group  $\operatorname{Spin}_{1,n-1} \subset \operatorname{Cliff}_{n-1,1}^0$  is a subset of the even Clifford algebra.<sup>11</sup> Spinor field data is a triple  $(\mathbb{S}, \Gamma, m)$  that consists of

$$\begin{array}{ll} \mathbb{S} & \text{real (ungraded) finite dimensional } \operatorname{Cliff}_{n-1,1}^{0}\text{-module} \\ (5.5) & \Gamma \colon \mathbb{S} \times \mathbb{S} \longrightarrow \mathbb{R}^{1,n-1} & \text{symmetric } \operatorname{Spin}_{1,n-1}\text{-invariant form; } \Gamma(s,s) \in \overline{C} \text{ for all } s \in \mathbb{S} \\ & m \colon \mathbb{S} \times \mathbb{S} \longrightarrow \mathbb{R} & \text{skew-symmetric } \operatorname{Spin}_{1,n-1}\text{-invariant } (mass) \text{ form} \end{array}$$

16

<sup>&</sup>lt;sup>11</sup>This Clifford algebra has n-1 generators that square to +1 and one generator that squares to -1.

The Clifford module S restricts to a spin representation of  $\text{Spin}_{1,n-1}$ . It is a remarkable theorem that the positive symmetric pairing  $\Gamma$  exists for all S; this is only true in Lorentz signature. Furthermore, if S is irreducible, then  $\Gamma$  is unique up to positive scale; in general there is a contractible space of possible  $\Gamma$ . The mass form m may vanish.

Several important algebraic facts are proved or referenced in [FH2, §9.2.4]. Given  $(\mathbb{S}, \Gamma)$ , there is a unique compatible  $\mathbb{Z}/2\mathbb{Z}$ -graded  $\operatorname{Cliff}_{n-1,1}$ -module structure on  $\mathbb{S} \oplus \mathbb{S}^*$ . Furthermore, every finite dimensional  $\operatorname{Cliff}_{n-1,1}$ -module has this form. Crucially, nondegenerate mass forms for  $(\mathbb{S}, \Gamma)$ correspond to  $\operatorname{Cliff}_{n-1,2}$ -module structures on  $\mathbb{S} \oplus \mathbb{S}^*$  that extend the  $\operatorname{Cliff}_{n-1,1}$ -module structure.

The anomaly theory. We give the conjectured formula for m = 0. In [CFLS] the theory with m as a scalar field is considered, and there is a conjectured formula for the anomaly in [CFLS, §7.4].

Since  $\Gamma$  is a contractible choice, we drop it from the notation. Atiyah-Bott-Shapiro [ABS] identify the abelian group of equivalence classes of  $\operatorname{Cliff}_{n-1,1}$ -modules modulo those that extend to  $\operatorname{Cliff}_{n-1,2}$ modules in terms of the KO-theory spectrum, and the module S determines a class in that group:

$$[\mathbb{S}] \in \pi_{2-n} KO \cong [S^0, \Sigma^{n-2} KO],$$

where  $S^0$  is the sphere spectrum.

**Claim 5.7.** The anomaly theory  $\alpha = \alpha_{(\mathbb{S},\Gamma)}$  of the massless free spinor field is a differential lift of

(5.8) 
$$M\operatorname{Spin} \xrightarrow{\phi \wedge [\mathbb{S}]} KO \wedge \Sigma^{n-2}KO \xrightarrow{\mu} \Sigma^{n-2}KO \xrightarrow{\operatorname{Pfaff}} \Sigma^{n+2}I\mathbb{Z}$$

Here  $\phi: MSpin \to KO$  is essentially the Thom class of a real spin bundle, also defined in [ABS]. The map  $\mu$  is multiplication in the ring spectrum KO, and Pfaff:  $KO \to \Sigma^4 I\mathbb{Z}$  is the map that enters the Anderson self-duality of KO; see [FMS].

Remark 5.9. As written (i.e., without a differential lift), (5.8) is the deformation class of the anomaly theory  $\alpha$ .

## Appendix A. On the definition of Proj

We make some comments on the definition of **Proj**, introduced in (3.8) as the codomain of a projective field theory (3.10). Our treatment is sparse since a detailed development is the subject of a forthcoming thesis of Chetan Vuppulury (a student of Domenico Fiorenza).

Warmup: Suppose S is a set equipped with the action of a group G. The naive quotient is the set S/G of orbits of the G-action. But it flouts the core ethos of categorical thinking to identify elements of S to form a quotient. Rather, one should remember the group element that effects the identification. This leads to the action groupoid S//G whose set of objects is S and whose set of morphisms is  $S \times G$ . Lesson: The quotient of a set by a group is a 1-category.

The sequence (3.8) exhibits **Proj** as the quotient of **Vect** by **Line**, and the warmup suggests that this quotient is a 2-category. A model for **Line** is the groupoid  $B\mathbb{C}^{\times}$ ; it has a single object

with automorphism group  $\mathbb{C}^{\times}$ . Hence the 2-category **Proj** has vector spaces as objects, linear maps as 1-morphisms, and diagrams

as 2-morphisms, where  $\lambda \in \mathbb{C}^{\times}$  and the linear maps  $T_0, T_1$  satisfy  $T_1 = \lambda T_0$ . (The same adjectives used to define **Vect**—certain *topological* vector spaces, *nuclear* linear maps—apply to **Proj**.)

More generally, suppose  $\mathcal{C}$  is a symmetric monoidal  $(\infty, n + 1)$ -category for some<sup>12</sup>  $n \in \mathbb{Z}^{\geq 0}$ . Let  $\mathcal{C}^{\times} \subset \mathcal{C}$  be the maximal Picard subgroupoid of  $\mathcal{C}$ . Heuristically, define the *projectivization*  $\mathbb{P}\Omega\mathcal{C}$  of  $\Omega\mathcal{C} = \operatorname{Hom}_{\mathcal{C}}(1, 1)$  as the pullback

(A.2) 
$$\begin{array}{c} \mathbb{P}\Omega\mathbb{C} & - \to \mathbb{C}^{\times} \\ & | \\ & | \\ & | \\ & \psi \\ & \ast \xrightarrow{} \mathbb{C} \end{array}$$

The special case that defines **Proj** is

where **Alg** is the Morita 2-category of algebras, bimodules, and intertwiners. The core issue is to *define* the pullback (A.2). In fact, more general constructions are given in [JS]. Namely, Johnson-Freyd and Scheimbauer define categories  $\mathbb{C}^{\downarrow}$  and  $\mathbb{C}^{\rightarrow}$ , each equipped with source and target homomorphisms  $s, t: \mathbb{C}^* \to \mathbb{C}$ , where  $* = \downarrow$  or  $* = \rightarrow$ . The pullback we seek is presumably the intersection  $s^{-1}(1) \cap t^{-1}(\mathbb{C}^{\times})$ . Finally, in terms of [JS, Definition 1.4] and the papers that inspired it, we see that a projective field theory is a field theory *relative* to its anomaly theory.

### References

[A]	M. F. Atiyah,	Topological	quantum	field	theories,	Inst.	Hautes	Études	Sci.	Publ.	Math.	(1988),	no.	68,
	175-186 (1989)	). 8												

- [ABS] M. F. Atiyah, R. Bott, and A. A. Shapiro, *Clifford modules*, Topology 3 (1964), 3–38. 17
- [Ad] Stephen L Adler, Axial-vector vertex in spinor electrodynamics, Physical Review **177** (1969), no. 5, 2426.

<sup>&</sup>lt;sup>12</sup>The construction also works for a symmetric monoidal  $(\infty, 0)$ -category, represented as an infinite loop space X. Then  $\mathbb{P}\Omega X$  is the space of paths in X emanating from the basepoint.

- [ADH] Araminta Amabel, Arun Debray, and Peter J. Haine, *Differential Cohomology: Categories, Characteristic Classes, and Connections*, arXiv:2109.12250. 16
- [AgVm] Luis Alvarez-Gaume and Miguel A. Vazquez-Mozo, Anomalies and the Green-Schwarz Mechanism, arXiv:2211.06467 [hep-th].1
- [BH] Dorje C. Brody and Lane P. Hughston, *Geometric quantum mechanics*, J. Geom. Phys. **38** (2001), no. 1, 19–53. 7
- [BJ] John Stewart Bell and Roman W Jackiw, A PCAC puzzle:  $\pi^0 \rightarrow \gamma\gamma$  in the  $\sigma$ -model, Nuovo cimento **60** (1969), no. CERN-TH-920, 47–61. 1
- [Bo] L. Bonolis, From the Rise of the Group Concept to the Stormy Onset of Group Theory in the New Quantum Mechanics. A saga of the invariant characterization of physical objects, events and theories., Nuovo Cimento Rivista Serie 27 (2004), no. 4, 040000–110. 7
- [CFLS] Clay Córdova, Daniel S. Freed, Ho Tat Lam, and Nathan Seiberg, Anomalies in the Space of Coupling Constants and Their Dynamical Applications I, SciPost Phys. 8 (2020), no. 1, 001, arXiv:1905.09315 [hep-th]. 17
- [CH] Curtis G. Callan, Jr. and Jeffrey A. Harvey, Anomalies and Fermion Zero Modes on Strings and Domain Walls, Nucl. Phys. B250 (1985), 427–436. 1
- [F1] Daniel S Freed, Anomalies and invertible field theories, Proc. Symp. Pure Math, Proc. Sympos. Pure Math., vol. 88, Amer. Math. Soc., Providence, RI, 2014, pp. 25–45. arXiv:1404.7224. 2
- [F2] Daniel S. Freed, On Wigner's theorem, Proceedings of the Freedman Fest (Vyacheslav Krushkal Rob Kirby and Zhenghan Wang, eds.), Geometry & Topology Monographs, vol. 18, Mathematical Sciences Publishers, 2012, pp. 83–89. arXiv:1211.2133. 7
- [Fa] L. D. Faddeev, Hamiltonian approach to the theory of anomalies, Recent Developments in Mathematical physics (H. Mitter and L. Pittner, eds.), Internationale Universitatswoche fur Kernphysik, Schladming, Austria, vol. 26, 1987, pp. 137–159. 14
- [FH1] Daniel S. Freed and Michael J. Hopkins, Chern-Weil forms and abstract homotopy theory, Bull. Amer. Math. Soc. (N.S.) 50 (2013), no. 3, 431–468, arXiv:1301.5959.8
- [FH2] \_\_\_\_\_, Reflection positivity and invertible topological phases, Geom. Topol. 25 (2021), no. 3, 1165–1330, arXiv:1604.06527.15, 17
- [FHT] Daniel S. Freed, Michael J. Hopkins, and Constantin Teleman, Consistent orientation of moduli spaces, The many facets of geometry, Oxford Univ. Press, Oxford, 2010, pp. 395–419. arXiv:0711.1909. 2
- [FMS] Daniel S. Freed, Gregory W. Moore, and Graeme Segal, The uncertainty of fluxes, Commun. Math. Phys. 271 (2007), 247–274, arXiv:hep-th/0605198. 17
- [FS] L.D. Faddeev and S. L. Shatashvili, Algebraic and Hamiltonian Methods in the Theory of Nonabelian Anomalies, Theor. Math. Phys. 60 (1985), 770–778. 14
- [FT] Daniel S. Freed and Constantin Teleman, Relative quantum field theory, Comm. Math. Phys. 326 (2014), no. 2, 459–476, arXiv:1212.1692. 2
- [JS] Theo Johnson-Freyd and Claudia Scheimbauer, (Op)lax natural transformations, twisted quantum field theories, and "even higher" Morita categories, Advances in Mathematics **307** (2017), 147–223, arXiv:1502.06526.18
- [KS] Maxim Kontsevich and Graeme Segal, Wick rotation and the positivity of energy in quantum field theory,
  Q. J. Math. 72 (2021), no. 1-2, 673–699, arXiv:2105.10161.8, 9
- [La] R. Lashof, Poincaré duality and cobordism, Trans. Amer. Math. Soc. 109 (1963), 257–277. 9
- [Lo] Ottmar Loos, Symmetric spaces. II: Compact spaces and classification, W. A. Benjamin, Inc., New York-Amsterdam, 1969. 7
- [M] John W. Milnor, *Lectures on the h-cobordism theorem*, Notes by L. Siebenmann and J. Sondow, Princeton University Press, Princeton, N.J., 1965. 9
- [NAg] Philip Nelson and Luis Alvarez-Gaumé, Hamiltonian interpretation of anomalies, Comm. Math. Phys. 99 (1985), no. 1, 103–114. 14
- [Po] Timothy Porter, Spaces as infinity-groupoids. https://ncatlab.org/nlab/files/Spaces+as+ infinity-groupoids.pdf. 15
- [S1] Graeme Segal, The definition of conformal field theory, Topology, geometry and quantum field theory, London Math. Soc. Lecture Note Ser., vol. 308, Cambridge Univ. Press, Cambridge, 2004, pp. 421–577.
   8, 11
- [S2] G. B. Segal, Faddeev's anomaly in Gauss's law. preprint. 14
- [St] J Steinberger, On the use of subtraction fields and the lifetimes of some types of meson decay, Physical Review 76 (1949), no. 8, 1180. 1
- [tH] Gerard 't Hooft, Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking, NATO Sci. Ser. B 59 (1980), 135–157. 1

[vNW]	J. v. Neumann and E. Wigner, Zur Erklärung einiger Eigenschaften der Spektren aus der Quanten-
	mechanik des Drehelektrons, Zeitschrift für Physik A Hadrons and Nuclei 49 (1928), 73–94. 7
[W1]	Edward Witten, World sheet corrections via D instantons, JHEP <b>02</b> (2000), 030, arXiv:hep-th/9907041.
[W2]	, Quantum field theory and the Jones polynomial, Comm. Math. Phys. <b>121</b> (1989), no. 3, 351–399. 12

Department of Mathematics, Harvard University, Cambridge, MA 02138  $\mathit{Email}\ address: \texttt{dafr@math.harvard.edu}$ 

20