A view of Coherent Elastic Neutrino-Nucleus Scattering

M. $CADEDDU^1$, F. $DORDEI^1$ and C. $GIUNTI^2$

¹ Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Cagliari, Complesso Universitario di Monserrato - S.P. per Sestu Km 0.700, 09042 Monserrato (Cagliari), Italy

² Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Torino, Via P. Giuria 1, 10125 Torino, Italy

Abstract –We review the physics of coherent elastic neutrino-nucleus scattering and the results and perspectives for the measurements of the radius of the neutron distribution of the nucleus, of the weak mixing angle, and of new neutrino interactions due to physics beyond the Standard Model.

Introduction. – Coherent elastic neutrino-nucleus scattering (CE ν NS) is a peculiar interaction process which was predicted theoretically in 1973 [1] (see also [2, 3]), but it was observed experimentally for the first time only in 2017 in the COHERENT experiment [4], with neutrinos produced by the Spallation Neutron Source (SNS) at the Oak Ridge National Laboratory. The cross-section of

CE ν NS is larger than the cross-sections of other neutrino interactions at the same neutrino energy. Thus, one may wonder why it took 44 years to observe CE ν NS after its theoretical prediction. The reason is that the only observable signature of CE ν NS is the recoil of the nucleus with an extremely small kinetic energy which is very difficult to observe. The recent development of new very sensitive detectors allowed this magnificent achievement.

The main characteristic of $CE\nu NS$ is that the interactions of the neutrino with the nucleons must be coherent, such that the nucleus recoils as a whole, without any change in its internal structure. In this case, the predicted Standard Model (SM) weak-interaction cross-section is approximately proportional to the square of the number of neutrons N in the nucleus. Therefore, the probability of $CE\nu NS$ interactions is approximately N times larger than the sum of the probabilities of neutrino interaction with every single nucleon in the same nucleus, which is only approximately proportional to N. The reason why the number of neutrons N in the nucleus is more relevant than the number of protons is that, as explained later, the crosssection of neutrino-proton interaction is much smaller than the cross-section of neutrino-neutron interaction.

 $CE\nu NS$ is a process that is not only interesting per se. It gives useful information on the nuclear structure, in particular on the neutron density distribution in the nucleus, which surprisingly is still unknown for most of the nuclei. It gives also information on the precise value of the neutrino neutral-current interaction at low energy, which is quantified by the so-called "weak mixing angle". Moreover, $CE\nu NS$ is sensitive to effects of physics Beyond the Standard Model (BSM) such as neutrino electromagnetic interactions, neutrino interactions mediated by new neutral BSM bosons, and, in general, neutrino neutral-current nonstandard interactions (NSI).

So far, all the results obtained from $CE\nu NS$ experiments are compatible with the SM prediction. After the first observation by the COHERENT Collaboration of $CE\nu NS$ in 2017 using a cesium-iodide (CsI) detector [4], a larger dataset of scattering events have been collected with the same experiment [5] and the same process has also been observed with a liquid argon detector [6]. Recently, the experimental collaboration working at the Dresden-II experiment reported the observation of $CE\nu NS$ produced by electron antineutrinos from a nuclear power reactor [7]. Other experiments (CONUS [8] and CONNIE [9]) have been able to obtain upper limits on $CE\nu NS$, which are compatible with the SM prediction, and are still working towards their first observation. Several other $CE\nu NS$ experiments are planned or under construction: new CO-HERENT detectors [10], MINER [11], NEON [12], NU-CLEUS [13], RED-100 [14], Ricochet [15], a project at the European Spallation Source [16], and others [17]. Therefore, we expect many new experimental results in the next years, which will bring very interesting information on nuclear physics, weak neutral-current neutrino interactions and BSM physics.

 $CE\nu NS$ in the Standard Model. – $CE\nu NS$ is the neutral-current process

$$\nu_{\alpha} + {}^{A}_{Z}\mathcal{N} \to \nu_{\alpha} + {}^{A}_{Z}\mathcal{N}, \tag{1}$$




Fig. 1: (a) Feynman diagram of $CE\nu NS$ in the Standard Model. ν_{α} represents a neutrino with flavor $\alpha = e, \mu, \text{ or } \tau, \frac{A}{Z}\mathcal{N}$ denotes a nucleus with A nucleons (Z protons and N = A - Z neutrons), and Z^0 is the Standard Model neutral vector boson which mediates neutral-current weak interactions. (b) Feynman diagram of contributions to $CE\nu NS$ of BSM physics in which the interaction is mediated by a photon γ , or a new massive vector boson Z', or scalar boson Φ .

where ν_{α} represents a neutrino with flavor $\alpha = e, \mu, \text{ or } \tau$, and ${}^{A}_{Z}\mathcal{N}$ denotes a nucleus with A nucleons, of which Zare protons and N = A - Z are neutrons. Neutral-current processes are characterized by the absence of an exchange of electric charge between the interacting particles. Hence, they are mediated by the exchange of a neutral boson. In the SM the mediator is the Z^{0} neutral boson, as illustrated in Fig. 1(a).

Figure 2 illustrates three types of interactions of a neutrino ν_{α} with a nucleus, which depend on the wavelength $\lambda_{Z^0} = h/|\vec{q}|$ of the Z^0 neutral vector boson which mediates SM neutral-current weak interactions (*h* is Planck's constant and \vec{q} is the Z^0 three-momentum). When $\lambda_{Z^0} \ll 2R$, where *R* is the radius of the nucleus, the Z^0 has a high probability to interact with a single nucleon in the nucleus (a neutron in the illustration), ejecting it. When $\lambda_{Z^0} \leq 2R$ the Z^0 has a high probability to interact with a group of nucleons in the nucleons, exciting the latter to the state \mathcal{N}^* . Elastic coherent neutrino-nucleus scattering can happen only when $\lambda_{Z^0} \gtrsim 2R$. In this case, the wavelength of the Z^0 covers all the nucleus, allowing a coherent nuclear recoil without any internal change of the nucleus.

For CE ν NS, the absolute value $|\vec{q}|$ of the Z^0 threemomentum is related to the measurable value of the kinetic energy of the recoiling nucleus T by $|\vec{q}| \simeq \sqrt{2 M T}$, where M is the mass of the nucleus. Therefore, the coherency condition is satisfied only for very small values of T, which are difficult to measure. For example, let us consider a heavy nucleus with $A \approx 100$ nucleons, for which the nuclear radius can be approximately estimated with the well-known semi-empirical formula $R \approx 1.2 A^{1/3}$ fm \approx 5 fm. Therefore, CE ν NS can happen for $|\vec{q}| \leq 40$ MeV, which implies $T \leq 10$ keV. This is a very low value of the recoil energy of the nucleus, which could not be measured before the recent technical developments of very sensitive detectors employed in the search of dark matter in the form of weakly interacting massive particles (WIMPs). Indeed, WIMPs with masses of $\mathcal{O}(10 \,\text{GeV}/c^2)$ induce nuclear recoils in the keV range which are very similar to those produced by CE ν NS processes.

Moreover, since the average momentum transfer depends on the neutrino energy, it is only possible to observe CE ν NS generated by low-energy neutrinos ($E_{\nu} \lesssim 30 \text{ MeV}$). There are various natural sources of low-energy neutrinos: solar neutrinos, geoneutrinos, supernova neutrinos, and the low-energy tail of atmospheric neutrinos [17]. Several experiments are sensitive to CE ν NS generated by these natural low-energy neutrino fluxes (especially experiments on the search of nuclear recoils produced by dark matter [18]), but so far these neutrino fluxes have been observed only through other interactions.

 $\mathrm{CE}\nu\mathrm{NS}$ has been observed only in two laboratory experiments:

- 1. The COHERENT experiment [4-6], with neutrinos produced by the Oak Ridge SNS. In this experiment, the neutrinos are generated by the decay at rest of positive pions (π^+) and muons (μ^+) . The positive pions are created by the interaction of a pulsed beam of protons with an energy of about 1 GeV with a mercury target (the produced negative pions are rapidly absorbed by the positive nuclei of the target before decay). They stop in the target and decay at rest with a lifetime of about 26 ns with the process $\pi^+ \to \mu^+ + \nu_{\mu}$, generating a prompt flux of muon neutrinos (ν_{μ}). The produced positive muons also stop in the target and decay at rest with a longer lifetime of about 2.2 μs with the process $\mu^+ \to e^+ + \bar{\nu}_{\mu} + \nu_e$, generating delayed fluxes of muon antineutrinos $(\bar{\nu}_{\mu})$ and electron neutrinos (ν_e). Since the energies of these neutrino fluxes are below about 53 MeV, they generate observable rates of $CE\nu NS$ in the COHERENT CsI [4,5] and LAr [6] detectors, where the total neutrino flux is about $4.3 \times 10^7 \,\nu \,\mathrm{cm}^{-2} \,s^{-1}$. It is remarkable that the detectors are very small in comparison with typical neutrino detectors, with mass scales of the order of multiples of tons. The CsI detector weighs only 14.6 kg and the LAr detector weighs 29 kg. The detection of $CE\nu NS$ with such small detectors is possible because of the large $CE\nu NS$ cross-section.
- 2. The Dresden-II experiment [7] with electron antineutrinos ($\bar{\nu}_e$) generated by the Dresden-II nuclear power reactor of the Dresden Generating Station located near Morris, Illinois, USA. The flux of electron antineutrinos at the very sensitive germanium detector is huge, about $4.8 \times 10^{13} \bar{\nu}_e \text{ cm}^{-2} s^{-1}$, with energy below about 10 MeV. It is well suited to generate an observable rate of CE ν NS in the extraordinarily small germanium detector weighing only 3 kg. We note that there is, however, some criticism of this result related to the anomalously large quenching factor, the reactor-related background and the incompatibility with the recent CONUS results [19] which must be clarified [20].




Fig. 2: Illustration of three types of interactions of a neutrino ν_{α} with a nucleus: inelastic incoherent scattering when $\lambda_{Z^0} \leq 2R$, elastic incoherent scattering when $\lambda_{Z^0} \leq 2R$, and elastic coherent scattering (CE ν NS) when $\lambda_{Z^0} \geq 2R$. Here R is the radius of the nucleus and $\lambda_{Z^0} = h/|\vec{q}|$ is the wavelength of the Z^0 neutral vector boson, where \vec{q} is its three-momentum and h is Planck's constant.

The measurable $CE\nu NS$ differential cross-section of a neutrino with a spin-zero nucleus \mathcal{N} is

$$\frac{d\sigma_{\nu\mathcal{N}}}{dT}(E_{\nu},T) = \frac{G_{\rm F}^2 M}{\pi} \left(1 - \frac{MT}{2E_{\nu}^2}\right) \left[Q_W^{\mathcal{N}}(|\vec{q}|)\right]^2, \quad (2)$$

where E_{ν} is the neutrino energy, and $G_{\rm F}$ is the Fermi constant. The interaction is characterized by the so-called "weak charge of the nucleus" $Q_W^{\mathcal{N}}(|\vec{q}|)$, a function of the three-momentum \vec{q} transferred from the neutrino to the nucleus that depends on the interaction type. In the SM

$$Q_W^{\mathcal{N}}(|\vec{q}|) = g_V^n N F_N^{\mathcal{N}}(|\vec{q}|) + g_V^p Z F_Z^{\mathcal{N}}(|\vec{q}|).$$
(3)

The functions $F_N^{\mathcal{N}}(|\vec{q}|)$ and $F_Z^{\mathcal{N}}(|\vec{q}|)$ are, respectively, the neutron and proton form factors of the nucleus \mathcal{N} , which characterize the amount of coherency of the interaction and depend on the neutron and proton distributions in the nucleus. The coefficients g_V^n and g_V^p quantify the weak neutral-current interactions of neutrons and protons, respectively. In the SM they are approximately given by¹

$$g_V^n \simeq -\frac{1}{2}$$
 and $g_V^p \simeq \frac{1}{2} - 2\sin^2\vartheta_W \simeq 0.022,$ (4)

where ϑ_W is the weak mixing angle, with $\sin^2 \vartheta_W \simeq 0.239$. Therefore, the neutron contribution is much larger than the proton contribution and the CE ν NS cross section is approximately proportional to N^2 .

Nuclear Physics. – The approximate increase of the total CE ν NS cross-section with the square of the number of neutrons in the nucleus is shown by the black curve in Fig. 3. However, one can see that the result of the CO-HERENT measurement of the total CE ν NS cross-section with the CsI detector lies below the black line and is compatible with the green line which takes into account a small amount of incoherency quantified by a realistic form factor $F_N^{\mathcal{N}}(|\vec{q}|)$. Indeed, the COHERENT CsI data show a 6σ evidence of the nuclear structure suppression of the full coherence [21].




Fig. 3: Approximate increase of the total $CE\nu NS$ cross-section with the square of the number of neutrons N in the nucleus \mathcal{N} (adapted from the Fig. 11 in Ref. [6]). The black curve shows the increase with N^2 in the case of full coherency (i.e. $F_N^{\mathcal{N}}(|\vec{q}|) = 1$ in Eq. (3)), whereas the green curve shows the expected increase which is slightly slower because of a small amount of incoherency quantified by a realistic form factor $F_N^{\mathcal{N}}(|\vec{q}|)$. The black dots correspond to the predictions for four selected nuclei. The blue dots with error bars show the measurements of the COHERENT experiment.

The dependence of the cross-section on the neutron form factor $F_N^N(|\vec{q}|)$ is a powerful tool for obtaining information on the neutron distribution in the nucleus, which is not well-known. Indeed, while the proton distribution of most nuclei is known from electromagnetic measurements [22], the nuclear neutron distribution can be probed only through measurements which employ the strong or weak forces. The interpretation of the results of experiments with hadron probes which explore the nuclear neutron distribution through the strong force is difficult, since the effects of strong-force interactions cannot be calculated with sufficient approximation and the interpretation can be done only by assuming a strong-interaction model with all its limitations [23]. On the other hand, the effects of the weak neutral-current interactions, embodied by $Q_W^{\mathcal{N}}(|\vec{q}|)$ in Eq. (3), are known with good approximation. Therefore, weak neutral-current processes as $CE\nu NS$ are ideal for probing the nuclear neutron distribu-

 $^{^{1}}$ The more accurate numerical values which incorporate the socalled radiative corrections can be found in [21].

tion. Besides $CE\nu NS$, the nuclear neutron distribution has been probed with neutral-current weak interactions through parity-violating electron scattering for only two nuclei: 208 Pb in the PREX experiment [24] and 48 Ca in the CREX experiment [25]. The quantity which in practice is measured is the radius R_n of the neutron distribution in the nucleus. Since in heavy nuclei there are more neutrons than protons, all nuclear models predict that R_n should be larger than the radius R_p of the proton distribution in the same nucleus. The excess $\Delta R_{np} = R_n - R_p$ is called "neutron skin". It is a quantity of great interest for nuclear physics and astrophysics, because it is the result of the competition between the Coulomb repulsion between the protons, the surface tension, that decreases when the excess neutrons are pushed to the surface, and the symmetry energy [26]. It gives information on the nuclear equation of state, which determines the size of neutron stars [27].

For CE ν NS, the neutron form factor $F_N^{\mathcal{N}}(|\vec{q}|)$, being the Fourier transform of the neutron distribution in the nucleus, depends on R_n . The CE ν NS measurements of the COHERENT collaboration with the CsI detector [4, 5]give information on the average radius R_n^{CsI} of the neutron distributions in ¹³³Cs and ¹²⁷I [21, 28–34]: $R_n^{\text{CsI}} =$ $5.47\pm0.38\,\mathrm{fm}$ [21]. This value is in agreement, within the uncertainty, with the recent Nuclear Shell Model (NSM) prediction $R_n^{\text{CsI}}(\text{NSM}) \simeq 5.06 \,\text{fm} [35, 36]$. The $\text{CE}\nu\text{NS}$ measurement of the COHERENT collaboration with the LAr detector [6] has larger uncertainties and leads only to an upper bound for the radius $R_n^{\rm Ar}$ of the neutron distributions in ⁴⁰Ar [37, 38]: $R_n^{\text{Ar}} < 4.2 \text{ fm}$ [38] at 1σ (i.e. 68% confidence level). This bound is compatible with the recent NSM prediction $R_n^{\rm Ar} \simeq 3.6 \, {\rm fm} \, [35]$. Further improvements in the determination of R_n^{CsI} and R_n^{Ar} are expected when new measurements of the COHERENT collaboration will be available [10], as illustrated in Fig. 4.

An important improvement in the determination of R_n^{CSI} can be achieved by combining the COHERENT CSI results with those of Atomic Parity Violation (APV) experiments with Cs atoms [39, 40], which are sensitive to R_n^{Cs} [21, 36, 41]. This method permits to disentangle the measurements of the radii of the neutron distributions of ¹³³Cs and ¹²⁷I: $R_n^{\text{Cs}} = 5.32^{+0.30}_{-0.23}$ fm and $R_n^{\text{I}} = 5.30^{+0.3}_{-0.6}$ fm [21]. Obviously R_n^{Cs} is better determined than R_n^{I} . Both are compatible with the NSM predictions $R_n^{\text{Cs}}(\text{NSM}) \simeq 5.09$ fm and $R_n^{\text{I}}(\text{NSM}) \simeq 5.03$ fm [35, 36].

Note that the results of reactor neutrino $CE\nu NS$ experiments, as those of the Dresden-II experiment [7], do not give information on the neutron distribution of the target nucleus, because the neutrino energy is low, of a few MeV, and there is an observable $CE\nu NS$ signal only for very small momentum transfers. Therefore, the interaction is fully coherent and the form factors in Eq. (3) are practically equal to one. On the other hand, in these experiments there is the advantage that the extraction from the data of other physical quantities does not depend on the poorly known neutron distribution of the nucleus.




Fig. 4: Current status and future projections for neutron distribution radii of different nuclei measured via $CE\nu NS$. The upper and lower plots show, respectively, the current and foreseen measurements of R_n^{CsI} from $CE\nu NS$ with CsI detectors [21,28,30,36] and those of R_n^{Ar} with Ar detectors [10,38].

Electroweak Physics. – The $CE\nu NS$ cross section depends on the value of $\sin^2 \vartheta_W$ through the proton coupling g_V^p in Eq. (4). Although the proton contribution to the $CE\nu NS$ cross section is subdominant with respect to the neutron one, it must be taken into account when performing accurate calculations. In quantum field theory, the value of $\sin^2 \vartheta_W$ (as well as other quantities) is not a constant, but depends on the energy scale at which it is measured, as illustrated in Fig. 5. This figure shows the SM prediction for $\sin^2 \vartheta_W$ obtained from high-precision measurements at a scale of about 100 GeV in experiments at the Tevatron, LEP, SLC and LHC high-energy colliders [42]. The figure shows also that the low-energy value of $\sin^2 \vartheta_W$ has been probed in several experiments with various results which have different degrees of compatibility with the SM prediction. The main interest in measuring the low-energy value of $\sin^2 \vartheta_W$ lies in the possibility of discovering a deviation from the SM due to new physics contributions [48,49], such as the mediation of the interaction by dark Z bosons [49-53]. The determination of the low-energy value of the weak mixing angle has been discussed in several studies [21,28,30-32,36-38,41,51,54-62], with the recent updated result $\sin^2 \vartheta_W = 0.231^{+0.027}_{-0.024}$ [21], obtained from the latest COHERENT CsI data [5]. As in the case of the radius of the neutron distribution in the nucleus, the measurement of $\sin^2 \vartheta_W$ can be improved with a combined analysis of the $CE\nu NS$ COHER-ENT CsI data and those of APV experiments with Cs atoms [39, 40], which are also sensitive to the weak mixing angle: $\sin^2 \vartheta_W = 0.2398^{+0.0016}_{-0.0015}$ [21]². This measure-

²We note that the APV $\sin^2 \vartheta_W$ determination depends on the value used for the theoretical parity non-conserving amplitude, for which two different determinations are available [43,63]. Here, only the value obtained with [63] has been reported. The particular choice




Fig. 5: Variation of $\sin^2 \vartheta_W$ with the energy scale. The SM prediction is shown by the blue solid curve, together with experimental determinations in black [39, 42–47]. The result obtained in [21] from the combination of COHERENT CE ν NS data and APV data is shown in red.

ment becomes $\sin^2 \vartheta_W = 0.2423^{+0.0032}_{-0.0029}$ [21], as reported in Fig. 5, when the neutron distribution is simultaneously determined. Clearly, the current combination is dominated by the APV result, but future measurements of the CO-HERENT collaboration [10] are expected to reduce significantly the uncertainty of the CE ν NS determination of the weak mixing angle, as illustrated in Fig. 6. As it can be seen, more $\sin^2 \vartheta_W$ measurements are also expected from other future CE ν NS experiments (CO ν US [64], TEX-ONO [65], CONNIE [66], and MINER [11]) that will be really powerful for further constraining such a quantity.

 $CE\nu NS$ beyond Standard Model. – $CE\nu NS$ measurements are powerful probes of physics beyond the SM which can induce neutral current interactions as those depicted in Fig. 1(b), which involve the mediation of a photon γ , or a new massive vector boson Z', or a new massive scalar boson Φ . Note that in these non-standard interactions the neutrino flavor can change: $\nu_{\alpha} + {}^{A}_{Z} \mathcal{N} \rightarrow \nu_{\beta} + {}^{A}_{Z} \mathcal{N}$, with a final neutrino flavor β which can be equal or different of the initial neutrino flavor α . This is different from the SM $CE\nu NS$ process in Fig. 1(a), where the initial and final neutrino flavors are equal, because the SM neutralcurrent weak interaction conserves flavor. In models BSM there can be flavor-changing neutral-current interactions. Unfortunately, the change of neutrino flavor is not observable in the current and foreseeable experiments, because the final neutrino cannot be detected (the only observable signal of $CE\nu NS$ is the recoil of the nucleus, as explained above). However, specific flavor-changing NSI can be probed by their contribution to $CE\nu NS$.

The possible mediators γ , Z', and Φ in Fig. 1(b) correspond to different scenarios of physics beyond the SM:

I) The photon mediates electromagnetic interactions. In the SM and in the common belief neutrinos are exactly neutral and do not interact electromagnetically. However, in models BSM neutrinos can have small electromag-




Fig. 6: Current status and future projections for low-energy weak mixing angle measurements. The gray points show the measurements from APV on Cs atoms during the years [67–71]. The brown measurements refer to the COHERENT only and the combination between COHERENT and APV as determined in 2020 [36], while the dark blue and red points refer to the updated measurements reported in [21]. The projections for future CE ν NS experiments (CONUS [64], TEXONO [65], CONNIE [66], and MINER [11]), shown by the light blue triangles and dashed error bars, are taken from [72]. The purple triangles are the projections for the future electron scattering experiment MOLLER [73] and P2@MESA [74, 75]. The dark blue triangles with dashed error bars are the projections for the future CryoCsI-I and CryoCsI-II determinations [21].

netic interactions [76] which contribute to $CE\nu NS$ [17,77], due to three non-standard neutrino electromagnetic properties: a very small electric charge (often called "millicharge") [78], a small magnetic moment, and a small anomalous charge radius, which is the only neutrino electromagnetic property that is non-zero already in the SM. So far there is no experimental indication in favour of BSM neutrino electromagnetic interactions, but the search is very active [77]. Several studies have probed neutrino electromagnetic interactions through the analysis of $CE\nu NS$ data [29, 30, 32, 37, 38, 50, 51, 54, 55, 79, 80], finding competitive upper limits. In particular, $CE\nu NS$ processes permit to improve the limit on the ν_e charge radius and represent the only existing laboratory bounds on the muonic and tauonic neutrino electric charges [79]. Thanks to the lower energy of reactor antineutrinos and the low energy threshold of semiconductor detectors, the data from Dresden-II provide complementary information with respect to $CE\nu NS$ processes observed with ν 's produced at SNSs, with negligible dependence on the neutron distribution even if with larger backgrounds and larger dependence on the so-called quenching factor that is not wellconstrained at low energies.

II) The existence of a new massive Z' vector boson is predicted in several BSM scenarios [81]. Its mass can be very high, much larger than the so-called "electroweak scale" of about 100 GeV, or even as low as about 100 MeV. The search for the possible effects of these bosons is very active in different types of experiments [82], including

can make a difference as large as 11% [21].

CE ν NS [32, 50–52, 54, 55, 62, 79, 80, 83–87]. Existing data permit to put limits on a variety of vector boson mediator models: the so-called universal, the B - L and other anomaly-free U(1)' gauge models with direct couplings of the new vector boson with neutrinos and quarks, and the anomaly-free $L_e - L_\mu$, $L_e - L_\tau$, and $L_\mu - L_\tau$ gauge models where the coupling of the new vector boson with the quarks is generated by kinetic mixing with the photon at the one-loop level. The constraints related to models where the muon couples to the new boson mediator can also be compared with the values that can explain the muon g - 2 anomaly [88] and further data will be needed to fully exclude these interpretations of the anomaly [52].

III) The existence of a new massive Φ scalar boson is more exotic, but we cannot miss the chance to probe it in CE ν NS. Interesting constraints have already been put using existing data [32, 50–52, 54, 55, 79, 80, 85].

Furthermore, even more exotic models as generalized neutrino interactions [51, 89, 90], scalar leptoquarks [91], and extra dimensions [92] can be probed in $CE\nu NS$.

Conclusions. – The $CE\nu NS$ era started with the first measurement of such a process by the COHERENT experiment in 2017 [4], 44 years after its theoretical prediction in 1973 [1]. Since then there is great interest in this field, with many experiments already operating or planned [8–17] that will provide complementary information. The aim is to observe $CE\nu NS$ in different nuclei, confirming the theoretical prediction, and to obtain from the data information on the radius of the neutron distribution of the nucleus, on the weak mixing angle, and on new neutrino interactions. The search for effects of physics beyond the Standard Model is the main goal of current theoretical and experimental research in particle physics, because we know that the Standard Model, albeit very successful, cannot be the final theory of everything: for example, it cannot explain the smallness of neutrino masses, the existence of Dark Matter and Dark Energy, the matter-antimatter asymmetry in the Universe, and does not include gravity in a consistent theory [93]. Moreover, $CE\nu NS$ is also important for the search for Dark Matter, because $CE\nu NS$ events produced by solar and atmospheric neutrinos are expected to be detected in future experiments with high sensitivity. This detection would be very interesting for the study of neutrino properties, but it constitutes a background for Dark Matter searches that can limit their sensitivity, the so-called "neutrino floor/fog" [94,95], making thus of utmost importance to precisely study $CE\nu NS$ in direct detection dedicated experiments.

For all these reasons, we expect interesting $CE\nu NS$ news in the next years and we look forward to the future.

* * *

The work of C. Giunti is supported by the research grant

"The Dark Universe: A Synergic Multimessenger Approach" number 2017X7X85K under the program "PRIN 2017" funded by the Italian Ministero dell'Istruzione, Università e della Ricerca (MIUR).

REFERENCES

- [1] D.Z. Freedman, Phys. Rev. D9 (1974) 1389.
- [2] D.Z. Freedman, D.N. Schramm and D.L. Tubbs, Ann. Rev. Nucl. Part. Sci. 27 (1977) 167.
- [3] A. Drukier and L. Stodolsky, Phys. Rev. D30 (1984) 2295.
- [4] COHERENT, D. Akimov et al., Science 357 (2017) 1123, arXiv:1708.01294.
- [5] COHERENT, D. Akimov et al., Phys.Rev.Lett. 129 (2022) 081801, arXiv:2110.07730.
- [6] COHERENT, D. Akimov et al., Phys.Rev.Lett. 126 (2021) 012002, arXiv:2003.10630.
- [7] J. Colaresi et al., Phys.Rev.Lett. 129 (2022) 211802, arXiv:2202.09672.
- [8] H. Bonet et al., Phys.Rev.Lett. 126 (2021) 041804, arXiv:2011.00210.
- [9] CONNIE, A. Aguilar-Arevalo et al., JHEP 22 (2020) 017, arXiv:2110.13033.
- [10] D. Akimov et al., arXiv:2204.04575.
- [11] MINER, G. Agnolet et al., Nucl.Instrum.Meth. A853 (2017) 53, arXiv:1609.02066.
- [12] J. Choi et al., Eur.Phys.J.C 83 (2023) 226, arXiv:2204.06318.
- [13] G. Angloher et al., Eur.Phys.J. C79 (2019) 1018, arXiv:1905.10258.
- [14] D.Y. Akimov et al., JINST 17 (2022) T11011, arXiv:2209.15516.
- [15] RICOCHET, J. Colas et al., J.Low Temp.Phys. (2022), arXiv:2111.12856.
- [16] D. Baxter et al., JHEP 2002 (2020) 123, arXiv:1911.00762.
- [17] M. Abdullah et al., arXiv:2203.07361.
- [18] D.A. Sierra et al., JHEP 1912 (2019) 124, arXiv:1910.12437.
- [19] W. Maneschg, Recent results from the CONUS experiment, presentation at the Magnificent $CE\nu NS$ workshop (2023).
- [20] E. Figueroa-Feliciano, Experimental summary, presentation at the Magnificent $CE\nu NS$ workshop (2023).
- [21] M.A. Corona et al., arXiv:2303.09360.
- [22] I. Angeli and K.P. Marinova, Atom. Data Nucl. Data Tabl. 99 (2013) 69.
- [23] M. Thiel et al., J. Phys. G 46 (2019) 093003, arXiv:1904.12269.
- [24] PREX, D. Adhikari et al., Phys. Rev. Lett. 126 (2021) 172502, arXiv:2102.10767.
- [25] CREX, D. Adhikari et al., Phys.Rev.Lett. 129 (2022) 042501, arXiv:2205.11593.
- [26] M. Baldo and G.F. Burgio, Prog.Part.Nucl. Phys. 91 (2016) 203, arXiv:1606.08838.
- [27] C.J. Horowitz and J. Piekarewicz, Phys. Rev. Lett. 86 (2001) 5647, astro-ph/0010227.
- [28] M. Cadeddu et al., Phys.Rev.Lett. 120 (2018) 072501, arXiv:1710.02730.
- [29] M. Cadeddu et al., Phys.Rev. D98 (2018) 113010, arXiv:1810.05606.

- [30] M. Cadeddu et al., Phys.Rev. D101 (2020) 033004, arXiv:1908.06045.
- [31] D.K. Papoulias et al., Phys.Lett. B800 (2020) 135133, arXiv:1903.03722.
- [32] A.N. Khan and W. Rodejohann, Phys. Rev. D 100 (2019) 113003, arXiv:1907.12444.
- [33] X.R. Huang and L.W. Chen, Phys.Rev. D100 (2019) 071301, arXiv:1902.07625.
- [34] P. Coloma et al., JHEP 2008 (2020) 030, arXiv:2006.08624.
- [35] M. Hoferichter, J. Menendez and A. Schwenk, Phys.Rev. D102 (2020) 074018, arXiv:2007.08529.
- [36] M. Cadeddu et al., Phys.Rev.C 104 (2021) 065502, arXiv:2102.06153.
- [37] O.G. Miranda et al., JHEP 2005 (2020) 130, arXiv:2003.12050.
- [38] M. Cadeddu et al., Phys.Rev. D102 (2020) 015030, arXiv:2005.01645.
- [39] C.S. Wood et al., Science 275 (1997) 1759.
- [40] J. Guena, M. Lintz and M.A. Bouchiat, Phys. Rev. A71 (2005) 042108, physics/0412017.
- [41] M. Cadeddu and F. Dordei, Phys.Rev. D99 (2019) 033010, arXiv:1808.10202.
- [42] Particle Data Group, M. Tanabashi et al., Phys. Rev. D98 (2018) 030001.
- [43] V.A. Dzuba et al., Phys. Rev. Lett. 109 (2012) 203003, arXiv:1207.5864.
- [44] SLAC E158, P.L. Anthony et al., Phys. Rev. Lett. 95 (2005) 081601, hep-ex/0504049.
- [45] PVDIS, D. Wang et al., Nature 506 (2014) 67.
- [46] NuTeV, G.P. Zeller et al., Phys. Rev. Lett. 88 (2002) 091802, hep-ex/0110059.
- [47] Qweak, D. Androic et al., Nature 557 (2018) 207, arXiv:1905.08283.
- [48] M. Safronova et al., Rev.Mod.Phys. 90 (2018) 025008, arXiv:1710.01833.
- [49] M. Cadeddu et al., Phys.Rev.D 104 (2021) L011701, arXiv:2104.03280.
- $[50]\,$ P. Coloma et al., JHEP 05 (2022) 037, arXiv:2202.10829.
- [51] V. De Romeri et al., JHEP 04 (2023) 035, arXiv:2211.11905.
- [52] M.A. Corona et al., JHEP 05 (2022) 109, arXiv:2202.11002.
- [53] R.D. Young et al., Phys. Rev. Lett. 99 (2007) 122003, arXiv:0704.2618.
- [54] D.K. Papoulias and T.S. Kosmas, Phys.Rev. D97 (2018) 033003, arXiv:1711.09773.
- [55] D.K. Papoulias, Phys.Rev. D102 (2020) 113004, arXiv:1907.11644.
- [56] B. Dutta et al., Phys.Rev.Lett. 123 (2019) 061801, arXiv:1903.10666.
- [57] D. Aristizabal Sierra, V. De Romeri and N. Rojas, Phys.Rev. D98 (2018) 075018, arXiv:1806.07424.
- [58] B. Dutta et al., Phys.Rev.Lett. 124 (2020) 121802, arXiv:1906.10745.
- [59] M. Abdullah et al., Phys.Rev. D98 (2018) 015005, arXiv:1803.01224.
- [60] S.F. Ge and I.M. Shoemaker, JHEP 11 (2018) 066, arXiv:1710.10889.
- [61] O.G. Miranda et al., JHEP 12 (2021) 191, arXiv:2109.09545.
- [62] L.J. Flores, N. Nath and E. Peinado, JHEP 06 (2020)

045, arXiv:2002.12342.

- [63] B.K. Sahoo, B.P. Das and H. Spiesberger, Phys. Rev. D 103 (2021) L111303, 2101.10095.
- [64] M. Lindner, W. Rodejohann and X.J. Xu, JHEP 1703 (2017) 097, arXiv:1612.04150.
- [65] TEXONO, H.T. Wong et al., Phys. Rev. D75 (2007) 012001, hep-ex/0605006.
- [66] CONNIE, A. Aguilar-Arevalo et al., JINST 11 (2016) P07024, arXiv:1604.01343.
- [67] Particle Data Group, C. Amsler et al., Phys.Lett. B667 (2008) 1.
- [68] Particle Data Group, K. Nakamura et al., J. Phys. G37 (2010) 075021.
- [69] Particle Data Group, B.D. Fields, P. Molaro and S. Sarkar, Chin.Phys. C38 (2014) 090001, arXiv:1412.1408.
- [70] Particle Data Group, P. Zyla et al., PTEP 2020 (2020) 083C01.
- [71] Particle Data Group, R.L. Workman, PTEP 2022 (2022) 083C01.
- [72] B.C. Canas et al., Phys.Lett. B784 (2018) 159, arXiv:1806.01310.
- [73] MOLLER, J. Benesch et al., arXiv:1411.4088.
- [74] D. Becker et al., Eur. Phys. J. A 54 (2018) 208, arXiv:1802.04759.
- [75] P.S.B. Dev et al., JHEP 2106 (2021) 039, arXiv:2103.09067.
- [76] C. Giunti and A. Studenikin, Rev.Mod.Phys. 87 (2015) 531, arXiv:1403.6344.
- [77] C. Giunti et al., arXiv:2209.03340.
- [78] A. Das et al., Phys. Rev. D 102 (2020) 115009, arXiv:2005.12304.
- [79] M.A. Corona et al., JHEP 09 (2022) 164, arXiv:2205.09484.
- [80] J. Liao, H. Liu and D. Marfatia, Phys.Rev.D 106 (2022) 031702, arXiv:2202.10622.
- [81] P. Langacker, Rev. Mod. Phys. 81 (2009) 1199, arXiv:0801.1345.
- [82] T. Han et al., JHEP 1911 (2019) 028, arXiv:1910.03272.
- [83] J. Liao and D. Marfatia, Phys.Lett. B775 (2017) 54, arXiv:1708.04255.
- [84] P. Coloma et al., Phys.Rev. D96 (2017) 115007, arXiv:1708.02899.
- [85] J. Billard, J. Johnston and B.J. Kavanagh, JCAP 1811 (2018) 016, arXiv:1805.01798.
- [86] C. Giunti, Phys.Rev. D101 (2020) 035039, arXiv:1909.00466.
- [87] M. Cadeddu et al., JHEP 2101 (2021) 116, arXiv:2008.05022.
- [88] Muon g-2, B. Abi et al., Phys. Rev. Lett. 126 (2021) 141801, 2104.03281.
- [89] W.F. Chang and J. Liao, Phys.Rev. D102 (2020) 075004, arXiv:2002.10275.
- [90] A. Majumdar et al., Phys.Rev.D 106 (2022) 093010, arXiv:2208.13262.
- [91] R. Calabrese et al., Phys.Rev.D 107 (2023) 055039, arXiv:2212.11210.
- [92] A.N. Khan, JHEP 01 (2023) 052, arXiv:2208.09584.
- [93] P. Coloma et al., arXiv:2209.10362.
- [94] J. Cooley et al., arXiv:2209.07426.
- [95] C.A.J. O'Hare, Phys. Rev. Lett. 127 (2021) 251802, 2109.03116.