On *P*-Interpolation in Local Theory Extensions and Applications to the Study of Interpolation in the Description Logics $\mathcal{EL}, \mathcal{EL}^+$

Dennis Peuter, Viorica Sofronie-Stokkermans, and Sebastian Thunert

University of Koblenz, Koblenz, Germany {dpeuter,sofronie}@uni-koblenz.de, s.thunert@gmx.de

Abstract. We study the problem of *P*-interpolation, where *P* is a set of binary predicate symbols, for certain classes of local extensions of a base theory. For computing the *P*-interpolating terms, we use a hierarchic approach: This allows us to compute the interpolating terms using a method for computing interpolating terms in the base theory. We use these results for proving \leq -interpolation in classes of semilattices with monotone operators; we show, by giving a counterexample, that \leq -interpolation does not hold if by "shared" symbols we mean just the *common* symbols. We use these results for the study of \sqsubseteq -interpolation in the description logics \mathcal{EL} and \mathcal{EL}^+ .

1 Introduction

In this paper we study the problem of *P*-interpolation, a problem strongly related to interpolation w.r.t. logical theories. The problem can be formulated as follows:

Let \mathcal{T} be a theory, A and B be conjunctions of ground literals in the signature of \mathcal{T} , possibly with additional constants, P a set of binary predicate symbols in the signature of \mathcal{T} , $R \in P$, a a constant occurring in A and b a constant occurring in B. Assume that $A \wedge B \models_{\mathcal{T}} aRb$. Can we find a ground term tcontaining only constants and function symbols "shared" by A and B, such that $A \wedge B \models_{\mathcal{T}} aRt \wedge tRb$?

Interpolation has been studied in classical and non-classical logics and in extensions and combinations of theories; and is very important in program verification and also in the area of description logics. The first algorithms for interpolant generation in program verification required explicit constructions and "separations" of proofs [15,17]. In [14] interpolants are computed using variants of resolution. For certain theories, the "separation" of proofs relied on the possibility of "separating" atoms, i.e. on *P*-interpolation, where *P* is a set of binary predicate symbols. Equality interpolation is used in [36] for devising an interpolation method in combinations of theories with disjoint signatures. In [23,25] and [20], for instance, we consider interpolation problems in certain classes of extensions $\mathcal{T}_0 \cup \mathcal{K}$ of a base theory \mathcal{T}_0 and use a hierarchical approach to compute interpolants. The method relies on the *P*-interpolation property of the base theory \mathcal{T}_0 . In most of the applications we considered, P contains the equality predicate \approx or a predicate \leq with the property that in all models of \mathcal{T}_0 , the interpretation of \leq is a partial ordering.

Since at that time our main interest was the study of *interpolation problems*, in [23,25] and [20] *P*-interpolation is only used in order to help in giving methods for interpolation and not as a goal in itself. However, in several papers in the area of description logics (cf. e.g. [33,9]) when defining the notion of interpolation in description logics the authors define in fact a notion of \Box -interpolation. In [9] (Theorem 4) it is proved that \mathcal{EL}^+ allows interpolation (in fact, the notion of \Box -interpolation mentioned above) for *safe* role inclusions – this is related to the notion of "sharing" considered in [25], cf. also Section 4. The proof technique in [9] uses simulations. In this paper, we analyze the property of *P*-interpolation in theory extensions, propose a method for solving it based on hierarchical reasoning and formulate the Ξ -interpolation problem for \mathcal{EL} and \mathcal{EL}^+ as a \leq -interpolation problem in a theory of semilattices with operators.

We first studied \leq -interpolation in [18] in the context of description logics; the \sqsubseteq -interpolating concept descriptions were regarded as a form of "high-level" explanations. In this paper we further extend the work in [18]. The general approach we propose opens the possibility of applying similar methods to more general classes of non-classical logics (including e.g. substructural logics or the logics with monotone operators studied in [28,29]) or in verification (to consider more general theory extensions than those with uninterpreted function symbols analyzed in [20]). The main results can be summarized as follows:

- We propose variants of the definitions of convexity, *P*-interpolation and Beth definability relative to a subsignature.
- We describe a hierarchical *P*-interpolation method in certain classes of local theory extensions.
- We illustrate the applicability of these results to prove that certain classes of semilattices with monotone operators have the property of \leq -interpolation for a certain interpretation of "shared" function symbols.
- We show, by giving a counterexample, that ≤-interpolation does not hold if by "shared" symbols we mean just the *common* symbols.
- We indicate how these results can be used to prove or disprove various notions of interpolation for the description logics \mathcal{EL} and \mathcal{EL}^+ .

Structure of the paper: In Section 2 basic notions in logic are briefly introduced, and some results on convex theory and the link between two versions of \approx interpolation and corresponding versions of Beth definability – needed later in the paper – are proved. In Section 3 we introduce some results on local theory extensions needed in the paper. In Section 4 we identify classes of local theory extensions allowing *P*-interpolation and propose a hierarchical method of computing *P*-interpolants. This is used in Section 5 to study the existence of \leq interpolation in classes of semilattices with monotone operators. In Section 6 we use the links between the theory of semilattices with operators and the description logics \mathcal{EL} and \mathcal{EL}^+ , and show how the results can be used in the study of these logics. Section 7 contains the conclusions and some plans for future work. This paper is the extended version of [19]: it provides details of proofs and additional examples.

Table of Contents

1	Intro	oduction	1					
2	Preliminaries							
	2.1	Convexity and <i>P</i> -convexity						
	2.2	2.2 Equality interpolation, <i>R</i> -interpolation						
	2.3	Beth definability	6					
3	Local Theory Extensions							
	3.1	.1 Hierarchical reasoning in local theory extensions						
	3.2	Partial structures, weak validity	9					
	3.3	Flat and linear clauses; Flattening and purification	9					
	3.4	Recognizing locality and Ψ -locality	10					
4	R-in	terpolation in local theory extensions	12					
5	Exa	mple: Semilattices with monotone operators						
	5.1	The theory SLat of semilattices	17					
	5.2	Semilattices with operators	20					
6	Applications to \mathcal{EL} and \mathcal{EL}^+ -Subsumption							
	6.1	Algebraic semantics for $\mathcal{EL}, \mathcal{EL}^+$ and \sqsubseteq -interpolation	25					
	6.2	Example: \sqsubseteq -Interpolation for \mathcal{EL}^+	26					
	6.3	Prototype implementation	30					
7	Con	clusions and future work	30					

2 Preliminaries

We assume known standard definitions from first-order logic such as Π -structures, models, homomorphisms, logical entailment, satisfiability, unsatisfiability.

We consider signatures of the form $\Pi = (\Sigma, \mathsf{Pred})$, where Σ is a family of function symbols and Pred a family of predicate symbols. If C is a fixed countable set of fresh constants, we denote by Π^C the extension of Π with constants in C. We denote "falsum" with \bot .

A theory \mathcal{T} is described by a set of closed formulae (the axioms of the theory). We call a theory axiomatized by a set of universally quantified equations an *equational theory*. We denote by $\mathsf{Mod}(\mathcal{T})$ the set of all models of \mathcal{T} .

If F and G are formulae we write $F \models G$ to express the fact that every model of F is a model of G; if \mathcal{T} is a theory, we write $F \models_{\mathcal{T}} G$ to express the fact that every model of F which is also a model of \mathcal{T} is a model of G. The definitions can be extended in a natural way to the case when F is a set of formulae; in this case, $F \models_{\mathcal{T}} G$ if and only if $\mathcal{T} \cup F \models G$. $F \models_{\perp}$ means that F is unsatisfiable; $F \models_{\mathcal{T}} \bot$ means that there is no model of \mathcal{T} which is also a model of F. If there is a model of \mathcal{T} which is also a model of F we say that F is \mathcal{T} -consistent.

In the next sections we introduce notions of convexity of a theory with respect to a subset of the predicate symbols, and establish links between two versions 4 Dennis Peuter, Viorica Sofronie-Stokkermans, Sebastian Thunert

of \approx -interpolation and corresponding versions of Beth definability, results which are needed later in the paper

2.1 Convexity and *P*-convexity

We can define a notion of convexity w.r.t. a subset P of the set of predicates.

Definition 1. A theory \mathcal{T} with signature $\Pi = (\Sigma, \mathsf{Pred})$ is convex with respect to a subset P of Pred (which may include also equality \approx) if for all conjunctions Γ of ground Π^C -atoms (with additional constants in a set C), relations $R_1, \ldots, R_m \in P$ and tuples of Π^C -terms of corresponding arity $\overline{t}_1, \ldots, \overline{t}_m$ such that $\Gamma \models_{\mathcal{T}} \bigvee_{i=1}^m R_i(\overline{t}_i)$ there exists $i_0 \in \{1, \ldots, m\}$ such that $\Gamma \models_{\mathcal{T}} R_{i_0}(\overline{t}_{i_0})$.

We will call a theory \mathcal{T} convex if it is $\mathsf{Pred} \cup \{\approx\}$ -convex. The following result is well-known (cf. e.g. [5,11,34]):

Theorem 1 Let \mathcal{T} be a theory and let $Mod(\mathcal{T})$ be the class of models of \mathcal{T} .

- (i) If $Mod(\mathcal{T})$ is closed under direct products then \mathcal{T} is convex.
- (ii) If \mathcal{T} is a universal theory and \mathcal{T} is convex, then \mathcal{T} has an axiomatization given by Horn clauses, hence $\mathsf{Mod}(\mathcal{T})$ is closed under direct products.

Corollary 2 Let \mathcal{T}_1 , \mathcal{T}_2 be two theories with signatures Π_1 , Π_2 . If $\mathsf{Mod}(\mathcal{T}_1)$ and $\mathsf{Mod}(\mathcal{T}_2)$ are closed under direct products, then $\mathcal{T}_1 \cup \mathcal{T}_2$ is convex.

Proof. $\operatorname{Mod}(\mathcal{T}_1 \cup \mathcal{T}_2) = \{\mathcal{A} \mid \mathcal{A} \text{ is a } (\Pi_1 \cup \Pi_2) \text{-structure with } \mathcal{A}_{\mid \Pi_1} \in \operatorname{Mod}(\mathcal{T}_1) \text{ and } \mathcal{A}_{\mid \Pi_2} \in \operatorname{Mod}(\mathcal{T}_2)\}.$ Let $\mathcal{A}_i \in \operatorname{Mod}(\mathcal{T}_1 \cup \mathcal{T}_2), i \in I$. This is the case iff $\mathcal{A}_{i\mid \Pi_1} \in \operatorname{Mod}(\mathcal{T}_1) \text{ and } \mathcal{A}_{i\mid \Pi_2} \in \operatorname{Mod}(\mathcal{T}_2) \text{ for all } i \in I$. Taking into account that, due to the definition of a direct product of structures, $(\prod_{i \in I} \mathcal{A}_i)_{\mid \Pi_j} = \prod_{i \in I} (\mathcal{A}_i \mid \Pi_j)$ for j = 1, 2, and the fact that $\operatorname{Mod}(\mathcal{T}_1)$ and $\operatorname{Mod}(\mathcal{T}_2)$ are closed under direct products, it follows that $\prod_{i \in I} \mathcal{A}_i \in \operatorname{Mod}(\mathcal{T}_1 \cup \mathcal{T}_2)$, so $\operatorname{Mod}(\mathcal{T}_1 \cup \mathcal{T}_2)$ is closed under products. From Theorem 1 it follows that $\mathcal{T}_1 \cup \mathcal{T}_2$ is convex.

Corollary 3 Let \mathcal{T}_1 and \mathcal{T}_2 be convex universal theories. Then $\mathcal{T}_1 \cup \mathcal{T}_2$ is convex.

Proof: By Theorem 1, since \mathcal{T}_i is a universal theory and convex, $\mathsf{Mod}(\mathcal{T}_i)$ is closed under direct products, for i = 1, 2. By Corollary 2 it follows that $\mathsf{Mod}(\mathcal{T}_1 \cup \mathcal{T}_2)$ is closed under products, hence $\mathcal{T}_1 \cup \mathcal{T}_2$ is also convex.

In particular, every extension of a convex universal theory \mathcal{T}_0 with a set of new function symbols axiomatized by a set \mathcal{K} of Horn clauses is convex.

2.2 Equality interpolation, *R*-interpolation

We now define various versions of interpolation w.r.t. certain predicates.

Definition 2. We say that a convex theory \mathcal{T} has the equality interpolation property if for every conjunction of ground Π^C -literals $A(\overline{c}, \overline{a_1}, a)$ and $B(\overline{c}, \overline{b_1}, b)$, if $A \wedge B \models_{\mathcal{T}} a \approx b$ then there exists a term $t(\overline{c})$ containing only the constants \overline{c} shared by $A(\overline{c}, \overline{a_1}, a)$ and $B(\overline{c}, \overline{b_1}, b)$ such that $A \wedge B \models_{\mathcal{T}} a \approx t(\overline{c}) \wedge t(\overline{c}) \approx b$. Sometimes, the theories and theory extensions we study contain interpreted symbols in a set $\Pi_0 = (\Sigma_0, \mathsf{Pred})$ and non-interpreted function symbols in a set Σ_1 . The classical definition for equality interpolation for a theory \mathcal{T} mentioned above allows the term $t(\overline{c})$ to contain all function symbols in the signature of \mathcal{T} – these symbols are in this case all seen as being interpreted. If we distinguish between interpreted and uninterpreted functions we might allow the intermediate term $t(\overline{c})$ to contain any interpreted function symbols, but require that $t(\overline{c})$ contains only "shared" uninterpreted functions and common constants.

If Σ_A and Σ_B are the uninterpreted function symbols occurring in A resp. B, and Θ is a closure operator, by "shared" uninterpreted functions we can mean:

- Intersection-shared symbols: \bigcap -Shared $(A, B) = \Sigma_A \cap \Sigma_B$, or
- Θ -shared symbols: Θ -Shared $(A, B) = \Theta(\Sigma_A) \cap \Theta(\Sigma_B)$.

Example 1. Let $\mathcal{T} = \mathcal{T}_0 \cup \mathcal{K}$ be the extension of a theory \mathcal{T}_0 with set of interpreted function symbols Σ_0 with a set \mathcal{K} of clauses containing new uninterpreted function symbols in a set Σ_1 .

- If A and B are sets of atoms in the signature of \mathcal{T} containing additional constants in a set C and uninterpreted function symbols Σ_A , Σ_B then the *intersection-shared* uninterpreted function symbols of A and B are $\Sigma_A \cap \Sigma_B$.
- For every $f, g \in \Sigma_1$ we define $f \sim_{\mathcal{K}} g$ iff there exists $C \in \mathcal{K}$ such that f, g both occur in C. Let $\sim_{\mathcal{K}}^*$ be the equivalence relation on Σ_1 induced by $\sim_{\mathcal{K}}$. Let $\Theta_{\mathcal{K}}$ be defined for every $\Sigma \subseteq \Sigma_1$ by $\Theta_{\mathcal{K}}(\Sigma) = \bigcup_{f \in \Sigma} \{g \in \Sigma_1 \mid f \sim_{\mathcal{K}}^* g\}$. Then the $\Theta_{\mathcal{K}}$ -shared symbols are $\Theta_{\mathcal{K}}(\Sigma_A) \cap \Theta_{\mathcal{K}}(\Sigma_B)$. In particular, if A contains a function symbol f and B contains a symbol g such that f, g occur both in a clause in \mathcal{K} , then f and g are considered to be $\Theta_{\mathcal{K}}$ -shared by A, B.

We are interested in similar properties for other binary relations than \approx . We define an *R*-interpolation property, where *R* is a binary predicate symbol in Π .

Definition 3. Let $R \in \mathsf{Pred} \cup \{\approx\}$ be a binary predicate symbol. An $\{R\}$ -convex theory \mathcal{T} with uninterpreted symbols Σ_1 has the R-interpolation property if for all conjunctions of ground atoms $A(\overline{c}, \overline{a_1}, a)$ and $B(\overline{c}, \overline{b_1}, b)$, if $A \wedge B \models_{\mathcal{T}} aRb$ then there exists a term $t(\overline{c})$ containing only common constants \overline{c} and only "shared" uninterpreted symbols in Σ_1 such that $A \wedge B \models_{\mathcal{T}} aRt(\overline{c}) \wedge t(\overline{c})Rb$.

If $P \subseteq \mathsf{Pred}$, we say that a theory has the P-interpolation property if it has the R-interpolation property for every $R \in P$.

In Section 5 we give examples of theories with this property and show that a theory may not have the *R*-interpolation property for a predicate symbol *R* if we use the notion of *intersection-shared symbols*, but has the *R*-interpolation property if we consider the less restrictive notion of Θ -shared symbols for a suitably defined closure operator Θ .

6 Dennis Peuter, Viorica Sofronie-Stokkermans, Sebastian Thunert

2.3 Beth definability

Let \mathcal{T} be a theory with signature $\Pi = (\Sigma_0 \cup \Sigma_1, \mathsf{Pred})$, where the function symbols in Σ_0 are regarded as interpreted function symbols and the function symbols in Σ_1 are regarded as uninterpreted function symbols, and let C be a set of additional constants. We define a notion of Beth definability relative to a subset $\Sigma_S \subseteq \Sigma_1 \cup C$ of non-interpreted function symbols and constants (similar to the one introduced in [33]), which we refer to as Σ_S -Beth definability. Let $\Sigma_S \subseteq \Sigma_1 \cup C$, let $\Sigma_r = \Sigma_1 \setminus \Sigma_S$, and let $\Pi' = (\Sigma_0 \cup (\Sigma_S \cap \Sigma_1) \cup \Sigma'_r, \mathsf{Pred})$, where $\Sigma'_r = \{f' \mid f \in \Sigma_1 \setminus \Sigma_S\}$ is the signature obtained by replacing all uninterpreted function symbols in Σ_1 which are not in Σ_S with new primed copies. If ϕ is a Π^C -formula, we will denote by ϕ' the formula obtained from ϕ by re-

placing all uninterpreted function symbols in $\Sigma_1 \setminus \Sigma_S$ and all constants in $C \setminus \Sigma_S$ with distinct, primed versions. The interpreted function symbols and the uninterpreted function symbols and constants in Σ_S are not changed. We regard the theory \mathcal{T} as a set of formulae; let $\mathcal{T}' := \{\phi' \mid \phi \in \mathcal{T}\}.$

Definition 4. Let A be a conjunction of ground Π^{C} -literals, and $a \in C$.

- We say that a is implicitly defined by A w.r.t. Σ_S and \mathcal{T} if, with the notations introduced before, $A \wedge A' \models_{\mathcal{T} \cup \mathcal{T}'} a \approx a'$.
- We say that a is explicitly defined by A w.r.t. Σ_S and \mathcal{T} if there exists a term t containing only symbols in Σ_0 , Pred and Σ_S such that $A \models_{\mathcal{T}} a \approx t$.

Definition 5. Let \mathcal{T} be a theory with uninterpreted function symbols in a set Σ_1 . Let $\Sigma_S \subseteq \Sigma_1 \cup C$. \mathcal{T} has the Beth definability property w.r.t. Σ_S (Σ_S -Beth definability), if for every conjunction of literals A and every $a \in C$, if A implicitly defines a w.r.t. Σ_S and \mathcal{T} then A explicitly defines a w.r.t. Σ_S and \mathcal{T} .

In [4,6] it was proved that if a convex theory \mathcal{T} has the \approx -interpolation property, then it has a version of Beth definability property which can be regarded as the extension of the classical Beth definability property [7] so as to take into account the theory \mathcal{T} . We give an analogous implication between \approx -interpolation and Beth definability w.r.t. a subsignature.

Theorem 4 Let \mathcal{T} be a convex theory with signature $\Pi = (\Sigma_0 \cup \Sigma_1, \mathsf{Pred}), C$ a set of constants, and $\Sigma_S \subseteq \Sigma_1 \cup C$. Let \mathcal{T}' be as defined above.

- (i) If $\mathcal{T} \cup \mathcal{T}'$ has the \approx -interpolation property with intersection-sharing, then \mathcal{T} has the Σ_S -Beth definability property.
- (ii) Assume that $\mathcal{T} = \mathcal{T}_0 \cup \mathcal{K}$ where all symbols in the signature of \mathcal{T}_0 are regarded as interpreted, and \mathcal{K} is a set of clauses also containing uninterpreted function symbols in Σ_1 . Let $\Theta_{\mathcal{K}}$ be the closure operator defined in Example 1. If $\mathcal{T} \cup \mathcal{T}'$ has the \approx -interpolation property with $\Theta_{\mathcal{K} \cup \mathcal{K}'}$ -sharing, then \mathcal{T} has the $\Theta_{\mathcal{K}}(\Sigma_S)$ -Beth definability property.

Proof: (i) Assume a is implicitly definable w.r.t. Σ_S , i.e. there exists a conjunction A of literals such that if A' is obtained by renaming the symbols in $(\Sigma_1 \cup C) \setminus \Sigma_S$

as explained at the beginning of this subsection, we have $A \wedge A' \models_{\mathcal{T} \cup \mathcal{T}'} a \approx a'$. Since $\mathcal{T} \cup \mathcal{T}'$ has the \approx -interpolation property, there exists a term t using only the functions and predicate symbols common to A and A' such that $A \wedge A' \models_{\mathcal{T} \cup \mathcal{T}'} a \approx t \wedge t \approx a'$. The only function symbols shared by A and A' are the symbols in $\Sigma_0 \cup \Sigma_S$. We show that $A \models_{\mathcal{T}} a \approx t$. Let \mathcal{A} be a model of \mathcal{T} in which A is true. We define new function symbols $\{f'_{\mathcal{A}} \mid f' \in \Sigma'_r\}$ by $f'_{\mathcal{A}} = f_{\mathcal{A}}$, and interpretations $c'_{\mathcal{A}} := c_{\mathcal{A}}$ for all constants not in Σ_S . The structure $\overline{\mathcal{A}}$ obtained this way is a model of $(\mathcal{T} \cup \mathcal{T}')$ and of $A \wedge A'$. Therefore, since $A \wedge A' \models_{\mathcal{T} \cup \mathcal{T}'} a \approx t \wedge t \approx a'$, we have $\overline{\mathcal{A}} \models a \approx t$, hence $\mathcal{A} \models a \approx t$. Therefore, $A \models_{\mathcal{T}} a \approx t$.

(ii) Assume *a* is implicitly definable w.r.t. $\Theta_{\mathcal{K}}(\Sigma_S)$, i.e. there exists a conjunction A of literals such that if A' is obtained from A and $\mathcal{T}' = \mathcal{T}_0 \cup \mathcal{K}'$ is obtained from $\mathcal{T} = \mathcal{T}_0 \cup \mathcal{K}$ by renaming as explained at the beginning of this subsection, then $A \wedge A' \models_{\mathcal{T} \cup \mathcal{T}'} a \approx a'$. The function symbols shared by A and A' are the symbols in $\Sigma_0 \cup \Theta_{\mathcal{K} \cup \mathcal{K}'}(\Sigma_S)$, where:

 $\begin{array}{l} \Theta_{\mathcal{K}\cup\mathcal{K}'}(\varSigma_S) = \bigcup_{f\in\varSigma_S} \{g\in\varSigma_1\cup\varSigma_1'\mid f\sim^*_{\mathcal{K}\cup\mathcal{K}'}g\} \\ \text{Note that:} \end{array}$

 $\begin{array}{l} \Theta_{\mathcal{K}}(\varSigma_S) = \bigcup_{f \in \varSigma_S} \{g \in \varSigma_1 \mid f \sim_{\mathcal{K}}^* g\} \text{ and } \Theta_{\mathcal{K}'}(\varSigma_S) = \bigcup_{f \in \varSigma_S} \{g \in \varSigma_1' \mid f \sim_{\mathcal{K}'}^* g\}. \\ \text{It is easy to see that for every } f \in \varSigma_1 \backslash \varSigma_S, \ f \in \Theta_{\mathcal{K}}(\varSigma_S) \text{ iff } f' = \Theta_{\mathcal{K}'}(\varSigma_S), \text{ and } \\ \Theta_{\mathcal{K} \cup \mathcal{K}'}(\varSigma_S) = \Theta_{\mathcal{K}}(\varSigma_S) \cup \Theta_{\mathcal{K}'}(\varSigma_S). \end{array}$

Since we assumed that $\mathcal{T} \cup \mathcal{T}'$ has the \approx -interpolation property with the notion of $\Theta_{\mathcal{K}\cup\mathcal{K}'}$ -sharing, there exists a term t over the signature $\Sigma_0\cup\Theta_{\mathcal{K}\cup\mathcal{K}'}(\Sigma_S)$ such that $A \wedge A' \models_{\mathcal{T} \cup \mathcal{T}'} a \approx t \wedge t \approx a$. The term t might contain primed versions of function symbols. We show that we can find a term \overline{t} containing only terms in $\Theta_{\mathcal{K}}(\Sigma_S)$ such that $A \models_{\mathcal{T}} a \approx \overline{t}$. Let \mathcal{A} be a Π^C -model of $\mathcal{T} = \mathcal{T}_0 \cup \mathcal{K}$ in which A is true. For every $f \in \Sigma_1 \setminus \Theta_{\mathcal{K}}(\Sigma_S)$ we define $f'_{\mathcal{A}} = f_{\mathcal{A}}$, and for every constant c not in Σ_S we define $c'_{\mathcal{A}} := c_{\mathcal{A}}$. The structure $\overline{\mathcal{A}}$ obtained this way is a model of $\mathcal{T}_0 \wedge \mathcal{K}$ and of A (because \mathcal{A} was a model of $\mathcal{T}_0 \cup \mathcal{K}$ and of A), a model of $\mathcal{T}_0 \cup \mathcal{K}'$ and of A' (because of the definition of the primed symbols), so $\overline{\mathcal{A}} \models a \approx t$. Since for $f \in \Sigma_1 \setminus \Theta_{\mathcal{K}}(\Sigma_S)$ the functions f and f' are defined in the same way in \mathcal{A} , and the primed and unprimed versions of constants in $C \setminus \Sigma_S$ are interpreted the same way in $\overline{\mathcal{A}}$, we can replace in t all primed function symbols with the non-primed versions, and its value does not change in $\overline{\mathcal{A}}$. Let \overline{t} be the term obtained this way. The value of \overline{t} in \mathcal{A} is equal to the value of t in $\overline{\mathcal{A}}$, i.e. is equal to the value of a in \mathcal{A} . Hence, $\mathcal{A} \models a \approx \overline{t}$.

3 Local Theory Extensions

Let $\Pi_0 = (\Sigma_0, \mathsf{Pred})$ be a signature, and \mathcal{T}_0 be a "base" theory with signature Π_0 . We consider extensions $\mathcal{T} := \mathcal{T}_0 \cup \mathcal{K}$ of \mathcal{T}_0 with new function symbols Σ_1 (*extension functions*) whose properties are axiomatized using a set \mathcal{K} of (universally closed) clauses in the extended signature $\Pi = (\Sigma_0 \cup \Sigma_1, \mathsf{Pred})$, which contain function symbols in Σ_1 .

If G is a finite set of ground Π^C -clauses, where C is an additional set of constants, and \mathcal{K} a set of Π -clauses, we will denote by $\mathsf{st}(\mathcal{K}, G)$ the set of all ground terms 8

which occur in G or \mathcal{K} , and by $est(\mathcal{K}, G)$ the set of all extension ground terms, i.e. terms starting with a function in Σ_1 which occur in G or \mathcal{K} .

In this paper we regard every finite set G of ground clauses as the ground formula $\bigwedge_{C \in G} C$. If T is a set of ground terms in the signature Π^C , we denote by $\mathcal{K}[T]$ the set of all instances of \mathcal{K} in which the terms starting with a function symbol in Σ_1 are in T. Let Ψ be a map associating with every finite set T of ground terms a finite set $\Psi(T)$ of ground terms containing T. For any set G of ground Π^C -clauses we write $\mathcal{K}[\Psi_{\mathcal{K}}(G)]$ for $\mathcal{K}[\Psi(\mathsf{est}(\mathcal{K},G))]$. We define the following condition:

 $(\mathsf{Loc}_{f}^{\Psi})$ For every finite set G of ground clauses in Π^{C} it holds that $\mathcal{T}_{0} \cup \mathcal{K} \cup G \models \bot$ if and only if $\mathcal{T}_{0} \cup \mathcal{K}[\Psi_{\mathcal{K}}(G)] \cup G$ is unsatisfiable.

Extensions satisfying condition $(\mathsf{Loc}_{f}^{\Psi})$ are called Ψ -local. If Ψ is the identity we obtain the notion of local theory extensions [22]; if in addition \mathcal{T}_{0} is the theory of pure equality we obtain the notion of local theories [16,10].

3.1 Hierarchical reasoning in local theory extensions

Consider a Ψ -local theory extension $\mathcal{T}_0 \subseteq \mathcal{T}_0 \cup \mathcal{K}$. Condition (Loc_f^{Ψ}) requires that for every finite set G of ground Π^C -clauses, $\mathcal{T}_0 \cup \mathcal{K} \cup G \models \bot$ iff $\mathcal{T}_0 \cup \mathcal{K}[\Psi_{\mathcal{K}}(G)] \cup G \models \bot$. In all clauses in $\mathcal{K}[\Psi_{\mathcal{K}}(G)] \cup G$ the function symbols in Σ_1 only have ground terms as arguments, so $\mathcal{K}[\Psi_{\mathcal{K}}(G)] \cup G$ can be flattened and purified by introducing, in a bottom-up manner, new constants $c_t \in C$ for subterms $t=f(c_1,\ldots,c_n)$ where $f\in\Sigma_1$ and c_i are constants, together with definitions $c_t=f(c_1,\ldots,c_n)$. We thus obtain a set of clauses $\mathcal{K}_0\cup G_0\cup \mathsf{Def}$, where \mathcal{K}_0 and G_0 do not contain Σ_1 -function symbols and Def contains clauses of the form $c=f(c_1,\ldots,c_n)$, where $f\in\Sigma_1, c, c_1,\ldots,c_n$ are constants.

Theorem 5 ([22,12,13]) Let \mathcal{K} be a set of clauses. Assume that $\mathcal{T}_0 \subseteq \mathcal{T}_0 \cup \mathcal{K}$ is a Ψ -local theory extension. For any finite set G of flat ground clauses (with no nestings of extension functions), let $\mathcal{K}_0 \cup G_0 \cup \mathsf{Def}$ be obtained from $\mathcal{K}[\Psi_{\mathcal{K}}(G)] \cup G$ by flattening and purification, as explained above. Then the following are equivalent to $\mathcal{T}_0 \cup \mathcal{K} \cup G \models \perp$:

(i)
$$\mathcal{T}_0 \cup \mathcal{K}[\Psi_{\mathcal{K}}(G)] \cup G \models \bot$$
.
(ii) $\mathcal{T}_0 \cup \mathcal{K}_0 \cup G_0 \cup \mathsf{Con}_0 \models \bot$, where $\mathsf{Con}_0 = \{\bigwedge_{i=1}^n c_i \approx d_i \rightarrow c \approx d | \substack{f(c_1, \dots, c_n) \approx c \in \mathsf{Def} \\ f(d_1, \dots, d_n) \approx d \in \mathsf{Def}} \}$

The locality of a theory extension can be recognized by proving embeddability of partial models into total models [22,29,12]. In [13] we showed that for extensions with sets of flat and linear clauses Ψ -locality can be checked by checking whether an embeddability condition of partial into total models holds. In [27] we mention (without proof) that the proof in [13] can be extended to situations in which the clauses in \mathcal{K} are not linear. A full proof of this result is given in Section 3.4. For presenting the proof we need to introduce notions such as partial structures and truth in partial structures.

P-interpolation in local theory extensions and applications to $\mathcal{EL}, \mathcal{EL}^+$

3.2 Partial structures, weak validity

We first introduce notions such as partial structures, truth in partial structures, and weak partial models for clauses resp. for sets of clauses.

Definition 6. A partial Π -structure is a structure $\mathcal{A} = (A, \{f_{\mathcal{A}}\}_{f \in \Sigma}, \{p_{\mathcal{A}}\}_{p \in \mathsf{Pred}}),$ where A is a non-empty set and for every $f \in \Sigma$ with arity $n, f_{\mathcal{A}}$ is a partial function from A^n to A and for every $p \in \mathsf{Pred}$ with arity $m, p_{\mathcal{A}} \subseteq A^m$. A (total) Π -structure is a partial Π -structure where all functions $f_{\mathcal{A}}$ are total.

The notion of evaluating a term t w.r.t. a variable assignment $\beta : X \to \mathcal{A}$ for its variables in a partial structure \mathcal{A} is the same as for a total structure, except that this evaluation is undefined if $t = f(t_1, \ldots, t_n)$ and either one of $\beta(t_i)$ is undefined, or else $(\beta(t_1), \ldots, \beta(t_n))$ is not in the domain of definition of $f_{\mathcal{A}}$.

Definition 7. Let D be a clause. We say that (\mathcal{A}, β) is a weak partial model of D (notation: $(\mathcal{A}, \beta) \models_w D$) if either at least one term in D is not defined in β or all terms are defined and at least one literal L of D holds in (\mathcal{A}, β) .

Definition 8. A total map $h: A \to B$ between partial Π -structures \mathcal{A} and \mathcal{B} is called a weak Π -homomorphism if whenever $f_A(a_1, \ldots, a_n)$ is defined in A, also $f_B(h(a_1), \ldots, h(a_n))$ is defined in B and $h(f_A(a_1, \ldots, a_n)) = f_B(h(a_1), \ldots, h(a_n))$. A partial structure \mathcal{A} weakly embeds into a (total) structure \mathcal{B} if there exists an injective weak Π -homomorphism h from \mathcal{A} to \mathcal{B} such that for every $R \in \mathsf{Pred}$ and for every $a, b \in \mathcal{A}$ we have: a Rb iff h(a)Rh(b).

3.3 Flat and linear clauses; Flattening and purification

We define flatness for non-ground clauses and for ground clauses.

Definition 9 (Flatness and linearity for non-ground clauses). A nonground clause is Σ_1 -flat if function symbols (including constants) do not occur as arguments of function symbols in Σ_1 . A Σ_1 -flat non-ground clause is called Σ_1 -linear if whenever a variable occurs in two terms in the clause which start with function symbols in Σ_1 , the two terms are identical, and if no term which starts with a function symbol in Σ_1 contains two occurrences of the same variable.

Definition 10 (Flatness and linearity for ground clauses). We say that a ground clause is Σ_1 -flat if only constants appear as arguments of function symbols in Σ_1 . A Σ_1 -flat ground clause is Σ_1 -linear if whenever a constant occurs in two terms in the clause starting with function symbols in Σ_1 , the two terms are identical, and if no term starting with a function symbol in Σ_1 contains two occurrences of the same constant.

Any set G of ground clauses in a signature Σ containing Σ_1 can be transformed into a set $G_{\text{flin}(\Sigma_1)}$ of ground clauses in which subterms starting with function symbols in Σ_1 are flat and linear. This can be done by introducing, in a bottomup manner, new constants for subterms occurring below functions in Σ_1 , and adding the corresponding definitions to the set of clauses.

9

10 Dennis Peuter, Viorica Sofronie-Stokkermans, Sebastian Thunert

A set G of ground clauses can be transformed into a *purified* set of clauses $G_{sep(\Sigma_1)}$ (i.e. the function symbols in Σ_1 are separated from the other symbols) by introducing, in a bottom-up manner, new constants c_t for subterms $t = f(g_1, \ldots, g_n)$ with $f \in \Sigma_1$, g_i ground $\Sigma_0 \cup \Sigma_c$ -terms (where Σ_c is a set of constants which contains the constants introduced by flattening), together with corresponding definitions $c_t \approx t$. These transformations preserves satisfiability and unsatisfiability with respect to total algebras, and also with respect to partial algebras in which all ground subterms which are flattened are defined.

3.4 Recognizing locality and Ψ -locality

We now give a semantic criterion for Ψ -locality.

Definition 11. With the above notations, let Ψ be a map associating with \mathcal{K} and a set of ground terms T a set $\Psi_{\mathcal{K}}(T)$ of ground terms. We call $\Psi_{\mathcal{K}}$ a term closure operator if the following holds for all sets of ground terms T, T':

- (i) $\operatorname{est}(\mathcal{K}, T) \subseteq \Psi_{\mathcal{K}}(T),$
- (*ii*) $T \subseteq T' \Rightarrow \Psi_{\mathcal{K}}(T) \subseteq \Psi_{\mathcal{K}}(T'),$
- (*iii*) $\Psi_{\mathcal{K}}(\Psi_{\mathcal{K}}(T)) \subseteq \Psi_{\mathcal{K}}(T),$
- (iv) for any map $h: C \to C$, $\bar{h}(\Psi_{\mathcal{K}}(T)) = \Psi_{\mathcal{K}}(\bar{h}(T))$, where \bar{h} is the canonical extension of h to extension ground terms.

We will use the following notation: If A is a set (for instance the support of a Π -structure), we denote by Π^A the extension of Π with the elements in A, seen as additional constants. Let $\mathcal{T} = \mathcal{T}_0 \cup \mathcal{K}$ be a theory extension with set of clauses \mathcal{K} and let $\Psi_{\mathcal{K}}$ be a closure operator.

In what follows, $\mathsf{PMod}^{\Psi}_{\mathsf{w}}(\Sigma_1, \mathcal{T})$ denotes the class of all partial structures $\mathcal{A} = (A, \{f_{\mathcal{A}}\}_{f \in \Sigma_0 \cup \Sigma_1}, \{p_{\mathcal{A}}\}_{p \in \mathsf{Pred}})$ in which the Σ_0 -functions are total, and $\mathcal{A}_{|\Pi_0}$ is a model of \mathcal{T}_0 , the Σ_1 -functions are partial and \mathcal{A} is a weak partial model of all the clauses in \mathcal{K} , and in which the set of Π^A terms

 $\mathsf{Def}(A) = \{ f(a_1, \dots, a_n) \mid a_1, \dots, a_n \in A, f_{\mathcal{A}}(a_1, \dots, a_n) \text{ defined} \}$

is closed under $\Psi_{\mathcal{K}}$.

For extensions $\mathcal{T}_0 \subseteq \mathcal{T} = \mathcal{T}_0 \cup \mathcal{K}$, where \mathcal{K} is a set of clauses, we consider the condition:

 $(\mathsf{Emb}_{\mathsf{w}}^{\Psi})$ Every $A \in \mathsf{PMod}_{\mathsf{w}}^{\Psi}(\Sigma_1, \mathcal{T})$ weakly embeds into a total model of \mathcal{T} .

Theorem 6 Let \mathcal{K} be a set of Σ_1 -flat clauses, and $\Psi_{\mathcal{K}}$ be a term closure operator, which satisfies conditions (i)–(iv) above and also satisfies the additional condition that for every set T of ground terms and for every clause D in \mathcal{K} , if a variable occurs in two terms in D then either the two terms are identical, or the variable occurs below two different unary function symbols f and g and, for every constant c, f(c) is in $\Psi_{\mathcal{K}}(T)$ iff g(c) is in $\Psi_{\mathcal{K}}(T)$. If the extension $\mathcal{T}_0 \subseteq \mathcal{T} = \mathcal{T}_0 \cup \mathcal{K}$ satisfies $(\mathsf{Emb}_{\Psi}^{\mathsf{W}})$ then the extension satisfies (Loc^{Ψ}) . Proof: Assume that $\mathcal{T}_0 \cup \mathcal{K}$ is not a Ψ -local extension of \mathcal{T}_0 . Then there exists a set G of ground clauses (with additional constants) such that $\mathcal{T}_0 \cup \mathcal{K} \cup G \models \bot$ but $\mathcal{T}_0 \cup \mathcal{K}[\Psi_{\mathcal{K}}(G)] \cup G$ has a model \mathcal{B} . We assume w.l.o.g. that $G = G_0 \cup G_1$, where G_0 contains no function symbols in Σ_1 and G_1 consists of ground unit clauses of the form $f(c_1, \ldots, c_n) \approx c$, where c_i, c are constants in $\Sigma_0 \cup C$ and $f \in \Sigma_1$.

We construct another structure, \mathcal{A} , having the same support $U_{\mathcal{A}}$ as \mathcal{B} , which inherits all relations in Pred and all maps in $\Sigma_0 \cup C$ from \mathcal{B} , but on which the domains of definition of the Σ_1 -functions are restricted as follows: for every $f \in \Sigma_1, f_{\mathcal{A}}(a_1, \ldots, a_n)$ is defined if and only if there exist constants c^1, \ldots, c^n such that $f(c^1, \ldots, c^n)$ is in $\Psi_{\mathcal{K}}(G)$ and $a_i = c_{\mathcal{B}}^i$ for all $i \in \{1, \ldots, n\}$. In this case we define $f_{\mathcal{A}}(a_1, \ldots, a_n) := f_{\mathcal{B}}(c_{\mathcal{B}}^1, \ldots, c_{\mathcal{B}}^n)$. The reduct of \mathcal{A} to $(\Sigma_0 \cup C, \operatorname{Pred})$ coincides with that of \mathcal{B} . Thus, \mathcal{A} is a model of $\mathcal{T}_0 \cup G_0$. By the way the operations in Σ_1 are defined in \mathcal{A} it is clear that \mathcal{A} satisfies G_1 , so \mathcal{A} satisfies G.

We show that $\mathcal{A} \models_w \mathcal{K}$. Let D be a clause in \mathcal{K} . If D is ground then all its terms are defined (and all terms starting with an extension function are contained in $\Psi_{\mathcal{K}}(G)$), i.e. $D \in \mathcal{K}[\Psi_{\mathcal{K}}(G)]$, so D is true in \mathcal{B} , hence it is also true in \mathcal{A} .

Now consider the case in which D is not ground. Let $\beta : X \to \mathcal{A}$ be an arbitrary valuation. Again, if there is a term t in D such that $\beta(t)$ is undefined, we immediately have that (\mathcal{A}, β) weakly satisfies D. So let us suppose that for all terms t occurring in D, $\beta(t)$ is defined. We associate with β a substitution σ as follows: Let x be a variable. We have the following possibilities:

Case 1: x does not occur below any extension function. This case is unproblematic. We can define $\sigma(x)$ arbitrarily.

Case 2: x occurs in a unique term $t = f(\dots x \dots y \dots)$ (which may occur more than once in the clause D). From the fact that $\beta(t)$ is defined, we know that there are ground terms (in fact constants) which we will denote by t_x, t_y, \dots such that $\beta(x) = (t_x)_{\mathcal{B}}, \beta(y) = (t_y)_{\mathcal{B}}, \dots, \beta(t) = f_A(\dots(t_x)_{\mathcal{B}} \dots (t_y)_{\mathcal{B}} \dots)$, and $f(\dots, t_x, \dots, t_y, \dots) \in \Psi_{\mathcal{K}}(G)$. We can define $\sigma(x) = t_x$.

Case 3: x occurs in two terms of the form f(x), g(x) in the clause D. By assumption, $\Psi_{\mathcal{K}}$ has the property that for every constant $c, f(c) \in \Psi_{\mathcal{K}}(G)$ iff $g(c) \in \Psi_{\mathcal{K}}(G)$. From the fact that $\beta(f(x))$ and $\beta(g(x))$ are defined, we know that there are ground terms which we will denote by t_x, s_x such that $\beta(x) = (t_x)_{\mathcal{B}} = (s_x)_{\mathcal{B}}$, $\beta(f(x)) = f(t_x)_{\mathcal{B}}$ and $\beta(g(x)) = g(s_x)_{\mathcal{B}}$, and $f(t_x), g(s_x) \in \Psi_{\mathcal{K}}(G)$. Since all terms in $\Psi_{\mathcal{K}}(G)$ are flat, t_x and s_x are constants. Assume that $t_x = c$. We know that if $f(c) \in \Psi_{\mathcal{K}}(G)$ then $g(c) \in \Psi_{\mathcal{K}}(G)$ and if $g(c) \in \Psi_{\mathcal{K}}(G)$ then $f(c) \in \Psi_{\mathcal{K}}(G)$, so we can choose $s_x = t_x = c$. Also in this case we can define $\sigma(x) = t_x = c$.

Thus, we can construct a substitution σ with $\sigma(D) \in \mathcal{K}[\Psi_{\mathcal{K}}(G)]$ and $\beta \circ \sigma = \beta$. As $(\mathcal{B}, \beta) \models \sigma(D)$ we can infer $(\mathcal{A}, \beta) \models_w D$. We now show that

$$\mathsf{Def}(\mathcal{A}) = \{ f(a_1, \dots, a_n) \mid a_1, \dots, a_n \in U_{\mathcal{A}}, f_{\mathcal{A}}(a_1, \dots, a_n) \text{ defined} \}$$

is closed under $\Psi_{\mathcal{K}}$. By definition, $f(a_1, \ldots, a_n) \in \mathsf{Def}(\mathcal{A})$ iff there exist constants c_1, \ldots, c_n with $c_{i\mathcal{A}} = a_i$ for all i and $f(c_1, \ldots, c_n) \in \Psi_{\mathcal{K}}(G)$. Thus,

$$\mathsf{Def}(\mathcal{A}) = \{f(a_1, \dots, a_n) \mid a_1, \dots, a_n \in U_{\mathcal{A}}, f_{\mathcal{A}}(a_1, \dots, a_n) \text{ defined} \}$$

= $\{f(c_{1,\mathcal{A}}, \dots, c_{n,\mathcal{A}}) \mid c_i \text{ constants with } f(c_1, \dots, c_n) \in \Psi_{\mathcal{K}}(G) \}$
= $\overline{h}(\Psi_{\mathcal{K}}(G))$ where $h(c_i) = a_i \text{ for all } i$
 $\Psi_{\mathcal{K}}(\mathsf{Def}(\mathcal{A})) = \Psi_{\mathcal{K}}(\overline{h}(\Psi_{\mathcal{K}}(G))) = \overline{h}(\Psi_{\mathcal{K}}(\Psi_{\mathcal{K}}(G)))$ by property (iv) of Ψ
 $\subseteq \overline{h}(\Psi_{\mathcal{K}}(G)) = \mathsf{Def}(\mathcal{A})$ by property (iii) of Ψ

As $\mathcal{A} \models_w \mathcal{K}$, \mathcal{A} weakly embeds into a total structure $\overline{\mathcal{A}}$ satisfying $\mathcal{T}_0 \cup \mathcal{K}$. But then $\overline{\mathcal{A}} \models G$, so $\overline{\mathcal{A}} \models \mathcal{T}_0 \cup \mathcal{K} \cup G$, which is a contradiction. \Box

4 *R*-interpolation in local theory extensions

In [25] we considered convex and *P*-interpolating theories \mathcal{T}_0 with signature $\Pi_0 = (\Sigma_0, \mathsf{Pred})$ (where $P \subseteq \mathsf{Pred}$). We studied Ψ -local extensions $\mathcal{T} = \mathcal{T}_0 \cup \mathcal{K}$ of a theory \mathcal{T}_0 with new function symbols in a set Σ_1 axiomatized by a set \mathcal{K} of clauses, with the property that all clauses in \mathcal{K} are of the form:

$$\begin{cases} x_1 R_1 s_1 \wedge \dots \wedge x_n R_n s_n \to f(x_1, \dots, x_n) R g(y_1, \dots, y_n) \\ x_1 R_1 y_1 \wedge \dots \wedge x_n R_n y_n \to f(x_1, \dots, x_n) R f(y_1, \dots, y_n) \end{cases}$$
(1)

where $n \geq 1, x_1, \ldots, x_n, y_1, \ldots, y_n$ are variables, $f, g \in \Sigma_1, R_1, \ldots, R_n, R$ are binary relations with $R_1, \ldots, R_n \in P$ and R transitive, and for each $i \in \{1, \ldots, n\}$, the term s_i is either a variable among the arguments of g, or a term of the form $h_i(z_1, \ldots, z_k)$, where $h_i \in \Sigma_1$ and all the arguments of h_i are variables occurring among the arguments of g.

Example 2. A set \mathcal{K} of axioms containing clauses of the form:

$$\begin{cases} x_1 \le h(y_1) \to f(x_1) \le g(y_1) \\ x_1 \le y_1 \to f(x_1) \le f(y_1) \end{cases}$$

satisfies the conditions above: n = 1, $R_1 = R = \leq$, $s_1 = h(y_1)$, $f, g, h \in \Sigma_1$.

We make the following assumptions (which were also made in [25]) on the theory \mathcal{T}_0 and its theory extension $\mathcal{T}_0 \cup \mathcal{K}$:

A1: \mathcal{T}_0 is convex and has the *P*-interpolation property. **A2:** Satisfiability of ground clauses w.r.t. \mathcal{T}_0 is decidable. **A3:** All clauses in \mathcal{K} are of the form (1). **A4:** $\mathcal{T}_0 \subseteq \mathcal{T}_0 \cup \mathcal{K}$ is a Ψ -local extension.

In [25], we proved that if \mathcal{T}_0 allows ground interpolation, then \mathcal{T} allows ground interpolation, and that the interpolants can be computed in a hierarchical way, using a method for ground interpolation in \mathcal{T}_0 .

We show that under the conditions above, the property of *P*-interpolation can be transferred from the theory \mathcal{T}_0 to the extension $\mathcal{T} = \mathcal{T}_0 \cup \mathcal{K}$ of \mathcal{T}_0 . The function

symbols in the signature of \mathcal{T}_0 are considered to be interpreted, and will always be considered to be shared. For the function symbols in the signature Σ_1 – considered to be "quasi"-interpreted – we use the notion of $\Theta_{\mathcal{K}}$ -sharing introduced in Section 2.

In order to show that \mathcal{T} has the *P*-interpolation property, we need to prove that if A, B are conjunctions of atoms and $A(\overline{c}, \overline{a}_1, a) \wedge B(\overline{c}, \overline{b}_1, b) \models_{\mathcal{T}} aRb$, where $R \in P$, then there exists a term t containing only the constants common to A and B and only function symbols which are $\Theta_{\mathcal{K}}$ -shared by A and B, such that $A(\overline{c}, \overline{a}_1, a) \wedge B(\overline{c}, \overline{b}_1, b) \models_{\mathcal{T}} aRt \wedge tRb.$

Note that

 $A(\overline{c}, \overline{a}_1, a) \land B(\overline{c}, \overline{b}_1, b) \models_{\mathcal{T}} aRb \text{ iff } A(\overline{c}, \overline{a}_1, a) \land B(\overline{c}, \overline{b}_1, b) \land \neg(aRb) \models_{\mathcal{T}} \bot.$

By Assumption A4, $\mathcal{T}_0 \cup \mathcal{K}$ is a Ψ -local extension of \mathcal{T}_0 . We can purify and flatten this conjunction and obtain a conjunction of unit clauses $A_0 \wedge B_0 \wedge \mathsf{Def} \wedge \neg(aRb)$, where Def is a set of definitions of newly introduced constants. Let T be the extension terms in Def. We introduce new constants and definitions also for all extension terms in $\Psi(T)$. This new set of definitions can be written as a conjunction $D_A \wedge D_B$ of its A-part and its B-part. By the Ψ -locality of the extension $\mathcal{T}_0 \subseteq \mathcal{T}_0 \cup \mathcal{K}$ and Theorem 5,

$$A_0 \wedge B_0 \wedge \mathsf{Def} \wedge \neg (aRb) \models_{\mathcal{T}} \bot$$
 iff $\mathcal{K}_0 \wedge A_0 \wedge B_0 \wedge \mathsf{Con}[D_A \wedge D_B]_0 \wedge \neg (aRb) \models_{\mathcal{T}_0} \bot$,

where \mathcal{K}_0 is obtained from $\mathcal{K}[D_A \wedge D_B]$ by replacing the Σ_1 -terms with the corresponding constants contained in the definitions $D_A \wedge D_B$ and

$$\mathsf{Con}[D_A \land D_B]_0 = \bigwedge \Big\{ \bigwedge_{i=1}^n c_i \approx d_i \to c \approx d \mid \frac{f(c_1, \dots, c_n) \approx c \in D_A \cup D_B}{f(d_1, \dots, d_n) \approx d \in D_A \cup D_B} \Big\}.$$

 $\operatorname{Con}[D_A \wedge D_B]_0 = \operatorname{Con}_0^A \wedge \operatorname{Con}_0^B \wedge \operatorname{Con}_{\operatorname{mix}} \operatorname{and} \mathcal{K}_0 = \mathcal{K}_0^A \wedge \mathcal{K}_0^B \wedge \mathcal{K}_{\operatorname{mix}}, \text{ where }$

- $\operatorname{Con}_0^A, \mathcal{K}_0^A$ only contain extension functions and constants which occur in A, $\operatorname{Con}_0^B, \mathcal{K}_0^B$ only contain extension functions and constants which occur in B, $\operatorname{Con}_{\text{mix}}, \mathcal{K}_{\text{mix}}$ contain mixed clauses with constants occurring in A and in B.

Our goal is to separate Con_{mix} and \mathcal{K}_{mix} into an A-part and a B-part, which would allow us to use the *P*-interpolation property of theory \mathcal{T}_0 .

Proposition 7 Assume that \mathcal{T}_0 is convex and P-interpolating and the intermediate terms can be effectively computed. Assume that ground satisfiability w.r.t. \mathcal{T}_0 is decidable.

Let \mathcal{H} be a set of Horn clauses $(\bigwedge_{i=1}^n c_i R_i d_i) \to cR_0 d$ in the signature Π_0^C (with R_0 transitive and $R_i \in P$) which are instances of flattened and purified clauses of type (1) and of congruence axioms. Let $\mathcal{H}_{\mathsf{mix}}$ be the mixed clauses in \mathcal{H} :

 $\mathcal{H}_{\mathsf{mix}} = \{ \bigwedge_{i=1}^{n} c_i R_i d_i \to c R_0 d \in \mathcal{H} \mid c_i, c \text{ constants in } A, d_i, d \text{ constants in } B \} \cup \\ \{ \bigwedge_{i=1}^{n} c_i R_i d_i \to c R_0 d \in \mathcal{H} \mid c_i, c \text{ constants in } B, d_i, d \text{ constants in } A \}$

Let A_0 and B_0 be conjunctions of ground literals in the signature Π_0^C such that $A_0 \wedge B_0 \wedge \mathcal{H} \wedge \neg(aRb) \models_{\mathcal{T}_0} \bot$. Then \mathcal{H} can be separated into an A- and a B-part by replacing the set \mathcal{H}_{mix} of mixed clauses with a separated set of formulae \mathcal{H}_{sep} :

14 Dennis Peuter, Viorica Sofronie-Stokkermans, Sebastian Thunert

(i) There exists a set T of $(\Sigma_0 \cup C)$ -terms containing only constants common to A_0 and B_0 such that $A_0 \wedge B_0 \wedge (\mathcal{H} \setminus \mathcal{H}_{mix}) \wedge \mathcal{H}_{sep} \wedge \neg (aRb) \models_{\mathcal{T}_0} \bot$, where

$$\begin{aligned} \mathcal{H}_{\mathsf{sep}} = & \{ (\bigwedge_{i=1}^{n} c_{i}R_{i}t_{i} \to cRc_{f(t_{1},...,t_{n})}) \land (\bigwedge_{i=1}^{n} t_{i}R_{i}d_{i} \to c_{f(t_{1},...,t_{n})}Rd) \mid \\ & \bigwedge_{i=1}^{n} c_{i}R_{i}d_{i} \to cRd \in \mathcal{H}_{\mathsf{mix}}, d_{i} \approx s_{i}(e_{1},\ldots,e_{n}), d \approx g(e_{1},\ldots,e_{n}) \in D_{B}, \\ & c \approx f(c_{1},\ldots,c_{n}) \in D_{A} \text{ or vice versa } \} = \mathcal{H}_{\mathsf{sep}}^{A} \land \mathcal{H}_{\mathsf{sep}}^{B} \end{aligned}$$

and $c_{f(t_1,\ldots,t_n)}$ are new constants in Σ_c (considered to be common) introduced for the corresponding terms $f(t_1,\ldots,t_n)$, where for $i \in \{1,\ldots,n\}$, t_i separates the atom $c_i R_i d_i$, which is entailed by the already deduced atoms.

(ii) $A_0 \wedge B_0 \wedge (\mathcal{H} \setminus \mathcal{H}_{mix}) \wedge \mathcal{H}_{sep} \wedge \neg(aRb)$ is logically equivalent with respect to \mathcal{T}_0 with the following separated conjunction of ground literals:

$$\overline{A}_0 \wedge \overline{B}_0 \wedge \neg (aRb) = A_0 \wedge B_0 \wedge \neg (aRb) \wedge \bigwedge \{cRd \mid \Gamma \rightarrow cRd \in \mathcal{H} \setminus \mathcal{H}_{\mathsf{mix}} \} \wedge \\ \wedge \{cRc_{f(\overline{t})} \wedge c_{f(\overline{t})}Rd \mid (\Gamma \rightarrow cRc_{f(\overline{t})}) \wedge (\Gamma \rightarrow c_{f(\overline{t})}Rd) \in \mathcal{H}_{\mathsf{sep}} \}.$$

Proof: The proof is similar to that of Proposition 5.7 in [25]. (i) and (ii) are proved simultaneously by induction on the number of clauses in \mathcal{H} . If $\mathcal{H} = \emptyset$, it is already separated into an A and a B part so we are done. Assume that \mathcal{H} contains at least one clause, and that for every \mathcal{H}' with fewer clauses and all conjunctions of literals A'_0, B'_0 with $A'_0 \wedge B'_0 \wedge \mathcal{H}' \wedge \neg aRb \models_{\mathcal{T}_0} \bot$, (i) and (ii) hold.

Let \mathcal{D} be the set of all atoms $c_i R_i d_i$ occurring in premises of clauses in \mathcal{H} . As every model of $A_0 \wedge B_0 \wedge \bigwedge_{(cR'd) \in \mathcal{D}} \neg (cR'd) \wedge \neg aRb$ is also a model for $\mathcal{H} \wedge A_0 \wedge B_0 \wedge \neg aRb$ and since $\mathcal{H} \wedge A_0 \wedge B_0 \wedge \neg aRb \models_{\mathcal{T}_0} \bot$, it follows that $A_0 \wedge B_0 \wedge \bigwedge_{(cR'd) \in \mathcal{D}} \neg (cR'd) \wedge \neg (aRb) \models_{\mathcal{T}_0} \bot$.

Let $(A_0 \wedge B_0)^+$ be the conjunction of all atoms in $A_0 \wedge B_0$, and $(A_0 \wedge B_0)^-$ be the set of all negative literals in $A_0 \wedge B_0$. Then

$$(A_0 \wedge B_0)^+ \models_{\mathcal{T}_0} \bigvee_{(cRd) \in \mathcal{D}} (cRd) \vee \bigvee_{\neg L \in (A_0 \wedge B_0)^-} L \quad \forall aRb.$$

We know that \mathcal{T}_0 is convex with respect to Pred. Moreover, $(A_0 \wedge B_0)^+$ is a conjunction of positive literals. Therefore, either

- (a) $(A_0 \wedge B_0)^+ \models aRb$, or
- (b) $(A_0 \wedge B_0)^+ \models L$ for some $L \in (A_0 \wedge B_0)^-$ (then $A_0 \wedge B_0$ is unsatisfiable and hence entails any atom $c_i R_i d_i$), or
- (c) there exists $(c_1R_1d_1) \in \mathcal{D}$ such that $(A_0 \wedge B_0)^+ \models_{\mathcal{T}_0} c_1R_1d_1$.

Case 1: $(A_0 \wedge B_0)^+ \models aRb$. Then we are done. **Case 2:** $A_0 \wedge B_0$ is unsatisfiable. In this case (i) and (ii) hold for $T = \emptyset$.

Case 3: $A_0 \wedge B_0$ is satisfiable, $(A_0 \wedge B_0)^+ \not\models aRb$, and there exists $(c_1R_1d_1) \in \mathcal{D}$ such that $(A_0 \wedge B_0)^+ \models_{\mathcal{T}_0} c_1R_1d_1$. Then $A_0 \wedge B_0$ is logically equivalent in \mathcal{T}_0 with $A_0 \wedge B_0 \wedge c_iR_id_i$. If it is not the case that by adding $c_iR_id_i$ all premises of some rule in \mathcal{H} become true we repeat the procedure for $\mathcal{D}_1 = \mathcal{D} \setminus (c_1R_1d_1)$: Again in

this case $A_0 \wedge B_0 \wedge \bigwedge_{(cR'd) \in \mathcal{D}_1} \neg (cR'd) \wedge \neg (aRb) \models_{\mathcal{T}_0} \bot$ (if it has a model then

 $A_0 \wedge B_0 \wedge \mathcal{H} \wedge \neg(aRb)$ has one), and as before, using convexity we infer that either $A_0 \wedge B_0 \models aRb$ (which cannot be the case) or $A_0 \wedge B_0$ is unsatisfiable (which cannot be the case) or there exists $c_2R_2d_2 \in \mathcal{D}_1$ with $A_0 \wedge B_0 \models_{\mathcal{T}_0} c_2R_2d_2$. We can repeat the process until all the premises of some clause in \mathcal{H} are proved to be entailed by $A_0 \wedge B_0$. Let $C = \bigwedge_{i=1}^n c_iR_id_i \to cRd$ be such a clause.

Case 3a. Assume that C contains only constants occurring in A or only constants occurring in B. Then $A_0 \wedge B_0 \wedge \mathcal{H}$ is equivalent with respect to \mathcal{T}_0 with $A_0 \wedge B_0 \wedge (\mathcal{H} \setminus C) \wedge cRd$. By the induction hypothesis for $A'_0 \wedge B'_0 = A_0 \wedge B_0 \wedge cRd$ and $\mathcal{H}' = \mathcal{H} \setminus \{C\}$, we know that there exists T' such that $A'_0 \wedge B'_0 \wedge (\mathcal{H}' \setminus \mathcal{H}'_{\text{mix}}) \wedge \mathcal{H}'_{\text{sep}} \wedge \neg (aRb) \models \bot$, and (ii) holds too.

Then, for T = T', $A'_0 \wedge B'_0 \wedge (\mathcal{H}' \setminus \mathcal{H}'_{\text{mix}}) \wedge \mathcal{H}'_{\text{sep}}$ is logically equivalent to $A_0 \wedge B_0 \wedge (\mathcal{H} \setminus \mathcal{H}_{\text{mix}}) \wedge \mathcal{H}_{\text{sep}}$, so (i) holds.

In order to prove (ii), note that, by definition, $\mathcal{H}'_{\mathsf{mix}} = \mathcal{H}_{\mathsf{mix}}$ and $\mathcal{H}'_{\mathsf{sep}} = \mathcal{H}_{\mathsf{sep}}$. By the induction hypothesis, $A'_0 \wedge B'_0 \wedge (\mathcal{H}' \setminus \mathcal{H}'_{\mathsf{mix}}) \cup \mathcal{H}'_{\mathsf{sep}}$ is logically equivalent to a corresponding conjunction $\overline{A}'_0 \wedge \overline{B}'_0$ containing as conjuncts all literals in A'_0 and B'_0 and all conclusions of rules in $\mathcal{H}' \setminus \mathcal{H}'_{\mathsf{mix}}$ and $\mathcal{H}'_{\mathsf{sep}}$. On the other hand, $A'_0 \wedge B'_0$ is logically equivalent to $A_0 \wedge B_0 \wedge (cRd)$, where (cRd) is the conclusion of the rule $C \in \mathcal{H} \setminus \mathcal{H}_{\mathsf{mix}}$. This proves (ii).

Case 3b. Assume now that C is mixed, for instance that c_1, \ldots, c_n, c are constants in A and d_1, \ldots, d_n, d are constants in B. Assume that C is obtained from an instance of a clause of the form $\bigwedge_{i=1}^n x_i R_i s_i(\overline{y}) \to f(x_1, \ldots, x_n) Rg(\overline{y})$. (The case when C corresponds to an instance of a monotonicity axiom is similar.) This means that there exist $c \approx f(c_1, \ldots, c_n) \in D_A$ and $d_i \approx s_i(\overline{e}), d \approx g(\overline{e}) \in D_B$. In particular, f is a A-function symbol and g a B-function symbol; since f and g occur together in an axiom in \mathcal{K} they are considered to be shared. The clause C was chosen such that for each premise $c_i R_i d_i$ of C, $A_0 \wedge B_0 \models_{\mathcal{T}_0} c_i R_i d_i$, and \mathcal{T}_0 is P-interpolating. Thus, there exist terms t_1, \ldots, t_n containing only constants common to A_0 and B_0 such that for all $i \in \{1, \ldots, n\}$

$$A_0 \wedge B_0 \models_{\mathcal{T}_0} c_i R_i t_i \wedge t_i R_i d_i. \tag{2}$$

Let $c_{f(t_1,\ldots,t_n)}$ be a new constant, denoting the term $f(t_1,\ldots,t_n)$, and let

$$C_A = \bigwedge_{i=1}^n c_i R_i t_i \rightarrow c R c_{f(t_1,\dots,t_n)} \quad \text{and} \quad C_B = \bigwedge_{i=1}^n t_i R_i d_i \rightarrow c_{f(t_1,\dots,t_n)} R d.$$

Thus, C_A corresponds to the monotonicity axiom

$$\bigwedge_{i=1}^{n} c_i R_i t_i \rightarrow f(c_1, \dots, c_n) R f(t_1, \dots, t_n)$$

whereas C_B corresponds to the rule

$$\bigwedge_{i=1}^{n} t_i R_i s_i(\overline{e}) \to f(t_1, \dots, t_n) Rg(\overline{e}).$$

As R is transitive, by (2) the following holds: $A_{0} \wedge B_{0} \wedge C_{A} \wedge C_{B} \wedge \neg aRb \models_{\mathcal{T}_{0}} A_{0} \wedge B_{0} \wedge (\bigwedge_{i=1}^{n} c_{i}R_{i}t_{i} \wedge C_{A}) \wedge (\bigwedge_{i=1}^{n} t_{i}R_{i}d_{i} \wedge C_{B})$ $\land \neg aRb \models_{\mathcal{T}_{0}} A_{0} \wedge B_{0} \wedge cRc_{f(t_{1},...,t_{n})} \wedge c_{f(t_{1},...,t_{n})}Rd \wedge \neg aRb$ $\models_{\mathcal{T}_{0}} A_{0} \wedge B_{0} \wedge cRd \wedge \neg aRb$

(where $\models_{\mathcal{T}_0}$ stands for logical equivalence with respect to \mathcal{T}_0).

Hence, $A_0 \wedge B_0 \wedge C_A \wedge C_B \wedge (\mathcal{H} \setminus C) \wedge \neg aRb \models_{\mathcal{T}_0} A_0 \wedge B_0 \wedge cRd \wedge (\mathcal{H} \setminus C) \wedge \neg aRb$. On the other hand, as $A_0 \wedge B_0 \models_{\mathcal{T}_0} \bigwedge_{i=1}^n c_i R_i d_i$, $A_0 \wedge B_0 \wedge \mathcal{H}$ is logically equivalent with $A_0 \wedge B_0 \wedge cRd \wedge (\mathcal{H} \setminus C)$, so $A_0 \wedge B_0 \wedge C_A \wedge C_B \wedge (\mathcal{H} \setminus C) \wedge \neg aRb \models_{\mathcal{T}_0} \bot$. By the induction hypothesis for $A_0 \wedge B_0 \wedge cRc_{f(t_1,...,t_n)} \wedge c_{f(t_1,...,t_n)}Rd$ and $\mathcal{H}' = \mathcal{H} \setminus C$ we know that there exists a set T' of terms such that

$$A_0 \wedge B_0 \wedge cRc_{f(t_1,\ldots,t_n)} \wedge c_{f(t_1,\ldots,t_n)} Rd \wedge (\mathcal{H}' \setminus \mathcal{H}'_{\mathsf{mix}}) \wedge \mathcal{H}'_{\mathsf{sep}} \wedge \neg aRb \models \bot,$$

and also (ii) holds. Then (i) holds for $T = T' \cup \{t_1, \ldots, t_n\}$. (ii) can be proved similarly using the induction hypothesis.

Theorem 8 Assume that \mathcal{T}_0 is convex and *P*-interpolating with respect to $P \subseteq$ **Pred**, and that the ground satisfiability w.r.t. \mathcal{T}_0 is decidable and the interpolating terms can be effectively computed. Assume that $\mathcal{T} = \mathcal{T}_0 \cup \mathcal{K}$ is a Ψ -local extension of \mathcal{T}_0 with a set of clauses \mathcal{K} which only contains combinations of clauses of type (1). Then \mathcal{T} is also *P*-interpolating and the interpolating terms can be effectively computed.

Proof: We prove that if A, B are conjunctions of literals and $A(\overline{c}, \overline{a}_1, a) \land B(\overline{c}, \overline{b}_1, b) \models_{\mathcal{T}} aRb$ where $R \in P$, then there exists a term t containing only the constants common to A and B and only function symbols which are shared by A and B, such that $A(\overline{c}, \overline{a}_1, a) \land B(\overline{c}, \overline{b}_1, b) \models_{\mathcal{T}} aRt \land tRb$. We can restrict w.l.o.g. to a purified and flattened conjunction of unit clauses $A_0 \land B_0 \land \mathsf{Def} \land \neg(aRb)$. With the notation used on page 13, by Theorem 5 we have:

 $A_0 \wedge B_0 \wedge \mathsf{Def} \wedge \neg (aRb) \models_{\mathcal{T}} \bot$ iff $\mathcal{K}_0 \wedge A_0 \wedge B_0 \wedge \mathsf{Con}[D_A \wedge D_B]_0 \wedge \neg (aRb) \models_{\mathcal{T}_0} \bot$. By Proposition 7 (ii), there exists a set T of $(\Sigma_0 \cup C)$ -terms containing only constants common to A_0 and B_0 such that $\mathcal{H} = \mathcal{K}_0 \wedge \mathsf{Con}[D_A \wedge D_B]_0$ can be separated as described in Proposition 7, $A_0 \wedge B_0 \wedge (\mathcal{H} \setminus \mathcal{H}_{\mathsf{mix}}) \wedge \mathcal{H}_{\mathsf{sep}} \wedge \neg aRb$ is logically equivalent w.r.t. \mathcal{T}_0 with a separated conjunction of ground literals $\overline{A}_0 \wedge \overline{B}_0 \wedge \neg aRb$, which is therefore unsatisfiable, so $\overline{A}_0 \wedge \overline{B}_0 \models aRb$. From the *P*-interpolation property in \mathcal{T}_0 , there exists a term containing the shared constants such that $\overline{A}_0 \wedge \overline{B}_0 \models_{\mathcal{T}_0} aRt \wedge tRb$. If we now replace all constants $c_{f(t_1,\ldots,t_n)}$ introduced in the purification process or in the separation process with the terms they denote, we obtain $A \wedge B \models_{\mathcal{T}} aRt \wedge tRb$. Since all intermediate terms t_i contain only shared symbols and the function symbol f is shared by A and B, all terms $f(t_1,\ldots,t_n)$, hence also t contain only symbols shared by A and B. *P*-interpolation in local theory extensions and applications to $\mathcal{EL}, \mathcal{EL}^+$ 17

We obtain the following procedure for *P*-interpolation if $A \wedge B \models_{\mathcal{T}} aRb$:

Step 1: Preprocess Using locality, flattening and purification we obtain a set $\mathcal{H} \wedge A_0 \wedge B_0$ of formulae in the base theory, where \mathcal{H} is as in Proposition 7.

Step 2: $\Delta := \mathsf{T}$. **Repeat as long as** $A_0 \wedge B_0 \wedge \Delta \not\models aRb$:

Let $C \in \mathcal{H}$ whose premise is entailed by $A_0 \wedge B_0 \wedge \Delta$.

If C is not mixed, move C to \mathcal{H}_{sep} and add its conclusion to Δ .

If C is mixed, compute terms t_i which separate the premises in C, and separate the clause into an instance C_1 of monotonicity and an instance C_2 of a clause in \mathcal{K} as in the proof of Proposition 7. Remove C from \mathcal{H} , and add C_1, C_2 to \mathcal{H}_{sep} and their conclusions to Δ .

Step 3: Compute separating term. Compute a separating term for $A_0 \wedge B_0 \wedge \Delta \models aRb$ in \mathcal{T}_0 , and construct an interpolant for the extension as explained in the proof of Theorem 8.

5 Example: Semilattices with monotone operators

We will now analyze \leq -interpolation properties for theories of semilattices with monotone operators.

5.1 The theory SLat of semilattices

We define the theory SLat of semilattices and show that conditions A1 and A2 are satisfied for SLat.

A semilattice (S, \sqcap) is set S with a binary operation \sqcap which is associative, commutative and idempotent. One can equivalently regard semilattices as partially ordered sets (S, \leq) , in which infima $a_1 \sqcap \ldots \sqcap a_n$ of finite non-empty subsets $\{a_1, \ldots, a_n\} \subseteq S$ exist; then $a \leq b$ iff $a \sqcap b = a$.

The theory SLat of semilattices can be axiomatized by equations (associativity, commutativity and idempotence of \sqcap), therefore it clearly is \approx -convex: Convexity w.r.t. \leq follows from the fact that $x \leq y$ iff $(x \sqcap y) \approx x$.

Lemma 1. Ground satisfiability w.r.t. SLat is decidable.

Proof: This is a consequence of the fact that the theory of semilattices SLat is a local extension of the theory of pure equality – this follows from a result on the locality of lattices by Skolem [21], or by results in [10], since every partial semilattice weakly embeds into a total one. ¹

The theory SLat is \leq -interpolating, therefore also \approx -interpolating (cf. [18]; we present the proof since it indicates how the intermediate terms can be computed):

¹ There are also other justifications for the decidability of ground satisfiability w.r.t. SLat, leading to different types of decision procedures: $SLat = ISP(S_2)$, where S_2 is the 2-element semilattice, i.e. every semilattice is isomorphic to a sublattice of a power of S_2 – or, alternatively, that every semilattice is isomorphic to a semilattice of sets. Thus, checking satisfiability w.r.t. SLat can be reduced to checking satisfiability w.r.t. S_2 and ultimately to propositional reasoning.

Lemma 9 The theory SLat of semilattices is \leq -interpolating.

Proof: This is a constructive proof based on the fact that $\mathsf{SLat} = ISP(S_2)$, where S_2 is the 2-element semilattice, i.e. every semilattice is isomorphic to a sublattice of a power of S_2 – or, alternatively, that every semilattice is isomorphic to a semilattice of sets. We prove that the theory of semilattices is \leq -interpolating, i.e. that if A and B are two conjunctions of literals and $A \wedge B \models_{\mathsf{SLat}} a \leq b$, where a is a constant occurring in A and b a constant occurring in B, then there exists a term containing only common constants in A and B such that $A \models_{\mathsf{SLat}} a \leq t$ and $A \wedge B \models_{\mathsf{SLat}} t \leq b$. We can assume without loss of generality that A and B consist only of atoms: Indeed, assume that $A \wedge B = A_1 \wedge \cdots \wedge A_n \wedge \neg A'_1 \wedge \cdots \wedge \neg A'_m$, where $A_1, \ldots, A_n, A'_1, \ldots, A'_m$ are atoms. Then the following are equivalent:

- $-A \wedge B \models_{\mathsf{SLat}} a \leq b$
- $-\models_{\mathsf{SLat}} A \land B \to a \leq b$
- $\models_{\mathsf{SLat}} \neg A_1 \lor \cdots \lor \neg A_n \lor A'_1 \lor \cdots \lor A'_m \lor a \le b \\ \models_{\mathsf{SLat}} (A_1 \land \cdots \land A_n) \to A'_1 \lor \cdots \lor A'_m \lor a \le b \\ A_1 \land \cdots \land A_n \models_{\mathsf{SLat}} A'_1 \lor \cdots \lor A'_m \lor a \le b$

Since the theory of semilattices is convex w.r.t. \leq and \approx , it follows that if $A \wedge B \models_{\mathsf{SLat}} a \leq b$ then either

(a) $A_1 \wedge \cdots \wedge A_n \models_{\mathsf{SLat}} A'_j$ for some $j \in \{1, \dots, m\}$ or (b) $A_1 \wedge \cdots \wedge A_n \models_{\mathsf{SLat}} a \leq b$.

It is easy to see that in case (a), $A \wedge B \models \perp$. Then the conclusion follows immediately. We therefore consider the case when A and B consist only of atoms.

As $SLat = ISP(S_2)$, in SLat the same Horn sentences are true as in the 2element semilattice S_2 . Thus, $A \wedge B \models_{\mathsf{SLat}} a \leq b$ iff $A \wedge B \models_{S_2} a \leq b$, so we can reduce such a test to entailment in propositional logic.

It follows that $A \wedge B \models_{\mathsf{SLat}} a \leq b$ if and only if the following conjunction of literals in propositional logic is unsatisfiable:

$$N_A: \begin{cases} \begin{array}{ccc} P_{e_1 \cap e_2} \leftrightarrow P_{e_1} \wedge P_{e_2} \\ P_{e_1} \leftrightarrow P_{e_2} & e_1 \approx e_2 \in A \\ P_{e_1} \rightarrow P_{e_2} & e_1 \leq e_2 \in A \\ \text{for all } e_1, e_2 \text{ subterms in } A \\ P_a \end{array} \qquad N_B: \begin{cases} \begin{array}{ccc} P_{g_1 \cap g_2} \leftrightarrow P_{g_1} \wedge P_{g_2} \\ P_{g_1} \leftrightarrow P_{g_2} & g_1 \approx g_2 \in B \\ P_{g_1} \rightarrow P_{g_2} & g_1 \leq g_2 \in B \\ \text{for all } g_1, g_2 \text{ subterms in } B \\ \neg P_b \end{cases} \end{cases}$$

i.e. $A \wedge B \models_{\mathsf{SLat}} a \leq b$ if and only if $(N_A \wedge P_a) \wedge (N_B \wedge \neg P_b) \models \bot$, where N_A and N_B are sets of Horn clauses in which each clause contains a positive literal. We show that if $A \wedge B \models_{\mathsf{SLat}} a \leq b$ holds, then for the term

$$t := \bigcap \{ e \mid A \models_{\mathsf{SLat}} a \le e, e \text{ common subterm of } A \text{ and } B \}$$

the following hold:

(i)
$$A \models_{\mathsf{SLat}} a \leq t$$
, and

(ii) $A \wedge B \models_{\mathsf{SLat}} t \leq b.$

This means that for the theory of semilattices we have a property stronger than \leq -interpolability.

Every $e \in T = \{e \mid A \models_{\mathsf{SLat}} a \leq e, e \text{ common subterm of } A \text{ and } B\}$ corresponds to the positive unit clause P_e (where P_e is a propositional variable common to N_A and N_B) which can be derived from N_A using ordered resolution (with the ordering described above).

It is clearly the case that $A \models_{\mathsf{SLat}} a \leq t$, because $N_A \wedge P_a \wedge \neg P_t \wedge (P_t \leftrightarrow \bigwedge_{e \in T} P_e)$ is unsatisfiable. Thus, (i) holds.

For proving (ii), observe that by saturating $N_A \wedge P_a$ under ordered resolution we obtain the following kinds of clauses which can possibly lead to \perp after inferences with $N_B \wedge \neg P_b$ (and thus to the consequence $a \leq b$ together with B):

- (a) P_{e_k} positive unit clauses s.t. e_k contains symbols common to A and B, for $k \in \{1, \ldots, l\}.$
- (b) $\bigwedge_{j=1}^{n_i} P_{c_{ij}} \to P_{d_i}$, where c_{ij} and d_i are common symbols, such that for all i, j and k we have $c_{ij} \neq e_k$ and $d_i \neq e_k$, for $i \in \{1, \ldots, p\}$.

Other types of clauses may appear too, but they can not be used to obtain $a \leq b$:

To see that clauses where some $c_{ij} = e_k$ are not necessary to derive the consequence $a \leq b$, note that if P_{e_k} is a positive unit literal and we have the clause $(P_{e_k} \wedge \bigwedge P_{c_{ij}}) \to P_{d_i}$, then by resolution we get as an inference $\bigwedge P_{c_{ij}} \to P_{d_i}$. It is easy to see that $(P_{e_k} \land \bigwedge P_{c_{ij}}) \to P_{d_i}$ is redundant in the presence of $\bigwedge P_{c_{ij}} \to P_{d_i}$. In the same way, clauses of the form $\bigwedge P_{c_{ij}} \to P_{e_k}$ (i.e. clauses of type (b) where $d_i = e_k$) are redundant in the presence of P_{e_1}, \ldots, P_{e_l} . For the proof of (ii) one needs to consider separately the case in which none of the P_{d_i} is needed to derive \perp together with N_B (and thus the consequence $a \leq b$) and the case when some P_{d_i} are needed.

Case 1: None of the P_{d_i} is needed to derive \perp together with N_B (and thus the consequence $a \leq b$). We know that $N_A \models P_a \rightarrow \bigwedge_{k=1}^l P_{e_k}$. From this it follows that $A \models a \leq \prod_{k=1}^{l} e_k$.

For $A \wedge B \models a \leq b$ to be true, $\bigwedge_{k=1}^{l} P_{e_k} \wedge N_B \wedge \neg P_b$ must be unsatisfiable, so there has to be a subset $S \subseteq \{1, ..., l\}$ such that $\bigwedge_{k \in S} P_{e_k} \wedge N_B \wedge \neg P_b$. This means that $B \models \prod_{s \in S} e_s \leq b$. But then, since $\prod_{k=1}^{l} e_k \leq \prod_{s \in S} e_s$, it follows that $B \models \prod_{k=1}^{l} e_k \leq b$, and therefore also $A \land B \models \prod_{k=1}^{l} e_k \leq b$.

Case 2: Some P_{d_i} are needed to derive \perp from $N_B \wedge \neg P_b$. Again, we know that

 $N_A \models P_a \rightarrow \bigwedge_{k=1}^{l} P_{e_k}$ (hence $A \models a \leq \prod_{k=1}^{l} e_k$). For $A \wedge B \models a \leq b$ to be true, i.e. $(N_A \wedge P_a) \wedge (N_B \wedge \neg P_b)$ to be unsatisfiable, there have to be subsets $S_1 \subseteq \{1, \ldots, l\}$ and $S_2 \subseteq \{1, \ldots, p\}$ such that $N_B \wedge I$ there have to be bubbles $\Sigma_1 \subseteq \mathbb{C}^+$, $\sum_{k \in S_1} P_{e_k} \wedge \bigwedge_{i \in S_2} ((\bigwedge_j P_{c_{ij}}) \to P_{d_i}) \wedge \neg P_b$ is unsatisfiable.

Let
$$N_{AB} := N_B \land \bigwedge_{k \in S_1} P_{e_k} \land \bigwedge_{i \in S_2} ((\bigwedge_j P_{c_{ij}}) \to P_{d_i})$$

We know that $N_B \wedge \bigwedge_{k \in S_1} P_{e_k} \wedge \neg P_b$ is satisfiable. Assume that there is no c_{ij} such that $N_B \wedge \bigwedge_{k \in S_1} P_{e_k} \wedge \neg P_b \models P_{c_{ij}}$. Then for every c_{ij} , $N_B \wedge \bigwedge_{k \in S_1} P_{e_k} \wedge \neg P_b \wedge \neg P_{c_{ij}}$ is satisfiable. Since all clauses in $N_b \wedge \bigwedge_{k \in S_1} P_{e_k} \wedge \neg P_b$ are Horn clauses, it follows that $N_B \wedge \bigwedge_{k \in S_1} P_{e_k} \wedge \neg P_b \wedge \bigwedge_{i,j} \neg P_{c_{ij}}$ is satisfiable. Every model of $N_B \wedge \bigwedge_{k \in S_1} P_{e_k} \wedge \neg P_b \wedge \bigwedge_{i,j} \neg P_{c_{ij}}$ is a model of $N_{AB} \wedge \neg P_b$. It would therefore follow that $N_{AB} \wedge \neg P_b$ is satisfiable, which is a contradiction.

Thus, there exists at least one c_{ij} such that $N_B \wedge \bigwedge_{k \in S_1} P_{e_k} \wedge \neg P_b \models P_{c_{ij}}$.

We can add $P_{c_{ii}}$ to this set of clauses and repeat the reasoning for the set of clauses obtained this way as long as we still have one clause of the form $((\bigwedge_i P_{c_{ij}}) \to P_{d_i})$ in N_{AB} such that there exists at least one c_{ij} such that $P_{c_{ij}}$ was not added to N_{AB} .

Then there has to be a sequence $(d_{i_1j})_{j \in J_1}, (d_{i_2j})_{j \in J_2}, ..., (d_{i_nj})_{j \in J_n}$ such that:

- $-P_{d_{i_1j_j}}$ can be derived from $N_{AB} \wedge \bigwedge P_{e_k}$, for all $j \in J_1$,
- $\begin{array}{l} P_{d_{i_{2}j}} \text{ can be derived from } N_{AB} \wedge \bigwedge P_{e_{k}} \wedge \bigwedge_{k \in J_{1}} P_{d_{i_{1}k}}, \text{ for all } j \in J_{2}, \\ P_{d_{i_{3}j}} \text{ can be derived from } N_{AB} \wedge \bigwedge P_{e_{k}} \wedge \bigwedge_{k \in J_{1}} P_{d_{i_{1}k}} \wedge \bigwedge_{k \in J_{2}} P_{d_{i_{2}k}}, \text{ for all } \end{array}$ $j \in J_3$,
- $P_{d_{i_nj}} \text{ can be derived from } N_{AB} \wedge \bigwedge_k P_{e_k} \wedge \bigwedge_{k \in J_1} P_{d_{i_1k}} \wedge \cdots \bigwedge_{k \in J_{n-1}} P_{d_{i_{n-1}k}},$ for all $j \in J_n$,
- $-P_b$ can be derived from $N_{AB} \wedge \bigwedge P_{e_k} \wedge \bigwedge_{k \in J_1} P_{d_{i_1k}} \wedge \ldots \wedge \bigwedge_{k \in J_n} P_{d_{i_nk}}$.

But then $A \wedge B \models \bigcap e_k \leq d_{i_1 l}$, for all $l \in J_1$, hence $A \wedge B \models \bigcap e_k \leq \bigcap_{l \in J_1} d_{i_1 l}$, hence $A \wedge B \models (\prod e_k \sqcap \prod_{l \in J_1} d_{i_1 l}) \approx \prod e_k$.

Therefore, as $A \wedge B \models (\prod e_k \sqcap \prod_{l \in J_1} d_{i_1 l}) \leq d_{i_2 j}$, for all $j \in J_2$, we conclude that $A \wedge B \models \prod e_k \leq \prod_j d_{i_2 j}.$

Similarly it can be proved that $A \wedge B \models \prod e_k \leq \prod_{j \in J_n} d_{i_n j}$, and finally that $A \wedge B \models \Box e_k \leq b.$ П

We illustrate the computation of intermediate terms on an example.

Example 3. Let $A = \{a_1 \leq c_1, c_2 \leq a_2, a_2 \leq c_3\}$ and $B = \{c_1 \leq b_1, b_1 \leq c_2\}$ $c_2, c_3 \leq b_2$. It is easy to see that $A \wedge B \models a_1 \leq b_2$. We can find an intermediate term by using the methods described in the proof of Lemma 9: We saturate the set of clauses

$$N_A \wedge P_{a_1} = (P_{a_1} \to P_{c_1}) \wedge (P_{c_2} \to P_{a_2}) \wedge (P_{a_2} \to P_{c_3}) \wedge P_a$$

under ordered resolution, in which the propositional variables P_{a_1}, P_{a_2} are larger than $P_{c_1}, P_{c_2}, P_{c_3}$. This yields the clauses P_{c_1} and $P_{c_2} \to P_{c_3}$ containing shared propositional variables. $(N_A \wedge P_{a_1}) \wedge (N_B \wedge \neg P_{b_2})$ is unsatisfiable iff $N_B \wedge \neg P_{b_2} \wedge$ $P_{c_1} \wedge (P_{c_2} \rightarrow P_{c_3})$ is unsatisfiable. Indeed $t = c_1$ is an intermediate term, as $A \models a_1 \leq c_1$ and $A \wedge B \models c_1 \leq b_2$. Note that $N_B \wedge \neg P_{b_2} \wedge P_{c_1}$ is satisfiable, so $B \not\models c_1 \leq b_2$. Moreover, we only need $P_{c_2} \rightarrow P_{c_3}$ in addition to $N_B \cup \neg P_{b_2}$ to derive \perp , thus $A \wedge B \models c_1 \leq b_2$ and the clause $P_{c_2} \rightarrow P_{c_3}$ obtained from N_A is really needed for this. \square

5.2Semilattices with operators

We define the theory of semilattices with monotone operators in a set Σ possibly satisfying additional properties and show that conditions A3 and A4 are satisfied for this theory.

Let Σ be a set of unary² function symbols. We consider the extension $SLat_{\Sigma} = SLat \cup Mon(\Sigma)$ of SLat with new function symbols in Σ satisfying the monotonicity axioms $Mon_{\Sigma} = \bigcup_{f \in \Sigma} Mon(f)$, where:

 $\mathsf{Mon}(f) \qquad \forall x, y(x \leq y \to f(x) \leq f(y))$

and also extensions $\mathsf{SLat} \cup \mathsf{Mon}(\Sigma) \cup \mathcal{K}$, where \mathcal{K} is a set of axioms of the form:

$$\forall x \qquad f(x) \le g(x) \tag{3}$$

21

$$\forall x, y \quad y \le g(x) \to f(y) \le h(x) \tag{4}$$

where $f, g, h \in \Sigma$, not necessarily all different.

Clearly, condition A3 is satisfied for the theory of semilattices with operators defined above. We show that condition A4 holds as well.

Lemma 10 The following extensions satisfy a locality property:

- (i) The theory of semilattices SLat is local.
- (*ii*) $\mathsf{SLat} \cup \mathsf{Mon}_{\Sigma}$ is a local extension of SLat .
- (iii) $\mathsf{SLat} \cup \mathsf{Mon}_{\Sigma} \cup \mathcal{K}$ is a Ψ -local extension of SLat , where Ψ is the closure operator on ground terms defined as follows:

$$\Psi(G) = \bigcup_{i \ge 0} \Psi^{i}(G), \text{ with } \Psi^{0}(G) = \mathsf{est}(G) \text{ (the set of ground terms in } G \text{ starting with extension functions), and}$$
$$\overset{i+1}{G} = \{h(c) \mid \forall x(a(x) \le h(x)) \in \mathcal{K} \text{ and } a(c) \in \Psi^{i}(G)\} \mid$$

$$\begin{split} \Psi^{i+1}(G) &= \{h(c) \mid \forall x(g(x) \le h(x)) \in \mathcal{K} \text{ and } g(c) \in \Psi^i(G)\} \cup \\ \{g(c) \mid \forall x(g(x) \le h(x)) \in \mathcal{K} \text{ and } h(c) \in \Psi^i(G)\} \cup \\ \{h(c) \mid \forall x, y(y \le g(x) \to f(y) \le h(x)) \in \mathcal{K} \text{ and } g(c) \in \Psi^i(G)\} \cup \\ \{g(c) \mid \forall x, y(y \le g(x) \to f(y) \le h(x)) \in \mathcal{K} \text{ and } h(c) \in \Psi^i(G)\}. \end{split}$$

Proof: (i) follows from a result on the locality of lattices by Skolem [21], or by results in [10], since every partial semilattice weakly embeds into a total one. (ii) follows from results in [28,29]. (iii) Since the axioms in \mathcal{K} are not always linear, we use the locality criterion for non-linear sets of clauses mentioned in Theorem 6, and the fact that every semilattice $P = (S, \sqcap, \{f\}_{f \in \Sigma})$ with partially defined monotone operators satisfying the axioms \mathcal{K} , and with the property that if a variable occurs in two terms g(x), h(x) in a clause in \mathcal{K} , then for every $s \in S$, g(s) is defined iff h(s) is defined, weakly embeds into a semilattice with totally defined operators satisfying \mathcal{K} , which was proved in Lemma 4.5 from [27]. \Box

Given two sets of conjunctions of ground literals A and B over the signature of semilattices with operators, we consider the lattice operation \sqcap to be interpreted and the function symbols in Σ to be uninterpreted. Let Σ_A be the function symbols in Σ occurring in A and Σ_B those occurring in B. We consider the following variants for "shared uninterpreted function symbols":

² We assume that the function symbols are unary to simplify the presentation, and because in the applications to description logics we need only unary function symbols. All the results can be extended to function symbols of higher arity.

- 22 Dennis Peuter, Viorica Sofronie-Stokkermans, Sebastian Thunert
 - Intersection-sharing: The shared function symbols of A and B are the function symbols in $\Sigma_A \cap \Sigma_B$.
 - $\Theta_{\mathcal{K}}$ -sharing (as defined in Example 1): $\Theta_{\mathcal{K}}(\Sigma_A) = \bigcup_{f \in \Sigma_A} \{g \in \Sigma \mid f \sim^*_{\mathcal{K}} g\},$ where $\sim^*_{\mathcal{K}}$ is the equivalence relation induced by $\sim_{\mathcal{K}}$ (with $f \sim_{\mathcal{K}} g$ iff f, goccur in the same clause in \mathcal{K}); $\Theta_{\mathcal{K}}(\Sigma_B)$ is defined analogously. The $\Theta_{\mathcal{K}}$ shared function symbols are the function symbols in $\Theta_{\mathcal{K}}(\Sigma_A) \cap \Theta_{\mathcal{K}}(\Sigma_B)$.

Theorem 11 For every set \mathcal{K} containing clauses of the form (3) and (4) above, the theory $\mathsf{SLat} \cup \mathsf{Mon}_{\Sigma} \cup \mathcal{K}$ of semilattices with monotone operators satisfying axioms \mathcal{K} is \leq -interpolating with the notion of $\Theta_{\mathcal{K}}$ -sharing for uninterpreted function symbols.

Proof. The clauses of type (3) and (4) satisfy the conditions in the statement of Proposition 7 and Theorem 8. The result is therefore a consequence of the fact that the theory extension $SLat \subseteq SLat \cup Mon_{\Sigma} \cup \mathcal{K}$ satisfies conditions A1, A2, A3, A4 and of Proposition 7 and Theorem 8.

We illustrate the way Theorem 5, Proposition 7 and Theorem 8 and the algorithm in Section 4 can be used for computing intermediate terms below:

Example 4. Consider the extension $\mathsf{SLO} = \mathsf{SLat} \cup \mathsf{Mon}_f \cup \mathsf{Mon}_g \cup \mathcal{K}$ of SLat with two monotone functions f, g satisfying: $\mathcal{K} = \{y \leq g(x) \rightarrow f(y) \leq g(x)\}$. Consider the following conjunctions of atoms: $A := d \leq g(a) \land a \leq c \land g(c) \leq a$ and $B := b \leq d \land b \leq f(b)$. It can be checked that $A \land B \models b \leq a$.

To obtain a separating term we proceed as follows: By the definition of SLO, $A \wedge B \models_{SLO} b \leq a$ iff SLat $\wedge \operatorname{Mon}_f \wedge \operatorname{Mon}_g \wedge \mathcal{K} \wedge A \wedge B \wedge \neg(b \leq a) \models \bot$. By Theorem 5, this is the case iff SLat $\wedge (\operatorname{Mon}_f \wedge \operatorname{Mon}_g \wedge \mathcal{K})[\Psi(G)] \wedge G \models \bot$, where $G = A \wedge B \wedge \neg(b \leq a)$, est $(G) = \{g(a), g(c), f(b)\}$ and $\Psi(G) = \{g(a), g(c), f(b)\}$.

- $\operatorname{\mathsf{Mon}}_{f}[\Psi(G)] = \{b \le b \to f(b) \le f(b)\} \text{ (redundant)}.$
- $\operatorname{Mon}_{g}[\Psi(G)] = \{ d_{1} \leq d_{2} \rightarrow g(d_{1}) \leq g(d_{2}) \mid d_{1}, d_{2} \in \{a, c\} \}.$
- $-\mathcal{K}[\Psi(G)] = \{b \le g(a) \to f(b) \le g(a), b \le g(c) \to f(b) \le g(c)\}.$

Step 1: We purify $(\mathsf{Mon}_f \wedge \mathsf{Mon}_g \wedge \mathcal{K})[\Psi(G)] \wedge G$, by introducing constants a_1 for $g(a), c_1$ for g(c) and b_1 for f(b) and obtain the formula $\mathsf{Def} \wedge A_0 \wedge B_0 \wedge \mathsf{Mon}_0 \wedge \mathcal{K}_0$:

Def	$A_0 \wedge B_0$		$Mon_0\wedge\mathcal{K}_0$
$\overline{D_A:a_1\approx g(a)\wedge c_1\approx g(c)}$	$A_0: d \le a_1 \land a \le c \land c_1 \le a$	Mon_A	$a \lhd c \to a_1 \lhd c_1$
$D_B: b_1 \approx f(b)$	$B_0: b \le d \land b \le b_1$	\mathcal{K}_{mix}	$b \le a_1 \to b_1 \le a_1$
	$\lhd \in \{\leq,\geq\}$		$b \le c_1 \to b_1 \le c_1$

The instances of the congruence axioms $\operatorname{Con}[D_A \wedge D_B]$ are redundant in the presence of the corresponding instances of the monotonicity axioms for f and g and can therefore be ignored.

Step 2. $\Delta := \top$. Find clauses in $\mathsf{Mon}_0 \land \mathcal{K}_0$ with premises entailed by $A_0 \land B_0 \land \Delta$.

 $C = a \leq c \rightarrow a_1 \leq c_1: C \text{ is not mixed. Since } A_0 \land B_0 \models_{\mathsf{SLat}} a \leq c, A_0 \land B_0 \land (a \leq c \rightarrow a_1 \leq c_1) \text{ is equivalent to } A_0 \land B_0 \land a_1 \leq c_1. \text{ Let } \Delta := \{a_1 \leq c_1\}.$

23

- $$\begin{split} C &= b \leq a_1 \rightarrow b_1 \leq a_1; \ C \ \text{is mixed. Since } A_0 \wedge B_0 \wedge a_1 \leq c_1 \models b \leq a_1 \ \text{we} \\ \text{find a separating term. For this we use the method described in the proof of } \\ \text{Lemma 9. We consider the encoding } N_B \wedge P_b := (P_b \rightarrow P_d) \wedge (P_b \rightarrow P_{b_1}) \wedge P_b. \\ \text{Using ordered resolution with an ordering in which } P_b, P_{b_1} \succ P_d \ \text{we derive} \\ \text{the unit clauses } P_d \ \text{and } P_{b_1}. \ \text{Since } d \ \text{is the only shared constant}, \ t = d \ \text{is the} \\ \text{separating term. Thus, } A_0 \wedge B_0 \wedge a_1 \leq c_1 \models b \leq d \ \wedge d \leq a_1. \ \text{We now can} \\ \text{separate the instance } b \leq a_1 \rightarrow b_1 \leq a_1 \ \text{of the clause in } \mathcal{K} \ \text{by introducing} \\ \text{a new shared constant } d_1 \ \text{as a name for } f(d) \ \text{and replacing the clause, as} \\ \text{described in the algorithm at the end of Section 4, with the conjunction of} \\ \text{(i) } b \leq d \rightarrow b_1 \leq d_1 \qquad (\text{corresponding to } b \leq d \rightarrow f(b) \leq f(b)) \end{split}$$
 - (ii) $d \le a_1 \to d_1 \le a_1$ (corresponding to $d \le g(a) \to f(d) \le g(a)$)
 - ((i) is an instance of a monotonicity axiom, (ii) is another instance of \mathcal{K}), and $A_0 \wedge B_0 \wedge a_1 \leq c_1 \wedge (b \leq d \rightarrow b_1 \leq d_1) \wedge (d \leq a_1 \rightarrow d_1 \leq a_1)$ is equivalent to $A_0 \wedge B_0 \wedge a_1 \leq c_1 \wedge b_1 \leq d_1 \wedge d_1 \leq a_1$. Let $\Delta := \Delta \wedge b_1 \leq d_1 \wedge d_1 \leq a_1$.

Step 3: The last conjunction entails $b \leq a$. To compute a separating term, we again use Lemma 9. We consider the encoding $N'_B \wedge P_b := (P_b \to P_d) \wedge (P_b \to P_{b_1}) \wedge (P_{b_1} \to P_{d_1}) \wedge P_b$ of the *B*-part of the conjunction, $B_0 \wedge b_1 \leq d_1$. Using ordered resolution with an ordering in which $P_b, P_{b_1} \succ P_d, P_{d_1}$ we derive the unit clauses P_d, P_{b_1} and P_{d_1} . Since d, d_1 are the shared constants, $t = d \sqcap d_1$ is the separating term w.r.t. SLat. Therefore, $d \sqcap f(d)$ is a separating term w.r.t. SLO. (it can in fact be seen that already d is a separating term).

Theorem 12 If \mathcal{K} contains axioms of type (4) then the theory of semilattices with operators is not \leq -interpolating when sharing is regarded as intersectionsharing.

Proof: Indeed, assume that for every \mathcal{K} containing axioms of type (4), $\mathsf{SLat}_{\Sigma}(\mathcal{K})$ is \leq -interpolating w.r.t. intersection-sharing. Then it would also be \approx -interpolating w.r.t. intersection-sharing. This cannot be the case, as can be seen from Example 5 which is presented in what follows.

Example 5. Consider the theory $\mathsf{SLat}_{\Sigma}(\mathcal{K})$ of semilattices with monotone operators f, g satisfying the axioms $\mathcal{K} = \{x \leq g(y) \to f(x) \leq g(y)\}$, and let C be a set of constants containing constants a, b, d, e. We show that this theory does not have the Σ_S -Beth-definability property, where $\Sigma_S = \{g, e\}$.

Consider the conjunction of literals $A = (a \leq f(e)) \land (e \leq g(b)) \land (g(b) \leq a)$. One can prove that a is implicitly definable w.r.t. $\{g, e\}$ by proving, using the hierarchical reduction for local theory extensions in Theorem 5, that:

 $\begin{array}{l} (a \leq f(e)) \wedge (e \leq g(b)) \wedge (g(b) \leq a) \wedge (a' \leq f'(e)) \wedge (e \leq g(b')) \wedge (g(b') \leq a') \models_{\mathsf{Slat}_{\Sigma}(\mathcal{K} \cup \mathcal{K}')} a \approx a'. \\ \text{We show that } a \text{ is not explicitly definable w.r.t. } \{g, e\}. \text{ If there exists a term } t \text{ containing only } g \text{ and } e \text{ such that } (a \leq f(e)) \wedge (e \leq g(b)) \wedge (g(b) \leq a) \models_{\mathsf{Slat}_{\Sigma}(\mathcal{K})} a \approx t, \\ \text{then the interpretations of } a \text{ and } t \text{ are equal in every model of } \mathsf{SLat}_{\Sigma}(\mathcal{K}) \text{ which is a model of } A. \\ \text{We show that this is not the case.} \end{array}$

Let $S = (\{a_S, e_S, b_S, d_S\}, \sqcap, f_S, g_S)$ be the semilattice where:

 $-d_S \leq e_S \leq a_S, d_S \leq b_S$ and $a_S \sqcap b_S = e_S \sqcap b_S = d_S$,

24 Dennis Peuter, Viorica Sofronie-Stokkermans, Sebastian Thunert

$$- f_S(a_S) = f_S(e_S) = a_S, f_S(b_S) = f_S(d_S) = d_S, - g_S(a_S) = g_S(e_S) = g_S(d_S) = d_S \text{ and } g_S(b_S) = a_S.$$

Then S satisfies A, f_S and g_S are monotone. We prove that S is a model of \mathcal{K} : Let $x, y \in S$. Assume that $x \leq g_S(y)$. We show that $f_S(x) \leq g_S(y)$.

- If $y \in \{a_S, e_S, d_S\}$ then $g_S(y) = d_S$ so $x = d_S$, and $f_S(d_S) = d_S \leq g_S(y)$.
- If $y = b_S$ then $g_S(b_S) = a_S$, so x can be a_S, e_S or d_S , and $f_S(a_S) = f_S(e_S) = a_S$, $f_S(d_S) = d_S$, so $f_S(x) \le g_S(b_S) = a_S$.

A term t containing only g and e can be e or can contain occurrences of g. If t = e then the interpretation of t in S is $e_S \neq a_S$. If t contains occurrences of g it can be proven that the interpretation of t in S is d_S , i.e. is again different from a_S .

Thus $\mathcal{T} = \mathsf{SLat}_{\Sigma}(\mathcal{K})$ does not have the Beth definability property w.r.t. Σ_S , hence, by Theorem 4, $\mathcal{T} \cup \mathcal{T}' = \mathsf{SLat}_{f,g}(\mathcal{K}) \cup \mathsf{SLat}_{f',g}(\mathcal{K}') = \mathsf{SLat}_{f,f',g}(\mathcal{K} \cup \mathcal{K}')$, where $\mathcal{K}' = \{y \leq g(x) \rightarrow f'(y) \leq g(x)\}$, does not have the \approx -interpolation property w.r.t. intersection-sharing, hence it does not have the \leq -interpolation property w.r.t. intersection-sharing.

Remark: By Theorem 11 and Theorem 4, \mathcal{T} has the $\Theta_{\mathcal{K}}(\Sigma_S)$ -Beth definability property, where $\Theta_{\mathcal{K}}(\Sigma_S) = \{f, g, e\}$. Indeed, then $A \models a \approx f(e)$. \Box

6 Applications to \mathcal{EL} and \mathcal{EL}^+ -Subsumption

We now explain how these results can be used in the study of the description logics \mathcal{EL} and \mathcal{EL}^+ . In any description logic a set N_C of concept names and a set N_R of roles is assumed to be given. Concept descriptions can be defined with the help of a set of concept constructors. The available constructors determine the expressive power of a description logic. If we only allow intersection and existential restriction as concept constructors, we obtain the description logic \mathcal{EL} [1], a logic used in terminological reasoning in medicine [31,30]. The table below shows the constructor names used in \mathcal{EL} and their semantics.

Constructor name	Syntax	Semantics
conjunction	$C_1 \sqcap C_2$	$C_1^\mathcal{I} \cap C_2^\mathcal{I}$
existential restriction	$\exists r.C$	$\{x \mid \exists y((x,y) \in r^{\mathcal{I}} \text{ and } y \in C^{\mathcal{I}})\}\$

The semantics is given by interpretations $\mathcal{I} = (\Delta, \cdot^{\mathcal{I}})$, where $C^{\mathcal{I}} \subseteq \Delta$ and $r^{\mathcal{I}} \subseteq \Delta^2$ for every $C \in N_C$, $r \in N_R$. The extension of $\cdot^{\mathcal{I}}$ to concept descriptions is inductively defined using the semantics of the constructors. In [3,2], the extension \mathcal{EL}^+ of \mathcal{EL} with role inclusion axioms is studied.

Definition 12. A TBox (or terminology) is a finite set consisting of general concept inclusions (GCI) of the form $C \sqsubseteq D$, where C and D are concept descriptions. A CBox consists of a TBox and a set of role inclusions of the form $r_1 \circ \cdots \circ r_n \sqsubseteq s$, so we view CBoxes as unions $GCI \cup \mathcal{R}$ of a set GCI of general

concept inclusions and a set \mathcal{R} of role inclusions of the form $r_1 \circ \cdots \circ r_n \sqsubseteq s$, with $n \ge 1.^3$

Definition 13. An interpretation \mathcal{I} is a model of the CBox $\mathcal{C} = GCI \cup \mathcal{R}$ if it is a model of GCI, i.e., $C^{\mathcal{I}} \subseteq D^{\mathcal{I}}$ for every $C \sqsubseteq D \in GCI$, and satisfies all role inclusions in \mathcal{C} , i.e., $r_1^{\mathcal{I}} \circ \cdots \circ r_n^{\mathcal{I}} \subseteq s^{\mathcal{I}}$ for all $r_1 \circ \cdots \circ r_n \subseteq s \in \mathcal{R}$. If \mathcal{C} is a CBox and C_1, C_2 are concept descriptions, then $\mathcal{C} \models C_1 \sqsubseteq C_2$ if and only if $C_1^{\mathcal{I}} \subseteq C_2^{\mathcal{I}}$ for every model \mathcal{I} of \mathcal{C} .

6.1 Algebraic semantics for $\mathcal{EL}, \mathcal{EL}^+$ and \sqsubseteq -interpolation

In [24] we studied the link between TBox subsumption in \mathcal{EL} and uniform word problems in the corresponding classes of semilattices with monotone functions. In [26], we showed that these results naturally extend to CBoxes and to the description logic \mathcal{EL}^+ . When defining the semantics of \mathcal{EL} or \mathcal{EL}^+ with role names N_R we use a class of \sqcap -semilattices with monotone operators of the form SLat_{Σ} , where $\Sigma = \{f_r \mid r \in N_R\}$. Every concept description C can be represented as a term \overline{C} ; the encoding is inductively defined:

- Every concept name $C \in N_C$ is regarded as a constant $\overline{C} = C$. - $\overline{C_1 \sqcap C_2} := \overline{C_1} \sqcap \overline{C_2}$, and

$$- \exists rC = f_r(C).$$

If \mathcal{R} is a set of role inclusions of the form $r \sqsubseteq s$ and $r_1 \circ r_2 \sqsubseteq s$, let \mathcal{K} be the set of all axioms of the form:

 $\begin{aligned} &\forall x \ (f_r(x) \leq f_s(x)) & \text{for all } r \sqsubseteq s \in \mathcal{R} \\ &\forall x \ (f_{r_1}(f_{r_2}(x)) \leq f_s(x)) & \text{for all } r_1 \circ r_2 \sqsubseteq s \in \mathcal{R} \end{aligned}$

Theorem 13 ([26]) Assume that the only concept constructors are intersection and existential restriction. Then for all concept descriptions D_1, D_2 and every \mathcal{EL}^+ CBox $\mathcal{C}=GCI\cup\mathcal{R}$ – where \mathcal{R} consists of role inclusions of the form $r \sqsubseteq s$ and $r_1 \circ r_2 \sqsubseteq s$ – with concept names $N_C = \{C_1, \ldots, C_n\}$ and set of roles N_R :

$$\mathcal{C} \models D_1 \sqsubseteq D_2$$
 iff $\left(\bigwedge_{C \sqsubseteq D \in GCI} \overline{C} \le \overline{D} \right) \models_{\mathsf{SLat}_{\Sigma}(\mathcal{K})} \overline{D_1} \le \overline{D_2},$

where Σ is associated with N_R and \mathcal{K} with \mathcal{R} as described above.

In [33,9] the following notion of interpolation, which we call \sqsubseteq -interpolation, is defined: A description logic has the \sqsubseteq -interpolation property if for any CBoxes $\mathcal{C}_A = GCI_A \cup \mathcal{R}_A, \ \mathcal{C}_B = GCI_B \cup \mathcal{R}_B$ and any concept descriptions C, Dsuch that $\mathcal{C}_A \cup \mathcal{C}_B \models C \sqsubseteq D$ there exists a concept description T containing only concept and role symbols "shared" by $\{\mathcal{C}_A, C\}$ and $\{\mathcal{C}_B, D\}$ such that

³ It can be shown that it is sufficient to consider role inclusions of the form $r \sqsubseteq s$ or $r_1 \circ r_2 \sqsubseteq s$, where r, s, r_1, r_2 are role names [3].

 $\mathcal{C}_A \cup \mathcal{C}_B \models C \sqsubseteq T$ and $\mathcal{C}_A \cup \mathcal{C}_B \models T \sqsubseteq D$. By Theorem 13, $\mathcal{C}_A \cup \mathcal{C}_B \models C \sqsubseteq D$ if and only if $A \land B \models_{\mathsf{SLat}_{\Sigma}(\mathcal{K})} \overline{C} \leq \overline{D}$, where $A = \bigwedge_{C_1 \sqsubseteq C_2 \in GCI_A} \overline{C_1} \leq \overline{C_2}$ and $B = \bigwedge_{C_1 \sqsubseteq C_2 \in GCI_B} \overline{C_1} \leq \overline{C_2}$, and $\mathcal{K} = \mathcal{K}_A \cup \mathcal{K}_B$, the union of the axioms associated with the set inclusions \mathcal{R}_A resp. \mathcal{R}_B . By Theorem 11, there exists a term containing only constants and function symbols $\Theta_{\mathcal{K}_A \cup \mathcal{K}_B}$ -shared by A and Bsuch that $A \land B \models_{\mathsf{SLat}_{\Sigma}(\mathcal{K}_A \cup \mathcal{K}_B)} \overline{C} \leq t \land t \leq \overline{D}$. From t we can construct a concept description T containing only concept names and roles shared by \mathcal{C}_A and \mathcal{C}_B , and by Theorem 13, $C_A \land C_B \models C \sqsubseteq T \land T \sqsubseteq D$. Therefore, the \sqsubseteq -interpolation problem studied for description logics in [33,9] can be expressed in the case of \mathcal{EL} and \mathcal{EL}^+ as a \leq -interpolation problem in the class of semilattices with operators, and the hierarchical method for \leq -interpolation can be used in this case. We distinguish between intersection-sharing and $\Theta_{\mathcal{R}}$ -sharing, where $\Theta_{\mathcal{R}}$ is the analogon of $\Theta_{\mathcal{K}}$ where \mathcal{K} is the translation of \mathcal{R} .

Corollary 14 \mathcal{EL} and \mathcal{EL}^+ have the \sqsubseteq -interpolation property w.r.t. $\Theta_{\mathcal{R}}$ -sharing.

Corollary 15 \mathcal{EL}^+ with role inclusions of the form $r_1 \circ r_2 \sqsubseteq s$ does not have \sqsubseteq -interpolation w.r.t. intersection-sharing.

6.2 Example: \Box -Interpolation for \mathcal{EL}^+

We now explain our method in detail and illustrate each step of the method for \leq -interpolation described in Section 4 on an example.

Example 6. Consider the ontology \mathcal{O}_{Med} in Figure 1. It is based on an example from [32], which we modified in some points. We changed the CBox in order to ensure that it only contains general concept inclusions and that conjunction only appears on the left hand side of an axiom. Furthermore we left out some axioms and concepts, but also added new concepts (LeftVentricle, RightVentricle, Ventricle) and changed some axioms accordingly.

We divided the CBox into three parts: The A-part is our main TBox, \mathcal{T}_A , which is supposed to be consistent. The B-part, TBox \mathcal{T}_B , is an extension of the main CBox and may introduce some new (and in the worst case even unwanted) consequences. The R-part contains only role axioms \mathcal{R} .

We have the following sets of symbols (we indicate also the abbreviations used in what follows):

$$\begin{split} N^A_C = & \{ & \text{Endocardium (Em), Tissue (T), HeartWall (HW),} \\ & & \text{LeftVentricle (LV), RightVentricle (RV), Ventricle (V),} \\ & & \text{Disease (D), Inflammation (I), Endocarditis (Es)} \\ N^B_C = & \{ & \text{Heart (H), HeartDisease (HD), Disease (D), Ventricle (V)} \} \end{split}$$

$$N_C^{AB} = \{ \text{Disease (D), Ventricle (V)} \}$$

Consider the subsumption Endocarditis \sqsubseteq HeartDisease. We have Endocarditis $\in N_C^A$ and HeartDisease $\in N_C^B$ and additionally the following hold:

 $\mathcal{T}_A \cup \mathcal{T}_B \cup \mathcal{R} \models \mathsf{Endocarditis} \sqsubseteq \mathsf{HeartDisease}$

$A_1:$	Endocardium	Tissue
$A_2:$	Endocardium	∃part-of.HeartWall
A_3 :	HeartWall	BodyWall
A_4 :	HeartWall	$\exists part-of.LeftVentricle$
A_5 :	HeartWall	$\exists part-of.RightVentricle$
A_6 :	LeftVentricle	Ventricle
$A_7:$	RightVentricle	Ventricle
A_8 :	Endocarditis	Inflammation
A_9 :	Endocarditis	∃has-location.Endocardium
$A_{10}:$	${\sf Inflammation} \sqcap \exists {\sf has}{\sf -location}.{\sf Endocardium}$	Endocarditis
$A_{11}:$	Inflammation	Disease
$A_{12}:$	Inflammation	∃acts-on.Tissue
B_1 :	Ventricle	∃part-of.Heart
B_2 :	HeartDisease	Disease
B_3 :	HeartDisease	∃has-location.Heart
B_4 :	$Disease \sqcap \exists has\text{-}location.Heart$	HeartDisease
R_1 :	part-of \circ part-of	part-of
R_2 :	has-location \circ part-of	has-location

P-interpolation in local theory extensions and applications to $\mathcal{EL}, \mathcal{EL}^+$ 27

Fig. 1: Ontology \mathcal{O}_{Med}

and, in addition:

 $\mathcal{T}_A \cup \mathcal{R} \not\models \mathsf{Endocarditis} \sqsubseteq \mathsf{HeartDisease} \quad \mathcal{T}_B \cup \mathcal{R} \not\models \mathsf{Endocarditis} \sqsubseteq \mathsf{HeartDisease}$

Therefore, we can use the method described in Section 4 (based on Proposition 7, Theorem 8 and Lemma 9) to compute an intermediate term containing only shared symbols for the subsumption Endocarditis \sqsubseteq HeartDisease, which serves as an explanation for the subsumption.

Step 1: We translate the original ontology to the theory of semilattices with operators. We now state the monotonicity axioms for each role explicitly. Figure 2 shows the ontology after the translation to the theory of semilattices with operators. Note that from here on we use the abbreviations for concept names indicated in the sets N_C^A, N_C^B and N_C^{AB} above and also abbreviations for role names, i.e. po for part-of, hl for has-location and ao for acts-on.

Step 2: Using unsat core computation we get the following minimal axiom set: $min_A = \{A_2, A_4, A_6, A_8, A_9, A_{11}, B_1, B_4, R_2\}$

$A_1:$	Em	\leq	Т				
A_2 :	Em	\leq	po(HW)	B_1 :	V	\leq	po(H)
A_3 :	HW	\leq	BW	B_2 :	HD	\leq	D
A_4 :	HW	\leq	po(LV)	B_3 :	HD	\leq	hl(H)
A_5 :	HW	\leq	po(RV)	B_4 :	$D\wedgehl(H)$	\leq	HD
A_6 :	LV	\leq	V				
$A_7:$	RV	\leq	V	R_1 :	∀X: po(po(X))	\leq	po(X)
A_8 :	Es	\leq	I	R_2 :	∀X: hl(po(X))	\leq	hl(X)
A_9 :	Es	\leq	hl(Em)				
$A_{10}:$	$I \wedge hl(Em)$	\leq	Es	M_1 :	$\forall X, Y: X \leq Y$	\rightarrow	$po(X) \leq po(Y)$
$A_{11}:$	I	\leq	D	M_2 :	$\forall X, Y: X \leq Y$	\rightarrow	$hl(X) \leq hl(Y)$
$A_{12}:$	I	\leq	ao(T)	M_3 :	$\forall X,Y: \ X \leq Y$	\rightarrow	$ao(X) \leq ao(Y)$

Fig. 2: \mathcal{O}_{Med} after translation to SLat with monotone operators

This means that for the following instantiation step we only have to consider the role axiom R_2 and none of the monotonicity axioms is needed.

Step 3: Let $\mathcal{T}_0 = \mathsf{SLat}$ and $T_1 = \mathsf{SLat} \cup R_2$ be the extension of T_0 with axiom R_2 . We know that it is a local theory extension, so we can use hierarchical reasoning. We first flatten the role axiom R_2 in the following way:

 $R_2^{\text{flat}}: \forall X, Y: X \leq \text{po}(Y) \rightarrow hl(X) \leq hl(Y)$ We have the following set of ground terms:

 $G = \text{est}(\mathcal{T}_A \cup \mathcal{T}_B \cup \mathcal{R}) = \{ \text{po}(HW), \text{po}(LV), \text{po}(H), \text{hl}(Em), \text{hl}(H) \}$ We use the closure operator Ψ described in Lemma 10 to extend our set of ground terms: For every term po(X) in G we have to add the term hl(X) and vice versa. This leads to the following extended set G' of ground terms:

 $G' = \{ po(Em), po(HW), po(LV), po(H), hl(Em), hl(HW), hl(LV), hl(H) \}$ From G' we get the following instances of the axiom R_2^{flat} :

I_1 :	$Em \le po(HW)$	\rightarrow	$hl(Em) \leq hl(HW)$
I_2 :	$Em \le po(LV)$	\rightarrow	$hl(Em) \leq hl(LV)$
I_3 :	$Em \leq po(H)$	\rightarrow	$hl(Em) \leq hl(H)$
I_4 :	$HW \le po(Em)$	\rightarrow	$hl(HW) \leq hl(Em)$
I_5 :	$HW \le po(LV)$	\rightarrow	$hl(HW) \leq hl(LV)$
I_6 :	$HW \leq po(H)$	\rightarrow	$hl(HW) \leq hl(H)$
I_7 :	$LV \le po(Em)$	\rightarrow	$hl(LV) \leq hl(Em)$
I_8 :	$LV \leq po(HW)$	\rightarrow	$hl(LV) \leq hl(HW)$
I_9 :	$LV \le po(H)$	\rightarrow	$hl(LV) \leq hl(H)$
$I_{10}:$	$H \leq po(Em)$	\rightarrow	$hl(H) \leq hl(Em)$
$I_{11}:$	$H \leq po(HW)$	\rightarrow	$hl(H) \leq hl(HW)$
$I_{12}:$	$H \leq po(LV)$	\rightarrow	$hI(H) \le hI(LV)$

We purify all formulae by introducing new constants for the terms starting with a function symbol, i.e. role names. We save the definitions in the following set:

$$\mathsf{Def} = \{\mathsf{po}_{\mathsf{HW}} = \mathsf{po}(\mathsf{HW}), \mathsf{po}_{\mathsf{LV}} = \mathsf{po}(\mathsf{LV}), \mathsf{po}_{\mathsf{H}} = \mathsf{po}(\mathsf{H}), \mathsf{hl}_{\mathsf{EM}} = \mathsf{hl}(\mathsf{EM}), \\ \mathsf{hl}_{\mathsf{HW}} = \mathsf{hl}(\mathsf{HW}), \mathsf{hl}_{\mathsf{LV}} = \mathsf{hl}(\mathsf{LV}), \mathsf{hl}_{\mathsf{HC}} = \mathsf{hl}(\mathsf{HC}), \mathsf{hl}_{\mathsf{H}} = \mathsf{hl}(\mathsf{H})\}$$

We then have the set $A_0 \wedge B_0 \wedge I_0$, where A_0 , B_0 and I_0 are the purified versions of $A = \{A_2, A_4, A_6, A_8, A_9, A_{11}\}, B = \{B_1, B_4\}$ and $I = \{I_1, ..., I_{10}\}$, respectively.

Step 4: To reduce the number of instances we compute an unsatisfiable core and obtain the following set of axioms:

 $min'_A = \{A_2, A_4, A_6, A_8, A_9, A_{11}, B_1, B_4, I_1, I_5, I_9\}$ So we have $\mathcal{H} = \{I_1, I_5, I_9\}$. Out of these instances the first two are pure A (meaning the premise contains only symbols in N_C^A), but I_9 is a mixed instance, since $\mathsf{LV} \in N_C^A \setminus N_C^B$ and $\mathsf{H} \in N_C^B \setminus N_C^A$, so $\mathcal{H}_{\mathsf{mix}} = \{I_9\}$.

Step 5: To separate the mixed instance $LV \leq po_H \rightarrow hl_{LV} \leq hl_H$ we use the construction in Proposition 7 to compute an intermediate term t in the common signature such that $LV \leq t$ and $t \leq po_{H}$. We obtain t = V. We get $\mathcal{H}_{sep} =$ $\{I_9^A, I_9^B\}$ where:

$$\begin{array}{rcl} I_9^A : & \mathsf{LV} \leq \mathsf{V} & \to & \mathsf{hl}_{\mathsf{LV}} \leq \mathsf{hl}_{\mathsf{V}} \\ I_9^B : & \mathsf{V} \leq \mathsf{po}_{\mathsf{H}} & \to & \mathsf{hl}_{\mathsf{V}} \leq \mathsf{hl}_{\mathsf{H}} \\ \end{array}$$

Note that I_9^A is an instance of the monotonicity axiom for the has-location role and I_9^B is an instance of axiom R_2^{flat} .

Step 6: Since for all the instances that are necessary to derive the consequence it must be true that $\mathcal{T}_A \cup \mathcal{T}_B$ entails its premise, it is sufficient to consider only the corresponding conclusions. Note that w.r.t. SLat the formula $A_0 \wedge I_1 \wedge I_5 \wedge I_9^A$ is equivalent to the following formula:

$$\begin{array}{rl} A_{0} = & \mathsf{Em} \leq \mathsf{po}_{\mathsf{HW}} \land \mathsf{HW} \leq \mathsf{po}_{\mathsf{LV}} \land \mathsf{LV} \leq \mathsf{V} \land \mathsf{Es} \leq \mathsf{I} \land \mathsf{Es} \leq \mathsf{hl}_{\mathsf{Em}} \land \mathsf{I} \leq \mathsf{D} \\ & \land \mathsf{hl}_{\mathsf{EM}} \leq \mathsf{hl}_{\mathsf{HW}} \land \mathsf{hl}_{\mathsf{HW}} \leq \mathsf{hl}_{\mathsf{LV}} \land \mathsf{hl}_{\mathsf{LV}} \leq \mathsf{hl}_{\mathsf{V}} \end{array}$$

This formula can be seen as a set of Horn clauses \overline{A}_0^h :

$$\overline{A}_0^n = \{ (\neg \mathsf{Em} \lor \mathsf{po}_{\mathsf{HW}}), (\neg \mathsf{HW} \lor \mathsf{po}_{\mathsf{LV}}), (\neg \mathsf{LV} \lor \mathsf{V}), (\neg \mathsf{Es} \lor \mathsf{I}), (\neg \mathsf{Es} \lor \mathsf{hl}_{\mathsf{Em}}), (\neg \mathsf{I} \lor \mathsf{D}), (\neg \mathsf{hl}_{\mathsf{EM}} \lor \mathsf{hl}_{\mathsf{HW}}), (\neg \mathsf{hl}_{\mathsf{HW}} \lor \mathsf{hl}_{\mathsf{LV}}), (\neg \mathsf{hl}_{\mathsf{LV}} \lor \mathsf{hl}_{\mathsf{V}}) \}$$

To obtain an explanation for $\mathcal{T}_A \cup \mathcal{T}_B \cup \mathcal{R} \models \mathsf{Endocarditis} \sqsubseteq \mathsf{HeartDisease}$ we saturate the set $\overline{A}_0^n \cup \{\mathsf{Es}\}$ under ordered resolution as described in the proof of Theorem 9, where symbols occurring in A but not in B are larger than common symbols:

- Resolution of Es and \neg Es \lor I yields I.
- Resolution of I and $\neg I \lor D$ yields D.
- Resolution of Es and $\neg \mathsf{Es} \lor \mathsf{hl}_{\mathsf{Em}}$ yields $\mathsf{hl}_{\mathsf{Em}}.$
- Resolution of hI_{Em} and $\neg hI_{\mathsf{EM}} \lor hI_{\mathsf{HW}}$ yields $hI_{\mathsf{HW}}.$
- Resolution of hl_{HW} and $\neg hl_{HW} \lor hl_{LV}$ yields hl_{LV} .

30 Dennis Peuter, Viorica Sofronie-Stokkermans, Sebastian Thunert

– Resolution of hl_{LV} and $\neg hl_{LV} \lor hl_V$ yields hl_V .

We obtained two resolvents containing only common symbols: D and hl_V . Taking the conjunction of these terms and translating the formula back to description logic yields the following formula:

 $J = \text{Disease} \ \sqcap \ \exists \text{has-location.Ventricle.}$ Indeed, the following properties hold:

$$\mathcal{T}_A \cup \mathcal{R} \models \mathsf{Endocarditis} \sqsubseteq J$$
$$\mathcal{T}_A \cup \mathcal{T}_B \cup \mathcal{R} \models \qquad J \sqsubseteq \mathsf{HeartDisease}$$

So J is the intermediate term we were looking for.

6.3 Prototype implementation

The ideas were implemented in a prototype implementation⁴ for the theory of semilattices with operators satisfying axioms of type (1) considered in this paper. The program is written in Python and uses Z3 [8] and SPASS [35] as external provers. The program implements Steps 1-3 in the algorithm presented at the end of Section 4 with the following optimization: In Step 1 after instantiation and purification, in order to reduce the size of the set of instances of axioms to be considered, an unsatisfiable core is computed with Z3. The program separates the mixed instances by computing intermediate terms for their premises using Theorem 9 and Proposition 7; for applying ordered resolution the prover SPASS is used. In Step 3, the intermediate term T for $C \leq D$ is computed using the method described in Theorem 9, again using SPASS.

For the use for interpolation in \mathcal{EL} and \mathcal{EL}^+ , the CBoxes \mathcal{C}_A and \mathcal{C}_B and the subsumption $C \sqsubseteq D$ are given as an input. A minimal subset of $\mathcal{C}_A \cup \mathcal{C}_B$ is computed from which $C \sqsubseteq D$ can be derived. (The user can choose between a precise translation to SPASS or a propositional translation to Z3 which is not always precise, but turned out to be a good approximation. Standard implementations available for computing justifications of entailments from description logic ontologies could be used as well.) The problem is then translated into a problem for \leq -interpolation in semilattices with operators. After computing the interpolating term, the result is expressed in the syntax of description logics.

7 Conclusions and future work

In this paper we gave a hierarchical method for *P*-interpolation in certain classes of local theory extensions $\mathcal{T}_0 \subseteq \mathcal{T}_0 \cup \mathcal{K}$. We used these results for proving \leq interpolation in classes of semilattices with monotone operators satisfying additional clauses \mathcal{K} with a suitable notion of $\Theta_{\mathcal{K}}$ -sharing we defined. We defined a form of Beth definability w.r.t. a subsignature Σ_S and used it to show that

 $^{^4}$ The implementation and some tests can be found here:

https://userpages.uni-koblenz.de/~sofronie/p-interpolation-and-el/

the class of semilattices with operators under consideration does not have the \leq -interpolation property if only the common function symbols and constants are considered to be "shared". We discussed how these results can be used for the study of interpolation in \mathcal{EL} and \mathcal{EL}^+ .

In future work we will explore other application areas of these results, both to classes of non-classical logics and to theories relevant in the verification. We will extend the implementation with possibilities of choosing the base theory and the methods for P-interpolation in the base theory. We will further investigate the links with Beth definability and possibilities of using Beth definability for computing explicit definitions for implicitly definable terms – and analyze the applicability of such results in description logics but also in verification.

Acknowledgments. We thank the reviewers for their helpful comments. The research reported here was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Projektnummer 465447331.

References

- F. Baader. Terminological cycles in a description logic with existential restrictions. In G. Gottlob and T. Walsh, editors, *IJCAI-03, Proceedings of the 18th International Joint Conference on Artificial Intelligence*, pages 325–330. Morgan Kaufmann, 2003.
- F. Baader, C. Lutz, and B. Suntisrivaraporn. Efficient reasoning in *EL*⁺. In B. Parsia, U. Sattler, and D. Toman, editors, *Proceedings of the 2006 International Workshop on Description Logics (DL2006)*, CEUR Workshop Proceedings 189. CEUR-WS.org, 2006.
- F. Baader, C. Lutz, and B. Suntisrivaraporn. Is tractable reasoning in extensions of the description logic *EL* useful in practice? *Journal of Logic, Language and Information*, 2007. Special issue on Method for Modality (M4M), 2007.
- R. Bruttomesso, S. Ghilardi, and S. Ranise. Quantifier-free interpolation in combinations of equality interpolating theories. ACM Trans. Comput. Log., 15(1):5:1– 5:34, 2014.
- 5. S. Burris and H. P. Sankappanavar. A course in universal algebra, volume 78 of Graduate texts in mathematics. Springer, 1981.
- D. Calvanese, S. Ghilardi, A. Gianola, M. Montali, and A. Rivkin. Combination of uniform interpolants via Beth definability. J. Autom. Reason., 66(3):409–435, 2022.
- 7. C. C. Chang and H. J. Keisler. Model Theory (3rd edition). North-Holland, 1990.
- L. M. de Moura and N. S. Bjørner. Z3: an efficient SMT solver. In C. R. Ramakrishnan and J. Rehof, editors, *Tools and Algorithms for the Construction and Analysis* of Systems, 14th International Conference, TACAS 2008, Proceedings, volume 4963 of Lecture Notes in Computer Science, pages 337–340. Springer, 2008.
- M. Fortin, B. Konev, and F. Wolter. Interpolants and explicit definitions in extensions of the description logic *EL*. In G. Kern-Isberner, G. Lakemeyer, and T. Meyer, editors, *Proceedings of the 19th International Conference on Principles* of Knowledge Representation and Reasoning, KR 2022, 2022.
- 10. H. Ganzinger. Relating semantic and proof-theoretic concepts for polynomial time decidability of uniform word problems. In 16th Annual IEEE Symposium on Logic

in Computer Science, Boston, Massachusetts, USA, June 16-19, 2001, Proceedings, pages 81–90. IEEE Computer Society, 2001.

- 11. W. Hodges. Model theory, volume 42 of Encyclopedia of mathematics and its applications. Cambridge University Press, 1993.
- C. Ihlemann, S. Jacobs, and V. Sofronie-Stokkermans. On local reasoning in verification. In C. R. Ramakrishnan and J. Rehof, editors, *Tools and Algorithms for* the Construction and Analysis of Systems, 14th International Conference, TACAS 2008, Proceedings, volume 4963 of Lecture Notes in Computer Science, pages 265– 281. Springer, 2008.
- C. Ihlemann and V. Sofronie-Stokkermans. On hierarchical reasoning in combinations of theories. In J. Giesl and R. Hähnle, editors, Automated Reasoning, 5th International Joint Conference, IJCAR 2010, Proceedings, volume 6173 of Lecture Notes in Computer Science, pages 30–45. Springer, 2010.
- L. Kovács and A. Voronkov. Interpolation and symbol elimination. In R. A. Schmidt, editor, Automated Deduction - CADE-22, 22nd International Conference on Automated Deduction, Proceedings, volume 5663 of Lecture Notes in Computer Science, pages 199–213. Springer, 2009.
- 15. J. Krajícek. Interpolation theorems, lower bounds for proof systems, and independence results for bounded arithmetic. J. Symb. Log., 62(2):457–486, 1997.
- D. A. McAllester. Automatic recognition of tractability in inference relations. J. ACM, 40(2):284–303, 1993.
- 17. K. L. McMillan. An interpolating theorem prover. In K. Jensen and A. Podelski, editors, Tools and Algorithms for the Construction and Analysis of Systems, 10th International Conference, TACAS 2004, Proceedings, volume 2988 of Lecture Notes in Computer Science, pages 16–30. Springer, 2004.
- 18. D. Peuter and V. Sofronie-Stokkermans. Finding high-level explanations for subsumption w.r.t. combinations of CBoxes in *EL* and *EL⁺*. In S. Borgwardt and T. Meyer, editors, *Proceedings of the 33rd International Workshop on Description* Logics (DL 2020) co-located with the 17th International Conference on Principles of Knowledge Representation and Reasoning (KR 2020), volume 2663 of CEUR Workshop Proceedings. CEUR-WS.org, 2020.
- D. Peuter, V. Sofronie-Stokkermans, and S. Thunert. On P-interpolation in local theory extensions and applications to the study of interpolation in the description logics \$\mathcal{EL}, \mathcal{EL}^+\$. Proceedings of the 29th International Conference on Automated Deduction (CADE-29), 2023. volume 14132 of Lecture Notes in Computer Science, Springer, to appear.
- A. Rybalchenko and V. Sofronie-Stokkermans. Constraint solving for interpolation. J. Symb. Comput., 45(11):1212–1233, 2010.
- T. Skolem. Logisch-kombinatorische Untersuchungen über die Erfüllbarkeit oder Beweisbarkeit mathematischer Sätze nebst einem Theorem über dichte Mengen. In Selected Works in Logic. Universitetsforlaget, 1920.
- V. Sofronie-Stokkermans. Hierarchic reasoning in local theory extensions. In R. Nieuwenhuis, editor, Automated Deduction - CADE-20, 20th International Conference on Automated Deduction, Proceedings, volume 3632 of Lecture Notes in Computer Science, pages 219–234. Springer, 2005.
- V. Sofronie-Stokkermans. Interpolation in local theory extensions. In U. Furbach and N. Shankar, editors, Automated Reasoning, Third International Joint Conference, IJCAR 2006, Proceedings, volume 4130 of Lecture Notes in Computer Science, pages 235–250. Springer, 2006.
- V. Sofronie-Stokkermans. Automated theorem proving by resolution in nonclassical logics. Ann. Math. Artif. Intell., 49(1-4):221–252, 2007.

- V. Sofronie-Stokkermans. Interpolation in local theory extensions. Log. Methods Comput. Sci., 4(4), 2008.
- V. Sofronie-Stokkermans. Locality and subsumption testing in *EL* and some of its extensions. In C. Areces and R. Goldblatt, editors, *Advances in Modal Logic 7*, pages 315–339. College Publications, 2008.
- 27. V. Sofronie-Stokkermans. Representation theorems and locality for subsumption testing and interpolation in the description logics \mathcal{EL} , \mathcal{EL}^+ and their extensions with *n*-ary roles and numerical domains. Fundamenta Informaticae, 156(3-4):361–411, 2017.
- V. Sofronie-Stokkermans and C. Ihlemann. Automated reasoning in some local extensions of ordered structures. In 37th International Symposium on Multiple-Valued Logic, ISMVL 2007, page 1. IEEE Computer Society, 2007.
- V. Sofronie-Stokkermans and C. Ihlemann. Automated reasoning in some local extensions of ordered structures. *Multiple-Valued Logic and Soft Computing*, 13(4-6):397–414, 2007.
- K. A. Spackman. Normal forms for description logic expressions of clinical concepts in SNOMED RT. In AMIA 2001, American Medical Informatics Association Annual Symposium. AMIA, 2001.
- K. A. Spackman, K. E. Campbell, and R. A. Côté. SNOMED RT: a reference terminology for health care. In AMIA 1997, American Medical Informatics Association Annual Symposium. AMIA, 1997.
- B. Suntisrivaraporn. Polynomial time reasoning support for design and maintenance of large-scale biomedical ontologies. PhD thesis, Dresden University of Technology, Germany, 2009.
- B. ten Cate, E. Franconi, and I. Seylan. Beth definability in expressive description logics. J. Artif. Intell. Res., 48:347–414, 2013.
- 34. C. Tinelli. Cooperation of background reasoners in theory reasoning by residue sharing. J. Autom. Reason., 30(1):1–31, 2003.
- C. Weidenbach, D. Dimova, A. Fietzke, R. Kumar, M. Suda, and P. Wischnewski. SPASS version 3.5. In R. A. Schmidt, editor, Automated Deduction - CADE-22, 22nd International Conference on Automated Deduction, Proceedings, volume 5663 of Lecture Notes in Computer Science, pages 140–145. Springer, 2009.
- 36. G. Yorsh and M. Musuvathi. A combination method for generating interpolants. In R. Nieuwenhuis, editor, Automated Deduction - CADE-20, 20th International Conference on Automated Deduction, Proceedings, volume 3632 of Lecture Notes in Computer Science, pages 353–368. Springer, 2005.