Higher Catoids, Higher Quantales and their Correspondences

Cameron Calk Philippe Malbos Damien Pous Georg Struth February 13, 2024

Abstract

This article is part of a programme on the formalisation of higher categories and the categorification of rewriting theory. Set-valued structures such as catoids are used in this context to formalise local categorical composition operations. We introduce ω -catoids as set-valued generalisations of (strict) ω -categories. We establish modal correspondences between ω -catoids and convolution ω -quantales. These are related to Jónsson-Tarski-style dualities between relational structures and lattices with operators. Convolution ω -quantales generalise the powerset ω -Kleene algebras recently proposed for algebraic coherence proofs in higher rewriting to weighted variants in the style of category algebras. In order to capture homotopic constructions and proofs in rewriting theory, we extend these correspondances to higher catoids with a groupoid structure above some dimension, which is reflected by an involution in higher quantales.

Keywords: higher catoids, higher quantales, multisemigroups, convolution algebras, categorification, higher rewriting.

1 Introduction

Rewriting systems are fundamental models of computation. Their rules generate computations as sequences or paths of rewriting steps. Computational structure, such as confluence or Church-Rosser properties, is modelled geometrically in terms of rewriting diagrams and algebraically via inclusions between rewriting relations, which form abstractions of sets of rewriting paths [Ter03]. Coherence properties of rewriting systems, like the Church-Rosser theorem, Newman's lemma or normal form theorems, can be derived in algebras that have binary relations among their models: Kleene algebras, quantales or relation algebras [DBvdW97, Str06, DS11, DMS11].

Higher rewriting generalises this approach to higher dimensions, using the structure of polygraphs (or computads) as higher rewriting systems and the higher paths they generate instead of rewriting relations, relations between rewriting relations, and so on [ABG⁺24]. This approach has been developed mainly in the context of (strict) ω -categories, where higher rewriting diagrams have globular shape [GM09]. They are filled with higher cells instead of relational inclusions, two-cells in **Rel**, as witnesses for the $\forall \exists$ -relationships between the higher paths that form their faces. Applications of higher rewriting range from string rewriting [GM18, GM12] to the computational analysis of coherence properties and cofibrant approximations in categorical algebra [GHM19, MR23].

It has recently been argued that algebras that admit sets of higher rewriting paths as models can once again be used for categorical coherence proofs in higher rewriting [CGMS22]. The axioms of such higher Kleene algebras must of course reflect the structure of globular cells of (ω, p) -categories and that of their pasting schemes. Yet a systematic construction of these algebras and a systematic justification of their axioms relative to the underlying (ω, p) -categories and polygraphs is so far missing.

Here we use the modal correspondences associated with the Jónsson-Tarski duality between relational structures and boolean algebras with operators for such a justification in the special case of ternary relations and binary modal operators. To apply the tools for these constructions, we generalise from ω -categories to ω -catoids, which are isomorphic to ternary-relational structures, and to ω -quantales, whose multiplications we consider as binary modalities. The correspondence between ω -catoids and ω -quantales then lifts properties

of the former (and thus of ω -categories) to identities in the latter. In the other direction, we can recover ω -catoids within the ω -quantales constructed. The ω -quantales considered are in general convolution quantales on spaces of functions from ω -catoids to ω -quantales. The power set ω -quantales suitable for higher rewriting arise as special cases. We extend these results to correspondences between (ω, p) -catoids and (ω, p) -quantales, and relate these quantales to the higher power set Kleene algebras previously used [CGMS22].

The techniques for proving such correspondences between catoids and quantales have been developed in a series of papers [DHS16, DHS21, CDS20, CDS21, FJSZ23]. They are inspired by group or category algebra constructions from representation theory or monoidal topology [HST14]. The notion of catoid, in particular, originates in [CDS20, FJSZ23]. Intuitively, catoids are (single set) categories in which arrow composition is relaxed to a multioperation, which maps pairs of arrows to sets of arrows that may be empty, and where compositions may be empty, hence undefined, even when ends of arrows match. See [CDS20, FJSZ23] for examples. Multioperations $X \times X \to \mathcal{P}X$ are obviously isomorphic to ternary relations $X \times X \times X \to 2$. In this context, we study correspondence triangles between catoids C, value quantales Q and convolution quantales Q^C on function spaces. In the simplest case, if C is a catoid and Q a quantale, then Q^C is a quantale; if Q^C and Q are quantales, then C is a catoid; and if Q^C is a quantale and C a catoid, then Q is a quantale [DHS21, CDS21]. The last two of these two-out-of-three properties require mild non-degeneracy conditions on C or Q explained in Section 8. In the construction of the convolution quantale Q^C , if $Q \times Q \to Q$ the composition in the value quantale Q and if Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q and if Q are maps in Q are maps in Q are maps

$$(f * g)(x) = \bigvee_{x \in y \odot z} f(y) \cdot g(z),$$

where the sup is taken with respect to y and z. In the powerset case Q=2, where 2 indicates the quantale of booleans, we can construct a catoid from the atom structure of any powerset quantale in the standard way for Stone- and Jónsson-Tarski-type duality. The atoms in $\mathcal{P}C$ are singleton sets; the relationship $x \in y \odot z \Leftrightarrow \{x\} \subseteq \{y\} * \{z\}$ and properties of compositions of singletons with the unit of the powerset generate the catoid. For arbitrary value quantales Q, functions $\delta_x^{\alpha}: C \to Q$, which map $y \in C$ to $\alpha \in Q$ if y = x and to $\alpha \in Q$ otherwise, where $\alpha \in Q$ is the minimal element of the quantale, replace singleton sets as atoms, and allow us to construct $\alpha \in Q$ and $\alpha \in Q$ and $\alpha \in Q$ and $\alpha \in Q$ and $\alpha \in Q$ from $\alpha \in Q$ and $\alpha \in Q$ from $\alpha \in Q$ and $\alpha \in Q$ from quantale, it is thus necessary and sufficient to understand the structure of the underlying catoid (if the value quantale is fixed).

Beyond this basic correspondence triangle, our results for ω -catoids and ω -quantales are based on two extensions. The first one is a correspondence triangle between concurrent catoids C, in isomorphic relational form, concurrent value quantales Q and concurrent convolution quantales Q^C [CDS21]. Catoids are then equipped with two multioperations and quantales with two monoidal structures that interact via interchange laws. For this extension, it remains to establish correspondences between interchange laws to the basic ones. This result justifies the axioms of concurrent semirings, Kleene algebras and quantales [HMSW11] relative to the concurrent catoid axioms and a fortiori to standard concurrency models such as shuffle and series-parallel pomset languages, including Q-weighted language models.

The second extension is a correspondence triangle between local catoids C, which exhibit the typical composition pattern of categories, modal value quantales Q and modal convolution quantales Q^C [FJSZ23]. This time, correspondences between the source and target maps of catoids and the domain and codomain maps of modal quantales must be added to the basic ones. This result justifies the laws of modal Kleene algebras [DS11] and modal quantales [FJSZ22a] relative to the catoid and category axioms from which they can be lifted. An important instance are relational modal Kleene algebras, which are algebraic variants of propositional dynamic logic, and can be lifted as powerset algebras from pair groupoids.

The correspondence triangles between ω -catoids and ω -quantales, the main results in this paper, require first of all definitions of these two structures. ω -Catoids are introduced in Section 6. They are obtained by adding globular shape axioms to those of local and concurrent catoids and then generalising beyond two dimensions in light of previous axiomatisations of (single-set) ω -categories [BH81, Str87, ML98, Ste04]. Based on the ω -catoid axioms and on previous axioms for globular n-Kleene algebras [CGMS22], we then

introduce ω -quantales in Section 7, and justified their axioms through the correspondence proofs in Section 8. Results for n-structures can obtained in the standard way by truncation; correspondence results for powerset ω -quantales are discussed in Section 10.

The ω -catoid axioms depend on rather delicate definedness conditions for compositions, captured multioperationally by mapping to non-empty sets. These are sensitive to Eckmann-Hilton-style collapses, as shown in Appendix A. They satisfy natural functorial properties, specialise to ω -category axioms and yield the globular cell structure expected, see the discussion in Section 6. The Isabelle/HOL proof assistant with its automated theorem provers and counterexample generators has allowed us to simplify these structural axioms quite significantly, which in turn simplified the development of ω -quantale axioms in Section 7 and the proofs of correspondence triangles in Section 8.

The extended correspondence triangles for (ω, p) -catoids and (ω, p) -quantales in Section 13 are compositional with respect to the ω -correspondences and those for groupoids and quantales with a suitable notion of converse. These are established in Section 9. They adapt Jónsson and Tarski's classical correspondence between groupoids and relation algebras [JT52] and extend it to convolution algebras. Catoids specialise automatically to groupoids when the two obvious inverse axioms are added, see Section 3. Yet we use Dedekind quantales instead of relation algebras. These are involutive quantales [MWP92] equipped with a variant of the Dedekind law from relation algebra, see Section 5. Apart from the interaction between the converse and the modal structure needed for (ω, p) -quantale, we also discuss weaker variants of converse useful for semirings or Kleene algebras in this section.

The content of the remaining sections of this paper is summarised as follows. In Section 2 and 4 we recall the basic properties of catoids and modal quantales. In Section 11 we develop the basic laws for modal box and diamond operators in ω -quantales to prepare for more advanced coherence proofs in higher rewriting. In Section 12 and 13 we specialise the ω - and (ω, p) -quantale axioms to those of ω -semirings and ω -Kleene algebras and their (ω, p) -variants. For general convolution algebras, this requires a finite decomposition property on ω - and (ω, p) -catoids. Section 12 also contains a detailed comparison of the ω -Kleene algebras and (ω, p) -Kleene algebras introduced in this paper with previous globular n-Kleene algebras [CGMS22] and their slightly different axioms. Overall, ω -quantales offer greater flexibility when reasoning about higher rewriting diagrams than ω -Kleene algebras. They admit additional operations such as residuals and support reasoning with least and greatest fixpoints beyond the Kleene star. Already the proof of the classical Newman's lemma in modal Kleene algebra [DMS11] assumes certain suprema that are not present in all Kleene algebras.

Most results in this paper have been verified with the Isabelle/HOL proof assistant, but the development of interactive proof support for higher categories, higher rewriting and categorical algebra remains part of a larger research programme outlined in the conclusion, which requires substantial additions. Our Isabelle components can currently be found in the Archive of Formal Proofs [Str23, CS23, CS24]. The PDF proof documents of these components cover more than 150 pages. The components contain specifications and basic libraries for 2-catoids, groupoids, 2-quantales and their ω -variants, as well as for Dedekind quantales, and they cover the basic properties of these structures that feature in this paper. All powerset liftings have been formalised with Isabelle/HOL, but neither the constructions of convolution algebras nor the full correspondences. In addition to the Isabelle proofs, we supply the most important proofs for this article in the relevant sections. All other proofs can be found in Appendix B or the references given.

2 Catoids

We recall the definitions and basic properties of catoids and related structures [FJSZ23]; see also the Isabelle theories for details. Catoids were originally called ℓr -multisemigroups, but this name became too unwieldy for the extensions introduced in this article. General background on multisemigrous can be found in [KM15] and the references given there.

A catoid is a structure (C, \odot, s, t) consisting of a set C, a multioperation $\odot: C \times C \to \mathcal{P}C$ and source

and target maps $s, t: C \to C$. These satisfy, for all $x, y, z \in C$,

$$\bigcup\{x\odot v\mid v\in y\odot z\}=\bigcup\{u\odot z\mid u\in x\odot y\},$$

$$x\odot y\neq\emptyset\Rightarrow t(x)=s(y), \qquad s(x)\odot x=\{x\}, \qquad x\odot t(x)=\{x\}.$$

The first catoid axiom expresses multirelational associativity. If we extend \odot from $C \times C \to \mathcal{P}C$ to $\mathcal{P}C \times \mathcal{P}C \to \mathcal{P}C$ such that, for all $X,Y \subseteq C$,

$$X\odot Y=\bigcup_{x\in X,y\in Y}x\odot y,$$

and write $x \odot X$ for $\{x\} \odot X$ and likewise, the associativity axiom becomes $x \odot (y \odot z) = (x \odot y) \odot z$. A multisemigroup can thus be defined as a set equipped with an associative multioperation [KM15]. From now on, we often write xy instead of $x \odot y$. We also write s(X) and t(X) for the direct images of X under s and t, respectively, for instance, $s(x \odot y)$ or s(xy) as well as $t(x \odot y)$ or t(xy).

The second catoid axiom, the *weak locality* axiom, states that the target t(x) of x and the source s(y) of y are equal whenever the composite of x and y is defined, that is, $x \odot y \neq \emptyset$. We write $\Delta(x,y)$ for $x \odot y \neq \emptyset$ and call Δ the *domain of definition* of \odot .

The third and fourth catoid axioms are left and right unit axioms.

A catoid C is functional if $x, x' \in yz$ imply x = x' for all $y, z \in C$, and local if $t(x) = s(y) \Rightarrow xy \neq \emptyset$ for all $x, y \in C$. A category can then be defined as a local functional catoid.

Catoids thus generalise categories beyond locality and functionality. They formalise categories in single-set style [Mac48], see also [ML98, Chapter XII], and with arrow composition in diagrammatic order. In functional catoids, composition is a partial operation that maps either to singleton sets or to the empty set. Locality imposes the standard composition pattern of arrows in categories: $\Delta(x, y) \Leftrightarrow t(x) = s(y)$.

Composition in a catoid is total if $\Delta = C \times C$. Total multioperations are known as hyperoperations. Total operations are therefore total functional multioperations.

Remark 2.1. In the definition of catoids and throughout this text, we are using "set" indifferently for small sets and classes and ignore the well known foundational issues, which do not arise in our setting. Distinctions can be made as for standard categories. See, for instance, Mac Lane's book [ML98] for a discussion.

Catoids form categories. A catoid morphism $f: C \to D$ between catoids C and D must preserve compositions, sources and targets:

$$f(x \odot_C y) \subseteq f(x) \odot_D f(y), \qquad f \circ s_C = s_D \circ f, \qquad f \circ t_C = t_D \circ f,$$

where, on the left-hand side of the left identity the image of the set $x \odot_C y$ with respect to f is taken. A morphism $f: C \to D$ is bounded if $f(x) \in u \odot_D v$ implies that $x \in y \odot_C z$, u = f(y) and v = f(z) for some $y, z \in C$. Here and henceforth, we write f(X) for the image of the set X under the map f.

Morphisms of categories, as local functional catoids, are functors. The inclusion in the definition of morphisms reflects that $x \odot_C y = \emptyset$ whenever the composition is undefined. Bounded morphisms are widely used in modal and substructural logics. They correspond to functional bisimulations or open maps.

Example 2.2. Bounded morphisms need not satisfy $f(x \odot_C y) = f(x) \odot_D f(y)$. Consider the discrete catoid on $C = \{a, b\}$ with $s = id_C = t$ and

$$xy = \begin{cases} \{x\} & \text{if } x = y, \\ \emptyset & \text{otherwise.} \end{cases}$$

The constant map $f_b: x \mapsto b$ on C is clearly a catoid endomorphism. It is bounded because every $x \in C$ satisfies $f_b(x) \in bb$, $x \in xx$ and $b = f_b(x)$. But $f_b(ab) = \emptyset \neq \{b\} = f_b(a)f_b(b)$.

The *opposite* of a catoid is defined as for categories. It is a structure in which s and t are exchanged and so are the arguments in \odot . The opposite of a (local, functional, total) catoid is again a (local, functional, total) catoid. Properties of catoids translate through this duality.

Properties of catoids and related structures have been collected in [FJSZ23] and our Isabelle theories. Here we list only some that are structurally interesting or needed in proofs below.

Lemma 2.3. In every catoid,

- 1. $s \circ s = s$, $t \circ t = t$, $s \circ t = t$ and $t \circ s = s$,
- 2. $s(x) = x \Leftrightarrow x = t(x)$,
- 3. $s(x)s(x) = \{s(x)\}\ and\ t(x)t(x) = \{t(x)\},\$
- 4. s(x)t(y) = t(y)s(x),
- 5. s(s(x)y) = s(x)s(y) and t(xt(y)) = t(x)t(y).

Note that direct images with respect to s and t are taken in the left-hand sides of the identities in (5). We define the sets of fixpoints of the source and target maps as

$$C_s = \{x \mid s(x) = x\}$$
 and $C_t = \{x \mid t(x) = x\}.$

The following fact implies that x is a unit, $x \in s(C) = t(C)$, if and only if x is a fixpoint of s.

Lemma 2.4. Let C be a catoid. Then $C_s = C_t = s(C) = t(C)$.

It is immediate from Lemma 2.3(1). By analogy with categories, we call units of catoids 0-cells and write C_0 instead of C_s or C_t . Similarly, the elements of a catoid C can be viewed as 1-cells, the elements of C_0 as degenerate 1-cells (s(x) = t(x) holds for all $x \in C_0$ by Lemma 2.3(2)) and those of C_0 as proper or non-degenerate 1-cells. Further, units of catoids can be seen as orthogonal idempotents.

Lemma 2.5. Let C be a catoid. For all $x, y \in C_0$, $\Delta(x, y) \Leftrightarrow x = y$ and

$$xy = \begin{cases} \{x\} & \text{if } x = y, \\ \emptyset & \text{otherwise.} \end{cases}$$

The next two lemmas recall an alternative equational characterisation of locality, which is important for the correspondence with modal quantales in Section 8.

Lemma 2.6. In every catoid,

- 1. $s(xy) \subseteq s(xs(y))$ and $t(xy) \subseteq t(t(x)y)$,
- 2. $\Delta(x,y)$ implies $s(xy) = \{s(x)\}\ and\ t(xy) = \{t(y)\}.$

Item (2) of the following lemma features the equational characterisation of locality mentioned.

Lemma 2.7. Let C be a catoid. The following statements are then equivalent:

- 1. C is local.
- 2. s(xy) = s(xs(y)) and t(xy) = t(t(x)y), for all $x, y \in C$,
- 3. $\Delta(x,y) \Leftrightarrow t(x)s(y) \neq \emptyset$, for all $x,y \in C$.

Many examples of catoids and related structures are listed in [FJSZ23]. Here we mention only a few.

Example 2.8. Categories, as local functional catoids, with (bounded) morphisms and single-set categories à la MacLane [ML98, Chapter XII] with the same morphisms are isomorphic as categories [FJSZ23]. The elements of single-set categories are arrows or 1-cells of categories. Their objects or 0-cells correspond bijectively to identity 1-cells and thus to units of catoids.

Example 2.9. The free category or *path category* generated by any digraph $s,t:E\to V$, for a set E of edges or 1-generators and a set V of vertices or 0-generators, is a forteriori a catoid. Recall that morphisms of path categories are finite paths between vertices, represented as alternating sequences of vertices and edges. Source and target maps extend from edges to paths, and we write $\pi:v\to w$ for a path π with source v and target w. Paths $\pi_1:u\to v$ and $\pi_2:v\to w$ can be composed to the path $vw:u\to w$ by gluing their ends. Identities are constant paths of length zero such as $v:v\to v$. See [ML98] for details.

Digraphs can be modelled in catoid-style as structures (X, s, t) satisfying $s \circ s = s$, $t \circ t = t$, $s \circ t = t$ and $t \circ s = s$. These conditions make $X_s = s(X) = t(X) = X_t$ the set of vertices.

Example 2.10. Let Σ^* be the free monoid generated by the finite set Σ . The *shuffle catoid* $(\Sigma^*, \|, s, t)$ on Σ^* has the total commutative multioperation $\|: \Sigma^* \times \Sigma^* \to \mathcal{P}\Sigma^*$ as its composition. For all $a, b \in \Sigma$ and $v, w \in \Sigma^*$,

$$\varepsilon \|v = \{v\} = v \|\varepsilon \qquad \text{and} \qquad (av)\|(bw) = a(v\|(bw)) \cup b((av)\|w),$$

where ε denotes the empty word. The source and target structure is trivial: $s(w) = \varepsilon = t(w)$ for all $w \in \Sigma^*$. The shuffle catoid is local because of totality and triviality of s and t. It is obviously not functional and hence not a category.

Remark 2.11. Catoids can be seen as multimonoids, that is, multisemigroups with multiple units, which generalise single-set categories with multiple units [ML98, Chapter I] to multioperations. Multimonoid morphisms are then unit-preserving multisemigroup morphisms, which ignore s- and t-preservation. Categories of catoids and multimonoids are isomorphic; the functional relationship between elements of multimonoids and their left and right units determines source and target maps [CDS20, FJSZ23]. Categories of local partial multimonoids are therefore isomorphic to categories. See [CDS20] for the appropriate notion of locality.

Remark 2.12. Multioperations $X \times X \to \mathcal{P}X$ are isomorphic to ternary relations $X \to X \to X \to 2$. Writing R^x_{yz} for $x \in y \odot z$, catoids can be axiomatised alternatively as relational structures (C, R, s, t) that satisfy the relational associativity axiom $\exists v. R^w_{xv} \wedge R^v_{yz} \Leftrightarrow \exists u. R^u_{xy} \wedge R^w_{uz}$, the weak locality axiom $\exists z. R^z_{xy} \Rightarrow t(x) = s(z)$ and the relational unit axioms $R^x_{s(x)x}$ and $R^x_{xt(x)}$. Such relational monoids are, in fact, monoids in the monoidal category Rel with the standard tensor. Functionality and locality translate to relations along the same lines. Relational variants of catoids have been studied in [CDS20]. Morphisms and strong morphisms of ternary relations are standard for modal and substructural logics [Gol89, Giv14]; categories of catoids and relational monoids are once again isomorphic [FJSZ23].

Ternary and more generally n+1-ary relations appear as duals of binary and more generally n-ary modal operators in Jónsson and Tarski's duality theory for boolean algebras with operators [JT51, JT52, Gol89, Giv14]. While multioperational catoids support set-theoretic algebraic reasoning, their relational siblings are more suitable for type-theoretic approaches.

Remark 2.13. A partial operation $\hat{\odot}: \Delta \to C$, where $y \hat{\odot} z$ denotes the unique element that satisfies $y \hat{\odot} z \in y \odot z$ whenever $\Delta(y, z)$, can be defined in any functional catoid. Using this partial operation, $x \in y \odot z$ if and only if $\Delta(y, z)$ and $x = y \hat{\odot} z$. While the adjunction of a zero to a multimonoid, a more conventional approach to partiality, collapses its units to a singleton (the original units survive in weakened form), catoids with zero admit multiple units [CDS20].

3 Groupoids

Groupoids are categories with invertible arrows, see [Bro87] for a survey. Particularly relevant to us is the work of Jónsson and Tarski [JT52] on the correspondence between groupoids and relation algebras, which we

revisit in the slightly different setting of Dedekind quantales. Several properties that feature in this section are theirs. Interestingly, catoids become groupoids and hence categories when the natural axioms for inverses are added

A groupoid is a catoid C with an inversion operation $(-)^-: C \to C$ such that, for all $x \in C$,

$$xx^{-} = \{s(x)\}\$$
 and $x^{-}x = \{t(x)\}.$

To justify this definition, we proceed in two steps to derive locality and functionality, showing selected proofs only. The remaining ones can be found in Appendix B.

Lemma 3.1. Let X be a catoid with operation $(-)^-: C \to C$ that satisfies $s(x) \in xx^-$ and $t(x) \in x^-x$ for all $x \in C$. Then

- 1. C is local.
- 2. $s(x^{-}) = t(x)$ and $t(x^{-}) = s(x)$,
- 3. $xy = \{s(x)\}$ implies $x^- = y$ and $yx = \{t(x)\}$ implies $x^- = y$,
- 4. $s(x)^- = s(x)$ and $t(x)^- = t(x)$.

Proof. For (1), suppose t(x) = s(y). Then $\{x\} = xt(x) = xs(y) = x(yy^-) = (xy)y^-$. Hence $x = uy^-$ and $u \in xy$ hold for some $u \in C$. Thus $\Delta(x,y)$ and C is local.

Lemma 3.2. In every groupoid,

- 1. $(x^{-})^{-} = x$,
- 2. $x \in yz \Leftrightarrow y \in xz^- \Leftrightarrow z \in y^-x$.

Proposition 3.3. Every groupoid is a category.

Proof. Relative to Lemma 3.1(1) it remains to check functionality. Suppose $x, x' \in yz$. Then $z \in y^-x$ by Lemma 3.2(2) and $x' \in yy^-x = s(y)x = s(x)x = \{x\}$ using the first assumption and $v \in xy \Rightarrow s(v) = s(x)$, which holds in any catoid by Lemma 2.7, in the second step. Thus x' = x.

Lemma 3.4. The following cancellation properties hold in every groupoid:

- 1. s(x) = t(z) = s(y) and zx = zy imply x = y,
- 2. t(x) = s(z) = t(y) and xz = yz imply x = y.

These properties correspond to the fact that every morphism in a groupoid is both epi and mono.

Lemma 3.5. In every groupoid,

- 1. $x \odot x^- \odot x = x$,
- 2. t(x) = s(y) implies $x^- \odot x = y^- \odot y$,
- 3. $(x \hat{\odot} y)^- = y^- \hat{\odot} x^-$.

The last identity follows from Lemma 3.2(2).

Example 3.6. In the pair groupoid $X \times X$ over the set X, s((a,b)) = (a,a), t((a,b)) = (b,b) and $(a,b)^- = (b,a)$ hold for every $(a,b) \in X \times X$. For all $(a,b),(c,d) \in X \times X$, composition is defined as

$$(a,b)\odot(c,d)= \begin{cases} \{(a,d)\} & \text{if } b=c,\\ \emptyset & \text{otherwise.} \end{cases}$$

Note that \odot is formalised as a multioperation. Pair groupoids give rise to algebras of binary relations, see Example 5.15 below.

Example 3.7. The free groupoid generated by a digraph is obtained by adding formal inverses to each edge and considering paths up to simplification. For instance, if a, b, c, d are appropriate edges, then $abc^{-1}cb^{-1}d = ad$, see also [Hig71, Chapter 4].

4 Modal Quantales

Recall that a quantale $(Q, \leq, \cdot, 1)$ is a complete lattice (Q, \leq) with an associative composition \cdot , which preserves all sups in both arguments, and has a unit 1. See [Ros90] for an introduction. We write \bigvee , \bigvee , \bigwedge and \bigwedge for sups, binary sups, infs and binary infs in a quantale, and \bot and \top for the smallest and greatest element of Q, respectively. A subidentity of Q is an element $\alpha \in Q$ such that $\alpha \leq 1$.

A quantale is *distributive* if its underlying lattice is distributive and *boolean* if its underlying lattice is a boolean algebra; in that case, we write – for boolean complementation.

We henceforth use greek letters for elements of quantales to distinguish them from elements of catoids, and we often write $\alpha\beta$ for $\alpha \cdot \beta$.

Example 4.1. We need the boolean quantale of booleans 2 with carrier set $\{0,1\}$, order 0 < 1, max as binary sup, min as binary inf and composition, 1 as its unit and $\lambda x.1 - x$ as complementation.

A Kleene star $(-)^*: Q \to Q$ can be defined on any quantale Q, for $\alpha^0 = 1$ and $\alpha^{i+1} = \alpha \alpha^i$, as

$$\alpha^* = \bigvee_{i \ge 0} \alpha^i.$$

A domain quantale [FJSZ22a] is a quantale Q with an operation $dom: Q \to Q$ such that, for all $\alpha, \beta \in Q$,

$$\alpha \leq dom(\alpha)\alpha, \qquad dom(\alpha dom(\beta)) = dom(\alpha\beta), \qquad dom(\alpha) \leq 1,$$

$$dom(\bot) = \bot, \qquad dom(\alpha \lor \beta) = dom(\alpha) \lor dom(\beta).$$

These domain axioms are known from domain semirings [DS11], see also Section 12. We call the first axiom the absorption axiom. The second expresses locality of domain. The third is the subidentity axiom, the fourth the bottom axiom and the final the (binary) sup axiom. Most properties of interest translate from domain semirings to domain quantales. An equational absorption law $dom(\alpha)\alpha = \alpha$ is derivable.

We define $Q_{dom} = \{\alpha \in Q \mid dom(\alpha) = \alpha\}$. As for catoids, $Q_{dom} = dom(Q)$ holds because $dom \circ dom = dom$ is derivable from the domain quantale axioms. Moreover, (Q_{dom}, \leq) forms a bounded distributive lattice with + as binary sup and \cdot as binary inf. We call Q_{dom} the lattice of domain elements or simply domain algebra. In a boolean quantale, Q_{dom} is the set of all subidentities and hence a complete boolean algebra.

Quantales are closed under opposition, which exchanges the arguments in compositions. A codomain quantale (Q, cod) is then a domain quantale (Q^{op}, dom) . Further, a modal quantale [FJSZ22a] is a domain and codomain quantale $(Q, \leq, \cdot, 1, dom, cod)$ that satisfies the compatibility axioms

$$dom \circ cod = cod$$
 and $cod \circ dom = dom$.

These guarantee that $Q_{dom} = Q_{cod}$, a set which we denote by Q_0 as for catoids.

Example 4.2. Let (C, \odot, s, t) be a local catoid. Then $\mathcal{P}C$ forms a modal catoid quantale over C. Its monoidal structure is given by $\odot: \mathcal{P}C \times \mathcal{P}C \to \mathcal{P}C$, the lifted composition of the catoid, and by C_0 ,. Its modal structure is given by dom(X) = s(X) and cod(X) = t(X), the images of any set $X \subseteq C$ with respect to s and t [FJSZ22a]. As a powerset quantale, $\mathcal{P}C$ is in fact an atomic boolean quantale. It is important to note that locality of C is needed for locality of $\mathcal{P}C$. This result has the following instances, among others.

- 1. Every category C lifts to a modal category quantale on PC.
- 2. The path category over any digraph, more specifically, lifts to a *modal path quantale* at powerset level. Domain and codomain elements are sets of vertices.
- 3. Every pair groupoid lifts to a modal relation quantale at powerset level with the standard relational domain and codomain maps $dom(R) = \{(x, x) \mid \exists y. \ (x, y) \in R\}$ and $cod(R) = \{(y, y) \mid \exists x. \ (x, y) \in R\}$, and with \odot lifted to relational composition. The quantalic unit is the identity relation.

4. The shuffle catoid on Σ^* lifts to the commutative shuffle language quantale on Σ . The quantalic composition is the shuffle product of languages, which are subsets of Σ^* . Its monoidal unit is $\{\varepsilon\}$, the modal structure is therefore trivial.

For further examples of modal quantales and their underlying catoids see Section 8 and [FJSZ22a].

Remark 4.3. Locality of catoids in the form $x \odot y \neq \emptyset \Leftrightarrow t(x)s(y) \neq \emptyset$, as in Lemma 2.7, corresponds to

$$\alpha \cdot \beta \neq \bot \Leftrightarrow cod(\alpha) \cdot dom(\beta) \neq \bot$$

in modal quantales. This is a consequence of locality of dom and cod in modal semirings [DS11] and modal quantales. In modal quantales, it is even equivalent to locality of dom and cod. Yet the more precise locality property $t(x)s(y) \neq \emptyset \Leftrightarrow t(x) = s(y)$ does not hold in all modal quantales.

Consider for instance the path category over the digraph $v_1 \stackrel{e_1}{\longleftarrow} v_2 \stackrel{e_2}{\longrightarrow} v_3 \stackrel{e_3}{\longrightarrow} v_4$ and the sets of paths $X = \{(v_2, e_1, v_1), (v_2, e_2, v_3)\}$ and $Y = \{(v_3, e_3, v_4)\}$. Then $cod(X) = \{v_1, v_3\} \neq \{v_3\} = dom(Y)$ whereas $X \cdot Y = \{(v_2, e_2, v_3, e_3, v_4)\} \neq \emptyset$.

Nevertheless, for any local catoid C with elements a and b,

$$cod(\{a\}) \cap dom(\{b\}) \neq \emptyset \Leftrightarrow \{t(a)\} \cap \{s(b)\} \neq \emptyset \Leftrightarrow \{t(a)\} = \{s(b)\} \Leftrightarrow cod(\{a\}) = dom(\{b\})$$

holds at least for the atoms $\{a\}$ and $\{b\}$ in the catoid quantale $\mathcal{P}C$.

5 Dedekind Quantales

Jónsson and Tarski have lifted groupoids to relation algebras [JT52]. To adapt these results to quantales, we consider Dedekind quantales and discuss their relationship with modal quantales. Dedekind quantales are single-object versions of the modular quantaloids studied by Rosenthal [Ros96]. We start from a weak form of converse. All proofs in this section can again be found in Appendix B and our Isabelle theories. Isabelle has also been instrumental in finding counterexamples in this section.

An involutive quantale [MWP92] is a quantale Q with an operation $(-)^{\circ}: Q \to Q$ that satisfies

$$\alpha^{\circ\circ} = \alpha, \qquad (\bigvee A)^{\circ} = \bigvee \{\alpha^{\circ} \mid \alpha \in A\}, \qquad (\alpha\beta)^{\circ} = \beta^{\circ}\alpha^{\circ}.$$

Involution makes quantalic opposition explicit.

Remark 5.1. Replacing the first two axioms by $\alpha^{\circ} \leq \beta \Leftrightarrow \alpha \leq \beta^{\circ}$ [DBvdW97] yields an equivalent axiomatisation. Involution is therefore self-adjoint.

Lemma 5.2. In every involutive quantale,

- 1. $\alpha \leq \beta \Rightarrow \alpha^{\circ} \leq \beta^{\circ}$,
- 2. $(\alpha \vee \beta)^{\circ} = \alpha^{\circ} \vee \beta^{\circ}$,
- 3. $(\bigwedge A)^{\circ} = \bigwedge \{\alpha^{\circ} \mid \alpha \in A\} \text{ and } (\alpha \wedge \beta)^{\circ} = \alpha^{\circ} \wedge \beta^{\circ},$
- 4. $\bot^{\circ} = \bot$, $1^{\circ} = 1$ and $\top^{\circ} = \top$,
- 5. $\alpha^{\circ} \wedge \beta = \bot \Leftrightarrow \alpha \wedge \beta^{\circ} = \bot$,
- 6. $\alpha^{*\circ} = \alpha^{\circ*}$.

The next lemma separates involutive quantales from relation algebras over complete lattices.

Lemma 5.3. Each of the following properties fails in some involutive quantale:

1. the strong Gelfand property $\alpha < \alpha \alpha^{\circ} \alpha$,

- 2. the modular law $\alpha\beta \wedge \gamma \leq (\alpha \wedge \gamma\beta^{\circ})\beta$,
- 3. the Dedekind law $\alpha\beta \wedge \gamma \leq (\alpha \wedge \gamma\beta^{\circ})(\beta \wedge \alpha^{\circ}\gamma)$.

Proof. In the involutive quantale $\bot < a < \top = 1$ with multiplication $aa = \bot$ (the rest being fixed) and $(-)^{\circ} = id$, the strong Gelfand property fails because $aa^{\circ}a = \bot < a$. The modular and Dedekind law fail because $(1 \land aa)a(1 \land aa)(a \land 1a) = 0 < 1a \land a = a$.

The strong Gelfand property (the name has been borrowed from [PR11]) has been used by Ésik can co-workers to axiomatise relational converse in semigroups and Kleene algebras, where infs are not available [BÉS95, ÉB95]. Similarly, and for the same reason, it has been used in globular n-Kleene algebras [CGMS22] and applications in higher rewriting. This is the main reason why we revisit this definition; see also Section 12.

Example 5.4. Adding the strong Gelfand property to the involutive quantale axioms implies neither the modular law nor the Dedekind law. The involutive quantale defined by

and $(-)^{\circ} = id$ satisfies the strong Gelfand property, but $1a \wedge \top = a > \bot = (1 \wedge \top a)a$.

Lemma 5.5. In every involutive quantale, the modular law holds if and only if the Dedekind law does.

A Dedekind quantale is an involutive quantale in which the Dedekind law (or the modular law) holds.

Lemma 5.6. The following properties hold in every Dedekind quantale:

- 1. the strong Gelfand property,
- 2. Peirce's law $\alpha\beta \wedge \gamma^{\circ} = \bot \Leftrightarrow \beta\gamma \wedge \alpha^{\circ} = \bot$,
- 3. the Schröder laws $\alpha\beta \wedge \gamma = \bot \Leftrightarrow \beta \wedge \alpha^{\circ}\gamma = \bot \Leftrightarrow \alpha \wedge \gamma\beta^{\circ} = \bot$.

Dedekind quantales bring us close to relation algebras [Tar41, HH02, Mad06], where domain and codomain operations can be defined explicitly as

$$dom(\alpha) = 1 \wedge \alpha\alpha^{\circ}$$
 and $cod(\alpha) = 1 \wedge \alpha^{\circ}\alpha$,

or alternatively as $dom(\alpha) = 1 \land \alpha \top$ and $cod(\alpha) = 1 \land \top \alpha$. In Dedekind quantales, we obtain the following properties.

Lemma 5.7. In every Dedekind quantale, $dom(\alpha) = 1 \wedge \alpha\alpha^{\circ}$ and $cod(\alpha) = 1 \wedge \alpha^{\circ}\alpha$ imply

- 1. $dom(\alpha) = 1 \wedge \alpha \top$,
- $2. dom(\alpha) < 1$,
- 3. $dom(\alpha)\alpha = \alpha$,
- 4. $dom(\alpha) \top = \alpha \top$,
- 5. $dom(\bot) = \bot$,
- 6. $\alpha \leq \beta \Rightarrow dom(\alpha) \leq dom(\beta)$,

- 7. $(dom(\alpha))^{\circ} = dom(\alpha)$,
- 8. $dom(\alpha dom(\beta)) = dom(\alpha\beta)$,
- 9. $dom(\alpha) \vee dom(\beta) = dom(\alpha \vee \beta)$,
- 10. $dom(cod(\alpha)) = cod(\alpha)$ and $cod(dom(\alpha)) = dom(\alpha)$,
- 11. $dom(\alpha^{\circ}) = cod(\alpha)$.

All domain quantale axioms are therefore derivable.

Proposition 5.8. Every Dedekind quantale is a modal quantale with $dom(\alpha) = 1 \wedge \alpha \alpha^{\circ}$ and $cod(\alpha) = 1 \wedge \alpha^{\circ} \alpha$.

Next we consider *modal Dedekind quantales*, Dedekind quantales that are also modal quantales, and ask about the relationship with the explicit relation-algebraic definition of *dom* and *cod*.

Lemma 5.9. In every modal Dedekind quantale, $dom(\alpha)^{\circ} = dom(\alpha)$ and $cod(\alpha)^{\circ} = cod(\alpha)$.

Dedekind quantales can admit several domain and codomain structures that satisfy the domain axioms, even in the quantales are distributive:

Lemma 5.10. There is a modal distributive Dedekind quantale in which $dom(\alpha) \neq 1 \wedge \alpha\alpha^{\circ}$ and $cod(\alpha) \neq 1 \wedge \alpha^{\circ}\alpha$.

Proof. In the modal distributive Dedekind quantale with $\bot < a < \top = 1$, multiplication aa = a and dom(a) = 1 = cod(a), we have $dom(a) = cod(a) = \top \neq a = 1 \land aa^{\circ} = 1 \land a^{\circ}a$.

This is related to the fact that the domain and codomain operations of modal semirings, even finite ones, need not not be uniquely determined [DS11]. (Finite modal semirings and finite modal quantales are the same structures). They are uniquely determined in the boolean case, however:

Remark 5.11. In any boolean modal quantale Q, $Q_0 = Q_{dom} = Q_{cod}$ is equal to the boolean subalgebra of all subidentities, and domain and codomain are uniquely defined. A proof for modal semirings [DS11] translates directly. Therefore, in any boolean modal Dedekind quantale, $dom(\alpha) = 1 \wedge \alpha\alpha^{\circ}$ and $cod(\alpha) = 1 \wedge \alpha^{\circ}\alpha$.

Remark 5.12. The law $dom(\alpha) = 1 \land \alpha \top$ fails in the modal path quantales from Example 4.2. In this model, 1 is the set V of all vertices of the digraph [FJSZ23]. Thus $V \cap P \top = \emptyset$ unless the set P of paths contains a path of length one and $d(P) = \emptyset$ if and only if $P = \emptyset$. Yet note that this identity is derivable in any Dedekind quantale (Lemma 5.7(1)).

Lemma 5.13. In any boolean Dedekind quantale, $(-\alpha)^{\circ} = -(\alpha^{\circ})$ and the following residual law is derivable:

$$\alpha^{\circ} - (\alpha \beta) \leq -\beta.$$

Such a law appears in Tarski's original axiomatisation of relation algebra [Tar41]. A boolean Dedekind quantale is thus a relation algebra over a complete lattice, and a relation algebra a boolean Dedekind quantale in which only finitary sups and infs are required to exist. Relation algebras are formed over boolean algebras that need not be complete.

Remark 5.14. The conditions $dom(\alpha) \leq \alpha\alpha^{\circ}$ and $cod(\alpha) \leq \alpha^{\circ}\alpha$ have been used for higher rewriting with globular 2-Kleene algebras [CGM21], see also Section 12. They are consequences of the explicit definition of domain and codomain in Dedekind quantales. In involutive modal quantales, each of them implies the strong Gelfand property, $dom(\alpha) \leq \alpha\alpha^{\circ}$ is equivalent to $dom(\alpha) = 1 \wedge \alpha\alpha^{\circ}$ and $cod(\alpha) = 1 \wedge \alpha^{\circ}\alpha$ is equivalent to $cod(\alpha) \leq \alpha^{\circ}\alpha$. Yet none of these laws need to hold in such quantales: in the modal distributive Dedekind quantale used in the proof of Lemma 5.10, $dom(a) = cod(a) = \top > a = aa^{\circ} = a^{\circ}a$. Finally, a modal

involutive quantale satisfying $dom(\alpha) = 1 \wedge \alpha\alpha^{\circ}$ and $cod(\alpha) = 1 \wedge \alpha^{\circ}\alpha$ need not be Dedekind: the modal involutive quantale with

dom(a) = 1 = cod(a) (the rest being fixed due to order-preservation and (co)domain axioms) and $(-)^{\circ} = id$ satisfies $dom(\alpha) = 1 \wedge \alpha\alpha^{\circ}$ and $cod(\alpha) = 1 \wedge \alpha^{\circ}\alpha$, but not the modular law:

$$ab \wedge b = \top \wedge b = b > (a \wedge bb^{\circ})b = (a \wedge b)b = \bot b = \bot.$$

Example 5.15. Let G be a groupoid. Then $\mathcal{P}G$ forms a relation algebra over G [JT52]. As a powerset algebra, the underlying lattice of $\mathcal{P}G$ is complete (even boolean and atomic). Hence every groupoid lifts to a Dedekind quantale, and in fact a modal Dedekind quantale in which $dom(X) = G_0 \cap XX^\circ$ and $cod(X) = G_0 \cap X^\circ X$. Again there are interesting instances.

- 1. Each free groupoid generated by some digraph lifts to a *modal path Dedekind quantale*, where converses are sets of formal inverses (formal inverses were not assumed in the path quantale in Remark 5.12).
- 2. The pair groupoid on X lifts to the modal relation Dedekind quantale on $\mathcal{P}(X \times X)$ with standard relational converse lifted from the inverse operation on the pair groupoid.

6 Higher catoids

Mac Lane has outlined axiomatisations of single-set 2-categories, n-categories and ω -categories, imposing a 2-category structure on each pair of single-set categories C_i and C_j for $0 \le i < j < \omega$ [ML98, Chapter XII]. Here, ω -category means strict globular ∞ -category. Similar single-set approaches appear, for instance, in [BH81, Str87, Ste04]. We adapt MacLane's axioms to catoids. We start from a uniform axiomatisation that includes the case of n or ω , but then focus mainly on ω -catoids.

For an ordinal $\alpha \in \{0, 1, \dots, \omega\}$, an α -catoid is a structure $(C, \odot_i, s_i, t_i)_{0 \le i < \alpha}$ such that each (C, \odot_i, s_i, t_i) is a catoid and these structures interact as follows:

• for all $i \neq j$,

$$s_i \circ s_j = s_j \circ s_i,$$
 $s_i \circ t_j = t_j \circ s_i,$ $t_i \circ t_j = t_j \circ t_i,$ $s_i(x \odot_j y) \subseteq s_i(x) \odot_j s_i(y),$ $t_i(x \odot_j y) \subseteq t_i(x) \odot_j t_i(y),$

• and for all i < j,

$$(w \odot_j x) \odot_i (y \odot_j z) \subseteq (w \odot_i y) \odot_j (x \odot_i z),$$

$$s_j \circ s_i = s_i, \qquad s_j \circ t_i = t_i, \qquad t_j \circ s_i = s_i, \qquad t_j \circ t_i = t_i.$$

An α -category is a local functional α -catoid, that is, each (C, \odot_i, s_i, t_i) is local and functional.

As the (C, \odot_i, s_i, t_i) are catoids, $C_{s_i} = C_{t_i} = s_i(C) = t_i(C)$ for each $i < \alpha$ by Lemma 2.4. We write C_i for C_{s_i} and call this set the set of *i-cells* of C. We also write Δ_i for the domain of definition of \odot_i and refer to the source and target cells of cells as (lower and upper) faces.

The axioms in the first item impose that source and target maps at each dimension i are catoid (endo)morphisms of the catoid (C, \odot_j, s_j, t_j) at each dimension $j \neq i$. In the local functional case of α -categories, these morphisms become functors, as expected. (In the ω -category of globular sets, the axioms in the first line are known as globular laws.)

The *interchange* axioms in the third line ensure that \odot_i is a catoid bi-morphism with respect to \odot_j , for all i < j.

The whisker axioms in the fourth line, together with the catoid laws in Lemma 2.3(1), imply that for $x \in C_i$, we have $s_j(x) = t_j(x) = x$, and thus $x \in C_j$, for all $j \ge i$. Lower dimensional cells thus remain units in higher dimensions and

$$C_0 \subseteq C_1 \subseteq C_2 \subseteq \cdots \subseteq C$$
.

We may thus regard *i*-cells as degenerate cells or whiskers in which sources and targets at each dimension greater than i-1 coincide. Further, Lemma 2.5 implies that all lower dimensional cells are orthogonal idempotents with respect to higher compositions. For all $i, j \leq k$,

$$s_i(x) \odot_k s_j(y) = \begin{cases} \{s_i(x)\} & \text{if } s_i(x) = s_j(y), \\ \emptyset & \text{otherwise.} \end{cases}$$

All higher dimensional compositions of lower dimensional cells are therefore trivial. The structure of units across dimensions can be seen in Example 6.6 below.

We now turn to ω -catoids. In this context, one often adds an axiom guaranteeing that for all $x \in C$ there exists and $i \leq \omega$ such that $s_i(x) = x = t_i(x)$ [Ste04]. It is needed in the following constructions, but not beyond. Because of the whisker axioms and the catoid axioms, the chain $C_0 \hookrightarrow C_1 \hookrightarrow C_2 \hookrightarrow \cdots \hookrightarrow C$ is in fact a filtration of ω -catoids (when $\alpha = \omega$). Catoid C is the (co)limit of an increasing chain of sub- ω -catoids C_n , which themselves are n-catoids. See [Ste04, Proposition 2.3] for a related discussion on ω -categories.

 ω -Catoids, and more generally α -catoids, form categories. Their morphisms are catoid morphisms that preserve all source and target maps s_i and t_i and all compositions \odot_i at each dimension $i < \omega$. These kind of categories are also categories with respect to bounded morphisms.

As in the one-dimensional case, ω -catoids and n-catoids generalise ω -categories and n-categories to multioperational compositions, and beyond functionality and locality. Yet the underlying ω -graphs or n-graphs, where compositions are forgotten, are the same for both. The globular cell shape of strict higher categories is therefore present in the corresponding catoids, too.

Lemma 6.1. In every ω -catoid, the following globular laws hold. For all $0 \le i < j < \omega$,

$$s_i \circ s_j = s_i, \qquad s_i \circ t_j = s_i, \qquad t_i \circ t_j = t_i, \qquad t_i \circ s_j = t_i.$$

Example 6.2. The globular cell shape of ω -catoids can be visualised, in two dimensions, as

$$s_0(x) \underbrace{\int_{t_1(x)}^{s_1(x)} t_0(x)}_{t_1(x)}$$

The relationships between 0-cells and 1-cells in this diagram can be calculated using Lemma 6.1. See [ML98, p280] for a detailed explanation for 2-categories, as well as a discussion of the category **Cat** of all small categories, the prototypical example of a 2-category.

In light of Example 2.2, the morphism laws of ω -categories are unusually strong.

Lemma 6.3. In every ω -category, for $0 \le i < j < \omega$, the following strong morphism laws hold:

$$s_j(x \odot_i y) = s_j(x) \odot_i s_j(y)$$
 and $t_j(x \odot_i y) = t_j(x) \odot_i t_j(y)$.

Proof. First we derive the law for s_j . Suppose the right-hand side is \emptyset . Then $s_j(x \odot_i y) = s_j(x) \odot_i s_j(y)$, using the morphism axiom $s_j(x \odot_i y) \subseteq s_j(x) \odot_i s_i(y)$. Otherwise, if $\Delta_i(s_j(x), s_j(y))$, then $t_i(s_j(x)) = s_i(s_j(x))$ and therefore $t_i(x) = s_i(y)$ by Lemma 6.1. Locality then implies that $\Delta_i(x, y)$, and $s_j(x \odot_i y) = s_j(x) \odot_i s_j(y)$ follows from the morphism axiom for s_j and functionality. The proof for t_j follows by opposition.

By contrast to the other morphism axioms, which require that if left-hand sides are defined, then so are right-hand sides, the strong morphism laws state that one side is defined if and only if the other is. Accordingly, an ω -catoid is strong if it satisfies the strong morphism laws.

Corollary 6.4. In every ω -category, $\Delta_i(x,y) \Leftrightarrow \Delta_i(s_j(x),s_j(y)) \Leftrightarrow \Delta_i(t_j(x),t_j(y))$ for all $0 \leq i < j < \omega$.

Example 6.5. The strong morphism laws of ω -categories can be explained diagrammatically for 2-categories and in particular in \mathbf{Cat} , where \odot_1 is the vertical composition of 2-cells or natural transformations, and \odot_0 the horizontal one. Lemma 6.3, Corollary 6.4 and locality imply that the horizontal composition $x \odot_0 y$ is defined if and only if the horizontal compositions $s_1(x) \odot_0 s_1(y)$ and $t_1(x) \odot_0 t_1(y)$ are defined. In all cases, this means that $t_0(x) = s_0(y)$.

$$s_0(x) \underbrace{ \int_{x}^{s_1(x)} t_0(x) }_{t_1(x)} \underbrace{ \int_{y}^{s_1(y)} t_0(y) }_{t_1(y)}$$


The morphism axioms $s_1(x \odot_0 y) \subseteq s_1(x) \odot_0 s_1(y)$ and $t_1(x \odot_0 y) \subseteq t_1(x) \odot_0 t_1(y)$ then become strong: the compositions $x \odot_0 y$, $s_1(x) \odot_0 s_1(y)$ and $t_1(x) \odot_0 t_1(y)$ are functional, each yields at most a single cell. The upper and lower faces $s_1(x \odot_0 y)$ and $t_1(x \odot_0 y)$ of $x \odot_0 y$ must therefore be equal to $s_1(x) \odot_0 s_1(y)$ and $t_1(x) \odot_0 t_1(y)$, respectively.

The next two examples explain the weak morphism laws in the absence of locality or functionality.

Example 6.6. In the local 2-catoid on the set $\{a, b, c\}$ with

	s_0					a	b	c		\odot_1	a	b	c
	b					$\{a,b\}$	<i>{a}</i>	{c}	<u>-</u> '		<i>{a}</i>		
b	b	b	b	b	b	$\{a\}$	$\{b\}$	$\{c\}$		b	Ø	$\{b\}$	Ø
c	b	b	a	a	c	$\{c\}$	$\{c\}$	$\{c\}$		c	$\{c\}$	Ø	$\{c\}$

 $s_1(a \odot_0 c) = s_1(\{c\}) = \{a\} \subset \{a,b\} = \{a\} \odot_0 \{a\} = s_1(a) \odot_0 s_1(a)$ and the same inequality holds for t_1 because $s_1 = t_1$. In this example, c is a proper 2-cell with 1-faces a and 0-faces b, a a proper 1-cell with 0-faces b and b is a 0-cell. So the strong morphism laws fail because \odot_0 may map to more than one cell. This highly degenerate situation can be depicted as



This 2-catoid is functional with respect to \bigcirc_1 , but not with respect to \bigcirc_0 .

It is also worth considering the unit structure given by the source and target maps and the whisker axioms, and its effect on \odot_0 and \odot_1 . They determine all compositions except $a \odot_0 a$, $c \odot_0 c$ and $c \odot_1 c$. The composition \odot_1 is trivial because of the whisker axioms.

Example 6.7. In the functional 2-catoid on the set $\{a, b\}$ with


 $s_1(a\odot_0 a)=s_1(\emptyset)=\emptyset\subset\{b\}=\{b\}\odot_0\{b\}=s_1(a)\odot s_1(a)$. The corresponding inequality holds for t_1 because $s_1=t_1$. This example is a monoid as a category with respect to \odot_1 , and a "broken monoid", hence simply a graph, with respect to \odot_0 , as $a\odot_0 a$ is undefined. Now, the strong morphism laws fail because the broken monoid is not local: $s_0(a)=b=t_0(a)$, but $a\odot_0 a=\emptyset$. So $x\odot_0 y$ and therefore $s_1(x\odot_0 y)$ may be \emptyset , whereas $s_1(x)\odot_0 s_1(y)$ or $t_1(x)\odot_0 t_1(y)$ are not.

Next we explain the weakness of the remaining morphism axioms. First, we need a lemma.

Lemma 6.8. In every ω -catoid, for all $0 \le i < j < \omega$, if $\Delta_j(x,y)$, then


$$s_i(x \odot_j y) = \{s_i(x)\} = \{s_i(y)\}$$
 and $t_i(x \odot_j y) = \{t_i(x)\} = \{t_i(y)\}.$

Example 6.9. In 2-catoids, for instance, weakness of $s_0(x \odot_1 y) \subseteq s_0(x) \odot_1 s_0(y)$ and $t_0(x \odot_1 y) \subseteq t_0(x) \odot_1 t_0(y)$ allows $x \odot_1 y$ to be undefined while $s_0(x) \odot_1 s_0(y)$ or $t_0(x) \odot_1 t_0(y)$ are defined. The last two compositions are defined if $s_0(x)$ is equal to $s_0(y)$, and $t_0(x)$ is equal to $t_0(y)$, respectively. For the first one, $t_1(x)$ must be equal to $s_1(y)$, from which $s_0(x) = s_0(t_1(x)) = s_0(s_1(y)) = s_0(y)$ and $t_0(x) = t_0(t_1(x)) = t_0(s_1(y)) = t_0(y)$ follow using the globular laws in Lemma 6.1. The left diagram below shows a situation where $s_0(x \odot_1 y) \subset s_0(x) \odot_1 s_0(y)$ because $s_0(x) = s_0(y)$ whereas $t_1(x) \neq s_1(y)$. The right one shows the opposite situation were $t_0(x \odot_1 y) \subset t_0(x) \odot_1 t_0(y)$ because $t_0(x) = t_0(y)$ whereas $t_1(x) \neq s_1(y)$. The middle diagram shows a situation where both sides are defined because $t_1(x) = s_1(y)$. The globular structure is imposed by Lemma 6.8.



The weak morphism axioms are thus consistent with vertical compositions of globes.

Example 6.10. In 2-catoids, weakness of $(w \odot_1 x) \odot_0 (y \odot_1 z) \subseteq (w \odot_0 y) \odot_1 (x \odot_0 z)$ allows the right-hand side of the interchange axiom to be defined while the left-hand side is undefined. The right-hand side is defined if $t_1(w \odot_0 y)$ is equal to $s_1(x \odot_0 z)$ and both sets are nonempty. Then, using the globular laws in Lemma 6.1 and Lemma 2.6, $s_0(w) = s_0(w \odot_0 y) = s_0(t_1(w \odot_0 y)) = s_0(s_1(x \odot_0 z)) = s_0(x \odot_0 z) = s_0(x)$ and $t_0(y) = t_0(z)$ for similar reasons. The left-hand side is defined if $t_0(w \odot_1 x)$ is equal to $s_0(y \odot_1 z)$ and both sets are nonempty. Therefore $s_0(w) = s_0(w \odot_1 x) = s_0(x)$ and $t_0(y) = t_0(y \odot_1 z) = t_0(z)$, but also $t_0(w) = t_0(w \odot_1 x) = t_0(x)$ and $s_0(y) = s_0(y \odot_1 z) = s_0(y)$ by Lemma 6.8. Thus in particular, $t_0(w) = t_0(x) = s_0(y) = s_0(z)$. The right diagram below shows a situation where $(w \odot_1 x) \odot_0 (y \odot_1 z) \subset (w \odot_0 y) \odot_1 (x \odot_0 z)$, because all compositions on the right are defined, but $t_0(w) = s_0(y) \neq t_0(x) = s_0(z)$. The left diagram below shows a situation where both sides are defined.



The interchange axiom is thus consistent with horizontal and vertical compositions of globes. The difference to the standard equational interchange laws of category theory is simply that, using multioperations, we express partiality by mapping to the empty set. \Box

The following example confirms that the interchange axiom and the morphism axioms for s_0 and t_0 remain inclusions in ω -categories.

Example 6.11. Consider the 2-category with $X = \{a, b\}$ and

	s_0					a				b
\overline{a}	b	b	a	a	\overline{a}	{ <i>b</i> }	<i>{a}</i>	\overline{a}	{ <i>a</i> }	Ø
b	b	b	b	b	b	{a}	$\{b\}$	b	Ø	$\{b\}$

It is actually a monoid as a category with 1-cell a, 0-cell b, and composition $a \odot_0 a = b$, where b is seen as a unit arrow. Further, \odot_1 is trivial because of the whisker axioms. Because of this, $(b \odot_1 a) \odot_0 (b \odot_1 a) = \emptyset \subset \{b\} = (b \odot_0 b) \odot_1 (a \odot_0 a)$ as well as $s_0(a \odot_1 b) = \emptyset \subset \{b\} = s_0(a) \odot_1 s_0(b)$ and $t_0(a \odot_1 b) = \emptyset \subset \{b\} = t_0(a) \odot_1 t_0(b)$. \square

Remark 6.12. The inclusions in the homomorphism axioms for s_i and t_i cannot be strengthened to equations. We show in Appendix A that this would collapse the entire structure. A similar collapse happens with an equational interchange law.

The α -catoid axioms contain redundancy. We have used Isabelle's SAT-solvers and automated theorem provers for an analysis. For irredundancy of a formula φ with respect to a set X of formulas, we ask the SAT-solvers for a model of $X \cup \{\neg \varphi\}$. For redundancy, we ask the automated theorem provers for a proof of $X \vdash \varphi$. This proofs-and-refutations game often succeeds in practice. Because of the set-up of ω -catoids as pairs of 2-catoids, an analysis of 2-catoids suffices.

Proposition 6.13. The following α -catoid axioms are irredundant and imply the other axioms shown above. For all $0 \le i < j \le \alpha$,

$$s_{j}(x \odot_{i} y) \subseteq s_{j}(x) \odot_{i} s_{j}(y), \qquad t_{j}(x \odot_{i} y) \subseteq t_{j}(x) \odot_{i} t_{j}(y),$$
$$(w \odot_{i} x) \odot_{i} (y \odot_{i} z) \subseteq (w \odot_{i} y) \odot_{i} (x \odot_{i} z).$$

This reduction is convenient for relating structures, and it streamlines our correspondence proofs below. More generally, the single-set approach makes n-categories accessible to SMT-solvers and first-order automated theorem provers, using Δ and $\hat{\odot}$ in specifications like in Section 2.

Remark 6.14. We cannot replace the morphism laws for s_j and t_j by those for s_i and t_i in the reduced axiomatisation for α -catoids. Otherwise the first morphism laws would no longer be derivable: Isabelle produces counterexamples. Likewise, if we use only the whisker axioms and the morphism axioms in the first line of the non-reduced axiomatisation that commute source and target maps, we can neither derive the interchange axiom nor the morphism axioms for s_1 and t_1 in the special case of 2-categories. Isabelle produces once again counterexamples. Coincidentally, the morphism laws for s_0 and t_0 are derivable; proofs can be found in our Isabelle theories.

Example 6.15. Let $(\Sigma^*, \odot_0, \varepsilon)$ denote the free monoid generated by the finite alphabet Σ , with word concatenation \odot_0 modelled as a multirelation and ε . It can be viewed as a category with $s_0(w) = \varepsilon = t_0(w)$ for all $w \in \Sigma^*$. Further, let $(\Sigma^* \odot_1, \varepsilon)$ be the shuffle multimonoid on Σ^* from Example 2.10 with $\odot_1 = \|.\|$ Then $(\Sigma^*, \odot_0, \odot_1, \{\varepsilon\})$ forms a *shuffle 2-catoid*. It has been modelled previously as a *relational interchange monoid* in [CDS21] based on ordered relational monoids. It has one single 0-cell, $\{\varepsilon\}$, which is also the only 1-cell. The source/target structure is therefore trivial, but an interchange law between word concatenation and word shuffle holds.

We are currently lacking examples of proper 2-catoids or ω -catoids that not only satisfy the interchange law, but also have a non-trivial source and target structure. Examples of 2-categories, such as **Cat**, or ω -categories, such as globular sets with their face maps, are of course well known, but they need not be repeated.

7 Higher Quantales

We now define the quantalic structures that match higher catoids in Jónsson-Tarski-type modal correspondences. Once again, we start with an axiomatisation of α -quantales, which correspond to α -catoids, but then turn our attention mainly to ω -quantales and n-quantales.

For an ordinal $\alpha \in \{0, 1, ..., \omega\}$, an α -quantale is a structure $(Q, \leq, \cdot_i, 1_i, dom_i, cod_i)_{0 \leq i < \alpha}$ such that each $(Q, \leq, \cdot_i, 1_i, dom_i, cod_i)$ is a modal quantale and the structures interact as follows:

• for all $i \neq j$,

$$dom_i(\alpha \cdot_j \beta) \leq dom_i(\alpha) \cdot_j dom_i(\beta)$$
 and $cod_i(\alpha \cdot_j \beta) \leq cod_i(\alpha) \cdot_j cod_i(\beta)$,

• and for all i < j

$$(\alpha \cdot_j \beta) \cdot_i (\gamma \cdot_j \delta) \leq (\alpha \cdot_i \gamma) \cdot_j (\beta \cdot_i \delta)$$
 and $dom_j(dom_i(\alpha)) = dom_i(\alpha)$.

An α -quantale is strong if for all i < j,

$$dom_j(\alpha \cdot_i \beta) = dom_j(\alpha) \cdot_i dom_j(\beta)$$
 and $cod_j(\alpha \cdot_i \beta) = cod_j(\alpha) \cdot_i cod_j(\beta)$.

These axiom systems are already reduced and irredundant in the sense of Section 6.

Example 7.1. In the double modal quantale on $\bot < a < \top$ with $\cdot_0 = \cdot_1 = \land$, $1_0 = 1_1 = \top$, $dom_0 = id$ and $dom_1(a) = \top = cod_1(a)$ (the rest being fixed), the first five globular 2-quantale axioms hold, but the last one does not: $dom_1(dom_0(a)) = \top \neq a = dom_0(a)$.

Irredundancy of the two weak morphism laws is established using similar counterexamples with 5-elements found by Isabelle. Their particular form is of little interest. \Box

Remark 7.2. The strong resemblance of the α -quantale axioms and the α -catoid ones is not accidental. It is caused by modal correspondences that are developed in the following sections. Nevertheless, there is a mismatch: $dom_j \circ dom_i = dom_i$ is an α -quantale axiom while $s_j \circ s_i = s_i$ is derivable in α -catoids and the same holds for the two morphism axioms of α -quantales. For $dom_j \circ dom_i = dom_i$ with i < j, this can be explained as follows. Our proof of $s_j \circ s_i = s_i$ relies on $\Delta_k(x,y) \Rightarrow t_k(x) = s_k(y)$, but Remark 4.3 shows that the corresponding property is not available for quantales. The related properties $\alpha \cdot_k \beta \neq \bot \Rightarrow cod_k(\alpha) \wedge dom_k(\beta) \neq \bot$ are available, but too weak to translate the proof of $s_j \circ s_i = s_i$ to quantales. Any formal proof is of course ruled out by Example 7.1.

We now turn to ω -quantales.

Lemma 7.3. In every ω -quantale Q, for $0 \le i < j < \omega$,

- 1. $dom_j \circ cod_i = cod_i$, $cod_j \circ dom_i = dom_i$ and $cod_j \circ cod_i = cod_i$,
- 2. $1_i \le 1_i \cdot_i 1_i$, $1_i \cdot_i 1_i = 1_i$ if Q is strong, and $1_i \cdot_i 1_i = 1_i$,
- $3. \ 1_i \leq 1_i$
- 4. $dom_j(1_i) = 1_i$, $dom_i(1_j) = 1_j$, $cod_j(1_i) = 1_i$ and $cod_i(1_j) = 1_j$,
- 5. $dom_i \circ dom_j = dom_j \circ dom_i$, $dom_i \circ cod_j = cod_j \circ dom_i$, $cod_i \circ dom_j = dom_j \circ cod_i$ and $cod_i \circ cod_j = cod_j \circ cod_i$,
- 6. $dom_i(\alpha \cdot_j \beta) = dom_i(\alpha \cdot_j dom_j(\beta))$ and $cod_i(\alpha \cdot_j \beta) = cod_i(cod_j(\alpha) \cdot_j \beta)$.

By (1), the sets $Q_i = Q_{dom_i}$ form a chain: $Q_0 \subseteq Q_1 \subseteq Q_2 \subseteq \cdots \subseteq Q$. Each Q_i is a complete distributive lattice with + as binary sup and \odot_i as binary inf according to the properties of domain quantales recalled in Section 4. Similarly to the situation for ω -catoids in Section 6, the elements of Q_i remain domain elements in all higher dimensions, hence each Q_i is a distributive sublattice of Q_j for all $j \geq i$ and of Q. We have

$$dom_i(x) \odot_k dom_i(y) = dom_i(x) \wedge dom_i(y)$$
 and $cod_i(x) \odot_k cod_i(y) = cod_i(x) \wedge cod_i(y)$

for all $i, j \leq k$. At the same time, all truncations Q_i of Q are i-quantales, so that the chain of the Q_i is a filtration of ω -quantales. The ω -quantale Q is the union of the quantales Q_i if we add an axiom guaranteeing that for all $x \in Q$ there exists and $i \leq \omega$ such that $dom_i(x) = x = cod_i(x)$. As for ω -catoids, we keep this optional and do not require this property in the considerations that follow. The same results hold for strong ω -quantales.

Remark 7.4. An interchange law $(\alpha \cdot_1 \beta) \cdot_0 (\gamma \cdot_1 \delta) \leq (\alpha \cdot_0 \gamma) \cdot_1 (\beta \cdot_0 \delta)$ features in concurrent semirings [Gis88] and concurrent Kleene algebras and quantales [HMSW11]. It has often been contrasted with the seemingly equational interchange laws of category theory. Yet this ignores the weak nature of equality in categories, which may depend on definedness conditions of terms, and which is captured explicitly and precisely by the multioperational language. Example 6.11 shows that the interchange laws of ω -categories are as weak as those of ω -quantales and ω -Kleene algebras (defined below), of which concurrent quantales and Kleene algebras are special cases. See Appendix A for pitfalls of strong interchange laws and related morphisms.

Example 7.5. The identity $1_i = 1_j$ need not hold for i < j in strong ω -quantales. There is a strong 2-quantale on $0 < 1_0 < 1_1$ in which $1_1 \cdot_0 1_1 = 1_1$ and $1_0 \cdot_1 1_1 = 1_1$ (the rest is fixed), and $dom_0(1_1) = 1_0 = cod_0(1_1)$ and $dom_1(1_0) = 1_0 = cod_1(1_0)$ (the rest is again fixed). This makes strong 2-quantales different from the original concurrent quantales mentioned and prevents smaller interchange laws with two or three variables. See [CDS21] for a discussion of how the condition $1_0 = 1_1$ leads to a partial Eckmann-Hilton-style collapse.

Next we consider the interactions of the Kleene stars with the ω -structure.

Lemma 7.6. In every ω -quantale Q, for $0 \le i < j < \omega$,

- 1. $dom_i(\alpha) \cdot_i \beta^{*_j} \leq (dom_i(\alpha) \cdot_i \beta)^{*_j}$ and $\alpha^{*_j} \cdot_i cod_i(\beta) \leq (\alpha \cdot_i cod_i(\beta))^{*_j}$,
- 2. $dom_j(\alpha) \cdot_i \beta^{*_j} \leq (dom_j(\alpha) \cdot_i \beta)^{*_j}$ and $\alpha^{*_j} \cdot_i cod_j(\beta) \leq (\alpha \cdot_i cod_j(\beta))^{*_j}$ if Q is strong,
- 3. $(\alpha \cdot_i \beta)^{*_i} \leq \alpha^{*_i} \cdot_i \beta^{*_i}$.

The properties in (1) and (2) feature as axioms of globular n-Kleene algebras in [CGMS22]. In sum, all axioms of these n-Kleene algebras have now been derived from our smaller, but slightly different set of axioms for ω -quantales and n-quantales. However, these quantales presuppose that arbitrary joins and meets exist, while globular n-quantales are based on globular n-semirings, where only finite sups are assumed to exist and meets are not part of the language. See Section 12 and in particular Remark 12.10 for a detailed discussion.

Finally we list further properties of domains and codomains that are useful below.

Lemma 7.7. In every ω -quantale, for $0 \le i < j < \omega$,

- 1. $dom_i(\alpha) \cdot_i dom_i(\alpha) = dom_i(\alpha)$ and $cod_i(\alpha) \cdot_i cod_i(\alpha) = cod_i(\alpha)$,
- 2. $dom_i(\alpha \cdot_i \beta) = dom_i(\alpha \cdot_i dom_i(\beta))$ and $cod_i(\alpha \cdot_i \beta) = cod_i(cod_i(\alpha) \cdot_i \beta)$,
- 3. $dom_i(\alpha \cdot_i \beta) = dom_i(cod_i(\alpha) \cdot_i \beta)$ and $cod_i(\alpha \cdot_i \beta) = cod_i(\alpha \cdot_i dom_i(\beta))$,
- 4. $dom_i(\alpha \cdot_i \beta) = dom_i(\alpha \cdot_i dom_j(\beta))$ and $cod_i(\alpha \cdot_i \beta) = cod_i(cod_j(\alpha) \cdot_i \beta)$,
- 5. $dom_i(\alpha \cdot_i \beta) \leq dom_i(cod_j(\alpha) \cdot_i \beta)$ and $cod_i(\alpha \cdot_i \beta) \leq cod_i(\alpha \cdot_i dom_j(\beta))$, and equalities hold if Q is strong,
- 6. $dom_i(\alpha) \cdot i(\beta \cdot j\gamma) \leq (dom_i(\alpha) \cdot i\beta) \cdot i(dom_i\alpha) \cdot i\gamma)$ and $(\alpha \cdot j\beta) \cdot i(dom_i(\gamma)) \leq (\alpha \cdot i(dom_i(\gamma)) \cdot i(\beta \cdot i(dom_i(\gamma)))$,
- 7. $dom_i(dom_j(\alpha) \cdot_j \beta) \leq dom_i(\alpha) \cdot_j dom_i(\beta)$ and $cod_i(\alpha \cdot_j cod_j(\beta)) \leq cod_i(\alpha) \cdot_j cod_i(\beta)$,
- 8. $dom_i(dom_i(\alpha) \cdot i\beta) = dom_i(\alpha) \cdot i dom_i(\beta)$ and $cod_i(\alpha \cdot i cod_i(\beta)) = cod_i(\alpha) \cdot i cod_i(\beta)$,
- 9. $dom_i(\alpha) \cdot_j dom_i(\beta) = dom_i(\alpha) \cdot_i dom_i(\beta)$ and $cod_i(\alpha) \cdot_j cod_i(\beta) = cod_i(\alpha) \cdot_i cod_i(\beta)$,
- 10. $(dom_i(\alpha) \cdot_j dom_i(\beta)) \cdot_i (dom_i(\gamma) \cdot_j dom_i(\delta)) = (dom_i(\alpha) \cdot_i dom_i(\gamma)) \cdot_j (dom_i(\beta) \cdot_i dom_i(\delta))$ and $(cod_i(\alpha) \cdot_j cod_i(\beta)) \cdot_i (cod_i(\gamma) \cdot_j cod_i(\delta)) = (cod_i(\alpha) \cdot_i cod_i(\gamma)) \cdot_j (cod_i(\beta) \cdot_i cod_i(\delta))$.

The laws in (2)-(5) are extended locality laws, those in (6) are weak distributivity laws for compositions, those in (7) and (8) extended export laws. Note that export laws $dom(dom(\alpha)\beta) = dom(\alpha)dom(\beta)$ and $cod(\alpha cod(\beta)) = cod(\alpha)cod(\beta)$ hold in any modal semiring and quantale. Finally, the laws in (9) are useful for proving the strong interchange laws in (10).

Example 7.8. The shuffle 2-catoid on Σ in Example 6.15 lifts to the shuffle language 2-quantale on Σ under the standard language product $XY = \{vw \mid v \in X, w \in Y\}$ and the shuffle product of languages discussed in Example 4.2 [CDS21]. The modal structures are trivial, as the empty word language $\{\varepsilon\}$ is the joint identity of the two underlying quantales, but an interchange law $(W||X) \cdot (Y||Z) \subseteq (W \cdot Y)||(X \cdot Z)$ at language level holds. 2-Quantales satisfying such more restrictive conditions have been called *interchange quantales* [CDS21].

8 Higher Convolution Quantales and their Correspondences

We can now extend modal correspondence triangles between local catoids C, modal quantales Q and modal convolution quantales Q^C [FJSZ23] as well as those for interchange multimonoids, interchange quantales and interchange convolution quantales [CDS21], to local ω -catoids, ω -quantales and convolution ω -quantales, as well as their truncations at dimension n. In Section 10 we specialise these results to modal powerset ω -quantales, which, at dimension n, have applications in higher rewriting [CGMS22]. We start with a summary of the 1-dimensional and 2-dimensional cases considered so far.

Let C be a catoid and Q a quantale, henceforth called value or weight quantale. We write Q^C for the set of functions from C to Q. We define, for $f,g:C\to Q$, the convolution operation $*:Q^C\times Q^C\to Q^C$ as

$$(f * g)(x) = \bigvee_{x \in y \odot z} f(y) \cdot g(z)$$

and the unit function $id_0: C \to Q$ as

$$id_0(x) = \begin{cases} 1 & \text{if } x \in Q_0, \\ \bot & \text{otherwise.} \end{cases}$$

We define $\bigvee F: C \to Q$ by point-wise extension, $(\bigvee F)(x) = \bigvee \{f(x) \mid f \in F\}$ for $F \subseteq Q^C$, and in particular $\bot: C \to Q$ by $\bot(x) = \bot$ for all $x \in C$, overloading notation. We also extend the order on Q point-wise to a relation on Q^C . It is consistent with the standard order on the lattice $C \to Q$.

Following [CDS21, FJSZ23] we introduce further notation. For any predicate P we define

$$[P] = \begin{cases} 1 & \text{if } P, \\ \bot & \text{otherwise,} \end{cases}$$

and then $\delta_x(y) = [x = y]$. Any $f: C \to Q$ can now be written as

$$f = \bigvee_{x \in C} f(x) \cdot \delta_x,$$

where \cdot is a module-style action between the "scalars" $f(x) \in Q$ and functions $\delta_x \in Q^C$. More generally, we often write δ_x^{α} for $\alpha \cdot \delta_x$. Then

$$id_{Q_0} = [e \in Q_0] = \bigvee_{e \in Q_0} \delta_e, \qquad \bigvee F = \bigvee_{x \in C} \bigvee \{f(x) \mid f \in F\} \cdot \delta_x, \qquad f \vee g = \bigvee_{x \in C} (f(x) \vee g(x)) \cdot \delta_x.$$

Also, for convolution,

$$(f*g)(x) = \bigvee_{y,z \in C} f(y) \cdot g(z) \cdot [x \in y \odot z], \qquad \text{ and } \qquad f*g = \bigvee_{x,y,z \in C} f(y) \cdot g(z) \cdot [x \in y \odot z] \cdot \delta_x.$$

Finally, whenever Q is a modal quantale, we define $Dom, Cod: Q^C \to Q^C$ as

$$Dom(f) = \bigvee_{x \in C} dom(f(x)) \cdot \delta_{s(x)}$$
 and $Cod(f) = \bigvee_{x \in C} cod(f(x)) \cdot \delta_{t(x)}$.

Correspondences for relational monoids and quantales as well as relational interchange monoids and interchange quantales are already known [CDS21]. They have been extended to local catoids and modal quantales [FJSZ23]. The resulting 2-out-of-3 laws between catoids C, value algebras Q and convolution algebras Q^C require mild non-degeneracy conditions on C or Q. They have previously been given at a fine level of granularity to explain correspondences between individual laws in C, Q and Q^C . Here we only summarise those results relevant to higher dimensional extensions.

Theorem 8.1 ([DHS21, CDS21]).

- 1. Let C be a catoid and Q a quantale. Then Q^C is a quantale.
- 2. Let Q^C and Q be quantales such that $\bot \neq 1$ and $\alpha \cdot (\beta \cdot \gamma) \neq \bot$ for some $\alpha, \beta, \gamma \in Q$. Then C is a catoid.
- 3. Let Q^C be a quantale and C a catoid such that $C_0 \neq \emptyset$ and $w \in (x \odot y) \odot z$ for some $w, x, y, z \in C$. Then Q is a quantale.

In the following lemma, *double catoid* refers to a set equipped with two catoid structures that do not interact. Likewise, *double quantale* refers to a complete lattice equipped with two monoidal structures that do not interact. In particular, there are no interchange laws.

Theorem 8.2 ([CDS21]). Let $(C, \cdot_0, s_0, t_0, \cdot_1, s_1, t_1)$ be a double catoid, and $(Q, \leq, \cdot_0, 1_0, \cdot_1, 1_1)$ be a double quantale. Let $(Q^C, \leq, *_0, id_0, *_1, id_1)$ be the associated double convolution quantale.

- 1. The interchange law holds in Q^C if it holds in C and Q.
- 2. The interchange law holds in C if it holds in Q and Q^C , and if $(\alpha \cdot_1 \beta) \cdot_0 (\gamma \cdot_1 \delta) \neq \bot$ for some $\alpha, \beta, \gamma, \delta \in Q$.
- 3. The interchange law holds in Q if it holds in C and Q^C , and if $y \in u \odot_1 v$, $z \in w \odot_1 x$ and $\Delta(y, z)$ hold for some $u, v, w, x, y, z \in C$.

The original statement in [CDS21] has been translated from relational monoids to catoids along the isomorphism between these structures.

Theorem 8.3 ([FJSZ23]). Let (C, \odot, s, t) be a catoid and $(Q, \leq, \cdot, 1)$ be a quantale. Let $(Q^C, \leq, *, id_0)$ be the associated convolution quantale.

- 1. Q^C is modal if C is local and Q modal.
- 2. C is local if Q and Q^C are modal, and if $1 \neq \bot$ in Q.
- 3. Q is modal if C is local and Q^C modal, and if $\Delta(\ell(x), r(y))$ and $\Delta(z, w)$ for some $w, x, y, z \in C$.

Example 8.4 ([CDS21, FJSZ23]). Theorems 8.2 and 8.3 have the following instances that generalise Example 4.2. Let Q be a modal value quantale.

- 1. Every category C lifts to a modal convolution quantale Q^C . It is similar to a category algebra, using a quantale as value algebra instead of a ring or field and lifting the source and target structure of the category, which category algebras ignore.
- 2. The modal powerset quantales from Example 4.2 are convolution quantales with Q=2.
- 3. Every path category over a digraph lifts to a modal convolution quantale of Q-valued paths.

- 4. Every pair groupoid lift to a modal convolution quantale of Q-valued relation.
- 5. Every shuffle 2-catoid lifts to a convolution 2-quantale of Q-weighted shuffle languages if Q is a 2-quantale.

Relative to Theorems 8.2 and 8.3, it remains to prove 2-out-of-3 correspondences for the globular structure in local ω -catoids and ω -quantales.

In the ω -setting, we use $[P]_i$, strictly speaking, which map to 1_i of the target quantale Q_i . But in practice, we can always pick and swap indices as well as the multiplication with which we integrate such expressions. Hence we usually suppress indices. In the same way we would have to distinguish δ -functions at different dimension formally in terms of the 1_i they map to, but do not do this in practice.

Theorem 8.5. Let C be a local ω -catoid and Q an ω -quantale. Then Q^C is an ω -quantale.

Proof. Relative to Theorem 8.2 and 8.3 it remains to lift the morphism axioms as well as the axiom $Dom_j \circ Dom_i = Dom_i$ for $0 \le i < j < \omega$. Nevertheless we start with the interchange axioms in the multioperational setting. We omit indices related to square brackets as mentioned. For $0 \le i < j < \omega$,

$$\begin{split} ((f *_{j} g) *_{i} (h *_{j} k))(x) &= \bigvee_{y,z} (f *_{j} g)(y) \cdot_{i} (h *_{j} k)(z) \cdot [x \in y \odot_{i} z] \\ &= \bigvee_{y,z} \left(\bigvee_{t,u} f(t) \cdot_{j} g(u) \cdot [y \in t \odot_{j} u] \right) \cdot_{i} \left(\bigvee_{v,w} h(v) \cdot_{j} k(w) \cdot [z \in v \odot_{j} w] \right) \cdot [x \in y \odot_{i} z] \\ &= \bigvee_{t,u,v,w} (f(t) \cdot_{j} g(u)) \cdot_{i} (h(v) \cdot_{j} k(w)) \cdot [x \in (t \odot_{j} u) \odot_{i} (v \odot_{j} w)] \\ &\leq \bigvee_{t,u,v,w} (f(t) \cdot_{i} h(v)) \cdot_{j} (g(u) \cdot_{i} k(w)) \cdot [x \in (t \odot_{i} v) \odot_{j} (u \odot_{i} w)] \\ &= \bigvee_{y,z} \left(\bigvee_{t,v} f(t) \cdot_{i} h(v) \cdot [y \in t \odot_{i} v] \right) \cdot_{j} \left(\bigvee_{u,w} g(u) \cdot_{i} k(w) \cdot [z \in u \odot_{i} w] \right) \cdot [x \in y \odot_{j} z] \\ &= ((f *_{i} h) *_{j} (g *_{i} k))(x). \end{split}$$

For the morphism axiom $Dom_j(f *_i g) \leq Dom_j(f) *_i Dom_j(g)$,

$$\begin{split} Dom_{j}(f*_{i}g)(x) &= \bigvee_{u} dom_{j} \left(\bigvee_{v,w} f(v) \cdot_{i} g(w)\right) \cdot [u \in v \odot_{i} w] \right) \cdot \delta_{s_{j}(u)}(x) \\ &= dom_{j} \left(\bigvee_{v,w} f(v) \cdot_{i} g(w)\right) \cdot [x \in s_{j}(v \odot_{i} w)] \\ &= \bigvee_{v,w} dom_{j}(f(v) \cdot_{i} g(w)) \cdot [x \in s_{j}(v \odot_{i} w)] \\ &\leq \bigvee_{v,w} dom_{j}(f(v)) \cdot_{i} dom_{j}(g(w)) \cdot [x \in s_{j}(v) \odot_{i} s_{j}(w)] \\ &= \bigvee_{t,u} \left(\bigvee_{v} dom_{j}(f(v)) \cdot_{j} \delta_{s_{j}(v)}(t)\right) \cdot_{i} \left(\bigvee_{w} dom_{j}(g(w)) \cdot_{j} \delta_{s_{j}(w)}(u)\right) \cdot [x \in t \odot_{i} u] \\ &= (Dom_{j}(f) *_{i} Dom_{j}(g))(x). \end{split}$$

The proof of $Cod_i(f *_i g) \leq Cod_i(f) *_i Cod_i(g)$ follows by opposition.

The proofs of $Dom_i(f *_j g) \leq Dom_i(f) *_j Dom_i(g)$ and $Cod_i(f *_j g) \leq Cod_i(f) *_j Cod_i(g)$ are obtained by re-indexing these proofs.

Finally, for $Dom_i \circ Dom_i = Dom_i$,

$$\begin{split} Dom_{j}(Dom_{i}(f)) &= \bigvee_{u} dom_{j} \left(\bigvee_{v} dom_{i}(f(v)) \cdot \delta_{s_{i}(v)}(u) \right) \cdot \delta_{s_{j}(u)} \\ &= \bigvee_{v} dom_{j}(dom_{i}(f(v))) \cdot \delta_{s_{j}(s_{i}(v))} \\ &= \bigvee_{v} dom_{i}(f(v)) \cdot \delta_{s_{i}(v)} \\ &= Dom_{i}(f). \end{split}$$

For ω -categories, or at least strong ω -catoids, we get a stronger result.

Corollary 8.6. Let C be a strong local ω -catoid and Q a strong ω -quantale. Then Q^C is a strong ω -quantale.

Proof. It suffices to replay the proofs for the two strong morphism laws for Dom_j and Cod_j with an equality step in the fourth proof step for Dom_j above. The proof for Cod_j follows by opposition.

For the following two theorems we tacitly assume the non-degeneracies needed for Theorem 8.2 and 8.3, calling the respective structures $sufficiently\ supported$, and mention those on C and Q explicitly in the proof. See [CDS21] for the general construction and more detailed explanations of the mechanics of proofs. First we recall three properties from [FJSZ23] that generalise readily beyond one dimension.

As in Theorem 8.2, an ω -fold catoid is a set equipped with ω catoid structures which do not interact. Similarly, an ω -fold quantale is a complete lattice equipped with ω (continuous) monoidal structures which do not interact.

Lemma 8.7. Let C be an ω -fold catoid and Q be an ω -fold modal quantale, all without globular structure. Let Q^C be the associated ω -fold modal quantale. Then, for $i \neq j$,

- 1. $Dom_i(\delta_x^{\alpha}) = \bigvee_y dom_i(\delta_x^{\alpha}(y))\delta_{s_i(y)} = dom_i(\alpha) \cdot \delta_{s_i(x)},$
- 2. $Dom_i(\delta_x^{\alpha} *_j \delta_y^{\beta})(z) = dom_i(\alpha \cdot_j \beta) \cdot_i [z \in s(x \odot_j y)],$
- 3. $(Dom_i(\delta_x^{\alpha}) *_i Dom_i(\delta_y^{\beta}))(z) = dom_i(\alpha) \cdot_i dom_i(\beta) \cdot_i [z \in s_i(x) \odot_i s_i(y)],$
- 4. $(\delta_x^{\alpha} * \delta_y^{\beta})(z) = \alpha \cdot \beta \cdot [z \in x \odot y].$

Theorem 8.8. Let Q^C and Q be ω -quantales with Q sufficiently supported. Then C is a local ω -catoid.

Proof. Relative to Theorem 8.2 and 8.3 we need to check the homomorphism axioms. As in Theorem 8.5, all proofs are similar and we show only one. Suppose $Dom_i(\delta_x^{\alpha} *_j \delta_y^{\beta})(z) \leq (Dom_i((\delta_x^{\alpha}) *_j Dom_i(\delta_y^{\beta}))(z)$ holds in Q^C and $dom_i(\alpha *_j \beta) \leq dom_i(\alpha) *_j dom_i(\beta)$ in Q. Suppose also, for non-degeneracy, that $dom_i(\alpha \cdot_j \beta) \neq \bot$. Then $s_i(x \odot_j y) \leq s_i(x) \odot_j s_i(y)$ using Lemma 8.7(2) and (3).

Once again we get stronger results for strong quantales. The proofs are obvious.

Corollary 8.9. Let Q^C and Q be strong ω -quantales with Q sufficiently supported. Then C is a strong local ω -catoid.

Additional assumptions are needed to obtain ω -categories. We do not investigate them in this article.

Theorem 8.10. Let Q^C be an ω -quantale and C a local ω -catoid that is sufficiently supported. Then Q is an ω -quantale.

Proof. Relative to Theorem 8.2 and 8.3 we need to check the homomorphism axioms and $dom_j \circ dom_i = dom_i$. As in Theorem 8.5, all proofs of homomorphism axioms are similar and we show only one. Suppose $Dom_i(\delta_x^{\alpha} *_j \delta_y^{\beta})(z) \leq (Dom_i(\delta_x^{\alpha}) *_j Dom_i(\delta_y^{\beta}))(z)$ holds in Q^C and $s_i(x \odot_j y) \leq s_i(x) \odot_j s_i(y)$ in C. Suppose also, for non-degeneracy, that $z \in s_i(x) \odot_j s_i(y)$. Then, using Lemma 8.7(2) and (3),

$$dom_i(\alpha \cdot_j \beta) = Dom_i(\delta_x^{\alpha} *_j \delta_y^{\beta})(z) \leq (Dom_i(\delta_x^{\alpha}) *_j Dom_i(\delta_y^{\beta}))(z) = dom_i(\alpha) \cdot_j dom_i(\beta).$$

Finally, using Lemma 8.7(1),

$$dom_j(dom_i(\alpha)) = Dom_j(Dom_i(\delta_x^{\alpha}))(s_j(s_i(x))) = Dom_i(\delta_x^{\alpha})(s_i(x)) = dom_i(\alpha).$$

Corollary 8.11. Let Q^C be a strong ω -quantale and C a strong local ω -catoid that is sufficiently supported. Then Q is a strong ω -quantale.

Example 8.12.

- 1. Cat lifts to a convolution 2-quantale Q^{Cat} for every value 2-quantale Q.
- 2. The ω -category of globular sets and their face maps lifts to a convolution ω -quantale for every value ω -quantale Q.

These convolution algebras may seem artificial, and we have no use for the general weighted case so far. Applications of the powerset case Q=2 relevant to higher rewriting are discussed in Sections 10, 12 and 13. Finally, a convolution 2-quantale relevant to concurrency theory based on certain weighted languages of isomorphism classes of labelled posets, equipped with a so-called serial and a parallel composition is discussed in [CDS21]. In this case, the only unit is the empty poset, and the modal structure is trivial, as for weighted languages with shuffle.

9 Dedekind Convolution Quantales and their Correspondences

In this section we study correspondences between groupoids, Dedekind value quantales and Dedekind convolution quantales, adapting the correspondence between groupoids and relation algebras established by Jónsson and Tarski [JT52].

We define, for every $f: C \to Q$ from a groupoid C into an involutive quantale Q,

$$f^{\circ}(x) = (f(x^{-}))^{\circ}$$
.

Using the previous notation, we can write $f^{\circ} = \bigvee_{x \in C} (f(x))^{\circ} \cdot \delta_{x^{-}}$.

Proposition 9.1. Let C be a groupoid and Q an involutive quantale. Then Q^C is an involutive quantale.

Proof. Relative to Theorem 8.1 it remains to check the three involution axioms. For sup-preservation,

$$(\bigvee F)^{\circ} = \bigvee_{x} ((\bigvee F)(x))^{\circ} \cdot \delta_{x^{-}}$$

$$= \bigvee_{x} ((\bigvee \{f(x) \mid f \in F\})^{\circ} \cdot \delta_{x^{-}}$$

$$= \bigvee \{\bigvee_{x} f(x)^{\circ} \cdot \delta_{x^{-}} \mid f \in F\}$$

$$= \bigvee \{f^{\circ} \mid f \in F\}.$$

For involution proper,

$$f^{\circ \circ} = \bigvee_{x} \left(\bigvee_{y} (f(y))^{\circ} \cdot \delta_{y^{-}}(x) \right)^{\circ} \cdot \delta_{x^{-}}$$

$$= \left(\bigvee_{y} (f(y))^{\circ} \cdot \delta_{y^{--}} \right)^{\circ}$$

$$= \bigvee_{y} (f(y))^{\circ \circ} \cdot \delta_{y^{--}}$$

$$= \bigvee_{y} (f(y)) \cdot \delta_{y}$$

$$= f.$$

For contravariance,

$$(f * g)^{\circ} = \bigvee_{x} \left(\bigvee_{y,z} f(y) \cdot g(z) \right)^{\circ} \cdot [x \in y \odot z] \cdot \delta_{x^{-}}$$

$$= \bigvee_{x,y,z} (f(y) \cdot g(z))^{\circ} \cdot [x \in (y \odot z)^{-}] \cdot \delta_{x}$$

$$= \bigvee_{x,y,z} g(z)^{\circ} \cdot f(y)^{\circ} \cdot [x \in z^{-} \odot y^{-}] \cdot \delta_{x}$$

$$= \bigvee_{x,y,z} g(z^{-})^{\circ} \cdot f(y^{-})^{\circ} \cdot [x \in z \odot y] \cdot \delta_{x}$$

$$= \bigvee_{x} \left(\bigvee_{z,y} g^{\circ}(z) \cdot f^{\circ}(y) \cdot [x \in z \odot y] \right) \cdot \delta_{x}$$

$$= \bigvee_{x} (g^{\circ} * f^{\circ})(x) \cdot \delta_{x}$$

$$= (g^{\circ} * f^{\circ}).$$

Theorem 9.2. Let C be a groupoid and Q a Dedekind quantale in which binary inf distributes over all sups. Then Q^C is a Dedekind quantale (in which binary inf distributes over all sups).

Proof.

$$f * g \wedge h = \left(\left(\bigvee_{y,z} f(y) \cdot g(z) \cdot [x \in y \odot z] \right) \wedge h(x) \right) \cdot \delta_{x}$$

$$= \bigvee_{x,y,z} (f(y) \cdot g(z) \wedge h(x)) \cdot [x \in y \odot z] \cdot \delta_{x}$$

$$\leq \bigvee_{y,z} (f(y) \wedge h(x) \cdot g(z)^{\circ}) \cdot g(z) \cdot [x \in y \odot z]$$

$$= \bigvee_{x,y,z} (f(y) \wedge h(x) \cdot g^{\circ}(z^{-})) \cdot g(z) \cdot [x \in y \odot z] \cdot \delta_{x}$$

$$\leq \bigvee_{x,y,z} \left(f(y) \wedge \bigvee_{v,w} h(v) \cdot g^{\circ}(w) \cdot [y \in v \odot w] \right) \cdot g(z) \cdot [x \in y \odot z] \cdot \delta_{x}$$

$$= \bigvee_{x,y,z} (f(y) \wedge (h * g^{\circ})(y)) \cdot g(z) \cdot [x \in y \odot z] \cdot \delta_{x}$$

$$= \bigvee_{x,y,z} (f \wedge h * g^{\circ})(y) \cdot g(z) \cdot [x \in y \odot z] \cdot \delta_{x}$$

$$= (f \wedge h * g^{\circ}) * g.$$

The distribution property is used in the first step of the proof. The 6th step works because $x \in y \odot z$ if and only if $y \in x \odot z^-$ by Lemma 3.2(2), so that the pair (x, z^-) is considered in the sup introduced.

The condition that binary infs distribute over all sups is well known from frames or locales, that is, complete Heyting algebras. It holds in every power set Dedekind quantale.

Theorem 9.3. Let Q^C and Q be Dedekind quantales such that $\bot \neq 1$ in Q. Then C is a groupoid.

Proof. We show that $x \odot x^- = \{s(x)\}$. Suppose $Dom(\delta_x^\alpha) \le \delta_x^\alpha * (\delta_x^\alpha)^\circ$ holds in Q^C and $dom(\alpha) \le \alpha \cdot \alpha^\circ$ holds in Q. Suppose also for non-degeneracy that $dom(\alpha) \ne \bot$; we can pick $\alpha = 1$ and simply require $\bot \ne 1$. By Lemma 8.7(1) and (4), $\delta_{s(x)}(y) = Dom(\delta_x)(y) \le (\delta_x * (\delta_x)^\circ)(y) = [y \in x \odot x^-]$. Thus $s(x) \in x \odot x^-$. Moreover, $\{s(x)\} = x \odot x^-$ because, in fact, $Dom(\delta_x) = \delta_x * (\delta_x)^\circ$, that is, δ_x is functional.

Theorem 9.4. Let Q^C be a Dedekind quantale and C a groupoid. Then Q is a Dedekind quantale.

Proof. We start with the involutive quantale axioms. First, suppose $f^{\circ\circ} = f$ in Q^C and $x^{--} = x$ in X. Then $\alpha^{\circ\circ} = (\delta_x^{\alpha})^{\circ\circ}(x^{--}) = \delta_x^{\alpha}(x) = \alpha$.

Second, suppose $(\bigvee \{\delta_x^{\alpha} \mid \alpha \in A)^{\circ} = \bigvee \{(\delta_x^{\alpha})^{\circ} \mid \alpha \in A\}$ in Q^X . Then

$$(\bigvee A)^{\circ} = (\bigvee \{\delta_x^{\alpha}(x^{-}) \mid \alpha \in A\})^{\circ} = \bigvee \{(\delta_x^{\alpha})^{\circ}(x^{-}) \mid \alpha \in A\} = \bigvee \{\alpha^{\circ} \mid \alpha \in A\}.$$

Third, suppose $(\delta_y^{\alpha} * \delta_z^{\beta})^{\circ} = (\delta_z^{\beta})^{\circ} * (\delta_y^{\alpha})^{\circ}$ in Q^C and $(y \odot z)^- = z^- \odot y^-$ in C. Also assume $z \in x \odot y$ for non-degeneracy. Then $(\alpha \cdot \beta)^{\circ} = (\delta_y^{\alpha} * \delta_z^{\beta})^{\circ} (x^-) = ((\delta_y^{\beta})^{\circ} * (\delta_z^{\alpha})^{\circ}))(x^-) = \beta^{\circ} \cdot \alpha^{\circ}$.

Finally, for the modular law, suppose $\delta_u^{\alpha} * \delta_v^{\beta} \wedge \delta_w^{\gamma} \leq (\delta_u^{\alpha} \wedge \delta_w^{\gamma} * (\delta_v^{\beta})^{\circ}) * \delta_v^{\beta}$. Assume, for non-degeneracy, that $x \in u \odot v$. Then

$$\alpha \cdot \beta \wedge \gamma = \alpha \cdot \beta \cdot [x \in u \odot v] \wedge \gamma \cdot \delta_x(x)$$

$$= (\delta_u^{\alpha} * \delta_v^{\beta} \wedge \delta_x^{\gamma})(x)$$

$$\leq ((\delta_u^{\alpha} \wedge \delta_x^{\gamma} * (\delta_v^{\beta})^{\circ}) * \delta_v^{\beta})(x)$$

$$= (\alpha \wedge \gamma \cdot \beta^{\circ} \cdot [u \in x \odot v^{-}]) \cdot \beta \cdot [x \in u \odot v]$$

$$= (\alpha \wedge \gamma \cdot \beta^{-}) \cdot \beta.$$

where $x \in u \odot v \Leftrightarrow u \in x \odot v^-$, which holds by Lemma 3.2(2), is used in the fourth step.

Theorem 9.2 and 9.4 clearly depend on the quantales Q or Q^C in their hypotheses being Dedekind, while Theorem 9.3 uses slightly more general properties of the shape $dom(x) \leq xx^{\circ}$ and $cod(x) \leq x^{\circ}x$, which are further discussed in the context of ω -semirings and ω -Kleene algebras in Section 12.

Example 9.5. Example 5.15 generalises from powerset quantales to convolution quantales. Let Q be a Dedekind value quantale.

1. Every free groupoid lifts to a modal Q-valued path Dedekind quantale in which formal inverses lift to converses.

2. Every pair groupoid lifts to a modal Dedekind quantale of Q-valued relations.

10 Correspondences for Powerset Quantales

The correspondences of Section 8 and 9 specialise to powerset algebras for the value quantale Q=2 of booleans (Example 4.1), using the isomorphism between $C \to 2$ and $\mathcal{P}C$. We have already seen several examples of powerset liftings in previous sections. Here we give standalone correspondence proofs for the globular and the converse structure, because they are interesting for higher rewriting.

The first corollary is an immediate instance of Theorems 8.5 and 8.8.

Corollary 10.1.

- 1. Let C be a local ω -catoid. Then $(\mathcal{P}C,\subseteq,\odot_i,C_i,s_i,t_i)_{0\leq i\leq\omega}$ is an ω -quantale.
- 2. Let PC be an ω -quantale in which $id_i \neq \emptyset$. Then C is a local ω -catoid.
- 3. If C is strong, then PC is strong and vice versa.

We emphasise the one-to-one relationship between laws of ω -catoids and those of ω -quantales in the proofs in Appendix B to highlight the underlying modal correspondence, including the proof of this corollary.

Remark 10.2. A classical result by Gautam [Gau57] shows that equations lift to the powerset level if each variable in the equation occurs precisely once in each side. These results have later been generalised by Grätzer and Whitney [GW84] (see also [Bri93] for an overview). It is therefore no surprise that all the (unreduced) axioms of ω -catoids lift directly to corresponding properties, which we have already derived from the ω -quantale axioms in Lemma 7.3. Nevertheless Gautam's result does not prima facie cover multioperations, let alone constructions of convolution algebras.

Next we show how groupoids lift to Dedekind quantales and relation algebras. This mainly reproduces Jónsson and Tarski's results [JT52] with groupoids based on catoids. In addition, we provide explicit liftings for dom and cod, which can otherwise be derived in powerset quantales or relation algebras. In any groupoid C, we write X^- for $X \subseteq C$. First we list some properties that are not directly related to the lifting.

Lemma 10.3. Let C be a groupoid. For all $X, Y \in \mathcal{P}(C)$ and all $X \in \mathcal{P}(\mathcal{P}(C))$.

- 1. $(\bigcup X)^- = \bigcup \{X^- \mid X \in X\} \text{ and } (X \cup Y)^- = X^- \cup X^-,$
- 2. $s(X) \subseteq XX^-$ and $t(X) \subseteq X^-X$,
- 3. $X \subseteq XX^-X$
- 4. $s(X) = C_0 \cap XX^- \text{ and } t(X) = C_0 \cap X^-X$,
- 5. $s(X) = C_0 \cap X \top$ and $r(X) = C_0 \cap \top X$,
- 6. $X \top = s(X) \top$ and $\top X = \top t(X)$.

Properties (2) and (3) are of course the conditions for involutive quantales from Section 5. Next we specialise the results of Section 9.

Corollary 10.4.

- 1. Let C be a groupoid. Then $(\mathcal{P}C, \odot, s, t, (-)^-)$ is a Dedekind quantale.
- 2. Let $\mathcal{P}C$ be a Dedekind quantale in which $C_0 \neq \emptyset$ and the atoms are functional. Then C is a groupoid.

The proofs are due to Jónsson and Tarski; we have also checked (1) with Isabelle. The functionality requirement on atoms, in particular, has been noticed and used by Jónsson and Tarski.

Remark 10.5. Jónsson and Tarski have considered relation algebras based on boolean algebras instead of complete lattices. They also use the residual law mentioned in Section 5 instead of the modular or Dedekind law. This makes no difference. See [HH02, Mad06] for details.

Examples of powerset liftings for groupoids have been given in Example 5.15. This section provides a formal development that subsumes these results.

11 Modal Operators and their Laws

Modal diamond operators can be defined in modal semirings or quantales [DS11, FJSZ22a] as

$$|\alpha\rangle\beta = dom(\alpha\beta)$$
 and $\langle\alpha|\beta = cod(\beta\alpha)$.

By domain and codomain locality, $|\alpha\rangle\beta = |\alpha\rangle dom(\beta)$ and $\langle\alpha|\beta = \langle\alpha|dom(\beta)$. Accordingly, we shall mostly use those modal operators with β a domain element. See [FJSZ22a, FJSZ23] for additional properties for modal operators on quantales. In particular, we can "demodalise" diamonds at left-hand sides of inequalities,

$$|\alpha\rangle p \le q \Leftrightarrow \alpha p \le q\alpha$$
 and $\langle \alpha|p \le q \Leftrightarrow p\alpha \le \alpha q$,

where $p, q \in Q_0$.

In boolean modal quantales, and hence in modal powerset quantales, we can define forward and backward modal box operators

$$[\alpha|p = \bigvee \{q \mid |\alpha\rangle q \le p\}$$
 and $|\alpha|p = \bigvee \{q \mid \langle \alpha|q \le p\},$

for all $\alpha \in Q$ and $p \in Q_0$. (In arbitrary modal quantales, we cannot guarantee that the sups, which feature in the definienda, are again elements of Q_0 .)

The modal box and diamond operators are then adjoints in the Galois connections

$$|\alpha\rangle p \le q \Leftrightarrow p \le |\alpha|q$$
 and $\langle\alpha|p \le q \Leftrightarrow p \le |\alpha|q$,

for all $\alpha \in Q$ and $p, q \in Q_0$. The additional demodalisation laws $p \leq [\alpha|q \Leftrightarrow \alpha p \leq q\alpha \text{ and } p \leq |\alpha|q \Leftrightarrow p\alpha \leq \alpha q$ are helpful in deriving them. Further, in boolean modal quantales, boxes and diamonds are related by De Morgan duality,

$$|\alpha|p = -|\alpha\rangle - p, \qquad |\alpha\rangle p = -|\alpha| - p, \qquad [\alpha|p = -\langle\alpha| - p, \qquad \langle\alpha|p = -[\alpha| - p.$$

Finally, in any modal Dedekind quantale, $\langle \alpha | = | \alpha^{\circ} \rangle$, $| \alpha \rangle = \langle \alpha^{\circ} |$, $[\alpha | = | \alpha^{\circ}]$ and $| \alpha | = [\alpha^{\circ} |$. Modal diamond operators satisfy the following module-style laws.

Lemma 11.1. In every modal quantale,

1.
$$|\alpha\beta\rangle p = |\alpha\rangle |\beta\rangle p$$
,

- 2. $|\alpha \vee \beta\rangle p = |\alpha\rangle p \vee |\beta\rangle p$.
- 3. $|\alpha\rangle(p\vee q) = |\alpha\rangle p\vee |\alpha\rangle q$ and $|\alpha\rangle\perp = \perp$,
- 4. $|\perp\rangle p = \perp$, $|1\rangle p = p$ and $|\alpha\rangle 1 = dom(\alpha)$.

These laws hold already in domain semirings [DS11]. Dual properties hold for backward diamonds, and for boxes, if the modal quantale is boolean.

Beyond these one-dimensional properties, we list some new modal laws for the ω -structure. We write $\langle \alpha \rangle_i$ for either $|\alpha \rangle_i$ or $\langle \alpha |_i$ in the following lemma.

Lemma 11.2. In every modal ω -quantale Q, for all $0 \le i < j < \omega$,

- 1. $\langle \alpha \rangle_i | \beta \rangle_j \gamma = \langle \alpha \rangle_i (\beta \cdot_j \gamma)$ and $\langle \alpha \rangle_i \langle \beta |_j \gamma = \langle \alpha \rangle_i (\gamma \cdot_j \beta)$,
- 2. $|\alpha\rangle_i\langle\beta\rangle_j\gamma \leq |\alpha\rangle_i(dom_i(\beta)\cdot_j dom_i(\gamma))$ and $|\alpha\rangle_i\langle\beta\rangle_j\gamma \leq |\alpha\rangle_i(cod_i(\gamma)\cdot_j cod_i(\beta))$,
- 3. If $\alpha = dom_k(\alpha)$ for some $k \leq j$, then $|\alpha\rangle_i |\beta\rangle_j \gamma \leq |\alpha\rangle_i \beta \cdot_j |\alpha\rangle_i \gamma$ and $|\alpha\rangle_i |\beta\rangle_j \gamma \leq |\alpha\rangle_i |\beta\rangle_i |\beta\rangle_j \gamma \leq |\alpha\rangle_i |\beta\rangle_i |\beta\rangle_i |\beta\rangle_j \gamma \leq |\alpha\rangle_i |\beta\rangle_i |\delta\rangle_i |\delta$
- 4. $|\alpha\rangle_j\langle\beta\rangle_i\gamma\leq |\alpha\rangle_j\langle dom_j(\beta)\rangle_i\gamma$ and $|\alpha\rangle_j\langle\beta\rangle_i\gamma\leq |\alpha\rangle_j\langle cod_j(\beta)\rangle_i\gamma$, and equality holds if Q is strong,
- 5. $dom_i(\alpha) \cdot_i \langle \beta \rangle_j \gamma \leq \langle dom_i(\alpha) \cdot_i \beta \rangle_j (dom_i(\alpha) \cdot_i \gamma)$ and $\langle \alpha \rangle_j \beta \cdot_i dom_i(\gamma) \leq \langle \alpha \cdot_i dom_i(\gamma) \rangle_j (\beta \cdot_i dom_i(\gamma))$, if Q is strong.

12 Higher Semirings and Kleene Algebras

A main reason for using quantales in our constructions so far is that convolutions

$$(f * g)(x) = \bigvee_{x \in y \odot z} f(y) \cdot g(z)$$

require arbitrary sups when infinitely many pairs (y, z) satisfy the ternary relation $x \in y \odot z$. Yet when C is *finitely decomposable*, when for each $x \in C$ the fibre $\odot^{-1}(x) = \{(y, z) \mid x \in y \odot z\}$ is finite, algebras with finite sups suffice. Modal value quantales can then be replaced by modal value semirings [DS11]. These are based on dioids or additively idempotent semirings instead of quantales. Similar adaptations can be made for concurrent quantales and concurrent semirings [CDS21].

Example 12.1 ([CDS21]). In the Q-valued shuffle language 2-quantale from Example 8.4, the fibres $\{(u,v) \mid w = uv\}$ and $\{(u,v) \mid w = u \mid v\}$ are finite for every word w. It therefore suffices that Q is a 2-semiring, as defined below.

We now we extend these special cases to higher dimensions. Recall that a modal semiring [DS11] is a semiring $(S, +, \cdot, 0, 1)$ equipped with operations $dom, cod : S \to S$ that satisfy precisely the modal quantale axioms up-to notational differences:

$$\alpha \leq dom(\alpha)\alpha, \qquad dom(\alpha dom(\beta)) = dom(\alpha\beta), \qquad dom(\alpha) \leq 1, \qquad dom(0) = 0, \\ dom(x+y) = dom(x) + dom(y),$$

the opposite axioms for cod and the compatibility axioms $dom \circ cod = cod$ and $cod \circ dom = dom$.

Every modal semiring is automatically a *dioid*, an additively idempotent semiring.

For an ordinal $\alpha \in \{0, 1, ..., \omega\}$, an α -semiring is a structure $(S, +, 0, \cdot_i, 1_i, dom_i, cod_i)_{0 \le i < \alpha}$ such that the $(S, +, 0, \cdot_i, 1_i, dom_i, cod_i)$ are modal semirings and the structures interact in the same way as for α -quantales

• for all $i \neq j$,

$$dom_i(\alpha \cdot_j \beta) \leq dom_i(\alpha) \cdot_j dom_i(\beta)$$
 and $cod_i(\alpha \cdot_j \beta) \leq cod_i(\alpha) \cdot_j cod_i(\beta)$,

• and for all i < j

$$(\alpha \cdot_j \beta) \cdot_i (\gamma \cdot_j \delta) \leq (\alpha \cdot_i \gamma) \cdot_j (\beta \cdot_i \delta)$$
 and $dom_j(dom_i(\alpha)) = dom_i(\alpha)$.

An α -semiring is strong if $dom_j(\alpha \cdot_i \beta) = dom_j(\alpha) \cdot_i dom_j(\beta)$ and $cod_j(\alpha \cdot_i \beta) = cod_j(\alpha) \cdot_i cod_j(\beta)$ hold for all i < j.

As before, we now turn to ω -semirings. Most properties derived for ω -quantales in previous sections are already available in this setting, in particular Lemma 7.3, 7.7, 11.1 and 11.1, the results on chains of C_i cells and ω -quantales as filtrations, and the construction of n-quantales by truncation.

Along the lines of Section 5 we can also define an *involutive dioid* S as a dioid S equipped with an operation $(-)^{\circ}: S \to S$ that satisfies $\alpha^{\circ \circ} = \alpha$, $(\alpha + \beta)^{\circ} = \alpha^{\circ} + \beta^{\circ}$ and $(\alpha\beta)^{\circ} = \beta^{\circ}\alpha^{\circ}$. These axioms imply in particular $0^{\circ} = 0$ and $1^{\circ} = 1$. Yet the interaction between domain, codomain and involution are weak.

Example 12.2. In the involutive modal semiring given by



with \cdot as inf, dom = id = cod and $a^{\circ} = b$, $b^{\circ} = a$ we have $dom(a^{\circ}) = dom(b) = b \neq a = cod(a)$ and likewise $cod(a^{\circ}) \neq dom(a)$. Domain and codomain are therefore not related via converse.

Similar examples show that $dom(\alpha)^{\circ}$ and $cod(\alpha)^{\circ}$ need not be equal to $dom(\alpha)$ and $cod(\alpha)$, respectively.

Remark 12.3. In an involutive domain semiring or domain semiring with converse S, one can of course define codomain explicitly as $cod(\alpha) = dom(\alpha^{\circ})$, from which $dom(\alpha) = cod(\alpha^{\circ})$ follows, but then S_{dom} need not be equal to S_{cod} : $dom \circ cod = cod$ may hold, but $cod \circ dom = dom$ may fail. We do not consider such alternative definitions of modal semirings with converse any further.

In the absence of infs or boolean complementation, the modular, Dedekind or residual law are out of reach. Instead, the strong Gelfand law $\alpha \leq \alpha \alpha^{\circ} \alpha$ is usually added to the involution axioms [BÉS95]. We call such a diod a dioid with converse. A modal semiring with converse is then a modal semiring that is also a dioid with converse.

Lemma 12.4. In every modal semiring with converse,

- 1. $dom(\alpha)^{\circ} = dom(\alpha)$ and $cod(\alpha)^{\circ} = cod(\alpha)$,
- 2. $dom(\alpha^{\circ}) = cod(\alpha)$ and $cod(\alpha^{\circ}) = dom(\alpha)$.

Stronger axioms, which link domain, codomain and converse more directly, have been proposed in [CGM21]: $dom(\alpha) \leq \alpha\alpha^{\circ}$ and $cod(\alpha) \leq \alpha^{\circ}\alpha$. These imply the strong Gelfand property, $\alpha = dom(\alpha)\alpha \leq \alpha\alpha^{\circ}x$, and hence link domain and codomain via converse as in Lemma 12.4. We call the resulting modal semirings modal semirings with strong converse.

Example 12.5. In the modal semiring with converse given by 0 < a < 1, aa = a, dom(a) = cod(a) = 1 and $(-)^{\circ} = id$, $dom(a) = cod(a)1 > a = aa^{\circ} = a^{\circ}a$. It is therefore not a modal semiring with strong converse.

For finitely decomposable ω -catoids C, our main theorems generalise to ω -semirings. First we present a corollary to Theorem 8.5.

Corollary 12.6. Let C be a finitely decomposable local ω -catoid and S an ω -semiring. Then S^C is an ω -semiring.

Proof. In the construction of the convolution algebra in the proof of Theorem 8.5, it is routine to check that all sups remain finite when C is finitely decomposable.

Dioids can sometimes be extended to Kleene algebras K, adding a Kleene star $(-)^*: K \to K$. By contrast to the quantalic Kleene star $\alpha^* = \bigvee_{i>0} \alpha^i$ in Section 4, it is modelled in terms of least fixpoints.

A Kleene algebra is a dioid K equipped with a star operation $(-)^* : K \to K$ that satisfies the star unfold and star induction axioms

$$1 + \alpha \cdot \alpha^* \le \alpha^*, \qquad \gamma + \alpha \cdot \beta \le \beta \Rightarrow \alpha^* \cdot \gamma \le \beta$$

and their opposites, where the arguments in compositions have been swapped.

Every quantale is a dioid, and in fact a Kleene algebra: the quantalic Kleene star satisfies the Kleene algebra axioms. An adaptation of the definition of ω -quantales to Kleene algebras is straightforward.

An α -Kleene algebra is an α -semiring K equipped with Kleene stars $(-)^{*i}: K \to K$ that satisfy the usual star unfold and star induction axioms for all $0 \le i < j < \alpha$

$$1_{i} + \alpha \cdot_{i} \alpha^{*_{i}} \leq \alpha^{*_{i}}, \qquad \gamma + \alpha \cdot_{i} \beta \leq \beta \Rightarrow \alpha^{*_{i}} \cdot_{i} \gamma \leq \beta,$$

$$1_{i} + \alpha^{*_{i}} \cdot_{i} \alpha \leq \alpha^{*_{i}}, \qquad \gamma + \beta \cdot_{i} \alpha \leq \beta \Rightarrow \gamma \cdot_{i} \alpha^{*_{i}} \leq \beta,$$

$$dom_{i}(\alpha) \cdot_{i} \beta^{*_{j}} \leq (dom_{i}(\alpha) \cdot_{i} \beta)^{*_{j}}, \qquad \alpha^{*_{j}} \cdot_{i} cod_{i}(\beta) \leq (\alpha \cdot_{i} cod_{i}(\beta))^{*_{j}}.$$

An α -Kleene algebra is *strong* if the underlying α -semiring is, and for all $0 \le i < j < \alpha$,

$$dom_j(\alpha) \cdot_i \beta^{*_j} \leq (dom_j(\alpha) \cdot_i \beta)^{*_j}, \qquad \alpha^{*_j} \cdot_i cod_j(\beta) \leq (\alpha \cdot_i cod_j(\beta))^{*_j}.$$

The axioms mentioning domain and codomain are derivable in (strong) ω -quantales, and more generally, α -quantales (Lemma 7.6). In α - or ω -Kleene algebras we have neither proofs nor counterexamples with respect to the remaining axioms, yet these laws are needed for coherence proofs in higher rewriting [CGMS22]. However, the following property, which was used as an axiom in [CGMS22], is derivable.

Lemma 12.7. In every
$$\omega$$
-Kleene algebra, $(\alpha \cdot_i \beta)^{*_i} \leq \alpha^{*_i} \cdot_i \beta^{*_i}$ for all $0 \leq i < j < \omega$.

Remark 12.8. A star on convolution Kleene algebras K^C can be defined for finitely decomposable catoids C with a single unit that satisfy a certain grading condition [CDS21]. This approach does not seem to scale to the present setting that requires multiple units.

The situation simplifies for powerset liftings. As in language or relational models of Kleene algebra, the Kleene star is simply a union of powers in this case and finite decomposibility is unnecessary. Corrollary 10.1 specialises as follows.

Corollary 12.9.

- 1. Let C be a local ω -catoid. Then $\mathcal{P}C$ is an ω -Kleene algebra.
- 2. Let PC be an ω -Kleene algebra in which $C_i \neq 0$. Then C is a local ω -catoid.
- 3. If C is strong, then PC is strong and vice versa.

Typical powerset ω -Kleene algebras are subalgebras of powerset ω -quantales that are generated from some finite set and closed under the operations of ω -Kleene algebras. In computer science, languages over a finite alphabet Σ , for instance, form a quantale, whereas the regular languages generated by Σ form a Kleene algebra, but not a quantale. In higher dimensions, this construction, with generators at each dimensions, can be based on computads or polygraphs [Str76, Bur91]. We will explore this as future work.

Remark 12.10. A previous axiomatisation of higher globular semirings and Kleene algebras [CGMS22, Section 3.2] has been based on slightly different axioms. These axioms have been proved sound with respect to a powerset model of path n-categories [CGMS22, Section 3.3], but axioms have been introduced in ad

hoc fashion with a view on higher rewriting proofs. Our modal correspondences with respect n-catoids and n-categories provide a more systematic structural justification of the ω -quantale, ω -semiring and ω -Kleene algebra axioms at powerset level, and hence the higher algebras in [CGMS22].

One difference between the previous axiomatisation and ours is that the previous one does not include the morphism laws $dom_i(\alpha \cdot_j \beta) \leq dom_i(\alpha) \cdot_j dom_i(\beta)$ and $cod_i(\alpha \cdot_j \beta) \leq cod_i(\alpha) \cdot_j cod_i(\beta)$, which are irredundant among the ω -quantale axioms, as mentioned in Example 7.1, but can be lifted from any ω -catoid. A second difference is that the previous axiomatisation contains redundant assumptions and axioms, for instance on the bounded distributive lattice structure of S_0 or K_0 , or the relationships between quantalic units at different dimensions, which are now derivable.

Next we consider (strong) converse in the context of dioids and Kleene algebras. An *involutive Kleene* algebra is an involutive dioid and a Kleene algebra. A *Kleene algebra with converse* is a dioid with converse and a Kleene algebra. A *modal Kleene algebra with strong converse* is a Kleene algebra and a modal semiring with strong converse.

Lemma 12.11. In every involutive Kleene algebra, $\alpha^{*\circ} = \alpha^{\circ*}$.

The following fact considers once again dioids instead of Kleene algebras in the context of convolution algebras.

Proposition 12.12. Let C be a finitely decomposable groupoid and S a dioid with (strong) converse. Then S^C is a dioid with (strong) converse.

Proof. First note that the proof of Proposition 9.1 needs finite sups only when C is finitely decomposable. Relative to this proposition, it remains to lift the strong Gelfand property and alternatively the axioms $dom(\alpha) \leq \alpha\alpha^{\circ}$ and $cod(\alpha) \leq \alpha^{\circ}\alpha$. In the proofs we still write \bigvee , but tacitly assume that all sups used are finite and can therefore be represented using +.

For dioids with converse,

$$f * (f^{\circ} * f) = \bigvee_{w,x,y,z \in C} f(x)f(y^{-})^{\circ}g(z)[w \in x \odot y \odot z]\delta_{w}$$

$$= \bigvee_{w,x,y,z \in C} f(x)f(y)^{\circ}g(z)[w \in x \odot y^{-} \odot z]\delta_{w}$$

$$\geq \bigvee_{w \in C} f(w)f(w)^{\circ}g(w)[w \in w \odot w^{-} \odot w]\delta_{w}$$

$$\geq \bigvee_{w \in C} f(w)[w \in s(w)w]\delta_{w}$$

$$= \bigvee_{w \in C} f(w)\delta_{w}$$

$$= f.$$

For dioids with strong converse,

$$Dom(f) = \bigvee_{x \in C} dom(f(x))\delta_{s(x)}$$

$$\leq \bigvee_{x \in C} f(x)f(x)^{\circ}[s(x) \in x \odot x^{-}]\delta_{x}$$

$$\leq \bigvee_{x,y,z \in C} f(y)f(z^{-})^{\circ}[x \in y \odot z^{-}]\delta_{x}$$

$$= \bigvee_{x \in C} (f * f^{\circ})(x)\delta_{x}$$

$$= f * f^{\circ}$$

and the proof for *Cod* follows by opposition.

For powersets and in the presence of (strong) converse, Corollary 10.4 specialises as follows.

Corollary 12.13.

- 1. Let C be a groupoid. Then PC is a modal Kleene algebra with strong converse.
- 2. Let PC be modal Kleene algebra with strong converse in which $C_0 \neq \emptyset$ and all atoms are functional. Then C is a groupoid.

Proof. For (1), it remains to lift the strong converse axioms, as the modular law is not available. This has been done in Lemma 10.3.

For (2), note that the strong converse axioms are used in the proof of Theorem 9.3 instead of the Dedekind or modular law, but the strong Gelfand law would be too weak. For a direct proof,

$$x \odot x^{-} = \{x\} \cdot \{x\}^{\circ} = dom(\{x\}) = \{s(x)\},\$$

and likewise for the target axiom, using the strong converse axiom for domain in the second step (as an equation because atoms are functional). \Box

Remark 12.14. McKinsey has shown in the 1040s that every group lifts to a relation algebra [JT52]. A group is a groupoid with a single unit, but our current tools still do not allow to extend this result to a convolution Kleene algebra. The technique introduced in [CDS21] depends on a grading function $|-|: G \to \mathbb{N}$ on the group G that needs to satisfy |x| > 0 for all $x \neq e$ and $|x \odot y| = |x| + |y|$. These definitions imply that |x| = 0 if and only if x = e so that $|x \odot x^-| = 0$, which is impossible. Extending the grading function to integers is also impossible as it is used as a basis of an induction.

13 (ω, p) -Catoids and (ω, p) -Quantales

In this final technical section we combine the results of Section 9 and 8 to the (n,p) Kleene algebras and related structures needed for higher rewriting [CGMS22]. We start from ω -catoids having a groupoid structure above some dimension $p < \omega$. These (ω, p) -catoids and the corresponding (ω, p) -quantales, which are Dedekind quantales above dimension p, are suitable for defining higher homotopies. They specialise to (ω, p) -dioids and (ω, p) -Kleene algebras and truncations at any dimension.

An (ω, p) -catoid is an ω -catoid C with operations $(-)^{-i}: C \to C$ for all $p < i \le \omega$ such that $(x)^{-i} \in C_i$ for all $x \in C_i$ and the groupoid axioms hold, for all $p < i \le \omega$ and $x \in C$:

$$x \odot_{i-1} x^{-i} = \{s_{i-1}(x)\}$$
 and $x^{-i} \odot_{i-1} x = \{t_{i-1}(x)\}.$

 $An(\omega, p)$ -category is a local functional (ω, p) -catoid.

Likewise, an (ω, p) -quantale is an ω -quantale Q with operations $(-)^{\circ_i}: Q \to Q$ for all $p < i < \omega$ such that $(\alpha)^{\circ_j} \in Q_j$ for all $\alpha \in Q_j$ and that satisfy the involution axioms and the modular law with respect to the composition \cdot_{j-1} . Strong (ω, p) -quantales are defined as usual.

All inverses in (ω, p) -catoids and all converses in (ω, p) -quantales are trivial on elements of lower dimensions. It follows from Lemma 3.1 that $x^{-j} = x$ for all $x \in C_i$ with i < j. Similarly, it follows from Lemma 5.9 that $\alpha^{\circ j} = \alpha$ for all $\alpha \in Q_i$ with i < j.

The results from Sections 8 and 9 can then be combined. We use the notion of sufficient support as for ω -catoids, ω -groupoids and ω -quantales defined in these sections.

Theorem 13.1.

- 1. Let C be a local (ω, p) -catoid and Q an (ω, p) -quantale. Then Q^C is an (ω, p) -quantale.
- 2. Let Q^C and Q be (ω, p) -quantales with Q sufficiently supported. Then C is a local (ω, p) -quantale.

3. Let Q^C be an (ω, p) -quantale and C a local (ω, p) -catoid that is sufficiently supported. Then Q is an (ω, p) -catoid.

These results specialise to correspondence triangles for strong (ω, p) -catoids and strong (ω, p) -quantales as usual. Further, we obtain correspondences for powerset quantales as instances.

Corollary 13.2.

- 1. Let C be a local (ω, p) -catoid. Then PC is an (ω, p) -quantale.
- 2. Let $\mathcal{P}C$ be an (ω, p) -quantale in which $C_i \neq \emptyset$ and the atoms are functional. Then C is a local (ω, p) -catoid.

Again there are special cases for strong structures.

For finitely decomposable catoids and groupoids, we may replace quantales with dioids or Kleene algebras. An (ω, p) -dioid is an ω -dioid S equipped with operations $(-)^{\circ_j}: S \to S$ for all $p < j \le n$ such that $(\alpha)^{\circ_j} \in S_j$ for all $\alpha \in S_j$ and that satisfy the involution axioms and $\alpha \le \alpha \alpha^{\circ_j} \alpha$ for $p < j \le n$. The definition of (ω, p) -Kleene algebra is then straightforward. The fact that $\alpha^{\circ_j} = \alpha$ for all $\alpha \in S_{dom_i}$ (or K_j) with i < j now follows from Lemma 12.4.

Corollary 13.2 then specialises further to (ω, p) -semirings and (ω, p) -Kleene algebras owing to Corollary 12.9. In particular, an (ω, p) -Kleene algebra with strong converse is needed to recover the (ω, p) -catoid among the atoms. For the more general convolution algebra constructions, the limitations mentioned in Section 12 remain. We can still lift to (ω, p) -semirings, that is, whenever C is a finitely decomposable local (ω, p) -catoid and S an (ω, p) -semiring, then S^C is an (ω, p) -semiring. Yet once again the extension to Kleene algebras requires further thought. Truncations at dimension n work as expected and the algebras introduced in [CGMS22] arise as special cases.

14 Conclusion

This paper combines two lines of research: one on higher globular algebras for higher rewriting and another one on the construction of convolution algebras on catoids and categories. More specifically, we have introduced ω -catoids and ω -quantales, and established modal correspondence results between them. These extend and justify the axioms of higher globular Kleene algebras [CGMS22], which have previously been used for coherence proofs in higher rewriting, up-to minor modifications. We have also introduced several extensions and specialisations of these constructions, in particular to (ω, p) -catoids and (ω, p) -quantales and to variants of such quantales based on semirings or Kleene algebras. While the technical focus has been on convolution algebras, which often lead to interesting applications in quantitative program semantics and verification in computer science, we currently have no use for convolution ω -quantales or (ω, p) -quantales with value quantales different from the quantale of booleans. In the latter case, however, we can use power set ω -quantales instead of globular Kleene algebras to reason about higher rewrite systems with greater flexibility and greater expressive power. A detailed introduction to the mechanics of higher rewriting, globular Kleene algebras and their use in coherence proofs such as higher Church-Rosser theorems and higher Newman's lemmas can be found in [CGMS22].

Our results add a new perspective to higher rewriting and yield new tools for reasoning with higher categories, but work remains to be done to apply them. A first line of research could consider categorical variations of the strict globular ω -catoids and ω -quantales introduced in this work to provide a categorical framework for the construction of coherence proofs and more generally of polygraphic resolutions, which are cofibrant replacement in higher categories:

 It is worth considering cubical versions of ω-catoids and ω-quantales, as confluence diagrams in higher rewriting have a cubical shape [Luc17, Luc20] and proofs of higher confluence properties can be developed more naturally in a cubical setting. A single-set axiomatisation of cubical categories has recently be developed [MMS24]. It remains to generalise it to cubical catoids, to introduce cubical quantales and to study their correspondences along the lines of the globular case. Cubical catoids and quantales may also be beneficial for explicit constructions of resolutions. Beyond that, we are interested in applications to precubical sets and higher-dimensional automata, where Kleene algebras or quantales describing their languages remain to be defined [FJSZ21, FJSZ22b], and to cubical ω -categories with connections [AABS02], where single-set formalisations might be of interest.

- To expand the constructions in this article to algebraic rewriting, such as string, term, linear or diagrammatic rewriting, the development of catoids and quantales internal to categories of algebras over an operad would be needed.
- The homotopic properties of algebraic rewriting require weakening the exchange law of higher categories [For21, FM22] and, in our context, Gray-variants of the ω -catoids and ω -quantales, where interchange law holds only up to coherent isomorphism [GPS95].

A second line of research could investigate the link of the correspondence triangles between catoids, value quantales and convolution quantales with duality results for convolution algebras beyond the Jónsson-Tarski case and similar structural results, and the categorification of the approach to obtain such results. Catoids, for instance, are (isomorphic to) monoids in the monoidal category Rel with the standard tensor and unit. Related to this are questions about free powerset or convolution ω -quantales generated by polygraphs, or computads, and about coherent rewriting properties of these internal monoids.

A third line of research could consider the formalisation of higher category theory, and higher rewriting support for these, in the single-set framework of catoids developed in [Str23, CS23, CS24] or by other means with proof assistants such as Coq, Lean or Isabelle. The formalisation of the theorems in [CGMS22] could serve as stepping stones towards coherence theorems like Squier's theorem in higher dimensions [GM09, GM18] or the computation of resolutions by rewriting in categorical and homological algebra [GM12, GHM19, MR23].

Last, but not least, the study of domain, codomain and converse in the setting of quantales leads to interesting questions about Dedekind quantales and the interplay of these operations in weak variants of allegories [FS90]. These will be addressed in successor articles.

Acknowledgement The authors would like to thank James Cranch, Uli Fahrenberg, Éric Goubault, Amar Hadzihasanovic, Christian Johanson, Tanguy Massacrier and Krzysztof Ziemiański for interesting discussions and the organisers of the GETCO 2022 conference and the Nordic Congress of Mathematicians 2023 for the opportunity to present some of the results in this article. The fourth author would like to thank the Plume team and the Computer Science Department of the ENS de Lyon for funding a short visit at the LIP laboratory in the final stages of this work.

References

- [AABS02] F. A. Al-Agl, R. Brown, and R. Steiner. Multiple categories: The equivalence of a globular and a cubical approach. *Advances in Mathematics*, 170:71–118, 2002.
- [ABG⁺24] D. Ara, A. Burroni, Y. Guiraud, P. Malbos, F. Métayer, and S. Mimram. Polygraphs: from rewriting to higher categories. London Mathematical Society Lecture Note Series, 666pp, arXiv:2312.00429, to appear, 2024.
- [BÉS95] S. L. Bloom, Z. Ésik, and G. Stefanescu. Notes on equational theories of relations. *Algebra Universalis*, 33, 1995.
- [BH81] R. Brown and P. J. Higgins. The equivalence of ∞-groupoids and crossed complexes. Cahiers de topologie et géométrie différentielle catégoriques, 22(4):371–383, 1981.
- [Bri93] C. Brink. Power structures. Algebra Universalis, 30:177–216, 1993.

- [Bro87] R. Brown. From groups to groupoids: A brief survey. Bulletin of the London Mathematical Society, 19:113–134, 1987.
- [Bur91] A. Burroni. Higher-dimensional word problems with applications to equational logic. *Theoretical Computer Science*, 115(1):43–62, 1991.
- [CDS20] J. Cranch, S. Doherty, and G. Struth. Relational semigroups and object-free categories. CoRR, abs/2001.11895, 2020.
- [CDS21] J. Cranch, S. Doherty, and G. Struth. Convolution and concurrency. *Mathematical Structures in Computer Science*, 31(8):918–949, 2021.
- [CGM21] C. Calk, E. Goubault, and P. Malbos. Abstract strategies and coherence. In RAMiCS 2021, volume 13027 of LNCS, pages 108–125. Springer, 2021.
- [CGMS22] C. Calk, E. Goubault, P. Malbos, and G. Struth. Algebraic coherent confluence and globular Kleene algebras. *Logical Methods in Computer Science*, 18(4):9:1–9:43, 2022.
- [CS23] C. Calk and G. Struth. Modal quantales, involutive quantales, dedekind quantales. Archive of Formal Proofs, July 2023. https://isa-afp.org/entries/Quantales_Converse.html, Formal proof development.
- [CS24] C. Calk and G. Struth. Higher globular catoids and quantales. Archive of Formal Proofs, January 2024. https://isa-afp.org/entries/OmegaCatoidsQuantales.html, Formal proof development.
- [DBvdW97] H. Doornbos, R. C. Backhouse, and J. van der Woude. A calculational approach to mathematical induction. *Theoretical Computer Science*, 179(1-2):103–135, 1997.
- [DHS16] B. Dongol, I. J. Hayes, and G. Struth. Convolution as a unifying concept: Applications in separation logic, interval calculi, and concurrency. *ACM TOCL*, 17(3):15:1–15:25, 2016.
- [DHS21] B. Dongol, I. J. Hayes, and G. Struth. Convolution algebras: Relational convolution, generalised modalities and incidence algebras. *Logical Methods in Computer Science*, 17(1), 2021.
- [DMS11] J. Desharnais, B. Möller, and G. Struth. Algebraic notions of termination. *Logical Methods in Computer Science*, 7(1), 2011.
- [DS11] J. Desharnais and G. Struth. Internal axioms for domain semirings. *Science of Computer Programming*, 76(3):181–203, 2011.
- [ÉB95] Zoltán Ésik and Laszlo Bernátsky. Equational properties of Kleene algebras of relations with conversion. *Theoretical Computer Science*, 137(2):237–251, 1995.
- [FJSZ21] U. Fahrenberg, C. Johansen, G. Struth, and Ziemiański. Languages of higher-dimensional automata. *Mathematical Structures in Computer Science*, 31(5):575–613, 2021.
- [FJSZ22a] U. Fahrenberg, C. Johansen, G. Struth, and K. Ziemiański. Domain semirings united. Acta Cybernetica, 25(3):575–583, 2022.
- [FJSZ22b] U. Fahrenberg, C. Johansen, G. Struth, and K. Ziemiański. A Kleene theorem for higher-dimensional automata. In CONCUR 2022, volume 243 of LIPIcs, pages 29:1–29:18. Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2022.
- [FJSZ23] U. Fahrenberg, C. Johansen, G. Struth, and K. Ziemiański. Catoids and modal convolution algebras. *Algebra Universalis*, 84:10, 2023.

- [FM22] S. Forest and S. Mimram. Rewriting in Gray categories with applications to coherence. *Mathematical Structures in Computer Science*, 32(5):574–647, 2022.
- [For21] S. Forest. Computational descriptions of higher categories. Phd thesis, Institut Polytechnique de Paris, 2021.
- [FS90] P. J. Freyd and A. Scedrov. Categories, Allegories. North-Holland, 1990.
- [Gau57] N. D. Gautam. The validity of equations in complex algebras. Archiv für mathematische Logik und Grundlagenforschung, 3:117–124, 1957.
- [GHM19] Y. Guiraud, E. Hoffbeck, and P. Malbos. Convergent presentations and polygraphic resolutions of associative algebras. *Mathematische Zeitschrift*, 293(1-2):113–179, 2019.
- [Gis88] J. L. Gischer. The equational theory of pomsets. Theoretical Computer Science, 61:199–224, 1988.
- [Giv14] S. Givant. Duality Theories for Boolean Algebras with Operators. Springer, 2014.
- [GM09] Y. Guiraud and P. Malbos. Higher-dimensional categories with finite derivation type. *Theory and Applications of Categories*, 22(18):420–478, 2009.
- [GM12] Y. Guiraud and P. Malbos. Higher-dimensional normalisation strategies for acyclicity. *Advances in Mathematics*, 231(3-4):2294–2351, 2012.
- [GM18] Y. Guiraud and P. Malbos. Polygraphs of finite derivation type. *Mathemathical Structures in Computer Science*, 28(2):155–201, 2018.
- [Gol89] R. Goldblatt. Varieties of complex algebras. Annals of Pure and Applied Logic, 44:173–242, 1989.
- [GPS95] R. Gordon, A. J. Power, and R. Street. Coherence for tricategories. *Memoirs of the American Mathematical Society*, 117(558):vi+81, 1995.
- [GW84] G. Grätzer and S. Whitney. Infinitary varieties of structures closed under the formation of complex structures. *Colloquium Mathematicum*, 48:1–5, 1984.
- [HH02] R. Hirsch and I. Hodkinson. Relation Algebras by Games. Elsevier, 2002.
- [Hig71] P. J. Higgins. Notes on Categories and Groupoids. van Nostrand Reynold Company, 1971.
- [HMSW11] T. Hoare, B. Möller, G. Struth, and I. Wehrman. Concurrent Kleene algebra and its foundations. Journal of Logic and Algebraic Programming, 80(6):266–296, 2011.
- [HST14] D Hoffmann, J Seal, G, and W. Tholen, editors. *Monoidal Topology: A Categorical Approach to Order, Metric and Topology.* Cambridge University Press, 2014.
- [JT51] B. Jónsson and A. Tarski. Boolean algebras with operators. Part I. American Journal of Mathematics, 73(4):891–939, 1951.
- [JT52] B. Jónsson and A. Tarski. Boolean algebras with operators. Part II. American Journal of Mathematics, 74(1):127–162, 1952.
- [KM15] G. Kudryavtseva and V. Mazorchuk. On multisemigroups. *Portugaliae Mathematica*, 71(1):47–80, 2015.
- [Luc17] M. Lucas. Cubical categories for homotopy and rewriting. Phd thesis, Université Paris 7, Sorbonne Paris Cité, 2017.

- [Luc20] M. Lucas. A cubical Squier's theorem. Mathematical Structures in Computer Science, 30(2):159–172, 2020.
- [Mac48] Saunders MacLane. Groups, categories and duality. PNAS, 34(6):263–267, 1948.
- [Mad06] R. D. Maddux. Relation Algebras. Elsevier, 2006.
- [ML98] S. Mac Lane. Categories for the Working Mathematician, volume 5. Springer, second edition, 1998.
- [MMS24] P. Malbos, T. Massacrier, and G. Struth. Single-set cubical categories and their formalisation with a proof assistant. arXiv:2401.10553, 2024.
- [MR23] P. Malbos and I. Ren. Shuffle polygraphic resolutions for operads. *Journal of the London Mathematical Society*, 107(1):61–122, 2023.
- [MWP92] C. J. Mulvey and J. Wick Pelletier. A quantisation of the calculus of relations. In Proceedings of the 1991 Summer Category Meeting, Montreal, Canada, volume 13 of Conference Proceedings, Canadian Mathematical Society, pages 345–360. AMS, 1992.
- [PR11] A. Palmigiano and R. Re. Groupoid quantales: A non-étale setting. *Journal of Pure and Applied Algebra*, 215(8):1945–1957, 2011.
- [Ros90] K. I. Rosenthal. Quantales and Their Applications. Longman Scientific & Technical, 1990.
- [Ros96] K. I. Rosenthal. The Theory of Quantaloids. Addison Wesley Longman Limited, 1996.
- [Ste04] R. Steiner. Omega-categories and chain complexes. *Homology, Homotopy and Applications*, 6(1):175–200, 2004.
- [Str76] R. Street. Limits indexed by category-valued 2-functors. *Journal of Pure and Applied Algebra*, 8(2):149–181, 1976.
- [Str87] R. Street. The algebra of oriented simplexes. *Journal of Pure and Applied Algebra*, 49:283–335, 1987.
- [Str06] G. Struth. Abstract abstract reduction. Journal of Logical and Algebraic Methods in Programming, 66(2):239–270, 2006.
- [Str23] G. Struth. Catoids, categories, groupoids. *Archive of Formal Proofs*, August 2023. https://isa-afp.org/entries/Catoids.html, Formal proof development.
- [Tar41] A. Tarski. On the calculus of relations. The Journal of Symbolic Logic, 6(3):73–89, 1941.
- [Ter03] Terese. Term Rewriting Systems. Cambridge University Press, 2003.

A Eckmann-Hilton-Style Collapses

Strengthening the weak homomorphism axioms $s_i(x \odot_j y) \subseteq s_i(x) \odot_j s_i(y)$ and $t_i(x \odot_j y) \subseteq t_i(x) \odot_j t_i(y)$, for $0 \le i < j < \omega$ to equations in the ω -catoid axioms collapses the structure. The equational homomorphism laws for s_j and t_j are catoid in which associativity of t_j has been forgotten. We extend this notion to t_j -streaming in the obvious way.

Lemma A.1. If the inclusions $s_i(x \odot_j y) \subseteq s_i(x) \odot_j s_i(y)$ and $t_i(x \odot_j y) \subseteq t_i(x) \odot_j t_i(y)$, for $0 \le i < j < \omega$ are replaced by equations in the axiomatisation of ω -st-multimagmas, then

1.
$$s_i = s_i$$
 and $t_i = t_i$,

2.
$$s_i = t_i$$
 and $s_j = t_j$.

Proof. For $s_i = s_j$,

$$s_i(s_j(x) \odot_j s_i(s_j(x))) = s_i(s_j(x)) \odot_j s_i(s_i(s_j(x))) = s_j(s_i(s_j(x))) \odot_j s_i(s_j(x)) = \{s_i(s_j(x))\},\$$

thus $s_i(s_i(x) \odot_i s_i(s_i(x))) \neq \emptyset$ and therefore $\Delta_i(s_i(x), s_i(s_i(x)))$. It follows that

$$s_i(x) = t_i(s_i(x)) = s_i(s_i(s_i(x))) = s_i(s_i(s_i(x))) = s_i(s_i(x)) = s_i(x).$$

By opposition, therefore, $t_i = t_j$. Also, $s_i(x) = t_j(s_i(x)) = s_i(t_j(x)) = s_i(t_i(x)) = t_i(x)$, and $s_j = t_j$ then follows from the previous properties.

Lemma A.2. If the same replacement is made for ω -categories, then $\odot_i = \odot_j$ for $0 \le i < j \le \omega$ and both operations commute.

Proof. If $x \odot_i y = \emptyset$, then $x \odot_i y \subseteq x \odot_j y$. Otherwise, if $x \odot_i y \neq \emptyset$, then $t_i(x) = s_i(y)$ and

$$\begin{aligned} \{x \odot_i y\} &= (x \odot_j t_j(x)) \odot_i (s_j(y) \odot_j y) \\ &\subseteq (x \odot_i s_j(y)) \odot_j (t_j(x) \odot_j y) \\ &= (x \odot_i s_i(y)) \odot_j (t_i(x) \odot_j y) \\ &= (x \odot_i t_i(x)) \odot_j (s_i(y) \odot_j y) \\ &= \{x \odot_j y\} \end{aligned}$$

and therefore $x \odot_i y = x \odot_i y$ by functionality.

Likewise, if $x \odot_i y = \emptyset$, then $x \odot_i y \subseteq y \odot_i x$. If $x \odot_i y \neq \emptyset$, then $t_i(x) = s_i(y)$ and

$$\begin{split} \{x\odot_i y\} &= (s_j(x)\odot_j x)\odot_i (y\odot_j t_j(x))\\ &\subseteq (s_j(x)\odot_i y)\odot_j (x\odot_i t_j(y))\\ &= (t_i(x)\odot_i y)\odot_j (x\odot_i s_i(y))\\ &= (s_i(y)\odot_i y)\odot_j (x\odot_i t_i(x))\\ &= \{y\odot_j x\}. \end{split}$$

Then $x \odot_i y = y \odot_i x$ and $x \odot_j y = y \odot_j x$ by functionality and the previous result.

Unlike the classical Eckmann-Hilton collapse, the resulting structure is *not* an abelian monoid: different elements can still have different units and \odot_i (and therefore \odot_j) need not be total. There are 2-element counterexamples for \odot_0 and \odot_1 in the case of 2-categories.

Finally we obtain a stronger collapse in the presence of an equational interchange law in the more general setting of multioperations and several units.

Lemma A.3. If the inclusions $(w \odot_j x) \odot_i (y \odot_j z) \subseteq (w \odot_i y) \odot_j (x \odot_i z)$, for $0 \le i < j < \omega$, are replaced by equations in the axiomatisation of ω -st-multimagmas, then \odot_i and \odot_j coincide, $s_i = s_j = t_i = t_j$ and \odot_i (as well as \odot_i) is associative and commutative.

We have verified this result with Isabelle in two dimensions, but leave a proof on paper to the reader. Once again, the resulting structure is not automatically an abelian monoid: different elements can have different units and \odot_i (and therefore \odot_j) need neither be total nor functional. There are again 2-element counterexamples.

B Proofs

Proof of Lemma 3.1(2)-(4). Item (2) is immediate from the axioms. For (3), suppose $xy = \{s(x)\}$. Then t(x) = s(y) and hence $\{y\} = t(x)y$. Thus $y \in x^-xy = x^-s(x) = x^-t(x^-) = x^-$ and therefore $x^- = y$. Finally, (4) is immediate from (3) because $s(x)t(x) = \{s(x)\}$ and t(x)t(x) = t(x) by Lemma 2.3(3).

Proof of Lemma 3.2. For (1), $x^-x = \{t(x)\} = \{s(x^-)\}$ implies $(x^-)^- = x$ by Lemma 3.1(2).

For (2), suppose $x \in yz$. Then t(y) = s(z) and $xz^- \subseteq yzz^- = ys(z) = yt(y) = \{y\}$. Moreover, by assumption, t(x) = t(z) and therefore $\Delta(x, z^-)$ by locality. It then follows that $xz^- = \{y\}$ and hence $y \in xz^-$. The converse implication follows from (1). The remaining equivalence follows by opposition. \square

Proof of Lemma 3.4. For (1), suppose s(x) = t(z) = s(y) and zx = zy. Then $z^-zx = z^-zy$, therefore t(z)x = t(z)y, s(x)x = s(y)y and finally x = y. (2) follows by opposition.

Proof of Lemma 5.2. (1) and (2) are immediate consequences of sup-preservation.

For (3) $\gamma \leq (\alpha \wedge \beta)^{\circ} \Leftrightarrow \gamma^{\circ} \leq \alpha \wedge \beta \Leftrightarrow \gamma^{\circ} \leq \alpha \wedge \gamma^{\circ} \leq \beta \Leftrightarrow \gamma \leq \alpha^{\circ} \wedge \gamma \leq \beta^{\circ} \Leftrightarrow \gamma \leq \alpha^{\circ} \wedge \beta^{\circ}$ implies the claim for \wedge , and the proof for \wedge is similar.

For (4), $\top^{\circ} \leq \top$ and hence $\top = \top^{\circ \circ} \leq \top^{\circ}$, and the other proofs are equally simple.

For (5), $\alpha^{\circ} \wedge \beta = \bot \Leftrightarrow (\alpha^{\circ} \wedge \beta)^{\circ} = \bot \Leftrightarrow \alpha \wedge \beta^{\circ} = \bot$.

For (6), we know that every quantale is a Kleene algebra an can use their induction axioms (see Section 12). First, $\alpha^{\circ *\circ} = (1 + \alpha^{\circ}\alpha^{\circ *})^{\circ} = 1 + \alpha\alpha^{\circ *\circ}$ and hence $\alpha^* \leq \alpha^{\circ *\circ}$ by star induction. Thus $\alpha^{*\circ} \leq \alpha^{\circ *}$. For the converse direction, $\alpha^{*\circ} = (1 + \alpha^*\alpha)^{\circ} = 1 + \alpha^{\circ}\alpha^{*\circ}$ and therefore $\alpha^{\circ *} \leq \alpha^{*\circ}$ by star induction. An alternative inductive proof uses the definition of the star in quantales.

Proof of Lemma 5.5. Let Q be an involutive quantale. Suppose the modular law holds. We derive two auxiliary properties before deriving the Dedekind law. First, the modular law can be strengthened to

$$\alpha\beta \wedge \gamma = (\alpha \wedge \gamma\beta^{\circ})\beta \wedge \gamma,$$

because $\alpha\beta \wedge \gamma = \alpha\beta \wedge \gamma \wedge \gamma \leq (\alpha \wedge \gamma\beta^{\circ})\beta \wedge \gamma$ using the modular law and $(\alpha \wedge \gamma\beta^{\circ})\beta \wedge \gamma \leq \alpha\beta \wedge \gamma$ by properties of infs and order preservation. Second, a dual equational modular law

$$\alpha\beta \wedge \gamma = \alpha(\beta \wedge \alpha^{\circ}\gamma) \wedge \gamma$$

then follows because $\alpha\beta \wedge \gamma = (\beta^{\circ}\alpha^{\circ} \wedge \gamma^{\circ})^{\circ} = ((\beta^{\circ} \wedge \gamma^{\circ}\alpha^{\circ\circ})\alpha^{\circ} \wedge \gamma^{\circ})^{\circ} = \alpha(\beta \wedge \alpha^{\circ}\gamma) \wedge \gamma$ using properties of converse and the first equational modular law in the second step. Therefore,

$$\alpha\beta \wedge \gamma = \alpha(\beta \wedge \alpha^{\circ}\gamma) \wedge \gamma$$

$$\leq (\alpha \wedge \gamma(\beta \wedge \alpha^{\circ}\gamma)^{\circ})(\beta \wedge \alpha^{\circ}\gamma)$$

$$= (\alpha \wedge \gamma(\beta^{\circ} \wedge \gamma^{\circ}\alpha))(\beta \wedge \alpha^{\circ}\gamma)$$

$$\leq (\alpha \wedge \gamma\beta^{\circ})(\beta \wedge \alpha^{\circ}\gamma),$$

using properties of converse and order preservation as well as the dual equational modular in the first and the modular law in the second step. This proves the Dedekind law.

Finally, $\alpha\beta \wedge \gamma \leq (\alpha \wedge \gamma\beta^{\circ})(\beta \wedge \alpha^{\circ}\gamma) \leq (\alpha \wedge \gamma\beta^{\circ})\beta$ yields the modular law from the Dedekind law using properties of inf and order preservation.

These proofs are standard in relation algebra.

Proof of Lemma 5.6. For the strong Gelfand property, $\alpha = 1\alpha \wedge \alpha \leq \top \alpha \wedge \alpha \leq (\top \wedge \alpha \alpha^{\circ})\alpha = \alpha \alpha^{\circ} \alpha$ using the modular law in the third step.

For Peirce's law, suppose $\alpha\beta \wedge \gamma^{\circ} = \bot$. Then $(\alpha\beta \wedge \gamma^{\circ})\beta^{\circ} \wedge \alpha = \bot$ and therefore $\beta\gamma \wedge \alpha^{\circ} = \bot$ using the first equational modular law from the proof of Lemma 5.5 and properties of convolution. The converse implication is similar, using the dual equational modular law from Lemma 5.5.

For the first Schröder law, $\alpha\beta \wedge \gamma = \bot \Leftrightarrow \beta \wedge \alpha^{\circ}\gamma = \bot$, $\alpha\beta \wedge \gamma = \bot$ implies $\beta^{\circ}\alpha^{\circ} \wedge \gamma^{\circ} = \bot$ by properties of converse and therefore $\beta \wedge \alpha^{\circ}\gamma = \bot$ by Peirce's law. The proof of the converse direction is similar.

The second Schröder law, $\alpha\beta \wedge \gamma = \bot \Leftrightarrow \alpha \wedge \gamma\beta^{\circ} = \bot$ follows in a similar way from Peirce's law, the first Schröder law and properties of converse.

These proofs are once again standard in relation algebra.

Proof of Lemma 5.7. For (1), $1 \wedge \alpha \top = 1 \wedge \alpha(\top \wedge \alpha^{\circ}1) = 1 \wedge \alpha\alpha^{\circ} = dom(\alpha)$ using the modular law.

Item (2) is trivial, and for (3), $\alpha = \alpha \wedge \alpha 1 \leq (1 \wedge \alpha \alpha^{\circ})\alpha = dom(\alpha)\alpha$ by the modular law, and the converse inequality follows from (2).

```
For (4), \alpha \top = dom(\alpha)\alpha \top \leq dom(\alpha)\top \top = dom(\alpha)\top and dom(\alpha)\top = (1 \land \alpha\top)\top \leq \top \land \alpha\top\top = \alpha\top.
```

Items (5)-(7) are trivial. For (8), $dom(\alpha dom(\beta)) = 1 \land \alpha dom(\beta) \top = 1 \land \alpha \beta \top = dom(\alpha \beta)$.

For (9), note that $dom(dom(\alpha)) = dom(\alpha)$ is immediate from (8). We first show that $dom(dom(\alpha) \vee dom(\beta)) = dom(\alpha) \vee dom(\beta)$. Indeed,

$$dom(dom(\alpha) \lor dom(\beta)) = 1 \land (dom(\alpha) \lor dom(\beta))(dom(\alpha) \lor dom(\beta))^{\circ}$$

$$= 1 \land (dom(\alpha) \lor dom(\beta))(dom(\alpha) \lor dom(\beta))$$

$$= 1 \land (dom(\alpha) \lor dom(\beta)) \land (dom(\alpha) \lor dom(\beta))$$

$$= 1 \land (dom(\alpha) \lor dom(\beta))$$

$$= (1 \land dom(\alpha)) \lor (1 \land dom(\beta))$$

$$= dom(\alpha) \lor dom(\beta),$$

where the third step uses the fact that multiplication of domain elements is commutative and idempotent. Proofs can be found in our Isabelle theories. Using this property with (1) and (4),

$$\begin{split} dom(\alpha \vee \beta) &= 1 \wedge (\alpha \vee \beta) \top \\ &= 1 \wedge (\alpha \top \vee \beta \top) \\ &= 1 \wedge (dom(\alpha) \top \vee dom(\beta) \top)) \\ &= 1 \wedge (dom(\alpha) \vee dom(\beta)) \top \\ &= dom(dom(\alpha) \vee dom(\beta)) \\ &= dom(\alpha) \vee dom(\beta). \end{split}$$

For (10) note that, by the strong Gelfand property, (2) and (7), $dom(\alpha)dom(\alpha)dom(\alpha) = dom(\alpha)$, therefore $dom(\alpha)dom(\alpha) = dom(\alpha)$ and likewise for $cod(\alpha)$. The proof of $cod(dom(\alpha)) = dom(\alpha)$ is similar. Finally, (11) is immediate from the definition of dom.

Proof of Proposition 5.8. Immediate from Lemma 5.7.

Proof of Lemma 5.9. By the strong Gelfand property,

$$dom(\alpha) \leq dom(\alpha)dom(\alpha)^{\circ}dom(\alpha) \leq 1dom(\alpha)^{\circ}1 = dom(\alpha)^{\circ},$$

from which $dom(\alpha)^{\circ} \leq dom(\alpha)$ follows using the adjunction in Remark 5.1. The proof for *cod* follows by duality.

Proof of Lemma 6.8. By the morphism axiom for s_i , $s_i(x \odot_j y) \subseteq s_i(x) \odot_j s_i(y)$. The right-hand side must not be empty whenever the left-hand side is defined, which is assumed. Hence it must be equal to $\{s_i(x)\}$ because compositions of lower cells in higher dimensions are trivial.

Proof of Proposition 6.13. Before the proof proper, we derive $s_i \circ t_j \circ s_i = s_i$ and $t_i \circ s_j \circ s_i = s_i$. We have

$$\begin{aligned} \{s_i(x)\} &= s_i(x) \odot_i s_i(x) \\ &= (s_j(s_i(x)) \odot_j s_i(x)) \odot_i (s_i(x) \odot_j t_j(s_i(x))) \\ &\subseteq (s_i(s_i(x)) \odot_i s_i(x)) \odot_i (s_i(x) \odot_i t_i(s_i(x))), \end{aligned}$$

hence $\Delta_j(s_j(s_i(x)) \odot_i s_i(x), s_i(x) \odot_i t_j(s_i(x)))$. Thus $\Delta_i(s_j(s_i(x)), s_i(x))$ and $\Delta_i(s_i(x), t_j(s_i(x)))$, and therefore $t_i(s_j(s_i(x))) = s_i(s_i(x)) = s_i(x)$ as well as $s_i(t_j(s_i(x))) = t_i(s_i(x)) = s_i(x)$.

Next we derive the missing n-catoid axioms. First we consider $s_j \circ s_i = s_i$, $s_j \circ t_i = t_i$, $t_j \circ s_i = s_i$, and $t_j \circ t_i = t_i$. For the first one,

$$\{s_{i}(x)\} = s_{i}(x) \odot_{i} s_{i}(x)$$

$$= (s_{j}(s_{i}(x)) \odot_{j} s_{i}(x)) \odot_{i} (s_{i}(x) \odot_{j} t_{j}(s_{i}(x)))$$

$$\subseteq (s_{j}(s_{i}(x)) \odot_{i} s_{i}(x)) \odot_{j} (s_{i}(x) \odot_{i} t_{j}(s_{i}(x)))$$

$$= (s_{j}(s_{i}(x)) \odot_{i} t_{i}(s_{j}(s_{i}(x)))) \odot_{j} (s_{i}(t_{j}(s_{i}(x))) \odot_{i} t_{i}(t_{j}(s_{i}(x))))$$

$$= s_{j}(s_{i}(x)) \odot_{j} t_{j}(s_{i}(x))$$

and therefore $s_i(x) = s_i(s_i(x))$ as well as $s_i(x) = t_i(s_i(x))$. The remaining identities hold by opposition.

Second, we derive $s_i \circ s_j = s_j \circ s_i$, $s_i \circ t_j = t_j \circ s_i$, $t_i \circ s_i = s_j \circ t_i$ and $t_i \circ t_j = t_j \circ t_i$. For the first identity, $\{s_j(x)\} = s_j(s_i(x)) \odot_i s_j(x) = s_i(x) \odot_i s_j(x)$ and therefore $\Delta_i(s_i(x), s_j(x))$. It follows that $s_j(s_i(x)) = s_i(x) = t_i(s_i(x)) = s_i(s_j(x))$. The remaining proofs are similar.

Note that none of the proofs so far require an associativity law.

It remains to derive $s_i(x \odot_j y) \subseteq s_i(x) \odot_j s_i(y)$ and $t_i(x \odot_j y) \subseteq t_i(x) \odot_j t_i(y)$. We prove the first inclusion by cases. If $s_i(x \odot_j y) = \emptyset$, then the claim is trivial. Otherwise, if $s_i(x \odot_j y) \neq \emptyset$, then $t_j(x) = s_i(y)$ and thus

$$s_{i}(x \odot_{j} y) = s_{i}(s_{j}(x \odot_{j} y))$$

$$= s_{i}(s_{j}(x \odot_{j} s_{j}(y)))$$

$$= s_{i}(s_{j}(x \odot_{j} t_{j}(y)))$$

$$= s_{i}(s_{j}(x))$$

$$= \{s_{i}(x)\}$$

$$= s_{i}(x) \odot_{j} t_{j}(s_{i}(x))$$

$$= s_{i}(x) \odot_{j} s_{i}(t_{j}(x))$$

$$= s_{i}(x) \odot_{j} s_{i}(s_{j}(x))$$

$$= s_{i}(x) \odot_{i} s_{i}(x).$$

This uses weak locality laws that are available in all catoids if the composition under the source operation is defined. See Lemma 2.6 and [FJSZ23] for details. The second inclusion follows by opposition. \Box

Proof of Lemma 7.3. For (1) $cod_j \circ dom_i = cod_j \circ dom_j \circ dom_i = dom_j \circ dom_i = dom_i$, and the remaining identities in (1) follow by opposition.

For (2),
$$1_j \cdot_i 1_j = d_j(1_j) \cdot_i d_j(1_j) = d_j(1_j \cdot_i 1_j) \le 1_j$$
 if Q is strong and

$$1_{j} = 1_{j} \cdot_{i} 1_{i} = (1_{j} \cdot_{j} 1_{j}) \cdot_{i} (1_{i} \cdot_{j} 1_{j}) \leq (1_{j} \cdot_{i} 1_{i}) \cdot_{j} (1_{j} \cdot_{i} 1_{j}) = 1_{j} \cdot_{j} (1_{j} \cdot_{i} 1_{j}) = 1_{j} \cdot_{i} 1_{j}.$$

Further, $1_i \cdot_i 1_i = d_i(1_i) \cdot_i d_i(1_i) = d_i(d_i(1_i)) \cdot_i d_i(1_i) = d_i(1_i) = 1_i$.

For (3), $1_i = 1_i \cdot_i 1_i = (1_j \cdot_j 1_i) \cdot_i (1_i \cdot_j 1_i) \le (1_j \cdot_i 1_i) \cdot_j (1_i \cdot_i 1_j) = 1_j \cdot_j 1_j = 1_j$.

For (4), $d_j(1_i) = d_j(d_i(1_i)) = d_i(1_i) = 1_i$, and $d_i(1_j) \le 1_i$ as well as $1_i = d_i(1_i) \le d_i(1_j)$. The other identities in (4) then follow by opposition.

For (5), $dom_i \circ dom_j = dom_j \circ dom_i$ follows from

$$\begin{split} dom_i(dom_j(\alpha)) &= dom_i(dom_j(dom_i(\alpha) \cdot_i \alpha)) \\ &= dom_i(dom_j(dom_i(\alpha)) \cdot_i dom_j(\alpha)) \\ &= dom_i(dom_i(\alpha) \cdot_i dom_j(\alpha)) \\ &= dom_i(\alpha) \cdot_i dom_i(dom_j(\alpha)) \\ &= dom_j(dom_i(\alpha)) \cdot_i dom_i(dom_j(\alpha)) \\ &\leq dom_j(dom_i(\alpha)) \cdot_i 1_i \\ &= dom_i(dom_i(\alpha)) \end{split}$$

and

$$dom_{j}(dom_{i}(\alpha)) = dom_{i}(\alpha)$$

$$= dom_{i}(dom_{j}(\alpha) \cdot_{j} \alpha)$$

$$\leq dom_{i}(dom_{j}(\alpha)) \cdot_{j} dom_{i}(\alpha)$$

$$\leq dom_{i}(dom_{j}(\alpha)) \cdot_{j} 1_{i}$$

$$\leq dom_{i}(dom_{j}(\alpha)) \cdot_{j} 1_{j}$$

$$= dom_{i}(dom_{j}(\alpha)).$$

Moreover, $dom_i \circ cod_j = cod_j \circ dom_i$ follows from

$$\begin{aligned} dom_i(cod_j(\alpha)) &= dom_i(cod_j(dom_i(\alpha) \cdot_i \alpha)) \\ &= dom_i(cod_j(dom_i(\alpha)) \cdot_i cod_i(\alpha)) \\ &= dom_i(dom_i(\alpha) \cdot_i cod_i(\alpha)) \\ &= dom_i(\alpha) \cdot_i dom_i(cod_i(\alpha)) \\ &= cod_j(dom_i(\alpha)) \cdot_i dom_i(cod_i(\alpha)) \\ &\leq cod_j(dom_i(\alpha)) \cdot_i 1_i \\ &= cod_j(dom_i(\alpha)) \end{aligned}$$

and

$$cod_{j}(dom_{i}(\alpha)) = dom_{i}(\alpha)$$

$$= dom_{i}(\alpha \cdot_{j} cod_{j}(\alpha))$$

$$\leq dom_{i}(\alpha) \cdot_{j} dom_{i}(cod_{j}(\alpha))$$

$$\leq 1_{i} \cdot_{j} dom_{i}(cod_{j}(\alpha))$$

$$\leq 1_{j} \cdot_{j} dom_{i}(cod_{j}(\alpha))$$

$$= dom_{i}(cod_{j}(\alpha)).$$

The remaining two identities in (6) then follow by opposition.

Finally, for (6), $dom_i(\alpha \cdot_j \beta) = dom_i(dom_j(\alpha \cdot_j \beta)) = dom_i(dom_j(\alpha \cdot_j dom_j(\beta))) = dom_i(\alpha \cdot_j dom_j(\beta))$ and the second identity in (7) then follows by opposition.

Proof of Lemma 7.6. For the first property in (1) and k = i, we first use induction on k to prove $dom_i(\alpha) \cdot_i \beta^{k_1} \leq (dom_i(\alpha) \cdot_i \beta)^{k_j}$, where $(-)^{k_j}$ indicates that powers are taken with respect to \cdot_j . The base case follows from

$$dom_i(\alpha) \cdot_i 1_j = dom_j(dom_i(\alpha)) \cdot_i 1_j \leq 1_j \cdot_i 1_j = 1_j$$
.

For the induction step, suppose $dom_i(\alpha) \cdot_i \beta^{k_j} \leq (dom_i(\alpha) \cdot_i \beta)^{k_j}$. Then

$$\begin{split} dom_{i}(\alpha) \cdot_{i} \beta^{(k+1)_{j}} &= dom_{j}(dom_{i}(\alpha)) \cdot_{i} (\beta \cdot_{j} \beta^{k_{j}}) \\ &= (dom_{j}(dom_{i}(\alpha)) \cdot_{j} dom_{j}(dom_{i}(\alpha))) \cdot_{i} (\beta \cdot_{j} \beta^{k_{j}}) \\ &= (dom_{i}(\alpha) \cdot_{j} dom_{i}(\alpha)) \cdot_{i} (\beta \cdot_{j} \beta^{k_{j}}) \\ &\leq (dom_{i}(\alpha) \cdot_{i} \beta) \cdot_{j} (dom_{i}(\alpha) \cdot_{i} \beta^{k_{j}}) \\ &\leq (dom_{i}(\alpha) \cdot_{i} \beta) \cdot_{j} (dom_{i}(\alpha) \cdot_{i} \beta)^{k_{j}} \\ &= (dom_{i}(\alpha) \cdot_{i} \beta)^{(k+1)_{j}}. \end{split}$$

Using this property,

$$dom_{i}(\alpha) \cdot_{i} \beta^{*_{j}} = dom_{i}(\alpha) \cdot_{i} \bigvee_{k \geq 0} \beta^{k_{j}}$$

$$= \bigvee \{ dom_{i}(\alpha) \cdot_{i} \beta^{k_{j}} \mid k \geq 0 \}$$

$$\leq \bigvee_{k \geq 0} (dom_{i}(\alpha) \cdot_{i} \beta)^{k_{j}}$$

$$= (dom_{i}(\alpha) \cdot_{i} \beta)^{*_{j}}.$$

The proof of the second property in (1) for k = i is dual. The proofs for k = j are very similar, but in the base case, $dom_j(x) \cdot_i 1_j \leq 1_j \cdot_i 1_j$ needs to be shown, which follows from Lemma 7.3(2) and needs a strong ω -quantale.

For (2), we first use induction on k to show that $(\alpha \cdot_j \beta)^{k_i} \leq \alpha^{k_i} \cdot_j \beta^{k_i}$. In the base case, $1_i = 1_i \cdot_j 1_i$. For the induction step, suppose $(\alpha \cdot_j \beta)^{k_i} \leq \alpha^{k_i} \cdot_j \beta^{k_i}$. Then

$$(\alpha \cdot_{i} \beta)^{(k+1)_{i}} = (\alpha \cdot_{i} \beta) \cdot_{i} (\alpha \cdot_{i} \beta)^{k_{i}} \leq (\alpha \cdot_{i} \beta) \cdot_{i} \alpha^{k_{i}} \cdot_{j} \beta^{k_{i}} \leq (\alpha \cdot_{i} \alpha^{k_{i}}) \cdot_{j} (\beta \cdot_{i} \beta^{k_{i}}) = \alpha^{k+1_{i}} \cdot_{j} \beta^{(k+1)_{i}}.$$

Using this property, $(\alpha \cdot_j \beta)^{k_i} \leq \alpha^{*_i} \cdot_j \beta^{*_i}$ for all $k \geq 0$ and thus $(\alpha \cdot_j \beta)^{*_i} \leq \alpha^{*_i} \cdot_j \beta^{*_i}$ by properties of sup.

Proof of Lemma 7.7. For (1), $dom_0(\alpha) \cdot_1 dom_0(\alpha) = dom_1(dom_0(\alpha)) \cdot_1 dom_1(dom_0(\alpha)) = dom_1(dom$

For (2), $dom_i(\alpha \cdot_j \beta) = dom_i(dom_j(\alpha \cdot_j \beta)) = dom_i(dom_j(\alpha \cdot_j dom_j(\beta))) = dom_j(\alpha \cdot_j dom_j(\beta))$, and the proof for codomain follows by opposition.

The proofs of (3) are similar to those of (2), inserting cod_j instead of dom_j in the first step. For (4),

$$dom_{i}(\alpha \cdot_{i} \beta) = dom_{i}(\alpha \cdot_{i} dom_{i}(\beta))$$

$$= dom_{i}(\alpha \cdot_{i} dom_{j}(dom_{i}(\beta)))$$

$$= dom_{i}(\alpha \cdot_{i} dom_{i}(dom_{j}(\beta)))$$

$$= dom_{i}(\alpha \cdot_{i} dom_{j}(\beta))$$

and the proof for codomain follows by opposition.

For (5),

$$dom_{i}(x \cdot_{i} y) = dom_{i}(cod_{j}(x \cdot_{i} y))$$

$$\leq dom_{i}(cod_{j}(x) \cdot_{i} cod_{j}(y))$$

$$= dom_{i}(cod_{j}(x) \cdot_{i} dom_{i}(cod_{j}(y)))$$

$$= dom_{i}(cod_{j}(x) \cdot_{i} dom_{i}(y))$$

$$= dom_{i}(cod_{j}(x) \cdot_{i} y).$$

The second step uses the morphism law for cod_j . If the quantale is strong, it can be replaced by an equational step. The proofs for cod_i follow by opposition.

Item (6) is immediate from (1) and interchange.

Items (7) and (8) are immediate consequences of the weak homomorphism laws and the homomorphism laws, respectively.

For (9),

```
\begin{split} dom_{i}(\alpha) \cdot_{j} dom_{i}(\beta) &= dom_{i}(dom_{i}(\alpha) \cdot_{j} dom_{i}(\beta)) \cdot_{0} (dom_{i}(\alpha) \cdot_{j} dom_{i}(\beta)) \\ &\leq (dom_{i}(dom_{i}(\alpha)) \cdot_{j} dom_{i}(dom_{i}(\beta)))) \cdot_{0} (dom_{i}(\alpha) \cdot_{j} dom_{i}(\beta)) \\ &\leq (dom_{i}(\alpha) \cdot_{j} 1_{i}) \cdot_{0} (1_{i} \cdot_{j} dom_{i}(\beta)) \\ &\leq (dom_{i}(\alpha) \cdot_{j} 1_{j}) \cdot_{0} (1_{i} \cdot_{j} dom_{j}(\beta)) \\ &\leq dom_{i}(\alpha) \cdot_{i} dom_{i}(\beta), \end{split}
```

where the first step uses domain absorption, the second a weak homomorphism law, and the remaining steps are straightforward approximations. For the converse direction,

```
dom_{i}\alpha) \cdot_{i} dom_{i}(\beta) = (dom_{i}(\alpha) \cdot_{j} dom_{i}(\alpha)) \cdot_{i} (dom_{i}(\beta) \cdot_{j} dom_{i}(\beta))
\leq (dom_{i}(\alpha) \cdot_{i} dom_{i}(\beta)) \cdot_{j} (dom_{i}(\alpha) \cdot_{i} dom_{i}(\beta))
\leq (dom_{i}(\alpha) \cdot_{i} 1_{i}) \cdot_{j} (1_{i} \cdot_{i} dom_{i}(\beta))
= dom_{i}(\alpha) \cdot_{j} dom_{i}(\beta),
```

where the first step uses (1), the second step the interchange law, and the remaining steps are obvious. For (10), using in particular (9),

```
 (dom_i(\alpha) \cdot_j dom_i(\beta)) \cdot_i (dom_i(\gamma) \cdot_j dom_i(\delta)) = (dom_i(\alpha) \cdot_i dom_i(\beta)) \cdot_i (dom_i(\gamma) \cdot_i dom_i(\delta)) 
= (dom_i(\alpha) \cdot_i dom_i(\gamma)) \cdot_i (dom_i(\beta) \cdot_i dom_i(\delta)) 
= dom_i (dom_i(\alpha) \cdot_i dom_i(\gamma)) \cdot_i dom_i (dom_i(\beta) \cdot_i dom_i(\delta)) 
= dom_i (dom_i(\alpha) \cdot_i dom_i(\gamma)) \cdot_j dom_i (dom_i(\beta) \cdot_i dom_i(\delta)) 
= (dom_i(\alpha) \cdot_i dom_i(\gamma)) \cdot_j (dom_i(\beta) \cdot_i dom_i(\delta)).
```

Proof of Corollary 10.1. For (1), we show explicit proofs for the globular structure, starting with the interchange laws. For $W, X, Y, Z \subseteq C$,

```
a \in (W \odot_j X) \odot_i (Y \odot_j Z) \Leftrightarrow \exists w \in W, x \in X, y \in Y, z \in Z. \ a \in (w \odot_j x) \odot_i (y \odot_j z)\Rightarrow \exists w \in W, x \in X, y \in Y, z \in Z. \ a \in (w \odot_i y) \odot_j (x \odot_i z)\Leftrightarrow a \in (W \odot_i X) \odot_j (Y \odot_i Z).
```

This requires only the interchange law in C. It remains to lift the globular laws.

For $dom_j(x \cdot_i y) \leq dom_j(x) \cdot_i dom_j(y)$ and $X, Y \subseteq X$,

```
a \in s_{j}(X \odot_{i} Y) \Leftrightarrow \exists b \in X \odot_{i} Y. \ a = s_{j}(b)
\Leftrightarrow \exists b, c \in X, d \in Y. \ a = s_{j}(b) \land b \in c \odot_{i} d
\Leftrightarrow \exists c \in X, d \in Y. \ a \in s_{j}(c \odot_{i} d)
\Rightarrow \exists c \in X, d \in Y. \ a \in s_{j}(c) \odot_{i} s_{j}(d)
\Leftrightarrow \exists c \in s_{j}(X), d \in s_{j}(Y). \ a \in c \odot_{i} d
\Leftrightarrow a \in s_{i}(X), \cdot i s_{j}(Y).
```

The implication in the fourth step can be replaced by \Leftrightarrow if C is an ω -category. The proofs of $cod_j(x \cdot_i y) \leq cod_j(x) \cdot_i cod_j(y)$ and its strong variant follows by opposition. These proofs require only the respective morphism laws in C.

For $dom_i(x \cdot_j y) \leq dom_i(x) \cdot_j dom_i(y)$, note that $s_i(x \odot_j y) \subseteq s_i(x) \odot_j s_i(y)$ is derivable even in the reduced axiomatisation of *n*-catoids (Proposition 6.13). The proof is then similar to the previous one.

Finally, for $dom_j \circ dom_i = dom_i$, note that $s_j \circ s_i = s_i$ is derivable in the reduced axiomatisation n-catoids (Proposition 6.13). Hence let $X \subseteq C$. Then

$$a \in s_i(s_i(X)) \Leftrightarrow \exists b \in X. \ a = s_i(s_i(b)) \Leftrightarrow \exists b \in X. \ a = s_i(b) \Leftrightarrow a \in s_i(X).$$

For (2), we consider the morphism laws for s_i and t_i . We consider these correspondences one-by-one between quantales and catoids and thus mention the laws for s_i and t_i for the context without locality. As for Theorem 8.5, we start with an explicit proof of interchange for multioperations and powersets.

$$(w \odot_j x) \odot_i (y \odot_j z) = (\{w\} \cdot_j \{x\}) \cdot_i (\{y\} \cdot_j \{z\}) \subseteq (\{w\} \cdot_i \{y\}) \cdot_j (\{x\} \cdot_i \{z\}) = (w \odot_i y) \odot_j (x \odot_i z).$$

For the morphism law for s_i ,

$$s_j(x\odot_i y) = dom_j(\{x\} \cdot_i \{y\}) \subseteq dom_j(\{x\}) \cdot_i dom_j(\{y\}) = s_j(x) \odot_i s_j(y).$$

The inclusion in the second step becomes an equality if Q is strong. The proofs for the morphism law for t_j are dual. The proofs of the laws for s_i and t_i are very similar.

Proof of Lemma 10.3. The proofs of (1) and (2) are obvious; (3) is immediate from (2): $X = s(X)X \subseteq XX^{-}X$. For (4),

$$a \in C_0 \cap XX^- \Leftrightarrow s(a) = a \land \exists b \in X, c \in X^- \ s(a) \in b \odot c$$

 $\Leftrightarrow s(a) = a \land \exists b \in X. \ s(a) = s(b)$
 $\Leftrightarrow a \in s(X)$

and the proof for t is dual. The proofs for (5) are very similar:

$$a \in C_0 \cap X \top \Leftrightarrow s(a) = a \land \exists b \in X, c \in C. \ s(a) \in b \odot c$$

$$\Leftrightarrow s(a) = a \land \exists b \in X. \ s(a) = s(b)$$

$$\Leftrightarrow a \in s(X)$$

and likewise for t. For (6), $X \top = s(X)X \top \subseteq s(X)\top \top \subseteq s(X)\top \top$ and $s(X)\top \subseteq XX^{-}\top \subseteq X\top \top \subseteq X\top$. \square

Proof of Lemma 11.2. The equations in (1) follow immediately from Lemma 7.7(2)-(5).

For (2), $dom_i(\alpha \cdot_i dom_j(\beta \cdot_j \gamma)) \leq dom_i(\alpha \cdot_i (dom_i(\beta) \cdot_j dom_i(\gamma)))$ follows from the above using the weak morphism axiom, and a dual property holds for cod_i and cod_j .

For (3),

```
\begin{split} dom_{i}(dom_{k}(\alpha) \cdot_{i} dom_{j}(\beta \cdot_{j} \gamma)) & \leq dom_{i}(dom_{k}(\alpha) \cdot_{i} (dom_{i}(\beta) \cdot_{j} dom_{i}(\gamma))) \\ & \leq dom_{i}((dom_{k}(\alpha) \cdot_{i} dom_{i}(\beta)) \cdot_{j} (dom_{k}(\alpha) \cdot_{i} dom_{i}(\gamma))) \\ & \leq dom_{i}(dom_{k}(\alpha) \cdot_{i} dom_{i}(\beta)) \cdot_{j} dom_{i}(dom_{k}(\alpha) \cdot_{i} dom_{i}(\gamma)) \\ & = dom_{i}(dom_{k}(\alpha) \cdot_{i} \beta) \cdot_{j} dom_{i}(dom_{k}(\alpha) \cdot_{i} \gamma). \end{split}
```

The first step uses an approximation from (2), the second Lemma 7.7(6), the third a weak morphism axiom and the last domain locality. The proof for codomains is similar.

For (4),

$$dom_{j}(\alpha \cdot_{j} dom_{j}(dom_{i}(\beta \cdot_{i} \gamma))) = dom_{j}(\alpha \cdot_{j} dom_{i}(dom_{j}(\beta \cdot_{i} \gamma)))$$

$$\leq dom_{j}(\alpha \cdot_{j} dom_{i}(dom_{j}(\beta) \cdot_{i} dom_{j}(\gamma)))$$

$$= dom_{j}(\alpha \cdot_{j} dom_{i}(dom_{j}(\beta) \cdot_{i} \gamma)).$$

The first step uses Lemma 7.3(5), the second applies an morphism axiom, the third 7.7(4). The second step becomes an equality if Q is strong. The remaining proofs are similar. For (5),

```
dom_0(\gamma) \cdot_0 dom_1(\alpha \cdot_1 \beta) = dom_1(dom_0(\gamma)) \cdot_0 dom_1(\alpha \cdot_1 \beta)
= dom_1(dom_0(\gamma) \cdot_0 (\alpha \cdot_1 \beta))
\leq dom_1((dom_0(\gamma) \cdot_0 \alpha) \cdot_1 (dom_0(\gamma) \cdot_0 \beta))
```

The first step uses an axiom, the second a strong morphism axiom and the third Lemma 7.7(6). The second proof is similar.

Proof of Lemma 12.4. For (1), $dom(\alpha) \leq dom(\alpha)dom(\alpha)^{\circ}dom(\alpha) \leq dom(\alpha)^{\circ}$, and thus $dom(\alpha)^{\circ} \leq dom(\alpha)^{\circ\circ} = dom(\alpha)$ as converse is order preserving. The property for cod then follows by opposition. For (2),

```
dom(\alpha^{\circ}) = dom((\alpha cod(\alpha))^{\circ})
= dom(cod(\alpha)^{\circ}x^{\circ})
= dom(cod(\alpha)\alpha^{\circ})
= cod(\alpha)dom(\alpha^{\circ})
= cod(\alpha dom(\alpha^{\circ}))
= cod(\alpha^{\circ\circ}dom(\alpha^{\circ})^{\circ})
= cod((dom(\alpha^{\circ})\alpha^{\circ})^{\circ})
= cod(\alpha^{\circ\circ})
= cod(\alpha).
```

The export steps in the fourth and fifth lines of the proof work because of the compatibility laws $dom \circ cod = cod$ and $cod \circ dom = dom$ of modal semirings. The second property then follows by opposition.

Proof of Lemma 12.7. It suffices to show that $1_i + (\alpha \cdot_j \beta) \cdot_i (\alpha^{*i} \cdot_j \beta^{*i}) \leq (\alpha^{*i} \cdot_j \beta^{*i})$ by star induction. Then $1_i \leq (\alpha^{*i} \cdot_j \beta^{*i})$ hold because $1_i = 1_i \cdot_j 1_i$ and $1_i \leq \alpha^{*i}$, $1_i \leq \beta^{*i}$ by standard Kleene algebra. Also, $(\alpha \cdot_j \beta) \cdot_i (\alpha^{*i} \cdot_j \beta^{*i}) \leq (\alpha^{*i} \cdot_j \beta^{*i})$ by interchange and $\alpha \cdot_i \alpha^{*i} \leq \alpha^{*i}$ and $\beta \cdot_i \beta^{*i} \leq \beta^{*i}$, again by standard Kleene algebra.

Proof of Lemma 12.11. First, $\alpha^{\circ *\circ} = (1 + \alpha^{\circ *}\alpha^{\circ})^{\circ} = 1 + \alpha \cdot \alpha^{\circ *\circ}$ by star unfold and properties of involution. Thus $\alpha^* \leq \alpha^{\circ *\circ}$ by star induction and therefore $\alpha^{*\circ} \leq \alpha^{\circ *}$.

Second, $\alpha^{*\circ} = (1 + \alpha \alpha^*)^{\circ} = 1 + \alpha^{\circ} \alpha^{*\circ}$ by star unfold and properties of involution. Thus $\alpha^{\circ *} \leq \alpha^{*\circ}$ by star induction.