

Bremsstrahlung Cross Section with Polarized Beams for Luminosity Determination at the EIC

Dhevan Gangadharan*
University of Houston
(Dated: August 1, 2023)

The bremsstrahlung cross section is calculated at leading order for polarized beams of electrons and ions, which is needed for luminosity measurements at the upcoming Electron Ion Collider (EIC). Analytic expressions, differential in the emitted photon energy and polar angle, are derived. The component of the cross section which depends on the beam polarizations is found to be highly suppressed with respect to the unpolarized Bethe-Heitler component, owing to the low q^2 that characterizes the bremsstrahlung process.

I. INTRODUCTION

The dominant contribution to the inelastic cross section in lepton-nucleus collisions is the bremsstrahlung process, where a photon is emitted into the final state: $e^- + N \rightarrow e^- + N + \gamma$. Owing to the QED nature of this process, it can be calculated to high precision, which allowed its use to measure the collider luminosity of HERA [1, 2]. The upcoming Electron Ion Collider (EIC) will similarly collide electrons on protons, as well as heavier ions, and will also make use of the bremsstrahlung process to measure luminosity. Unlike HERA, the EIC will accelerate polarized electrons *and* polarized nuclei. As the leading-order cross section only depends on the beam polarization if *both* beams are polarized, the EIC requires calculation of the polarized component of the cross section. The process has been considered before and numerically calculated for q^2 larger than those relevant for luminosity programs [3, 4]. Analytic expressions are presented here for appropriately low q^2 .

II. FORMALISM

The two Feynman diagrams that contribute to the leading-order bremsstrahlung amplitude are shown in Fig. 1. The corresponding unpolarized cross section was




FIG. 1. Leading-order bremsstrahlung Feynman diagrams for ep scattering. Incoming momenta denoted by p (electron) and r (proton). Outgoing momenta denoted by p' (scattered electron), r' (scattered proton), and k (emitted photon).

first calculated by Bethe and Heitler in 1934 [5]. A more complete calculation of the cross section that goes beyond the Born approximation (leading order) and is applicable for high atomic number Z , was made by Bethe and Maximon in 1954 [6–8]. Higher-order contributions enter at the subpercent level for ep scattering [9]. The amplitude shown in Fig. 1 is given by Eq. 1:

$$M = e^3 \frac{g_{\mu\nu}}{q^2} \epsilon_\sigma^*(k) [\bar{u}(r') \gamma^\mu u(r)] \times \quad (1)$$

$$\bar{u}(p') \left[\gamma^\sigma \frac{\not{p}' + \not{k} + m_e}{2p'k} \gamma^\nu - \gamma^\nu \frac{\not{p} - \not{k} + m_e}{2pk} \gamma^\sigma \right] u(p).$$

Incoming 4-momenta (energy, momentum) in the laboratory frame are denoted as $p = (\varepsilon, \mathbf{p})$ (electron) and $r = (\varepsilon_p, \mathbf{r})$ (proton). Outgoing momenta are given by $p' = (\varepsilon', \mathbf{p}')$ (scattered electron), $r' = (\varepsilon'_p, \mathbf{r}')$ (scattered proton), and $k = (\omega, \mathbf{k})$ (emitted photon). The electron charge is given by e , m_e is the electron mass, $\epsilon^*(k)$ is the photon polarization vector, $u(p)$ and $u(r)$ are the spinors of the incoming electron and proton, respectively.

As in the Bethe-Heitler calculation, a suitable set of approximations is applied to calculate the polarized contribution to the cross section. Owing to the smallness of the electron mass with respect to the large energies of most experimental measurements, ultra-relativistic approximations ($p \approx \varepsilon - m_e^2/(2\varepsilon)$) are applied to the final expressions. The “no-recoil” approximation [8] is applied, where the energy transferred by the exchange photon is neglected: $q = (0, \mathbf{q})$. Since the photon polar-angle distribution of the Bethe-Heitler expression is sharply peaked near zero, it follows that the virtuality q^2 of the exchange photon is also very small, $q^2 \sim m_e^2$ (see section 97 of Ref. [8]). The structure of the nucleus is neglected, which is justified at low q^2 . It additionally follows that this process occurs coherently with all charged nucleons in a large nucleus, and so the amplitude scales with the atomic number Z . For simplicity, ep scattering is considered and Z is set to unity.

The modulus square of the amplitude in Eq. 1 takes

* dhevanga@gmail.com

on the following form:

$$|M|^2 \equiv -\frac{e^6}{4q^4} W^{\mu\alpha} w_{\mu\alpha}, \quad (2)$$

$$W^{\mu\alpha} = Tr[u(r')\bar{u}(r')\gamma^\mu u(r)\bar{u}(r)\gamma^\alpha], \quad (3)$$

$$w_{\mu\alpha} = Tr[u(p')\bar{u}(p')Q_\mu^\sigma u(p)\bar{u}(p)Q_{\sigma\alpha}], \quad (4)$$

$$Q_\mu^\sigma = \gamma^\sigma \frac{\not{p}' + \not{k} + m_e}{p'k} \gamma_\mu - \gamma_\mu \frac{\not{p} - \not{k} + m_e}{pk} \gamma^\sigma, \quad (5)$$

where $W^{\mu\alpha}$ is the proton tensor and $w_{\mu\alpha}$ is the electron tensor, and both are expressed as a trace over products of gamma matrices. Typically, the final-state particle polarizations are not experimentally measurable. Accordingly, the photon, scattered electron, and scattered proton polarizations are summed over: $\sum_{pol} \varepsilon_a^*(k)\varepsilon_b(k) \rightarrow -g_{ab}$, $\sum_{spin} u(p')\bar{u}(p') = \not{p}' + m_e$, $\sum_{spin} u(r')\bar{u}(r') = \not{r}' + m_p$, where m_p is the proton mass. The incoming beam polarizations are measurable, for which the electron and proton spinor products are expressed as [8]

$$\begin{aligned} u(p)\bar{u}(p) &= \frac{1}{2}(\not{p} + m_e)(1 - \gamma^5 \not{\phi}^{(e)}), \\ u(r)\bar{u}(r) &= \frac{1}{2}(\not{r} + m_e)(1 - \gamma^5 \not{\phi}^{(p)}). \end{aligned} \quad (6)$$

The electron and proton spin 4-vectors (Pauli–Lubanski pseudovectors) are $a^{(e)}$ and $a^{(p)}$, respectively. They have the form $(0, \vec{\xi})$ in the particle's rest frame, for which $\vec{\xi}$ depends on the beam polarization. For the remaining expressions, longitudinal beam polarizations are assumed. In the target (proton) rest frame, $a^{(e)} = 2\mathbb{P}_e \frac{\varepsilon \varepsilon_p}{m_e m_p} (-1, 0, 0, +1)$. The parameter \mathbb{P}_e is the electron beam polarization, which is more conveniently defined as that measured in the laboratory frame (like ε and ε_p) along the electron's momentum. Terms of $\mathcal{O}(m_e/\varepsilon)$ and $\mathcal{O}(m_p/\varepsilon_p)$ are neglected in $a^{(e)}$. The proton spin 4-vector has the following form in the target rest frame, $a^{(p)} = \mathbb{P}_p (0, 0, 0, +1)$. The proton beam polarization, \mathbb{P}_p , is also defined as that measured in the laboratory frame along the proton's momentum.

The proton tensor, $W^{\mu\alpha}$, and the electron tensor, $w_{\mu\alpha}$, are expressed in terms of their unpolarized ($\mathcal{U}^{\mu\alpha}$, $\mathbf{u}_{\mu\alpha}$) and polarized ($\mathcal{P}^{\mu\alpha}$, $\mathbf{p}_{\mu\alpha}$) parts:

$$W^{\mu\alpha} \equiv \mathcal{U}^{\mu\alpha} + \mathcal{P}^{\mu\alpha}, \quad (7)$$

$$w_{\mu\alpha} \equiv \mathbf{u}_{\mu\alpha} + \mathbf{p}_{\mu\alpha}, \quad (8)$$

where the polarized parts arise from the γ^5 terms in Eq. 6. Evaluating the tensor traces yields the following expres-

sions for the polarized parts:

$$\mathcal{P}^{\mu\alpha} = 2i m_p q_a a^{(p)} \varepsilon^{a\mu\alpha}, \quad (9)$$

$$\mathbf{p}_{\mu\alpha} \equiv \mathbf{p}_{\mu\alpha}^{(1)} + \mathbf{p}_{\mu\alpha}^{(2)} + \mathbf{p}_{\mu\alpha}^{(3)} + \mathbf{p}_{\mu\alpha}^{(4)}, \quad (10)$$

$$\mathbf{p}_{\mu\alpha}^{(1)} = 8i \frac{m_e a^{(e),\lambda}}{(p'k)^2} \varepsilon_{a\mu\lambda\alpha} [m_e^2 q^a - p'k(p^a + k^a)], \quad (11)$$

$$\begin{aligned} \mathbf{p}_{\mu\alpha}^{(2)} &= 8i \frac{m_e a^{(e),\lambda}}{(pk)^2} \left[(m_e^2 q^a - p'^a pk) \varepsilon_{a\mu\lambda\alpha} \right. \\ &\quad \left. - p'^a k_\lambda (k^b - p^b) \varepsilon_{a\mu b\alpha} \right], \end{aligned} \quad (12)$$

$$\begin{aligned} \mathbf{p}_{\mu\alpha}^{(3)} &= 8i \frac{m_e a^{\lambda,(e)}}{(pk)(p'k)} \left[\frac{q^2}{2} p'^a \varepsilon_{\lambda a \mu \alpha} + p'^a p^\mu (k^b - p^b) \varepsilon_{\lambda b \alpha} \right. \\ &\quad \left. + (p'^a k^a (p^b - p'^b) \right. \\ &\quad \left. + p^b (p^\alpha - k^\alpha) (p'^a + k^a)) \varepsilon_{\lambda a b \mu} \right] \end{aligned} \quad (13)$$

$$\mathbf{p}_{\mu\alpha}^{(4)} = -\mathbf{p}_{\alpha\mu}^{(3)}. \quad (14)$$

The polarized tensors, $\mathcal{P}^{\mu\alpha}$ and $\mathbf{p}_{\mu\alpha}$, are antisymmetric, while the unpolarized tensors, $\mathcal{U}^{\mu\alpha}$ and $\mathbf{u}_{\mu\alpha}$, are symmetric. Thus, only $\mathcal{U}^{\mu\alpha} \mathbf{u}_{\mu\alpha}$ and $\mathcal{P}^{\mu\alpha} \mathbf{p}_{\mu\alpha}$ contribute to the cross section. However, at higher orders with loop corrections, single-spin asymmetries emerge [10]. We have:

$$|M|^2 = -\frac{e^6}{4q^4} [\mathcal{U}^{\mu\alpha} \mathbf{u}_{\mu\alpha} + \mathcal{P}^{\mu\alpha} \mathbf{p}_{\mu\alpha}] \quad (15)$$

$$= -\frac{(4\pi)^3 m_e^2 \alpha r_e^2}{4q^4} [\mathcal{U}^{\mu\alpha} \mathbf{u}_{\mu\alpha} + \mathcal{P}^{\mu\alpha} \mathbf{p}_{\mu\alpha}] \quad (16)$$

where the fine-structure constant is given by $\alpha = \frac{e^2}{4\pi}$ and the classical electron radius is given by $r_e = \frac{e^2}{4\pi m_e}$.

III. DIFFERENTIAL CROSS SECTIONS

The fully differential cross section expressed in the target rest frame is

$$d\sigma = \frac{1}{(4\pi)^5} |M|^2 \frac{|\mathbf{p}'| \omega d\omega}{|\mathbf{p}| m_p^2} d\Omega' d\Omega_k, \quad (17)$$

$$\equiv d\sigma_{\mathcal{U}} + d\sigma_{\mathcal{P}}. \quad (18)$$

Inserting the polarized part of Eq. 16 into Eq. 17 gives

$$d\sigma_{\mathcal{P}} = \frac{-\alpha r_e^2 m_e^2}{4(4\pi)^2 q^4 m_p^2} \mathcal{P}^{\mu\alpha} \mathbf{p}_{\mu\alpha} \frac{|\mathbf{p}'| \omega d\omega}{|\mathbf{p}|} d\Omega' d\Omega_k. \quad (19)$$

The unpolarized term, $d\sigma_{\mathcal{U}}$, corresponds to the usual Bethe-Heitler expression and was re-derived as a cross check. The angular phase space of the scattered electron and emitted photon are denoted by $d\Omega'$ and $d\Omega_k$, respectively. In order to provide a practical expression, integration over the angles in the final state is needed.

Evaluating the contraction of Levi-Civita symbols in $\mathcal{P}^{\mu\nu}\mathbf{p}_{\mu\nu}$ leads to the following:

$$\begin{aligned} \mathcal{P}^{\mu\nu}\mathbf{p}_{\mu\nu} = & -32m_e m_p \left[\frac{1}{(pk)^2} \left(\frac{q^2}{2} k a^{(e)} (q a^{(p)} - 2p' a^{(p)}) \right) \right. \\ & + \left(\frac{1}{(pk)^2} + \frac{1}{(p'k)^2} \right) \left(q a^{(e)} q a^{(p)} m_e^2 - a^{(e)} a^{(p)} q^2 m_e^2 \right) \\ & + \frac{1}{(pk)(p'k)} \left(-q^4 a^{(e)} a^{(p)} - 2q a^{(e)} q a^{(p)} m_e^2 \right. \\ & \quad \left. + \frac{q^2}{2} \left(2p' a^{(e)} (p' a^{(p)} - p a^{(p)}) \right. \right. \\ & \quad \left. \left. + k a^{(e)} (3q a^{(p)} + 2p a^{(p)}) \right. \right. \\ & \quad \left. \left. + 4m_e^2 a^{(e)} a^{(p)} \right) \right] \\ & + \left(\frac{1}{p'k} - \frac{1}{pk} \right) \left((q a^{(e)} + k a^{(e)}) q a^{(p)} - 2q^2 a^{(e)} a^{(p)} \right) \Big]. \end{aligned} \quad (20)$$

Integration of the differential cross section over the scattered electron angles, $d\Omega'$, can be performed analytically [11] and is done first. For longitudinally polarized beams, all 4-vector products involving the scattered electron momentum, p' , in Eq. 20 can be expressed in terms of two basis vectors, \mathbf{a} and \mathbf{b} , defined through q^2 and $p'k$:

$$\begin{aligned} q^2 &= -\mathbf{q}^2 = -(\mathbf{p}' - \mathbf{p} + \mathbf{k})^2, \\ &\equiv -(\mathbf{p}' - \mathbf{T})^2 = -(\mathbf{p}'^2 + T^2)(1 - \mathbf{p}'\mathbf{a}), \end{aligned} \quad (21)$$

$$p'k = \omega \varepsilon' - \mathbf{p}'\mathbf{k} \equiv \omega \varepsilon' (1 - \mathbf{p}'\mathbf{b}), \quad (22)$$

$$\mathbf{a} = \frac{2\mathbf{T}}{\mathbf{p}'^2 + T^2}, \quad (23)$$

$$\mathbf{b} = \frac{\mathbf{k}}{\omega \varepsilon'}. \quad (24)$$

The other 4-vector products containing p' , $p' a^{(e)}$ and $p' a^{(p)}$, are also expressed with terms containing $(1 - \mathbf{p}'\mathbf{a})$ and $(1 - \mathbf{p}'\mathbf{b})$. Ultimately, the integral over the scattered electron angles is of the form:

$$I_{m,n} = \int d\Omega' (1 - \mathbf{p}'\mathbf{a})^{-m} (1 - \mathbf{p}'\mathbf{b})^{-n}. \quad (25)$$

Note that by convention a factor of $1/(2\pi)$ as defined for $I_{m,n}$ in Ref. [11] is not included here. The array of integrals present is: $I_{0,0}, I_{0,1}, I_{1,0}, I_{-1,1}, I_{1,-1}, I_{1,1}, I_{2,0}, I_{0,2}, I_{2,1}, I_{1,2}, I_{2,-1}, I_{2,-2}, I_{2,2}$. Integrals with negative indices can be expressed in terms of the other integrals. For the non-trivial integrals, $I_{1,1}, I_{2,1}, I_{1,2}, I_{2,2}$, Feynman parameters are first used to combine denominators. A strategically chosen polar angle axis is then chosen to make the azimuthal integrations trivial. Finally, a table of integrals [12] is used for the final integration over the Feynman parameter. Integration over $d\Omega_k$ similarly starts with a strategically chosen polar axis ($\hat{\mathbf{p}}$) that makes the azimuthal integration trivial. Assembly of all terms resulting from each integral is quite algebraically laborious. Mathematica is used to assemble and simplify the algebra.

The expressions for the double- and single-differential cross sections are given in the laboratory frame. In the

small-angle limit that characterizes the Bethe-Heitler expression and for ultra-relativistic particles, the Lorentz transformation from the lab frame (LF) to the target rest frame (TRF) is especially simple: $E_{TRF} = \frac{2\varepsilon_p}{m_p} E_{LF}$ for $E \in \{\varepsilon, \varepsilon', \omega\}$.

The resulting polarized cross section, double-differential in the photon polar angle and energy up to $\mathcal{O}(m_e^2)$, in the laboratory frame, is given by

$$\begin{aligned} \frac{d\sigma_{\mathcal{P}}}{d\omega d\delta} = & \mathbb{P}_e \mathbb{P}_p \frac{\alpha r_e^2 m_e^2 \delta}{\omega \varepsilon^3 \varepsilon' \varepsilon_p (1 + \delta^2)^2} \left[2\omega^3 + 2(-1 + 2L_2)\varepsilon'^3 \right. \\ & + (8 + 6L_2)\varepsilon'^2 \omega + 2(1 + L_2)\varepsilon' \omega^2 \\ & - 2(L_1 + L_2 - L_\theta)(\varepsilon^2 + \varepsilon'^2 - \varepsilon \omega \delta^2) \varepsilon \\ & - (4\varepsilon'^3 + 2(-1 + L_2)\varepsilon'^2 \omega + 5\varepsilon' \omega^2 + 2\omega^3) \delta^2 \\ & \left. + \frac{2\varepsilon \varepsilon' \omega}{1 + \delta^2} + 2L_1 \varepsilon \varepsilon' (\varepsilon + \varepsilon') (1 + \delta^2) \right], \end{aligned} \quad (26)$$

where $L_1 = \ln \frac{4\varepsilon' \varepsilon_p}{m_e m_p}$, $L_2 = \ln \frac{4\varepsilon \varepsilon' \varepsilon_p}{\omega m_e m_p}$, $L_\theta = \ln(1 + \delta^2)$. One observes that, like the Bethe-Heitler spectrum, the angular spectrum is sharply peaked for $\delta = \theta_k \frac{\varepsilon}{m_e} \lesssim 1$.

Finally, integrating over the photon polar angle yields the single-differential polarized cross section

$$\frac{d\sigma_{\mathcal{P}}}{d\omega} = \mathbb{P}_e \mathbb{P}_p \frac{4\alpha r_e^2 \varepsilon' m_e^2}{\omega \varepsilon \varepsilon' \varepsilon_p} \left(F_1 + \frac{\varepsilon}{4\varepsilon'} F_2 + \frac{\varepsilon'}{8\varepsilon} F_3 + \frac{\varepsilon^2}{2\varepsilon'^2} F_4 \right), \quad (27)$$

where

$$F_1 = \frac{1}{8} (5 + L_1(-2 + 4L_3) - 4L_3(-3 + L_2) + 2L_2),$$

$$F_2 = L_1 + (-1 + L_3)(3 + 2L_3 - 2L_2),$$

$$F_3 = -1 - 2L_2 + 2L_3(-9 + 2L_2),$$

$$F_4 = (-1 + L_3)(-2 + L_1 + L_2 - L_3).$$

and $L_3 = \ln \frac{\pi \varepsilon \varepsilon_p}{m_e m_p}$. The polarized cross section is to be compared to the unpolarized Bethe-Heitler expression [5]:

$$\frac{d\sigma_{\mathcal{U}}}{d\omega} = \frac{4\alpha r_e^2 \varepsilon'}{\omega \varepsilon} \left(\frac{\varepsilon}{\varepsilon'} + \frac{\varepsilon'}{\varepsilon} - \frac{2}{3} \right) \left[L_2 - \frac{1}{2} \right]. \quad (28)$$

It is clear that the polarized component of the bremsstrahlung cross section is highly suppressed with respect to the unpolarized component by a factor of $m_e^2/(\varepsilon \varepsilon_p)$. Figures 2 and 3 show the energy spectra of the polarized and unpolarized components, respectively, for $\varepsilon = 18$ GeV and $\varepsilon_p = 275$ GeV (top EIC energies). The sharp turn in the spectrum near the photon energy limit (18 GeV) in Fig. 2 occurs when logs with different signs change in magnitude in F_4 . It was also observed in a numerical calculation at larger q^2 [3].

For the case of longitudinally polarized electrons and *transversely* polarized protons, as expected for the EIC, the polarized component vanishes exactly when integrating over azimuthal angles of the final-state particles. Additionally, the vectors \mathbf{a} and \mathbf{b} do not form a complete basis to express the 4-vector products in Eq. 20, which




FIG. 2. Polarized component of the bremsstrahlung cross section versus photon energy for $\varepsilon = 18$ GeV and $\varepsilon_p = 275$ GeV (top EIC energies). The product of beam polarizations is set to unity: $\mathbb{P}_e\mathbb{P}_p = 1$.




FIG. 3. Unpolarized component of the bremsstrahlung cross section (Bethe-Heitler) versus photon energy for $\varepsilon = 18$ GeV and $\varepsilon_p = 275$ GeV (top EIC energies).

further complicates analytic integration. Transverse po-

larization is not considered further here.

In regards to the general luminosity program at the EIC, where high Z heavy-ions will be accelerated, it should be noted that the Born approximation used in the Bethe-Heitler expression is expected to be inadequate. Thus, the Bethe-Heitler expression should be replaced by the Bethe-Maximon expression [6–8]:

$$\frac{d\sigma_U}{d\omega} = \frac{4Z^2\alpha r_e^2 \varepsilon'}{\omega \varepsilon} \left(\frac{\varepsilon}{\varepsilon'} + \frac{\varepsilon'}{\varepsilon} - \frac{2}{3} \right) \left[L_2 - \frac{1}{2} - f(\alpha Z) \right], \quad (29)$$

$$f(\alpha Z) = (\alpha Z)^2 \sum_{n=1}^{\infty} \frac{1}{n(n^2 + (\alpha Z)^2)}. \quad (30)$$

For heavy nuclei such as uranium, the Bethe-Maximon and Bethe-Heitler expressions differ by about 2%. For light nuclei, $\alpha Z \ll 1$ and $f(\alpha Z) \approx 1.2(\alpha Z)^2$.

IV. CONCLUSION

The polarized component of the bremsstrahlung cross section has been calculated. An analytic expression has been derived, which shows that the polarized component is highly suppressed with respect to the usual unpolarized Bethe-Heitler expression. The suppression is linked to the low q^2 that characterizes the bremsstrahlung process, which is $\sim m_e^2$.

ACKNOWLEDGMENTS

I would like to thank Andrei Afanasev, Wim Cosyn, and Katarzyna Wichmann for useful discussions. This work is supported by US DOE Nuclear Physics Grant No. DE-FG02-07ER41521.

-
- [1] J. Andruszkow *et al.* (ZEUS Luminosity Group), Luminosity measurement in the ZEUS experiment, *Acta Phys. Polon. B* **32**, 2025 (2001).
 - [2] L. Adamczyk *et al.* (ZEUS), Measurement of the Luminosity in the ZEUS Experiment at HERA II, *Nucl. Instrum. Meth. A* **744**, 80 (2014), arXiv:1306.1391 [physics.ins-det].
 - [3] I. V. Akushevich, A. N. Ilyichev, and N. M. Shumeiko, Radiative electroweak effects in deep inelastic scattering of polarized leptons by polarized nucleons, *J. Phys. G* **24**, 1995 (1998), arXiv:hep-ph/9804361.
 - [4] A. Afanasev, I. Akushevich, and N. Merenkov, Model independent radiative corrections in processes of polarized electron nucleon elastic scattering, *Phys. Rev. D* **64**, 113009 (2001), arXiv:hep-ph/0102086.
 - [5] H. Bethe and W. Heitler, On the Stopping of fast particles and on the creation of positive electrons, *Proc. Roy. Soc. Lond. A* **146**, 83 (1934).
 - [6] H. A. Bethe and L. C. Maximon, Theory of Bremsstrahlung and Pair Production. 1. Differential Cross Section, *Phys. Rev.* **93**, 768 (1954).
 - [7] H. Davies, H. A. Bethe, and L. C. Maximon, Theory of Bremsstrahlung and Pair Production. 2. Integral Cross Section for Pair Production, *Phys. Rev.* **93**, 788 (1954).
 - [8] V. B. Berestetskii, E. M. Lifshitz, and L. P. Pitaevskii, *QUANTUM ELECTRODYNAMICS*, Course of Theoretical Physics, Vol. 4 (Pergamon Press, Oxford, 1982).
 - [9] T. Haas and V. Makarenko, Precision calculation of processes used for luminosity measurement at the ZEUS experiment, *Eur. Phys. J. C* **71**, 1574 (2011), arXiv:1009.2451 [hep-ph].
 - [10] A. V. Afanasev, M. I. Konchatnij, and N. P. Merenkov, Single-spin asymmetries in the Bethe-Heitler process $e + p \rightarrow e + \gamma + p$ from QED radiative corrections, *J. Exp. Theor. Phys.* **102**, 220 (2006), arXiv:hep-ph/0507059.
 - [11] R. L. Gluckstern and M. H. Hull, Polarization Dependence of the Integrated Bremsstrahlung Cross Section, *Phys. Rev.* **90**, 1030 (1953).
 - [12] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products* (1943).