Uncertainty relations for metric adjusted skew information and Cauchy-Schwarz inequality

Xiaoli Hu^{*}

School of Artificial Intelligence, Jianghan University, Wuhan, Hubei 430056, China

Naihuan Jing[†]

Department of Mathematics, North Carolina State University, Raleigh, NC 27695, USA

(Dated: August 1, 2023)

Skew information is a pivotal concept in quantum information, quantum measurement, and quantum metrology. Further studies have lead to the uncertainty relations grounded in metric-adjusted skew information. In this work, we present an in-depth investigation using the methodologies of sampling coordinates of observables and convex functions to refine the uncertainty relations in both the product form of two observables and summation form of multiple observables.

PACS numbers:

I. INTRODUCTION

Quantum mechanics is a fundamental theory that describes the behavior of matter and energy at the atomic and subatomic scale [1]. One of its most distinctive features is the Heisenberg uncertainty principle, which states that certain pairs of physical properties, such as position and momentum, cannot be simultaneously measured with arbitrary precision [2, 3]. The generalization of Heisenberg's uncertainty principle to encompass any two non-commuting observables was accomplished by Robertson [4] and later refined by Schrödinger [5]. Although uncertainty relations play a crucial role in the study of quantum systems, there are other types of uncertainty in quantum systems that are not captured by the standard uncertainty relations, for instance the uncertainty of skew information. This concept has important applications in quantum information processing, such as in the design of more accurate and efficient quantum measurements.

Skew information was first introduced by Wigner and Yanase in 1963 as a measure of the information content of quantum states in the presence of conserved quantities [6]. For a composite system, such as a measuring device consisting of a system of interest and a pointer, Wigner demonstrated that it is impossible to perform an ideal measurement of an observable that does not commute with a conserved quantity in the presence of the conservation law [7]. Luckily, performing an approximate measurement of the observable is still feasible, and the error diminishes as the size of the system grows [8, 9]. Such a trade-off between the measuring accuracy and the size of the measuring apparatus is known as Wigner-Araki-Yanase theorem in literature. Further remarkable studies were later presented in the references [10–15].

In recent years, there has been significant interest in developing new techniques for quantifying and understanding skew information in quantum systems. One important direction of research has been to take into account the geometry of the underlying quantum state space. This has led to the development of the metric-adjusted skew information, which is a measure of skew information that takes into account the curvature of the quantum state space [16]. Uncertainty relations based on metric-adjusted skew information can establish fundamental limits on quantum measurement accuracy and information processing speed. Additionally, skew information can help understand quantum correlations and classical behavior in quantum systems. Recent developments in uncertainty relations include the introduction of a new type of uncertainty relation using metric-adjusted skew information [17], which extended the existing Wigner-Yanase-Dyson (WYD) skew information and demonstrated its practical importance. The results indicate that the skew information provides a new way to quantify Bohr's complementary principle and Heisenberg's uncertainty principle. The Wigner-Yanase-Dyson metric adjusted skew information was introduced as a generalization of the Wigner-Yanase skew information in [18]. In [19], it was demonstrated that the sum uncertainty relation for the Wigner-Yanase skew information, introduced in [20], also holds for arbitrary metric-adjusted skew information, and the related uncertainty relations for quantum channels were given there. An equivalent formulation for arbitrary two observables and two channels was also presented in [21], and a generalization to multi-observables was given in [22, 23]. Recently, product

^{*}Electronic address: xiaolihumath@jhun.edu.cn

[†]Electronic address: jing@ncsu.edu

and summation forms of the uncertainty relation were proposed in [24] for the WYD-metric adjusted skew information based on two different refinements of the Cauchy-Schwarz inequality.

In this work, we introduce novel and improved uncertainty relations in both product and summation forms for two and N quantum observables, respectively. Our approach involves utilizing carefully chosen convex functions to enhance the Cauchy-Schwarz inequality and measuring the observables using sampled measurement coordinates. This work presents new uncertainty relations for the WYD-metric adjusted skew information, which provide a tighter bound on the amount of uncertainty in a quantum system. Our motivation stems from an information-theoretic perspective, where several information measures are crucial to the theory's structural properties. Overall, the study of uncertainty and skew information in quantum systems is an important and active area of research, with many potential applications in quantum information processing and other areas of science and technology.

II. PRELIMINARIES

Recall that the Wigner-Yanase skew information [6] is a measure of non-commutativity between the square root of the state ρ and the conserved observable A:

$$I_{\rho}(A) = -\frac{1}{2} \operatorname{Tr}([A,\sqrt{\rho}])^2 = \operatorname{Tr}(\rho A^2) - \operatorname{Tr}(\sqrt{\rho}A\sqrt{\rho}A),$$
(1)

where Tr means the trace operation, $[A, \sqrt{\rho}]$ is the commutator of A and $\sqrt{\rho}$, so it vanishes when ρ commutes with A. I_{ρ} is clearly homogeneous in ρ . When $\rho = |\psi\rangle\langle\psi|$ is a pure state, $I_{\rho}(A) = \text{Tr}(\rho A^2) - (\text{Tr}(\rho A))^2 = \Delta_{\rho}(A)$ is the variance of the observable A with respect to ρ . This means that the Wiger-Yanase skew information generalizes the variance. Furthermore, when ρ is a mixed state, we have $I_{\rho}(A) > \Delta_{\rho}(A)$ in general. The Heisenberg uncertainty principle in terms of the skew information states that [17]:

$$I_{\rho}(A)I_{\rho}(B) \ge \frac{1}{4}|\mathrm{Tr}(\rho[A,B])|^2.$$
 (2)

The Fisher-Rao metric for classical information is the unique contracting Riemannian metric under the Markov morphism [18]. The symmetric monotone metric for quantum case is constructed through the so-called Morozova-Chentsov functions. Such a function $f: (0, +\infty) \to \mathbb{R}$ is a positive operator monotonic function (i.e. 0 < f(A) < f(B)holds when 0 < A < B) such that f is symmetric (i.e. $f(x) = xf(x^{-1})$), and normalized (i.e. f(1) = 1). The Morozova-Chentsov function corresponding to a positive operator-monotone and symmetric function f is defined by

$$c(x,y) = \frac{1}{yf(xy^{-1})}, \quad x,y > 0.$$
(3)

According to Morozova, Chentsov and Petz, the monotone metric on the state space of a quantum system is given by (cf. [25])

$$K^c_{\rho}(A,B) = \operatorname{Tr}(A^{\dagger}c(L_{\rho},R_{\rho})B),\tag{4}$$

where $L_{\rho}(A) = \rho A$ and $R_{\rho}(A) = A\rho$ be left and right multiplication operators, then $[\rho, A] = \rho A - A\rho = (L_{\rho} - R_{\rho})A$. This metric form depends on the choice of the Morozova-Chentsov function, so it is not unique. The metric constant is denoted by $m(c) = \lim_{t\to 0} c(t, 1)^{-1} = \lim_{t\to 0} f(t)$. A symmetric monotone metric is called regular if m(c) > 0. Because f(t) is a positive operator monotone function on $[0, \infty)$, we have m(c) = f(0) for a regular metric. Let c be the Morozova-Chentsov function of a regular metric. Then the so-called metric adjusted skew information $I_{\rho}^{c}(A)$ for observable A and quantum state ρ is defined by [11, 18]:

$$I_{\rho}^{c}(A) = \frac{m(c)}{2} K_{\rho}^{c}(\mathbf{i}[\rho, A], \mathbf{i}[\rho, A]) = \frac{m(c)}{2} \operatorname{Tr}(\mathbf{i}[\rho, A]c(L_{\rho}, R_{\rho})\mathbf{i}[\rho, A]),$$
(5)

where $i^2 = -1$. If we consider the following positive operator monotone and symmetric function

$$f_p(x) = p(1-p)\frac{(x-1)^2}{(x^p-1)(x^{1-p}-1)}, \qquad 0
(6)$$

then the corresponding Morozova-Chentsov function is of the form

$$c^{WYD}(x,y) = \frac{1}{p(1-p)} \frac{(x^p - y^p)(x^{1-p} - y^{1-p})}{(x-y)^2}, \quad 0
⁽⁷⁾$$

Therefore $m(c^{WYD}) = \lim_{t\to 0} c^{WYD}(t,1) = p(1-p)$ and the WYD-metric adjusted skew information is given by

$$I_{\rho}^{c^{WYD}}(A) = -\frac{1}{2} \operatorname{Tr}([\rho^{p}, A][\rho^{1-p}, A]), \quad 0
(8)$$

Equation (8) is the Dyson generalization of the Wigner-Yanase skew information when $p = \frac{1}{2}$. The famous Wigner-Yanase-Dyson (WYD) conjecture on the convexity of $I_{\rho}^{c^{WYD}}(A)$ was proved by Lieb [26].

When c is a regular Morozova-Chentsov function, the metric-adjusted correlation of observables A and B is given by

$$\operatorname{Corr}_{\rho}^{c}(A,B) = \frac{m(c)}{2} K_{\rho}^{c}(\mathbf{i}[\rho,A],\mathbf{i}[\rho,B]).$$
(9)

The metric adjusted correlation is a measure of statistical dependence between two observables in quantum mechanics that takes into account their non-commutativity. It's a useful tool for detecting entanglement and studying quantum phase transitions. However, it's not a real-valued function and can take complex values, so it can't directly provide positive bounds. We present a novel approach that addresses this issue by utilizing coordinate sampling of observables and a carefully chosen convex function, in combination with the Cauchy-Schwarz inequality, to derive robust uncertainty relations for the WYD-metric adjusted correlation. Our method yields stronger bounds than several previous uncertainty relations.

The paper is organized as follows. In Section II, we provide a review of two recent uncertainty relations that offer a "fine-grained" analysis of the product-form uncertainty relation for two observables using the Cauchy-Schwarz inequality. We then introduce our technique of coordinate sampling and convex function selection to further enhance the lower bound. In Section III, we present a set of practical and straightforward uncertainty relations for the sumform skew information based on the fundamental inequality and well-chosen convex functions. We conclude in Section IV.

III. IMPROVED UNCERTAINTY RELATIONS FOR PRODUCT-FORM SKEW INFORMATION

Consider a Hilbert space H of dimensional d over the complex field \mathbb{C} . Let $M_d(\mathbb{C})$ denote by the set of complex matrices and $D_d(\mathbb{C})$ the set of complex density matrices on H. We use $\langle X, Y \rangle = \text{Tr}(X^{\dagger}Y)$ to denote the Hilbert-Schmidt scalar product, where $X, Y \in M_d(\mathbb{C})$. The computational basis of H is given by $\{|i\rangle, i = 1, 2, \dots, d\}$, and $\{E_{ij} = |i\rangle\langle j|, i, j = 1, 2, \dots, d\}$ forms a complete set of local orthogonal matrices. Specifically, $\langle E_{ij}, E_{kl} \rangle = \text{Tr}(E_{ji}E_{kl}) = \delta_{ik}\delta_{jl}$, and they form an orthogonal basis for all observables in $M_d(\mathbb{C})$. Given a matrix $M = [m_{ij}] \in M_d(\mathbb{C})$, we define $\vec{M} = (m_{11}, m_{12}, \dots, m_{1d}, m_{21}, \dots, m_{dd})^T$ as its d^2 -dimensional coordinate vector with respect to the standard basis $\{E_{ij}\}$. Let $\vec{\mathbf{E}} = (E_{11}, E_{12}, \dots, E_{1d}, E_{21}, \dots, E_{dd})^T$, then any observable A can be expressed as a linear combination of these basis observables, i.e.

$$A = \sum_{i,j=1}^{d} a_{ij} E_{ij} = \langle \vec{A}, \vec{E} \rangle = \vec{A}^{\dagger} \cdot \vec{E}$$
(10)

with $a_{ij} = \langle E_{ij}, A \rangle = \operatorname{Tr}(E_{ji}A).$

The metric adjusted skew information of A is given by

$$I_{\rho}^{c}(A) = \frac{m(c)}{2} K_{\rho}^{c}(\mathbf{i}[\rho, A], \mathbf{i}[\rho, A])$$

$$= \frac{m(c)}{2} \sum_{i,j=1}^{d} \sum_{k,l=1}^{d} a_{ij}^{*} a_{kl} K_{\rho}^{c}(\mathbf{i}[\rho, E_{ij}], \mathbf{i}[\rho, E_{kl}])$$

$$= \vec{A}^{\dagger} \cdot \Gamma \cdot \vec{A},$$

(11)

where $\Gamma = (\Gamma_{i_j,k_l})$ and $\Gamma_{i_j,k_l} = \frac{m(c)}{2} K_{\rho}^c(\mathbf{i}[\rho, E_{i_j}], \mathbf{i}[\rho, E_{k_l}]) = \operatorname{Corr}_{\rho}^c(E_{i_j}, E_{k_l})$ with $i_j = d(i-1) + j, k_l = d(k-1) + l$.

In fact, under the Wigner-Yanase-Dyson-metric we have

$$\Gamma_{ij,k_l} = \frac{m(c^{WYD})}{2} \operatorname{Tr}((\mathbf{i}[\rho, E_{ij}])^{\dagger} c^{WYD}(L_{\rho}, R_{\rho})\mathbf{i}[\rho, E_{kl}])
= -\frac{1}{2} \operatorname{Tr}[(L_{\rho} - R_{\rho})E_{ji} \frac{(L_{\rho}^{p} - R_{\rho}^{p})(L_{\rho}^{1-p} - R_{\rho}^{1-p})}{(L_{\rho} - R_{\rho})^{2}} (L_{\rho} - R_{\rho})E_{kl})]
= -\frac{1}{2} \operatorname{Tr}([\rho^{p}, E_{ij}^{\dagger}][\rho^{1-p}, E_{kl}])
= \frac{1}{2} \langle [\rho^{p}, E_{ij}], [\rho^{1-p}, E_{kl}] \rangle.$$
(12)

It follows from (11) that Γ is semi-positive, so there is a matrix C such that $\Gamma = C^{\dagger}C$. Then $I_{\rho}^{c}(A) = \vec{A}^{\dagger}C^{\dagger}C\vec{A} = (C\vec{A})^{\dagger}(C\vec{A}) = |\vec{\alpha}|^{2}$, where $\vec{\alpha} = C\vec{A} = (\alpha_{1}, \cdots, \alpha_{d^{2}})^{\dagger}$. Let \vec{B} be the coordinate vector of the quantum observable B. Similarly $I_{\rho}^{c}(B) = |\vec{\beta}|^{2}$ with $\vec{\beta} = C\vec{B} = (\beta_{1}, \cdots, \beta_{d^{2}})^{\dagger}$. Therefore, the product of the Wigner-Yanase-Dyson-metric adjusted skew information for observables A and B can be written as

$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) = |\vec{\alpha}|^{2}|\vec{\beta}|^{2} = \sum_{i,j=1}^{d^{2}} |\alpha_{i}|^{2}|\beta_{j}|^{2} \ge |\sum_{i=1}^{d^{2}} \alpha_{i}^{*}\beta_{i}|^{2} = |(\vec{\alpha},\vec{\beta})|^{2} = |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^{2},$$
(13)

which is the usual form of the adjusted metric skew information based uncertainty relation.

Consider an *n*-dimensional real vector space \mathbb{R}^n equipped with the classical inner product (,). Let $\vec{v} = (v_1, \dots, v_n)^T \in \mathbb{R}^n$ be a vector, and let $\vec{v}_k = (v_1, \dots, v_k, 0, \dots, 0)^T$ denote a partial vector of \vec{v} with the first k components. We define the complement vector $\vec{v}_k^c = \vec{v} - \vec{v}_k$. The symmetric group \mathfrak{S}_n acts on \mathbb{R}^n by permutation, i.e. for $\sigma \in \mathfrak{S}_n$, the action is given by $\sigma(\vec{v}) = (v_{\sigma(1)}, \dots, v_{\sigma(n)})^T$. In particular, for any $\sigma \in \mathfrak{S}_n$ and for any $k \leq n$, we have $\sigma(\vec{v}_k) = (v_{\sigma(1)}, \dots, v_{\sigma(k)}, 0, \dots, 0)^T$, then $\sigma(\vec{v}_k) = \sigma(v) - \sigma(\vec{v}_k)$.

In the sequel, we set $n = d^2$, where dim(H) = d. We define $x_i = |\alpha_i|$ and $y_j = |\beta_j|$ for $i, j = 1, \dots, n$ in (13), so that $I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) = \sum_{i,j=1}^{n} x_i^2 y_j^2$. The n^2 elements $x_i^2 y_j^2$ can be stored in a real matrix $M_n = [x_i^2 y_j^2]_{1 \le i,j \le n}$. Let M_k be the k-th sequential principal matrix of M_n . More generally, let $M_I = [x_i^2 y_j^2]_{i,j \in I}$ for $I \subset \{1, 2, \dots, n\}$. The norm of M_k is denoted by $|M_k|_{\infty} = \sum_{i,j=1}^{k} x_i^2 y_j^2$. In particular,

$$|M_n|_{\infty} = \sum_{i,j=1}^n x_i^2 y_j^2 = I_{\rho}^{c^{WYD}}(A) I_{\rho}^{c^{WYD}}(B).$$
(14)

For partial vectors $\vec{x_k} = (x_1, \dots, x_k, 0, \dots, 0)^T$, $\vec{y_k} = (y_1, \dots, y_k, 0, \dots, 0)^T \in \mathbb{R}^n$, let f_{CS} be the function defined by $f_{CS}(M_k) = (\vec{x_k}, \vec{y_k})^2$, and similarly defined for a general principal matrix. By the Cauchy-Schwarz inequality, we have

$$|M_k|_{\infty} = \sum_{i,j=1}^k x_i^2 y_j^2 = (\vec{x}_k, \vec{x}_k) (\vec{y}_k, \vec{y}_k) \ge (\vec{x}_k, \vec{y}_k)^2 = f_{CS}(M_k).$$
(15)

The quantity proposed by Xiao et al. in [27] and Yu et al. in [28] is given by this expression for $k \leq n$, i.e.

$$I_k = |M_n|_{\infty} - |M_k|_{\infty} + f_{CS}(M_k) = \sum_{i=1}^n x_i^2 y_i^2 + \sum_{1 \le i < j \le n, k < j} (x_i^2 y_j^2 + x_j^2 y_i^2) + \sum_{1 \le i < j \le k} 2x_i y_i x_j y_j.$$
(16)

We will mainly use I_2 , I_3 and I_4 in our later examples:

$$I_{2} = \sum_{i,j=1}^{n} x_{i}^{2} y_{j}^{2} - x_{1}^{2} y_{2}^{2} - x_{2}^{2} y_{1}^{2} + 2x_{1} x_{2} y_{1} y_{2};$$

$$I_{3} = \sum_{i,j=1}^{n} x_{i}^{2} y_{j}^{2} - x_{1}^{2} y_{2}^{2} - x_{2}^{2} y_{1}^{2} - x_{1}^{2} y_{3}^{2} - x_{3}^{2} y_{1}^{2} - x_{2}^{2} y_{3}^{2} - x_{3}^{2} y_{2}^{2} + 2x_{1} x_{2} y_{1} y_{2} + 2x_{1} x_{3} y_{1} y_{3} + 2x_{2} x_{3} y_{2} y_{3};$$

$$I_{4} = \sum_{i,j=1}^{n} x_{i}^{2} y_{j}^{2} - x_{1}^{2} (y_{2}^{2} + y_{3}^{2} + y_{4}^{2}) - x_{2}^{2} (y_{3}^{2} + y_{4}^{2}) - x_{3}^{2} y_{4}^{2} - y_{1}^{2} (x_{2}^{2} + x_{3}^{2} + x_{4}^{2}) - y_{2}^{2} (x_{3}^{2} + x_{4}^{2}) - y_{3}^{2} x_{4}^{2} + 2x_{1} x_{2} y_{1} y_{2} + 2x_{1} x_{3} y_{1} y_{3} + 2x_{1} x_{4} y_{1} y_{4} + 2x_{2} x_{3} y_{2} y_{3} + 2x_{2} x_{4} y_{2} y_{4} + 2x_{3} x_{4} y_{3} y_{4}.$$

$$(17)$$

$$\begin{bmatrix} x_1^2 y_1^2 & \cdots & x_1^2 y_k^2 & x_1^2 y_{k+1}^2 & \cdots & x_1^2 y_n^2 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ x_k^2 y_1^2 & \cdots & x_k^2 y_k^2 & x_k^2 y_{k+1}^2 & \cdots & x_k^2 y_n^2 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ x_n^2 y_1^2 & \cdots & x_n^2 y_k^2 & x_n^2 y_{k+1}^2 & \cdots & x_n^2 y_n^2 \end{bmatrix}$$

FIG. 1: The matrix M_k and quantity I_k . The part framed in orange and blue are the sequential principal matrices M_k and M_{k+1} respectively. I_k is equal to $f_{CS}(M_k)$ plus the elements outside of the k-th sequential principal matrix M_k .

See FIG.1 for the meaning of M_k and I_k . In practice, the storage of n^2 elements $x_i^2 y_j^2$ $(i, j = 1, \dots, n)$ in a matrix is random. The randomness can be generated by the action of the symmetry group \mathfrak{S}_n . For $\sigma, \tau \in \mathfrak{S}_n$, we define $M_n^{\sigma,\tau} = (\sigma, \tau)M_n = (m_{\sigma(i),\tau(j)})_{n \times n}$, then $|M_n^{\sigma,\tau}|_{\infty} = |M_n|_{\infty}$ and

$$\sigma,\tau) \cdot I_{k} = |M_{n}^{\sigma,\tau}|_{\infty} - |M_{k}^{\sigma,\tau}|_{\infty} + f_{CS}(M_{k}^{\sigma,\tau})$$

$$= \sum_{i=1}^{n} x_{\sigma(i)}^{2} y_{\tau(i)}^{2} + \sum_{1 \le i < j \le n, k < j} (x_{\sigma(i)}^{2} y_{\tau(j)}^{2} + x_{\sigma(j)}^{2} y_{\tau(i)}^{2}) + \sum_{1 \le i < j \le k} 2x_{\sigma(i)} y_{\tau(j)} x_{\sigma(j)} y_{\tau(i)}.$$
(18)

Let $T_1 = \{\vec{t} = (t_1, \cdots, t_{d^2}) | 0 \le t_i \le 1, \sum_{i=1}^{d^2} t_i = 1\}$ be the principal region of the unit ball. For $\vec{t} \in T_1$, we denote by $I^{\vec{t}} = \sum_{i=1}^{d^2} t_i I_i$ the convex combination of $I_1, I_2, \cdots, I_{d^2}$, then we have the following result.

Theorem 1. For any two observables A and B with a state ρ on a d-dimensional Hilbert space H, the product of the Wigner-Yanase-Dyson-metric adjusted skew information of A and B obeys the following uncertainty relations:

(i)
$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) \ge \max_{\vec{t}\in T_{1}}\{I^{\vec{t}}\} \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^{2};$$

(ii) $I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) \ge qI_{1} + (1-q)\max_{\sigma,\tau\in\mathfrak{S}_{d^{2}}}\{(\sigma,\tau)I_{k}\} \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^{2}.$
(19)

where $\vec{t} \in T_1$ and $0 \le q \le 1$.

Proof. From FIG.1, we can see that $I_k - I_{k-1} = -(\sum_{i=1}^{k-1} x_i y_k + y_i x_k)^2 \leq 0$. Then the following descending sequence of inequalities holds:

$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) = I_1 \ge I_2 \ge \dots \ge I_{d^2} = |\text{Corr}_{\rho}^{c^{WYD}}(A,B)|^2.$$
(20)

Moreover, $I_1 \ge \max_{\sigma, \tau \in \mathfrak{S}_{d^2}} \{ (\sigma, \tau) I_k \}$ for any permutation pairs $(\sigma, \tau) \in \mathfrak{S}_{d^2} \times \mathfrak{S}_{d^2}$. Then we have the following

$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) \ge \max_{\sigma,\tau \in \mathfrak{S}_{d^2}} \{(\sigma,\tau)I_k\} \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^2.$$

$$(21)$$

The theorem follows from (20), (21) and the properties of the convex functions.

Next, we introduce the second uncertainty relation for the WYD-metric adjusted skew information. Recall the quantities proposed by Li et al. in [29]:

$$S_{pq} = S_{p,q-1} - \sum_{i,j \in \{p,q\}} x_i^2 y_j^2 + f_{CS}(M_{i,j \in \{p,q\}}),$$

$$S_{p+1,1} = S_{p,p-1} - \sum_{i,j \in \{p+1,1\}} x_i^2 y_j^2 + f_{CS}(M_{i,j \in \{p+1,1\}}),$$
(22)



FIG. 2: Explanation of S_{pq} . The Cauchy-Schwarz inequality is used sequentially from top to bottom in the direction of the arrow. Set $|M_n|_{\infty} = S_{10} = \sum_{i,j=1}^n x_i^2 y_j^2$ as the initial value, the second is $S_{21} = S_{10} - \sum_{i,j \in \{2,1\}} x_i^2 y_j^2 + f_{CS}(M_{\{2,1\}})$; the third is $S_{31} = S_{21} - \sum_{i,j \in \{3,1\}} x_i^2 y_j^2 + f_{CS}(M_{\{3,1\}})$; the fourth is $S_{32} = S_{31} - \sum_{i,j \in \{3,2\}} x_i^2 y_j^2 + f_{CS}(M_{\{3,2\}})$; the fifth is $S_{41} = S_{32} - \sum_{i,j \in \{4,1\}} x_i^2 y_j^2 + f_{CS}(M_{\{4,1\}})$, and the last term $S_{n,n-1} = S_{n,n-2} - \sum_{i,j \in \{n,n-1\}} x_i^2 y_j^2 + f_{CS}(M_{\{n,n-1\}})$.

where $0 \le q \le p-1$ (see FIG. 2 for an explanation). For example,

$$S_{21} = \sum_{i,j=1}^{n} x_i^2 y_j^2 - x_1^2 y_2^2 - x_2^2 y_1^2 + 2x_1 x_2 y_1 y_2;$$

$$S_{31} = \sum_{i,j=1}^{n} x_i^2 y_j^2 - x_1^2 y_2^2 - x_2^2 y_1^2 - x_1^2 y_3^2 - x_3^2 y_1^2 + 2x_1 x_2 y_1 y_2 + 2x_1 x_3 y_1 y_3.$$
(23)

By the randomness of the entries in M_n along the horizontal direction in FIG.2, we have

$$(\sigma,\tau) \cdot S_{pq} = (\sigma,\tau) \cdot S_{p,q-1} - \sum_{i,j \in \{\sigma(p),\tau(q-1)\}} x_i y_j + f_{CS}(M_{i,j \in \{\sigma(p),\tau(q-1)\}}),$$
(24)

along the left slash direction in FIG.2 we also have

$$(\sigma,\tau) \cdot S_{p+1,1} = (\sigma,\tau) \cdot S_{p,p-1} - \sum_{i,j \in \{\sigma(p+1),\tau(1)\}} x_i y_j + f_{CS}(M_{\{\sigma(p+1),\tau(1)\}}).$$
(25)

Let $S^{\vec{t}} = \sum_{1 \le j < i \le n} t_{ij} S_{ij}$ for $\vec{t} = (t_{10}, t_{21}, t_{31}, t_{32}, t_{41}, t_{42} \cdots)$, then we have the following result.

Theorem 2. For any two observables A and B associated to a state ρ on a d-dimensional Hilbert space H, the following uncertainty relations hold for the Wigner-Yanase-Dyson-metric adjusted skew information:

(i)
$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) \ge \max_{t \in T_2} \{S^t\} \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^2,$$

(ii) $I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) \ge qS_{10} + (1-q)\max_{\sigma,\tau \in \mathfrak{S}_{d^2}} \{(\sigma,\tau)S_{pq}\} \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^2,$
(26)

where $T_2 = \{ \vec{t} = (t_{10}, t_{21}, \cdots, t_{pq}, \cdots) | 0 \le t_{ij} \le 1, \sum_{1 \le j < i \le d^2} t_{ij} = 1 \}.$

Proof. By (22) and the Cauchy-Schwarz inequality, we have the following sequence of inequalities (cf. [29]):

$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) = S_{10} \ge S_{21} \ge \dots \ge S_{d^2, d^2 - 1} \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A, B)|^2,$$
(27)

where the subscript (k, l) of $S_{k,l}$ satisfies k > l.

As $S_{pq} \leq \max_{\sigma, \tau \in \mathfrak{S}_{d^2}} \{ (\sigma, \tau) \cdot S_{pq} \}$, we get the following inequalities:

$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) \ge \max_{\sigma,\tau \in \mathfrak{S}_{d^2}} \{(\sigma,\tau)S_{pq}\} \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^2.$$

$$(28)$$

The results of the theorem follow easily from (27), (28) and the properties of convex functions.

Note that Theorem 2 is a refinement of Theorem 1, as $S_{k,k-1} = I_k$.

We will now derive a third uncertainty relation for the WYD-metric adjusted skew information. The quantities obtained by sampling observable coordinates in [30] are given as follows:

$$K_k = (|\vec{x}_k| |\vec{y}_k| + |\vec{x}_k^c| |\vec{y}_k^c|)^2, \quad k = 1, \cdots, n.$$
⁽²⁹⁾

Explicitly

$$K_{n} = |\vec{x}_{n}|^{2} |\vec{y}_{n}|^{2} = I_{\rho}^{c^{WYD}}(A) I_{\rho}^{c^{WYD}}(B);$$

$$K_{1} = \left(\sqrt{x_{1}^{2}y_{1}^{2}} + \sqrt{(x_{2}^{2} + \dots + x_{n}^{2})(y_{2}^{2} + \dots + y_{n}^{2})}\right)^{2};$$

$$K_{2} = \left(\sqrt{(x_{1}^{2} + x_{2}^{2})(y_{1}^{2} + y_{2}^{2})} + \sqrt{(x_{3}^{2} + \dots + x_{n}^{2})(y_{3}^{2} + \dots + y_{n}^{2})}\right)^{2}.$$
(30)

Due to randomness of data stored, we consider

$$\sigma(K_k) = (|\sigma(\vec{x}_k)||\sigma(\vec{y}_k)| + |\sigma(\vec{x}_k^c)||\sigma(\vec{y}_k^c)|)^2.$$

$$(31)$$

Let $T_3 = \{\vec{t} = (t_1, \cdots, t_{d^2}) | \sum_{k=1}^{d^2} t_k = 1, t_k \ge 0\}$. For $\vec{t} \in T_3$, we define the convex combination

$$K^{\vec{t}} = \sum_{k=1}^{d^2} t_k K_k.$$
 (32)

Theorem 3. For arbitrary observables A and B, and a state ρ on a d-dimensional Hilbert space H, let $\tilde{K}_k = \max_{\sigma \in \mathfrak{S}_{d^2}} \{\sigma(K_k)\}$ and $\tilde{K} = \max_k \{\tilde{K}_k\}$ for $k \in \{1, \dots, d^2\}$. Then, we can obtain the following inequality:

(i)
$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) \ge \tilde{K} \ge \tilde{K}_{k} \ge K_{k} \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^{2};$$

(ii) $I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) \ge \max_{\vec{t} \in T_{3}} \{K^{\vec{t}}\} \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^{2}.$
(33)

Proof. It follows from the Cauchy-Schwarz inequality that

$$K_{k} = |\vec{x}_{k}|^{2} \cdot |\vec{y}_{k}|^{2} + |\vec{x}_{k}^{c}|^{2} \cdot |\vec{y}_{k}^{c}|^{2} + 2|\vec{x}_{k}| \cdot |\vec{y}_{k}| \cdot |\vec{x}_{k}^{c}| \cdot |\vec{y}_{k}^{c}|$$

$$\geq (\vec{x}_{k}, \vec{y}_{k})^{2} + (\vec{x}_{k}^{c}, \vec{y}_{k}^{c})^{2} + 2(\vec{x}_{k}, \vec{y}_{k})(\vec{x}_{k}^{c}, \vec{y}_{k}^{c}) = (\vec{x}, \vec{y})^{2},$$
(34)

where the equality holds if and only if \vec{x}_k and \vec{y}_k are proportional. Also by the fundamental equality

$$K_k \le |\vec{x}_k|^2 \cdot |\vec{y}_k|^2 + |\vec{x}_k^c|^2 \cdot |\vec{y}_k^c|^2 + (|\vec{x}_k| \cdot |\vec{y}_k^c|)^2 + (|\vec{y}_k| \cdot |\vec{x}_k^c|)^2 = (\vec{x}, \vec{x})(\vec{y}, \vec{y}),$$
(35)

where the equality holds if and only if $|\vec{x}_k||\vec{y}_k^c| = |\vec{x}_k^c||\vec{y}_k|$. Therefore,

$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) = (\vec{x}, \vec{x}) \cdot (\vec{y}, \vec{y}) \ge K_k \ge (\vec{x}, \vec{y})^2 = |\text{Corr}_{\rho}^{c^{WYD}}(A, B)|^2.$$
(36)

Similarly, we also have

$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) \ge \sigma(K_k) \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^2.$$
(37)

This proves (a), from which (b) follows easily in view of convex functions.

Note that $\tilde{K}_k = \tilde{K}_{n-k}$ by (31), so it is enough to consider \tilde{K}_k for $k = 1, \dots, [\frac{n}{2}]$. Set $\tilde{K}_k^q = q\tilde{K}_k + (1-q)\tilde{K}_n$ for $q \in [0, 1]$, then we have

$$I_{\rho}^{c^{WYD}}(A)I_{\rho}^{c^{WYD}}(B) \ge \tilde{K}_{k}^{q} \ge \tilde{K}_{k} \ge |\operatorname{Corr}_{\rho}^{c^{WYD}}(A,B)|^{2}.$$
(38)

The arguments presented above demonstrate that our new uncertainty bounds, formulated in terms of convex functions, are tighter than previous ones. This is illustrated clearly in the following examples. We note that our uncertainty relations are state-dependent and apply to specific states in the quantum system. However, the comparison provided below is valid in general, and we have selected the state and observables solely for the purpose of illustration.



FIG. 3: The lower bounds (LB) of $I_{\rho}^{c}(A)I_{\rho}^{c}(B)$ for the mixed state $\rho(\theta)$ in Example 1. $I_{\rho}^{c}(A)I_{\rho}^{c}(B)$, $|\operatorname{Corr}_{\rho}^{c}(A,B)|^{2} = I_{2} = S_{31}$ and our bound K_{2} are shown in bold green, dashed blue and red curves respectively.

Example 1. Let $\rho := \rho(\theta) = \frac{1}{2}(I + \vec{r} \cdot \vec{\sigma})$ be a mixed state with $\vec{r} = (\frac{\sqrt{3}}{3}\cos\theta, \frac{\sqrt{3}}{3}\sin\theta, 0)$, where $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ is the vector of Pauli matrices. Consider observables $A = \sigma_x - \frac{1}{2}\sigma_z$ and $B = \sigma_x + \sigma_y + \sigma_z$. Under the standard orthogonal basis $\{e_1 = E_{11}, e_2 = E_{12}, e_3 = E_{21}, e_4 = E_{22}\}$, we have $\Gamma_{ij} = \operatorname{Corr}_{\rho}^c(e_i, e_j) = \frac{1}{2}\sigma_z$.

Under the standard orthogonal basis $\{e_1 = E_{11}, e_2 = E_{12}, e_3 = E_{21}, e_4 = E_{22}\}$, we have $\Gamma_{ij} = \operatorname{Corr}^c_\rho(e_i, e_j) = -\frac{1}{2}\operatorname{Tr}([\rho^{\frac{1}{4}}, e_i^T][\rho^{\frac{3}{4}}, e_j])$, where $c = c^{WYD}$ with p = 1/4 (see (7)). The lower bounds I_k , S_{pq} and K_2 can be calculated by (16), (22) and (30) respectively. Explicitly $I^c_\rho(A)I^c_\rho(B) = I_1 = S_{41} = K_2$ and $|\operatorname{Corr}^c_\rho(A, B)|^2 = I_2 = I_3 = I_4 = S_{21} = S_{31} = S_{32} = S_{42} = S_{43}$. The bounds are drawn in figure 3 which shows that our bound K_2 is the best.

To further demonstrate that our bounds are sharper than the previously available results, we chart the differences of our bounds with the previous bounds in the following example.

Example 2. Consider the pure state $|\psi(\theta)\rangle = \cos \theta |0\rangle - \sin \theta |2\rangle$ on a 3-dimensional Hilbert space. Let observables A and B be

$$A = \begin{bmatrix} 1 & 1-i & 0\\ 1+i & -1 & i\\ 0 & -i & 0 \end{bmatrix}, \qquad B = \begin{bmatrix} 0 & i & 1-i\\ -i & 0 & 1\\ 1+i & 1 & 0 \end{bmatrix}.$$

The bounds I_2 , S_{31} and $K^{(0,0.1,0,0.9)}$ for the uncertainty relations (for p = 1/3) are computed by (17), (23) and (??), and are shown in Figure 4. Their differences from the other bounds are drawn in Figure 5, which clearly shows that ours are the strongest.



FIG. 4: The lower bounds of $I_{\rho}^{c}(A)I_{\rho}^{c}(B)$ for the quantum state $|\psi(\theta)\rangle$ in Example 2. $I_{\rho}^{c}(A)I_{\rho}^{c}(B)$ and $|\operatorname{Corr}_{\rho}^{c}(A,B)|^{2}$ are shown in green and black bold curves respectively. The bounds I_{2} , S_{31} and $K^{(0,0.1,0,0.9)}$ are shown in blue, purple and red dashed curves respectively, and are seen to be the closest to the product.



FIG. 5: The differences of the lower bounds (DLB) in FIG.4. The blue and purple curves represent the differences $K_2^{(0,0.1,0,0.9)} - I_2$ and $K_2^{(0,0.1,0,0.9)} - S_{21}$ respectively, thus $K_2^{(0,0.1,0,0.9)}$ is the best bound.

IV. IMPROVED UNCERTAINTY RELATIONS FOR SUM-FORM SKEW INFORMATION

In this section, we consider *m* observables $\{A^i\}_{i=1}^m$ and a quantum state ρ on a *d*-dimensional Hilbert space *H*. Let $n = d^2$ as before. For each observable A^i , set $\vec{\alpha}^i = C_{\rho}\vec{A}^i = (\alpha_{i1}, \cdots, \alpha_{in})^{\dagger}$ and $\vec{X}^i = (x_{i1}, \cdots, x_{in})^T$ with $x_{ij} = |\alpha_{ij}|$, where C_{ρ} is the matrix of the WYD-metric with respect to the state ρ given as in (12). Then,

$$\sum_{i=1}^{m} I_{\rho}^{c^{WYD}}(A^{i}) = \sum_{i=1}^{m} |\vec{X}_{i}|^{2} = \sum_{i=1}^{m} \sum_{k=1}^{n} x_{ik}^{2}.$$
(39)

Similarly, the elements $x_{ij}^2(i = 1, \dots, m, j = 1, \dots, n)$ can be stored in a matrix. Let $M_{mn} = [x_{ij}^2]_{m \times n}$, and $B_2((a,b), (c,d)) = |M_{mn}|_{\infty} - (x_{ab}^2 + x_{cd}^2) + 2x_{ab}x_{cd}$. Set $B_2 = \max\{B_2((a,b), (c,d))|a, c \in \{1, \dots, m\}, b, d \in \{1, \dots, n\}$ and $(a,b) \neq (c,d)\}$. For $q \in [0,1]$, we define $B_2^q = qB_2 + (1-q)\sum_{i=1}^m |\vec{X}^i|^2$. Then we have the following strong uncertainty relation for the sum form WYD-metric adjusted skew information.

Theorem 4. For arbitrary m observables A^i $(i = 1, \dots, m)$ and a quantum state ρ on a d-dimensional Hilbert space H, then the following sum form uncertainty relation holds for the WYD-metric adjusted skew information:

$$\sum_{i=1}^{m} I_{\rho}^{c^{WYD}}(A^i) \ge B_2^q \ge B_2 \ge B_2((a,b),(c,d)),$$
(40)

$$\begin{bmatrix} x_{11}^2 & x_{12}^2 & x_{13}^2 & x_{14}^2 & \cdots & x_{1n}^2 \\ x_{21}^2 & x_{22}^2 & x_{23}^2 & x_{24}^2 & \cdots & x_{2n}^2 \\ x_{31}^2 & x_{32}^2 & x_{33}^2 & x_{34}^2 & \cdots & x_{3n}^2 \\ x_{41}^2 & x_{42}^2 & x_{43}^2 & x_{44}^2 & \cdots & x_{4n}^2 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ x_{m1}^2 & x_{m2}^2 & x_{m3}^2 & x_{m4}^2 & \cdots & x_{mn}^2 \end{bmatrix}$$

FIG. 6: Explanation of bound $B_2((a,b),(c,d))$. The FIG.6 shows $B_2((1,1),(4,1)) = \sum_{i=1}^m \sum_{j=1}^n x_{ij}^2 - (x_{11}^2 + x_{41}^2) + 2x_{11}x_{41}$.

Proof: First it is clear that

$$B_2((a,b),(c,d)) = \sum_{i=1}^m \sum_{j=1}^n x_{ij}^2 - (x_{ab}^2 + x_{cd}^2) + 2x_{ab}x_{cd} \le \sum_{i=1}^m \sum_{j=1}^n x_{ij}^2 = \sum_{i=1}^m I_{\rho}^{c^{WYD}}(A^i).$$
(41)

10

For N real non-negative numbers a_i^2 $(i = 1, \dots, N)$, according to the fundamental inequality,

$$N(\sum_{i=1}^{N} a_i^2) = \sum_{i=1}^{N} a_i^2 + \sum_{1 \le i < j \le N} (a_i^2 + a_j^2) \ge (\sum_{i=1}^{N} a_i)^2,$$
(42)

Therefore

$$\frac{m(m-1)}{2} \sum_{1 \le i < j \le m} (|\vec{X}^i - \vec{X}^j|^2) \ge (\sum_{1 \le i < j \le m} |\vec{X}^i - \vec{X}^j|)^2.$$
(43)

So we have the following uncertainty relation (cf. [24])

$$\sum_{i=1}^{m} |\vec{X}^{i}|^{2} = \frac{1}{2(m-1)} \sum_{1 \le i < j \le m} (|\vec{X}^{i} + \vec{X}^{j}|^{2} + |\vec{X}^{i} - \vec{X}^{j}|^{2})$$

$$\geq \frac{1}{2(m-1)} \Big[\sum_{1 \le i < j \le m} |\vec{X}^{i} + \vec{X}^{j}|^{2} + \frac{2}{m(m-1)} (\sum_{1 \le i < j \le m} |\vec{X}^{i} - \vec{X}^{j}|)^{2} \Big].$$
(44)

For simplicity, set $L_{Ma} = \frac{1}{2(m-1)} \left[\sum_{1 \le i < j \le m} |\vec{X}^i + \vec{X}^j|^2 + \frac{2}{m(m-1)} \left(\sum_{1 \le i < j \le m} |\vec{X}^i - \vec{X}^j| \right)^2 \right].$

Example 3. Let us consider the mixed state $\rho := \rho(\theta) = \frac{I_2 + \vec{r} \cdot \vec{\sigma}}{2}$ with $\vec{r} = (\frac{3}{4}\sin\theta, 0, \frac{3}{4}\cos\theta)$ and the following four observables A^i $(i = 1, \dots, 4)$

$$A^{1} = \begin{bmatrix} 1 & 2+i \\ 2-i & -1 \end{bmatrix}, \quad A^{2} = \begin{bmatrix} 1 & i \\ -i & -1 \end{bmatrix}, A^{3} = \begin{bmatrix} 0 & 1+\frac{1}{2}i \\ 1-\frac{1}{2}i & 0 \end{bmatrix}, \quad A^{4} = \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}.$$

The bounds $B_2((3,1),(4,1))$ and L_{Ma} can be calculated by (41) and (44) respectively. They are drawn in FIG. 7 and their differences are shown in FIG. 8, all are done with p = 1/3 in defining the WYD-metric adjusted skew information. It happens that $B_2((3,1),(1,1)) = \max\{B_2((a,b),(c,d))|a,c \in \{1,\cdots,m\}, b,d \in \{1,\cdots,n\} \text{ and } (a,b) \neq (c,d)\}$ in this case. The graph shows that our bound is tighter than the lower bound L_{Ma} in [24].



FIG. 7: The lower bounds of $\sum_{k=1}^{4} I_{\rho}^{c}(A^{i})$ for the quantum state $\rho(\theta)$ in Example 3. The quantity $\sum_{k=1}^{4} I_{\rho}^{c}(A^{i})$, the bound L_{Ma} from [24] and our bound $B_{2}((3,1),(4,1))$ are in green, black and red-dashed curves respectively.

Example 4. Let $|\psi(\theta)\rangle = \cos \theta |0\rangle - \sin \theta |2\rangle$ be a pure state on a 3-dimensional Hilbert space. Consider the four observables A^i $(i = 1, \dots, 4)$:

$$A^{1} = \begin{bmatrix} 1 & 1-i & 0\\ 1+i & -1 & i\\ 0 & -i & 0 \end{bmatrix}, \quad A^{2} = \begin{bmatrix} 0 & i & 1-i\\ -i & 0 & 1\\ 1+i & 1 & 0 \end{bmatrix}, A^{3} = \begin{bmatrix} 0 & 0 & 1-i\\ 0 & 0 & 1\\ 1+i & 1 & 0 \end{bmatrix}, \quad A^{4} = \begin{bmatrix} 2 & 1-i & 0\\ 1+i & -2 & 0\\ 0 & 0 & 0 \end{bmatrix}$$

Similarly, the bounds $B_2((2,3), (3,3))$, $B_2((3,2), (4,3))$ and L_{Ma} (with p = 1/3) can be calculated by (41) and (44). $B_2((2,3), (3,3)) = \max\{B_2((a,b), (c,d))|a, c \in \{1, \dots, m\}, b, d \in \{1, \dots, n\}$ and $(a,b) \neq (c,d)\}$. The bounds are drawn in FIG. 9 and the differences are drawn in FIG. 10. It is seen that our bound is tighter than the lower bound L_{Ma} in [24] and almost approximates the summation WYD-metric adjusted skew information.



FIG. 8: The difference of the lower bounds in FIG.7. The solid red curve represents $B_2((3,1),(4,1)) - L_{Ma}$, the difference of our new bound and the recent lower bound in [24], which reveals that our lower bound $B_2((3,1),(4,1))$ is tighter than that of L_{Ma} .



FIG. 9: The lower bounds of $\sum_{i=1}^{4} I_{\rho}^{c}(A^{i})$ for quantum state $|\psi(\theta)\rangle$ in Example 4. The quantity $\sum_{i=1}^{4} I_{\rho}^{c}(A^{i})$, the bound L_{Ma} from [24], our bounds $B_{2}((2,3), (3,3))$ and $B_{2}((3,2), (4,3))$ are shown in green, black, red (dashed) and blue (dashed) respectively.



FIG. 10: The differences of the lower bounds in FIG.9. The solid red and blue curves are the differences $B_2((2,3),(3,3)) - L_{Ma}$ and $B_2((3,2),(4,3)) - L_{Ma}$, which shows that our bounds are tighter.

V. CONCLUSION

The Wigner-Yanase-Dyson-metric adjusted skew information was introduced by Hansen [18] as a measure of quantum information, based on the geometrical formulation of quantum statistics by Chentsov and Morozova, and earlier work by Wigner and Yanase. This quantity is non-negative and bounded by the variance of an observable in a state, and vanishes for observables that commute with the state. In this work, we have reviewed and improved the "finegrained" bounds for the product form of the Wigner-Yanase-Dyson-metric adjusted skew information, using sampling observable coordinates and appropriate convex functions. Additionally, we have proposed uncertainty relations for the sum form of N quantum observables. Our method relies on straightening the key mathematical inequality underlying the uncertainty principle, resulting in new and stronger uncertainty relations in both the sum and product forms. Our results are competitive with the recent sharp bound presented in [24], as demonstrated by several examples.

Acknowledgments

We would like to thank Dr. Yunlong Xiao for helpful discussions. The research is supported in part by NSFC grants 12226321, 12126351, 12126314 and 11871325, and by Foundation of Jianghan University as well as Simons Foundation grant no. 523868.

Data Availability Statement. All data generated during the study are included in the article. **Conflict of Interest Statement.** The authors declare no competing interests.

- [1] P. A. M. Dirac, Proc. Royal Soc. London. Ser. A 117, 610-624 (1928).
- [2] W. Heisenberg, Zeit. für Physik, 43, 172–198 (1927).
- [3] E. H. Kennard, Zeit. für Physik, 44, 326–352 (1927).
- [4] H. P. Robertson, Phys. Rev. **34**, 163 (1929).
- [5] E. Schrödinger, Königliche Berlin Akad. Wissenschften 24, 296(1930).
- [6] E. P. Wigner and M. M. Yanase, Proc. Natl. Acad. Sci. USA 49, 910 (1963).
- [7] E. P. Wigner, Zeit. für Physik A Hadrons and nuclei 133, 101–108 (1952).
- [8] H. Araki and M. M. Yanase, Phys. Rev. 120, 622 (1960).
- [9] M. M. Yanase, Phys. Rev. **123**, 666 (1961).
- [10] M. Ozawa, Phys. Rev. Lett. 67, 1956 (1991).
- [11] K. Kakazu, S. Pascazio, Phys. Rev. A 51, 3469 (1995).
- [12] S. Matsumoto, Prog. Theor. Phys. 90, 35 (1993).
- [13] M. Ozawa, Phys. Rev. Lett. 88, 050402 (2002).
- [14] M. Ozawa, Phys. Rev. Lett. 89, 057902 (2002).
- [15] B. R. Frieden, Cambridge University, Cambridge, England, (1998).
- [16] M. Hayashi, Quantum Information: An Introduction (Springer Science & Business Media, 2006).
- [17] S. Luo, Phys. Rev. Lett. **91**, 180403 (2003).
- [18] F. Hansen, Proc. Natl. Acad. Sci. USA 105, 9909-9916(2008).
- [19] L. Cai, Quant. Inf. Process. 20, 72 (2021).
- [20] B. Chen, S. M. Fei, and G. L. Long, Quant. Inf. Process. 15, 2639-2648 (2016).
- [21] R. Ren, P. Li, M. Ye, and Y. Li, Phys. Rev. A 104, 052414 (2021).
- [22] Z. Chen, H. Wang, J. Li, et al. Sci. Rep. 9, 5687 (2019).
- [23] Q. H. Zhang and S. M. Fei, Quant. Inf. Process. 20, 384 (2021).
- [24] X. Ma, Q. H. Zhang, and S. M. Fei, Laser Phys. Lett. 19, 055205 (2022).
- [25] D. Petz, Linear. Algebra Applied 244, 81-96 (1996).
- [26] E. H.Lieb, Adv. Math. 11, 267-288 (1973).
- [27] Y. Xiao, N. Jing, B. Yu, S. M. Fei, and X. Li-Jost, Front. Phys. 10, 846330 (2022).
- [28] B. Yu, N. Jing, and X. Li-Jost, Phys. Rev. A 100, 022116 (2019).
- [29] J. Li, S. Zhang, L. Liu, and C. M. Bai, Laser Phys. Lett. 17, 015201 (2020).
- [30] X. Hu, N. Jing, Quant. Inf. Process. 21, 52 (2022).