Gonality of curves whose normalizations are one or two copies of \mathbb{P}^1

Juliana Coelho

julianacoelhochaves@id.uff.br

October 27, 2023

Abstract

We study the gonality of curves C over \mathbb{C} whose normalization consist of one or two copies of \mathbb{P}^1 . In the first case, C is a nodal curve with g(C)nodes, and in the second case C is a so-called binary curve. In any case we show that the usual bound $gon(C) \leq \left\lfloor \frac{g(C)+3}{2} \right\rfloor$ holds if $g(C) \geq 2$, with equality holding generically.

1 Introduction

A smooth curve C is said to be k-gonal if it admits a degree-k map to \mathbb{P}^1 . The gonality gon(C), that is, the minimum k such that C is k-gonal, is an important numerical invariant in the study of algebraic curves. When considering smooth curves in families, one feels the need to consider singular curves as well, since smooth curves can degenerate to singular ones.

There are a few possible ways to extend the notion of gonality to singular curves. One way is to simply take the same definition as in the smooth case, as is done in [16] and [14]. The drawback here is that this definition does not work very well in families, since not all curves that are limits of smooth k-gonal ones have maps of degree k to \mathbb{P}^1 . One solution is to consider limits of maps from smooth curves to \mathbb{P}^1 . This can be done using stable maps, as in [1], or using Harris and Mumford's admissible covers [12]. In this paper we adopt the latter point of view, as was done previously in [4], where admissible covers were used to obtain bounds on the gonality of stable curves from the gonality of its components, and [5], where two-component nodal curves of gonality 2 or 3 were characterized, with the simplifying hypothesis that the components are not rational.

It is worth mentioning other approaches to the problem. For instance, since a map from a smooth curve to \mathbb{P}^1 is given by a linear series on the curve, one can consider Eisenbud and Harris' limit linear series introduced in [7] (see for instance [8] and [15]). Furthermore, some authors have been working on relating the gonality of stable curves to the gonality of graphs and tropical curves (see [3]).

1.1 Main Results

In this paper we always work over the field of complex numbers \mathbb{C} , and we study the gonality of nodal curves C whose normalizations are one (in Subsection 3.1) or two (in Subsection 3.2) copies of \mathbb{P}^1 . In the first case, C is an irreducible curve with g(C) nodes, and in the second case, C is a so-called binary curve.

Both types of curves have been extensively used as degeneration targets when proving results that hold generically, since they are combinatorically easier to handle. For instance, in [10] the Brill-Noether theory of a general curve was studied by degenerating it to a irreducible curve of genus g with g nodes. In the same way, Prym maps we studied in [6] via degeneration to binary curves. Moreover, these curves were also used as test models for results that hold for smooth curves but are yet unknown for singular ones, as in [2], [9] and [13].

In Section 2.1 we review some background, and we prove some auxiliary results on admissible covers (see Subsection 2.2) and maps between \mathbb{P}^1 (see Subsection 2.3). In Section 3 we prove the main results of the paper, which are Theorems 6 and 8.

Denote by I_g (resp. B_g) the locus of irreducible curves with g nodes in the moduli of genus-g stable curves \overline{M}_g .

Theorem. Let C be an irreducible curve with g nodes (resp. a binary curve) having genus $g \ge 2$. Then

$$\operatorname{gon}(C) \le \left\lfloor \frac{g+3}{2} \right\rfloor$$

and equality holds if C is general in I_g (resp. B_g).

2 Preliminaries

2.1 Technical background

In this paper we always work over the field of complex numbers \mathbb{C} .

A curve C is a connected, projective and reduced scheme of dimension 1 over \mathbb{C} . The genus of C is $g(C) := h^1(C, \mathcal{O}_C)$. A subcurve Y of C is a reduced subscheme of pure dimension 1, or equivalently, a reduced union of irreducible components of C. If $Y \subseteq C$ is a subcurve, we say $Y^c := \overline{C \setminus Y}$ is the complement of Y in C.

A nodal curve C is a curve with at most ordinary double points, called nodes. A node is said to be separating if there is a subcurve Y of C such that $Y \cap Y^c = \{n\}$. A rational chain is a nodal curve of genus 0. We remark that a curve is a rational chain if and only if its nodes are all separating and its components are all rational.

Let g and n be non negative integers such that 2g - 2 + n > 0. A *n*-pointed stable curve of genus g is a curve C of genus g together with n distinct marked points $p_1, \ldots, p_n \in C$ such that for every smooth rational component E of C,

the number of points in the intersection $E \cap E^c$ plus the number of indices *i* such that p_i lies on E is at least three. A stable curve is a 0-pointed stable curve. We denote by \overline{M}_g the moduli space of stable curves of genus g.

Moreover, we denote by $M_{0,n}$ the moduli space of n-pointed smoth curves of genus 0. Such a pointed curve is simply \mathbb{P}^1 with a choice a n-uple of distinct points, where two n-uples give the same pointed curve in $M_{0,n}$ if there is an automorphism of \mathbb{P}^1 sending one to the other. More preciselly, if Δ is the set of n-uples $(p_1, \ldots, p_n) \in (\mathbb{P}^1)^n$ such that $n_i \neq n_j$ for all $i \neq j$, then there is a surjective map

$$P_n := (\mathbb{P}^1)^n \setminus \Delta \to M_{0,n} \tag{1}$$

whose fibers have dimension 3. In particular, $\dim(M_{0,n}) = n - 3$.

2.2 Gonality and admissible covers

A smooth curve is k-gonal if it admits a map of degree k to \mathbb{P}^1 . A stable curve C is k-gonal if it is a limit of smooth k-gonal curves in the moduli space \overline{M}_g of stable curves. More precisely, a *smoothing* of a curve C is a proper and flat morphism $f: \mathcal{C} \to \operatorname{Spec}(\mathbb{C}[[t]])$ whose fibers are curves, with special fiber C and such that C is regular. A stable curve C is then said to be k-gonal if it admits a smoothing $f: \mathcal{C} \to S$ whose general fiber is a k-gonal smooth curve and the special fiber is C.

For the sake of completeness, we remark that some authors consider any curve to be k-gonal if it admits a degree-k map to \mathbb{P}^1 . Altough this is a simpler definition, it does not necessarily work well in families.

The gonality of a stable curve C, denoted gon(C), is the smallest k such that C is k-gonal. We say that C is hyperelliptic if gon(C) = 2. It is a well-known fact that all stable curves of genus 2 are hyperelliptic.

Alternatively, k-gonal stable curves can be characterized in terms of admissible covers. A k-sheeted admissible cover consists of a finite morphism $\pi : C \to B$ of degree k, such that B and C are nodal curves and:

- 1. $\pi^{-1}(B^{\text{sing}}) = C^{\text{sing}};$
- 2. π is simply branched away from C^{sing} , that is, over each smooth point of B there exists at most one point of C where π is ramified and this point has ramification index 2;
- 3. *B* is a stable pointed curve of genus 0, when considered with its smooth points on the branch locus of π ;
- 4. for every node q of B and every node n of C lying over it, the two branches of C over n map to the branches of B over q with the same ramification index.

Let C be a nodal curve. We say that a nodal curve C' is stably equivalent to C if C can be obtained from C' by contracting to a point some of the smooth rational components of C' meeting the other components of C' in only one or two

points. In that case, g(C) = g(C') and there is a contraction map $\tau: C' \to C$. We say that a point $p' \in C'$ lies over a point $p \in C$ if $\tau(p') = p$.

We remark that, if C' is stably equivalent to a curve C then, by [4, Lemma

3.2], we can consider the smooth components of C as components of C' as well. The following result is a consequence of [12, Thm. 4, p. 58] and relates the notion of admissible covers to that of gonality of stable curves.

Theorem 1 (Harris-Mumford). A stable curve C is k-gonal if and only if there exists a k-sheeted admissible cover $C' \to B$, where C' is stably equivalent to C.

Proof. Cf. [11, Thm. 3.160, p. 185].

The next two thechnical results are similar to [5, Proposition 10] and will be used in Subsections 3.1 and 3.2.

Lemma 2. Let C be an irreducible nodal curve with nodes n_1, \ldots, n_{δ} . Let $\pi: C' \to B$ be an admissible cover, where C' is stably equivalent to C. Let C^{ν} be the normalization of C and let $n_j^{(1)}, n_j^{(2)} \in C^{\nu}$ be the branches of n_j , for $j = 1, \ldots, \delta$. Then C^{ν} is a component of C' and

$$\deg(\pi) \ge \deg(\pi|_{C^{\nu}}) + \ell,$$

where ℓ is the cardinality of $\{1 \le j \le \delta \mid \pi(n_j^{(1)}) \ne \pi(n_j^{(2)})\}$.

Proof. By [4, Lemma 3.2], C^{ν} can be considered as a component of C'. Set $B' = \pi(C^{\nu})$ and let k' be the degree of $\pi|_{C^{\nu}} : C^{\nu} \to B'$.

Now, for each $1 \leq j \leq \delta$ let $W_j := \tau^{-1}(n_j)$, where $\tau : C' \to C$ is the contraction map. Then W_j is a rational subcurve of C' containing both $n_j^{(1)}$ and $n_j^{(2)}$ and thus $\pi(W_j)$ contains both $\pi(n_j^{(1)})$ and $\pi(n_j^{(2)})$.

If $\pi_1(n_j^{(1)}) \neq \pi_2(n_j^{(2)})$, since $\pi(W_j)$ contains two distinct points $\pi(n_j^{(1)})$ and $\pi(n_j^{(2)})$ on B', then it must contain the entire line B'. So, for a general point $q \in B', \pi^{-1}(q)$ contains at least one point on W_j . Therefore $\pi^{-1}(q)$ contains k' points on C^{ν} , counted with multiplicity, and at least one point on W_j for each $1 \leq j \leq \delta$ such that $\pi_1(n_j^{(1)}) \neq \pi_2(n_j^{(2)})$, and the result is proven.

Lemma 3. Let C be a nodal curve with irreducible smooth components C_1 and C_2 and nodes n_1, \ldots, n_{δ} . For i = 1, 2 let $n_j^{(i)}$ be the branch of n_j at C_i , for $j = 1, \ldots, \delta$. Let $\pi: C' \to B$ be an admissible cover, where C' is stably equivalent to C.

- i. If $\pi(C_1) \neq \pi(C_2)$ then $\deg(\pi) \geq \delta$.
- ii. If $\pi(C_1) = \pi(C_2)$, let ℓ be the cardinality of the set

$$\{1 \le j \le \delta \mid \pi(n_j^{(1)}) \ne \pi(n_j^{(2)})\}.$$

Then $\deg(\pi) \ge \deg(\pi|_{C_1}) + \deg(\pi|_{C_2}) + \ell$.

Proof. For each $1 \leq j \leq \delta$ let $W_j := \tau^{-1}(n_j)$, where $\tau : C' \to C$ is the contraction map. Then W_j is a rational subcurve of C' containing both $n_j^{(1)}$ and $n_j^{(2)}$ and thus $\pi(W_j)$ contains both $\pi(n_j^{(1)})$ and $\pi(n_j^{(2)})$. For each i = 1, 2, let k_i be the degree of $\pi_i := \pi|_{C_i}$ and set $B_i = \pi(C_i)$.

First we prove (i), so assume $B_1 \neq B_2$. By [5, Lemma 9], either $B_1 \cap B_2$ is a point or $B_0 = B_1^c \cap B_2^c$ is a rational chain and $B_i \cap B_0$ is a unique point. In any case, we thus determine a unique point $q_i \in B_i$.

For each $1 \leq j \leq \delta$ such that $n_j^{(i)} \neq q_i$, the image $\pi(W_j)$ contains B_i . Indeed, $\pi(W_j)$ contains $\pi(n_j^{(1)}) \in B_1$ and $\pi(n_j^{(2)}) \in B_2$ and hence it must also contain q_i . Since $\pi(W_j)$ contains two distinct points of B_i , namely $\pi(n_j^{(i)})$ and q_i , it must contain the entire line B_i .

Note that there are at most k_i indexes j such that $\pi(n_j^{(i)}) = q_i$, that is, there at least $\delta - k_i$ indexes j such that $\pi(n_j^{(i)}) \neq q_i$. Moreover, for a general point $q \in B_i$, $\pi^{-1}(q)$ contains k_i points on C_i , counted with multiplicity, and at least one pont on W_j for each $1 \leq j \leq \delta$ such that $\pi(n_j^{(i)}) \neq q_i$. Hence $\deg(\pi) \geq k_i + \delta - k_i = \delta$.

Now we show (ii), so assume $B_1 = B_2$. Let $1 \le j \le \delta$ be such that $\pi_1(n_j^{(1)}) \ne \pi_2(n_j^{(2)})$. Since $\pi(W_j)$ contains two distinct points $\pi_1(n_j^{(1)})$ and $\pi_2(n_j^{(2)})$ on B_i , then it must contain the entire line B_i . So, for a general point $q \in B_i$, $\pi^{-1}(q)$ contains at least one point on W_j . Therefore $\pi^{-1}(q)$ contains k_1 points on C_1 and k_2 points on C_2 , counted with multiplicity, and at least one point on W_j for each $1 \le j \le \delta$ such that $\pi_1(n_j^{(1)}) \ne \pi_2(n_j^{(2)})$, and the result is proven.

The following result is a slight improvement upon [4, Proposition 3.11], and will be used in Subsection 3.2.

Proposition 4. Let C be a stable curve with p smooth irreducible components C_1, \ldots, C_p . For each node n_j of C, let $C_{j(1)}$ and $C_{j(2)}$ be the irreducible components of C containing n_j and denote by $n_j^{(1)}$ and $n_j^{(2)}$ the branches of n_j on $C_{j(1)}$ and $C_{j(2)}$, respectively. Assume that there is a degree- k_s map $\pi_s \colon C_s \to \mathbb{P}^1$ for each component C_s of C such that $\pi_{j(1)}(n_j^{(1)}) = \pi_{j(2)}(n_j^{(2)})$ for every node n_j . Then C is $(k_1 + \ldots + k_p)$ -gonal.

Proof. Let n_1, \ldots, n_{δ} be the nodes of C and let

$$Q = \{\pi_{j(1)}(n_j^{(1)}) \mid j = 1, \dots, \delta\}.$$

For each $q \in Q$, fix one $1 \leq j(q) \leq \delta$ such that $\pi_{j(q)(1)}(n_{j(q)}^{(1)}) = q$.

Let $J = \{1, \ldots, \delta\} \setminus \{j(q) \mid q \in Q\}$ and let C_J be the curve obtained by normalizing C at the nodes n_j such that $j \in J$. Note that C_J may be disconnected. However, applying [4, Proposition 3.11] to C_1, \ldots, C_p and the nodes $n_{j(q)}$ for $q \in Q$, we obtain a finite map $\pi'_J \colon C'_J \to B$ of degree $k_1 + \ldots + k_p$, where C'_J is stably equivalent to C_J , satisfying conditions 1 to 4 of the definition of an admissible covering. By construction, $\pi'_J|_{C_s} = \pi_s$ for every component C_s of C. Furthermore, it is easy to see that [4, Proposition 3.4] can be applied to a map such as π_J , even if C'_J is not connected. We thus proceed in the following manner. Write $J = \{j_1, \ldots, j_r\}$ and for each $\ell = 1, \ldots, r$ let $C(\ell)$ be the curve obtained by normalizing C at the nodes $n_{j_\ell}, \ldots, n_{j_r}$, and let C(r+1) = C. We will show by induction that for every $1 \leq \ell \leq r+1$ there is a finite map $\pi'_\ell \colon C'(\ell) \to B_\ell$ of degree $k_1 + \ldots + k_p$ satisfying conditions 1 to 4 of the definition of an admissible covering, and such that $C'(\ell)$ is stably equivalent to $C(\ell)$ and $\pi'_\ell|_{C_s} = \pi_s$ for every component C_s of C. Note that π'_J satisfies the statement for $\ell = 1$.

Assume that the statement holds for $\ell - 1$. Choose points $n'_{j_{\ell}}^{(1)}$ and $n'_{j_{\ell}}^{(2)}$ on $C'(\ell - 1)$ lying over $n_{j_{\ell}}^{(1)}$ and $n_{j_{\ell}}^{(2)}$, respectively, such that $\pi'_{\ell-1}(n'_{j_{\ell}}^{(1)}) = \pi'_{\ell-1}(n'_{j_{\ell}}^{(2)})$. Then [4, Theorem 3.4 (b)] gives the required admissible cover π'_{ℓ} as stated.

The case $\ell = r + 1$ shows that C is $(k_1 + \ldots + k_p)$ -gonal.

2.3 Maps between \mathbb{P}^1

In the following section we will study the gonality of nodal curves whose nornalizations are one or two copies of \mathbb{P}^1 . For that end, we will first examine endomorphisms of \mathbb{P}^1 .

Recall that set of non-constant morphisms $\varphi \colon \mathbb{P}^1 \to \mathbb{P}^1$ of degree at most k can be identified with the Grassmanian $\mathcal{G}_k := Gr(2, H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(k)))$ of 2dimensional subspaces of $H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(k))$. Indeed, such a morphism corresponds to a linear system of degree k and (projective) dimension 1, which, up to change of projective coordinates, corresponds to a linear subspaces of $H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(k))$ of dimension 2. Since $H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(k))$ is a vector space of dimension k + 1, we get in particular that $\dim(\mathcal{G}_k) = 2k - 2$.

Lemma 5. Let $k, \ell \geq 1$ be integers such that $\ell \leq 2k-2$ and let $n_1^{(i)}, \ldots, n_\ell^{(i)} \in \mathbb{P}^1$, for i = 1, 2.

- *i.* There exists $\varphi \in \mathcal{G}_k$ such that $\varphi(n_j^{(1)}) = n_j^{(2)}$ for every $j = 1, \ldots, \ell$.
- ii. There exists $\varphi \in \mathcal{G}_k$ such that $\varphi(n_j^{(1)}) = \varphi(n_j^{(2)})$ for every $j = 1, \ldots, \ell$.

Proof. For $j = 1, \ldots, \ell$ we let V_j (resp. W_j) be the set of $\varphi \in \mathcal{G}_k$ such that $\varphi(n_j^{(1)}) = n_j^{(2)}$ (resp. $\varphi(n_j^{(1)}) = \varphi(n_j^{(2)})$). Then V_j (resp. W_j) is a hyperplane (resp. quadric hypersurface) in \mathcal{G}_k . Since $\ell \leq \dim(\mathcal{G}_k)$, the intersection $V_1 \cap \cdots \cap V_\ell$ (resp. $W_1 \cap \cdots \cap W_\ell$) is non-empty and any point on this intersection gives the desired φ , thus showing (i) (resp. (ii)).

3 Results on gonality

In [5], the gonality of two-component nodal curves is studied, focusing on the case where both components are non-rational. In this work we study the case of

nodal curves whose normalization are one (Subsection 3.1) or two (Subsection 3.2) copies of \mathbb{P}^1 .

3.1 Irreducible curves of genus g with g nodes

In this subsection we study the gonality of irreducible curves C having g(C) nodes. We note that the normalization of such a curve is \mathbb{P}^1 . Indeed, the curve C is obtained from \mathbb{P}^1 by identifying g(C) pairs of distinct points on \mathbb{P}^1 .



An irreducible curve of genus 3 with 3 nodes.

Denote by I_g the locus in \overline{M}_g of irreducible curves of genus g with g nodes. **Theorem 6.** Let C be an irreducible curve with g(C) nodes. Then

$$\operatorname{gon}(C) \le \left\lceil \frac{g(C) + 3}{2} \right\rceil$$

and equality holds if C is general in I_g .

Proof. Let C be an irreducible curve of genus g = g(C) and nodes n_1, \ldots, n_g . Then the branches of the nodes give two g-uples of points $\underline{n}^{(i)} = (n_1^{(i)}, \ldots, n_g^{(i)}) \in (\mathbb{P}^1)^g$, such that

$$n_j^{(i)} \neq n_{j'}^{(i')}$$
 unless $(i,j) = (i',j')$ (2)

for i = 1, 2 and j = 1, ..., g.

By Lemma 5 (ii), there exists a non-constant morphism $\varphi \colon \mathbb{P}^1 \to \mathbb{P}^1$ of degree at most $k = \lfloor \frac{g+2}{2} \rfloor$ such that $\varphi(n_j^{(1)}) = \varphi(n_j^{(2)})$ for every $j = 1, \ldots, g$. Hence, by [4, Corollary 3.5], C is k-gonal and $gon(C) \leq k = \lfloor \frac{g+2}{2} \rfloor = \lceil \frac{g+3}{2} \rceil$, showing the first statement.

To show the second statement, we note that $P := P_{2g}$ is the set of $(\underline{n}^{(1)}, \underline{n}^{(2)}) \in (\mathbb{P}^1)^g \times (\mathbb{P}^1)^g$ satisfying (2) (see Subsection 2.1). Since any $(\underline{n}^{(1)}, \underline{n}^{(2)}) \in P$ gives

an irreducible curve of genus g with g nodes by identifying $n_j^{(1)}$ and $n_j^{(2)}$ for every $j = 1, \ldots, g$, then there is a surjective map $\Phi \colon P \to I_g$.

For each $k, r \in \mathbb{N}$, denote by P(k, r) the set of the pairs $(\underline{n}^{(1)}, \underline{n}^{(2)})$ in P such that there exists $I \subset \{1, \ldots, g\}$ of cardinality r and a non-constant map $\varphi \colon \mathbb{P}^1 \to \mathbb{P}^1$ of degree at most k satisfying $\varphi(n_j^{(1)}) = \varphi(n_j^{(2)})$ for all $j \in I$. Then $P(k, r) \subset P(k + 1, r), P(k, r)$ is closed in P and, by Lemma 5 (ii), we have $P(\lceil \frac{r+3}{2} \rceil, r) = P$ for every $1 \le r \le g$.

$$U := P \setminus \bigcup_{r=1}^{g} P\left(\left\lceil \frac{r+3}{2} \right\rceil - 1, r\right).$$

Since U is open and dense in P, its image $\Phi(U)$ contains an open an dense subscheme of I_g . Let C be an irreducible curve with g(C) = g nodes corresponding to a point $(\underline{n}^{(1)}, \underline{n}^{(2)}) \in U$. We will show that $\operatorname{gon}(C) = \lceil \frac{g+3}{2} \rceil$. Let $\pi \colon C' \to B$ be an admissible cover of degree $k := \operatorname{gon}(C)$, where C' is

Let $\pi: C' \to B$ be an admissible cover of degree k := gon(C), where C' is stably equivalent to C. Restricting π to the normalization $C^{\nu} = \mathbb{P}^1$ of C, we obtain a non-constant map $\varphi := \pi|_{C^{\nu}} : \mathbb{P}^1 \to \mathbb{P}^1$ of degree $k' := \text{deg}(\varphi)$. Let J be the set of $j \in \{1, \ldots, g\}$ such that $\varphi(n_j^{(1)}) \neq \varphi(n_j^{(2)})$ and let $\ell = |J|$. By Lemma 2, we have $k \geq k' + \ell$.

On the other hand, let γ be the cardinality of $I := \{1, \ldots, g\} \setminus J$, then $\ell + \gamma = g$. Since $(\underline{n}^{(1)}, \underline{n}^{(2)}) \in U$, from the existence of the degree- $k' \operatorname{map} \varphi$ such that $\varphi(n_j^{(1)}) = \varphi(n_j^{(2)})$ for every $j \in I$, we get that $k' \geq \lceil \frac{\gamma+3}{2} \rceil = \lceil \frac{g-\ell+3}{2} \rceil$. Since $k \leq \lceil \frac{g+3}{2} \rceil$, then

$$\left\lceil \frac{g-\ell+3}{2} \right\rceil + \ell \le k' + \ell \le k \le \left\lceil \frac{g+3}{2} \right\rceil$$

which implies that $\ell = 0$ and we must have $k = \lceil \frac{g+3}{2} \rceil$.

3.2 Binary curves

A binary curve C is a nodal curve made of two smooth rational components meeting at g + 1 points. Note that the genus of C is g(C) = g and its normalization is two disjoint copies of \mathbb{P}^1 .

More preciselly, if $n_1^{(i)}, \ldots, n_{g+1}^{(i)} \in \mathbb{P}^1$ are distinct points, we set $\underline{n}^{(i)} = (n_1^{(i)}, \ldots, n_{g+1}^{(i)})$, for i = 1, 2, and we denote by $C(\underline{n}^{(1)}, \underline{n}^{(2)})$ the binary curve obtained by gluing two copies of \mathbb{P}^1 at $n_j^{(1)}$ and $n_j^{(2)}$ for every j.



A binary curve of genus 2.

Two binary curves $C(\underline{n}^{(1)}, \underline{n}^{(2)})$ and $C(\underline{m}^{(1)}, \underline{m}^{(2)})$ are isomorphic if there are isomorphisms ψ_i of \mathbb{P}^1 such that $\psi_i(n_j^{(i)}) = m_j^{(i)}$ for every $j = 1, \ldots, n$ and i = 1, 2. Hence, if $B_g \subset \overline{M}_g$ is the locus of binary curves of genus $g \ge 2$, there exists a surjective morphism with finite fibers

$$M_{0,g+1} \times M_{0,g+1} \to B_g. \tag{3}$$

Proposition 7. Let $C = C(\underline{n}^{(1)}, \underline{n}^{(2)})$ be a binary curve of genus $g \ge 2$. Then C is hyperelliptic if and only if there exists an automorphism $\psi \colon \mathbb{P}^1 \to \mathbb{P}^1$ such that $\psi(n_j^{(1)}) = n_j^{(2)}$ for every $j = 1, \ldots, \delta$.

Proof. First assume there is such an automorphism. Set $C_1 := \mathbb{P}^1 =: C_2$ and $\pi_1 := \psi$ and let π_2 be the identity on \mathbb{P}^1 . Then, for $i = 1, 2, \pi_i : C_i \to \mathbb{P}^1$ are maps of degree 1 that satisfy the conditions of Proposition 4, which implies that C is 2-gonal. Since C is not rational, we have gon(C) = 2.

Now assume that C is hyperelliptic and let $\pi: C' \to B$ be an admissible cover of degree 2 where C' is stably equivalent to C. For i = 1, 2 let k_i be the degree of $\pi_i := \pi|_{C_i}$, then $k_i \leq \deg(\pi) = 2$.

Let $B_i := \pi_i(C_i)$ for i = 1, 2. If $B_1 \neq B_2$, by Lemma 3, we have $3 \leq g+1 = \delta \leq \deg(\pi) = 2$, a contradiction. Then $B_1 = B_2$ and, in the notation of Lemma 3, we have $2 = \deg(\pi) \geq k_1 + k_2 + \ell$. Hence $k_1 = k_2 = 1$, $\ell = 0$ and $\psi := \pi_2^{-1} \circ \pi_1$ is the required automorphism of \mathbb{P}^1 .

In particular, since there is always an automorphism of \mathbb{P}^1 taking any set of three distinct points to another one, we see that the previous result recovers, in the case of binary curves, the already know fact that all stable curves of genus 2 are hyperelliptic.

The first part of the following result was already proved in [9, Lemma 5.3], with the same argument. We include a proof here, for the sake of completeness.

Theorem 8. Let C be a binary curve of genus $g(C) \ge 2$. Then

$$\operatorname{gon}(C) \le \left\lfloor \frac{g(C) + 3}{2} \right\rfloor$$

and equality holds if C is general in B_g .

Proof. Set $C = C(\underline{n}^{(1)}, \underline{n}^{(2)})$ and set g = g(C). By Lemma 5 (i) with $\ell = g - 2$, there exists a morphism $\varphi \colon \mathbb{P}^1 \to \mathbb{P}^1$ of degree $k = \lceil \frac{g}{2} \rceil$ such that $\varphi(n_j^{(1)}) = n_j^{(2)}$ for every $j = 1, \ldots, g - 2$.

Now note that $C(\underline{n}^{(1)}, \underline{n}^{(2)})$ is isomorphic to the binary curve $C(\underline{n}^{(1)}, \underline{m}^{(2)})$ where $m_j^{(2)} = n_j^{(2)}$ for $j = 1, \ldots, g-2$ and $m_j^{(2)} = \varphi(n_j^{(1)})$ for j = g-1, g, g+1. Hence we may assume that $\varphi(n_j^{(1)}) = n_j^{(2)}$ for every $j = 1, \ldots, g+1$. Therefore, by Proposition 4 applied to φ and the identity of \mathbb{P}^1 , C is (k+1)-gonal and

$$\operatorname{gon}(C) \le k+1 = \left\lceil \frac{g+2}{2} \right\rceil = \left\lfloor \frac{g+3}{2} \right\rfloor.$$

Now we prove the second statement. As in the proof of Theorem 6, for each $k, \gamma \in \mathbb{N}$, we denote by $P(k, \gamma)$ the set of the pairs $(\underline{n}^{(1)}, \underline{n}^{(2)})$ in $P_{g+1} \times P_{g+1}$ (see Subsection 2.1) such that there exists $I \subset \{1, \ldots, g+1\}$ of cardinality γ and, for i = 1, 2, a map $\psi_i \colon \mathbb{P}^1 \to \mathbb{P}^1$ satisfying $\psi_1(n_j^{(1)}) = \psi_2(n_j^{(2)})$ for all $j \in I$ and such that $\deg(\psi_1) + \deg(\psi_2)$ is at most k. Then $P(k, \gamma) \subset P(k+1, \gamma)$, $P(k, \gamma)$ is closed in $P_{g+1} \times P_{g+1}$ and, by Lemma 5 (i), we have $P(\lfloor \frac{\gamma+2}{2} \rfloor, \gamma) = P$ for every $\gamma \geq 4$.

Set

$$U := P \setminus \bigcup_{\gamma=1}^{g+1} P\left(\left\lfloor \frac{\gamma+2}{2} \right\rfloor - 1, \gamma\right)$$

and note that U is open and dense in $P_{g+1} \times P_{g+1}$. If $\Phi: P_{g+1} \times P_{g+1} \to M_{0,g+1} \times M_{0,g+1} \to B_g$ is the composition of the maps (1) and (3), then the image $\Phi(U)$ contains an open dense subscheme of B_g . Let C be a binary curve of genus g corresponding to a point $(\underline{n}^{(1)}, \underline{n}^{(2)}) \in U$. We will show that $\operatorname{gon}(C) = \lfloor \frac{g+3}{2} \rfloor$.

Let $\pi: C' \to B$ be an admissible cover of degree $k := \operatorname{gon}(C)$, where $\overline{C'}$ is stably equivalent to C. By the first part of the result, we have $k \leq \lfloor \frac{g+3}{2} \rfloor < g+1$. Restricting π to the components $C_i = \mathbb{P}^1$ of C, we obtain maps $\psi_i := \pi|_{C_i} \colon \mathbb{P}^1 \to \mathbb{P}^1$ of degree $k_i := \operatorname{deg}(\psi_i)$. Let J be the set of $j \in \{1, \ldots, g+1\}$ such that $\psi_1(n_j^{(1)}) \neq \psi_2(n_j^{(2)})$ and let $\ell = |J|$. By Lemma 3, we have $\pi(C_1) = \pi(C_2)$ and thus $k \geq k_1 + k_2 + \ell$.

On the other hand, let γ be the cardinality of $I := \{1, \ldots, g+1\} \setminus J$, then $\gamma = g+1-\ell$. Since $(\underline{n}^{(1)}, \underline{n}^{(2)}) \in U$, from the existence of ψ_1 and ψ_2 such that $\psi_1(n_j^{(1)}) = \psi_2(n_j^{(2)})$ for every $j \in I$, we get that $k_1 + k_2 \geq \lfloor \frac{\gamma+2}{2} \rfloor = \lfloor \frac{g+3-\ell}{2} \rfloor$. Since by Proposition ?? we have $k \leq \lfloor \frac{g+3}{2} \rfloor$, then

$$\left\lfloor \frac{g+3+\ell}{2} \right\rfloor = \left\lfloor \frac{g+3-\ell}{2} \right\rfloor + \ell \le k_1 + k_2 + \ell \le k \le \left\lfloor \frac{g+3}{2} \right\rfloor$$

which implies that $\ell = 0$. Thus $\gamma = g + 1$ and $k = \lfloor \frac{g+3}{2} \rfloor$.

References

- D. Abramovich and A. Vistoli, Compactifying the space of stable maps, J. Amer. Math. Soc. 15 (2002), p. 27–75.
- [2] L. Caporaso, Brill-Noether theory of binary curves, Math. Res. Letters 17 n. 2 (2010), p. 243–262.
- [3] L. Caporaso, Gonality of algebraic curves and graphs, Algebraic and Complex Geometry: In honour of Klaus Hulek's 60th Birthday, Springer Proceedings in Mathematics and Statistics 71 (2014), p. 77–109.
- [4] J. Coelho and F. Sercio, On the gonality of stable curves, Mathematische Nachrichten 292 n. 11 (2019), p. 2338–2351.
- [5] J. Coelho and F. Sercio, *Characterizing gonality for two-component stable curves*, Geom Dedicata 214 n. 1 (2021), p. 157–176.
- [6] E. Colombo and P. Frediani, Prym map and second gaussian map for Prymcanonical line bundles, Advances in Mathematics 239 (2013), p. 47–71.
- [7] D. Eisebud and J Harris, *Limit linear series: Basic theory* Invent. math. 85 (1986), p. 337–371.
- [8] E. Esteves and N. Medeiros, *Limit canonical systems on curves with two components*, Inventiones Mathematicae 149 n. 2 (2002), p. 267–338.
- M.Franciosi, Scrolls containing binary curves, Rend. Circ. Mat. Palermo II. 69 (2020), p. 325–338.
- [10] P. Griffiths and J. Harris, On the variety of special linear systems on a general algebraic curve, Duke Math. J 47 (1980) n. 1, p. 233–272.
- [11] J. Harris and I. Morrison, *Moduli of curves*. Springer, New York, 1998.
- [12] J. Harris and D. Mumford, On the Kodaira dimension of the moduli space of curves, Invent. Math. 67 (1982), p. 23–86.
- [13] X. He, Brill-Noether Generality of Binary Curves, Canadian Math. Bul. 64 n. 4 (2021), p. 787–807.
- [14] D. Lara, J. Menezes Souza and R. Vidal Martins, On gonality, scrolls, and canonical models of non-Gorenstein curves, Geometriae Dedicata 198 (2019), p. 1–23.
- [15] B. Osserman, Limit linear series for curves not of compact type, Journal für die reine und angewandte Mathematik (Crelles Journal) 753 (2019), p. 57–88.
- [16] R. Rosa, K.-O. Stöhr, *Trigonal Gorenstein curves*, Journal of Pure and Applied Algebra 174 (2002), p. 187–205.

JULIANA COELHO, UNIVERSIDADE FEDERAL FLUMINENSE (UFF), INSTITUTO DE MATEMÁTICA E ESTATÍSTICA - RUA PROF. MARCOS WALDEMAR DE FREITAS REIS, S/N, GRAGOATÁ, 24210-201 - NITERÓI - RJ, BRASIL *E-mail address:* julianacoelhochaves@id.uff.br