

LIE ALGEBRAS ASSOCIATED WITH LABELED DIRECTED GRAPHS

MAURICIO GODOY MOLINA AND DIEGO LAGOS

ABSTRACT. We present a construction of 2-step nilpotent Lie algebras using labeled directed simple graphs, which allows us to give a criterion to detect certain ideals and subalgebras by finding special subgraphs. We prove that if a label occurs only once, then reversing the orientation of that edge leads to an isomorphic Lie algebra. As a consequence, if every edge is labeled differently, the Lie algebra depends only on the underlying undirected graph. In addition, we construct the labeled directed graphs of all 2-step nilpotent Lie algebras of dimension ≤ 6 and we compute the algebra of strata preserving derivations of the Lie algebra associated with the complete bipartite graph $K_{m,n}$ with two different labelings.

1. INTRODUCTION

Nilpotent Lie algebras have been an important object of study in many areas of mathematics, such as geometric analysis, representation theory and control theory for many years, for example see [1, 4, 5, 16, 19, 22]. In particular, nilpotent Lie algebras of step 2 have attracted much interest for some time, not only because they are the simplest ones in most senses, but due to several close connections with Clifford algebras, dynamics and differential geometry [12, 13, 14, 15, 17, 18].

In 2005, S. G. Dani and M. G. Mainkar [10] proposed a method for constructing 2-step nilpotent Lie algebras using graphs in a very natural manner. This construction is sufficiently rigid so that one can prove very interesting results relating the combinatorial structure of the graph and the algebraic structure of the Lie algebra, for example see [2] for a deep connection with the notion of degeneration. Moreover, this point of view has been generalized by A. Ray in [26] in a very original manner to certain 3-step nilpotent Lie algebras.

In the present paper, we develop further the construction of the 2-step nilpotent Lie algebra structures associated to labeled directed graphs introduced in [6]. This construction is a generalization from the one presented in [10]. From the definition of the associated Lie bracket, the structure constants are 0, -1 and 1 in an appropriate basis. Examples of these Lie algebras are very important in geometric analysis, see [9, 15], since by a well-known result by Mal'cev if a real or complex Lie algebra \mathfrak{g} admits a basis with respect to which the structure constants are rational, then the unique connected and simply connected Lie group \mathfrak{G} that has \mathfrak{g} as its Lie algebra has a lattice, that is, a discrete co-compact subgroup, see [8, 21]. These objects provide natural settings for the study of compact nilmanifolds, such as in [3].

The structure of the paper is the following. We present the construction of the 2-step nilpotent Lie algebra associated to a labeled directed graph in Section 2. Afterward, in Section 3, we prove a combinatorial relation between certain special subgraphs and subalgebras and ideals of the corresponding Lie algebra. In Section 4, we summarize

2010 *Mathematics Subject Classification.* 17B30; 17B40; 05C20; 05C25.

Key words and phrases. 2-step nilpotent Lie algebras, labeled directed graph, ideals, degree preserving derivations, complete bipartite graphs.

This research is partially supported by Fondecyt #1181084.

explicitly some of the ideals and subalgebras presented in the previous section in all 2-step nilpotent Lie algebras of dimensions 4, 5 and 6, according to the classification in [20]. We conclude the paper with Section 5, in which we use the combinatorial description of certain Lie algebras associated to complete bipartite graphs to study the structure of the algebra of strata preserving derivations in these examples.

2. CONSTRUCTING A LIE ALGEBRA FROM A LABELED DIRECTED GRAPH

A finite *directed graph* (V, E) consists of a finite set $V = \{x_1, \dots, x_n\}$ of vertices and a finite set $E = \{\overrightarrow{x_i x_j} \mid x_i, x_j \in V\}$ of directed edges. We say that (V, E) is *simple* if the underlying undirected graph is simple, that is, it has neither loops nor multiple edges.

The directed graphs we will be interested in this paper are edge-labeled, that is, given a finite simple directed graph (V, E) we will consider a *labeling function*

$$c: E \rightarrow \mathcal{C}$$

where $\mathcal{C} = \{c_1, \dots, c_m\}$ is the set of labels. We denote the label of edge \overrightarrow{xy} by $c(x, y)$.

Definition 2.1. *Given a finite simple directed graph (V, E) with a surjective labeling function $c: E \rightarrow \mathcal{C}$, we will denote it by $G = (V, E, c)$ and call it a labeled directed graph.*

Given a labeled directed graph $G = (V, E, c)$, we can associate to it a finite dimensional 2-step nilpotent Lie algebra $\text{Lie}(G)$ over a field F of characteristic $\text{char}(F) \neq 2$ by means of the following procedure:

- $\text{Lie}(G) = \text{span}_F\{x_1, \dots, x_n, c_1, \dots, c_m\}$ as a vector space.
- Define the Lie bracket on the basis $\{x_1, \dots, x_n, c_1, \dots, c_m\}$ as follows

$$[x_i, x_j]_{\text{Lie}(G)} = \begin{cases} c_l, & \text{if } c(x_i, x_j) = c_l, \\ -c_l, & \text{if } c(x_j, x_i) = c_l, \\ 0, & \text{otherwise,} \end{cases} \quad [x_i, c_l]_{\text{Lie}(G)} = [c_l, c_\ell]_{\text{Lie}(G)} = 0,$$

and then extend it linearly. For the rest of the paper, the indices i, j correspond to vertices and l, ℓ correspond to labels.

Taking into account the fact that G is simple, this definition includes the identity $[x, x] = 0$ needed for Lie algebras over fields of characteristic 2. In all other characteristics, this is equivalent to the skew-symmetry of the Lie bracket. Note that the Jacobi identity is satisfied trivially. We consider only fields of characteristic different from 2 due to technical reasons in some of the proofs below.

According to the construction described before, these algebras are naturally *stratified*, that is, they admit a grading

$$\text{Lie}(G) = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1}, \quad \mathfrak{g}_{-2} = \text{span}_F\{c_1, \dots, c_m\}, \quad \mathfrak{g}_{-1} = \text{span}_F\{x_1, \dots, x_n\},$$

for which $[\mathfrak{g}_{-1}, \mathfrak{g}_{-1}] = \mathfrak{g}_{-2}$. In the cases $F = \mathbb{R}$ and $F = \mathbb{C}$, Lie algebras satisfying this condition are particularly important, either for geometric, algebraic or analytic reasons, for example, see [4, 24, 29].

From the definition of $\text{Lie}(G)$, with respect to the basis $V \cup \mathcal{C}$ the structure constants are 0, -1 and 1. As mentioned in the introduction, in the real and complex cases, these Lie algebras admit a lattice. Moreover, given such an algebra with a basis as before, then one can easily construct an appropriate labeled directed graph.

Remark 2.2. *The real Lie algebra $N(d)$ in [27] with d irrational, is an example of a 2-step nilpotent Lie algebra cannot be obtained from a labeled directed graph because its structure constants are not rational in any basis.*




FIGURE 1. Graph in example 2.3

Example 2.3. Consider the labeled directed graph

According to the construction, this graph corresponds to the 5-dimensional 2-step nilpotent Lie algebra $\mathfrak{g} = \text{span}\{x_1, x_2, x_3, x_4, c_1\}$ with non-trivial Lie brackets

$$[x_1, x_2]_{\mathfrak{g}} = [x_1, x_3]_{\mathfrak{g}} = [x_1, x_4]_{\mathfrak{g}} = [x_2, x_3]_{\mathfrak{g}} = [x_2, x_4]_{\mathfrak{g}} = [x_3, x_4]_{\mathfrak{g}} = c_1.$$

It is important to observe that, in fact, this Lie algebra is isomorphic to the Lie algebra $\mathfrak{g}_{5,1}$ in [20], which corresponds to the graph




FIGURE 2. Isomorphic algebra with a simpler graph

This graph will appear again in this paper in Table 1, when listing the graphs of all low-dimensional real nilpotent Lie algebras. An explicit graded isomorphism $\varphi : \mathfrak{g}_{5,1} \rightarrow \mathfrak{g}$ is given in the bases by

$$\varphi(y_1) = x_1, \quad \varphi(y_2) = x_1 + x_2 - x_4, \quad \varphi(y_3) = x_1 + x_2 + x_3, \quad \varphi(y_4) = x_2, \quad \varphi(k) = c_1.$$

Remark 2.4. The construction above is different from the ones that can be found in [10, 26]. The main advantage is that our construction allows for a much broader class of Lie algebras. The main drawback is that we lose some uniqueness results, as seen in the previous example.

3. GRAPH-IDEALS OF $\text{Lie}(G)$

In this section we will study how certain aspects of the combinatorics of the graph G are related to the algebraic structure of the Lie algebra $\text{Lie}(G)$. In order to proceed initially, we need to recall a couple of concepts of graph theory.

Definition 3.1. Assume the vertices of an undirected graph (V, E) are numbered as $V = \{x_1, \dots, x_n\}$.

- The adjacency matrix, given by

$$\mathcal{A}(V, E) = (a_{ij})_{1 \leq i, j \leq n}$$

where $a_{ij} = 1$ if x_i and x_j are connected, and $a_{ij} = 0$ otherwise.

- The valency matrix, given by

$$\mathcal{B}(V, E) = (b_{ij})_{1 \leq i, j \leq n}$$

is a diagonal matrix, where the diagonal entry b_{ii} corresponds to the degree of the vertex x_i , that is, the number of vertices connected to x_i .

- The Laplacian matrix $\mathcal{L}(V, E)$ is the difference $\mathcal{A}(V, E) - \mathcal{B}(V, E)$.

Following well-known results in spectral graph theory, the first natural question to ask concerns the connectedness of the graph (see [7]).

Proposition 3.2. *Let $\mathfrak{g} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1}$ be a stratified 2-step nilpotent Lie algebra admitting a stratified basis $\{c_1, \dots, c_m\} \cup \{x_1, \dots, x_n\}$ with structure constants 0, -1 and 1 . The associated labeled directed graph $G = (V, E, c)$ has $\dim \ker \mathcal{L}(V, E)$ connected components.*

Proof. The connected components of a graph are in one-to-one correspondence with the linearly independent eigenvectors of $\mathcal{L}(V, E)$ with eigenvalue zero (see [23]).

The Laplacian matrix $\mathcal{L}(V, E)$ can be easily found from \mathfrak{g} , since the adjacency matrix $\mathcal{A}(V, E)$ of the graph can be directly found from the structure constants of \mathfrak{g} , forgetting the signs. The result follows. \square

We study the relation between certain special subgraphs of G and the subalgebras and ideals of $\text{Lie}(G) = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1}$. Note that any vector subspace of \mathfrak{g}_{-2} is an ideal of $\text{Lie}(G)$, in particular, the algebra $\text{Lie}(G)$ is not simple.

Definition 3.3. *The following subalgebras and ideals of $\text{Lie}(G)$ are considered trivial:*

- The Lie algebra $\text{Lie}(G)$.
- Vector subspaces of \mathfrak{g}_{-2} .
- Abelian factors, that is, spanned by disconnected single vertices in G .

Recall that a subgraph G' of a directed graph G is *induced* if it is formed from a subset of the vertices of G and all of the edges of G connecting pairs of vertices in that subset.

Proposition 3.4. *Let $G = (V, E, c)$ be a labeled directed graph and $G' = (V', E', c)$ a induced subgraph of G . Then*

- (1) $\text{Lie}(G')$ is a subalgebra of $\text{Lie}(G)$.
- (2) If for each pair of adjacent vertices x, y of G' we have $[x, y]_{\text{Lie}(G)}$ or $[y, x]_{\text{Lie}(G)}$ is a label of G' , then $\text{Lie}(G')$ is an ideal of $\text{Lie}(G)$.

Proof. (1) Let $a, b \in V'$ be two connected vertices in G' . If G' is a induced subgraph, then $[a, b]_{\text{Lie}(G)} \neq 0$ and $[a, b]_{\text{Lie}(G)} = \pm c(a, b) \in \text{Lie}(G')$.

- (2) We need to prove that if x is a vertex of G' and y is a vertex of G , then $[x, y]_{\text{Lie}(G)} \in \text{Lie}(G')$.

- If x, y are not adjacent, then $[x, y]_{\text{Lie}(G)} = 0 \in \text{Lie}(G')$.
- If x, y are adjacent, then $[x, y]_{\text{Lie}(G)} = \pm c(x, y) \in \text{Lie}(G')$.

Therefore, $\text{Lie}(G')$ is an ideal of $\text{Lie}(G)$. \square

For the rest of the paper we will refer to the ideals in point (2) above as graph-ideals.

Corollary 3.5. *Let $G = (V, E, c)$ be a labeled graph and H a connected component of G . Then $\text{Lie}(H)$ is a graph-ideal of $\text{Lie}(G)$.*

The following technical lemma gives a condition under which a change of the direction of a single directed edge of G leads to isomorphic Lie algebras. Recall that the *neighborhood* $N(v)$ of a vertex v of a directed graph G is the set of all vertices of G connected to or from v .

Lemma 3.6. *Let $G = (V, E, c)$ be a labeled directed graph and G' the graph obtained changing the direction of an edge $\overrightarrow{ab} \in E$. Assume the edge \overrightarrow{ab} is the only edge with label $c(a, b) \in \mathcal{C}$. Then $\text{Lie}(G)$ is isomorphic to $\text{Lie}(G')$.*

Proof. Define $C : V \times V \rightarrow \text{span } \mathcal{C}$ by

$$C(x, y) = \begin{cases} c(x, y) & \text{if } \overrightarrow{xy} \in E \\ -c(x, y) & \text{if } \overrightarrow{yx} \in E \\ 0 & \text{if } x \notin N(y) \end{cases}$$

Consider the function $\varphi : \text{Lie}(G) \rightarrow \text{Lie}(G')$ by

$$\varphi(x) = \begin{cases} x & \text{if } x \in V \text{ and } x = a \\ -x & \text{if } x \in V \text{ and } x \neq a \end{cases}$$

and

$$\varphi(k) = \begin{cases} -k & \text{if } x \in \mathcal{C} \text{ and } k = \pm C(a, x), \quad x \neq b \text{ and } x \in N(a) \\ k & \text{otherwise} \end{cases}$$

We need to prove that $\varphi[x, y]_{\text{Lie}(G)} = [\varphi(x), \varphi(y)]_{\text{Lie}(G')}$.

Case 1: $x = a$ and $y = b$.

Let $\alpha = c(a, b)$. Then $\varphi[a, b]_{\text{Lie}(G)} = \varphi(\alpha) = \alpha$ and

$$\varphi[a, b]_{\text{Lie}(G)} = [-a, b]_{\text{Lie}(G')} = -[a, b]_{\text{Lie}(G')} = [b, a]_{\text{Lie}(G')} = \alpha.$$

Case 2: $x = a$ and $y \in N(a) \setminus \{b\}$.

In this case

$$\varphi[a, y]_{\text{Lie}(G)} = \varphi(C(a, y)) = -C(a, y)$$

and

$$[\varphi(a), \varphi(y)]_{\text{Lie}(G')} = [-a, y]_{\text{Lie}(G')} = -[a, y]_{\text{Lie}(G')} = -C(a, y).$$

Case 3: $x \notin N(a)$ and $x \notin N(a)$

In this case $[x, y]_{\text{Lie}(G)} = 0$ and $[\varphi(x), \varphi(y)]_{\text{Lie}(G')} = 0$.

We conclude that $\varphi : \text{Lie}(G) \rightarrow \text{Lie}(G')$ is an isomorphism. \square

Remark 3.7. Notice that if there is another edge in G with the same label as the edge $\overrightarrow{ab} \in E$, then the previous proof does not work. As can be easily seen from Example 2.3, this is by no means a sufficient condition.

Corollary 3.8. Let $G = (V, E, c)$ be a labeled directed graph such that each $u \in \mathcal{C}$ appears at most once in each connected component of G . Let $G' = (V, E', c)$ be the graph obtained inverting the direction of all the edges with the same label simultaneously. Then $\text{Lie}(G) \cong \text{Lie}(G')$.

Proof. Define φ as before in each connected component of G . \square

The following is the main theorem of this paper. We use inductively the technical Lemma 3.6 to find a general criterion to perform direction changes and obtain isomorphic Lie algebras.

Theorem 3.9. Let $G = (V, E, c)$ be a labeled directed and connected graph such that $|\mathcal{C}| = |E|$, that is, each edge is uniquely labeled. Let $G' = (V, E', c)$ be the labeled directed graph obtained by changing the direction of any subset of edges in G . Then $\text{Lie}(G) \cong \text{Lie}(G')$.

Proof. Enumerate the edges that change direction by $\overrightarrow{a_1 b_1}, \overrightarrow{a_2 b_2}, \dots, \overrightarrow{a_q b_q}$.

For $p = 0, 1, \dots, q$ define $G_p = (V, E_p, c)$ inductively by $G_0 = G$ and G_p as the graph obtained by changing the direction of $\overrightarrow{a_p b_p}$ in G_{p-1} . Using the Proposition 3.6 we have $\text{Lie}(G_p) \cong \text{Lie}(G_{p-1})$ and by construction $G_q = G'$.

We conclude that $\text{Lie}(G) = \text{Lie}(G_0) \cong \text{Lie}(G_1) \cong \dots \cong \text{Lie}(G_q) = \text{Lie}(G')$. \square

This theorem connects our construction to the one in [10]. Since in that reference all labels are considered different, then the Lie algebras are defined regardless of the direction of the edges of the graph.

Example 3.10. *The Lie algebras associated to the complete graphs K_p , with $\binom{p}{2}$ different labels correspond to the free 2-step nilpotent Lie algebras generated by p elements. See [25].*

4. GRAPHS FOR MAGNIN'S CLASSIFICATION AND THEIR GRAPH-IDEALS





In [20] it is possible to find a careful description of all real nilpotent Lie algebras of dimensions ≤ 6 and a classification of certain nilpotent algebras of dimension 7. The classification of real nilpotent algebras is still undergoing active research.

Applying directly the results from the previous section, we can present complete tables for the graphs of the 2-step nilpotent real Lie algebras shown in [20] in dimension. For the sake of clarity, we divide this classification in dimensions 4 and 5, and dimension 6.

For brevity, we will not include as separate cases the inclusion of Abelian factors of \mathfrak{g}_{-1} , that is, those Abelian factors coming from disconnected vertices.








For the sake of notation, we denote by \mathfrak{h} the 3-dimensional Heisenberg Lie algebra and by \mathfrak{g}_1 the 1-dimensional Abelian Lie algebra. The Lie algebras $\mathfrak{g}_{5,1}$ and $\mathfrak{g}_{5,2}$ in Table 1 appear named as such in [20].

TABLE 1. Graphs, non-trivial subalgebras and non-trivial graph-ideals in dimensions 4 and 5

| Dimension | Lie algebra | Graph | Subalgebras | Graph-ideals |
|-----------|--|---|--|--|
| 4 | $\mathfrak{h} \times \mathfrak{g}_1$ |  | $\langle x_1, x_2, x_3 \rangle$ | $\langle x_1, x_2, x_3 \rangle$ |
| 5 | $\mathfrak{g}_{5,1}$ |  | $\langle x_1, x_2, x_5 \rangle$ $\langle x_3, x_4, x_5 \rangle$ | $\langle x_1, x_2, x_5 \rangle$ $\langle x_3, x_4, x_5 \rangle$ |
| | $\mathfrak{g}_{5,2}$ |  | $\langle x_1, x_3, x_5 \rangle$ $\langle x_1, x_2, x_4 \rangle$ | — |
| | $\mathfrak{h} \times \mathfrak{g}_1^2$ |  | $\langle x_1, x_2, x_3 \rangle$ | $\langle x_1, x_2, x_3 \rangle$ |

The Lie algebras $\mathfrak{g}_{6,1}$, $\mathfrak{g}_{6,2}$ and $\mathfrak{g}_{6,3}$ in Table 2 only appear listed in [20], and we named them as such to make both tables coherent. The Lie algebra $\mathfrak{g}_{6,3}$ does not exactly fit our theory, since it corresponds to a family of Lie algebras depending on a parameter $\gamma \neq 0$ and not a square. As such, we do not mention either subalgebras nor graph-ideals, but we add it to the table for completeness.

TABLE 2. Graphs, non-trivial subalgebras and non-trivial graph-ideals in dimension 6

| Lie algebra | Graph | Subalgebras | Graph-ideals |
|---|---|---|---|
| $\mathfrak{g}_{6,1}$ |  | $\langle x_1, x_3, x_4 \rangle$ $\langle x_1, x_2, x_6 \rangle$ $\langle x_2, x_3, x_5 \rangle$ | — |
| $\mathfrak{g}_{6,2}$ |  | $\langle x_1, x_2, x_6 \rangle$ $\langle x_1, x_2, x_5 \rangle$ $\langle x_2, x_4, x_6 \rangle$ $\langle x_1, x_2, x_3, x_5, x_6 \rangle$ $\langle x_1, x_2, x_4, x_5, x_6 \rangle$ | $\langle x_1, x_2, x_3, x_5, x_6 \rangle$ $\langle x_1, x_2, x_4, x_5, x_6 \rangle$ |
| $\mathfrak{g}_{6,3}$ $\gamma \neq 0, \alpha^2$ |  | — | — |
| $\mathfrak{h} \times \mathfrak{h}$ |  | $\langle x_1, x_2, x_3 \rangle$ $\langle x_4, x_5, x_6 \rangle$ | $\langle x_1, x_2, x_3 \rangle$ $\langle x_4, x_5, x_6 \rangle$ |
| $\mathfrak{h} \times \mathfrak{g}_1^3$ |  | $\langle x_1, x_2, x_3 \rangle$ | $\langle x_1, x_2, x_3 \rangle$ |
| $\mathfrak{g}_{5,1} \times \mathfrak{g}_1$ |  | $\langle x_1, x_2, x_5 \rangle$ $\langle x_3, x_4, x_5 \rangle$ $\langle x_1, x_2, x_3, x_4, x_5 \rangle$ | $\langle x_1, x_2, x_5 \rangle$ $\langle x_3, x_4, x_5 \rangle$ $\langle x_1, x_2, x_3, x_4, x_5 \rangle$ |
| $\mathfrak{g}_{5,2} \times \mathfrak{g}_1$ |  | $\langle x_1, x_3, x_5 \rangle$ $\langle x_1, x_2, x_4 \rangle$ $\langle x_1, x_2, x_3, x_4, x_5 \rangle$ | $\langle x_1, x_2, x_3, x_4, x_5 \rangle$ |

5. THE ALGEBRA OF GRADED DERIVATIONS FOR $K_{m,n}$

When studying Lie algebras in different contexts, it is common to ask oneself firstly about isomorphisms, and immediately after about derivations. If the Lie algebra \mathfrak{g} being studied is graded, then the natural question is to try to compute the space of graded derivations $\text{Der}_0(\mathfrak{g})$. In this section, we present some computations associated to the complete bipartite graph $K_{m,n}$. As it will be mentioned at the end of this section, this space is related to infinitesimal symmetries in the case of real and complex Lie algebras, according to the theory of Tanaka prolongation, see [28].

Recall that the complete bipartite graph $K_{m,n} = (V, E)$ is the undirected graph in which all vertices from the set $X = \{x_1, \dots, x_m\}$ are connected to all vertices from the set $Y = \{y_1, \dots, y_n\}$. As such, the set V is the disjoint union of X and Y , and E contains mn edges.

In what follows, we will direct all edges going from X to Y . In subsection 5.2, this orientation is irrelevant, according to Theorem 3.9.

5.1. Labeling $K_{m,n}$ with a single label. Let G_1 be the complete bipartite graph with edges

$$E = \{\overrightarrow{xy} \mid x \in X, y \in Y\}$$

and labeled by a unique label u . As before, recall that $\text{Lie}(G_1)$ is naturally stratified as $\mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1}$, where $\mathfrak{g}_{-1} = \text{span}_F(X \cup Y)$ and $\mathfrak{g}_{-2} = \text{span}_F\{u\}$.

In order to study the space of graded derivations of the Lie algebra associated to G_1 , it is convenient to think of an element $\varphi \in \text{Der}_0(\text{Lie}(G_1))$ as a matrix in the obvious basis, namely

$$\varphi = \left(\begin{array}{c|c|c} \alpha & \gamma & 0 \\ \beta & \delta & \\ \hline 0 & & \lambda \end{array} \right)$$

where we have the linear maps

$$\begin{aligned} \alpha &: \text{span}_F(X) \rightarrow \text{span}_F(X), & \beta &: \text{span}_F(X) \rightarrow \text{span}_F(Y), \\ \gamma &: \text{span}_F(Y) \rightarrow \text{span}_F(X), & \delta &: \text{span}_F(Y) \rightarrow \text{span}_F(Y), \end{aligned}$$

and $\lambda \in F$ is a scalar.

Proposition 5.1. *The dimension of $\text{Der}_0(\text{Lie}(G_1))$ is*

$$\frac{(m+n)(m+n+1)}{2} + 1.$$

Proof. Let $\varphi \in \text{Der}_0(\text{Lie}(G_1))$. Then there are constants $\lambda, \alpha_{ri}, \beta_{si}, \gamma_{rj}, \delta_{sj} \in F$ (where $r, i = 1, \dots, m$ and $s, j = 1, \dots, n$), such that

$$\varphi(x_i) = \sum_{r=1}^m \alpha_{ri} x_r + \sum_{s=1}^n \beta_{si} y_s, \quad \varphi(y_j) = \sum_{r=1}^m \gamma_{rj} x_r + \sum_{s=1}^n \delta_{sj} y_s, \quad \varphi(u) = \lambda u.$$

Thus, for $i, i' = 1, \dots, m$ and $j, j' = 1, \dots, n$, we have

$$\begin{aligned} \varphi[x_i, x_{i'}]_{\text{Lie}(G_1)} &= [\varphi(x_i), x_{i'}]_{\text{Lie}(G_1)} + [x_i, \varphi(x_{i'})]_{\text{Lie}(G_1)} = \left(-\sum_{s=1}^n \beta_{si} + \sum_{s=1}^n \beta_{si'} \right) u = 0, \\ \varphi[y_j, y_{j'}]_{\text{Lie}(G_1)} &= [\varphi(y_j), y_{j'}]_{\text{Lie}(G_1)} + [y_j, \varphi(y_{j'})]_{\text{Lie}(G_1)} = \left(\sum_{r=1}^m \gamma_{rj} - \sum_{r=1}^m \gamma_{rj'} \right) u = 0, \\ \varphi[x_i, y_j]_{\text{Lie}(G_1)} &= [\varphi(x_i), y_j]_{\text{Lie}(G_1)} + [x_i, \varphi(y_j)]_{\text{Lie}(G_1)} = \left(\sum_{r=1}^m \alpha_{ri} + \sum_{s=1}^n \delta_{sj} \right) u = \lambda u. \end{aligned}$$

As a consequence, we have the following homogeneous system

$$-\sum_{s=1}^n \beta_{si} + \sum_{s=1}^n \beta_{si'} = 0, \quad \sum_{r=1}^m \gamma_{rj} - \sum_{r=1}^m \gamma_{rj'} = 0, \quad \sum_{r=1}^m \alpha_{ri} + \sum_{s=1}^n \delta_{sj} - \lambda = 0$$

of $\frac{n(n-1)}{2} + \frac{m(m-1)}{2} + mn$ independent linear equations. Considering the following simple computation

$$(m+n)^2 + 1 - \frac{m(m-1)}{2} - \frac{n(n-1)}{2} - mn = \frac{(m+n)(m+n+1)}{2} + 1.$$

we conclude with the expected result. \square

5.2. Labeling $K_{m,n}$ with mn different labels. Let G_2 be the complete bipartite graph with edges

$$E = \{\vec{xy} \mid x \in X, y \in Y\}$$

and labeled by the set $\mathcal{C} = \{c_{ij} \mid 1 \leq i \leq m, 1 \leq j \leq n\}$. We will consider that all labels are different, that is, we assume \mathcal{C} has mn elements. Once again, we stratify $\text{Lie}(G_2)$ as $\mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1}$, where $\mathfrak{g}_{-1} = \text{span}_F(X \cup Y)$ and $\mathfrak{g}_{-2} = \text{span}_F(\mathcal{C})$.

As in the previous subsection, we write an element $\varphi \in \text{Der}_0(\text{Lie}(G_2))$ as a matrix in the obvious basis, namely

$$\varphi = \left(\begin{array}{c|c|c} \alpha & \gamma & 0 \\ \beta & \delta & 0 \\ \hline 0 & & \epsilon \end{array} \right)$$

where we have the linear maps

$$\begin{aligned} \alpha &: \text{span}_F(X) \rightarrow \text{span}_F(X), & \beta &: \text{span}_F(X) \rightarrow \text{span}_F(Y), \\ \gamma &: \text{span}_F(Y) \rightarrow \text{span}_F(X), & \delta &: \text{span}_F(Y) \rightarrow \text{span}_F(Y), \\ \epsilon &: \text{span}_F(\mathcal{C}) \rightarrow \text{span}_F(\mathcal{C}). \end{aligned}$$

Proposition 5.2. *The dimension of $\text{Der}_0(\text{Lie}(G_2))$ is $m^2 + n^2 + m^2n^2 - mn$.*

Proof. Let $\varphi \in \text{Der}_0(\text{Lie}(G_2))$. Then there are constants $\alpha_{ri}, \beta_{si}, \gamma_{rj}, \delta_{sj}, \epsilon_{ij}^{rs} \in F$ (where $r, i = 1, \dots, m$ and $s, j = 1, \dots, n$), such that

$$\varphi(x_i) = \sum_{r=1}^m \alpha_{ri} x_r + \sum_{s=1}^n \beta_{si} y_s, \quad \varphi(y_j) = \sum_{r=1}^m \gamma_{rj} x_r + \sum_{s=1}^n \delta_{sj} y_s, \quad \varphi(c_{ij}) = \sum_{r=1}^m \sum_{s=1}^n \epsilon_{ij}^{rs} c_{rs}.$$

Thus, for $i, i' = 1, \dots, m$ and $j, j' = 1, \dots, n$, we have

$$\begin{aligned} 0 &= \varphi[x_i, x_{i'}]_{\text{Lie}(G_2)} = [\varphi(x_i), x_{i'}]_{\text{Lie}(G_2)} + [x_i, \varphi(x_{i'})]_{\text{Lie}(G_2)} = - \sum_{s=1}^n \beta_{si} c_{i's} + \sum_{s=1}^n \beta_{si'} c_{is}, \\ 0 &= \varphi[y_j, y_{j'}]_{\text{Lie}(G_2)} = [\varphi(y_j), y_{j'}]_{\text{Lie}(G_2)} + [y_j, \varphi(y_{j'})]_{\text{Lie}(G_2)} = \sum_{r=1}^m \gamma_{rj} c_{rj'} - \sum_{r=1}^m \gamma_{rj'} c_{rj}. \end{aligned}$$

If $i = i'$, then the first equality reduces to

$$- \sum_{s=1}^n \beta_{si} c_{i's} + \sum_{s=1}^n \beta_{si'} c_{is} = - \sum_{s=1}^n \beta_{si} c_{is} + \sum_{s=1}^n \beta_{si} c_{is} = 0,$$

that is, it is satisfied trivially for any $i = 1, \dots, m$.

On the other hand, if $i \neq i'$ then $c_{i's} \neq c_{is}$ and as $\mathcal{C} = \{c_{ij}\}$ is a linearly independent set, then $\beta_{si} = \beta_{si'} = 0$ for all $i, i' = 1, \dots, m$ and $s = 1, \dots, n$. Similarly, we obtain $\gamma_{rj} = \gamma_{rj'} = 0$ for all $j, j' = 1, \dots, n$ and $r = 1, \dots, m$.

Note that

$$\varphi(c_{ij}) = \sum_{r=1}^m \sum_{s=1}^n \epsilon_{ij}^{rs} c_{rs} = [\varphi(x_i), y_j]_{\text{Lie}(G_2)} + [x_i, \varphi(y_j)]_{\text{Lie}(G_2)} = \sum_{r=1}^m \alpha_{ri} c_{rj} + \sum_{s=1}^n \delta_{sj} c_{is}$$

and we have mn linearly independent equations. The result follows. \square

Remark 5.3. For a real or complex stratified Lie algebra $\mathfrak{n} = \mathfrak{n}_{-s} \oplus \cdots \oplus \mathfrak{n}_{-1}$, there is an important object called its Tanaka prolongation

$$\text{Prol}(\mathfrak{n}) = \mathfrak{n}_{-s} \oplus \cdots \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \cdots$$

which helps computing infinitesimal symmetries of certain differential systems, see [28]. Each of the subspaces \mathfrak{n}_r for $r \geq 0$ can be computed explicitly using a simple induction procedure. In particular, it follows directly from the construction that

$$\mathfrak{n}_0 = \text{Der}_0(\mathfrak{n}).$$

This equality has been successfully in [29] to completely characterize the Tanaka prolongations of free Lie algebras.

6. ACKNOWLEDGMENTS

The second author would like to thank professor María Alejandra Álvarez from Universidad de Antofagasta, Chile, for her hospitality and her comments regarding earlier versions of this paper. Both authors would like to thank professor Andrew Clarke from Universidade Federal do Rio de Janeiro, Brazil, for his hospitality.

REFERENCES

- [1] Agrachev, Andrei; Barilari, Davide; Boscain, Ugo. *A comprehensive introduction to sub-Riemannian geometry. From the Hamiltonian viewpoint.* With an appendix by Igor Zelenko. Cambridge Studies in Advanced Mathematics, 181. Cambridge University Press, Cambridge, 2020.
- [2] Alfaro Arancibia, B.; Alvarez, M. A.; Anza, Y. *Degenerations of graph Lie algebras.* Linear Multilinear Algebra 70 (2022), no. 1, 91–100.
- [3] W. Bauer, K. Furutani, C. Iwasaki, A. Laaroussi, *Spectral theory of a class of nilmanifolds attached to Clifford modules.* Mathematische Zeitschrift, volume 297, pages 557–583 (2021)
- [4] A. Bonfiglioli, E. Lanconelli and F. Uguzzoni. *Stratified Lie groups and potential theory for their sub-Laplacians.* Springer Monogr. Math. Springer, Berlin, 2007.
- [5] Čap, A., Slovak, J. *Parabolic geometries. I. Background and general theory.* Mathematical Surveys and Monographs, 154. American Mathematical Society, Providence, RI, 2009.
- [6] D. Chakrabarti, M. Mainkar, S. Swiatlowski, *Automorphism groups of nilpotent Lie algebras associated to certain graphs.* Comm. Algebra 48 (2020), no. 1, 263–273.
- [7] F. Chung. *Spectral Graph Theory.* American Mathematical Society (1997).
- [8] L. J. Corwin, F. P. Greenleaf, *Representations of nilpotent Lie groups and their applications. Part I.* Cambridge Stud. Adv. Math., 18, 1990,
- [9] G. Crandall, J. Dodziuk, *Integral structures on H-type Lie algebras.* J. Lie Theory 12 (2002), no.1, 69–79.
- [10] S. G. Dani and M. G. Mainkar. *Anosov automorphisms on compact nilmanifolds associated with graphs.* Trans. Amer. Math. Soc., vol. 357, no. 6, 2235–2251, 2005.
- [11] R. DeCoste, L. DeMeyer, M. Mainkar and A. Ray. *Abelian factors in 2-step nilpotent Lie algebras constructed from graphs.* Comm. Algebra, vol. 51, no. 5, pp. 2155–2175, 2023.
- [12] P. Eberlein. *Geometry of 2-step nilpotent groups with a left invariant metric.* Ann. Sci. École Norm. Sup. (4), vol. 27, no. 5, pp. 611–660, 1994.
- [13] P. Eberlein. *Geometry of 2-step nilpotent groups with a left invariant metric. II.* Trans. Amer. Math. Soc., vol. 343, no. 2, pp. 805–828, 1994.
- [14] K. Furutani, M. Godoy Molina, I. Markina, T. Morimoto and A. Vasilev. *Lie algebras attached to Clifford modules and simple graded Lie algebras.* J. Lie Theory 28 (2018), no. 3, 843–864.
- [15] K. Furutani and I. Markina. *Complete classification of pseudo H-type Lie algebras: I.* Geom. Dedicata, vol. 190, pp. 23–51, 2017.
- [16] V. Jurdjevic. *Geometric control theory.* Cambridge Studies in Advanced Mathematics, 52. Cambridge University Press, Cambridge, 1997.
- [17] A. Kaplan. *Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms.* Trans. Amer. Math., vol. 258, no. 1, pp. 147–153, 1980.

- [18] A. Kaplan. *Riemannian nilmanifolds attached to Clifford modules*. *Geom. Dedicata*, vol. 11, no. 2, pp. 127–136, 1981.
- [19] A. Kushner, V. Lychagin and V. Rubtsov. *Contact geometry and non-linear differential equations*. *Encyclopedia of Mathematics and its Applications*, 101. Cambridge University Press, Cambridge, 2007.
- [20] L. Magnin. *Sur les algèbres de Lie nilpotentes de dimension ≤ 7* . *Journal of Geometry and Physics*, vol. 3, no. 1, 1986.
- [21] A.I. Malcev, *On a class of homogeneous spaces*, Amer. Math. Soc. Translation, no. 39, 1951.
- [22] R. Montgomery. *A tour of subriemannian geometries, their geodesics and applications*. *Mathematical Surveys and Monographs*, 91. American Mathematical Society, Providence, RI, 2002.
- [23] B. Nica, *A Brief Introduction to Spectral Graph Theory*. EMS Textbooks in Mathematics (2018).
- [24] S. Nicolussi, A. Ottazzi, *Polarised Lie groups contactomorphic to stratified groups*. arXiv:1807.03854.
- [25] C. Reutenauer. *Free Lie algebras*. *Handbook of algebra*, Vol. 3, 887–903, *Handb. Algebr.*, 3, Elsevier/North-Holland, Amsterdam, 2003.
- [26] A. Ray. *Two-step and three-step nilpotent Lie algebras constructed from Schreier graphs*. *J. Pure Appl. Algebra*, vol. 28, no. 2, pp. 479–495, 2016.
- [27] J. Scheuneman. *Two-step nilpotent Lie algebras*. *J. Algebra*, vol. 7, pp. 152–159, 1967.
- [28] N. Tanaka. *On differential systems, graded Lie algebras and pseudogroups*. *J. Math. Kyoto Univ.* 10, (1970), 1–82
- [29] B. Warhurst. *Tanaka prolongation of free Lie algebras*. *Geom. Dedicata* 130 (2007), 59–69.

DEPARTAMENTO DE MATEMÁTICA Y ESTADÍSTICA, UNIVERSIDAD DE LA FRONTERA, AV. FRANCISCO SALAZAR 01145 TEMUCO, CHILE.

Email address: mauricio.godoy@ufrontera.cl

Email address: diego.lagos@ufrontera.cl