On the properties of the linear conjugate gradient method

Zexian Liu $\,\cdot\,$ Qiao Li

Received: date / Accepted: date

Abstract The linear conjugate gradient method is an efficient iterative method for the convex quadratic minimization problems $\min_{x \in \mathbb{R}^n} f(x) = \frac{1}{2}x^T A x + b^T x$, where $A \in \mathbb{R}^{n \times n}$ is symmetric and positive definite and $b \in \mathbb{R}^n$. It is generally agreed that the gradients g_k are not conjugate with respective to A in the linear conjugate gradient method (see page 111 in Numerical optimization (2nd, Springer, 2006) by Nocedal and Wright). In the paper we prove the conjugacy of the gradients g_k generated by the linear conjugate gradient method, namely,

$$g_k^T A g_i = 0, \ i = 0, 1, \cdots, k - 2$$

In addition, a new way is exploited to derive the linear conjugate gradient method based on the conjugacy of the search directions and the orthogonality of the gradients, rather than the conjugacy of the search directions and the exact stepsize.

Keywords Conjugacy · Orthogonality · Conjugate gradient method · Convex quadratic optimization

Mathematics Subject Classification (2000) 90C06 · 65K

1 Introduction

We consider the following convex quadratic optimization problem:

$$\min_{x \in R^n} f(x) = \frac{1}{2} x^T A x + b^T x,$$
(1.1)

where $A \in \mathbb{R}^{n \times n}$ is symmetric and positive definite and $b \in \mathbb{R}^n$.

The linear conjugate gradient method for solving (1.1) has the following form

$$x_{k+1} = x_k + \alpha_k d_k, \tag{1.2}$$

where d_k is the search direction given by

$$d_k = \begin{cases} -g_0, & \text{if } k = 0, \\ -g_k + \beta_k d_{k-1}, & \text{if } k > 0, \end{cases}$$
(1.3)

Zexian Liu, e-mail: liuzexian2008@163.com(☉); Qiao Li, e-mail:liqiaomuzi@163.com School of Mathematics and Statistics, Guizhou University, Guiyang, 550025, China

and α_k is the stepsize determined by the exact line search, namely,

$$\alpha_k = -\frac{g_k^T d_k}{d_k^T A d_k} = \arg\min_{\alpha>0} f\left(x_k + \alpha d_k\right).$$
(1.4)

It follows from (1.4) that

$$g_{k+1}^T d_k = 0. (1.5)$$

Here the parameter β_k is often derived by making it satisfy the following condition:

$$d_{k+1}^T A d_k = 0, (1.6)$$

which together with (1.5) can yield the conjugacy of the whole search directions, as well as the global convergence. Some well-known formulae for β_k are called the Fletcher-Reeves (FR) [5], Hestenes-Stiefel (HS) [6], Polak-Ribière-Polyak (PRP) [8,7] and Dai-Yuan (DY) [4] formulae, and are given by

$$\beta_k^{FR} = \frac{\|g_k\|^2}{\|g_{k-1}\|^2}, \quad \beta_k^{HS} = \frac{g_k^T y_{k-1}}{d_{k-1}^T y_{k-1}}, \quad \beta_k^{PRP} = \frac{g_k^T y_{k-1}}{\|g_{k-1}\|^2}, \quad \beta_k^{DY} = \frac{\|g_k\|^2}{d_{k-1}^T y_{k-1}},$$

where $y_{k-1} = g_k - g_{k-1}$, and $\|\cdot\|$ denotes the Euclidean norm. In the case that f is given by (1.1) and the exact line search (1.4) is performed, all β_k should be the same.

It is well-known that the linear conjugate gradient method enjoys the following nice properties.

Theorem 1.1 [1, 2] Suppose that the iterate $\{x_k\}$ is generated by the linear conjugate gradient method for solving (1.1), and x_k is not the solution point x^* . Then,

$$g_i^T d_i = -\|g_i\|^2, \quad i = 0, 1, 2, \cdots,$$

$$d_i^T A d_j = 0, \quad j = 0, 1, 2, \cdots, i - 1$$

$$g_i^T d_j = 0, \quad j = 0, 1, 2, \cdots, i - 1,$$

$$g_i^T g_j = 0, \quad j = 0, 1, 2, \cdots, i - 1.$$

Further, the sequence $\{x_k\}$ converges to x^* in at most n steps.

It follows from Theorem 1.1 that the gradients g_k are mutually orthogonal in the linear conjugate gradient method. However, it is generally agreed that the gradients g_k generated by the linear conjugate gradient method are not conjugate with respective to A. For example, one can find the well-known description "Since the gradients r_k are mutually orthogonal, the term "conjugate gradient method" is actually a **misnomer**. It is the search directions, not the gradients, that are conjugate with respect to A. " in Page 111 of Nocedal and Wright's monograph [3]. Must the gradients generated by the linear conjugate gradient method not be conjugate with respective to the matrix A? In the paper we will give a negative answer to this question, as well as exploiting a new way for deriving the linear conjugate gradient method based on the conjugacy of the search direction and the orthogonality of the gradient, rather than the conjugacy of the search direction and the exact stepsize.

2 The conjugacy of the gradients

In the section we will establish the conjugacy of the gradients generated by the linear conjugate gradient method.

Theorem 2.1 Suppose that $\{x_k\}$ and $\{g_k\}$ are generated by linear conjugate gradient method for solving (1.1). Then,

$$g_{k+1}^T A g_k = -\frac{\|g_{k+1}\|_2^2}{\alpha_k} ,$$

$$g_{k+1}^T A g_i = 0, \ i = 0, 1, \cdots, k-1.$$
(2.1)

Proof We first prove the first equality in (2.1). If k = 0, then it follows from (1.3) and Theorem 1.1 that

$$g_1^T A g_0 = -g_1^T A d_0 = -\frac{1}{\alpha_0} g_1^T (g_1 - g_0) = -\frac{\|g_1\|^2}{\alpha_0}.$$

When k > 0, we also obtain from the orthogonality of gradients and (1.3) that

$$g_{k+1}^T A g_k = g_{k+1}^T A (-d_k + \beta_k d_{k-1})$$

= $-\frac{g_{k+1}^T (g_{k+1} - g_k)}{\alpha_k} + \frac{\beta_k g_{k+1}^T (g_k - g_{k-1})}{\alpha_{k-1}}$
= $-\frac{\|g_{k+1}\|_2^2}{\alpha_k}.$

We then prove the second equality in (2.1). If k = 1, by (1.3) and Theorem 1.1, we obtain

$$g_2^T A g_0 = -g_2^T A d_0 = -\frac{1}{\alpha_0} g_2^T (g_1 - g_0) = 0.$$
(2.2)

When k > 1, it follows from (1.3) and Theorem 1.1 that

$$g_{k+1}^T A g_i = g_{k+1}^T A (-d_i + \beta_i d_{i-1})$$

= $-\frac{g_{k+1}^T (g_{i+1} - g_i)}{\alpha_i} + \frac{\beta_i g_{k+1}^T (g_i - g_{i-1})}{\alpha_{i-1}}$
= 0,

where $i = 0, 1, \dots, k - 1$. It completes the proof.

Remark 1. It follows from Theorem 2.1 that g_{k+1} is conjugate to g_i $(i = 0, 1, \dots, k-1)$ with respective to A. Due to the conjugacy of the gradients, it seems that it is also reasonable to name the iterative method (1.2) and (1.3) with (1.4) the linear conjugate gradient method.

3 New way for deriving the linear conjugate gradient method

In the section, we will exploit a new way for deriving the linear conjugate gradient method.

The orthogonality of gradients is crucial to the linear conjugate gradient method. Different from the stepsize (1.4) which makes f reduce mostly along the search direction d_k , the new stepsize α_k in (1.3) is choose such that the resulting new gradient g_{k+1} is orthogonal to the latest gradient g_k , namely,

$$g_{k+1}^T g_k = 0, (3.1)$$

which implies that

$$\alpha_k = -\frac{g_k^T g_k}{g_k^T A d_k}.\tag{3.2}$$

Therefore, the new iterative method has the form (1.2) and (1.3) with (3.2), which is derived based on the conditions:

$$g_{k+1}^T g_k = 0 \text{ and } d_{k+1}^T A d_k = 0.$$
 (3.3)

Note that it is different from the conditions (1.5) and (1.6) used to derive the linear conjugate gradient method.

In the following theorem, we will prove the equivalence of the iterative method (1.2) and (1.3) with (3.2) and the linear conjugate gradient method.

Theorem 3.1 The new stepsize (3.2) is also the exact stepsize, namely, $\alpha_k = \arg\min_{\alpha>0} f(x_k + \alpha d_k)$.

Proof It follows from $d_k^T A d_{k-1} = 0$ that $\alpha_k = \frac{g_k^T g_k}{d_k^T A d_k}$. Thus, we have $\begin{aligned} g_{k+1}^T d_k &= (g_k + \alpha_k A d_k)^T d_k \\ &= g_k^T d_k + \frac{g_k^T g_k}{d_k^T A d_k} d_k^T A d_k \\ &= \beta_k g_k^T d_{k-1} \\ &= \beta_k \beta_{k-1} \cdots \beta_1 g_1^T d_0 \\ &= -\beta_k \beta_{k-1} \cdots \beta_1 g_1^T g_0 \\ &= 0, \end{aligned}$

which togetcher with (1.1) yields that

$$\alpha_k = \arg\min_{\alpha>0} f\left(x_k + \alpha d_k\right)$$

It completes the proof.

Remark 2. It follows from Theorem 3.1 that the iterative method (1.2) and (1.3) with (3.2) is indeed the linear conjugate gradient method. Thus we can easily obtain the following corollary.

Remark 3. The above new way for deriving the linear conjugate gradient method might motivate us to design a new criterion that the new trial gradient satisfies, which might make the resulting conjugate gradient method enjoy finite temination property without the exact line search, as well as nice numerical performance.

Corollary 3.1 Suppose that the iterate $\{x_k\}$ is generated by the iterative method (1.2) and (1.3) with (3.2) for solving (1.1), and x_k is not the solution point x^* . Then, the sequence $\{x_k\}$ converges to x^* in at most n steps.

4 Conclusion and Discussion

In the paper we establish the conjugacy of the gradients generated by the linear conjugate gradient method, and give a new way for deriving the linear conjugate gradient method based on the orthogonality of the gradients and the conjugacy of the search dirction at the most recent iterations.

However, there are still many questions under investigation. For example, how to design a new design that the new trial gradient satisfies, which might make the resulting conjugate gradient method enjoy nice theoretical property, as well as good numerical performance.

Acknowledgements This research is supported by National Science Foundation of China (No. 12261019), Guizhou Provincial Science and Technology Projects (No. QHKJC-ZK[2022]YB084).

Data availability. The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Conflict of interest. The authors declare no competing interests.

References

- 1. Dai Y. H., Yuan, Y. X. Nonlinear conjugate gradient method. Shanghai Scientific and Technical Publishers, 1999.
- 2. Yuan, Y. X., Sun, W. Y.: Theory and Methods of Optimization. Science Press of China, 1999.
- 3. Nocedal J., Wright S. J. Numerical optimization (2nd). Springer, 2006.
- Dai Y. H., Yuan Y. X.: A nonlinear conjugate gradient method with a strong global convergence property. SIAM J. Optim. 10(1), 177-182(1999).
- 5. Fletcher R., Reeves C.: Function minimization by conjugate gradients. Comput. J. 7(2), 149-154(1964).
- Hestenes M. R., Stiefel E. L.: Methods of conjugate gradients for solving linear systems. J. Res. Natl. Bur. Stand. 49(6), 409-436(1952).
- 7. Polak, E., Ribière, G.: Note sur la convergence de méthodes de directions conjugées. Rev. Francaise Informat. Recherche Opértionelle, **3**, 35-43(1969).
- 8. Polyak B. T.: The conjugate gradient method in extreme problems. USSR Comput. Math. Math. Phys. 9, 94-112(1969).