

# Dirac operators with torsion, spectral Einstein functionals and the noncommutative residue

Jian Wang<sup>a</sup>, Yong Wang<sup>b,\*</sup>, Tong Wu<sup>b</sup>

<sup>a</sup>*School of Science, Tianjin University of Technology and Education, Tianjin, 300222, P.R.China*

<sup>b</sup>*School of Mathematics and Statistics, Northeast Normal University, Changchun, 130024, P.R.China*

---

## Abstract

Recently Dabrowski etc. [1] obtained the metric and Einstein functionals by two vector fields and Laplace-type operators over vector bundles, giving an interesting example of the spinor connection and square of the Dirac operator. Pfäffle and Stephan [2] considered orthogonal connections with arbitrary torsion on compact Riemannian manifolds and computed the spectral action. Motivated by the spectral functionals and Dirac operators with torsion, we give some new spectral functionals which is the extension of spectral functionals to the noncommutative realm with torsion, and we relate them to the noncommutative residue for manifolds with boundary. Our method of producing these spectral functionals is the noncommutative residue and Dirac operators with torsion.

*Keywords:* Dirac operators with torsion; noncommutative residue; orthogonal connection with torsion, spectral functional.

---

## 1. Introduction

An eminent spectral scheme that generates geometric objects on manifolds such as residue, scalar curvature, and other scalar combinations of curvature tensors is the small-time asymptotic expansion of the (localised) trace of heat kernel[3, 4]. The theory has very rich structures both in physics and mathematics. In the recent paper[1], Dabrowski etc. defined bilinear functionals of vector fields and differential forms, the densities of which yield the metric and Einstein spectral functionals on even-dimensional Riemannian manifolds, and they obtained certain values or residues of the (localised) zeta function of the Laplacian arising from the Mellin transform and the coefficients of this expansion.

Let  $E$  be a finite-dimensional complex vector bundle over a closed compact manifold  $M$  of dimension  $n$ , the noncommutative residue of a pseudo-differential operator  $P \in \Psi DO(E)$  can be defined by

$$res(P) := (2\pi)^{-n} \int_{S^*M} \text{Tr}(\sigma_{-n}^P(x, \xi)) dx d\xi, \quad (1.1)$$

where  $S^*M \subset T^*M$  denotes the co-sphere bundle on  $M$  and  $\sigma_{-n}^P$  is the component of order  $-n$  of the complete symbol

$$\sigma^P := \sum_i \sigma_i^P \quad (1.2)$$

of  $P$ , cf. [5–8], and the linear functional  $res : \Psi DO(E) \rightarrow \mathbb{C}$  is in fact the unique trace (up to multiplication by constants) on the algebra of pseudo-differential operators  $\Psi DO(E)$ . In [9], Connes computed a conformal four-dimensional Polyakov action analogy using the noncommutative residue. Connes proved

---

\*Corresponding author.

*Email addresses:* wangj484@nenu.edu.cn (Jian Wang), wangy581@nenu.edu.cn (Yong Wang)

that the noncommutative residue on a compact manifold  $M$  coincided with Dixmier's trace on pseudodifferential operators of order  $-\dim M$ [10]. More precisely, Connes made a challenging observation that the Wodzicki residue of the inverse square of the Dirac operator yields the Einstein-Hilbert action of general relativity[10][11]. Kastler[12] gave a brute-force proof of this theorem, and Kalau and Walze[13] proved this theorem in the normal coordinates system simultaneously, which is called the Kastler-Kalau-Walze theorem now. Based on the theory of the noncommutative residue introduced by Wodzicki, Fedosov etc.[14] constructed a noncommutative residue on the algebra of classical elements in Boutet de Monvel's calculus on a compact manifold with boundary of dimension  $n > 2$ . With elliptic pseudodifferential operators and noncommutative residue, it is natural way to study the Kastler-Kalau-Walze type theorem and operator-theoretic explanation of the gravitational action for manifolds with boundary. For Dirac operators and signature operators, Wang computed the noncommutative residue and proved Kastler-Kalau-Walze type theorem for manifolds with boundary[15–17].

Earlier Jean-Michel Bismut [18] proved a local index theorem for Dirac operators on a Riemannian manifold  $M$  associated with connections on  $TM$  which have non zero torsion. In [19], Ackermann and Tolksdorf proved a generalized version of the well-known Lichnerowicz formula for the square of the most general Dirac operator with torsion  $D_T$  on an even-dimensional spin manifold associated to a metric connection with torsion. In [20], Pfüfle and Stephan considered compact Riemannian spin manifolds without boundary equipped with orthogonal connections, and investigated the induced Dirac operators. Moreover, Pfüfle and Stephan considered orthogonal connections with arbitrary torsion on compact Riemannian manifolds, and for the induced Dirac operators, twisted Dirac operators and Dirac operators of Chamseddine-Connes type they computed the spectral action[2]. In [21], we computed the lower dimensional volume  $\widetilde{\text{Wres}}[\pi^+(D_T^*)^{-p_1} \circ \pi^+ D_T^{-p_2}]$  and got a Kastler-Kalau-Walze type theorems associated with Dirac operators with torsion on compact manifolds with boundary. The purpose of this paper is to generalize the results in [1],[16], [20] and get spectral functionals associated with Dirac operators with torsion on compact manifolds with boundary. For lower dimensional compact Riemannian manifolds with boundary, we compute the lower dimensional residue of  $\widetilde{\nabla}_X \widetilde{\nabla}_Y D_T^{-4}$  and get the Kastler-Kalau-Walze theorems.

## 2. Spectral functionals for Dirac operator with Torsion

In this section we consider an  $n$ -dimensional oriented Riemannian manifold  $(M, g^M)$  equipped with some spin structure. The Levi-Civita connection  $\nabla : \Gamma(TM) \rightarrow \Gamma(T^*M \otimes TM)$  on  $M$  induces a connection  $\nabla^S : \Gamma(S) \rightarrow \Gamma(T^*M \otimes S)$ . By adding a additional torsion term  $t \in \Omega^1(M, \text{End}TM)$  we obtain a new covariant derivative

$$\widetilde{\nabla} := \nabla + t \quad (2.1)$$

on the tangent bundle  $TM$ . Since  $t$  is really a one-form on  $M$  with values in the bundle of skew endomorphism  $Sk(TM)$  in [22],  $\nabla$  is in fact compatible with the Riemannian metric  $g$  and therefore also induces a connection  $\widetilde{\nabla}^S := \nabla^S + T$  on the spinor bundle. Here  $T \in \Omega^1(M, \text{End}S)$  denotes the 'lifted' torsion term  $t \in \Omega^1(M, \text{End}TM)$ .

Next, we will briefly discuss the construction of this connection. Again, we write  $\widetilde{\nabla}_X Y = \nabla_X Y + A(X, Y)$  with the Levi-Civita connection  $\nabla$ . For any  $X \in T_p M$  the endomorphism  $A(X, \cdot)$  is skew-adjoint and hence it is an element of  $\mathfrak{so}(T_p M)$ , we can express it as

$$A(X, \cdot) = \sum_{i < j} \alpha_{ij} E_i \wedge E_j. \quad (2.2)$$

Here  $E_i \wedge E_j$  is meant as the endomorphism of  $T_p M$  defined by  $E_i \wedge E_j$ . For any  $X \in T_p M$  one determines the coefficients in (2.2) by

$$\alpha_{ij} = \langle A(X, E_i), E_j \rangle = A_{X E_i E_j}. \quad (2.3)$$

Each  $E_i \wedge E_j$  lifts to  $\frac{1}{2} E_i \cdot E_j$  in  $spin(n)$ , and the spinor connection induced by  $\widetilde{\nabla}$  is locally given by

$$\widetilde{\nabla}_X \psi = \nabla_X \psi + \frac{1}{2} \sum_{i < j} \alpha_{ij} E_i \cdot E_j \psi = \nabla_X \psi + \frac{1}{2} \sum_{i < j} A_{X E_i E_j} E_i \cdot E_j \psi. \quad (2.4)$$

The connection given by (2.4) is compatible with the metric on spinors and with Clifford multiplication. Then, the Dirac operator associated to the spinor connection from (2.4) is defined as

$$\begin{aligned} D_T \psi &= \sum_{i=1}^n E_i \tilde{\nabla}_{E_i} \psi = D\psi + \frac{1}{2} \sum_{i=1}^n \sum_{j < k} A_{E_i E_j E_k} E_i \cdot E_j \cdot E_k \psi \\ &= D\psi + \frac{1}{4} \sum_{i,j,k=1}^n A_{E_i E_j E_k} E_i \cdot E_j \cdot E_k \psi \end{aligned} \quad (2.5)$$

where  $D$  is the Dirac operator induced by the Levi-Civita connection and “ $\cdot$ ” is the Clifford multiplication. Let  $c(e)$  be the Clifford operators acting on  $S(TM)$ , satisfying

$$c(e_i)c(e_j) + c(e_j)c(e_i) = -2\langle e_i, e_j \rangle. \quad (2.6)$$

For any orthogonal connection  $\tilde{\nabla}$  on the tangent bundle of  $M$  there exist a unique vector field  $V$ , we have

$$D_T \psi = D\psi + \frac{3}{2}T \cdot \psi - \frac{n-1}{2}V \cdot \psi, \quad D_T^* \psi = D\psi + \frac{3}{2}T \cdot \psi + \frac{n-1}{2}V \cdot \psi, \quad (2.7)$$

where Clifford multiplication by any 3-form is self-adjoint. As the Clifford multiplication by the vector field  $V$  is skew-adjoint we get that  $D_T$  is symmetric with respect to the natural  $L^2$ -scalar product on spinors if and only if the vectorial component of the torsion vanishes,  $V \equiv 0$ . Note that the Cartan type torsion  $S$  does not contribute to the Dirac operator  $D_T$ . As  $D_T^* D_T$  is a generalized Laplacian, one has the following Lichnerowicz formula.

**Lemma 2.1.** [20] *For the Dirac operator  $D_T$  associated to the orthogonal connection  $\tilde{\nabla}$ , we have*

$$\begin{aligned} D_T^* D_T \psi &= \Delta \psi + \frac{1}{4}R^g \psi + \frac{3}{2}dT \cdot \psi - \frac{3}{4} \| T \|^2 \psi \\ &\quad + \frac{n-1}{2} \operatorname{div}^g(V) \psi + \left(\frac{n-1}{2}\right)^2 (2-n) |V|^2 \psi \\ &\quad + 3(n-1)(T \cdot V \cdot \psi + (V \lrcorner T) \cdot \psi), \end{aligned} \quad (2.8)$$

for any spinor field  $\psi$ , where  $\Delta$  is the Laplacian associated to the connection

$$\tilde{\nabla}_X \psi = \nabla_X \psi + \frac{3}{2}(X \lrcorner T) \cdot \psi - \frac{n-1}{2}V \cdot X \cdot \psi - \frac{n-1}{2}\langle V, X \rangle \psi. \quad (2.9)$$

The following lemma of Dabrowski etc.’s Einstein functional play a key role in our proof of the Einstein functional with torsion. Let  $V, W$  be a pair of vector fields on a compact Riemannian manifold  $M$ , of dimension  $n = 2m$ . Using the Laplace operator  $\Delta_T^{-1} = (D_T^* D_T)^{-1} = \Delta + E$  acting on sections of a vector bundle  $S(TM)$  of rank  $2^m$  which may contain both some nontrivial connections and torsion, the spectral functionals over vector fields defined by

**Lemma 2.2.** [1] *The Einstein functional equal to*

$$Wres(\tilde{\nabla}_V \tilde{\nabla}_W \Delta_T^{-m}) = \frac{v_{n-1}}{6} 2^m \int_M G(V, W) \operatorname{vol}_g + \frac{v_{n-1}}{2} \int_M F(V, W) \operatorname{vol}_g + \frac{1}{2} \int_M (\operatorname{Tr} E) g(V, W) \operatorname{vol}_g, \quad (2.10)$$

where  $G(V, W)$  denotes the Einstein tensor evaluated on the two vector fields,  $F(V, W) = \operatorname{Tr}(V_a W_b F_{ab})$  and  $F_{ab}$  is the curvature tensor of the connection  $T$ ,  $\operatorname{Tr} E$  denotes the trace of  $E$  and  $v_{n-1} = \frac{2\pi^m}{\Gamma(m)}$ .

The aim of this section is to prove the following.

**Theorem 2.3.** For the Laplace (type) operator with torsion  $\Delta_T$ , the Einstein functional equal to

$$\begin{aligned} Wres(\tilde{\nabla}_V \tilde{\nabla}_W \Delta_{T,E}^{-m}) &= \frac{2^{m+1} \pi^m}{6\Gamma(m)} \int_M (Ric(V, W) - \frac{1}{2} sg(V, W)) vol_g \\ &\quad + \int_M 2^{m-1} \left( -\frac{1}{4} R^g - \frac{3}{2} div^g(V) + \frac{3}{2} \|T\|^2 + \frac{9}{2} \|V\|^2 \right) g(V, W) vol_g, \end{aligned} \quad (2.11)$$

where  $\int_M div(V) dVol_M = -\int_{\partial M} g(n, V) dVol_{\partial M}$ ,  $R_g$  denotes the curvature tensor and  $s$  is the scalar curvature.

*Proof.* Let  $T(X, \cdot, \cdot) = \sum_{1 \leq i < j \leq n} T(X, e_i, e_j) e_i^* \wedge e_j^*$  and  $c(T(X, \cdot, \cdot)) = \sum_{1 \leq i < j \leq n} T(X, e_i, e_j) c(e_i) c(e_j)$ , repeated application of (2.11) yields that

$$\begin{aligned} \tilde{\nabla}_X \psi &= \nabla_X^{S(TM)} \psi + \frac{3}{2} T(X, \cdot, \cdot) \psi - \frac{n-1}{2} c(V) c(X) \psi - \frac{n-1}{2} \langle V, X \rangle \psi \\ &= X \psi + \sigma(X) \psi + \frac{3}{2} \sum_{1 \leq i < j \leq n} T(X, e_i, e_j) c(e_i) c(e_j) \\ &\quad - \frac{n-1}{2} c(V) c(X) \psi - \frac{n-1}{2} \langle V, X \rangle \psi \\ &= X \psi + \bar{A}(X) \psi, \end{aligned} \quad (2.12)$$

where

$$\sigma(X) = -\frac{1}{4} \sum_{s,t} \omega_{s,t}(X) c(e_s) c(e_t). \quad (2.13)$$

Let  $V = \sum_{a=1}^n V^a e_a, W = \sum_{b=1}^n W^b e_b$ , in view of that

$$F(V, W) = Tr(V_a W_b F_{ab}) = \sum_{a,b=1}^n V^a W^b Tr^{S(TM)}(F_{e_a, e_b}), \quad (2.14)$$

we obtain

$$\begin{aligned} F_{e_a, e_b} &= (e_a + \bar{A}(e_a))(e_b + \bar{A}(e_b)) - (e_b + \bar{A}(e_b))(e_a + \bar{A}(e_a)) - ([e_a, e_a] + \bar{A}([e_a, e_b])) \\ &= e_a \circ \bar{A}(e_b) + \bar{A}(e_a) \circ e_b + \bar{A}(e_a) \bar{A}(e_b) - e_b \circ \bar{A}(e_a) - \bar{A}(e_b) \circ e_a \\ &\quad - \bar{A}(e_b) \bar{A}(e_a) - \bar{A}([e_a, e_b]) \\ &= \bar{A}(e_b) e_a + e_a \bar{A}(e_b) + \bar{A}(e_a) \circ e_b + \bar{A}(e_a) \bar{A}(e_b) - \bar{A}(e_a) \circ e_b - e_b \bar{A}(e_a) \\ &\quad - \bar{A}(e_b) e_a - \bar{A}(e_b) \bar{A}(e_a) - \bar{A}([e_a, e_b]) \\ &= e_a \bar{A}(e_b) - e_b \bar{A}(e_a) + \bar{A}(e_a) \bar{A}(e_b) - \bar{A}(e_b) \bar{A}(e_a) - \bar{A}([e_a, e_b]). \end{aligned} \quad (2.15)$$

Also, straightforward computations yield

$$\begin{aligned}
\mathrm{Tr}^{S(TM)}(e_a(\overline{A}(e_b))) &= \mathrm{Tr}^{S(TM)} \left[ e_a \left( -\frac{1}{4} \sum_{s,t} \omega_{s,t}(e_b) c(e_s) c(e_t) + \frac{3}{2} \sum_{1 \leq i < j \leq n} T(e_b, e_i, e_j) c(e_i) c(e_j) \right. \right. \\
&\quad \left. \left. - \frac{n-1}{2} c(V) c(e_b) - \frac{n-1}{2} \langle V, e_b \rangle \right) \right] \\
&= \mathrm{Tr}^{S(TM)} \left[ -\frac{1}{4} \sum_{s,t} e_a(\omega_{s,t}(e_b)) c(e_s) c(e_t) + \frac{3}{2} \sum_{1 \leq i < j \leq n} e_a(T(e_b, e_i, e_j)) c(e_i) c(e_j) \right. \\
&\quad \left. - \frac{n-1}{2} \sum_{k=1}^n e_a(V^k) c(e_k) c(e_b) - \frac{n-1}{2} e_a(V^b) \right] \\
&= \mathrm{Tr}^{S(TM)} \left[ -\frac{1}{8} \sum_{s=t} R_{abst}^{TM} c(e_s) c(e_t) + \frac{3}{2} \sum_{1 \leq i < j \leq n} e_a(T(e_b, e_i, e_j)) c(e_i) c(e_j) \right. \\
&\quad \left. - \frac{n-1}{2} \sum_{k=1}^n e_a(c(V^k)) c(e_k) c(e_b) - \frac{n-1}{2} e_a(V^b) \right] \\
&= -\frac{1}{8} \sum_{s=t} R_{abst}^{TM} (-\delta_{st}) \mathrm{Tr}^{S(TM)}(Id) - \frac{n-1}{2} e_a(V^b) \mathrm{Tr}^{S(TM)}(Id) \\
&\quad - \frac{n-1}{2} \sum_{k=1}^n e_a(c(V^k)) (-\delta_{kb}) \mathrm{Tr}^{S(TM)}(Id) \\
&= 0,
\end{aligned} \tag{2.16}$$

where

$$e_a(\omega_{s,t}(e_b))(x_0) = \frac{\partial}{\partial y^a} (\omega_{s,t}(e_b))(x_0) = \frac{\partial}{\partial y^a} (H_{b,s,t})(x_0) = \frac{1}{2} R_{abst}^{TM}(x_0), \tag{2.17}$$

where we take the normal coordinate about  $x_0$ , it follows that

$$\begin{aligned}
&\mathrm{Tr}^{S(TM)}(\overline{A}(e_a)\overline{A}(e_b) - \overline{A}(e_b)\overline{A}(e_a)) \\
&= \mathrm{Tr}^{S(TM)} \left[ \left( \frac{3}{2} \sum_{1 \leq i < j \leq n} T(e_a, e_i, e_j) c(e_i) c(e_j) - \frac{n-1}{2} c(V) c(e_a) - \frac{n-1}{2} \langle V, e_a \rangle \right) \right. \\
&\quad \times \left. \left( \frac{3}{2} \sum_{1 \leq i < j \leq n} T(e_b, e_i, e_j) c(e_i) c(e_j) - \frac{n-1}{2} c(V) c(e_b) - \frac{n-1}{2} \langle V, e_b \rangle \right) \right] \\
&\quad - \mathrm{Tr}^{S(TM)} \left[ \left( \frac{3}{2} \sum_{1 \leq i < j \leq n} T(e_b, e_i, e_j) c(e_i) c(e_j) - \frac{n-1}{2} c(V) c(e_b) - \frac{n-1}{2} \langle V, e_b \rangle \right) \right. \\
&\quad \times \left. \left( \frac{3}{2} \sum_{1 \leq i < j \leq n} T(e_a, e_i, e_j) c(e_i) c(e_j) - \frac{n-1}{2} c(V) c(e_a) - \frac{n-1}{2} \langle V, e_a \rangle \right) \right] \\
&= 0,
\end{aligned} \tag{2.18}$$

and

$$\begin{aligned}
&\mathrm{Tr}^{S(TM)}(\overline{A}([e_a, e_b]))(x_0) \\
&= \mathrm{Tr}^{S(TM)} \left( \sigma([e_a, e_b]) + \frac{3}{2} \sum_{1 \leq i < j \leq n} T([e_a, e_b], e_i, e_j) c(e_i) c(e_j) - \frac{n-1}{2} c(V) c([e_a, e_b]) \right. \\
&\quad \left. - \frac{n-1}{2} \langle V, [e_a, e_b] \rangle \right)(x_0) \\
&= 0.
\end{aligned} \tag{2.19}$$

Let  $\Delta_T = \Delta + E$ , by (53) in [2], we have

$$E = \left( -\frac{1}{4}R^g - \frac{3}{2}\text{div}^g(V) + \frac{3}{2}\|T\|^2 + \frac{9}{2}\|V\|^2 \right) Id_{S(TM)} - \frac{3}{2}dT - 9T \cdot V - 9V \lrcorner T, \quad (2.20)$$

Now, what is left is to show that

$$\begin{aligned} \text{Tr}^{S(TM)}(E) &= 2^{\frac{n}{2}} \left( -\frac{1}{4}R^g - \frac{3}{2}\text{div}^g(V) + \frac{3}{2}\|T\|^2 + \frac{9}{2}\|V\|^2 \right) \\ &\quad - \text{Tr}^{S(TM)}\left(\frac{3}{2}dT\right) - 9\text{Tr}^{S(TM)}(T \cdot V) - 9\text{Tr}^{S(TM)}(V \lrcorner T) \\ &= 2^{\frac{n}{2}} \left( -\frac{1}{4}R^g - \frac{3}{2}\text{div}^g(V) + \frac{3}{2}\|T\|^2 + \frac{9}{2}\|V\|^2 \right) \\ &\quad - \frac{3}{2}\text{Tr}^{S(TM)}\left(\sum_{1 \leq \alpha < j < k < l \leq n} dT(e_\alpha, e_j, e_k, e_l)c(e_\alpha)c(e_j)c(e_k)c(e_l)\right) \\ &\quad - 9\text{Tr}^{S(TM)}\left(\sum_{1 \leq \alpha < j < k \leq n} T(e_\alpha, e_j, e_k)c(e_\alpha)c(e_j)c(e_k)c(V)\right) \\ &\quad - 9\text{Tr}^{S(TM)}\left(\sum_{1 \leq i < j \leq n} T(V, e_i, e_j)c(e_i)c(e_j)\right) \\ &= 2^{\frac{n}{2}} \left( -\frac{1}{4}R^g - \frac{3}{2}\text{div}^g(V) + \frac{3}{2}\|T\|^2 + \frac{9}{2}\|V\|^2 \right) \\ &\quad - 9 \sum_{l=1}^n V_l \sum_{1 \leq \alpha < j < k \leq n} T(e_\alpha, e_j, e_k) \text{Tr}^{S(TM)}(c(e_\alpha)c(e_j)c(e_k)c(e_l)) \\ &= 2^{\frac{n}{2}} \left( -\frac{1}{4}R^g - \frac{3}{2}\text{div}^g(V) + \frac{3}{2}\|T\|^2 + \frac{9}{2}\|V\|^2 \right). \end{aligned} \quad (2.21)$$

Summing up (2.18)-(2.21) leads to the desired equality (2.11), and the proof of the Theorem is complete.  $\square$

### 3. Noncommutative residue for manifold with boundary

In this section, to define the noncommutative residue for Dirac operator with torsion, some basic facts and formulae about Boutet de Monvel's calculus can be find in Sec.2 in [15]. Let  $M$  be an  $n$ -dimensional compact oriented manifold with boundary  $\partial M$ . Some basic facts and formulae about Boutet de Monvel's calculus are recalled as follows.

Let

$$F : L^2(\mathbf{R}_t) \rightarrow L^2(\mathbf{R}_v); F(u)(v) = \int e^{-ivt} u(t) dt$$

denote the Fourier transformation and  $\varphi(\overline{\mathbf{R}^+}) = r^+ \varphi(\mathbf{R})$  (similarly define  $\varphi(\overline{\mathbf{R}^-})$ ), where  $\varphi(\mathbf{R})$  denotes the Schwartz space and

$$r^+ : C^\infty(\mathbf{R}) \rightarrow C^\infty(\overline{\mathbf{R}^+}); f \rightarrow f|_{\overline{\mathbf{R}^+}; \overline{\mathbf{R}^+} = \{x \geq 0; x \in \mathbf{R}\}. \quad (3.1)$$

We define  $H^+ = F(\varphi(\overline{\mathbf{R}^+}))$ ;  $H_0^- = F(\varphi(\overline{\mathbf{R}^-}))$  which are orthogonal to each other. We have the following property:  $h \in H^+$  ( $H_0^-$ ) iff  $h \in C^\infty(\mathbf{R})$  which has an analytic extension to the lower (upper) complex half-plane  $\{\text{Im}\xi < 0\}$  ( $\{\text{Im}\xi > 0\}$ ) such that for all nonnegative integer  $l$ ,

$$\frac{d^l h}{d\xi^l}(\xi) \sim \sum_{k=1}^{\infty} \frac{d^l}{d\xi^l} \left( \frac{c_k}{\xi^k} \right) \quad (3.2)$$

as  $|\xi| \rightarrow +\infty, \text{Im}\xi \leq 0$  ( $\text{Im}\xi \geq 0$ ).

Let  $H'$  be the space of all polynomials and  $H^- = H_0^- \oplus H'$ ;  $H = H^+ \oplus H^-$ . Denote by  $\pi^+$  ( $\pi^-$ ) respectively the projection on  $H^+$  ( $H^-$ ). For calculations, we take  $H = \tilde{H} = \{\text{rational functions having no poles on the real axis}\}$  ( $\tilde{H}$  is a dense set in the topology of  $H$ ). Then on  $\tilde{H}$ ,

$$\pi^+ h(\xi_0) = \frac{1}{2\pi i} \lim_{u \rightarrow 0^-} \int_{\Gamma^+} \frac{h(\xi)}{\xi_0 + iu - \xi} d\xi, \quad (3.3)$$

where  $\Gamma^+$  is a Jordan close curve included  $\text{Im}\xi > 0$  surrounding all the singularities of  $h$  in the upper half-plane and  $\xi_0 \in \mathbf{R}$ . Similarly, define  $\pi'$  on  $\tilde{H}$ ,

$$\pi' h = \frac{1}{2\pi} \int_{\Gamma^+} h(\xi) d\xi. \quad (3.4)$$

So,  $\pi'(H^-) = 0$ . For  $h \in H \cap L^1(\mathbf{R})$ ,  $\pi' h = \frac{1}{2\pi} \int_{\mathbf{R}} h(v) dv$  and for  $h \in H^+ \cap L^1(\mathbf{R})$ ,  $\pi' h = 0$ . Denote by  $\mathcal{B}$  Boutet de Monvel's algebra (for details, see Section 2 of [15]).

An operator of order  $m \in \mathbf{Z}$  and type  $d$  is a matrix

$$A = \begin{pmatrix} \pi^+ P + G & K \\ T & S \end{pmatrix} : \begin{array}{c} C^\infty(X, E_1) \\ \oplus \\ C^\infty(\partial X, F_1) \end{array} \longrightarrow \begin{array}{c} C^\infty(X, E_2) \\ \oplus \\ C^\infty(\partial X, F_2) \end{array}.$$

where  $X$  is a manifold with boundary  $\partial X$  and  $E_1, E_2$  ( $F_1, F_2$ ) are vector bundles over  $X$  ( $\partial X$ ). Here,  $P : C_0^\infty(\Omega, \overline{E_1}) \rightarrow C^\infty(\Omega, \overline{E_2})$  is a classical pseudodifferential operator of order  $m$  on  $\Omega$ , where  $\Omega$  is an open neighborhood of  $X$  and  $\overline{E_i}|_X = E_i$  ( $i = 1, 2$ ).  $P$  has an extension:  $\mathcal{E}'(\Omega, \overline{E_1}) \rightarrow \mathcal{D}'(\Omega, \overline{E_2})$ , where  $\mathcal{E}'(\Omega, \overline{E_1})$  ( $\mathcal{D}'(\Omega, \overline{E_2})$ ) is the dual space of  $C^\infty(\Omega, \overline{E_1})$  ( $C_0^\infty(\Omega, \overline{E_2})$ ). Let  $e^+ : C^\infty(X, E_1) \rightarrow \mathcal{E}'(\Omega, \overline{E_1})$  denote extension by zero from  $X$  to  $\Omega$  and  $r^+ : \mathcal{D}'(\Omega, \overline{E_2}) \rightarrow \mathcal{D}'(\Omega, E_2)$  denote the restriction from  $\Omega$  to  $X$ , then define

$$\pi^+ P = r^+ P e^+ : C^\infty(X, E_1) \rightarrow \mathcal{D}'(\Omega, E_2).$$

In addition,  $P$  is supposed to have the transmission property; this means that, for all  $j, k, \alpha$ , the homogeneous component  $p_j$  of order  $j$  in the asymptotic expansion of the symbol  $p$  of  $P$  in local coordinates near the boundary satisfies:

$$\partial_{x_n}^k \partial_{\xi'}^\alpha p_j(x', 0, 0, +1) = (-1)^{j-|\alpha|} \partial_{x_n}^k \partial_{\xi'}^\alpha p_j(x', 0, 0, -1),$$

then  $\pi^+ P : C^\infty(X, E_1) \rightarrow C^\infty(X, E_2)$  by Section 2.1 of [15].

Let  $M$  be a compact manifold with boundary  $\partial M$ . We assume that the metric  $g^M$  on  $M$  has the following form near the boundary

$$g^M = \frac{1}{h(x_n)} g^{\partial M} + dx_n^2, \quad (3.5)$$

where  $g^{\partial M}$  is the metric on  $\partial M$ . Let  $U \subset M$  be a collar neighborhood of  $\partial M$  which is diffeomorphic  $\partial M \times [0, 1)$ . By the definition of  $h(x_n) \in C^\infty([0, 1))$  and  $h(x_n) > 0$ , there exists  $\tilde{h} \in C^\infty((-\varepsilon, 1))$  such that  $\tilde{h}|_{[0, 1)} = h$  and  $\tilde{h} > 0$  for some sufficiently small  $\varepsilon > 0$ . Then there exists a metric  $\hat{g}$  on  $\hat{M} = M \cup_{\partial M} \partial M \times (-\varepsilon, 0]$  which has the form on  $U \cup_{\partial M} \partial M \times (-\varepsilon, 0]$

$$\hat{g} = \frac{1}{\tilde{h}(x_n)} g^{\partial M} + dx_n^2, \quad (3.6)$$

such that  $\hat{g}|_M = g$ . We fix a metric  $\hat{g}$  on the  $\hat{M}$  such that  $\hat{g}|_M = g$ . Now we recall the main theorem in [14].

**Theorem 3.1. (Fedosov-Golse-Leichtnam-Schrohe)** *Let  $X$  and  $\partial X$  be connected,  $\dim X = n \geq 3$ ,  $A = \begin{pmatrix} \pi^+ P + G & K \\ T & S \end{pmatrix} \in \mathcal{B}$ , and denote by  $p$ ,  $b$  and  $s$  the local symbols of  $P$ ,  $G$  and  $S$  respectively. Define:*

$$\begin{aligned} \widetilde{\text{Wres}}(A) &= \int_X \int_{\mathbf{S}} \text{Tr}_E [p_{-n}(x, \xi)] \sigma(\xi) dx \\ &\quad + 2\pi \int_{\partial X} \int_{\mathbf{S}'} \{ \text{Tr}_E [(\text{Tr} b_{-n})(x', \xi')] + \text{Tr}_F [s_{1-n}(x', \xi')] \} \sigma(\xi') dx', \end{aligned} \quad (3.7)$$

Then a)  $\widetilde{\text{Wres}}([A, B]) = 0$ , for any  $A, B \in \mathcal{B}$ ; b) It is a unique continuous trace on  $\mathcal{B}/\mathcal{B}^{-\infty}$ .

Let  $p_1, p_2$  be nonnegative integers and  $p_1 + p_2 \leq n$ , denote by  $\sigma_l(A)$  the  $l$ -order symbol of an operator  $A$ , an application of (3.5) and (3.6) in [15] shows that

**Definition 3.2.** Lower-dimensional volumes of spin manifolds with boundary with torsion are defined by

$$\text{Vol}_n^{\{p_1, p_2\}} M := \widetilde{\text{Wres}}[\pi^+(\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-p_1}) \circ \pi^+((D_T^* D_T)^{-1})^{p_2}]. \quad (3.8)$$

where  $\pi^+(\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-p_1})$ ,  $\pi^+((D_T^* D_T)^{-1})^{p_2}$  are elements in Boutet de Monvel's algebra[16].

For Dirac operators with torsion  $\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-1}$  and  $(D_T^* D_T)^{-1}$ , denote by  $\sigma_l(A)$  the  $l$ -order symbol of an operator  $A$ . An application of (2.1.4) in [15] shows that

$$\begin{aligned} & \widetilde{\text{Wres}}[\pi^+(\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-1})^{p_1} \circ \pi^+(D_T^* D_T)^{-p_2}] \\ &= \int_M \int_{|\xi|=1} \text{Tr}_{S(TM)}[\sigma_{-n}(\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-p_1} \circ (D_T^* D_T)^{-p_2})] \sigma(\xi) dx + \int_{\partial M} \Phi, \end{aligned} \quad (3.9)$$

where

$$\begin{aligned} \Phi &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+\ell}}{\alpha!(j+k+1)!} \text{Tr}_{S(TM)}[\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+((\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-p_1})(x', 0, \xi', \xi_n)) \\ & \times \partial_x^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l((D_T^* D_T)^{-p_2})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx', \end{aligned} \quad (3.10)$$

and the sum is taken over  $r - k + |\alpha| + \ell - j - 1 = -n$ ,  $r \leq -p_1$ ,  $\ell \leq -p_2$ .

For Dirac operators with torsion  $\widetilde{\nabla}_X \widetilde{\nabla}_Y D_T^{-1}$  and  $(D_T^* D_T D_T^*)^{-1}$ , similarly we have

$$\begin{aligned} & \widetilde{\text{Wres}}[\pi^+(\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T)^{-1})^{p_1} \circ \pi^+(D_T^* D_T D_T^*)^{-p_2}] \\ &= \int_M \int_{|\xi|=1} \text{Tr}_{S(TM)}[\sigma_{-n}(\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T)^{-p_1} \circ (D_T^* D_T D_T^*)^{-p_2})] \sigma(\xi) dx + \int_{\partial M} \widetilde{\Phi}, \end{aligned} \quad (3.11)$$

where

$$\begin{aligned} \widetilde{\Phi} &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+\ell}}{\alpha!(j+k+1)!} \text{Tr}_{S(TM)}[\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+((\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T)^{-p_1})(x', 0, \xi', \xi_n)) \\ & \times \partial_x^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l((D_T^* D_T D_T^*)^{-p_2})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx', \end{aligned} \quad (3.12)$$

and the sum is taken over  $r - k + |\alpha| + \ell - j - 1 = -n$ ,  $r \leq -p_1$ ,  $\ell \leq -p_2$ .

#### 4. Residue for Dirac operators with torsion $\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-1}$ and $(D_T^* D_T)^{-1}$

In this section, we compute the lower dimensional volume for 4-dimension compact manifolds with boundary and get a Kastler-Kalau-Walze type formula in this case. We will consider  $D_T^* D_T$  (since  $D_T$  is not self-adjoint in general). Since  $[\sigma_{-4}(\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-1} \circ (D_T^* D_T)^{-1})]_M$  has the same expression as  $[\sigma_{-4}(\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-1} \circ (D_T^* D_T)^{-1})]_M$  in the case of manifolds without boundary, so locally we can use Theorem 2.4 to compute the first term.

**Theorem 4.1.** Let  $M$  be a 4-dimensional compact manifold without boundary and  $\widetilde{\nabla}$  be an orthogonal connection with torsion. Then we get the volumes associated to  $\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-1}$  and  $D_T^* D_T$  on compact manifolds without boundary

$$\begin{aligned} & \text{Wres}[\sigma_{-4}(\widetilde{\nabla}_X \widetilde{\nabla}_Y (D_T^* D_T)^{-1} \circ (D_T^* D_T)^{-1})] \\ &= \frac{4\pi^2}{3} \int_M \left( \text{Ric}(X, Y) - \frac{1}{2} \text{sg}(X, Y) \right) \text{vol}_g \\ & \quad + \int_M \left( -\frac{1}{2} R^g - 3 \text{div}^g(X) + 3 \|T\|^2 + 9 \|X\|^2 \right) g(X, Y) \text{vol}_g, \end{aligned} \quad (4.1)$$



where  $R_g$  denotes the curvature tensor and  $s$  is the scalar curvature.

So we only need to compute  $\int_{\partial M} \Phi$ . Recall the definition of the Dirac operator  $D$  in [24]. Denote by  $\sigma_l(A)$  the  $l$ -order symbol of an operator  $A$ . In the local coordinates  $\{x_i; 1 \leq i \leq n\}$  and the fixed orthonormal frame  $\{\tilde{e}_1, \dots, \tilde{e}_n\}$ , the connection matrix  $(\omega_{s,t})$  is defined by

$$\tilde{\nabla}(\tilde{e}_1, \dots, \tilde{e}_n) = (\tilde{e}_1, \dots, \tilde{e}_n)(\omega_{s,t}). \quad (4.2)$$

The Dirac operator

$$D = \sum_{i=1}^n c(\tilde{e}_i) \left[ \tilde{e}_i - \frac{1}{4} \sum_{s,t} \omega_{s,t}(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) \right], \quad (4.3)$$

where  $c(\tilde{e}_i)$  denotes the Clifford action. Then

$$\begin{aligned} D_T &= \sum_{i=1}^n c(\tilde{e}_i) \left[ \tilde{e}_i - \frac{1}{4} \sum_{s,t} \omega_{s,t}(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) \right] + \frac{1}{4} \sum_{i \neq s \neq t} A_{ist} c(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) \\ &\quad + \frac{1}{4} \sum_{i,s,t} [-A_{iit} c(\tilde{e}_t) + A_{isi} c(\tilde{e}_s) - A_{iss} c(\tilde{e}_i) + 2A_{iii} c(\tilde{e}_i)], \end{aligned} \quad (4.4)$$

$$\begin{aligned} D_T^* &= \sum_{i=1}^n c(\tilde{e}_i) \left[ \tilde{e}_i - \frac{1}{4} \sum_{s,t} \omega_{s,t}(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) \right] + \frac{1}{4} \sum_{i \neq s \neq t} A_{ist} c(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) \\ &\quad - \frac{1}{4} \sum_{i,s,t} [-A_{iit} c(\tilde{e}_t) + A_{isi} c(\tilde{e}_s) - A_{iss} c(\tilde{e}_i) + 2A_{iii} c(\tilde{e}_i)], \end{aligned} \quad (4.5)$$

and

$$\sigma_1(D_T) = \sigma_1(D_T^*) = \sqrt{-1}c(\xi); \quad (4.6)$$

$$\begin{aligned} \sigma_0(D_T) &= -\frac{1}{4} \sum_{i,s,t} \omega_{s,t}(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) + \frac{1}{4} \sum_{i \neq s \neq t} A_{ist} c(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) \\ &\quad + \frac{1}{4} \sum_{i,s,t} [-A_{iit} c(\tilde{e}_t) + A_{isi} c(\tilde{e}_s) - A_{iss} c(\tilde{e}_i) + 2A_{iii} c(\tilde{e}_i)], \end{aligned} \quad (4.7)$$

$$\begin{aligned} \sigma_0(D_T^*) &= -\frac{1}{4} \sum_{i,s,t} \omega_{s,t}(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) + \frac{1}{4} \sum_{i \neq s \neq t} A_{ist} c(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) \\ &\quad - \frac{1}{4} \sum_{i,s,t} [-A_{iit} c(\tilde{e}_t) + A_{isi} c(\tilde{e}_s) - A_{iss} c(\tilde{e}_i) + 2A_{iii} c(\tilde{e}_i)]. \end{aligned} \quad (4.8)$$

We define  $\nabla_X^{S(TM)} := X + \frac{1}{4} \sum_{ij} \langle \nabla_X^L e_i, e_j \rangle c(e_i) c(e_j)$ , which is a spin connection. Set

$$A(X) = \frac{1}{4} \sum_{ij} \langle \nabla_X^L e_i, e_j \rangle c(e_i) c(e_j). \quad (4.9)$$

Let  $\tilde{\nabla}_X = X + A(X) + \bar{T}(X, \cdot, \cdot)$ ,  $\bar{T}(X, \cdot, \cdot) = \frac{3}{2} \sum_{1 \leq i < j \leq n} T(X, e_i, e_j) c(e_i) c(e_j) - \frac{1}{2} c(V) c(X) - \frac{1}{2} \langle V, X \rangle$  and  $\tilde{\nabla}_Y = Y + A(Y) + \bar{T}(Y, \cdot, \cdot)$ , by (2.11), we obtain

$$\begin{aligned} \tilde{\nabla}_X \tilde{\nabla}_Y &= (X + A(X) + \bar{T}(X, \cdot, \cdot))(Y + A(Y) + \bar{T}(Y, \cdot, \cdot)) \\ &= XY + X[A(Y)] + A(Y)X + A(X)Y + A(X)A(Y) + \bar{T}(X, \cdot)Y + \bar{T}(X, \cdot, \cdot)A(Y) \\ &\quad + X[\bar{T}(Y, \cdot, \cdot)] + \bar{T}(Y, \cdot, \cdot)X + A(X)\bar{T}(X, \cdot, \cdot) + \bar{T}(Y, \cdot, \cdot)\bar{T}(Y, \cdot, \cdot), \end{aligned} \quad (4.10)$$

where  $X = \sum_{j=1}^n X_j \partial_{x_j}$ ,  $Y = \sum_{l=1}^n Y_l \partial_{x_l}$ .

Let  $g^{ij} = g(dx_i, dx_j)$ ,  $\xi = \sum_k \xi_j dx_j$  and  $\nabla_{\partial_i}^L \partial_j = \sum_k \Gamma_{ij}^k \partial_k$ , we get

$$\sigma_i = -\frac{1}{4} \sum_{s,t} \omega_{s,t}(e_i) c(e_s) c(e_t); \quad \xi^j = g^{ij} \xi_i; \quad \Gamma^k = g^{ij} \Gamma_{ij}^k; \quad \sigma^j = g^{ij} \sigma_i. \quad (4.11)$$

Then we have the following lemmas.

**Lemma 4.2.** *The following identities hold:*

$$\begin{aligned} \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y) &= X[A(Y)] + A(X)A(Y) + \bar{T}(X, \cdot, \cdot)A(Y) \\ &\quad + X[\bar{T}(Y, \cdot, \cdot)] + A(X)\bar{T}(X, \cdot, \cdot) + \bar{T}(Y, \cdot, \cdot)\bar{T}(Y, \cdot, \cdot); \end{aligned} \quad (4.12)$$

$$\begin{aligned} \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y) &= \sqrt{-1} \sum_{j,l=1}^n X_j \frac{\partial Y_l}{\partial x_j} \sqrt{-1} \xi_l + \sqrt{-1} \sum_j A(Y) X_j \xi_j + \sqrt{-1} \sum_l A(Y) Y_l \xi_l \\ &\quad + \sum_j \bar{T}(X, \cdot) Y_j \sqrt{-1} \xi_j + \sum_j \bar{T}(Y, \cdot) X_j \sqrt{-1} \xi_j; \end{aligned} \quad (4.13)$$

$$\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y) = - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l. \quad (4.14)$$

Hence by Lemma 2.1 in [16], we have

**Lemma 4.3.** *The symbol of the Dirac operator*

$$\sigma_{-1}(D_T^{-1}) = \sigma_{-1}((D_T^*)^{-1}) = \frac{\sqrt{-1}c(\xi)}{|\xi|^2}; \quad (4.15)$$

$$\sigma_{-2}(D_T^{-1}) = \frac{c(\xi)\sigma_0(D_T)c(\xi)}{|\xi|^4} + \frac{c(\xi)}{|\xi|^6} \sum_j c(dx_j) \left[ \partial_{x_j}(c(\xi))|\xi|^2 - c(\xi)\partial_{x_j}(|\xi|^2) \right]; \quad (4.16)$$

$$\sigma_{-2}((D_T^*)^{-1}) = \frac{c(\xi)\sigma_0(D_T^*)c(\xi)}{|\xi|^4} + \frac{c(\xi)}{|\xi|^6} \sum_j c(dx_j) \left[ \partial_{x_j}(c(\xi))|\xi|^2 - c(\xi)\partial_{x_j}(|\xi|^2) \right]. \quad (4.17)$$

Let  $u = \frac{1}{4} \sum_{i \neq s \neq t} A_{ist} c(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t)$ ,  $v = \frac{1}{4} \sum_{i,s,t} [-A_{iit} c(\tilde{e}_t) + A_{isi} c(\tilde{e}_s) - A_{iss} c(\tilde{e}_i) + 2A_{iii} c(\tilde{e}_i)]$ , we get

**Lemma 4.4.** *The following identities hold:*

$$\sigma_{-2}((D_T^* D_T)^{-1}) = |\xi|^2 = \sigma_{-2}(D^{-2}) = |\xi|^{-2}; \quad (4.18)$$

$$\begin{aligned} \sigma_{-3}((D_T^* D_T)^{-1}) &= -\sqrt{-1} |\xi|^{-4} \xi_k (\Gamma^k - 2\delta^k) - \sqrt{-1} |\xi|^{-6} 2\xi^j \xi_\alpha \xi_\beta \partial_j g^{\alpha\beta} \\ &\quad - ((u-v)\sqrt{-1}c(\xi) + \sqrt{-1}c(\xi)(u+v)) |\xi|^{-4}. \end{aligned} \quad (4.19)$$

By Lemma 4.2, Lemma 4.4 and  $\sigma(p_1 \circ p_2) = \sum_\alpha \frac{1}{\alpha!} \partial_\xi^\alpha [\sigma(p_1)] D_x^\alpha [\sigma(p_2)]$ , then

**Lemma 4.5.** *The following identities hold:*

$$\sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) = - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l |\xi|^{-2}; \quad (4.20)$$

$$\begin{aligned} \sigma_{-1}(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) &= \sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-3}((D_T^* D_T)^{-1}) + \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-2}((D_T^* D_T)^{-1}) \\ &\quad + \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] D_{x_j} [\sigma_{-2}((D_T^* D_T)^{-1})]. \end{aligned} \quad (4.21)$$

Since  $\Phi$  is a global form on  $\partial M$ , so for any fixed point  $x_0 \in \partial M$ , we can choose the normal coordinates  $U$  of  $x_0$  in  $\partial M$  (not in  $M$ ) and compute  $\Phi(x_0)$  in the coordinates  $\tilde{U} = U \times [0, 1)$  and the metric  $\frac{1}{h(x_n)}g^{\partial M} + dx_n^2$ . The dual metric of  $g^{\partial M}$  on  $\tilde{U}$  is  $\frac{1}{h(x_n)}g^{\partial M} + dx_n^2$ . Write  $g_{ij}^M = g^M(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j})$ ;  $g_M^{ij} = g^M(dx_i, dx_j)$ , then

$$[g_{i,j}^M] = \begin{bmatrix} \frac{1}{h(x_n)}[g_{i,j}^{\partial M}] & 0 \\ 0 & 1 \end{bmatrix}; \quad [g_M^{i,j}] = \begin{bmatrix} h(x_n)[g_{\partial M}^{i,j}] & 0 \\ 0 & 1 \end{bmatrix}, \quad (4.22)$$

and

$$\partial_{x_s} g_{ij}^{\partial M}(x_0) = 0, \quad 1 \leq i, j \leq n-1; \quad g_{i,j}^M(x_0) = \delta_{ij}. \quad (4.23)$$

Let  $\{e_1, \dots, e_{n-1}\}$  be an orthonormal frame field in  $U$  about  $g^{\partial M}$  which is parallel along geodesics and  $e_i = \frac{\partial}{\partial x_i}(x_0)$ , then  $\{\tilde{e}_1 = \sqrt{h(x_n)}e_1, \dots, \tilde{e}_{n-1} = \sqrt{h(x_n)}e_{n-1}, \tilde{e}_n = dx_n\}$  is the orthonormal frame field in  $\tilde{U}$  about  $g^M$ . Locally  $S(TM)|_{\tilde{U}} \cong \tilde{U} \times \wedge_C^*(\frac{n}{2})$ . Let  $\{f_1, \dots, f_n\}$  be the orthonormal basis of  $\wedge_C^*(\frac{n}{2})$ . Take a spin frame field  $\sigma: \tilde{U} \rightarrow Spin(M)$  such that  $\pi\sigma = \{\tilde{e}_1, \dots, \tilde{e}_n\}$  where  $\pi: Spin(M) \rightarrow O(M)$  is a double covering, then  $\{[\sigma, f_i], 1 \leq i \leq 4\}$  is an orthonormal frame of  $S(TM)|_{\tilde{U}}$ . In the following, since the global form  $\Phi$  is independent of the choice of the local frame, so we can compute  $\text{tr}_{S(TM)}$  in the frame  $\{[\sigma, f_i], 1 \leq i \leq 4\}$ . Let  $\{E_1, \dots, E_n\}$  be the canonical basis of  $R^n$  and  $c(E_i) \in cl_C(n) \cong Hom(\wedge_C^*(\frac{n}{2}), \wedge_C^*(\frac{n}{2}))$  be the Clifford action. By [24], then

$$c(\tilde{e}_i) = [(\sigma, c(E_i))]; \quad c(\tilde{e}_i)[(\sigma, f_i)] = [\sigma, (c(E_i)f_i)]; \quad \frac{\partial}{\partial x_i} = [(\sigma, \frac{\partial}{\partial x_i})], \quad (4.24)$$

then we have  $\frac{\partial}{\partial x_i}c(\tilde{e}_i) = 0$  in the above frame. By Lemma 2.2 in [16], we have

**Lemma 4.6.** *With the metric  $\frac{1}{h(x_n)}g^{\partial M} + dx_n^2$  on  $M$  near the boundary*

$$\begin{aligned} \partial x_j(|\xi|_{g^M}^2)(x_0) &= 0, \quad \text{if } j < n; \quad = h'(0)|\xi'|_{g^{\partial M}}^2, \quad \text{if } j = n. \\ \partial x_j(c(\xi))(x_0) &= 0, \quad \text{if } j < n; \quad = \partial x_n(c(\xi'))(x_0), \quad \text{if } j = n. \end{aligned} \quad (4.25)$$

where  $\xi = \xi' + \xi_n dx_n$

Now we need to compute  $\int_{\partial M} \Phi$ . When  $n = 4$ , then  $\text{tr}_{S(TM)}[\mathbf{id}] = \dim(\wedge^*(\mathbb{R}^2)) = 4$ , the sum is taken over  $r + l - k - j - |\alpha| = -3$ ,  $r \leq 0$ ,  $l \leq -2$ , then we have the following five cases:

**case a) I)**  $r = 0$ ,  $l = -2$ ,  $k = j = 0$ ,  $|\alpha| = 1$ .

By (3.10), we get

$$\Phi_1 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{Tr}[\partial_{\xi'}^\alpha \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \times \partial_{x'}^\alpha \partial_{\xi_n} \sigma_{-2}((D_T^* D_T)^{-1})](x_0) d\xi_n \sigma(\xi') dx'. \quad (4.26)$$

By Lemma 2.2 in [16], for  $i < n$ , then

$$\partial_{x_i} \sigma_{-2}((D_T^* D_T)^{-1})(x_0) = \partial_{x_i} (|\xi|^{-2})(x_0) = - \frac{\partial_{x_i} (|\xi|^2)(x_0)}{|\xi|^4} = 0, \quad (4.27)$$

so  $\Phi_1 = 0$ .

**case a) II)**  $r = 0$ ,  $l = -2$ ,  $k = |\alpha| = 0$ ,  $j = 1$ .

By (3.10), we get

$$\Phi_2 = - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \times \partial_{\xi_n}^2 \sigma_{-2}((D_T^* D_T)^{-1})](x_0) d\xi_n \sigma(\xi') dx'. \quad (4.28)$$

By Lemma 4.4, we have

$$\partial_{\xi_n}^2 \sigma_{-2}((D_T^* D_T)^{-1})(x_0) = \partial_{\xi_n}^2 (|\xi|^{-2})(x_0) = \frac{6\xi_n^2 - 2}{(1 + \xi_n^2)^3}. \quad (4.29)$$

It follows that

$$\partial_{x_n} \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1})(x_0) = \partial_{x_n} \left( - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l |\xi|^{-2} \right) = \frac{\sum_{j,l=1}^n X_j Y_l \xi_j \xi_l h'(0) |\xi'|^2}{(1 + \xi_n^2)^2}. \quad (4.30)$$

By integrating formula, we obtain

$$\begin{aligned} \pi_{\xi_n}^+ \partial_{x_n} \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1})(x_0) &= \partial_{x_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \\ &= - \frac{i \xi_n}{4(\xi_n - i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) + \frac{2 - i \xi_n}{4(\xi_n - i)^2} X_n Y_n h'(0) \\ &\quad - \frac{i}{4(\xi_n - i)^2} \sum_{j=1}^{n-1} X_j Y_n \xi_j - \frac{i}{4(\xi_n - i)^2} \sum_{l=1}^{n-1} X_n Y_l \xi_l. \end{aligned} \quad (4.31)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case a) II**. From (4.29) and (4.31), we obtain

$$\begin{aligned} &\text{Tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \times \partial_{\xi_n}^2 \sigma_{-2}((D_T^* D_T)^{-1})](x_0) \\ &= 2 \frac{1 + \xi_n i - 3\xi_n^3 i - i}{(\xi_n - i)^5 (\xi_n + i)^3} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) + 2 \frac{1 + \xi_n i - 3\xi_n^3 i - i}{(\xi_n - i)^5 (\xi_n + i)^3} X_n Y_n h'(0) \\ &\quad + 2 \frac{(1 - 3\xi_n^2) i}{(\xi_n - i)^5 (\xi_n + i)^3} \sum_{j=1}^{n-1} X_j Y_n \xi_j + 2 \frac{(1 - 3\xi_n^2) i}{(\xi_n - i)^5 (\xi_n + i)^3} \sum_{l=1}^{n-1} X_n Y_l \xi_l. \end{aligned} \quad (4.32)$$

Therefore, we get

$$\begin{aligned} \Phi_2 &= - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left\{ 2 \frac{1 + \xi_n i - 3\xi_n^3 i - i}{(\xi_n - i)^5 (\xi_n + i)^3} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) + 2 \frac{1 + \xi_n i - 3\xi_n^3 i - i}{(\xi_n - i)^5 (\xi_n + i)^3} X_n Y_n h'(0) \right\} d\xi_n \sigma(\xi') dx' \\ &= - \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \int_{|\xi'|=1} \int_{\Gamma^+} \frac{1 + \xi_n i - 3\xi_n^3 i - i}{(\xi_n - i)^5 (\xi_n + i)^3} \xi_j \xi_l d\xi_n \sigma(\xi') dx' \\ &\quad - X_n Y_n h'(0) \Omega_3 \int_{\Gamma^+} \frac{1 + \xi_n i - 3\xi_n^3 i - i}{(\xi_n - i)^5 (\xi_n + i)^3} d\xi_n dx' \\ &= - \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \frac{4\pi}{3} \frac{2\pi i}{4!} \left[ \frac{1 + \xi_n i - 3\xi_n^3 i - i}{(\xi_n + i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' \\ &\quad - X_n Y_n h'(0) \Omega_3 \frac{2\pi i}{4!} \left[ \frac{1 + \xi_n i - 3\xi_n^3 i - i}{(\xi_n + i)^3} \right]^{(4)} \Big|_{\xi_n=i} dx' \\ &= \frac{13\pi^2}{24} \sum_{j=1}^{n-1} X_j Y_j h'(0) dx' + \frac{13}{32} X_n Y_n h'(0) \pi \Omega_3 dx', \end{aligned} \quad (4.33)$$

where  $\Omega_3$  is the canonical volume of  $S^2$ .

**case a) III**)  $r = 0$ ,  $l = -2$ ,  $j = |\alpha| = 0$ ,  $k = 1$ .

By (3.10), we get

$$\begin{aligned} \Phi_3 &= - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-2}(D_T^* D_T)^{-1}](x_0) d\xi_n \sigma(\xi') dx' \\ &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \times \partial_{x_n} \sigma_{-2}(D_T^* D_T)^{-1}](x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (4.34)$$

By Lemma 4.4, we have

$$\partial_{x_n} \sigma_{-2}(D_T^* D_T)^{-1}(x_0)|_{|\xi'|=1} = -\frac{h'(0)}{(1 + \xi_n^2)^2}. \quad (4.35)$$

An easy calculation gives

$$\begin{aligned} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1})(x_0)|_{|\xi'|=1} &= \frac{i}{2(\xi_n - i)} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{1}{2(\xi_n - i)} X_n Y_n \\ &\quad - \frac{1}{2(\xi_n - i)} \sum_{j=1}^{n-1} X_j Y_n \xi_j - \frac{1}{2(\xi_n - i)} \sum_{l=1}^{n-1} X_n Y_l \xi_l. \end{aligned} \quad (4.36)$$

Also, straightforward computations yield

$$\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1})(x_0)|_{|\xi'|=1} = \frac{i}{(\xi_n - i)^3} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{1}{(\xi_n - i)^3} X_n Y_n. \quad (4.37)$$

From (4.35) and (4.37), we obtain

$$\begin{aligned} &\text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-2}(D_T^* D_T)^{-1}](x_0) \\ &= -4 \frac{h'(0)i}{(\xi_n - i)^5 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l + 4 \frac{h'(0)}{(\xi_n - i)^5 (\xi_n + i)^2} X_n Y_n. \end{aligned} \quad (4.38)$$

Therefore, we get

$$\begin{aligned} \Phi_3 &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left( -4 \frac{h'(0)i}{(\xi_n - i)^5 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l + 4 \frac{h'(0)}{(\xi_n - i)^5 (\xi_n + i)^2} X_n Y_n \right) d\xi_n \sigma(\xi') dx' \\ &= -2 \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \int_{\Gamma^+} \frac{i}{(\xi_n - i)^5 (\xi_n + i)^2} \xi_j \xi_l d\xi_n dx' + 2 X_n Y_n h'(0) \Omega_3 \int_{\Gamma^+} \frac{1}{(\xi_n - i)^5 (\xi_n + i)^2} d\xi_n dx' \\ &= -2 \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \frac{2\pi i}{4!} \left[ \frac{i}{(\xi_n + i)^2} \right]^{(4)} \Big|_{\xi_n=i} dx' + 2 X_n Y_n h'(0) \Omega_3 \frac{2\pi i}{4!} \left[ \frac{1}{(\xi_n + i)^2} \right]^{(4)} \Big|_{\xi_n=i} dx' \\ &= \frac{5\pi^2}{12} \sum_{j=1}^{n-1} X_j Y_j h'(0) dx' + \frac{5i}{16} X_n Y_n h'(0) \pi \Omega_3 dx'. \end{aligned} \quad (4.39)$$

**case b)**  $r = 0$ ,  $l = -3$ ,  $k = j = |\alpha| = 0$ .

By (3.10), we get

$$\begin{aligned} \Phi_4 &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \times \partial_{\xi_n} \sigma_{-3}(D_T^* D_T)^{-1}](x_0) d\xi_n \sigma(\xi') dx' \\ &= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \times \sigma_{-3}(D_T^* D_T)^{-1}](x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (4.40)$$

By Lemma 4.4, we have

$$\begin{aligned} \sigma_{-3}(D_T^* D_T)^{-1}(x_0)|_{|\xi'|=1} &= -\frac{i}{(1 + \xi_n^2)^2} \left( -\frac{1}{2} h'(0) \sum_{k < n} \xi_n c(e_k) c(e_n) + \frac{5}{2} h'(0) \xi_n \right) - \frac{2ih'(0)\xi_n}{(1 + \xi_n^2)^3} \\ &\quad - ((u - v)\sqrt{-1}c(\xi) + \sqrt{-1}c(\xi)(u + v))|\xi|^{-4}. \end{aligned} \quad (4.41)$$

$$\begin{aligned} \partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1})(x_0)|_{|\xi'|=1} &= -\frac{i}{2(\xi_n - i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{1}{2(\xi_n - i)^2} X_n Y_n \\ &+ \frac{1}{2(\xi_n - i)^2} \sum_{j=1}^{n-1} X_j Y_n \xi_j + \frac{1}{2(\xi_n - i)^2} \sum_{l=1}^{n-1} X_n Y_l \xi_l. \end{aligned} \quad (4.42)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , and  $\sum_{i \neq s \neq t} \text{Tr}[A_{ist} c(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) c(\xi')] = 0$ , so we omit some items that have no contribution for computing **case b**). Then, we have

$$\begin{aligned} &\text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \times \sigma_{-3}(D_T^* D_T)^{-1}](x_0) \\ &= -\frac{h'(0)(5\xi_n^2 - 5 + 4\xi_n)}{(\xi_n - i)^5(\xi_n + i)^3} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l + \frac{h'(0)i(5\xi_n^3 - \xi_n)}{(\xi_n - i)^5(\xi_n + i)^3} X_n Y_n. \end{aligned} \quad (4.43)$$

Therefore, we get

$$\begin{aligned} \Phi_4 &= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left( -\frac{h'(0)(5\xi_n^2 - 5 + 4\xi_n)}{(\xi_n - i)^5(\xi_n + i)^3} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l + \frac{h'(0)i(5\xi_n^3 - \xi_n)}{(\xi_n - i)^5(\xi_n + i)^3} X_n Y_n \right) d\xi_n \sigma(\xi') dx' \\ &= -i \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \int_{\Gamma^+} \frac{5\xi_n^2 - 5 + 4\xi_n}{(\xi_n - i)^5(\xi_n + i)^2} \xi_j \xi_l d\xi_n dx' + i X_n Y_n h'(0) \Omega_3 \int_{\Gamma^+} \frac{5\xi_n^3 - \xi_n}{(\xi_n - i)^5(\xi_n + i)^2} d\xi_n dx' \\ &= -i \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \frac{2\pi i}{4!} \left[ \frac{5\xi_n^2 - 5 + 4\xi_n}{(\xi_n + i)^2} \right]^{(4)} \Big|_{\xi_n=i} dx' + i X_n Y_n h'(0) \Omega_3 \frac{2\pi i}{4!} \left[ \frac{5\xi_n^3 - \xi_n}{(\xi_n + i)^2} \right]^{(4)} \Big|_{\xi_n=i} dx' \\ &= \frac{(1-5i)\pi^2}{12} \sum_{j=1}^{n-1} X_j Y_j h'(0) dx' + \frac{11i}{16} X_n Y_n h'(0) \pi \Omega_3 dx'. \end{aligned} \quad (4.44)$$

**case c**)  $r = -1$ ,  $\ell = -2$ ,  $k = j = |\alpha| = 0$ .

By (3.10), we get

$$\Phi_5 = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\pi_{\xi_n}^+ \sigma_{-1}(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1}) \times \partial_{\xi_n} \sigma_{-2}(D_T^* D_T)^{-1}](x_0) d\xi_n \sigma(\xi') dx'. \quad (4.45)$$

By Lemma 4.4, we have

$$\partial_{\xi_n} \sigma_{-2}(D_T^* D_T)^{-1}(x_0)|_{|\xi'|=1} = -\frac{2\xi_n}{(\xi_n^2 + 1)^2}. \quad (4.46)$$

Since

$$\begin{aligned} \sigma_{-1}(\tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1})(x_0)|_{|\xi'|=1} &= \sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-3}((D_T^* D_T)^{-1}) + \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-2}((D_T^* D_T)^{-1}) \\ &+ \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] D_{x_j} [\sigma_{-2}((D_T^* D_T)^{-1})]. \end{aligned} \quad (4.47)$$

Explicit representation the first item of (4.47),

$$\begin{aligned} &\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-3}((D_T^* D_T)^{-1})(x_0)|_{|\xi'|=1} \\ &= -\sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \times \left( -\sqrt{-1} |\xi|^{-4} \xi_k (\Gamma^k - 2\delta^k) - \sqrt{-1} |\xi|^{-6} 2\xi^j \xi_\alpha \xi_\beta \partial_j g^{\alpha\beta} \right. \\ &\quad \left. - ((u-v)\sqrt{-1}c(\xi) + \sqrt{-1}c(\xi)(u+v)) |\xi|^{-4} \right) \\ &= -\sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \times \left( -\frac{i}{(1+\xi_n^2)^2} \left( -\frac{1}{2} h'(0) \sum_{k < n} \xi_n c(e_k) c(e_n) + \frac{5}{2} h'(0) \xi_n \right) - \frac{2ih'(0)\xi_n}{(1+\xi_n^2)^3} \right. \\ &\quad \left. - ((u-v)\sqrt{-1}c(\xi) + \sqrt{-1}c(\xi)(u+v)) |\xi|^{-4} \right). \end{aligned} \quad (4.48)$$

Explicit representation the second item of (4.47),

$$\begin{aligned}
& \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-2}((D_T^* D_T)^{-1})(x_0)|_{|\xi'|=1} \\
&= \left( \sqrt{-1} \sum_{j,l=1}^n X_j \frac{\partial Y_l}{\partial x_j} \sqrt{-1} \xi_l + \sqrt{-1} \sum_j A(Y) X_j \xi_j + \sqrt{-1} \sum_l A(Y) Y_l \xi_l \right. \\
&\quad \left. + \sum_j \bar{T}(X, \cdot) Y_j \sqrt{-1} \xi_j + \sum_j \bar{T}(Y, \cdot) X_j \sqrt{-1} \xi_j \right) \times |\xi|^{-2}.
\end{aligned} \tag{4.49}$$

Explicit representation the third item of (4.47),

$$\begin{aligned}
& \sum_{j=1}^n \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] D_x^{\alpha} [\sigma_{-2}((D_T^* D_T)^{-1})](x_0)|_{|\xi'|=1} \\
&= \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] (-\sqrt{-1}) \partial_{x_j} [\sigma_{-2}((D_T^* D_T)^{-1})] \\
&= \sum_{j=1}^n \partial_{\xi_j} \left[ - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \right] (-\sqrt{-1}) \partial_{x_j} [|\xi|^{-2}] \\
&= \sum_{j=1}^n \sum_{l=1}^n \sqrt{-1} (x_j Y_l + x_l Y_j) \xi_l \partial_{x_j} (|\xi|^{-2}).
\end{aligned} \tag{4.50}$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , and  $\sum_{i \neq s \neq t} Tr[A_{ist} c(\tilde{e}_i) c(\tilde{e}_s) c(\tilde{e}_t) c(\xi')] = 0$ , so we omit some items that have no contribution for computing **case c**). An easy calculation gives

$$\begin{aligned}
& Tr[\pi_{\xi_n}^+ \sigma_{-1}(\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-3}((D_T^* D_T)^{-1})) \times \partial_{\xi_n} \sigma_{-2}(D^{-2})](x_0)|_{|\xi'|=1} \\
&= \frac{h'(0)(2\xi_n^2 - \xi_n - 2\xi_n i)}{(\xi_n - i)^4 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \sum_{k < n} \xi_k c(e_k) c(e_n) + \frac{h'(0)(17\xi_n i - \xi_n^2 + 4\xi_n^3 i)}{(\xi_n - i)^5 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l,
\end{aligned} \tag{4.51}$$

and

$$Tr(\bar{T}(X, \cdot, \cdot)) = Tr\left(\frac{3}{2} \sum_{1 \leq i < j \leq n} T(X, e_i, e_j) c(e_i) c(e_j) - \frac{1}{2} c(V) c(X) - \frac{1}{2} (V, X)\right) = 0. \tag{4.52}$$

Also, straightforward computations yield

$$\begin{aligned}
& Tr[\pi_{\xi_n}^+ \sigma_{-1}(\sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-2}((D_T^* D_T)^{-1})) \times \partial_{\xi_n} \sigma_{-2}(D^{-2})](x_0)|_{|\xi'|=1} \\
&= X_n \frac{\partial Y_n}{\partial x_n} \frac{-2\xi_n}{(\xi_n^2 + 1)^2},
\end{aligned} \tag{4.53}$$

and

$$\begin{aligned}
& Tr[\pi_{\xi_n}^+ \sigma_{-1} \left( \sum_{j=1}^n \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] D_x^{\alpha} [\sigma_{-2}((D_T^* D_T)^{-1})] \right) \times \partial_{\xi_n} \sigma_{-2}(D^{-2})](x_0)|_{|\xi'|=1} \\
&= 2i X_n Y_n h'(0) \xi_n \times \frac{-2\xi_n}{(\xi_n^2 + 1)^2}.
\end{aligned} \tag{4.54}$$

Substituting (4.51), (4.53) and (4.54) into (4.45) yields

$$\begin{aligned}
\Phi_5 &= \left( \frac{5i - 13}{6} \sum_{j=1}^{n-1} X_j Y_j + \frac{3 - 96i}{8} X_n Y_n \right) h'(0) \pi^2 dx' \\
&\quad - X_n \frac{\partial Y_n}{\partial x_n} \frac{\pi}{2} \Omega_3 dx'
\end{aligned} \tag{4.55}$$

Let  $X = X^T + X_n \partial_n$ ,  $Y = Y^T + Y_n \partial_n$ , then we have  $\sum_{j=1}^{n-1} X_j Y_j = g(X^T, Y^T)$ . Now  $\Phi$  is the sum of the cases (a), (b) and (c). Therefore, we get

$$\Phi = \sum_{i=1}^5 \Phi_i = \frac{15 - 362i}{32} X_n Y_n h'(0) \pi \Omega_3 dx' + \frac{(10i - 27)\pi^2}{24} g(X^T, Y^T) h'(0) dx' - X_n \frac{\partial Y_n}{\partial x_n} \frac{\pi}{2} \Omega_3 dx' \quad (4.56)$$

Then we obtain following theorem

**Theorem 4.7.** *Let  $M$  be a 4-dimensional compact manifold without boundary and  $\tilde{\nabla}$  be an orthogonal connection with torsion. Then we get the volumes associated to  $\tilde{\nabla}_X \tilde{\nabla}_Y (D_F^* D_F)^{-1}$  and  $D_T^* D_T$  on compact manifolds with boundary*

$$\begin{aligned} & \widetilde{\text{Wres}}[\pi^+(\tilde{\nabla}_X \tilde{\nabla}_Y (D_F^* D_F)^{-1}) \circ \pi^+(D_T^* D_T)] \\ &= \frac{4\pi^2}{3} \int_M \left( Ric(X, Y) - \frac{1}{2} sg(X, Y) \right) vol_g \\ &+ \int_M \left( -\frac{1}{2} R^g - 3 \text{div}^g(X) + 3 \|T\|^2 + 9 \|X\|^2 \right) g(X, Y) vol_g \\ &+ \int_{\partial M} \left[ \left( \frac{15 - 362i}{32} X_n Y_n \pi h'(0) - X_n \frac{\partial Y_n}{\partial x_n} \frac{\pi}{2} \right) \Omega_3 + \frac{(10i - 27)\pi^2}{24} g(X^T, Y^T) h'(0) \right] vol_{\partial M}, \end{aligned} \quad (4.57)$$

where  $R_g$  denotes the curvature tensor and  $s$  is the scalar curvature.

## 5. Residue for Dirac operators with torsion $\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}$ and $(D_T^* D_T D_T^*)^{-1}$

In this section, we compute the 4-dimension volume for Dirac operators with torsion  $\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}$  and  $(D_T^* D_T D_T^*)^{-1}$ . Since  $[\sigma_{-4}(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1} \circ (D_T^* D_T D_T^*)^{-1})]_M$  has the same expression as  $[\sigma_{-4}(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1} \circ (D_T^* D_T D_T^*)^{-1})]_M$  in the case of manifolds without boundary, so locally we can use Theorem 2.4 to compute the first term.

**Theorem 5.1.** *Let  $M$  be a four dimensional compact manifold without boundary and  $\tilde{\nabla}$  be an orthogonal connection with torsion. Then we get the volumes associated to  $\tilde{\nabla}_X \tilde{\nabla}_Y (D_F^* D_F)^{-1}$  and  $D_T^* D_T$  on compact manifolds without boundary*

$$\begin{aligned} & \text{Wres}[\sigma_{-4}(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1} \circ (D_T^* D_T D_T^*)^{-1})] \\ &= \frac{4\pi^2}{3} \int_M \left( Ric(X, Y) - \frac{1}{2} sg(X, Y) \right) vol_g \\ &+ \int_M \left( -\frac{1}{2} R^g - 3 \text{div}^g(X) + 3 \|T\|^2 + 9 \|X\|^2 \right) g(X, Y) vol_g, \end{aligned} \quad (5.1)$$

where  $R_g$  denotes the curvature tensor and  $s$  is the scalar curvature.

From lemma 4.2 and lemma 4.3, we have

**Lemma 5.2.** *The following identities hold:*

$$\sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) = -\sqrt{-1} \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l c(\xi) |\xi|^{-2}; \quad (5.2)$$

$$\begin{aligned} \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) &= \sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-2}(D_T^{-1}) + \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-1}(D_T^{-1}) \\ &+ \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] D_{x_j} [\sigma_{-1}(D_T^{-1})]. \end{aligned} \quad (5.3)$$



Write

$$D_x^\alpha = (-i)^{|\alpha|} \partial_x^\alpha; \quad \sigma(D_t^3) = p_3 + p_2 + p_1 + p_0; \quad (\sigma(D_t)^{-3}) = \sum_{j=3}^{\infty} q_{-j}. \quad (5.4)$$

By the composition formula of pseudodifferential operators, we have

$$\begin{aligned} 1 &= \sigma(D^3 \circ D^{-3}) = \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma(D) D_x^{\alpha} [\sigma(D^{-3})]] \\ &= (p_3 + p_2 + p_1 + p_0)(q_{-3} + q_{-4} + q_{-5} + \cdots) \\ &\quad + \sum_j (\partial_{\xi_j} p_3 + \partial_{\xi_j} p_2 + \partial_{\xi_j} p_1 + \partial_{\xi_j} p_0)(D_{x_j} q_{-3} + D_{x_j} q_{-4} + D_{x_j} q_{-5} + \cdots) \\ &= p_3 q_{-3} + (p_3 q_{-4} + p_2 q_{-3} + \sum_j \partial_{\xi_j} p_3 D_{x_j} q_{-3}) + \cdots, \end{aligned} \quad (5.5)$$

so

$$q_{-3} = p_3^{-1}; \quad q_{-4} = -p_3^{-1} [p_2 p_3^{-1} + \sum_j \partial_{\xi_j} p_3 D_{x_j} (p_3^{-1})]. \quad (5.6)$$

Then it is easy to check that

**Lemma 5.3.** *The following identities hold:*

$$\sigma_{-2}((D_T^* D_T)^{-1}) = |\xi|^2; \quad (5.7)$$

$$\begin{aligned} \sigma_{-3}((D_T^* D_T)^{-1}) &= -\sqrt{-1} |\xi|^{-4} \xi_k (\Gamma^k - 2\delta^k) - \sqrt{-1} |\xi|^{-6} 2\xi^j \xi_{\alpha} \xi_{\beta} \partial_j g^{\alpha\beta} \\ &\quad - ((u-v)\sqrt{-1}c(\xi) + \sqrt{-1}c(\xi)(u+v)) |\xi|^{-4} \end{aligned} \quad (5.8)$$

$$\sigma_{-3}((D_T^* D_T D_T^*)^{-1}) = \sqrt{-1} c(\xi) |\xi|^{-4}; \quad (5.9)$$

$$\begin{aligned} \sigma_{-4}((D_T^* D_T D_T^*)^{-1}) &= \frac{c(\xi) \sigma_2((D_T^* D_T D_T^*)^{-1}) c(\xi)}{|\xi|^8} + \frac{\sqrt{-1} c(\xi)}{|\xi|^8} \left( |\xi|^4 c(dx_n) \partial_{x_n} c(\xi') \right. \\ &\quad \left. - 2h'(0) c(dx_n) c(\xi) + 2\xi_n c(\xi) \partial_{x_n} c(\xi') + 4\xi_n h'(0) \right), \end{aligned} \quad (5.10)$$

where

$$\sigma_2((D_T^* D_T D_T^*)^{-1}) = c(\xi) (4\sigma^k - 2\Gamma^k) \xi_k - \frac{1}{4} |\xi|^2 \sum_{s,t} \omega_{s,t}(\tilde{e}_l) c(e_l) c(\tilde{e}_s) c(\tilde{e}_t) + (3u-v) |\xi|^2. \quad (5.11)$$

Now we need to compute  $\int_{\partial M} \tilde{\Phi}$ . When  $n = 4$ , then  $\text{tr}_{S(TM)}[\text{id}] = \dim(\wedge^*(\mathbb{R}^2)) = 4$ , the sum is taken over  $r+l-k-j-|\alpha| = -3$ ,  $r \leq 0$ ,  $l \leq -2$ , then we have the following five cases:

**case a) I)**  $r = 1$ ,  $l = -2$ ,  $k = j = 0$ ,  $|\alpha| = 1$ .

By (3.12), we get

$$\tilde{\Phi}_1 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{Tr}[\partial_{\xi'}^{\alpha} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \partial_{x'}^{\alpha} \partial_{\xi_n} \sigma_{-3}((D_T^* D_T D_T^*)^{-1})](x_0) d\xi_n \sigma(\xi') dx'. \quad (5.12)$$

By Lemma 2.2 in [16], for  $i < n$ , then

$$\partial_{x_i} \sigma_{-3}((D_T^* D_T D_T^*)^{-1})(x_0) = \partial_{x_i} (\sqrt{-1} c(\xi) |\xi|^{-4})(x_0) = \sqrt{-1} \frac{\partial_{x_i} c(\xi)}{|\xi|^4} (x_0) + \sqrt{-1} \frac{c(\xi) \partial_{x_i} (|\xi|^4)}{|\xi|^8} (x_0) = 0, \quad (5.13)$$

so  $\tilde{\Phi}_1 = 0$ .

**case a) II)**  $r = 1, l = -3, k = |\alpha| = 0, j = 1$ .

By (3.12), we get

$$\tilde{\Phi}_2 = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \partial_{\xi_n}^2 \sigma_{-3}((D_T^* D_T D_T^*)^{-1})](x_0) d\xi_n \sigma(\xi') dx'. \quad (5.14)$$

By Lemma 5.3, we have

$$\partial_{\xi_n}^2 \sigma_{-3}((D_T^* D_T D_T^*)^{-1})(x_0) = \partial_{\xi_n}^2 (c(\xi) |\xi|^{-4})(x_0) = \sqrt{-1} \frac{(20\xi_n^2 - 4)c(\xi') + 12(\xi^3 - \xi)c(dx_n)}{(1 + \xi_n^2)^4}, \quad (5.15)$$

and

$$\begin{aligned} \partial_{x_n} \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1})(x_0) &= \partial_{x_n} (-\sqrt{-1} \Sigma_{j,l=1}^n X_j Y_l \xi_j \xi_l c(\xi) |\xi|^{-2}) \\ &= \Sigma_{j,l=1}^n X_j Y_l \xi_j \xi_l \left[ \frac{\partial_{x_n} c(\xi')}{1 + \xi_n^2} + \frac{c(\xi) h'(0) |\xi'|^2}{(1 + \xi_n^2)^2} \right]. \end{aligned} \quad (5.16)$$

Then, we have

$$\begin{aligned} \pi_{\xi_n}^+ \partial_{x_n} \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1})(x_0) &= \partial_{x_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1})(x_0) \\ &= \sqrt{-1} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) |\xi'|^2 \left[ \frac{ic(\xi')}{4(\xi_n - i)} + \frac{c(\xi') + ic(dx_n)}{4(\xi_n - i)^2} \right] \\ &\quad - \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \frac{\partial_{x_n} c(\xi')}{2(\xi_n - i)} \\ &\quad - \sqrt{-1} X_n Y_n \left\{ \frac{\partial_{x_n} c(\xi')}{2(\xi_n - i)} + h'(0) |\xi'| \left[ -\frac{2ic(\xi') - 3c(dx_n)}{4(\xi_n - i)} \right. \right. \\ &\quad \left. \left. + \frac{[c(\xi') + ic(dx_n)][i(\xi_n - i) + 1]}{4(\xi_n - i)^2} \right] \right\} \\ &\quad - \Sigma_{j=1}^{n-1} X_j Y_n \xi_j \left[ \sqrt{-1} \frac{\partial_{x_n} c(\xi')}{2(\xi_n - i)} - \frac{\sqrt{-1} h'(0) |\xi'| [c(\xi') + 2ic(dx_n)]}{4(\xi_n - i)} \right. \\ &\quad \left. - \frac{[ic(\xi') - c(dx_n)][i(\xi_n - i) + 1]}{(\xi_n - i)^2} \right] \\ &\quad - \Sigma_{l=1}^{n-1} X_n Y_l \xi_l \left[ \sqrt{-1} \frac{\partial_{x_n} c(\xi')}{2(\xi_n - i)} - \frac{\sqrt{-1} h'(0) |\xi'| [c(\xi') + 2ic(dx_n)]}{4(\xi_n - i)} \right. \\ &\quad \left. - \frac{[ic(\xi') - c(dx_n)][i(\xi_n - i) + 1]}{(\xi_n - i)^2} \right]. \end{aligned} \quad (5.17)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case a) II)**. Then there is the following formula

$$\begin{aligned} &\text{Tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \partial_{\xi_n}^2 \sigma_{-3}((D_T^* D_T D_T^*)^{-1})](x_0) \\ &= \Sigma_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \left[ 8i \frac{5\xi_n^2 - 1}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \\ &+ X_n Y_n h'(0) \left[ \frac{(4i - 4)(\xi^2 - 1) + 48(\xi_n^3 - \xi_n)}{(\xi_n - i)^5 (\xi_n + i)^4} - \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \\ &+ 8\Sigma_{j=1}^{n-1} X_j Y_n \xi_j \left[ \frac{(6 - 3ih'(0))(\xi_n^3 - \xi_n) - 2i(5\xi_n^2 - 1)}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{2(5\xi_n^2 - 1) + 6i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \\ &+ 8\Sigma_{l=1}^{n-1} X_n Y_l \xi_l \left[ \frac{(6 - 3ih'(0))(\xi_n^3 - \xi_n) - 2i(5\xi_n^2 - 1)}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{2(5\xi_n^2 - 1) + 6i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right]. \end{aligned} \quad (5.18)$$

Therefore, we get

$$\begin{aligned}
\tilde{\Phi}_2 &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left\{ \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \left[ 8i \frac{5\xi_n^2 - 1}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \right. \\
&\quad \left. + X_n Y_n h'(0) \left[ \frac{(4i - 4)(\xi^2 - 1) + 48(\xi_n^3 - \xi_n)}{(\xi_n - i)^5 (\xi_n + i)^4} - \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \right\} d\xi_n \sigma(\xi') dx' \\
&= \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \int_{\Gamma^+} \left[ 8i \frac{5\xi_n^2 - 1}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \xi_j \xi_l d\xi_n dx' \\
&\quad + X_n Y_n h'(0) \Omega_3 \int_{\Gamma^+} \left[ \frac{(4i - 4)(\xi^2 - 1) + 48(\xi_n^3 - \xi_n)}{(\xi_n - i)^5 (\xi_n + i)^4} - \frac{4(5\xi_n^2 - 1) + 12i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] d\xi_n dx' \\
&= -\frac{592}{3} \pi^2 \sum_{j=1}^{n-1} X_j Y_j h'(0) dx' - \left( \frac{461}{4} + \frac{23}{4} i \right) X_n Y_n h'(0) \pi \Omega_3 dx', \tag{5.19}
\end{aligned}$$

where  $\Omega_3$  is the canonical volume of  $S^2$ .

**case a) III)**  $r = 1$ ,  $l = -3$ ,  $j = |\alpha| = 0$ ,  $k = 1$ .

By (3.12), we get

$$\begin{aligned}
\tilde{\Phi}_3 &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-3}((D_T^* D_T D_T^*)^{-1})](x_0) d\xi_n \sigma(\xi') dx' \\
&= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \partial_{x_n} \sigma_{-3}((D_T^* D_T D_T^*)^{-1})](x_0) d\xi_n \sigma(\xi') dx'. \tag{5.20}
\end{aligned}$$

By Lemma 5.3, we have

$$\partial_{x_n} \sigma_{-3}((D_T^* D_T D_T^*)^{-1})(x_0)|_{|\xi'|=1} = \frac{\sqrt{-1} \partial_{x_n} [c(\xi')]}{(1 + \xi_n^2)^4} - \frac{2\sqrt{-1} h'(0) c(\xi') |\xi'|_{g^{\partial M}}^2}{(1 + \xi_n^2)^6}. \tag{5.21}$$

By integrating formula we obtain

$$\begin{aligned}
\pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) &= -\frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)} X_n Y_n \\
&\quad - \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)} \sum_{j=1}^{n-1} X_j Y_n \xi_j - \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)} \sum_{l=1}^{n-1} X_n Y_l \xi_l. \tag{5.22}
\end{aligned}$$

Then, we have

$$\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) = -\frac{c(\xi') + ic(dx_n)}{(\xi_n - i)^3} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{c(\xi') + ic(dx_n)}{(\xi_n - i)^3} X_n Y_n. \tag{5.23}$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case a) III)**, then

$$\begin{aligned}
&\text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-3}((D_T^* D_T D_T^*)^{-1})](x_0) \\
&= -2 \frac{h'(0)}{(\xi_n - i)^5 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - 2 \frac{h'(0)}{(\xi_n - i)^5 (\xi_n + i)^2} X_n Y_n. \tag{5.24}
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
\tilde{\Phi}_3 &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left( -2 \frac{h'(0)}{(\xi_n - i)^5 (\xi_n + i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - 2 \frac{h'(0)}{(\xi_n - i)^5 (\xi_n + i)^2} X_n Y_n \right) d\xi_n \sigma(\xi') dx' \\
&= - \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \int_{\Gamma^+} \frac{1}{(\xi_n - i)^5 (\xi_n + i)^2} \xi_j \xi_l d\xi_n dx' - X_n Y_n h'(0) \Omega_3 \int_{\Gamma^+} \frac{1}{(\xi_n - i)^5 (\xi_n + i)^2} d\xi_n dx' \\
&= - \sum_{j,l=1}^{n-1} X_j Y_l h'(0) \Omega_3 \frac{2\pi i}{4!} \left[ \frac{1}{(\xi_n + i)^2} \right]^{(4)} \Big|_{\xi_n=i} dx' + 2 X_n Y_n h'(0) \Omega_3 \frac{2\pi i}{4!} \left[ \frac{1}{(\xi_n + i)^2} \right]^{(4)} \Big|_{\xi_n=i} dx' \\
&= \frac{5i\pi^2}{6} \sum_{j=1}^{n-1} X_j Y_j h'(0) dx' + \frac{5i}{8} X_n Y_n h'(0) \pi \Omega_3 dx'. \tag{5.25}
\end{aligned}$$

**case b)**  $r = 0$ ,  $l = -3$ ,  $k = j = |\alpha| = 0$ .

By (3.12), we get

$$\tilde{\Phi}_4 = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \partial_{\xi_n} \sigma_{-3}((D_T^* D_T D_T^*)^{-1})(x_0)] d\xi_n \sigma(\xi') dx'. \tag{5.26}$$

By Lemma 5.3, we obtain

$$\partial_{\xi_n} \sigma_{-3}((D_T^* D_T D_T^*)^{-1})(x_0)|_{|\xi'|=1} = \frac{ic(dx_n)}{(1 + \xi_n^2)^2} - \frac{4\sqrt{-1}\xi_n c(\xi)}{(1 + \xi_n^2)^3}. \tag{5.27}$$

By Lemma 5.2, we have

$$\begin{aligned}
\sigma_0(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) &= \sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-2}(D_T^{-1}) + \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-1}(D_T^{-1}) \\
&\quad + \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] D_{x_j} [\sigma_{-1}(D_T^{-1})]. \tag{5.28}
\end{aligned}$$

(1) Explicit representation the first item of (5.28)

$$\begin{aligned}
&\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-2}(D_T^{-1})(x_0)|_{|\xi'|=1} \\
&= - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \left[ \frac{c(\xi) \sigma_0(D_T) c(\xi)}{|\xi|^4} + \frac{c(\xi)}{|\xi|^6} \sum_j c(dx_j) (\partial_{x_j} [c(\xi)] |\xi|^2 - c(\xi) \partial_{x_j} (|\xi|^2)) \right], \tag{5.29}
\end{aligned}$$

By integrating formula, we obtain

$$\pi_{\xi_n}^+ \left[ \frac{c(\xi) \sigma_0(D_T)(x_0) c(\xi) + c(\xi) c(dx_n) \partial_{x_n} [c(\xi')](x_0)}{(1 + \xi_n^2)^2} \right] = -\frac{A_1}{4(\xi_n - i)} - \frac{A_2}{4(\xi_n - i)^2} + \frac{A_3}{4(\xi_n - i)^2}, \tag{5.30}$$

where

$$A_1 = ic(\xi') p_0 c(\xi') + ic(dx_n) \left( -\frac{3}{4} h'(0) c(dx_n) \right) c(dx_n) + ic(\xi') c(dx_n) \partial_{x_n} [c(\xi')], \tag{5.31}$$

$$A_2 = [c(\xi') + ic(dx_n)] p_0 [c(\xi') + ic(dx_n)] + c(\xi') c(dx_n) \partial_{x_n} c(\xi') - i \partial_{x_n} [c(\xi')], \tag{5.32}$$

$$\begin{aligned}
A_3 &= \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \left( (-2 - i\xi_n) c(\xi') (u + v) c(\xi') - ic(dx_n) (u + v) c(\xi') - ic(\xi') (u + v) c(dx_n) \right. \\
&\quad \left. - i\xi_n c(dx_n) (u + v) c(dx_n) \right) + X_n Y_n \left( -i\xi_n c(\xi') (u + v) c(\xi') - ic(dx_n) (u + v) c(\xi') \right. \\
&\quad \left. - ic(\xi') (u + v) c(dx_n) + c(dx_n) (u + v) c(dx_n) \right). \tag{5.33}
\end{aligned}$$

In the same way we get

$$\pi_{\xi_n}^+ \left[ \frac{c(\xi)c(dx_n)c(\xi)}{(1+\xi_n)^3} (x_0)|_{|\xi'|=1} \right] = \frac{1}{2} \left[ \frac{c(dx_n)}{4i(\xi_n-i)} + \frac{c(dx_n)-ic(\xi')}{8(\xi_n-i)^2} + \frac{3\xi_n-7i}{8(\xi_n-i)^3} [ic(\xi')-c(dx_n)] \right]. \quad (5.34)$$

Then adding these identities gives

$$\begin{aligned} & \text{Tr} \left( \pi_{\xi_n}^+ \left( \sum_{j=1}^{n-1} \sqrt{-1} T(X, \cdot, \cdot) Y_j \xi_j + \sum_{j=1}^{n-1} \sqrt{-1} T(Y, \cdot, \cdot) X_j \xi_j \right) \times \partial_{\xi_n} \sigma_{-3} ((D_T^* D_T D_T^*)^{-1}) \right) (x_0) \\ &= \sum_{j=1}^{n-1} Y_j \xi_j \frac{2i\xi_n}{(\xi_n-i)(1+\xi_n^2)^3} \text{trace} \left( T(X, \cdot, \cdot) c(dx_n) c(\xi') \right) \\ & \quad + \sum_{j=1}^{n-1} Y_j \xi_j \frac{3\xi_n^2-1}{2(\xi_n-i)(1+\xi_n^2)^3} \text{trace} \left( T(X, \cdot, \cdot) c(\xi') c(dx_n) \right) \\ & \quad + \sum_{j=1}^{n-1} X_j \xi_j \frac{2i\xi_n}{(\xi_n-i)(1+\xi_n^2)^3} \text{trace} \left( T(Y, \cdot, \cdot) c(dx_n) c(\xi') \right) \\ & \quad + \sum_{j=1}^{n-1} Y_j \xi_j \frac{3\xi_n^2-1}{(\xi_n-i)(1+\xi_n^2)^3} \text{trace} \left( T(Y, \cdot, \cdot) c(dx_n) c(\xi') \right) \\ &= \sum_{j=1}^{n-1} Y_j \xi_j \frac{2i\xi_n}{(\xi_n-i)(1+\xi_n^2)^3} \sum_{i=1}^{n-1} T(X, e_i, e_n) \xi_i - \sum_{j=1}^{n-1} Y_j \xi_j \frac{3\xi_n^2-1}{2(\xi_n-i)(1+\xi_n^2)^3} \sum_{i=1}^{n-1} T(X, e_i, e_n) \xi_i \\ & \quad + \sum_{j=1}^{n-1} X_j \xi_j \frac{2i\xi_n}{(\xi_n-i)(1+\xi_n^2)^3} \sum_{i=1}^{n-1} T(Y, e_i, e_n) \xi_i - \sum_{j=1}^{n-1} Y_j \xi_j \frac{3\xi_n^2-1}{(\xi_n-i)(1+\xi_n^2)^3} \sum_{i=1}^{n-1} T(Y, e_i, e_n) \xi_i \end{aligned} \quad (5.35)$$

Substituting (5.35) into (5.26) yields

$$\begin{aligned} & -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr} \left[ \pi_{\xi_n}^+ \left( \sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y) \frac{c(\xi)(u+v)c(\xi)}{(1+\xi_n)^3} \right) \times \partial_{\xi_n} \sigma_{-3} ((D_T^* D_T D_T^*)^{-1}) \right] (x_0) d\xi_n \sigma(\xi') dx' \\ &= \left( \sum_{j,l=1}^{n-1} X_j Y_l \frac{2\pi}{3} + \frac{3}{8} X_n Y_n \right) \sum_{i=1}^{n-1} A_{iin} \pi \Omega_3 dx' + \sum_{j=1}^{n-1} \left( X_j \text{Tr}(T(X, e_j, e_n)) Y_n + Y_j \text{Tr}(T(Y, e_j, e_n)) \right) \frac{4i\pi^2}{3} \Omega_3 dx' \\ &= \sum_{j,l=1}^{n-1} X_j Y_l \frac{2\pi^2}{3} \sum_{i=1}^{n-1} A_{iin} dx' + \frac{3}{8} X_n Y_n \sum_{i=1}^{n-1} A_{iin} \pi \Omega_3 dx'. \end{aligned} \quad (5.36)$$

(2) Explicit representation the second item of (5.28)

$$\begin{aligned} & \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-1}(D_T^{-1})(x_0)|_{|\xi'|=1} \\ &= \left( \sqrt{-1} \sum_{j,l=1}^n X_j \frac{\partial Y_l}{\partial x_j} \partial_{x_l} + \sqrt{-1} \sum_j A(Y) X_j \xi_j + \sqrt{-1} \sum_l A(Y) Y_l \xi_l \right. \\ & \quad \left. + \sum_j \sqrt{-1} T(X, \cdot) Y_j \xi_j + \sum_j \sqrt{-1} T(Y, \cdot) X_j \xi_j \right) \frac{\sqrt{-1} c(\xi)}{|\xi|^2}; \end{aligned} \quad (5.37)$$

By integrating formula we get

$$\begin{aligned}
& \pi_{\xi_n}^+ \left( \left( \sum_j^n \sqrt{-1}T(X, \cdot)Y_j\xi_j + \sum_j^n \sqrt{-1}T(Y, \cdot)X_j\xi_j \right) \frac{\sqrt{-1}c(\xi)}{|\xi|^2} \right) \\
&= \pi_{\xi_n}^+ \left( \left( \sum_j^{n-1} \sqrt{-1}T(X, \cdot)Y_j\xi_j + \sum_j^{n-1} \sqrt{-1}T(Y, \cdot)X_j\xi_j \right) \frac{\sqrt{-1}c(\xi)}{|\xi|^2} \right) \\
&\quad + \pi_{\xi_n}^+ \left( \left( \sqrt{-1}T(X, \cdot)Y_n\xi_n + \sqrt{-1}T(Y, \cdot)X_n\xi_n \right) \frac{\sqrt{-1}c(\xi)}{|\xi|^2} \right) \\
&= \left( \sum_j^{n-1} \sqrt{-1}T(X, \cdot)Y_j\xi_j + \sum_j^{n-1} \sqrt{-1}T(Y, \cdot)X_j\xi_j \right) \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)} \\
&\quad + \left( \sqrt{-1}T(X, \cdot)Y_n\xi_n + \sqrt{-1}T(Y, \cdot)X_n\xi_n \right) \frac{-c(\xi') - ic(dx_n)}{2(\xi_n - i)} \tag{5.38}
\end{aligned}$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1}\xi_{i_2}\cdots\xi_{i_{2d+1}}\sigma(\xi') = 0$ , and  $\sum_{i \neq s \neq t} Tr[A_{ist}c(\tilde{e}_i)c(\tilde{e}_s)c(\tilde{e}_t)c(\xi')] = 0$ , then

$$\text{Tr} \left( \pi_{\xi_n}^+ \left( \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y) \sigma_{-1}(D_T^{-1}) \right) \times \partial_{\xi_n} \sigma_{-3}((D_T^* D_T D_T^*)^{-1}) \right) (x_0) = 0. \tag{5.39}$$

(3) Explicit representation the third item of (5.28)

$$\begin{aligned}
\sum_{j=1}^n \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] D_x^{\alpha} [\sigma_{-1}(D_T^{-1})] (x_0)|_{|\xi'|=1} &= \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] (-\sqrt{-1}) \partial_{x_j} [\sigma_{-1}(D_T^{-1})] \\
&= \sum_{j=1}^n \partial_{\xi_j} \left[ - \sum_{l=1}^n X_j Y_l \xi_j \xi_l \right] (-\sqrt{-1}) \partial_{x_j} \left[ \frac{\sqrt{-1}c(\xi)}{|\xi|^2} \right] \\
&= \sum_{j=1}^n \sum_{l=1}^n \sqrt{-1} (x_j Y_l + x_l Y_j) \xi_l \partial_{x_j} \left( \frac{\sqrt{-1}c(\xi)}{|\xi|^2} \right). \tag{5.40}
\end{aligned}$$

By integrating formula we obtain

$$\begin{aligned}
& \pi_{\xi_n}^+ \left( \sum_{j=1}^n \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] D_x^{\alpha} [\sigma_{-1}(D_T^{-1})] \right) \\
&= \pi_{\xi_n}^+ \left( \sum_{l=1}^{n-1} \sqrt{-1} (x_n Y_l + x_l Y_n) \xi_l \partial_{x_n} \left( \frac{\sqrt{-1}c(\xi)}{|\xi|^2} \right) \right) + \pi_{\xi_n}^+ \left( \sqrt{-1} (x_n Y_n + x_n Y_n) \xi_n \partial_{x_n} \left( \frac{\sqrt{-1}c(\xi)}{|\xi|^2} \right) \right) \\
&= \sum_{l=1}^{n-1} (x_n Y_l + x_l Y_n) \xi_l \left( \frac{i \partial_{x_n} (c(\xi'))}{2(\xi_n - i)} + h'(0) \frac{(-2 - i \xi_n) c(\xi')}{4(\xi_n - i)^2} - h'(0) \frac{ic(dx_n)}{4(\xi_n - i)^2} \right) \\
&\quad + x_n Y_n \left( \frac{-\partial_{x_n} (c(\xi'))}{(\xi_n - i)} + h'(0) \frac{(-i) c(\xi')}{2(\xi_n - i)^2} - h'(0) \frac{-i \xi_n c(dx_n)}{2(\xi_n - i)^2} \right) \tag{5.41}
\end{aligned}$$

Substituting (5.41) into (5.26) yields

$$\begin{aligned}
& -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr} \left[ \pi_{\xi_n}^+ \left( \sum_{j=1}^n \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] D_x^{\alpha} [\sigma_{-1}(D_T^{-1})] \right) \right. \\
&\quad \times \left. \partial_{\xi_n} \sigma_{-3}((D_T^* D_T D_T^*)^{-1}) \right] (x_0) d\xi_n \sigma(\xi') dx' \\
&= \frac{7-15i}{8} X_n Y_n \pi h'(0) \Omega_3 dx'. \tag{5.42}
\end{aligned}$$

Summing up (1), (2) and (3) leads to the desired equality

$$\tilde{\Phi}_4 = \left( \frac{55\pi^2}{3} \sum_{j=1}^{n-1} X_j Y_j + \frac{4-15i}{8} X_n Y_n \pi \Omega_3 \right) h'(0) dx' + \left( \frac{2\pi^2}{3} \sum_{j,l=1}^{n-1} X_j Y_l + \frac{3}{8} X_n Y_n \pi \Omega_3 \right) \sum_{i=1}^{n-1} A_{iin} dx'. \quad (5.43)$$

**case c)**  $r = 1$ ,  $\ell = -4$ ,  $k = j = |\alpha| = 0$ .

By (3.12), we get

$$\begin{aligned} \tilde{\Phi}_5 &= - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \partial_{\xi_n} \sigma_{-4}(D_T^* D_T D_T^*)^{-1}](x_0) d\xi_n \sigma(\xi') dx' \\ &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \sigma_{-4}(D_T^* D_T D_T^*)^{-1}](x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (5.44)$$

By Lemma 5.3, we have

$$\begin{aligned} \sigma_{-4}(D_T^* D_T D_T^*)^{-1}(x_0)|_{|\xi'|=1} &= \frac{1}{(\xi_n^2 + 1)^4} \left[ \left( \frac{11}{2} \xi_n (1 + \xi_n^2) + 8i \xi_n \right) h'(0) c(\xi') \right. \\ &\quad + \left[ -2i + 6i \xi_n^2 - \frac{7}{4} (1 + \xi_n^2) + \frac{15}{4} \xi_n^2 (1 + \xi_n^2) \right] h'(0) c(dx_n) \\ &\quad - 3i \xi_n (1 + \xi_n^2) \partial_{x_n} c(\xi') + i(1 + \xi_n^2) c(\xi') c(dx_n) \partial_{x_n} c(\xi') \\ &\quad \left. + \frac{c(\xi)(3u-v)|\xi|^2 c(\xi)}{|\xi|^8} \right], \end{aligned} \quad (5.45)$$

and

$$\begin{aligned} \partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1})(x_0)|_{|\xi'|=1} &= \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2} X_n Y_n \\ &\quad + \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)^2} \sum_{j=1}^n X_j Y_n \xi_j + \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)^2} \sum_{l=1}^n X_n Y_l \xi_l. \end{aligned} \quad (5.46)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2a+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case c)**. Here

$$\begin{aligned} \text{tr}[c(\xi') c(\xi') c(dx_n) \partial_{x_n} c(\xi')] &= 0; \\ \text{tr}[c(dx_n) c(\xi') c(dx_n) \partial_{x_n} c(\xi')] &= -2h'(0). \end{aligned} \quad (5.47)$$

Also, straightforward computations yield

$$\begin{aligned} &\text{tr} \left[ \partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \frac{1}{(\xi_n^2 + 1)^4} \left( \left( \frac{11}{2} \xi_n (1 + \xi_n^2) + 8i \xi_n \right) h'(0) c(\xi') \right. \right. \\ &\quad + \left( -2i + 6i \xi_n^2 - \frac{7}{4} (1 + \xi_n^2) + \frac{15}{4} \xi_n^2 (1 + \xi_n^2) \right) h'(0) c(dx_n) \\ &\quad \left. \left. - 3i \xi_n (1 + \xi_n^2) \partial_{x_n} c(\xi') + i(1 + \xi_n^2) c(\xi') c(dx_n) \partial_{x_n} c(\xi') \right) \right] (x_0)|_{|\xi'|=1} \\ &= \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \frac{h'(0)(7 + 6i - (20 - 15i)\xi_n - (7 - 6i)\xi_n^2 + 15i\xi_n^3)}{(\xi_n - i)^5 (\xi_n + i)^4} \\ &\quad + X_n Y_n \frac{(3i - 11)\xi_n(1 - \xi_n^2) - 16i\xi_n + (13 + \frac{7}{2}i)(1 + \xi_n^2) - 16 - \frac{15}{2}\xi_n^2(1 + \xi_n^2)}{(\xi_n - i)^2 (\xi_n + i)^4}, \end{aligned} \quad (5.48)$$

and

$$\begin{aligned} & \text{tr}\left(\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \times \frac{c(\xi)(3u-v)|\xi|^2 c(\xi)}{|\xi|^8}\right)(x_0)|_{|\xi|=1} \\ &= \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \frac{-3i\pi}{8} \sum_{i=1}^n A_{iin} + X_n Y_n \frac{-3i\pi}{8} \sum_{i=1}^n A_{iin}. \end{aligned} \quad (5.49)$$

From (5.44),(5.48) and (5.49), we get

$$\tilde{\Phi}_5 = \left( \left( -\frac{35}{3} + \frac{50}{3}i \right) \sum_{j=1}^{n-1} X_j Y_j \pi^2 + \left( 5 - \frac{137}{32}i \right) X_n Y_n \pi \Omega_3 \right) h'(0) dx'. \quad (5.50)$$

Let  $X = X^T + X_n \partial_n$ ,  $Y = Y^T + Y_n \partial_n$ , then we have  $\sum_{j=1}^{n-1} X_j Y_j = g(X^T, Y^T)$ . Now  $\Phi$  is the sum of the cases (a), (b) and (c). Combining with the five cases, this yields

$$\begin{aligned} \tilde{\Phi} = \sum_{i=1}^5 \tilde{\Phi}_i &= \left[ \left( -\frac{2801}{12} - \frac{33i}{32} \right) X_n Y_n \pi \Omega_3 + \left( -\frac{572}{3} + \frac{35i}{2} \right) \pi^2 g(X^T, Y^T) \right] h'(0) dx' \\ &+ \left( \frac{3}{8} - \frac{3i}{8} \right) X_n Y_n \pi \Omega_3 + \frac{4+3i}{6} \pi^2 g(X^T, Y^T) \sum_{i=1}^n A_{iin} dx'. \end{aligned} \quad (5.51)$$

So, we are reduced to prove the following.

**Theorem 5.4.** *Let  $M$  be a 4-dimensional compact manifold with boundary  $\partial M$  and  $\tilde{\nabla}$  be an orthogonal connection with torsion. Then we get the noncommutative residue associated to  $\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}$  and  $(D_T^* D_T D_T^*)^{-1}$  on compact manifolds with boundary*

$$\begin{aligned} & \widetilde{\text{Wres}}[\pi^+(\tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1}) \circ \pi^+((D_T^* D_T D_T^*)^{-1})] \\ &= \frac{4\pi^2}{3} \int_M \left( \text{Ric}(X, Y) - \frac{1}{2} s g(X, Y) \right) \text{vol}_g \\ &+ \int_M \left( -\frac{1}{2} R^g - 3 \text{div}^g(X) + 3 \|T\|^2 + 9 \|X\|^2 \right) g(X, Y) \text{vol}_g \\ &+ \int_{\partial M} \left( \left( \frac{-2801}{24} - \frac{33i}{32} \right) X_n Y_n \pi \Omega_3 + \left( \frac{35i}{2} - \frac{572}{3} \right) \pi^2 g(X^T, Y^T) \right) h'(0) \\ &+ \left( \left( \frac{3}{8} - \frac{3i}{8} \right) X_n Y_n \pi \Omega_3 + \frac{4+3i}{6} \pi^2 g(X^T, Y^T) \right) \sum_{i=1}^n A_{iin} \text{vol}_{\partial M}, \end{aligned} \quad (5.52)$$

where  $R_g$  denotes the curvature tensor and  $s$  is the scalar curvature.

## Acknowledgements

The first author was supported by NSFC. 11501414. The second author was supported by NSFC. 11771070. The authors also thank the referee for his (or her) careful reading and helpful comments.

## References

- [1] Dabrowski L, Sitarz A, Zalecki P. Spectral Metric and Einstein Functionals. arXiv:2206.02587, (2022).
- [2] F. Pfäffle and C. A. Stephan.: Chiral Asymmetry and the Spectral Action. Commun. Math. Phys. 321, 283-310 (2013).
- [3] P. B. Gilkey.: Invariance theory, the Heat equation, and the Atiyah-Singer Index theorem. Inc., USA, (1984).
- [4] H. Figueroa, J. Gracia-Bondía, and J. Várilly.: Elements of Noncommutative Geometry. Birkhäuser Boston, (2001).
- [5] T. Ackermann.: A note on the Wodzicki residue. J. Geom. Phys. 20, 404-406, (1996).



- [6] M. Wodzicki.: Local invariants of spectral asymmetry. *Invent. Math.* 75(1), 143-178, (1984).
- [7] M. Wodzicki.: Non-commutative residue I, *Lecture Notes in Math.*, Springer, New York, Vol. 1289 320-399, (1987).
- [8] V. W. Guillemin.: A new proof of Weyl's formula on the asymptotic distribution of eigenvalues. *Adv. Math.* 55, no. 2, 131-160, (1985).
- [9] A. Connes.: Quantized calculus and applications. XIth International Congress of Mathematical Physics(Paris,1994), Internat Press, Cambridge, MA, 15-36, (1995).
- [10] A. Connes.: The action functional in Noncommutative geometry. *Comm. Math. Phys.* 117, 673-683, (1988).
- [11] A. Connes, J. Lott.: Particle models and Non-commutative geometry. *Nucl. Phys. B Proc.Supp* 18B, 29-47, (1990).
- [12] D. Kastler.: The Dirac Operator and Gravitation. *Comm. Math. Phys.* 166, 633-643, (1995).
- [13] W. Kalau and M. Walze.: Gravity, Noncommutative geometry and the Wodzicki residue. *J. Geom. Physics.* 16, 327-344,(1995).
- [14] B. V. Fedosov, F. Gölse, E. Leichtnam, E. Schrohe.: The noncommutative residue for manifolds with boundary. *J. Funct. Anal.* 142, 1-31, (1996).
- [15] Y. Wang.: Differential forms and the Wodzicki residue for Manifolds with Boundary. *J. Geom. Physics.* 56, 731-753, (2006).
- [16] Y. Wang.: Gravity and the Noncommutative Residue for Manifolds with Boundary. *Letters in Mathematical Physics.* 80, 37-56, (2007).
- [17] Y. Wang.: Lower-Dimensional Volumes and Kastler-kalau-Walze Type Theorem for Manifolds with Boundary . *Commun. Theor. Phys.* Vol 54, 38-42, (2010).
- [18] J. M. Bismut.: A local index theorem for non Kähler manifolds. *Math. Ann.* 284, 681-699 (1989).
- [19] T. Ackermann and J. Tolksdorf.: A generalized Lichnerowicz formula, the Wodzicki residue and gravity. *J. Geom. Physics.* 19, 143-150,(1996).
- [20] F. Pfäffle and C. A. Stephan.: On gravity, torsion and the spectral action principle. *J. Funct. Anal.* 262, 1529-1565,(2012).
- [21] J. Wang, Y. Wang, C.L. Yang, Dirac operators with torsion and the noncommutative residue for manifolds with boundary, *J. Geom. Phys.* 81,92-111 (2014).
- [22] W. Greub, S. Halperin and R.Vanstone.: *Connections, Curvature and Cohomology, Vo.1* (Academic press, New York) (1976).
- [23] J. Wang and Y. Wang.: Nonminimal operators and non-commutative residue, *J. Math. Phys.* 53, 072503 (2012).
- [24] Y. Yu.: *The Index Theorem and The Heat Equation Method*, Nankai Tracts in Mathematics-Vol.2, World Scientific Publishing, (2001).
- [25] G. Grubb, E. Schrohe.: Trace expansions and the noncommutative residue for manifolds with boundary. *J. Reine Angew. Math.* 536, 167-207, (2001).
- [26] S. Alexandrov and D. Vassilevich.: Heat kernel for nonminimal operators on a Kähler manifold. *J. Math. Phys.* 37, 5715-5718, (1996).