THE MILLIONS OF OPTICAL - RADIO/X-RAY ASSOCIATIONS (MORX) CATALOGUE, v2

ERIC WIM FLESCH
PO Box 15, Dannevirke 4942, New Zealand; eric@flesch.org
Version January 23, 2024

ABSTRACT

Announcing the release v2 of the MORX (Millions of Optical-Radio/X-ray Associations) catalogue which presents probable (40%-100% likelihood) radio/X-ray associations, including double radio lobes, to optical objects over the whole sky. Detections from all the largest radio/X-ray surveys to June 2023 are evaluated, those surveys being VLASS, LoTSS, RACS, FIRST, NVSS, and SUMSS radio surveys, and Chandra, XMM-Newton, Swift, and ROSAT X-ray surveys. The totals are 3115575 optical objects of all classifications (or unclassified) so associated. The MORX v2 catalogue is available on multiple sites.

Subject headings: catalogs — x-rays: general — radio continuum: general

1. INTRODUCTION

This is an update to the 2016 MORX catalogue of radio/X-ray associations onto optical objects (MORXv1: Flesch 2016), using all the largest radio and X-ray source surveys available to 30 June 2023. This update, version 2, has three times the number of optical objects so associated, and has a simpler format, just giving cumulative likelihoods and the radio/X-ray detection identifiers without details of their optical solutions, for ease of use. Figure 1 shows MORXv2 coverage over the sky, and explains the features seen thereon.

MORX v2 can be downloaded from CDS¹ or from the MORX home page² which also provides a FITS file. CDS and NASA HEASARC³ provide query pages. The MORX ReadMe gives essential information about the data, and the file "MORX-references.txt" gives the legend of citations for the data. Many issues involved in the production of MORX are explained in the paper for MORXv1 (Flesch 2016), and the reader can consult there for in-depth topics. Updates and changes are as presented below, starting with a listing of the input radio/X-ray catalogues for MORX v2.

2. INPUT RADIO/X-RAY CATALOGUES

In 2016 I published the Million Optical - Radio/X-ray Associations Catalogue (MORXv1: Flesch 2016) as a quick guide to all such associations able to be calculated from the largest optical, radio, and X-ray surveys over the whole sky. That paper gives the methods used to produce this catalogue, including the optical field solutions and calculation of association likelihoods, and probabilistic classification of unclassified objects; the reader is referred there for those topics. In the seven years since,

eric@flesch.org

1 https://cdsarc.cds.unistra.fr/viz-bin/cat/V/158

² https://quasars.org/morx.htm

3 https://heasarc.gsfc.nasa.gov/W3Browse/all/morx.html

there have of course been new editions of these surveys and new such surveys published; following is the list of input radio/X-ray surveys used for MORX v2.

For radio sources there are the high-resolution VLASS, LoTSS, and FIRST surveys, the medium-resolution RACS survey, and the low-resolution NVSS and SUMSS surveys:

- The Very Large Array Sky Survey (VLASS: Gordon et al. 2020) Quick Look catalogue⁴ is their first complete publication of VLASS radio sources. VLASS covers the whole sky north of declination -40° in 3GHz to a depth⁵ of 1mJy and astrometric accuracy of ≈0.5 arcsec in most places. I accept only their Gaussian detections with S/N≥ 4 and duplicate_flag<2 and not flagged as probable sidelobes. The VLASS Quick Look detection prefix of "VLASS1QLCIR" is here shortened to "VLA" for brevity.
- The LOFAR Two-metre Sky Survey second data release (LoTSS: Shimwell et al. 2022) which is a 120-168 MHz survey which covers 27% of the Northern sky as is seen in dark on Figure 1. It has a depth of 0.5 mJy and astrometric accuracy of 0.2 arcsec, and comprises 52% of all core radio associations presented in MORX v2. LoTSS comes in two primary tables, the Source 'island' catalogue and the 'Gaussian component' (i.e., functional detections) catalogue. MORX processing is done on the detections catalogue which is architecturally many-to-one with the sources, although most sources have only the one detection. The source catalogue provides source names prefixed

⁴ at https://cirada.ca/catalogues

 $^{^5}$ "depth" = expected 67% completeness at the given point source flux




Fig. 1.— Sky coverage of MORX v2, darker is denser. The Galaxy winds through with X-ray stars mostly, and the Magellanic Clouds are seen at lower right. The dark footprints in the North show LoTSS radio coverage. VLASS radio sources cover sky of $\delta > -40^{\circ}$ as does NVSS, RACS radio covers sky of $\delta < 30^{\circ}$, and FIRST radio covers sky of $\delta > -10^{\circ}$ away from the Galaxy. X-ray source coverage is broadly uniform over the sky, following where pointed investigations were done. SDSS extensions radiate from their North sky optical coverage. The thin strip along the equator shows the deep-sky investigations there. The dense bead at lower left shows the XXL-South AAOmega X-ray field. (chart produced with TOPCAT (Taylor 2005))

with 'ILT' (for International LOFAR Telescope), and that name is used for any MORX optically-associated detection which is the only detection for its source island. If that detection is one of many for its source island, then LoTSS does not provide a name for it, so MORX constructs the detection name with the prefix 'ILD' (for ILT Detection) and the J2000 of the detection location. LoTSS is the largest input catalogue and took a week for me to process.

- The Faint Images of the Radio Sky at Twenty-cm survey catalogue (FIRST: Helfand/White/Becker 2015) which is a 1.4GHz survey with a Northernsky footprint away from the Galaxy, with a depth of 1mJy and astrometric accuracy of ≈1 arcsec.
- The Rapid ASKAP Continuum Survey (RACS: Hale et al. 2021) which is an 887.5 MHz survey which maps all sky south of declination +41°, although this first catalogue of sources spans only $(-80^{\circ} < \delta < +30^{\circ})$ for quality reasons. It has a point-source depth of 2 mJy and astrometric accuracy of ≈ 2 arcsec. RACS uses the same source extraction software as LoTSS, so also presents a source catalogue and a 'Gaussian' functional detections catalogue. So, as with LoTSS, MORX uses the RACS-provided source name when an optically-associated detection is the only detection for its source, and otherwise constructs a detection name with the prefix 'RACD' (for RACS Detection) and the J2000 of the detection location. I did not use the small Galaxy RACS files which

give only low-confidence associations due to optical crowding on the Galactic plane.

- The NRAO VLA Sky Survey catalogue (NVSS: Condon et al. 1998) which covers the whole sky north of declination -40° in 1.4GHz to a depth of 2.5mJy and astrometric accuracy of ≈5 arcsec.
- The Sydney University Molonglo Sky Survey catalogue (SUMSS: Murphy et al. 2007) which is analogous to NVSS but covers the sky south of declination -30° in 843MHz. It includes those associations marked as "MGPS" (Molonglo Galactic Plane survey) which is the Galactic plane component of the SUMSS survey.

X-ray surveys are collected from pointed observations by X-ray detecting satellites, so their data are not characterized by completeness and depth as are the sky-sweeping radio surveys. For X-ray sources there are 12 input catalogues from the high-resolution *Chandra* and *XMM-Newton* surveys, the medium-resolution *Swift* survey, and lower-resolution XMM-Slew and *ROSAT* satellite surveys; these are:

• For Chandra data, the Chandra ACIS source catalog (CXOG: Wang et al. 2016), the Chandra Source Catalog v1.1 (CXO: Evans et al. 2010) and v2.0 (2CXO: Evans et al. 2020), and the XAssist Chandra source list (CXOX: Ptak & Griffiths 2001a). The CXO v1.1 data is preferred over the CXO v2.0 data because CXO v1.1 provides the observation ID for each detection, needed by MORX to calculate the optical solution, whereas CXO v2.0 provides stacked detections only. Chandra data has

the best astrometric accuracy of the X-ray surveys, but published data is only to 2014. CXO v2.1⁶ will publish more recent data when released, but may provide only stacked X-ray sources aligned to optical sources which, in my view, would prejudge the identification of true sources of X-ray emission.

- For XMM-Newton data, catalogues used are the XMM-Newton DR13 (4XMM: Webb et al. 2020), the XMM-Newton DR3 (2XMMi: Watson et al. 2009) which gives ≈20K valid clean detections dropped by its successors (Rosen et al. 2016, Appendix D: "mostly real sources"), the XAssist XMM-Newton source list (XMMX: Ptak & Griffiths 2001b), and the XMM-Newton Slew survey release 2 (XMMSL2: Saxton et al. 2008). The Slew data is of much lower astrometric accuracy than the others, so is treated as a separate survey of low resolution in the MORX processing.
- Swift data is taken from the Living Swift X-ray Point Source catalogue (SXPS: Evans et al. 2023) as at 01-July-2023. The SXPS data are given both as observation-identified and stacked detections, so where both are present for a single source, MORX processing selects the observation-identified detections, as with the CXO processing.
- Lastly, the legacy ROSAT data was of a lesser astrometric accuracy; catalogues used are the High Resolution Imager (HRI: Voges et al. 1999a), Position Sensitive Proportional Counter (PSPC: Voges et al. 1999b), and the WGACAT (WGA: White/Giommi/Angelini 1994, May 2000 edition) which used different processing over the PSPC survey data. The ROSAT All-Sky Survey (RASS) is no longer used by MORX because its resolution was too coarse to yield confident optical associations in isolation.

Counts of associations from all these input catalogues are given in Table 1, plus a count of "primary associations" which are one-to-one with optical sources for each of radio and X-ray. The radio associations are calculated independently from the X-ray associations; an optical source showing both radio and X-ray associations has earned each one separately, and so can tally both as a primary radio association and a primary X-ray association in Table 1.

3. THE OPTICAL BACKGROUND USED IN MORX V2

The optical astrometry for MORX v2 hails from the "best" sources which comprise a mix of 23.7% *Gaia*-EDR3 (Lindegren et al. 2021), 46.2% Pan-STARRS (Chambers et al. 2016), 19.0% SDSS-Sweeps⁷ supplemented with SDSS XDQSO (Bovy et al. 2011) optical data, 0.6% from DES (Abbott et al. 2021) or discovery authors' papers, and 10.5% from the 0.1-arcsec resolution All-Sky Portable optical catalogue (ASP: Flesch 2017) which includes legacy optical data. Most MORX objects are optically too faint for *Gaia* coverage, but SDSS is

TABLE 1 Summary of Radio/X-ray Associations in MORX

Source	# total core	# primary* core	# double	
Catalogue(s)	associations	associations	radio lobes	
Radio surveys				
VLASS	439283	439283	15763	
LoTSS	1804886	1710297	73142	
FIRST	275552	69766	9000	
RACS	582668	337540	12009	
NVSS	316039	78759	675	
SUMSS	47549	15891	42	
total Radio Survey	s 3465977	2651536	110631	
X-ray Surveys				
CXOG	63503	63503		
CXO v1.1	13376	13376		
CXO v2.0	6807	6807		
CXOX	18966	18966		
$total\ Chandra$	102652	102652		
4XMM-DR13	250592	221701		
2XMM-DR3	4541	4086		
XMMX	8516	8040		
total XMM-Newton	n 263649	233827		
Swift LSXPS	120647	95484		
XMM Slew v2.0	11428	8778		
ROSAT HRI	11177	5146		
ROSAT PSPC	17813	12155		
ROSAT WGA	2760	2321		
$total\ ROSAT$	31750	19622		
total X-ray Surveys	s 530126	460363		

^{*} For each radio source, the 'primary' association is the first occurence on this list from top to bottom, e.g., if a radio source has associations from the RACS and NVSS surveys, then RACS has the primary association because it is topmost on this list. X-ray sources are tallied similarly.

used as available, Pan-STARRS gives almost complete coverage for $\delta > -30^{\circ}$, and ASP covers south of there.

By contrast, for optical photometry MORX gives only 2 bands, red and blue, so no attempt is made to give "best" values for these, as researchers will source those elsewhere, anyway. Instead, MORX presents historically-valuable calibrated POSS-I (1950's epoch) & POSS-II/UKST (1980's epoch) photometry sourced from ASP. In particular, the priority is to present POSS-I magnitudes because the blue POSS-I O, centred at violet 4050Å, is well-separated from the red Cousins 6400Å, and the two plates were always taken on the same night thus giving accurate red-blue colour even for variable objects. In total, MORX gives photometry comprised of 37.4% POSS-I, 26.7% POSS-II/UKST, 31.6% SDSS-Sweeps, 1.9% Pan-STARRS, 0.6% DES, and 1.9% other.

For MORX v2 (this edition), I have modified the optical field solutions documented in MORXv1, in that I have constrained the optical field shifts to be no more than 8 arcsec in each of RA and DEC. This is because of the concise presentation of this edition where each radio/X-ray association is displayed simply with its J2000-based name, without elaboration. Thus each such association needs to be "seen to be" correct, lest it be "seen to be" wrong. Thus I drop any association found to be positionally too far removed from its own J2000-based name (that name showing its original astrometry, in principle); in practice, this drops about 2% of the X-ray associations but does not disturb the radio associations. Spot checks on the removed X-ray field shifts show most

⁶ https://asc.harvard.edu/csc2/about2.1.html

⁷ at https://data.sdss.org/sas/dr9/boss/sweeps/dr9/

were only marginally aligned to the optical background, but some were quite clear, e.g., the ROSAT-HRI field US700009H.N1 (centred over NGC 3628) shifted 15 arcsec which revealed X-ray emission for 5 quasars in a closely-aligned fit. However, usually such places have since been resurveyed anyway, thus clarifying the true optical sources, as in this case which was resurveyed by Chandra which confirmed the field-shifted HRI-found associations, and so preserving those optical sources in MORX.

4. ASSOCIATION LIKELIHOODS OF RADIO/X-RAY SOURCES

The calculations of the likelihood of core radio or X-ray associations to an optical object are as presented in MORX v1, but now those likelihoods are calculated in 0.1-arcsec offset bins instead of the previous 1-arcsec bins. Smoothing is now done onto a logarithmic profile instead of linear, to avoid over-representing likelihoods. While the likelihood lower-limit cutoff for MORX remains at 40%, past analysis has shown performance at that lower limit to be better than that; a Y2009 test of this against the predecessor QORG catalogue (Flesch & Hardcastle 2004) is given at https://quasars.org/docs/Testing-QORG-via-SDSS-DR7.txt.

The best quasar candidates ($66\,026$ with pQSO $\approx 99\%$ in bulk) are also presented in the Milliquas v8 catalogue (Flesch 2023) which was extracted alongside MORX out of a large frozen database holding data to 30 June 2023. Milliquas used to hold many more quasar candidates; those are now included here in this MORX v2 catalogue.

5. COMPARISON AGAINST LoTSS OPTICAL IDENTIFICATIONS

A few weeks after this MORX paper was first submitted, a large optical identifications catalogue for LoTSS-DR2 radio sources appeared (Hardcastle et al. 2023, hereinafter: LOPTS) which contains 3519018 optical identifications (952 more than reported in their Table 5) over 4116934 LoTSS sources for a hit rate of 85.48%. MORX, on the other hand, shows "only" 1829471 optical identifications for LoTSS sources. It behoves us to cross-check these catalogues to see what can be learned about the nature of their data.

Three substantive differences between the two catalogues' handling of the LoTSS radio data are:

- LOPTS uses optical data wholly from the DESI-DR9 (Dey et al. 2019) "sweeps" catalogues whilst MORX uses the optical data outlined in Section 3. DESI photometry reaches 1-2 magnitudes deeper than surveys used by MORX, so LOPTS will have more optical objects to match to. The DESI data covers most but not all of the LoTSS-covered sky (which is ≈14% of all-sky), so the remaining ≈300 deg² of LoTSS sky is not covered by LOPTS, but MORX does cover it, being an all-sky catalogue.
- LOPTS matches opticals primarily to the LoTSS 'source' table (described in Section 2, above), whilst MORX matches only to the LoTSS 'Gaussian' (detections) table. However, often a radio source manifests as multiple detections, 2 or 3 in the case of double lobes, more for extended or saturating emission. Thus LOPTS supplements the

- groupings provided by the source extraction software with further visually-ascertained groupings. In contrast, MORX groups only raw detections via algorithms previously used by MORXv1 and earlier works, including algorithmic identification of double lobes. All this will impact optical selection.
- Both catalogues give a confidence-of-association figure, MORX as a simple percentage, LOPTS as a Likelihood Ratio (LR) which converts to confidence-of-association by conf = LR/(1+LR). MORX gives optical identifications down to a confidence of 40%, and no lower. LOPTS reports down to about 24% confidence (i.e., LR=0.3) for those not identified visually. (MORX has no flagged visual identifications, but thousands of visual checks have been done over the life of the catalogue development and use.)

Let's be clear that not all these identifications are equal; if the reported confidence-of-associations are accurate, then one 100%-confidence association is worth two 50%-confidence associations. If the confidence pcts are totalled, then MORX will deliver 1695614 true associations out of 1829471 reported associations, and LOPTS will deliver 3246715 true associations out of 3451555 reported associations for which an LR is given. This confidence-driven "yield" is an essential part of any probabilistic analysis, and is presented alongside count numbers below and in Table 2.

Now let's see how well the catalogues agree, the better to gauge any points of disagreement. Matching by LoTSS source name and accepting only those where the MORX and LOPTS optical co-ordinates are within one arcsec of each other (although arbitrarily far from the radio source), we obtain 1494158 agreeing radio-optical associations. For these, MORX gives a mean confidenceof-association of 95.084%, while LOPTS gives 97.268%. Thus MORX is more pessimistic, but the data-driven MORX calculations evaluate and so incorporate the contribution of unseen background sources. Since LOPTS uses deeper DESI optical data, the chance of unseen background sources is less, so LOPTS is justified to display more confidence. Also, over these data, the mean of the absolute difference between the MORX and LOPTS confidence-of-association is 4.244%, and the median absolute difference is just 1.332%. Thus the two catalogues align pleasingly well, for those LoTSS-optical associations that they share.

With the two catalogues thus showing equivalent throughputs, we now list where they differ. Counts are supplemented with the yields after factoring in confidence-of-association:

• LOPTS has 1936725 LoTSS-optical associations not presented by MORX. Of these, 42718 are visually evaluated without an LR, so crediting those as 100% confident, the total yield of confidences is 1777414 LOPTS-only associations. As we have established the similar performance of LOPTS and MORX above, the expectation is that the LOPTS LRs are assessing these with good accuracy. Inspection of those data show dominantly faint opticals, assuredly from DESI, which LOPTS carries and MORX does not.

 ${\rm TABLE~2}$ Optical selections for LoTSS sources – MORX vs LOPTS (refer Section 5)

MORX Counts & Confidence-Based Yields

	IN DESI I	FOOTPRINT	OUTSIDE	OUTSIDE OF DESI		ΓAL			
	Counts	Yields	Counts	Yields	Counts	Yields			
MATCHES ACROSS MORX AND LOPTS (opticals within 2 arcsec of eachother):									
Isolated Sources from both MORX & LOPTS	1441336	1372317	6801	6393	1448137	1378710			
MORX isolated sources / LOPTS blobs-lobes	58577	51359	358	307	58935	51666			
MORX cored lobes [†] / LOPTS isolated sources	25892	24912	89	84	25981	24996			
MORX blank lobes [†] / LOPTS isolated sources	1541	1295	10	8	1551	1303			
MORX cored lobes / LOPTS blobs-lobes	9046	8311	64	58	9110	8369			
MORX blank lobes / LOPTS blobs-lobes	3734	3140	25	20	3759	3160			
TOTAL MORX-LOPTS MATCHES:	1540126	1461334	7347	6870	1547473	1468204			
MORX OPTICALS VALID AND UNMATCHED TO LOPTS:									
Isolated Sources	85466	70039	59711	54033	145177	124072			
Cored Lobes	10968	9869	1854	1671	12822	11540			
Blank Lobes	9037	6750	804	591	9841	7341			
TOTAL MORX VALID UNMATCHED:	105471	86658	62369	56295	167840	142953			
OPTICALS DISAGREE, LOPTS OPTICAL SELECTED OVER MORX* (opticals 2+ arcsec apart):									
Isolated Sources	103359	76333	721	560	104080	76893			
Cored Lobes	3252	2726	36	29	3288	2755			
Blank Lobes	6770	4794	20	15	6790	4809			
TOTAL MORX SUPERSEDED:	113381	83853	777	604	114158	84457			
TOTAL MORX OPTICAL SELECTIONS FOR LoTSS:	1758978	1631845	70493	63769	1829471	1695614			

LOPTS Counts & Confidence-Based Yields (no-LR visuals counted as 100% confidence)

	IN DESI I	FOOTPRINT	OUTSIDE	OUTSIDE OF DESI		Γ AL			
	Counts	Yields	Counts	Yields	Counts	Yields			
MATCHES ACROSS LOPTS AND MORX (opticals within 2 arcsec of eachother):									
Isolated Sources from both LOPTS & MORX	1441336	1404857	6801	6579	1448137	1411436			
LOPTS isolated sources / MORX lobes	27433	26742	99	95	27532	26837			
LOPTS blobs-lobes / MORX isolated sources	58577	54095	358	306	58935	54401			
LOPTS blobs-lobes / MORX lobes	12780	12653	89	83	12869	12736			
TOTAL LOPTS-MORX MATCHES:	1540126	1498347	7347	7063	1547473	1505410			
LOPTS OPTICALS VALID AND UNMATCHED TO M	ORX:								
Isolated Sources with Likelihood Ratio (LR):	1758919	1624109	10084	9292	1769003	1633401			
Visual Isolated Sources without LR:	37140	37140	22	22	37162	37162			
Blobs / Lobes with LR:	123793	100376	1211	919	125004	101295			
Visual Blobs / Lobes without LR:	5551	5551	5	5	5556	5556			
TOTAL LOPTS VALID UNMATCHED:	1925403	1767176	11322	10238	1936725	1777414			
DISAGREEMENT, LOPTS OPTICALS SUPERSEDED BY MORX DOUBLE LOBES (in this exercise):									
Isolated Sources	32054	29564	91	83	32145	29647			
Blobs / Lobes	2649	1566	26	13	2675	1579			
TOTAL LOPTS SUPERSEDED:	34703	31130	117	96	34820	31226			
TOTAL LOPTS OPTICAL SELECTIONS FOR LoTSS:	3500232	3296653	18786	17397	3519018	3314050			

^{*} this decision-branch used for this exercise. There are a mix of objects on each side of the 2-arcsec dividing line, but visual inspection of all those objects is beyond the scope of this paper.

- MORX has 167840 (yield 142953) valid LoTSSoptical associations not in LOPTS. Of those, 62 369 (yield 56295) are outside of the DESI footprint, MORX being an all-sky catalogue. I visually inspected some within the DESI footprint to find why LOPTS did not match those. Among a variety of situations, some stood out: (1) Galaxies can have multiple LoTSS sources not consolidated by the source extraction software, so sometimes LOPTS and MORX select different nominal radio sources for the same optical association, thus evading the matching method of this exercise, (2) Some faint reddish galaxies appear to be in Pan-STARRS or SDSS data but not in the DESI data, and (3) When a LoTSS source has multiple components (detections), one of those
- components may match well to a prominent optical, but somehow this can get missed by the LOPTS method. An example is the LoTSS source ILTJ000412.62+325711.8 which has a component MORX-named as ILDJ000412.17+325709.5, which matches with 100% confidence to the SDSS-DR16Q quasar SDSS J000412.15+325709.2, z=1.386 but missed by LOPTS. Browsing the LOPTS components table shows this component to be absent, with only the other one component reported for this source, for some reason. Querying the whole LOPTS components table finds 28 841 sources with only one component reported, curiously. A more thorough evaluation of this overall topic is out of scope for this paper.
- MORX uses a unique algorithm to identify dou-

^{† &}quot;cored lobes": double radio lobes with a radio centroid. "blank lobes": same but no radio centroid in the data.

TABLE 3 Sample lines from the MORX catalogue (left half placed on top of right half)

RA (J2000) DECL IN DEGREES	NAME OR J2000 (IF NO NAME)	R B OPT			SSOC QSO GAL STR XRAY		XMM-NEWTON X-RAY SOURCE ID	SWIFT X-RAY SOURCE ID	
228.7634445 61.5889250 228.7634696 42.0498599 228.7635214 36.9288189		QR2 18.81 18.77 gG ARX 6.96 7.91 p+ X 19.63 21.27 j+N	1 1 0.008 1592	1761			4XMM J151503.2+420259	LSXPSJ151503.2+420259	
228.7635418 36.8302012 228.7636380 50.6961568	NPM1+37.1259	SX 8.05 8.63 b+G R 17.52 19.61 pG QR2 18.44 18.40 gG	0982 1 - 0.200 MQ	MQ	100 1 1 98 100 C 100 7 93 0		4XMM J151503.2+364948	LSXPSJ151503.2+364949	
228.7638054 62.3565436 228.7638674 66.9552929 228.7639118 36.4723147	J151503.31+622123.5	R 19.63 20.99 jN R 22.32 23.43 gN X 14.92 17.21 pmG	1 1 0.300 MO	MO	100 18 82 0 98 23 74 1 69 19 21 29 69				
228.7639208 46.7413589 228.7639867 38.5757067	J151503.34+464428.8 J151503.35+383432.5 SDSS J151503.38+013147.7	R 19.70 21.02 pN R 20.30 21.68 j+m	1 1 0.500 MQ 1 - MQ	MQ	99 35 64 0 99 32 67 0 100 96 3 1				
	DIO VLASS ONF RADIO CORE SOURCE ID		LoTSS RADIO CORE SOURC			NVSS RADIO CORE SOURCE ID	DOUBLE RADIO LOBE 1 ID	DOUBLE RADIO LOBE 2 ID	LOBE MAX"
	00 VLAJ151503.04+613519.8 00 VLAJ151503.26+420259.4		ILDJ151503.20+61 ILDJ151503.24+42			NVSS J151503.8+613520) ILDJ151502.23+613519	.5 ILDJ151504.49+613521	.2 9
	00		ILTJ151503.27+50						
1	97 VLAJ151503.38+223145.8 00 98		ILTJ151503.39+62 ILDJ151503.28+66	2123.7	KACD J151503.5+223145	NVSS J151503.4+22314	FIRST J151503.4+2231	39 FIRST J151503.3+22314	19 6
	99 99 00 VLAJ151503.37+013147.9	FIRSTJ151503.3+013147	ILTJ151503.40+46 ILTJ151503.35+38	3432.8	RACS J151503.3+013147	NVSS J151503.2+013144	1		

Notes on columns (see ReadMe for full descriptions):

- TYPE: R=core radio detection, X=X-ray detection, 2=double radio lobes (calculated), Q=QSO, A=AGN, G=galaxy, S=star.
- REF & ZREF: citations for name and redshift: citations are indexed in the file "MORX-references.txt".
- OPT COMM: comment on photometry: p=POSS-I magnitudes so blue is POSS-I O, j=POSS-II/UKST Bj, g=SDSS g & r, +=optically variable, m=nominal proper motion, G=Gaia astrometry, N=Pan-STARRS astrometry.
- R/B PSFs: '-'=stellar, 1=fuzzy, n=no PSF available, x=not seen in this band.
- ASSOC/RADIO/X-RAY CONF: calculated percentage confidence that this source is truly associated to this optical object in total/radio/X-ray, respectively.
- QSO/GAL/STR PCT: based on its photometry and the radio/X-ray association(s), the calculated percentage confidence that this optical object is a QSO/galaxy/star.
- LOBE MAX": offset of the longest radio lobe from the centroid, in arcsec. This calculation uses detection centroids, so visual lobe length can be longer.

The full table can be downloaded from http://quasars.org/morx.htm, also available in FITS.

ble radio lobes, including ones without radio centroids which require visual follow-up. LOPTS finds these from the source extraction software, and visual evaluation. In general, double radio lobes need a radio centroid confidently associated to an optical object, to be regarded as fully confirmed (hereinafter: "cored lobes"). The MORX-presented lobes without a radio centroid (hereinafter: "blank lobes") are assigned onto the "best" candidate optical centroid; visual inspection there often reveals a faint radio signature which confirms those lobes, but sometimes the faint radio signature (if one is found at all) is onto another nearby optical, thus falsifying the MORX candidate. In this exercise, as shown in Table 2 (4^{th} and 6^{th} lines of data), 5310 blank MORX lobes match their optical-only centroids to LOPTS radio-optical sources which thus provide radio-optical centroids for those MORX lobes, provisionally converting them from "blank lobes" to "cored lobes".

Of 32 741 MORX-only LoTSS double lobes, 10 078 get falsified in this exercise by a lobe radio source matching to a LOPTS optical association, because the faint DESI optical was not in the MORX optical data. But also 34 820 LOPTS associations get superseded (in this exercise) by qualifying MORX double lobes, either because the lobes have a supporting core radio-optical association which yields a high-confidence lobe configuration, or because the LOPTS optical was offset ≥ 2 arcsec from the radio detection, thus confirming that radio detection as

a lobe signature, in this exercise.

Thus, this exercise has shown the MORX and LOPTS catalogues to be well-aligned in their throughput, with LOPTS having twice as many LoTSS-optical associations due to its use of deep DESI optical data absent from MORX, whilst MORX associations are generally confirmed by LOTSS where not overtaken by those same deep DESI opticals. There are of course some disagreeing optical selections inevitable in such large data. The one contentious arena is that of double radio lobes, which can be settled only by individual visual inspections.

6. CATALOGUE LAYOUT

The MORX catalogue presents one line per optical object; Table 3 displays 12 sample lines which are wrapped with the left half of the block of lines shown on top of the right half. Names are given as found in the literature; else, if anonymous, the J2000 location is given as a convenience to the user. The ReadMe gives full details and indexes to the columns.

7. CONCLUSION

The MORX v2 catalogue is presented which gives 3 115 575 probable radio/X-ray associations onto optical objects over the whole sky, including double radio lobes, using all the largest radio/X-ray source surveys to June 2023. Identifications are included to provide an informative map for pointed investigations.

ACKNOWLEDGEMENTS

Thanks to Heinz Andernach for steering me onto VLASS, RACS, and LoTSS, although processing the lat-

ter almost broke my computer. Also thanks to the referee for good suggestions. This work was not funded.

DATA AVAILABILITY

MORX v2 can be downloaded from CDS at https:// cdsarc.cds.unistra.fr/viz-bin/cat/V/158 or from its home page at https://quasars.org/morx.htm which also provides a FITS file. Both sites also provide the ReadMe and the references list. Query pages are provided by CDS and NASA HEASARC at https: //heasarc.gsfc.nasa.gov/W3Browse/all/morx.html.

REFERENCES

Abbott T. M. C., Adamów M., Aguena M., et al., 2021, ApJS, 255, 20, DES DR2 Alam S., Albareti F. D., Allende Prieto C., et al., 2015, ApJS, 219, 12, SDSS DR12 pipeline Bovy J., Hennawi J. F., Hogg D. W., et al., 2011, ApJ, 729, 141, SDSS-XDQSO Chambers K. C., Magnier E. A., Metcalfe N., 2015, arXiv:1612.05560, Pan-STARRS DR1 Condon J. J., Cotton W. D., Greisen E. W., et al., 1998, AJ, 115, 1693, NRAO VLA sky survey (NVSS), https://www.cv.nrao.edu/nvss Dey A., Schlegel D. J., Lang D., et al., 2019, AJ, 157, 168, DESI Legacy Imaging Surveys Evans I. N., et al., 2010, ApJS, 189, 37, Chandra Source Catalog release 1.1 (CXO), https://asc.harvard.edu/csc1/ Evans I. N., Primini F. A., Miller J. B., et al., 2020, AAS, 235, 154.05, Chandra Source Catalog v2.0 (2CXO), https://asc.harvard.edu/csc2/ Evans P. A., Page K. L., Beardmore A. P., et al., 2023, MNRAS, 518, 175, Living Swift X-ray Point Source catalog (LSXPS),

http://www.swift.ac.uk/LSXPS, as at 1-July-2023. Flesch E. & Hardcastle M. J., 2004, A&A, 427, 387, Quasars.Org catalogue (QORG)

Flesch E. W., 2016, PASA, 33, 52, MORX v1 catalogue Flesch E. W., 2017, PASA, 34, 25, All-Sky Portable Optical catalogue (ASP)

Flesch E. W., 2023, OJAp, 6, 49, Milliquas v8, $\verb|https://doi.org/10.21105/astro.2308.01505|, home page|$ https://quasars.org/milliquas.htm

Gordon Y. A., Boyce M. M., O'Dea C. P., et al., 2020, RNAAS, 4, 175, VLASS Epoch 1 Quick Look, https://science.nrao.edu/vlass

Hale C. L., McConnell D., Thomson A. J. M., et al., 2021, PASA, 38, 58, Rapid ASKAP Continuum Survey (RACS), https://data.csiro.au/collection/csiro:52217 and click on "Files 1602".

Hardcastle M. J., Horton M. A., Williams W. L., et al., 2023, A&A, 678, A151, Optical Identifications for LoTSS DR2 (LOPTS)

Helfand D. J., White R. L., & Becker R. H., 2015, ApJ, 801, 26, VLA FIRST survey, 13Jun05 version, https://sundog.stsci.edu

Lindegren L., Klioner S. A., Hernández J., et al., 2021, A&A, 649, A2, Gaia EDR3

This paper was built using the Open Journal of Astrophysics LATEX template. The OJA is a journal which Murphy T., Mauch T., Green A., Hunstead R. W., Piestrzynska B., Kels A. P., Sztajer P., 2007, MNRAS, 382, 382, Sydney U. Molonglo Survey (SUMSS), http://www.astrop.physics.usyd.edu.au/sumsscat. Molonglo galactic plane (MGPS) data at http://www.astrop.physics.usyd.edu.au/mgpscat. Ptak A. & Griffiths R., 2001, AAS, 198, 311, XAssist Chandra source list (CXOX), https://asd.gsfc.nasa.gov/xassist/pipeline4/chandra/ Ptak A. & Griffiths R., 2001, AAS, 198, 311, XAssist XMM-Newton source list, https://asd.gsfc.nasa.gov/xassist/pipeline5/xmm/ Rosen S. R., Webb N. A., Watson M. G., et al., 2016, A&A, 590, A1, 3XMM-DR6 Saxton R. D., Read A. M., Esquej P., Freyberg M. J., Altieri B., & Bermejo D., 2008, A&A, 480, 611, XMM-Newton Slew Survey Release 2.0 (XMMSL2), https://www.cosmos.esa.int/web/xmm-newton/xmmsl2-ug Shimwell T. W., Hardcastle M. J., Tasse C., et al., 2022, A&A, 659, A1, LOFAR Two-metre Sky Survey second data release (LoTSS-DR2), https://www.lofar-surveys.org Taylor M. B., 2005, ASPC, 347, 29, TOPCAT software Voges W. & MPE-ROSAT Team, 1999DTRP.conf..179V, ROSAT high resolution imager (1RXH), http://cdsarc.cds.unistra.fr/cat/?IX/28A Voges W. & MPE-ROSAT Team, 1999DTRP.conf..179V, ROSAT position sensitive proportional counter (2RXP), http://cdsarc.cds.unistra.fr/cat/?IX/30 Wang S., Liu J., Qiu Y., et al., 2016, ApJS, 224, 40, Chandra ACIS source catalog (CXOG) Watson M. G., Schr'oder A. C., Fyfe D., et al., 2009, A&A, 493, 339, XMM-Newton 2XMMi-DR3, http://cdsarc.cds.unistra.fr/cat/?IX/41 Webb N. A., Coriat M., Traulsen I., et al., 2020, A&A, 641, 136, XMM-Newton 4XMM-DR13 (2023 update),

provides fast and easy peer review for new papers in the astro-ph section of the arXiv, making the reviewing process simpler for authors and referees alike. Learn more at http://astro.theoj.org.

https://www.cosmos.esa.int/web/xmm-newton/xsa

http://cdsarc.cds.unistra.fr/cat/?IX/31

WGACAT May-2000,

White N. E., Giommi P., & Angelini L., 1994, AAS, 185, 4111,