

REPRESENTATION DIMENSION OF SOME FINITE GROUPS

GURLEEN KAUR, AMIT KULSHRESTHA, AND ANUPAM SINGH

ABSTRACT. For a finite group G , the representation dimension is the smallest integer realizable as the degree of a complex faithful representation of G . In this article, we compute representation dimension for some p -groups, their direct products, and groups with certain conditions on nonlinear irreducible characters. We also make similar computations for the smallest integer realizable as the degree of an irreducible complex faithful representation of G , if one exists. In the appendix, we present GAP codes to compute these numbers.

1. INTRODUCTION

Throughout this paper, we restrict our attention to finite groups and complex representations. Let G be a finite group. One of the classical and challenging problems in the representation theory of finite groups is to determine the smallest positive integer n such that G embeds into $\mathrm{GL}(n, \mathbb{C})$, i.e., G is isomorphic to a linear group of degree n . The study of the linear groups has been of natural interest with intensive development by O'Brien, Flannery, Green, James, Jansen, Lusztig, Martino, Steinberg, Tamburini, Tiep, and Zalesskii (see [11, 15, 17, 21, 23, 28, 35, 37]) to name a few. The *representation dimension* $\delta(G)$ of a finite group G is defined as follows:

$$\delta(G) := \min\{\deg(\rho) : \rho \text{ is a complex faithful representation of } G\},$$

where $\deg(\rho)$ denotes the degree of complex faithful representation ρ of G .

Representation dimension has been extensively studied for finite groups of Lie type (see for example Lubeck [27], Tiep and Zalesskii [37]), and for finite p -groups (see for example Bardestani, Mallahi-Karai and Salmasian [1], Martino and Tamburini [12], Karpenko and Merkurjev [24]).

In literature, another related number $\delta_{irr}(G)$, as defined below, is studied.

$$\delta_{irr}(G) := \min\{\deg(\rho) : \rho \text{ is an irreducible complex faithful representation of } G\}.$$

We note that $\delta_{irr}(G)$ does not exist for all groups. However, when it exists, we have the inequality $\delta(G) \leq \delta_{irr}(G)$. Szechtman provided a brief history of determining $\delta_{irr}(G)$ in §2 of [36].

2020 *Mathematics Subject Classification.* 20D15, 20C15.

Key words and phrases. irreducible representations, faithful representations, characters, representation dimension.

In this article, we begin with listing several examples in §2 and we prove in a series of theorems that $\delta(G)$ equals $\delta_{irr}(G)$ if G is one of the following:

- (i) direct product of simple groups with relatively coprime order;
- (ii) direct product of monolithic groups with relatively coprime order;
- (iii) finite nilpotent group with cyclic center;
- (iv) nonabelian finite group whose all nonlinear irreducible characters have distinct degrees (in particular, finite group with unique nonlinear irreducible character);
- (v) direct product of nonabelian groups of relatively coprime order, whose all nonlinear irreducible characters have distinct degrees.

We determine representation dimension for some finite p -groups in Theorem 3.1, for certain direct products of finite groups in Theorem 4.1, for odd order groups with exactly two nonlinear irreducible characters of each degree in Theorem 5.1 (following the classification of same from [10]), for finite groups whose all nonlinear irreducible characters are of distinct degree in Theorem 6.1 (following the classification of the same in [5]). In Appendix A, we present GAP codes to compute $\delta(G)$ and $\delta_{irr}(G)$.

2. PRELIMINARIES

We begin with some examples to understand the representation dimension, $\delta(G)$, of a finite group G .

Example 2.1. When G is a cyclic group, $\delta(G) = \delta_{irr}(G) = 1$. However, if $G = C_2 \times C_2$, then $\delta(G)$ equals 2 which shows that understanding the character table of a group is not enough to determine its representation dimension. It is evident that for a group G , $\delta(G) = 1$ if, and only if, G is cyclic. We note that $\delta_{irr}(C_2 \times C_2)$ does not exist.

Example 2.2. If G is a dihedral group, quaternion group, or $C_n \times C_m$, where m is not coprime to n , then $\delta(G) = 2$. Determining all finite groups with representation dimension 2 is equivalent to determining all the finite subgroups of $\text{GL}(2, \mathbb{C})$. It is pertinent to mention that Brauer, Feit and their successors classified finite linear groups of degree upto 11 over \mathbb{C} (for more details, see the survey article [37] by Tiep and Zalesskii).

Example 2.3. The representation dimension of a finite abelian group is the rank of the group (see Lemma 3.4 of [31]). Thus, the representation dimension $\delta(G)$ for finite groups G may be arbitrarily large.

If a group G has a faithful irreducible character χ , then $\delta(G) \leq \deg(\chi)$, however $\delta(G)$ may be different from $\delta_{irr}(G)$.

Example 2.4. For the group $G = A_4 \times D_{10}$, we have $\delta(A_4 \times D_{10}) = 5$ and $\delta_{irr}(A_4 \times D_{10}) = 6$. This is a particular case of a more general result given in Theorem 4.1.

In the case of nonabelian finite simple groups, every nontrivial character is faithful. Hence, for these groups, the representation dimension is equal to the minimal degree of a nonlinear irreducible character.

Example 2.5. From the character table of the alternating group A_4 , we obtain that $\delta(A_4) = 3$. We remark that the representation dimension of $2.A_4$, the double cover of A_4 , is 2. This is because it is a subgroup of $SU(2)$. Along similar lines, $\delta(A_5) = 3$, as A_5 is a subgroup of $SO(3)$.

Based on the work of Bessenrodt-Tong-Viet-Zhang [6, Lemma 3.1], $\delta(A_n) = n - 1$, when $n \geq 15$. Using the GAP code given in Appendix A, we obtain that $\delta(A_n)$, for $6 \leq n \leq 14$, is $n - 1$. Hence, the representation dimension of A_n , for $n \geq 6$, is $n - 1$.

Example 2.6. Due to Result 1 of [34], we have that $\delta_{irr}(S_n) = n - 1$, when $n \geq 5$. Indeed, the standard character of S_n is faithful and has degree $n - 1$. Consequently, $\delta(S_n) = n - 1$, for $n \geq 5$.

Example 2.7. The minimal nonlinear faithful irreducible character degree of $GL(2, q)$, $GL(3, q)$ and $GL(4, q)$ is $q - 1$, $q^2 + q$ and $(q + 1)(q^2 + 1)$ respectively (see table II, V, IX in [35]). Since these are minimal nonlinear irreducible character degrees in the character table of their respective groups, these are also the representation dimensions.

Example 2.8. We have that $\delta(SL(2, q)) = q - 1$, when q is even, and $\delta(SL(2, q)) = \frac{q-1}{2}$, when q is odd (see [26]). The representation dimension for finite groups of Lie type is well studied, see [27, 37] for further details.

Example 2.9. Let G be a finite group with a non-normal Sylow p -subgroup P such that the intersection of two distinct conjugates of P is trivial. Then, from [3, Theorem 3.2], it follows that $\delta(G) > \sqrt{|P|} - 1$.

Example 2.10. If G and H are isoclinic groups with $|G| = |H|$, then $\delta(G) = \delta(H)$, due to [7, Theorem 2.2].

The essential dimension $\text{ed}(G)$ of a finite group G over \mathbb{C} , is also related to the representation dimension of G . The notion of the essential dimension of G was introduced by Buhler and Reichstein [8]. It has been proved in [15, Proposition 4.15] that $\text{ed}(G) \leq \delta(G)$. One is interested to see that when $\text{ed}(G)$ coincides with $\delta(G)$.

Example 2.11. In Theorem 4.1 of [24], Karpenko and Merkurjev proved that if G is a finite p -group, then $\text{ed}(G)$ coincides with $\delta(G)$.

3. REPRESENTATION DIMENSION OF p -GROUPS

Let G be a finite group. Suppose it has a faithful irreducible character. This information can be read from the character table of G . Then $\delta(G) \leq \delta_{irr}(G)$. One is interested in

understanding when $\delta(G)$ and $\delta_{irr}(G)$ are equal. In the next theorem, we prove that for p -groups whose center $\mathcal{Z}(G)$ is cyclic, $\delta(G)$ equals $\delta_{irr}(G)$. We also determine $\delta(G)$ for certain p -groups in the following theorem. We note that for part (4) of the theorem, we depend on the classification of non abelian groups of order p^5 , up to isoclinism, into classes $\Phi_2, \Phi_3, \dots, \Phi_{10}$, as given in [22].

Theorem 3.1. *The representation dimension $\delta(G)$ of certain p -groups G is as follows:*

- (1) $\delta_{irr}(G)$, if G is a finite p -group with $\mathcal{Z}(G)$ cyclic;
- (2) p , if G is a nonabelian p -group of order p^3 ;
- (3) p , if G is a nonabelian p -group of order p^4 with $\mathcal{Z}(G)$ cyclic, and is $p+1$ otherwise;
- (4) if G is a nonabelian p -group of order p^5 , then the following cases arise, using the classification of groups via their isoclinism classes given in [22];
 - (i) p , if $G \in \Phi_2$ and its center is cyclic, $p+1$ if $\mathcal{Z}(G) \cong C_{p^2} \times C_p$ and is $p+2$ if $\mathcal{Z}(G) \cong C_p \times C_p \times C_p$;
 - (ii) $p+1$, if $G \in \Phi_3$;
 - (iii) $2p$, if $G \in \Phi_4$;
 - (iv) p^2 , if $G \in \Phi_5$;
 - (v) $2p$, if $G \in \Phi_6$;
 - (vi) p^2 , if $G \in \{\Phi_7, \Phi_8\}$;
 - (vii) p , if $G \in \Phi_9$;
 - (viii) p^2 , if $G \in \Phi_{10}$.
- (5) $\sqrt{[G : \mathcal{Z}(G)]}$, if G is a p -group with $\mathcal{Z}(G)$ cyclic and $|G'| = p$;
- (6) $\sqrt{[G : \mathcal{Z}(G)]}$, if G is isoclinic to semi-extraspecial p -group with $\mathcal{Z}(G)$ cyclic (in particular, p^r , if G is an extraspecial p -group of order p^{2r+1});
- (7) $[G : A]$, if G is a (nonabelian) normally monomial p -group with $\mathcal{Z}(G)$ cyclic, where A is a normal abelian subgroup of maximal order (in particular, groups of order p^5 with its center cyclic);
- (8) $\sqrt{[G : \mathcal{Z}(G)]} + \delta(\mathcal{Z}(G)) - 1$, if G is a p -group of nilpotency class 2 with its derived subgroup cyclic (in particular, $\sqrt{[G : \mathcal{Z}(G)]}$, if G is a p -group of nilpotency class 2 with its center cyclic).

Proof.

- (1) Let G be a p -group with cyclic center. Suppose χ is a faithful character of minimal degree. Then, due to [1, Lemma 3.5], it follows that χ can be written as a direct sum of exactly r irreducible characters, where r is the minimal number of generators of $\mathcal{Z}(G)$. Since $\mathcal{Z}(G)$ is cyclic, it follows that $r = 1$. Consequently, χ is an irreducible character of G . Hence, $\delta(G)$ equals $\delta_{irr}(G)$.
- (2) Suppose G is a nonabelian p -group of order p^3 . Since G is nonabelian, there exists an irreducible nonlinear character of G . Further, no linear character of G

is faithful, otherwise, the group G would be isomorphic to a finite subgroup of \mathbb{C}^* and is therefore cyclic. If the order of $\mathcal{Z}(G)$ is p^2 , then G is abelian. Hence, the order of $\mathcal{Z}(G)$ must be p and is therefore cyclic. By [20, Theorem 2.32], G has a faithful irreducible character. Also, in this case, $\delta(G)$ equals $\delta_{irr}(G)$. So, we need to determine $\delta_{irr}(G)$. As the quotient group $G/\mathcal{Z}(G)$ is abelian, we have that $G' \leq \mathcal{Z}(G)$. Since the order of $\mathcal{Z}(G)$ is p , it follows that G' equals $\mathcal{Z}(G)$. Due to [20, Corollary 2.30], it follows that the minimal faithful irreducible character degree, $\delta_{irr}(G)$, equals p and hence $\delta(G) = p$.

- (3) Let G be a nonabelian p -group of order p^4 and $\mathcal{Z}(G)$ be cyclic. Then $[G : \mathcal{Z}(G)] \leq p^3$. Consequently, the degree of an arbitrary nonlinear irreducible character must be p , due to [20, Corollary 2.30]. Since $\mathcal{Z}(G)$ is cyclic, $\delta(G) = \delta_{irr}(G)$. Hence $\delta(G) = p$.

Suppose that $\mathcal{Z}(G)$ is non cyclic. Then the order of $\mathcal{Z}(G)$ is p^2 . From [20, Theorem 2.32], the group G does not have a faithful irreducible character. Also, any nonlinear irreducible character of G must be of degree p , due to Corollary 2.30 of [20]. Suppose χ is a faithful character of G . From [1, Lemma 3.5], it follows that χ is a sum of two irreducible characters of G . Consequently, the degree of χ is either $p + 1$ or $2p$. However, in view of Lemma 15 of [9], if p is an odd prime, then the $\deg(\chi) \leq p + 1$; therefore, $\deg(\chi) = p + 1$. For an arbitrary group G of order 16, one can verify from the character table of G that $\delta(G) = p + 1$. Hence, the result follows.

- (4) (i) If $G \in \Phi_2$, then the set of all the irreducible character degrees of G , i.e. $\text{cd}(G)$, is $\{1, p\}$. Also, $|\mathcal{Z}(G)| = p^3$ and $|G'| = p$, due to Lemma 5.1 of [32]. Clearly $\delta(G) = p$, if $\mathcal{Z}(G)$ is cyclic. Suppose $\mathcal{Z}(G)$ is non cyclic. Now, G is p -group of nilpotency class 2 and its derived subgroup is cyclic. Therefore, in view of Theorem 1.3 of [1], it follows that $\delta(G) = p + 1$, if $\mathcal{Z}(G) \cong C_{p^2} \times C_p$ and $\delta(G) = p + 2$, if $\mathcal{Z}(G) \cong C_p \times C_p \times C_p$.
- (ii) If $G \in \Phi_3$, then $|\mathcal{Z}(G)| = p^2$ and $\Omega_1(\mathcal{Z}(G))$ is not contained in G' , where $\Omega_1(\mathcal{Z}(G))$ is the subgroup generated by elements $g \in \mathcal{Z}(G)$ such that $g^p = 1$. From Lemma 14 of [9], it follows that $\delta(G) \leq p + 1$. As $\mathcal{Z}(G)$ is non cyclic, the group G does not have an irreducible faithful character. Consequently, $\delta(G) = p + 1$.
- (iii) Let $G \in \Phi_4$. Lemma 3.5 of [1] yields that $\delta(G)$ is at most $2p$. By the description of irreducible characters of G given in Lemma 5.2 of [32], it follows that $\delta(G) = 2p$.
- (iv) Let $G \in \Phi_5$. Then G is an extraspecial p -group and hence $\delta(G) = p$ [25, Theorem 21.2.18].

- (v) If $G \in \Phi_6$, then it follows from the description of irreducible characters given in Lemma 5.4 of [32] that $\delta(G) = p$.
 - (vi) If $G \in \Phi_7$, then $\delta_{irr}(G) = p^2$ and hence $\delta(G) = p$. Similar argument holds when $G \in \Phi_8$.
 - (vii) Let $G \in \Phi_9$. Then $\mathcal{Z}(G)$ is cyclic and $\text{cd}(G) = \{1, p\}$. Consequently, $\delta(G) = p$.
 - (viii) If $G \in \Phi_{10}$, then $\delta_{irr}(G) = p^2$, in view of Lemma 5.8 of [32]. Hence $\delta(G) = p^2$.
- (5) Lemma 1 of [11] yields that all non linear irreducible characters of G are faithful of degree $\sqrt{[G : \mathcal{Z}(G)]}$. Hence, the result follows.
- (6) Suppose G is isoclinic to semi-extraspecial p -group with cyclic center. It follows from [14, Theorem A] that the degree of any nonlinear irreducible character is $\sqrt{[G : \mathcal{Z}(G)]}$. Since $\mathcal{Z}(G)$ is cyclic, G has a faithful irreducible character. Therefore, $\delta(G) = \sqrt{[G : \mathcal{Z}(G)]}$.
- (7) The group G has a faithful irreducible character χ , since $\mathcal{Z}(G)$ is cyclic. By [29, Lemma 4], $\delta_{irr}(G) = [G : A]$, where A is an abelian normal subgroup of G of maximal order. Consequently, $\delta(G) = [G : A]$.
- (8) This is a direct consequence of Theorem 1.3 of [1].

□

From the above theorem, we note that for p -groups G with cyclic center and order at most p^5 , $\delta_{irr}(G)$ equals the maximum degree of all the irreducible characters of G . Further, the p -groups of order at most p^5 are metabelian and hence normally monomial (see [2, 18] for the definition of normally monomial groups). This observation can also be obtained as a special case of the following:

Remark 3.2. In view of [30, Proposition 3], if G is a normally monomial group and $\delta_{irr}(G)$ exists, then $\delta_{irr}(G)$ equals the maximum degree of all the irreducible characters of G .

Let G be a finite p -group. In the next theorem, we provide a connection between $\delta(G)$ and $\text{cd}(G)$.

Theorem 3.3. *Let G be a finite nonabelian p -group with cyclic center.*

- (i) $\delta(G) = p$ if, and only if, $\text{cd}(G) = \{1, p\}$;
- (ii) if for some integer $a > 1$, $\text{cd}(G) = \{1, p^a\}$ or $\text{cd}(G) = \{1, p, p^a\}$, then $\delta(G) = p^a$.

Proof. (i) Let $\delta(G) = p$. In view of Theorem 3.1, $\delta(G)$ equals $\delta_{irr}(G)$. Consequently, $\delta_{irr}(G) = p$. Hence, the derived length of G is at most 2, due to Theorem 22.25 of [4]. This implies that G is a metabelian group. Since metabelian groups are normally monomial, it follows as a consequence of Remark 3.2 that the set of

all possible irreducible character degrees is $\{1, p\}$. Now, suppose that $\text{cd}(G) = \{1, p\}$. Since G is a finite p -group with cyclic center, it follows from Theorem 3.1 that $\delta_{\text{irr}}(G)$ exists and $\delta(G) = \delta_{\text{irr}}(G)$. Consequently, $\delta(G) = \delta_{\text{irr}}(G) = p$.

- (ii) If $\text{cd}(G) = \{1, p^a\}$, then it follows from Theorem 3.1 that $\delta(G) = \delta_{\text{irr}}(G) = p$. If $\text{cd}(G) = \{1, p, p^a\}$, then it immediately follows from Lemma 21 of [33] that $\delta(G) = p$.

□

4. REPRESENTATION DIMENSION OF THE DIRECT PRODUCT OF GROUPS

In section 3, we have proved that if G is a p -group with cyclic center, then $\delta(G) = \delta_{\text{irr}}(G)$. For a nilpotent group, a necessary and sufficient condition for the existence of a faithful irreducible character is that its center is cyclic. Using the GAP codes given in Appendix A, we have checked that for all nilpotent groups with cyclic center and of order up to 100, $\delta(G) = \delta_{\text{irr}}(G)$. This raises a question: *For a nilpotent group G with cyclic center, do we necessarily have $\delta(G) = \delta_{\text{irr}}(G)$?* In this section, we provide an affirmative answer to this question.

Theorem 4.1. *Let G_1 and G_2 be two finite nonabelian groups. Suppose one of the following holds:*

- (1) G_1 and G_2 are simple groups;
- (2) G_1 and G_2 are p -groups with cyclic center;
- (3) G_1 and G_2 are monolithic groups, i.e., groups with a unique minimal normal subgroup;
- (4) G_1 and G_2 have unique nonlinear irreducible character.

If $Z(G_1)$ and $Z(G_2)$ are of coprime order, then $\delta(G_1 \times G_2) \leq \delta(G_1) + \delta(G_2)$ and $\delta_{\text{irr}}(G_1 \times G_2) = \delta_{\text{irr}}(G_1)\delta_{\text{irr}}(G_2)$. Further, if G_1 and G_2 are of coprime order, then $\delta(G_1 \times G_2) = \delta_{\text{irr}}(G_1 \times G_2) = \delta_{\text{irr}}(G_1)\delta_{\text{irr}}(G_2)$.

Proof. We first prove that in each of the following cases, $\delta(G_i) = \delta_{\text{irr}}(G_i)$, for $i = 1, 2$.

Case (1). G_1 and G_2 are simple groups.

Since G_i is a simple group, for $i = 1, 2$, we have that any nonlinear character must be faithful. Hence $\delta(G_i) = \delta_{\text{irr}}(G_i)$.

Case (2). G_1 and G_2 are p -groups with cyclic center.

It immediately follows from Theorem 3.1(1) that $\delta(G_i) = \delta_{\text{irr}}(G_i)$, for $i = 1, 2$.

Case (3). G_1 and G_2 are monolithic groups.

Let G be a monolithic group and let χ be a faithful character of G with $\deg(\chi) = \delta(G)$. We write $\chi = \sum \eta_i \psi_i$, where ψ_i are irreducible characters of G . Now, $\{1\} = \ker(\chi) = \cap \ker(\psi_i)$, where $\ker(\chi)$ denotes the kernel of the character χ . We observe that

$\ker(\psi_j) = \{1\}$ for some j , else $\ker(\chi)$ would contain the minimal normal subgroup of G , contradicting that χ is faithful. Thus, $\delta(G) \leq \deg(\psi_j) \leq \deg(\chi) = \delta(G)$. Hence, χ must be a faithful irreducible character of G . Consequently, $\delta(G_i) = \delta_{irr}(G_i)$, for $i = 1, 2$.

Case (4). G_1 and G_2 have unique nonlinear irreducible character.

In view of Lemma 1.4 of [19], $\delta(G_i)$, for $i = 1, 2$, is the degree of the unique nonlinear irreducible character and hence the result follows.

In all the above cases, it is proved that $\delta(G_i) = \delta_{irr}(G_i)$, for $i = 1, 2$. Defining a representation on $G_1 \times G_2$ by block diagonal matrices yields that $\delta(G_1 \times G_2) \leq \delta(G_1) + \delta(G_2)$. If the orders of $Z(G_1)$ and $Z(G_2)$ are relatively coprime, then $\delta_{irr}(G_1 \times G_2) = \delta_{irr}(G_1)\delta_{irr}(G_2)$, due to Problem 4.3 of [20].

Suppose that G_1 and G_2 are of coprime order. If χ is a minimal faithful character of $G_1 \times G_2$, then $\chi = \chi_1 + \chi_2 + \cdots + \chi_n$, where $\chi_i \in \text{Irr}(G_1 \times G_2)$, for $1 \leq i \leq n$. Let $\chi_i = \tau_i\psi_i$, for $1 \leq i \leq n$, where $\tau_i \in \text{Irr}(G_1)$ and $\psi_i \in \text{Irr}(G_2)$. Since G_1 and G_2 are of coprime order, $\langle (g_1, g_2) \mid g_1 \in \bigcap_{1 \leq i \leq n} \ker(\tau_i), g_2 \in \bigcap_{1 \leq i \leq n} \ker(\psi_i) \rangle \leq \bigcap_{1 \leq i \leq n} \ker(\tau_i\psi_i) = \ker(\chi) = \{1\}$. This implies that $\langle (g_1, g_2) \mid g_1 \in \bigcap_{1 \leq i \leq n} \ker(\tau_i), g_2 \in \bigcap_{1 \leq i \leq n} \ker(\psi_i) \rangle = \{1\}$. Hence $\tau_1 + \tau_2 + \cdots + \tau_n$ is a faithful character of G_1 and $\psi_1 + \psi_2 + \cdots + \psi_n$ is a faithful character of G_2 . If $\tau \in \text{Irr}(G_1)$ and $\psi \in \text{Irr}(G_2)$ such that $\deg(\tau) = \delta_{irr}(G_1) = \delta(G_1)$ and $\deg(\psi) = \delta_{irr}(G_2) = \delta(G_2)$, then $\tau\psi \in \text{Irr}(G_1 \times G_2)$ is a minimal faithful character of $G_1 \times G_2$. Therefore, $\delta(G_1 \times G_2) = \delta_{irr}(G_1)\delta_{irr}(G_2)$. \square

Corollary 4.2. *If G is a finite nilpotent group with cyclic center, then $\delta(G)$ equals $\delta_{irr}(G)$.*

Proof. Since G is a nilpotent group, it is a direct product of Sylow p -subgroups. Due to Theorem 4.1, it follows that $\delta(G) = \delta_{irr}(G)$. \square

Remark 4.3. In view of Theorem 4.1, it follows that $\delta(A_4 \times S_3) = 5$ and $\delta_{irr}(A_4 \times S_3) = 6$. This is the smallest example of a finite group G with cyclic center such that $\delta(G) \neq \delta_{irr}(G)$.

Let G be a finite nilpotent group with cyclic center. We can write $G = P_1 \times P_2 \times \cdots \times P_n$, where P_i 's are Sylow subgroups of G . Due to Theorem 4.1, $\delta(G) = \delta_{irr}(P_1)\delta_{irr}(P_2) \cdots \delta_{irr}(P_n)$. Consequently, from Theorem 3.3, we have the following:

Theorem 4.4. *Let G be a finite nilpotent group with cyclic center.*

- (i) $\delta(G) = p$ if and only if $\text{cd}(G) = \{1, p\}$.
- (ii) If for some integer $a > 1$, $\text{cd}(G) = \{1, p^a\}$ or $\text{cd}(G) = \{1, p, p^a\}$, then $\delta(G) = p^a$.

5. REPRESENTATION DIMENSION OF GROUPS WITH FEW IRREDUCIBLE CHARACTERS

Chillag and Herzog [10] provided a complete classification of nonabelian groups of odd order having exactly two nonlinear irreducible characters of each degree. We can use this classification to determine their representation dimensions.

Theorem 5.1. *Let G be a finite nonabelian group of odd order with exactly two nonlinear irreducible characters of each degree. Then,*

- (i) $\delta(G) = \delta_{irr}(G) = 3^r$, if G is an extraspecial 3-group of order 3^{2r+1} , for some integer $r \geq 1$.
- (ii) $\delta(G) = \delta_{irr}(G) = \frac{p^n-1}{2}$, if G is a Frobenius group of odd order $\frac{(p^n-1)p^n}{2}$ for some odd prime p and some integer n , with cyclic kernel K of order p^n and cyclic complement. In case, K is abelian but non cyclic, then $\delta(G) = \frac{(p^n-1)}{2} \text{rank}(K)$, whereas $\delta_{irr}(G)$ does not exist.

Proof. From Theorem 1 of [10], a non abelian group G of odd order which has exactly two non linear irreducible characters of each degree, is one of the following:

- (i) an extraspecial 3-group of order 3^{2r+1} , for some integer $r \geq 1$; or
- (ii) a Frobenius group of odd order $\frac{(p^n-1)p^n}{2}$ for some odd prime p and some integer n , with abelian Frobenius kernel K of order p^n and cyclic Frobenius complement.

Clearly, (i) follows from Theorem 3.1(6). For (ii), first consider the case when the Frobenius kernel K of the Frobenius group G is cyclic, and so is its Frobenius complement. We denote the Frobenius complement by H . It follows from Theorem 37.5.1 of [25] that the irreducible characters of G are $\chi_1^G, \chi_2^G, \dots, \chi_r^G, \lambda_1, \lambda_2, \dots, \lambda_s$, where $\{\chi_1, \chi_2, \dots, \chi_r\}$ is a complete set of representatives of G -conjugacy classes of non principal irreducible characters of K , χ_i^G are their induced representations on G , for $1 \leq i \leq n$, and $\lambda_1, \lambda_2, \dots, \lambda_s$ are extensions of irreducible characters of H to G . Clearly, λ_i 's are linear non faithful characters, for $1 \leq i \leq s$. Note that the degree of each nonlinear irreducible character of G is $[G : K]$. Since K is cyclic, G has a nonlinear irreducible character which is faithful of degree $[G : K]$, i.e., $\frac{p^n-1}{2}$. Since it is the minimal nonlinear irreducible character degree, it immediately follows that $\delta(G) = \delta_{irr}(G) = \frac{p^n-1}{2}$.

Now, suppose that the Frobenius kernel K of G is abelian but non cyclic and its complement H is cyclic. Since K is abelian, it is a direct product of cyclic groups. Hence, each subgroup of K is normal in G . Consequently, for $1 \leq i \leq n$, $\ker(\chi_i^G) = \text{core}_G(\ker(\chi_i)) = \ker(\chi_i)$, where $\text{core}_G(\ker(\chi_i))$ denotes the largest normal subgroup of G contained in $\ker(\chi_i)$. As K is non cyclic, the Frobenius kernel K has no faithful irreducible character. Hence, the group G does not have a faithful irreducible character. Further, $K \leq \ker(\lambda_i)$, for $1 \leq i \leq s$. Therefore, $\delta(G) = [G : K]\delta(K) = \frac{(p^n-1)}{2} \text{rank}(K)$. This completes the proof of the theorem. \square

Theorem 2.3 of [13] provides the classification of finite metabelian groups (nonabelian) of odd order with the property that any two non-principal irreducible characters of the same degree are Galois conjugate. Once again, we can compute the representation dimension for such groups using this classification.

Theorem 5.2. *Let G be a finite metabelian group of odd order with the property that any two non-principal irreducible characters of the same degree are Galois conjugate. Then,*

- (i) $\delta(G) = \delta_{irr}(G) = p$, where $[G : K] = p$ is a prime, if the group G is a Frobenius group with Frobenius complement of order p and Frobenius kernel K is cyclic group of prime order q , where $q \neq p$.
- (ii) $\delta(G) = \text{rank}(K) \cdot p$ and $\delta_{irr}(G)$ does not exist, if G is a Frobenius group with complement of prime order p and the Frobenius kernel K is an elementary abelian q -group of order q^b , where $q \neq p$ is a prime and b is also a prime with $p = \frac{q^b - 1}{q - 1}$.

6. REPRESENTATION DIMENSION FOR GROUPS IN WHICH THE DEGREES OF THE NONLINEAR IRREDUCIBLE CHARACTERS ARE DISTINCT

Let G be a finite group whose all nonlinear irreducible characters have distinct degrees. In this section, we compute the representation dimensions for such groups. These groups are classified by Berkovich, Chillag, and Herzog in ([5], pg.955) and the explicit description is as follows.

Theorem (Berkovich-Chillag-Herzog). *If G is a nonabelian finite group whose all non-linear irreducible characters have distinct degrees. Then, one of the following holds:*

- (1) G is an extraspecial 2-group.
- (2) G is a Frobenius group of order $p^n(p^n - 1)$ for some prime power p^n with an abelian Frobenius kernel of order p^n and a cyclic Frobenius complement.
- (3) G is a Frobenius group of order 72 in which the Frobenius complement is isomorphic to the quaternion group of order 8.

Now, we have the following:

Theorem 6.1. *Let G be a nonabelian finite group whose all non-linear irreducible characters have distinct degrees. Then, $\delta(G) = \delta_{irr}(G)$, and it equals,*

- (i) 2^r , if G is an extraspecial 2-group of order 2^{2r+1} , for some integer $r \geq 1$.
- (ii) $p^n - 1$, if G is a Frobenius group of order $p^n(p^n - 1)$ for some prime power p^n with an abelian Frobenius kernel of order p^n and a cyclic Frobenius complement.
- (iii) 8, if G is a Frobenius group of order 72 in which the Frobenius complement is isomorphic to the quaternion group.

Proof. Let G be a nonabelian finite group whose all nonlinear irreducible characters have distinct degrees. Then, due to the above theorem, we can deal with such groups case by case.

Case 1. G is an extraspecial 2-group.

In this case, $\delta(G) = \delta_{irr}(G) = 2^r$, in view of Theorem 3.1(6).

Case 2. G is a Frobenius group of order $p^n(p^n - 1)$ for some prime power p^n with an abelian Frobenius kernel K of order p^n and a cyclic Frobenius complement H .

It follows from Theorem 5.1 of [25] that all the irreducible characters of G are obtained from the extensions of the irreducible characters of H , and the rest irreducible characters of G are induced from the representatives of G -conjugacy classes of non principal irreducible characters of K . As K is abelian, the degree of each nonlinear irreducible character is $[G : K]$. By assumption, all non-linear irreducible characters of the group G have distinct degrees. This yields that there is a unique non-linear irreducible character of degree $[G : K]$, which is faithful, due to Lemma 1.4 of [19]. Consequently, $\delta(G) = \delta_{irr}(G) = p^n - 1$.

Case 3. G is a Frobenius group of order 72 in which the Frobenius complement is isomorphic to the quaternion group.

From Lemma 1.4 of [19], it follows that G has a unique nonlinear irreducible faithful character of degree 8. This completes the proof of the theorem. \square

Iranmanesh and Saiedi [19, Lemma 1.4] proved that if G is a finite group with a unique non-linear irreducible character, then it is faithful. In view of the above theorem, we have the following:

Corollary 6.2. *If G is a finite group with unique non-linear irreducible character, then $\delta(G) = \delta_{irr}(G)$ is*

- (i) 2^r , if G is an extraspecial 2-group of order 2^{2r+1} .
- (ii) $p^n - 1$, if G is a Frobenius group of order $p^n(p^n - 1)$ for some prime power p^n with an abelian Frobenius kernel of order p^n and a cyclic Frobenius complement.

Using the same arguments as in Theorem 4.1, the following corollary arises:

Corollary 6.3. *The representation dimension of the direct product of finite groups G of relatively coprime order with the property that all non-linear irreducible characters of G have distinct degree equals the minimal faithful irreducible character degree.*

APPENDIX A. GAP CODE TO COMPUTE $\delta(G)$ AND $\delta_{irr}(G)$

In this section, we present a GAP code [16] to determine the representation dimension and minimal faithful irreducible character degree.

Given a finite group G , we have a GAP code in Algorithm 1 to compute $\delta_{irr}(G)$. The first step of the algorithm calls in the character table of G and the number of conjugacy classes of G . Denote by M , the list of irreducible characters of G with their values on conjugacy classes. In this list, for each irreducible character of G , we compute its kernel K and add $[i, K]$ to the list L . Initially, L is taken to be $[\]$. From L , list those irreducible characters whose kernel is $\{e\}$ and denote the list by F . Consider the list D of degrees of irreducible characters in F , i.e., the degrees of irreducible faithful characters of G . If D is empty, then output is 0, i.e., $\delta_{irr}(G)$ does not exist. If D is nonempty, then the minimum of all the irreducible faithful character degrees $\delta_{irr}(G)$ is returned.

```

Data: A finite group  $G$ .
 $c :=$ CharacterTable( $G$ );
 $CC :=$  ConjugacyClasses( $G$ );
 $s :=$  Size( $CC$ );
 $M :=$  List(Irr( $c$ ), ValuesOfClassFunction);
 $L := [\ ]$ ;
for  $i$  in  $[1..s]$  do
   $K := [\ ]$ ;
  for  $j$  in  $[1..s]$  do
    if  $M[i][j] = M[i][1]$  then
      | Add( $K, j$ );
    end
  end
  Add( $L, [i, K]$ );
   $F :=$  Filtered( $L, x \rightarrow$  Length( $x[2]$ ) = 1);
   $D := [\ ]$ ;
   $d := 0$ ;
  for  $j$  in  $F$  do
    | Add( $D, M[j[1]][1]$ );
  end
  if not(IsEmpty( $D$ )) then
    |  $d :=$  Minimum( $D$ );
  end
  return  $d$ ;
end
Result:  $\delta_{irr}(G)$ 

```

Algorithm 1: Minimal irreducible faithful character degree of G

Next, we provide GAP code Algorithm 5 to determine the representation dimension of G . For this, we first give Algorithm 2 to compute a list of kernels of all the irreducible complex characters of G . In this algorithm, the first step is to call its character table which lists all the irreducible characters of G with its values on conjugacy classes. Denote this list by M . Suppose there are s irreducible characters in the list, i.e., s number of conjugacy classes. For the i th irreducible character $M[i]$ mentioned in the list, look for that $j \in [1 \cdots s]$ for which the value of $M[i]$ on j th conjugacy class coincides with its

degree, i.e., $M[i][j] = M[i][1]$. The union of all these conjugacy classes is the kernel K of $M[i]$. Repeat this process for each irreducible character. Suppose $L = []$. In the list L , add $[i, K]$ for each $i \in [1 \cdots s]$. The list L is the desired one.

```

Data: A finite group  $G$ .
ListKernels:= function( $M$ );
 $s := \text{Size}(M)$ ;
 $L := [ ]$ ;
for  $i$  in  $[1..s]$  do
   $K := [ ]$ ;
  for  $j$  in  $[1..s]$  do
    if  $M[i][j] = M[i][1]$  then
      |  $\text{Add}(K, j)$ ;
    end
  end
   $\text{Add}(L, [i, K])$ ;
end
return  $L$ ;
Result: Returns a list consisting of  $[\chi, \ker(\chi)]$ , where  $\chi$  is an irreducible character of  $G$ 

```

Algorithm 2: List of all the irreducible characters of G and their kernel

The next step is to provide Algorithm 3 for determining all the possible kernels of characters of G and Algorithm 4 to determine the degree corresponding to each character of G in the above list.

```

Data: A finite group  $G$ .
KernelIntersection := function( $L, S$ );
 $S := \text{Size}(L)$ ;
IntKer:=  $[1..S]$ ;
for  $i$  in  $S$  do
  | IntKer := Intersection(IntKer,  $L[i][2]$ );
end
return IntKer;
Result: Returns the intersection of  $\ker(\chi)$  of characters  $\chi$ , where  $i$  lies in the set  $S$ .

```

Algorithm 3: The set of all possible kernels of characters of a group G

```

Data: A finite group  $G$ .
DegreeSum := function( $M, S$ );
 $d := 0$ ;
for  $i$  in  $S$  do
  |  $d := d + M[i][1]$ ;
end
return  $d$ ;
Result: Returns the sum of degrees of characters  $\chi_i$ , where  $i$  lies in the set  $S$ .

```

Algorithm 4: Degrees of all possible characters of group G

```

Data: A finite group  $G$ .
Representation dimension:= function( $G$ );
 $c :=$  CharacterTable( $G$ );
 $CC:=$  Conjugacy classes( $G$ );
 $s:=$ Size( $CC$ );
 $M :=$  List(Irr( $c$ ),ValuesOfClassFunction);
 $L :=$  ListKernels( $M$ );
minDeg := Sum(TransposedMat( $M$ )[1]);
minK := [1.. $s$ ];
rList := [1.. $s - 1$ ];
 $r := 1$ ;
while  $r \leq$  minDeg do
  rCombos := Combinations([1.. $s$ ],  $r$ );
  for  $x$  in rCombos do
    DegSumx := DegreeSum( $M$ ,  $x$ );
    if DegSumx < minDeg and  $r <$  minDeg then
      if Length(KernelIntersection( $L$ , $x$ )) = 1 then
        minDeg := DegSumx;
        minK :=  $x$ ;
      end
    end
  end
   $r := r+1$ ;
  Print("Representation dimension is : ");
end
return minDeg;
Result:  $\delta(G)$ 

```

Algorithm 5: Representation dimension of G

In the above algorithm, we use previous algorithms to determine the representation dimension of a group G . Clearly, the representation dimension of group G is less than or equal to the degree of the regular character of G , denoted by MinDeg. Now, we look at the possible combination of irreducible characters which yield a faithful character whose degree is less than MinDeg. Suppose a given combination is providing a faithful character χ with its degree being less than MinDeg. Then, repeat the process with MinDeg being replaced by $\deg(\chi)$. This process will stop at a finite number of steps. Finally, the representation dimension of G is determined.

Acknowledgment. We thank B. Sury for his feedback on this paper which helped us improve it. The first named author acknowledges the research support of the National Board for Higher Mathematics, Department of Atomic Energy, Govt. of India (NBHM Reference No.: 0204/16(7)/2022/R&D-II/ 11978). The third named author is funded by NBHM through 02011/23/2023/NBHM(RP)/RDII/5955 for this research.

REFERENCES

1. Mohammad Bardestani, Keivan Mallahi-Karai, and Hadi Salmasian, *Minimal dimension of faithful representations for p -groups*, J. Group Theory **19** (2016), no. 4, 589–608. MR 3518394
2. B. G. Basmaji, *Monomial representations and metabelian groups*, Nagoya Math. J. **35** (1969), 99–107. MR 244394
3. T. R. Berger, P. Landrock, and G. O. Michler, *Minimal degrees of faithful characters of finite groups with a T.I. Sylow p -subgroup*, Proc. Amer. Math. Soc. **99** (1987), no. 1, 15–21. MR 866421
4. Yakov Berkovich, *Groups of prime power order. Vol. 1*, De Gruyter Expositions in Mathematics, vol. 46, Walter de Gruyter GmbH & Co. KG, Berlin, 2008, With a foreword by Zvonimir Janko. MR 2464640
5. Yakov Berkovich, David Chillag, and Marcel Herzog, *Finite groups in which the degrees of the nonlinear irreducible characters are distinct*, Proc. Amer. Math. Soc. **115** (1992), no. 4, 955–959. MR 1088438
6. Christine Bessenrodt, Hung P. Tong-Viet, and Jiping Zhang, *Huppert's conjecture for alternating groups*, J. Algebra **470** (2017), 353–378. MR 3565438
7. J. C. Bioch, *On n -isoclinic groups*, Nederl. Akad. Wetensch. Proc. Ser. A **79**=Indag. Math. (1976), no. no. 5., 400–407. MR 424937
8. J. Buhler and Z. Reichstein, *On the essential dimension of a finite group*, Compositio Math. **106** (1997), no. 2, 159–179. MR 1457337
9. Shane Cernele, Masoud Kamgarpour, and Zinovy Reichstein, *Maximal representation dimension of finite p -groups*, J. Group Theory **14** (2011), no. 4, 637–647. MR 2818953
10. David Chillag and Marcel Herzog, *Finite groups with almost distinct character degrees*, J. Algebra **319** (2008), no. 2, 716–729. MR 2381804
11. L. Di Martino and M. C. Tamburini, *Some remarks on the degrees of faithful irreducible representations of a finite p -group*, Geom. Dedicata **41** (1992), no. 2, 155–164. MR 1153979
12. ———, *Some remarks on the degrees of faithful irreducible representations of a finite p -group*, Geom. Dedicata **41** (1992), no. 2, 155–164. MR 1153979
13. Silvio Dolfi, Gabriel Navarro, and Pham Huu Tiep, *Finite groups whose same degree characters are Galois conjugate*, Israel J. Math. **198** (2013), no. 1, 283–331. MR 3096641
14. Gustavo A. Fernández-Alcober and Alexander Moretó, *Groups with two extreme character degrees and their normal subgroups*, Trans. Amer. Math. Soc. **353** (2001), no. 6, 2171–2192. MR 1814066
15. D. L. Flannery and E. A. O'Brien, *Linear groups of small degree over finite fields*, Internat. J. Algebra Comput. **15** (2005), no. 3, 467–502. MR 2151423
16. The GAP Group, *Gap – groups, algorithms, and programming, version 4.12.2*, 2022.
17. J. A. Green, *The characters of the finite general linear groups*, Trans. Amer. Math. Soc. **80** (1955), 402–447. MR 72878
18. Guan Aun How, *Special classes of monomial groups. III*, Chinese J. Math. **12** (1984), no. 3, 199–211. MR 764985
19. Ali Iranmanesh and Amin Saeidi, *Finite groups with a unique nonlinear nonfaithful irreducible character*, Arch. Math. (Brno) **47** (2011), no. 2, 91–98. MR 2813535
20. I. Martin Isaacs, *Character theory of finite groups*, Pure and Applied Mathematics, No. 69, Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1976. MR 0460423
21. Gordon Douglas James, *Representations of general linear groups*, vol. 94, Cambridge University Press, 1984.

22. Rodney James, *The groups of order p^6 (p an odd prime)*, Math. Comp. **34** (1980), no. 150, 613–637. MR 559207
23. Christoph Jansen, *The minimal degrees of faithful representations of the sporadic simple groups and their covering groups*, LMS J. Comput. Math. **8** (2005), 122–144. MR 2153793
24. Nikita A. Karpenko and Alexander S. Merkurjev, *Essential dimension of finite p -groups*, Invent. Math. **172** (2008), no. 3, 491–508. MR 2393078
25. Gregory Karpilovsky, *Group representations. Vol. 1. Part B*, North-Holland Mathematics Studies, vol. 175, North-Holland Publishing Co., Amsterdam, 1992, Introduction to group representations and characters. MR 1183468
26. Gustav Isaac Lehrer, *The characters of the finite special linear groups*, J. Algebra **26** (1973), 564–583. MR 354889
27. Frank Lübeck, *Smallest degrees of representations of exceptional groups of Lie type*, Comm. Algebra **29** (2001), no. 5, 2147–2169. MR 1837968
28. G. Lusztig, *Irreducible representations of finite classical groups*, Invent. Math. **43** (1977), no. 2, 125–175. MR 463275
29. Avinoam Mann, *Minimal characters of p -groups*, J. Group Theory **2** (1999), no. 3, 225–250. MR 1696312
30. ———, *Minimal characters of p -groups*, J. Group Theory **2** (1999), no. 3, 225–250. MR 1696312
31. Alexander Moretó, *On the minimal dimension of a faithful linear representation of a finite group*, arXiv preprint arXiv:2102.01463 (2021).
32. S. K. Prajapati, M. R. Darafsheh, and M. Ghorbani, *Irreducible characters of p -group of order $\leq p^5$* , Algebr. Represent. Theory **20** (2017), no. 5, 1289–1303. MR 3707916
33. Sunil Kumar Prajapati and Ayush Udeep, *Minimal faithful quasi-permutation representation degree of p -groups with cyclic center*, arXiv preprint arXiv:2302.06257 (2023).
34. Richard Rasala, *On the minimal degrees of characters of S_n* , J. Algebra **45** (1977), no. 1, 132–181. MR 427445
35. Robert Steinberg, *The representations of $GL(3, q)$, $GL(4, q)$, $PGL(3, q)$, and $PGL(4, q)$* , Canad. J. Math. **3** (1951), 225–235. MR 41851
36. Fernando Szechtman, *Groups having a faithful irreducible representation*, J. Algebra **454** (2016), 292–307. MR 3473429
37. Pham Huu Tiep and Alexander E. Zalesskii, *Minimal characters of the finite classical groups*, Comm. Algebra **24** (1996), no. 6, 2093–2167. MR 1386030

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH MOHALI, KNOWLEDGE CITY, SECTOR 81, MOHALI 140 306, INDIA

Email address: gurleen@iisermohali.ac.in

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH MOHALI, KNOWLEDGE CITY, SECTOR 81, MOHALI 140 306, INDIA

Email address: amitk@iisermohali.ac.in

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH PUNE, DR. HOMI BHABHA ROAD, PASHAN, PUNE 411 008, INDIA

Email address: anupamk18@gmail.com