

# Isomeric pair $^{95\text{m,g}}\text{Nb}$ in photonuclear reactions on $^{\text{nat}}\text{Mo}$ at end-point bremsstrahlung energy of 35–95 MeV

I.S. Timchenko<sup>1,2,\*</sup> and O.S. Deiev<sup>2</sup>, S.M. Olejnik<sup>2</sup>, S.M. Potin<sup>2</sup>,  
V.A. Kushnir<sup>2</sup>, V.V. Mytrochenko<sup>2,3</sup>, S.A. Perezhogin<sup>2</sup>, V.A. Bocharov<sup>2</sup>

<sup>1</sup> *Institute of Physics, Slovak Academy of Sciences, SK-84511 Bratislava, Slovakia*

<sup>2</sup> *National Science Center "Kharkov Institute of Physics and Technology",  
1, Akademichna St., 61108, Kharkiv, Ukraine and*

<sup>3</sup> *CNRS/IJCLAB, 15 Rue Georges Clemenceau Str., Orsay, 91400, France*

(Dated: August 7, 2023)

The  $^{\text{nat}}\text{Mo}(\gamma, xnp)^{95\text{m,g}}\text{Nb}$  photonuclear reaction was studied using the electron beam from the NSC KIPT linear accelerator LUE-40. Experiment was performed using the activation and off-line  $\gamma$ -ray spectrometric technique. The experimental isomeric yield ratio  $d(E_{\gamma\text{max}}) = Y_{\text{m}}(E_{\gamma\text{max}})/Y_{\text{g}}(E_{\gamma\text{max}})$  was determined for the reaction products  $^{95\text{m,g}}\text{Nb}$  at the end-point bremsstrahlung energy  $E_{\gamma\text{max}}$  range of 35–95 MeV. The obtained values of  $d(E_{\gamma\text{max}})$  are in satisfactory agreement with the results of other authors and extend the range of previously known data. The theoretical values of the yields  $Y_{\text{m,g}}(E_{\gamma\text{max}})$  and the isomeric yield ratio  $d(E_{\gamma\text{max}})$  for the isomeric pair  $^{95\text{m,g}}\text{Nb}$  from the  $^{\text{nat}}\text{Mo}(\gamma, xnp)$  reaction were calculated using the partial cross-sections  $\sigma(E)$  from the TALYS1.95 code for six different level density models  $LD$ . The comparison showed a noticeable excess (more than 3.85 times) of the experimental isomeric yield ratio over all theoretical estimates. At the investigated range of  $E_{\gamma\text{max}}$  the theoretical dependence of  $d(E_{\gamma\text{max}})$  on energy was confirmed – the isomeric yield ratio smoothly decreases with increasing energy.

Keywords: photonuclear reactions,  $^{\text{nat}}\text{Mo}$ , isomeric pair  $^{95\text{m,g}}\text{Nb}$ , isomeric yield ratio, reaction yield, end-point bremsstrahlung energies of 35–95 MeV, activation and off-line  $\gamma$ -ray spectrometric technique, TALYS1.95, level density model, GEANT4.9.2.

## I. INTRODUCTION

Photonuclear reactions are accompanied by the emission of a nucleon or a group of nucleons from a compound nucleus. This leads to an excited state of the final nucleus, the discharge of excitation energy of which occurs in a time of  $10^{-12}$ – $10^{-17}$  sec. However, in some cases, at low energy of the excitation level and a high degree of forbidden transition, long-lived excited states of atomic nuclei occur. Such states are called isomeric or metastable states, and their half-lives can vary from nanoseconds to many years [1].

Nuclei with isomeric (m) and unstable ground (g) states are of particular interest, since they allow to study of the metastable state population of this nucleus relative to its ground state, i.e. obtain the isomeric ratio of the reaction products. This characteristic is defined as the ratio of the cross-sections for the formation of the reaction product in the metastable  $\sigma_{\text{m}}(E)$  and in the ground  $\sigma_{\text{g}}(E)$  states:  $d(E) = \sigma_{\text{m}}(E)/\sigma_{\text{g}}(E)$ . The  $d(E)$  values in the literature can be found as the ratio of the cross-sections for the formation of a product nucleus in the high-spin state (as a rule, this is a metastable state)  $\sigma_{\text{H}}(E)$  to the cross-section for the low-spin state  $\sigma_{\text{L}}(E)$ :  $d(E) = \sigma_{\text{H}}(E)/\sigma_{\text{L}}(E)$  [2–4]. In the case of research with the use of bremsstrahlung flux, this ratio  $d(E_{\gamma\text{max}})$  is expressed through the bremsstrahlung flux-averaged cross-

sections or yields ( $\langle\sigma(E_{\gamma\text{max}})\rangle$  or  $Y(E_{\gamma\text{max}})$ ) of the reactions under study [4–6].

Data on isomeric ratios of reaction products make it possible to investigate issues related to nuclear reactions and nuclear structure, such as the spin dependence of the nuclear level density, angular momentum transfer, nucleon pairing, shell effects, refine the theory of gamma transitions, and test theoretical models of the nucleus [7–11]. The study of isomeric ratios using photonuclear reactions has an advantage since the  $\gamma$ -quantum introduces a small angular momentum and does not change the nucleon composition of the compound nucleus.

The studies in the energy range above giant dipole resonance (GDR) and before the pion production threshold (30–145 MeV) are of interest, because the mechanism of the nuclear reaction changes here: from dominance of GDR to dominance of the quasideuteron mechanism [12]. However, in this energy range, there is still a lack of experimental data on the cross-sections of multiparticle photonuclear reactions and isomeric ratios of the products of a nuclear reaction [13, 14]. This complicates the analysis of the mechanism of the nuclear reaction, as well as the systematization and analysis of the dependences of these quantities on various characteristics of the nucleus.

Experiments on the photodisintegration of stable isotopes of the Mo nucleus and determination of the isomeric ratio values  $d(E_{\gamma\text{max}})$  of the nuclide products  $^{95\text{m,g}}\text{Nb}$  from the  $^{\text{nat}}\text{Mo}(\gamma, xnp)$  reaction were performed in works [15–22], using beams of bremsstrahlung photons and the residual  $\gamma$ -activity method.

In the energy region of GDR, the formation of the isomeric pair  $^{95\text{m,g}}\text{Nb}$  in the photonuclear reaction

\* timchenko@kipt.kharkov.ua;  
iryana.timchenko@savba.sk

on  $^{\text{nat}}\text{Mo}$  has been studied in works [15–19]. Thus, in [15, 16] the experimental result for  $d(E_{\gamma_{\text{max}}}) = Y_{\text{m}}(E_{\gamma_{\text{max}}})/(Y_{\text{g}}(E_{\gamma_{\text{max}}})+Y_{\text{m}}(E_{\gamma_{\text{max}}}))$  was obtained at an energy of 30 MeV. The authors of [17] defined the values of  $d(E_{\gamma_{\text{max}}})$  as the ratio of  $Y_{\text{m}}(E_{\gamma_{\text{max}}})/Y_{\text{g}}(E_{\gamma_{\text{max}}})$  at energies of 25 and 30 MeV. In [18, 19] the range of investigated energies was extended to 14–24 MeV and the isomeric ratios of the reaction products of the  $^{\text{nat}}\text{Mo}(\gamma, xnp)^{95\text{m,g}}\text{Nb}$  reaction were also obtained by the  $\gamma$ -activation method. However, the results of [18, 19] contradict each other. It should be noted in works [15–19] in obtained experimental results the main role in the formation of the isomeric pair  $^{95\text{m,g}}\text{Nb}$  on the  $^{\text{nat}}\text{Mo}$  plays the isotope  $^{96}\text{Mo}$  due to the reaction on the  $^{97}\text{Mo}$  nucleus has the higher threshold energy.

At an intermediate energy region the  $^{\text{nat}}\text{Mo}(\gamma, xnp)^{95\text{m,g}}\text{Nb}$  reaction was investigated in works [20–22]. In [20] the values of  $d(E_{\gamma_{\text{max}}}) = Y_{\text{H}}(E_{\gamma_{\text{max}}})/Y_{\text{L}}(E_{\gamma_{\text{max}}})$  were obtained based on  $(\gamma, p)$ ,  $(\gamma, np)$ , and  $(\gamma, 2np)$  reactions, which was the first measurement with  $^{\text{nat}}\text{Mo}$  targets at  $E_{\gamma_{\text{max}}} = 50, 60, 70$  MeV. In [21] authors extended the range of the study up to 45–70 MeV and attempted to compare the independent isomeric yield ratios of  $^{95\text{m,g}}\text{Nb}$  for different reactions:  $^{\text{nat}}\text{Mo}(\gamma, xnp)$ ,  $^{\text{nat}}\text{Mo}(p, \alpha xn)$ , and  $^{\text{nat}}\text{Zr}(p, xn)$ .

In work [22] the experimentally and theoretically studying the photodisintegration of molybdenum isotopes were performed. The yields of various photonuclear reactions on stable molybdenum isotopes were determined by the  $\gamma$ -activation method for the end-point bremsstrahlung energies of 19.5, 29.1, and 67.7 MeV. For the energy of 67.7 MeV, authors were able to determine the yields  $Y_{\text{m}}(E_{\gamma_{\text{max}}})$  and  $Y_{\text{g}}(E_{\gamma_{\text{max}}})$ , respectively, for the formation of the  $^{95\text{m,g}}\text{Nb}$  nucleus in the isomeric and ground states, whose ratio is 5.5/5.1.

The experimental results from [20–22] agree with each other, while the data from [15–19] obtained in different representations of the value  $d(E_{\gamma_{\text{max}}})$  contradict each other. This does not allow us to estimate the energy dependence of  $d(E_{\gamma_{\text{max}}})$  in the wide region  $E_{\gamma_{\text{max}}} = 14$ –70 MeV.

The present work is devoted to the study of the formation of the isomeric pair  $^{95\text{m,g}}\text{Nb}$  in photonuclear reactions on  $^{\text{nat}}\text{Mo}$  at the end-point bremsstrahlung energy range of  $E_{\gamma_{\text{max}}} = 35$ –95 MeV. This will extend the range of previously known experimental isomeric yield ratios  $d(E_{\gamma_{\text{max}}})$  and improve the reliability of these data.

## II. EXPERIMENTAL SETUP AND PROCEDURE

An experimental study of the  $^{95\text{m,g}}\text{Nb}$  isomeric pair formation in the photonuclear reactions on  $^{\text{nat}}\text{Mo}$  was performed using the electron beam of the linac LUE-40 RDC "Accelerator" NSC KIPT with the activation and off-line  $\gamma$ -ray spectrometric technique. The experimental

procedure is described in detail, for example, in [23–26].

The experimental complex for the study of photonuclear reactions is presented as a block diagram in Fig. 1. The linac LUE-40 provides an electron beam with an average current  $I_e \approx 3 \mu\text{A}$  and an energy spectrum with full width at half maximum (FWHM)  $\Delta E_e/E_e \approx 1\%$ . The range of initial energies of electrons  $E_e = 35$ –95 MeV. A detailed description and parameters of the LUE-40 linac are given in works [27–30].

On the axis of the electron beam, there are a converter, an absorber, and a reaction chamber. The converter, made of natural tantalum, is a  $20 \times 20$  mm plate with thickness  $l = 1.05$  mm, and is attached to an aluminum absorber, shaped as a cylinder, with dimensions  $\varnothing 100$  mm and a thickness of 150 mm. The thickness of the aluminum absorber was calculated to clean the beam of  $\gamma$ -quanta from electrons with energies up to 100 MeV.

In this experiment, targets were made of natural molybdenum, which were thin discs with a diameter of 8 mm and a thickness of  $\approx 0.11$  mm, which corresponded to a mass of  $\approx 57$ –60 mg. Natural molybdenum consists of 7 stable isotopes, with isotope abundance, %:  $^{92}\text{Mo} - 14.84$ ,  $^{94}\text{Mo} - 9.25$ ,  $^{95}\text{Mo} - 15.92$ ,  $^{96}\text{Mo} - 16.68$ ,  $^{97}\text{Mo} - 9.55$ ,  $^{98}\text{Mo} - 24.13$ ,  $^{100}\text{Mo} - 9.63$  (according to [31, 32]).

The targets were placed in an aluminum capsule and using a pneumatic transport system were taken to the reaction chamber for irradiation and back to the measurement room to record the induced  $\gamma$ -activity of reaction products in the target substance.

The  $\gamma$ -quanta of the reaction products were detected using a Canberra GC-2018 semiconductor HPGe detector. At energy  $E_{\gamma} = 1332$  keV its efficiency was 20% relative to the NaI(Tl) scintillator with dimensions 3 inches in diameter and 3 inches in thickness. The resolution




FIG. 1. Schematic block diagram of the experiment. The upper part shows the measurement room, where the irradiated target is extracted from the capsule and is arranged before the HPGe detector for induced  $\gamma$ -activity measurements. The lower part shows the linac LUE-40, the Ta converter, the Al absorber, and the exposure reaction chamber.



in the metastable state are higher than in the ground state at an excitation energy of 235.7 keV.

The calculation of theoretical cross-sections  $\sigma(E)$  of studied reactions for monochromatic photons was performed using the TALYS1.95 code [31], which is installed on Linux Ubuntu-20.04. The calculations were performed for different level density models *LD* 1-6. There are three phenomenological level density models and three options for microscopic level densities:

*LD1*: Constant temperature + Fermi gas model, introduced by Gilbert and Cameron [34].

*LD2*: Back-shifted Fermi gas model [35].

*LD3*: Generalized superfluid model (GSM) [36, 37].

*LD4*: Microscopic level densities (Skyrme force) from Goriely's tables [38].

*LD5*: Microscopic level densities (Skyrme force) from Hilaire's combinatorial tables [39].

*LD6*: Microscopic level densities based on temperature-dependent Hartree-Fock-Bogoliubov calculations using the Gogny force from Hilaire's combinatorial tables [40].

Fig. 3 shows the total (metastable + ground) cross-sections  $\sigma(E)$  for the formation of the  $^{95}\text{Nb}$  nucleus on 4 stable isotopes of molybdenum (96, 97, 98, 100) calculated in the TALYS1.95 code, *LD1*. The cross-sections are given taking into account the abundance of isotopes. As can be seen, in the energy range above 35 MeV, the cross-section of the  $^{98}\text{Mo}(\gamma, 2n)^{95}\text{Nb}$  reaction has the highest values. This is due both to the  $^{98}\text{Mo}$  isotopic abundance of 24.13% and to insignificant differences in the cross-sections for the formation of the  $^{95}\text{Nb}$  nucleus on different molybdenum isotopes.

Fig. 4 shows the cross-sections  $\sigma(E)$  for the formation of the  $^{95}\text{Nb}$  nucleus in the metastable and ground states in the  $^{\text{nat}}\text{Mo}(\gamma, xnp)^{95}\text{Nb}$  reactions and the total cross-section calculated in the TALYS1.95 code, *LD1*. As can be seen from figure, the contribution of the metastable state to the total cross-section does not exceed 20% at energies above 30 MeV.

Using the theoretical cross-sections  $\sigma(E)$ , one can obtain the reaction yield, which is determined by the formula:

$$Y(E_{\gamma_{\text{max}}}) = N_n \int_{E_{\text{thr}}}^{E_{\gamma_{\text{max}}}} \sigma(E) \cdot W(E, E_{\gamma_{\text{max}}}) dE, \quad (1)$$

where  $N_n$  is the number of atoms of the element under study,  $W(E, E_{\gamma_{\text{max}}})$  is the bremsstrahlung  $\gamma$ -flux,  $E_{\text{thr}}$  – an energy of the reaction threshold,  $E_{\gamma_{\text{max}}}$  – the endpoint bremsstrahlung energy.

To estimate the  $i$ -th reaction contribution in the total production of a studied nuclide (for example, the  $^{96}\text{Mo}(\gamma, p)$  reaction in the production of the  $^{95}\text{Nb}$  nucleus on  $^{\text{nat}}\text{Mo}$ ), the relative reaction yield  $Y_i(E_{\gamma_{\text{max}}})$  were used. To calculate the  $Y_i(E_{\gamma_{\text{max}}})$  we used the cross-sections from the TALYS1.95 code and expression:




FIG. 3. Theoretical total (metastable + ground) cross-sections  $\sigma(E)$  for the formation of the  $^{95}\text{Nb}$  nucleus on stable isotopes of molybdenum  $^{96,97,98,100}\text{Mo}$  (color curves) and on  $^{\text{nat}}\text{Mo}$  (black curve). The calculations were performed for the level density model *LD1*.




FIG. 4. Theoretical cross-sections  $\sigma(E)$  of the  $^{\text{nat}}\text{Mo}(\gamma, xnp)^{95\text{m,g}}\text{Nb}$  reactions. The calculations were performed for the level density model *LD1*.

$$Y_i(E_{\gamma_{\text{max}}}) = \frac{A_i \int_{E_{\text{thr}}}^{E_{\gamma_{\text{max}}}} \sigma_i(E) \cdot W(E, E_{\gamma_{\text{max}}}) dE}{\sum_{k=1}^4 A_k \int_{E_{\text{thr}}}^{E_{\gamma_{\text{max}}}} \sigma_k(E) \cdot W(E, E_{\gamma_{\text{max}}}) dE}, \quad (2)$$

where  $\sigma_k(E)$  is the cross-section for the formation of the  $^{95}\text{Nb}$  nucleus on the  $k$ -th isotope with isotopic abundance  $A_k$ . Summation over  $k$  was carried out for 4 stable molybdenum isotopes  $^{96,97,98,100}\text{Mo}$ .




FIG. 5. Theoretical reaction yields of the  $^{95}\text{Nb}$  formation on various molybdenum isotopes relative to the total yield on  $^{\text{nat}}\text{Mo}$  according to Eq. 2. The sum of the relative yield is equal to 1.0. The calculations were performed for the level density model *LD1*.

Fig. 5 shows the contributions of the yields of various isotopes to the total yield of the  $^{\text{nat}}\text{Mo}(\gamma, xnp)^{95}\text{Nb}$  reaction according to Eq. 2 with the use cross-sections from TALYS1.95 code, *LD1*. The contribution of the reaction yield on a given isotope is determined by the cross-section, the reaction threshold, and the isotope abundance.

As a rule, in the presence of several isotopes, there is one whose contribution to the reaction yield dominates ( $> 90\%$ ), for example, as shown in [41, 42]. In the case of the reaction under study at energies up to 20 MeV, the contribution of  $^{96}\text{Mo}$  is 100%. But at the end-point bremsstrahlung energy above 30 MeV, it is difficult to determine the dominant reaction. Thus, for natural molybdenum, it is necessary to take into account the contribution of all stable isotopes to the yield of the  $^{\text{nat}}\text{Mo}(\gamma, xnp)^{95}\text{Nb}$  reaction.

Figures 3–5 show the results of calculations only for the level density model *LD1*. This was done so as not to clutter up the figures with many curves.

In this work, the values of the isomeric yield ratio are calculated as the ratio of the yield  $Y_m(E_{\gamma_{\text{max}}})$  of the formation of the nucleus in the metastable state to the yield  $Y_g(E_{\gamma_{\text{max}}})$  of the formation of the nucleus in the ground state:

$$d(E_{\gamma_{\text{max}}}) = Y_m(E_{\gamma_{\text{max}}})/Y_g(E_{\gamma_{\text{max}}}). \quad (3)$$

Fig. 6 shows the theoretical prediction of the isomeric yield ratio  $d(E_{\gamma_{\text{max}}})$  calculated using the cross-sections from the TALYS1.95 code, *LD1* model. As can be seen, the calculated isomeric yield ratios differ for different isotopes 96–100. The value  $d(E_{\gamma_{\text{max}}})$  calculated for natural Mo is between them.




FIG. 6. Theoretical isomeric yield ratios  $d(E_{\gamma_{\text{max}}})$  for the reaction products  $^{95\text{m,g}}\text{Nb}$  on different molybdenum isotopes and  $^{\text{nat}}\text{Mo}$ . The calculations were performed for the level density model *LD1*.




FIG. 7. Theoretical isomeric yield ratios  $d(E_{\gamma_{\text{max}}})$  for the reaction products from the  $^{\text{nat}}\text{Mo}(\gamma, xnp)^{95\text{m,g}}\text{Nb}$  reaction for different level density models *LD 1-6*.

The theoretical estimations of the isomeric yield ratio  $d(E_{\gamma_{\text{max}}})$  were calculated using the cross-sections from the TALYS1.95 code for six level density models *LD* and shown in Fig. 7. As can be seen, the theoretical isomeric yield ratios differ on (57–37)% at the range of  $E_{\gamma_{\text{max}}} = 35\text{--}100$  MeV. The  $d(E_{\gamma_{\text{max}}})$  calculated for the *LD3* model has the highest value at the energy range of  $E_{\gamma_{\text{max}}} = 30\text{--}100$  MeV.






FIG. 8. Simplified representation of formation and decay scheme of the isomeric pair  $^{95\text{m,g}}\text{Nb}$ . The nuclear level energies are in keV. The red color shows the emission  $\gamma$ -lines that were used in this work for the analysis.

#### IV. EXPERIMENTAL RESULTS

A simplified diagram of the decay of the niobium nucleus from the ground  $^{95\text{g}}\text{Nb}$  and metastable  $^{95\text{m}}\text{Nb}$  states is shown in Fig. 8. Nuclear spectroscopic data of the reactions  $^{\text{nat}}\text{Mo}(\gamma, xn\text{p})^{95\text{m,g}}\text{Nb}$  are presented according to [33].

The isomeric state  $^{95\text{m}}\text{Nb}$  (low-spin state,  $J^\pi = 1/2^-$ ) with a half-life  $T_{1/2}$  of  $86.6 \pm 0.08$  h decays to the unstable ground state  $^{95\text{g}}\text{Nb}$  (high-spin state,  $J^\pi = 9/2^+$ ) by emitting  $\gamma$ -quanta with the energy of 235.7 keV through an internal transition with a branching ratio  $p$  of  $94.4 \pm 0.6\%$ . Meanwhile, 5.6% of the isomeric state decays to the various energy levels of stable  $^{95}\text{Mo}$  by a  $\beta^-$ -process. The unstable ground state  $^{95\text{g}}\text{Nb}$  with a half-life  $T_{1/2}$  of  $34.975 \pm 0.007$  d decays to the 765.8 keV energy level of  $^{95}\text{Mo}$  by a  $\beta^-$ -process.

This decay pattern leads to the ground state radionuclide  $^{95\text{g}}\text{Nb}$  can be formed in two ways, directly from the target nuclide and/or indirectly through the decay of the metastable radionuclide.

As a result, to find the experimental values of the isomeric yield ratio  $d(E_{\gamma\text{max}})$ , it is necessary to solve the system of equations describing the radioactive decay of the isomeric state and decay with accumulation for the ground state. The solution of such a system of equations is given in several works, for example, [43–46], in different analytical representations.

In this work to determine the experimental values of

the isomeric yield ratio  $d(E_{\gamma\text{max}})$ , the following expression was used according to [43]:

$$d(E_{\gamma\text{max}}) = Y_{\text{m}}(E_{\gamma\text{max}})/Y_{\text{g}}(E_{\gamma\text{max}}) = \left[ \frac{\lambda_{\text{g}} F_{\text{m}}(t)}{\lambda_{\text{m}} F_{\text{g}}(t)} \left( \frac{\Delta A_{\text{g}} I_{\text{m}} \varepsilon_{\text{m}}}{\Delta A_{\text{m}} I_{\text{g}} \varepsilon_{\text{g}}} - p \frac{\lambda_{\text{g}}}{\lambda_{\text{g}} - \lambda_{\text{m}}} \right) + p \frac{\lambda_{\text{m}}}{\lambda_{\text{g}} - \lambda_{\text{m}}} \right]^{-1}, \quad (4)$$

$$F_{\text{m}}(t) = (1 - e^{-\lambda_{\text{m}} t_{\text{irr}}}) e^{-\lambda_{\text{m}} t_{\text{cool}}} (1 - e^{-\lambda_{\text{m}} t_{\text{meas}}}), \quad (5)$$

$$F_{\text{g}}(t) = (1 - e^{-\lambda_{\text{g}} t_{\text{irr}}}) e^{-\lambda_{\text{g}} t_{\text{cool}}} (1 - e^{-\lambda_{\text{g}} t_{\text{meas}}}), \quad (6)$$

where  $\lambda_{\text{g}}$  and  $\lambda_{\text{m}}$  are the decay constants for the ground and isomeric states, respectively;  $\Delta A_{\text{g}}$  and  $\Delta A_{\text{m}}$  are the number of counts in the peaks at the energies of  $\gamma$ -quanta corresponding to the decays of the isomeric and ground states, respectively;  $\varepsilon_{\text{g}}$  and  $I_{\text{g}}$  ( $\varepsilon_{\text{m}}$  and  $I_{\text{m}}$ ) are the detector efficiency and the absolute intensity of a  $\gamma$ -quantum with an energy corresponding to the decay of the ground state (isomeric state);  $p$  – the branching ratio for the decay of the isomeric to the ground state (94.4%);  $t_{\text{irr}}$ ,  $t_{\text{cool}}$  and  $t_{\text{meas}}$  are the irradiation time, cooling time, and measurement time, respectively.

In natural molybdenum targets, as a result of the  $^{\text{nat}}\text{Mo}(\gamma, xn2\text{p})$  reaction, the  $^{95}\text{Zr}$  nucleus can also be formed. In the decay scheme of  $^{95}\text{Zr}$  ( $T_{1/2} = 65.02 \pm 0.05$  d) there is a  $\gamma$ -transition with the energy  $E_{\gamma} = 235.7$  keV and intensity  $I_{\gamma} = 0.294 \pm 0.016\%$ . The decay of the  $^{95}\text{Zr}$  nucleus can contribute to the observed value of  $\Delta A_{\text{m}}$ . To take this contribution into account, calculations were performed in the TALYS1.95 code, level density model *LD1*. It was found that the estimated activity of  $^{95}\text{Zr}$  by  $\gamma$ -line with 235.7 keV is negligible. The contribution of  $^{95}\text{Zr}$  was also experimentally verified by the  $\gamma$ -lines corresponding to the decay of the  $^{95}\text{Zr}$  nucleus, namely,  $E_{\gamma} = 724.2$  keV ( $I_{\gamma} = 44.17 \pm 0.13\%$ ) and  $E_{\gamma} = 756.7$  keV ( $I_{\gamma} = 54\%$ ). No such peaks were found in the measured spectra of the induced  $\gamma$ -activity of the targets.

For the  $\gamma$ -quanta with an energy of 235.7 keV and the thicknesses of the molybdenum targets used, the self-absorption coefficients were calculated using the GEANT4.9.2 code. It was found that the value of the self-absorption coefficient did not exceed 0.8%, which was taken into account when processing the experimental results.

The experimental values of the isomeric yield ratio  $d(E_{\gamma\text{max}})$  of the nuclei products from the  $^{\text{nat}}\text{Mo}(\gamma, xn\text{p})^{95\text{m,g}}\text{Nb}$  reaction were determined at the end-point bremsstrahlung energy of 35–95 MeV (see Fig. 9 and Table I).

The calculation of the experimental error of the values  $d(E_{\gamma\text{max}})$  was performed taking into account statistical and systematic errors, the description of which can be found, for example, in [23, 24]. The uncertainties related to the  $d(E_{\gamma\text{max}})$  were calculated from Eq. 4 by using the




FIG. 9. Isomeric yield ratio  $d(E_{\gamma_{\max}})$  for the reaction products from the  $^{\text{nat}}\text{Mo}(\gamma, xnp)^{95\text{m,g}}\text{Nb}$  reaction. Experimental results of this work – blue circles, data of other authors: square – [22], red empty circles – [20], black circles – [21]. The curve – calculation in the code TALYS1.95, *LD3*, multiplied by a factor of 3.85.

error propagation principle, which indicates the maximum uncertainty of the measured values in Fig. 9 and Table I.

TABLE I. Isomeric yield ratio  $d(E_{\gamma_{\max}})$  for the nuclei products  $^{95\text{m,g}}\text{Nb}$  from the  $^{\text{nat}}\text{Mo}(\gamma, xnp)$  reaction.

$E_{\gamma_{\max}}$ , MeV	$d(E_{\gamma_{\max}}) \pm \Delta d(E_{\gamma_{\max}})$
37.50	$1.34 \pm 0.19$
41.10	$1.27 \pm 0.16$
44.20	$1.10 \pm 0.09$
49.00	$1.25 \pm 0.09$
53.70	$1.03 \pm 0.10$
53.70*	$0.97 \pm 0.10$
59.60	$1.05 \pm 0.08$
59.60*	$1.24 \pm 0.10$
63.30	$0.95 \pm 0.11$
63.30*	$1.02 \pm 0.13$
65.50	$0.86 \pm 0.09$
65.50*	$0.89 \pm 0.10$
68.25	$0.95 \pm 0.11$
68.25*	$1.00 \pm 0.13$
73.10	$1.20 \pm 0.15$
77.50	$0.91 \pm 0.08$
82.50	$0.78 \pm 0.08$
86.90	$0.97 \pm 0.11$
92.50	$0.93 \pm 0.10$
92.90	$0.90 \pm 0.10$

\* an additional series of measurements.

## V. DISCUSSION

Two main representations of the isomeric ratio  $d(E_{\gamma_{\max}})$  are used in the literature. One of them is defined as the ratio of the yields  $Y_{\text{m}}(E_{\gamma_{\max}})$  and  $Y_{\text{g}}(E_{\gamma_{\max}})$  states:  $d(E_{\gamma_{\max}}) = Y_{\text{m}}(E_{\gamma_{\max}})/Y_{\text{g}}(E_{\gamma_{\max}})$  (for example, see in [17]). The  $d(E_{\gamma_{\max}})$  values also can be found as the ratio of the yield for the formation of a product nucleus in the high-spin state  $Y_{\text{H}}(E_{\gamma_{\max}})$  to the yield for the low-spin state  $Y_{\text{L}}(E_{\gamma_{\max}})$ :  $d(E_{\gamma_{\max}}) = Y_{\text{H}}(E_{\gamma_{\max}})/Y_{\text{L}}(E_{\gamma_{\max}})$  [20, 21]. These values will be equal to each other if the nucleus in the isomeric state has a larger spin. However, the ground state of the  $^{95}\text{Nb}$  nucleus has a high spin  $J^{\pi} = 9/2^{+}$ , while the spin of the  $^{95\text{m}}\text{Nb}$  isomeric state is  $J^{\pi} = 1/2^{-}$ . This means that the values of  $d(E_{\gamma_{\max}})$  described above are inversely proportional to each other and are presented differently in different works.

In our work the experimental isomeric yield ratios of the pair  $^{95\text{m,g}}\text{Nb}$  were obtained by Eq. 4. A comparison of our results with the data of other authors [20, 21] shows satisfactory agreement within the experimental error. The estimate of the value of  $d(E_{\gamma_{\max}})$ , obtained using the yield values from [22], is in agreement with both our results and the data from [20, 21].

The range of changes in the experimental values  $d(E_{\gamma_{\max}})$  indicates the yield for formation of the  $^{95\text{m}}\text{Nb}$  nucleus on  $^{\text{nat}}\text{Mo}$  differs on 1.5–0.8 times from the yield for production  $^{95\text{g}}\text{Nb}$ . This contradicts to theoretical prediction. At the same time, there is a gradual decrease in the values of  $d(E_{\gamma_{\max}})$  with an increase in the energy  $E_{\gamma_{\max}}$ , which may be associated with a change in the contributions from different molybdenum isotopes to the formation of the  $^{95\text{m,g}}\text{Nb}$  isomeric pair.

The experimental results obtained in this work and data [20–22] significantly exceed theoretical estimates obtained with the cross-sections from the TALYS1.95 code, level density models *LD* 1–6. Difference between theory and experiment is from 3.85 up to 5.80 times. These factors were obtained by the  $\chi^2$  method with using all experimental values shown in Fig. 9. The result of the theoretical calculation  $d(E_{\gamma_{\max}})$ , obtained using the level density model *LD3*, is closest to experimental data.

The observed difference may be due to the underestimation of the calculation for the cross-section of the formation of the nucleus in the isomeric state. Alternatively, it may be the result of incorrect calculation of both  $\sigma_{\text{m}}(E)$  and  $\sigma_{\text{g}}(E)$ . However, the theoretical prediction of the energy dependence  $d(E_{\gamma_{\max}})$  describes well the gradual decrease in the measured values with increasing  $E_{\gamma_{\max}}$  in the energy range under study.

## VI. CONCLUSIONS

In the present work, the experiment was performed using the beam from the NSC KIPT linear accelerator LUE-40 and activation and off-line  $\gamma$ -ray spectrometric

technique. The isomeric yield ratios  $d(E_{\gamma_{\max}})$  of the  $^{95\text{m,g}}\text{Nb}$  reaction products from photonuclear reactions on natural Mo targets were determined. The region of end-point energy of bremsstrahlung  $\gamma$ -quanta spectra was  $E_{\gamma_{\max}} = 35\text{--}95$  MeV.

The data from [20–22] are in satisfactory agreement with obtained results.

The calculation of  $d(E_{\gamma_{\max}})$  was performed using the cross-sections  $\sigma(E)$  for studied reactions from the TALYS1.95 code, for six level density model  $LD$  1–6.

Comparison of the experimental and calculated values of  $d(E_{\gamma_{\max}})$  for the reaction  $^{\text{nat}}\text{Mo}(\gamma, xn)^{95\text{m,g}}\text{Nb}$  showed a significant excess of the experimental results over the computations. The result of the theoretical calculation  $d(E_{\gamma_{\max}})$ , obtained using the level density model  $LD3$ , is closest to experimental data. The calculated energy dependence  $d(E_{\gamma_{\max}})$  in the energy range under study describes well the gradual decrease in the measured

values with increasing energy  $E_{\gamma_{\max}}$ .

## ACKNOWLEDGMENT

The authors would like to thank the staff of the linear electron accelerator LUE-40 NSC KIPT, Kharkiv, Ukraine, for their cooperation in the realization of the experiment.

## DECLARATION OF COMPETING INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

- 
- [1] S.Y.F. Chu, L.P. Ekstrom, R.B. Firestone, The Lund/LBNL, Nuclear Data Search, Version 2.0, February 1999, WWW Table of Radioactive Isotopes, available from <http://nucleardata.nuclear.lu.se/toi/>.
- [2] R. Volpel, Bestimmung von Isomerenverhältnissen bei Verschiedenen Kernphoto- und (n, 2n)-Reaktionen // Nucl Phys A182 (1972) 411, doi.org/10.1016/0375-9474(72)90287-4.
- [3] H. Bartsch, K. Huber, U. Kneissl, and H. Krieger, Critical consideration of the statistical model analysis of photonuclear isomeric cross-section ratios // Nucl. Phys. A256 (1976) 243, doi.org/10.1016/0375-9474(76)90106-8.
- [4] D. Kolev, E. Dobрева, N. Nenov, and V. Todorov, A convenient method for experimental determination of yields and isomeric ratios in photonuclear reactions measured by the activation technique // Nucl. Instrum. Method. Phys. Research A356 (1995) 390, doi.org/10.1016/0168-9002(94)01319-5.
- [5] I.B. Haller, G. Rudstam, Relative yields of the isomeric pairs  $^{69\text{g}}\text{Zn}$   $^{69\text{m}}\text{Zn}$  and  $^{52\text{g}}\text{Mn}$   $^{52\text{m}}\text{Mn}$  in some spallation reactions induced by 20–153 MeV protons // J. Inorg. Nucl. Chem. 19 (1961) 1, doi.org/10.1016/0022-1902(61)80038-9.
- [6] D. Kolev, Studies of some isomeric yield ratios produced with bremsstrahlung // Appl. Radiat. Isot. 49 (1998) 989, doi.org/10.1016/S0969-8043(97)10094-X.
- [7] J.R. Huizenga and R. Vandenbosch, Interpretation of Isomeric Cross-Section Ratios for (n, $\gamma$ ) and ( $\gamma$ ,n) Reactions // Phys. Rev. 120 (1960) 1305, doi.org/10.1103/PhysRev.120.1305.
- [8] R. Vandenbosch and J.R. Huizenga, Isomeric Cross Section Ratios for Reactions Producing the Isomeric Pair  $\text{Hg}^{197,197\text{m}}$ , Phys Rev 12: (1960) 1313, doi.org/10.1103/PhysRev.120.1313.
- [9] H. A. Bethe, Nuclear Physics B. Nuclear Dynamics, Theoretical // Rev. Mod. Phys. 9 (1937) 84, doi.org/10.1103/RevModPhys.9.69.
- [10] C. Bloch, Theory of Nuclear Level Density // Phys. Rev. 93 (1954) 1094, doi.org/10.1103/PhysRev.93.1094.
- [11] K.J. Le Couteur, D.W. Lang, Neutron evaporation and level densities in excited nuclei // Nucl. Phys. 13 (1959) 32, doi.org/10.1016/0029-5582(59)90136-1.
- [12] B.S. Ishkhanov, V.N. Orlin, and S.Yu. Troschiev, Photodisintegration of Tantalum // Phys. Atom. Nucl. 75 (2012) 253, doi.org/10.1134/S1063778812020093. (Yadernaya Fizika 75 (2012) 283).
- [13] Experimental Nuclear Reaction Data (EXFOR) // <https://www-nds.iaea.org/exfor/>
- [14] Data Center of Photonuclear Experiments, <http://cdfc.sinp.msu.ru/>.
- [15] T. Kato and O. Yoshinaga. Yields of photonuclear reactions for photon-activation analysis with high energy bremsstrahlung // Talanta. Pergamon Press 19 (1972) p. 515.
- [16] T. Kato. Quantitative evaluation of the photon-activation analysis // J. Radioanal. Chem. 16 (1973) 307.
- [17] S.R. Palvanov and O. Razhabov. Isomeric yield ratios of photonuclear reactions at  $E_{\gamma_{\max}}$  25 and 30 MeV // At. Energy 87 (1999) 533.
- [18] T.D. Thiep, T.T. An, N.T. Khai et al. The isomeric ratios in some photonuclear reactions ( $\gamma$ ,n), ( $\gamma$ ,p), ( $\gamma$ ,2n) and ( $\gamma$ ,np) induced by bremsstrahlung with end-point energies in the giant dipole resonance region // Phys. Part. Nucl. Lett. 6 (2009) 126, doi.org/10.1134/S1547477109020058.
- [19] Tran Duc Thiep, Truong Thi An, Phan Viet Cuong et al. Isomeric Ratios in Photonuclear Reactions of Molybdenum Isotopes Induced by Bremsstrahlung in the Giant Dipole Resonance Region // Phys. Part. Nucl. Lett. 14 (2017) 102, doi.org/10.1134/S1547477117010241.
- [20] K.S. Kim, M.D. Sh. Rahman, M. Lee, G. Kim et al. Measurement of isomeric yield ratios for  $^{93}\text{Nb}(\gamma, 4n)^{89\text{m,g}}\text{Nb}$  and  $^{\text{nat}}\text{Mo}(\gamma, xn)^{95\text{m,g}}\text{Nb}$  reactions with 50-, 60-, and 70-MeV bremsstrahlung // J. Radioanal. Nucl. Chem. 287 (2011) 869, doi.org/10.1007/s10967-010-0839-3.
- [21] H. Naik, G. Kim, Kw. Kim, M. Zaman et al. Independent isomeric yield ratios of  $^{95\text{m,g}}\text{Nb}$  in the  $^{\text{nat}}\text{Mo}(\gamma, xn)$  and  $^{\text{nat}}\text{Zr}(p, xn)$  reactions // J. Radioanal. Nucl. Chem. 300 (2014) 1121, doi.org/10.1007/s10967-014-3045-x.



- [22] B.S. Ishkhanov, I.M. Kapitonov, A.A. Kuznetsov, V.N. Orlin, and H.D. Yoon, Photonuclear Reactions on Molybdenum Isotopes // *Phys. Atom. Nucl.* 77 (2014) 1362, doi.org/10.1134/S106377881410007X. (*Yadernaya Fizika* 77 (2014) 1427)
- [23] O.S. Deiev, I.S. Timchenko, S.N. Olejnik, V.A. Kushnir, V.V. Mytrochenko, S.A. Perezhogin. Isomeric ratio of the  $^{181}\text{Ta}(\gamma, 3n)^{178\text{m,g}}\text{Ta}$  reaction products at energy  $E_{\gamma\text{max}}$  up to 95 MeV // *Chin. Phys. C*46 (2022) 014001, doi.org/10.1088/1674-1137/ac2a95.
- [24] O.S. Deiev, I.S. Timchenko, S.M. Olejnik, V.A. Kushnir, V.V. Mytrochenko, S.A. Perezhogin. Cross-sections of photoneutron reactions on  $^{181}\text{Ta}$  at  $E_{\gamma\text{max}}$  up to 95 MeV // *Phys. Rev. C*106 (2022) 024617, doi.org/10.1103/PhysRevC.106.024617.
- [25] A.N. Vodin, O.S. Deiev, I.S. Timchenko, S.N. Olejnik, N.I. Ayzatsky, V.A. Kushnir, V.V. Mytrochenko, S.A. Perezhogin. Photoneutron cross-sections for the reactions  $^{181}\text{Ta}(\gamma, xn; x = 1 - 8)^{181-x}\text{Ta}$  at  $E_{\gamma\text{max}} = 80-95$  MeV // *Eur. Phys. J. A*57 (2021) 208, doi.org/10.1140/epja/s10050-021-00484-x, arXiv:2103.09859.
- [26] O.S. Deiev, I.S. Timchenko, S.N. Olejnik et al. Cross-sections for the  $^{27}\text{Al}(\gamma, x)^{22}\text{Na}$  multichannel reaction with the 28.3 MeV difference of the reaction thresholds // *Chin. Phys. C*46 (2022) 064002, doi.org/10.1088/1674-1137/ac5733, arXiv:2105.12658.
- [27] A.N. Dovbnya, M.I. Aizatsky, V.N. Boriskin et al. Beam parameters of an S-band electron linac with beam energy of 30–100 MeV // *Probl. At. Sci. Technol.* 2 (2006) 11.
- [28] M. I. Aizatsky, V. I. Beloglazov, V. N. Boriskin et al., State and prospects of the linac of nuclear-physics complex with energy of electrons up to 100 MeV // *Probl. At. Sci. Technol.* 3 (2014) 60.
- [29] V.V. Mytrochenko, L.I. Selivanov, V.Ph. Zhyglo et al. Magnetic system for cleaning the gamma beam at the LUE-40 linac output // *Probl. Atom. Scien. Techn.* 3 (2022) 62, doi.org/10.46813/2022-139-062.
- [30] O.S. Deiev, I.S. Timchenko, S.M. Olejnik et al. Photonuclear reactions cross-section at energies up to 100 MeV for different experimental setups // *Probl. Atom. Scien. Techn.* 5 (2022) 11, doi.org/10.46813/2022-141-011.
- [31] A. Koning and D. Rochman. // *Nucl. Data Sheets* 2012, V. 113, p. 2841, TALYS – based evaluated nuclear data library. <https://tendl.web.psi.ch/tendl2019/tendl2019.html>.
- [32] S. Agostinelli et al. Electron and Positron Incident // *Methods Phys.* A506 (2003) 250, <http://GEANT4.9.2.web.cern.ch/GEANT4.9.2/>.
- [33] S.Y.F. Chu, L.P. Ekstrom, R.B. Firestone. The Lund/LBNL, Nuclear Data Search, Version 2.0, February 1999, WWW Table of Radioactive Isotopes // <http://nucleardata.nuclear.lu.se/toi/>.
- [34] A. Gilbert, and A.G.W. Cameron, A composite nuclear-level density formula with shell corrections // *Canad. Journ. Phys.* 43 (1965) 1446; doi.org/10.1139/p65-139
- [35] W. Dilg, W. Schantl, H. Vonach, and M. Uhl. Level density parameters for the back-shifted fermi gas model in the mass range  $40 < A < 250$  // *Nucl. Phys. A*217 (1973) 269; doi.org/10.1016/0375-9474(73)90196-6.
- [36] A.V. Ignatyuk, K.K. Istekov, and G.N. Smirenkin. Role of collective effects in the systematics of nuclear level densities // *Sov. J. Nucl. Phys.* 29 (1979) 450.
- [37] A. V. Ignatyuk, J. L. Weil, S. Raman, and S. Kahane. Density of discrete levels in  $^{116}\text{Sn}$  // *Phys. Rev. C*47 (1993) 1504; doi.org/10.1103/PhysRevC.47.1504.
- [38] S. Goriely, F. Tondeur, and J.M. Pearson. A Hartree-Fock nuclear mass table // *Atom. Data Nucl. Data Tables* 77 (2001) 311; doi.org/10.1006/adnd.2000.0857.
- [39] S. Goriely, S. Hilaire, and A.J. Koning. Improved microscopic nuclear level densities within the Hartree-Fock-Bogoliubov plus combinatorial method // *Phys. Rev. C*78 (2008) 064307; doi.org/10.1103/PhysRevC.78.064307.
- [40] S. Hilaire, M. Girod, S. Goriely, and A. J. Koning. Temperature-dependent combinatorial level densities with the D1M Gogny force // *Phys. Rev. C*86 (2012) 064317; doi.org/10.1103/PhysRevC.86.064317.
- [41] O.S. Deiev, I.S. Timchenko, S.N. Olejnik et al. Photonuclear reactions  $^{65}\text{Cu}(\gamma, n)^{64}\text{Cu}$  and  $^{63}\text{Cu}(\gamma, xn)^{63-x}\text{Cu}$  cross-sections in the energy range  $E_{\gamma\text{max}} = 35-94$  MeV // *Chin. Phys. C*46 (2022) 124001, doi.org/10.1088/1674-1137/ac878a.
- [42] O.S. Deiev, I.S. Timchenko, S.M. Olejnik et al. Photonuclear reactions  $^{nat}\text{Ni}(\gamma, xn)^{57}\text{Ni}$  and  $^{nat}\text{Ni}(\gamma, xn)^{56}\text{Ni}$  in the energy range  $E_{\gamma\text{max}} = 35-94$  MeV // *Nucl. Phys. A*1028 (2022) 122542, doi.org/10.1016/j.nuclphysa.2022.122542.
- [43] S. R. Palvanov and O. Razhabov, M. Kajumov, M.I. Mamayusupova, O. Zhuraev. Isomeric yield ratios of the  $(\gamma, n)$  and  $(\gamma, 2n)$  reactions on nuclei of  $^{110}\text{Pb}$ ,  $^{142}\text{Nd}$ , and  $^{144}\text{Sm}$  // *Bulletin Rus. Acad. Sciences. Physics* 75 (2011) 222. (*Izvestiya Ross. Akad. Nauk.* 75 (2011) 239).
- [44] R. Vanska, R. Rieppo. The experimental isomeric cross-section ratio in the nuclear activation technique // *Nucl. Instr. Meth.* 179 (1981) 525, doi.org/10.1016/0029-554X(81)90178-6.
- [45] Tran Duc Thiep, Truong Thi An, Phan Viet Cuong et al. Isomeric Ratios in Photonuclear Reactions of Molybdenum Isotopes Induced by Bremsstrahlung in the Giant Dipole Resonance Region // *Phys. Part. Nucl. Lett.* 14 (2017) 102, doi.org/10.1134/S1547477117010241.
- [46] D. Kolev, E. Dobreva, N. Nenov, V. Todorov. A convenient method for experimental determination of yields and isomeric ratios in photonuclear reactions measured by the activation technique // *Nucl. Instr. Meth. Phys. Research A*356 (1995) 390, doi.org/10.1016/0168-9002(94)01319-5.