

Exact invariants for a class of three-dimensional time-dependent classical Hamiltonians

Jürgen Struckmeier and Claus Riedel

Gesellschaft für Schwerionenforschung (GSI), Planckstr. 1, 64291 Darmstadt, Germany

(Dated: Received 26 April 2000; revised manuscript received 8 June 2000)

An exact invariant is derived for three-dimensional Hamiltonian systems of N particles confined within a general velocity-independent potential. The invariant is found to contain a time-dependent function $f_2(t)$, embodying a solution of a linear third-order differential equation whose coefficients depend on the explicitly known trajectories of the particle ensemble. Our result is applied to a one-dimensional time-dependent non-linear oscillator, and to a system of Coulomb interacting particles in a time-dependent quadratic external potential.

Published in: Phys. Rev. Lett. **85**, No. 18, 3830 (2000)

PACS numbers: 41.85.-p, 45.50.Jf

We consider a system of a non-relativistic ensemble of N particles of the same species moving in an explicitly time-dependent and velocity-independent potential, whose Hamiltonian H takes the form

$$H = \sum_{i=1}^N \frac{1}{2} \left[p_{x,i}^2 + p_{y,i}^2 + p_{z,i}^2 \right] + V(\mathbf{x}, \mathbf{y}, \mathbf{z}, t), \quad (1)$$

with \mathbf{x} , \mathbf{y} , and \mathbf{z} the N component vectors of the spatial coordinates of all particles. It is hereby assumed that the system may be completely described within $6N$ -dimensional Cartesian phase-space spanned by the $3N$ particle coordinates and their conjugate momenta. From the canonical equations, we

derive for each particle i the equations of motion

$$\dot{x}_i = p_{x,i}, \quad \dot{p}_{x,i} = -\frac{\partial V(\mathbf{x}, \mathbf{y}, \mathbf{z}, t)}{\partial x_i}, \quad (2)$$

and likewise for the y and z degree of freedom. The solution functions $\mathbf{x}(t)$, $\mathbf{y}(t)$, $\mathbf{z}(t)$, and $\mathbf{p}_x(t)$, $\mathbf{p}_y(t)$, $\mathbf{p}_z(t)$ define a path within the $6N$ -dimensional phase-space that completely describes the system's time evolution. A quantity

$$I = I(\mathbf{x}(t), \mathbf{p}_x(t), \mathbf{y}(t), \mathbf{p}_y(t), \mathbf{z}(t), \mathbf{p}_z(t), t) \quad (3)$$

constitutes an invariant of the particle motion if its total time derivative vanishes:

$$\frac{dI}{dt} = \frac{\partial I}{\partial t} + \sum_{i=1}^N \left[\frac{\partial I}{\partial x_i} \dot{x}_i + \frac{\partial I}{\partial y_i} \dot{y}_i + \frac{\partial I}{\partial z_i} \dot{z}_i + \frac{\partial I}{\partial p_{x,i}} \dot{p}_{x,i} + \frac{\partial I}{\partial p_{y,i}} \dot{p}_{y,i} + \frac{\partial I}{\partial p_{z,i}} \dot{p}_{z,i} \right] = 0.$$

We examine the existence of a conserved quantity (3) for a system described by (1) with a special ansatz for I being at most quadratic in the momenta[1]

$$I = \sum_i \left[f_2(t) (p_{x,i}^2 + p_{y,i}^2 + p_{z,i}^2) + f_1(x_i, t) p_{x,i} + g_1(y_i, t) p_{y,i} + h_1(z_i, t) p_{z,i} \right] + f_0(\mathbf{x}, \mathbf{y}, \mathbf{z}, t). \quad (4)$$

The set of functions $f_2(t)$, $f_1(x_i, t)$, $g_1(y_i, t)$, $h_1(z_i, t)$, and $f_0(\mathbf{x}, \mathbf{y}, \mathbf{z}, t)$ that render I invariant are to be determined. With the single particle equations of motion (2), a vanishing total time derivative of Eq. (4) means explicitly

$$\sum_i \left[(\dot{x}_i^2 + \dot{y}_i^2 + \dot{z}_i^2) \frac{df_2}{dt} + \dot{x}_i \frac{\partial f_1}{\partial t} + \dot{y}_i \frac{\partial g_1}{\partial t} + \dot{z}_i \frac{\partial h_1}{\partial t} + \dot{x}_i^2 \frac{\partial f_1}{\partial x_i} + \dot{y}_i^2 \frac{\partial g_1}{\partial y_i} + \dot{z}_i^2 \frac{\partial h_1}{\partial z_i} + \dot{x}_i \frac{\partial f_0}{\partial x_i} + \dot{y}_i \frac{\partial f_0}{\partial y_i} + \dot{z}_i \frac{\partial f_0}{\partial z_i} - (2f_2 \dot{x}_i + f_1) \frac{\partial V}{\partial x_i} - (2f_2 \dot{y}_i + g_1) \frac{\partial V}{\partial y_i} - (2f_2 \dot{z}_i + h_1) \frac{\partial V}{\partial z_i} \right] + \frac{\partial f_0}{\partial t} = 0. \quad (5)$$

We may arrange the terms of this equation with regard to their power in the velocities \dot{x}_i , \dot{y}_i , and \dot{z}_i . Eq. (5) must hold independently of the specific phase-space location of each individual particle i . Therefore, the coefficients pertaining to the velocity powers must vanish separately for each index i . The

condition for the terms proportional to \dot{x}_i^2 is

$$\frac{\partial f_1(x_i, t)}{\partial x_i} + \frac{df_2(t)}{dt} = 0,$$

and similarly for the functions g_1 and h_1 . It follows that $f_1(x_i, t)$, $g_1(y_i, t)$, and $h_1(z_i, t)$ must be linear functions in

x_i , y_i , and z_i , respectively

$$f_1(x_i, t) = -\dot{f}_2(t) x_i + b_{x,i}(t) \quad (6a)$$

$$g_1(y_i, t) = -\dot{f}_2(t) y_i + b_{y,i}(t) \quad (6b)$$

$$h_1(z_i, t) = -\dot{f}_2(t) z_i + b_{z,i}(t), \quad (6c)$$

with $b_{x,i}(t)$, $b_{y,i}(t)$, and $b_{z,i}(t)$ defined as arbitrary functions of time only.

The terms of Eq. (5) that are linear in \dot{x}_i sum up to

$$\frac{\partial f_1}{\partial t} + \frac{\partial f_0}{\partial x_i} - 2f_2(t) \frac{\partial V}{\partial x_i} = 0. \quad (7)$$

In order to eliminate $\partial f_1/\partial t$ from (7), we calculate the partial time derivative of $f_1(x_i, t)$ from Eq. (6a)

$$\frac{\partial f_1}{\partial t} = -\ddot{f}_2(t) x_i + \dot{b}_{x,i}(t). \quad (8)$$

Again, similar expressions apply to $\partial g_1/\partial t$ and $\partial h_1/\partial t$. Inserting (8) into (7), and solving for the terms containing the partial derivatives of the yet unknown but arbitrary ancillary function $f_0(\mathbf{x}, \mathbf{y}, \mathbf{z}, t)$, one obtains the following three differential equations for f_0 :

$$\frac{\partial f_0}{\partial x_i} = \ddot{f}_2(t) x_i - \dot{b}_{x,i}(t) + 2f_2(t) \frac{\partial V}{\partial x_i} \quad (9a)$$

$$\frac{\partial f_0}{\partial y_i} = \ddot{f}_2(t) y_i - \dot{b}_{y,i}(t) + 2f_2(t) \frac{\partial V}{\partial y_i} \quad (9b)$$

$$\frac{\partial f_0}{\partial z_i} = \ddot{f}_2(t) z_i - \dot{b}_{z,i}(t) + 2f_2(t) \frac{\partial V}{\partial z_i}. \quad (9c)$$

A function $f_0(\mathbf{x}, \mathbf{y}, \mathbf{z}, t)$ with partial derivatives (9) is obviously given by

$$f_0(\mathbf{x}, \mathbf{y}, \mathbf{z}, t) = 2f_2(t) V(\mathbf{x}, \mathbf{y}, \mathbf{z}, t) + \sum_i \left[\frac{1}{2} \ddot{f}_2(t) (x_i^2 + y_i^2 + z_i^2) - \dot{b}_{x,i} x_i - \dot{b}_{y,i} y_i - \dot{b}_{z,i} z_i \right]. \quad (10)$$

The remaining terms of Eq. (5) do not depend on the velocities \dot{x}_i , \dot{y}_i , and \dot{z}_i . With (6), these terms impose the following condition for I to embody an invariant of the particle motion

$$\sum_i \left[\left(\dot{f}_2 x_i - b_{x,i} \right) \frac{\partial V}{\partial x_i} + \left(\dot{f}_2 y_i - b_{y,i} \right) \frac{\partial V}{\partial y_i} + \left(\dot{f}_2 z_i - b_{z,i} \right) \frac{\partial V}{\partial z_i} \right] + \frac{\partial f_0}{\partial t} = 0. \quad (11)$$

In order to express Eq. (11) in a closed form for $f_2(t)$, one has to eliminate $\partial f_0/\partial t$. To this end, we calculate the partial time derivative of Eq. (10), i.e., the time derivative at fixed particle coordinates x_i , y_i , and z_i :

$$\frac{\partial f_0(\mathbf{x}, \mathbf{y}, \mathbf{z}, t)}{\partial t} = 2\dot{f}_2(t) V + 2f_2(t) \frac{\partial V}{\partial t} + \sum_i \left[\frac{1}{2} \ddot{f}_2(t) (x_i^2 + y_i^2 + z_i^2) - \ddot{b}_{x,i} x_i - \ddot{b}_{y,i} y_i - \ddot{b}_{z,i} z_i \right]. \quad (12)$$

Inserting Eq. (12) into Eq. (11), we finally get a linear third-order differential equation for $f_2(t)$ and the $b_{x,y,z,i}(t)$ that only depends on the spatial variables of the particle ensemble

$$2\dot{f}_2(t) V + 2f_2(t) \frac{\partial V}{\partial t} + \sum_i \left[\frac{1}{2} \ddot{f}_2(t) (x_i^2 + y_i^2 + z_i^2) + \dot{f}_2(t) \left(x_i \frac{\partial V}{\partial x_i} + y_i \frac{\partial V}{\partial y_i} + z_i \frac{\partial V}{\partial z_i} \right) + b_{x,i} \ddot{x}_i - \ddot{b}_{x,i} x_i + b_{y,i} \ddot{y}_i - \ddot{b}_{y,i} y_i + b_{z,i} \ddot{z}_i - \ddot{b}_{z,i} z_i \right] = 0. \quad (13)$$

At this point, it is helpful to review our derivation made so far. Speaking of an invariant I of the particle motion means explicitly to pinpoint a quantity (3) that is conserved along the phase-space path representing the system's time evolution. This path is defined as the subset of the $6N$ -dimensional phase-space on which the equations of motion (2) are fulfilled. In order to work out the invariant I of the particle motion, the equations of motion (2) have been inserted into the expression for $dI/dt = 0$ in Eq. (5). This means that the domain of (5), and hence the physical significance of the subsequent equa-

tions (13) and (14), is restricted to the actual phase-space path. Along the phase-space path, all terms of Eq. (13) that depend on the particle trajectories are in fact functions of the parameter t only. Accordingly, the potential $V(\mathbf{x}(t), \mathbf{y}(t), \mathbf{z}(t), t)$ and its derivatives are time-dependent coefficients of an ordinary differential equation for $f_2(t)$. In contrast to Ref. [1], Eq. (13) is not conceived as a partial differential equation for V in our context.

With $f_2(t)$ representing a solution of (13), the invariant I then follows from (4), (6), and (10) together with the Hamil-

tonian (1) as

$$I = 2f_2(t)H + \sum_i \left[-\dot{f}_2(t)(x_i p_{x,i} + y_i p_{y,i} + z_i p_{z,i}) + \frac{1}{2}\ddot{f}_2(t)(x_i^2 + y_i^2 + z_i^2) + b_{x,i} p_{x,i} - \dot{b}_{x,i} x_i + b_{y,i} p_{y,i} - \dot{b}_{y,i} y_i + b_{z,i} p_{z,i} - \dot{b}_{z,i} z_i \right]. \quad (14)$$

The invariant (14) is easily shown to embody a time integral of Eq. (13) by calculating the total time derivative of (14), and inserting the single particle equations of motion (2). Hence, Eq. (14) provides a time integral of Eq. (13) if and only if the system's evolution is governed by the equations of motion (2).

From their definition in Eqs. (6), $b_{x,i}(t)$, $b_{y,i}(t)$, and $b_{z,i}(t)$ are arbitrary functions of time that do not depend on $f_2(t)$. As a consequence, the sums over the terms containing the respective functions in Eq. (13) must vanish separately:

$$\dot{f}_2(t) \left(2V + \sum_{i=1}^N \left[x_i \frac{\partial V}{\partial x_i} + y_i \frac{\partial V}{\partial y_i} + z_i \frac{\partial V}{\partial z_i} \right] \right) + 2f_2(t) \frac{\partial V}{\partial t} + \ddot{f}_2(t) \sum_{i=1}^N \frac{1}{2}(x_i^2 + y_i^2 + z_i^2) = 0, \quad (15)$$

$$b_{x,i} \ddot{x}_i - \dot{b}_{x,i} \dot{x}_i = 0, \quad b_{y,i} \ddot{y}_i - \dot{b}_{y,i} \dot{y}_i = 0, \quad b_{z,i} \ddot{z}_i - \dot{b}_{z,i} \dot{z}_i = 0, \quad i = 1, \dots, N. \quad (15')$$

We thus obtain the following distinct invariants

$$I_{f_2} = 2f_2(t)H - \dot{f}_2(t) \sum_{i=1}^N (x_i p_{x,i} + y_i p_{y,i} + z_i p_{z,i}) + \ddot{f}_2(t) \sum_{i=1}^N \frac{1}{2}(x_i^2 + y_i^2 + z_i^2), \quad (16)$$

$$I_{b_{x,i}} = b_{x,i} p_{x,i} - \dot{b}_{x,i} x_i, \quad I_{b_{y,i}} = b_{y,i} p_{y,i} - \dot{b}_{y,i} y_i, \quad I_{b_{z,i}} = b_{z,i} p_{z,i} - \dot{b}_{z,i} z_i, \quad i = 1, \dots, N. \quad (16')$$

Regarding Eq. (15), one finds that for the particular case $\partial V/\partial t \equiv 0$, hence for autonomous systems, $f_2(t) = \text{const.}$ is always a solution of Eq. (15). For this case, the invariant (16) reduces to $I_{f_2} \propto H$, thus provides the system's total energy, which is a known invariant for Hamiltonian systems with no explicit time-dependence. Nevertheless, Eq. (15) also allows for solutions $f_2(t) \neq \text{const.}$ for these systems. We thereby obtain other non-trivial invariants for autonomous systems that exist in addition to the invariant representing the energy conservation law.

Eq. (15) can be significantly simplified for potentials V that may be expressed as a sum of homogeneous functions $V = \sum_m V_m$. By definition, V_m is referred to as homogeneous if for every real $\lambda > 0$ and all \mathbf{x} , \mathbf{y} , and \mathbf{z} the condition

$$V_m(\lambda \mathbf{x}, \lambda \mathbf{y}, \lambda \mathbf{z}, t) = \lambda^{k_m} V_m(\mathbf{x}, \mathbf{y}, \mathbf{z}, t)$$

is satisfied, k_m specifying the degree of homogeneity of V_m . With V a sum of homogeneous functions, Euler's relation may be written as

$$\sum_{i=1}^N \left[x_i \frac{\partial V}{\partial x_i} + y_i \frac{\partial V}{\partial y_i} + z_i \frac{\partial V}{\partial z_i} \right] = \sum_m k_m V_m. \quad (17)$$

Using (17), the differential equation (15) for $f_2(t)$ finally reads for homogeneous potential functions V_m

$$2f_2 \frac{\partial V}{\partial t} + \dot{f}_2 \sum_m (k_m + 2)V_m + \ddot{f}_2 \sum_{i=1}^N \frac{1}{2}(x_i^2 + y_i^2 + z_i^2) = 0. \quad (18)$$

As a simple example, we investigate the one-dimensional non-linear Hamiltonian system of a time-dependent ‘‘asymmetric spring’’, defined by

$$H = \frac{1}{2}p^2 + \frac{1}{2}\omega^2(t)x^2 + a(t)x^3. \quad (19)$$

The related equation of motion follows as

$$\ddot{x} + \omega^2(t)x + 3a(t)x^2 = 0. \quad (20)$$

The invariant I_{f_2} is immediately found writing down the general invariant (16) for one dimension and one particle with the Hamiltonian H given by (19)

$$I_{f_2} = f_2(p^2 + \omega^2 x^2 + 2ax^3) - \dot{f}_2 xp + \frac{1}{2}\ddot{f}_2 x^2. \quad (21)$$

The function $f_2(t)$ for this particular case is given as a solution of the linear third-order differential equation

$$\ddot{f}_2 + 4\dot{f}_2\omega^2 + 4f_2\omega\dot{\omega} + x(t)(4f_2\dot{a} + 10\dot{f}_2a) = 0, \quad (22)$$

which follows from (15) or, equivalently, from (18). Since the particle trajectory $x = x(t)$ is explicitly contained in Eq. (22), it must be known prior to integrating Eq. (22). The trajectory is obtained integrating the equation of motion (20).

We may easily convince ourselves that I_{f_2} is indeed a conserved quantity. Calculating the total time derivative of Eq. (21), and inserting the equation of motion (20), we end up with Eq. (22), which is fulfilled by definition of $f_2(t)$ for the given trajectory $x = x(t)$.

The third-order equation (22) may be converted into a coupled set of first- and second-order equations. With the substitution $\rho_x^2(t) \equiv f_2(t)$, the second-order equation writes

$$\ddot{\rho}_x + \omega^2(t) \rho_x - \frac{g_x(t)}{\rho_x^3} = 0. \quad (23)$$

Eq. (23) is equivalent to (22), provided that the time derivative of the function $g_x(t)$, introduced in (23), is given by

$$\dot{g}_x(t) = -x(t) (2\dot{a}\rho_x^4 + 10a\rho_x^3\dot{\rho}_x). \quad (24)$$

Expressing the invariant (21) in terms of $\rho_x(t)$, we get inserting the auxiliary equation (23)

$$I_{\rho_x} = (\rho_x p - \dot{\rho}_x x)^2 + \frac{x^2}{\rho_x^2} g_x(t) + 2a(t) \rho_x^2 x^3. \quad (25)$$

The invariant (25) reduces to the well-known Lewis invariant [2, 3] for the time-dependent harmonic oscillator if $a(t) \equiv 0$, which means that $g_x(t) = \text{const.}$ For this particular case, Eq. (24), and hence Eq. (23), no longer depends on the specific particle trajectory $x = x(t)$. Consequently, the solution functions $\rho_x(t)$ and $\dot{\rho}_x(t)$ apply to all trajectories that follow from $\ddot{x} + \omega^2(t)x = 0$. With regard to Eq. (15), we conclude that a decoupling from the equations of motion (2) may occur for linear systems only.

A more challenging example is constituted by an ensemble of Coulomb interacting particles of the same species moving

in a time-dependent quadratic external potential, as typically given in the co-moving frame for charged particle beams that propagate through linear external focusing devices

$$V(\mathbf{x}, \mathbf{y}, \mathbf{z}, t) = \sum_i \left[\frac{1}{2} \omega_x^2(t) x_i^2 + \frac{1}{2} \omega_y^2(t) y_i^2 + \frac{1}{2} \omega_z^2(t) z_i^2 + \frac{1}{2} \sum_{j \neq i} \frac{c_1}{r_{ij}} \right], \quad (26)$$

with $r_{ij}^2 = (x_i - x_j)^2 + (y_i - y_j)^2 + (z_i - z_j)^2$ and $c_1 = q^2/4\pi\epsilon_0 m$, q and m denoting the particles' charge and mass, respectively. The equations of motion that follow from (2) with (26) are

$$\ddot{x}_i + \omega_x^2(t) x_i - c_1 \sum_{j \neq i} \frac{x_i - x_j}{r_{ij}^3} = 0, \quad (27)$$

and likewise for the y and z degrees of freedom. We note that the factor 1/2 in front of the Coulomb interaction term in (26) disappears since each term occurs twice in the symmetric form of the double sum.

Equation (26) may be split into a sum of two homogeneous functions, namely the focusing potential part with the degree of homogeneity $k_1 = 2$, and the Coulomb interaction part, the latter with the degree $k_2 = -1$. Consequently, the third-order differential equation (18) for $f_2(t)$ reads (after rearranging)

$$\sum_i \left[\dot{f}_2 \sum_{j \neq i} \frac{c_1}{r_{ij}} + x_i^2 \left(\ddot{f}_2 + 4\dot{f}_2 \omega_x^2 + 4f_2 \omega_x \dot{\omega}_x \right) + y_i^2 \left(\ddot{f}_2 + 4\dot{f}_2 \omega_y^2 + 4f_2 \omega_y \dot{\omega}_y \right) + z_i^2 \left(\ddot{f}_2 + 4\dot{f}_2 \omega_z^2 + 4f_2 \omega_z \dot{\omega}_z \right) \right] = 0. \quad (28)$$

Of course, the same result is obtained if the potential function (26) is directly inserted into Eq. (15).

The invariant I_{f_2} for a system determined by the Hamiltonian (1) containing the potential (26) is given by (16), provided that $f_2(t)$ is a solution of (28).

Equation (28) may be cast into a compact form if the sums over the particle coordinates are written in terms of "second beam moments," denoted as $\langle x^2 \rangle$ for the x -direction. The double sum over the Coulomb interaction terms constitutes the electric field energy $W(t)$ of all particles

$$\langle x^2 \rangle(t) = \frac{1}{N} \sum_i x_i^2(t), \quad W(t) = \frac{m}{2} \sum_i \sum_{j \neq i} \frac{c_1}{r_{ij}}.$$

Substituting $\rho^2(t) \equiv f_2(t)$ and defining the function $g = g(t)$ according to

$$g(t) = \langle x^2 \rangle \rho^3 (\ddot{\rho} + \omega_x^2(t)\rho) + \langle y^2 \rangle \rho^3 (\ddot{\rho} + \omega_y^2(t)\rho) + \langle z^2 \rangle \rho^3 (\ddot{\rho} + \omega_z^2(t)\rho), \quad (29)$$

the third-order equation (28) can be transformed into an equivalent coupled system of a second-order equation for $\rho(t)$, and a first order equation for $g(t)$, thereby eliminating the derivatives $\dot{\omega}_{x,y,z}(t)$ of the lattice functions. Solving (29) for $\ddot{\rho}(t)$ means to express it in the form of an "envelope equation"

$$\ddot{\rho} + \omega^2(t) \rho - \frac{g(t)}{\rho^3 (\langle x^2 \rangle + \langle y^2 \rangle + \langle z^2 \rangle)} = 0, \quad (30)$$

with the "average focusing function" $\omega^2(t)$ defined as

$$\omega^2(t) = \frac{\omega_x^2 \langle x^2 \rangle + \omega_y^2 \langle y^2 \rangle + \omega_z^2 \langle z^2 \rangle}{\langle x^2 \rangle + \langle y^2 \rangle + \langle z^2 \rangle}.$$

Eq. (30) is equivalent to (28) if the derivative of $g(t)$ satisfies

$$\dot{g}(t) = 2\rho^3 \left(\langle xp_x \rangle (\ddot{\rho} + \omega_x^2 \rho) + \langle yp_y \rangle (\ddot{\rho} + \omega_y^2 \rho) + \langle zp_z \rangle (\ddot{\rho} + \omega_z^2 \rho) - \frac{W}{mN} \dot{\rho} \right). \quad (31)$$

Expressed in terms of $\rho(t)$ and $g(t)$, the invariant (16) writes

$$I_\rho/N = \left\langle (\rho p_x - \dot{\rho} x)^2 \right\rangle + \left\langle (\rho p_y - \dot{\rho} y)^2 \right\rangle + \left\langle (\rho p_z - \dot{\rho} z)^2 \right\rangle + \rho^2 \frac{2W}{mN} + \frac{g(t)}{\rho^2}.$$

Figure 1 displays the function $\rho(t)$ resulting from a numerical integration of the coupled set of differential equations (30) and (31) for given focusing functions $\omega_x^2(t)$, $\omega_y^2(t)$, $\omega_z^2(t)$ and initial conditions $g(0)$, $\rho(0)$, $\dot{\rho}(0)$. The time-dependent coef-




FIG. 1. ρ versus time as obtained for 2500 simulation particles in a 3D simulation of a periodic focusing lattice with strong Coulomb interaction. τ denotes the focusing period common to all three directions.

ficients contained herein, namely the second-order beam moments as well as the field energy $W(t)$ were obtained from a three-dimensional simulation of a charged particle beam propagating through a periodic focusing lattice with non-negligible

Coulomb interaction, as described by the potential function (26). We observe that for appropriate initial conditions the obtained evolution of $\rho(t)$ is approximately periodic, as imposed by the cell length of the periodic focusing lattice.

With regard to Eq. (30), we may interpret the function $\rho(t)$ as a “generalized beam envelope”. Since the individual inter-particle forces are included in (26), non-Liouvillian effects [4] emerging from the granular nature of charge distributions are also covered.

As an outlook, we point out that a major benefit of our result may be derived in the realm of numerical simulations of systems described by (1). Eq. (16) embodies a time integral of Eq. (15), provided that the phase-space flow of the particle ensemble is *strictly* consistent with the equations of motion (2). This strict consistency can never be accomplished if the time evolution of the particle ensemble is obtained from a computer simulation because of the generally limited accuracy of numerical methods. Under these circumstances, the quantity I_{f_2} as given by Eq. (16) — with $f_2(t)$, $\dot{f}_2(t)$, and $\ddot{f}_2(t)$ following from (15) — can no longer be expected to be strictly constant. The deviation of a numerically obtained I_{f_2} from a constant of motion may thus be used as *a posteriori* error estimation for the respective simulation.

We finally note that the procedure to derive a quantity I that is conserved along a system’s phase-space path can straightforwardly be generalized on the basis of (4) to potentials with quadratic velocity dependence.

-
- [1] H. R. Lewis and P. G. L. Leach, J. Math. Phys. **23**, 2371 (1982).
 - [2] H. R. Lewis, Phys. Rev. Lett. **18**, 510 (1967); J. Math. Phys. **9**, 1976–1986 (1968).
 - [3] E.D. Courant and H.S. Snyder, Ann. Phys. (N.Y.) **3**, 1–48 (1958).
 - [4] J. Struckmeier, Phys. Rev. ST-AB **3**, 034202 (2000).