Cluster radioactivity from trans-tin to superheavy region using an improved empirical formula

G. Saxena^{1,2} and A. Jain^{3,4}

¹ Department of Physics (H&S), Govt. Women Engineering College, Ajmer-305002, India

² Department of Physics, Faculty of Science, University of Zagreb, Bijenička c. 32, 10000 Zagreb, Croatia

³ Department of Physics, School of Basic Sciences, Manipal University Jaipur, Jaipur-303007, India

⁴ Department of Physics, S. S. Jain Subodh P.G.(Autonomous) College, Jaipur-302004, India

Received: date / Revised version: date

Abstract A simple relation $(aZ_c + b)(Z_d/Q)^{1/2} + (cZ_c + d)$ of estimation of the halflife of cluster emission is further improved for cluster and α -decays, separately, by incorporating isospin of parent nucleus as well as angular momentum taken away by the emitted particle. This improved version is not only found robust in producing experimental half-lives belonging to the trans-tin and trans-lead regions but also elucidates cluster emission in superheavy nuclei over the usual α -decay. Considering daughter nuclei around the doubly magic ¹⁰⁰Sn and ²⁰⁸Pb nuclei for trans-tin and trans-lead (including superheavy) parents, respectively, a systematic and extensive study of $56 \le Z \le 120$ isotopes is performed for the light and heavy cluster emissions. A fair competition among cluster emission, α -decay, spontaneous fission, and β decay is observed in this broad range resulting in a substantial probability of C to Sr clusters from several nuclei, which demonstrates the adequacy of shell effects. The present article proposes a single, improved, latest-fitted, and effective formula of cluster radioactivity that can be used to estimate precise half-lives for a wide range of the periodic chart from trans-tin to superheavy nuclei.

PACS. 2 1.10.-k, 21.10.Ft, 21.10.Dr, 21.10.Gv, 21.10.-n, 21.60.Jz

1 Introduction

An exotic decay mode, in which a heavy nucleus fragmentizes itself by emitting a nucleus with the mass between the fission product and α -particle, is envisioned already in 1980 [1] and named 'cluster decay'. After its first experimental confirmation [2] by emission of ¹⁴C from actinide parent nucleus ²²³Ra, this decay mode has been observed from many trans-lead nuclei as the emission of light ¹⁴C to heavy ³²Si clusters [3]. From the last two decades it has gained healthy attention [4–13] due to the direct signature of the crucial role of shell effects as all known cluster emissions have daughter nuclei ²⁰⁸Pb.

Except for the trans-lead region, there are two potential and new islands of cluster emitters that are still less explored and require a comprehensive investigation. The first one is the trans-tin region, regulated by doubly magic nucleus ¹⁰⁰Sn, where only one experimental observation of ¹²C cluster decay from ¹¹⁴Ba exists [14–16], though it is not observed in the later measurement [15]. The other region is the superheavy region in which many heavy clusters are speculated to compete with α -decay [17–27], however, still requires experimental confirmation [28–30]. Therefore, an exhaustive study of cluster radioactivity including trans-tin, trans-lead as well as superheavy regions can lead to important structure information for the detection of heavy and superheavy nuclei in future experimental investigations.

The estimation of the half-life of cluster decay was made in the same decade of its experimental confirmation by several theoretical models which are primarily based on the quantum tunneling effect through a potential barrier [31–35] or cold asymmetric fission process [36, 37]. Alternatively and concomitantly, the very first estimation of the half-life of heavy ion (cluster) emission was given by Poenaru et al. using an analytical formula based on an analytical superheavy asymmetric fission model (ASAFM) [38, 39]. Since then, cluster radioactivity has been studied by several models out of which a few recent ones are the preformedcluster-decay model (PCM) [40, 41], Coulomb and proximity potential model for deformed nuclei [42], Wentzel-Kramers-Brilliouin (WKB) method [43], and modified Gamow-like model [44]. Similarly, by fitting available experimental data, many (semi)empirical/analytical formulas related to cluster decay were proposed from time to time such as Horoi formula [45], BKAG formula [46], RenA formula [47], NRDX formula [48], UDL formula [49], and UNIV formula [50]. Later on, Tavares and Medeiros proposed a semiempirical relationship for the accurate estimation of exotic decay from translead nuclei [51]. Recently, a few of above mentioned semiempirical relationships are modified in Refs. [52–57] which endorse the crucial contributions of asymmetry of parent nucleus and centrifugal barrier effect in the estimation of half-lives of cluster radioactivity. It is, however, important to point out here that almost all the formulas (older or modified) work fairly well in the trans-lead region, but act somewhat unfavourable for the simultaneous estimation of half-lives of cluster emitters in trans-tin and superheavy regions [26,58]. As a matter of fact, especially for the trans-tin region, the UDL formula overestimates the half-life of 114 Ba by four orders of magnitude whereas it works well in the superheavy region due to its fission-like mechanisms [26]. On the other hand, Horoi and NRDX formulas are capable to reproduce the half-life of ¹¹⁴Ba [58,59] but are found not suitable for the heavy clusters in the superheavy region [26]. Therefore, there is a need for a

unique formula that can be applied for the accurate estimation of cluster decay half-life emulated with α -decay half-life for the whole periodic chart (from trans-tin to superheavy nuclei), which is precisely the aim of the present investigation.

In this article, we present a single formula that can be applied for the accurate estimation of half-lives of cluster emission for a systematic study of nuclei in the range $56 \le Z \le 120$ considering daughter nuclei around the doubly magic nuclei ¹⁰⁰Sn and ²⁰⁸Pb for the trans-tin and trans-lead (including superheavy region) regions, respectively. The most probable clusters are found as a result of its contest with α decay, β -decay (referring for β^- , β^+ , and EC throughout the paper), and spontaneous fission (SF) for this wide region of the periodic chart.

2 Formalism

In 2013, Tavares and Medeiros found that experimental cluster decay half-lives show straight lines when plotted as a function of $\xi = (Z_d/Q)^{1/2}$, where Z_d is the atomic number of daughter and Q is the energy of the two-body disintegrating system [51]. A formula (TM), shown below, was proposed for cluster decay half-lives dependent on only 4 parameters and was able to reproduce the majority of the available experimental half-life data reasonably well which belong to the trans-lead region.

$$log_{10}T_{1/2}^{TM}(s) = (aZ_c + b)\sqrt{\frac{Z_d}{Q}} + (cZ_c + d),(1)$$

In our investigation, this relationship while refitted, is also found to reproduce α -decay half-lives of the latest evaluated database NUBASE2020 [60] with a good accuracy for the wide range of the periodic chart. Similar to the crucial roles of (i) angular momentum (l) taken away by the emitted particle, and (ii) isospin (I = (N - Z)/A) of parent nucleus in α -emission [61], these two quantities are reported equally important in determining the cluster decay half-lives by new UDL formula of Soylu and Qi [53] as well as by improved NRDX formula (named as improved unified formula (IUF)) of Ismail *et al.* [54]. In addition, the crucialness of the isospin effect is

Table 1. The coefficients of ITM formula proposed in the present work. The formula is fitted separately for trans-tin and tans-lead regions (see the text for details).

Coefficient	Tra	ns-tin	Trans-lead				
	ITM-CT	ITM-AT	ITM-CL	ITM-AL			
a	$14.5427 {\pm} 0.8701$	294.8314 ± 52.3194	$11.4570 {\pm} 0.0185$	217.4129 ± 42.6150			
b	-1.5239 ± 0.5639	-578.2390 ∓ 104.6389	-8.9167 ± 0.0446	-420.0786 ∓ 85.2188			
с	-3.9917 ± 0.0639	-10.4338 ± 0.1024	-4.2778 ∓ 0.0131	-12.5941 ∓ 0.1005			
d	-109.0040 ∓ 10.4162	-22.8685 ± 0.1952	-103.9499 ∓ 0.2137	-27.1875 ± 0.2003			
e	-10.8146 ∓ 3.6985	$11.8588 \mp 1E-6$	93.2694 ± 0.4313	4.2401 ∓ 0.0845			
f	$0.6419 {\pm} 0.1954$	$0.1495 \pm 1E-6$	$0.3527 {\pm} 0.0009$	0.2630 ± 0.0009			

also probed by improving the semi-empirical formula (ISEM) for the cluster radioactivity in Ref. [56]. Recently, a few semiempirical relationships are modified for the trans-lead region in Ref. [52] which endorse the crucial contributions of asymmetry of parent nucleus and centrifugal barrier effect in estimating half-lives of cluster radioactivity. In order to elucidate the effect of these terms in Eqn. (1), we have initially fitted our data set for (i) cluster decay and (ii) α -decay in the trans-tin region together with (iii) cluster decay and (iv) α -decay in the trans-lead region by incorporating only the *I*-dependent term $(\sqrt{I}(I+1))$, which accounts for the asymmetry. This addition of term has yielded a reduction of RMSE values from 1.53, 1.21, 0.97, and 0.92 to 1.37, 0.74, 0.84, and 0.89, respectively. Subsequently, when we have introduced only the l-dependent term $(\sqrt{l(l+1)})$, which reflects the hindrance effect of the centrifugal barrier, then the RMSE values reduce to 1.01, 1.09, 0.75, and 0.78, respectively. Finally, when both the I and l dependent terms are incorporated in Eqn. (1) then RMSE values significantly reduced to 0.82, 0.71, 0.62, and 0.76, respectively. Hence, the Eqn. (1) can be improvised by adding isospin dependent term as well as a centrifugal barrier term which lead to the following improved formula (named as improved Tavares and Medeiros formula or ITM formula):

$$log_{10}T_{1/2}^{ITM}(s) = (aZ_c + b)\sqrt{\frac{Z_d}{Q}} + (cZ_c + d) + e\sqrt{I(I+1)} + f\sqrt{l(l+1)},$$
(2)

where, a, b, c, d, e, and f are fitting coefficients, tabulated in Table 1. Z_c and Z_d are the proton number of emitted cluster and daughter nucleus, respectively. Q and I are the disintegration energy and isospin, respectively. l is the minimum angular momentum of the cluster particle, which is obtained by following selection rules:

$$l = \begin{cases} \triangle_j & \text{for even } \triangle_j \text{ and } \pi_i = \pi_f \\ \triangle_j + 1 & \text{for even } \triangle_j \text{ and } \pi_i \neq \pi_f \\ \triangle_j & \text{for odd } \triangle_j \text{ and } \pi_i \neq \pi_f \\ \triangle_j + 1 & \text{for odd } \triangle_j \text{ and } \pi_i = \pi_f, \end{cases}$$
(3)

here, $\Delta_j = |j_p - j_d - j_c|$ with j_p , π_i , are the spin and parity values of the parent nucleus, respectively. j_d is the spin of the daughter nucleus. $\pi_f = (\pi_d)(\pi_c)$, in which, π_d and π_c are the parities of the daughter nucleus and cluster, respectively. For the purpose of fitting, the data of spin and parity are taken from NUBASE2020 [60].

Before discussing the results obtained by the ITM formula, there are two important facts that are worth mentioning here. The first fact is related to the inclusion of deformation which is an important ingredient for the estimation of half-lives as also has been described in our recent article [62]. We have followed a similar analysis by incorporating the deformation of the parent nucleus but the reduction in RMSE values is found very insignificant on the cost of the addition of one more parameter. In several articles [42,63], it is demonstrated that in addition to the deformation of the parent nucleus, the deformation of cluster and daughter nuclei needs to be considered to visualize the complete picture of this exotic decay. Such kind of investigation requires a comprehensive and detailed analysis with the inclusion of quadrupole (β_2) and hexadecapole (β_4) deformations, which will

be reported in our subsequent work on the deformation effect of α and cluster decays along with the effect of shape-coexistence on halflives. However, when compared to the half-lives of cluster emission considering the deformation effect from Refs. [42, 63], the present form of the ITM formula results reasonably well with the RMSE value of 0.62 instead of RMSE values 2.30 of Ref. [63] and 0.74 of Ref. [42] on the provided experimental data.

4

Another fact is related to the internal structure of the heavy cluster. For simplicity, in general, cluster formation is assumed similar to the formation of α -particle inside the nucleus but this approximation is valid for the light clusters only. For the heavy clusters, one needs to consider their internal structure together with the neutron-to-proton ratio which eventually can affect cluster preformation in the parent nuclei and consequently the half-life [64]. With this in view, a separate study considering internal structure is required for the clusters with the mass number $A_c>28$.

3 Results and discussion

Considering the role of the shell effect on cluster decay, we divide our region of interest $(56 \le Z \le 120)$ into two parts: (i) the trans-tin region in which daughter nuclei are reckoned around doubly magic nucleus ¹⁰⁰Sn, and (ii) the trans-lead and superheavy regions where daughter nuclei are counted around doubly magic nucleus ²⁰⁸Pb. For the trans-tin region only one data for the cluster (^{114}Ba) is available [58,59], therefore, to fit the formula for this region we include 199 GLDM data [58] for cluster decay. For the α -decay fitting, we use 107 experimental data taken from NUBASE2020 [60]. On the other side, for the trans-lead region, we fit the formula for 37 parent-cluster combinations (total of 61 data points due to multiple Q-value corresponding to various detection systems) [3, 7–10] and 308 α -decay data from NUBASE2020 [60], separately. In this way, the same ITM formula can be utilized to estimate both clusters and α -decay half-lives for both regions (trans-tin and trans-lead) which are speculated to show different decay properties of nuclei [65, 66]. The corresponding coefficients are mentioned in Table 1 for trans-tin

region (ITM-CT and ITM-AT) and for translead region (ITM-CL and ITM-AL), separately. For the nomenclature of the formula 'C' and 'A' refer to cluster and alpha decay whereas last letters 'T' and 'L' refer to trans-tin and trans-lead region, respectively (please see Table 1). Since, in heavy and superheavy regions, uncertainties play a crucial role therefore we have calculated all the fitting parameters with their respective uncertainties by taking into account experimental uncertainties [60]. These uncertainties in parameters are mentioned in Table 1 for all the present formulas i.e. ITM-CT, ITM-AT, ITM-CL, and ITM-AL which lead to uncertainties in the theoretical half-live with ± 0.71 , ± 0.40 , ± 0.56 , and ± 0.41 , respectively.

Table 2. The RMSE and χ^2 of various formulas viz. ITM (present work), IUF [54], ISEF [56], MBKAG [52], UDL [49], TM [51], and New UDL [53] for 62 cluster decay data.

Formula	No. of	RMSE	χ^2
	coefficient		
ITM (present work)	6+6	0.66	0.55
IUF [54]	7	0.87	0.85
ISEF [56]	4	0.99	1.04
MBKÅG [52]	5	1.14	1.41
UDL [49]	3	1.52	2.44
TM [51]	4	1.81	3.51
New UDL [53]	6	1.94	4.16

The applicability of this kind of separate fitting for trans-tin and trans-lead regions can be judged by comparing the results of ITM formula with the few latest and well-established formulas of cluster decay. In Table 2, we list the calculated values of root mean square error (RMSE) as well as χ^2 (formulas are mentioned in Ref. [52]) for ITM and other similar formulas viz. improved unified formula (IUF) [54], improved semi-empirical formula (ISEF) [56], MBKAG [52], UDL [49], TM formula [51], and New UDL [53] formulas. The lowest value of RMSE of ITM formula among all the considered formulas clearly demonstrates its accuracy and improvement over the original TM formula (as RMSE improves from 1.81 to 0.61). In addition to this, the least value of χ^2 exhibits pertinency of ITM formula on the ground of the number of parameters. Further, to analyze the estimation of half-lives using the ITM formula with the available experimental data, we



Figure 1. (Colour online) Comparison of logarithmic half-lives of cluster decay, calculated by using ITM (present work), New UDL [53], and UDL [49] formulas with available experimental data. X-axis refers to the clusters for which experimental data are available. Inset shows one data for trans-tin region i.e. decay of 12 C cluster from 114 Ba.

have plotted half-lives of several clusters calculated by using the ITM formula, New UDL [53], and UDL formula [49] along with their experimental half-lives [3, 7–10] in Fig. 1. The figure contains all the experimental C, N, O, Ne, Mg, and Si clusters (in different colour bands) that belong to the trans-lead region. In the inset, we have depicted the half-life of ¹²C cluster from the only trans-tin parent nucleus ¹¹⁴Ba. A reasonable match of theoretical half-lives from ITM formula with the experimental half-lives endorses the utilization of ITM formula for the unknown region to find the possibility of cluster decay for a wide range of periodic chart from trans-tin to trans-lead as well as the superheavy regions. For C clusters in Fig. 1, the theoretical half-lives of all formulas are found the same for all the points due to the same Q-value (31.85) MeV) for the single parent-cluster combination $^{223}Ra \longrightarrow ^{209} Pb + ^{14}C.$ To explore the full periodic chart, we perform a systematic and comprehensive analysis for all the parent isotopes in the range $56 \le Z \le 120$ using ITM formula by varying their corresponding daughters in such a way that cluster decay leads daughter nuclei around 100 Sn for the trans-tin region and around 208 Pb for rest of the chart (including trans-lead and superheavy regions). As per Eqn. (2), half-lives of cluster decay mainly depend on *Q*-value which is calculated by using the following relation:

$$Q(MeV) = B.E.(d) + B.E.(c) - B.E.(p) +k[Z_n^{\beta} - Z_d^{\beta}], \qquad (4)$$

here, the term $k[Z_p^{\epsilon} - Z_d^{\epsilon}]$ indicates the screening effect caused by the surrounding electrons around the nucleus [67] where k=8.7 eV and ϵ =2.517 for Z \geq 60 and k=13.6 eV and ϵ =2.408 for Z < 60 have been deducted from the data



Figure 2. (Colour online) Cluster decay half-lives for various isotopes of Ba (N_p=56-62), Ce (N_p=61-71), Ra (N_p=113-128), Th (N_p=120-132), Og (N_p=162-192), and Z=120 (N_p=166-192) calculated by the ITM formula.

6

shown by Huang *et al.* [68]. For accurate prediction of theoretical *Q*-values, we have used binding energies (for daughter(d), cluster(c), and parent(p) nuclei) from Weizsacker-Skyrme mass model (WS4) [69] which was found more precise [52] compared to other theories viz. Finite Range Droplet Model (FRDM) [70], nonrelativistic Skyrme Hartree-Fock-Bogoliubov (HFB) [71], and relativistic mean-field theory (RMF) [72–75].

As representative examples for the trans-tin region, in Fig. 2 (a) and (b), we have shown the estimated half-lives of Ba and Ce isotopes by choosing experimentally known isotopes of C and O as clusters, where each line represents half-lives of all considered clusters from the corresponding one parent isotope. Similarly, for the trans-lead region, we have shown plots (c) and (d) for Ra and Th isotopes for the emission of isotopes of C and O, respectively. Towards the superheavy region, heavy clusters Kr and Sr isotopes are estimated in plots (e) and (f) for Z=118 (Og) and Z=120 isotopes, respectively. A parabolic trend is clearly visible from all the plots and with all the extensive analysis in the range $56 \le Z \le 120$ (not shown here), from which one can select a minimum (indicating least halflife) of each parabola corresponding to the potential cluster nucleus for a particular parent nucleus. From this selection, we have extracted the most probable cluster for each considered isotope in the chosen range.

This systematic study indeed provides the most probable cluster for each considered isotope, however, its lowest half-life does not guarantee the chances of cluster decay due to its contest with other probable decay modes. For example, as per experimental systematics, β decay and α -decay are the principal decay modes for the trans-tin region whereas α -decay and spontaneous fission (SF) play the primary roles in the trans-lead and superheavy region. As a result, the chances of cluster decay are always dependent on its competition with other primary decay modes. With this in view, we have calculated half-lives of other decay modes using (i) ITM-AT and ITM-AL formulas (mentioned in Table 1) for the estimation of halflives of α -decay in trans-tin and trans-lead region, respectively, (ii) Fisset and Nix formula [76] for β -decay in trans-tin region, and (iii) Modified Bao Formula (MBF) [77] for spontaneous fission in trans-lead and superheavy regions. These formulas for β -decay and SF are recently been applied and found very successful in determining the respective half-lives in the heavy and superheavy regions [30, 75, 77–79]. The competition can be quantized in the form of branching ratios (BR) which are defined as:

$$BR = \frac{T_{1/2}^{Th.}}{T_{1/2}^{Cluster/\alpha/\beta/SF}},$$
(5)

where, $T_{1/2}^{Th}$ is the total half-life calculated by considering half-lives of all decay modes (cluster, α , and β /SF), and the relation is given by:

$$\frac{1}{T_{1/2}^{Th.}} = \frac{1}{T_{1/2}^{Cluster}} + \frac{1}{T_{1/2}^{\alpha}} + \frac{1}{T_{1/2}^{\beta/SF}},$$
 (6)

where the superscripts refer to the half-lives of concerned decay modes. The superscript β /SF refers to the consideration of β -decay (for transtin region) or SF (for trans-lead and superheavy regions). A good test of our predictions of all the used formulas can be performed using Eqn. (5) from which we can compare values of BR of various decay modes from our theoretical half-lives with the BR available in NUBASE2020 [60]. Taking this into consideration, we have listed the theoretical and experimental BR (in percentage) for several known cluster decay [3, 7, 53, 60] in Table 3. The dominance of α -decay can be easily verified by Table 3 because the value of concern BR is near to 100%. In contrast, the values of BR for cluster decay are very less, however, the experimental values of BR for cluster decay are indeed reproduced very well from the half-life obtained by using the ITM formula. This excellent match endorses our approach of determining BR by using ITM and other formulas for our considered unknown domain of the periodic chart.

In Table 4, a worthful competition among various decay modes viz. cluster decay, α -decay, and β -decay is listed for the trans-tin region in the range 56 \leq Z \leq 81. As mentioned earlier this whole systematic study provides many clusters i.e. one probable cluster for each considered isotope (as obtained by the minimum of each parabola of Fig. 2) but only those are mentioned in the Table 4 which are found with reasonable BR as compared to other decay modes.

Parent	Emitted	Daughter	Expt. BR			Theoretical BR			
nucleus	cluster	nucleus	Cluster	α	SF	Cluster	α	SF	
^{221}Fr	$^{14}\mathrm{C}$	207 Tl	8.8e-11	100.00	0.0048	8e-15	100.00	0.00	
221 Ra	$^{14}\mathrm{C}$	207 Pb	1.2e-10	100.00	-	9e-09	100.00	0.00	
222 Ra	$^{14}\mathrm{C}$	208 Pb	3.0e-8	100.00	-	9e-09	100.00	0.00	
223 Ra	$^{14}\mathrm{C}$	209 Pb	8.9e-8	100.00	-	4e-08	100.00	0.00	
^{223}Ac	$^{14}\mathrm{C}$	²⁰⁹ Bi	-	99.00	-	4e-07	100.00	0.00	
^{223}Ac	^{15}N	208 Pb	-	99.00	-	8e-11	100.00	0.00	
224 Ra	$^{14}\mathrm{C}$	210 Pb	4.0e-9	100.00	-	6e-07	100.00	0.00	
^{225}Ac	$^{14}\mathrm{C}$	211 Bi	5.3e-10	100.00	-	5e-09	100.00	0.00	
226 Ra	$^{14}\mathrm{C}$	212 Pb	2.6e-9	100.00	-	2e-05	100.00	0.00	
$^{226}{ m Th}$	$^{18}\mathrm{O}$	208 Pb	< 3.2e-12	100.00	-	3e-11	100.00	0.00	
$^{228}{ m Th}$	^{20}O	208 Pb	1.13e-11	100.00	-	3e-08	100.00	0.00	
$^{230}{ m Th}$	24 Ne	206 Hg	5.8e-11	100.00	4e-12	4e-10	100.00	0.00	
^{230}U	22 Ne	208 Pb	4.8e-12	100.00	?	6.18e-13	100.00	0.00	
231 Pa	24 Ne	207 Tl	13.4e-10	100.00	<3e-10	6.18e-13	100.00	0.00	
231 Pa	24 Ne	207 Tl	13.4e-10	100.00	< 3e-10	4.91e-11	100.00	0.00	
232 Th	24 Ne	208 Hg	$<\!\!2.78\text{e-}10$	100.00	1.1e-9	3.02e-09	100.00	0.00	
232 Th	26 Ne	206 Hg	$<\!\!2.78\text{e-}10$	100.00	1.1e-9	3.02e-09	100.00	0.00	
^{232}U	24 Ne	$^{208}\mathrm{Pb}$	8.9e-10	100.00	2.7e-12	6.18e-10	100.00	0.00	
^{233}U	24 Ne	209 Pb	7.2e-11	100.00	$<\!\!6e-\!11$	5.21e-10	100.00	0.00	
^{233}U	25 Ne	208 Pb	7.2e-13	100.00	$<\!\!6e-\!11$	5.21e-10	100.00	0.00	
^{233}U	^{28}Mg	205 Hg	< 1.3e-13	100.00	$<\!\!6e-\!11$	2.85e-12	100.00	0.00	
234 U	24 Ne	210 Pb	9e-12	100.00	1.64e-9	5.85e-11	100.00	0.00	
^{234}U	26 Ne	208 Pb	9e-12	100.00	1.64e-9	2.69e-10	99.97	0.03	
234 U	^{28}Mg	206 Hg	1.4e-11	100.00	1.64e-9	3.07e-11	99.97	0.03	
${}^{235}U$	24 Ne	211 Pb	8e-10	100.00	7e-9	2.88e-12	100.00	0.00	
${}^{235}U$	25 Ne	210 Pb	8e-10	100.00	7e-9	1.78e-12	100.00	0.00	
${}^{235}U$	^{28}Mg	207 Hg	8e-10	100.00	7e-9	1.01e-12	100.00	0.00	
^{235}U	^{29}Mg	206 Hg	8e-10	100.00	7e-9	4.61e-13	100.00	0.00	
^{235}U	24 Ne	211 Pb	8e-10	100.00	7e-9	2.87e-12	100.00	0.00	
${}^{236}U$	^{28}Mg	208 Hg	2e-13	100.00	9.4e-8	2.23e-12	99.58	0.42	
${}^{236}U$	^{30}Mg	206 Hg	2e-13	100.00	9.4e-8	1.31e-11	99.58	0.42	
²³⁶ Pu	^{28}Mg	208 Pb	2e-12	100.00	1.9e-7	6.69e-12	100.00	0.00	
^{237}Np	^{30}Mg	207 Tl	$<\!\!4e - 12$	100.00	$<\!2e-10$	1.28e-12	100.00	0.00	
²³⁸ Pu	^{28}Mg	210 Pb	$\approx 6e-15$	100.00	1.9e-7	1.66e-14	99.87	0.13	
²³⁸ Pu	^{30}Mg	208 Pb	$\approx 6e-15$	100.00	1.9e-7	1.50e-13	99.87	0.13	
²³⁸ Pu	^{32}Si	206 Hg	$\approx 1.4 \text{e-} 14$	100.00	1.9e-7	3.28e-14	99.87	0.13	
²⁴⁰ Pu	^{34}Si	206 Hg	$<\!\!1.3e\text{-}11$	100.00	5.796e-6	2.25e-13	93.28	6.72	
$^{241}\mathrm{Am}$	^{34}Si	207 Tl	$<\!\!2.6e-13$	100.00	3.6e-10	2.08e-14	100.00	0.00	
242 Cm	³⁴ Si	208 Pb	1.1e-14	100.00	6.2e-6	$5.54e{-}14$	99.75	0.25	

Table 3. The comparison of branching ratio (BR) with available experimental results [3, 7, 53, 60] for various types of decay viz. cluster decay, α -decay, and SF.

More than 15 cluster decays are found probable in the trans-tin region, and in the future, looking into the experimental progress one may expect more observation of cluster decay in the trans-tin region where so far only one candidate i.e. ¹¹⁴Ba is detected. It is noticeable from Table 4 that two mentioned clusters have values of BR more than 60% or more than α or β decay modes and at the same time the values of BR for a few nuclei are several orders of magnitude larger than the BR of experimentally known clusters (please see Table 3). Therefore, this comparison certainly anticipates cluster decay

Table 4. Competition among various decay modes viz. cluster decay, α -decay, and β -decay for trans-tin nuclei in the range 56 \leq Z \leq 81. Experimental BR and half-life are taken from NUBASE2020 [60] and [59], respectively. Cluster decay, α -decay half-lives are calculated by using ITM-CT and ITM-AT formulas, respectively (present work), and for the β -decay half-lives formula given by Fiset and Nix [76] has been used.

Parent	t Expt. Data		Theoretical Estimates for Cluster Emission									
Nucleus	$\log T_{1/2}$	BR	Emitted	Daughter	Q-value	l	$\log T_{1/2}$ (s)					
	(s)	(in %)	Cluster	Nucleus	(MeV)		Cluster	α	β	Cluster	α	β
^{114}Ba	>4.10	$\beta^+ \approx 100; \beta^+ p = 20;$	^{12}C	^{102}Sn	19.00	0	4.68	2.48	-0.34	0.0010	0.1431	99.8560
		$\alpha = 0.9$; ¹² C<0.0034										
¹¹¹ La	-	-	¹³ N	98 Sn	25.64	1	1.27	-5.00	2.19	0.0001	99.9999	0.0000
¹¹² La	-	-	¹³ N	99 Sn	26.15	3	0.26	-3.09	1.84	0.0438	99.9550	0.0012
¹¹³ La	-	-	¹³ N	100 Sn	26.61	1	-0.51	-1.33	2.41	13.1268	86.8573	0.0160
^{112}Ce	-	-	¹⁷ O	95 Sn	32.55	2	0.85	-5.12	2.77	0.0001	99.9999	0.0000
^{113}Ce	-	-	¹⁷ O	⁹⁶ Sn	33.58	2	-1.05	-3.41	2.13	0.4308	99.5689	0.0003
114 Ce	-	-	¹⁷ O	97 Sn	34.73	2	-3.04	-2.79	2.95	63.9793	36.0206	0.0001
^{115}Ce	-	-	¹⁷ O	98 Sn	35.88	2	-4.84	-2.76	2.36	99.1812	0.8188	0.0000
^{116}Ce	-	-	¹⁶ O	100 Sn	32.30	0	1.92	-2.65	3.29	0.0027	99.9972	0.0001
^{117}Ce	-	-	¹⁶ O	101 Sn	31.45	3	5.06	-0.52	2.58	0.0003	99.9208	0.0790
^{118}Ce	-	-	¹⁶ O	^{102}Sn	30.07	0	5.71	3.22	3.48	0.2104	64.2798	35.5098
114 Pr	-	-	^{15}F	99 Sn	37.38	4	2.62	-2.90	1.63	0.0003	99.9967	0.0030
115 Pr	-	-	^{15}F	100 Sn	37.04	1	4.50	-2.44	2.11	0.0001	99.9953	0.0046
116 Pr	-	-	^{17}F	99 Sn	36.65	6	4.74	-1.51	1.75	0.0001	99.9460	0.0540
117 Pr	-	-	¹⁷ F	100 Sn	36.92	1	5.53	-1.10	2.33	0.0003	99.9383	0.0614
¹¹⁶ Nd	-	-	¹⁸ Ne	⁹⁸ Sn	43.32	0	3.63	-0.54	2.70	0.0067	99.9357	0.0576
¹¹⁷ Nd	-	-	¹⁸ Ne	99 Sn	43.11	2	5.87	1.22	2.08	0.0020	87.8970	12.1011
¹¹⁸ Nd	-	-	¹⁸ Ne	100 Sn	42.68	0	5.42	2.30	2.86	0.0584	78.4906	21.4510
$^{121}\mathrm{Sm}$	-	-	²² Mg	99 Sn	54.21	0	7.51	4.23	1.94	0.0003	0.5094	99.4903

as a prominent mode of decay in several nuclei of the trans-tin region.

In a similar way, the study has been performed for trans-lead nuclei together with superheavy nuclei, and the probable clusters are mentioned in Table 5. Various heavy clusters are found with comparable BR in this region of the periodic chart and show a great probability of decay compared to α -decay and SF, especially for Z=118 and 120 isotopes similar to what was found in Refs. [21–24, 26, 28–30]. In both the tables (Tables 4 and 5), the theoretical half-lives of various decay modes are mentioned in columns 8, 9, and 10 which by using Eqn. (6)provide total half-lives with an excellent agreement with the available experimental half-lives (mentioned in column 2). Hence, the prediction of various cluster decay in the present article is reliable and of great importance due to its sizeable probability over other prominent decay modes of heavy and superheavy regions.

4 Conclusions

An extensive analysis of cluster decay within the range $56 \le Z \le 120$ is carried out using an improved version of a semi-empirical formula

in which isospin and angular momentum effects are included. This improved Tavares and Medeiros formula (ITM) is found with greater accuracy when compared with the available data of cluster decay in the trans-lead region. The applicability of ITM formula is demonstrated by considering daughter nuclei around the double magic nucleus ¹⁰⁰Sn for trans-tin nuclei and ^{208}Pb for trans-lead and superheavy nuclei. The formula is indeed found suitable for the trans-tin and trans-lead regions including the superheavy domain of the periodic chart. Competition of cluster decay mode with other probable decay modes is visualized in the form of branching ratios (BR) which are also found in an excellent match with the available experimental data. Various clusters with substantial BR are reported separately for trans-tin, translead, and superheavy regions. This kind of comprehensive analysis is expected to stimulate the wide region of the periodic chart $(56 \le Z \le 120)$ in view of cluster decay and to enforce theoretical and experimental studies eyeing heavy-ion or cluster decays.

Parent	Expt. Data		Theoretical Estimates for Cluster Emission									
Nucleus	$\log T_{1/2}$	BR	Emitted	Daughter	Q-value	l	lo	${\rm pgT}_{1/2}$ (s)		В	R (in %)	
	(s)	(in %)	Cluster	Nucleus	(MeV)		Cluster	α	SF	Cluster	α	SF
^{285}Mc	-	-	⁸¹ As	204 Pb	272.66	2	2.95	-0.91	6.02	0.01	99.99	0.00
^{289}Mc	-	_	⁸¹ As	²⁰⁸ Pb	272.95	2	4.33	0.16	7.91	0.01	99.99	0.00
^{275}Lv	_	_	⁸⁰ Se	¹⁹⁵ Pb	284 92	6	-3.26	-5.81	-5.24	0.22	78 64	21.14
276 L V			80 So	¹⁹⁶ Pb	284.87	õ	-5.00	-7.47	-6.05	0.40	05.02	3.60
277 L V	_		80 50	197 Dh	204.01	2	3.65	6 10	1.04	0.40	00.71	0.00
278 T	-	-	80 g	198 DI	204.40	4	-3.05	-0.19	-1.04	0.29	99.71	0.00
279 I	-	-	80 g	199 DI	284.28	0	-4.02	-0.77	-5.04	0.10	92.93	0.91
280-LV	-	-	°°Se	200 Pb	284.06	6	-1.23	-4.93	-1.86	0.02	99.89	0.09
200 Lv	-	-	⁸⁰ Se	²⁰⁰ Pb	283.90	0	-3.04	-6.71	-2.25	0.02	99.98	0.00
^{281}Lv	-	-	⁸⁰ Se	201 Pb	283.53	1	-2.00	-3.64	1.02	2.25	97.75	0.00
^{282}Lv	-	-	80 Se	202 Pb	283.39	0	-2.04	-5.68	-0.45	0.02	99.98	0.00
^{283}Lv	-	-	82 Se	201 Pb	283.49	1	-1.18	-4.30	3.33	0.07	99.93	0.00
284 Lv	-	-	82 Se	202 Pb	283.64	0	-1.33	-4.63	1.44	0.05	99.95	0.00
^{285}Lv	-	-	82 Se	203 Pb	283.48	1	-0.39	-3.19	4.69	0.16	99.84	0.00
²⁸⁶ Ly	-	_	⁸² Se	204 Pb	283.41	0	-0.47	-3.55	2.97	0.08	99.92	0.00
²⁸⁷ Ly	_	_	⁸² Se	²⁰⁵ Pb	283.44	1	0.40	-3.47	5.86	0.00	90.02	0.00
288 L V	_		82So	206 pb	200.44	0	0.40	3.47	3.40	0.01	00.08	0.00
289 T	1.00	- 2	82 C -	207 DL	203.00	2	1.00	-3.47	6.64	0.02	99.90	0.00
290 T	-1.60	α:	84 C	206 DI	283.43	3	1.88	-2.55	0.04	0.01	99.99	0.00
291 T	-2.05 ± 0.15	$\alpha \approx 100; SF?$	^{s-} Se	-** PD	283.57	0	0.97	-3.00	4.41	0.01	99.99	0.00
251 Lv	-1.59 ± 0.22	$\alpha \approx 100$; SF?	° ⁴ Se	207 Pb	283.77	1	1.77	-2.20	7.59	0.01	99.99	0.00
279 Ts	-	-	⁸³ Br	196Pb	294.22	0	-6.32	-6.37	-0.88	47.04	52.96	0.00
281 Ts	-	-	^{83}Br	198 Pb	293.88	0	-5.35	-6.13	2.40	14.32	85.68	0.00
^{283}Ts	-	-	^{83}Br	200 Pb	293.58	0	-4.42	-5.54	4.00	7.13	92.87	0.00
285 Ts	-	-	^{83}Br	202 Pb	293.17	0	-3.48	-4.73	5.98	5.25	94.75	0.00
$^{287}\mathrm{Ts}$	-	-	^{85}Br	202 Pb	292.75	0	-2.54	-3.97	7.35	3.58	96.42	0.00
289 Ts	-	-	85 Br	²⁰⁴ Pb	292.92	0	-1.84	-3.81	8.50	1.04	98.96	0.00
^{291}Ts	-2.70	α ?·SF?	⁸⁵ Br	²⁰⁶ Pb	293.18	Ő	-1.18	-3.26	10.10	0.82	99.18	0.00
²⁹³ Ts	-1.60 ± 0.11	$\alpha = 100$	^{85}Br	²⁰⁸ Pb	200.10	0	-0.33	-3.04	9.45	0.02	99.80	0.00
295 Tc	-1.00±0.11	α=100	87Br	208 pb	202.00	2	1.83	0.04	10.12	0.20	00.00	0.00
280	-	-	84 IZ-	196 DL	291.24	2	0.10	-2.34	10.15	0.01	99.99	0.00
281 O	-	-	8617	195 DI	303.72	1	-9.19	-7.07	-1.14	97.09	2.91	0.00
201 Og	-	-	⁸⁴ Kr	100 Pb	304.59	1	-7.87	-5.80	-1.02	99.16	0.84	0.00
²⁸² Og	-	-	° ⁴ Kr	¹⁹⁸ Pb	304.93	0	-8.06	-7.29	-1.80	85.57	14.43	0.00
²⁸³ Og	-	-	⁸⁴ Kr	¹⁹⁹ Pb	304.54	1	-7.01	-4.37	1.54	99.77	0.23	0.00
^{284}Og	-	-	84 Kr	200 Pb	304.28	0	-7.01	-6.82	-0.25	60.71	39.29	0.00
285 Og	-	-	⁸⁶ Kr	199 Pb	304.17	1	-6.06	-6.54	3.42	24.99	75.01	0.00
²⁸⁶ Og	-	-	86 Kr	200 Pb	304.14	0	-6.15	-6.26	1.59	43.80	56.20	0.00
²⁸⁷ Og	-	-	86 Kr	201 Pb	303.98	1	-5.20	-6.04	5.05	12.50	87.50	0.00
288 Og	-	_	84 Kr	204 Pb	302.62	0	-4.83	-5.70	3.17	11.90	88.10	0.00
289 Og	_	_	⁸⁶ Kr	²⁰³ Pb	303 76	1	-4.34	-5.64	6.65	4 75	95.25	0.00
290 Og			86 Kr	204 pb	303.60	0	1.01	5.64	4.50	5.92	04.17	0.00
291 Og	-	-	86 IZ n	205 DL	202.54	2	-4.43	-0.04	4.00	0.00	07.84	0.00
292 O	-	-	8612	206 DI	303.34	0	-2.10	-4.45	1.95	2.10	97.04	0.00
203 O	-	-	87 Kr	206 DI	303.53	0	-3.62	-4.95	0.14	4.44	95.50	0.00
200 Og	-3.00	α:	^{or} Kr	200 Pb	302.51	0	-2.89	-4.07	9.89	6.21	93.79	0.00
204 Og	-3.15 ± 0.20	$\alpha \approx 100; SF?$	^{oo} Kr	200 Pb	302.91	0	-2.66	-4.84	6.15	0.67	99.33	0.00
²⁹⁵ Og	-0.175 ± 0.47	$\alpha \approx 100$	°' Kr	²⁰⁸ Pb	301.96	0	-1.97	-4.26	8.45	0.51	99.49	0.00
²⁹⁶ Og	-	-	⁸⁸ Kr	²⁰⁸ Pb	301.70	0	-1.52	-3.95	5.65	0.37	99.63	0.00
299 Og	-	-	91 Kr	208 Pb	298.62	0	0.62	-3.61	7.57	0.01	99.99	0.00
287120	-	-	88 Sr	199 Pb	325.93	3	-10.67	-5.90	3.70	100.00	0.00	0.00
²⁸⁸ 120	-	-	88 Sr	200 Pb	325.54	0	-11.35	-7.27	2.03	99.99	0.01	0.00
$^{289}120$	-	-	89 Sr	200 Pb	324.18	0	-10.50	-7.23	5.48	99.95	0.05	0.00
²⁸⁹ 120	-	-	90 Sr	$^{199}\mathrm{Pb}$	323.31	1	-9.71	-7.23	5.48	99.67	0.33	0.00
290120	_	_	⁸⁸ Sr	202 Pb	324 70	0	-10.27	-7.19	3 33	99.92	0.08	0.00
291 120			88 Sr	²⁰³ Pb	324.10	1	-0.17	-6.00	7 30	00.02	0.00	0.00
292120	_		88 G.	204 Db	2024.10	0	0.12	6.78	5.30	00.55	0.01	0.00
293 1 20	-	-	90 C	203 DL	220.00	1	- 3.10	-0.78	0.53	00.99	0.40	0.00
294100	-	-	880	206 DI	322.00	1	-7.69	-5.79	9.00	99.22	0.70	0.00
295120	-	-	89 G	206 D	322.79	0	-8.08	-0.38	1.69	98.07	1.93	0.00
200 120	-	-	^{oo} Sr	-~~Pb	322.26	0	-7.53	-5.54	10.85	98.97	1.03	0.00
²³⁰ 120	-	-	⁵⁰ Sr	²⁰⁰ Pb	322.29	0	-7.16	-6.51	7.61	81.86	18.14	0.00
²⁹ ¹ 120	-	-	⁸⁹ Sr	²⁰⁸ Pb	321.13	0	-6.40	-6.16	10.10	63.71	36.29	0.00
298120	-	-	90 Sr	208 Pb	321.34	0	-6.11	-5.91	5.25	61.19	38.81	0.00
$^{299}120$	-	-	91 Sr	208 Pb	320.72	2	-4.68	-6.32	7.91	2.22	97.78	0.00
$^{300}120$	-	-	92 Sr	208 Pb	320.87	0	-5.23	-6.40	4.72	6.34	93.66	0.00
$^{301}120$	-	-	^{93}Sr	$^{208}\mathrm{Pb}$	320.00	0	-4.59	-5.10	7.08	23.54	76.46	0.00
$^{302}120$	_	-	94 Sr	²⁰⁸ Pb	319.86	0	-4.19	-5.65	3.41	3.38	96.62	0.00
303120	_	-	^{95}Sr	²⁰⁸ Ph	318.35	1	-2.84	-4.63	5.39	1.58	98.42	0.00
304120	_	_	96 Sr	²⁰⁸ Pb	317 35	ñ	-2.65	-5.30	1.90	0.18	99.82	0.00
305120	_	_	^{94}Sr	²¹¹ Ph	315.96	0	-1.84	-5.39	0.60	0.03	99.97	0.00

Table 5. Same as Table 4, but for trans-lead and superheavy regions in the range $88 \le Z \le 120$. Cluster decay and α -decay half-lives are calculated by using ITM-CL and ITM-AL formulas, respectively (present work). For the SF half-lives MBF formula [77] has been used.

5 Acknowledgement

GS acknowledges the support provided by SERB (DST), Govt. of India under SIR/2022/000566, and would like to thank Prof. Nils Paar for his kind hospitality at the University of Zagreb, Croatia. AJ is indebted to Prof. S. K. Jain, Manipal University, Jaipur, India for his guidance.

References

- A. Sandulescu, D. Poenaru, W. Greiner, Sov. J. Part. Nucl. II. **11**, 528 (1980).
- 2. H.J. Rose, G.A. Jones, Nature ${\bf 307},\,245$ (1984).
- R. Bonetti, A. Guglielmetti, Rom. Rep. Phys. 59, 301 (2007).
- S. Kumar *et al.*, J. Phys. G: Nucl. Part. Phys. 29, 625 (2003).
- 5. S. Kumar, R. Rani, R. Kumar, J. Phys. G: Nucl. Part. Phys. **36**, 015110 (2009).
- R.K. Gupta, et al., Phys. Rev. C 68, 034321 (2003).
- R.K. Gupta, W. Greiner, Int. J. Mod. Phys. E 03, 335 (1994).
- P.B. Price, Annu. Rev. Nucl. Part. Sci. 39, 19 (1989).
- K.P. Santhosh, B. Priyanka, M.S. Unnikrishnan, Nucl. Phys. A 889, 29 (2012).
- E. Hourani, M. Hussonnois, D.N. Poenaru, Ann. Phys. (Paris) 14, 311 (1989).
- G. Royer, R. Moustabchir, Nucl. Phys. A 683, 182 (2001).
- A. Singh *et al.*, J. Phys. G: Nucl. Part. Phys. 49, 025101 (2022).
- S.N. Kuklin, G.G. Adamian, N.V. Antonenko, Phys. Rev. C 71, 014301 (2005).
- Y.T. Oganessian *et al.*, Z. Fur Phys. A Hadron. Nucl. **349**, 341 (1994).
- A. Guglielmetti *et al.*, Phys. Rev. C. **52**, 740 (1995).
- A. Guglielmetti *et al.*, Nucl. Phys. A 583, 867 (1995).
- K. Prathapan, R.K. Biju, Int. J. Mod. Phys. E 30, 2150106 (2021).
- K.P. Santhosh, T.A. Jose, N. K. Deepak, Phys. Rev. C 105, 054605 (2022).
- G. Royer, Q. Ferrier, M. Pineau, Nucl. Phys. A 1021, 122427 (2022).
- C. Nithya, K.P. Santhosh, Nucl. Phys. A **1020**, 122400 (2022).
- D.N. Poenaru, R.A. Gherghescu, Phys. Rev. C 97, 044621 (2018).

- 22. D.N. Poenaru, H. Stöcker, R.A. Gherghescu, Eur. Phys. J. A 54, 14 (2018).
- Z. Matheson, S.A. Giuliani, W. Nazarewicz, J. Sadhukhan, N. Schunck, Phys. Rev. C 99, 041304 (2019).
- 24. M. Warda, A. Zdeb, L.M. Robledo, Phys. Rev. C 98, 041602(R) (2018).
- D.N. Poenaru, R.A. Gherghescu, W. Greiner, Phys. Rev. C 85, 034615 (2012).
- 26. Y.L. Zhang, Y.Z. Wang, Phys. Rev. C 97, 014318 (2018).
- 27. K.P. Santhosh, R.K. Biju, J. Phys. G: Nucl. Part. Phys. 36, 015107 (2009).
- K.P. Santhosh, C. Nithya, Phys. Rev. C 97, 064616 (2018).
- 29. A. Soylu, F. Koyuncu, Eur. Phys. J. A 55, 118 (2019).
- 30. A. Jain, P.K. Sharma, S.K. Jain, Dashty T. Akrawy, G. Saxena, Phys. Scr. 98, 085304 (2023).
- D.N. Poenaru, M. Ivascu, D. Mazilu, Comput. Phys. Commun. 25, 297 (1982).
- 32. D.S. Delion, J. Suhonen, Phys. Rev. C. 64, 064302 (2001).
- D.N. Poenaru, W. Greiner, Clarendon Press, Oxford, (1996).
- D.N. Poenaru, Institute of Physics Publishing, Bristol, (1996).
- 35. S.K. Arun *et al.*, Phys. Rev. C **79**, 064616 (2009).
- 36. M. Brack *et al.*, Rev. Mod. Phys. **44**, 320 (1972).
- 37. A.C. Wahl, At. Data Nucl. Data Tables **39**, 1 (1988).
- D.N. Poenaru *et al.*, Phys. Rev. C **32**, 572 (1985).
- D.N. Poenaru *et al.*, At. Data Nucl. Data Tables 34, 423 (1986).
- 40. Joshua T. Majekodunmi *et al.*, Phys. Rev. C 105, 044617 (2022).
- W.A Yahya, T.T. Ibrahim, Eur. Phys. J. A 58, 48 (2022).
- 42. S.S. Hosseini, S.M. Motevalli, Phys. Rev. C 107, 034611 (2023).
- R. Dagtas, O. Bayrak, Phys. Scr. 97, 105301 (2022).
- 44. Hong-Ming Liu *et al.*, Phys. Scr. **96**, 125322 (2021).
- 45. Mihai Horoi, J. Phys. G: Nucl. Part. Phys. 30, 945 (2004).
- M. Balasubramaniam *et al.*, Phys. Rev. C 70, 017301 (2004).
- 47. Z. Ren, C. Xu, Z. Wang, Phys. Rev. C 70, 034304 (2004).
- 48. D. Ni, Z. Ren, T. Dong, C. Xu, Phys. Rev. C 78, 044310 (2008).

12G. Saxena and A. Jain: Cluster radioactivity from trans-tin to superheavy region....

- Lett. 103, 072501 (2009).
- 49. C. Qi, F.R. Xu, R.J. Liotta, R. Wyss, Phy. Rev. 76. E.O. Fiset, J.R. Nix, Nucl. Phys. A 193, 647 (1972).
- 50. D.N. Poenaru, R.A. Gherghescu, W. Greiner, Phys. Rev. C 83, 014601 (2011).
- 51. O.A.P. Tavares, E.L. Medeiros, Eur. Phys. J. A 49, 6 (2013).
- 52. A. Jain, P.K. Sharma, S.K. Jain, J.K. Deegwal, G. Saxena, Nucl. Phys. A 1031, 122597 (2023).
- 53. A. Soylu, C. Qi, Nucl. Phys. A **1013**, 122221 (2021).
- 54. M. Ismail, A.Y. Ellithi, A. Adela, M.A. Abbas, Eur. Phys. J. A 58, 225 (2022).
- 55. Y. Wang et al., Chin. Phys. C 45, 044111 (2021).
- 56. S. Cheng et al., Eur. Phys. J. A 58, 168 (2022).
- 57. Lin-Jing Qi et al., Chin. Phys. C 47, 064107 (2023).
- 58. Y. Gao, J. Cui, Y. Wang, J. Gu, Sci Rep 10, 9119 (2020).
- 59. A. Guglielmetti et al., Phys. Rev. C. 56, R2912 (1997)
- 60. F.G. Kondev et al., Chin. Phys. C 45, 030001 (2021).
- 61. P.K. Sharma, A. Jain, G. Saxena, Nucl. Phys. A **1016**, 122318 (2021).
- 62. G. Saxena et al., J. Phys. G: Nucl. Part. Phys. **50**, 015102 (2023).
- 63. A. Soylu et al., Eur. Phys. J. A 48, 128 (2012).
- 64. K. Wei, H.F. Zhang, Phys. Rev. C 96, 021601(R) (2017).
- 65. G. Saxena, A. Jain, P.K. Sharma, Prafulla Saxena, Communicated to Scientific Reports (2023).
- 66. Jun-Gang Deng, Hong-Fei Zhang, Phys. Lett. B 816, 136247 (2021).
- 67. V.Yu. Denisov, A.A. Khudenko, Phys. Rev. C 79, 054614 (2009).
- 68. K.N. Huang et al., At. Data Nucl. Data Tables 18, 243 (1976).
- 69. N. Wang, M. Liu, X. Wu, J. Meng, Phys. Lett. B 734, 215 (2014).
- 70. P. Möller, M.R. Mumpower, T. Kawano, W.D. Myers, At. Data Nucl. Data Tables 125, 1 (2019).
- 71. J. Dobaczewski, M.V. Stoitsov, W. Nazarewicz, AIP Conference Proceedings 726, 51 (2004).
- 72. G. Saxena, M. Kumawat, M. Kaushik, S.K. Jain, Mamta Aggarwal, Phys. Lett. B 788, 1 (2019).
- 73. G. Saxena, M. Kumawat, M. Kaushik, U.K. Singh, S.K. Jain, S. Somorendro Singh, M. Aggarwal, Int. J. Mod. Phys. E 26, 1750072 (2017).
- 74. G. Saxena, M. Kumawat, M. Kaushik, S.K. Jain, M. Aggarwal, Phys. Lett. B 775, 126 (2017)
- 75. U.K. Singh, P.K. Sharma, M. Kaushik, S.K. Jain, Dashty T Akrawy, G. Saxena, Nucl. Phys. A 1004, 122035 (2020).

- 77. G. Saxena, P.K. Sharma, P. Saxena, J. Phys. G: Nucl. Part. Phys. 48, 055103 (2021).
- 78. R. Sharma, A. Jain, P.K. Sharma, S.K. Jain, G. Saxena, Phys. Scr. 97, 045307 (2022).
- 79. G. Saxena, A. Jain, P.K. Sharma, Phys. Scr. 96, 125304 (2021).