DISCRETE WEAK DUALITY OF HYBRID HIGH-ORDER METHODS FOR CONVEX MINIMIZATION PROBLEMS

NGOC TIEN TRAN

ABSTRACT. This paper derives a discrete dual problem for a prototypical hybrid high-order method for convex minimization problems. The discrete primal and dual problem satisfy a weak convex duality that leads to a priori error estimates with convergence rates under additional smoothness assumptions. This duality holds for general polyhedral meshes and arbitrary polynomial degrees of the discretization. A novel postprocessing is proposed and allows for a posteriori error estimates on regular triangulations into simplices using primal-dual techniques. This motivates an adaptive mesh-refining algorithm, which performs superiorly compared to uniform mesh refinements.

1. INTRODUCTION

1.1. **Overview.** Given an open bounded polyhedral Lipschitz domain $\Omega \subset \mathbb{R}^n$ with the boundary $\partial \Omega$ and unit normal vector ν , we consider the convex minimization problem of the energy

(1.1)
$$E(v) \coloneqq \int_{\Omega} (\Psi(\mathrm{D}v) + \psi(x, v)) \,\mathrm{d}x \quad \text{among } v \in V_{\mathrm{D}} \coloneqq W_{\mathrm{D}}^{1, p}(\Omega; \mathbb{R}^m),$$

 $p \in (1,\infty)$. Here, $\Psi : \mathbb{M} \to \mathbb{R} \cup \{+\infty\}$ with $\mathbb{M} := \mathbb{R}^{m \times n}$ denotes a proper lower semicontinuous convex energy density, $\psi : \Omega \times \mathbb{R}^m \to \mathbb{R} \cup \{+\infty\}$ is measurable such that $\psi(x, \bullet)$ is a proper lower semicontinuous convex function for a. e. $x \in \Omega$, and $V_{\mathrm{D}} = W_{\mathrm{D}}^{1,p}(\Omega; \mathbb{R}^m)$ is the space of functions in the Sobolev space $W^{1,p}(\Omega; \mathbb{R}^m)$ with homogenous boundary data on a non-empty closed part $\Gamma_{\mathrm{D}} \subset \partial\Omega$ of $\partial\Omega$. Let Ψ^* (resp. ψ^*) denote the convex conjugate of Ψ (resp. ψ a.e. in Ω) and $p' := p/(p-1) \in$ $(1,\infty)$ with 1/p + 1/p' = 1 is the Hölder conjugate of p. The dual problem of (1.1) maximizes the dual energy

(1.2)
$$E^*(\tau) \coloneqq -\int_{\Omega} (\Psi^*(\tau) + \psi^*(x, \operatorname{div} \tau)) \, \mathrm{d}x$$

among $\tau \in W_{\mathrm{N}} := W_{\mathrm{N}}^{p'}(\operatorname{div},\Omega;\mathbb{M})$, the Sobolev space of functions $\tau \in L^{p'}(\Omega;\mathbb{M})$ with a distributional divergence $\operatorname{div}\tau$ belonging to $L^{p'}(\Omega;\mathbb{R}^m)$ and the normal trace $\tau\nu$ vanishes along the Neumann boundary $\Gamma_{\mathrm{N}} := \partial\Omega \setminus \Gamma_{\mathrm{D}}$ in a weak sense. The weak duality $E^*(\tau) \leq E(v)$ holds for any $v \in V_{\mathrm{D}}$ and $\tau \in W_{\mathrm{N}}$.

It is well-established that duality relations on the discrete level can be utilized in the error analysis of nonconforming methods [14, 5, 6, 18]. Only a few methods in the literature are known to preserve the weak duality on the discrete level,

(1.3)
$$\max E_h^*(W_N(\mathcal{M})) \le \min E_h(V_D(\mathcal{M}))$$

Date: 9th April 2024.

²⁰²⁰ Mathematics Subject Classification. 65N12, 65N30, 65Y20.

Key words and phrases. discrete weak convex duality, hybrid high-order, convex minimization, a priori, a posteriori, adaptive mesh-refining.

This work has been supported by the European Union's Horizon 2020 research and innovation programme (project DAFNE, grant agreement No. 891734, and project RandomMultiScales, grant agreement No. 865751.

for an appropriate discretization $V_{\rm D}(\mathcal{M})$ of $V_{\rm D}$ (resp. $W_{\rm N}(\mathcal{M})$ of $W_{\rm N}$) and discrete energy E_h (resp. discrete dual energy E_h^*). Historically, (1.3) was first known for a Crouzeix-Raviart finite element method (FEM) for the primal problem and a discrete Raviart-Thomas FEM for the dual problem [41, 14, 6]. The generalized Marini identity even leads to the strong duality max $E_h^*(W_{\rm N}(\mathcal{M})) = \min E_h(V_{\rm D}(\mathcal{M}))$ under additional assumptions on Ψ and ψ . For higher-order approximations, the unstabilized hybrid high-order (HHO) method from [1] and the classical Raviart-Thomas FEM satisfy (1.3) without a discrete strong duality in general [18]. Finally, (1.3) holds for the class of lowest-order discontinuous Galerkin (dG) and Raviart-Thomas FEM from [5]. The analysis of all three aforementioned methods requires a regular triangulation of Ω into simplices (i.e., no hanging nodes are allowed) because they heavily utilize the structure of Crouzeix-Raviart or Raviart-Thomas functions.

1.2. Motivation and description of main results. The method from [5] stands out as the only one with a stabilization. The later is required for the design of hybridizable methods, e.g., HHO or hybrid discontinuous Galerkin (HDG) methods, on general polyhedral meshes. The foundation for establishing (1.3) is an integration by parts on the discrete level and the resulting integrals on the skeleton of the mesh determine the stabilizations. In [5], the dG methodology leads to stabilizations that penalize the average of the trace on the primal and the normal trace on the dual level. These terms do not possess any convergence rates for smooth functions – an undesired effect that requires overpenalization and projections onto the Crouzeix-Raviart and the Raviart-Thomas finite element space in the error analysis. Therefore, it only applies to the lowest-order case on regular triangulations into simplices. Hybridizable methods, on the other hand, provide further flexibility due to additional degrees of freedom on the skeleton, but can they overcome these restrictions?

This paper provides a clear affirmative answer. In fact, we consider a prototypical HHO method [28, 27, 24], which is also related to the HDG methodology [22]. The ansatz space $V(\mathcal{M}) \coloneqq P_{k+1}(\mathcal{M}; \mathbb{R}^m) \times P_k(\Sigma; \mathbb{R}^m)$ features a split of degrees of freedom into piecewise polynomials of degree at most k + 1 on the (polyhedral) mesh \mathcal{M} and piecewise polynomials of degree at most k on the skeleton Σ , the set of faces of \mathcal{M} . The discrete scheme replaces the gradient in (1.1) by a reconstruction operator $D_h : V(\mathcal{M}) \to P_k(\mathcal{M}; \mathbb{M})$. In contrast to [18], this operator has a non-trivial kernel and so, a stabilization $s_h : V(\mathcal{M}) \to \mathbb{R}_{\geq 0}$ is required. This paper utilizes the nonlinear version of the Lehrenfeld-Schöberl stabilization, but the analysis can be extended to the classical HHO stabilization as well. Given $r \in (1, \infty)$ with Hölder conjugate r' = r/(r-1), the discrete primal problem minimizes

(1.4)
$$E_h(v_h) \coloneqq \int_{\Omega} (\Psi(\mathbf{D}_h v_h) + \psi_h(x, \Pi^k_{\mathcal{M}} v_{\mathcal{M}})) \, \mathrm{d}x + \mathbf{s}_h(v_h)/r$$

among $v_h = (v_{\mathcal{M}}, v_{\Sigma}) \in V_{\mathcal{D}}(\mathcal{M})$, where $V_{\mathcal{D}}(\mathcal{M}) \subset V(\mathcal{M})$ is a discrete analogue to $V_{\mathcal{D}}$ and $\psi_h : \Omega \times \mathbb{R}^m \to \mathbb{R} \cup \{+\infty\}$ is an approximation of ψ . Our analysis departs from a discrete integration by parts involving three different operators: a gradient reconstruction $D_h : V(\mathcal{M}) \to P_k(\mathcal{M}; \mathbb{M})$, a divergence reconstruction $\operatorname{div}_h : W(\mathcal{M}) \to P_k(\mathcal{M}; \mathbb{R}^m)$, which maps from the ansatz space $W(\mathcal{M}) \coloneqq P_k(\mathcal{M}; \mathbb{M}) \times P_k(\Sigma; \mathbb{R}^m)$ of $W^{p'}(\operatorname{div}, \Omega; \mathbb{M})$ onto the space of piecewise polynomials of degree k, and a potential reconstruction $\mathcal{R}_h^* : W(\mathcal{M}) \to P_k(\mathcal{M}; \mathbb{M})$ in the spirit of [30, 29] on the dual level. This leads to a discrete dual problem that maximizes

(1.5)
$$E_h^*(\tau_h) \coloneqq -\int_{\Omega} (\Psi^*(\mathcal{R}_h^*\tau_h) + \psi_h^*(x, \operatorname{div}_h \tau_h)) \,\mathrm{d}x - \gamma_h(\tau_h)/r'$$

among $\tau_h = (\tau_M, \tau_\Sigma) \in W_N(\mathcal{M})$ with a suitable stabilization $\gamma_h : W(\mathcal{M}) \to \mathbb{R}_{\geq 0}$ and a discrete analogue $W_N(\mathcal{M}) \subset W(\mathcal{M})$ of W_N . We establish that E_h from (1.4) and E_h^* from (1.5) satisfy (1.3). This duality leads to a priori error estimates with convergence rates under suitable smoothness assumptions. Apart from a priori results, this paper proposes a novel $W^{p'}(\operatorname{div}, \Omega, \mathbb{M})$ conforming postprocessing σ_0 for the approximation of the dual variable on regular triangulations into simplices. Its construction is based on the discrete Euler-Lagrange equations, which allow for the concept of equilibrated tractions in the HHO methodology [23, 1]. This is a cheap alternative to equilibrium techniques [11, 35, 36] in the linear case and can be used to obtain a posteriori error estimates via primal-dual gap similar to [42, 4, 14, 18, 8].

1.3. Notation. Standard notation for Sobolev and Lebesgue spaces applies throughout this paper with the abbreviation $V \coloneqq W^{1,p}(\Omega; \mathbb{R}^m)$, $W \coloneqq W^{p'}(\operatorname{div}, \Omega; \mathbb{M})$, and $\| \bullet \|_p \coloneqq \| \bullet \|_{L^p(\Omega)}$ for any $p \in (1, \infty)$. In particular, the gradient of scalarvalued (resp. vector- or matrix-valued) functions is denoted by ∇ (resp. D). For any open bounded domain ω , $c_{\mathrm{P}}(\omega)$ is the constant in the Poincaré inequality $\|v\|_{L^p(\omega)} \leq c_{\mathrm{P}}(\omega)\operatorname{diam}(\omega)\|\nabla v\|_{L^p(\omega)}$ for any $v \in W^{1,p}(\omega)$ with $\int_{\omega} v \, dx = 0$. If ω is convex, then $c_{\mathrm{P}}(\omega) \leq 2(p/2)^{1/p}$ [21, Theorem 1.2] for $1 and <math>c_{\mathrm{P}}(\omega) = 1/\pi$ for p = 2 [9]. Given $A, B \in \mathbb{M} = \mathbb{R}^{m \times n}$, A : B denotes the Euclidean scalar product of A and B, which induces the Frobenius norm $|A| \coloneqq (A : A)^{1/2}$ in \mathbb{M} . The subdifferential $\partial \varphi(A)$ of a convex function $\varphi : \mathbb{M} \to \mathbb{R} \cup \{+\infty\}$ at $A \in \mathbb{M}$ is

(1.6)
$$\partial \varphi(A) \coloneqq \{G \in \mathbb{M} : 0 \le \varphi(B) - \varphi(A) - G : (B - A) \text{ for any } B \in \mathbb{M}\}.$$

The convex conjugate $\varphi^* : \mathbb{M} \to \mathbb{R} \cup \{+\infty\}$ of φ is defined by

$$\varphi^*(G) \coloneqq \sup_{A \in \mathbb{M}} (A : G - \varphi(A)) \quad \text{for any } G \in \mathbb{M}.$$

Given $g \in \mathbb{R}^m$, the (convex) indicator function $\chi_g(a)$ is 0 if a = g and $+\infty$ if $a \neq g$. The notation $A \leq B$ abbreviates $A \leq CB$ for a generic constant C independent of the mesh-size and $A \approx B$ abbreviates $A \leq B \leq A$.

1.4. **Outline.** The remaining parts of this paper are organized as follows. Section 2 provides detailed information on the discretization including reconstruction operators and stabilizations utilized in (1.4)–(1.5) and throughout this paper. The discrete weak duality (1.3) is established in Section 3. Section 4 is devoted to the application of (1.3) in the a priori error analysis, while a posteriori error estimates are established in Section 5. Some numerical benchmarks conclude this paper.

2. The HHO methodology

This section recalls the HHO methodology from [28, 27, 24].

2.1. **Polytopal Mesh.** Let \mathcal{M} be a finite collection of closed polytopes of positive volume with overlap of measure zero that covers $\overline{\Omega} = \bigcup_{K \in \mathcal{M}} K$. A face S of the mesh \mathcal{M} is a closed connected subset of a hyperplane H_S with positive (n-1)dimensional surface measure such that either (a) there exist $K_+, K_- \in \mathcal{M}$ with $S \subset H_S \cap K_+ \cap K_-$ (interior face) or (b) there exists $K_+ \in \mathcal{M}$ with $S \subset H_S \cap K_+ \cap \partial \Omega$ (boundary face). Let Σ be a finite collection of faces with overlap of (n-1)dimensional surface measure zero that covers the skeleton $\partial \mathcal{M} := \bigcup_{K \in \mathcal{M}} \partial K =$ $\bigcup_{S \in \Sigma} S$ with the split $\Sigma = \Sigma(\Omega) \cup \Sigma(\partial \Omega)$ into the set of interior faces $\Sigma(\Omega)$ and the set of boundary faces $\Sigma(\partial \Omega)$. Let $\Sigma_D := \{S \in \Sigma : S \subset \Gamma_D\}$ (resp. $\Sigma_N :=$ $\Sigma(\partial \Omega) \setminus \Sigma_D$) denote the set of Dirichlet (resp. Neumann) faces. For $K \in \mathcal{M}, \Sigma(K)$ is the set of all faces of K. The normal vector ν_S of an interior face $S \in \Sigma(\Omega)$ is fixed beforehand and set $\nu_S := \nu|_S$ for boundary faces $S \in \Sigma(\partial \Omega)$. For $S \in$

N. T. TRAN

$$\begin{split} &\Sigma(\Omega), \ K_+(S) \in \mathcal{M} \ (\text{resp. } K_-(S) \in \mathcal{M}) \ \text{denotes the unique cell with } S \subset \partial K_+(S) \\ &(\text{resp. } S \subset \partial K_-(S)) \ \text{and } \nu_{K_+(S)}|_S = \nu_S \ (\text{resp. } \nu_{K_-(S)}|_S = -\nu_S). \ \text{For } S \in \Sigma(\partial \Omega), \\ &K_+(S) \in \mathcal{M} \ \text{is the unique cell with } S \subset \partial K_+(S). \ \text{The explicit reference to } S \\ &\text{is omitted if there is no likelihood of confusion. The jump } [v]_S \ \text{and the average} \\ &\{v\}_S \ \text{of any function } v \in W^{1,1}(\text{int}(T_+ \cup T_-); \mathbb{R}^m) \ \text{along } S \in \Sigma(\Omega) \ \text{are defined by} \\ &[v]_S \coloneqq v|_{K_+} - v|_{K_-} \in L^1(S; \mathbb{R}^m) \ \text{and} \ \{v\}_S \coloneqq (v|_{K_+} + v|_{K_-})/2 \in L^1(S; \mathbb{R}^m). \ \text{If} \\ &S \in \Sigma(\partial \Omega), \ \text{then } [v]_S \coloneqq v|_S = v_{K_+}|_S \eqqcolon \{v\}_S. \end{split}$$

For theoretical purposes, let ϑ denote the mesh regularity parameter of \mathcal{M} associated with a matching simplicial submesh, cf. [26, Definition 1.38] for a precise definition. The constants in discrete inequalities such as the trace or inverse inequality depend on this parameter. The notation $c_{\rm P} := \max_{K \in \mathcal{M}} c_{\rm P}(K)$ denotes the best possible local Poincaré constant on \mathcal{M} . For general polyhedral meshes, $c_{\rm P}$ may depend on ϑ . The differential operators $D_{\rm pw}$ and div_{pw} denote the piecewise version of D and div without explicit reference to the underlying mesh.

2.2. Finite element spaces. Given a subset $M \subset \mathbb{R}^n$ of diameter h_M , let $P_k(M)$ denote the space of polynomials of degree at most k. The vector- and matrix-valued version of $P_k(M)$ read $P_k(M; \mathbb{R}^m) \coloneqq P_k(M)^m$ and $P_k(M; \mathbb{M}) \coloneqq P_k(M; \mathbb{R}^n)^m$. This notation applies to other spaces as well. For any $v \in L^1(M)$, $\prod_M^k v \in P_k(M)$ denotes the L^2 projection of v onto $P_k(M)$. The space of piecewise polynomials of degree at most k with respect to the mesh \mathcal{M} or the faces Σ is denoted by $P_k(\mathcal{M})$ or $P_k(\Sigma)$. Given $v \in L^1(\Omega)$, we define the L^2 projection $\prod_{\mathcal{M}}^k v$ of v onto $P_k(\mathcal{M})$ by $(\prod_{\mathcal{M}}^k v)|_K = \prod_K^k v|_K$ in any cell $K \in \mathcal{M}$. We recall the stability of the L^2 projection $\|\prod_{\mathcal{M}}^k \xi\|_p \lesssim \|\xi\|_p$ for any $\xi \in L^p(\Omega)$ from, e.g., [23, Lemma 3.2]. A consequence of this is the quasi-best approximation

(2.1)
$$\|(1 - \Pi_{\mathcal{M}}^{k})\xi\|_{p} \leq \|\xi - \varphi_{k}\|_{p} + \|\Pi_{\mathcal{M}}^{k}(\varphi_{k} - \xi)\|_{p} \lesssim \|\xi - \varphi_{k}\|_{p}$$

for any $\varphi_k \in P_k(\mathcal{M})$, where $\varphi_k = \prod_{\mathcal{M}}^k \varphi_k$ is utilized in the penultimate inequality.

For a simplex T, $\operatorname{RT}_k(T) \coloneqq P_k(T; \mathbb{R}^n) + xP_k(T)$ denotes the space of Raviart-Thomas finite element functions. The piecewise version of this on a regular triangulation \mathcal{T} into simplices is called $\operatorname{RT}_k^{\operatorname{pw}}(\mathcal{T})$, while $\operatorname{RT}_k(\mathcal{T}) \coloneqq \operatorname{RT}_k^{\operatorname{pw}}(\mathcal{T}) \cap W$ denotes the space of globally $W^{p'}(\operatorname{div}, \Omega)$ Raviart-Thomas functions. The mesh \mathcal{M} gives rise to the piecewise constant function $h_{\mathcal{M}} \in P_0(\mathcal{M})$ with $h_{\mathcal{M}}|_K = h_K$; $h_{\max} \coloneqq \max_{K \in \mathcal{M}} h_K$ is the maximal mesh-size of \mathcal{M} . For any $f \in L^{p'}(\Omega; \mathbb{R}^m)$, the oscillation of f reads $\operatorname{osc}(f, \mathcal{M}) \coloneqq \|h_{\mathcal{M}}(1 - \Pi_{\mathcal{M}}^k)f\|_{p'}$.

2.3. **Discrete ansatz spaces.** Given a fixed $k \in \mathbb{N}_0$, let $V(\mathcal{M}) \coloneqq P_{k+1}(\mathcal{M}; \mathbb{R}^m) \times P_k(\Sigma; \mathbb{R}^m)$ (resp. $W(\mathcal{M}) \coloneqq P_k(\mathcal{M}; \mathbb{M}) \times P_k(\Sigma; \mathbb{R}^m)$) denote the discrete ansatz space for V (resp. W). For any $v_h = (v_{\mathcal{M}}, v_{\Sigma})$, we use the notation $v_K \coloneqq v_{\mathcal{M}}|_K$ and $v_S \coloneqq v_{\Sigma}|_S$ to abbreviate the restriction of $v_{\mathcal{M}}$ in a cell $K \in \mathcal{M}$ and v_{Σ} along a face $S \in \Sigma$. The same notation applies to discrete objects in $W(\mathcal{M})$. The set of Dirichlet faces Σ_D (resp. Neumann faces Σ_N) gives rise to the subspace $P_k(\Sigma \setminus \Sigma_D; \mathbb{R}^m)$ (resp. $P_k(\Sigma \setminus \Sigma_N; \mathbb{R}^m)$) of $P_k(\Sigma; \mathbb{R}^m)$ with the convention $v_{\Sigma} \in P_k(\Sigma \setminus \Sigma_D; \mathbb{R}^m)$ (resp. $\tau_{\Sigma} \in P_k(\Sigma \setminus \Sigma_N; \mathbb{R}^m)$) if $v_S \equiv 0$ (resp. $\tau_S \equiv 0$) on Dirichlet faces $S \in \Sigma_D$ (resp. Neumann faces $S \in \Sigma_N$) to model homogenous Dirichlet boundary condition (resp. vanishing normal traces). This defines the discrete spaces $V_D(\mathcal{M}) \coloneqq P_{k+1}(\mathcal{M}; \mathbb{R}^m) \times P_k(\Sigma \setminus \Sigma_D; \mathbb{R}^m)$ and $W_N(\mathcal{M}) \coloneqq P_k(\mathcal{M}; \mathbb{M}) \times P_k(\Sigma \setminus \Sigma_N; \mathbb{R}^m)$ as subspaces of $V(\mathcal{M})$ and $W(\mathcal{M})$. The interpolation operators $I_V : V \to V(\mathcal{M})$ and $I_W : W^{1,1}(\Omega; \mathbb{M}) \to W(\mathcal{M})$ map $v \in V$ to $I_V v \coloneqq (\Pi_{\mathcal{M}}^{\mathcal{H}+1}v, (\Pi_S^k v)_{S\in\Sigma}) \in V(\mathcal{M})$ and $\tau \in W^{1,1}(\Omega; \mathbb{M})$ to $I_W \tau \coloneqq (\Pi_{\mathcal{M}}^k \tau, (\Pi_S^k \tau v_S)_{S\in\Sigma}) \in W(\mathcal{M})$.

2.4. Reconstruction operators and stabilization. The reconstruction operators D_h and div_h provide an approximation of the differential operators D and div.

Gradient reconstruction. The gradient reconstruction $D_h v_h \in P_k(\mathcal{M}; \mathbb{M})$ of $v_h = (v_{\mathcal{M}}, v_{\Sigma}) \in V(\mathcal{M})$ is the unique solution to, for all $\tau_k \in P_k(\mathcal{M}; \mathbb{M})$,

(2.2)
$$\int_{\Omega} \mathcal{D}_h v_h : \tau_k \, \mathrm{d}x = -\int_{\Omega} v_{\mathcal{M}} \cdot \operatorname{div}_{\mathrm{pw}} \tau_k \, \mathrm{d}x + \sum_{S \in \Sigma} \int_S v_S \cdot [\tau_k \nu_S]_S \, \mathrm{d}s.$$

Divergence reconstruction. The divergence reconstruction $\operatorname{div}_h \tau_h \in P_k(\mathcal{M}; \mathbb{R}^m)$ of $\tau_h = (\tau_{\mathcal{M}}, \tau_{\Sigma}) \in W(\mathcal{M})$ is the unique solution to, for all $v_k \in P_k(\mathcal{M}; \mathbb{R}^m)$,

(2.3)
$$\int_{\Omega} \operatorname{div}_{h} \tau_{h} \cdot \varphi_{k} \, \mathrm{d}x = -\int_{\Omega} \operatorname{D}_{\mathrm{pw}} \varphi_{k} : \tau_{\mathcal{M}} \, \mathrm{d}x + \sum_{S \in \Sigma} \int_{S} [\varphi_{k}]_{S} \cdot \tau_{S} \, \mathrm{d}s.$$

Potential reconstruction. The potential reconstruction $\mathcal{R}_h v_h \in P_{k+1}(\mathcal{M}; \mathbb{R}^m)$ of $v_h = (v_{\mathcal{M}}, v_{\Sigma}) \in V(\mathcal{M})$ satisfies, for any $\varphi_{k+1} \in P_{k+1}(\mathcal{M}; \mathbb{R}^m)$, that

(2.4)
$$\int_{\Omega} D_{pw} \mathcal{R}_h v_h : D_{pw} \varphi_{k+1} \, dx$$
$$= -\int_{\Omega} v_{\mathcal{M}} \cdot \Delta_{pw} \varphi_{k+1} \, dx + \sum_{S \in \Sigma} \int_S v_S \cdot [D_{pw} \varphi_{k+1} \nu_S]_S \, ds.$$

This defines $\mathcal{R}_h v_h$ uniquely up to piecewise constants, which are fixed by

(2.5)
$$\int_{K} \mathcal{R}_{h} v_{h} \, \mathrm{d}x = \int_{K} v_{K} \, \mathrm{d}x \quad \text{for any } K \in \mathcal{M}.$$

The following lemma states the characteristic commuting property of the reconstruction operators in the HHO methodology [28, 27, 24].

Lemma 2.1 (commuting property). Any $v \in V$ and $\tau \in W^{1,1}(\Omega; \mathbb{M})$ satisfy $D_h I_V v = \Pi^k_{\mathcal{M}} Dv$, $\operatorname{div}_h I_W \tau = \Pi^k_{\mathcal{M}} \operatorname{div} \tau$, and the orthogonality $D_{pw}(v - \mathcal{R}_h I_V v) \perp D_{pw} P_{k+1}(\mathcal{M}; \mathbb{R}^m)$ with respect to the L^2 scalar product.

The key for establishing the discrete weak duality (1.3) is the design of a reconstruction operator on the dual level in the spirit of [30, 29]. The latter can be computed explicitly, although in this paper, it is only used for theoretical purposes.

Dual potential reconstruction. Let $Z(\mathcal{M})$ denote the L^2 orthogonal complement of $D_{pw}P_{k+1}(\mathcal{M};\mathbb{R}^m)$ in $P_k(\mathcal{M};\mathbb{M})$,

(2.6)
$$P_k(\mathcal{M};\mathbb{M}) = \mathcal{D}_{\mathrm{pw}} P_{k+1}(\mathcal{M};\mathbb{R}^m) \oplus Z(\mathcal{M}),$$

and $\Pi_{Z(\mathcal{M})}$ is the L^2 projection that maps from $P_k(\mathcal{M}; \mathbb{M})$ onto $Z(\mathcal{M})$. The dual potential reconstruction $\mathcal{R}_h^* \tau_h \in P_k(\mathcal{M}; \mathbb{M})$ of $\tau_h = (\tau_{\mathcal{M}}, \tau_{\Sigma}) \in W(\mathcal{M})$ is the unique function in $P_k(\mathcal{M}; \mathbb{M})$ that satisfies

(2.7)
$$\int_{\Omega} \mathcal{R}_h^* \tau_h : \mathcal{D}_{pw} v_{k+1} \, \mathrm{d}x = -\int_{\Omega} v_{k+1} \cdot \operatorname{div}_h \tau_h \, \mathrm{d}x + \sum_{S \in \Sigma} \int_S [v_{k+1}]_S \cdot \tau_S \, \mathrm{d}s$$

for any $v_{k+1} \in P_{k+1}(\mathcal{M}; \mathbb{R}^m)$ and the projection property

(2.8)
$$\Pi_{Z(\mathcal{M})} \mathcal{R}_h^* \tau_h = \Pi_{Z(\mathcal{M})} \tau_{\mathcal{M}}$$

Lemma 2.2 (well-definedness of \mathcal{R}_h^*). For any $\tau_h = (\tau_{\mathcal{M}}, \tau_{\Sigma}) \in W(\mathcal{M})$, there exists a unique function $\mathcal{R}_h^* \tau_h \in P_k(\mathcal{M}; \mathbb{M})$ with (2.7)–(2.8).

Proof. Let $\Phi(\tau_h) \in P_{k+1}(\mathcal{M}; \mathbb{R}^m) / P_0(\mathcal{M}; \mathbb{R}^m)$ be the unique solution to

(2.9)
$$\int_{\Omega} \mathcal{D}_{pw} \Phi(\tau_h) : \mathcal{D}_{pw} v_{k+1} \, \mathrm{d}x = -\int_{\Omega} v_{k+1} \cdot \operatorname{div}_h \tau_h \, \mathrm{d}x + \sum_{S \in \Sigma} \int_S [v_{k+1}]_S \cdot \tau_S \, \mathrm{d}s$$

for any $v_{k+1} \in P_{k+1}(\mathcal{M}; \mathbb{R}^m)/P_0(\mathcal{M}; \mathbb{R}^m)$. In fact, the right-hand side of (2.7) vanishes for any piecewise constant test function $v_{k+1} \in P_0(\mathcal{M}; \mathbb{R}^m)$ thanks to the definition (2.3) of div_h. Therefore, it is a (well-defined) linear functional in the

quotient space $P_{k+1}(\mathcal{M}; \mathbb{R}^m)/P_0(\mathcal{M}; \mathbb{R}^m)$. Thus, $\Phi(\tau_h)$ is the Riesz representation of this linear functional in $P_{k+1}(\mathcal{M}; \mathbb{R}^m)/P_0(\mathcal{M}; \mathbb{R}^m)$ equipped with the piecewise energy scalar product. Hence, $\mathcal{R}_h^* \tau_h \coloneqq D_{pw} \Phi(\tau_h) + \prod_{Z(\mathcal{M})} \tau_{\mathcal{M}}$ satisfies (2.7)–(2.8). The uniqueness follows from the uniqueness of $\Phi(\tau_h)$. \Box

Stabilization. The kernel of D_h is non-trivial because any $v_h = (v_{\mathcal{M}}, 0) \in V_D(\mathcal{M})$ with $v_{\mathcal{M}} \perp P_{k-1}(\mathcal{M}; \mathbb{R}^m)$ satisfies $D_h v_h \equiv 0$, but v_h may not be zero. Hence, $\|D_h \bullet\|_p$ is not a norm in $V_D(\mathcal{M})$ and so, a stabilization s_h is required. Fix the parameters $r \in (1, \infty)$ and $s \in \mathbb{R}$. Given any $u_h = (u_{\mathcal{M}}, u_{\Sigma}), v_h = (v_{\mathcal{M}}, v_{\Sigma}) \in$ $V(\mathcal{M})$ and $\tau_h = (\tau_{\mathcal{M}}, \tau_{\Sigma}) \in W(\mathcal{M})$, define

(2.10)
$$\mathbf{s}_h(u_h; v_h) \coloneqq \sum_{K \in \mathcal{M}} \sum_{S \in \Sigma(K)} h_S^{-s} \int_S T_{K,S} u_h \cdot (v_S - \Pi_S^k v_K) \, \mathrm{d}s$$

(2.11)
$$\gamma_h(\tau_h) \coloneqq \sum_{K \in \mathcal{M}} \sum_{S \in \Sigma(K)} h_S^{s/(r-1)} \| \tau_S - (\mathcal{R}_h^* \tau_h)|_K \nu_K \|_{L^{r'}(S)}^{r'}$$

with the notation, for any $K \in \mathcal{M}$ and $S \in \Sigma(K)$,

(2.12)
$$T_{K,S}u_h \coloneqq \Pi_S^k(|u_S - \Pi_S^k u_K|^{r-2}(u_S - \Pi_S^k u_K))$$

and the abbreviation $s_h(v_h; v_h) \coloneqq s_h(v_h)$. In the quadratic case r = 2, s_h is known under the label *Lehrenfeld-Schöberl stabilization* in the HDG methodology. While the reconstruction operators and stabilization of this section are defined globally, they can be computed cell-wise and their computation is carried out in parallel.

3. DISCRETE WEAK CONVEX DUALITY

This section establishes the discrete weak duality (1.3).

Theorem 3.1 (weak duality). The discrete energy E_h from (1.4) and the discrete dual energy E_h^* from (1.5) satisfy (1.3).

The point of departure is the following result.

Lemma 3.2 (discrete integration by parts). Any $v_h = (v_M, v_\Sigma) \in V_D(\mathcal{M})$ and $\tau_h = (\tau_M, \tau_\Sigma) \in W_N(\mathcal{M})$ satisfy

(3.1)
$$\int_{\Omega} \mathcal{D}_{h} v_{h} : \mathcal{R}_{h}^{*} \tau_{h} \, \mathrm{d}x = \sum_{S \in \Sigma \setminus \Sigma_{\mathrm{D}}} \int_{S} (v_{S} - \Pi_{S}^{k} \{v_{\mathcal{M}}\}_{S}) \cdot [\mathcal{R}_{h}^{*} \tau_{h} \nu_{S}]_{S} \, \mathrm{d}s$$
$$+ \sum_{S \in \Sigma \setminus \Sigma_{\mathrm{N}}} \int_{S} \Pi_{S}^{k} [v_{\mathcal{M}}]_{S} \cdot (\tau_{S} - \{\mathcal{R}_{h}^{*} \tau_{h} \nu_{S}\}_{S}) \, \mathrm{d}s - \int_{\Omega} v_{\mathcal{M}} \cdot \operatorname{div}_{h} \tau_{h} \, \mathrm{d}x.$$

Proof. A piecewise integration by parts shows

$$(3.2) \qquad -\int_{\Omega} v_{\mathcal{M}} \cdot \operatorname{div}_{\mathrm{pw}} \mathcal{R}_{h}^{*} \tau_{h} \, \mathrm{d}x = \int_{\Omega} \mathrm{D}_{\mathrm{pw}} v_{\mathcal{M}} : \mathcal{R}_{h}^{*} \tau_{h} \, \mathrm{d}x$$
$$(3.2) \qquad -\sum_{S \in \Sigma \setminus \Sigma_{\mathrm{D}}} \int_{S} \{v_{\mathcal{M}}\}_{S} \cdot [\mathcal{R}_{h}^{*} \tau_{h} \nu_{S}]_{S} \, \mathrm{d}s - \sum_{S \in \Sigma \setminus \Sigma_{\mathrm{N}}} \int_{S} [v_{\mathcal{M}}]_{S} \cdot \{\mathcal{R}_{h}^{*} \tau_{h} \nu_{S}\}_{S} \, \mathrm{d}s.$$

This and the definition (2.2) of the gradient reconstruction D_h prove

$$\int_{\Omega} \mathcal{D}_{h} v_{h} : \mathcal{R}_{h}^{*} \tau_{h} \, \mathrm{d}x = \int_{\Omega} \mathcal{D}_{pw} v_{\mathcal{M}} : \mathcal{R}_{h}^{*} \tau_{h} \, \mathrm{d}x + \sum_{S \in \Sigma \setminus \Sigma_{D}} \int_{S} (v_{S} - \Pi_{S}^{k} \{v_{\mathcal{M}}\}_{S}) \cdot [\mathcal{R}_{h}^{*} \tau_{h} \nu_{S}]_{S} \, \mathrm{d}s - \sum_{S \in \Sigma \setminus \Sigma_{N}} \int_{S} [v_{\mathcal{M}}]_{S} \cdot \{\mathcal{R}_{h}^{*} \tau_{h} \nu_{S}\}_{S} \, \mathrm{d}s.$$

Since $\tau_S - \{\mathcal{R}_h^* \tau_h \nu_S\}_S \in P_k(S; \mathbb{R}^m)$ along $S \in \Sigma$, the combination of this with the definition (2.7) of \mathcal{R}_h^* concludes the proof.

Proof of Theorem 3.1. Let any $v_h = (v_{\mathcal{M}}, v_{\Sigma}) \in V_{\mathcal{D}}(\mathcal{M})$ and $\tau_h = (\tau_{\mathcal{M}}, \tau_{\Sigma}) \in W_{\mathcal{N}}(\mathcal{M})$ be given. Since Ψ is convex, the Fenchel-Young inequality [43, p. 105] proves $D_h v_h : \mathcal{R}_h^* \tau_h \leq \Psi(D_h v_h) + \Psi^*(\mathcal{R}_h^* \tau_h)$ pointwise a.e. in Ω . An integration of this over the domain Ω provides

(3.3)
$$0 \leq \int_{\Omega} (\Psi(\mathbf{D}_h v_h) + \Psi^*(\mathcal{R}_h^* \tau_h) - \mathbf{D}_h v_h : \mathcal{R}_h^* \tau_h) \, \mathrm{d}x.$$

Elementary algebra shows the identities

$$(v_S - \Pi_S^k \{ v_{\mathcal{M}} \}_S) \cdot [\mathcal{R}_h^* \tau_h \nu_S]_S + \Pi_S^k [v_{\mathcal{M}}]_S \cdot (\tau_S - \{ \mathcal{R}_h^* \tau_h \nu_S \}_S) = (v_S - \Pi_S^k v_{K_-}) \cdot (\tau_S - (\mathcal{R}_h^* \tau_h)|_{K_-} \nu_S) - (v_S - \Pi_S^k v_{K_+}) \cdot (\tau_S - (\mathcal{R}_h^* \tau_h)|_{K_+} \nu_S)$$

on any interior face $S \in \Sigma(\Omega)$,

$$\Pi_S^k[v_{\mathcal{M}}]_S \cdot (\tau_S - \{\tau_{\mathcal{M}}\nu_S\}_S) = -(v_S - \Pi_S^k v_{\mathcal{M}}) \cdot (\tau_S - \mathcal{R}_h^* \tau_h \nu_S)$$

on any Dirichlet face $S \in \Sigma_D$, and

$$(v_S - \Pi_S^k \{ v_{\mathcal{M}} \}_S) \cdot [\mathcal{R}_h^* \tau_h \nu_S]_S = -(v_S - \Pi_S^k v_{\mathcal{M}}) \cdot (\tau_S - \mathcal{R}_h^* \tau_h \nu_S)$$

on any Neumann face $S \in \Sigma_N$. Thus, the first two integrals on the right-hand side of (3.1) are equal to

$$-\sum_{K\in\mathcal{M}}\sum_{S\in\Sigma(K)}(\nu_S\cdot\nu_K)\int_S(v_S-\Pi_S^kv_K)\cdot(\tau_S-(\mathcal{R}_h^*\tau_h)|_K\nu_S)\,\mathrm{d}s.$$

This is bounded by $s_h(v_h)/r + \gamma_h(\tau_h)/r'$ due to the Hölder, Cauchy, and Young inequalities. Hence, (3.1) and $\operatorname{div}_h \tau_h \in P_k(\mathcal{M}; \mathbb{R}^m)$ provide

$$-\int_{\Omega} \mathcal{D}_h v_h : \mathcal{R}_h^* \tau_h \, \mathrm{d}x \le \int_{\Omega} \Pi_{\mathcal{M}}^k v_{\mathcal{M}} \cdot \operatorname{div}_h \tau_h \, \mathrm{d}x + \mathbf{s}_h(v_h)/r + \gamma_h(v_h)/r'.$$

The combination of this with (3.3) results in

$$0 \leq \int_{\Omega} (\Psi(\mathbf{D}_h v_h) + \Psi^*(\mathcal{R}_h^* \tau_h) + \Pi_{\mathcal{M}}^k v_{\mathcal{M}} \cdot \operatorname{div}_h \tau_h) \, \mathrm{d}x + \frac{1}{r} \mathbf{s}_h(v_h) + \frac{1}{r'} \gamma_h(\tau_h).$$

This and $\Pi^k_{\mathcal{M}} v_{\mathcal{M}} \cdot \operatorname{div}_h \tau_h \leq \psi_h(x, \Pi^k_{\mathcal{M}} v_{\mathcal{M}}) + \psi^*_h(x, \operatorname{div}_h \tau_h)$ for a.e. $x \in \Omega$ from the Fenchel-Young inequality [43, p. 105] conclude the proof.

The general idea of this paper can be applied to different choices of stabilizations, e.g., the classical HHO stabilization from [28, 27, 23, 24].

Example 3.3 (classical stabilization). The HHO method from [23] utilizes the ansatz space $V(\mathcal{M}) = P_k(\mathcal{M}; \mathbb{R}^m) \times P_k(\Sigma; \mathbb{R}^m)$ and a stabilization of the form

(3.4)
$$\widetilde{\mathbf{s}}_h(v_h) \coloneqq \sum_{K \in \mathcal{M}} \sum_{S \in \Sigma(K)} h_S^{-s} \| \Pi_S^k(v_S - (\mathcal{P}_h v_h)|_K) \|_{L^r(S)}^r$$

for any $v_h = (v_{\mathcal{M}}, v_{\Sigma}) \in V(\mathcal{M})$ with $\mathcal{P}_h v_h \coloneqq v_{\mathcal{M}} + (1 - \Pi_{\mathcal{M}}^k)\mathcal{R}_h v_h \in P_{k+1}(\mathcal{M}; \mathbb{R}^m)$ and the potential reconstruction \mathcal{R}_h from (2.4)–(2.5). Observe that $v_{\mathcal{M}} - \mathcal{P}_h v_h \perp P_k(\mathcal{M}; \mathbb{R}^m)$ and so, it is possible to replace $v_{\mathcal{M}}$ by $\mathcal{P}_h v_h$ throughout the proof of Lemma 3.2. This leads to the discrete integration by parts formula

$$\int_{\Omega} \mathcal{D}_{h} v_{h} : \mathcal{R}_{h}^{*} \tau_{h} \, \mathrm{d}x = \sum_{S \in \Sigma \setminus \Sigma_{D}} \int_{S} (v_{S} - \Pi_{S}^{k} \{\mathcal{P}_{h} v_{h}\}_{S}) \cdot [\mathcal{R}_{h}^{*} \tau_{h} \nu_{S}]_{S} \, \mathrm{d}s$$
$$+ \sum_{S \in \Sigma \setminus \Sigma_{N}} \int_{S} \Pi_{S}^{k} [\mathcal{P}_{h} v_{h}]_{S} \cdot (\tau_{S} - \{\mathcal{R}_{h}^{*} \tau_{h} \nu_{S}\}_{S}) \, \mathrm{d}s - \int_{\Omega} v_{\mathcal{M}} \cdot \operatorname{div}_{h} \tau_{h} \, \mathrm{d}x$$

From this, we deduce that (1.3) holds for E_h from (1.4) with s_h replaced by \tilde{s}_h and E_h^* from (1.3) (without any modification).

N. T. TRAN

4. A priori

This section establishes error estimates using the weak duality (1.3). For simplicity, we assume the explicit representation of the lower-order term

(4.1)
$$\psi(x,a) \coloneqq -f(x) \cdot a \quad \text{and} \quad \psi_h(x,a) \coloneqq -f_h(x) \cdot a$$

with a volume force $f \in L^{p'}(\Omega; \mathbb{R}^m)$ and $f_h \coloneqq \Pi^k_{\mathcal{M}} f \in P_k(\mathcal{M}; \mathbb{R}^m)$. Then

$$E^*(\tau) = -\int_{\Omega} \Psi^*(\tau) \, \mathrm{d}x - \chi_{-f}(\mathrm{div}\tau) \qquad \text{for any } \tau \in W_{\mathrm{N}},$$
$$E^*_h(\tau_h) = -\int_{\Omega} \Psi^*(\mathcal{R}^*_h\tau_h) \, \mathrm{d}x - \chi_{-f_h}(\mathrm{div}_h\tau_h) \qquad \text{for any } \tau_h \in W_{\mathrm{N}}(\mathcal{M}).$$

Let $u \in \arg\min E(V_{\rm D})$ (resp. $\sigma \in \arg\max E^*(W_{\rm N})$) be a minimizer (resp. maximizer) of E (resp. E^*) in $V_{\rm D}$ (resp. $W_{\rm N}$). The error analysis of this paper applies to convex minimization problems without a duality gap

(4.2)
$$\max E^*(W_N) = E^*(\sigma) = E(u) = \min E(V_D) < \infty.$$

Recall the subdifferential of a convex function from (1.6).

Theorem 4.1 (a priori). Suppose that (4.1)–(4.2) hold. If $\sigma \in W^{1,1}(\Omega; \mathbb{M})$, then any $\xi \in L^{p'}(\Omega; \mathbb{M})$ with $\xi \in \partial \Psi(D_h I_V u)$ and $\varrho \in L^p(\Omega; \mathbb{M})$ with $\varrho \in \partial \Psi^*(\mathcal{R}_h^* I_W \sigma)$ a.e. in Ω satisfy

$$E_{h}(\mathbf{I}_{V}u) - \min E_{h}(V_{\mathbf{D}}(\mathcal{M})) \leq -\int_{\Omega} (1 - \Pi_{\mathcal{M}}^{k})\xi : (1 - \Pi_{\mathcal{M}}^{k})\mathbf{D}u \,\mathrm{d}x + \int_{\Omega} (\mathbf{D}_{h}\mathbf{I}_{V}u - \varrho) : (\sigma - \mathcal{R}_{h}^{*}\mathbf{I}_{W}\sigma) \,\mathrm{d}x + \int_{\Omega} (f - f_{h}) \cdot (u - \mathcal{R}_{h}\mathbf{I}_{V}u) \,\mathrm{d}x (4.3) + s_{h}(\mathbf{I}_{V}u)/r + \gamma_{h}(\mathbf{I}_{W}\sigma)/r' - \sum_{S \in \Sigma \setminus \Sigma_{N}} \int_{S} [\mathcal{R}_{h}\mathbf{I}_{V}u]_{S} \cdot (1 - \Pi_{S}^{k})\sigma\nu_{S} \,\mathrm{d}x.$$

Proof. The proof departs from the split

(4.4)
$$E_h(\mathbf{I}_V u) - E_h(u_h) \le E_h(\mathbf{I}_V u) - E(u) + E^*(\sigma) - E_h^*(\mathbf{I}_W \sigma).$$

Since $0 \leq \Psi(\mathrm{D}u) - \Psi(\mathrm{D}_h\mathrm{I}_V u) - \xi : (\mathrm{D}u - \mathrm{D}_h\mathrm{I}_V u)$ a.e. in Ω from $\xi \in \partial \Psi(\mathrm{D}_h\mathrm{I}_V u)$, the projection property $\mathrm{D}_h\mathrm{I}_V u = \Pi^k_{\mathcal{M}}\mathrm{D}u$ from Lemma 2.1 imply

(4.5)
$$E_h(\mathbf{I}_V u) - E(u) \le -\int_{\Omega} (\xi : (1 - \Pi_{\mathcal{M}}^k) \mathbf{D}u - (f - f_h) \cdot u) \, \mathrm{d}x + \mathbf{s}_h(\mathbf{I}_V u)/r.$$

The property $\rho \in \partial \Psi^*(\mathcal{R}_h^* \mathbf{I}_W \sigma)$ provides $0 \leq \Psi^*(\sigma) - \Psi^*(\mathcal{R}_h^* \mathbf{I}_W \sigma) - \rho : (\sigma - \mathcal{R}_h^* \mathbf{I}_W \sigma)$ a.e. in Ω . Since $\operatorname{div}_h \mathbf{I}_W \sigma = \prod_{\mathcal{M}}^k \operatorname{div} \sigma = -f_h$ from Lemma 2.1, this shows

(4.6)
$$E^*(\sigma) - E_h^*(\mathbf{I}_W \sigma) \le -\int_{\Omega} \varrho : (\sigma - \mathcal{R}_h^* \mathbf{I}_W \sigma) \, \mathrm{d}x + \gamma_h(\mathbf{I}_W \sigma)/r'.$$

The L^2 orthogonality $D_h I_V u - D_{pw} \mathcal{R}_h I_V u \perp D_{pw} P_{k+1}(\mathcal{M}; \mathbb{M})$ from Lemma 2.1, $D_h I_V u \in P_k(\mathcal{M}; \mathbb{M})$, and $\Pi_{Z(\mathcal{M})}(\Pi^k_{\mathcal{M}} \sigma - \mathcal{R}^*_h I_W \sigma) = 0$ from (2.8) lead to

(4.7)
$$\int_{\Omega} (\sigma - \mathcal{R}_h^* \mathbf{I}_W \sigma) : \mathbf{D}_h \mathbf{I}_V u \, \mathrm{d}x = \int_{\Omega} (\sigma - \mathcal{R}_h^* \mathbf{I}_W \sigma) : \mathbf{D}_{\mathrm{pw}} \mathcal{R}_h \mathbf{I}_V u \, \mathrm{d}x.$$

The definition (2.7) of \mathcal{R}_h^* , a piecewise integration by parts, div $\sigma = -f$, and div_hI_W $\sigma = -f_h$ prove

$$\int_{\Omega} (\sigma - \mathcal{R}_{h}^{*} \mathbf{I}_{W} \sigma) : \mathbf{D}_{h} \mathbf{I}_{V} u \, \mathrm{d}x$$

$$(4.8) \qquad = \int_{\Omega} (f - f_{h}) \cdot \mathcal{R}_{h} \mathbf{I}_{V} u \, \mathrm{d}x + \sum_{S \in \Sigma \setminus \Sigma_{N}} \int_{S} [\mathcal{R}_{h} \mathbf{I}_{V} u]_{S} \cdot (\sigma - \Pi_{S}^{k} \sigma) \nu_{S} \, \mathrm{d}x.$$

The combination of this with (4.4)–(4.6) concludes the proof.

Remark 4.2 (lowest-order case). If k = 0, then it is possible to choose $\xi \in P_0(\mathcal{M}; \mathbb{M})$ (because $D_h I_V u \in P_0(\mathcal{M}; \mathbb{M})$). In this case, the L^2 orthogonality $u - \mathcal{R}_h I_V u \perp P_0(\mathcal{M}; \mathbb{R}^m)$ from (2.5) imply the simplified version

$$E_{h}(\mathbf{I}_{V}u) - \min E_{h}(V_{\mathrm{D}}(\mathcal{M})) \leq \int_{\Omega} (\mathbf{D}_{h}\mathbf{I}_{V}u - \varrho) : (\sigma - \mathcal{R}_{h}^{*}\mathbf{I}_{W}\sigma) \,\mathrm{d}x + s_{h}(\mathbf{I}_{V}u)/r + \gamma_{h}(\mathbf{I}_{W}\sigma)/r' - \sum_{S \in \Sigma \setminus \Sigma_{\mathrm{N}}} \int_{S} [\mathcal{R}_{h}\mathbf{I}_{V}u]_{S} \cdot (1 - \Pi_{S}^{k})\sigma\nu_{S} \,\mathrm{d}x.$$

Theorem 4.1 applies to minimization problems (1.1) with energy densities Ψ that satisfy a two-sided growth condition, i.e., there exist positive constants $0 < c_1 \leq c_2$ and nonnegative constants c_3 , $c_4 \geq 0$ such that

(4.9)
$$c_1|A|^p - c_3 \le \Psi(A) \le c_2|A|^p + c_4 \text{ for any } A \in \mathbb{M}.$$

An immediate consequence of (4.9) is the a priori bound $\|Du\|_p + \|\sigma\|_{p'} \leq 1$ on the continuous level. (For instance, $\|Du\|_p \leq c_5$ with the positive root c_5 of the scalar-valued function $t \mapsto c_1 t^p - c_P t \|f\|_{p'} - c_3 |\Omega|$ [16].) Lemma 4.2 from [18] implies that a selection of $\xi \in L^{p'}(\Omega; \mathbb{M})$ and $\varrho \in L^p(\Omega; \mathbb{M})$ in Theorem 4.1 with $\xi \in \partial \Psi(D_h I_V u)$ and $\varrho \in \partial \Psi^*(\mathcal{R}_h^* I_W \sigma)$ a.e. in Ω is always possible. The two-sided growth (4.9) provides the bound

(4.10)
$$\|\xi\|_{p'}^{p} \lesssim 1 + \|\mathbf{D}_h\mathbf{I}_V u\|_p^p \text{ and } \|\varrho\|_p^p \lesssim 1 + \|\mathcal{R}_h\mathbf{I}_W\sigma\|_{p'}^{p}$$

cf. [18, Lemma 2.1(c)] for explicit constants hidden in the notation \leq .

Corollary 4.3 (convergence rates). Suppose that (4.1)–(4.2) hold and Ψ satisfy (4.9). Let r = p and $p - 1 - (k + 1) \leq s \leq (k + 2)(p - 1)$. Under the smoothness assumptions $u \in V_{\mathrm{D}} \cap W^{k+2,p}(\mathcal{M};\mathbb{R}^m)$ and $\sigma \in W^{1,1}(\Omega;\mathbb{M}) \cap W^{k+1,p'}(\mathcal{M};\mathbb{M})$, it holds $E_h(\mathbf{I}_V u) - E_h(u_h) \leq h_{\max}^{k+1}$.

Proof. The proof of $\|\sigma - \mathcal{R}_h^* \mathrm{I}_W \sigma\|_{p'} + \gamma(\mathrm{I}_W \sigma) \lesssim h_{\max}^{k+1}$ can follow the arguments of [29, Section 6]. For the convenience of the reader, an alternative proof is provided below. Define $\sigma_k := \Pi_{\mathcal{M}}^k \sigma - \mathcal{R}_h^* \mathrm{I}_W \sigma \in P_k(\mathcal{M}; \mathbb{M})$. Given $\Phi_k \in P_k(\mathcal{M}; \mathbb{M})$, let $\mathrm{D}_{\mathrm{pw}} v_{k+1}$ for some $v_{k+1} \in P_{k+1}(\mathcal{M}; \mathbb{R}^m)$ denote the L^2 projection of Φ_k onto $\mathrm{D}_{\mathrm{pw}} P_{k+1}(\mathcal{M}; \mathbb{R}^m)$. Since $\Pi_{Z(\mathcal{M})} \sigma_k = 0$ by the definition (2.8) of \mathcal{R}_h^* , it holds $\int_{\Omega} \sigma_k : \Phi_k \, \mathrm{d}x = \int_{\Omega} \sigma_k : \mathrm{D}_{\mathrm{pw}} v_{k+1} \, \mathrm{d}x$. By replacing $\mathrm{D}_h \mathrm{I}_V u$ by Φ_k in (4.7)–(4.8), we deduce from the Hölder and the Poincaré inequalities that

$$\int_{\Omega} \sigma_k : \Phi_k \, \mathrm{d}x = -\int_{\Omega} (1 - \Pi_{\mathcal{M}}^k) \mathrm{div} \sigma \cdot v_{k+1} \, \mathrm{d}x + \sum_{S \in \Sigma \setminus \Sigma_N} \int_S [v_{k+1}]_S \cdot (1 - \Pi_S^k) \sigma \nu_S \, \mathrm{d}s$$
$$\lesssim \operatorname{osc}(\operatorname{div}\sigma, \mathcal{M}) \| \mathcal{D}_{pw} v_{k+1} \|_p + \sum_{S \in \Sigma \setminus \Sigma_N} \int_S [v_{k+1}]_S \cdot (1 - \Pi_S^k) \sigma \nu_S \, \mathrm{d}s.$$

For any $S \in \Sigma \setminus \Sigma_N$ and $K \in \mathcal{M}$ with $S \in \Sigma(K)$, (2.1), a discrete trace, and a Poincaré inequality reveal $\|(1 - \Pi_S^k)\sigma\|_{L^{p'}(S)} \lesssim \|\sigma - (\Pi_K^k\sigma)|_S\|_{L^{p'}(S)} \lesssim h_K^{1/p}\|\mathcal{D}(1 - \Pi_K^k)\sigma\|_{L^{p'}(K)}$. This and the same argument applied to $(1 - \Pi_S^k)[v_{k+1}]_S$ lead to

$$\sum_{S \in \Sigma \setminus \Sigma_{\mathbf{N}}} \int_{S} [v_{k+1}]_{S} \cdot (1 - \Pi_{S}^{k}) \sigma \nu_{S} \, \mathrm{d}s \lesssim \|\mathbf{D}_{\mathrm{pw}} v_{k+1}\|_{p} \|h_{\mathcal{M}} \mathbf{D}_{\mathrm{pw}} (1 - \Pi_{\mathcal{M}}^{k}) \sigma\|_{p'}$$

Since $\|\sigma_k\|_{p'} = \sup_{\Phi \in L^p(\Omega; \mathbb{M}) \setminus \{0\}} \int_{\Omega} \sigma_k : \Pi^k_{\mathcal{M}} \Phi \, dx / \|\Phi\|_p$, the combination of the two previously displayed formula with $\|D_{pw} v_{k+1}\|_p \lesssim \|\Phi_k\|_p \lesssim \|\Phi\|_p$ for $\Phi_k := \Pi^k_{\mathcal{M}} \Phi$ from the stability of L^2 projections yields

(4.11)
$$\|\sigma_k\|_{p'} \lesssim \operatorname{osc}(\operatorname{div}\sigma, \mathcal{M}) + \|h_{\mathcal{M}} \mathcal{D}_{\mathrm{pw}}(1 - \Pi_{\mathcal{M}}^k)\sigma\|_{p'} \lesssim h_{\max}^{k+1}.$$

The triangle inequality and the stability of the L^2 projection Π_S^k show $\|\Pi_S^k(\tau - (\mathcal{R}_h^* \mathbf{I}_W \tau)|_K) \nu_S\|_{L^{p'}(S)} \lesssim \|\tau - \Pi_K^k \tau\|_{L^{p'}(S)} + \|\sigma_k|_K \nu_S\|_{L^{p'}(S)}$ for any $K \in \mathcal{M}$ and $S \in \Sigma(K)$. Therefore, the trace inequality, its discrete version, and the Poincaré inequality provide

(4.12)
$$\gamma_h(\mathbf{I}_W \sigma) \lesssim h_{\max}^{s/(p-1)+p'-1} (\|\mathbf{D}_{pw}(1-\Pi_{\mathcal{M}}^k)\sigma\|_{p'}^{p'} + \|h_{\mathcal{M}}^{-1}\sigma_k\|_{p'}^{p'}) \\ \lesssim h_{\max}^{s/(p-1)+p'(k+1)-1}.$$

A triangle inequality and (4.11) show

(4.13)
$$\|\sigma - \mathcal{R}_{h}^{*}\mathbf{I}_{W}\sigma\|_{p'} \le \|(1 - \Pi_{\mathcal{M}}^{k})\sigma\|_{p'} + \|\sigma_{k}\|_{p'} \le h_{\max}^{k+1}$$

The convergence rates of the remaining terms in the error estimate (4.3) are well established in the HHO literature [26, 28, 27, 23, 24] and briefly recalled below. The Hölder inequality, (4.10), and $\|D_hI_V u\|_p + \|\mathcal{R}_h^*I_W\sigma\|_{p'} \lesssim \|Du\|_p + \|h_\mathcal{M}D_{pw}\sigma\|_{p'}$ from (4.11) and (4.13) imply

$$(4.14) - \int_{\Omega} (1 - \Pi_{\mathcal{M}}^{k}) \xi : (1 - \Pi_{\mathcal{M}}^{k}) \mathrm{D}u \,\mathrm{d}x + \int_{\Omega} (\mathrm{D}_{h} \mathrm{I}_{V} u - \varrho) : (\sigma - \mathcal{R}_{h}^{*} \mathrm{I}_{W} \sigma) \,\mathrm{d}x$$
$$\lesssim \| (1 - \Pi_{\mathcal{M}}^{k}) \mathrm{D}u \|_{p} + \| \sigma - \mathcal{R}_{h}^{*} \mathrm{I}_{W} \sigma \|_{p'} \lesssim h_{\max}^{k+1}.$$

The convergence rates

(4.15)
$$s_h(I_V u) \lesssim h_{\max}^{-s+p-1} \|D_{pw}(1-\Pi_{\mathcal{M}}^{k+1})u\|_p^p \lesssim h_{\max}^{-s+(k+2)p-1}$$

follow from the stability of the L^2 projection, the trace, and the Poincaré inequality. These arguments, $[u]_S = 0$ along $S \in \Sigma \setminus \Sigma_N$, and the Hölder inequality prove

(4.16)
$$-\sum_{S\in\Sigma\setminus\Sigma_{N}}\int_{S} [\mathcal{R}_{h}\mathbf{I}_{V}u]_{S} \cdot (1-\Pi_{S}^{k})\sigma\nu_{S} \,\mathrm{d}s$$
$$\lesssim h_{\max}\|\mathbf{D}_{pw}(u-\mathcal{R}_{h}\mathbf{I}_{V}u)\|_{p}\|\mathbf{D}_{pw}(1-\Pi_{\mathcal{M}}^{k})\sigma\|_{p'} \lesssim h_{\max}^{2(k+1)}.$$

The regularity assumption on σ ensures $f = -\operatorname{div} \sigma \in W^{k,p'}(\mathcal{M}; \mathbb{M})$. This and a piecewise application of the Poincaré inequality show

(4.17)
$$\int_{\Omega} (f - f_h) \cdot (u - \mathcal{R}_h \mathbf{I}_V u) \, \mathrm{d}x \lesssim \operatorname{osc}(f, \mathcal{M}) \| \mathbf{D}_{\mathrm{pw}}(u - \mathcal{R}_h \mathbf{I}_V u) \|_p \lesssim h_{\max}^{2(k+1)}.$$

Since $\min\{-s + (k+2)p - 1, s/(p-1) + p'(k+1) - 1\} \ge k+1$ for $p-1 - (k+1) \le s \le (k+2)(p-1)$, the combination of (4.13)–(4.17) concludes the proof. \Box

If E_h is coercive, then the error estimate (4.3) also bounds the error arising from the coercivity of E_h . Suppose that $\Psi \in C^1(\mathbb{M})$ and $\psi(x, \bullet) \in C^1(\mathbb{R}^m)$ for a.e. $x \in \Omega$. Let $\Phi : L^p(\Omega; \mathbb{M}) \times L^p(\Omega; \mathbb{M}) \to \mathbb{R}_{\geq 0}$ and $\varphi : L^p(\Omega; \mathbb{R}^m) \times L^p(\Omega; \mathbb{R}^m) \to \mathbb{R}_{\geq 0}$ be given such that any $\alpha, \beta \in L^p(\Omega; \mathbb{M})$ and $a, b \in L^p(\Omega; \mathbb{R}^m)$ satisfy

(4.18)
$$\Phi(\alpha,\beta) \le \int_{\Omega} (\Psi(\alpha) - \Psi(\beta) - D\Psi(\beta) : (\alpha - \beta)) \, \mathrm{d}x,$$

(4.19)
$$\varphi(a,b) \le \int_{\Omega} (\psi_h(x,a) - \psi_h(x,b) - \nabla_u \psi_h(x,b) \cdot (a-b)) \, \mathrm{d}x$$

The discrete Euler-Lagrange equations associated with the minimization of (1.4) read, for any $v_h = (v_{\mathcal{M}}, v_{\Sigma}) \in V_{\mathrm{D}}(\mathcal{M})$,

(4.20)
$$\int_{\Omega} (\sigma_{\mathcal{M}} : \mathbf{D}_h v_h + \mathbf{D}_u \psi_h(x, \Pi^k_{\mathcal{M}} u_{\mathcal{M}}) \cdot \Pi^k_{\mathcal{M}} v_{\mathcal{M}}) \, \mathrm{d}x + \mathbf{s}_h(u_h; v_h) = 0.$$

The choice $\alpha := D_h v_h$ and $\beta := D_h u_h$ in (4.18) provides

$$\Phi(\mathbf{D}_h u_h, \mathbf{D}_h v_h) \le \int_{\Omega} (\Psi(\mathbf{D}_h v_h) - \Psi(\mathbf{D}_h u_h) - \sigma_{\mathcal{M}} : \mathbf{D}_h (v_h - u_h)) \, \mathrm{d}x$$

with the abbreviation $\sigma_{\mathcal{M}} \coloneqq \Pi^k_{\mathcal{M}} \mathrm{D}\Psi(\mathrm{D}_h u_h)$. This and (4.20) imply

(4.21)

$$\Phi(\mathbf{D}_h u_h, \mathbf{D}_h v_h) \le E_h(v_h) - E_h(u_h) - \mathbf{s}_h(v_h)/r + \mathbf{s}_h(u_h)/r + \mathbf{s}_h(u_h; v_h - u_h) - \int_{\Omega} (\psi_h(x, \Pi^k_{\mathcal{M}} v_{\mathcal{M}}) - \psi_h(x, \Pi^k_{\mathcal{M}} u_{\mathcal{M}}) - \nabla_u \psi_h(x, \Pi^k_{\mathcal{M}} u_{\mathcal{M}}) \cdot \Pi^k_{\mathcal{M}}(v_{\mathcal{M}} - u_{\mathcal{M}})) \, \mathrm{d}x.$$

The convexity of the continuously differentiable function $x \in \mathbb{R}^m \mapsto |x|^r$ leads to

(4.22)
$$0 \le s_h(v_h)/r - s_h(u_h)/r - s_h(u_h; v_h - u_h).$$

At this point, we note that s_h is strongly convex [19, Lemma 5.4(e)]. This allows for additional error control in the stabilization, which is neglected because it is not required in the analysis of this paper. The combination of (4.21)–(4.22) with (4.19) results in

(4.23)
$$e_h(u_h, v_h) \le E_h(v_h) - E_h(u_h) \quad \text{for any } v_h \in V_D(\mathcal{M})$$

with $e_h(u_h, v_h) \coloneqq \Phi(\mathbf{D}_h u_h, \mathbf{D}_h v_h) + \varphi(\Pi_{\mathcal{M}}^k u_{\mathcal{M}}, \Pi_{\mathcal{M}}^k v_{\mathcal{M}})$. If (4.1) holds, then $\varphi \equiv 0$. In this case, the choice $v_h \coloneqq \mathbf{I}_V u$ in (4.23) and Corollary 4.3 lead to the convergence rates $e_h(u_h, \mathbf{I}_V u) \lesssim h_{\max}^{k+1}$ under the assumptions of Corollary 4.3. It is well known [38, 44, 20] that the convergence rates in Corollary 4.3 can be improved under additional coercivity assumptions on Ψ . In the following, the application of the arguments from the aforementioned references will be briefly outlined.

Example 4.4 (strongly convex with Lipschitz continuous gradient). Suppose that $\Psi \in C^1(\mathbb{M})$ satisfy (4.9) with $p \geq 2$ and there exist positive constants c_6, c_7 with

(4.24)
$$c_{6}|D\Psi(A) - D\Psi(B)|^{2} \leq (1 + |A|^{p-2} + |B|^{p-2}) \times (\Psi(A) - \Psi(B) - D\Psi(B) : (A - B)),$$

(4.25)
$$c_7 |A - B|^p \le \Psi(A) - \Psi(B) - D\Psi(B) : (A - B)$$

for any $A, B \in \mathbb{M}$. Then (4.18) holds with $\Phi(\alpha, \beta) \coloneqq \frac{c_8}{2} \| D\Psi(\alpha) - D\Psi(\beta) \|_{p'}^2 / (1 + \|\alpha\|_p^p + \|\beta\|_p^p)^{(2-p')/p'} + \frac{c_7}{2} \|\alpha - \beta\|_p^p$ and $c_8 \coloneqq c_6/3$ [16, 19]. Recall ρ and $\xi = D\Psi(D_h I_V u)$ from Theorem 4.1. Let the assumptions of Corollary 4.3 hold. The combination of (4.18) with $\Psi(\alpha) - \Psi(\beta) \leq -D\Psi(\alpha) : (\beta - \alpha)$ a.e. in Ω from the convexity of Ψ implies the monotonicity

(4.26)
$$\Phi(\alpha,\beta) \le \int_{\Omega} (D\Psi(\alpha) - D\Psi(\beta)) : (\alpha - \beta) \, \mathrm{d}x \quad \text{for any } \alpha, \beta \in L^p(\Omega;\mathbb{M})$$

with two immediate consequences. On the one hand, (4.26) with $\alpha \coloneqq \mathrm{D}u$, $\beta \coloneqq \mathrm{D}_h \mathrm{I}_V u$, $\xi = \mathrm{D}\Psi(\mathrm{D}_h \mathrm{I}_V u)$, a Hölder inequality, and $\|\mathrm{D}u\|_p + \|\mathrm{D}_h \mathrm{I}_V u\|_p \lesssim 1$ lead to $\|\sigma - \xi\|_{p'} \lesssim \|\mathrm{D}u - \mathrm{D}_h \mathrm{I}_V u\|_p$. This, $\|(1 - \Pi^k_{\mathcal{M}})\xi\|_{p'} \lesssim \|\Pi^k_{\mathcal{M}}\sigma - \xi\|_{p'}$ from (2.1), and a triangle inequality provide

(4.27)
$$\begin{aligned} \|(1 - \Pi_{\mathcal{M}}^{k})\xi\|_{p'} &\lesssim \|(1 - \Pi_{\mathcal{M}}^{k})\sigma\|_{p'} + \|\sigma - \xi\|_{p'} \\ &\lesssim \|(1 - \Pi_{\mathcal{M}}^{k})\sigma\|_{p'} + \|\mathrm{D}u - \mathrm{D}_{h}\mathrm{I}_{V}u\|_{p} \lesssim h_{\mathrm{max}}^{k+1}. \end{aligned}$$

On the other hand, (4.26) with $\alpha \coloneqq \mathrm{D}u$, $\beta \coloneqq \varrho$, a Hölder inequality, and $\mathrm{D}\Psi(\varrho) = \mathcal{R}_h^* \mathrm{I}_W \sigma$ from $\varrho \in \partial \Psi^*(\mathcal{R}_h^* \mathrm{I}_W \sigma)$ [43, Theorem 23.5 (a^{*})] imply $\|\mathrm{D}u - \varrho\|_p \lesssim \|\sigma - \mathcal{R}_h^* \mathrm{I}_W \sigma\|_{p'}^{1/(p-1)}$. This and a triangle inequality provide

The bounds (4.27)-(4.28) lead to the convergence rates

(4.29)
$$-\int_{\Omega} (1-\Pi_{\mathcal{M}}^k)\xi : (1-\Pi_{\mathcal{M}}^k)\mathrm{D}u\,\mathrm{d}x \lesssim h_{\max}^{2(k+1)},$$

(4.30)
$$\int_{\Omega} (\mathbf{D}_h \mathbf{I}_V u - \varrho) : (\sigma - \mathcal{R}_h^* \mathbf{I}_W \sigma) \, \mathrm{d}x \lesssim h_{\max}^{p'(k+1)}.$$

This, (4.12), and (4.15)–(4.17) conclude $e_h(u_h, \mathbf{I}_V u) \leq E_h(\mathbf{I}_V u) - E_h(u_h) \lesssim h_{\max}^{p'(k+1)}$ for any $p-1 \leq s \leq (k+1)(p-p') + p-1$. In the case $1 , suppose that the coercivity (4.18) holds for <math>\Phi(\alpha, \beta) \coloneqq c_9 \|\alpha - \beta\|_p^2/(\|\alpha\|_p^p + \|\beta\|_p^p)^{(2-p)/2} + c_{10} \|\mathbf{D}\Psi(\alpha) - \mathbf{D}\Psi(\beta)\|_{p'}^{p'}$. Then, by similar arguments, $\|\sigma - \xi\|_{p'} \lesssim \|\mathbf{D}u - \mathbf{D}_h\mathbf{I}_V u\|_p^{1/(p'-1)}$ and $\|\mathbf{D}u - \varrho\|_p \lesssim \|\sigma - \mathcal{R}_h^*\mathbf{I}_W \sigma\|_{p'}$. Thus, the convergence rates $e_h(u_h, \mathbf{I}_V u) \leq E_h(\mathbf{I}_V u) - E_h(u_h) \lesssim h_{\max}^{p(k+1)}$ hold for any $p(k+1)(p-2) + p - 1 \leq s \leq p - 1$.

Remark 4.5 (p-Laplace). Given $f \in L^{p'}(\Omega; \mathbb{R}^m)$, the p-Laplace problem seeks the unique solution $u \in V_D$ to $-\operatorname{div}\sigma = f$ in Ω with $\sigma \coloneqq |\mathrm{D}u|^{p-2}\mathrm{D}u$. Then u minimizes E in V_D with the energy density $\Psi(A) \coloneqq |A|^p/p$ for any $A \in \mathbb{M}$. Since Ψ and Ψ^* are strictly convex, the minimizer u of E in V_D and the maximizer $\sigma \coloneqq \mathrm{D}\Psi(\mathrm{D}u)$ of E^* in W_N are unique. The coercivity assumptions of Example 4.4 are satisfied and so, the convergence rates therein hold. They coincide with [38] for $p \ge 2$ and [44, 20] for $1 . The consequence <math>\|\mathrm{D}_h(\mathrm{I}_V u - u_h)\|_p \lesssim h^{(k+1)p/2}$ of the latter provides an improvement to existing rates in the HHO literature [18, 25]. The derivation of quadratic convergence rates for $E_h(\mathrm{I}_V u) - E_h(u_h)$ appears possible using the coercivity of Ψ with respect to the so-called quasi-norm [3, 32], cf. [31] for the dG method and [6, 39] for the Crouzeix-Raviart FEM. This approach, however, is currently restricted to the lowest-order case due to the lack of higher convergence rates for interpolation in Orlicz-Sobolev spaces [33].

5. A posteriori

The main goal of this section is the design of a $W^{p'}(\operatorname{div}, \Omega; \mathbb{M})$ conforming postprocessing from a computed discrete minimizer $u_h = (u_{\mathcal{M}}, u_{\Sigma})$ of (1.4) for the derivation of primal-dual error estimates. Recall the abbreviation $\sigma_{\mathcal{M}} = \prod_{\mathcal{M}}^k D\Psi(D_h u_h)$ for $\Psi \in C^1(\mathbb{M})$. In contrast to the unstabilized HHO method on regular triangulations into simplices from [18], $\sigma_{\mathcal{M}} \notin W_N$ has to be expected. In fact, the discrete Euler-Lagrange equations (4.20) imply the following result also known as equilibrium of traces [10, 1].

Corollary 5.1 (normal jump of $\sigma_{\mathcal{M}}$). Suppose that $\Psi \in C^1(\mathbb{M})$ and $\psi_h(x, \bullet) \in C^1(\mathbb{R}^m)$ for a.e. $x \in \Omega$. For any $S \in \Sigma \setminus \Sigma_D$, $\sigma_{\mathcal{M}} = \prod_{\mathcal{M}}^k D\Psi(D_h u_h) \in P_k(\mathcal{M}; \mathbb{M})$ satisfies, for $T_{K,S}$ from (2.12), that

(5.1)
$$[\sigma_{\mathcal{M}}\nu_S]_S = -\sum_{K \in \mathcal{M}, S \in \Sigma(K)} h_S^{-s} T_{K,S} u_h.$$

r

Proof. The discrete Euler-Lagrange equations (4.20) and the definition (2.2) of D_h imply, for any $v_h = (v_M, v_\Sigma) \in V_D(\mathcal{M})$, that

(5.2)
$$\int_{\Omega} \Pi^{k}_{\mathcal{M}} v_{\mathcal{M}} \cdot (\nabla_{u} \psi_{h}(x, \Pi^{k}_{\mathcal{M}} u_{\mathcal{M}}) - \operatorname{div}_{\mathrm{pw}} \sigma_{\mathcal{M}}) \, \mathrm{d}x + \sum_{S \in \Sigma \setminus \Sigma_{\mathrm{D}}} \int_{S} v_{S} \cdot [\sigma_{\mathcal{M}} \nu_{S}]_{S} \, \mathrm{d}s + \mathrm{s}(u_{h}; v_{h}) = 0.$$

Fix a face $S \in \Sigma \setminus \Sigma_{\mathrm{D}}$. The choice $v_{\mathcal{M}} \equiv 0$ and $v_{\Sigma} \in P_k(\Sigma \setminus \Sigma_{\mathrm{D}}; \mathbb{R}^m)$ with $v_{\Sigma}|_E \equiv 0$ for all $E \in \Sigma$, $E \neq S$ in (5.2) provides

$$\int_{S} v_{S} \cdot \left([\sigma_{\mathcal{M}} \nu_{S}]_{S} + \sum_{K \in \mathcal{M}, S \in \Sigma(K)} h_{S}^{-s} T_{K,S} u_{h} \right) \mathrm{d}s = 0$$

for all $v_S \in P_k(S; \mathbb{R}^m)$. This orthogonality concludes (5.1).

Although $\sigma_{\mathcal{M}} \notin W_{N}$ holds in general, it can be utilized to construct an appropriate postprocessing $\sigma_0 \in \operatorname{RT}_k(\mathcal{T}; \mathbb{M}) \cap W_N$ in a matching simplicial submesh \mathcal{T} of \mathcal{M} as follows. (We refer to [26, Definition 1.37] for a precise definition of a matching submesh.)

Lemma 5.2 (Raviart-Thomas reconstruction). Suppose that $\Psi \in C^1(\mathbb{M})$ and $\psi_h(x, \bullet) \in C^1(\mathbb{R}^m)$ for a.e. $x \in \Omega$. Then the discrete dual variable $\sigma_h = (\sigma_{\mathcal{M}}, \sigma_{\Sigma}) \in$ $W_{\rm N}(\mathcal{M})$ with $\sigma_{\mathcal{M}} = \Pi^k_{\mathcal{M}} \mathrm{D}\Psi(\mathrm{D}_h u_h) \in P_k(\mathcal{M}; \mathbb{M})$ and

$$\sigma_S \coloneqq \begin{cases} \{\sigma_{\mathcal{M}}\nu_S\}_S + h_S^{-s}(T_{K_+,S}u_h - T_{K_-,S}u_h)/2 \in P_k(S;\mathbb{R}^m) & \text{if } S \in \Sigma(\Omega), \\ \sigma_{\mathcal{M}}\nu_S + h_S^{-s}T_{K_+,S}u_h \in P_k(S;\mathbb{R}^m) & \text{if } S \in \Sigma_D \end{cases}$$

satisfies $\operatorname{div}_h \sigma_h = \prod_{\mathcal{M}}^k \nabla_u \psi_h(x, \prod_{\mathcal{M}}^k u_{\mathcal{M}}) \in P_k(\mathcal{M}; \mathbb{R}^m)$. Given a matching simplicial submesh \mathcal{T} of \mathcal{M} with the set \mathcal{F} of faces, let $\sigma_0 \in \mathrm{RT}_k(\mathcal{T}; \mathbb{M}) \cap W_{\mathrm{N}}$ be the unique Raviart-Thomas finite element function defined by the weights

(5.3)
$$\Pi_{\mathcal{T}}^{k-1}\sigma_{0} = \Pi_{\mathcal{T}}^{k-1}\sigma_{\mathcal{M}},$$
$$\sigma_{0}\nu_{F} = \begin{cases} \sigma_{S} & \text{if } F \in \mathcal{F}, F \subset S \text{ for some } S \in \Sigma, \\ \sigma_{K}\nu_{F} & \text{if } F \in \mathcal{F}, F \subset K \text{ for some } K \in \mathcal{M} \text{ but } F \notin \partial K. \end{cases}$$

Then $\Pi^k_{\mathcal{M}} \operatorname{div} \sigma_0 = \Pi^k_{\mathcal{M}} \nabla_u \psi_h(x, \Pi^k_{\mathcal{M}} u_{\mathcal{M}}).$

1 1

The weights in (5.3) are the degrees of freedom for Raviart-Thomas finite element functions [34, Section 14.3] and therefore, σ_0 is uniquely defined by (5.3).

Proof. Given any $v_h = (v_{\mathcal{M}}, v_{\Sigma}) \in V_{\mathcal{D}}(\mathcal{M})$ with $v_{\mathcal{M}} \in P_k(\mathcal{M}; \mathbb{R}^m)$, a piecewise integration by parts followed by the definition (2.3) of div_h leads to

$$-\int_{\Omega} \operatorname{div}_{\mathrm{pw}} \sigma_{\mathcal{M}} \cdot v_{\mathcal{M}} \, \mathrm{d}x = -\int_{\Omega} \operatorname{div}_{h} \sigma_{h} \cdot v_{\mathcal{M}} \, \mathrm{d}x - \sum_{S \in \Sigma \setminus \Sigma_{\mathrm{D}}} \{v_{\mathcal{M}}\}_{S} \cdot [\sigma_{\mathcal{M}} \nu_{S}]_{S} \, \mathrm{d}s + \sum_{S \in \Sigma \setminus \Sigma_{\mathrm{N}}} [v_{\mathcal{M}}]_{S} \cdot (\sigma_{S} - \{\sigma_{\mathcal{M}} \nu_{S}\}_{S}) \, \mathrm{d}s.$$

This and the definition (2.2) of D_h imply

$$\int_{\Omega} \sigma_{\mathcal{M}} : \mathcal{D}_{h} v_{h} \, \mathrm{d}x = -\int_{\Omega} \operatorname{div}_{h} \sigma_{h} \cdot v_{\mathcal{M}} \, \mathrm{d}x + \sum_{S \in \Sigma \setminus \Sigma_{\mathrm{D}}} (v_{S} - \{v_{\mathcal{M}}\}_{S}) \cdot [\sigma_{\mathcal{M}} \nu_{S}]_{S} \, \mathrm{d}s + \sum_{S \in \Sigma \setminus \Sigma_{\mathrm{N}}} [v_{\mathcal{M}}]_{S} \cdot (\sigma_{S} - \{\sigma_{\mathcal{M}} \nu_{S}\}_{S}) \, \mathrm{d}s.$$

The combination of this with the discrete Euler-Lagrange equations (4.20) yields

$$\int_{\Omega} v_{\mathcal{M}} \cdot (\nabla_u \psi_h(x, \Pi_{\mathcal{M}}^k u_{\mathcal{M}}) - \operatorname{div}_h \sigma_h) \, \mathrm{d}x + \sum_{S \in \Sigma \setminus \Sigma_D} \int_{S} (v_S - \{v_{\mathcal{M}}\}_S) \cdot [\sigma_{\mathcal{M}} \nu_S]_S \, \mathrm{d}s$$
(5.4)
$$+ \sum_{S \in \Sigma \setminus \Sigma_N} \int_{S} [v_{\mathcal{M}}]_S \cdot (\sigma_S - \{\sigma_{\mathcal{M}} \nu_S\}_S) \, \mathrm{d}s + \mathrm{s}(u_h; v_h) = 0.$$

Let v_{Σ} be chosen such that $v_{S} \coloneqq \prod_{S}^{k} \{v_{\mathcal{M}}\}_{S} = \{v_{\mathcal{M}}\}_{S}$ along $S \in \Sigma \setminus \Sigma_{\mathrm{D}}$ in (5.4). For any $K \in \mathcal{M}$ and $S \in \Sigma(K)$, $v_{S} - v_{K}$ equals $-[v_{\mathcal{M}}]_{S}$ if $S \in \Sigma_{\mathrm{D}}$, 0 if $S \in \Sigma_{\mathrm{N}}$, and $-\frac{v_{S} \cdot v_{K}|_{S}}{2} [v_{\mathcal{M}}]_{S}$ if $S \in \Sigma(\Omega)$. Therefore,

$$s(u_h; v_h) = -\sum_{S \in \Sigma(\Omega)} \frac{h_S^{-s}}{2} \int_S (T_{K_+, S} u_h - T_{K_-, S} u_h) \cdot \Pi_S^k [v_\mathcal{M}]_S \, \mathrm{d}s$$
$$-\sum_{S \in \Sigma_D} h_S^{-s} \int_S T_{K_+, S} u_h \cdot \Pi_S^k v_\mathcal{M} \, \mathrm{d}s.$$

This, the definition of σ_{Σ} in Lemma 5.2, and (5.4) conclude

$$\int_{\Omega} v_{\mathcal{M}} \cdot (\nabla_u \psi_h(x, \Pi_{\mathcal{M}}^k u_{\mathcal{M}}) - \operatorname{div}_h \sigma_h) \, \mathrm{d}x = 0 \quad \text{for any } v_{\mathcal{M}} \in P_k(\mathcal{M}; \mathbb{R}^m),$$

whence $\operatorname{div}_h \sigma_h = \prod_{\mathcal{M}}^k \nabla_u \psi_h(x, \prod_{\mathcal{M}}^k u_{\mathcal{M}})$. Given any $v_k \in P_k(\mathcal{M}; \mathbb{R}^m)$, a piecewise integration by parts and $D_{pw} v_k \in P_{k-1}(\mathcal{M}; \mathbb{M})$ show

$$\int_{\Omega} \operatorname{div} \sigma_0 \cdot v_k \, \mathrm{d}x = -\int_{\Omega} \sigma_{\mathcal{M}} : \mathcal{D}_{\mathrm{pw}} v_k + \sum_{F \in \mathcal{F}, F \subset \partial \mathcal{M}} \int_F [v_k]_F \cdot \sigma_0 \nu_F \, \mathrm{d}s$$

because the function v_k only jumps on the skeleton $\partial \mathcal{M} = \bigcup_{K \in \mathcal{M}} \partial K$ of \mathcal{M} . This, the definition of σ_0 from (5.3), and of div_h from (2.3) imply

$$\int_{\Omega} \operatorname{div} \sigma_0 \cdot v_k \, \mathrm{d}x = -\int_{\Omega} \sigma_{\mathcal{M}} : \mathcal{D}_{\mathrm{pw}} v_k + \sum_{S \in \Sigma \setminus \Sigma_{\mathrm{N}}} \int_{S} [v_k]_S \cdot \sigma_S \, \mathrm{d}s = \int_{\Omega} \operatorname{div}_h \sigma_h \cdot v_k \, \mathrm{d}x.$$

The assertion then follows from $\operatorname{div}_h \sigma_h = \prod_{\mathcal{M}}^k \nabla_u \psi_h(x, \prod_{\mathcal{M}}^k u_{\mathcal{M}}).$

In order to derive a posteriori error estimate, we consider throughout the remaining parts of this paper the explicit representation (4.1) of the lower-order terms. In particular, $\varphi \equiv 0$ in (4.19). The following arguments are well known in the literature [42, 4, 14, 18, 8] and sketched below for later reference. Let \tilde{u} minimize

$$E_{\sigma_0}(v) \coloneqq \int_{\Omega} (\Psi(\mathrm{D}v) + \mathrm{div}\sigma_0 \cdot v) \,\mathrm{d}x \quad \text{among } v \in V_{\mathrm{D}}.$$

The corresponding dual energy reads

$$E_{\sigma_0}^*(\tau) \coloneqq -\int_{\Omega} \Psi^*(\tau) \,\mathrm{d}x - \chi_{\mathrm{div}\sigma_0}(\mathrm{div}\tau) \quad \text{for any } \tau \in W_{\mathrm{N}}.$$

Abbreviate $\tilde{f} \coloneqq f + \operatorname{div} \sigma_0$. The L^2 orthogonality $\tilde{f} \perp P_k(\mathcal{M}; \mathbb{R}^m)$ from Lemma 5.2, a Hölder inequality, and a piecewise application of the Poincaré inequality imply

$$E(u) - E_{\sigma_0}(u) = -\int_{\Omega} \widetilde{f} \cdot u \, \mathrm{d}x \le c_{\mathrm{P}} c_5 \mathrm{osc}(\widetilde{f}, \mathcal{M})$$

Given a conforming postprocessing $v_0 \in V_D$, this and $E^*_{\sigma_0}(\sigma_0) \leq E_{\sigma_0}(\widetilde{u})$ prove

(5.5)
$$\text{LEB} \coloneqq E^*_{\sigma_0}(\sigma_0) - c_{\mathrm{P}}c_5 \mathrm{osc}(\tilde{f}, \mathcal{M}) \le E(u) \le E(v_0).$$

In practice, $v_0 \in S_{\mathrm{D}}^{k+1}(\mathcal{T};\mathbb{R}^m) \coloneqq P_{k+1}(\mathcal{T};\mathbb{R}^m) \cap V_{\mathrm{D}}$ is the nodal average of $u_{\mathcal{M}}$ in a matching simplicial submesh \mathcal{T} of \mathcal{M} . If Ψ satisfies (4.18), then the error arising from the coercivity of Ψ can be bounded as follows. The arguments, that lead to (4.23), apply to the continuous level and prove $e(u, v_0) \leq E(v_0) - E(u)$ with $e(u, v_0) \coloneqq \Phi(\mathrm{D}u, \mathrm{D}v_0)$. This and (5.5) imply the a posteriori error control

(5.6)
$$e(u, v_0) \le E(v_0) - E(u) \le E(v_0) - \text{LEB}.$$

Here, we follow [8] and consider the conforming postprocessing v_0 as the approximation for u. Therefore, the quantity of interest is $e(u, v_0)$ instead of $\Phi(Du, D_h u_h)$, which avoids additional consistency terms. In fact, the duality gap (5.6) is, among other terms, part of the a posteriori error estimators in [15, 18]. Thus, this approach leads to a more accurate error control. In practice, a posteriori error estimates for $\Phi(Du, D_h u_h)$ can be obtained from (5.6) and a triangle inequality. If \mathcal{M} is a regular triangulation into simplices, then $\operatorname{osc}(\tilde{f}, \mathcal{M}) = \operatorname{osc}(f, \mathcal{M})$. Notice that the data oscillation $\operatorname{osc}(\tilde{f}, \mathcal{M})$ in (5.6) may dominate the error for higher-order methods. Since pointwise control over the divergence of the postprocessing σ_0 from Lemma 5.2 is lost on polyhedral meshes, it will not vanish even if $f \in P_k(\mathcal{M}; \mathbb{R}^m)$. Therefore, the usefulness of this reconstruction is restricted in the latter case unless $\operatorname{osc}(\tilde{f}, \mathcal{M})$ scales quadratically. This is guaranteed for uniformly convex W or in the case of the *p*-Laplace problem.

Remark 5.3 (data oscillation). If Ψ satisfies (4.25), then (5.6) proves

$$e(u, v_0) \le E(v_0) - E(u) \le E_{\sigma_0}(v_0) - E_{\sigma_0}(u) + \int_{\Omega} \widetilde{f} \cdot (u - v_0) \, \mathrm{d}x$$

This, a Hölder, a Poincaré, and a Young inequality imply

$$e(u, v_0)/p' \le E_{\sigma_0}(v_0) - E_{\sigma_0}(u) + c_{11} \operatorname{osc}(\widetilde{f}, \mathcal{M})^{p'}$$

with the constant $c_{11} \coloneqq c_{\mathrm{P}}^{p'} c_7^{-1/(p-1)}/p'$. For the *p*-Laplace equation from Remark 4.5, the energy density W satisfies (5.6) with $\Phi(\alpha,\beta) \coloneqq c_{12} \|(|\alpha|+|\beta|)^{(p-2)/2} \times (\alpha-\beta)\|_2^2$ for some positive constant $c_{12} > 0$ [44, 20, 3]. This is known under the label *quasi-norm* with different equivalent versions in [13, 32]. The properties of the N-shift functions [32] lead to, for any $\delta > 0$,

$$\int_{\Omega} \widetilde{f} \cdot (u - v_0) \, \mathrm{d}x \lesssim c_{\delta} \mathrm{osc}_p(\widetilde{f}, v_0, \mathcal{M})^2 + \delta \Phi(\mathrm{D}u, \mathrm{D}v).$$

with $\operatorname{osc}_p(\widetilde{f}, v_0, \mathcal{M})^2 \coloneqq \sum_{K \in \mathcal{M}} h_K^2 \int_K (|\nabla v_0|^{p-1} + h_K|\widetilde{f}|)^{p'-2} |\widetilde{f}|^2 \, dx$ and a constant c_{δ} depending on δ , cf. [39, Ineq. (3.29)]. By choosing δ sufficiently small, we obtain the a posteriori error estimate

$$e(u, v_0) \lesssim E_{\sigma_0}(v_0) - E^*_{\sigma_0}(\sigma_0) + \operatorname{osc}_p(f, v_0, \mathcal{M})^2.$$

6. Numerical examples

This section tests the performance of the a posteriori error control from Section 5 in three numerical benchmarks on the two dimensional L-shaped domain $\Omega := (-1,1)^2 \setminus ([0,1) \times (-1,0])$ with pure Dirichlet boundary $\Gamma_{\rm D} = \partial \Omega$. The initial triangulation in all benchmarks is displayed in Figure 1(a). The computer experiments are carried out on regular triangulations into *simplices*.

6.1. Adaptive mesh-refining algorithm. Up to data oscillation, the a posteriori error estimator (5.6) consists of the duality gap $E_{\sigma_0}(v_0) - E^*_{\sigma_0}(\sigma_0)$. The localization of this has been discussed in [8] and we only restate some relevant details to the benchmarks below. An integration by parts with div $\sigma_0 = -f_h$ implies

$$E_{\sigma_0}(v_0) - E^*_{\sigma_0}(\sigma_0) = \int_{\Omega} (\Psi(\nabla v_0) - \sigma_0 \cdot \nabla v_0 + \Psi^*(\sigma_0)) \, \mathrm{d}x \rightleftharpoons \sum_{K \in \mathcal{M}} \eta(K)$$

with the local refinement indicator

(6.1)
$$\eta(K) \coloneqq \int_{K} (\Psi(\nabla v_0) - \sigma_0 \cdot \nabla v_0 + \Psi^*(\sigma_0)) \, \mathrm{d}x.$$

Notice from the Fenchel-Young inequality that $\Psi(\nabla v_0) - \sigma_0 \cdot \nabla v_0 + \Psi^*(\sigma_0) \ge 0$ holds pointwise a.e. in Ω and so, $\eta(K) \ge 0$. The data oscillation is ignored unless it scales quadratically. In this case, its local contribution will be added to (6.1). Adaptive computations utilize the refinement indicator (6.1) in the standard adaptive meshrefining loop [12] with the Dörfler marking strategy, i.e., at each refinement step, a subset $\mathfrak{M} \subset \mathcal{M}$ with minimal cardinality is selected such that

$$\sum_{K \in \mathcal{M}} \eta(K) \le \frac{1}{2} \sum_{K \in \mathfrak{M}} \eta(K).$$

The convergence history plots display the quantities of interest against the number of degrees of freedom ndof in a log-log plot. (Recall the scaling ndof $\approx h_{\rm max}^{-2}$ for uniform meshes.) Solid lines indicate adaptive, while dashed lines are associated with uniform mesh refinements.

The discrete minimization problem (1.4) is solved by an iterative solver fminunc from the MATLAB standard library in an extension of the data structures and the short MATLAB programs [2]. The first and (piecewise) second derivatives of Whave been provided for the trust-region quasi-Newton scheme with parameters of fminunc set to FunctionTolerance = OptimalityTolerance = StepTolerance = 10^{-15} and MaxIterations = 10^3 for improved accuracy. The numerical integration of piecewise polynomials is carried out exactly. For non-polynomial functions such as $W(D_h v_h)$ with $v_h \in V_D(\mathcal{M})$, the number of chosen quadrature points allows for exact integration of polynomials of degree at most 2pk + 1 with the growth p of Wand the polynomial order k of the discretization.




FIGURE 1. (a) Initial triangulation of the L-shaped domain into 6 triangles and (b) material distribution in the optimal design problem of Subsection 6.2




FIGURE 2. (a) Convergence history plot of $E_{\sigma_0}(v_0) - E^*_{\sigma_0}(\sigma_0)$ for various k and (b) adaptive triangulation into 2013 triangles obtained with k = 2 in Subsection 6.2

6.2. **Optimal design problem.** The optimal design problem seeks the optimal distribution of two materials with fixed amounts to fill a given domain for maximal torsion stiffness [40, 7]. Given parameters $0 < t_1 < t_2$ and $0 < \mu_1 < \mu_2$ with $t_1\mu_2 = \mu_1t_2$, the energy density $\Psi(a) \coloneqq w(|a|), a \in \mathbb{R}^n$, with

$$w(t) \coloneqq \begin{cases} \mu_2 t^2 / 2 & \text{if } 0 \le t \le t_1, \\ t_1 \mu_2 (t - t_1 / 2) & \text{if } t_1 \le t \le t_2, \\ \mu_1 t^2 / 2 + t_1 \mu_2 (t_2 / 2 - t_1 / 2) & \text{if } t_2 \le t \end{cases}$$

satisfies (4.9) with p = 2 and (4.24) with $c_6 = 1/(2\mu_2)$ [7]. This benchmark considers the parameters $\mu_1 = 1$, $\mu_2 = 2$, $t_1 = \sqrt{2\lambda\mu_1/\mu_2}$ for $\lambda = 0.0084$, $t_2 = \mu_2 t_1/\mu_1$ from [7] and the input r = 2, s = 1 for the stabilization s_h . Since the data oscillation vanishes, the a posteriori error estimate from (5.6) reads

$$e(u, v_0) \coloneqq \frac{1}{2\mu} \|\nabla \Psi(\nabla u) - \nabla \Psi(\nabla v_0)\|_2^2 \le E_{\sigma_0}(v_0) - E_{\sigma_0}^*(\sigma_0) = E(v_0) - E^*(\sigma_0).$$

The approximated material distribution obtained by adaptive computation with k = 0 is displayed in Figure 1(b) using the volume fraction plot from [7, Section 5]. We observe two homogenous phases and a transition layer with a fine mixture of the two materials. On uniform meshes, the convergence rate 2/3 for $E_{\sigma_0}(v_0) - E^*_{\sigma_0}(\sigma_0)$ is observed in Figure 2(a). The adaptive mesh-refining algorithm refines towards the singularity at the origin and the transition layer as displayed Figure 2(b). This leads to slightly improved convergence rates for $E_{\sigma_0}(v_0) - E^*_{\sigma_0}(\sigma_0)$.




FIGURE 3. Convergence history plot of $E_{\sigma_0^{\varepsilon}}(v_0^{\varepsilon}) - E_{\sigma_0^{\varepsilon}}^*(\sigma_0^{\varepsilon})$ for (a) various k and $\varepsilon = 10^{-4}$ and (b) k = 2 and various ε in Subsection 6.3

6.3. Bingham flow in a pipe. Given fixed positive parameters $\mu, g > 0$, the modelling of a uni-directional flow through a pipe with cross-section $\Omega \subset \mathbb{R}^2$ leads to the minimization problem (1.1) with the energy density

$$\Psi(a) \coloneqq \mu |a|^2 / 2 + g|a|$$
 for any $a \in \mathbb{R}^2$,

cf. [17] for further details. Given $\alpha \in \mathbb{R}^2 \setminus \{0\}$, the radial symmetry of Ψ leads to

$$\Psi^*(\alpha) = \sup_{a \in \mathbb{R}^2} (\alpha \cdot a - \Psi(a)) = \sup_{t \ge 0} (t|\alpha|^2 - \mu t^2 |\alpha|^2 / 2 - gt|\alpha|) \eqqcolon \varphi_\alpha(t).$$

If $|\alpha| \leq g$, then $\varphi_{\alpha}(t) \leq 0$ and so, $\Psi^*(\alpha) = \sup_{t \geq 0} \varphi_{\alpha}(t) = \varphi_{\alpha}(0) = 0$. If $|\alpha| > g$, then the maximum of $\varphi_{\alpha}(t)$ in $[0, \infty)$ is attained at $t = (|\alpha| - g)/(\mu|\alpha|)$. Hence,

$$\Psi^*(\alpha) = \begin{cases} 0 & \text{if } |\alpha| \le g\\ (|\alpha| - g)^2/(2\mu) & \text{if } |\alpha| > g. \end{cases}$$

The strict convexity of Ψ leads to a unique the minimizer u of E in $V_{\rm D}$. Although Ψ is not differentiable, there exists $\sigma \in H(\operatorname{div}, \Omega) = W^2(\operatorname{div}, \Omega)$ such that $\sigma \in \partial \Psi(\nabla u)$ and $\operatorname{div} \sigma = -f$ pointwise a.e. in Ω [37, Chapter II, Theorem 6.3]. Thus, there is no duality gap $E(u) = E^*(\sigma)$. Furthermore, the minimizer u and any $v \in V_{\rm D}$ satisfy $e(u, v) \leq E(v) - E(u)$ for any $v \in V_{\rm D}$ with $e(u, v) := \frac{\mu}{2} \|\nabla(u - v)\|_2^2$ [17, Lemma 1]. Thus, Remark 5.3 applies and leads, for any $\tau \in W_{\rm N}$, to

(6.2)
$$e(u,v)/2 \le E_{\tau}(v) - E_{\tau}^{*}(\tau) + c_{11} \operatorname{osc}(\tilde{f},\mathcal{M})^{2}$$

The postprocessings for (6.2) are obtained from a regularized discrete problem as in [17]. Given $\varepsilon > 0$, define $\Psi_{\varepsilon} \in C^1(\mathbb{R}^2)$ by

$$\Psi_{\varepsilon}(a)\coloneqq \mu |a|^2/2 + g(\sqrt{|a|^2+\varepsilon^2}) \quad \text{for any } a\in \mathbb{R}^2.$$

The unique minimizer $u_h^{\varepsilon} = (u_{\mathcal{M}}^{\varepsilon}, u_{\Sigma}^{\varepsilon})$ of the discrete energy

$$E_h^{\varepsilon}(v_h) \coloneqq \int_{\Omega} (\Psi_{\varepsilon}(\mathbf{D}_h v_h) - f_h v_{\mathcal{M}}) \, \mathrm{d}x + \mathbf{s}_h(v_h)/2 \quad \text{among } v_h = (v_{\mathcal{M}}, v_{\Sigma}) \in V_{\mathrm{D}}(\mathcal{M})$$

allows for the postprocessings $\sigma_0^{\varepsilon} \in W_N$ with $\operatorname{div} \sigma_0^{\varepsilon} = -f_h$ from Lemma 5.2 and $v_0^{\varepsilon} \in S_D^{k+1}(\mathcal{M})$ as the nodal average of $u_{\mathcal{M}}^{\varepsilon}$. From (6.2), we deduce

$$e(u, v_0^{\varepsilon})/2 \leq E_{\sigma_0^{\varepsilon}}(v_0^{\varepsilon}) - E_{\sigma_0^{\varepsilon}}^*(\sigma_0^{\varepsilon}) + c_{11} \operatorname{osc}(f, \mathcal{M})^2.$$

In the following computer experiment, we set $\mu = 1$, g = 0.2, $f \equiv 10$ with the reference minimal energy E(u) = -9.32049 obtained from the computable bounds (5.5) in adaptive computations and r = 2, s = 1 for the stabilization s_h . On uniform meshes, the convergence rate 2/3 for $E(v_0^{\varepsilon}) - E^*(\sigma_0^{\varepsilon})$ is observed in Figure 3(a) for all polynomial degrees and $\varepsilon = 10^{-4}$. The adaptive algorithm refines towards the expected singularity at the origin and leads to the optimal convergence rates k + 1 for all displayed polynomial degree k until stagnation due to the regularization. The effect of the latter is displayed in Figure 3(b) for the polynomial degree k = 2 and parameters $\varepsilon = 10^0, 10^{-1}, \ldots, 10^{-4}$.




FIGURE 4. Convergence history plot of (a) RHS and (b) $\|\sqrt{\rho}\nabla(u-v_0)\|_2^2$ for various k in Subsection 6.4

6.4. *p*-Laplace problem. In this final benchmark, we consider the 4-Laplace problem from Remark 4.5 with the exact solution

$$u(r,\varphi) = r^{7/8} \sin(7\varphi/8) \in W^{1,4}(\Omega)$$

and the right-hand side

$$f(r,\varphi) = (7/8)^{3/4} r^{-11/8} \sin(7\varphi/8) \in L^{16/11-\varepsilon}(\Omega)$$




FIGURE 5. Convergence history plot of (a) $\|\sigma - \nabla \Psi(\nabla v_0)\|_{4/3}^2$ and (b) $\|\nabla (u - v_0)\|_4^2$ for various k in Subsection 6.4

for any $\varepsilon > 0$ from [13]. Some remarks are in order due to the inhomogenous Dirichlet data prescribed by u. The discrete problem minimizes (1.4) in the affine space $I_V u + V_D(\mathcal{M})$. Since the construction of σ_0 in Lemma 5.2 only relies on the discrete Euler-Lagrange equations, it applies verbatim. To obtain a postprocessing $v_0 \in u + V_D$, let $\tilde{v}_0 \in S_D^{k+1}(\mathcal{M})$ denote the nodal average of $u_{\mathcal{M}} - \Pi_{\mathcal{M}}^{k+1}u$. We set $v_0 \coloneqq \tilde{v}_0 + u$. (Notice that, in the computation of v_0 , u can be replaced by any extension of the given Dirichlet data $u|_{\partial\Omega}$.) The coercivity of Ψ from Remark 4.5 and Remark 5.3 proves

$$e(u, v_0) \coloneqq \|\nabla(u - v_0)\|_4^4 + \|\sigma - \nabla\Psi(\nabla v_0)\|_{4/3}^2 + \|\sqrt{\varrho}\nabla(u - v_0)\|_2^2 \lesssim E_{\sigma_0}(v_0) - E_{\sigma_0}^*(\sigma_0) + \operatorname{osc}_4(f, v_0, \mathcal{M})^2 \eqqcolon \operatorname{RHS}$$

with $\rho := (|\nabla u| + |\nabla v_0|)^2$ and the dual energy

$$E_{\sigma_0}^*(\sigma_0) = -\int_{\Omega} \Psi^*(\sigma_0) \,\mathrm{d}x + \int_{\Gamma_D} u\sigma_0 \cdot \nu \,\mathrm{d}s$$

In view of the analysis in [32, 39] for lowest-order methods, we expect $h_{\max}^{2(k+1)}$ as the best possible convergence rate for the primal-dual gap $E_{\sigma_0}(v_0) - E_{\sigma_0}^*(\sigma_0)$. To achieve this, the stabilization s_h should – at least in theory – scale quadratically. This motivates the input r = 2 and s = 1 for s_h in this benchmark. Figure 4– 5 display a convergence rate between 0.45 and 2/3 for RHS, $\|\sqrt{\rho}\nabla(u-v_0)\|_{4/3}^2$, and $\|\sigma - \nabla\Psi(\nabla v_0)\|_{4/3}^2$, but only a marginal improvement of $\|\nabla(u-v_0)\|_4^2$ in the preasymtotic range on uniform meshes. Adaptive computation refines towards the singularity at the origin and recovers the optimal convergence rates k + 1 for RHS, $\|\sqrt{\rho}\nabla(u-v_0)\|_{2}^2$, and $\|\sigma - \nabla W(\nabla v_0)\|_{4/3}^2$ with all displayed k. For the displacement error $\|\nabla(u-v_0)\|_4^2$, we observe the empirical convergence rates (k+1)/2 for $k \geq$ 2. In undisplayed computer experiments with the input r = p = 4, the optimal convergence rates 2(k+1) is observed for all displayed errors of this subsection for the choice $s \geq (p-2)(k+1) + p - 1$. (Notice from (4.12) and (4.15) that s = (p-2)(k+1) + p - 1 is the unique parameter with $s_h(I_V u) + \gamma_h(I_W \sigma) \lesssim h_{\max}^{2(k+1)}$ for smooth u and σ).

6.5. Conclusions. In all computer experiments, adaptive computation leads to improved convergence rates for the a posteriori error estimator compared to uniform mesh refinements. Higher polynomial degrees provide additional improvement, which depends on the model problem. For the Bingham flow and the p-Laplace

N. T. TRAN

problem, we observe the optimal convergence rates $ndof^{-(k+1)}$. This appears to be sensitive to the choice of the parameters r and s in the stabilization s_h if $p \neq 2$.

References

- M. Abbas, A. Ern, and N. Pignet, Hybrid high-order methods for finite deformations of hyperelastic materials, Comput. Mech. 62 (2018), no. 4, 909–928.
- J. Alberty, C. Carstensen, and S. A. Funken, Remarks around 50 lines of Matlab: short finite element implementation, Numer. Algorithms 20 (1999), no. 2-3, 117–137.
- J. W. Barrett and W. B. Liu, Finite element approximation of the p-Laplacian, Math. Comp. 61 (1993), no. 204, 523–537.
- 4. S. Bartels, *Numerical methods for nonlinear partial differential equations*, Springer Series in Computational Mathematics, vol. 47, Springer, Cham, 2015.
- 5. _____, Error estimates for a class of discontinuous Galerkin methods for nonsmooth problems via convex duality relations, Math. Comp. **90** (2021), no. 332, 2579–2602.
- _____, Nonconforming discretizations of convex minimization problems and precise relations to mixed methods, Comput. Math. Appl. 93 (2021), 214–229.
- S. Bartels and C. Carstensen, A convergent adaptive finite element method for an optimal design problem, Numer. Math. 108 (2008), no. 3, 359–385.
- S. Bartels and A. Kaltenbach, Explicit and efficient error estimation for convex minimization problems, Math. Comp. 92 (2023), no. 343, 2247–2279.
- M. Bebendorf, A note on the Poincaré inequality for convex domains, Z. Anal. Anwendungen 22 (2003), no. 4, 751–756.
- Michele Botti, Daniele A. Di Pietro, and Pierre Sochala, A hybrid high-order method for nonlinear elasticity, SIAM J. Numer. Anal. 55 (2017), no. 6, 2687–2717. MR 3721565
- D. Braess and J. Schöberl, Equilibrated residual error estimator for edge elements, Math. Comp. 77 (2008), no. 262, 651–672.
- C. Carstensen, M. Feischl, M. Page, and D. Praetorius, Axioms of adaptivity, Comput. Math. Appl. 67 (2014), no. 6, 1195–1253.
- C. Carstensen and R. Klose, A posteriori finite element error control for the p-Laplace problem, SIAM J. Sci. Comput. 25 (2003), 792–814.
- C. Carstensen and D. J. Liu, Nonconforming FEMs for an optimal design problem, SIAM J. Numer. Anal. 53 (2015), no. 2, 874–894.
- Monconforming FEMs for an optimal design problem, SIAM J. Numer. Anal. 53 (2015), no. 2, 874–894.
- C. Carstensen and P. Plecháč, Numerical solution of the scalar double-well problem allowing microstructure, Math. Comp. 66 (1997), no. 219, 997–1026.
- C. Carstensen, B. D. Reddy, and M. Schedensack, A natural nonconforming FEM for the Bingham flow problem is quasi-optimal, Numer. Math. 133 (2016), no. 1, 37–66.
- C. Carstensen and N. T. Tran, Unstabilized Hybrid High-order Method for a Class of Degenerate Convex Minimization Problems, SIAM J. Numer. Anal. 59 (2021), no. 3, 1348–1373.
- <u>Convergent</u> adaptive hybrid higher-order schemes for convex minimization, Numer. Math. 151 (2022), no. 2, 329–367.
- S.-S. Chow, Finite element error estimates for non-linear elliptic equations of monotone type, Numer. Math. 54 (1989), 373–393.
- S.-K. Chua and R. L. Wheeden, Estimates of best constants for weighted Poincaré inequalities on convex domains, Proc. London Math. Soc. (3) 93 (2006), no. 1, 197–226.
- B. Cockburn, D. A. Di Pietro, and A. Ern, Bridging the hybrid high-order and hybridizable discontinuous Galerkin methods, ESAIM Math. Model. Numer. Anal. 50 (2016), no. 3, 635– 650.
- D. A. Di Pietro and J. Droniou, A hybrid high-order method for Leray-Lions elliptic equations on general meshes, Math. Comp. 86 (2017), no. 307, 2159–2191.
- 24. _____, The hybrid high-order method for polytopal meshes, MS&A. Modeling, Simulation and Applications, vol. 19, Springer, Cham, 2020, Design, analysis, and applications.
- D. A. Di Pietro, J. Droniou, and A. Harnist, Improved error estimates for hybrid high-order discretizations of Leray-Lions problems, Calcolo 58 (2021), no. 2, Paper No. 19, 24.
- 26. D. A. Di Pietro and A. Ern, Mathematical aspects of discontinuous Galerkin methods, Mathématiques & Applications, vol. 69, Springer, Heidelberg, 2012.
- _____, A hybrid high-order locking-free method for linear elasticity on general meshes, Comput. Methods Appl. Mech. Engrg. 283 (2015), 1–21.
- D. A. Di Pietro, A. Ern, and S. Lemaire, An arbitrary-order and compact-stencil discretization of diffusion on general meshes based on local reconstruction operators, Comput. Methods Appl. Math. 14 (2014), no. 4, 461–472.

- Daniele A. Di Pietro and Jérôme Droniou, An arbitrary-order discrete de Rham complex on polyhedral meshes: exactness, Poincaré inequalities, and consistency, Found. Comput. Math. 23 (2023), no. 1, 85–164. MR 4546145
- Daniele A. Di Pietro, Jérôme Droniou, and Francesca Rapetti, Fully discrete polynomial de Rham sequences of arbitrary degree on polygons and polyhedra, Math. Models Methods Appl. Sci. 30 (2020), no. 9, 1809–1855. MR 4151796
- L. Diening, D. Köner, M. Růžička, and I. Toulopoulos, A local discontinuous Galerkin approximation for systems with p-structure, IMA J. Numer. Anal. 34 (2014), no. 4, 1447–1488.
- L. Diening and C. Kreuzer, Linear convergence of an adaptive finite element method for the p-Laplacian equation, SIAM J. Numer. Anal. 46 (2008), no. 2, 614–638.
- L. Diening and M. Růžička, Interpolation operators in Orlicz-Sobolev spaces, Numer. Math. 107 (2007), no. 1, 107–129.
- A. Ern and J.-L. Guermond, *Finite elements I—Approximation and interpolation*, Texts in Applied Mathematics, vol. 72, Springer, Cham, 2021.
- 35. A. Ern and M. Vohralík, Polynomial-degree-robust a posteriori estimates in a unified setting for conforming, nonconforming, discontinuous Galerkin, and mixed discretizations, SIAM J. Numer. Anal. 53 (2015), no. 2, 1058–1081.
- Stable broken H¹ and H(div) polynomial extensions for polynomial-degree-robust potential and flux reconstruction in three space dimensions, Math. Comp. 89 (2020), no. 322, 551–594.
- 37. R. Glowinski, Numerical methods for nonlinear variational problems, Scientific Computation, Springer-Verlag, Berlin, 2008, Reprint of the 1984 original.
- 38. R. Glowinski and A. Marrocco, Sur l'approximation, par éléments finis d'ordre un, et la résolution, par pénalisation-dualité, d'une classe de problèmes de Dirichlet non linéaires, Rev. Française Automat. Informat. Recherche Opérationnelle Sér. Rouge Anal. Numér. 9 (1975), no. no. R-2, 41–76.
- A. Kaltenbach, Error analysis for a crouzeix-raviart approximation of the p-dirichlet problem, arXiv:2210.12116 (2022).
- R. V. Kohn and G. Strang, Optimal design and relaxation of variational problems. I, Comm. Pure Appl. Math. 39 (1986), no. 1, 113–137.
- L. D. Marini, An inexpensive method for the evaluation of the solution of the lowest order Raviart-Thomas mixed method, SIAM J. Numer. Anal. 22 (1985), no. 3, 493–496.
- S. I. Repin, A posteriori error estimation for variational problems with uniformly convex functionals, Math. Comp. 69 (2000), no. 230, 481–500.
- R. T. Rockafellar, *Convex analysis*, Princeton Mathematical Series, No. 28, Princeton University Press, Princeton, N.J., 1970.
- 44. V. B. Tyukhtin, The rate of convergence of approximation methods for solving one-sided variational problems. I, Teoret. Mat. Fiz. **51** (1982), no. 2, 111–113, 121.

(N. T. Tran) INSTITUT FÜR MATHEMATIK, UNIVERSITÄT AUGSBURG, 86159 AUGSBURG Email address: ngoc1.tranQuni-a.de